

Table 15.5 from (1976AJ04): Radiative decays in  $^{15}\text{N}$  <sup>a</sup>

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Mult. mixing ratio $\delta$	Refs.
5.27	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	100	$-0.131 \pm 0.013$	(1975BE23, 1976BE1B)
5.30	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	100		
6.32	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100	$+0.122 \pm 0.006$ <sup>b</sup>	(1975MO28, 1976BE1B)
		5.27	$\frac{5}{2}^+$	$< 1$		(1965WA16, 1975MO28)
		5.30	$\frac{1}{2}^+$	$< 0.5$		(1975MO28)
7.16	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	$< 0.1$		(1976BE1B)
		5.27	$\frac{5}{2}^+$	$100 \pm 0.4$ <sup>c</sup>	$-0.014^{+0.012}_{-0.015}$	(1976BE1B)
		5.30	$\frac{1}{2}^+$	$< 4$		(1966AL18, 1968GI11)
		6.32	$\frac{3}{2}^-$	$< 0.5$		(1965WA16)
7.30 <sup>c</sup>	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	$99.3 \pm 0.7$	$-0.017^{+0.005}_{-0.008}$	(1976BE1B)
		5.27	$\frac{5}{2}^+$	$0.6 \pm 0.1$	$+0.18 \pm 0.15$ , or $+2.5 \pm 1.0$	(1976BE1B)
		5.30	$\frac{1}{2}^+$	$0.2 \pm 0.1$	$-0.31 \pm 0.15$ , or $+4.6 \pm 3.4$	(1976BE1B)
		6.32	$\frac{3}{2}^-$	$< 0.25$		(1965WA16)
7.57	$\frac{7}{2}^+$	0	$\frac{1}{2}^-$	$1.3 \pm 0.6$		(1975BE23, 1976BE1B)
		5.27	$\frac{5}{2}^+$	$98.7 \pm 1.0$	$-0.028 \pm 0.012$	(1976BE1B)
		5.30	$\frac{1}{2}^+$	$< 4$		(1966AL18)
		6.32	$\frac{3}{2}^-$	$< 0.6$		(1965WA16)
8.31	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	$79 \pm 2$		(1965WA16, 1966WA08, 1967PH03)
		5.27	$\frac{5}{2}^+$	$< 3$		(1965WA16)
		5.30	$\frac{1}{2}^+$	$10 \pm 2$		(1965WA16)
		6.32	$\frac{3}{2}^-$	$7.8 \pm 2$		(1965WA16)
				$4.4 \pm 1.0$		(1967PH03)
		7.16	$\frac{5}{2}^+$	$1.2 \pm 0.6$		(1967PH03)
		7.30	$\frac{3}{2}^+$	$2.2 \pm 0.4$		(1965WA16)
				$4.4 \pm 0.7$		(1967PH03)
8.57	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	$33 \pm 2$	$-0.085^{+0.005}_{-0.009}$	(1965WA16, 1966WA08, 1967PH03)
		5.27	$\frac{5}{2}^+$	$65 \pm 3$		(1976BE1B)
					$-0.091 \pm 0.007$	(1966WA08)
		5.30	$\frac{1}{2}^+$	$< 12$		(1976BE1B)
		6.32	$\frac{3}{2}^-$	$3 \pm 1$		(1965WA16)
				$1.4 \pm 0.6$		(1967PH03)
		7.16	$\frac{5}{2}^+$	$3.6 \pm 0.5$		(1967PH03)
		7.30	$\frac{3}{2}^+$	$< 0.7$		(1965WA16)
		7.57	$\frac{7}{2}^+$	$< 3$		(1965WA16, 1966WA08)

Table 15.5 from (1976AJ04):Radiative decays in  $^{15}\text{N}^a$  (continued)

$E_i$ (MeV)	$J_1^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Mult. mixing ratio $\delta$	Refs.		
9.05	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	$92 \pm 3$		(1965WA16, 1966WA08)		
				$91.6 \pm 0.9$		(1967PH03)		
		5.27	$\frac{5}{2}^+$	$3.5 \pm 1$		(1966WA08)		
				$4.7 \pm 0.7$		(1967PH03)		
		6.32	$\frac{3}{2}^-$	$4.5 \pm 1$		(1966WA08)		
				$3.7 \pm 0.5$		(1967PH03)		
		7.16	$\frac{5}{2}^+$	$< 10$		(1965WA16)		
		7.30	$\frac{3}{2}^+$	$1.2 \pm 0.4$		(1965WA16)		
		$7.57$	$\frac{7}{2}^+$	$< 2$	(1965WA16)			
		$8.31$	$\frac{1}{2}^+$	$< 0.5$	(1965WA16)			
9.152	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	$100 \pm 3$	$+0.015^{+0.041}_{-0.034}$	(1969SI04, 1976BE1B) <sup>d</sup>		
						(1976BE1B)		
9.155	$\frac{5}{2}$	0	$\frac{1}{2}^-$	$9 \pm 9$		(1968ST10)		
		5.27	$\frac{3}{2}^+$	$\approx 8$		(1967TH05)		
		5.30	$\frac{1}{2}^+$	$\approx 10$		(1967TH05)		
		6.32	$\frac{3}{2}^-$	$20 \pm 2$		(1967TH05, 1968ST10, 1969SI04)		
		7.16	$\frac{5}{2}^+$	$\approx 50$		(1967TH05, 1968ST10, 1969SI04)		
		$7.30$	$\frac{3}{2}^+$	$8 \pm 1$	(1968ST10)			
9.23	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	$< 30$		(1965WA16)		
				$41.5 \pm 2.2$		(1967PH03)		
		5.27	$\frac{5}{2}^+$	$< 25$		(1965WA16)		
		5.30	$\frac{1}{2}^+$	100		(1965WA16)		
		5.27 + 5.30		$31.2 \pm 1.7$		(1967PH03)		
		6.32	$\frac{3}{2}^-$	$\leq 25$		(1965WA16)		
				$24.7 \pm 1.5$		(1967PH03)		
		7.16	$\frac{5}{2}^+$	$< 30$		(1965WA16)		
				$< 1$		(1967PH03)		
		7.30	$\frac{3}{2}^+$	$< 30$		(1965WA16)		
		$7.57$	$\frac{7}{2}^+$	$2.6 \pm 0.7$	(1967PH03)			
		$8.31$	$\frac{1}{2}^+$	$< 20$	(1965WA16)			
		$8.31$	$\frac{1}{2}^+$	$< 5$	(1965WA16)			
9.76 <sup>e</sup>	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	$81.5 \pm 2.8$		(1967PH03)		
		5.27 + 5.30		$7.5 \pm 1.5$		(1967PH03)		
		6.32	$\frac{3}{2}^-$	$3.7 \pm 0.8$		(1967PH03)		
		7.16	$\frac{5}{2}^+$	$2.3 \pm 0.5$		(1967PH03)		
		7.30	$\frac{3}{2}^+$	$< 2$		(1967PH03)		
				$7.57$		$\frac{7}{2}^+$	$5.0 \pm 0.6$	(1967PH03)
								(1967PH03)

Table 15.5 from (1976AJ04):Radiative decays in  $^{15}\text{N}^a$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Mult. mixing ratio $\delta$	Refs.		
9.83	$\frac{7}{2}$	8.31	$\frac{1}{2}^+$	< 1		(1967PH03)		
		8.57	$\frac{3}{2}^+$	< 2		(1965WA16, 1967PH03)		
		0	$\frac{1}{2}^-$	< 4		(1967PH03)		
		5.27	$\frac{3}{2}^+$	$\approx 85$		(1965WA16, 1967PH03)		
		5.30	$\frac{1}{2}^+$	< 15		(1965WA16)		
		6.32	$\frac{3}{2}^-$	$2.2 \pm 0.9$		(1967PH03)		
		7.16	$\frac{3}{2}^+$	$2.4 \pm 1.1$		(1967PH03)		
		7.30	$\frac{3}{2}^+$	$3.7 \pm 0.9$		(1967PH03)		
		7.57	$\frac{7}{2}^+$	$7.3 \pm 1.0$		(1967PH03)		
		9.93 <sup>e</sup>	$(\frac{1}{2}, \frac{3}{2})^+$	0	$\frac{1}{2}^-$	$77.6 \pm 1.9$		(1967PH03)
5.27 + 5.30				$15.4 \pm 1.5$		(1967PH03)		
6.32	$\frac{3}{2}^-$			$4.9 \pm 1.2$		(1967PH03)		
7.16	$\frac{3}{2}^+$			< 1		(1967PH03)		
7.30	$\frac{3}{2}^+$			$2.1 \pm 0.8$		(1967PH03)		
7.57	$\frac{7}{2}^+$			< 1		(1967PH03)		
8.31	$\frac{1}{2}^+$			< 1		(1967PH03)		
8.57	$\frac{3}{2}^+$			< 1		(1967PH03)		
10.07 <sup>e</sup>	$\frac{3}{2}^+$			0	$\frac{1}{2}^-$	$96.0 \pm 0.7$		(1967PH03)
				5.27 + 5.30		$4.0 \pm 0.7$		(1967PH03)
		6.32, 7.16, 7.30, 7.57		< 2		(1966WA08)		
		8.31	$\frac{1}{2}^+$	< 2		(1965WA16)		
		8.57	$\frac{3}{2}^+$	< 3		(1965WA16)		
		10.45	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	< 12		(1976BE1B)
				5.27	$\frac{3}{2}^+$	$55.0 \pm 0.8$	$+0.021 \pm 0.033$	(1976BE1B)
				5.30	$\frac{1}{2}^+$	< 2		(1976BE1B)
				6.32	$\frac{3}{2}^-$	$31.3 \pm 1.7$	$-0.59 \pm 0.13$	(1976BE1B)
				7.16	$\frac{3}{2}^+$	$5.2 \pm 0.1$	$+0.13^{+0.03}_{-0.04}$	(1976BE1B)
8.57	$\frac{3}{2}^+$			$3.8 \pm 0.6$	$-0.3 \pm 0.4$	(1976BE1B)		
9.152	$\frac{3}{2}^-$			$4.7 \pm 0.1$	$-0.32^{+0.09}_{-0.10}$	(1976BE1B)		
9.83	$\frac{7}{2}^-$			< 0.1		(1976BE1B)		
10.53	$\frac{5}{2}^+$			0	$\frac{1}{2}^-$	< 0.1		(1976BE1B)
				5.27	$\frac{3}{2}^+$	$38.7 \pm 0.2$	$-0.27 \pm 0.03$	(1976BE1B)
		6.32	$\frac{3}{2}^-$	$7.7 \pm 0.1$	$-0.028 \pm 0.004$	(1976BE1B)		
		7.16	$\frac{3}{2}^+$	$19.4 \pm 0.2$	$+0.007^{+0.010}_{-0.008}$	(1976BE1B)		
		7.30	$\frac{3}{2}^+$	$31.4 \pm 0.5$	$+0.066 \pm 0.005$	(1976BE1B)		
		8.57	$\frac{3}{2}^+$	$2.4 \pm 0.1$	$+0.012^{+0.006}_{-0.005}$	(1976BE1B)		

Table 15.5 from (1976AJ04):Radiative decays in  $^{15}\text{N}$  <sup>a</sup> (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Mult. mixing ratio $\delta$	Refs.
10.69	$\frac{9}{2}^+$	9.152	$\frac{3}{2}^-$	$0.3 \pm 0.1$	$-0.20^{+0.03}_{-0.02}$	(1976BE1B)
		5.27	$\frac{5}{2}^+$	$61.6 \pm 0.3$	$\equiv 0$	(1975BE23, 1976BE1B)
		7.16	$\frac{5}{2}^+$	$2.1 \pm 0.1$	$-0.03 \pm 0.07$	(1975BE23, 1976BE1B)
		7.57	$\frac{7}{2}^+$	$36.3 \pm 0.6$	$+0.118 \pm 0.008$	(1975BE23, 1976BE1B)
10.70 <sup>f</sup>	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	$52.6 \pm 0.8$	$+0.180^{+0.006}_{-0.002}$	(1976BE1B)
		5.27	$\frac{3}{2}^+$	$37.4 \pm 0.6$	$-0.024^{+0.004}_{-0.008}$	(1976BE1B)
		5.30	$\frac{1}{2}^+$	$0.8 \pm 0.1$	$-0.13 \pm 0.07$	(1976BE1B)
		6.32	$\frac{3}{2}^-$	$3.8 \pm 0.1$	$+0.135 \pm 0.015$	(1976BE1B)
		7.16	$\frac{3}{2}^+$	$0.4 \pm 0.1$	$0.3 \pm 0.3$	(1976BE1B)
		7.30	$\frac{3}{2}^+$	$2.3 \pm 0.1$	$-0.027 \pm 0.023$	(1976BE1B)
		8.31	$\frac{1}{2}^+$	$0.8 \pm 0.1$	$-0.017^{+0.018}_{-0.016}$	(1976BE1B)
		9.05	$\frac{1}{2}^+$	$0.2 \pm 0.1$	$-0.007 \pm 0.12$	(1976BE1B)
		9.152	$\frac{3}{2}^-$	$0.2 \pm 0.1$	$-0.11 \pm 0.03$	(1976BE1B)
		9.23	$\frac{1}{2}^-$	$1.5 \pm 0.1$	$+0.049^{+0.006}_{-0.005}$	(1976BE1B)
		10.80 <sup>g</sup>	$\frac{3}{2}^{(+)}$	0	$\frac{1}{2}^-$	$51.5 \pm 0.4$
5.27	$\frac{3}{2}^+$			$4.9 \pm 0.1$	$-0.63 \pm 0.04$ <sup>f</sup>	(1976BE1B)
5.30	$\frac{1}{2}^+$			$15.5 \pm 0.2$	$-0.55 \pm 0.02$ <sup>f</sup>	(1976BE1B)
6.32	$\frac{3}{2}^-$			$5.4 \pm 0.2$	$-0.07 \pm 0.05$ <sup>f</sup>	(1976BE1B)
7.16	$\frac{3}{2}^+$			$7.8 \pm 0.1$	$+0.14 \pm 0.03$ <sup>f</sup>	(1976BE1B)
7.30	$\frac{3}{2}^+$			$5.8 \pm 0.1$	$-0.12 \pm 0.02$ <sup>f</sup>	(1976BE1B)
8.31	$\frac{1}{2}^+$			$3.6 \pm 0.1$	$+0.12 \pm 0.03$ <sup>f</sup>	(1976BE1B)
9.05	$\frac{1}{2}^+$			$0.3 \pm 0.1$		(1976BE1B)
9.152	$\frac{3}{2}^-$			$0.9 \pm 0.1$		(1976BE1B)
9.155	$\frac{3}{2}^-$			$4.2 \pm 0.1$		(1976BE1B)
11.62 <sup>h</sup>	$\frac{1}{2}^+; T = \frac{3}{2}$			0	$\frac{1}{2}^-$	$90.7 \pm 3.0$
		5.27	$\frac{3}{2}^+$	$< 1$		(1971KU01)
		5.30	$\frac{1}{2}^+$	$7.4 \pm 1.5$		(1971KU01)
		6.32	$\frac{3}{2}^-$	$1.9 \pm 1.5$		(1971KU01)
12.52	$\frac{5}{2}^+; T = \frac{3}{2}$	0	$\frac{1}{2}^-$	$< 1$		(1971KU01)
		5.27	$\frac{3}{2}^+$	$94.2 \pm 0.6$ <sup>i</sup>		(1971YO03)
		5.30	$\frac{1}{2}^+$	$< 1$		(1971KU01)
13.42 <sup>k</sup>	$\frac{3}{2}^+$	6.32	$\frac{3}{2}^-$	$5.8 \pm 0.6$ <sup>j</sup>		(1971YO03)
		0	$\frac{1}{2}^-$	100		(1976KU01)

- <sup>a</sup> See also [Table 15.10 in \(1970AJ04\)](#).
- <sup>b</sup>  $\Gamma_{\gamma_0} = 3.1 \pm 0.3$  eV,  $\delta(\text{E2/M1}) = 0.137 \pm 0.005$  ([1975MO28](#)); see also [Table 15.16](#).
- <sup>c</sup> See also ([1965WA16](#), [1966PE04](#), [1968GI11](#)).
- <sup>d</sup> See also ([1968ST06](#)) and [reaction 44](#).
- <sup>e</sup> See also ([1965WA16](#)).
- <sup>f</sup> See ([1969SI04](#)).
- <sup>g</sup> See also ([1965WA16](#), [1966WA08](#)).
- <sup>h</sup>  $\Gamma_{\gamma} = 49 \pm 20, 4 \pm 2, 1.0 \pm 0.8$  for transitions to  $^{15}\text{N}^*(0, 5.30, 6.32)$  ([1971KU01](#)); see also [Table 15.12](#).
- <sup>i</sup>  $\Gamma_{\gamma} = 4.3 \pm 0.7$  eV ([1971YO03](#));  $\delta = -0.02 \pm 0.04$  (E2/M1) ([1971KU01](#)).
- <sup>j</sup>  $\Gamma_{\gamma} = 0.27 \pm 0.05$  eV ([1971YO03](#));  $\delta = -0.02 \pm 0.04$  (E2/M1) ([1971KU01](#)).
- <sup>k</sup>  $\Gamma_{\gamma_0} = 3.0 \pm 0.9$  eV,  $\Gamma_p \Gamma_{\gamma_0} / \Gamma = 1.70 \pm 0.5$  eV;  $\delta = 0.00 \pm 0.04$  (M2/E1);  $B(\text{E1}) = 1.2 \pm 0.4 \times 10^{-3} e^2 \cdot \text{fm}^2$ . Transitions to  $^{15}\text{N}^*(5.27, 5.30) < 8\%$  and to  $^{15}\text{N}^*(6.32, 7.16, 7.30) < 5\%$  ([1976KU01](#)). See also ([1975HA39](#)).