

Table 15.10 from (1976AJ04): Resonances in $^{13}\text{C} + \text{d}$

E_d (MeV)	Particles out	Γ_{lab} (keV)	$^{15}\text{N}^*$ (MeV)	Refs.
0.37	p		16.48	(1956VA17)
0.64	n, p ₀ , t ₀	≈ 100	16.71	(1950CU13, 1950RI57, 1953KO42, 1956MA46, 1956VA17, 1971PU01)
0.85	n, p ₀	≈ 400	16.90	(1950RI57, 1971PU01)
1.10	α_0	broad	17.11	(1956MA35, 1966KL06, 1966KL1A)
1.24 ± 0.04	t ₀ , (α_0)	≈ 200	17.23	(1956MA46, 1966KL1A)
1.40 ± 0.04	p ₀ , t ₀ , α_0	≈ 400	17.37	(1956MA35, 1956MA46, 1966KL1A)
1.55 ^a	n	≈ 100		(1941BE1A, 1950RI57, 1966KL06)
1.64 ± 0.04	t ₀	≈ 200	17.58	(1956MA35)
1.75 ± 0.04	γ_0 , n, α_0	≈ 600	17.67	(1950RI57, 1955MA76, 1956MA35, 1973WE12)
1.80 ± 0.01	(p ₀), t ₀ , α_1	55 ± 10	17.72	(1956MA35, 1956MA46)
2.20 ± 0.01	(n), α_0 , α_1	22 ± 4	18.06	(1956MA35, 1963AL21)
2.23 ± 0.02	(n), p ₀ , t	≈ 50	18.09	(1956MA35, 1956MA46, 1963AL21)
2.45 ± 0.03	n, p ₀ , α_0	270 ± 70	18.28	(1955MA76, 1956MA35, 1956MA46)
3.46 ± 0.03	n	≈ 150	19.16	(1955MA76, 1963DE19)
5.1	n ₁ , p ₀	≈ 50	20.6	(1973BO10)
6.63	γ_0		21.90	(1975DE1M)

^a Possibly to be identified with 1.40 MeV resonance (1956MA35).