

Table 6. 10 from (2002TI10): Levels of  ${}^6\text{Li}$  from  ${}^6\text{Li}(e, e')$  and  ${}^6\text{Li}(\gamma, \gamma')$  <sup>a</sup>

$E_x$ (MeV)	$J^\pi; T$	$\Gamma_{\gamma_0}$ (eV)	Multipolarity
$2.183 \pm 0.009$	$3^+; 0$	$(4.40 \pm 0.34) \times 10^{-4}$ <sup>b</sup>	E2
$3.56288 \pm 0.00010$ <sup>c</sup>	$0^+; 1$	$8.19 \pm 0.17$ <sup>d</sup>	M1
$4.27 \pm 0.04$	$2^+; 0$	$(5.4 \pm 2.8) \times 10^{-3}$	E2
$5.379 \pm 17$ <sup>d, e</sup>	$2^+; 1$	$0.27 \pm 0.05$	M1

<sup>a</sup> See Tables 6.4 in (1979AJ01, 1984AJ01) for references and for the earlier work.

<sup>b</sup> (1969EI06),  $B(E2)\uparrow = 25.6 \pm 2.0 e^2 \cdot \text{fm}^4$ . The value given in (1988AJ01) was incorrect.

We are grateful to Dr. John Millener for pointing out this error.

<sup>c</sup> (1981RO02).

<sup>d</sup> Weighted mean of values shown in Table 6.4 in (1979AJ01).

<sup>e</sup>  $\Gamma = 540 \pm 20$  keV.