

Energy Levels of Light Nuclei $A = 5$

F. Ajzenberg-Selove^a and T. Lauritsen^b

^a *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

^b *California Institute of Technology, Pasadena, California*

Abstract: An evaluation of $A = 5-24$ was published in *Nuclear Physics* 11 (1959), p. 1. This version of $A = 5$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 01, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 5$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^5\text{H}\$](#) , [\$^5\text{He}\$](#) , [\$^5\text{Li}\$](#)

B. Tables of Recommended Level Energies:

[Table 5.1](#): Energy levels of ^5He

[Table 5.3](#): Energy levels of ^5Li

C. [References](#)

D. Figures: [\$^5\text{He}\$](#) , [\$^5\text{Li}\$](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

${}^5\text{H}$

(Not illustrated)

The possible existence of a particle-stable ${}^5\text{H}$ is discussed by (1957BL1A) who point out that a $T = \frac{3}{2}$ level of ${}^5\text{He}$ - ${}^5\text{Li}$ might plausibly be formed by combination of ${}^3\text{H}$ or ${}^3\text{He}$ and a deuteron in the singlet ($T = 1$) state at an energy ≈ 2.3 MeV higher than the known 16.7 – 16.8 MeV level. If such a level exists, calculation of Coulomb corrections and n - ${}^1\text{H}$ mass difference suggests a mass excess of 32.22 MeV for ${}^5\text{H}$, which would be 0.35 MeV stable against ${}^3\text{H} + 2n$. Presumably ${}^5\text{H}$ would then decay by β -emission (≈ 19 MeV) followed by neutron emission (1957BL1A). A search for delayed neutrons from ${}^7\text{Li}(\gamma, 2p){}^5\text{H}$ with 320-MeV bremsstrahlung yielded no evidence of formation of ${}^5\text{H}$ (1958TA1A). It is concluded that less than 1% of the expected yield of ${}^7\text{Li}(\gamma, 2p){}^5\text{H}$ leads to a particle-stable product (1958TA1A); see also (1958CE1A). A reaction yielding ${}^5\text{H}$ might be ${}^3\text{H}(t, p){}^5\text{H}$ with $Q \approx -8.1$ MeV, assuming the mass of ${}^5\text{H}$ given by (1957BL1A).

${}^5\text{He}$

(Fig. 1)

1. ${}^3\text{H}(d, \gamma){}^5\text{He}$ $Q_m = 16.629$

At $E_d = 160$ keV, the capture cross section is less than 0.5 mb. This limit is not inconsistent with $\Gamma_\gamma \approx 11$ eV as estimated from the mirror reaction ${}^3\text{He}(d, \gamma){}^5\text{Li}$ (1955SA52).

2. (a) ${}^3\text{H}(d, n){}^4\text{He}$ $Q_m = 17.586$ $E_b = 16.629$
(b) ${}^3\text{H}(d, 2n){}^3\text{He}$ $Q_m = -2.991$
(c) ${}^3\text{H}(d, pn){}^3\text{H}$ $Q_m = -2.226$
 $Q_0 = 17.580 \pm 0.025$ (1957MA1C).

Excitation curves and angular distributions for reaction (a) from $E_d = 8$ keV to 10 MeV are summarized by (1956FO1A, 1957JA37). Additional data are given for $E_d = 0.04$ to 0.73 MeV by (1957BA1F, 1957BA1G), for $E_d = 1.0$ to 5.8 MeV by (1956GA51) and for $E_d = 0.25$ to 7.0 MeV by (1957BA21). Below $E_d = 100$ keV, the cross section follows the Gamow function, $\sigma = (A/E)\exp(-44.40E^{-1/2})$ (1953JA1A, 1954AR02). A strong resonance, $\sigma(\text{peak}) = 5.0$ b, appears at $E_d = 107$ keV. A precision measurement ($\pm 5\%$) of the cross section at $E_t = 1.50$ MeV gives 20.0 and 19.6 mb/sr (c.m.) at 30° and 60° (lab), respectively (1958AL05). There is some question as to whether a broad maximum exists between 4 and 8 MeV (1956GA51, 1957BA21).

In the region $E_d = 10$ to 500 keV, the cross section is closely fitted with the assumption of s-wave formation of a $J = \frac{3}{2}^+$ state, with the parameters given in Table 5.2. For a given interaction

Table 5.1: Energy levels of ${}^5\text{He}$

E_x in ${}^5\text{He}$ (MeV)	J^π	Γ (MeV)	Decay	Reactions
0	$\frac{3}{2}^-$	0.55 ± 0.03	n, α	5, 6, 8, 9, 10, 11, 12, 13, 14, 16, 17
3 – 6	$\frac{1}{2}^-$	3 – 5	n, α	8, 14
16.69	$\frac{3}{2}^+$	0.08	n, d, t, α	2, 8
(≈ 20)		> 1	n, d, t, α	2

radius, two sets of parameters are obtained, depending upon whether Γ_n/Γ_d is assumed > 1 or < 1 (1952AR30, 1952CO35). Agreement with the mirror reaction, ${}^3\text{He}(d, p){}^4\text{He}$, is obtained with the second choice (1955KU03). The fact that the proton width is relatively small suggests that this level arises from excitation of the ${}^4\text{He}$ core. See also (1951FL1A, 1954JO1C, 1955JO1A, 1957DA1B, 1958LE1A) and (1955HA1B; theor.).

The angular distribution of neutrons is isotropic at and below resonance, and shows increasing forward peaking at higher energies (1957JA37). Angular distributions at $E_d = 6$ MeV are almost identical to those of ${}^3\text{H}(d, p){}^4\text{H}$ (1957BR23). At $E_d = 10$ MeV, the distribution is dominated by the stripping process, with $l_p = 0$ (1951BU1B). Again, close correspondence is found with the mirror reaction. See also (1955HE89, 1956BA1D, 1958LE1A, 1958PA09).

At $E_d = 12$ to 14 MeV, reactions (b) and (c) are observed (1956BO44, 1958BR14). The ${}^3\text{He}$ distributions from (b) show no evidence for a bound dineutron or for a well-defined virtual state, although some interaction between the neutrons does appear to occur. Absolute differential cross sections are reported (1958BR14).

For $E_d > 3.71$ MeV, deuteron breakup (reaction (c)) is energetically possible. The cross section for this process has been studied for $E_d = 3.8$ to 6.0 MeV by (1955HE90). At $E_d = 14$ MeV, the number of low-energy (4 to 10 MeV) neutrons is about three times as large as observed in the corresponding reaction ${}^3\text{He}(d, pn){}^3\text{He}$. The difference may indicate formation of a $T = 0$, 22-MeV excited state of ${}^4\text{He}$ via ${}^3\text{H}(d, n){}^4\text{He}^*$ (1956BO1F, 1956BO43, 1956BO44: see, however, (1958BR14)). See also (1954CO1B, 1955BA1G, 1955BE1B, 1956FO1A, 1957BL1A) and (1955LA1B, 1956BL1A, 1958PO1A; theor.).

3. ${}^3\text{H}(d, p){}^4\text{H}$

Assuming the atomic mass excess of ${}^4\text{H} = 26$ MeV (if first $T = 1$ state in ${}^4\text{He}$ has $E_x = 22$ MeV), Q_m for this reaction would be -4 MeV. This reaction has not been observed (1951MC37). An attempt has also been made by (1955RE44) to observe the β -decay of ${}^4\text{H}$ formed in the 300-MeV proton bombardment of ${}^{12}\text{C}$. The results are negative: if $\tau_{1/2} = (2 \text{ to } 4) \times 10^{-3}$ sec, $\sigma < 1 \mu\text{b}$; if $\tau_{1/2} = (4 \text{ to } 10) \times 10^{-3}$ sec, $\sigma < 10 \mu\text{b}$. See also (1957NO17: ${}^{10}\text{B}({}^7\text{Li}, {}^4\text{H}){}^{13}\text{N}$).

Table 5.2: Resonance parameters for ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$ and ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$

E_{r} (keV)	Γ_{lab} (keV)	l_{d}	J^{π}	$l_{\text{n,p}}$	R ($\times 10^{-13}$ cm)	E_{λ} (keV)	γ_{d}^2 (keV)	$\gamma_{\text{n,p}}^2$ (keV)	θ_{d}^2 ^c	$\theta_{\text{n,p}}^2$ ^c	E_{x} (MeV)
107 ^a	135	0	$\frac{3}{2}^{+}$	2	5.0	-464	2000	56	1.0	0.018	16.69
					7.0	-126	715	17	0.7	0.011	
430 ^b	≈ 450	0	$\frac{3}{2}^{+}$	2	5.0	-391	2930	42	1.4	0.013	16.81
					7.0	129	780	12	0.7	0.008	

^a ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$: (1952AR30, 1952CO35, 1955KU03). See also (1957BA1F, 1957BA1G).

^b ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$: (1955KU03).

^c Units of $3\hbar^2/2MR^2$.

4. ${}^3\text{H}(\text{d}, \text{d}){}^3\text{H}$

$$E_{\text{b}} = 16.629$$

Differential cross sections for $E_{\text{d}} = 0.96$ to 3.2 MeV are tabulated by (1952ST69) and for $E_{\text{d}} = 10$ MeV by (1952AL36); see also (1957BR23, 1957JA37). The distributions are closely similar to those for ${}^3\text{He}(\text{d}, \text{d}){}^3\text{He}$. See also (1958BA82).

5. ${}^3\text{H}(\text{t}, \text{n}){}^5\text{He}$

$$Q_{\text{m}} = 10.371$$

The ground state of ${}^5\text{He}$ has been observed at $E_{\text{t}} = 1.48$ MeV (1957BA10) and 1.9 MeV (1958JA06). The neutron spectrum contains an excess of medium-energy neutrons, attributed to direct three-body reaction or to a broad excited state of ${}^5\text{He}$. An earlier reported peak corresponding to a 2.6 -MeV excited state (1951LE1A) is not confirmed (1957BA10). The alphas show a double peaking, reflecting the influence of the $\text{P}_{3/2}$ ground state, superposed on a distribution arising from the $\text{P}_{1/2}$ state and direct three-body disintegration (1958JA06); see ${}^6\text{He}$.

6. ${}^3\text{H}({}^3\text{He}, \text{p}){}^5\text{He}$

$$Q_{\text{m}} = 11.136$$

$$Q_0 = 11.18 \pm 0.07 \text{ (1953AL1A);}$$

$$Q_0 = 11.13 \pm 0.07 \text{ (1953MO61).}$$

The spectrum shows a well-defined proton peak corresponding to the ground state, superposed on a background attributed to the three-body breakup (1953AL1A, 1953MO61). See also (1954GO18).

7. ${}^3\text{H}(\alpha, \text{d}){}^5\text{He}$ $Q_{\text{m}} = -7.215$

Not observed.

8. ${}^4\text{He}(\text{n}, \text{n}){}^4\text{He}$ $E_{\text{b}} = -0.957$

Total cross sections for $E_{\text{n}} = 0.0004$ eV to 20 MeV are given in (1955HU1B, 1957HU1D, 1958HU18); angular distributions of (1952AD09) and (1953SE29) are given in (1956HU1A); additional data for $E_{\text{n}} = 2.6$ to 4.1 MeV are given by (1957ST1B), for $E_{\text{n}} = 14$ MeV by (1954SM97, 1955SH1D) and at $E_{\text{n}} = 15.7$ MeV by (1954AL28). The current experimental and theoretical situation is surveyed by (1958HO1B). The total cross section has a peak of 7.4 b at $E_{\text{n}} = 1.15 \pm 0.05$ MeV, $E_{\text{c.m.}} = 0.95$, with a width of ≈ 1.7 MeV (1951BA85), $\Gamma_{\text{c.m.}} = 1.4 \pm 0.2$ MeV (1958HU18). The thermal cross section is 0.78 b (1951HI1A), 0.71 ± 0.1 b (1951MC63), 0.74 ± 0.04 b (1955SO56).

Both the total cross sections and the angular distributions are well accounted for by the phase shifts determined by (1949CR1A, 1952DO30) for ${}^4\text{He}(\text{p}, \text{p}){}^4\text{He}$ with a shift in E_{λ} of about 1 MeV: see also (1955CL1A; theor.). Earlier discrepancies in the range 3 to 4 MeV (1952HU1A) appear to have been resolved by (1957ST1B). In a polarization measurement, (1957LE1B, 1957LE1C) find $\delta(\text{P}_{1/2}) = 12 \pm 1^\circ$ at $E_{\text{n}} = 2.45$ MeV, in disagreement with the value $\delta = 20^\circ$ derived from ${}^4\text{He}(\text{p}, \text{p}){}^4\text{He}$, but agreeing with a low cross section point at $E_{\text{n}} = 2.61$ MeV reported by (1953SE29). The s-wave phase shift decreases monotonically with increasing energy, and can be accounted for by hard-sphere scattering, with $R = 2.9 \times 10^{-13}$ cm (1952AD09, 1952DO30: see, however, (1954BR1B, 1954HO1B, 1955HO1C, 1956VA1C)). The $\text{P}_{3/2}$ shift shows strong resonance behavior near 1 MeV, while the $\text{P}_{1/2}$ shift changes more slowly, possibly indicating a broad $\text{P}_{1/2}$ level at several-MeV excitation (1952DO30). At $E_{\text{n}} = 15.7$ MeV the angular distribution is best accounted for with $\delta(\text{D}_{3/2}) = -14^\circ$, $\delta(\text{D}_{5/2}) = -7^\circ$, the latter being somewhat less than the hard-sphere value, suggesting a higher resonance (1954AL28). Theory: see ${}^5\text{Li}$. (For present purpose, the ground state of ${}^5\text{He}$ is assumed to correspond to the maximum in the total cross section; $E_{\text{c.m.}} = 0.95$ MeV).

A resonance is reported at $E_{\text{n}} = 22.15 \pm 0.13$ MeV, $\Gamma \leq 120$ keV, corresponding to the 16.7-MeV level (see ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$) (1957BO14). See also (1957IN1A).

Polarization of neutrons scattered in He has been discussed by (1952AD09, 1953SE29, 1953SI1A, 1955LE1D, 1956LE1B, 1957LE1B, 1957LE1C, 1957WH1A, 1958PA09) and others.

9. (a) ${}^4\text{He}(\text{d}, \text{p}){}^5\text{He}$ $Q_{\text{m}} = -3.184$

(b) ${}^4\text{He}(\text{d}, \text{pn}){}^4\text{He}$ $Q_{\text{m}} = -2.226$

$Q = -3.10 \pm 0.05$ (1954FR22) for reaction (a).

The proton spectrum observed at $E_d = 14.8$ MeV, $\theta = 19.5^\circ$, shows a prominent peak, of width $\Gamma_{c.m.} = 550 \pm 30$ keV, and a monotonic continuum of lower energy protons, attributed to reaction (b). There is no evidence of structure corresponding to possible sharp excited states of ${}^5\text{He}$. Cross sections for the ground-state group are $d\sigma/d\Omega = 25 \pm 5$ mb/sr at $\theta_{c.m.} = 18^\circ$ and 15 ± 4 mb/sr at $\theta_{c.m.} = 24^\circ$. The dimensionless reduced width of the ground-state group, analyzed by stripping theory is $\theta^2 = 0.05$, more than a factor of 10 smaller than is indicated by ${}^4\text{He}(n, n){}^4\text{He}$ (see ${}^4\text{He}(p, p){}^4\text{He}$ in ${}^5\text{Li}$) (1956WA1B, 1957WA01). For reaction (b), see ${}^6\text{Li}$.

$$10. {}^6\text{Li}(\gamma, p){}^5\text{He} \quad Q_m = -4.655$$

The threshold is $E_\gamma = 4.64 \pm 0.08$ MeV (1958RY77); see ${}^6\text{Li}$.

$$11. {}^6\text{Li}(n, d){}^5\text{He} \quad Q_m = -2.428$$

At $E_n = 14$ MeV, a well-defined ground-state group ($\Gamma_{c.m.} = 0.8$ MeV) is observed, as is a continuum extending to $E_x \approx 4$ MeV in ${}^5\text{He}$. Angular distributions of the ground-state group and of the continuum indicate $l_p = 1$ for the ground state transition and are not inconsistent with $l_p = 1$ for the continuum (1954FR03): see ${}^6\text{Li}$ and reaction 8 in ${}^7\text{Li}$.

$$12. {}^6\text{Li}(d, {}^3\text{He}){}^5\text{He} \quad Q_m = 0.839$$

At $E_d = 14.5$ MeV, the ground-state group is observed: $Q_0 = 0.91 \pm 0.09$ MeV, $\Gamma_{c.m.} = 0.69 \pm 0.2$ MeV (1955LE24).

$$13. {}^6\text{Li}(t, \alpha){}^5\text{He} \quad Q_m = 15.158$$

$$Q_0 = 15.15 \pm 0.04 \text{ (1956CR47).}$$

The width of the ground state $\Gamma_{c.m.} = 0.7 \pm 0.2$ MeV (1956CR47). See also (1954AL35, 1956BA1E, 1956MA09).

$$14. {}^7\text{Li}(n, t){}^5\text{He} \quad Q_m = -3.423$$

The angular distribution exhibits a forward maximum at $E_n = 14$ MeV. The total cross section is 55 ± 8 mb (1954FR03). At $E_n = 14$ MeV, events corresponding to transitions to the ground state and possibly to a level at 2.4 ± 0.6 MeV are observed in Li-loaded photoplates; the latter group may actually be due to a level at 9.25 MeV in ${}^7\text{Li}$ (1954AL24): see ${}^7\text{Li}(n, n'){}^7\text{Li}^*$.

15. ${}^7\text{Li}(p, {}^3\text{He}){}^5\text{He}$ $Q_m = -4.188$

Not reported.

16. ${}^7\text{Li}(d, \alpha){}^5\text{He}$ $Q_m = 14.163$
 $Q_0 = 14.26 \pm 0.09$ (1955LE1C);
 $Q_0 = 13.719 \pm 0.02$ (1955KH31, 1955KH35);
 $Q_0 = 14.15 \pm 0.22$ (1957FA10);
 $Q_0 = 14.11 \pm 0.08$ (1958WE27).

The angular correlation of ground state alpha particles and those resulting from the break up of ${}^5\text{He}$ is $W(\theta) = 1 + 1.2 \sin^2 \theta$, excluding a $J = \frac{1}{2}$ assignment, but consistent with $J = \frac{3}{2}$ (1951FR1A). (1956RI37) reports $W(\theta) = \sin^2 \theta$, also consistent with $J = \frac{3}{2}^-$. The $(\alpha - n)$ correlation observed at $E_d = 0.16$ MeV, yields $W(\theta) = 1 + (0.75 \pm 0.05) \sin^2 \theta$ and is again consistent with the assignment $J = \frac{3}{2}^-$ for the ground state of ${}^5\text{He}$ (1957FA10). The coincidence α -spectrum agrees in shape with a computed spectrum based on the ${}^4\text{He}(n, n){}^4\text{He}$ cross sections (1956RI37).

The width of the ground state of ${}^5\text{He}$ is 0.3 ± 0.1 MeV (1953CU20), 0.66 ± 0.2 MeV (1955LE24). The work of (1953CU20, 1956JU1B) at $E_d = 0.6$ to 1.5 MeV appears to indicate an excited state at $E_x = 2.5 \pm 0.2$ MeV, $\Gamma \approx 1.5 \pm 0.3$ MeV. High-resolution magnetic spectra, observed for $E_d = 1.0$ to 2.2 MeV show only the ground state peak, superposed on a continuous distribution, with no evidence of an excited-state group. The shape of the ground state peak is well accounted for with the parameters $\gamma^2 = 17.6 \times 10^{-13}$ MeV-cm, $R = 2.9 \times 10^{-13}$ cm, taken from (n, α) scattering data (1958WE27).

17. ${}^9\text{Be}(\gamma, \alpha){}^5\text{He}$ $Q_m = -2.529$

See ${}^9\text{Be}$.

Table 5.3: Energy levels of ${}^5\text{Li}$

E_x (MeV)	J^π	Γ (MeV)	Decay	Reactions
0	$\frac{3}{2}^-$	≈ 1.5	p, α	2, 5, 6, 9, 10, 11, 12, 13
5 – 10	$\frac{1}{2}^-$	3 – 5	p, α	6
16.81	$\frac{3}{2}^+$	≈ 0.3	d, ${}^3\text{He}$, p, α , γ	2, 3, 4

${}^5\text{Li}$
(Fig. 2)

1. ${}^3\text{H}({}^3\text{He}, n){}^5\text{Li}$ $Q_m = 10.297$

Not reported.

2. ${}^3\text{He}(d, \gamma){}^5\text{Li}$ $Q_m = 16.555$

The excitation curve measured from $E_d = 0.2$ to 2.85 MeV shows a broad maximum at $E_d = 0.45 \pm 0.04$ MeV ($E_\gamma = 16.6 \pm 0.2$, $\sigma = 50 \pm 10 \mu\text{b}$, $\Gamma_\gamma = 11 \pm 2$ eV). Above this maximum, non-resonant capture is indicated by a slow rise of the cross section. The radiation appears to be isotropic to $\pm 10\%$ at $E_d = 0.58$ MeV, consistent with s-wave capture (1954BL89). See also (1955KU1B), (1955BA1G; theor.).

3. (a) ${}^3\text{He}(d, p){}^4\text{He}$ $Q_m = 18.351$ $E_b = 16.555$
 (b) ${}^3\text{He}(d, np){}^3\text{He}$ $Q_m = -2.226$

Cross sections and angular distributions for reaction (a) from $E_d = 35$ keV to 10 MeV are given in (1957JA37); see also (1957BO79, 1957FI1A). Below 100 keV the cross section follows the simple Gamow form: $\sigma = (18.2 \times 10^3/E)\exp(-91E^{-1/2})$ b (E in keV) (1953JA1A, 1954AR02). A pronounced resonance occurs at $E_d = 430$ keV, of about 450-keV width. The peak cross section is given as 0.695 ± 0.014 b by (1952BO68, 1955KU03) and 0.92 ± 0.07 b by (1953YA02, 1954FR01). The resonance is closely fitted with the one-level dispersion formula, using the parameters listed in Table 5.2 (see ${}^3\text{H}(d, n){}^4\text{He}$). See also (1955BA1G, 1955HA1B, 1955JO1A, 1956KL1A), (1958PO1A; theor.).

The angular distribution of protons is isotropic near resonance and shows forward peaking at higher energies; the similarity to ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$ is very close. See also (1955KU1B, 1957BO79, 1957BR23).

Above $E_{\text{d}} = 3.71$ MeV, deuteron breakup (reaction (b)) is observed (1955HE90).

4. ${}^3\text{He}(\text{d}, \text{d}){}^3\text{He}$

$$E_{\text{b}} = 16.555$$

Differential cross sections for $E_{\text{d}} = 0.4$ to 3 MeV are plotted in (1957JA37); see also (1952AL36, 1957BR23, 1957FI1A). In the range $E_{\text{d}} = 380$ to 570 keV, $\theta_{\text{c.m.}} = 65^\circ$, the scattering cross section is considerably below Rutherford scattering and is consistent with s-wave formation of a $J = \frac{3}{2}^+$ state. Above $E_{\text{d}} = 2$ MeV, the distributions are quite similar to those observed in ${}^3\text{H}(\text{d}, \text{d}){}^3\text{H}$ (1952AL36, 1954BR05, 1957BR23).

5. ${}^3\text{He}({}^3\text{He}, \text{p}){}^5\text{Li}$

$$Q_{\text{m}} = 11.062$$

The proton spectrum at $E({}^3\text{He}) = 360$ keV shows an unresolved ground-state group superposed on a broad continuum. No evidence is found for well-defined proton groups of lower energy than the ground-state group (1954GO18). See also (1953AL1A), (1957BR18) and ${}^6\text{Be}$.

6. ${}^4\text{He}(\text{p}, \text{p}){}^4\text{He}$

$$E_{\text{b}} = -1.796$$

Differential elastic scattering cross sections have been measured at numerous energies from 0.95 to 95 MeV, as indicated in Table 5.4; curves at several energies are given by (1957JA37) and (1957BR28). Phase shifts derived from the experimental data are listed in the table. At $E_{\text{p}} = 40$ MeV, the differential cross section shows “diffraction” maxima and minima characteristic of the optical model (1957BR24: see also (1956BU95, 1957GI14, 1957HO1C, 1958GA13)). Recent surveys of the experimental and theoretical situation are reported by (1958GA13, 1958HO1B, 1958MI93).

Even at low energies, the phase shift analysis clearly requires a splitting of $\text{P}_{1/2}$ and $\text{P}_{3/2}$ levels, generally attributed to spin-orbit effect. Either order of the P-doublet can be used to fit the cross section data (1949CR1A, 1952DO30): that the $\text{P}_{3/2}$ state is the lower is established by measurements of the polarization of scattered protons (1952HE15, 1955SC1A, 1956JU10, 1958SC1A). The tabulated phase shifts apply to this case.

The $\text{P}_{3/2}$ phase shift shows a pronounced resonance effect, passing through 90° at $E_{\text{p}} = 2.8$ MeV (1949CR1A), while the $\text{P}_{1/2}$ changes only slowly over a range of several MeV. Analysis by (1952DO30), based on resonance theory, yields for the $\text{P}_{3/2}$ level (ground state of ${}^5\text{Li}$) $E_{\lambda}(\text{c.m.}) = -4.1$ MeV, $\gamma_{\lambda}^2 = 25 \times 10^{-13}$ MeV-cm and for the $\text{P}_{1/2}$, $E_{\lambda}(\text{c.m.}) = 3.4$ MeV, $\gamma_{\lambda}^2 = 105 \times 10^{-13}$ MeV-cm (see also (1952AD09)). (These widths correspond to $\theta_{\text{p}}^2 = 0.9$ and 3.9 times the sum-rule

limit, respectively, using $R = 2.9 \times 10^{-13}$ cm.) It thus appears that the $P_{1/2}$ state, if resonance theory is at all appropriate here, is extremely broad and located 5 – 10 MeV above the ground state. A new analysis of all data < 18 MeV yields $E_{\text{res}}(\text{lab}) = 2.6$ MeV, $\gamma_p^2 = 12 \times 10^{-13}$ MeV-cm, $\theta_p^2 = 0.40$ for the ground state and $E_{\text{res}}(\text{lab}) = 10.8$ MeV, $\gamma_p^2 = 30 \times 10^{-13}$ MeV-cm, $\theta_p^2 = 1.0$ for the excited state, using $R = 2.6 \times 10^{-13}$ cm (1958MI93). The s-wave phase shifts are well accounted for by hard-sphere scattering, with $R = 2.9 \times 10^{-13}$ cm (1952AD09, 1952DO30: see, however, (1956VA1C)), $R = 2.0 \times 10^{-13}$ cm (1958MI93). Semi-empirical phase shifts and polarizations for $E_p = 10$ to 40 MeV are given by (1958GA13).

Proton and neutron scattering in helium are discussed in terms of a central potential with spin-orbit interaction by (1954BR1B, 1954HO1B, 1954JA1A, 1954SA1B, 1955HO1C, 1958GA13). (1955HO1C) obtain a good account of the s- and p-wave interactions with a Gaussian potential and Serber exchange force. The effect of tensor forces is discussed by (1956FE1A). See also (1956AB1A, 1956LE1C, 1958HO1B).

Polarization of the scattered protons is discussed by (1949CR1A, 1952AD09, 1952DO30, 1952HE15, 1955SC1A, 1956BR1D, 1956JU10, 1957RO1A, 1957SC1B, 1957SE40, 1958BR24, 1958SC1A, 1958SC27) and others.

For α -scattering in hydrogen: see (1954JU1B, 1954RU1A, 1957RO1A). See also (1956EI05, 1957TY27).

7. (a) ${}^4\text{He}(p, d){}^3\text{He}$	$Q_m = -18.351$	$E_b = -1.796$
(b) ${}^4\text{He}(p, pn){}^3\text{He}$	$Q_m = -20.577$	

Angular distributions are reported at $E_p = 27.9$ MeV (1957WI22), 31.6 MeV (1952BE1A) and 95 MeV (1955TE1A) for reaction (a). The cross section at $E_p = 40$ MeV, $\theta = 30^\circ$, is 10 ± 1 mb/sr (1956EI05). For reaction (b), see (1957WI22) and (1956EI05).

8. ${}^4\text{He}(p, 2p){}^3\text{H}$	$Q_m = -19.812$	$E_b = -1.796$
---------------------------------------	-----------------	----------------

See (1957WI22) and (1956EI05).

9. ${}^4\text{He}(d, n){}^5\text{Li}$	$Q_m = -4.023$
---------------------------------------	----------------

At $E_d = 13$ MeV, a broad, asymmetric neutron group corresponding to the ground state is observed. There is no evidence for structure corresponding to the $P_{1/2}$ excited state (1956BO1F, 1956BO43).

10. ${}^6\text{Li}(\gamma, n){}^5\text{Li}$	$Q_m = -5.494$
---	----------------

Table 5.4: Phase shifts in ${}^4\text{He}(p, p){}^4\text{He}$ ^a

E_p (lab) (MeV)	$S_{1/2}$ (deg)	$P_{1/2}$ (deg)	$P_{3/2}$ (deg)	$D_{3/2}$ (deg)	$D_{5/2}$ (deg)	References
0.95	-12.0	3.3	3.3	0	0	(1949CR1A, 1949FR20)
1.49	-18.1	4.1	20.4			(1949CR1A, 1949FR20)
1.70	-17.6	4.2	31.1			(1949CR1A, 1949FR20)
2.02	-24.6	8.1	47.8			(1949CR1A, 1949FR20)
2.22	-26.7	9.4	60.6			(1949CR1A, 1949FR20)
2.53	-28.2	13.1	78.8			(1949CR1A, 1949FR20)
3.03	-27.3	10.3	99.3	0.7	0.1	(1958MI93)
3.04	-32.0	15.7	96.6			(1949CR1A, 1949FR20)
3.51	-30.9	18.8	107.6	1.3	-1.1	(1958MI93)
3.58	-35.2	20.3	105.4			(1949CR1A, 1949FR20)
4.02	-32.9	23.6	112.4	2.1	-1.7	(1958MI93)
4.50	-43.1	29.1	111.6	-3.1	-0.5	(1958MI93)
5.00	-51.8	35.5	109.9	-4.7	0.1	(1958MI93)
5.1	(no analysis)					(1951BR93)
5.78	-47.9	38.7	112.9	-1.3	-0.49	(1952DO30, 1954KR1B, 1955LU60)
7.50	-57.95	52.51	112.1	-1.87	+0.44	(1956PU41)
9.48	-65.36	54.72	109.2	-5.73	-3.21	(1952PU1A, 1956PU41)
9.55	(optical model)					(1954FR22, 1957GI14: see (1955WI26, 1957HO1C))
9.76	(no analysis)					(1955WI26)
9.8	(no analysis)					(1954CO69)
12.0	-66.9	60.1	108.7			(1957BR28) ^b
14.0	-76.7	50.2	92.7			(1957BR28) ^b
16.0	-81.1	49.1	89.5			(1957BR28) ^b
17.0 ^c				(-14)	(-17)	(1954AL28)
17.45	-85.7	53.2	94.8			(1956BR29, 1957BR28)
18	-85.8	46.4	85.4			(1956BR29, 1957BR28)
19.4	(no analysis)					(1956VA1B, 1957VA1B)
27.9	(no analysis)					(1957WI22)
31.6	(no analysis)					(1953CO62)
39.85	(optical model)					(1957BR24)
95	(no analysis)					(1955TE1A)

^a Phase shifts for $E_p = 1$ to 18 MeV are plotted by (1958MI93); extrapolated values from 10 to 40 MeV are tabulated by (1958GA13).

^b (1957BR28) also tabulates values for $E_p = 13, 15$ and 17 MeV; D-wave $< 8^\circ$.

^c From ${}^4\text{He}(n, n){}^4\text{He}$.

The photoneutron threshold is 5.4 ± 0.2 MeV (1951TI06), 5.35 ± 0.2 MeV (1951SH63), 5.6 ± 0.1 MeV (1955TI1A), 5.73 ± 0.05 MeV (1958RY77). An earlier reported higher state is not confirmed (1955TI1A).

$$\begin{aligned} 11. \quad {}^6\text{Li}(p, d){}^5\text{Li} & \quad Q_m = -3.267 \\ & \quad Q_0 = -3.0 \pm 0.15 \text{ (1955LI09)}. \end{aligned}$$

At $E_p = 18.6$ MeV, the ground-state group appears as a broad, asymmetric peak ($\Gamma = 1.8$ MeV) which tails off to a low continuum at lower energies. The angular distribution of the ground-state group has a maximum at 15° (c.m.) and, at small angles, conforms with stripping theory ($l_n = 1$) (1955LI09).

$$\begin{aligned} 12. \quad {}^6\text{Li}(d, t){}^5\text{Li} & \quad Q_m = 0.765 \\ & \quad Q_0 = 0.80 \pm 0.15 \text{ (1958FR52)}. \end{aligned}$$

At $E_d = 1$ MeV a broad ground-state triton group is observed ($\Gamma_{\text{c.m.}} = 1.5$ MeV). The distribution of protons from the ${}^5\text{Li}$ breakup indicates a pronounced forward distribution (1958FR52). See also ${}^8\text{Be}$.

$$13. \quad {}^6\text{Li}({}^3\text{He}, \alpha){}^5\text{Li} \quad Q_m = 15.084$$

See (1955AL57) and (1953KU24).

$$14. \quad {}^7\text{Li}(p, t){}^5\text{Li} \quad Q_m = -4.262$$

This reaction has been observed at $E_p = 17.5$ MeV (1957MA04).

References

(Closed December 01, 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1949CR1A Critchfield and Dodder, Phys. Rev. 76 (1949) 602
- 1949FR20 G. Freier, E. Lampi, W. Sleator and J.H. Williams, Phys. Rev. 75 (1949) 1345
- 1951BA85 S. Bashkin, F.P. Mooring and B. Petree, Phys. Rev. 82 (1951) 378
- 1951BR93 C.H. Braden, Phys. Rev. 84 (1951) 762
- 1951BU1B Butler and Symonds, Phys. Rev. 83 (1951) 858
- 1951FL1A Flowers, Proc. Roy. Soc. A204 (1951) 503
- 1951FR1A French and Treacy, Proc. Phys. Soc. (London) A64 (1951) 452
- 1951HI1A Hibdon and Muehlhause, A. N. L. 4680 (1951)
- 1951LE1A Leland and Agnew, Phys. Rev. 82 (1951) 559
- 1951MC63 A.W. McReynolds, Phys. Rev. 84 (1951) 969
- 1951MC37 K.G. McNeill and W. Rall, Phys. Rev. 83 (1951) 1244
- 1951SH63 R. Sher, J. Halpern and A.K. Mann, Phys. Rev. 84 (1951) 387
- 1951TI06 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A64 (1951) 212
- 1952AD09 R.K. Adair, Phys. Rev. 86 (1952) 155
- 1952AL36 J.C. Allred, A.H. Armstrong, A.M. Hudson, R.M. Potter, E.S. Robinson, L. Rosen and E.J. Stovall Jr., Phys. Rev. 88 (1952) 425
- 1952AR30 H.V. Argo, R.F. Taschek, H.M. Agnew, A. Hemmendinger and W.T. Leland, Phys. Rev. 87 (1952) 612
- 1952BE1A Benveniste, Thesis, UCRL 1689 (1952)
- 1952BO68 T.W. Bonner, J.P. Conner and A.B. Lillie, Phys. Rev. 88 (1952) 473
- 1952CO35 J.P. Conner, T.W. Bonner and J.R. Smith, Phys. Rev. 88 (1952) 468
- 1952DO30 D.C. Dodder and J.L. Gammel, Phys. Rev. 88 (1952) 520
- 1952HE15 M. Heusinkveld and G. Freier, Phys. Rev. 85 (1952) 80
- 1952HU1A Huber and Baldinger, Helv. Phys. Acta 25 (1952) 435
- 1952PU1A Putnam, Phys. Rev. 87 (1952) 932

1952ST69 W.R. Stratton, G.D. Freier, G.R. Keepin, D. Rankin and T.F. Stratton, Phys. Rev. 88 (1952) 257

1953AL1A Almqvist, Allen, Dewan and Pepper, Phys. Rev. 91 (1953) 1022

1953CO62 B. Cork, Phys. Rev. 89 (1953) 78

1953CU20 P. Cuer and J.J. Jung, Compt. Rend. 236 (1953) 1252

1953JA1A Jarvis and Roaf, Proc. Roy. Soc. 218 (1953) 432

1953KU24 W.E. Kunz, C.D. Moak and W.D. Good, Phys. Rev. 91 (1953) 676

1953MO61 C.D. Moak, Phys. Rev. 92 (1953) 383

1953SE29 J.D. Seagrave, Phys. Rev. 92 (1953) 1222

1953SI1A Simon and Welton, Phys. Rev. 90 (1953) 1036

1953YA02 J.L. Yarnell, R.H. Lovberg and W.R. Stratton, Phys. Rev. 90 (1953) 292

1954AL24 D.L. Allan, Nature 174 (1954) 267

1954AL28 M.H. Alston, A.V. Crewe, W.H. Evans, L.L. Green and J.C. Willmott, Proc. Phys. Soc. (London) A67 (1954) 657

1954AL35 K.W. Allen, E. Almqvist, J.T. Dewan and T. Pepper, Phys. Rev. 96 (1954) 684

1954AR02 W.A. Arnold, J.A. Phillips, G.A. Sawyer, E.J. Stovall Jr. and J.L. Tuck, Phys. Rev. 93 (1954) 483

1954BL89 J.M. Blair, N.M. Hintz and D.M. Van Patter, Phys. Rev. 96 (1954) 1023

1954BR05 R.J.S. Brown, K.F. Famularo, H.D. Holmgren, D. Rankin and T.F. Stratton, Phys. Rev. 96 (1954) 80

1954BR1B Bransden and McKee, Phil. Mag. 45 (1954) 869

1954CO1B Cooper, AECU 3204 (1954)

1954CO69 B. Cork and W. Hartsough, Phys. Rev. 96 (1954) 1267; UCRL 2590 (1954)

1954FR01 G. Freier and H. Holmgren, Phys. Rev. 93 (1954) 825

1954FR03 G.M. Frye Jr., Phys. Rev. 93 (1954) 1086

1954FR22 R.G. Freemantle, T. Grotdal, W.M. Gibson, R. McKeague, D.J. Prowse and J. Rotblat, Phil. Mag. 45 (1954) 1090

1954GO18 W.M. Good, W.E. Kunz and C.D. Moak, Phys. Rev. 94 (1954) 87

1954HO1B Hochberg, Massey and Underhill, Proc. Phys. Soc. (London) A67 (1954) 957

1954JA1A B. Jancovici and I. Talmi, Phys. Rev. 95 (1954) 289

1954JO1C Johnson and Jones, Phys. Rev. 93 (1954) 1286

1954JU1B Juric, Atomics and Atomic Tech. 5 (1954) 333

1954KR1B Kravtsov, Usp. Fiz. Nauk 54 (1954) 3

1954RU1A Ruhla, J. Physique 15 (1954) 451
1954SA1B Sack, Biedenharn and Breit, Phys. Rev. 93 (1954) 321
1954SM97 J.R. Smith, Phys. Rev. 95 (1954) 730
1955AL57 E. Almqvist, K.W. Allen and C.B. Bigham, Phys. Rev. 99 (1955) 631A
1955BA1G Baz, Izv. Akad. Nauk 19 (1955) 360
1955BE1B Benveniste and Zenger, UCRL 4266 (1955)
1955CL1A Clementel and Villi, Nuovo Cim. 2 (1955) 1121
1955HA1B Hale and Present, Phys. Rev. 99 (1955) 1366
1955HE89 A. Hemmendinger and H.V. Argo, Phys. Rev. 98 (1955) 70
1955HE90 R.L. Henkel, J.E. Perry, Jr. and R.K. Smith, Phys. Rev. 99 (1955) 1050
1955HO1C Hochberg, Massey, Robertson and Underhill, Proc. Phys. Soc. (London) A68 (1955) 746
1955HU1B Hughes and Harvey, BNL-325 (1955)
1955JO1A Johnson and Porter, Phys. Rev. 99 (1955) 630A
1955KH31 L.M. Khromchenko and V.A. Blinov, Zh. Eksp. Teor. Fiz. 28 (1955) 741; JETP (Sov. Phys.) 1 (1955) 596
1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252
1955KU03 W.E. Kunz, Phys. Rev. 97 (1955) 456
1955KU1B Kunz, Butler and Holmgren, Phys. Rev. 100 (1955) 1252A
1955LA1B Larsson, Ark. Fys. 9 (1955) 293
1955LE1C Lehmann, Leveque and Fiehrer, Compt. Rend. 241 (1955) 700
1955LE1D Levintov, Miller and Samsev, Dokl. Akad. Nauk SSSR 104 (1955) 803
1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
1955LI09 J.G. Likely, Phys. Rev. 98 (1955) 1538
1955LU60 H. Lustig and J.M. Blatt, Phys. Rev. 100 (1955) 777
1955RE44 A.A. Reut, S.M. Korenchenko, V.V. Yurev and B.M. Pontecorvo, Dokl. Akad. Nauk. SSSR 102 (1955) 723; AERE Lib/Trans-600; AEC-tr-2205 (1955)
1955SA52 G.A. Sawyer and L.C. Burkhardt, Phys. Rev. 98 (1955) 1305
1955SC1A Scott and Segel, Phys. Rev. 100 (1955) 1244A
1955SH1D Shaw, Proc. Phys. Soc. (London) A68 (1955) 43
1955SO56 H.S. Sommers, Jr., J.G. Dash and L. Goldstein, Phys. Rev. 97 (1955) 855
1955TE1A Teem, Selove and Kruse, Phys. Rev. 98 (1955) 259A

1955TI1A Titterton, Prog. Nucl. Phys. 4 (1955) 1
 1955WI26 J.H. Williams and S.W. Rasmussen, Phys. Rev. 98 (1955) 56
 1956AB1A Abraham, Nucl. Phys. 1 (1956) 415
 1956BA1D Bame and Perry, Bull. Amer. Phys. Soc. 1 (1956) 93
 1956BA1E Barjon, Magnac-Valette and Schmouker, Compt. Rend. 242 (1956) 896
 1956BL1A Blumberg and Schlesinger, AECU 3118 (1956)
 1956BO1F Bogdanov, Vlasov, Kalinin, Rybakov and Sidorov, Physica 22 (1956) 1150
 1956BO43 G.F. Bogdanov, N.A. Vlasov, S.P. Kalinin, B.V. Rybakov and V.A. Sidorov, Zh. Eksp. Teor. Fiz. 30 (1956) 981; JETP (Sov. Phys.) 3 (1956) 793
 1956BO44 G.F. Bogdanov, N.A. Vlasov, S.P. Kalinin, B.V. Rybakov and V.A. Sidorov, Zh. Eksp. Teor. Fiz. 30 (1956) 185; JETP (Sov. Phys.) 3 (1956) 113
 1956BR1D Brinkworth and Rose, Nuovo Cim. 3 (1956) 195
 1956BR29 K.W. Brockman, Jr, Phys. Rev. 102 (1956) 391
 1956BU95 E.J. Burge, Y. Fujimoto and A. Hossain, Phil. Mag. 1 (1956) 19
 1956CR47 D.S. Craig, W.G. Gross and R.G. Jarvis, Phys. Rev. 103 (1956) 1427
 1956EI05 R.M. Eisberg, Phys. Rev. 102 (1956) 1104
 1956FE1A Feingold, Phys. Rev. 101 (1956) 258
 1956FO1A Fowler and Brolley, Rev. Mod. Phys. 28 (1956) 103
 1956GA51 A. Galonsky and C.H. Johnson, Phys. Rev. 104 (1956) 421
 1956HU1A Hughes and Carter, BNL-400 (1956)
 1956JU10 A.C. Juveland and W. Jentschke, Z. Physik 144 (1956) 521
 1956JU1B Jung and Cuer, Physica 22 (1956) 1159A
 1956KL1A Kliucharev, Eselson and Valter, Dokl. Akad. Nauk. SSSR 109 (1956) 737
 1956LE1B Levintov, Miller, Shamshev and Tarumov, Physica 22 (1956) 1181A
 1956LE1C Levintov, Zh. Eksp. Teor. Fiz. 30 (1956) 987; JETP (Sov. Phys.) 3 (1956) 796
 1956MA09 D. Magnac-Valette, Compt. Rend. 242 (1956) 760
 1956PU41 T.M. Putnam, J.E. Brolley Jr. and L. Rosen, Phys. Rev. 104 (1956) 1303
 1956RI37 A.C. Riviere, Nucl. Phys. 2 (1956) 81
 1956VA1B Vanetsian and Fedchenko, Physica 22 (1966) A1124
 1956VA1C Van der Spuy, Nucl. Phys. 1 (1956) 381
 1956WA1B Warburton, Ph.D. Thesis, Univ. of Pittsburgh (1956)
 1957BA10 S.J. Bame Jr. and W.T. Leland, Phys. Rev. 106 (1957) 1257

- 1957BA1F Balabanov, Barit, Katsaurov, Frank and Stranikh, Sov. J. At. Energy, Suppl. 5 (1957) 37
- 1957BA1G Balabanov, Barit, Katsaurov, Frank and Stranikh, Sov. J. At. Energy, Suppl. 5 (1957) 43
- 1957BA21 S.J. Bame Jr. and J.E. Perry Jr., Phys. Rev. 107 (1957) 1616
- 1957BL1A Blanchard and Winter, Phys. Rev. 107 (1957) 774
- 1957BO14 T.W. Bonner, F.W. Prosser Jr. and J.H. Slattey, Bull. Amer. Phys. Soc. 2 (1957) 180, E3
- 1957BO79 D.L. Booth, R.S. Hill, F.V. Price and D. Roaf, Proc. Phys. Soc. (London) A70 (1957) 863
- 1957BR18 D.A. Bromley, E. Almqvist, H.E. Gove, A.E. Litherland, E.B. Paul and A.J. Ferguson, Phys. Rev. 105 (1957) 957
- 1957BR23 J.E. Brolley Jr., W.S. Hall and L. Rosen, Bull. Amer. Phys. Soc. 2 (1957) 207, P6
- 1957BR24 M.K. Brussell and J.H. Williams, Phys. Rev. 106 (1957) 286
- 1957BR28 K.W. Brockman, Phys. Rev. 108 (1957) 1000
- 1957DA1B Davidenko, Pogrebov and Saukov, Sov. J. At. Energy 2 (1957) 474
- 1957FA10 F.J.M. Farley and R.E. White, Nucl. Phys. 3 (1957) 561; Erratum Nucl. Phys. 3 (1957) 692
- 1957FI1A Fischer, Bull. Amer. Phys. Soc. 2 (1957) 28
- 1957GI14 W.M. Gibson, D.J. Prowse and J. Rotblat, Proc. Roy. Soc. A243 (1957) 237
- 1957HO1C Hossain and Syeduzzaman, Nucl. Phys. 4 (1957) 88
- 1957HU1D Hughes and Schwartz, BNL-325, Suppl. 1 (1957)
- 1957IN1A Innes, UCRL 8040 (1957)
- 1957JA37 N. Jarmie, J.D. Seagrave et al., LA-2014 (1957)
- 1957LE1B Levintov, Miller and Shamshev, Nucl. Phys. 3 (1957) 221
- 1957LE1C Levintov, Miller and Shamshev, Zh. Eksp. Teor. Fiz. 32 (1957) 274; JETP (Sov. Phys.) 5 (1957) 258
- 1957MA04 D.R. Maxson and E.F. Bennett, Bull. Amer. Phys. Soc. 2 (1957) 180, E5
- 1957MA1C Mazari, Castro, Alba and Vazquez, Rev. Mex. Fis. 6 (1957) 9
- 1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
- 1957RO1A Rosen and Brolley, Phys. Rev. 107 (1957) 1454
- 1957SC1B Scott, Bull. Amer. Phys. Soc. 2 (1957) 349
- 1957SE40 R.E. Segel and S.S. Hanna, Phys. Rev. 106 (1957) 536
- 1957ST1B Striebel and Huber, Helv. Phys. Acta 30 (1957) 67

1957TY27 H. Tyren, G. Tibell and T.A.J. Maris, Nucl. Phys. 4 (1957) 277
1957VA1B Vanetsian and Fedchenko, Sov. J. At. Energy 2 (1957) 141
1957WA01 E.K. Warburton and J.N. McGruer, Phys. Rev. 105 (1957) 639
1957WH1A White and Farley, Nucl. Phys. 3 (1957) 476
1957WI22 A.F. Wickersham Jr., Phys. Rev. 107 (1957) 1050
1958AL05 R.C. Allen and N. Jarmie, Phys. Rev. 111 (1958) 1129
1958BA82 Y.G. Balashko and I.Y. Barit, Zh. Eksp. Teor. Fiz. 34 (1958) 1034; JETP (Sov. Phys.) 7 (1958) 715
1958BR14 J.E. Brolley Jr., W.S. Hall, L. Rosen and L. Stewart, Phys. Rev. 109 (1958) 1277
1958BR24 K.W. Brockman Jr., Phys. Rev. 110 (1958) 163
1958CE1A Cence, Moyer and Waddell, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)
1958FR52 R.T. Frost and S.S. Hanna, Phys. Rev. 110 (1958) 939
1958GA13 J.L. Gammel and R.M. Thaler, Phys. Rev. 109 (1958) 2041
1958HO1B Hodgson, Adv. Phys. 7 (1958) 1
1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
1958JA06 N. Jarmie and R.C. Allen, Phys. Rev. 111 (1958) 1121
1958LE1A Levintov, Miller and Shamshev, JETP (Sov. Phys.) 7 (1958) 712
1958MI93 P.D. Miller and G.C. Phillips, Phys. Rev. 112 (1958) 2043
1958PA09 P.J. Pasma, Nucl. Phys. 6 (1958) 141
1958PO1A Porter, Roth and Johnson, Phys. Rev. 111 (1958) 1578
1958RY77 T.W. Rybka and L. Katz, Phys. Rev. 110 (1958) 1123
1958SC1A Scott, Ph.D. Thesis, Johns Hopkins Univ. (1958)
1958SC27 M.J. Scott, Phys. Rev. 110 (1958) 1398
1958TA1A Tautfest, Phys. Rev. 111 (1958) 1162
1958WE27 G. Weber, Phys. Rev. 110 (1958) 529

