

Energy Levels of Light Nuclei $A = 5$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A227* (1974), p. 1. This version of $A = 5$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1973)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 5$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^5\text{n}\$](#) , [\${}^5\text{H}\$](#) , [\${}^5\text{He}\$](#) , [\${}^5\text{Li}\$](#) , [\${}^5\text{Be}\$](#)

B. Tables of Recommended Level Energies:

[Table 5.1](#): Energy levels of ${}^5\text{He}$

[Table 5.5](#): Energy levels of ${}^5\text{Li}$

C. [References](#)

D. Figures: [\${}^5\text{He}\$](#) , [\${}^5\text{Li}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

⁵n
(Not illustrated)

⁵n has not been observed in the interaction of π^- and ¹⁴N and ¹⁶O: see (1972AG01).

⁵H
(Fig. 3)

From the work of (1967AD05)[†] on the ³He(³He, n)⁵Be reaction (see ⁵Be) it follows that ⁵H is unstable by more than 2.1 MeV to decay into ³H + 2n (using Coulomb corrections based on the 16.7 MeV states in ⁵He–⁵Li). [With the “conventional” correction of (1966LA04, p.2) ⁵H is unbound by > 0.7 MeV.] A study of ⁹Be(α , ⁸B)⁵H at $E_\alpha = 129$ MeV shows no evidence for sharp ⁵H states for several MeV above ³H + 2n (1968MC02). In ³H(t, p)⁵H, at $E_t = 22.25$ MeV, a broad peak appears in the proton spectrum which may correspond to a ⁵H state at ³H + 2n + 1.8 MeV, but which could also result from a four-body breakup (1968YO06). Several recent attempts to observe ⁵H formation in ⁷Li(π^- , pn) and ⁷Li(π^- , d) have been unsuccessful: see (1965CO1D, 1968BO32, 1969MI10). See (1966LA04) for a review of the earlier work, (1965AR04), and (1965AR08, 1966DE1E, 1968CE1A, 1970WA1G, 1971WA08).

⁵He
(Figs. 1 and 3)

GENERAL: (See also (1966LA04).):

Shell model calculations: (1966FR1B, 1967BO1C, 1967EL03, 1967YA1A, 1968EL1A, 1968GO01, 1969EL1B, 1969GO1G, 1970IR01, 1970RA1D, 1970ZO1A, 1971RA15, 1971WA08, 1972KA38, 1972LE1L, 1973HA49).

Cluster calculations: (1965NE1B, 1966HO06, 1967BE1G, 1969ME1C, 1969WI1C, 1971LE1N, 1972DE30).

Special levels: (1971RA15, 1973FE1J).

Electromagnetic transitions: (1966HO06, 1973HA49).

General reviews: (1966DE1E).

Reactions involving pions: (1969GO1C, 1969KO30, 1972GA11, 1973EI01).

Special reactions: (1971CH31, 1972CH31, 1972PN1A, 1973GR22).

[†] References are collected at the end of this article.

Other topics: (1967BO1C, 1967EL03, 1968GO01, 1968HE1D, 1969GI1A, 1969HE1L, 1969SC1J, 1970SU1B, 1970ZO1A, 1971BA2Y, 1971CH50, 1971ZA1D, 1972CA37, 1972GI11, 1972JA14, 1972KA38, 1972LE1L, 1972MA57, 1972RI10, 1973MA20, 1973RA1E, 1973RO1R).

States of ${}^5\text{He}^\ddagger$:

Ground state: The ground and first excited states of ${}^5\text{He}$ comprise the components ${}^2P_{3/2}$ and ${}^2P_{1/2}$ formed by $l = 1$ scattering of a neutron by an α -particle. Shape parameters of the state are best described by tabulated phase shifts, but it is sufficiently narrow to appear as a distinct group in many reactions. The central energy of this group is generally taken as defining the “mass” of ${}^5\text{He}$.

First excited state: Ordinarily appears as a broad continuum in particle spectra, centered several MeV (3–5) above the ground state. The resonant phase shift goes through $\frac{1}{2}\pi$ at $E_{\text{cm}} = 6.43$ MeV, about 5.5 MeV above the ground state.

16.8 MeV: Conspicuous in the ${}^3\text{H} + d$ reactions, the state is almost entirely of a $3 + 2$ configuration, with relative angular momentum zero, and a quite small reduced width for neutron ($l_n = 2$) emission. Properties of the state are listed in Table 5.2.

19.9 MeV: In the range $E_x = 20 - 25$ MeV, there appears to be a number of broad overlapping states, principally of $3 + 2$ configuration, with even parity. There is some indication from ${}^7\text{Li}(p, {}^3\text{He})$, [${}^7\text{Li}(p, t)$] that they have mainly $S = \frac{3}{2}$. Among this manifold, a level at 19.9 ± 0.4 MeV is reported to form a well-defined group ($\Gamma = 3$ MeV) in ${}^7\text{Li}(p, {}^3\text{He})$. There are indications of another structure near $E_x = 24 - 25$ MeV.

1. ${}^3\text{H}(d, \gamma){}^5\text{He}$ $Q_m = 16.70$

At low energies the reaction is dominated by a resonance at $E_d = 107$ keV; the mirror reaction shows resonance at $E_d = 430$ keV. (1970BE1A) have measured the cross section for emission of 16.7 MeV γ -rays for $E_d = 25$ to 100 keV: the ratio $\sigma(d, \gamma)/\sigma(d, n)$ is approximately constant at $(2.1 \pm 0.6) \times 10^{-4}$, leading to $\Gamma_\gamma = 14 \pm 4$ eV, where Γ_n is taken as 66 keV. (1963BU07) have measured thick target yields from $E_d = 150$ to 1300 keV. The derived cross sections are analyzed into resonant and direct-capture contributions: the cross section at resonance is here reported as $60 \mu\text{b}$ [vs. 1 mb reported by (1970BE1A)]. At $E_d = 1025 \pm 47$ keV, the differential cross section is $0.44 \pm 0.12 \mu\text{b/sr}$ (90°) and the γ to n branching ratio is an order of magnitude smaller: 2.3×10^{-5} . The angular distribution of the γ -rays is forward peaked and the total cross section is estimated to be $4.8 \mu\text{b}$ (1970KO09). A cluster model calculation of major even parity states predicts a broad peak in dipole strength due mainly to a $\frac{5}{2}^+$ state in a cluster of positive parity states some 4 MeV above the $\frac{3}{2}^+$ state (1971WA08).

\ddagger Similar considerations prevail for ${}^5\text{Li}$.

Table 5.1: Energy levels of ${}^5\text{He}$ ^a

E_x (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	0.60 ± 0.02 ^b	n, α	1, 5, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26
4 ± 1.5	$\frac{1}{2}^-; \frac{1}{2}$	4 ± 1.5	n, α	5, 7, 11, 12, 14, 20
16.76 ± 0.13	$\frac{3}{2}^+; \frac{1}{2}$	0.10 ± 0.05	γ , n, d, t, α	1, 2, 6, 7, 9, 12, 14, 19, 20
19.9 ± 0.4 ^a	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	3 ± 0.6	n, d, t, α	2, 4, 6, 11, 17, 19, 20
24 – 25		broad		19, 20

^a See also the discussion ‘States of ${}^5\text{He}$ ’ section in ${}^5\text{He}$.

^b See Table 5.2 in (1966LA04).

 Table 5.2: Resonance parameters for ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$ and ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$

E_r (keV)	Γ_{lab} (keV)	l_d	J^π	$l_{\text{n,p}}$	R (fm)	E_λ (keV)	γ_d^2 (keV)	$\gamma_{\text{n,p}}^2$ (keV)	θ_d^2 ^c	$\theta_{\text{n,p}}^2$ ^c	E_x (MeV)
107 ^a	135	0	$\frac{3}{2}^+$	2	5.0	−464	2000 ± 500	50 ± 10	1.0	0.018	16.76
					7.0	−126	715	17	0.7	0.011	
450 ^b	≈ 450	0	$\frac{3}{2}^+$	2	5.0	−391	2930	42	1.4	0.013	16.66
					7.0	129	780	12	0.7	0.008	

^a ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$.

^b ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$.

^c Units of $3\hbar^2/2MR^2$.

2. (a) ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$	$Q_{\text{m}} = 17.5900$	$E_{\text{b}} = 16.70$
(b) ${}^3\text{H}(\text{d}, 2\text{n}){}^3\text{He}$	$Q_{\text{m}} = -2.9885$	
(c) ${}^3\text{H}(\text{d}, \text{pn}){}^3\text{H}$	$Q_{\text{m}} = -2.2246$	

Below $E_{\text{d}} = 100$ keV, the cross section for reaction (a) follows the Gamow function, $\sigma = (A/E) \exp(-44.40 E^{-1/2})$ (1953JA1A, 1954AR02). A strong resonance, $\sigma(\text{peak}) = 5.0$ b, appears at $E_{\text{d}} = 107$ keV: see Table 5.2. From $E_{\text{d}} = 10$ to 500 keV, the cross section is well fitted with the assumption of s-wave formation of a $J^{\pi} = \frac{3}{2}^{+}$ state (1952AR1A, 1952CO35, 1955KU03). Excitation curves and angular distributions for reaction (a) have been measured from $E_{\text{d}} = 8$ keV to 21 MeV: see (1966LA04) for earlier references and (1966KO19: differential cross section (90°); $E_{\text{d}} = 0.115$ to 1.65 MeV), (1968IV01: excitation functions at 3 angles; $E_{\text{d}} = 3.97$ to 10.99 MeV), (1973MC05: angular distributions, at ≈ 1 MeV intervals; absolute 0° cross sections to within $\pm 3.1\%$; $E_{\text{d}} = 5.0$ to 15.7 MeV) and (1968SI1B: differential cross sections (0°), angular distributions; $E_{\text{d}} = 7.0, 11.4, 15.0$ and 19.0 MeV; relative cross sections at 1 MeV intervals, $E_{\text{d}} = 7.0$ to 21.0 MeV). There is some evidence of resonant behavior between $E_{\text{d}} = 3$ and 9 MeV (1960ST25). See also (1968SI1B, 1973MC05).

A study of reaction (a) with polarized deuterons at $E_{\text{d}} = 0.2$ to 1.0 MeV indicates intervention of the s-wave, $J^{\pi} = \frac{1}{2}^{+}$ channel, as well as possible p-waves above $E_{\text{d}} = 0.3$ MeV (1965TR01, 1971GR32). This effect is evidently very small below $E_{\text{d}} = 100$ keV (1971OH1B). At higher energies, the neutron polarization $P_1(\theta_1)$ shows an angular distribution that peaks typically at $\theta_1 = 30^{\circ}$ lab, then goes negative with increasing angle and has a minimum between 90° and 130° , depending upon energy. (1971MU04) have made an extensive study of $P_1(30^{\circ})$ for $E_{\text{d}} = 5$ to 15 MeV and, with deuterium as target, for $E_{\text{t}} = 4.5$ to 19.5 MeV (neutrons near 135° lab). The polarization increases monotonically from 0.03 at $E_{\text{d}} = 3$ MeV to ≈ 0.5 at $E_{\text{d}} = 6.5$ MeV and then with a lower slope to 0.69 at $E_{\text{d}} = 13$ MeV. The change in the slope may be caused by excited states of ${}^5\text{He}$ near 20 MeV. Comparison with the ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$ mirror reaction at corresponding cm energies shows excellent agreement between the polarization values in the two reactions up to $E_{\text{d}} = 6$ MeV, but then the proton polarization becomes $\approx 15\%$ higher, converging back to the neutron values at $E_{\text{d}} \approx 12-13$ MeV. This may be due to experimental factors (1971MU04). Using polarized deuterons (1971HI07) find that the average ratio of vector analyzing power of ${}^3\text{He}(\text{d}, \text{p})$ to ${}^3\text{H}(\text{d}, \text{n})$ is 1.016 ± 0.015 at $E_{\text{d}} = 6$ MeV and 1.035 ± 0.020 at 10 MeV. The vector analyzing power of the two reactions agree to within ± 0.025 at all angles. This agreement between the mirror reactions is in apparent conflict with the result of (1971MU04). See also reaction 3 in ${}^5\text{Li}$. For other polarization measurements see Table 5.3.

(1968BO35) have examined the parity non-conserving effects in this strong reaction with polarized deuterons ($E_{\text{d}} = 0.14$ MeV). The experiment measured the ratio of the number of neutrons emitted parallel and anti-parallel to the direction of the initial polarization vector. The magnitude of the real part of the parity violating amplitude, F , is $\lesssim 3.8 \times 10^{-3}$.

The energy spectrum of ${}^3\text{He}$ particles in reaction (b) has been studied at $E_{\text{d}} = 10.9$ MeV at several angles: no evidence was found for a bound dineutron [$< 5 \mu\text{b}/\text{sr}$ at 6° and 7.5°]. Strong variations of the spectral shape with angle indicate that the Watson-Migdal approximation is inadequate and that several other first-order processes must be included. The determination of the

Table 5.3: ${}^3\text{H}(d, n){}^4\text{He}$ polarization studies

E_d (MeV)	Refs.
51, 76, 93 keV	(1971OH1B)
0.1, 0.17 MeV	(1961RU1A, 1962SE09)
0.1 – 7.7	(1961PE13, 1964PE14)
0.125	(1968BE05)
0.2 – 1.0	(1971GR32)
0.3 – 1.8	(1968HE16, 1969HE1H)
0.4 – 1.7	(1969WI1D)
0.6, 1.2	(1965BO13)
1.0 – 5.0	(1972SM05)
2.1 – 2.9	(1965CH15)
2.1 – 5.8	(1967BU11)
$E_d = 5 - 15, E_t = 4.5 - 19.5$	(1971MU04, 1970BR30)
6 – 10	(1971HI07)
6 – 11	(1964WA22)
8 – 20	(1964AL1E)
10	(1961TR05)
17	(1972RY01)

nn scattering length is ambiguous (1970LA10, 1970LA1K). This problem is also discussed by (1971GR45: $E_d = 13.4$ MeV), (1970SL1A: $E_d = 31.9$ MeV), (1966BA1H: $E_d = 29.8, 32.5$ MeV), (1971GR12: $E_t = 22$ MeV) and (1972BA32: $E_d = 83$ MeV). A discussion and comparison of attempts to measure the 1S_0 nn scattering length is presented by (1971GR45): based on kinematically complete measurements and using the Watson-Migdal treatment, they find $a_{nn} = -16.0 \pm 1.0$ fm. Intervention of n - ${}^3\text{He}$ final state interactions is believed to be small in this work (1971GR45). See also (1973GR1L, 1973JE02). At $E_t = 22$ MeV no clear evidence is seen for sequential processes via excited states of ${}^4\text{He}$ (1971GR12).

Analysis of reaction (c) at $E_d = 20$ MeV is consistent with the existence of a broad resonance in the ${}^3\text{H}+p$ system corresponding to ${}^4\text{He}^*(22.)$ (1968NE1D): see (1973FI04) for a general discussion of the states of ${}^4\text{He}$. n-p final-state interaction enhancement has been studied by (1973SL03) at $E_d = 35$ MeV. They find that only when $E_{t,p} \leq 1$ MeV is agreement with the Watson-Migdal theory obtained, even if the data far from dominant n-t final-state interaction and p-t quasi-free scattering is chosen. See also (1966LA04).

See also (1965FL1A, 1966BR02, 1966BR1F, 1966BR1E, 1966BU09, 1966VA1A, 1967BO1D, 1967NE1B, 1968OH1A, 1969BU19, 1969ME1B, 1970HA1L, 1971MC1L, 1971OH1A, 1971PO1C,

1971SC1L, 1973GE1G), (1968TO1F, 1970SL1B, 1971HA2G, 1971WA1D, 1972OH1D, 1972SC1R) and (1965DU1A, 1965GO1B, 1966BA1J, 1966HU16, 1967OH1A, 1968HA1F, 1969HA1M, 1969HE1K, 1970PO1A, 1970SE1B, 1971DE20, 1971OH1C, 1971SI30, 1972SE09; theor.).

$$3. \text{}^3\text{H}(\text{d}, \text{p})\text{}^4\text{H} \rightarrow \text{}^3\text{H} + \text{n} \quad Q_{\text{m}} = -2.2246 \quad E_{\text{b}} = 16.70$$

See (1967JA07, 1973SL03) and (1973FI04).

$$4. \text{}^3\text{H}(\text{d}, \text{d})\text{}^3\text{H} \quad E_{\text{b}} = 16.70$$

The elastic scattering has been studied for $E_{\text{d}} = 2.6$ to 11.0 MeV. The excitation curves show an interference at $E_{\text{x}} \approx 19$ MeV and a broad ($\Gamma > 1$ MeV) resonance corresponding to $E_{\text{x}} = 20.0 \pm 0.5$ MeV, also seen in ${}^3\text{He}(\text{d}, \text{d})$ [see ${}^7\text{Li}$]. Together with data from ${}^3\text{H}(\text{d}, \text{n})\text{}^4\text{He}$, this work favors an assignment $D_{3/2}$ or $D_{5/2}$ with a mixture of doublet and quartet components (channel spin $\frac{1}{2}$ and $\frac{3}{2}$) if only one state is involved (1967TO02, 1968IV01) [any appreciable doublet component would, however, be in conflict with results from ${}^7\text{Li}(\text{p}, \text{}^3\text{He})\text{}^5\text{He}$]. Measurements of differential cross section and analyzing power using polarized deuterons with $E_{\text{d}} = 3.2$ to 12.3 MeV show resonance-like behavior in the vector analyzing power near $E_{\text{d}} = 5$ MeV. The anomaly appears in the odd Legendre coefficients and is interpreted in terms of a $(\frac{1}{2}, \frac{3}{2})^-$ excited state of ${}^5\text{He}$ with $E_{\text{x}} \approx 19.6$ MeV. Broad structure in the differential cross section near 6 MeV, principally in the even Legendre coefficients, corresponds to an even parity state ${}^5\text{He}^*(20.0)$ (1971KI02)[§]. See also (1966BA1J, 1966BA1M, 1966DE1F, 1969BE1M, 1970HE1C, 1973CH27, 1974CH02; theor.) and (1966LA04).

$$5. \text{(a) } \text{}^3\text{H}(\text{t}, \text{n})\text{}^5\text{He} \quad Q_{\text{m}} = 10.44$$

$$\text{(b) } \text{}^3\text{H}(\text{t}, 2\text{n})\text{}^4\text{He} \quad Q_{\text{m}} = 11.3324$$

At $E_{\text{t}} = 0.5$ MeV, the reaction appears to proceed via three channels: (i) direct breakup into ${}^4\text{He} + 2\text{n}$, the three-body breakup shape being modified by the n-n interaction; (ii) sequential decay via ${}^5\text{He}(0)$; (iii) sequential decay via a broad excited state of ${}^5\text{He}$. The branching ratios at $\theta = 90^\circ$ are $0.7: 0.2: 0.1$. The width of ${}^5\text{He}(0)$ is estimated to be 0.74 ± 0.18 MeV. Some evidence is also shown for ${}^5\text{He}^*$ at $E_{\text{x}} \approx 2$ MeV, $\Gamma \approx 2.4$ MeV (1965WO03). For reaction (b), see ${}^6\text{He}$.

$$6. \text{(a) } \text{}^3\text{He}(\text{t}, \text{p})\text{}^5\text{He} \quad Q_{\text{m}} = 11.20$$

$$\text{(b) } \text{}^3\text{He}(\text{t}, \text{pn})\text{}^4\text{He} \quad Q_{\text{m}} = 12.0963$$

[§] Tensor analyzing powers have been measured for $E_{\text{d}} = 5.0$ to 11.5 MeV by (1973DE51).

Table 5.4: Recent ${}^4\text{He}(n, n){}^4\text{He}$ polarization studies

E_n (MeV)	Refs.
0.262	(1966JE01)
1.02, 2.44	(1968SA25)
3.4, 7.8	(1970ST15)
6.0, 10.0, 16.4, 23.1, 23.7	(1966HO07)
11 – 30	(1972BR10)
12.0, 16.2	(1966BU09)
14, 17.1	(1973LI1K)
15	(1973TO06)
21 – 29	(1971MU04)
25, 28, 34	(1965AR10, 1967HO1D)

Some evidence is reported at $E_t = 22.25$ MeV in reaction (a) for a broad state of ${}^5\text{He}$ at $E_x \approx 20$ MeV, in addition to a sharp peak corresponding to ${}^5\text{He}^*(16.7)$ (1968YO06). For reaction (b) see ${}^6\text{Li}$. See also (1966LA04).

7. ${}^4\text{He}(n, n){}^4\text{He}$

$$E_b = -0.89$$

The coherent scattering length (thermal, bound) is 3.0 ± 0.1 fm (1973MU14). The thermal scattering cross section is 0.773 ± 0.009 b and the absorption cross section at 2200 m/sec $\sigma_{n,\gamma}^0$ is < 0.05 b (1969RO16). (1973MU14) adopt 0.76 ± 0.01 b for $\bar{\sigma}_s$. Total cross sections for $E_n = 4 \times 10^{-4}$ eV to 147 MeV are summarized in (1958HU18, 1960HU1A, 1964ST25, 1966LA04). Recent measurements include those by (1969RO16: $E_n = 0.187$ to 6.195 eV), (1973GO38: $E_n = 0.7$ to 20.0 MeV), (1966ME14: 77.2, 88.2, 110.0, 129.4, 150.9 MeV) and (1968EN1A: 10 GeV/c).

Earlier angular distribution studies are summarized in (1970GA1A). Other recent work is reported by (1968MO26, 1970MO31: $E_n = 0.2$ to 7.0 MeV), (1969CR1B, 1972CR01: $E_n = 0.55, 0.84, 1.16, 1.33$ MeV) and (1971NI06: 17.6, 20.9 and 23.7 MeV). Recent polarization studies are listed in Table 5.4.

The total cross section has a peak of 7.6 b (1973GO38) [see also (1960VA04)] at $E_n = 1.15 \pm 0.05$ MeV, $E_{cm} = 0.92 \pm 0.04$ MeV, with a width of about 1.2 MeV (1964ST25). A second resonance is observed at $E_n = 22.15 \pm 0.12$ MeV, corresponding to the 16.7 MeV $J^\pi = \frac{3}{2}^+$ state (1959BO54, 1964SH1A): $\Gamma_{c.m.} = 100 \pm 50$ keV, $\Gamma_n = \Gamma_d = 50 \pm 35$ keV (1960HU1A) [(1966HO07) find that the data are fitted best by $\Gamma_n < \Gamma_d$ although $\Gamma_n > \Gamma_d$ is not excluded]. Attempts to detect additional resonances in the total cross section have been unsuccessful: see (1966LA04).

The $P_{3/2}$ phase shift shows strong resonance behavior near 1 MeV, while the $P_{1/2}$ shift changes more slowly, indicating a broad $P_{1/2}$ level at several MeV excitation (1952DO30, 1966HO07, 1970AR1B, 1972ST01). (1966HO07) have constructed a set of phase shifts for $E_n = 0$ to 31 MeV, $l = 0, 1, 2, 3$ using largely p- α phase shifts. They have measured differential cross sections from 6 to 30 MeV, with special attention to the region near 22.15 MeV and have fitted the data with the assumed phase shifts. At the $\frac{3}{2}^+$ state the best fit to all data is given by $E_{\text{res}} = 17.669 \text{ MeV} \pm 10 \text{ keV}$, $\gamma_d^2 = 2.0 \text{ MeV} \pm 25\%$, $\gamma_n^2 = 50 \text{ keV} \pm 25\%$ (see Table 5.2). The polarization has been calculated from the phase shifts and is presented as a contour plot (1966HO07).

(1968MO26) have carried out accurate angular distribution measurements for $E_n = 0.2$ to 7.0 MeV and deduced an independent set of p-wave phase shifts, assuming that s-wave shifts are those of a hard sphere, $R = 2.44 \text{ fm}$. The derived total cross sections fit existing data satisfactorily, and polarization results of (1968SA25) and (1963MA29) are well accounted for. Both single-level dispersion formula and Woods-Saxon potential parameters are deduced. A more extensive optical model analysis for $E_n = 0$ to 20 MeV has been carried out by (1968SA1B).

(1970AR1B, 1973AR1N, 1973AR1P) have analyzed existing differential cross sections and polarization data with phase shifts constrained to analytic (effective range) energy dependence, for $E_n = 0$ to 21 MeV. As compared with the (1966HO07) values, the $P_{1/2}$ phase shift is higher in the range 3 to 10 MeV, and both $S_{1/2}$ and $P_{3/2}$ are larger for $E_n = 10$ to 16 MeV. D- and F-wave phase shifts are also determined (1973AR1P).

A single-level-with-background R -matrix parameterization has been carried out by (1972ST01), fitting differential cross section data for $E_n = 0$ to 20 MeV and polarization analyzing power for 0 to 15 MeV. Parameters for $P_{3/2}$ and $P_{1/2}$ levels are $E_{\text{res}} = 0.97$ and 6.43 MeV, and $\gamma^2 = 7.55$ and 12.30 MeV, respectively. See also (1973NI1B). Polarization measurements at six energies from $E_n = 11$ to 30.3 MeV are reported by (1972BR10) together with derived phase shifts. Partial waves as high as $l = 4$ are required, but none shows resonance, other than the $D_{3/2}$. See also (1969RO20).

See also (1966BR1E, 1969MA1H, 1973GA17), (1966BA1K, 1971RH1A, 1971ST1J, 1971WA1D, 1972SC1T) and (1966BA1J, 1966DE1F, 1966PE1C, 1966TO04, 1967SC1E, 1967SW1B, 1969ER1B, 1968SE1A, 1968BA1U, 1969BE1M, 1969CO1E, 1969GE1A, 1969HE1L, 1969KU1D, 1968TH1C, 1969VO1E, 1970ER1A, 1970FE1C, 1970GN1A, 1970HE1C, 1970HO32, 1970OM1A, 1970RE1A, 1971BA1Y, 1971BA74, 1971EF01, 1971GI1B, 1971PI04, 1971PL06, 1971TH09, 1972AD02, 1972IK01, 1972KI1F, 1972LO1K, 1972ZH1B, 1973AD01, 1973PL02, 1974CH02; theor.).

$$8. \text{}^4\text{He}(n, d)\text{}^3\text{H} \qquad Q_m = -17.5900 \qquad E_b = -0.89$$

See (1964DE1B, 1966ME03) and (1966LA04).

$$9. \text{(a) } \text{}^4\text{He}(d, p)\text{}^5\text{He} \qquad Q_m = -3.12$$

$$\text{(b) } \text{}^4\text{He}(d, pn)\text{}^4\text{He} \qquad Q_m = -2.2246$$

A typical proton spectrum consists of a peak corresponding to formation of the ground state of ${}^5\text{He}$, plus a lower continuum of protons ascribed to deuteron breakup (reaction (b)). Ground-state protons show pronounced azimuthal asymmetry when the reaction is induced by 8.5, 10 and 11 MeV vector polarized deuterons. A DWBA calculation is in only qualitative agreement (1967TR05, 1971KE16). See also (1972AV1E).

Coincidence studies of the final state interactions ($\alpha + p$) and ($\alpha + n$) have been carried out at $E_d = 14.2$ MeV (1967FU1D), $E_\alpha = 18.0$ and 24.0 MeV (1970AS02), 42 MeV (1967WA08, 1968WA01), $E_\alpha = 70$ MeV (1973TR04) and $E_d = 78$ MeV (1968BO1M). Spectra of α -particles in coincidence with protons exhibit peaks corresponding to ${}^5\text{He}_{\text{g.s.}}$ as well as broad spectator peaks which occur where the neutron laboratory energy is a minimum (1967WA08, 1968WA01): see ${}^5\text{Li}$. [For earlier work see (1966LA04)]. See also (1971LE02).

At $E_\alpha = 70$ MeV, a kinematically complete experiment shows evidence for sequential decay in reaction (a), proceeding through excited states of ${}^5\text{He}$. Peaks in the coincident yield of protons and deuterons are ascribed to narrow states at $E_x = 16.7 \pm 0.1$ MeV, $\Gamma = 80 \pm 30$ keV, at $E_x = 18.6 \pm 0.1$, 18.8 ± 0.1 and 19.2 ± 0.1 MeV, all with $\Gamma = 180 \pm 60$ keV (1973TR04).

See also (1968TA11, 1971AL1B, 1972LA1L) and (1966HE1B, 1967NA1D, 1969NA09, 1972BL1F, 1972NA1D, 1973DE2B; theor.).

$$10. {}^4\text{He}(t, d){}^5\text{He} \quad Q_m = -7.15$$

Not reported.

$$11. {}^6\text{Li}(\gamma, p){}^5\text{He} \quad Q_m = -4.59$$

At $E_\gamma = 60$ MeV, the proton spectrum shows prominent peaks attributed to ${}^5\text{He}^*(0 + 4.0, 20 \pm 2)$ (1973GA16). See also (1973DE17) and ${}^6\text{Li}$.

$$12. {}^6\text{Li}(e, ep){}^5\text{He} \quad Q_m = -4.59$$

At $E_e = 1180$ MeV, two values of Q are observed: $Q = -3.5 \pm 1.0$ MeV, corresponding to ${}^5\text{He}(0)$, and $Q = -19.0 \pm 1.0$ MeV, corresponding to ${}^5\text{He}^*(16.7)$. Angular distributions at $E_p = 1158$ MeV show that the first represents ejection of a proton with $l > 0$ and the second, an s-proton (1972AN24, 1972AN27, 1972AN29).

$$13. \text{(a) } {}^6\text{Li}(n, d){}^5\text{He} \quad Q_m = -2.37$$

$$\text{(b) } {}^6\text{Li}(n, nd){}^4\text{He} \quad Q_m = -1.4737$$

The angular distribution of ground state deuterons has been studied at $E_n = 14.4$ MeV (1965VA05). DWBA analysis does not reproduce either its shape or the cross section: this may be due to neglect of the deuteron knock-on process (1971MI12). At $E_n = 152$ MeV, two broad structures in the forward spectrum are interpreted as being due to transitions to ${}^5\text{He}^*(0, 14)$. The $Q = -17$ MeV peak is the stronger of the two (1966ME03, 1967ME11). For reaction (b) see (1967VA12). See also (1970VA1L, 1971HE1M, 1973BO1Y) and (1966WE1B). See also (1966LA04).

$$14. {}^6\text{Li}(p, 2p){}^5\text{He} \quad Q_m = -4.59$$

At $E_p = 100$ to 460 MeV, the summed proton spectra show two peaks [$Q = -4.9 \pm 0.3$ and -22.7 ± 0.3 MeV (1966TY01)]. The lower peak corresponds to ejection of an $l = 0$ proton, presumably leaving ${}^5\text{He}$ in the $\frac{3}{2}^+$ state at 16.7 MeV. See also (1966LA04). At $E_p = 100$ MeV, the $\frac{1}{2}^-$ first excited state of ${}^5\text{He}$ is weakly excited (1972MA61). At $E_p = 100$ and 155 MeV, transitions to states above 20 MeV are reported by (1965RO15, 1967RO06, 1972MA61). See also (1965CO1E, 1966BE1B, 1968ZU1A, 1969RU1A, 1973BH1A), (1967KO1B, 1967SH1C, 1968SA1C, 1972KO13, 1973CH1Q; theor.) and ${}^6\text{Li}$.

$$15. (a) {}^6\text{Li}(d, {}^3\text{He}){}^5\text{He} \quad Q_m = 0.90$$

$$(b) {}^6\text{Li}(d, n{}^3\text{He}){}^4\text{He} \quad Q_m = 1.795$$

At $E_d = 14.5$ MeV, the ground state group is observed (1955LE24). See also (1960HA14, 1971IN1C). A study of the proximity process ${}^3\text{He}(n, p){}^3\text{H}$ in the final state of reaction (b) has led to an estimate of the lifetime of ${}^5\text{He}_{\text{g.s.}}$, $\tau = 1.4 \times 10^{-21}$ sec (to within a factor of 2 or 3) (1972KA44). See also (1968LE15).

$$16. {}^6\text{Li}(t, \alpha){}^5\text{He} \quad Q_m = 15.22$$

The width of the ground state is 605 ± 45 keV (1966BI1A). See also (1967BE13) and (1966LA04).

$$17. {}^7\text{Li}(\pi^+, 2p){}^5\text{He} \quad Q_m = 128.51$$

The two-proton spectrum shows broad structures attributed to the ground state of ${}^5\text{He}$, and to $p^{-1}s^{-1}$ and s^{-2} states at ≈ 20 and (≈ 35) MeV excitation (1965CH12).

18. (a) ${}^7\text{Li}(n, t){}^5\text{He}$ $Q_m = -3.36$
 (b) ${}^7\text{Li}(n, tn){}^4\text{He}$ $Q_m = -2.4668$

The angular distribution of ground state tritons has been measured at $E_t = 14.4$ MeV: it is fairly well reproduced by DWBA (1970MI05). Reaction (b) mainly proceeds as a sequential process through ${}^5\text{He}_{\text{g.s.}}$ (1967VA12). See also (1972AN1Q, 1973LI02), (1967BA1E; theor.) and (1966LA04).

19. ${}^7\text{Li}(p, {}^3\text{He}){}^5\text{He}$ $Q_m = -4.12$

At $E_p = 43.7$ MeV, angular distributions of the ${}^3\text{He}$ groups to the ground state of ${}^5\text{He}$ ($\Gamma = 0.80 \pm 0.04$ MeV) and to levels at 16.7 and 19.9 ± 0.4 MeV ($\Gamma = 2.7$ MeV) have been determined. The angular distribution of the ${}^5\text{He}$ ground state group indicates substantial mixing of $L = 0$ and $L = 2$ transfer. The distribution to ${}^5\text{He}^*(16.7)$ is consistent with $L = 1$. Since no transitions are observed in the ${}^7\text{Li}(p, t){}^5\text{Li}$ reaction to the analogue 20 MeV state in ${}^5\text{Li}$ [see ${}^5\text{Li}$], the transition is presumably S-forbidden and the putative states in ${}^5\text{He}$ - ${}^5\text{Li}$ near 20 MeV are ${}^4D_{3/2}$ or ${}^4D_{5/2}$ [compare ${}^3\text{H}(d, d)$] (1966CE05). Particle-particle coincidence data have been obtained at $E_p = 43.7$ MeV. They suggest the existence of ${}^5\text{He}^*(20.0)$ with $\Gamma = 3.0 \pm 0.6$ MeV and of a broad state at ≈ 25 MeV. No $T = \frac{3}{2}$ states decaying via $T = 1$ states in ${}^4\text{He}$ were observed (1968MC02). See also (1967JO1B, 1969DE04, 1970CO1M).

20. (a) ${}^7\text{Li}(d, \alpha){}^5\text{He}$ $Q_m = 14.23$
 (b) ${}^7\text{Li}(d, n){}^4\text{He}{}^4\text{He}$ $Q_m = 15.1233$

At $E_d = 24$ MeV, the α -particle spectrum from reaction (a) shows structures corresponding to the ground and 16.7 MeV states and to states at $E_x \approx 20.2$ and 23.8 MeV with $\Gamma \approx 2$ MeV and ≈ 1 MeV, respectively (1972BA30). See also (1966B11A, 1966MI09, 1966PO1D) and ${}^9\text{Be}$.

Spectra measured in reaction (b) suggest the involvement of a ${}^5\text{He}$ state with $E_x = 2.6 \pm 0.4$ MeV, $\Gamma = 4.0 \pm 1.0$ MeV (1964FE01), $E_x = 5.2 \pm 0.2$ MeV, $\Gamma = 5.6_{-0.6}^{+0.3}$ MeV (1965AS06, 1966AS04), $E_x = 2.6 \pm 0.2$ MeV, $\Gamma = 7 \pm 2$ MeV (1972GI10). See also (1972ST08, 1973HE26). Study of n- α coincidences for $E_d = 2.6$ to 4.0 MeV shows that reaction (b) proceeds mainly by sequential decay, primarily via ${}^5\text{He}(0)$ and excited states of ${}^8\text{Be}$ (1967VA11). See also (1966AS04, 1966MI09, 1967JE01, 1972DE44, 1973HE26). Attempts to observe n- α rescattering effects, following formation of ${}^8\text{Be}^*(16.63, 16.91)$ have been unsuccessful at $E_d = 1.90$ to 2.10 MeV (1972BR08), 2.07 and 2.09 MeV (1971SW10) and 3.0, 3.2 and 3.6 MeV (1968VA12). The upper limit for the rescattering yield is 1% of the yield from sequential decay via intermediate states in ${}^5\text{He}$ and ${}^8\text{Be}$ (1968VA12). Positive results are reported for $E_d = 2.07$ to 2.25 MeV by (1968TH1D). See also (1967BE13, 1967FL12, 1967WI1C, 1968GA1E, 1968WI1E, 1971HU1H, 1973HE06, 1973KA32), ${}^8\text{Be}$, ${}^9\text{Be}$ and (1966LA04).

21. ${}^7\text{Li}({}^6\text{Li}, 2\alpha){}^5\text{He}$ $Q_m = 12.76$

See (1965BE1C).

22. ${}^9\text{Be}(\gamma, \alpha){}^5\text{He}$ $Q_m = -2.46$

See ${}^9\text{Be}$.

23. ${}^9\text{Be}(p, p\alpha){}^5\text{He}$ $Q_m = -2.46$

Studies of this reaction have been carried out at $E_p = 26.0, 35.0$ and 46.8 MeV (1972QU01) and at 57 MeV (1968RO19). Observation of protons and alphas in coincidence at selected angles shows quasifree scattering of the incident protons by S-state α -clusters in ${}^9\text{Be}$ (1968RO19, 1972QU01): see ${}^9\text{Be}$. See also (1969YA1B, 1970GO12).

24. ${}^9\text{Be}(\alpha, 2\alpha){}^5\text{He}$ $Q_m = -2.46$

The population of ${}^5\text{He}(0)$ has been observed at $E_\alpha = 28$ MeV (1965YA02) and at 32.3 and 37.4 MeV (1968YA02). See also (1965HI1B, 1965KU1B, 1966HI1A, 1967ME1C) and ${}^9\text{Be}$.

25. ${}^{11}\text{Be}(d, n){}^4\text{He}{}^4\text{He}{}^4\text{He}$ $Q_m = 6.4577$

At $E_d = 10.4$ and 12.0 MeV, this reaction involves ${}^5\text{He}(0)$ and states in ${}^8\text{Be}$ and ${}^9\text{Be}$ (1971RE19).

26. ${}^{12}\text{C}(n, n){}^4\text{He}{}^4\text{He}{}^4\text{He}$ $Q_m = -7.2748$

See ${}^{12}\text{C}$ in (1968AJ02).

⁵Li
(Figs. 2 and 3)

GENERAL: (See also (1966LA04).):

Shell model calculations: (1966FR1B, 1968GO01, 1969GO1G, 1970RA1D, 1971RA15, 1972LE1L, 1973HA49).

Cluster calculations: (1965NE1B, 1971HE05).

Special levels: (1970HE1D, 1971HE05, 1971RA15, 1973JO1J).

Electromagnetic transitions: (1973HA49).

General reviews: (1966DE1E).

Special reactions: (1971CH31).

Other topics: (1968GO01, 1970RA1J, 1971CH50, 1971ZA1D, 1972CA37, 1972HA57, 1972JA14, 1972LE1L, 1973RO1R).

1. ³He(d, γ)⁵Li

$$Q_m = 16.39$$

Excitation curves and angular distributions have been measured for $E_d = 0.2$ to 2.85 MeV (1954BL89), $E_d = 0.2$ to 1.2 MeV (1968BU09), $E_d = 1$ to 5 and $E(^3\text{He}) = 2$ to 5.5 MeV (1968KR03), $E(^3\text{He}) = 2$ to 26 MeV (1972KI01), $E(^3\text{He}) = 2.3$ to 11.2 MeV (1968DE14) and $E(^3\text{He}) = 3.7$ to 12 MeV (1970SC18).

A broad maximum in the cross section is observed at $E_d = 0.45 \pm 0.04$ MeV [⁵Li*(16.66)]: $\sigma = 50 \pm 10 \mu\text{b}$, $\Gamma_\gamma = 11 \pm 2$ eV (1954BU06: $\gamma_0 + \gamma_1$); $\sigma_{\gamma_0} = 21 \pm 4 \mu\text{b}$, $\Gamma_{\gamma_0} = 5 \pm 1$ eV (1968BU09). See also (1968KR03). The radiation at resonance is isotropic, consistent with s-wave capture: see (1954BU06, 1968BU09, 1968KR03). Study of γ_0 and γ_1 yield $\Gamma = 2.6 \pm 0.4$ MeV for the ground state width, and $E_x = 7.5 \pm 1.0$ MeV, $\Gamma = 6.6 \pm 1.2$ MeV for the $\frac{1}{2}^-$ state. The ratio of γ_0 to γ_1 is 1.00 ± 0.2 and 1.9 ± 0.4 at $E_d = 480$ and 1025 keV (1968BU09).

An excess in the cross section at higher bombarding energies is interpreted by (1972KI01) as being due to a state at $E_x \approx 18$ MeV: even parity is deduced from the relative intensity of γ_0 and γ_1 . It is presumed to be the $\frac{1}{2}^+$ state reported in reactions 3 and 7. A broad peak is also observed at $E_x \approx 20.7$ MeV in the γ_0 cross section. The cross section for γ_1 is ≈ 0 . The observations are consistent with $J^\pi = \frac{5}{2}^+$: angular distributions appear to require at least one other state with significant strength near 19 MeV (1972KI01). (1970SC18) also report this state but find $E_x = 19.7 \pm 0.2$ MeV [$\Gamma = 5.0$ MeV]: $(2J + 1) \Gamma_{\gamma_0} = 1.32$ keV ($\pm 50\%$), suggesting E1 (if only one state is involved). See also (1971WA08; theor.).

Table 5.5: Energy levels of ${}^5\text{Li}$ ^a

E_x (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	≈ 1.5	p, α	1, 5, 6, 7, 9, 10, 11, 12, 13, 14, 15, 16
5 – 10	$\frac{1}{2}^-; \frac{1}{2}$	5 ± 2	p, α	1, 7, 12, 14, 15, 16
16.66 ± 0.07	$\frac{3}{2}^+; \frac{1}{2}$	≈ 0.3	γ , p, d, ${}^3\text{He}$, α	1, 3, 4, 5, 7, 12, 14
(18 ± 1)	$(\frac{1}{2}^+); \frac{1}{2}$	broad	γ , p, d, ${}^3\text{He}$, α	1, 3, 7
(20.0 ± 0.5)	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	≈ 5	γ , p, d, ${}^3\text{He}$, α	1, 3, 4, 7, 14
^a				

^a See also discussions in reactions 3, 4, 7, 14 and 15.

$$2. \quad {}^3\text{He}(d, n){}^4\text{Li} \quad Q_{\text{m}} = -5.324$$

This reaction has not been observed: see (1966LA04) and reaction 7 in ${}^4\text{Li}$ in (1968ME03).

$$\begin{aligned}
 3. \text{ (a) } & {}^3\text{He}(d, p){}^4\text{He} & Q_{\text{m}} &= 18.3538 & E_{\text{b}} &= 16.39 \\
 \text{ (b) } & {}^3\text{He}(d, np){}^3\text{He} & Q_{\text{m}} &= -2.22464 \\
 \text{ (c) } & {}^3\text{He}(d, 2p){}^3\text{H} & Q_{\text{m}} &= -1.4608 \\
 \text{ (d) } & {}^3\text{He}(d, 2d){}^1\text{H} & Q_{\text{m}} &= -5.4938
 \end{aligned}$$

Below 100 keV the cross section follows the simple Gamow form:

$\sigma = (18.2 \pm 10^3/E) \exp(-91E^{-1/2})$ b (E in keV) (1953JA1A, 1954AR02). The zero-energy cross section factor $S_0 = 6700$ keV · b (1964PA1A). A pronounced resonance occurs at $E_{\text{d}} = 430$ keV, $\Gamma \approx 450$ keV. The peak cross section is given as 695 ± 14 mb (1952BO68, 1955KU03): see Table 5.2. See also (1970CA1K, 1972PL02). Excitation functions for ground state protons have also been reported by (1966NE1A: $E({}^3\text{He}) = 0.39$ to 1.46 MeV), (1971GR47: $E_{\text{d}} = 2.8$ to 11.5 MeV), (1972BA30: $E_{\text{d}} = 9.4$ to 17.8 MeV) and (1972KI02: $E({}^3\text{He}) = 18.7$ to 44.1 MeV). Angular distribution and polarization measurements are summarized in Table 5.6. Below $E_{\text{d}} = 1.1$ MeV, the polarization analyzing power of the reaction is minimally affected by p-wave contributions (less than a few percent). However, at $E_{\text{d}} = 500$ keV, the analyzing power is reduced by $12 \pm 5\%$ by intervention of the s-wave $J^\pi = \frac{1}{2}^+$ channel (1966BR02, 1967MC01, 1971LE27). See also (1971RO35, 1972SI1F, 1973OH02). (1971LE13) have evaluated matrix elements for $l = 1$ at $E_{\text{d}} = 0.43$ MeV. Vector polarization effects at $E_{\text{d}} > 1$ MeV disagree with the predictions of stripping theory (1966BR02). See also (1971ZA1C). Contour maps, T_{ab} vs. θ , E_{d} for the vector and tensor analyzing powers are presented for $E_{\text{d}} = 0$ to 12 MeV by (1971GR47). See also

(1971KL02, 1973CL13). Energy dependence of the angular distributions indicate resonance-like behavior at $E_x = 16.6, 17.5, 20.0, 20.9$ and 22.4 MeV in ${}^5\text{Li}$ (1971GR47). See also (1972SE09, 1973CL13). At $E_d \approx 6$ MeV (1971KL02) find the angular distribution more complex than could be accounted for by a single ${}^2D_{3/2}$ or ${}^2D_{5/2}$ level at $E_x \approx 20$ MeV [see ${}^3\text{He}(d, d)$]. For $E_d(\text{cm}) = 7$ to 18 MeV, differential cross sections fall monotonically with no indication of states of ${}^5\text{Li}$ with $23.4 < E_x < 34$, with $\Gamma > 0.8$ MeV and with appreciable widths for decay into ${}^3\text{He} + d$ (1972KI02). See also (1972BA30). The analyzing power for ${}^3\text{He}$ polarization has been studied by (1971HU1B, 1971LE13, 1971LE27, 1971HA02): see (1971ST1J). For a discussion of excitation of ${}^4\text{He}$ excited states in reaction (a) see (1973FI04).

Reaction (b) has been studied at $E_d = 10.92$ MeV (1972NI02), 11.0 MeV (1970LA10, 1970LA1K), $E({}^3\text{He}) = 16.5$ MeV (1971LI04), 18 MeV (1965ZU02), 18.0 and 24.0 MeV (1970AS02), $E({}^3\text{He}) = 27$ MeV (1971WA18, 1973WA13) and $E_d = 27.5$ MeV (1973CH05). See also (1966LA04). The spectra show the influence of neutron-proton final state interaction. Sequential decay via excited states of ${}^4\text{He}$ is also reported: see (1970AS02, 1972NI02) and (1973FI04).

Reaction (c) has been investigated at $E_d = 10.9$ MeV (1972NI02), 11 MeV (1965TO01, 1970LA10, 1970LA1K), 15.2 and 20.2 MeV (1967NE1B, 1968NE1D), $E({}^3\text{He}) = 16.5$ MeV (1971LI04), 18 MeV (1965ZU02), 18.0 and 24.0 MeV (1970AS02), 19.9 MeV (1971GR12), $E_d = 20.7$ MeV (1970WA15), 24 to 33 MeV (1963BI14, 1964CO1A), 29.8 MeV (1970SL1A), $E({}^3\text{He}) = 27$ MeV (1971WA18, 1973WA13), 30 MeV (1973TR1H), $E_d = 39.1$ MeV (1971RU07, 1971RU1E), $E_d = 36$ MeV and $E({}^3\text{He}) = 53$ MeV (1967MO08, 1967MO1E, 1968MO10) and at $E_d = 83$ MeV (1972BA32). See also (1972SC1R, 1973SL03). The triton spectra are characterized by p-p final state interaction and by sequential decay via ${}^4\text{He}^*$, as in reaction (b). (1970LA10, 1970LA1K) have analyzed the data of (1968MO10) using all first order interactions, in PWBA. The calculations indicate that complex exchange processes cannot be ignored, and that estimates of the nucleon-nucleon scattering length from such experiments are highly ambiguous (1970LA10, 1970LA1K).

Under kinematic conditions such that two outgoing particles have small relative kinetic energy, a DWBA analysis with two-body interactions has some success in accounting for spectral shapes: see (1967HE1B, 1971CH31, 1971WA18, 1972WA22, 1973WA13, 1973SL03). Reaction (d) has been studied at $E({}^3\text{He}) = 26.8$ MeV (1972WA22).

See also (1965DA05, 1967ME14, 1967RO07, 1968FI1D, 1968GO1N, 1968HO1D, 1969ME1B, 1971FO1J, 1970HO1L, 1970RO1L, 1970WI1G, 1971AL1B, 1971WA18, 1972RO1M, 1974PR1D), (1968TO1F, 1971BA1U, 1971HA2G, 1972OH1D) and (1965DU1A, 1965HE1A, 1966BA1J, 1966CS1A, 1966DE1D, 1966DU1A, 1967DE1G, 1968DE1K, 1968HA1F, 1968TA1J, 1969HE1K, 1969SE1B, 1970BR1L, 1970KH03, 1970PO1A, 1971DE20, 1971HA1L, 1971KH03, 1971SE1G, 1972OT05, 1973GA1H, 1973GR1L, 1973LI19; theor.).

4. ${}^3\text{He}(d, d){}^3\text{He}$

$$E_b = 16.39$$

In the range $E_d = 380$ to 570 keV, the scattering cross section is consistent with s-wave formation of the $J^\pi = \frac{3}{2}^+$ state at 16.66 MeV (1954BR05). The excitation curves for $E_d = 1.96$

Table 5.6: Measurements of angular distributions and polarization in ${}^3\text{He}(d, p){}^4\text{He}$ ^a

Angular distributions to ${}^4\text{He}(0)$		Polarization ^b	
Energy (MeV)	Refs.	E_d (MeV)	Refs.
$E_d = 0.25 - 0.85$	(1970CA1K)	0.19 - 0.50	(1972OH1C)
$E_d = 2.8 - 11.5$	(1971GR47)	0.30 - 2.5	(1971RO35)
$E_d = 5.9, 7.5, 10.4, 12, 12.3, 13.7$	(1960ST25)	0.31 - 2.94	(1966BR02)
		0.34 - 0.73	(1972GA1N)
$E({}^3\text{He}) = 18.7 - 44.1$	(1972KI02)	0.37 - 1.09	(1972SI1F)
		0.43	(1971LE13, 1971LE27)
$E_d = 23.2 - 27.0$	(1964BI06)	0.45	(1970AD1C)
		0.45, 0.70, 0.98, 1.5, 2.0, 2.5	(1971HU1B)
		2	(1968SL1A)
		2.0, 2.8, 3.9, 6.0	(1973CL13)
		2 - 11.2	(1971GR15)
		2.23 - 13.0	(1971KL02)
		2.8 - 10.0	(1971LE1G)
		2.8 - 11.5	(1971GR47)
		3.0 - 12.0	(1963BR10)
		4, 6, 8, 10	(1969PL01)
		4.0 - 14.0	(1973HA51)
		4.78 - 11.88	(1971HA02, 1970HA1M)
		5.55, 7.97, 10.68	(1966WE03)
		6.0, 7.7, 10	(1968ZA1A, 1971ZA1C)
		6, 8, 10	(1970WA1L)
		6, 10	(1971HI07)
		6.49, 9.30	(1971KO09)
		6.7 - 15.8	(1973LI1L)
		9	(1973KA08)
		10, 12, 14	(1969ST1F, 1968ST1H)
		52	(1969BR22)

^a See also (1957JA37).

^b See also (1966LA04).

to 10.99 MeV show a broad resonance ($\Gamma > 1$ MeV) corresponding to $E_x = 20.0 \pm 0.5$ MeV. From the behavior of the angular distributions an assignment of ${}^2D_{3/2}$ or $({}^2D, {}^4D)_{5/2}$ is favored, if only one state is involved [see, however, ${}^7\text{Li}(p, t){}^5\text{Li}$] (1967TO02). [There is some evidence that there is more than one D-wave state in this E_x region: see reaction 3]. See also (1969DA1F, 1972AL1P). In the range $E_d(\text{cm}) = 7$ to 18 MeV differential cross sections show only a monotonic variation with energy. There is no evidence for any other resonances from $E_x = 23$ to 34 MeV (1972BA30, 1972KI02). See also (1970WI1G). Angular distributions have been measured at a number of energies from $E({}^3\text{He}) = 18.7$ to 44.1 MeV (1972KI02). Polarization measurements are reported for $E_d = 4.0$ to 11.5 MeV (1972KO07), 4 to 12 MeV (1973BA2P), 4.78 to 11.88 MeV (1970HA1M, 1971HA02), 5.7, 7.1, 8.4 and 9.4 MeV (1973LO1D), 6, 8 and 10 MeV (1969PL01), 9.9 and 11.9 MeV (1970WA32) and 10 and 12 MeV (1971DO1C). Strong variations of the angular distributions for all components of the analyzing power are observed between $E_d = 4.0$ and 8.0 MeV: it is suggested that one or more positive parity states are involved ($18.8 < E_x < 21.2$ MeV). At higher energies potential scattering is the dominant mode (1972KO07). See also (1971BA1U, 1971ST1J) and (1966BA1J, 1966BA1M, 1967HE1B, 1970BR1L, 1970HE1D, 1971HA1L, 1972TA1E, 1973CH27, 1974CH02; theor.).

5. (a) ${}^3\text{He}(t, n){}^5\text{Li}$ $Q_m = 10.13$
 (b) ${}^3\text{He}(t, np){}^4\text{He}$ $Q_m = 12.0963$

For reaction (a) see (1966LA04) and (1971KL04). For reaction (b) see (1970GR17) and ${}^6\text{Li}$.

6. (a) ${}^3\text{He}({}^3\text{He}, p){}^5\text{Li}$ $Q_m = 10.89$
 (b) ${}^3\text{He}({}^3\text{He}, 2p){}^4\text{He}$ $Q_m = 12.8601$
 (c) ${}^3\text{He}({}^3\text{He}, 3p){}^3\text{H}$ $Q_m = -6.9546$

The spectrum of protons shows a pronounced peak corresponding to ${}^5\text{Li}_{g.s.}$ superposed on a continuum (1965BA1D, 1965BA1E, 1972DE46: $E({}^3\text{He}) = 3$ to 18 MeV): the ${}^5\text{Li}$ breaks up with a strong angular asymmetry (1968BA1N). The α -spectra $E({}^3\text{He}) = 6.9, 7.9, 9.1$ MeV have been analyzed in terms of final state interactions including both p-p and p- α interactions (1972DE46). At $E({}^3\text{He}) = 43.7$ and 53.0 MeV, the spectra show a prominent peak at the high energy end whose angular distributions exhibit a pronounced diffraction pattern (1967SL01, 1968MO10). See also (1965ZU02, 1966NE1A, 1966BL02, 1967MO1E, 1968BL06) and (1973LI1M; theor.). A search for a three-proton enhancement (reaction (c)) was unsuccessful at $E({}^3\text{He}) = 44$ MeV (1968TO01) and 53 MeV (1968MO10). See also (1966LA04).

7. ${}^4\text{He}(p, p){}^4\text{He}$ $E_b = -1.97$

Table 5.7: Measurements of elastic scattering and polarization in ${}^4\text{He}(p, p){}^4\text{He}$ ^a

Elastic scattering angular distribution		Polarization	
E_p (MeV)	Refs.	E_p (MeV)	Refs.
11.157, 12.040, 13.600, 14.230	(1973JA1M)	0.2 – 0.5	(1967AD1B)
12.0, 14.2, 17.5	(1969GA12)	0.9 – 3.2	(1967BR02, 1967BR03)
19.9 – 39.8	(1972BA24)	2.1 – 4.3	(1965BL02)
20.6, 23.3, 26.1, 27.7	(1968AL1B)	2.38 – 4.46	(1966DR01)
22.2 – 24.8	(1968DA04)	4.58, 5.95, 7.89, 9.89, 11.94	(1971SC04)
25.0 – 29.2	(1971PL07)	6.00, 7.89, 9.89, 11.16	(1967JA09)
39.8 – 47.7	(1973DA1M)	7.9, 9.9, 11.9	(1971OH04)
46	(1969BU10)	10	(1968PL02)
48.8	(1967DA08)	12.0	(1972KE17)
85	(1973VO1M)	12.0, 14.2, 17.5	(1969GA12)
100	(1970GO36)	14.5	(1965RO22)
141	(1972NI04)	17.5 – 27	(1966WE03)
144	(1972JA07)	20 – 45	(1972BA24)
587	(1968BO1K, 1972BO29)	22.2 – 24.8	(1968DA04)
1000	(1967PA25)	25.0 – 29.2	(1971PL07)
		48	(1966GR08)
		70, 80	(1969PE01)
		544	(1968GO1M, 1972BO29)

^a See also Table 5.5 (1966LA04) for a summary of earlier work.

Differential cross sections and polarizations have been measured at many energies: see Table 5.5 in (1966LA04) and Table 5.7 here.

Phase shifts below $E_p = 3.2$ MeV have been determined by (1967BR03) based on polarization measurements of (1967BR02) and available differential cross sections. The results are in essential agreement with earlier phase shifts of (1949CR1A) but are considerably more precise. In this range, the s-wave phase shift is that of a hard sphere with $R = 2.48$ fm; d-wave shifts are zero. The $P_{3/2}$ phase shift shows a pronounced resonance corresponding to ${}^5\text{Li}_{\text{g.s.}}$ while the $P_{1/2}$ shift changes slowly over a range of several MeV, suggesting that the first excited state is very broad and located 5–10 MeV above the ground state: see (1959AJ76, 1966LA04) for reviews of earlier results.

From $E_p = 3$ to 18 MeV (1971SC04) have analyzed available polarization and differential cross section data to produce a set of phase shifts constrained to an analytic (effective range expansion) energy dependence. d- and f-wave phase shifts are positive and become of some importance above 8 MeV. A polarization contour plot is presented for $E_p = 1$ to 18 MeV (1971SC04). See also (1968MO26, 1971AR20, 1973AR1N). An R -matrix (single level plus background) formalism is used by (1972ST01). For $P_{1/2}$ and $P_{3/2}$ the resonance energies are given by $E_R = 8.10$ and 2.06 MeV and the reduced widths $\gamma^2 = 12.30$ and 8.02 MeV, respectively (1972ST01). See also (1973NI1B).

A resonance is observed at $E_p = 23$ MeV, corresponding to the known $\frac{3}{2}^+$ state at $E_x = 16.7$ MeV (1968AL1B, 1968DA04). An anomaly in the polarization is also observed at this energy (1966WE03, 1968DA04, 1972BA24). A further broad feature in the polarization excitation function ($\theta = 102^\circ$) is observed at $E_p = 30$ MeV ($E_x = 22$ MeV) (1972BA24). Cross sections for $23 < E_p < 45$ MeV show no further evidence of excited states (1969BU10).

An extensive phase shift analysis, using complex phases with $l \leq 4$ over a range $E_p = 20$ to 40 MeV was made by (1972PL02), using mainly polarization and differential cross section data of (1972BA24). The $D_{3/2}$ level is fit with R -matrix formalism, including background interference, with the following parameters: $E_x = 16.68$ MeV, $\gamma_p^2 = 122$ keV, $\gamma_d^2(l=0) = 1.58$ MeV (negative sign), $\gamma_d^2(l=2) = 1.58$ MeV, $\theta_p^2 = 0.014$, $\theta_d^2 = 0.765$.

The $E_p = 30$ MeV structure is not reflected in anomalous behavior of any single phase shift. Strong absorption of even partial waves may indicate broad overlapping positive-parity levels, $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$ of d + ${}^3\text{He}$ character near $E_x = 22$ MeV, but there is no unambiguous identification of excited states other than the $\frac{3}{2}^+$ state in the p- α results (1972PL02). In an analysis of data of (1971PL07) for $E_p = 25$ to 29 MeV, (1971RA27) report evidence for a $\frac{5}{2}^+$ level at $E_x = 20$ MeV [$\Gamma_{\text{el}}/\Gamma_{\text{total}} = 0.15$, $\Gamma_{\text{tot}} = 8$ MeV] and a $\frac{1}{2}^+$ level at $E_x = 18 \pm 1$ MeV.

Other phase-shift analyses are reported by (1971AR20, 1973AR1N: 0 to 23 MeV), (1972ST01: 0.3 to 20 MeV), (1967BR02, 1967BR03: 0.9 to 3.2 MeV), (1966WE03: 14.3 to 31.0 MeV), (1967DA08: 29, 40, 48 MeV) and (1969PE01: 63, 70, 80 and 94 MeV). See also (1968PL02, 1973AR1P, 1973HO1U).

Elastic scattering at $E_p = 587$ MeV (1972BO29) and at 1 GeV (1967PA25) reflect the mass distribution and multibody correlations within the α -particle.

Alpha-proton bremsstrahlung is observed at $E_p = 7.0$ to 12.0 MeV (1971WO07, 1972WO1E), and at $E_\alpha = 20$ MeV (1967BO1B). See also (1972AN1P). Cross section measurements have

also been carried out at $E_p = 23.4$ to 48.5 MeV (1973DA1N, 1973DA1P), 180 to 560 MeV (1972SC1M) and at 1 GeV (1967IG1A). For inelastic scattering to excited states of ^4He , see (1973FI04). For pion production, see (1972GU1D).

See also (1965JA1B, 1965KU13, 1966BL1B, 1967AU1A, 1967CA1E, 1968KO1F, 1968PA1J, 1969KO1M, 1970LI06, 1970RO1M, 1973CO1Y, 1973EK1A), (1966BA1K, 1966HA1F, 1966RO1B, 1970BO1P, 1971HA2G, 1971ST1J) and (1965HO1A, 1965HO1B, 1966RA1B, 1966TO04, 1967BA1F, 1967BA1G, 1967CZ1A, 1967SA1C, 1967TA1B, 1968BA1K, 1968CH35, 1968FR1F, 1968GL1A, 1968SA1B, 1968SC1D, 1968SE1A, 1968TA1E, 1968BA1U, 1969CO1E, 1969CR1A, 1969GE1A, 1969HE15, 1969LE03, 1968TH1C, 1970BA1F, 1970HE1D, 1970KU1C, 1970RE1A, 1970SA01, 1970TH1D, 1971BA1Y, 1971BA61, 1971BA74, 1971FE10, 1971HA1L, 1971IK01, 1971KR1A, 1971LO24, 1971MU1H, 1971PI04, 1971PL06, 1971RA15, 1971TH09, 1972BB19, 1972GR06, 1972IK01, 1972TH06, 1973AD01, 1973CL01, 1973FE1H, 1973LA14, 1973LY02, 1973PL02, 1973SA09, 1973UL1C, 1973YO1F, 1974CH02; theor.).

8. (a) $^4\text{He}(p, d)^3\text{He}$	$Q_m = -18.3538$	$E_b = -1.97$
(b) $^4\text{He}(p, pn)^3\text{He}$	$Q_m = -20.5785$	
(c) $^4\text{He}(p, 2p)^3\text{H}$	$Q_m = -19.8147$	

Angular distributions of ^3He ions (reaction (a)) have been measured at $E_p = 27.9$ MeV (1957WI22), 31 MeV (1953BE14, 1964BU1B), 46.8 MeV (1969RO24, 1969RO1L), 49.5 MeV (1969HA16, 1970HA36), 53 MeV (1964CA1B), 55 MeV (1964HA13, 1964HA1P, 1964HA49), 94 MeV (1958SE74) and 155.4 MeV (1967BE35, 1970BE49). See also (1973VO1M). The excitation function shows no indication of resonances for $E_p = 38.5$ to 44.6 MeV (1969BU10). Polarization measurements have been carried out at $E_p = 32, 40, 50$ and 52.5 MeV (1973SA1W) and 55 and 63 MeV (1966BO1A). At $E_p = 141$ MeV the total reaction cross section (not including elastic scattering) is 79.5 ± 2.0 mb (1972NI04).

Reactions (b) and (c) have been studied at $E_p = 46.8$ MeV (1969RO24, 1969RO1L, 1970RO17, 1970SA01), 49.5 MeV (1969HA16, 1970HA36) and 156 MeV (1967BE35, 1970BE49). See also (1973KI1L, 1973RO1Z). Angular distributions have been measured in the region of the final state interactions. The t and ^3He spectra show peaks due to sequential reaction processes via continuum resonance states in ^4He : see (1973FI04). See also (1967AU1A, 1968PA1J) and (1969DO1G, 1969NA1F, 1973HA46, 1973JU02; theor.). For $^4\text{He}(p, 2N + \chi\pi)$ see (1965KU1C).

9. (a) $^4\text{He}(d, n)^5\text{Li}$	$Q_m = -4.19$
(b) $^4\text{He}(d, np)^4\text{He}$	$Q_m = -2.22464$

Reaction (b) has been studied at $E_d = 14.2$ MeV (1967FU1D), $E_\alpha = 29.2$ MeV (1968TA11), 38.8 MeV (1971LE02), 42 MeV (1967WA08, 1968WA01) and at $E_\alpha = 18.0$ and 24.0 MeV

(1970AS02). See also (1971CH1T). The data indicate that at 42 MeV, direct breakup, with quasi-free α -p scattering taking place and the n acting as a spectator, is at least as important a mechanism as the final state interaction in the ${}^5\text{Li}$ ground state (1967WA08, 1968WA01): see ${}^5\text{He}$. See also (1967NA1D, 1969NA09, 1972NA1D; theor.) and (1966LA04) for the earlier references.

$$10. {}^4\text{He}({}^3\text{He}, \text{np}){}^5\text{Li} \quad Q_m = -9.68$$

See (1973HA50).

$$11. {}^6\text{Li}(\gamma, \text{n}){}^5\text{Li} \quad Q_m = -5.66$$

See ${}^6\text{Li}$ and (1965BA16).

$$12. \text{(a) } {}^6\text{Li}(\text{p}, \text{d}){}^5\text{Li} \quad Q_m = -3.44$$

$$\text{(b) } {}^6\text{Li}(\text{p}, \text{pd}){}^4\text{He} \quad Q_m = -1.4737$$

Deuteron groups are observed to ${}^5\text{Li}^*(0, 16.7)$. Angular distributions have been measured at $E_p = 18.6$ MeV (1955LI09; d_0), 33.6 MeV (1967KU10, 1970KU1D; d_0, d_2), 100 MeV (1969LI02; d_0) and 156 MeV (1968BE72, 1969BA05, 1969TO1A; d_0, d_2). See (1969TO1A) for spectroscopic factors. In addition, the excitation of a state with $E_x = 3.72 \pm 0.10$, $\Gamma = 6.3 \pm 0.5$ MeV is reported by (1966SE1C). [See, however, (1967KU10, 1970KU1D)]. (1969BA05) confirm the population of the first excited state. See also (1966SE1C, 1972AZ03) and (1968JA1D; theor.).

At $E_p = 9$ and 10 MeV (1968VA02) and at $E_p = 45$ MeV (1971BR12), the p- α final state interaction corresponding to ${}^5\text{Li}(0)$ is observed.

$$13. \text{(a) } {}^6\text{Li}(\text{d}, \text{t}){}^5\text{Li} \quad Q_m = 0.59$$

$$\text{(b) } {}^6\text{Li}(\text{d}, \text{pt}){}^4\text{He} \quad Q_m = 2.5592$$

Angular distributions of the tritons to the ground state of ${}^5\text{Li}$ have been measured at $E_d = 15$ and 20 MeV (1959VL24, 1960HA14). See also (1971IN1C). For reaction (b) see (1968LE15).

$$14. \text{(a) } {}^6\text{Li}({}^3\text{He}, \alpha){}^5\text{Li} \quad Q_m = 14.91$$

$$\text{(b) } {}^6\text{Li}({}^3\text{He}, \text{p}\alpha){}^4\text{He} \quad Q_m = 16.880$$

At $E(^3\text{He}) = 25.5$ MeV, the spectra show $^5\text{Li}^*(0, 16.7)$ and two broad peaks at $E_x \approx 19.8$ and 22.7 MeV with $\Gamma_{\text{cm}} = 2$ and 1 MeV, respectively (1972BA30).

Cylindrical asymmetry observed in the breakup of $^5\text{Li}(0)$ is attributed to the short lifetime of the ^5Li intermediate state and to the memory retained by the proton of its localization at the time of formation of ^5Li (1967RE03, 1968RE10). The first excited state of ^5Li also appears to be involved (1972TH08, 1972TH1B). See also (1964MA57, 1966LO10, 1967HO1C, 1968VI03, 1969VI05, 1970GA1G), (1966LA04), (1970DE41, 1972TH04; theor.) and ^8Be .

$$15. \text{ (a) } ^7\text{Li}(\text{p}, \text{t})^5\text{Li} \quad Q_{\text{m}} = -4.43$$

$$\text{ (b) } ^7\text{Li}(\text{p}, \text{p}\alpha)^3\text{H} \quad Q_{\text{m}} = -2.467$$

At $E_{\text{p}} = 43.7$ MeV, a triton group is observed to $^5\text{Li}(0)$ ($\Gamma = 1.55 \pm 0.15$ MeV): the angular distribution is consistent with a substantial mixing of $L = 0$ and 2 transfer. There is some evidence for a very broad excited state between $E_x = 2$ and 5 MeV. $^5\text{Li}^*(16.7, 20.0)$ were not observed. The formation of $^5\text{Li}^*(16.7)$ ($^4S_{3/2}$) would be S -forbidden: the absence of $^5\text{Li}^*(20.0)$ would indicate that this state(s) is also of quartet character [see reaction 19 in ^5He] (1966CE05). See also (1966BA1L, 1967PO1C). Weak, broad states at $E_x = 22.0 \pm 0.5$ MeV and 25.0 ± 0.5 MeV and possibly 34 MeV are reported by (1968MC02) in a coincidence experiment in which 3- and 4-particle breakup was analyzed. The t_0 angular distribution has also been studied at $E_{\text{p}} = 16.6$ MeV (1965OG03) and 30.3 MeV (1969DE04).

For reaction (b) see (1967JO1C).

$$16. \text{ } ^{10}\text{B}(^3\text{He}, \text{p}\alpha)^4\text{He}^4\text{He} \quad Q_{\text{m}} = 12.420$$

At $E(^3\text{He}) = 2.45$ and 6.00 MeV the reaction proceeds in part via the first two states of ^5Li (1966WA16).

$$17. \text{ } ^{12}\text{C}(\text{p}, 2\alpha)^5\text{Li} \quad Q_{\text{m}} = -9.24$$

Not observed: see (1972MA62).

^5Be (Fig. 3)

The absence of any group structure in the neutron spectrum in the reaction $^3\text{He}(^3\text{He}, \text{n})^5\text{Be}$ at $E(^3\text{He}) = 18.0$ to 26.0 MeV indicates that $^5\text{Be}(0)$ is at least 4.2 MeV unstable with respect to $^3\text{He} + 2\text{p}$ [$(M - A) > 33.7$ MeV]. With Coulomb corrections adjusted to match the 16.7 MeV states of ^5He - ^5Li , this observation places the first $T = \frac{3}{2}$ level in these nuclei above $E_x = 21.4$ MeV (1967AD05). [With the “conventional” correction (1966LA04, p.2) $E_x > 20.3$ MeV].

See also (1968CE1A, 1970WA1G).

References

(Closed December 31, 1973)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1949CR1A Critchfield and Dodder, Phys. Rev. 76 (1949) 602
- 1952AR1A Argo, Taschek, Agnew, Hemmendinger and Leland, Phys. Rev. 87 (1952) 612
- 1952BO68 T.W. Bonner, J.P. Conner and A.B. Lillie, Phys. Rev. 88 (1952) 473
- 1952CO35 J.P. Conner, T.W. Bonner and J.R. Smith, Phys. Rev. 88 (1952) 468
- 1952DO30 D.C. Dodder and J.L. Gammel, Phys. Rev. 88 (1952) 520
- 1953BE14 J. Benveniste and B. Cork, Phys. Rev. 89 (1953) 422
- 1953JA1A Jarvis and Roaf, Proc. Roy. Soc. 218 (1953) 432
- 1954AR02 W.A. Arnold, J.A. Phillips, G.A. Sawyer, E.J. Stovall, Jr. and J.L. Tuck, Phys. Rev. 93 (1954) 483
- 1954BL89 J.M. Blair, N.M. Hintz and D.M. Van Patter, Phys. Rev. 96 (1954) 1023
- 1954BR05 R.J.S. Brown, K.F. Famularo, H.D. Holmgren, D. Rankin and T.F. Stratton, Phys. Rev. 96 (1954) 80
- 1954BU06 W.H. Burke, J.R. Risser and G.C. Phillips, Phys. Rev. 93 (1954) 188
- 1955KU03 W.E. Kunz, Phys. Rev. 97 (1955) 456
- 1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
- 1955LI09 J.G. Likely, Phys. Rev. 98 (1955) 1538
- 1957JA37 N. Jarmie and J.D. Seagrave, Eds., LA-2014 (1957)
- 1957WI22 A.F. Wickersham, Jr., Phys. Rev. 107 (1957) 1050
- 1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
- 1958SE74 W. Selove and J.M. Teem, Phys. Rev. 112 (1958) 1658
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959BO54 T.W. Bonner, F.W. Prosser, Jr. and J. Slattery, Phys. Rev. 115 (1959) 398
- 1959VL24 N.A. Vlasov and A.A. Ogloblin, Zh. Eksp. Teor. Fiz. 37 (1959) 54; JETP (Sov. Phys.) 10 (1960) 39
- 1960HA14 E.W. Hamburger and J.R. Cameron, Phys. Rev. 117 (1960) 781

- 1960HU1A Hughes, Magurno and Brussel, BNL 325 (2nd Edition) Suppl. 1 (1960)
- 1960ST25 L. Stewart, J.E. Brolley, Jr. and L. Rosen, Phys. Rev. 119 (1960) 1649
- 1960VA04 F.J. Vaughn, W.L. Imhof, R.G. Johnson and M. Walt, Phys. Rev. 118 (1960) 683
- 1961PE13 R.B. Perkins and J.E. Simmons, Phys. Rev. 124 (1961) 1153
- 1961RU1A Rudin, Streibel, Baumgartner, Brown and Huber, Helv. Phys. Acta 34 (1961) 58
- 1961TR05 I.S. Trostin, V.A. Smotryaev and I.I. Levintov, Zh. Eksp. Teor. Fiz. 41 (1961) 725; JETP (Sov. Phys.) 14 (1962) 524
- 1962SE09 F. Seiler, E. Baumgartner, W. Haeberli, P. Huber and H.R. Striebel, Helv. Phys. Acta 35 (1962) 385
- 1963BI14 O.M. Bilaniuk and R.J. Slobodrian, Phys. Lett. 7 (1963) 77
- 1963BR10 R.I. Brown and W. Haeberli, Phys. Rev. 130 (1963) 1163
- 1963BU07 W. Buss, H. Waffler and B. Ziegler, Phys. Lett. 4 (1963) 198
- 1963MA29 T.H. May, R.L. Walter and H.H. Barschall, Nucl. Phys. 45 (1963) 17
- 1964AL1E Alekseev, Arifkhanov, Vlasov, Davidiv and Samoilov, Zh. Eksp. Teor. Fiz. 47 (1964) 433; JETP (Sov. Phys.) 20 (1965) 287
- 1964BI06 O.M. Bilaniuk and R.J. Slobodrian, Nucl. Phys. 50 (1964) 585
- 1964BU1B Bunch, Forster and Kim, Nucl. Phys. 53 (1964) 241
- 1964CA1B Cairns, Griffith, Lush, Metheringham and Thomas, Nucl. Phys. 60 (1964) 369
- 1964CO1A Conzett, Shield, Slobodrian and Yamabe, Phys. Rev. Lett. 13 (1964) 625
- 1964DE1B De Facio, UCRL 7961 T (1964)
- 1964FE01 P. Fessenden and D.R. Maxson, Phys. Rev. 133 (1964) B71
- 1964HA13 S. Hayakawa, N. Horikawa, R. Kajikawa, K. Kikuchi, H. Kobayakawa, K. Matsuda, S. Nagata and Y. Sumi, Phys. Lett. 8 (1964) 333
- 1964HA1P S. Hayakawa, N. Horikawa, R. Kajikawa, K. Kikuchi, H. Kobayakawa, K. Matsuda, S. Nagata and Y. Sumi, Phys. Lett. 8 (1964) 330
- 1964HA49 S. Hayakawa, N. Horikawa, R. Kajikawa, K. Kikuchi, H. Kobayakawa, K. Matsuda, S. Nagata and Y. Sumi, J. Phys. Soc. Jpn. 19 (1964) 2004
- 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed. W.H. Johnson, Jr. (Springer Verlag, Vienna; 1964) p. 305
- 1964PA1A Parker, Bahcall and Fowler, Astrophys. J. 139 (1964) 602
- 1964PE14 R.B. Perkins and C. Glashauser, Nucl. Phys. 60 (1964) 433
- 1964SH1A Shamu and Jenkin, Phys. Rev. 135 (1964) B99

- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Ed., Suppt. 2, Vol. 1 (1964)
- 1964WA22 R.L. Walter, W. Benenson, T.H. May and C.A. Kelsey, Nucl. Phys. 59 (1964) 235
- 1965AR04 P.E. Argan, L. Meneghetti and S. Vitale, Nuovo Cim. 38 (1965) 1489
- 1965AR08 P.E. Argan, G.C. Mantovani, P. Marazzini, A. Piazzoli and D. Scannicchio, Suppl. Nuovo Cim. 3 (1965) 245
- 1965AR10 U.R. Arifkhanov, N.A. Vlasov, V.V. Davydov and L.N. Samoilo, Yad. Fiz. 2 (1965) 239; Sov. J. Nucl. Phys. 2 (1966) 170
- 1965AS06 P.A. Assimakopoulos, N.H. Gangas, S. Kossionides and N.R.C. Democritus, Phys. Lett. 19 (1965) 316
- 1965BA16 E.B. Bazhanov, A.P. Komar, A.V. Kulikov and E.D. Makhnovsky, Nucl. Phys. 68 (1965) 191
- 1965BA1D Bacher and Tombrello, Revs. Mod. Phys. 37 (1965) 433
- 1965BA1E Bacher and Tombrello, Bull. Amer. Phys. Soc. 10 (1965) 693
- 1965BE1C Berkowitz, Carlson and Carpenter, Bull. Amer. Phys. Soc. 10 (1965) 694
- 1965BL02 R.A. Blue and W. Haeberli, Phys. Rev. 137 (1965) B284
- 1965BO13 F. Boreli, V. Lazarevic and N. Radisic, Nucl. Phys. 66 (1965) 301
- 1965CH12 G. Charpak, G. Gregoire, L. Massonnet, J. Saudinos, J. Favier, M. Gusakow and M. Jean, Phys. Lett. 16 (1965) 54
- 1965CH15 J. Christiansen, F.W. Busser, F. Niebergall and G. Sohngen, Nucl. Phys. 67 (1965) 133
- 1965CO1D Cohen, Kanaris, Margulies and Rosen, Phys. Lett. 14 (1965) 242
- 1965CO1E Cortellessa, Suppl. Nuovo Cim. 3 (1965) 820
- 1965DA05 S.E. Darden and A.J. Froelich, Phys. Rev. 140 (1965) B69
- 1965DU1A Dumtrescu and Sandalescu, Rev. Phys. 10 (1965) 77
- 1965FL1A Fleischmann, Fritsch and Graw, Z. F. Phys. 186 (1965) 485
- 1965GO1B Goldfarb and Hug, Helv. Phys. Acta 38 (1965) 541
- 1965HE1A Henley, Richards and Yu, Phys. Lett. 15 (1965) 331
- 1965HI1B Hiura and Shimodaya, Prog. Theor. Phys. 34 (1965) 861
- 1965HO1A Holdeman and Thaler, Phys. Rev. 139 (1965) B1186
- 1965HO1B Horikawa and Kanada, J. Phys. Soc. Jpn. 20 (1965) 1758
- 1965JA1B Jarmie et al, Bull. Amer. Phys. Soc. 10 (1965) 1204
- 1965KU13 V.N. Kuzmin and R.M. Yakovlev, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 1242; Bull. Acad. Sci. USSR Phys. Ser. 29 (1966) 1245

- 1965KU1B Kudo, Prog. Theor. Phys. 34 (1965) 942
- 1965KU1C Kuzmin and Yakovlev, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 236
- 1965NE1B Neudachin and Smirnov, At. Energy Rev. 3 (1965) 3-157
- 1965OG03 A.A. Ogloblin and V.I. Chuev, Yad. Fiz. 2 (1965) 670; Sov. J. Nucl. Phys. 2 (1966) 480
- 1965RO15 J.C. Roynette, C. Ruhla, M. Arditi, J.C. Jacmart and M. Riou, Phys. Lett. 19 (1965) 497
- 1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, Ann. Phys. 34 (1965) 96
- 1965TO01 T.A. Tombrello and A.D. Bacher, Phys. Lett. 17 (1965) 37
- 1965TR01 W. Trachslin, H. Burgisser, P. Huber, G. Michel and H.R. Striebel, Helv. Phys. Acta 38 (1965) 523
- 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331
- 1965WO03 C. Wong, J.D. Anderson and J.W. McClure, Nucl. Phys. 71 (1965) 106
- 1965YA02 T. Yanabu, S. Yamashita, K. Takimoto and K. Ogino, J. Phys. Soc. Jpn. 20 (1965) 1303
- 1965ZU02 R.W. Zurmuhle, Nucl. Phys. 72 (1965) 225
- 1966AS04 P.A. Assimakopoulos, N.H. Gangas and S. Kossionides, Nucl. Phys. 81 (1966) 305
- 1966BA1H Baumgartner et al, Phys. Rev. Lett. 16 (1966) 105
- 1966BA1J Balashko, Proc. Conf. Nucl. Rxns, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 398
- 1966BA1K Barschall, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 393
- 1966BA1L Bachelier et al, J. Phys. C1-70 (1966)
- 1966BA1M Balashko, Trudy of the P.N. Lebedev Phys., Vol. 33 (1965) 51 (Transl. by Consultant Bureau, NY, 1966)
- 1966BE1B Berggren and Tyren, Ann. Rev. Nucl. Sci. 16 (1966) 153
- 1966BI1A Bilwes, Bourotte and Magnac-Valette, J. Phys. C1-68 (1966)
- 1966BL02 E.W. Blackmore and J.B. Warren, Phys. Rev. Lett. 16 (1966) 520
- 1966BL1B Blue and Stout, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 149
- 1966BO1A Boschitz, Chabre, Conzett and Slobodrian, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 354
- 1966BR02 L. Brown, H.A. Christ and H. Rudin, Nucl. Phys. 79 (1966) 459

- 1966BR1E Brunning et al, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 146
- 1966BR1F Breunlich, Acta Phys. Aust. 23 (1966) 517
- 1966BU09 F.W. Busser, F. Niebergall, G. Sohngen and J. Christiansen, Nucl. Phys. 88 (1966) 593
- 1966CE05 J. Cerny, C. Detraz and R.H. Pehl, Phys. Rev. 152 (1966) 950
- 1966CS1A Csonka, Moravcsik and Scadron, Phys. Rev. 143 (1966) 775
- 1966DE1D De Facio, Umberjee and Gammel, Phys. Lett. 21 (1966) 666
- 1966DE1E Detraz, J. Phys. C1-64 (1966)
- 1966DE1F De Facio, Umberjee and Gammel, Phys. Rev. 151 (1966) 819
- 1966DR01 L. Drigo, C. Manduchi, G.C. Nardelli, M.T. Russo-Manduchi, G. Tornielli and G. Zannoni, Nuovo Cim. B42 (1966) 363
- 1966DU1A Duck, Nucl. Phys. 80 (1966) 617
- 1966FR1B Fraser and Spicer, Aust. J. Phys. 19 (1966) 893
- 1966GR08 T.C. Griffith, D.C. Imrie, G.J. Lush and L.A. Robbins, Phys. Rev. 146 (1966) 626
- 1966HA1F Hanna, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 280
- 1966HE1B Henneberg, Z. Phys. 193 (1966) 23
- 1966HI1A Hiura and Shimodaya, Prog. Theor. Phys. Jpn. 36 (1966) 977
- 1966HO06 T.D. Hoang and R.C. Smith, Nucl. Phys. 83 (1966) 9
- 1966HO07 B. Hoop, Jr. and H.H. Barschall, Nucl. Phys. 83 (1966) 65
- 1966HU16 A. Huq, Helv. Phys. Acta 39 (1966) 507
- 1966JE01 R.W. Jewell, W. John, J.E. Sherwood and D.H. White, Phys. Rev. 142 (1966) 687
- 1966KO19 A.P. Kobzev, V.I. Salatskii and S.A. Telezhnikov, Yad. Fiz. 3 (1966) 1060; Sov. J. Nucl. Phys. 3 (1966) 774
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966LO10 T. Lorenz, Z. Naturforsch. A21 (1966) 1196
- 1966ME03 D.F. Measday, Phys. Lett. 21 (1966) 62
- 1966ME14 D.F. Measday and J.N. Palmieri, Nucl. Phys. 85 (1966) 129
- 1966MI09 C. Milone and R. Potenza, Nucl. Phys. 84 (1966) 25
- 1966NE1A Neng-Ming et al, Yad. Fiz. 3 (1966) 1064
- 1966PE1C W.A. Pearce and P. Swan, Nucl. Phys. 78 (1966) 433

- 1966PO1D Potenza, Proc. Int. School Enrico Fermi, Course 36, Ed. C. Bloch (Academic Press, 1966) 584
- 1966RA1B Rahman, Khan and Sen Gupta, Nuovo Cim. 44 (1966) 36
- 1966RO1B Rosen, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 253
- 1966SE1C Seifert, NYO GEN 72 48 (1966)
- 1966TO04 T.A. Tombrello, Phys. Lett. 23 (1966) 134
- 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692
- 1966VA1A Vasilev et al, At. Energ. 20 (1966) 341
- 1966WA16 M.A. Waggoner, J.E. Etter, H.D. Holmgren and C. Moazed, Nucl. Phys. 88 (1966) 81
- 1966WE03 W.G. Weitkamp and W. Haeberli, Nucl. Phys. 83 (1966) 46
- 1966WE1B Weinberg, Antwerp 1965 Neutron Conf. (Noth Holland, 1966) 37
- 1967AD05 E.G. Adelberger, A.B. McDonald, T.A. Tombrello, F.S. Dietrich and A.V. Nero, Phys. Lett. B25 (1967) 595
- 1967AD1B Adyasevich, Antonenko, Polunin and Fomenko, Yad. Fiz. 5 (1967) 933
- 1967AU1A Audouze, Epherre and Reeves, High Energy Nucl. Rxns in Astrophys., Ed. B.S.P. Shen (W.A. Benjamin, 1967) 255
- 1967BA1E Bang, Zelemskaya, Magzumov and Neudachin, Sov. J. Nucl. Phys. 4 (1967) 688
- 1967BA1F Baz, Sov. J. Nucl. Phys. 4 (1967) 182
- 1967BA1G R.H. Bassel and C. Wilkin, Phys. Rev. Lett. 18 (1967) 871
- 1967BE13 M. Berrada, J.-P. Laugier, C. Lemeille, N. Saunier and L. Marquez, J. Phys. (Paris) 28 (1967) 135
- 1967BE1G H.C. Benoehr and K. Wildermuth, Phys. Lett. B26 (1967) 68
- 1967BE35 M. Bernas, J.K. Lee, D. Bachelier, I. Brissaud, C. Detraz, P. Radvanyi and M. Roy, Phys. Lett. B25 (1967) 260
- 1967BO1B Boyd et al, Bull. Amer. Phys. Soc. 12 (1967) 465
- 1967BO1C M. Bouten, P. Van Leuven, H. Depuydt and L. Schotsmans, Nucl. Phys. A100 (1967) 90
- 1967BO1D Bodart and Deconninck, Ann. Soc. Sci. Bruxelles 81 (1967) 137
- 1967BR02 L. Brown and W. Trachslin, Nucl. Phys. A90 (1967) 334
- 1967BR03 L. Brown, W. Haeberli and W. Trachslin, Nucl. Phys. A90 (1967) 339
- 1967BU11 W. Busse, J. Christiansen, D. Hilscher, U. Morfeld, J.A. Scheer and W.U. Schroder, Nucl. Phys. A100 (1967) 490

1967CA1E Cameron et al, Bull. Amer. Phys. Soc. 12 (1967) 1140
1967CZ1A W. Czyz and L. Lesniak, Phys. Lett. B24 (1967) 227
1967DA08 B.W. Davies, M.K. Craddock, R.C. Hanna, Z.J. Moroz and L.P. Robertson, Nucl. Phys. A97 (1967) 241
1967DE1G B. Defacio, R.K. Umerjee and J L. Gammel, Phys. Lett. B25 (1967) 449
1967EL03 J.P. Elliott, H.A. Mavromatis and E.A. Sanderson, Phys. Lett. B24 (1967) 358
1967FL12 Y. Flamant, Y. Chanut and R. Ballini, J. Phys. (Paris) 28 (1967) 622
1967FU1D Fukunga et al, J. Phys. Soc. Jpn. 22 (1967) 28
1967HE1B E.M. Henley, F.C. Richards and D.U.L. Yu, Nucl. Phys. A103 (1967) 361
1967HO1C Holmgren, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 213
1967HO1D Hoop and Huber, Helv. Phys. Acta 40 (1967) 710
1967IG1A G.J. Igo, J.L. Friedes, H. Palevsky, R. Sutter, G. Bennett, W.D. Simpson, D.M. Corley and R.L. Stearns, Nucl. Phys. B3 (1967) 181
1967JA07 N. Jarmie, R.H. Stokes, G.G. Ohlsen and R.W. Newsome, Jr., Phys. Rev. 161 (1967) 1050
1967JA09 M.F. Jahns and E.M. Bernstein, Phys. Rev. 162 (1967) 871
1967JE01 H. Jeremie, P. Martin and A. Calamand, Nucl. Phys. A105 (1967) 689
1967JO1B Joseph et al, Bull. Amer. Phys. Soc. 12 (1967) 189
1967JO1C Joseph et al, Bull. Amer. Phys. Soc. 12 (1967) 465
1967KO1B Kolybasov and Smorodinskaya, Yad. Fiz. 5 (1967) 777
1967KU10 L.A. Kull, Phys. Rev. 163 (1967) 1066
1967MC01 L.C. McIntyre and W. Haeberli, Nucl. Phys. A91 (1967) 369
1967ME11 D.F. Measday and J.N. Palmieri, Phys. Rev. 161 (1967) 1071
1967ME14 H. Meiner, E. Baumgartner, S.E. Darden, P. Huber and G.R. Plattner, Helv. Phys. Acta 40 (1967) 483
1967ME1C Meboniya, Bull. Moscow Univ., Phys. Astron. 1 (1967) 114
1967MO08 B.J. Morton, E.E. Gross, J.J. Malanify and A. Zucker, Phys. Rev. Lett. 18 (1967) 1007
1967MO1E Morton, ORNL TM 1914 (1967)
1967NA1D H. Nakamura, Phys. Lett. B24 (1967) 509
1967NE1B Newsome, Bull. Amer. Phys. Soc. 12 (1967) 17
1967OH1A G.G. Ohlsen, Phys. Rev. 164 (1967) 1268

- 1967PA25 H. Palevsky, J.L. Friedes, R.J. Sutter, G.W. Bennett, G.J. Igo, W.D. Simpson, G.C. Phillips, D.M. Corley, N.S. Wall, R.L. Stearns et al, Phys. Rev. Lett. 18 (1967) 1200
- 1967PO1C Portner and Moore, Bull. Amer. Phys. Soc. 12 (1967) 632
- 1967RE03 M.A. Reimann, P.W. Martin and E.W. Vogt, Phys. Rev. Lett. 18 (1967) 246
- 1967RO06 J.C. Roynette, M. Arditi, J.C. Jacmart, F. Mazloum, M. Riou and C. Ruhla, Nucl. Phys. A95 (1967) 545
- 1967RO07 D.C. Robinson, Nucl. Phys. A95 (1967) 663
- 1967SA1C Sannes and Stairs, Can. J. Phys. 45 (1967) 1497
- 1967SC1E Schenter, Bull. Amer. Phys. Soc. 12 (1967) 11
- 1967SH1C Shapiro, Proc. Prob. Symp. on Nucl. Phys., Tbilisi, April 1967 (Moscow, 1967) 307
- 1967SL01 R.J. Slobodrian, J.S.C. McKee, W.F. Tivol, D.J. Clark and T.A. Tombrello, Phys. Lett. B25 (1967) 19
- 1967SW1B Swan, Private Communication (1967)
- 1967TA1B B. Tatischeff, Nucl. Phys. A98 (1967) 384
- 1967TO02 T.A. Tombrello, R.J. Spiger and A.D. Bacher, Phys. Rev. 154 (1967) 935
- 1967TR05 A. Trier and W. Haerberli, Phys. Rev. Lett. 18 (1967) 915
- 1967VA11 V. Valkovic, W.R. Jackson, Y.S. Chen, S.T. Emerson and G.C. Phillips, Nucl. Phys. A96 (1967) 241
- 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, Nucl. Phys. A98 (1967) 305
- 1967WA08 R.E. Warner and R.W. Bercaw, Phys. Lett. B24 (1967) 517
- 1967WI1C Wise, Knowles and Bunch, Bull. Amer. Phys. Soc. 12 (1967) 633
- 1967YA1A Yazaki, Prog. Theor. Phys. 37 (1967) 353
- 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
- 1968AL1B Allison and Smythe, Nucl. Phys. A121 (1968) 97
- 1968BA1K R.H. Bassel and C. Wilkin, Phys. Rev. 174 (1968) 1179
- 1968BA1N Bacher and Tombrello, Bull. Amer. Phys. Soc. 13 (1968) 564
- 1968BA1U Barashenkov and Eliseev, Joint Inst. Nucl. Res., Lab. Theor. Phys., USSR, Rept. No. P2 4333 (1969)
- 1968BE05 A.F. Behof, T.H. May and W.I. McGarry, Nucl. Phys. A108 (1968) 250
- 1968BE72 M. Bernas and D. Smet, Ann. Phys. (Paris) 3 (1968) 213
- 1968BL06 E.W. Blackmore and J.B. Warren, Can. J. Phys. 46 (1968) 233
- 1968BO1K E.T. Boschitz, W.K. Roberts, J.S. Vincent, K. Gotow, P.C. Gugelot, C.F. Perdrisat and L.W. Swenson, Phys. Rev. Lett. 20 (1968) 1116

- 1968BO1M Bonbright et al, Bull. Amer. Phys. Soc. 13 (1968) 1367
- 1968BO32 N.E. Booth, A. Beretvas, R.E.P. Davis, C. Dolnick, R.E. Hill, M. Raymond and D. Sherden, Nucl. Phys. A119 (1968) 233
- 1968BO35 D.C. Bonar, C.W. Drake, R.D. Headrick and V.W. Hughes, Phys. Rev. 174 (1968) 1200
- 1968BU09 W. Buss, W. Del Bianco, H. Waffler and B. Ziegler, Nucl. Phys. A112 (1968) 47
- 1968CE1A Cerny, Ann. Rev. Nucl. Sci. 18 (1968) 27
- 1968CH35 J.S. Chalmers and A.M. Saperstein, Phys. Rev. 168 (1968) 1145; Addendum Phys. Rev. C2 (1970) 2034
- 1968DA04 P. Darriulat, D. Garreta, A. Tarrats and J. Testoni, Nucl. Phys. A108 (1968) 316
- 1968DE14 W. Del Bianco, F. Lemire, R.J.A. Levesque and J.M. Poutissou, Can. J. Phys. 46 (1968) 1585
- 1968DE1K Del Alba, Jaidar and Martinez Negrete, Rev. Mex. Fis. 17 (1968) 143
- 1968EL1A Elliott, 3rd Symp. on the Struct. of Low-Medium Mass Nucl., Ed. Davidson (Univ. Press of Kansas, 1968) 48
- 1968EN1A J. Engler, K. Horn, J. Konig, F. Monnig, P. Schludecker, H. Schopper, P. Sievers, H. Ullrich and K. Runge, Phys. Lett. B28 (1968) 64
- 1968FI1D Findley, Baker, Carter and Hatfield, Bull. Amer. Phys. Soc. 13 (1968) 558
- 1968FR1F V. Franco, Phys. Rev. Lett. 21 (1968) 1360
- 1968GA1E Gagne, Lambert and Teado, Bull. Amer. Phys. Soc. 13 (1968) 698
- 1968GL1A Glauber, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 41, 60
- 1968GO01 P. Goldhammer, J.R. Hill and J. Nachamkin, Nucl. Phys. A106 (1968) 62
- 1968GO1M K. Gotow, E.T. Boschitz, W.K. Roberts, J.S. Vincent, P.C. Gugelot, C.F. Perdrisat and L.W. Swenson, Phys. Rev. Lett. 21 (1968) 1816
- 1968GO1N Gofman et al, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 690
- 1968HA1F R.M. Haybron, Nucl. Phys. A112 (1968) 594
- 1968HE16 G. Hentschel, G. Mack and G. Mertens, Z. Naturforsch. A23 (1968) 1401
- 1968HE1D R.C. Herndon and Y.C. Tang, Phys. Rev. 165 (1968) 1093
- 1968HO1D Hojvat and Jones, Nucl. Instrum. Meth. 66 (1968) 13
- 1968IV01 M. Ivanovich, P.G. Young and G.G. Ohlsen, Nucl. Phys. A110 (1968) 441
- 1968JA1D Jackson, Advance Phys. 17 (1968) 481
- 1968KO1F Komarov and Savchenko, Joint Inst. Nucl. Res., Lab. Nucl. Probs., USSR, Rept. No. P1 3720 (1968)
- 1968KR03 L. Kraus, M. Suffert and D. Magnac-Valette, Nucl. Phys. A109 (1968) 593

1968LE15 J.C. Legg, W.D. Simpson and S.T. Emerson, Nucl. Phys. A119 (1968) 209
 1968MC02 R.L. McGrath, J. Cerny and S.W. Cosper, Phys. Rev. 165 (1968) 1126
 1968ME03 W.E. Meyerhof and T.A. Tombrello, Nucl. Phys. A109 (1968) 1
 1968MO10 B.J. Morton, E.E. Gross, E.V. Hungerford, J.J. Malanify and A. Zucker, Phys. Rev. 169 (1968) 825
 1968MO26 G.L. Morgan and R.L. Walter, Phys. Rev. 168 (1968) 1114; Erratum Phys. Rev. 185 (1969) 1598
 1968NE1D Newsome, LA 3997 (1968)
 1968OH1A Ohlsen, McKibben and Lawrence, Bull. Amer. Phys. Soc. 13 (1968) 1443
 1968PA1J Palevsky, Proc. of Symp. on Use of Nimrod, 1968, RHEL-R166 (1968) 19
 1968PL02 D.J. Plummer, T.A. Hodges, K. Ramavataram, D.G. Montague and N.S. Chant, Nucl. Phys. A115 (1968) 253
 1968RE10 M.A. Reimann, P.W. Martin and E.W. Vogt, Can. J. Phys. 46 (1968) 2241
 1968RO19 P.G. Roos, H.G. Pugh, M. Jain, H.D. Holmgren, M. Epstein and C.A. Ludemann, Phys. Rev. 176 (1968) 1246
 1968SA1B G.R. Satchler, L.W. Owen, A.J. Elwyn, G.L. Morgan and R.L. Walter, Nucl. Phys. A112 (1968) 1
 1968SA1C Saito, Hiura and Tanaka, Prog. Theor. Phys. 39 (1968) 635
 1968SA25 J.R. Sawers, Jr., G.L. Morgan, L.A. Schaller and R.L. Walter, Phys. Rev. 168 (1968) 1102
 1968SC1D L.I. Schiff, Phys. Rev. 176 (1968) 1390
 1968SE1A Sergeev, Geradin, Wery and Magnac-Valette, Compt. Rend. 261 (1965) 291
 1968SI1B Simmons and Malanify, Bull. Amer. Phys. Soc. 13 (1968) 564
 1968SL1A Slabospitskii et al, Sov. J. Nucl. Phys. 6 (1968) 723
 1968ST1H Starkovich et al, Bull. Amer. Phys. Soc. 13 (1968) 1448
 1968TA11 T. Tanabe, J. Phys. Soc. Jpn. 25 (1968) 21
 1968TA1E B. Tatischeff and A. Willis, Nucl. Phys. A115 (1968) 593
 1968TA1J Tanifuji and Yazaki, Prog. Theor. Phys. 40 (1968) 1023
 1968TH1C D.R. Thompson, I. Reichstein, W. McClure and Y.C. Tang, Phys. Rev. 185 (1969) 1351
 1968TH1D G. Thievent, J. Lang, R. Muller and P. Marmier, Phys. Lett. B30 (1969) 23
 1968TO01 T.A. Tombrello and R.J. Slobodrian, Nucl. Phys. A111 (1968) 236
 1968TO1F Tombrello, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl., J. Phys. Soc. Jpn. 24 (1968) 63

- 1968VA02 V. Valkovic, C. Joseph, S.T. Emerson and G.C. Phillips, Nucl. Phys. A106 (1968) 138
- 1968VA12 V. Valkovic, C. Joseph, A. Niiler and G.C. Phillips, Nucl. Phys. A116 (1968) 497
- 1968VI03 B. Vignon, J.-F. Cavaignac and J.-P. Longequeue, Compt. Rend. B266 (1968) 878
- 1968WA01 R.E. Warner and R.W. Bercaw, Nucl. Phys. A109 (1968) 205
- 1968WI1E Wise, Thesis, Washington State Univ. (1968); Phys. Abs. 27627 (1969)
- 1968YA02 T. Yanabu, S. Yamashita, K. Hosono, S. Matsuki, T. Tanabe, K. Takimoto, Y. Okuma, K. Ogino, S. Okumura and R. Ishiwari, J. Phys. Soc. Jpn. 24 (1968) 667
- 1968YO06 P.G. Young, R.H. Stokes and G.G. Ohlsen, Phys. Rev. 173 (1968) 949
- 1968ZA1A Zaika, Mokhnach, Shmarin and Yasnogorodskii, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 257
- 1968ZU1A Zupancic, Proc. Symp. on Use of Nimrod, 1968; RHEL/R166 (1968) 67
- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60
- 1969BE1M H. C. Benohr and K. Wildermuth, Nucl. Phys. A128 (1969) 1
- 1969BR22 H. Bruckmann and F.K. Schmidt, Nucl. Phys. A136 (1969) 81
- 1969BU10 S.N. Bunker, J.M. Cameron, M.P. Epstein, G. PaiImage, J.R. Richardson, J.G. Rogers, P. TomaImage and J.W. Verba, Nucl. Phys. A133 (1969) 537
- 1969BU19 F.W. Busser, H. Dubenkropp, F. Niebergall and K. Sinram, Nucl. Phys. A129 (1969) 666
- 1969CO1E G. Cocho, A. Mondragon and M. Colon-Vela, Nucl. Phys. A125 (1969) 425
- 1969CR1A A.H. Cromer, Nucl. Phys. B11 (1969) 152
- 1969CR1B Cramer, Thesis, Univ. of Virginia (1969); Phys. Abs. 39036 (1970)
- 1969DA1F Dauchy et al, Lett. Nuovo Cim. 2 (1969) 263
- 1969DE04 D.W. Devins, S.M. Bunch, H.H. Forster, J. Hokinian and C.C. Kim, Nucl. Phys. A126 (1969) 261
- 1969DO1G Dolinski and Turovtsev, Sov. J. Nucl. Phys. 9 (1969) 445
- 1969EL1B Elliott, Proc. Int. Conf. Montreal (1969) 277
- 1969ER1B Ericson and Locher, Nucl. Struct., Dubna Symp. 1968, IAEA (1968) 389
- 1969GA12 D. Garreta, J. Sura and A. Tarrats, Nucl. Phys. A132 (1969) 204
- 1969GE1A George and Shamu, Bull. Amer. Phys. Soc. 14 (1969) 535
- 1969GI1A B.F. Gibson, A. Goldberg and M.S. Weiss, Phys. Rev. 181 (1969) 1486
- 1969GO1C Golovanova and Zelenskaya, Sov. J. Nucl. Phys. 8 (1969) 158
- 1969GO1G Gogsadze and Kopaleishvili, Sov. J. Nucl. Phys. 8 (1969) 509

- 1969HA16 S.A. Harbison, R.J. Griffiths, F.G. Kingston, A.R. Johnston and G.T.A. Squier, Nucl. Phys. A130 (1969) 513
- 1969HA1M Halbleib and Scott, Nucl. Sci. Eng. 37 (1969) 271
- 1969HE15 P. Heiss and H.H. Hackenbroich, Phys. Lett. B30 (1969) 373
- 1969HE1H Hentschel, Z. Phys. 219 (1969) 32
- 1969HE1K Henley, Isospin in Nucl. Phys., Ed. D.H. Wilkinson (North Holland, 1969) 15
- 1969HE1L Heiss, BMWF FBK 69 38 (1969)
- 1969KO1M Komarov, Kosarev and Savchenko, Joint. Inst. Nucl. Res., Lab. Nucl. Probs., USSR, Rept. No. P1 4876 (1969)
- 1969KO30 T.I. Kopaleishvili and I.Z. Machabeli, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 69; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 65
- 1969KU1D Kujawski, Tech. Rept. No. 70-048, Univ. of Maryland (1969)
- 1969LE03 L. Lesniak and H. Wolek, Nucl. Phys. A125 (1969) 665
- 1969LI02 T.Y. Li and S.K. Mark, Nucl. Phys. A123 (1969) 147
- 1969MA1H Manduchi, Moschini, Torielli and Zannoni, Nucl. Instrum. Meth. 67 (1969) 267
- 1969ME1B Meiner et al, Bull. Amer. Phys. Soc. 14 (1969) 512
- 1969ME1C Meyer, Amer. J. Phys. 37 (1969) 296
- 1969MI10 R.C. Minehart, L. Coulson, W.F. Grubb, III and K. Ziock, Phys. Rev. 177 (1969) 1455, 1464
- 1969NA09 H. Nakamura, Nucl. Phys. A137 (1969) 51
- 1969NA1F Nagatani, Herling and Kolltveit, Bull. Amer. Phys. Soc. 14 (1969) 511
- 1969PE01 V. Perez-Mendez, J.M. Sperinde and A.W. Stetz, Phys. Lett. B28 (1969) 648
- 1969PL01 G.R. Plattner and L.G. Keller, Phys. Lett. B29 (1969) 301
- 1969RO16 D.C. Rorer, B.M. Ecker and R.O. Akyuz, Nucl. Phys. A133 (1969) 410
- 1969RO1L Rogers, UCLA P-84 (1969)
- 1969RO20 L.P. Robertson, R.C. Hanna, K. Ramavataram, D.W. Devins, T.A. Hodges, Z.J. Moroz, S.J. Hoey and D.J. Plummer, Nucl. Phys. A134 (1969) 545
- 1969RO24 J.G. Rogers, J.M. Cameron, M.B. Epstein, G. Paic, P. Tomas, J.R. Richardson, J.W. Verba and P. Doherty, Nucl. Phys. A136 (1969) 433
- 1969RU1A Ruhla, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (Academic Press, 1969) 701
- 1969SC1J R. Schrials and H.M. Darley, Phys. Lett. B29 (1969) 291
- 1969SE1B Seiler, Baumgartner, Huber and Leeman, Helv. Phys. Acta 42 (1969) 567
- 1969ST1F Starkovich, LA 4191 (1969)

1969TO1A I.S. Towner, Nucl. Phys. A126 (1969) 97
 1969VI05 B. Vignon, E. Ligeon and J.P. Longequeue, J. Phys. (Paris) 30 (1969) 913
 1969VO1E Vogt, Proc. Int. Conf., Montreal (1969) 5
 1969WI1C Wildermuth, Bochum Conf. STI/PUB/232 IAEA (1969) 3
 1969WI1D Wilhelmi et al, Contrib. Montreal (1969) 284
 1969YA1B Yamashita et al, J. Phys. Soc. Jpn. 26 (1969) 1078
 1970AD1C Adeyasevich and Antonenko, Sov. J. Nucl. Phys. 10 (1970) 278
 1970AR1B R.A. Arndt and L.D Roper, Phys. Rev. C1 (1970) 903
 1970AS02 P.A. Assimakopoulos, E. Beardsworth, D.P. Boyd and P.F. Donovan, Nucl. Phys. A144 (1970) 272
 1970BA1F Barashenkov and Toneev, Uspekhi Fiz. Nauk 100 (1970) 425
 1970BE1A Bezotosnyi, Zhmailo, Surov and Shvetsov, Sov. J. Nucl. Phys. 10 (1970) 127
 1970BE49 M. Bernas, D. Bachelier, J.K. Lee, P. Radvanyi, M. Roy-Stephan, I. Brissaud and C. Detraz, Nucl. Phys. A156 (1970) 289
 1970BO1P Boschitz, Proc. 3rd Int. Conf. on High-Energy Phys. and Nucl. Struct., New York, 1969 (Plenum, 1970) 288; Phys. Abs. 43588 (1971)
 1970BR1L Bruno and Duck, Bull. Amer. Phys. Soc. 15 (1970) 1652
 1970BR30 W.B. Broste, G.P. Lawrence, J.L. McKibben, G.G. Ohlsen and J.E. Simmons, Phys. Rev. Lett. 25 (1970) 1040
 1970CA1K Carlton, Thesis, Univ. Georgia (1970); Phys. Abs. 67885 (1971)
 1970CO1M Cominos, Crawford and Li, Bull. Amer. Phys. Soc. 15 (1970) 768
 1970DE41 P.A. Deutchman, Lett. Nuovo Cim. 4 (1970) 61
 1970ER1A Ericson and Locher, Nucl. Phys. A148 (1970) 1
 1970FE1C H. Federsel, E.-J. Kanellopoulos, W. Sunkel and K. Wildermuth, Phys. Lett. B33 (1970) 140
 1970GA1A Garber et al, BNL 400, 3rd Edition, Vol. 1 (1970)
 1970GA1G Gagne, Lambert and Treado, Bull. Amer. Phys. Soc. 15 (1970) 1695
 1970GN1A Gnedin, Dolginov, Silantsev and Shibanov, Yad. Fiz. 12 (1970) 815
 1970GO12 B. Gottschalk and S.L. Kannenberg, Phys. Rev. C2 (1970) 24
 1970GO36 N.P. Goldstein, A. Held and D.G. Stairs, Can. J. Phys. 48 (1970) 2629
 1970GR17 E.E. Gross, E.V. Hungerford, J.J. Malanify and R. Woods, Phys. Rev. C1 (1970) 1365
 1970HA1L Haussen, Schweitzer, Seeliger and Seidel, Nucl. Instrum. Meth. 88 (1970) 251
 1970HA1M Hardy, Thesis, Rice Univ. (1970); Phys. Abs. 51261 (1971)

- 1970HA36 S.A. Harbison, R.J. Griffiths, N.M. Stewart, A.R. Johnston and G.T.A. Squier, Nucl. Phys. A152 (1970) 503
- 1970HE1C Heiss and Hackenbroich, Z. Phys. 231 (1970) 230
- 1970HE1D Heiss and Hackenbroich, Z. Phys. 235 (1970) 422
- 1970HO1L Hojvat, Thesis, Univ. British Columbia (1970); Phys. Abs. 6316 (1970)
- 1970HO32 H.M. Hofmann and D. Fick, Z. Phys. 240 (1970) 420
- 1970IR01 J.M. Irvine and V.F.E. Pucknell, Nucl. Phys. A159 (1970) 513
- 1970KH03 V.A. Khangulyan, Yad. Fiz. 11 (1970) 967; Sov. J. Nucl. Phys. 11 (1970) 538
- 1970KO09 A. Kosiara and H.B. Willard, Phys. Lett. B32 (1970) 99
- 1970KU1C E. Kujawski, Phys. Rev. C1 (1970) 1651
- 1970KU1D Kull, Thesis, Michigan State Univ. (1970); Phys. Abs. 6667 (1970)
- 1970LA10 H.T. Larson, A.D. Bacher, K. Nagatani and T.A. Tombrello, Nucl. Phys. A149 (1970) 161
- 1970LA1K Larson, Thesis, CIT (1970); Phys. Abs. 51262 (1971)
- 1970LI06 W.-K. Lin, F. Scheibling and R.W. Kavanagh, Phys. Rev. C1 (1970) 816
- 1970MI05 D. Miljanic, M. Furic and V. Valkovic, Nucl. Phys. A148 (1970) 312
- 1970MO31 G.L. Morgan and R.L. Walter, Phys. Rev. C2 (1970) 2034; Addendum to 1968CH35
- 1970OM1A Omojola, J. Phys. A3 (1970) 630
- 1970PO1A Porter, Yale Univ. (1970); Phys. Abs. 3490 (1970)
- 1970RA1D Ramavataram and Ramavataram, Nucl. Phys. A147 (1970) 293
- 1970RA1J Rao, Swami, Gurtu and Singh, Proc. Indian Acad. Sci. A71 (1970) 257; Phys. Abs. 73791 (1970)
- 1970RE1A Reichstein and Tang, Nucl. Phys. A158 (1970) 529
- 1970RO17 J.G. Rogers, G. Paic, J.R. Richardson and J.W. Verba, Phys. Rev. C2 (1970) 828
- 1970RO1L Rohrer et al, Helv. Phys. Acta 43 (1970) 746
- 1970RO1M Roberts et al, Rept. NASA-TN-D-5961 (1970); Phys. Abs. 39304 (1971)
- 1970SA01 T. Sawada, G. Paic, M.B. Epstein and J.G. Rogers, Nucl. Phys. A141 (1970) 169
- 1970SC18 H. Schroder and W. Mausberg, Z. Phys. 235 (1970) 234
- 1970SE1B F. Seiler and E. Baumgartner, Nucl. Phys. A153 (1970) 193
- 1970SL1A Slobodrian et al, Proc. 1st Int. Conf. on Three Body Problems in Nucl. & Particle Phys., Birmingham, England, 1969 (London: North-Holland, 1970) p. 390
- 1970SL1B Slaus, Theory of Nucl. Struct. - Trieste 1969, IAEA STI/PUB/249 (1970) 17
- 1970ST15 T. Stammbach, J. Taylor, G. Spalek and R.L. Walter, Phys. Rev. C2 (1970) 434

- 1970SU1B Sukhoruchkin, *Sov. J. Nucl. Phys.* 10 (1970) 144
- 1970TH1D Thompson, Epstein and Sawada, *Nucl. Phys.* A142 (1970) 571
- 1970VA1L Valkovic, Furic, Miljanic and Tomas, *Proc. 1st Int. Conf. on Three Body Problem in Nucl. & Particle Phys., Birmingham, England, 1969* (London: North-Holland, 1970) p. 436
- 1970WA15 R.E. Warner, B.E. Corey, E.L. Petersen, R.W. Bercaw and J.E. Poth, *Nucl. Phys.* A148 (1970) 503
- 1970WA1G Walker and Stokes, LA-DC 11224 (1970)
- 1970WA1L B.E. Watt and W.T. Leland, *Phys. Rev.* C2 (1970) 1680
- 1970WA32 B.E. Watt and W.T. Leland, *Phys. Rev.* C2 (1970) 1677
- 1970WI1G Willmes, Shadoan and Cahill, *Bull. Amer. Phys. Soc.* 15 (1970) 1694
- 1970ZO1A Zofka, *Czech. J. Phys.* 20 (1970) 926
- 1971AL1B Allas et al, *Bull. Amer. Phys. Soc.* 16 (1971) 512
- 1971AR20 R.A. Arndt, L.D. Roper and R.L. Shotwell, *Phys. Rev.* C3 (1971) 2100
- 1971BA1U Baker, *Polarization, Madison, 1970* (Univ. Wisconsin Press, 1971) p. 899
- 1971BA1Y Baguil et al, *Nuovo Cim.* A1 (1971) 285
- 1971BA2Y Baus-Baghdikian, *U. Libre Bruxelles, Bull. No. 45* (1971)
- 1971BA61 A.G. Baryshnikov and L.D. Blokhintsev, *Phys. Lett.* B36 (1971) 205
- 1971BA74 C.J. Batty, E. Friedman and D.F. Jackson, *Nucl. Phys.* A175 (1971) 1
- 1971BR12 K.H. Bray, S.N. Bunker, M. Jain, K.S. Jayaraman, G.A. Moss, W.T.H. van Oers, D.O. Wells and Y.I. Wu, *Phys. Rev.* C3 (1971) 1771
- 1971CH1T Chamberlin, *Thesis, Texas A and M Univ.* (1971); *Phys. Abs.* 67322 (1972)
- 1971CH31 G. Chenevert, N.S. Chant, I. Halpern, C. Glashauser and D.L. Hendrie, *Phys. Rev. Lett.* 27 (1971) 434
- 1971CH50 N.S. Chant and E.M. Henley, *Phys. Rev. Lett.* 27 (1971) 1657
- 1971DE20 B. De Facio and R.K. Umerjee, *Nucl. Phys.* A169 (1971) 257
- 1971DO1C Dodder et al, *Polarization, Madison, 1970* (Univ. Wisconsin Press, 1971) p. 520
- 1971EF01 V.D. Efros and M.V. Zhukov, *Phys. Lett.* B37 (1971) 18
- 1971FE10 H. Feshbach, A. Gal and J. Hufner, *Ann. Phys. (New York)* 66 (1971) 20
- 1971FO1J Forssmann, Graf, Schober and Jochim, *Polarization, Madison, 1970* (Univ. Wisconsin Press, 1971) p. 537
- 1971GI1B B. Giraud and D. Zaikine, *Phys. Lett.* B37 (1971) 25
- 1971GR12 E.E. Gross, E.V. Hungerford and J.J. Malanify, *Nucl. Phys.* A164 (1971) 376

- 1971GR15 W. Gruebler, V. Konig, A. Ruh, R.E. White, P.A. Schmelzbach, R. Risler and P. Marmier, Nucl. Phys. A165 (1971) 505
- 1971GR32 H. Grunder, R. Gleyvod, G. Lietz, G. Morgan, H. Rudin, F. Seiler and A. Stricker, Helv. Phys. Acta 44 (1971) 662
- 1971GR45 R. Grotzschel, B. Kuhn, K. Moller, J. Mosner and G. Schmidt, Nucl. Phys. A176 (1971) 261
- 1971GR47 W. Gruebler, V. Konig, A. Ruh, P.A. Schmelzbach, R.E. White and P. Marmier, Nucl. Phys. A176 (1971) 631
- 1971HA02 D.M. Hardy, S.D. Baker, D.H. McSherry and R.J. Spiger, Nucl. Phys. A160 (1971) 154
- 1971HA1L Hale, Dodder and Witte, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 573
- 1971HA2G Haeberli, Proc. 2nd Int. Conf. on Polarized Targets LBL 500 (1971) p. 329
- 1971HE05 P. Heiss and H.H. Hackenbroich, Nucl. Phys. A162 (1971) 530
- 1971HE1M Hertzog, Thesis, Brown Univ. (1971) ; Phys. Abs. 70452 (1972)
- 1971HI07 D. Hilscher, P.A. Quin and J.C. Davis, Nucl. Phys. A173 (1971) 216
- 1971HU1B Huber et al, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 546
- 1971HU1H Hudomalj, Valkovic and Tomas, Fizika (Yugoslavia) 4 (1971) 36
- 1971IK01 M. Ikeda, Prog. Theor. Phys. 46 (1971) 1088
- 1971IN1C Ingalls, Thesis, Princeton Univ. (1971); Phys. Abs. 67321 (1972)
- 1971KE16 L.G. Keller and W. Haeberli, Nucl. Phys. A172 (1971) 625
- 1971KH03 V.A. Khangulyan, Yad. Fiz. 13 (1971) 525; Sov. J. Nucl. Phys. 13 (1971) 295
- 1971KI02 K. Kilian, H. Treiber, R. Strausz and D. Fick, Phys. Lett. B34 (1971) 283
- 1971KL02 W. Klinger, F. Dusch and R. Fleischmann, Nucl. Phys. A166 (1971) 253
- 1971KL04 J.T. Klopocic and S.E. Darden, Phys. Rev. C3 (1971) 2171; Erratum Phys. Rev. C4 (1971) 1494
- 1971KO09 V. Konig, W. Gruebler, A. Ruh, R.E. White, P.A. Schmelzbach, R. Risler and P. Marmier, Nucl. Phys. A166 (1971) 393
- 1971KR1A Kraus, Linck and Magnac-Valette, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 564
- 1971LE02 C.W. Lewis, D.P. Saylor and L.C. Northcliffe, Phys. Rev. C3 (1971) 28
- 1971LE13 C. Leemann, H. Burgisser, P. Huber, U. Rohrer, H. Schieck and F. Seiler, Helv. Phys. Acta 44 (1971) 141
- 1971LE1G Leemann et al, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 548

- 1971LE1N Leclercq-Villain, Conf. on Certain Microscopic Aspects of Nucl. Rxns, La Toursinei, 1971 (Vieurbanne, France: Inst. Nucl. Phys. 1971); Phys. Abs. 47250 (1971)
- 1971LE27 C. Leemann, H. Burgisser, P. Huber, U. Rohrer, H. Paetz gen.Schieck and F. Seiler, Ann. Phys. (New York) 66 (1971) 810
- 1971LI04 W.-K. Lin and F. Scheibling, Phys. Rev. C3 (1971) 20
- 1971LO24 W.G. Love and L.W. Owen, Phys. Lett. B37 (1971) 463
- 1971MC1L McKibben, Proc. 2nd Int. Conf. on Polarized Targets LBL 500 (1971) 307
- 1971MI12 D. Miljanic and V. Valkovic, Nucl. Phys. A176 (1971) 110
- 1971MU04 G.S. Mutchler, W.B. Broste and J.E. Simmons, Phys. Rev. C3 (1971) 1031
- 1971MU1H Mullensiefen, Nucl. Phys. B28 (1971) 368
- 1971NI06 A. Niiler, M. Drosig, J.C. Hopkins, J.D. Seagrave and E.C. Kerr, Phys. Rev. C4 (1971) 36
- 1971OH04 G.G. Ohlsen, J.L. McKibben, G.P. Lawrence, P.W. Keaton, Jr. and D.D. Armstrong, Phys. Rev. Lett. 27 (1971) 599
- 1971OH1A Ohlsen, Keaton and Simmons, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 415
- 1971OH1B Ohlsen, McKibben and Lawrence, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 503
- 1971OH1C Ohlsen, Keaton and Gammel, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 512
- 1971PI04 J. Pigeon, J. Barguil, C. Fayard, G-H. Lamot and E. El Baz, Phys. Rev. C4 (1971) 704
- 1971PL06 G.R. Plattner and A.D. Bacher, Phys. Lett. B36 (1971) 211
- 1971PL07 D.J. Plummer, K. Ramavataram, T.A. Hodges, D.G. Montague, A. Zucker and N.K. Ganguly, Nucl. Phys. A174 (1971) 193
- 1971PO1C Poutissou and Jeremie, Bull. Amer. Phys. Soc. 16 (1971) 133
- 1971RA15 S. Ramavataram and K. Ramavataram, Can. J. Phys. 49 (1971) 1798
- 1971RA27 K. Ramavataram, D.J. Plummer, T.A. Hodges and D.G. Montague, Nucl. Phys. A174 (1971) 204
- 1971RE19 D. Rendic, N.D. Gabitzsch, V. Valkovic, W. von Witsch and G.C. Phillips, Nucl. Phys. A178 (1971) 49
- 1971RH1A Rhea, Stammbach and Walter, Polariz. Phenom. in Nucl. Rxns, Madison, 1970, Eds. H.H. Barscahl and W. Haeberli (Univ. Wisconsin Press, 1971) 556
- 1971RO35 U. Rohrer, P. Huber, C. Leemann, H. Meiner and F. Seiler, Helv. Phys. Acta 44 (1971) 846
- 1971RU07 R.W. Rutkowski and E.E. Gross, Phys. Lett. B35 (1971) 151

- 1971RU1E Rutkowski, Thesis, Univ. Tennessee (1971); Phys. Abs. 42582 (1972)
- 1971SC04 P. Schwandt, T.B. Clegg and W. Haerberli, Nucl. Phys. A163 (1971) 432
- 1971SC1L Scuster and Hagengruber, Polarization Phenomena in Nucl. Rxns, Madison, 1970, Eds. H.H. Barscahl and W. Haerberli (Univ. Wisconsin press, 1971) p. 865
- 1971SE1G Seiler and Baumgartner, Polarization Phenomena in Nucl. Rxns, Madison, 1970, Eds. H.H. Barscahl and W. Haerberli (Univ. Wisconsin Press, 1971) p. 518
- 1971SI30 M. Sieminski, Z. Wilhelmi and P. Zupranski, Acta Phys. Pol. B2 (1971) 503
- 1971ST1J Striebel, Polarization Phenomena in Nucl. Rxns., Madison, 1970, Eds. H.H. Barscahl and W. Haerberli (Univ. Wisconsin Press, 1971) p. 95
- 1971SW10 W.E. Sweeney, Jr., V. Valkovic, D. Rendic and G.C. Phillips, Phys. Lett. B37 (1971) 183
- 1971TH09 D.R. Thompson and Y.C. Tang, Phys. Rev. C4 (1971) 306
- 1971WA08 R.F. Wagner and C. Werntz, Phys. Rev. C4 (1971) 1
- 1971WA18 R.E. Warner, G.C. Ball, W.G. Davies, A.J. Ferguson and J.S. Forster, Phys. Rev. Lett. 27 (1971) 961
- 1971WA1D Walter, Polarization Phenomena in Nucl. Reactions, Madison, 1970, Eds. H.H. Barscahl and W. Haerberli (Univ. Wisconsin Press, 1971) p. 317
- 1971WO07 W. Wolfli, J. Hall and R. Muller, Phys. Rev. Lett. 27 (1971) 271
- 1971ZA1C Zaika et al, Polarization Phenomena in Nucl. Rxns, Madison, 1970, Eds. H.H. Barscahl and W. Haerberli (Univ. Wisconsin, 1971) 531
- 1971ZA1D D.A. Zaikin, Nucl. Phys. A170 (1971) 584
- 1972AD02 M.L. Adelberg and A.M. Saperstein, Phys. Rev. C5 (1972) 1180
- 1972AG01 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, G.P. Gulkanyan, V.M. Sidorov and V.A. Yarba, Yad. Fiz. 15 (1972) 18; Sov. J. Nucl. Phys. 15 (1972) 10
- 1972AL1P Allab, Beaumevielle and Dauchy, Bull. Amer. Phys. Soc. 17 (1972) 929
- 1972AN1P Anzelon et al, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 54
- 1972AN1Q Antolkovic, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 695
- 1972AN24 Y.N. Antufev, V.L. Agranovich, V.S. Kuzmenko, I.I. Miroshnichenko and P.V. Sorokin, Pisma Zh. Eksp. Teor. Fiz. 16 (1972) 77; JETP Lett. (USSR) 16 (1972) 52
- 1972AN27 Y.P. Antoufi ev, V.L. Agranovich, V.S. Kuzmenko and P.V. Sorokin, Phys. Lett. B42 (1972) 347
- 1972AN29 Y.P. Antufev, V.L. Agranovich, V.S. Kuzmenko and P.V. Sorokin, Pisma Zh. Eksp. Teor. Fiz. 16 (1972) 339; JETP Lett. (USSR) 16 (1972) 240
- 1972AV1E Avida, Chang, Boyd and Glavish, Bull. Amer. Phys. Soc. 17 (1972) 562

- 1972AZ03 L.S. Azhgirei, Z. Cisek, Z.V. Krumstein, Y.P. Merekov, Z. Moroz, Ngo Quang Zui, V.I. Petrukhin, A.I. Ronzhin, G.A. Shelkov and O.D. Dalkarov, Nucl. Phys. A195 (1972) 581
- 1972BA24 A.D. Bacher, G.R. Plattner, H.E. Conzett, D.J. Clark, H. Grunder and W.F. Tivol, Phys. Rev. C5 (1972) 1147
- 1972BA30 M.P. Baker, J.M. Cameron, N.S. Chant and N.F. Mangelson, Nucl. Phys. A184 (1972) 97
- 1972BA32 D. Bachelier, M. Bernas, H.L. Harney, J.C. Jourdain, P. Radvanyi and M. Roy-Stephan, Nucl. Phys. A184 (1972) 641
- 1972BB19 A.G. Baryshnikov, L.D. Blokhintsev, A.N. Safronov and V.V. Turovtsev, Pisma Zh. Eksp. Teor. Fiz. 16 (1972) 414; JETP Lett. 16 (1972) 294
- 1972BL1F Blokhintsev and Safronov, Nucl. Phys. A180 (1972) 363
- 1972BO29 E.T. Boschitz, W.K. Roberts, J.S. Vincent, M. Blecher, K. Gotow, P.C. Gugelot, C.F. Perdrisat, L.W. Swenson and J.R. Priest, Phys. Rev. C6 (1972) 457
- 1972BR08 K.H. Bray, J.M. Cameron, G.C. Neilson and T.C. Sharma, Nucl. Phys. A181 (1972) 319
- 1972BR10 W.B. Broste, G.S. Mutchler, J.E. Simmons, R.A. Arndt and L.D. Roper, Phys. Rev. C5 (1972) 761
- 1972CA37 P. Camiz, E. Olivieri, M. Scalia and A. D'Andrea, Nuovo Cim. A12 (1972) 71
- 1972CH31 E. Cheifetz, B. Eylon, E. Fraenkel and A. Gavron, Phys. Rev. Lett. 29 (1972) 805
- 1972CR01 D.S. Cramer and L. Cranberg, Nucl. Phys. A180 (1972) 273
- 1972DE30 J. Deenen, Nucl. Phys. A189 (1972) 73
- 1972DE44 J. Decharge, G. Surget, G. Bruno and M.Y. Decharge, J. Phys. (Paris) 33 (1972) 485
- 1972DE46 M.L.V.L. de Slobodrian, B. Cujec, R. Roy and R.J. Slobodrian, Nucl. Phys. A194 (1972) 577
- 1972GA11 K. Gabathuler, J. Rohlin, J.J. Domingo, C.H.Q. Ingram, S. Rohlin and N.W. Tanner, Nucl. Phys. B40 (1972) 32
- 1972GA1N Garrett, 4th Ainsel Nucl. Phys. Conf., Sydney, 1972 (Sutherland, Australia, Austr. Inst. Nucl. Sci. Eng., 1972) 445
- 1972GI10 P. Gil, L. Marquez, J.L. Quebert and H. Sztark, J. Phys. (Paris) 33 (1972) 315
- 1972GI11 B.F. Gibson, A. Goldberg and M.S. Weiss, Phys. Rev. C6 (1972) 741
- 1972GR06 A.M. Green and A. Prodon, Nucl. Phys. A183 (1972) 225
- 1972GU1D Guy, Eisenberg, Noble and Weber, Bull. Amer. Phys. Soc. 17 (1972) 918
- 1972HA57 H.H. Hackenbroich and T.H. Seligman, Phys. Lett. B41 (1972) 102
- 1972IK01 M. Ikeda, Phys. Rev. C6 (1972) 1608

1972JA07 O.N. Jarvis, C. Whitehead and M. Shah, Nucl. Phys. A184 (1972) 615
 1972JA14 J. Janecke, Phys. Rev. C6 (1972) 467
 1972KA38 N.I. Kassis, Nucl. Phys. A194 (1972) 205
 1972KA44 S. Kato, H. Orihara, S. Kubono, J. Kasagi, H. Ueno, T. Nakagawa and T. Tohei, Nucl. Phys. A195 (1972) 534
 1972KE17 P.W. Keaton, Jr., D.D. Armstrong, R.A. Hardekopf, P.M. Kurjan and Y.K. Lee, Phys. Rev. Lett. 29 (1972) 880
 1972KI01 H.T. King, W.E. Meyerhof and R.G. Hirko, Nucl. Phys. A178 (1972) 337
 1972KI02 T.R. King and R. Smythe, Nucl. Phys. A183 (1972) 657
 1972KI1F Kisslinger, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 593
 1972KO07 V. Konig, W. Gruebler, R.E. White, P.A. Schmelzbach and P. Marmier, Nucl. Phys. A185 (1972) 263
 1972KO13 V.M. Kolybasov and N.Y. Smorodinskaya, Yad. Fiz. 15 (1972) 483; Sov. J. Nucl. Phys. 15 (1972) 269
 1972LA1L Lambert et al, Few Body Problems., UCLA, 1972 (North Holland, 1972) 725
 1972LE1L Lee and Cusson, Ann. Phys. 72 (1972) 353
 1972LO1K Locher, Nucl. Phys. B36 (1972) 634
 1972MA57 H.A. Mavromatis and B. Singh, Phys. Lett. B41 (1972) 251
 1972MA61 I.A. Mackenzie, S.K. Mark and T.Y. Li, Nucl. Phys. A195 (1972) 609
 1972MA62 A.M. MacLeod and G.R. Milne, J. Phys. (London) A5 (1972) 1252
 1972NA1D Nakamura, Prog. Theor. Phys. 48 (1972) 695
 1972NI02 A. Niiler, R.J. Spiger, W. von Witsch and G.C. Phillips, Nucl. Phys. A179 (1972) 263
 1972NI04 J.E. Nicholls, A. Craig, T.C. Griffith, D.C. Imrie, C.J. Lush and A.J. Metheringham, Nucl. Phys. A181 (1972) 329
 1972OH1C Ohlsen and Mitchell, Bull. Amer. Phys. Soc. 17 (1972) 153
 1972OH1D Ohlsen, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 585
 1972OT05 I. Ots, Eesti NSV Tead. Akad. Toim. Fuus. Mat. (USSR) 21 (1972) 289; Izv. Akad. Nauk Est. SSR Fiz. Mat. 21 (1972) 289; Phys. Abs. 17220 (1973)
 1972PL02 G.R. Plattner, A.D. Bacher and H.E. Conzett, Phys. Rev. C5 (1972) 1158
 1972PN1A Pniewski, Few Particle Problems, UCLA, 1972 (North-Holland, 1972) 145
 1972QU01 J.R. Quinn, M.B. Epstein, S.N. Bunker, J.W. Verba and J.R. Richardson, Nucl. Phys. A181 (1972) 440
 1972RI10 A.S. Rinat, L.P. Kok and M. Stingl, Nucl. Phys. A190 (1972) 328
 1972RO1M Rohrig, Knutson, Goddard and Haeberli, Bull. Amer. Phys. Soc. 17 (1972) 153

- 1972RY01 N. Ryu, J. Sanada, H. Hasai, D.C. Worth, M. Nishi, T. Hasegawa, H. Ueno, M. Seki, K. Iwatani and Y. Nojiri, Nucl. Phys. A180 (1972) 657
- 1972SC1M Schwaller et al, CERN 72-13 (1972)
- 1972SC1R Schroder et al, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 1002
- 1972SC1T Schmidt, Wiss. Z. Tech. Univ. Dres. (Germany) 21 (1972) 700; Phys. Abs. 60156 (1973)
- 1972SE09 F. Seiler, Nucl. Phys. A187 (1972) 379
- 1972SI1F Simon, Mitchell and Ohlsen, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 735
- 1972SM05 J.R. Smith and S.T. Thornton, Nucl. Phys. A187 (1972) 433
- 1972ST01 T. Stambach and R.L. Walter, Nucl. Phys. A180 (1972) 225
- 1972ST08 L. Strauss and E. Friedland, Z. Phys. 250 (1972) 370
- 1972TA1E Tanifuji and Yazaki, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 739
- 1972TH04 D.T. Thompson, G.E. Tripard and D.H. Ehlers, Phys. Rev. C5 (1972) 1174
- 1972TH06 D.R. Thompson, Y.C. Tang and R.E. Brown, Phys. Rev. C5 (1972) 1939
- 1972TH08 D.T. Thompson and G.E. Tripard, Phys. Rev. C6 (1972) 452
- 1972TH1B Thompson, Thesis, Washington State Univ. (1972); Phys. Abs. 984 (1973)
- 1972WA22 R.E. Warner, G.R. Flierl, W.G. Davies, G.C. Ball, A.J. Ferguson, J.S. Forster and S.A. Gottlieb, Nucl. Phys. A192 (1972) 341
- 1972WO1E Wolfli et al, Few Particle Problems, UCLA, 1972 (North Holland, 1972) 96
- 1972ZH1B Zhukov and Efros, Sov. J. Nucl. Phys. 14 (1972) 322
- 1973AD01 M.L. Adelberg and A.M. Saperstein, Phys. Rev. C7 (1973) 63
- 1973AR1N Arndt, Long and Roper, Nucl. Phys. A209 (1973) 429
- 1973AR1P Arndt and Roper, Nucl. Phys. A209 (1973) 447
- 1973BA2P Baker, May, Hardekopf and Ohlsen, Bull. Amer. Phys. Soc. 18 (1973) 1394
- 1973BH1A Bhowmik, Chang and Holmgren, Bull. Amer. Phys. Soc. 18 (1973) 78
- 1973BO1Y Bollen et al, Bull. Amer. Phys. Soc. 18 (1973) 1600
- 1973CH05 M. Chemarin, J.P. Burq, J.C. Cabrillat and B. Ille, Nucl. Phys. A202 (1973) 71
- 1973CH1Q Chang, Roos, Bhowik and Chant, Bull. Amer. Phys. Soc. 18 (1973) 1401
- 1973CH27 F.S. Chwieroth, R.E. Brown, Y.C. Tang and D.R. Thompson, Phys. Rev. C8 (1973) 938
- 1973CL01 B.C. Clark, R.L. Mercer, D.G. Ravenhall and A.M. Saperstein, Phys. Rev. C7 (1973) 466

1973CL13 J.F. Clare, Nucl. Phys. A217 (1973) 342
 1973CO1Y Collaboration Clermont-FD-Lyon-Strasbourg, in Contrib., Uppsala (1973) 36
 1973DA1M Davison et al, Bull. Amer. Phys. Soc. 18 (1973) 553
 1973DA1N Davison et al, Phys. in Canada 29 (1973) 3
 1973DA1P Davison et al, Bull. Amer. Phys. Soc. 18 (1973) 1381
 1973DE17 V.P. Denisov, L.A. Kulchitskii and I.Y. Chubukov, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 107; Bull. Acad. Sci. USSR Phys. Ser. 37:1 (1973) 94
 1973DE2B De Facio and Wuller, Bull. Amer. Phys. Soc. 18 (1973) 1392
 1973DE51 A.A. Debenham, V. Konig, W. Gruebler, P.A. Schmelzbach, R. Risler and D.O. Boerma, Nucl. Phys. A216 (1973) 42
 1973EI01 J.M. Eisenberg, R. Guy, J.V. Noble and H.J. Weber, Phys. Lett. B43 (1973) 20; Erratum Phys. Lett. B45 (1973) 93
 1973EK1A Ekelof et al, in Contrib., Uppsala (1973) 35
 1973FE1H Feshbach and Ullo, Bull. Amer. Phys. Soc. 18 (1973) 22
 1973FE1J Federsel, Schraner, Teufel and Wildermuth, in Munich 1 (1973) 499
 1973FI04 S. Fiarman and W.E. Meyerhof, Nucl. Phys. A206 (1973) 1
 1973GA16 S.N. Gardiner, J.L. Matthews and R.O. Owens, Phys. Lett. B46 (1973) 186
 1973GA17 R.B. Galloway and R.M.A. Maayouf, Nucl. Phys. A212 (1973) 182
 1973GA1H Gari, Phys. Rept. C6 (1973) 317
 1973GE1G Geisler et al, Bull. Amer. Phys. Soc. 18 (1973) 580
 1973GO38 C.A. Goulding, P. Stoler and J.D. Seagrave, Nucl. Phys. A215 (1973) 253
 1973GR1L Grotzschel et al, in Munich 1 (1973) 6
 1973GR22 A.P. Graevskii, B.A. Bochagov and Y.A. Chestnov, Pisma Zh. Eksp. Teor. Fiz. 17 (1973) 204; JETP Lett. (USSR) 17 (1973) 145
 1973HA46 R.D. Haracz and T.K. Lim, Phys. Rev. Lett. 31 (1973) 1263
 1973HA49 P.S. Hauge and S. Maripuu, Phys. Rev. C8 (1973) 1609
 1973HA50 M.L. Halbert, A. van der Woude and N.M. O'Fallon, Phys. Rev. C8 (1973) 1621
 1973HA51 R.A. Hardekopf, D.D. Armstrong, W. Gruebler, P.W. Keaton, Jr. and U. Meyer-Berkhout, Phys. Rev. C8 (1973) 1629
 1973HE06 J.C.P. Heggie and P.W. Martin, Phys. Lett. B43 (1973) 289
 1973HE26 J.C.P. Heggie and P.W. Martin, Nucl. Phys. A212 (1973) 78
 1973HO1U Houdayer and Van Oers, Phys. in Canada 29 (1973) 3
 1973JA1M Jarnie, in Munich 1 (1973) 332

- 1973JE02 H. Jeremie and R. Larose-Poutissou, Phys. Lett. B44 (1973) 68
- 1973JO1J John and Seigelman, in Munich 1 (1973) 500
- 1973JU02 M. Jung, Y. Sakamoto, J.N. Suren, C. Jacquot, L. Girardin and R. Schmitt, Phys. Rev. C7 (1973) 2209
- 1973KA08 S. Kato and Y. Aoki, J. Phys. Soc. Jpn. 34 (1973) 1128
- 1973KA32 D. Kamke, Z. Phys. 263 (1973) 251
- 1973KI1L Kirkby et al, in Contrib., Uppsala (1973) 176
- 1973LA14 E. Lambert and H. Feshbach, Ann. Phys. (New York) 76 (1973) 80
- 1973LI02 R.H. Lindsay, W. Toews and J.J. Veit, Nucl. Phys. A199 (1973) 513
- 1973LI19 T.K. Lim, Phys. Rev. Lett. 31 (1973) 1258
- 1973LI1K Lisowski et al, Bull. Amer. Phys. Soc. 18 (1973) 697
- 1973LI1L Lisowski, Trainor, Busch and Clegg, Bull. Amer. Phys. Soc. 18 (1973) 699
- 1973LI1M Ligatto de Slobodrian, Phys. in Canada 29 (1973) 4
- 1973LO1D Lovoi, Ohlsen, Salzman and Gruebler, Bull. Amer. Phys. Soc. 18 (1973) 699
- 1973LY02 G.I. Lykasov and A.V. Tarasov, Yad. Fiz. 17 (1973) 301; Sov. J. Nucl. Phys. 17 (1974) 153
- 1973MA20 H.A. Mavromatis, Nucl. Phys. A206 (1973) 477
- 1973MC05 D.K. McDaniels, M. Drog, J.C. Hopkins and J.D. Seagrave, Phys. Rev. C7 (1973) 882
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973NI1B Nisley et al, Bull. Amer. Phys. Soc. 18 (1973) 552
- 1973OH02 G.G. Ohlsen, G.C. Salzman, C.K. Mitchell, W.G. Simon and W. Gruebler, Phys. Rev. C8 (1973) 1639
- 1973PL02 G.R. Plattner, R.D. Viollier, D. Trautmann and K. Alder, Nucl. Phys. A206 (1973) 513
- 1973RA1E Rayet, Nucl. Phys. B57 (1973) 269
- 1973RO1R Robson, Nucl. Phys. A204 (1973) 523
- 1973RO1Z Roos et al, Bull. Amer. Phys. Soc. 18 (1973) 20
- 1973SA09 A.M. Saperstein, Phys. Rev. Lett. 30 (1973) 1257
- 1973SA1W Sagle et al, Bull. Amer. Phys. Soc. 18 (1973) 699
- 1973SL03 I. Slaus, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, J.M. Lambert and D.L. Shannon, Phys. Rev. C8 (1973) 444
- 1973TO06 W. Tornow, G. Mack, G. Mertens and H. Spiegelhauer, Phys. Lett. B44 (1973) 53

- 1973TR04 P.A. Treado, J.M. Lambert, R.J. Kane, L.A. Beach, E.L. Petersen and R.B. Theus, Phys. Rev. C7 (1973) 1742
- 1973TR1H Treado et al, Bull. Amer. Phys. Soc. 18 (1973) 1381
- 1973UL1C Ullo and Feshbach, in Contrib., Uppsala (1973) 95
- 1973VO1M Votta et al, Bull. Amer. Phys. Soc. 18 (1973) 19
- 1973WA13 R.E. Warner and E.W. Vogt, Nucl. Phys. A204 (1973) 433
- 1973YO1F Young and Redish, Bull. Amer. Phys. Soc. 18 (1973) 548
- 1974CH02 F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Rev. C9 (1974) 56
- 1974PR1D P.P. Pronko and J.G. Pronko, Phys. Rev. B9 (1974) 2870