

Energy Levels of Light Nuclei $A = 5$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A320* (1979), p. 1. This version of $A = 5$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed in 1978)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 5$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^5\text{n}\$](#) , [\${}^5\text{H}\$](#) , [\${}^5\text{He}\$](#) , [\${}^5\text{Li}\$](#) , [\${}^5\text{Be}\$](#)

B. Tables of Recommended Level Energies:

[Table 5.1](#): Energy levels of ${}^5\text{He}$

[Table 5.3](#): Energy levels of ${}^5\text{Li}$

C. [References](#)

D. Figures: [\${}^5\text{He}\$](#) , [\${}^5\text{Li}\$](#) , [Isobar diagram](#)

⁵n
(Not illustrated)

⁵n has not been observed: see (1972AG01) and (1977DE08).

⁵H
(Fig. 3)

From the work of (1967AD05) on the ³He(³He, n)⁵Be reaction (see ⁵Be) it follows that ⁵H is unstable by more than 2.1 MeV to decay into ³H + 2n (using Coulomb corrections based on the 16.7 MeV states in ⁵He–⁵Li). A study of ⁹Be(α, ⁸B)⁵H at $E_\alpha = 129$ MeV shows no evidence for sharp ⁵H states for several MeV above ³H + 2n (1968MC02). In ³H(t, p)⁵H, at $E_t = 22.25$ MeV, a broad peak appears in the proton spectrum which may correspond to a ⁵H state at ³H + 2n + 1.8 MeV, but which could also result from a four-body breakup (1968YO06). A recent study of the ⁷Li(⁶Li, ⁸B)⁵H reaction at $E(^6\text{Li}) = 93.3$ MeV shows no evidence for the formation of a narrow ⁵He ground state: at $\theta_{\text{lab}} = 14.7^\circ$ the yield near the predicted location of the ⁵H_{g.s.}, above the phase-space curve, is ≈ 100 nb/sr. MeV compared to an order of magnitude greater yield for the final state interaction in ⁴H (1977WE1B, 1977WE03). Attempts to observe ⁵H formation in ⁷Li + π⁻ have also been unsuccessful: see (1974AJ01). See also (1975BE31; theor.).

⁵He
(Figs. 1 and 3)

GENERAL: (See also (1974AJ01).)

Model calculations: (1974JA30, 1974LE22, 1975DI04).

Special states: (1974GO13, 1974IR04, 1974JA30, 1974LE22, 1975DI04, 1976IR1B).

Special reactions: (1973FE1A, 1974FE1A, 1975FE1A, 1976VA29, 1978ME1C).

Reactions involving pions: (1973BA30, 1978FI1E).

Other topics: (1974GO13, 1974IR04, 1976BI1A, 1976IR1B, 1977SH1B, 1978GO1D).

Ground state of ⁵He: (1975BE31, 1977HI09).

1. ³H(d, γ)⁵He $Q_m = 16.70$

At low energies the reaction is dominated by a resonance at $E_d = 107$ keV; the mirror reaction shows resonance at $E_d = 430$ keV. (1970BE1A) have measured the cross section for emission of 16.7 MeV γ-rays for $E_d = 25$ to 100 keV: the ratio $\sigma(d, \gamma)/\sigma(d, n)$ is approximately constant at

Table 5.1: Energy levels of ${}^5\text{He}$ ^a

E_x (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	0.60 ± 0.02 ^a	n, α	1, 4, 6, 8, 9, 10, 11, 12, 13, 14, 15, 18, 19, 20, 21, 22, 23, 26, 27, 29, 30
4 ± 1	$\frac{1}{2}^-; \frac{1}{2}$	4 ± 1	n, α	4, 6, 10, 18, 22
16.76 ± 0.13	$\frac{3}{2}^+; \frac{1}{2}$	0.10 ± 0.05	γ , n, d, t, α	1, 2, 5, 6, 8, 9, 11, 13, 14, 15, 21, 22
19.8 ± 0.4 ^b	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	2.5 ± 0.5	n, d, t, α	2, 3, 5, 8, 9, 10, 13, 14, 15, 19, 21, 22
$24 - 25$ ^b		broad		21, 22

^a See Table 5.2 in (1966LA04) and reactions 6 and 20 here.

^b See (1974AJ01), pp. 7 – 8.

$(2.1 \pm 0.6) \times 10^{-4}$, leading to $\Gamma_\gamma = 14 \pm 4$ eV, where Γ_n is taken as 66 keV. (1963BU07) have measured thick target yields from $E_d = 150$ to 1300 keV. The derived cross sections are analyzed into resonant and direct-capture contributions: the cross section at resonance is here reported as 60 μb [versus 1 mb reported by (1970BE1A)]. At $E_d = 1025 \pm 47$ keV, the differential cross section is 0.44 ± 0.12 $\mu\text{b}/\text{sr}$ (90°) and the γ to n branching ratio is an order of magnitude smaller: 2.3×10^{-5} . The angular distribution of the γ -rays is forward peaked and the total cross section is estimated to be 4.8 μb (1970KO09). The cross section for γ_0 has been measured (at 90°) for $E_d = 2.0$ to 12.0 MeV and angular distributions were obtained at $E_d = 4, 8, 10$ and 12 MeV for γ_0 (1975BA1E; and C.A. Barnes, private communication). See also (1974AJ01).

2. (a) ${}^3\text{H}(d, n){}^4\text{He}$ $Q_m = 17.5894$ $E_b = 16.70$
 (b) ${}^3\text{H}(d, 2n){}^3\text{He}$ $Q_m = -2.9884$
 (c) ${}^3\text{H}(d, pn){}^3\text{H}$ $Q_m = -2.2246$

Below $E_d = 100$ keV, the cross section for reaction (a) follows the Gamow function, $\sigma = (A/E)\exp(-44.40E^{-1/2})$ (1953JA1A, 1954AR02). A strong resonance, $\sigma(\text{peak}) = 5.0$ b, appears at $E_d = 107$ keV: see Table 5.2. From $E_d = 10$ to 500 keV, the cross section is well fitted with the assumption of s-wave formation of a $J^\pi = \frac{3}{2}^+$ state (1952AR1A, 1952CO35, 1955KU03). Excitation curves and angular distributions for reaction (a) have been measured from $E_d = 8$ keV to 21 MeV: see (1974AJ01) for earlier references and (1976DR1A, 1976DR1B; preliminary work; angular distributions at $E_d = 7.00$ and 10.00 MeV, 13.36 and 16.50 MeV (partial) and at

Table 5.2: Resonance parameters for the $\frac{3}{2}^+$ states observed in ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$ and ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$ ^a

E_{r} (keV)	Γ_{lab} (keV)	l_{d}	J^{π}	l_{np}	R (fm)	E_{λ} (keV)	γ_{d}^2 (keV)	γ_{np}^2 (keV)	θ_{d}^2 ^d	θ_{np}^2 ^d	E_{x} (MeV)
107 ^b	135	0	$\frac{3}{2}^+$	2	5.0	-464	2000 ± 500	50 ± 10	1.0	0.018	16.76
450 ^c	≈ 450	0	$\frac{3}{2}^+$	2	7.0	-126	715	17	0.7	0.011	16.66
					5.0	-391	2930	42	1.4	0.013	
					7.0	129	780	12	0.7	0.008	

^a See also (1974KR07, 1976AH1A, 1977BO24, 1977DO01).

^b ${}^3\text{H}(\text{d}, \text{n}){}^4\text{He}$.

^c ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$.

^d Units of $3\hbar^2/2MR^2$.

$E_{\text{t}} = 10.48$ and 14.98 MeV; 0° excitation function for $E_{\text{d}} = 7$ to 16.5 MeV), (1975MA28; angular distributions at $E_{\text{d}} = 8$ to 16 MeV, in 1 MeV steps; 0° excitation function for same interval) and (1977JA07; $d\sigma/d\Omega$ at $E_{\text{t}} = 20.00$ MeV; very accurate).

A study of reaction (a) with polarized deuterons at $E_{\text{d}} = 0.2$ to 1.0 MeV indicates intervention of the s-wave, $J^{\pi} = \frac{1}{2}^+$ channel, as well as possible p-waves above $E_{\text{d}} = 0.3$ MeV (1965TR01, 1971GR32). At higher energies, the neutron polarization $P_1(\theta_1)$ shows an angular distribution that peaks typically at $\theta_1 = 30^\circ$ lab, then goes negative with increasing angle and has a minimum between 90° and 130° , depending upon energy. (1971MU04) have made an extensive study of $P_1(30^\circ)$ for $E_{\text{d}} = 5$ to 15 MeV and, with deuterium as target, for $E_{\text{t}} = 4.5$ to 19.5 MeV (neutrons near 135° lab). The polarization increases monotonically from 0.03 at $E_{\text{d}} = 3$ MeV to ≈ 0.5 at $E_{\text{d}} = 6.5$ MeV and then with a lower slope to 0.69 at $E_{\text{d}} = 13$ MeV. The change in the slope may be caused by excited states of ${}^5\text{He}$ near 20 MeV. Comparison with the ${}^3\text{He}(\text{d}, \text{p}){}^4\text{He}$ mirror reaction at corresponding c.m. energies shows excellent agreement between the polarization values in the two reactions up to $E_{\text{d}} = 6$ MeV, but then the proton polarization becomes $\approx 15\%$ higher, converging back to the neutron values at $E_{\text{d}} \approx 12 - 13$ MeV. This may be due to experimental factors (1971MU04). Using polarized deuterons (1971HI07) find that the average ratio of vector analyzing power of ${}^3\text{He}(\text{d}, \text{p})$ to ${}^3\text{H}(\text{d}, \text{n})$ is 1.016 ± 0.015 at $E_{\text{d}} = 6$ MeV and 1.035 ± 0.020 at 10 MeV. The vector analyzing power of the two reactions agree to within ± 0.025 at all angles. This agreement between the mirror reactions is in apparent conflict with the result of (1971MU04). For earlier polarization measurements see Table 5.3 in (1974AJ01). Polarization studies are also reported by (1976SU06; $E_{\text{d}} = 7$ MeV), (1976LI15; $E_{\text{d}} = 3.5$ to 12.8 MeV), (1976OH02; $E_{\text{t}} = 5.37$ to 10.50 MeV) and (1975SA1A; preliminary; $E_{\text{d}} = 38$ MeV; looked at pol. of 50 MeV neutrons). See also (1977DR1B). The $K_{\text{y}}^{\text{y}'}(0^\circ)$ determined for $E_{\text{t}} = 5.37$ to 10.50 MeV show a minimum at $E_{\text{t}} = 9.6$ MeV (1976OH02). See also (1978CA13; $E_{\text{d}} = 0.7$ MeV).

Reaction (b) has been studied for $E_{\text{d}} = 10.9$ to 83 MeV: see (1974AJ01). A discussion and comparison of attempts to measure ${}^1\text{S}_0$ nn scattering length is presented by (1971GR45; $E_{\text{d}} = 13.4$

MeV): based on kinematically complete measurements and using the Watson-Migdal treatment, they find $a_{nn} = -16.0 \pm 1.0$ fm. Intervention of n - ${}^3\text{He}$ final state interactions is believed to be small in this work (1971GR45). At $E_t = 22$ MeV no clear evidence is seen for sequential processes via excited states of ${}^4\text{He}$ (1971GR12).

A reanalysis of the work of (1963PO02) on reaction (c) by (1974SC04) leads to the suggestion of a resonance at $E_{c.m.} = 2.9 \pm 0.3$ MeV [$E_x = 19.6$ MeV], $\Gamma_{c.m.} = 1.9 \pm 0.2$ MeV, consistent with $J^\pi = \frac{3}{2}^-$ [see, however, Table 5.1]. n - p final-state interaction enhancement has been studied by (1973SL03) at $E_d = 35$ MeV. They find that only when $E_{np} \leq 1$ MeV is in agreement with the Watson-Migdal theory obtained, even if data far from dominant n - t final-state interaction and p - t quasi-free scattering are chosen. See also (1974AL01).

See also (1974AD11), (1974GE1B, 1974WI1D, 1975CR1A, 1975ST1A, 1975ST1B, 1976BL1A, 1976BR1B, 1976JA1C, 1976MO1A, 1977BE1H, 1977KI1C, 1977MC1C; applied topics), (1973LI1B, 1975KU1C, 1975SE07, 1975SO1C, 1976HA1C, 1976SE1B, 1976WA1C, 1977SE09; reviews), (1975FO19; astrophysics) and (1974KE1B, 1974ST14, 1975DO1B; theor.).

3. ${}^3\text{H}(d, d){}^3\text{H}$

$$E_b = 16.70$$

The elastic scattering has been studied for $E_d = 2.6$ to 11.0 MeV. The excitation curves show an interference at $E_x \approx 19$ MeV and a broad ($\Gamma > 1$ MeV) resonance corresponding to $E_x = 20.0 \pm 0.5$ MeV, similar to that seen in ${}^3\text{He}(d, d)$ [see ${}^5\text{Li}$]. Together with data from ${}^3\text{H}(d, n){}^4\text{He}$, this work favors an assignment $D_{3/2}$ or $D_{5/2}$ with a mixture of doublet and quartet components (channel spin $\frac{1}{2}$ and $\frac{3}{2}$) if only one state is involved (1967TO02, 1968IV01) [any appreciable doublet component would, however, be in conflict with results from ${}^7\text{Li}(p, {}^3\text{He}){}^5\text{He}$]. Measurements of differential cross section and analyzing power using polarized deuterons with $E_d = 3.2$ to 12.3 MeV show resonance-like behavior in the vector analyzing power near $E_d = 5$ MeV. The anomaly appears in the odd Legendre coefficients and is interpreted in terms of a $(\frac{1}{2}, \frac{3}{2})^-$ excited state of ${}^5\text{He}$ with $E_x \approx 19.6$ MeV. Broad structure in the differential cross section near 6 MeV, principally in the even Legendre coefficients, corresponds to an even-parity state ${}^5\text{He}^*(20.0)$ (1971KI02). Tensor analyzing powers have been measured for $E_d = 5.0$ to 11.5 MeV (1973DE51), at $E_t = 10.5$ MeV (1976OH02) and at $E_d = 11.4$ and 14.4 MeV (1974DO05; angular distributions of tensor and vector polarizations). See also (1974ST14, 1975AB1C, 1975DO1B, 1976BA1E; theor.).

$$4. (a) {}^3\text{H}(t, n){}^5\text{He} \quad Q_m = 10.44$$

$$(b) {}^3\text{H}(t, 2n){}^4\text{He} \quad Q_m = 11.3321$$

At $E_t = 0.5$ MeV, the reaction appears to proceed via three channels: (i) direct breakup into ${}^4\text{He} + 2n$, the three-body breakup shape being modified by the n - n interaction; (ii) sequential decay via ${}^5\text{He}(0)$; (iii) sequential decay via a broad excited state of ${}^5\text{He}$. The branching ratios at $\theta = 90^\circ$ are 0.7 : 0.2 : 0.1. The width of ${}^5\text{He}(0)$ is estimated to be 0.74 ± 0.18 MeV. Some evidence is

also shown for ${}^5\text{He}^*$ at $E_x \approx 2$ MeV, $\Gamma \approx 2.4$ MeV (1965WO03). For reaction (b), see ${}^6\text{He}$ and (1974LA02).

5. (a) ${}^3\text{He}(t, p){}^5\text{He}$ $Q_m = 11.20$
 (b) ${}^3\text{He}(t, pn){}^4\text{He}$ $Q_m = 12.0959$

Some evidence is reported at $E_t = 22.25$ MeV in reaction (a) for a broad state of ${}^5\text{He}$ at $E_x \approx 20$ MeV, in addition to a sharp peak corresponding to ${}^5\text{He}^*(16.7)$ (1968YO06). For reaction (b) see ${}^6\text{Li}$ and (1975SC28, 1977JO1A).

6. ${}^4\text{He}(n, n){}^4\text{He}$ $E_b = -0.89$

The coherent scattering length (thermal, bound) is 3.0 ± 0.1 fm (1973MU14). The thermal scattering cross section is 0.773 ± 0.009 b and the absorption cross section at 2200 m/sec $\sigma_{n,\gamma}^0$ is < 0.05 b (1969RO16). (1973MU14) adopt 0.76 ± 0.01 b for $\bar{\sigma}_s$. Total cross sections for $E_n = 4 \times 10^{-4}$ eV to 150.9 MeV and at 10 GeV/c are summarized in (1974AJ01, 1976GAYV). Angular distribution measurements are summarized in (1970GA1A, 1974AJ01). Polarization studies are displayed in Table 5.4 of (1974AJ01): recent work includes that of (1976BO05: $E_n = 1.5$ to 6.0 MeV; used an absolutely calibrated source of polarized neutrons), (1976LI15: $E_n = 20$ to 30 MeV; $A_y(\theta)$ is the same for ${}^4\text{He} + n$ and ${}^4\text{He} + p$) and (1978YO1A: $E_n = 50$ MeV). See also (1974TO03).

The total cross section has a peak of 7.6 b (1973GO38) at $E_n = 1.15 \pm 0.05$ MeV, $E_{c.m.} = 0.92 \pm 0.04$ MeV, with a width of about 1.2 MeV (1964ST25). A second resonance is observed at $E_n = 22.15 \pm 0.12$ MeV, corresponding to the 16.7 MeV $J^\pi = \frac{3}{2}^+$ state (1959BO54, 1964SH1A): $\Gamma_{c.m.} = 100 \pm 50$ keV, $\Gamma_n = \Gamma_d = 50 \pm 35$ keV (1960HU1A). [(1966HO07) find that the data are fitted best by $\Gamma_n < \Gamma_d$ although $\Gamma_n < \Gamma_d$ is not excluded]. Attempts to detect additional resonances in the total cross section have been unsuccessful: see (1966LA04).

The $P_{3/2}$ phase shift shows strong resonance behavior near 1 MeV, while the $P_{1/2}$ shift changes more slowly, indicating a broad $P_{1/2}$ level at several MeV excitation (1952DO30, 1966HO07, 1970AR1B, 1972ST01). (1966HO07) have constructed a set of phase shifts for $E_n = 0$ to 31 MeV, $l = 0, 1, 2, 3$, using largely p- α phase shifts. They have measured differential cross sections from 6 to 30 MeV, with special attention to the region near 22.15 MeV and have fitted the data with the assumed phase shifts. At the $\frac{3}{2}^+$ state the best fit to all data is given by $E_{\text{res}} = 17.669$ MeV ± 10 keV, $\gamma_d^2 = 2.0$ MeV $\pm 25\%$, $\gamma_n^2 = 50$ keV $\pm 25\%$ (see Table 5.2). The polarization has been calculated from the phase shifts and is presented as a contour plot (1966HO07). The work of (1976LI15) indicates some discrepancies with the results of (1966HO07) [below $E_n = 22$ MeV] and with the measurements of (1972BR10).

A recent R -function analysis of the ${}^4\text{He} + n$ data below 21 MeV (including the absolute neutron analyzing power measurement of (1976BO05) and the accurate cross section measurements of

(1973GO38)) has led to a set of phase shifts and analyzing powers which are based on the ${}^4\text{He} + n$ data alone (rather than also including the ${}^4\text{He} + p$ data). At $r = 3.3$ fm the values obtained for the $P_{1/2}$ and $P_{3/2}$ resonances are, respectively, $E_{c.m.} = 1.97$ and 0.77 MeV, $\Gamma_{c.m.} = 5.22$ and 0.64 MeV (1977BO24). See also (1974KR07). For earlier analyses, see (1974AJ01).

See also (1974MA1C, 1977YO1B), (1975FI1B, 1976OH1B, 1976WA1B, 1976WA1C) and (1974BA34, 1974DO10, 1974GH01, 1974JA1F, 1974ST14, 1975BA76, 1975CA05, 1975LO1A, 1975TH12, 1976AH1A, 1976AL1E, 1976CH1B, 1976SP1A, 1977BA1N, 1977BE50, 1977CH1C, 1977LA1C, 1977LE1D, 1977PH01, 1977ST1D, 1977TH09, 1978AH01, 1978FO1G, 1978FR02, 1978KO13, 1978LE1H, 1978MA37; theor.).

$$7. {}^4\text{He}(n, d){}^3\text{H} \qquad Q_m = -17.5894 \qquad E_b = -0.89$$

See (1976BR1B), (1974AJ01) and (1976BA1E; theor.).

$$8. \text{(a) } {}^4\text{He}(d, p){}^5\text{He} \qquad Q_m = -3.12$$

$$\text{(b) } {}^4\text{He}(d, pn){}^4\text{He} \qquad Q_m = -2.2246$$

A typical proton spectrum consists of a peak corresponding to formation of the ground state of ${}^5\text{He}$, plus a lower continuum of protons ascribed to deuteron breakup (reaction (b)). Ground-state protons show pronounced azimuthal asymmetry when the reaction is induced by 8.5, 10 and 11 MeV vector polarized deuterons. A DWBA calculation is in only qualitative agreement (1967TR05, 1971KE16).

Reaction (b) has been studied for $E = 9$ to 165 MeV: see (1974AJ01) and $E_d = 6.8$ and 7.8 MeV (1978KA11), 8.9 MeV (1977SA21), at $E_d^- = 15$ MeV (1978NA08), $E_\alpha = 21.9$ and 23.7 MeV (1974RA10), 22.5 MeV (1977PR06), 27.2 MeV (1974GR40) and 165 MeV (1974HO11). See also ${}^6\text{Li}$.

At $E_d = 8.9$ MeV, the FSI (final state interaction) is very important (1977SA21). In the kinematically complete experiment of (1977PR06) at $E_\alpha = 22.5$ MeV, the results are in quite good agreement with the predictions of the sequential decay model: the slight deviations are assumed to be due to the omission of Coulomb interaction effects and to the restriction to partial waves with $l \leq 2$. At $E_\alpha = 70$ MeV, a kinematically complete experiment shows evidence for sequential decay, proceeding through excited states of ${}^5\text{He}$. Peaks in the coincident yield of protons and deuterons are ascribed to narrow states at $E_x = 16.7 \pm 0.1$ MeV, $\Gamma = 80 \pm 30$ keV, at $E_x = 18.6 \pm 0.1$, 18.8 ± 0.1 and 19.2 ± 0.1 MeV, all with $\Gamma = 180 \pm 60$ keV (1973TR04). The fine structure near 19 MeV is not confirmed in other experiments [see, however, reaction 12].

See also (1974HE21, 1974NA10, 1975HE09, 1976KO21, 1976LI1E, 1977BR25, 1978FI1E, 1978KO13), ${}^5\text{Li}$ and ${}^6\text{Li}$.

9. ${}^4\text{He}(t, \text{nd}){}^4\text{He}$ $Q_m = -0.4105$

At $E_\alpha = 70$ MeV, there is indication of the population of ${}^5\text{He}^*(0, 16.7)$ and of a structure with $E_x = 18 - 20$ MeV (1975AL1A, 1974LA1A; preliminary results).

10. ${}^6\text{Li}(\gamma, \text{p}){}^5\text{He}$ $Q_m = -4.59$

At $E_\gamma = 60$ MeV, the proton spectrum shows two prominent peaks attributed to ${}^5\text{He}^*(0 + 4.0, 20 \pm 2)$ (1973GA16, 1976MA34). See also (1973DE17), (1975MA1E) and ${}^6\text{Li}$.

11. ${}^6\text{Li}(\text{e}, \text{ep}){}^5\text{He}$ $Q_m = -4.59$

At $E_p = 1180$ MeV, the excitation of ${}^5\text{He}^*(0, 16.7)$ is reported: the latter state is formed with the ejection of an s-proton (1972AN27, 1972AN24, 1972AN29). See also (1975ME27; theor.).

12. (a) ${}^6\text{Li}(\text{n}, \text{d}){}^5\text{He}$ $Q_m = -2.37$

(b) ${}^6\text{Li}(\text{n}, \text{nd}){}^4\text{He}$ $Q_m = -1.4735$

Angular distributions of ground state deuterons have been studied at $E_n = 6.57$ and 6.77 MeV (1977RO01), 14.4 MeV (1965VA05) and 56.3 MeV (1977BR17). In the latter experiment, angular distributions have also been obtained to ${}^5\text{He}^*(16.7, 18.5 \pm 0.5, 20.5 \pm 0.5)$. The observation of the two highest states is not certain: if they exist their widths are less than the instrumental width, 1.6 MeV (1977BR17). For reaction (b) see (1967VA12). See also (1974AJ01).

13. ${}^6\text{Li}(\text{p}, \text{d}\pi^+){}^5\text{He}$ $Q_m = -142.72$

See (1976HU1A).

14. ${}^6\text{Li}(\text{p}, 2\text{p}){}^5\text{He}$ $Q_m = -4.59$

At $E_p = 100$ MeV the population of ${}^5\text{He}^*(0, 16.7)$ and possibly of a broad structure at $E_x \approx 19$ MeV is observed: momentum distributions for ${}^5\text{He}^*(0, 16.7)$ and angular correlation measurements are also reported. The main features of the data are reasonably well described by DWIA (1974BH03). For the earlier work see (1974AJ01). See also (1975VO04, 1977RO1E) and (1974GH03, 1974PR10, 1975CH1C, 1976OH04; theor.).

15. (a) ${}^6\text{Li}(\text{d}, {}^3\text{He}){}^5\text{He}$ $Q_{\text{m}} = 0.90$
 (b) ${}^6\text{Li}(\text{d}, \text{n}{}^3\text{He}){}^4\text{He}$ $Q_{\text{m}} = 1.795$

At $E_{\text{d}} = 14.5$ MeV, ${}^5\text{He}_{\text{g.s.}}$ is populated (1955LE24). At $E_{\text{d}} = 80$ MeV, the population of ${}^5\text{He}^*(16.7)$ as well as a broad state near $E_{\text{x}} = 19$ MeV is reported (1975DI1A, 1975DI1B; abstracts). For reaction (b) see (1974AJ01).

16. ${}^6\text{Li}(\text{t}, \alpha){}^5\text{He}$ $Q_{\text{m}} = 15.22$

See (1974AJ01).

17. (a) ${}^6\text{Li}(\alpha, {}^5\text{Li}){}^4\text{He} + \text{n}$ $Q_{\text{m}} = -5.66$
 (b) ${}^6\text{Li}(\alpha, \alpha\text{p}){}^5\text{He}$ $Q_{\text{m}} = -4.59$

See (1975GR42) for reaction (a) and (1978CA1E) for reaction (b).

18. ${}^7\text{Li}(\gamma, \text{d}){}^5\text{He}$ $Q_{\text{m}} = -9.62$

See (1975DE37) in ${}^7\text{Li}$.

19. (a) ${}^7\text{Li}(\pi^+, 2\text{p}){}^5\text{He}$ $Q_{\text{m}} = 128.51$
 (b) ${}^7\text{Li}(\pi^-, 2\text{n}){}^5\text{He}$ $Q_{\text{m}} = 126.94$

The two-proton spectrum shows broad structures attributed to the ground state of ${}^5\text{He}$, and to $\text{p}^{-1}\text{s}^{-1}$ and s^{-2} states at ≈ 20 and (≈ 35) MeV excitation (1965CH12). For reaction (b) see (1977BA1M).

20. (a) ${}^7\text{Li}(\text{n}, \text{t}){}^5\text{He}$ $Q_{\text{m}} = -3.36$
 (b) ${}^7\text{Li}(\text{n}, \text{tn}){}^4\text{He}$ $Q_{\text{m}} = -2.467$

The angular distribution of the t_0 group has been measured at $E_{\text{t}} = 14.4$ MeV: it is fairly well reproduced by DWBA (1970MI05). Reaction (b) at $E_{\text{n}} = 14.4$ MeV proceeds as a sequential process primarily involving ${}^7\text{Li}^*(4.63)$ and ${}^5\text{He}_{\text{g.s.}}$ (1974AN02). See also (1974TU1A) and (1974AJ01) for the earlier work.

21. (a) ${}^7\text{Li}(p, {}^3\text{He}){}^5\text{He}$ $Q_m = -4.12$
 (b) ${}^7\text{Li}(p, pd){}^5\text{He}$ $Q_m = -9.618$

At $E_p = 43.7$ MeV, angular distributions of the ${}^3\text{He}$ groups to the ground state of ${}^5\text{He}$ ($\Gamma = 0.80 \pm 0.04$ MeV) and to levels at 16.7 and 19.9 ± 0.4 MeV ($\Gamma = 2.7$ MeV) have been determined. The angular distribution of the ${}^5\text{He}$ ground state group indicates substantial mixing of $L = 0$ and $L = 2$ transfer. The distribution to ${}^5\text{He}^*(16.7)$ is consistent with $L = 1$. Since no transitions are observed in the ${}^7\text{Li}(p, t){}^5\text{Li}$ reaction to the analogue 20 MeV state in ${}^5\text{Li}$ [see ${}^5\text{Li}$], the transition is presumably S -forbidden and the states in ${}^5\text{He}$ - ${}^5\text{Li}$ near 20 MeV are ${}^4D_{3/2}$ or ${}^4D_{5/2}$ [compare ${}^3\text{H}(d, d)$] (1966CE05). Particle-particle coincidence data have been obtained at $E_p = 43.7$ MeV. They suggest the existence of ${}^5\text{He}^*(20.0)$ with $\Gamma = 3.0 \pm 0.6$ MeV and of a broad state at ≈ 25 MeV. No $T = \frac{3}{2}$ states decaying via $T = 1$ states in ${}^4\text{He}$ were observed (1968MC02). In reaction (b) at $E_p = 100$ MeV ${}^5\text{He}^*(0, 19.9)$ are strongly populated while ${}^5\text{He}^*(16.7)$ is very weak. Positive parity is suggested for ${}^5\text{He}^*(19.9)$ (1975CH1B; preliminary).

22. (a) ${}^7\text{Li}(d, \alpha){}^5\text{He}$ $Q_m = 14.23$
 (b) ${}^7\text{Li}(d, n){}^4\text{He}{}^4\text{He}$ $Q_m = 15.123$

At $E_d = 24$ MeV, the α -particle spectrum from reaction (a) shows structures corresponding to the ground and 16.7 MeV states and to states at $E_x \approx 20.2$ and 23.8 MeV with $\Gamma \approx 2$ MeV and ≈ 1 MeV respectively (1972BA30). The width of the ground state of ${}^5\text{He}$ is 615 ± 30 keV (1973GA28). Angular distributions are reported for $E_d = 0.6$ to 1.25 MeV.

Spectra measured in reaction (b) suggest the involvement of the $P_{1/2}$ state: the values suggested are inconsistent with each other because of the difficulty of evaluating the contribution of other reactions: see (1974AJ01) for the earlier values and see (1976FO21) for a discussion of some of the problems involved in these. (1976FO21) suggest a width for ${}^5\text{He}_{g.s.} = 0.6$ MeV and $E_x = 4.1 \pm 0.2$, $\Gamma_{c.m.} = 4.4 \pm 0.2$ MeV for the $P_{1/2}$ state. Reaction (b) proceeds mainly via excited states of ${}^8\text{Be}$ and ${}^5\text{He}_{g.s.}$: see, e.g., (1967VA11). See also (1973HU12, 1974DA28, 1974GR44, 1978AR10, 1978SP03). Attempts to observe n- α rescattering effects following formation of ${}^8\text{Be}^*(16.63, 16.91)$ have been unsuccessful: an upper limit of 1% of the yield from sequential decay is given by (1968VA12). See (1974AJ01) for more complete references to the older work. See also ${}^8\text{Be}$ and ${}^9\text{Be}$.

23. ${}^7\text{Li}({}^3\text{He}, d{}^3\text{He}){}^5\text{He}$ $Q_m = -9.62$

See (1976WA12).

24. ${}^7\text{Li}(\alpha, {}^6\text{Li}){}^4\text{He} + n$ $Q_m = -7.251$

See (1975GR42).

$$25. \text{}^9\text{Be}(\gamma, \alpha)^5\text{He} \quad Q_m = -2.47$$

See ${}^9\text{Be}$.

$$26. \text{}^9\text{Be}(p, p\alpha)^5\text{He} \quad Q_m = -2.47$$

Studies of this reaction have been carried out at $E_p = 26.0$ to 57 MeV [see (1974AJ01)] and at $E_p = 100$ MeV (1977RO02). In the latter experiment good agreement is found with DWIA calculations: the average value of $S_\alpha = 0.45 \pm 0.02$ (1977RO02). See also ${}^9\text{Be}$.

$$27. \text{}^9\text{Be}(t, {}^7\text{Li})^5\text{He} \quad Q_m = 0.096$$

See (1976VO1A).

$$28. \text{}^9\text{Be}({}^3\text{He}, \alpha){}^5\text{He} \quad Q_m = -2.47$$

See ${}^9\text{Be}$.

$$29. \text{(a) } \text{}^9\text{Be}(\alpha, 2\alpha)^5\text{He} \quad Q_m = -2.47$$
$$\text{(b) } \text{}^9\text{Be}(\alpha, {}^8\text{Be})^5\text{He} \quad Q_m = -2.56$$

Reaction (a) has been studied at $E_\alpha = 28$ to 37.4 MeV: see (1974AJ01). See also (1974GR42, 1977BR1E). Reaction (b) has been studied at $E_\alpha = 65$ MeV: only the ground state of ${}^5\text{He}$ is observed for $E_x \lesssim 25$ MeV; $S_\alpha = 0.53$ (1976WO11).

$$30. \text{}^{11}\text{B}(d, n){}^4\text{He}{}^4\text{He} \quad Q_m = 6.4575$$

At $E_d = 3.75$ MeV (1978GR07), 10.4 and 12.0 MeV (1971RE19), this reaction involves ${}^5\text{He}_{\text{g.s.}}$ and states in ${}^8\text{Be}$ and ${}^9\text{Be}$ (1971RE19, 1978GR07) and ${}^{12}\text{C}$ (1978GR07).

$$31. \text{}^{12}\text{C}(n, n){}^4\text{He}{}^4\text{He} \quad Q_m = -7.2748$$

See ${}^{12}\text{C}$ in (1975AJ02).

⁵Li
(Figs. 2 and 3)

GENERAL: (See also (1974AJ01).)

Model calculations: (1975KR1A).

Special states: (1974GO13, 1974IR04, 1976IR1B).

Astrophysical questions: (1974RA1C, 1978ME1C).

Special reactions: (1975BR1A, 1976VA29, 1978ME1C).

Reactions involving pions: (1973AR1B, 1974AM01).

Applied topics: (1975HU1A).

Other topics: (1974GO13, 1974IR04, 1976IR1B, 1978GO1D).

Ground state of ⁵Li: (1975BE31).

1. ³He(d, γ)⁵Li $Q_m = 16.39$

Excitation curves and angular distributions have been measured for $E_d = 0.2$ to 5 MeV and $E(^3\text{He}) = 2$ to 26 MeV: see (1974AJ01).

A broad maximum in the cross section is observed at $E_d = 0.45 \pm 0.04$ MeV [⁵Li*(16.66)]: $\sigma = 50 \pm 10 \mu\text{b}$, $\Gamma_\gamma = 11 \pm 2$ eV (1954BU06: $\gamma_0 + \gamma_1$); $\sigma_{\gamma_0} = 21 \pm 4 \mu\text{b}$, $\Gamma_{\gamma_0} = 5 \pm 1$ eV (1968BU09). The radiation at resonance is isotropic, consistent with s-wave capture: see (1954BU06, 1968BU09, 1968KR03). Study of γ_0 and γ_1 yield $\Gamma = 2.6 \pm 0.4$ MeV for the ground state width, and $E_x = 7.5 \pm 1.0$ MeV, $\Gamma = 6.6 \pm 1.2$ MeV for the $\frac{1}{2}^-$ state. The ratio of γ_0 to γ_1 is 1.00 ± 0.2 and 1.9 ± 0.4 at $E_d = 480$ and 1025 keV (1968BU09).

An excess in the cross section at higher bombarding energies is interpreted by (1972KI01) as being due to a state at $E_x \approx 18$ MeV: even parity is deduced from the relative intensity of γ_0 and γ_1 . It is presumed to be the $\frac{1}{2}^+$ state reported in reactions 3 and 7. A broad peak is also observed at $E_x \approx 20.7$ MeV in the γ_0 cross section. The cross section for γ_1 is ≈ 0 . The observations are consistent with $J^\pi = \frac{5}{2}^+$: angular distributions appear to require at least one other state with significant strength near 19 MeV (1972KI01). (1970SC18) also report this state but find $E_x = 19.7 \pm 0.2$ MeV [$\Gamma = 5.0$ MeV]: $(2J + 1)\Gamma_{\gamma_0} = 1.32$ keV ($\pm 50\%$), suggesting E1 (if only one state is involved).

2. ³He(d, n)⁴Li $E_b = 16.39$

Table 5.3: Energy levels of ${}^5\text{Li}$ ^a

E_x (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	≈ 1.5	p, α	1, 5, 6, 7, 9, 13, 14, 15, 16, 17
5 – 10	$\frac{1}{2}^-; \frac{1}{2}$	5 ± 2	p, α	1, 5, 7, 13, 15, 16, 17
16.66 ± 0.07	$\frac{3}{2}^+; \frac{1}{2}$	≈ 0.3	γ , p, d, ${}^3\text{He}$, α	1, 3, 4, 7, 13, 15
(18 ± 1)	$(\frac{1}{2}^+); \frac{1}{2}$	broad	γ , p, d, ${}^3\text{He}$, α	1, 3, 7
(20.0 ± 0.5)	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	≈ 5	γ , p, d, ${}^3\text{He}$, α	1, 3, 4, 5, 7, 13, 15
^a				

^a See also discussion in reactions 3, 4, 7, 14 and 15.

This reaction has not been observed: see (1966LA04) and (1974POZN; abstract). See also (1978MC1C).

3. (a) ${}^3\text{He}(d, p){}^4\text{He}$	$Q_m = 18.3532$	$E_b = 16.39$
(b) ${}^3\text{He}(d, np){}^3\text{He}$	$Q_m = -2.2246$	
(c) ${}^3\text{He}(d, 2p){}^3\text{H}$	$Q_m = -1.4608$	
(d) ${}^3\text{He}(d, 2d){}^1\text{H}$	$Q_m = -5.4936$	

Below 100 keV the cross section follows the simple Gamow form: $\sigma = (18.2 \times 10^3 / E) \exp(-91E^{-1/2})$ b (E in keV) (1954AR02, 1953JA1A). The zero-energy cross section factor $S_0 = 6700$ keV · b (1964PA1A). A pronounced resonance occurs at $E_d = 430$ keV, $\Gamma \approx 450$ keV. The peak cross section is given as 695 ± 14 mb (1955KU03, 1952BO68): see Table 5.2. Excitation functions for ground state protons have also been reported for $E({}^3\text{He}) = 0.39$ to 1.46 MeV and 18.7 to 44.1 MeV and for $E_d = 2.8$ to 17.8 MeV: see (1974AJ01). Angular distributions have been measured for $E_d = 0.25$ to 27 MeV and $E({}^3\text{He}) = 18.7$ to 44.1 MeV [see Table 5.6 in (1974AJ01)]. At $E_d = 10.02$ MeV differential cross sections have been measured at four angles to better than 3% (1974JA15). Polarization measurements have been carried out at many energies: see Table 5.6 in (1974AJ01) and Table 5.4 here.

Below $E_d = 1.1$ MeV, the polarization analyzing power of the reaction is minimally affected by p-wave contributions (less than a few percent). However, at $E_d = 500$ keV, the analyzing power is reduced by $(12 \pm 5)\%$ by intervention of the s-wave $J^\pi = \frac{1}{2}^+$ channel (1967MC01, 1966BR02, 1971LE27). Contour maps, T_{ab} versus θ , E_d , for the vector and tensor analyzing powers are presented for $E_d = 0$ to 12 MeV by (1971GR47). Energy dependence of the angular distributions indicate resonance-like behavior at $E_x = 16.6$, 17.5, 20.0, 20.9 and 22.4 MeV in ${}^5\text{Li}$ (1971GR47). At $E_d \approx 6$ MeV (1971KL02) find the angular distribution more complex than could be accounted for by a single ${}^2D_{3/2}$ or ${}^2D_{5/2}$ level at $E_x \approx 20$ MeV [see ${}^3\text{He}(d, d)$]. For

Table 5.4: Polarization measurements in ${}^3\text{He}(d, p){}^4\text{He}$ ^a

$E_{\vec{d}}$ (MeV)	Refs.
0.344, 0.465, 0.727	(1974GA21)
0.34 – 11.60	(1976SC15)
0.75, 1.5 ^b	(1976BR1C) ^c
1 – 6	(1976DR1C, 1977DR1B) ^c
6.44 ^d	(1977ST06)
6.6 – 15.8	(1974TR02)
8.5 – 10.5 ^e	(1976GR10)
12 ^{b,f}	(1974BE67)
15 ^g	(1976ME13)
15 – 40	(1975RO1C, 1976RO1G) ^c
27 ^h	(1976OK1A) ^c
30, 35, 40 ⁱ	(1976RO1H) ^c

^a See Table 5.6 in (1974AJ01) for a listing of the earlier work.

^b $E_{\vec{d}}$.

^c Preliminary results.

^d Measurement to better than 1% of iT_{11} .

^e Maximum value of the tensor analyzing power (A_{yy}) at 9.28 MeV is nearly 1.

^f ${}^3\vec{\text{He}}$ target.

^g Reactions 5(b) and 5(c).

^h $E({}^3\vec{\text{He}})$.

ⁱ Reaction 5(b).

$E_{\vec{d}}(\text{c.m.}) = 7$ to 18 MeV, differential cross sections fall monotonically with no indication of states of ${}^5\text{Li}$ with $23.4 < E_x < 34$, with $\Gamma > 0.8$ MeV and with appreciable widths for decay into ${}^3\text{He} + d$ (1972KI02).

Reactions (b), (c) and (d) have been studied at $E_{\vec{d}} = 22.3$ and 35 MeV and at $E({}^3\text{He}) = 30$, 33.5 and 52.5 MeV by (1977SL04) and analyzed with a PWIA: Fourier transforms of the wave functions were obtained. At $E({}^3\text{He}) = 35.9$ MeV, the spectra in reactions (b) and (c) are dominated by the nucleon-nucleon FSI: the results were fitted with a fully antisymmetrized PWBA and with DWBA (1974WA06, 1975WA31). At $E_{\vec{d}} = 15$ MeV (1976ME13) have measured the vector and the two tensor analyzing powers for reactions (b) and (c). (1976SC26, 1974SC04) have studied the excitation function for reaction (c) for $E_{\vec{d}} = 2.2$ to 6 MeV in a kinematically complete experiment. They have extracted the $p + t$ FSI going via ${}^4\text{He}^*(20.1) [J^\pi = 0^+]$ and suggest that the reaction

Table 5.5: Measurements of elastic scattering and polarization in ${}^4\text{He}(p, p){}^4\text{He}$ ^a

Elastic scattering angular distribution and cross sections		Polarization	
E_p (MeV)	Refs.	$E_{\bar{p}}$ (MeV)	Refs.
0.5 – 3.0	(1974KR07)	3.6, 6.35, 12.2	(1974LI1C, 1976LI1G)
11.16, 12.04, 13.60, 14.23	(1977DO01)	3.74 – 8.94	(1976BR17)
85.0	(1974VO05)	6.36	(1977ME06)
156	(1975CO1B)	11.93, 17.00	(1977HA06)
185 – 500	(1978MC06)	45.04, 52.34	(1977SA1E)
200, 350, 500	(1977ST30)	185 – 500	(1978MC06)
350, 650, 1050, 1150	(1977AS01)	560, 800, 1030,	(1977KL03)
438, 648, 1036	(1976BE45)	1270, 1730	
$[E_\alpha = 4.00, 5.08, 6.09 \text{ GeV}/c]$			
560, 800, 1030, 1270, 1730	(1977KL08)		
580, 720	(1975VE09)		
600	(1976FA09)		
788	(1978FO33)		
1000	(1977AL1E)		
1050	(1974BA14, 1978BE30)		
$E_\alpha = 7.0 \text{ GeV}/c$	(1977GE01)		
$[E_p = 1050]$			
2680	(1975IG1B, 1977IG1A, 1977IG1B, 1977IG1C)		
4.89 GeV	(1978NA1H)		
50 – 500 GeV/c	(1978SM1C)		

^a See also Tables 5.5 in (1966LA04) and 5.7 in (1974AJ01) for a listing of the earlier work.

goes primarily via a $J^\pi = \frac{3}{2}^-, T = \frac{1}{2}$ state of ${}^5\text{Li}$ located $0.8 \pm 0.2 \text{ MeV}$ above threshold [i.e., $E_x = 18.9 \pm 0.2 \text{ MeV}$]. This suggests that the attraction of a $p_{3/2}$ nucleon to ${}^4\text{He}^*(0^+)$ is stronger than is the attraction of such a nucleon to ${}^4\text{He}_{\text{g.s.}}$ (1976SC26). See also (1975RU04) for reaction (c). See (1974AJ01) for the earlier work.

See also (1974TA1A, 1975GL08, 1976GO1B), (1974LA1C, 1974PR1B, 1975HE1C, 1975HU1A; applied topics), (1975SE1D, 1975SE07, 1975SE1E, 1976SE03, 1976SE1B, 1976SE1C, 1977SE1C, 1977SE09, 1978FI1E, 1978SL1B; reviews) and (1974AL01, 1974CH02, 1974DO10, 1974KE1B, 1975DO1B, 1975NE11, 1975YA12, 1977CH1D, 1977CO1B, 1978KO13, 1978SE01; theor.).

4. ${}^3\text{He}(d, d){}^3\text{He}$

$$E_b = 16.39$$

In the range $E_d = 380$ to 570 keV, the scattering cross section is consistent with s-wave formation of the $J^\pi = \frac{3}{2}^+$ state at 16.66 MeV (1954BR05). The excitation curves for $E_d = 1.96$ to 10.99 MeV show a broad resonance ($\Gamma > 1$ MeV) corresponding to $E_x = 20.0 \pm 0.5$ MeV. From the behavior of the angular distributions an assignment of ${}^2D_{3/2}$ or $({}^2D, {}^4D)_{5/2}$ is favored, if only one state is involved (1967TO02). [There is some evidence that there is more than one D -wave state in this E_x region: see reaction 3]. See also (1976AL22: $E_d = 1.01$ to 2.95 MeV). In the range $E_d(\text{c.m.}) = 7$ to 18 MeV differential cross sections show only a monotonic variation with energy. There is no evidence for any other resonances from $E_x = 23$ to 34 MeV (1972BA30, 1972KI02). Angular distributions have been measured at a number of energies from $E({}^3\text{He}) = 18.7$ to 44.1 MeV (1972KI02) and at $E({}^3\text{He}) = 10.00$ MeV (1974JA15, $d\sigma/d\Omega$ to $\pm 1\%$) and $E_d = 39$ MeV (1975RU04).

Polarization measurements have been reported for $E_d = 4.0$ to 12 MeV [see (1974AJ01)] and polarization analyzing powers have been studied at $E_{\vec{d}} = 2.0$ to 11.5 MeV (1975JE1A; abstract), $E_{\vec{d}} = 3.73$ to 11.89 MeV (1974OH05; ${}^3\vec{\text{He}}$; also spin correlation parameters), $E_d = 5.35$ to 9.2 MeV (1974LO06; polarization transfer coefficients), $E_{\vec{d}} = 15$ to 40 MeV (1976RO1C, 1976RO1B; vector analyzing powers) and $E({}^3\vec{\text{He}}) = 33$ MeV (1977KA10).

See also (1975TO1A) and (1973HA1F, 1974CH02, 1974CH1E, 1974LY03, 1975DO1B, 1975HA1E, 1975KA1D, 1975NE11, 1975TA1A, 1976DO1B, 1976LY1A, 1977KA1H, 1977SA1C; theor.).

5. (a) ${}^3\text{He}(t, n){}^5\text{Li}$ $Q_m = 10.13$
 (b) ${}^3\text{He}(t, np){}^4\text{He}$ $Q_m = 12.0959$

Angular distributions have been measured at $E({}^3\text{He}) = 2.0$ to 5.5 MeV for n_0 ($\Gamma_{\text{g.s.}} = 2.03 \pm 0.15$ MeV) and at 3.5 to 5.5 MeV (partial distributions) for n_1 to an excited state at $E_x = 10.21 \pm 0.28$ MeV, $\Gamma = 1.49 \pm 0.61$ MeV (1975AB11). At $E({}^3\text{He}) = 14$ to 26 MeV, the spectra show the n_0 group and a broad resonance with $E_x = 20.5 \pm 0.8$ MeV (1974CH15). Reaction (b) has been studied at $E({}^3\text{He}) = 0.31$ to 2.80 to attempt to detect a possible resonant energy dependence of the spin-singlet ($T = 1$) n-p FSI: a marked effect is not present (1975SC28). See also (1977JO1A) and ${}^6\text{Li}$.

6. (a) ${}^3\text{He}({}^3\text{He}, p){}^5\text{Li}$ $Q_m = 10.89$
 (b) ${}^3\text{He}({}^3\text{He}, 2p){}^4\text{He}$ $Q_m = 12.8596$
 (c) ${}^3\text{He}({}^3\text{He}, 3p){}^3\text{H}$ $Q_m = -6.9544$

The spectrum of protons shows a pronounced peak at $E({}^3\text{He}) = 3$ to 18 MeV corresponding to ${}^5\text{Li}_{\text{g.s.}}$ superposed on a continuum: see (1974AJ01). At $E({}^3\text{He}) = 13.6$ MeV differential cross sections and polarizations have been measured for the p_0 group (1976IR02). A deuteron cluster transfer process may be involved (1976AS05). At $E({}^3\text{He}) = 43.7$ and 53.0 MeV, the spectra

show a prominent peak at the high energy end whose angular distributions exhibit a pronounced diffraction pattern (1967SL01, 1968MO10). Searches for three-proton enhancement (reaction (c)) have been unsuccessful: see (1974AJ01). See also (1976HE07; theor.).

7. ${}^4\text{He}(p, p){}^4\text{He}$

$$E_b = -1.97$$

Differential cross sections and polarizations have been measured at many energies: see Tables 5.5 in (1966LA04), 5.7 in (1974AJ01) and 5.5 here.

Phase shifts below $E_p = 18$ MeV have been determined by (1977DO01) based on all the available cross section and polarization measurements, using an R -matrix analysis program. The $P_{3/2}$ phase shift shows a pronounced resonance corresponding to ${}^5\text{Li}_{g.s.}$ while the $P_{1/2}$ shift changes slowly over a range of several MeV, suggesting that the first excited state is very broad and located 5 – 10 MeV above the ground state. (1977DO01) find that the reduced widths of the P -wave resonance states are nearly the same. See also (1974KR07: $E_x = 2.3 \pm 0.5$ MeV, $\Gamma = 9.0 \pm 1.5$ MeV for the $P_{1/2}$ state). The $D_{5/2}$, $D_{3/2}$, $F_{7/2}$ and $F_{5/2}$ phase shift become greater than 1° at $E_p \approx 11, 13, 14$ and 16 MeV, respectively (1977DO01). A contour plot of A_y is shown in (1973AR1N): the results are not significantly different from those of (1977DO01). The work of (1977DO01) relies heavily on the very accurate elastic scattering cross sections which they measured (better than 1%) and on the polarization measurements of (1977HA06) [absolute precision of ± 0.01]. It should be noted that $A_y = 0.984 \pm 0.005$ at $E_p = 11.93$ MeV, $\theta_{c.m.} = 128.28^\circ$ (1977HA06). For other phase-shift analyses, see (1976BR17) and (1974AJ01).

A resonance is observed at $E_p = 23$ MeV, corresponding to the known $\frac{3}{2}^+$ state at $E_x = 16.7$ MeV (1968DA04, 1968AL1B). An anomaly in the polarization is also observed at this energy (1966WE03, 1968DA04, 1972BA24). A further broad feature in the polarization excitation function ($\theta = 102^\circ$) is observed at $E_p = 30$ MeV ($E_x = 22$ MeV) (1972BA24). Cross sections for $23 < E_p < 45$ MeV show no further evidence of excited states (1969BU10). An extensive phase-shift analysis, using complex phases with $l \leq 4$ over the range $E_p = 20$ to 40 MeV has been made by (1972PL02), using mainly the polarization and differential cross-section data of (1972BA24). The $D_{3/2}$ level is fit with R -matrix formalism, including background interference, with the following parameters: $E_x = 16.68$ MeV, $\gamma_p^2 = 122$ keV, $\gamma_d^2(l = 0) = 1.58$ MeV (negative sign), $\gamma_d^2(l = 2) = 1.58$ MeV, $\theta_p^2 = 0.014$, $\theta_d^2 = 0.765$.

The $E_p = 30$ MeV structure is not reflected in anomalous behavior of any single phase shift. Strong absorption of even partial waves may indicate broad overlapping positive-parity levels, $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$, of $d + {}^3\text{He}$ character near $E_x = 22$ MeV, but there is no unambiguous identification of excited states other than the $\frac{3}{2}^+$ state in the p - α results (1972PL02). In an analysis of the data of (1971PL07) for $E_p = 25$ to 29 MeV, (1971RA27) report evidence for a $\frac{5}{2}^+$ level at $E_x = 20$ MeV [$\Gamma_{el}/\Gamma_{total} = 0.15$, $\Gamma_{tot} = 8$ MeV] and a $\frac{1}{2}^+$ level at $E_x = 18 \pm 1$ MeV.

A recent analysis by (1975PL01) of the phase shifts below 50 MeV suggests a small D -state admixture to the dominant S -state configuration of ${}^4\text{He}_{g.s.}$.

Total reaction cross sections have been measured at $E_p = 18$ to 48 MeV (1974SO06, 1976SO01) and at 0.87 and 2.1 GeV (1975JA1A). At an equivalent E_p of 1.05 GeV, the elastic distribution

is characterized by a shallow first minimum (1977GE01) and is said to be in excellent agreement with the predictions of multiple-diffraction theory (1977WA06). Alpha-proton bremsstrahlung is observed at $E_p = 7.0$ to 12.0 MeV, at $E_\alpha = 20$ MeV [see (1974AJ01)] and at $E_p = 22$, 40 and 45 MeV (1975AN18). Coherent single pion production has been studied at $E_p = 18.6$ GeV/c (1977BR1C). See also (1973BE1A, 1978GL1B), (1975IG1A, 1975RO1B, 1975TO1A, 1976GR1D, 1977BR1E, 1977RI1B, 1978BR1E, 1978IG1B), (1973GU1A, 1973HA1F, 1974BA34, 1974BA38, 1974CH02, 1974CH1E, 1974CI02, 1974DO10, 1974HA43, 1974JA1F, 1974LY02, 1974SA1G, 1974UL1A, 1974WA02, 1975AH07, 1975BA05, 1975BA1G, 1975BA76, 1975BL04, 1975CA05, 1975CO1A, 1975DU1A, 1975GR1C, 1975GU17, 1975KA1D, 1975NA07, 1975RU01, 1975RU07, 1975TA1A, 1975WA16, 1976AH1A, 1976AL1E, 1976AR12, 1976AU04, 1976CH1B, 1976DO1C, 1976DU06, 1976LE22, 1976LE1E, 1976NA04, 1976RU04, 1976SP1A, 1977AL06, 1977AR01, 1977CH1C, 1977DY01, 1977JA12, 1977KA1E, 1977KA1F, 1977KA1H, 1977LA1C, 1977LE1E, 1977PH01, 1977TH07, 1977TH09, 1977WA1C, 1977WA1E, 1977YO04, 1978CL1C, 1978HE2B, 1978LA02, 1978LE23, 1978MA37, 1978ME05, 1978YE1A, 1978YO1B; theor.).

8. (a) ${}^4\text{He}(p, d){}^3\text{He}$	$Q_m = -18.3532$	$E_b = -1.97$
(b) ${}^4\text{He}(p, pn){}^3\text{He}$	$Q_m = -20.5778$	
(c) ${}^4\text{He}(p, 2p){}^3\text{H}$	$Q_m = -19.8140$	
(d) ${}^4\text{He}(p, pd){}^2\text{H}$	$Q_m = -23.8467$	

Angular distributions of ${}^3\text{He}$ ions (reaction (a)) have been measured for $E_p = 27.9$ to 155.4 MeV [see (1974AJ01)] and at $E_p = 85$ MeV (1974VO05), 275 to 500 MeV (1978CA05: back angles), 770 MeV (1977BA19, 1976CO1C) and at $E_\alpha = 3.98$ GeV/c (1977BE34). The 85 MeV data are in agreement with DWBA at forward angles but the back angle behavior probably results from the need to include two-nucleon pickup in the analysis (1974VO05). The differential cross section is strongly backward peaked at $E_p = 275$ and 500 MeV and the differential cross section is similar in slope and magnitude to that for ${}^3\text{He}(p, p)$, suggesting that deuteron exchange may be the major component of the reaction mechanism (1978CA05). The excitation function shows no indication of resonance for $E_p = 38.5$ to 44.6 MeV (1969BU10). Polarization measurements have been reported for $E_p = 32$ to 63 MeV [see (1974AJ01)] and at 45.04 and 52.34 MeV (1977SA1E) and 350 , 400 and 500 MeV (1978CA05). At $E_\alpha = 6.85$ GeV/c, the integrated cross section for ${}^3\text{He}$ production (reactions (a) + (b)) is 24.1 ± 1.9 mb, suggesting an important component of ${}^3\text{He}$ in ${}^4\text{He}$ (1977BI05).

Reaction (b) and (c) have also been reported at $E_p = 46.8$ to 156 MeV [see (1974AJ01)]. The (p, 2p) reaction (reaction (c)) has also been studied at 350 and 500 MeV (1978EP1C, 1978KO1F) and 600 MeV (1969PE15), while both that reaction and reaction (d) have been studied at $E_p = 156$ MeV (1975FR14). See also (1973KR1A, 1978GL1B), (1977TE1A, 1978IG1B), (1976CO1B; astrophysics) and (1974BA34, 1974CH02, 1974CH29, 1974DO10, 1974HA36, 1974KA1D, 1974RO19, 1974YO01, 1975HA03, 1975KA1D, 1977CL1A, 1977KA1D, 1977KA1G, 1977RE1A, 1978BI1C, 1978RO06; theor.).

9. (a) ${}^4\text{He}(d, n){}^5\text{Li}$ $Q_m = -4.19$
 (b) ${}^4\text{He}(d, np){}^4\text{He}$ $Q_m = -2.2246$

Reaction (b) has been studied at $E_d = 14.2$ MeV and at $E_\alpha = 18.0$ to 42 MeV [see (1974AJ01)] and at $E_\alpha = 18$ MeV (1978SA07) and 21.9 and 23.7 MeV (1974RA10). The data show that at 42 MeV, direct breakup, with quasi-free α -p scattering taking place and the n acting as a spectator, is at least as important a mechanism as the FSI in the ${}^5\text{Li}_{g.s.}$ (1968WA01). See also (1974HE21, 1976KO21, 1976LI1E, 1977BR25; theor.) and ${}^6\text{Li}$.

10. ${}^4\text{He}(t, 2np){}^4\text{He}$ $Q_m = -8.3820$

See (1974LA1A, 1975AL1A; $E_\alpha = 70$ MeV; abstracts).

11. ${}^4\text{He}({}^3\text{He}, np){}^5\text{Li}$ $Q_m = -9.68$

See (1973HA50).

12. ${}^6\text{Li}(\gamma, n){}^5\text{Li}$ $Q_m = -5.66$

See ${}^6\text{Li}$.

13. (a) ${}^6\text{Li}(p, d){}^5\text{Li}$ $Q_m = -3.44$
 (b) ${}^6\text{Li}(p, pd){}^4\text{He}$ $Q_m = -1.4735$
 (c) ${}^6\text{Li}(p, pn){}^5\text{Li}$ $Q_m = -5.66$

Angular distributions have been measured at $E_p = 18.6$ to 156 MeV [see (1974AJ01)] and at $E_p = 185$ MeV (1976FA03, 1974KA28). In the latter experiment the spectra are characterized by a broad, asymmetric peak corresponding to ${}^5\text{Li}_{g.s.}$, a narrow peak [${}^5\text{Li}^*(16.7)$] and a broad peak at $E_x \approx 20$ MeV. DWBA analysis leads to $C^2S = 0.64$ and 0.57 for ${}^5\text{Li}^*(0, 16.7)$ (1976FA03). The first excited state of ${}^5\text{Li}$ is reported to be populated by (1969BA05; $E_p = 156$ MeV). See also (1978IG1A: 0.65 and 0.8 GeV).

Reaction (b) has been studied at $E_p = 9$ and 10 MeV, at 45 MeV [see (1974AJ01)] and 40, 45 and 50 MeV (1974BE46): the p- α FSI corresponding to ${}^5\text{Li}_{g.s.}$ is observed. See also (1975MI1A). For reaction (c) see (1977WA05). See also (1975ST1C, 1975VO04, 1978CH1G) and ${}^6\text{Li}$.

14. (a) ${}^6\text{Li}(\text{d}, \text{t}){}^5\text{Li}$ $Q_{\text{m}} = 0.59$
 (b) ${}^6\text{Li}(\text{d}, \text{pt}){}^4\text{He}$ $Q_{\text{m}} = 2.5592$

Angular distributions of the t_0 group have been measured at $E_{\text{d}} = 15$ and 20 MeV: see (1974AJ01). For reaction (b) see (1977RO18: $E_{\text{d}} = 0.47$ MeV), (1974MI10, 1977MI13; $E_{\text{d}} = 7.5$ to 10.5 MeV) and (1975KO1A; theor.).

15. (a) ${}^6\text{Li}({}^3\text{He}, \alpha){}^5\text{Li}$ $Q_{\text{m}} = 14.91$
 (b) ${}^6\text{Li}({}^3\text{He}, \text{p}\alpha){}^4\text{He}$ $Q_{\text{m}} = 16.880$

At $E({}^3\text{He}) = 25.5$ MeV, the spectra show ${}^5\text{Li}^*(0, 16.7)$ and two broad peaks at $E_{\text{x}} \approx 19.8$ and 22.7 MeV with $\Gamma_{\text{c.m.}} = 2$ and 1 MeV, respectively (1972BA30). In a kinematically complete experiment at $E({}^3\text{He}) = 36$ MeV the population of ${}^5\text{Li}^*(16.7, 20.2, 22.6)$ is reported (1976RA1B). The decay of the ground state has also been studied at $E({}^3\text{He}) = 1.25$ MeV (1974LI10) and at 1.47 to 1.75 MeV (1978GU15).

Cylindrical asymmetry observed in the breakup of ${}^5\text{Li}(0)$ is attributed to the short lifetime of the ${}^5\text{Li}$ intermediate state and to the memory retained by the proton of its localization at the time of formation of ${}^5\text{Li}$ (1967RE03, 1968RE10). The first excited state of ${}^5\text{Li}$ also appears to be involved (1972TH08, 1972TH1B): its parameters are given as $E_{\text{x}} = 3.2 \pm 0.2$ MeV, $\Gamma = 1.5 \pm 0.5$ MeV (1975GA14). See also (1975GL08, 1976ST1B, 1977AR09) and ${}^8\text{Be}$.

16. (a) ${}^7\text{Li}(\text{p}, \text{t}){}^5\text{Li}$ $Q_{\text{m}} = -4.43$
 (b) ${}^7\text{Li}(\text{p}, \text{p}\alpha){}^3\text{H}$ $Q_{\text{m}} = -2.467$

At $E_{\text{p}} = 43.7$ MeV, a triton group is observed to ${}^5\text{Li}(0)$ ($\Gamma = 1.55 \pm 0.15$ MeV): the angular distribution is consistent with a substantial mixing of $L = 0$ and 2 transfer. There is some evidence also for a very broad excited state between $E_{\text{x}} = 2$ and 5 MeV. ${}^5\text{Li}^*(16.7, 20.0)$ were not observed. The formation of ${}^5\text{Li}^*(16.7)$ (${}^4S_{3/2}$) would be S -forbidden: the absence of ${}^5\text{Li}^*(20.0)$ would indicate that this state(s) is also of quartet character [see reaction 20 in ${}^5\text{He}$] (1966CE05). Weak, broad states at $E_{\text{x}} = 22.0 \pm 0.5$ MeV and 25.0 ± 0.5 MeV and possibly 34 MeV are reported by (1968MC02) in a coincidence experiment in which three- and four-particle breakup was analyzed. The t_0 angular distribution has also been studied at $E_{\text{p}} = 16.6$ and 30.3 MeV: see (1974AJ01).

17. ${}^{10}\text{B}({}^3\text{He}, \text{p}\alpha){}^4\text{He}{}^4\text{He}$ $Q_{\text{m}} = 12.4192$

At $E({}^3\text{He}) = 2.45$ and 6.00 MeV the reaction proceeds in part via the first two states of ${}^5\text{Li}$ (1966WA16).

18. $^{12}\text{C}(\text{p}, 2\alpha)^5\text{Li}$

$$Q_m = -9.24$$

Not observed: see (1972MA62).

^5Be

(Fig. 3)

The absence of any group structure in the neutron spectrum in the reaction $^3\text{He}(^3\text{He}, \text{n})^5\text{Be}$ at $E(^3\text{He}) = 18.0$ to 26.0 MeV indicates that $^5\text{Be}(0)$ is at least 4.2 MeV unstable with respect to $^3\text{He} + 2\text{p}$ [$(M - A) > 33.7$ MeV]. With Coulomb corrections adjusted to match the 16.7 MeV states of ^5He - ^5Li , this observation places the first $T = \frac{3}{2}$ level in these nuclei above $E_x = 21.4$ MeV (1967AD05). See also (1975BE31; theor.).

References

(Closed 1978)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1952AR1A Argo, Taschek, Agnew, Hemmendinger and Leland, Phys. Rev. 87 (1952) 612
- 1952BO68 T.W. Bonner, J.P. Conner and A.B. Lillie, Phys. Rev. 88 (1952) 473
- 1952CO35 J.P. Conner, T.W. Bonner and J.R. Smith, Phys. Rev. 88 (1952) 468
- 1952DO30 D.C. Dodder and J.L. Gammel, Phys. Rev. 88 (1952) 520
- 1953JA1A Jarvis and Roaf, Proc. Roy. Soc. 218 (1953) 432
- 1954AR02 W.A. Arnold, J.A. Phillips, G.A. Sawyer, E.J. Stovall, Jr. and J.L. Tuck, Phys. Rev. 93 (1954) 483
- 1954BR05 R.J.S. Brown, K.F. Famularo, H.D. Holmgren, D. Rankin and T.F. Stratton, Phys. Rev. 96 (1954) 80
- 1954BU06 W.H. Burke, J.R. Risser and G.C. Phillips, Phys. Rev. 93 (1954) 188
- 1955KU03 W.E. Kunz, Phys. Rev. 97 (1955) 456
- 1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
- 1959BO54 T.W. Bonner, F.W. Prosser, Jr. and J. Slattery, Phys. Rev. 115 (1959) 398
- 1960HU1A Hughes, Magurno and Brussel, BNL 325 (2nd Edition) Suppl. 1 (1960)
- 1963BU07 W. Buss, H. Waffler and B. Ziegler, Phys. Lett. 4 (1963) 198
- 1963PO02 C.H. Poppe, C.H. Holbrow and R.R. Borchers, Phys. Rev. 129 (1963) 733
- 1964PA1A Parker, Bahcall and Fowler, Astrophys. J. 139 (1964) 602
- 1964SH1A Shamu and Jenkin, Phys. Rev. 135 (1964) B99
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL 325, 2nd Ed., Suppt. 2, Vol. 1 (1964)
- 1965CH12 G. Charpak, G. Gregoire, L. Massonnet, J. Saudinos, J. Favier, M. Gusakov and M. Jean, Phys. Lett. 16 (1965) 54
- 1965TR01 W. Trachslin, H. Burgisser, P. Huber, G. Michel and H.R. Striebel, Helv. Phys. Acta 38 (1965) 523
- 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331

1965WO03 C. Wong, J.D. Anderson and J.W. McClure, Nucl. Phys. 71 (1965) 106
 1966BR02 L. Brown, H.A. Christ and H. Rudin, Nucl. Phys. 79 (1966) 459
 1966CE05 J. Cerny, C. Detraz and R.H. Pehl, Phys. Rev. 152 (1966) 950
 1966HO07 B. Hoop, Jr. and H.H. Barschall, Nucl. Phys. 83 (1966) 65
 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
 1966WA16 M.A. Waggoner, J.E. Etter, H.D. Holmgren and C. Moazed, Nucl. Phys. 88 (1966) 81
 1966WE03 W.G. Weitkamp and W. Haeberli, Nucl. Phys. 83 (1966) 46
 1967AD05 E.G. Adelberger, A.B. McDonald, T.A. Tombrello, F.S. Dietrich and A.V. Nero, Phys. Lett. B25 (1967) 595
 1967MC01 L.C. McIntyre and W. Haeberli, Nucl. Phys. A91 (1967) 369
 1967RE03 M.A. Reimann, P.W. Martin and E.W. Vogt, Phys. Rev. Lett. 18 (1967) 246
 1967SL01 R.J. Slobodrian, J.S.C. McKee, W.F. Tivol, D.J. Clark and T.A. Tombrello, Phys. Lett. B25 (1967) 19
 1967TO02 T.A. Tombrello, R.J. Spiger and A.D. Bacher, Phys. Rev. 154 (1967) 935
 1967TR05 A. Trier and W. Haeberli, Phys. Rev. Lett. 18 (1967) 915
 1967VA11 V. Valkovic, W.R. Jackson, Y.S. Chen, S.T. Emerson and G.C. Phillips, Nucl. Phys. A96 (1967) 241
 1967VA12 V. Valkovic, I. Slaus, P. Tomas and M. Cerineo, Nucl. Phys. A98 (1967) 305
 1968AL1B Allison and Smythe, Nucl. Phys. A121 (1968) 97
 1968BU09 W. Buss, W. Del Bianco, H. Waffler and B. Ziegler, Nucl. Phys. A112 (1968) 47
 1968DA04 P. Darriulat, D. Garreta, A. Tarrats and J. Testoni, Nucl. Phys. A108 (1968) 316
 1968IV01 M. Ivanovich, P.G. Young and G.G. Ohlsen, Nucl. Phys. A110 (1968) 441
 1968KR03 L. Kraus, M. Suffert and D. Magnac-Valette, Nucl. Phys. A109 (1968) 593
 1968MC02 R.L. McGrath, J. Cerny and S.W. Cosper, Phys. Rev. 165 (1968) 1126
 1968MO10 B.J. Morton, E.E. Gross, E.V. Hungerford, J.J. Malanify and A. Zucker, Phys. Rev. 169 (1968) 825
 1968RE10 M.A. Reimann, P.W. Martin and E.W. Vogt, Can. J. Phys. 46 (1968) 2241
 1968VA12 V. Valkovic, C. Joseph, A. Niiler and G.C. Phillips, Nucl. Phys. A116 (1968) 497
 1968WA01 R.E. Warner and R.W. Bercaw, Nucl. Phys. A109 (1968) 205
 1968YO06 P.G. Young, R.H. Stokes and G.G. Ohlsen, Phys. Rev. 173 (1968) 949
 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60

- 1969BU10 S.N. Bunker, J.M. Cameron, M.P. Epstein, G. PailImage, J.R. Richardson, J.G. Rogers, P. TomaImage and J.W. Verba, Nucl. Phys. A133 (1969) 537
- 1969PE15 C.F. Perdrisat, L.W. Swenson, P.C. Gugelot, E.T. Boschitz, W.K. Roberts, J.S. Vincent and J.R. Priest, Phys. Rev. 187 (1969) 1201
- 1969RO16 D.C. Rorer, B.M. Ecker and R.O. Akyuz, Nucl. Phys. A133 (1969) 410
- 1970AR1B R.A. Arndt and L.D Roper, Phys. Rev. C1 (1970) 903
- 1970BE1A Bezotosnyi, Zhmailo, Surov and Shvetsov, Sov. J. Nucl. Phys. 10 (1970) 127
- 1970GA1A Garber et al, BNL 400, 3rd Edition, Vol. 1 (1970)
- 1970KO09 A. Kosiara and H.B. Willard, Phys. Lett. B32 (1970) 99
- 1970MI05 D. Miljanic, M. Furic and V. Valkovic, Nucl. Phys. A148 (1970) 312
- 1970SC18 H. Schroder and W. Mausberg, Z. Phys. 235 (1970) 234
- 1971GR12 E.E. Gross, E.V. Hungerford and J.J. Malanify, Nucl. Phys. A164 (1971) 376
- 1971GR32 H. Grunder, R. Gleyvod, G. Lietz, G. Morgan, H. Rudin, F. Seiler and A. Stricker, Helv. Phys. Acta 44 (1971) 662
- 1971GR45 R. Grotzschel, B. Kuhn, K. Moller, J. Mosner and G. Schmidt, Nucl. Phys. A176 (1971) 261
- 1971GR47 W. Gruebler, V. Konig, A. Ruh, P.A. Schmelzbach, R.E. White and P. Marmier, Nucl. Phys. A176 (1971) 631
- 1971HI07 D. Hilscher, P.A. Quin and J.C. Davis, Nucl. Phys. A173 (1971) 216
- 1971KE16 L.G. Keller and W. Haeberli, Nucl. Phys. A172 (1971) 625
- 1971KI02 K. Kilian, H. Treiber, R. Strausz and D. Fick, Phys. Lett. B34 (1971) 283
- 1971KL02 W. Klinger, F. Dusch and R. Fleischmann, Nucl. Phys. A166 (1971) 253
- 1971LE27 C. Leemann, H. Burgisser, P. Huber, U. Rohrer, H. Paetz gen.Schieck and F. Seiler, Ann. Phys. (N.Y.) 66 (1971) 810
- 1971MU04 G.S. Mutchler, W.B. Broste and J.E. Simmons, Phys. Rev. C3 (1971) 1031
- 1971PL07 D.J. Plummer, K. Ramavataram, T.A. Hodges, D.G. Montague, A. Zucker and N.K. Ganguly, Nucl. Phys. A174 (1971) 193
- 1971RA27 K. Ramavataram, D.J. Plummer, T.A. Hodges and D.G. Montague, Nucl. Phys. A174 (1971) 204
- 1971RE19 D. Rendic, N.D. Gabitzsch, V. Valkovic, W. von Witsch and G.C. Phillips, Nucl. Phys. A178 (1971) 49
- 1972AG01 N.M. Agababyan, Y.A. Batusov, S.A. Bunyatov, G.P. Gulkanyan, V.M. Sidorov and V.A. Yarba, Yad. Fiz. 15 (1972) 18; Sov. J. Nucl. Phys. 15 (1972) 10

- 1972AN24 Y.N. Antufev, V.L. Agranovich, V.S. Kuzmenko, I.I. Miroschnichenko and P.V. Sorokin, *Pisma Zh. Eksp. Teor. Fiz.* 16 (1972) 77; *JETP Lett. (USSR)* 16 (1972) 52
- 1972AN27 Y.P. Antoufi ev, V.L. Agranovich, V.S. Kuzmenko and P.V. Sorokin, *Phys. Lett.* B42 (1972) 347
- 1972AN29 Y.P. Antufev, V.L. Agranovich, V.S. Kuzmenko and P.V. Sorokin, *Pisma Zh. Eksp. Teor. Fiz.* 16 (1972) 339; *JETP Lett. (USSR)* 16 (1972) 240
- 1972BA24 A.D. Bacher, G.R. Plattner, H.E. Conzett, D.J. Clark, H. Grunder and W.F. Tivol, *Phys. Rev. C*5 (1972) 1147
- 1972BA30 M.P. Baker, J.M. Cameron, N.S. Chant and N.F. Mangelson, *Nucl. Phys.* A184 (1972) 97
- 1972BR10 W.B. Broste, G.S. Mutchler, J.E. Simmons, R.A. Arndt and L.D. Roper, *Phys. Rev. C*5 (1972) 761
- 1972KI01 H.T. King, W.E. Meyerhof and R.G. Hirko, *Nucl. Phys.* A178 (1972) 337
- 1972KI02 T.R. King and R. Smythe, *Nucl. Phys.* A183 (1972) 657
- 1972MA62 A.M. MacLeod and G.R. Milne, *J. Phys. (London)* A5 (1972) 1252
- 1972PL02 G.R. Plattner, A.D. Bacher and H.E. Conzett, *Phys. Rev. C*5 (1972) 1158
- 1972ST01 T. Stammbach and R.L. Walter, *Nucl. Phys.* A180 (1972) 225
- 1972TH08 D.T. Thompson and G.E. Tripard, *Phys. Rev. C*6 (1972) 452
- 1972TH1B Thompson, Thesis, Washington State Univ. (1972); *Phys. Abs.* 984 (1973)
- 1973AR1B Arthur, Thesis LA 5230 T (1973)
- 1973AR1N Arndt, Long and Roper, *Nucl. Phys.* A209 (1973) 429
- 1973BA30 Y.D. Bayukov, L.S. Vorobev, V.M. Kolybasov, G.A. Leksin, V.L. Stolin, V.B. Fedorov and V.D. Khovanskii, *Yad. Fiz.* 17 (1973) 916; *Sov. J. Nucl. Phys.* 17 (1974) 479
- 1973BE1A Berthot et al, U. Clajde Bernard de Lyon, Rpt. No. Lycen 7338 (1973)
- 1973DE17 V.P. Denisov, L.A. Kulchitskii and I.Y. Chubukov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 107; *Bull. Acad. Sci. USSR Phys. Ser.* 37:1 (1973) 94
- 1973DE51 A.A. Debenham, V. Konig, W. Gruebler, P.A. Schmelzbach, R. Risler and D.O. Boerma, *Nucl. Phys.* A216 (1973) 42
- 1973FE1A N. Feather and D.G. Vass, *Proc. Roy. Soc. Edinburgh* A71 (1973) 233
- 1973GA16 S.N. Gardiner, J.L. Matthews and R.O. Owens, *Phys. Lett.* B46 (1973) 186
- 1973GA28 H.P. Garnir, G. Robaye and J.M. Delbrouck-Habaru, *Bull. Soc. Roy. Sci. Liege (Belgium)* 42 (1973) 195
- 1973GO38 C.A. Goulding, P. Stoler and J.D. Seagrave, *Nucl. Phys.* A215 (1973) 253

- 1973GU1A Guardiola, in Nucl. Many-Body Problem, Vol. 2 (1972) (Ed. Compositori, Bologna, 1973) p. 433
- 1973HA1F Hackenbroich, in Nucl. Many-Body Problem, Vol. 1 (1972) (Ed. Compositori, Bologna, 1973) p. 706
- 1973HA50 M.L. Halbert, A. van der Woude and N.M. O'Fallon, Phys. Rev. C8 (1973) 1621
- 1973HU12 J. Hudomalj and P. Tomas, Fizika (Yugoslavia) 5 (1973) 203
- 1973KR1A Krebs et al, Phys. in Canada 29:4 (1973) 22; Phys. Abs. 40019 (1974)
- 1973LI1B Liskien and Paulsen, Nucl. Data Tables 11 (1973) 569
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973SL03 I. Slaus, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, J.M. Lambert and D.L. Shannon, Phys. Rev. C8 (1973) 444
- 1973TR04 P.A. Treado, J.M. Lambert, R.J. Kane, L.A. Beach, E.L. Petersen and R.B. Theus, Phys. Rev. C7 (1973) 1742
- 1974AD11 A. Adam and J. Cabe, Nucl. Instrum. Meth. 121 (1974) 339
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974AL01 R.G. Allas, L.A. Beach, R.O. Bondelid, E.M. Diener, E.L. Petersen, J.M. Lambert, P.A. Treado and I. Slaus, Phys. Rev. C9 (1974) 787
- 1974AM01 J. Amato, R.L. Burman, R. Macek, J. Oostens, W. Shlaer, E. Arthur, S. Sobottka and W.C. Lam, Phys. Rev. C9 (1974) 501
- 1974AN02 B. Antolkovic, Nucl. Phys. A219 (1974) 332
- 1974BA14 S.D. Baker, R. Beurtey, G. Bruge, A. Chaumeaux, J.-M. Durand, J.C. Faivre, J.M. Fontaine, D. Garreta, D. Legrand, J. Saudinos et al, Phys. Rev. Lett. 32 (1974) 839
- 1974BA34 A.G. Baryshnickov, L.D. Blokhintsev, A.N. Safronov and V.V. Turovtsev, Nucl. Phys. A224 (1974) 61
- 1974BA38 R.H. Bassel and M. Rosen, Phys. Rev. C10 (1974) 928
- 1974BE46 J.L. Beveridge, R. Corfu, J.P. Egger, F. Foroughi, J. Piffaretti, J. Rossel, J. Arvieux, J.L. Durand and C. Perrin, Helv. Phys. Acta 47 (1974) 211
- 1974BE67 R. Beckmann, U. Holm and K. Lorenzen, Z. Phys. 271 (1974) 89
- 1974BH03 R.K. Bhowmik, C.C. Chang, P.G. Roos and H.D. Holmgren, Nucl. Phys. A226 (1974) 365
- 1974CH02 F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Rev. C9 (1974) 56
- 1974CH15 C.C. Chang and E. Ventura, Phys. Rev. C9 (1974) 1671
- 1974CH1E F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Lett. B46 (1974) 301
- 1974CH29 F.S. Chwieroth, Y.C. Tang and D.R. Thompson, Phys. Rev. C10 (1974) 406

- 1974CI02 C. Ciofi degli Atti and R. Guardiola, Phys. Lett. B51 (1974) 26
- 1974DA28 V. D'Amico, G. Fazio, S. Jannelli, F. Mezzanares and R. Potenza, Nuovo Cim. A22 (1974) 313
- 1974DO05 T.R. Donoghue, G.G. Ohlsen, N. Jarmie and P.W. Keaton, Jr., Phys. Rev. C10 (1974) 571
- 1974DO10 E.I. Dolinsky and A.M. Mukhamedzhanov, Phys. Lett. B52 (1974) 17
- 1974FE1A Feather and Vass, Proc. Royal Soc. (Edinburgh) A71 (1974) 293
- 1974GA21 R. Garrett and W.W. Lindstrom, Nucl. Phys. A224 (1974) 186
- 1974GE1B Geisler et al, Science 186 (1974) 361
- 1974GH01 A. Ghovanlou and D.R. Lehman, Phys. Rev. C9 (1974) 1730
- 1974GH03 A. Ghovanlou and F. Prats, Phys. Rev. C10 (1974) 1309
- 1974GO13 G.S. Gogsdaze and T.I. Kopaleishvili, Yad. Fiz. 19 (1974) 539; Sov. J. Nucl. Phys. 19 (1974) 272
- 1974GR40 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2148; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 115
- 1974GR42 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2543; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 69
- 1974GR44 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, G.B. Lyubanskii, O.F. Nemets, R.G. Ofengenden, V.M. Pugach, B.A. Rudenko, M.V. Sokolov, B.G. Struzhko et al, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2547; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 72
- 1974HA36 R.D. Haracz and T.K. Lim, Phys.Rev. C10 (1974) 431
- 1974HA43 H.H. Hackenbroich, P. Heiss and Le-Chi-Niem, Phys. Lett. B51 (1974) 334
- 1974HE21 P. Heiss, H.H. Hackenbroich and K. Prescher, Phys. Lett. B52 (1974) 411
- 1974HO11 E. Hourany, H. Nakamura, F. Takeutchi and T. Yuasa, Nucl. Phys. A222 (1974) 537
- 1974IR04 J.M. Irvine, G.S. Mani and M. Vallieres, Czech. J. Phys. B24 (1974) 1269
- 1974JA15 N. Jarmie and J.H. Jett, Phys. Rev. C10 (1974) 54
- 1974JA1F Jackson, Prog. in Phys. 37 (1974) 55
- 1974JA30 V.K. Jain and B.B. Srivastava, Indian J. Pure Appl. Phys. 12 (1974) 117; Phys. Abs. 66134 (1974)
- 1974KA1D Kanada, Kaneko and Nomoto, Prog. Theor. Phys. 52 (1974) 725
- 1974KA28 J. Kallne, B. Fagerstrom, O. Sundberg and G. Tibell, Phys. Lett. B52 (1974) 313
- 1974KE1B Keaton, Gammel and Ohlsen, Ann. Phys. 85 (1974) 152

- 1974KR07 L. Kraus and I. Linck, Nucl. Phys. A224 (1974) 45
- 1974LA02 R. Larose-Poutissou and H. Jeremie, Nucl. Phys. A218 (1974) 559
- 1974LA1A Lambert et al, Bull. Amer. Phys. Soc. 19 (1974) 506
- 1974LA1C Langley, Picrajx and Vook, J. Nucl. Materials 53 (1974) 257
- 1974LE22 C. Leclercq-Willain and M. Libert-Heinemann, Nucl. Phys. A229 (1974) 15
- 1974LI10 D.L. Livesey and C.J. Piluso, Can. J. Phys. 52 (1974) 1167
- 1974LI1C Lisowski, Walter, Jacobs and Clegg, Bull. Amer. Phys. Soc. 19 (1974) 1075
- 1974LO06 P.A. Lovoi, G.C. Salzman, G.G. Ohlsen, C.K. Mitchell and W. Gruebler, Phys. Rev. C9 (1974) 1336
- 1974LY02 G.I. Lykasov and A.V. Tarasov, Yad. Fiz. 20 (1974) 489; Sov. J. Nucl. Phys. 20 (1975) 263
- 1974LY03 E.B. Lyovshin, O.F. Nemets and A.M. Yasnogorodsky, Phys. Lett. B52 (1974) 392
- 1974MA1C Maayouf and Galloway, Nucl. Instrum. Meth. 118 (1974) 343
- 1974MI10 D. Miljanic, J. Hudomalj, G.S. Mutchler, E. Andrade and G.C. Phillips, Phys. Lett. B50 (1974) 330
- 1974NA10 H. Nakamura, Nucl. Phys. A223 (1974) 599
- 1974OH05 G.G. Ohlsen, R.A. Hardekopf, D.P. May, S.D. Baker and W.T. Armstrong, Nucl. Phys. A233 (1974) 1
- 1974POZN C.H. Poppe, J.C. Davis, J.D. Anderson, S.M. Grimes and C. Wong, Bull. Amer. Phys. Soc. 19 (1974) 1035
- 1974PR10 F. Prats, Nucl. Phys. A227 (1974) 469
- 1974PR1B Pronko, J. Nucl. Materials 53 (1974) 252
- 1974RA10 T. Rausch, H. Zell, D. Wallenwein and W. Von Witsch, Nucl. Phys. A222 (1974) 429
- 1974RA1C Ramaty and Kozlovsky, Astrophys. J. 193 (1974) 729
- 1974RO19 P.G. Roos, Phys. Rev. C9 (1974) 2437
- 1974SA1G Saudinos and Wilkin, Ann. Rev. Nucl. Science 24 (1974) 341
- 1974SC04 H. Schroder, K.K. Kern and D. Fick, Phys. Lett. B48 (1974) 206
- 1974SO06 A.M. Sourkes, N.E. Davison, S.A. Elbaker, J.L. Horton, A. Houdayer, W.T.H. van Oers and R.F. Carlson, Phys. Lett. B51 (1974) 232
- 1974ST14 M. Stingl and A.S. Rinat, Phys. Rev. C10 (1974) 1253
- 1974TA1A Tanabe et al, Bull. Inst. Chem. Res. Kyoto U. 52 (1974) 284
- 1974TO03 W. Tornow, Z. Phys. 266 (1974) 357
- 1974TR02 T.A. Trainor, T.B. Clegg and P.W. Lisowski, Nucl. Phys. A220 (1974) 533

- 1974TU1A Turk, Antolkovic, Dolenc and Winterhalter, in Proc. of 1973 Gaussig Conf., ZFK-271 (1974) 115
- 1974UL1A Ullo and Feshbach, Ann. Phys. 82 (1974) 156
- 1974VO05 L.G. Votta, P.G. Roos, N.S. Chant and R. Woody, III, Phys. Rev. C10 (1974) 520
- 1974WA02 W.L. Wang and R.G. Lipes, Phys. Rev. C9 (1974) 814
- 1974WA06 R.E. Warner, S.A. Gottlieb, G.C. Ball, W.G. Davies, A.J. Ferguson and J.S. Forster, Nucl. Phys. A221 (1974) 593
- 1974WI1D Wiemer, Atomkernenergie 23 (1974) 261
- 1974YO01 S.K. Young and E.F. Redish, Phys. Rev. C10 (1974) 498
- 1975AB11 U. Abbondanno, F. Demanins, C. Tuniz and G. Nardelli, Nuovo Cim. A29 (1975) 187
- 1975AB1C Abulaffio and Peres, Nucl. Cross Sect. and Tech., Pt. 3, Washington, D.C., 1975 (Washington, DC Nat. Bur. Standards 1975) 701; Phys. Abs. 22996 (1977)
- 1975AH07 A.A.Z. Ahmad, S. Ali, N. Ferdous and M. Ahmed, Nuovo Cim. A30 (1975) 385
- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
- 1975AL1A Allas et al, in Few Body Problems, Quebec, 1974 (U. Laval, 1975) p. 422
- 1975AN18 G.A. Anzelon, I. Slaus, S.Y. Tin, W.T.H. Van Oers, R.M. Eisberg, M. Makino, C.N. Waddell and M.B. Epstein, Nucl. Phys. A255 (1975) 250
- 1975BA05 S. Barshay, C.B. Dover and J.P. Vary, Phys. Rev. C11 (1975) 360
- 1975BA1E Batay-Csorba and Barnes, Bull. Amer. Phys. Soc. 20 (1975) 829
- 1975BA1G Baldin, in LASL, AIP Conf. Proc. 26 (1975) 621
- 1975BA76 A.I. Baz, V.F. Demin and M.V. Zhukov, Fiz. Elem. Chastits At. Yadra 6 (1975) 515; Sov. J. Part. Nucl. 6 (1975) 207
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BL04 M. Bleszynski and T. Jaroszewicz, Phys. Lett. B56 (1975) 427
- 1975BR1A Brown et al, Bull. Amer. Phys. Soc. 20 (1975) 1157
- 1975CA05 G. Cattapan, G. Pisent and V. Vanzani, Nucl. Phys. A241 (1975) 204
- 1975CH1B Chang and Bhowmik, in Clustering Phenomena in Nuclei, II, ORO-4856-26 (1975) p. 317
- 1975CH1C Chang, Bhowmik, Chant and Roos, ORO-4856-32 (1975)
- 1975CO1A Coker and Hoffmann, Bull. Amer. Phys. Soc. 20 (1975) 1166
- 1975CO1B V. Comparat, R. Frascaria, N. Fujiwara, N. Marty, M. Morlet, P.G. Roos and A. Willis, Phys. Rev. C12 (1975) 251
- 1975CR1A Cranberg, Bull. Amer. Phys. Soc. 20 (1975) 1182

- 1975DE37 V.P. Denisov and I.Y. Chubukov, *Yad. Fiz.* 22 (1975) 897; *Sov. J. Nucl. Phys.* 22 (1976) 466
- 1975DI04 H. Dirim, J.P. Elliott and J.A. Evans, *Nucl. Phys.* A244 (1975) 301
- 1975DI1A Didelez et al, *Bull. Amer. Phys. Soc.* 20 (1975) 597
- 1975DI1B Didelez et al, in *JUL-Conf-16* (1975) 90
- 1975DO1B Dodder, *Bull. Amer. Phys. Soc.* 20 (1975) 821
- 1975DU1A Dubnicka and Dumbrajs, *Phys. Rept.* 19 (1975) 141
- 1975FE1A Feather, *Commun. R. Soc. Edinb. (Phys. Sci.)* 2 (1975) 19
- 1975FI1B Firk et al, *Nucl. Cross Sections and Tech., Pt. II, Wasington, D.C. 1975* (Wash., D.C., Nat. Bur. Standards 1975) p. 875; *Phys. Abs.* 23088 (1977)
- 1975FO19 W.A. Fowler, G.R. Caughlan and B.A. Zimmerman, *Ann. Rev. Astron. Astrophys.* 13 (1975) 69
- 1975FR14 R. Frascaria, P.G. Roos, M. Morlet, N. Marty, A. Willis, V. Comparat and N. Fujiwara, *Phys. Rev.* C12 (1975) 243
- 1975GA14 R.M. Gagne, C.M. Fou and S. Ward, *J. Phys. (Paris)* 36 (1975) 759
- 1975GL08 Y.A. Glukhov, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and V.I. Chuev, *Yad. Fiz.* 22 (1975) 679; *Sov. J. Nucl. Phys.* 22 (1976) 351
- 1975GR1C Green, *Few Body Problems, Quebec, 1974* (Univ. Laval, 1975) p. 352
- 1975GR42 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 39 (1975) 2117; *Bull. Acad. Sci. USSR Phys. Ser.* 39 (1975) 95
- 1975GU17 S.A. Gurvits, Y. Alexander and A.S. Rinat, *Phys. Lett.* B59 (1975) 22
- 1975HA03 R.D. Haracz and T.K. Lim, *Phys. Rev.* C11 (1975) 634
- 1975HA1E Hackenbroich, in *Clustering Phenom. in Nucl., II, ORO-4856-26* (1975) 107
- 1975HE09 P. Heiss, *Z. Phys.* A272 (1975) 267
- 1975HE1C J. Henley and H.W. Meldner, *Phys. Rev.* C12 (1975) 407
- 1975HU1A Hufschmidt, Heintze, Moller and Kamke, *Nucl. Instrum. Meth.* 124 (1975) 573
- 1975IG1A Igo, in *Lasl, AIP Conf. Proc.* 26 (1975) 63
- 1975IG1B Igo et al, *Bull. Amer. Phys. Soc.* 20 (1975) 1166
- 1975JA1A Jaros et al, *Bull. Amer. Phys. Soc.* 20 (1975) 666
- 1975JE1A Jenny et al, *Helv. Phys. Acta* 48 (1975) 558
- 1975KA1D Kanada, Kaneko and Nomoto, *Prog. Theor. Phys.* 54 (1975) 1707
- 1975KO1A S. Kohmoto and Y. Sakamoto, *Phys. Rev. Lett.* 34 (1975) 550
- 1975KR1A Kramer, in *Clustering Phenom. in Nucl., II, ORO-4856-26* (1975) 56

1975KU1C Kuhn, Few Body Prob., Quebec, 1974 (Univ. Laval, 1975) 122

1975LO1A M.P. Locher, Nucl. Phys. A251 (1975) 493

1975MA1E Matthews, in LASL, AIP Conf. Proc. 26 (1975) 437

1975MA28 E. Magiera, M. Bormann, W. Scobel and P. Heiss, Nucl. Phys. A246 (1975) 413

1975ME27 Y.P. Melnik and E.V. Inopin, Ukr. Fiz. Zh. 20 (1975) 2032; Phys. Abs. 20829 (1976)

1975MI1A Miljanic, Zabel and Phillips, Few Body Prob., Quebec, 1974 (Univ. Laval, 1975) 669

1975NA07 M.A. Nagarajan, W.L. Wang, D.J. Ernst and R.M. Thaler, Phys. Rev. C11 (1975) 1167

1975NE11 O.F. Nemets, Y.S. Stryuk and A.M. Yasnogorodskii, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2066; Bull. Acad. Sci. USSR Phys. Ser. 39:10 (1975) 49

1975PL01 G.R. Plattner, R.D. Viollier and K. Alder, Phys. Rev. Lett. 34 (1975) 830

1975RO1B Roos and Chant, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 242, 265

1975RO1C Roy et al, Bull. Amer. Phys. Soc. 20 (1975) 693

1975RU01 D.W. Rule and Y. Hahn, Phys. Rev. Lett. 34 (1975) 332

1975RU04 R.W. Rutkowski and E.E. Gross, Phys. Rev. C12 (1975) 362

1975RU07 D.W. Rule and Y. Hahn, Phys. Rev. C12 (1975) 1616

1975SA1A Sagle et al, Few Body Problems, Quebec, 1974 (Univ. Laval, 1975) p. 621

1975SC28 G. Schrieder, H. Genz, A. Richter and W. Von Witsch, Nucl. Phys. A247 (1975) 203

1975SE07 F. Seiler, Nucl. Phys. A244 (1975) 236

1975SE1D Seiler, in Clustering Phenomena in Nuclei, II, ORO-4856-26 (1975) p. 191

1975SE1E Seiler, Few Body Problems, Quebec, 1974 (Univ. Laval, 1975) p. 595

1975SO1C Soleilhac, CEA-N-1812 (1975)

1975ST1A Stengl, Vonach and Fabian, Nucl. Instrum. Meth. 126 (1975) 235

1975ST1B Stewart and Hale, USNDC-CTR-2, LA-5828-MS (1975)

1975ST1C Strobel, JUL-CONF-16 (1975) p. 89

1975TA1A Tang and Thompson, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 119

1975TH12 D.R. Thompson and Y.C. Tang, Phys. Rev. C12 (1975) 1432; Erratum Phys. Rev. C13 (1976) 2597

1975TO1A Tombrello, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 167

1975VE09 S.L. Verbeck, J.C. Fong, G. Igo, C.A. Whitten, Jr., D.L. Hendrie, Y. Terrien, V. Perez-Mendez and G.W. Hoffmann, Phys. Lett. B59 (1975) 339

1975VO04 W. Von Witsch, G.S. Mutchler and D. Miljanic, Nucl. Phys. A248 (1975) 485

1975WA16 S.J. Wallace, Phys. Rev. C12 (1975) 179

- 1975WA31 R.E. Warner, R.L. Ruyle, W.G. Davies, G.C. Ball, A.J. Ferguson and J.S. Forster, Nucl. Phys. A255 (1975) 95
- 1975YA12 A.M. Yasnogorodsky, Phys. Lett. B58 (1975) 139
- 1976AH1A M.U. Ahmed and P.E. Shanley, Phys. Rev. Lett. 36 (1976) 25
- 1976AL1E Ali, Proc. 7th Int. Conf. on Few Body Problems in Nucl. and Part. Phys., Delhi 1975 (Amsterdam, North-Holland 1976) p. 438; Phys. Abs. 18501 (1977)
- 1976AL22 M. Allab and H. Beaumevieille, Nuovo Cim. A36 (1976) 95
- 1976AR12 L.G. Arnold, B.C. Clark, R.L. Mercer, D.G. Ravenhall and A.M. Saperstein, Phys. Rev. C14 (1976) 1878
- 1976AS05 J. Asai and R.J. Slobodrian, Phys. Lett. B64 (1976) 257
- 1976AU04 J.P. Auger, J. Gillespie and R.J. Lombard, Nucl. Phys. A262 (1976) 372
- 1976BA1E Baz and Filippov, in Proc. Dubna Conf. JINR-9920, Vol. II (1976) 268
- 1976BE45 J. Berger, J. Duflo, L. Goldzahl, F. Plouin, J. Oostens, M. Van Den Bossche, L. Vu Hai, G. Bizard, C. Le Brun, F.L. Fabbri et al, Phys. Rev. Lett. 37 (1976) 1195
- 1976BI1A A. Billoire and M. Chemtob, Phys. Lett. B64 (1976) 389
- 1976BL1A Block, Bull. Amer. Phys. Soc. 21 (1976) 1342
- 1976BO05 J.E. Bond and F.W.K. Firk, Nucl. Phys. A258 (1976) 189
- 1976BR17 M.E. Brandan, G.R. Plattner and W. Haeberli, Nucl. Phys. A263 (1976) 189
- 1976BR1B Brady et al, in Lowell Conf., CONF-760715-P2 (1976) p. 1273
- 1976BR1C Brook et al, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 546
- 1976CH1B Chuu, J. Natl. Chiao Tung Univ. (Taiwan) 1 (1976) 295; Phys. Abs. 18488 (1977)
- 1976CO1B Colgate, Audouze and Fowler, OAP-458 (1976)
- 1976CO1C Couvert, Meson-Nucl. Phys., 1976 (AIP, 1976) p. 310
- 1976DO1B Dodder, Polarization, Zurich, 1975 (Birkhauser Verlag, 1976) 167
- 1976DO1C Dodder et al, LA 6186 MS (1976)
- 1976DR1A Drogg, in Lowell Conf., CONF-760715-P2 (1976) p. 1384
- 1976DR1B Drogg, Smith and Woods, LA 6262 MS (1976)
- 1976DR1C Dries et al, Bull. Amer. Phys. Soc. 21 (1976) 926
- 1976DU06 O. Dumbrajs and M. Staszal, J. Phys. (London) G2 (1976) 795
- 1976FA03 B. Fagerstrom, J. Kallne, O. Sundberg and G. Tibell, Phys. Scr. 13 (1976) 101
- 1976FA09 J. Fain, J. Gardes, A. Lefort, L. Meritet, J.F. Pauty, G. Peynet, M. Querrou, F. Vazeille and B. Ille, Nucl. Phys. A262 (1976) 413
- 1976FO21 C.-M. Fou, Y.-C. Liu, C.-C. Hsu and S.-L. Huang, J. Phys. G2 (1976) 847

1976GAYV D.I. Garber and R.R. Kinsey, BNL 325 (1976)
 1976GO1B Goddard, Rohrig, Knutson and Haeberli, Nucl. Instrum. Meth. 137 (1976) 451
 1976GR10 W. Gruebler, P.A. Schmelzbach, V. Konig, B. Jenny, R. Risler, H.R. Burgi and J. Nurzynski, Nucl. Phys. A271 (1976) 29
 1976GR1D Gruebler, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 307
 1976HA1C Hackenbroich, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1975) 133
 1976HE07 P. Heiss and H.H. Hackenbroich, Phys. Lett. B61 (1976) 339
 1976HU1A Hudomalj-Gabitzsch et al, Bull. Amer. Phys. Soc. 21 (1976) 998
 1976IR02 M. Irshad, J. Asai, S. Sen, R. Pigeon and R.J. Slobodrian, Nucl. Phys. A265 (1976) 349
 1976IR1B Irvine et al, Ann. Phys. 102 (1976) 129
 1976JA1C Jarvis, J. Phys. G2 (1976) 603
 1976KO21 Y. Koike, Prog. Theor. Phys. 55 (1976) 2016
 1976LE1E Levin and Strikman, Sov. J. Nucl. Phys. 23 (1976) 216
 1976LE22 H. Lesniak, L. Lesniak and A. Tekou, Nucl. Phys. A267 (1976) 503
 1976LI15 P.W. Lisowski, R.L. Walter, G.G. Ohlsen and R.A. Hardekopf, Phys. Rev. Lett. 37 (1976) 809
 1976LI1E Lindner, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 817
 1976LI1G Lisowski et al, Bull. Amer. Phys. Soc. 21 (1976) 534
 1976LY1A Lyovshin, Nemets and Yasnogorodsky, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 542
 1976MA34 J.L. Matthews, D.J.S. Findlay, S.N. Gardiner and R.O. Owens, Nucl. Phys. A267 (1976) 51
 1976ME13 H.O. Meyer, G.G. Ohlsen, R.A. Hardekopf, R.V. Poore and J.W. Sunier, Nucl. Phys. A265 (1976) 280
 1976MO1A Motonaga et al, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 837
 1976NA04 N. Nakamura, Y. Takahashi and T. Kohmura, Phys. Lett. B60 (1976) 439
 1976OH02 G.G. Ohlsen, R.A. Hardekopf, N. Jarmie, R.V. Poore, R.L. Walter and P.W. Lisowski, Phys. Rev. C14 (1976) 1688
 1976OH04 G. Ohlen and T. Berggren, Nucl. Phys. A272 (1976) 21
 1976OH1B Ohlsen, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 315
 1976OK1A Okumusoglu, Blyth and Dahme, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 560
 1976RA1B Ramavataram et al, Bull. Amer. Phys. Soc. 21 (1976) 774

- 1976RO1B Roy et al, Bull. Amer. Phys. Soc. 21 (1976) 774
- 1976RO1C Roy et al, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 540
- 1976RO1G Roy et al, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 548
- 1976RO1H Roy, Rad, Conzett and Seiler, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 554
- 1976RU04 D.W. Rule and Y. Hahn, Phys. Rev. C14 (1976) 1102
- 1976SC15 P.A. Schmelzbach, W. Gruebler, V. Konig, R. Risler, D.O. Boerma and B. Jenny, Nucl. Phys. A264 (1976) 45
- 1976SC26 H. Schroder, K.K. Kern, K. Schmidt and D. Fick, Nucl. Phys. A269 (1976) 74
- 1976SE03 F.Seiler, Phys.Lett. B61 (1976) 144
- 1976SE1B Seiler, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 159
- 1976SE1C Seiler, Roy, Conzett and Rad, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 550
- 1976SO01 A.M. Sourkes, A. Houdayer, W.T.H. van Oers, R.F. Carlson and R.E. Brown, Phys. Rev. C13 (1976) 451
- 1976SP1A D.A. Sparrow and W.J. Gerace, Nucl. Phys. A260 (1976) 474
- 1976ST1B Stahel, Jahn, Wozniak and Cerny, Bull. Amer. Phys. Soc. 21 (1976) 1295
- 1976SU06 J.W. Sunier, R.V. Poore, R.A. Hardekopf, L. Morrison, G.C. Salzman and G.G. Ohlsen, Phys. Rev. C14 (1976) 8
- 1976VA29 G.V. Valskii, Yad. Fiz. 24 (1976) 270; Sov. J. Nucl. Phys. 24 (1976) 140
- 1976VO1A Von Oertzen and Flynn, Ann. Phys. 95 (1976) 326
- 1976WA12 R.E. Warner, G.C. Ball, W.G. Davies, A.J. Ferguson and J.S. Forster, Nucl. Phys. A269 (1976) 286
- 1976WA1B Walter, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 377
- 1976WA1C Walter and Lisowski, in Lowell Conf., CONF-760715-P2 (1976) p. 1061
- 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, Phys. Rev. C14 (1976) 815
- 1977AL06 G.D. Alkhazov, Nucl. Phys. A280 (1977) 330
- 1977AL1E Alkhazov et al, Pisma V. Zh. Eksp. Teor. Fiz. 26 (1977) 110
- 1977AR01 L.G. Arnold, B.C. Clark and R.L. Mercer, Lett. Nuovo Cim. 18 (1977) 151
- 1977AR09 N. Arena, S. Cavallaro, M. Lattuada, R. Potenza, M.L. Sperduto, V. D'Amico, G. Fazio, S. Femino and S. Jannelli, Lett. Nuovo Cim. 19 (1977) 125
- 1977AS01 E. Aslanides, T. Bauer, R. Bertini, R. Beurtey, A. Boudard, F. Brochard, G. Bruge, A. Chaumeaux, H. Catz, J.M. Fontaine et al, Phys. Lett. B68 (1977) 221
- 1977BA19 T. Bauer, A. Boudard, H. Catz, A. Chaumeaux, P. Couvert, M. Garcon, J. Guyot, D. Legrand, J.C. Lugol, M. Matoba et al, Phys. Lett. B67 (1977) 265
- 1977BA1M Bassalleck et al, in Proc., Zurich-Sin (1977) 35

- 1977BA1N D. Baye, J. Deenen and Y. Salmon, Nucl. Phys. A289 (1977) 511
- 1977BE1H Beynon and Constantine, J. Phys. G3 (1977) 81
- 1977BE34 J. Berger, J. Duflo, L. Goldzahl, F. Plouin, G. Bizard, C. Le Brun, J. Oostens, M. Van Den Bossche, L. Vu Hai, F.L. Fabbri et al, Lett. Nuovo Cim. 19 (1977) 287
- 1977BE50 F. Becker and I. Strykowski, Nucl. Phys. A289 (1977) 425
- 1977BI05 G. Bizard, C. le Brun, J. Berger, J. Duflo, L. Goldzahl, F. Plouin, J. Oostens, M. van den Bossche, L. Vu Hai, F.L. Fabbri et al, Nucl. Phys. A285 (1977) 461
- 1977BO24 J.E. Bond and F.W.K. Firk, Nucl. Phys. A287 (1977) 317
- 1977BR17 F.P. Brady, N.S.P. King, B.E. Bonner, M.W. McNaughton, J.C. Wang and W.W. True, Phys. Rev. C16 (1977) 31
- 1977BR1C Bruton et al, in Proc., Zurich-Sin (1977) 158
- 1977BR1E Bruge, High Energy Phys. and Nucl. Phys. Structure, Zurich 1977, Birkhauser Verlag (1977) p. 69
- 1977BR25 W. Bretfeld, W. Burgmer, H. Eichner, D. Gola, C. Heinrich, H.J. Helten, H. Kretzer, K. Prescher and W. Schnorrenberg, Z. Phys. A282 (1977) 365
- 1977CH1C Chuu and Han, J. Phys. G3 (1977) 555
- 1977CH1D Chwioroth, Tang and Thompson, Fizika (Yugoslavia) 9, Suppl. 2 (1977) 13; Phys. Abs. 581 (1978)
- 1977CL1A W.L. Clinton, K.P. Mandal, and J.A. McClure, Phys. Rev. Lett. 38 (1977) 143
- 1977CO1B H.E. Conzett and F. Seiler, Nucl. Phys. A290 (1977) 93
- 1977DE08 C. Detraz, Phys. Lett. B66 (1977) 333
- 1977DO01 D.C. Dodder, G.M. Hale, N. Jarmie, J.H. Jett, P.W. Keaton, Jr., R.A. Nisley and K. Witte, Phys. Rev. C15 (1977) 518
- 1977DR1B Dries et al, Bull. Amer. Phys. Soc. 22 (1977) 1016
- 1977DY01 R. Dymarz and A. Malecki, Phys. Lett. B66 (1977) 413
- 1977GE01 J.V. Geaga, M.M. Gazzaly, G.J. Igo, J.B. McClelland, M.A. Nasser, A.L. Sagle, H. Spinka, J.B. Carroll, V. Perez-Mendez and E.T.B. Whipple, Phys. Rev. Lett. 38 (1977) 1265
- 1977HA06 R.A. Hardekopf and G.G. Ohlsen, Phys. Rev. C15 (1977) 514
- 1977HI09 R.-P. Hirsekorn and L.J. Weigert, Nucl. Phys. A288 (1977) 439
- 1977IG1A Igo et al, Bull. Amer. Phys. Soc. 22 (1977) 562
- 1977IG1B Igo et al, in Tokyo (1977), Contrib. Papers p. 13, 15
- 1977IG1C Igo et al, in Proc., Zurich-Sin. (1977) 219
- 1977JA07 N. Jarmie and J.H. Jett, Phys. Rev. C16 (1977) 15

- 1977JA12 N. Jarmie, Phys. Rev. C16 (1977) 2090
- 1977JO1A de Jong et al, in Tokyo (1977), Contrib. Papers p. 18
- 1977KA10 O. Karban, A.K. Basak, C.O. Blyth, W. Dahme, J.B.A. England, J.M. Nelson, N.T. Okumusoglu, S. Roman, G.G. Shute and R. Vlastou, J. Phys. (London) G3 (1977) 571
- 1977KA1D Kaneko and Kamada, Prog. Theor. Phys. 57 (1977) 1277
- 1977KA1E Kanada and Kaneko, in Tokyo (1977), Contrib. Papers p. 39
- 1977KA1F Kanada, Kaneko, Nagata and Nomoto, in Tokyo (1977), Contrib. Papers p. 40
- 1977KA1G Kanada and Kaneko, in Tokyo (1977), Contrib. Papers p. 471
- 1977KA1H Kanada and Kaneko, in Tokyo (1977), Contrib. Papers p. 472
- 1977KI1C Kim, Nucl. Instrum. Meth. 145 (1977) 9
- 1977KL03 R. Klem, G. Igo, R. Talaga, A. Wriekat, H. Courant, K. Einsweiler, T. Joyce, H. Kagan, Y. Makdisi, M. Marshak et al, Phys. Rev. Lett. 38 (1977) 1272
- 1977KL08 R. Klem, G. Igo, R. Talaga, A. Wriekat, H. Courant, K. Einsweiler, T. Joyce, H. Kagan, Y. Makdisi, M. Marshak et al, Phys. Lett. B70 (1977) 155
- 1977LA1C M. Lassaut and N. Vinh Mau, Phys. Lett. B70 (1977) 14
- 1977LE1D D.R. Lehman and B.F. Gibson, Phys. Rev. C16 (1977) 1275
- 1977LE1E Lesniak and Lesniak, in Proc., Zurich-Sin (1977) 193
- 1977MC1C McNally, Bull. Amer. Phys. Soc. 22 (1977) 581
- 1977ME06 H.O. Meyer, T.P. Cleary and W. Haeberli, Nucl. Instrum. Meth. 140 (1977) 85
- 1977MI13 D. Miljanic, E. Andrade, J. Hudomalj-Gabitzsch, G.S. Mutchler and G.C. Phillips, Nucl. Phys. A290 (1977) 27
- 1977PH01 R.J. Philpott, Nucl. Phys. A289 (1977) 109
- 1977PR06 K. Prescher, W. Bretfeld, W. Burgmer, H. Eichner, H.J. Helten, H. Klein, H. Kretzer, H. Stehle and W.A. Wohlfarth, Nucl. Phys. A286 (1977) 142
- 1977RE1A Redish, in Momentum Wave Functions, 1976, Ed. Devins, AIP Conf. Proc. 37 (AIP, New York, 1977) 111
- 1977RI1B Rinat, High Energy Phys. and Nucl. Structure, Zurich, 1977 (Birkhauser Verlag, 1977) p. 303
- 1977RO01 E. Rosario-Garcia and R.E. Benenson, Nucl. Phys. A275 (1977) 453
- 1977RO02 P.G. Roos, N.S. Chant, A.A. Cowley, D.A. Goldberg, H.D. Holmgren and R. Woody, III, Phys. Rev. C15 (1977) 69
- 1977RO18 J. Roth, W. Bretfeld, W. Burgmer, H. Eichner, D. Gola, C. Heinrich, H.J. Helten, H. Kretzer, K. Prescher and W. Schnorrenberg, Z. Phys. A283 (1977) 345

- 1977RO1E Roos, in Momentum Wave Functions, 1976, Ed. Devins, AIP Conf. Proc. 37 (AIP, New York, 1977) 32
- 1977SA1C Saito, Prog. Theor. Phys. Suppl. 62 (1977) 11
- 1977SA1E Saito et al., in Tokyo (1977), Contrib. Papers p. 12
- 1977SA21 K. Sagara, M. Hara, N. Takahashi, T. Motobayashi, F. Takeuchi, F. Soga and Y. Nogami, J. Phys. Soc. Jpn. 42 (1977) 732
- 1977SE09 F. Seiler, Nucl. Phys. A286 (1977) 1
- 1977SE1C F. Seiler and H.W. Roser, Phys. Lett. B69 (1977) 411
- 1977SH1B Shimodaya, in Tokyo (1977), Contributed Papers p. 43
- 1977SL04 I. Slaus, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, J.M. Lambert, P.A. Treado and R.A. Moyle, Nucl. Phys. A286 (1977) 67
- 1977ST06 K. Stephenson, L.D. Knutson and W. Haeberli, Nucl. Phys. A277 (1977) 365
- 1977ST1D Stubeda and Tang, Proc. Int. Conf. Nucl. Structure, Tokyo, (1977), J. Phys. Soc. Jpn. Suppl. 44 (1978) 236
- 1977ST30 A.W. Stetz, J.M. Cameron, D.A. Hutcheon, R.H. McCamis, C.A. Miller, G.A. Moss, G. Roy, J.G. Rogers, C.A. Goulding and W.T.H. van Oers, Nucl. Phys. A290 (1977) 285
- 1977TE1A Terrien, in Tokyo (1977), Contrib. Papers p. 843
- 1977TH07 D.R. Thompson, R.E. Brown, M. LeMere and Y.C. Tang, Phys. Rev. C16 (1977) 1
- 1977TH09 D.R. Thompson, M. LeMere and Y.C. Tang, Nucl. Phys. A286 (1977) 53
- 1977WA05 C.N. Waddell, E.M. Diener, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, A.G. Pieper, R.B. Theus, C.C. Chang and N.S. Chant, Nucl. Phys. A281 (1977) 418
- 1977WA06 S.J. Wallace and Y. Alexander, Phys. Rev. Lett. 38 (1977) 1269
- 1977WA1C Wallace, Bull. Amer. Phys. Soc. 22 (1977) 561
- 1977WA1E Wallace, Bull. Amer. Phys. Soc. 22 (1977) 1264
- 1977WE03 R.B. Weisenmiller, N.A. Jelley, D. Ashery, K.H. Wilcox, G.J. Wozniak, M.S. Zisman and J. Cerny, Nucl. Phys. A280 (1977) 217
- 1977WE1B Weisenmiller, Thesis, Univ. California, Berkeley (1976) - LBL-5077 (1977)
- 1977YO04 S.K. Young and C.W. Wong, Phys. Rev. C15 (1977) 2146
- 1977YO1B York et al, Bull. Amer. Phys. Soc. 22 (1977) 532
- 1978AH01 M.U. Ahmed and P.E. Shanley, Phys. Rev. C17 (1978) 1505
- 1978AR10 N. Arena, S. Cavallaro, R. Potenza, V. D'Amico, G. Fazio, S. Femino, G. Giardina, S. Jannelli and M. Lombardi, Lett. Nuovo Cim. 23 (1978) 1

- 1978BE30 J. Berger, J. Duflo, L. Goldzahl, J. Oostens, F. Plouin, F.L. Fabbri, P. Picozza, L. Satta, C. Le Brun and D. Legrand, Phys. Rev. Lett. 41 (1978) 152
- 1978BI1C Bizard and Tekou, in Proc., Cluster, Winnipeg (1978) D1
- 1978BR1E Brown, Invited Paper, 3rd Int. Conf. on Clustering Aspects of Nucl. Structure and Nucl. reactions, Winnipeg (1978)
- 1978CA05 J.M. Cameron, L.G. Greeniaus, D.A. Hutcheon, R.H. McCamis, M.S. de Jong, B.T. Murdoch, W.T.H. van Oers, J.G. Rogers and A.W. Stetz, Phys. Lett. B74 (1978) 31
- 1978CA13 G. Calvi, S. Cavallaro, A.S. Figuera, R. Fonte and C. Milone, Lett. Nuovo Cim. 22 (1978) 536
- 1978CA1E Carey et al, Bull. Amer. Phys. Soc. 23 (1978) 594
- 1978CH1G Chant, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. & Nucl. Rxns, Winnipeg (1978)
- 1978CL1C Clark, Arnold and Mercer, Bull. Amer. Phys. Soc. 23 (1978) 964
- 1978EP1C Epstein et al, Bull. Amer. Phys. Soc. 23 (1978) 625
- 1978FI1E Fick, Invited Talk, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. & Nucl. Rxns, Winnipeg (1978)
- 1978FO1G Fonseca, Revai and Matveenko, in Proc. Cluster, Winnipeg (1978) B1
- 1978FO33 J. Fong, T.S. Bauer, G.J. Igo, G. Pauletta, R. Ridge, R. Rolfe, J. Soukup, C.A. Whitten, Jr., G.W. Hoffmann, N. Hintz et al, Phys. Lett. B78 (1978) 205
- 1978FR02 H. Friedrich, Nucl. Phys. A294 (1978) 81
- 1978GL1B V.V. Glagolev, R.M. Lebedev, I.S. Saitov, V.N. Streltsov, L.I. Zuravleva, A.N. Gorbunov, K.U. Khayretdinov, G. Martinska, I. Patocka, M. Seman et al, Phys. Rev. C18 (1978) 1382
- 1978GO1D J. Gosset, J.I. Kapusta and G.D. Westfall, Phys. Rev. C18 (1978) 844
- 1978GR07 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.V. Ostashko, V.M. Pugach, V.A. Pilipchenko, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 147; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 126
- 1978GU15 O.P. Gupta, A. Bond and P.M.S. Lesser, Phys. Rev. C18 (1978) 1075
- 1978HE2B E.M. Henley, Nucl. Phys. A300 (1978) 273
- 1978IG1A Igo et al, Bull. Amer. Phys. Soc. 23 (1978) 47
- 1978IG1B Igo, Revs. Mod. Phys. 50 (1978) 523
- 1978KA11 T. Kambara, M. Takai, M. Nakamura and S. Kobayashi, J. Phys. Soc. Jpn. 44 (1978) 704
- 1978KO13 Y. Koike, Nucl. Phys. A301 (1978) 411

- 1978KO1F Koene et al, in Proc., Cluster, Winnipeg (1978) D9
- 1978LA02 V. Layly and R. Schaeffer, Phys. Rev. C17 (1978) 1145
- 1978LE1H Le Mere and Tang, in Proc., Cluster, Winnipeg (1978) A21
- 1978LE23 S.W-L. Leung and H.S. Sherif, Can. J. Phys. 56 (1978) 1116
- 1978MA37 Z. Majka, Phys. Lett. B76 (1978) 161
- 1978MC06 R.H. McCamis, J.M. Cameron, L.G. Greeniaus, D.A. Hutcheon, C.A. Miller, G.A. Moss, G. Roy, M.S. de Jong, B.T. Murdoch, W.T.H. van Oers et al, Nucl. Phys. A302 (1978) 388
- 1978MC1C McNally, Nucl. Fusion 18 (1978) 133
- 1978ME05 G. Mechttersheimer, G. Buche, U. Klein, W. Kluge, H. Matthay, D. Munchmeyer and A. Moline, Phys. Lett. B73 (1978) 115
- 1978ME1C A.Z. Mekjian, Phys. Rev. C17 (1978) 1051
- 1978NA08 H. Nakamura, H. Noya, S.E. Darden and S. Sen, Nucl. Phys. A305 (1978) 1
- 1978NA1H M.A. Nasser, M.M. Gazzaly, J V. Geaga, B. Hoistad, G. Igo, J.G. McClelland, A.L. Sagle, H. Spinka, J.B. Carroll, V. Perez-Mendez et al, Phys. Rev. C17 (1978) 1748
- 1978RO06 E. Rost, J.R. Shepard and D.A. Sparrow, Phys. Rev. C17 (1978) 1513
- 1978SA07 K. Sagara, T. Motobayashi, N. Takahashi, Y. Hashimoto, M. Hara, Y. Nogami, H. Nakamura and H. Noya, Nucl. Phys. A299 (1978) 77
- 1978SE01 F. Seiler, F.N. Rad, H.E. Conzett and R. Roy, Nucl. Phys. A296 (1978) 205
- 1978SL1B Slaus, Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 57
- 1978SM1C Smirnov et al, Bull. Amer. Phys. Soc. 23 (1978) 51
- 1978SP03 C. Spitaleri, M. Lattuada, F. Riggi, N. Arena and D. Vinciguerra, Lett. Nuovo Cim. 21 (1978) 345
- 1978YE1A Yeh, Bull. Amer. Phys. Soc. 23 (1978) 553
- 1978YO1A York et al, Bull. Amer. Phys. Soc. 23 (1978) 602
- 1978YO1B S.K. Young and C.W. Wong, Phys. Lett. B77 (1978) 41