

Energy Levels of Light Nuclei $A = 6$

T. Lauritsen ^a and F. Ajzenberg-Selove ^b

^a *California Institute of Technology, Pasadena, California*

^b *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics* 78 (1966), p. 1. This version of $A = 6$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed July 01, 1965)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 6$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^6\text{He}\$](#) , [\${}^6\text{Li}\$](#) , [\${}^6\text{Be}\$](#)

B. Tables of Recommended Level Energies:

[Table 6.1](#): Energy levels of ${}^6\text{He}$

[Table 6.4](#): Energy levels of ${}^6\text{Li}$

[Table 6.9](#): Energy levels of ${}^6\text{Be}$

C. [References](#)

D. Figures: [\${}^6\text{He}\$](#) , [\${}^6\text{Li}\$](#) , [\${}^6\text{Be}\$](#) , [Isobar diagram](#)

${}^6\text{He}$
(Figs. 4 and 7)

GENERAL:

See ((1960PH1A, 1960TA1C, 1961AH1A, 1961BA1E, 1962IN02, 1962IN1A, 1963BO1K, 1963MO1F, 1963VL1A, 1964GR1J, 1964WA1E, 1965BO1C, 1965LO1G).

Ground state : $J = 0$ ((1958CO68).

1. ${}^6\text{He}(\beta^-){}^6\text{Li}$ $Q_m = 3.510$

The decay proceeds to the ground state of ${}^6\text{Li}(1^+)$ and is a super-allowed Gamow-Teller transition. Recently reported end points are: $E_\beta = 3.50 \pm 0.05$ ((1952WU22), 3.50 ± 0.02 ((1956SC40), 3.508 ± 0.015 ((1963VI06), 3.508 ± 0.004 MeV ((1963JO04). The weighted mean decay energy, including recoil energy of 1.4 keV, leads to ${}^6\text{He}-{}^6\text{Li} = 3509.8 \pm 3.8$ keV ((1963JO04). Half lives are listed in Table 6.2. With a half life of 802 msec and $E_{\beta^-}(\text{max}) = 3508.4$ keV, $ft = 802$ ((1965BA2C).

The electron-neutrino correlation has the form $W(\theta) = 1 + \alpha(\nu/c) \cos \theta$ with $\alpha = -0.39 \pm 0.05$ ((1958HE46, 1959AL10), $\alpha = -0.353^{+0.033}_{-0.053}$ ((1961RI03), $\alpha = -0.319 \pm 0.028$ ((1963VI06), $\alpha = -0.3343 \pm 0.0030$ ((1963JO15), in good agreement with the expected value $\alpha = -\frac{1}{3}$ for pure axial vector interaction. An upper limit to the possible contribution of tensor interaction is 0.4% ((1963JO15). See also ((1958CS88, 1959AL1E, 1959UB1A, 1961KO1H, 1961YU01, 1963CA1H, 1963DA04, 1963KI1D, 1964DE1G, 1965LO1H).

2. (a) ${}^3\text{H}(t, n){}^5\text{He}$ $Q_m = 10.374$ $E_b = 12.302$
(b) ${}^3\text{H}(t, \alpha)2n$ $Q_m = 11.332$

At $E_t = 1.9$ MeV, the α -spectrum observed at 30° extends from 1 to 7 MeV, with peaks at $E_\alpha = 2$ and 5 MeV. The same general shape is observed at other angles and for $E_t = 0.95$ to 2.1 MeV. These peaks are attributed to a two-stage process involving formation and breakup of ${}^5\text{He}$ in the $P_{\frac{3}{2}}$ and $P_{\frac{1}{2}}$ states and are superposed on the three-body spectrum, reaction (b). Structure observed near the end point may indicate a correlation between the two neutrons ((1958JA06). Alpha spectra observed at $\theta = 90^\circ$, $E_t = 0.2$ to 1.0 MeV, likewise exhibit a structure indicating appreciable contribution of ${}^5\text{He}_{g.s.}$; a peak ascribed to $\alpha + n^2$ is reported ((1961GO18).

At $E_t = 1.48$ MeV, the neutron spectrum shows a continuum from 0 to 12 MeV with a broad peak at 11.3 MeV, corresponding to formation of ${}^5\text{He}$ in the ground state ((1957BA10).

The cross section for neutron production rises monotonically from 0.1 to 2.2 MeV ((1951AG30, 1957JA37, 1958JA06). The total cross section rises monotonically from 10 mb at $E_t = 60$ keV to 82 mb at 1.14 MeV ((1962GO1Q). At $E_t = 1.90$ MeV, the total cross section for production

Table 6.1: Energy levels of ${}^6\text{He}$

E_x (MeV \pm keV)	$J^\pi; T$	Γ (keV)	Decay	Reactions
g.s.	$0^+; 1$		β^-	1, 5, 7, 8, 9, 10, 11, 12
1.797 ± 25	$(2)^+$	113 ± 20	n	9, 10, 11, 12
(≈ 15)	$(1^-, 2^-)$			9

of α -particles is 106 ± 5 mb ((1958JA06). The zero-energy cross section factor $S_0 \approx 300$ keV \cdot b ((1964PA1A). See also ((1964BA2B).

3. ${}^3\text{H}(t, t){}^3\text{H}$

$$E_b = 12.302$$

Differential scattering cross sections have been measured at $E_t = 1.58$ to 2.01 MeV by ((1956HO12). At $E_t = 1.90$ MeV, $\theta = 30^\circ$, $\sigma(\theta) = 286$ mb/sr ($\pm 5\%$) ((1958AL05). A phase shift analysis shows only ${}^1\text{S}$ hard-sphere scattering, with $R = 2.35$ fm ((1955FR1C).

4. ${}^4\text{He}(t, p){}^6\text{He}$

$$Q_m = -7.512$$

Not reported.

5. ${}^6\text{Li}(n, p){}^6\text{He}$

$$Q_m = -2.727$$

See ${}^7\text{Li}$ and ((1960VA1D).

6. ${}^6\text{Li}(t, {}^3\text{He}){}^6\text{He}$

$$Q_m = -3.491$$

Not reported.

7. ${}^6\text{Li}(\gamma, p){}^6\text{He}$

$$Q_m = -9.980$$

See ((1954TH16) and ${}^7\text{Li}$.

Table 6.2: Half life of ${}^6\text{He}$ ^a

$\tau_{1/2}$ (msec)	Reference	$\tau_{1/2}$ (msec)	Reference
850 ± 50	(1946SO05)	799 ± 3	(1954KL36)
870 ± 60	(1947CA15)	850 ± 30	(1955RU06)
820 ± 60	(1948KN13)	852 ± 16	(1956VE10)
823 ± 13	(1949HO24)	830 ± 20	(1958HE46)
860 ± 30	(1952SH44)	797 ± 3 ^a	(1962BI14)
840 ± 30	(1952VE1A)	862 ± 17	(1962MA38)
830 ± 30	(1953BA04)	830 ± 20	(1963VI06)

^a It is suggested that the longer half lives found in the older measurements may be due to impurities (1962BI14). We adopt here the mean of all cited values, weighted by the square of stated errors: 802 ± 3 msec.

8. ${}^7\text{Li}(n, d){}^6\text{He}$ $Q_m = -7.755$

See ${}^8\text{Li}$.

9. ${}^7\text{Li}(p, 2p){}^6\text{He}$ $Q_m = -9.980$

The summed proton spectrum at $E_p = 155$ to 450 MeV shows two peaks with $Q \approx -11.6$ and -25.4 MeV corresponding to ${}^6\text{He}^*(0, 1.80)$, and an excited state with $J = 1^-$ or 2^- at $E_x \approx 15$ MeV: see Table 6.3. Angular distributions indicate that the higher energy peak corresponds to ejection of a p-proton while the lower results from removal of an s-proton: see ((1957TY35, 1958MA1B, 1958TY49, 1961GA09, 1961PU1A, 1962GA09, 1962GA23, 1962GO1P, 1962TI01, 1964TI02, 1965RI1A, 1965TY1A)). See also ${}^7\text{Li}$.

10. ${}^7\text{Li}(d, {}^3\text{He}){}^6\text{He}$ $Q_m = -4.486$

At $E_d = 14.4$ MeV, the ground state and the 1.80 MeV level are observed. The angular distributions analyzed by pick-up theory indicate even parity for both states (1955LE24); $\theta^2 = 0.025$ and 0.008 ((1960MA32). See also ((1960HA14).

11. ${}^7\text{Li}(t, \alpha){}^6\text{He}$ $Q_m = 9.834$

Table 6.3: ${}^6\text{He}$ levels from ${}^7\text{Li}(p, 2p){}^6\text{He}$

Q ($l = 1$) (MeV)	Γ (MeV)	Q ($l = 0$) (MeV)	Γ (MeV)	Reference
10.5 ± 1.6	5.0	24.9 ± 1.6	5.5	(1958TY49)
10.2 ± 1.6		23 ± 1.5		(1962GA09)
11.5 ± 2.5		25 ± 2.5		(1962GO1P)
10.1 ± 1.4		24.1 ± 1.5		(1961PU1A)
11.8 ± 0.3		25.5 ± 0.4		(1965TY1A)
11.3 ± 0.5		25.8 ± 0.6		(1964TI02)
11.6 ± 0.2		25.4 ± 0.3		mean

The energy of the first excited state is 1.71 ± 0.01 MeV, $\Gamma \lesssim 100$ keV ((1954AL35, (1960AL10); 1.797 ± 0.025 MeV, $\Gamma = 113 \pm 20$ keV ((1965AJ01). Evidence has also been reported for states at 3.4 ± 0.2 MeV ((1960AL10), (6.0 ± 0.9) and for one or more states at 9.3 ± 0.7 MeV ((1956MA1R, 1956MA50): see, however ((1965AJ01). The width of the proton group corresponding to the 1.8 MeV state and the absence of ${}^6\text{He}^*$ recoils implies that this state decays predominantly into ${}^4\text{He} + 2n$ ((1954AL35, 1963MA1P, 1965AJ01). The branching ratio Γ_γ/Γ_n is $< 4 \times 10^{-4}$ ((1964HU1A). Angular distributions at $E_t = 0.24$ keV are consistent with $J = 0$ and 2 for the ground state and 1.8 MeV level, respectively ((1954AL38). At $E_t = 13$ MeV, the two angular distributions are strongly peaked forward ((1965AJ01). See also ((1956MA09, 1961HO01, 1961HO23) and ${}^{10}\text{Be}$.

$$12. {}^9\text{Be}(n, \alpha){}^6\text{He} \quad Q_m = -0.601$$

((1964GA11) report α -groups to excited states at $E_x = 1.66 \pm 0.26$, 3.29 ± 0.38 and 6.05 ± 0.26 MeV. See also ((1963AL18) and ${}^{10}\text{Be}$.

⁶Li
(Figs. 5 and 7)

GENERAL:

See (1955AU1A, 1955LA1C, 1956ME1A, 1957FR1B, 1957HU1C, 1957LE1E, 1958PI1A, 1959BA1M, 1959BR1E, 1959FE1B, 1959SK1A, 1959UB1A, 1960AN1B, 1960JA1G, 1960KO1D, 1960PH1A, 1960TA1C, 1960WA1F, 1961BA1E, 1961KO1G, 1961SH1B, 1961TA05, 1961VA1G, 1962CO1D, 1962CR09, 1962DI1B, 1962FO08, 1962GA09, 1962IN02, 1962IN1C, 1962IN1A, 1962JA06, 1962ME1C, 1962NA1B, 1962SA1F, 1962ST1E, 1962WA1E, 1963BO1K, 1963BU1C, 1963DA04, 1963EL1D, 1963HA1H, 1963JA1C, 1963JO07, 1963KL1A, 1963KU03, 1963KU1B, 1963MO1F, 1963OL1B, 1963SA1F, 1963SC1J, 1963SC30, 1963VL1A, 1963WA1H, 1964GR1J, 1964JI1A, 1964MA1G, 1964MI16, 1964NE1E, 1964OL1A, 1964SA1F, 1964SH05, 1964ST1B, 1964WA1E, 1965BE1R, 1965BE1H, 1965BO1C, 1965DA1H, 1965EL1B, 1965GO1L, 1965HA1L, 1965JA04, 1965JA1L, 1965LO1G, 1965LO1H, 1965MA1N, 1965MU1A, 1965NE1C, 1965RA1E, 1965SA1J).

Ground state :

$$\mu = +0.82201 \text{ nm (1965FU1G: see also (1962SC15)).}$$

$$Q = (-)0.5 \text{ mb (1958TO34);}$$

$$Q = -(0.80 \pm 0.08) \text{ mb (1964WH01).}$$

1. (a) ${}^3\text{He}({}^3\text{H}, \gamma){}^6\text{Li}$	$Q_m = 15.793$	$E_b = 15.793$
(b) ${}^3\text{He}({}^3\text{H}, p){}^5\text{He}$	$Q_m = 11.138$	
(c) ${}^3\text{He}({}^3\text{H}, p){}^4\text{He} + n$	$Q_m = 12.096$	
(d) ${}^3\text{He}({}^3\text{H}, n){}^5\text{Li}$	$Q_m = 10.131$	
(e) ${}^3\text{He}({}^3\text{H}, d){}^4\text{He}$	$Q_m = 14.321$	
(f) ${}^3\text{He}({}^3\text{H}, {}^3\text{H}){}^3\text{He}$		

Capture γ -rays (reaction (a)) to the first three states of ${}^6\text{Li}$ have been observed for $E({}^3\text{He}) = 1$ to 3 MeV: the differential cross section for the ground state transition (90°) varies from $1.4 \mu\text{b/sr}$ at $E({}^3\text{He}) = 1.1$ MeV to $8 \mu\text{b/sr}$ at 3.0 MeV. At $E({}^3\text{He}) = 1.1$ MeV the transitions to the 2.18 and 3.56 MeV states each have intensities $\approx 15\%$ that of the ground-state transition (1963KO04).

Angular distributions and total cross sections for reactions (b), (c) and (e) are given by (1963KU18) for $E_t = 0.46$ to 1.09 MeV. The total cross section has also been determined for $E({}^3\text{He}) = 0.1$ to 0.8 MeV (1953MO61), $E_t = 0.15$ to 0.97 MeV (1960YO06), and at $E_t = 1.9$ MeV (1963SM03). In the range $E_t = 0.46$ to 1.09 MeV, the cross section increases from 27 to 53 mb (1963KU18). At

Table 6.4: Energy levels of ${}^6\text{Li}$

E_x (MeV \pm keV)	$J^\pi; T$	Γ (keV)	Decay	Reactions
g.s.	$1^+; 0$		stable	1, 4, 5, 6, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24
2.184 ± 2	$3^+; 0$	25 ± 1	γ, d, α	1, 4, 5, 8, 9, 10, 12, 13, 14, 15, 17, 18, 19, 21
3.562 ± 4	$0^+; 1$	< 5	γ	1, 4, 7, 9, 12, 17, 18, 19, 21
4.57 ± 34	$2^+; 0$	350 ± 150	d, α	4, 9, 10, 12, 14, 19
5.36 ± 15	$; (1)$	320 ± 50		9, 12, 18, 19, 21
6.0 ± 200	$1^+; 0$		d, α	4
(6.77)	$(2^-); 0$		d, α	4, 8, 12
(7.8)	$(1^-); 0$		d, α	4, 8, 12
(9)	$(0^-); 0$		d, α	4, 9
(12.5)			γ, n	8
(14.0)	$+$			9
(15.8)	$+$			9

$E_t = 1.9$ MeV, it is 53 mb (1963SM03). The zero-energy cross section factor $S_0 \approx 1000$ keV \cdot b (1964BA2B, 1964PA1A). See also (1961BA40).

The angular distribution of deuterons is isotropic at $E_t = 360$ keV (1953MO61), but from 0.5 to 1 MeV it shows an increasingly strong maximum at 90° . It is suggested that the reaction proceeds via compound nucleus formation and that $l = 1$ is required (1963KU18).

For reaction (d), see (1963SM03). For reaction (f), see (1965LE02). See also (1963IN1A, 1963KU1H).

2. ${}^4\text{He}(d, \gamma){}^6\text{Li}$

$$Q_m = 1.472$$

An upper limit for capture radiation at $E_d = 1.06$ MeV (${}^6\text{Li}^* = 2.18$ MeV) is 0.1 mb (1954SI07). A search for resonant capture radiation at $E_d = 3.1$ MeV (${}^6\text{Li}^* = 3.56$ MeV) yields $\Gamma_{d\alpha} < 0.2$ eV. It is concluded that the intensity of parity non-conserving parts of the wave functions, $F^2 \lesssim 10^{-7}$ (1958WI15).

3. (a) ${}^4\text{He}(d, n){}^5\text{Li}$	$Q_m = -4.190$	$E_b = 1.472$
(b) ${}^4\text{He}(d, p){}^5\text{He}$	$Q_m = -3.182$	
(c) ${}^4\text{He}(d, pn){}^4\text{He}$	$Q_m = -2.225$	

Neutron spectra have been measured at 0° for $E_d = 7.9, 8.9$ and 9.9 MeV, and at several angles at the higher energy. The neutrons show forward peaking, indicative of stripping via reaction (a). The differential cross section (0°) plotted for $E_d = 4.5$ to 18.6 MeV shows a monotonic rise from 0.5 to 150 mb/sr (1962LE12: see also (1964RO1D)). Neutron spectra at $E_d = 18.6$ MeV ($\theta = 0^\circ$ and 180°) indicate participation of reactions (a) (0°) and (b) (180°) (1961RY01). The proton spectra are also forward peaked at $E_d = 20.2$ MeV (1960AR1A) and 24 and 27 MeV (1963ER02). The proton yield over the range from 6.5 to 8.7 MeV excitation in ${}^6\text{Li}$ gives no evidence of a 7.4 MeV, $T = 0$ state (see Table 6.8) (1964OH01).

At $E_d = 20.2$ MeV, the (d, pn) cross section is nearly as large as that for the elastic scattering at $\theta \approx 45^\circ$ (1960AR1A). Reaction (c) is presumably forbidden by isospin selection rules near threshold (1955HE1B). See also (1965TO01).

4. ${}^4\text{He}(d, d){}^4\text{He}$	$E_b = 1.472$
---------------------------------------	---------------

Elastic scattering studies have been carried out by (1949BL66: 0.9 to 3.5 MeV), (1953LA28: 1.0 to 1.2 MeV), (1955GA26: 0.3 to 4.6 MeV), (1947GU1A: 6.5 MeV), (1951BU1C: 7.9 MeV), (1964SE1G, 1964SE07: 2.9 to 11.5 MeV), (1964OH1B: 3.5 to 10 MeV), (1962ST19: 6.0 to 13.7 MeV), (1963RO23, 1964RO1D: 5.7 to 14.3 MeV), (1951AL26: 10.3 MeV), (1954FR22: 13.7 and 19.0 MeV), (1960AR1A: 20.2 MeV), (1964BR1R: 21 MeV), (1963VA14: 24.9 MeV), (1963ER02: 21.3 to 27.3 MeV).

Phase shift analyses have been carried out for the range $E_d = 0.3$ to 4.2 MeV by (1955GA74) and for 3 to 10 MeV by (1964SE07); see also (1964MC1F). Extrapolated phase shifts based on an optical model have been calculated by (1960GA08). The experimental s-wave phase shift decreases monotonically from $E_d = 0.3$ to 10 MeV: in the range 0 to 4 MeV it can be accounted for by hard-sphere scattering with $r = 5$ fm; if a radius of 3.5 fm is chosen, a contribution from the ground state is required, with $\theta_\alpha^2(\text{g.s.}) = 0.51$ (1955GA74, 1960GA08). The d-wave phase shifts are split and exhibit resonance at $E_x = 2.18$ (${}^3\text{D}_3$), 4.87 (${}^3\text{D}_2$) and 6.24 MeV (${}^3\text{D}_1$): see Table 6.5 (1955GA74, 1964SE07). Single level analysis of the p-wave phase shifts indicate the possible presence of 2^- , 1^- and 0^- levels at 6.8 , 7.8 and 9 MeV (1964SE07: see, however, (1964MC1F)). Preliminary analysis of data in the range $E_d = 5.6$ to 12.5 MeV shows no effect from the presumptive $T = 1$ levels at $E_x = 5.35$, 6.63 and 8.37 MeV (1963RO23), nor from the suggested $T = 0$ state at 7.4 MeV (1964MC1F). At $E_d = 21$ to 27 MeV, increasing complexity of the angular distributions suggest $l = 3$ wave contributions (1963ER02). See also (1964MC1F).

A (d+ α) model has been constructed by (1960GA08) to reproduce the phase shifts of (1955GA74); with this model, phase shifts and polarization parameters have been extrapolated to 14 MeV. Polarization parameters have also been calculated for $E_d = 1.07$ MeV (1959GO1H, 1959PO1C),

Table 6.5: Levels of ${}^6\text{Li}$ from ${}^4\text{He}(d, d){}^4\text{He}$

E_{res} (MeV)	Γ (keV)	l_d	J^π	E_x^a (MeV)	$\theta_d^2{}^b$	Reference
		0	1^+	0	0.51	1955GA74
1.070 ± 0.003	35 ± 5	2	3^+	2.184	0.80	1955GA74
4.57 ± 0.08		2	2^+	4.6	1.0	1955GA74
5.10				4.87	1.59	1964SE07
5.1 – 6.5		2	1^+	5.7		1955GA74
7.16				6.24	1.59	1964SE07
7.95		1	2^-	6.77	0.27	1964SE07
9.5		1	1^-	7.8	0.21	1964SE07
11.6		1	0^-	9	0.21	1964SE07

^a See also (1964MC1F).

^b $R = 3.5$ fm.

at 2 and 3.5 MeV by (1959PH1B). A measurement at $E_d = 1.07$ MeV is consistent with expectation (1961PO01). Tensor polarization parameters for $E_d = 4$ to 7.5 MeV, measured by (1964MC1E, 1964SE1F), are in fair agreement with calculations of (1960GA08, 1964SE07). See also (1960DU1C, 1965GR1Q, 1965SE1E).

5. ${}^4\text{He}({}^3\text{He}, p){}^6\text{Li}$ $Q_m = -4.021$

Differential cross sections for formation of the ground and 2.18 MeV states of ${}^6\text{Li}$ have been measured for $E_\alpha = 27.5$ to 41 MeV. Protons to the 3.56 MeV state were not observed: intensity $< \frac{1}{5}$ of ground state group (1961CH09).

6. ${}^6\text{He}(\beta^-){}^6\text{Li}$ $Q_m = 3.510$

See ${}^6\text{He}$.

7. ${}^6\text{Li}(\gamma, \gamma){}^6\text{Li}$

The width, Γ_γ , of the 3.56 MeV state is in good agreement with that expected from the intermediate coupling shell model (8.7 eV): see Table 6.6 (1959CO67, 1963SK02). See also (1961VA1G, 1962BO17, 1962SE02).

Table 6.6: Levels of ${}^6\text{Li}$ from ${}^6\text{Li}(e, e')$ and ${}^6\text{Li}(\gamma, \gamma)$

E_x (MeV)	J^π	Γ_γ (eV)	$B(E2\uparrow)^a$ (fm 4)	Reference
2.18	3^+	$4_{-1.5}^{+3} \times 10^{-4}$ 3×10^{-5}	32 ± 3	(1960BA47) (1959DA04) ^b
3.56 ^e	0^+	6.2 ± 0.6 4.7 ± 0.9 9 ± 2 $9.1_{-1.5}^{+2.0}$ $8.8_{-1.3}^{+1.9}$		(1960BA47) (1963BA19) ^c (1963BE25) (1959CO67) ^d (1963SK02) ^d
4.52	2^+		18.2 ± 1.5	
5.35	(2^+)	0.16	7 ± 2	(1963BA19)
(7.40)	(1^+)		6 ± 2	
(8.37)	(1^+)		4 ± 2	
(9.3)	$0^+/1^+/2^+$	15/5/3		(1963BA19)
(14.0)	$0^+/1^+/2^+$	24/8/5		(1963BA19)
(15.8)	$0^+/1^+/2^+$	37.5/12.5/7.5		(1963BA19)

^a (1963BE25, 1963BE53): E2 transition assumed; preliminary values.

^b From ${}^6\text{Li}(\gamma, d)$.

^c (1963BA19): M1 transitions assumed.

^d From ${}^6\text{Li}(\gamma, \gamma)$.

^e See also (1964GO15).

8. (a) ${}^6\text{Li}(\gamma, n){}^5\text{Li}$ $Q_m = -5.662$
 (b) ${}^6\text{Li}(\gamma, p){}^5\text{He}$ $Q_m = -4.655$
 (c) ${}^6\text{Li}(\gamma, d){}^4\text{He}$ $Q_m = -1.472$
 (d) ${}^6\text{Li}(\gamma, t){}^3\text{He}$ $Q_m = -15.793$

The total cross section for photoneutrons, measured with $E(\text{brems}) = 5$ to 20 MeV exhibits a peak at $E_\gamma = 12.5$ MeV, $\sigma = 2.8 \pm 0.53$ mb (1959RO62). Measurements with monochromatic γ -rays have been made for $E_\gamma = 5.4$ to 9.0 MeV by (1964GR1A) and from 10 to 32 MeV by (1965BE1D). The former work shows a smooth rise, with a possible peak at 6.75 MeV, while the latter yields a maximum at ≈ 12 MeV, $\sigma \approx 1.6$ mb and a gentle decrease to 0.6 mb at 32 MeV. See also (1951TI06, 1956ED15, 1964SH27). Another peak, at $E_\gamma \approx 26$ MeV is reported by (1963CO15): it is suggested that this peak corresponds to giant resonance excitation of the α -particle core (see also (1963BI10, 1963CO1D)); however, (1965BE1D) find no evidence for this

peak. At $E(\text{brems}) = 17.3$ MeV, 58% of the neutrons are ascribed to reaction (a), 31% to (b). A search for n-p correlations expected on the quasi-deuteron model for (γ, np) yielded a negative result (1960PR06). The integrated cross section (total neutrons) to 40 MeV is about 40 MeV · mb (1963CO15). See also (1964ER1B).

Photoprotons have been observed with E_γ up to 90 MeV: see (1959AJ76) and (1960BA45, 1962CH1B, 1962VO1C). A peak in the proton spectrum corresponding to $E_x = 6.63$ or $E_x = 7.94$ MeV is reported by (1965SH1F). See also (1960KO14, 1961MA29, 1964BA2C, 1964ER1B, 1964MA1X).

The cross section for reaction (c) is $\lesssim 5 \mu\text{b}$ in the range $E_\gamma = 2.6$ to 17 MeV (see (1959AJ76)) consistent with the expected inhibition of dipole absorption by isospin selection rules. In common with other (γ, d) reactions in light nuclei, an effective threshold first appears when sufficient energy is available to release at least one particle in addition to the deuteron (1962CH1B, 1962VO1D, 1962VO1C). Resonant capture at the 2.18 MeV state is observed by (1959DA04): a value $\Gamma_\gamma = 30 \mu\text{eV}$ is obtained. See also (1964KO1F).

For reaction (d), see (1962VO1C, 1964KO1F, 1965SH1F).

9. (a) ${}^6\text{Li}(e, e){}^6\text{Li}$

(b) ${}^6\text{Li}(e, p){}^5\text{He}$ $Q_m = -4.655$

(c) ${}^6\text{Li}(e, d){}^4\text{He}$ $Q_m = -1.472$

Elastic scattering has been studied at $E_e = 187$ (1955ST85, 1956HO93) and 426 MeV (1958BU17). Several different charge distributions provide acceptable fits to the observed form factors. The maximum charge density is 0.064 protons · fm $^{-3}$, $R_{\text{rms}} = 2.70 \pm 0.15$ fm (1958BU17, 1959ME24, 1960JA1G: see also (1958CA1B, 1963BO1K, 1963SC1J)). Magnetic elastic scattering at $E_e = 41.5$ MeV, $\theta = 180^\circ$, has been studied by (1963GO04).

Inelastic spectra have been reported by (1960BA47: $E_e = 40$ MeV; $\theta = 132^\circ, 160^\circ$), (1963BA19: $E_e = 41.5$ MeV; $\theta = 180^\circ$), (1964GO15: 75 MeV), (1963BI10: 101.4 MeV; $\theta = 60^\circ$) and (1963BE25: 180 MeV). Table 6.6 summarizes the results. [For a review of the inelastic work to 1962, see (1962BA1D).] The inelastic spectrum of (1963BI10) shows two major absorption regions, from 9 to 17.5 MeV and from 17.5 to 32 MeV. Analysis of form factors on the assumption of E1 transitions leads to quite different values of $\langle r^2 \rangle$ for the two regions, supporting an $\alpha + d$ model. Fine structure is observed in both regions, possibly corresponding to discrete ${}^6\text{Li}$ levels (1963BE25, 1963BI10, 1964BI04). Form factors for ${}^6\text{Li}^*(2.2)$ and (3.6) have been studied as a function of momentum transfer by (1963BE25, 1963BE53, 1964GO15). Interpretation in terms of nuclear models is discussed by (1963BE53, 1964KU1G, 1965DA1H). See also (1959ME1D, 1959UB1A, 1961PA1A, 1962JA06, 1965LO1J, 1965RA1D).

For reactions (b) and (c), see (1957KE1A).

10. ${}^6\text{Li}(n, n'){}^6\text{Li}^*$

Inelastic neutron groups have been observed corresponding to the 2.18 MeV state with incident neutron energies up to 14 MeV. Differential cross sections are reported from $E_n = 3.4$ to 7.5 MeV (1963BA50, 1963BA1V), and at 14 MeV (1962WO07, 1964AR1C, 1965ME05). Angular distributions up to $E_n = 7.5$ MeV show no obvious indication of direct interaction; at $E_n = 14$ MeV, both elastic and inelastic neutrons (n_0 and n_1) show sharp forward peaking. At both energies, the excitation of ${}^6\text{Li}^*(3.56)$ is weak (1962WO07, 1963BA50). See also (1960KO1C, 1962EL1D, 1962JA1B, 1963EL1C, 1964PE1C, 1965WE1E).

11. ${}^6\text{Li}(n, t){}^4\text{He}$ $Q_m = 4.785$

(1956LI37) calculates $\theta_d^2 = 0.5$ for ${}^6\text{Li}_{\text{g.s.}}$ from the data of (1954FR03). See also ${}^7\text{Li}$.

12. (a) ${}^6\text{Li}(p, p){}^6\text{Li}$

(b) ${}^6\text{Li}(p, 2p){}^5\text{He}$ $Q_m = -4.655$

(c) ${}^6\text{Li}(p, pd){}^4\text{He}$ $Q_m = -1.472$

(d) ${}^6\text{Li}(p, {}^3\text{He}){}^4\text{He}$ $Q_m = 4.021$

Elastic scattering has been studied at $E_p = 31$ MeV by (1963DE01), at 40 MeV by (1960CH1B), and at 155 MeV by (1964TA02). At the latter energy, the polarization at high momentum transfers, $q > 1.7 \text{ fm}^{-1}$ is strikingly different for ${}^6\text{Li}$ and ${}^7\text{Li}$ (1964TA02). Optical model analysis is discussed by (1963DE01, 1963JA1C). See also (1960SA28). Polarization of the scattered protons has been studied by (1962RO20, 1963HW01, 1964MA1Y).

Inelastic groups corresponding to several ${}^6\text{Li}$ states are observed: see Table 6.7. Angular distributions have been determined for $E_p = 4$ to 9 MeV (1963HA49), 31 MeV (1963DE01), 39.8 MeV (1959CH1B: see also (1963JA1C)), 155 MeV (1964JA03) and 185 MeV (1965HA17). The distributions at $E_p = 155$ to 185 MeV indicate dominant E2 transitions to ${}^6\text{Li}^*(2.2, 4.6, 5.4)$ and M1 to ${}^6\text{Li}^*(3.6)$ (1964JA03, 1965HA17). At $E_p = 149$ MeV, a 3.56 MeV γ -ray is observed with a cross section of $0.7 \pm 0.3 \text{ mb}$ (1961CL09). See also (1957LE1E, 1959SY1A).

Reaction (b) has been studied at $E_p = 155$ MeV (1961GA09, 1962GA09, 1962GA23), 185 MeV (1962TI01, 1964TI02) and 460 MeV (1965TY1A). The spectra of summed proton energies ($E_{p1} + E_{p2}$) show two peaks, with $Q = -4.8$ and $Q = -22.4$ MeV. As a function of angle of emission $\theta_1 = -\theta_2$, the higher energy peak exhibits a maximum at the angle corresponding to zero momentum of the target proton ($\theta_1 + \theta_2 \approx 90^\circ$), while the lower has a minimum at this point. The higher peak is thus interpreted as a quasi-free scattering from an $l = 1$ proton ${}^6\text{Li} \rightarrow {}^5\text{He}_{\text{g.s.}} + p$ and the lower corresponding to a $l = 0$ proton, leaving ${}^5\text{He}^*$ in a positive parity configuration with an excitation of about 16 – 17 MeV (see ${}^5\text{He}^* = 16.70 \text{ MeV}$).

Calculations of expected momentum distributions with various wave functions are discussed by (1962BE1J, 1962BE1K, 1962DI1A, 1962IN1C, 1962IN1A, 1962SA1F, 1963BE42, 1963EL1C, 1963JO07, 1963RI1B, 1964LI1D, 1965JA1J, 1965JA04, 1965JA1M, 1965MC1F, 1965RI1A, 1965WI1G).

Table 6.7: Levels of ${}^6\text{Li}$ from (p, p'), (d, d'), ${}^9\text{Be}(p, \alpha)$, ${}^7\text{Li}(d, t)$

Reaction	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	Reference
${}^6\text{Li}(p, p'){}^6\text{Li}^*$	2.188	25.4	(1957BR12)
${}^6\text{Li}(d, d'){}^6\text{Li}^*$	2.186	24.5	(1957BR12)
${}^9\text{Be}(p, \alpha){}^6\text{Li}$	2.192	29	(1957BR12)
${}^9\text{Be}(p, \alpha){}^6\text{Li}$	2.19 ± 0.02	< 35	(1963GR29)
${}^7\text{Li}(d, t){}^6\text{Li}$	(2.18)	< 27	(1957BR12)
mean	2.188 ± 0.006	26.3	(1957BR12)
${}^6\text{Li}(p, p'){}^6\text{Li}^*$	3.559	< 5	(1957BR12)
${}^9\text{Be}(p, \alpha){}^6\text{Li}$	3.561		(1957BR12)
${}^9\text{Be}(p, \alpha){}^6\text{Li}$	3.55 ± 0.02	< 35	(1963GR29)
mean	3.560 ± 0.006	< 5	(1957BR12)
${}^9\text{Be}(p, \alpha){}^6\text{Li}$	4.40 ± 0.12	350 ± 150	(1963GR29)
${}^6\text{Li}(p, p'){}^6\text{Li}^*$	4.4 ± 0.2	≈ 600	(1965HA17)
${}^9\text{Be}(p, \alpha){}^6\text{Li}$	5.32 ± 0.06	280 ± 60	(1963GR29)
${}^6\text{Li}(p, p'){}^6\text{Li}^*$	5.4 ± 0.2	≈ 1000	(1965HA17)
${}^6\text{Li}(p, p'){}^6\text{Li}^*$	≈ 6.5		(1965HA17)
${}^6\text{Li}(p, p'){}^6\text{Li}^*$	≈ 7.5		(1965HA17)

As compared with the analogous case of ${}^7\text{Li}$, there appears to be a difficulty here in accounting for the small width of the observed angular distribution.

At $E_p = 155$ MeV, the angular correlation of outgoing protons and deuterons (reaction (c)) give evidence for a substructure ${}^6\text{Li} = {}^4\text{He} + \text{d}$ in a relative $l = 0$ state with a probability of 20 to 31% (1963RU05). See also (1962RU04, 1963RI1B, 1963SA1F, 1963TA1D, 1964BA1P, 1964DE1F, 1964SA1H, 1965DE1Q, 1965JA1L).

For reaction (d) see (1963TE1B, 1963RU05) and ${}^7\text{Be}$. (1956LI37) gives θ_d^2 for the ground state of ${}^6\text{Li} = 0.30$ ($E_p = 15$ MeV), 0.45 (18.5 MeV).

13. ${}^6\text{Li}(\text{d}, \text{d}'){}^6\text{Li}^*$

Deuteron groups have been observed corresponding to the ground state and the 2.18 MeV level (1957BR12: $E_d = 7.0$ and 7.5 MeV): see Table 6.7. The 3.56 MeV state is not observed at $E_d = 7.5$ MeV, $\theta = 60^\circ$, consistent with its $T = 1$ character. The shape of the angular distribution of the deuterons to the 2.18 MeV state ($E_d = 14.7$ MeV) does not fit simple direct interaction theory (1960HA14: see also (1956HA90, 1964HA05, 1965JO09)). See also (1956SO21, 1956SO33, 1960EL09).

14. ${}^6\text{Li}(\alpha, \alpha){}^6\text{Li}$

Angular distributions of elastic and inelastic scattering (to the 2.18 MeV state) have been studied at $E_\alpha = 10.0$ and 12.5 MeV (1963BL20), at 24 MeV (1964GR39) and at 31.5 MeV (1956WA29). At $E_\alpha = 31.5$ MeV, a group is also observed to the 4.57 MeV state but not to the $T = 1$, 3.56 MeV level (1956WA29: see also (1964GR39)). See also (1962TE1D, 1963TE1B).

15. ${}^6\text{Li}({}^6\text{Li}, {}^6\text{Li}'){}^6\text{Li}^*$

Angular distributions ($\theta_{\text{c.m.}} = 12^\circ$ to 40°) to the ground and 2.18 MeV states have been measured at $E({}^6\text{Li}) = 63$ MeV: they show diffraction-type structure. No evidence of the excitation of the 3.56 MeV $T = 1$ state is seen (1964GA01).

16. ${}^7\text{Li}(\gamma, \text{n}){}^6\text{Li}$ $Q_m = -7.253$

See ${}^7\text{Li}$.

17. ${}^7\text{Li}(\text{p}, \text{d}){}^6\text{Li}$ $Q_m = -5.028$

At $E_p = 17.5$ MeV, angular distributions of deuteron groups corresponding to ${}^6\text{Li}(\text{g.s.})$, (2.18) and (3.56), analyzed by Butler theory give $l_n = 1$ for all these states. The ratios of observed reduced widths (see ${}^7\text{Li}$) are consistent with predictions of the shell model with pure, or nearly pure, LS coupling (1956RE04, 1959BE84, 1960MA32). See also (1961CL09). The bearing of this reaction and its inverse on tests of invariance under time reversal is discussed by (1959HE1C). See also (1964SH07).

$$18. {}^7\text{Li}(\text{d}, \text{t}){}^6\text{Li} \quad Q_m = -0.995$$

At $E_d \approx 15$ MeV, the angular distributions of the tritons indicate $l_n = 1$ (even parity) for the first three states of ${}^6\text{Li}$ (1955LE24, 1957FR1B, 1959HA29, 1960HA14): reported reduced widths are listed in Table 7.7. (1960HA14) report a broad asymmetric bump at $E_x \approx 5.4$ MeV, $\Gamma \approx 600$ keV; no sharp states are reported for $4.4 < E_x < 8.5$ MeV with $\sigma > 0.3$ mb/sr ($E_d = 14.9$ MeV, $\theta = 14^\circ, 25^\circ$). See also (1956SO21, 1956SO33, 1959KU1C, 1959VL24, 1961OG1A, 1961SL06, 1962SL04, 1963OG1A, 1964BL1C).

$$19. {}^7\text{Li}({}^3\text{He}, \alpha){}^6\text{Li} \quad Q_m = 13.325$$

Alpha particle groups have been observed with $E({}^3\text{He}) = 0.6$ to 0.9 MeV (1960AL10), 1.5 MeV (1963CA02), 0.8 to 3 MeV (1965PA03), 4 MeV (1961WO05, 1963KN1C), 0.8 to 5.1 MeV (1963LI16) and 1.3 to 5.5 MeV (1965FO07). (1963LI16) find no evidence for reported levels at 4.3 or 6.6 MeV; (1965CO1F) confirms the levels at 4.3 and 5.35 MeV but finds no evidence for the higher states reported by (1960AL10). The results are summarized in Table 6.8.

$$20. {}^9\text{Be}(\gamma, \text{t}){}^6\text{Li} \quad Q_m = -17.688$$

See (1955AJ61).

$$21. {}^9\text{Be}(\text{p}, \alpha){}^6\text{Li} \quad Q_m = 2.126$$

$$Q_0 = 2.1254 \pm 0.0018 \text{ (1965BR28).}$$

Alpha particle groups have been observed with $E_p = 7.2$ to 7.5 MeV (1957BR12), 10 MeV (1963GR29), 3.50 to 12.50 MeV (1963BL20), 15.6 and 18.6 MeV (1962MA40). The results are summarized in Table 6.7. Angular distributions of the α_0 and α_1 groups are reported for $E_p = 6$ to 11.5 MeV (1963BL20, 1963TE1B), 5.9 and 7 MeV (1964YA1A) and 15.6 and 18.6 MeV (1962MA40). See also ${}^9\text{Be}$. The 3.56 MeV state is observed to decay by γ -radiation: $E_\gamma =$

Table 6.8: States in ${}^6\text{Li}$ from ${}^7\text{Li}({}^3\text{He}, \alpha){}^6\text{Li}$

(1960AL10)		(1963LI16)		(1963CA02, 1965MA1E)	(1965CO1F)	
E_x (MeV \pm keV)	Γ (keV)	E_x (MeV \pm keV)	Γ (keV)	E_x (MeV \pm keV)	E_x (MeV \pm keV)	Γ (keV)
0		0		0	0	≤ 30
2.19		2.179 ± 8	< 40	2.184 ± 10	2.17 ± 20	≤ 30
3.56 ± 60		3.568 ± 8	< 40	3.564 ± 11	3.55 ± 20	≤ 30
4.3 ± 200				4.59 ± 40	4.3 ± 150	≈ 500
5.35 ± 70	< 100	5.47 ± 40	≈ 600	5.34 ± 30	5.35 ± 20	360 ± 40
(5.6 ± 200)	(2000)					
6.63 ± 800	200					
7.40 ± 100	600					
(8.37 ± 80)	(200)					
(9.3 ± 200)	(≈ 600)					

3.572 ± 0.012 MeV; the internal pair spectrum is consistent with an M1 transition (1954MA26: see also (1960GO13)).

The γ -ray angular distribution and the $(\alpha - \gamma)$ correlation are isotropic at $E_p = 2.56$ MeV, consistent with $J = 0$ for ${}^6\text{Li}^*(3.56)$ (1956ST93). See also (1959LE1B, 1962HA23, 1963ED1A, 1964BA1C) and ${}^{10}\text{B}$.

22. ${}^9\text{Be}(d, {}^5\text{He}){}^6\text{Li}$ $Q_m = -1.056$

Not reported: see (1964BL1C).

23. (a) ${}^{10}\text{B}(\gamma, \alpha){}^6\text{Li}$ $Q_m = -4.461$
 (b) ${}^{10}\text{B}(p, p\alpha){}^6\text{Li}$ $Q_m = -4.461$
 (c) ${}^{10}\text{B}(\alpha, 2\alpha){}^6\text{Li}$ $Q_m = -4.461$
 (d) ${}^{10}\text{B}(n, dn2\alpha)$ $Q_m = -5.933$
 (e) ${}^{10}\text{B}(d, {}^6\text{Li}){}^6\text{Li}$ $Q_m = -2.989$

For reaction (a) see ${}^{10}\text{B}$; for (b) see (1964BA1C); for (c) see (1963ME01); for (d) see (1956FR18); for (e) see (1964GE1A).

$$24. \text{ (a) } {}^{11}\text{B}(\text{d}, {}^7\text{Li}){}^6\text{Li} \quad Q_{\text{m}} = -7.192$$

$$\text{ (b) } {}^{11}\text{B}({}^3\text{He}, {}^8\text{Be}){}^6\text{Li} \quad Q_{\text{m}} = 4.566$$

Reaction (a) has not been reported: see (1964BL1C). For reaction (b), see (1964YO06).

Table 6.9: Energy levels of ${}^6\text{Be}$

E_x (MeV)	J^π	Γ (keV)	Decay	Reactions
g.s.		92 ± 6	α, p	3, 4
(1.5 ± 0.2)				2, 3
$(18 - 24)$	$(-)$		${}^3\text{He}$	1

${}^6\text{Be}$
(Figs. 6 and 7)

GENERAL:

See (1960TA1C, 1961GO1D, 1964GR1J, 1965BO1C, 1965LO1G, 1965JA1C).

Mass of ${}^6\text{Be}$: From the Q -values of the ${}^6\text{Li}(p, n){}^6\text{Be}$ reaction ($Q_0 = -5.069 \pm 0.010$ MeV) and the ${}^6\text{Li}({}^3\text{He}, t){}^6\text{Be}$ reaction ($Q_0 = -4.306 \pm 0.006$ MeV), we adopt ${}^6\text{Be}-{}^6\text{Li} = 4.2872 \pm 0.005$ MeV. The mass excess of ${}^6\text{Be}$ is then 18.376 ± 0.005 MeV, based on ${}^{12}\text{C} \equiv 0$.

- | | | |
|--|----------------|----------------|
| 1. (a) ${}^3\text{He}({}^3\text{He}, 2p){}^4\text{He}$ | $Q_m = 12.860$ | $E_b = 11.487$ |
| (b) ${}^3\text{He}({}^3\text{He}, p){}^5\text{Li}$ | $Q_m = 10.895$ | |
| (c) ${}^3\text{He}({}^3\text{He}, {}^3\text{He}){}^3\text{He}$ | | |

The total cross section for proton production shows a monotonic increase for $E({}^3\text{He}) = 100$ to 800 keV ((1954GO18). The zero-energy cross section factor $S_0 \approx 1100$ keV · b ((1964PA1A). The elastic yield (reaction (c)) does not show any structure for $E({}^3\text{He}) = 3$ to 12 MeV: the yields at $\theta = 31^\circ, 55^\circ, 70^\circ$ and 90° all decrease with increasing energy in a smooth fashion. The angular distributions at $E({}^3\text{He}) = 3.0, 5.9, 7.9, 9.9$ and 11.9 MeV disagree with the theoretical predictions based on the resonating group structure method ((1963TO03). The 90° and 45° cross sections continue to decrease smoothly up to $E({}^3\text{He}) = 18$ MeV, implying that an odd-parity partial wave (probably $l = 3$) is beginning to contribute strongly to the cross section in a region corresponding to ${}^6\text{Be}^*$ ($E_x > 18$ MeV) (((1965BA2B) and private communication). The 65° differential cross section increases by a factor of 2.6 between $E({}^3\text{He}) = 12$ and 18 MeV ((1965BA2B). This sharp upward trend is continued to $E({}^3\text{He}) = 25$ MeV ((1960GA1C). Elastic $\sigma(\theta)$ have also been obtained at $E({}^3\text{He}) = 12, 16, 19, 22$ and 25 MeV ((1965LE02), 29 MeV ((1960BR19) and 30 MeV ((1960MC1E). See also ((1960BR27, 1964AR08, 1965BA1E, 1965TO01).

- | | |
|---|----------------|
| 2. ${}^4\text{He}({}^3\text{He}, n){}^6\text{Be}$ | $Q_m = -9.091$ |
|---|----------------|

At $E(^3\text{He}) = 24.2 \pm 0.3$ MeV, two groups are reported corresponding to the ground state of ^6Be and to an excited state at ≈ 1.5 MeV ((1964BR13).

3. $^6\text{Li}(p, n)^6\text{Be}$

$$Q_m = -5.070$$

$$Q_0 = -5.2 \pm 0.2 \text{ MeV ((1957BO1F);$$

$$Q_0 = -5.05 \pm 0.2 \text{ MeV ((1959AJ81);$$

$$Q_0 = -5.05 \pm 0.05 \text{ MeV ((1964BA16);$$

$$Q_0 = -5.08 \pm 0.04 \text{ MeV ((1963GU07);$$

$$Q_0 = -5.074 \pm 0.013 \text{ MeV ((1965HO1E);$$

$$Q_0 = -5.061 \pm 0.017 \text{ MeV ((1963FR1E).$$

At $E_p = 9$ ((1957BO1F), 10 ((1963GU07) and 10.5 MeV ((1959AJ81), the ground-state neutron group is observed: $\Gamma < 300$ keV ((1957BO1F), $\lesssim 150$ keV ((1959AJ81), 140 ± 40 keV ((1963GU07). The ground-state neutron threshold function indicates participation of both s- and p-wave neutrons ((1965HO1E). The ground-state width is 126 ± 15 keV ((1963FR1E), 95 ± 28 keV ((1965HO1E). ((1959AJ81) report evidence for an excited state at $E_x = 1.5 \pm 0.2$ MeV, $\Gamma < 100$ keV. However ((1965HO1E) does not observe any but the ground-state threshold up to $E_x \approx 3.8$ MeV.

4. $^6\text{Li}(^3\text{He}, t)^6\text{Be}$

$$Q_m = -4.306$$

At $E(^3\text{He}) = 12$ MeV ($\theta = 2^\circ, 10^\circ$), a triton group is observed corresponding to the ground state of ^6Be ($\Gamma = 89 \pm 6$ keV, $Q_0 = -4.306 \pm 0.006$ MeV). No other states of ^6Be are seen with $\Gamma < 1$ MeV up to $E_x = 2.8$ MeV ((1964WH06).

References

(Closed July 01, 1965)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1946SO05 H.S. Sommers, Jr. and R. Sherr, Phys. Rev. 69 (1946) 21
- 1947CA15 J.M. Cassels and R. Latham, Nature 159 (1947) 367
- 1947GU1A Guggenheimer, Heitler and Powell, Proc. Roy. Soc. A190 (1947) 196
- 1948KN13 W.J. Knox, Phys. Rev. 74 (1948) 1192
- 1949BL66 J.M. Blair, G. Freier, E.E. Lampi and W. Sleator, Jr., Phys. Rev. 75 (1949) 1678
- 1949HO24 J.E.R. Holmes, Proc. Phys. Soc. (London) A62 (1949) 293
- 1951AG30 H.M. Agnew, W.T. Leland, H.V. Argo, R.W. Crews, A.H. Hemmendinger, W.E. Scott and R.F. Taschek, Phys. Rev. 84 (1951) 862
- 1951AL26 J.C. Allred, D.K. Froman, A.M. Hudson and L. Rosen, Phys. Rev. 82 (1951) 786
- 1951BU1C Burge, Burrows, Gibson and Rotblat, Proc. Roy. Soc. A210 (1951) 534
- 1951TI06 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A64 (1951) 212
- 1952SH44 R.K. Sheline, Phys. Rev. 87 (1952) 557
- 1952VE1A Vendryes, Thesis, Univ. of Paris (1952)
- 1952WU22 C.S. Wu, B.M. Rustad, V. Perez-Mendez and L. Lidofsky, Phys. Rev. 87 (1952) 1140
- 1953BA04 M.E. Battat and F.L. Ribe, Phys. Rev. 89 (1953) 80
- 1953LA28 T. Lauritsen, T. Huus and S.G. Nilsson, Phys. Rev. 92 (1953) 1501
- 1953MO61 C.D. Moak, Phys. Rev. 92 (1953) 383
- 1954AL35 K.W. Allen, E. Almqvist, J.T. Dewan and T. Pepper, Phys. Rev. 96 (1954) 684
- 1954AL38 E. Almqvist, T.P. Pepper and P. Lorrain, Can. J. Phys. 32 (1954) 621
- 1954FR03 G.M. Frye, Jr., Phys. Rev. 93 (1954) 1086
- 1954FR22 R.G. Freemantle, T. Grotdal, W.M. Gibson, R. McKeague, D.J. Prowse and J. Rotblat, Phil. Mag. 45 (1954) 1090
- 1954GO18 W.M. Good, W.E. Kunz and C.D. Moak, Phys. Rev. 94 (1954) 87
- 1954KL36 R.M. Kline and D.J. Zaffarano, Phys. Rev. 96 (1954) 1620
- 1954MA26 R.J. Mackin, Jr., Phys. Rev. 94 (1954) 648

- 1954SI07 R.M. Sinclair, Phys. Rev. 93 (1954) 1082
- 1954TI16 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A67 (1954) 469
- 1955AJ61 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 27 (1955) 77
- 1955AU1A T. Auerbach and J.B. French, Phys. Rev. 98 (1955) 1276
- 1955FR1C Frank and Gammel, Phys. Rev. 100 (1955) 973A
- 1955GA26 A. Galonsky, R.A. Douglas, W. Haeberli, M.T. McEllistrem and H.T. Richards, Phys. Rev. 98 (1955) 586
- 1955GA74 A. Galonsky and M.T. McEllistrem, Phys. Rev. 98 (1955) 590
- 1955HE1B Henkel, Perry and Smith, Phys. Rev. 99 (1955) 1050
- 1955LA1C Lane and Wilkinson, Phys. Rev. 97 (1955) 1199
- 1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
- 1955RU06 B.M. Rustad and S.L. Ruby, Phys. Rev. 97 (1955) 991
- 1955ST85 J.F. Streib, Phys. Rev. 100 (1955) 1797A
- 1956ED15 R.D. Edge, Aust. J. Phys. 9 (1956) 429
- 1956FR18 G.M. Frye, Jr. and J.H. Gammel, Phys. Rev. 103 (1956) 328
- 1956HA90 J.W. Haffner, Phys. Rev. 103 (1956) 1398
- 1956HO12 D.M. Holm and H.V. Argo, Phys. Rev. 101 (1956) 1772
- 1956HO93 R. Hofstadter, Revs. Mod. Phys. 28 (1956) 214
- 1956LI37 J.G. Likely and F.P. Brady, Phys. Rev. 104 (1956) 118
- 1956MA09 D. Magnac-Valette, Compt. Rend. 242 (1956) 760
- 1956MA1R D. Magnac-Valette and P. Cuer, Physica 22 (1956) 1156A
- 1956MA50 D. Magnac-Valette and P. Cuer, J. Phys. Rad. 17 (1956) 553
- 1956ME1A Meshkov and Ufford, Phys. Rev. 101 (1956) 734
- 1956RE04 J.B. Reynolds and K.G. Standing, Phys. Rev. 101 (1956) 158
- 1956SC40 A. Schwarzschild, B.M. Rustad and C.S. Wu, Bull. Amer. Phys. Soc. 1 (1956) 336, N4
- 1956SO21 I.L. Sokolov, M.M. Sulkovskaia, E.I. Karpushkina and E.A. Albitskaia, Zh. Eksp. Teor. Fiz. 30 (1956) 1007; JETP (Sov. Phys.) 3 (1956) 740
- 1956SO33 I.L. Sokolov, M.M. Sulkovskaia, E.A. Albitskaia and E.I. Karpushkina, Dokl. Akad. Nauk SSSR 111 (1956) 1219; Sov. Phys. Doklady 1 (1957) 773
- 1956ST93 J. Stoltzfus, J. Friichtenicht and E.B. Nelson, Bull. Amer. Phys. Soc. 1 (1956) 329, J2; Oral Report
- 1956VE10 N. Veeraraghavan, Proc. Ind. Acad. Sci. A43 (1956) 319

1956WA29 H.J. Watters, Phys. Rev. 103 (1956) 1763
1957BA10 S.J. Bame, Jr. and W.T. Leland, Phys. Rev. 106 (1957) 1257
1957BO1F Bogdanov, Vlasov, Kalinin, Rybakov and Sidorov, Sov. J. At. Energ. 3 (1957) 987
1957BR12 C.P. Browne and C.K. Bockelman, Phys. Rev. 105 (1957) 1301
1957FR1B French and Fujii, Phys. Rev. 105 (1957) 652
1957HU1C Huper, Z. Naturforsch. A12 (1957) 295
1957JA37 N. Jarmie, J.D. Seagrave et al, LA-2014 (1957)
1957KE1A Kendall and Meyer-Berkhout, Bull. Amer. Phys. Soc. 2 (1957) 378
1957LE1E Levinson and Banerjee, Ann. Phys. 2 (1957) 471
1957TY35 H. Tyren, T.A.J. Maris and P. Hillman, Nuovo Cim. 6 (1957) 1507
1958AL05 R.C. Allen and N. Jarmie, Phys. Rev. 111 (1958) 1129
1958BU17 G.R. Burlinson and R. Hofstadter, Phys. Rev. 112 (1958) 1282
1958CA1B Cazzola and Foglia, Nuovo Cim. 10 (1958) 913
1958CO68 E.D. Commins and P. Kusch, Phys. Rev. Lett. 1 (1958) 208
1958CS88 J. Csikai and A. Szalay, Zh. Eksp. Teor. Fiz. 35 (1958) 1074; JETP (Sov. Phys.) 8 (1959) 749
1958HE46 W.B. Herrmannsfeldt, R.L. Burman, P. Stahelin, J.S. Allen and T.H. Braid, Phys. Rev. Lett. 1 (1958) 61
1958JA06 N. Jarmie and R.C. Allen, Phys. Rev. 111 (1958) 1121
1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1
1958PI1A Pinkston and Brennan, Phys. Rev. 109 (1958) 499
1958TO34 C.H. Townes, Handbuch der Physik 38-1; Ed. S. Flugge (Springer Verlag, Gottingen, 1958) 377
1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10
1958WI15 D.H. Wilkinson, Phys. Rev. 109 (1958) 1603
1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
1959AJ81 F. Ajzenberg-Selove, C.F. Osgood and C.P. Baker, Phys. Rev. 116 (1959) 1521
1959AL10 J.S. Allen, R.L. Burman, W.B. Herrmannsfeldt, P. Stahelm and T.H. Braid, Phys. Rev. 116 (1959) 134
1959AL1E Allen, Revs. Mod. Phys. 31 (1959) 791
1959BA1M Balashov, Zh. Eksp. Teor. Fiz. 36 (1959) 1123; JETP (Sov. Phys.) 9 (1959) 798
1959BE84 E.F. Bennett and D.R. Maxson, Phys. Rev. 116 (1959) 131
1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314

1959CH1B Chen and Hintz, *Congres Int. de Phys. Nucl.*, Paris, 1958 (Dunod, Paris, 1959) 387
 1959CO67 L. Cohen and R.A. Tobin, *Nucl. Phys.* 14 (1959) 243
 1959DA04 F. Daublin, F. Berthold and P. Jensen, *Z. Naturforsch. A*14 (1959) 208
 1959FE1B Feingold, *Phys. Rev.* 114 (1959) 540
 1959GO1H Goldfarb, *Nucl. Phys.* 12 (1959) 494
 1959HA29 E.W. Hamburger, Thesis, Univ. of Pittsburgh (1959)
 1959HE1C Henley and Jacobsohn, *Phys. Rev.* 113 (1959) 225
 1959KU1C Kurepin and Neudachin, *Zh. Eksp. Teor. Fiz.* 36 (1959) 1725; *JETP (Sov. Phys.)* 9 (1959) 1229
 1959LE1B Levin and MacDonald, *Bull. Amer. Phys. Soc.* 4 (1959) 58
 1959ME1D Meyer-Berkhout, *Phys. Rev.* 115 (1959) 1300
 1959ME24 U. Meyer-Berkhout, K.W. Ford and A.E.S. Green, *Ann. Phys.* 8 (1959) 119
 1959PH1B Phillips, *Phys. Rev. Lett.* 3 (1959) 101
 1959PO1C Pondrom, *Phys. Rev. Lett.* 2 (1959) 346
 1959RO62 T.A. Romanowski and W.H. Voelker, *Phys. Rev.* 113 (1959) 886
 1959SK1A Skyrme, *Nucl. Phys.* 9 (1959) 641
 1959SY1A Sytenko and Kharchenko, *Ukr. Fiz. Zh. USSR* 4 (1959) 569; *Phys. Abs.* 13780 (1961)
 1959UB1A Uberall, *Phys. Rev.* 116 (1959) 218
 1959VL24 N.A. Vlasov and A.A. Ogloblin, *Zh. Eksp. Teor. Fiz.* 37 (1959) 54; *JETP (Sov. Phys.)* 10 (1960) 39
 1960AL10 K.W. Allen, E. Almqvist and C.B. Bigham, *Proc. Phys. Soc. (London)* 75 (1960) 913
 1960AN1B Anderson, Knox and Quinton, *Bull. Amer. Phys. Soc.* 5 (1960) 292
 1960AR1A Artemov and Vlasov, *Zh. Eksp. Teor. Fiz.* 39 (1960) 1612; *JETP (Sov. Phys.)* 12 (1961) 1124
 1960BA45 E.B. Bazhanov and L.A. Kulchitskii, *Zh. Eksp. Teor. Fiz.* 38 (1960) 1685; *JETP (Sov. Phys.)* 11 (1960) 1215
 1960BA47 W.C. Barber, F. Berthold, G. Fricke and F.E. Gudden, *Phys. Rev.* 120 (1960) 2081
 1960BR19 D.J. Bredin, J.B.A. England, D. Evans, J.S. McKee, P.V. March, E.M. Mosinger and W.T. Toner, *Proc. Roy. Soc. A*258 (1960) 202
 1960BR27 B.H. Bransden and R.A.H. Hamilton, *Proc. Phys. Soc. (London)* 76 (1960) 987
 1960CH1B Chen and Hintz, *Nucl. Forces and the Few Nucleon Problem*, Pergamon (1960) 683
 1960DU1C Dul-Kova, *Zh. Eksp. Teor. Fiz.* 39 (1960) 1008; *JETP (Sov. Phys.)* 12 (1961) 701
 1960EL09 M. El-Nadi and M. Wafik, *Proc. Phys. Soc. (London)* 76 (1960) 185

- 1960GA08 J.L. Gammel, B.J. Hill and R.M. Thaler, Phys. Rev. 119 (1960) 267
- 1960GA1C Gammel, Brolley, Rosen and Stewart, Proc. Int. Conf. on Nucl. Structure (1960) 215
- 1960GO13 S. Gorodetzky, G. Sutter, F. Scheibling, P. Mennrath, P. Chevallier and R. Armbruster, Compt. Rend. 250 (1960) 3308
- 1960HA14 E.W. Hamburger and J.R. Cameron, Phys. Rev. 117 (1960) 781
- 1960JA1G Jackson, Proc. Phys. Soc. (London) 76 (1960) 949
- 1960KO14 A.P. Komar and E.D. Makhnovskii, Dokl. Akad. Nauk SSSR 135 (1960) 52; Sov. Phys. Doklady 5 (1960) 1229
- 1960KO1C Kotin, Rev. Mex. Fisica 9 (1960) 73
- 1960KO1D Kopaleishvili, Vashakidze, Mamasakhlishov and Chilashvili, Zh. Eksp. Teor. Fiz. 38 (1960) 1758; JETP (Sov. Phys.) 11 (1960) 1268
- 1960MA32 R.D. Macfarlane and J.B. French, Revs. Mod. Phys. 32 (1960) 567
- 1960MC1E McKee, Nucl. Forces and the Few Nucleon Problem, Pergamon (1960) 609
- 1960PH1A Phillips and Tombrello, Nucl. Phys. 19 (1960) 555
- 1960PR06 D.G. Proctor and W.H. Voelker, Phys. Rev. 118 (1960) 217
- 1960SA28 Y. Sakamoto and T. Takemiya, Prog. Theor. Phys. 23 (1960) 172
- 1960TA1C Talmi and Unna, Ann. Revs. Nucl. Sci. 10 (1960) 353
- 1960VA1D Vashakidze, Kopaleishvili and Chilashvili, Zh. Eksp. Teor. Fiz. 39 (1960) 393; JETP (Sov. Phys.) 12 (1961) 278
- 1960WA1F Wackman, Thesis, Univ. of Pittsburgh (1960)
- 1960YO06 L.G. Youn, G.M. Osetinskii, N. Sodnom, A.M. Govorov, I.V. Sizov and V.I. Salatskii, Zh. Eksp. Teor. Fiz. 39 (1960) 225; JETP (Sov. Phys.) 12 (1961) 163
- 1961AH1A Ahrens, J. Inorg. Nucl. Chem. 16 (1961) 368
- 1961BA1E Balashov, Neudachin and Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 170; Bull. Acad. Sci. USSR Phys. 25 (1961) 165
- 1961BA40 J.F. Barry, R. Batchelor and B.E.F. Macefield, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press, New York, 1961) 543
- 1961CH09 R. Chiba, H.E. Conzett, H. Morinaga, N. Mutsuro, K. Shoda and M. Kimura, J. Phys. Soc. Jpn. 16 (1961) 1077
- 1961CL09 A.B. Clegg, K.J. Foley, G.L. Salmon and R.E. Segel, Proc. Phys. Soc. (London) 78 (1961) 681
- 1961GA09 J.P. Garron, J.C. Jacmart, M. Riou, C. Ruhla, J. Teillac, C. Caverzasio and K. Strauch, Phys. Rev. Lett. 7 (1961) 261
- 1961GO18 A.M. Govorov, L.H. Youn, G.M. Osetinskii, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teor. Fiz. 41 (1961) 703; JETP (Sov. Phys.) 14 (1962) 508

1961GO1D Golsansky, Nucl. Phys. 27 (1961) 648

1961HO01 H.D. Holmgren and L.M. Cameron, Bull. Amer. Phys. Soc. 6 (1961) 36, MA2

1961HO23 H.D. Holmgren and L.M. Cameron, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 537

1961KO1G Kopaleishvili, Vashakidze, Mamasakhlisov and Chilashvili, Nucl. Phys. 23 (1961) 430

1961KO1H Kolos, Acta Phys. Pol. 20 (1961) 175

1961MA29 M. Masuda, J. Phys. Soc. Jpn. 16 (1961) 1801

1961OG1A Ogloblin and Chuen, Instr. Expt. Tech. 5 (1962) 859; Pribory Tekhnika Eksp. 5 (1961) 37

1961PA1A Patel and Bhavsar, Current Sci. (India) 30 (1961) 51

1961PO01 L.G. Pondrom and J.W. Daughtry, Phys. Rev. 121 (1961) 1192

1961PU1A Pugh and Riley, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 195

1961RI03 B.W. Ridley, Nucl. Phys. 25 (1961) 483

1961RY01 B.V. Rybakov, V.A. Sidorov and N.A. Vlasov, Nucl. Phys. 23 (1961) 491

1961SH1B Shah and Pandya, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 393

1961SL06 R.J. Slobodrian, Bol. Acad. Nac. Cienc. (Cordoba, Rep. Arg.) 42 (1961) 75; Phys. Abs. 67, 1989, Phys. Abs. 22164 (1964)

1961TA05 Y.C. Tang, K. Wildermuth and L.D. Pearlstein, Phys. Rev. 123 (1961) 548

1961VA1G Vashakidze, Kopaleishvili and Chilashvili, Nucl. Phys. 23 (1961) 694

1961WO05 E.A. Wolicki and A.R. Knudson, Bull. Amer. Phys. Soc. 6 (1961) 415, B2

1961YU01 T. Yuasa, J. Phys. Rad. 22 (1961) 169

1962BA1D Barber, Ann. Rev. Nucl. Sci. 12 (1962) 1

1962BE1J Berggren and Jacob, Phys. Lett. 1 (1962) 258

1962BE1K Berggren, Brown and Jacob, Phys. Lett. 1 (1962) 88

1962BI14 J.K. Bienlein and F. Pleasonton, Nucl. Phys. 37 (1962) 529; Erratum Nucl. Phys. 40 (1963) 694

1962BO17 E.C. Booth and K.A. Wright, Nucl. Phys. 35 (1962) 472

1962CH1B Chizhov et al, Nucl. Phys. 34 (1962) 562

1962CO1D Cohen, Nucl. Phys. 33 (1962) 52

1962CR09 C.L. Critchfield, LA-2714 (1962)

1962DI1A Dietrich, Phys. Lett. 2 (1962) 139

1962DI1B Dietrich, Z. Phys. 167 (1962) 563
1962EL1D Elton and Jackson, Nucl. Phys. 35 (1962) 209
1962FO08 W.F. Ford, Nucl. Phys. 37 (1962) 119
1962GA09 J.P. Garron, J.C. Jacmart, M. Riou, C. Ruhla, J. Teillac and K. Strauch, Nucl. Phys. 37 (1962) 126
1962GA23 J.P. Garron, Ann. Phys. (Paris) 7 (1962) 301
1962GO1P Gottschalk, Unpublished Thesis, Harvard Univ. (1962)
1962GO1Q Govorov, Ka-Yeng, Osetinsky, Salatsky and Sizov, Zh. Eksp. Teor. Fiz. 42 (1962) 383; JETP (Sov. Phys.) 15 (1962) 266
1962HA23 S.S. Hanna, Bull. Amer. Phys. Soc. 7 (1962) 470, LA4
1962IN02 D.R. Inglis, Nucl. Phys. 30 (1962) 1
1962IN1A Inglis, Revs. Mod. Phys. 34 (1962) 165
1962IN1C Inglis, Phys. Rev. 126 (1962) 1789
1962JA06 D.F. Jackson, Proc. Phys. Soc. (London) 79 (1962) 1041
1962JA1B Jackson, Nucl. Phys. 35 (1962) 194
1962LE12 H.W. Lefevre, R.R. Borchers and C.H. Poppe, Phys. Rev. 128 (1962) 1328
1962MA38 S. Malmskog and J. Konijn, Nucl. Phys. 38 (1962) 196
1962MA40 D.R. Maxson, Phys. Rev. 128 (1962) 1321
1962ME1C Meshkov, Nucl. Phys. 35 (1962) 485
1962NA1B Narasimhan, Shah and Pandya, Nucl. Phys. 33 (1962) 529
1962RO20 L. Rosen and W.T. Leland, Phys. Rev. Lett. 8 (1962) 379
1962RU04 C. Ruhla, M. Riou, J.P. Garron, J.C. Jacmart and L. Massonnet, Phys. Lett. 2 (1962) 44
1962SA1F Sakamoto, Phys. Lett. 1 (1962) 256; Nuovo Cim. 26 (1962) 461
1962SC15 R. Schlecht, D. McColm and I. Maleh, Bull. Amer. Phys. Soc. 7 (1962) 604, G1
1962SE02 F.D. Seward, Phys. Rev. 125 (1962) 335
1962SL04 R.J. Slobodrian, Phys. Rev. 126 (1962) 1059
1962ST19 L. Stewart, J.E. Brolley, Jr. and L. Rosen, Phys. Rev. 128 (1962) 707
1962ST1E Strnad, Phys. Rev. 125 (1962) 1639
1962TE1D Temmer, Bull. Amer. Phys. Soc. 7 (1962) 569
1962TI01 G. Tibell, O. Sundberg and U. Miklavzic, Phys. Lett. 1 (1962) 172
1962VO1C Volkov and Kulchitskii, Zh. Eksp. Teor. Fiz. 42 (1962) 108; JETP (Sov. Phys.) 15 (1962) 77

1962VO1D Volkov, Kulikov and Chizhov, Zh. Eksp. Teor. Fiz. 42 (1962) 61; JETP (Sov. Phys.) 15 (1962) 42

1962WA1E Wackman and Austern, Nucl. Phys. 30 (1962) 529

1962WO07 C. Wong, J.D. Anderson and J.W. McClure, Nucl. Phys. 33 (1962) 680

1963AL18 D.E. Alburger, Phys. Rev. 132 (1963) 328

1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527

1963BA1V Batchelor and Towle, AWRE NR 6/63 (1963)

1963BA50 R. Batchelor and J.H. Towle, Nucl. Phys. 47 (1963) 385

1963BE25 M. Bernheim and G.R. Bishop, Phys. Lett. 5 (1963) 270

1963BE42 T. Berggren and G. Jacob, Nucl. Phys. 47 (1963) 481

1963BE53 M. Bernheim and G.R. Bishop, J. Phys. 24 (1963) 970

1963BI10 G.R. Bishop and M. Bernheim, Phys. Lett. 5 (1963) 140

1963BL20 H.R. Blieden, G.M. Temmer and K.L. Warsh, Nucl. Phys. 49 (1963) 209

1963BO1K Bodmer and Ali, Nucl. Phys. 40 (1963) 463

1963BU1C Bunakov, Phys. Lett. 7 (1963) 14

1963CA02 J.A. Careaga and M. Mazari, Bull. Amer. Phys. Soc. 8 (1963) 124, U5, and Private Communication (1963)

1963CA1H Carlson, Pleasonton and Johnson, Phys. Rev. 129 (1963) 2220

1963CO15 S. Costa, S. Ferroni, V. Wataghin and R. Malvano, Phys. Lett. 4 (1963) 308

1963CO1D Costa et al, Phys. Lett. 6 (1963) 226

1963DA04 J.F. Dawson and J.D. Walecka, Ann. Phys. 22 (1963) 133

1963DE01 D.W. Devins and H.H. Forster, Bull. Amer. Phys. Soc. 8 (1963) 12, BA13

1963ED1A Edwards, Padua (1963) 469

1963EL1C Elton, Padua (1963) 1093

1963EL1D El-Nadi, Rabie and Sherif, Nucl. Phys. 48 (1963) 569

1963ER02 H.J. Erramuspe and R.J. Slobodrian, Nucl. Phys. 49 (1963) 65

1963FR1E Freeman, West and Montague, AERE Prog. Rept. PR/NP 3, 4 (1963), and Private Communication (1963)

1963GO04 J. Goldemberg and Y. Torizuka, Phys. Rev. 129 (1963) 312

1963GR29 D.E. Groce and W. Whaling, Phys. Rev. 132 (1963) 2614

1963GU07 M. Gulyamov, B.V. Rybakov and V.A. Sidorov, Zh. Eksp. Teor. Fiz. 44 (1963) 1829; JETP (Sov. Phys.) 17 (1963) 1230

1963HA1H Hansteen and Kanestrom, Nucl. Phys. 46 (1963) 303
 1963HA49 W.D. Harrison, Bull. Amer. Phys. Soc. 8 (1963) 597, F2
 1963HW01 C.F. Hwang, G. Clausnitzer, D.H. Nordby, S. Suwa and J.H. Williams, Phys. Rev. 131 (1963) 2602
 1963IN1A Inopin, Nucl. Phys. 42 (1963) 666
 1963JA1C Jackson and Elton, Nucl. Phys. 43 (1963) 136
 1963JO04 C.H. Johnson, F. Pleasonton and T.A. Carlson, Nucl. Phys. 41 (1963) 167
 1963JO07 A. Johansson and Y. Sakamoto, Nucl. Phys. 42 (1963) 625
 1963JO15 C.H. Johnson, F. Pleasonton and T.A. Carlson, Phys. Rev. 132 (1963) 1149
 1963KI1D Kim, Nucl. Phys. 49 (1963) 651
 1963KL1A Klapisch, Rameau, Epherre and Gradsztajn, J. Phys. 24 (1963) 839
 1963KN1C Knudson and Wolicki, Padua (1963) 981
 1963KO04 D. Kohler and S.M. Austin, Bull. Amer. Phys. Soc. 8 (1963) 290, A2
 1963KU03 D. Kurath, Phys. Rev. 130 (1963) 1525
 1963KU18 B. Kuhn and B. Schlenk, Nucl. Phys. 48 (1963) 353
 1963KU1B Kunz, Can. J. Phys. 41 (1963) 2187
 1963KU1H Kuhn and Schlenk. Joint. Inst. Nucl. Res., Lab. Neutron Phys. USSR Rept. No. P1197 (1963)
 1963LI16 J. Linck, I. Nicolas-Linck, R. Bilwes and D. Magnac-Valette, J. Phys. 24 (1963) 983
 1963MA1P Magnac-Valette, Seltz, Bilwes and Spyns, Padua (1963) 1088A
 1963ME01 J.B. Mead, L.B. Geesaman and H.B. Knowles, Bull. Amer. Phys. Soc. 8 (1963) 292, A15
 1963MO1F Morpurgo, Nucl. Spectroscopy; Ed. Racah (Academic Press, 1962)
 1963OG1A Ogloblin, Nucl. Phys. 47 (1963) 408
 1963OL1B Ollerhead, Chasman and Bromley, Padua (1963) 984
 1963RI1B Riou, Padua (1963) 18
 1963RO23 R.E. Rothe, L. Jacobson and J. Jobst, Bull. Amer. Phys. Soc. 8 (1963) 537, S6
 1963RU05 C. Ruhla, M. Riou, M. Gusakow, J.C. Jacmart, M. Liu and L. Valentin, Phys. Lett. 6 (1963) 282
 1963SA1F Sakamoto, Nuovo Cim. 28 (1963) 206
 1963SC1J Schmid, Tang and Wildermuth, Phys. Lett. 7 (1963) 263
 1963SC30 R.G. Schlecht, UCRL-11047 (1963)
 1963SK02 S.J. Skorka, R. Hubner, T.W. Retz-Schmidt and H. Wahl, Nucl. Phys. 47 (1963) 417

1963SM03 D.B. Smith, N. Jarmie and A.M. Lockett, Phys. Rev. 129 (1963) 785
 1963TA1D Takemiya, Prrog. Theor. Phys. 30 (1963) 191
 1963TE1B Temmer, Pauda (1963) 1013
 1963TO03 T.A. Tombrello and A.D. Bacher, Phys. Rev. 130 (1963) 1108
 1963VA14 W.T.H. Van Oers and K.W. Brockman, Jr., Nucl. Phys. 44 (1963) 546
 1963VI06 J.B. Vise and B.M. Rustad, Phys. Rev. 132 (1963) 2573
 1963VL1A Vlasov, Zh. Eksp. Teor. Fiz. 45 (1963) 160; JETP (Sov. Phys.) 18 (1964) 160
 1963WA1H Wackman and Austern, Bull. Amer. Phys. Soc. 8 (1963) 56
 1964AR08 K.P. Artjomov, V.J. Chuev, V.Z. Goldberg, A.A. Ogloblin, V.P. Rudakov and I.N. Serikov, Phys. Lett. 12 (1964) 53
 1964AR1C Armstrong, Gammel, Rosen and Frye, Nucl. Phys. 52 (1964) 505
 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209
 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
 1964BA1P Balashov and Boyarkina, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1964) 359
 1964BA2B Bahcall and Wolf, Astrophys. J. 139 (1964) 622
 1964BA2C Bazhanov, Komar and Kulikov, Zh. Eksp. Teor. Fiz. 46 (1964) 1497; JETP (Sov. Phys.) 19 (1964) 1014
 1964BI04 G.R. Bishop and M. Bernheim, Phys. Lett. 8 (1964) 128
 1964BL1C Blieden, Phys. Lett. 9 (1964) 176
 1964BR13 H.C. Bryant, J.G. Beery, E.R. Flynn and W.T. Leland, Nucl. Phys. 53 (1964) 97
 1964BR1R Broek and Yntema, Phys. Rev. 135 (1964) B678
 1964DE1F Devins, Forster, Bunch and Kim, Phys. Lett. 9 (1964) 35
 1964DE1G Delorme, Congres Int. de Phys. Nucl., Paris (1964)
 1964ER1B Eramjian, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1181
 1964GA01 G.T. Garvey and J.C. Hiebert, Bull. Amer. Phys. Soc. 9 (1964) 44, DC9
 1964GA11 N. Gangas, S. Kossionides, R. Rigopoulos and M.L. Ahmad, Phys. Lett. 12 (1964) 233
 1964GE1A Gemmell, Erskine and Schiffer, Phys. Rev. 134 (1964) B110
 1964GO15 J. Goldemberg, W.C. Barber, F.H. Lewis, Jr. and J.D. Walecka, Phys. Rev. 134 (1964) B1022
 1964GR1A Green and Donahue, Phys. Rev. 135 (1964) B701
 1964GR1J Green, Nucl. Phys. 54 (1964) 505
 1964GR39 G. Gregoire and P.C. Macq, Phys. Lett. 8 (1964) 328

1964HA05 E.W. Hamburger, Nucl. Phys. 50 (1964) 66
 1964HU1A Hunchen, Kropf and Waffler, Nucl. Phys. 58 (1964) 477
 1964JA03 J.C. Jacmart, J.P. Garron, M. Riou and C. Ruhla, Phys. Lett. 8 (1964) 269
 1964JI1A Jibuti, Kopaleishvili and Mamasakhlisov, Nucl. Phys. 52 (1964) 345
 1964KO1F Komar and Makhnovskii, Dokl. Akad. Nauk SSSR 156 (1964) 774
 1964KU1G Kurath, Phys. Rev. 134 (1964) B1025
 1964LI1D Lim and McCarthy, Phys. Rev. 133 (1964) B1006
 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
 1964MA1X Makhnovskii, Zh. Eksp. Teor. Fiz. 46 (1964) 1136; JETP (Sov. Phys.) 19 (1964) 769
 1964MA1Y Marty, Geoffrion, Rolland, Tatischeff and Willis, Congres Int. de Phys. Nucl., Paris (1964)
 1964MC1E McIntyre and Haeberli, Bull. Amer. Phys. Soc. 9 (1964) 390
 1964MC1F McIntyre and Haeberli, Bull. Amer. Phys. Soc. 9 (1964) 627
 1964MI16 F.C. Michel, Phys. Rev. 133 (1964) B329
 1964NE1E Neudachin, Shevchenko and Yudin, Phys. Lett. 10 (1964) 180
 1964OH01 G.G. Ohlsen and P.G. Young, Phys. Rev. 136 (1964) B1632
 1964OH1B Ohlsen and Young, Nucl. Phys. 52 (1964) 134
 1964OL1A Ollerhead, Chasman and Bromley, Phys. Rev. 134 (1964) B74
 1964PA1A Parker, Bahcall and Fowler, Astrophys. J. 139 (1964) 602
 1964PE1C Perkin, Nucl. Phys. 60 (1964) 561
 1964RO1D Rothe, Private Communication (1964)
 1964SA1F Sandulescu and Dumitrescu, Phys. Lett. 11 (1964) 420
 1964SA1H Sakamoto, Phys. Rev. 134 (1964) B1211
 1964SE07 L.S. Senhouse, Jr. and T.A. Tombrello, Nucl. Phys. 57 (1964) 624
 1964SE1F Seiler, Darden, McIntyre and Weitakmp, Nucl. Phys. 53 (1964) 65
 1964SE1G Senhouse, Thesis, Cal Tech (1964)
 1964SH05 J.H. Shafer, Phys. Rev. 133 (1964) B920
 1964SH07 T.H. Short and N.M. Hintz, Bull. Amer. Phys. Soc. 9 (1964) 391, BA16
 1964SH27 A.K. Shardanov, V.G. Shevchenko and B.A. Yurev, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 60; Bull. Acad. Sci. USSR Phys. Ser. 28 (1965) 60
 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
 1964TA02 B. Tatischeff, N. Marty, X. De Bouard, J.G. Fox, B. Geoffrion and C. Rolland, Phys. Lett. 8 (1964) 54

1964TI02 G. Tibell, O. Sundberg and P.U. Renberg, Ark. Fys. 25 (1964) 433
 1964WA1E Walecka, Congres Int. de Phys. Nucl., Paris (1964)
 1964WH01 L.Wharton, L.P.Gold and W. Klemperer, Phys. Rev. 133 (1964) B270
 1964WH06 W. Whaling, Bull. Amer. Phys. Soc. 9 (1964) 627, E7
 1964YA1A Yanabu et al, J. Phys. Soc. Jpn. 19 (1964) 1818
 1964YO06 F.C. Young, P.D. Forsyth, M.L. Roush, W.F. Hornyak and J.B. Marion, Phys. Lett. 13 (1964) 50
 1965AJ01 F. Ajzenberg-Selove, J.W. Watson and R. Middleton, Phys. Rev. 139 (1965) B592
 1965BA1E Bacher and Tombrello, Bull. Amer. Phys. Soc. 10 (1965) 693
 1965BA2B Bacher and Tombrello, Bull. Amer. Phys. Soc. 10 (1965) 423
 1965BA2C Bahcall, Nucl. Phys. (1965)
 1965BE1D Berman, Bramblett, Caldwell, Harvey and Fultz, Phys. Rev. Lett. 15 (1965) 727
 1965BE1H Beregi, Zelenskaja, Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 513
 1965BE1R Bernas, Epherre, Gradsztajn, Klapisch and Yiou, Phys. Lett. 15 (1965) 147
 1965BO1C Bodmer and Murphy, Nucl. Phys. 73 (1965) 664
 1965BR28 C.P. Browne, W.E. Dorenbusch and F.H. O'Donnell, Nucl. Phys. 72 (1965) 194
 1965CO1F Cocke, Private Communication (1965)
 1965DA1H Da Providencia and Shakin, Nucl. Phys. 65 (1965) 54
 1965DE1Q Devins, Forster and Scott, Revs. Mod. Phys. 37 (1965) 396
 1965EL1B Elton and Lodhi, Nucl. Phys. 66 (1965) 209
 1965FO07 P.D. Forsyth and R.R. Perry, Nucl. Phys. 67 (1965) 517
 1965FU1G Fuller and Cohen, Appendix I, Nucl. Data Sheets 6-5 (1965)
 1965GO1L Gogny and Jean, Compt. Rend. 260 (1965) 510
 1965GR1Q Griffith and Roman, Phys. Lett. 14 (1965) 42
 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81
 1965HA1L Hansteen and Wittern, Phys. Rev. 137 (1965) B524
 1965HO1E Honsaker, Unpublished Thesis, Cal Tech (1965)
 1965JA04 D.F. Jackson and J. Mahalanabis, Nucl. Phys. 64 (1965) 97
 1965JA1C Janecke, Nucl. Phys. 61 (1965) 383
 1965JA1J Jackson and Berggren, Nucl. Phys. 62 (1965) 353
 1965JA1L Jackson and Elton, Proc. Phys. Soc. (London) 85 (1965) 659
 1965JA1M Jackson, Revs. Mod. Phys. 37 (1965) 393

1965JO09 G. Johnson, H. Langevin-Joliot and P. Roussel, J. Phys. (Paris) 26 (1965) 161
1965LE02 W.T. LeLand, J.E. Brolley, Jr. and L. Rosen, Bull. Amer. Phys. Soc. 10 (1965) 51,
DC7
1965LO1G Lovitch, Nucl. Phys. 62 (1965) 653
1965LO1H Lodder and Jonker, Phys. Lett. 15 (1965) 245
1965LO1J Lodhi, Bull. Amer. Phys. Soc. 10 (1965) 583
1965MA1E Mazari, Private Communication (1965)
1965MA1N Malvano and Ricco, Nuovo Cim. 35 (1965) 484
1965MC1F McCarthy, Revs. Mod. Phys. 37 (1965) 388
1965ME05 F. Merchez, N. Van Sen, V. Regis and R. Bouchez, Compt. Rend. 260 (1965) 3922
1965MU1A Murphy and Rosati, Nucl. Phys. 63 (1965) 625
1965NE1C Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 25
1965PA03 P. Paul, S.L. Blatt and D. Kohler, Phys. Rev. 137 (1965) B499
1965RA1D Rand, Frosch and Yearian, Bull. Amer. Phys. Soc. 10 (1965) 542
1965RA1E Rahman, Proc. Phys. Soc. (London) 85 (1965) 653
1965RI1A Riou, Revs. Mod. Phys. 37 (1965) 375
1965SA1J Sakamoto, Nuovo Cim. 37 (1965) 774
1965SE1E Sergeev, Izv. Akad. Nauk 29 (1965) 105
1965SH1F Sherman, Morrison and Stewart, Bull. Amer. Phys. Soc. 10 (1965) 541
1965TO01 T.A. Tombrello and A.D. Bacher, Phys. Lett. 17 (1965) 37
1965TY1A Tyren, Kullander, Sundberg, Ramachandran, Isacson and Berggren (1965), Un-
known source
1965WE1E Weidenmuller, Nucl. Phys. (1965)
1965WI1G Wittern and Hansteen, Bull. Amer. Phys. Soc. 10 (1965) 424