

Energy Levels of Light Nuclei $A = 7$

T. Lauritsen ^a and F. Ajzenberg-Selove ^b

^a *California Institute of Technology, Pasadena, California*

^b *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics* 78 (1966), p. 1. This version of $A = 7$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed July 01, 1965)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 7$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^7\text{He}\$](#) , [\${}^7\text{Li}\$](#) , [\${}^7\text{Be}\$](#) , [\${}^7\text{B}\$](#)

B. Tables of Recommended Level Energies:

[Table 7.1](#): Energy levels of ${}^7\text{Li}$

[Table 7.9](#): Energy levels of ${}^7\text{Be}$

C. [References](#)

D. Figures: [\${}^7\text{Li}\$](#) , [\${}^7\text{Be}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

⁷He

(Not illustrated)

From the known location of the ⁷Li $T = \frac{3}{2}$ state, the mass excess of ⁷He is calculated as 26.03 ± 0.15 MeV (calculation of the Coulomb energy difference based on the pair ⁶He-⁶Li* ($T = 1$)); ⁷He is then unstable with respect to decay into ⁶He + n by 0.36 MeV (1965DE08). See also (1953PE1A, 1960BA51, 1960GO1B, 1961YA04, 1962EL1E, 1963NE02). Some reactions leading to ⁷He are ⁷Li(n, p)⁷He, ⁷Li(t, ³He)⁷He and ⁹Be(n, ³He)⁷He.

⁷Li

(Figs. 8 and 10)

GENERAL: See (1957HU1C, 1959BA1M, 1959BA1D, 1959BR1E, 1959FE1B, 1959MA1F, 1959MA1G, 1960KU1B, 1960PE11, 1960PH1A, 1960SH1A, 1960TA1C, 1961BA1D, 1961BA1E, 1961BL1C, 1961CL10, 1961KH03, 1961TA05, 1961TO04, 1962CL1F, 1962CR09, 1962IN02, 1963CH08, 1963CL1B, 1963KL1A, 1963SC30, 1964BE1N, 1964GR1J, 1964MA1G, 1964NE1E, 1964OL1A, 1964SA1F, 1965BE1R, 1965FA1B, 1965JA1L, 1965NE1C, 1965PR1F).

Ground state :

$$Q = -45 \pm 5 \text{ mb (1961KA29, 1963VA19, 1964WH01);}$$
$$\mu = +3.2564 \text{ nm (1965FU1G).}$$

1. ⁴He(t, γ)⁷Li $Q_m = 2.467$

Excitation functions and angular distributions have been studied in the range $E_\alpha = 0.5$ to 1.9 MeV by (1961GR27), $E_\alpha = 0.5$ to 1.3 MeV by (1959HO03). The cross section rises smoothly, as expected for a direct capture process; at $E_\alpha = 1.32$ MeV, $\sigma = 3.58 \pm 0.06 \mu\text{b}$, and the corresponding reduced cross section factor $S = 0.064 \pm 0.016 \text{ keV} \cdot \text{b}$ (1961GR27) in good agreement with the value 0.05 calculated by (1961CH1C). Cross sections of (1961GR27) are 2 to 2.5 times higher than those of (1959HO03).

The branching ratio of γ_0/γ_1 (to ⁷Li(0) and ⁷Li*(0.48)) is $1/(0.4 \pm 0.05)$, essentially independent of energy and angle. At $E_\alpha = 0.56$ MeV, the angular distribution of γ 's is isotropic ($\pm 6\%$), while some preference for forward emission appears at $E_\alpha = 1.6$ MeV. Assuming $\theta_\alpha^2 = 1.25$ and 1.05 for the ground and 0.48 MeV states of ⁷Li, and taking into account only E1 capture from s- and d-waves, (1963TO06) have calculated the total cross section from 0 to 7 MeV and the γ -ray intensity ratios: the calculations are in excellent agreement with the data of (1961GR27). See also (1961TO04).

Table 7.1: Energy levels of ${}^7\text{Li}$

E_x (MeV \pm keV)	$J^\pi; T$	Γ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$		stable	1, 3, 4, 9, 10, 22, 26, 28, 29
0.4779 ± 0.3	$\frac{1}{2}^-; \frac{1}{2}$	6.2 ± 0.3 meV	γ	1, 3, 4, 9, 10, 13, 15, 17, 20, 21, 22, 26, 27, 28, 29
4.629 ± 8	$\leq \frac{7}{2}^-; \frac{1}{2}$	93 ± 8	α, t	9, 15, 16, 17, 20, 21, 26, 28
(6.56 ± 120)	$(\frac{5}{2}^-); \frac{1}{2}$	≈ 1000		15, 16, 17
7.475 ± 4	$\frac{5}{2}^-; \frac{1}{2}$	89 ± 7	n, α, t	5, 8, 9, 14, 15, 16, 17, 20, 26, 28
(9.7)			γ, n, t	14, 16, 17
11.13 ± 0.05	$(\frac{3}{2}^-; \frac{3}{2})$	268 ± 30	γ, n, t	14, 15, 26
(12.5)			γ, n, p	14, 15
(14)			γ, n, p, t	14, 15
(15.5) ^a			γ, n, p	14
(16.8) ^a			γ, n	14
(17.3)			γ, n	14
(19.6) ^a			γ, t	14

^a Giant resonance.

2. ${}^4\text{He}(t, t){}^4\text{He}$

$$E_b = 2.467$$

Angular distributions have been measured for $E_t = 1.2$ to 2.2 MeV (1956HE16), 1.7 MeV (1958AL05), $E_\alpha = 11$ to 28 MeV (1960BR1J). See also (1961TO04).

3. ${}^4\text{He}(\alpha, p){}^7\text{Li}$

$$Q_m = -17.347$$

At $E_\alpha = 38.5$ MeV, two groups of protons are observed leading to the ground and 0.48 MeV states of ${}^7\text{Li}$ (1958BU38).

4. ${}^6\text{Li}(n, \gamma){}^7\text{Li}$

$$Q_m = 7.253$$

Two γ -rays with $E_\gamma = 7.26 \pm 0.03$ and 6.78 ± 0.05 MeV, and relative intensities 10 and 7.5 ± 2.0 , corresponding to transitions to the first two states of ${}^7\text{Li}$ are observed (1957BA18). The total radiative capture cross section is 45 ± 10 mb (1964ST25). See also (1961JA19, 1961TO04).

5. ${}^6\text{Li}(n, n){}^6\text{Li}$

$$E_b = 7.253$$

The total cross section has been measured for $E_n = 4$ eV to 29 MeV (1958HU18, 1960HU08, 1960PE25, 1963BA50, 1964AR25, 1964ST25). A pronounced resonance occurs at $E_n = 262$ keV with a peak cross section of 11.2 b (1960HU08). The elastic contribution is 7.2 b (1961LA1A). No other clearly defined resonance is observed, although the cross section exhibits a broad maximum at $E_n \approx 5$ MeV (1954JO17, 1960HU08). The coherent scattering length (thermal, bound) is 1.8 fm (1964ST25).

Angular distributions are tabulated by (1963GO1M): see also (1961LA1A, 1962BA1W, 1963BA50). All observations near the 0.262 MeV resonance are consistent with p-wave formation of a $J^\pi = \frac{5}{2}^-$ level (${}^7\text{Li}^*(7.48)$). Table 7.2 gives the resonance parameters compared with those for ${}^7\text{Be}^*(7.18)$. These states are believed to have a ${}^4\text{P}_{5/2}$ character, in agreement with their large θ_n^2 and θ_p^2 (see ${}^7\text{Be}$ and (1956ME1A, 1957MA57)).

The scattering of polarized neutrons on ${}^6\text{Li}$ has been studied for $E_n = 0.19$ to 0.42 MeV (1961DA04), 0.14 to 0.66 MeV (1962EL01), and 0.2 to 2.0 MeV (1964LA19). The data agree reasonably well with polarizations calculated from the resonance parameters of Table 7.2 with a background of s-wave potential scattering in which channel spin $J_c = \frac{1}{2}$ dominates (1961DA04, 1962EL01, 1964LA19).

See also (1960KO1C, 1960LA1C, 1962MA1R, 1963AL1J, 1964PE1E).

6. (a) ${}^6\text{Li}(n, p){}^6\text{He}$

$$Q_m = -2.727$$

$$E_b = 7.253$$

(b) ${}^6\text{Li}(n, d){}^5\text{He}$

$$Q_m = -2.430$$

The cross section for reaction (a) at $E_n = 14$ MeV is 6.7 mb (1953BA04, 1954FR03). See also (1963BA56).

For reaction (b) see reaction 8 and (1954FR03, 1956RI34).

7. ${}^6\text{Li}(n, 2n){}^5\text{Li}$

$$Q_m = -5.662$$

The cross section is 33 ± 15 mb at $E_n = 10.2$ MeV, 70 ± 6 mb at $E_n = 14.1$ MeV (1963AS01).

8. ${}^6\text{Li}(n, \alpha){}^3\text{H}$

$$Q_m = 4.785$$

$$E_b = 7.253$$

Table 7.2: Resonance parameters for 7.2 – 7.5 MeV levels in ${}^7\text{Li}$ and ${}^7\text{Be}$

Reaction	${}^6\text{Li} + \text{n}$	${}^6\text{Li} + \text{p}$
References	a	b
E_r (keV, lab)	262	1840
$\Gamma(E_\gamma)$ (keV, c.m.)	154	836
E_λ (keV above g.s.)	7700	7580
$\Gamma_{\text{n, p}}(E_r)$ (keV, c.m.)	118	798
radius (n, p) in fm	3.94	4.08
$\gamma_{\text{n, p}}^2$ (MeV · fm)	4.85	5.02
$\theta_{\text{n, p}}^2$	0.26	0.28
$\Gamma_\alpha(E_r)$ (keV, c.m.)	36	38
radius (α) in fm	4.39	4.39
γ_α^2 (MeV · fm)	0.101	0.101
θ_α^2	0.012	0.012

^a (1959GA08: see (1963MC09)).

^b (1963MC09).

Excitation functions and angular distributions are summarized in (1958HU18, 1960HU08, 1963GO1M, 1964ST25). Recent measurements are reported for $E_n < 30$ keV (1961BE24) and for $E_n = 0.19$ MeV (1963BA2A), 1 to 600 keV (1965SC07), 0.1 to 0.3 MeV (1959PA02), 9 to 340 keV (1959BA46), 1.2 to 8 MeV (1959MU25), 2.0 to 2.65 MeV (1959PU75), 8 to 14 MeV (1963MA61), 14.4 MeV (1964VA19). See also (1960PE24, 1961BE24, 1962BE1P, 1962CA1F, 1963AL1J, 1963CH20, 1963WA1K).

The isotropic thermal cross section is 949 b: in the eV-keV range, the cross section falls off somewhat more slowly than $1/v$ (1958HU18); for $E_n = 9$ to 90 keV, $\sigma = 3.96E^{-0.367}$ b (E in keV) (1959BA46). The failure to follow the $1/v$ law may reflect a broad s-wave resonance (1959BA46, 1959GA08, 1960LA1C, 1961BE24). Results of (1965SC07) are consistent with strict $1/v$ dependence for $E_n < 20$ keV. (1956MA83) have analyzed cross section data for $E_n = 20$ to 565 keV and find no need for an s-wave resonance. The s-wave background contribution is mainly (75%) in the $J_c = \frac{1}{2}^+$ channel.

A resonance occurs at $E_n = 258$ keV, with $\sigma_{\text{max}} = 2.75$ b (1959BA46), 2.80 ± 0.22 b (1959GA08): see also (1965MA1Y). The resonance is formed by p-waves, $J^\pi = \frac{5}{2}^-$, and has a large neutron width and a small α -width (Table 7.2) (1959GA08). Above the resonance the cross section decreases monotonically to $E_n = 18$ MeV, except for a slight bump near $E_n = 1.6$ to 2.1 MeV (1959GA08, 1959MU25). A careful search in the range $E_n = 2.0$ to 2.65 MeV

Table 7.3: ${}^7\text{Li}$ levels from ${}^6\text{Li}(d, p){}^7\text{Li}$

E_x ^a (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	θ_n^2 ^b
g.s.		0.048
0.477 ± 2 ^c		0.063
4.630 ± 9 (6.54 ± 20) ^d	93 ± 8 ^e	(isotropic)
7.464 ± 10	91 ± 8 ^e	0.040

^a (1952GE07, 1955KH35, 1957BR97, 1961JA23) and C.P. Browne, private communication.

^b (1960MA32): data of (1955LE24, 1960HA14); PWBA. See also (1962RO23, 1963ME09).

^c See (1955AJ61).

^d (1955KH35; see, however, (1960HA14)).

^e (1957BR97).

($E_x = 9$ to 9.55 MeV) revealed no evidence for a level reported in ${}^6\text{Li}(\gamma, n)$ (1959PU75). See also (1964GI1F).

In the range $E_n = 8$ to 14 MeV, the reaction ${}^6\text{Li}(n, dn){}^4\text{He}$ shows a large cross section, reaching 0.6 b at $E_n \approx 6$ MeV (1962RO12).

9. ${}^6\text{Li}(d, p){}^7\text{Li}$

$$Q_m = 5.028$$

Proton spectra and angular distributions have been studied at $E_d = 1.5$ MeV (1964RI1E), 1.6 MeV (1960AN02), 3.4 to 5.2 MeV (1963ME09), 7 to 7.5 MeV (1957BR97), 8.0 MeV (1953HO48), 14 to 15 MeV (1955LE24, 1959HA29, 1960HA14). Groups corresponding to the ground state and to the states at 0.48, 4.6 and 7.5 MeV have been identified: see Table 7.3. The first two and the last show stripping patterns with $l_n = 1$, while the angular distribution of p_2 ($E_x = 4.6$ MeV) is isotropic. It is noted that stripping to this level is forbidden if it has the character ${}^{22}\text{F}$. Ratio of observed θ_n^2 are consistent with assignments ${}^{22}\text{P}$ to the g.s. and 0.48 and ${}^{24}\text{P}$ to the 7.5 MeV state (1960HA14, 1960MA32).

The angular correlation between the protons and the 0.48 MeV γ -rays is isotropic (see (1955AJ61)) indicating $J = \frac{1}{2}$ for ${}^7\text{Li}^*$. The mean lifetime is reported to be $(7.7 \pm 0.8) \times 10^{-2}$ psec (1956BU83; see also ${}^9\text{Be}(d, \alpha){}^7\text{Li}$) see Table 7.4. See also (1959BO1C, 1959HE1C, 1960NE1C, 1961HA1F, 1963BI27). The polarization of p_0 and p_1 protons has been observed at $E_d = 1.6$ MeV by (1961VA03). The circular polarization of the 0.48 MeV γ -rays has been studied by (1964SC1J). See also (1964PA16, 1965HE1B).

10. ${}^6\text{Li}(t, d){}^7\text{Li}$ $Q_m = 0.995$

The reaction has been observed to the ground and 0.48 MeV states at $E_t = 0.24$ MeV (1954AL35) and 1.5 and 1.9 MeV (1961HO21). See also (1955CU17, 1961HO1F, 1963BA2B).

11. ${}^6\text{Li}(\alpha, {}^3\text{He}){}^7\text{Li}$ $Q_m = -13.325$

Not reported.

12. (a) ${}^6\text{Li}({}^6\text{Li}, p^4\text{He}){}^7\text{Li}$ $Q_m = 3.556$

(b) ${}^6\text{Li}({}^9\text{Be}, {}^8\text{Be}){}^7\text{Li}$ $Q_m = 5.587$

${}^9\text{Be}({}^6\text{Li}, {}^8\text{Be}){}^7\text{Li}$

See (1960MA1H, 1961LE1K, 1962BE16, 1962MC12, 1963BA1Q, 1963CO35, 1963NO02, 1965SA1L), ${}^{12}\text{C}$ and ${}^{15}\text{N}$.

13. ${}^7\text{Li}(\gamma, \gamma){}^7\text{Li}$

Resonance scattering and absorption by ${}^7\text{Li}^*(0.48)$ has been studied by a number of observers: the derived mean lifetimes are listed in Table 7.4. See also (1960VA1G).

14. (a) ${}^7\text{Li}(\gamma, n){}^6\text{Li}$ $Q_m = -7.253$

(b) ${}^7\text{Li}(\gamma, p){}^6\text{He}$ $Q_m = -9.980$

(c) ${}^7\text{Li}(\gamma, d){}^5\text{He}$ $Q_m = -9.683$

(d) ${}^7\text{Li}(\gamma, t){}^4\text{He}$ $Q_m = -2.467$

Reports on the structure of the (γ, n) cross section differ widely. According to (1954GO1A, 1958RY77: $E(\text{brems.}) = 24$ MeV), a broad maximum appears at $E_\gamma = 16.8$ MeV, with $\sigma_{\text{max}} = 2.3$ mb and a width $\Gamma = 9.3$ MeV. Fine structure corresponding to levels at $E_x = 9.66 \pm 0.04, 10.8, 12.4, 14.0$ and 17.5 MeV is also reported (1954GO39, 1958RY77). Up to $E(\text{brems.}) = 19$ MeV, the work of (1964AL08) confirms the gross structure of (1958RY77): additional levels at $E_x = 13.6, 15.3$ and 16.5 MeV are found: see Table 7.5. (1955HE51) and (1959RO62) find, on the other hand, a weak rise at 8 MeV, a pronounced narrow peak at 14 MeV, followed by a deep minimum at 17 MeV and a sharp rise thereafter. It is suggested by (1964AL08) that the discrepancies are to be traced to different spectral sensitivities in the neutron detectors. Evidence is found for enhanced

Table 7.4: Mean life of ${}^7\text{Li}^*(0.48)$

τ_m (sec $\times 10^{13}$)	Reaction	Reference
0.77 ± 0.08	${}^6\text{Li}(d, p)$	1956BU83
0.75 ± 0.25	${}^{10}\text{B}(n, \alpha)$	1949EL07
1.1 ± 0.3	${}^7\text{Li}(\gamma, \gamma)$	1958BE10
1.15 ± 0.14	ibid	1959SW63
$\geq 1.09 \pm 0.07$	ibid	1959SW63
1.4 ± 0.7	ibid	1960BO23
1.0 ± 0.5	ibid	1960BO23
1.25 ± 0.06	ibid	1962MO17
1.48 ± 0.35	ibid	1963MO02
0.93 ± 0.13	ibid	1964BO22
1.07 ± 0.05		mean

emission of slow neutrons from a level near $E_x = 17.25$ MeV ([1964AL08](#)). With $E(\text{brems.}) = 57$ MeV, ([1960FA06](#)) find a broad maximum near 19 MeV, $\sigma = 3.2 \pm 0.8$ mb, and a slow tailing off to 57 MeV: see also ([1963CO1D](#)). Reported integral cross sections, $\int \sigma dE$, are 0 – 24 MeV: 33 MeV · mb ([1954GO1A](#)), 18 MeV · mb ([1958RY77](#)); 0 – 25 MeV: 39 MeV · mb ([1960FA06](#)); 0 – 50 MeV: 93 MeV · mb ([1960FA06](#)). ([1964GR40](#)) report cross sections for $E_\gamma = 7.4$ to 10.8 MeV: a value $\Gamma_\gamma = 0.9 \pm 0.4$ eV is obtained for ${}^7\text{Li}^*(7.48)$. See also ([1960KU1C](#)).

The (γ, p) cross section determined by ${}^6\text{He}$ production shows a splitting of the giant resonance into 2 components, at 15.5 and 19.6 MeV: smaller resonances are reported at 12.7, 25.4, 32.0 and 38.0 MeV ([1963CL03](#): see also ([1954RU27](#))). Peaks at 12.5, 13.5 and 14.3 MeV are reported by ([1962SH24](#)), while ([1963KU25](#)) report 6 peaks in the range $E_x = 11$ to 24.5 MeV. According to ([1954TI16](#), [1962GR08](#)) only a single maximum occurs, at ≈ 15.6 MeV, with a width of ≈ 4 MeV. Polarization of photoprotons at $E_\gamma = 335$ MeV is small and consistent with zero ([1962LI13](#)). See also ([1963FU1D](#), [1963KI1C](#)).

The ratio $\sigma_{\gamma p}/\sigma_{\gamma d}$ has been investigated by ([1962BE1N](#), [1962CH26](#), [1962VO1C](#)). See also ([1963KU16](#)).

Peaks reported in the (γ, t) cross section are listed in Table 7.5. See also ([1961KO1J](#), [1963WA07](#)). See also ([1964BI03](#), [1964MA2B](#)).

15. ${}^7\text{Li}(e, e){}^7\text{Li}$

Elastic scattering has been studied at $E_e = 41.5$ MeV ([1963GO04](#): $\theta = 180^\circ$) and 187 MeV ([1955ST85](#)). At the lower energy the magnetic scattering is consistent with that expected from a

Table 7.5: Levels of ${}^7\text{Li}(\gamma, n){}^6\text{Li}$, ${}^7\text{Li}(\gamma, p){}^6\text{He}$ and ${}^7\text{Li}(\gamma, t){}^4\text{He}$

(γ, n)		(γ, p)			(γ, t)						
(1954GO39)	(1964AL08)	(1962SH24)	(1963CL03)	(1963KU25)	(1953TI02)	(1953ST27)	(1954ST89)	(1955MI55)	(1960MI02)	(1961SH11, 1961SH21)	(1963KU16) ^d
						5.25	4.7 5.5 6.8			5.3 6.6 (7.5) 8.3	
9.6 ^a					9.3	(9.25)		7.6 8.6 9.6	7.8 8.9 9.8		
10.8				11.0				11.7			
12.4		12.5	12.7								
	13.6	13.5						(13.5)	13.5		
14.0		14.3		14.0							14.1
	15.3		15.5 ^b								
16.8 ^b	16.5			16.0	(16.7)			16.2			16.2
17.5	17.3 ^c			17.9							18.0
			19.6 ^b								19.6
				21.4	(21.5)						(21.5)
					(23.5)						(23.5)
			25.4	24.5							(25.3)
			32.0								
			38.0								

^a 9.66 ± 0.04 MeV (1958RY77).

^b Giant resonance.

^c Resonance for $E_{\text{n}} < 6$ MeV.

^d And (1963KU25); ± 0.2 MeV, except 25.3 ± 0.3 MeV.

point dipole (1963GO04). The higher energy results yield an r.m.s. charge radius of about 2.1 fm (1957HO1E).

In inelastic scattering studies at $E_e = 41.5$ MeV, $\theta = 180^\circ$, weak broad peaks, ascribed to M1 transfer, are reported at $Q = -6.9, -10.5, -14.0$ MeV (1963BA19: see Table 7.6). At $E_e = 102$ to 177 MeV, excitation of states at 0.48, 4.63 ± 0.05 , (5.7 ± 0.1) , 6.8 ± 0.1 and 7.5 ± 0.08 MeV is reported. Transitions to the $E_x = 0.48$ MeV state show both longitudinal and transverse E2 contributions (see (1963WI1B)), while the others are mainly longitudinal E2: reported $B(E2)$ are listed in Table 7.6 (1963BE26, 1963BE53, 1964BI03). See also (1959ME1D, 1962BA1D, 1964GU1B, 1964MA2C, 1965CH11).

16. ${}^7\text{Li}(n, n){}^7\text{Li}$

Elastic angular distributions have been measured for $E_n = 1.5$ to 7.5 MeV (1963BA1V, 1963BA50) and 14 MeV (1964AL1N) and compared with optical model scattering.

At $E_n = 14$ MeV, evidence is reported for states at $E_x = 4.6 \pm 0.25, \approx 6.5, 7.5 \pm 0.25$, and (9.25) MeV (1954AL24). For $E_n = 1.5$ to 7.5 MeV, the excitation of the 4.6 MeV state shows no evidence of direct interaction (1963BA50). A DWBA analysis of the distribution of this group at $E_n = 14$ MeV has been carried out by (1960PE1A). See also (1960HE1F, 1962WO07).

A tabulation of various partial cross sections is given by (1963BA50) and (1964AL1N). See also (1963GL1F, 1963OP1A, 1964VA19).

17. ${}^7\text{Li}(p, p){}^7\text{Li}$

Elastic scattering and polarization has been studied at $E_p = 40$ MeV (1959CH1B, 1960CH1B), 150 MeV (1962NE12), 155 MeV (1964TA02), 156 MeV (1964JA03) and 160 to 180 MeV (1959JO43, 1960JO14, 1961JO18, 1962RO1F). Analysis in terms of optical parameters is reported by (1961JO18, 1964SA1L).

Inelastic proton groups have been observed corresponding to the excited states of ${}^7\text{Li}$ at 0.48, 4.63, 6.56 and 7.48 MeV: see (1952AJ38). At $E_p = 185$ MeV, proton groups are observed to these states, as well as to states at 5.5 ± 0.3 MeV ($\Gamma \approx 0.4$ MeV) and 9.6 ± 0.2 MeV. The width of the 6.5 MeV state is reported to be ≈ 1 MeV. Angular distributions have been measured for the 4.6 and 6.5 MeV states (1965HA17). A check of the isotropy of the 477 keV radiation ($J = \frac{1}{2} \rightarrow \frac{3}{2}$) yields an upper limit $F^2 < 10^{-4}$ for the intensity of a parity non-conserving part of the wave function (1958WI38). Relative intensities of the $Q = -4.6, -6.6$ and -7.5 MeV groups, observed at $E_p = 17.5$ MeV, are consistent with the assignments ${}^{22}\text{F}_{7/2}, {}^{22}\text{F}_{5/2}, {}^{24}\text{P}_{5/2}$, respectively (1957LE1E, 1957MA04). At $E_p = 150$ MeV, the $E_x = 4.6$ MeV state is strongly excited compared to the $E_x = 0.48$ and 6.5 MeV states, consistent with the assumed rotational character of these levels (1962NE12, 1964JA03). See also (1960HA21, 1960HE1F, 1962RU04, 1963CH08, 1963RI1B, 1964ST15, 1965WE1E).

Table 7.6: Electromagnetic transitions in ${}^7\text{Li}$ from ${}^7\text{Li}(e, e')$ and Coulomb excitation

E_x (MeV)	J^π	$B(E2\uparrow)$ (fm^4)	$(2J + 1)\Gamma_\gamma$	References
0.48 ^a	$\frac{1}{2}^-$	6.8 ± 1 7.3 ± 1.5 7.6 ± 1.1		(1963BE26) (1960ST17) (1962RI09)
4.63 (5.7) ^b	$(\frac{7}{2}^-)$	15.5 ± 0.8 (4.1 ± 2)		(1963BE26, 1963BE53) (1963BE26, 1963BE53)
6.8	$(\frac{5}{2}^-)$	12.5 ± 1.2 ^c		(1963BE26, 1963BE53)
7.5 (10.5) (12.5) (14.0)	$\frac{5}{2}^-$	$2.5^{+0.5}_{-1.0}$	3.6 ± 2 ^d 38 ± 10 62 ± 25	(1963BA19, 1963BE26) (1963BA19) (1963BA19) (1963BA19)

^a See also Table 7.4.

^b Possibly due to ${}^7\text{Li} \rightarrow \alpha + t$ (1964BI03).

^c $B(M1\uparrow) = 6.9 \times 10^{-3} \text{ fm}^2$ (1963BE26).

^d $\Gamma_\gamma = 0.9 \pm 0.4 \text{ eV}$ (1964GR40).

18. (a) ${}^7\text{Li}(p, 2p){}^6\text{He}$ $Q_m = -9.980$
 (b) ${}^7\text{Li}(p, pd){}^5\text{He}$ $Q_m = -9.683$
 (c) ${}^7\text{Li}(p, \alpha){}^4\text{He}$ $Q_m = 17.347$

The summed proton energy spectrum, observed at $E_p = 155$ to 450 MeV, shows two peaks, with $Q \approx -11.6$ and $Q \approx -25.4$ MeV, corresponding to formation of ${}^6\text{He}(\text{g.s.} + 1.80)$ and an excited state near 15 MeV (see ${}^6\text{He}$). Angular distributions indicate that the higher energy peak corresponds to the removal of a p-proton while the lower results from removal of an s-proton (1958MA1B, 1959MA1F, 1960HI10, 1961GA09, 1962BE1J, 1962BE1K, 1962DI1A, 1962GA09, 1962GO1P, 1962IN02, 1962IN1A, 1962ST1E, 1962ST1F, 1962TI01, 1963BE1A, 1963BE42, 1963EL1C, 1963JO07, 1963RI1B, 1963TA1D, 1964BA1C, 1964LI1D, 1964TI02, 1965RI1A, 1966TY01).

For reaction (b), see (1962RU04, 1963SH1A, 1964SA1H, 1965JA1L). For reaction (c), see (1962MA40, 1965ZH1A) and ${}^8\text{Be}$.

19. (a) ${}^7\text{Li}(p, d){}^6\text{Li}$ $Q_m = -5.028$

Table 7.7: ${}^7\text{Li}(p, d){}^6\text{Li}$, ${}^7\text{Li}(d, t){}^6\text{Li}$ reduced widths ^a

E_x in ${}^6\text{Li}$ (MeV)	J^π	l	$\theta_{(p, d)}^2$ ^b	$\theta_{(p, d)}^2$ ^c	$\theta_{(d, t)}^2$ ^d	$\theta_{(d, t)}^2$ ^e
0	1^+	1	0.053	(0.053)	(0.048)	0.11
2.18	3^+	1	0.036	0.027	(0.036)	0.061
3.56	0^+	1		0.028	0.032	0.083
4.57	2^+					
5.36	(2^+)	1			0.025	0.017

^a PWBA.

^b (1956RE04, 1960MA32).

^c (1959BE84, 1960MA32).

^d (1955LE24, 1960HA14, 1960MA32).

^e (1959VL24); see also (1962SL04).

$$\begin{aligned} \text{(b) } {}^7\text{Li}(d, t){}^6\text{Li} & \quad Q_m = -0.995 \\ \text{(c) } {}^7\text{Li}(d, {}^3\text{He}){}^6\text{He} & \quad Q_m = -4.486 \\ \text{(d) } {}^7\text{Li}(t, \alpha){}^6\text{He} & \quad Q_m = 9.834 \end{aligned}$$

Reduced widths derived from PWBA analysis of pickup reactions (a) and (b) are listed in Table 7.7. For reaction (c), $\theta^2 = 0.025$ and 0.008 for ${}^6\text{He}(0)$ and ${}^6\text{He}^*(1.80)$ (1957FR1B, 1960MA32).

20. ${}^7\text{Li}(d, d'){}^7\text{Li}^*$

Inelastic deuteron groups are observed corresponding to the ground and 0.48, 4.6 and 7.5 MeV states. At $E_d = 14.8$ MeV, the $Q = -7.46$ MeV group is only weakly excited (1960HA14). At $E_d = 28$ MeV, the $Q = -4.6$ MeV group is strong: the angular distribution is strongly forward. A fit with PWBA requires $l = 0 + 2$ (1962SL02). See also (1952AJ38, 1959AJ76).

Elastic scattering at 28 MeV has been analyzed in the black disc approximation, yielding $R = 4.1$ fm (1962SL02). See also ${}^9\text{Be}$ and (1958EL45, 1958RO49, 1959HA29, 1959SI1A, 1961SL06, 1965JU1A).

21. (a) ${}^7\text{Li}(\alpha, \alpha'){}^7\text{Li}^*$

$$\text{(b) } {}^7\text{Li}({}^7\text{Li}, {}^7\text{Li}'){}^7\text{Li}^*$$

$$\text{(c) } {}^7\text{Li}({}^{20}\text{Ne}, {}^{20}\text{Ne}'){}^7\text{Li}^*$$

Table 7.8: Branching fraction in ${}^7\text{Be}(\epsilon){}^7\text{Li}$

$\gamma/\text{disintegration (\%)}$	References
10.7 ± 2	(1949WI13)
11.8 ± 1.2	(1949TU06, 1949TU1B)
12.3 ± 0.6	(1949TU1B, 1951DI12)
10.32 ± 0.16	(1962TA11)

Inelastic alpha groups are observed corresponding to the 0.48 and 4.6 MeV states: see (1955AJ61). At $E_\alpha = 13.2, 31.8$ and 48 MeV, the angular distributions of the $Q = -4.6$ MeV group show a prominent peak in the forward hemisphere (1956CO61, 1957SI36, 1960MA15). Analysis by PWBA yields $R = 5.6$ fm, $l = 2$ (1960MA15). See also (1962MA59). For reaction (b), see (1960BL1B).

Coulomb excitation of ${}^7\text{Li}^*(0.48)$ in reaction (c) has been studied at $E({}^{20}\text{Ne}) = 9$ to 11 (1960ST17), 15.4 (1962RI09) and 16 MeV (1961AN07). The observed intensity leads to a value of the reduced transition matrix element $B(E2) = 7.6 e^2\text{fm}^4$ (1962RI09), $7.3 e^2\text{fm}^4$ (1960ST17) corresponding to a partial half life $\tau = 1.5$ nsec. Comparison with the quoted $\tau_{1/2}$ of the state, 0.08 psec, yields $\Gamma(E2)/\Gamma(M1) = 5 \times 10^{-5}$ (1960ST17): see Tables 7.4 and 7.6. See also (1963BE1R).

22. ${}^7\text{Be}(\epsilon){}^7\text{Li}$ $Q_m = 0.862$

The decay proceeds to the ground and 0.48 MeV states. Reported branching ratios are listed in Table 7.8. The γ -ray energy is 477.8 ± 0.3 keV (see (1955AJ61, 1957DU37)). The weighted mean value of the half life is 53.37 ± 0.11 days (1949SE20, 1953KR16, 1956BO36, 1957WR37); $ft = 2.00 \times 10^3$ for the ground state transition and 3.45×10^3 for the excited state (1965BA2C). Both transitions are super-allowed (1954MA1D, 1956CH1B). The internal conversion coefficient of the 0.48 MeV γ is $5.8 - 16.6 \times 10^{-7}$ indicating an E2/M1 ratio between 0 and 0.8 (1959LE30). Calculations of electron capture and nuclear matrix elements support the conclusion that ${}^7\text{Li}$ is close to LS coupling (1962BA1X). See also (1959BL1C, 1963CH08, 1963KI1D).

23. ${}^7\text{Be}(n, p){}^7\text{Li}$ $Q_m = 1.644$

See ${}^8\text{Be}$.

24. (a) ${}^9\text{Be}(\gamma, d){}^7\text{Li}$ $Q_m = -16.693$

(b) ${}^9\text{Be}(\gamma, np){}^7\text{Li}$ $Q_m = -18.917$

See (1955AJ61) and ${}^9\text{Be}$.

$$25. {}^9\text{Be}(n, t){}^7\text{Li} \quad Q_m = -10.435$$

See (1957SC12, 1957VA12) and ${}^{10}\text{Be}$.

$$26. {}^9\text{Be}(p, {}^3\text{He}){}^7\text{Li} \quad Q_m = -11.199$$

At $E_p = 43.7$ MeV, ${}^3\text{He}$ groups are observed corresponding to the ${}^7\text{Li}$ levels at 0, 0.48, 4.6 and 7.5 MeV and to a new state at 11.13 ± 0.05 MeV, with $\Gamma = 268 \pm 30$ keV. From the similarity of the angular distribution and cross section to that in the (p, t) mirror reaction, it is concluded that the level has $J^\pi = \frac{3}{2}^-$; $T = \frac{3}{2}$ (1965DE08).

$$27. \text{(a) } {}^9\text{Be}(p, \text{pd}){}^7\text{Li} \quad Q_m = -16.693$$
$$\text{(b) } {}^{10}\text{B}(p, p{}^3\text{He}){}^7\text{Li} \quad Q_m = -17.786$$

See (1963SH1A, 1964BA1C, 1964BA1P, 1964SH1C).

$$28. {}^9\text{Be}(d, \alpha){}^7\text{Li} \quad Q_m = 7.154$$

A number of α -groups have been observed with deuteron energies up to 27.5 MeV. These correspond to levels at 480 ± 2 keV (1948BU31, 1953CO02, 1961JA23), 4.62 ± 0.02 MeV (1953GE01: see also (1964MA04)), and 7.5 MeV (1951GO47, 1964MA04). Angular distributions have been studied for the ground and $E_x = 0.48$ MeV states at $E_d = 0.4$ to 2.4 MeV (1962BI11), 10 MeV (1962WE04) and 13.6 MeV (1962IV1A); those to the 4.6 and 7.5 MeV states at $E_d = 27.5$ MeV (1964MA04). The widths of the 4.6 and 7.5 MeV states are, respectively, 93 ± 25 and 80 ± 20 keV (1966HA09). The upper limit to the intensity of an α -particle group to a state at $E_x \approx 5.5$ MeV is 5% of the intensity of the group to ${}^7\text{Li}^*(7.48)$ (1966HA09: $E_d = 11.1$ MeV). See also (1964YA1A).

The (α, γ) angular correlation has been observed for $E_d = 0.40$ and 0.84 MeV, (1953UE01, 1954CO17). There is no significant departure from isotropy, in agreement with $J = \frac{1}{2}$ for the 0.48 MeV level. The mean life of this state is reported to be $(7.7 \pm 0.8) \times 10^{-2}$ psec (1956BU83): see Table 7.4.

See also (1959AJ76) and ${}^{11}\text{B}$.

29. $^{10}\text{B}(\text{n}, \alpha)^7\text{Li}$

$$Q_m = 2.792$$

With thermal neutrons, two groups of α -particles are observed, corresponding to $^7\text{Li}^*(0, 0.48)$; the fraction of transitions leading to the ground state is about 6%: see ^{11}B . The γ -ray energy is 478.5 ± 1.5 keV (1948EL1A), 478 ± 4 keV (1956DA23); the mean life is $(7.5 \pm 2.5) \times 10^{-2}$ psec (1949EL07): see Table 7.4.

See also (1952AJ38, 1955AJ61, 1959AJ76, 1960AN14, 1962LA07, 1963DE1F).

30. (a) $^{11}\text{B}(\alpha, 2\alpha)^7\text{Li}$

$$Q_m = -8.664$$

(b) $^{11}\text{B}(\gamma, \alpha)^7\text{Li}$

(c) $^{11}\text{B}(\text{p}, \text{p}\alpha)^7\text{Li}$

For reaction (a) see (1963ME01). For reaction (b) see ^{11}B . For reaction (c) see (1964BA1C).

31. $^{18}\text{O}(\text{d}, ^{13}\text{C})^7\text{Li}$

$$Q_m = -5.678$$

See (1963DR1B).

⁷Be
(Figs. 9 and 10)

GENERAL: See (1957FR1B, 1960PH1A, 1960SH1A, 1960TA1C, 1961KU1C, 1961TA05, 1961TO04, 1962GL1A, 1962IN02, 1964AR22, 1964BA2A, 1964GR1J, 1964HO31, 1964LI1F, 1964MO1K, 1964NE1H, 1964PA1K, 1964PH1A, 1964RA1B, 1964SA1F, 1964ST1B).

1. ⁷Be(ϵ)⁷Li $Q_m = 0.862$

The decay is complex: see ⁷Li.

2. ⁴He(³He, γ)⁷Be $Q_m = 1.587$

In the range $E_\alpha = 0.42$ to 5.80 MeV the cross section rises from 0.02 to 4 μ b. The branching ratio γ_0 (to g.s.)/ γ_1 (to 0.4 MeV state) remains at 73/27 for $E_\alpha = 1.1$ to 3.2 MeV (1963PA12). The zero-energy intercept of the cross section factor $S = 0.47 \pm 0.05$ keV \cdot b (1963PA12). See also (1958BA59, 1959HO03, 1961FA02, 1964PA1A). A direct capture calculation, assuming an $\alpha + ^3\text{He}$ model with phase shifts obtained from ⁴He(³He, ³He), gives good agreement with the excitation function of (1963PA12) with $R = 2.8$ fm, $\theta_\alpha^2 = 1.25$ and 1.05 for ⁷Be(0) and ⁷Be*(0.43), respectively. The capture proceeds mainly by E1, with both s- and d-waves contributing above $E_\alpha = 1$ MeV (1963TO06). See also (1961CH1C, 1961TO04, 1962HE1C). The bearing of this reaction on ⁴He production in stars is discussed by (1958FO1A, 1964PA1A).

Table 7.9: Energy levels of ⁷Be

E_x (MeV \pm keV)	$J^\pi; T$	Γ (keV)	Decay	Reactions
g.s.	$\leq \frac{3^-}{2}; \frac{1}{2}$	$\tau_{1/2} = 53.37 \pm 0.11$ d	ϵ	1, 3, 5, 10, 14, 15, 17, 19
0.431 \pm 1	$\frac{1^-}{2}; \frac{1}{2}$	$\tau_m = 0.27 \pm 0.10$ ps	γ	5, 10, 14, 15, 17, 19
4.55 \pm 20	$\frac{7^-}{2}; \frac{1}{2}$	100	³ He, α	3, 17, 19
6.51 \pm 40	$\frac{5^-}{2}; \frac{1}{2}$	1200	³ He, α	3, 10, 19
7.185 \pm 20	$(\frac{5^-}{2}); \frac{1}{2}$	836	p, ³ He, α	3, 7, 9, 10, 19
(9.2 \pm 0.5)		broad	p, ³ He	3
9.9	$(\frac{3^-}{2}; \frac{1}{2})$	≈ 1800	p, p ₁ , p ₂	7
10.79 \pm 40	$(\frac{3^-}{2}; \frac{3}{2})$	298 \pm 25		17
(14.6 \pm 300)				19

3. (a) ${}^4\text{He}({}^3\text{He}, {}^3\text{He}){}^4\text{He}$

$$E_b = 1.587$$

(b) ${}^4\text{He}({}^3\text{He}, p){}^6\text{Li}$

$$Q_m = -4.021$$

Elastic scattering studies have been reported for $E({}^3\text{He}) = 3$ to 5.5 MeV (1958MI92), 2.5 to 5.7 MeV (1964BA09), 4 to 12 MeV (1963TO04), 8 to 18 MeV (1964SP04), 29 to 30 MeV (1960BR19, 1960MC1E), $E_\alpha = 11$ to 28 MeV (1960BR1J), 28 to 41 MeV (1961CH09). Two resonances are reported in the f-wave phase shifts: at $E_x = 4.55$ MeV (${}^2F_{7/2}$) and $E_x = 6.51$ MeV (${}^2F_{5/2}$); see Table 7.10. In the range $E({}^3\text{He}) = 2.5$ to 12 MeV, the s-wave and d-wave phase shifts reflect hard-sphere scattering with $R = 2.8$ fm, although some systematic deviation seems to appear in the s-waves for $E({}^3\text{He}) > 5$ MeV (1963TO04, 1964BA09). The $p_{3/2}$ phase shift exhibits a contribution from the ground state, with $\theta_\alpha^2(\text{g.s.}) = 0.15$, but the $p_{1/2}$ phase shift cannot be adequately accounted for by ${}^7\text{Be}^*(0.43)$ (1961TO04, 1964BA09). The ${}^4P_{5/2}$ level ($E_x = 7.18$ MeV) seems to have no influence on the scattering: $\theta_p^2 < 0.02$ (1963TO04). See also (1964PH1A). A broad resonance has been observed at $E_x = 9.2 \pm 0.5$ MeV. It is not clear whether it can be identified with the ($\frac{3}{2}^-$) state at 9.9 MeV (1964SP04).

At the higher energies, distinct minima are observed in the angular distribution at $\theta = 45^\circ$, 100° and 140° . No evidence of sharp resonances is observed for $E_x > 13$ MeV (1960BR19, 1960BR1J, 1961CH09). Optical model calculations are reported by (1963SQ1A), resonating group calculations by (1963TA10). See also (1963SC1M).

Polarizations have been calculated from the observed phase shifts by (1959PH37, 1963TO04, 1964BA09). Measurement of scattering of α -particles from a polarized ${}^3\text{He}$ target at $E_\alpha = 6.53$ and 7.33 MeV confirms the expected reversal of polarization between these two energies (1962PH1B).

Reaction (b) has been studied for $E({}^3\text{He}) = 8$ to 12 MeV by (1963TO04) and at 29 MeV by (1960BR19, 1960MC1E). A peak appears in the excitation function at $E({}^3\text{He}) = 9.8$ MeV, corresponding to the ${}^4P_{5/2}$ level at $E_x = 7.18$ MeV (1963TO04). The resonance corresponding to ${}^7\text{Be}^*(9.2)$, observed in the elastic scattering, is strongly present in the yield of ${}^6\text{Li}^*(2.18)$ protons but not in the yield of ground state protons (1964SP04). See also (1962TE1D).

4. ${}^4\text{He}(\alpha, n){}^7\text{Be}$

$$Q_m = -18.991$$

See ${}^8\text{Be}$.

5. ${}^6\text{Li}(p, \gamma){}^7\text{Be}$

$$Q_m = 5.608$$

Gamma transitions are observed to the ground and 0.43 MeV states. The yield shows no evidence of resonance for $E_p = 0.4$ to 1.0 MeV and the branching ratio remains approximately constant at $(62 \pm 5)\%$ to the ground state, 38% to the 0.43 MeV state, $< 4\%$ to the 4.6 MeV state (1955BA59, 1956WA03). The 90° differential cross section at 750 keV is $0.02 \mu\text{b/sr}$ (1956WA03).

Table 7.10: ${}^7\text{Be}$ levels from ${}^4\text{He}({}^3\text{He}, {}^3\text{He}){}^4\text{He}$ and ${}^6\text{Li}(p, p){}^6\text{Li}$

E_x (MeV)	J^π	Term	R (fm)	θ_α^2	θ_p^2	Refs.
0	$\frac{3}{2}^-$	${}^2\text{P}_{\frac{3}{2}}$	3.8	0.15		(1964BA09)
0.43	$\frac{1}{2}^-$	${}^2\text{P}_{\frac{1}{2}}$				
4.54 ± 0.02	$\frac{7}{2}^-$	${}^2\text{F}_{\frac{7}{2}}$	4.4	0.36		(1958MI92)
4.57 ± 0.04				0.38		(1964BA09)
6.51 ± 0.04	$\frac{5}{2}^-$	${}^2\text{F}_{\frac{5}{2}}$	4.4	0.48	< 0.02	(1963TO04)
7.18	$\frac{5}{2}^-$	${}^4\text{P}_{\frac{5}{2}}$	4.1	0.012	0.28	(1963MC09)
9.2 ± 0.5						(1964SP04)

The angular distributions of γ_0 and γ_1 are the same at $E_p = 0.75$ MeV, $W(\theta) = 1 + (1.05 \pm 0.15) \cos^2 \theta$. Neither s- nor d-wave capture yields such a distribution, and p-wave, $J = \frac{3}{2}^-$, is indicated (see, however, ${}^4\text{He}({}^3\text{He}, {}^3\text{He}){}^4\text{He}$); a direct non-resonant capture process seems also possible (1956WA03). For $E_p = 1$ to 2 MeV, large $\cos^2 \theta$ terms appear: the yield shows no resonance behavior in this range (1963MC09). See also (1959GR1A, 1961TO04, 1965LA03).

6. ${}^6\text{Li}(p, n){}^6\text{Be}$

$$Q_m = -5.066$$

$$E_b = 5.608$$

The yield of neutrons increases approximately monotonically from threshold to $E_p = 14.3$ MeV (1964BA16). The excitation function for slow neutrons near threshold indicates that both s- and p-wave neutrons contribute significantly (1964HOZZ). The cross section for production of ground state neutrons is 5 ± 1 mb at $E_p = 9$ MeV (1957BO1F), 0.5 mb/sr at $E_p = 10.5$ MeV, $\theta_{c.m.} = 104^\circ$ (1959AJ81).

7. ${}^6\text{Li}(p, p){}^6\text{Li}$

$$E_b = 5.608$$

Elastic scattering has been studied from $E_p = 1$ to 3 MeV by (1951BA79), from 0.5 to 2.9 MeV by (1963MC09), from 1.2 to 5.6 MeV by (1964FA1D), from 2.4 to 12 MeV by (1963HA53), and at 31 MeV by (1963DE01). Two resonances are reported at $E_p = 1.84$ and 5 MeV. In the range 0.5 to 2.9 MeV, the data are consistent with p-wave formation of a $\frac{5}{2}^-$ or $\frac{3}{2}^-$ state, with $\Gamma_{c.m.} = 836$ keV, $\theta_p^2 = 0.28$, $\theta_\alpha^2 = 0.012$ (see Table 7.2); an s-wave background is evident, possibly reflecting a $\frac{1}{2}^+$ state at higher energies (${}^7\text{Be}^* > 8$ MeV). No evidence is found for a previously reported $\frac{3}{2}^+$

state near $E_p = 1$ MeV (1963MC09). The 5 MeV resonance, corresponding to $E_x = 9.9$ MeV, has a width of ≈ 1.8 MeV and exhibits a behavior much like that of the lower resonance, suggesting that it too is formed by p-wave: on this assumption, the reduced width $\gamma_p^2 = 3 \pm 2$ MeV \cdot fm. A weak rise near $E_p = 8$ to 9 MeV may indicate a further level, ${}^7\text{Be}^* \approx 13$ MeV (1963HA53). See also (1964LE1E).

Polarization of elastically scattered protons has been studied by (1962RO20: 15 MeV), (1963HW01: 39 MeV), while polarization of inelastically scattered protons (to ${}^6\text{Li}^*(3.56)$) has been studied by (1964MA1Y: 150 MeV). See also (1964VE1A).

Inelastic protons (to ${}^6\text{Li}^*(2.18)$) studied from $E_p = 3.6$ to 9.4 MeV show the resonance at $E_p = 5.5$ MeV, Angular distributions suggest p-wave formation with $J^\pi = \frac{3}{2}^-$ or $\frac{5}{2}^-$ (1963HA49). The yield of 3.6 MeV γ -rays (from ${}^6\text{Li}^*(3.56)$) shows a broad maximum at ≈ 6 MeV, probably associated with ${}^7\text{Be}^*(9.9)$: $J^\pi = \frac{3}{2}^-$ is suggested. At $E_p \approx 6.3$ MeV there is an abrupt decrease in the inelastic cross section, which is either due to the onset of the (${}^6\text{Be}+n$) channel or to interference with an other broad state with $J^\pi = \frac{3}{2}^-$ (1964HA37).

$$8. {}^6\text{Li}(p, d){}^5\text{Li} \qquad Q_m = -3.438 \qquad E_b = 5.608$$

See ${}^5\text{Li}$.

$$9. {}^6\text{Li}(p, \alpha){}^3\text{He} \qquad Q_m = 4.021 \qquad E_b = 5.608$$

The cross section exhibits a broad, low maximum near $E_p = 1$ MeV and a pronounced resonance at $E_p = 1.85$ MeV (1951BA79, 1956MA91: see (1963MC09)). No other structure is reported up to $E_p = 5.6$ MeV (1963JE03, 1964FA03). From $E_p = 3$ to 12 MeV, $\theta = 70^\circ$, the excitation function shows only a smooth decrease (1962HE03, 1963TE1B). In the range $E_p = 0.5$ to 2.5 MeV, a strong $\cos \theta$ term is observed, indicating interference between the p-wave resonance and s-wave background (1956MA91, 1963JE03). At $E_p > 8$ MeV, the angular distributions are characterized by forward and backward peaks (1956LI37, 1960AL18, 1962HE03, 1963TE1B). See also (1957JA37, 1960BO13, 1960SA28, 1961KH01, 1963BE08, 1964BE37) and ${}^6\text{Li}$.

$$10. {}^6\text{Li}(d, n){}^7\text{Be} \qquad Q_m = 3.384$$

Two neutron groups are reported, corresponding to the ground and 0.43 MeV states. The γ -ray energy is 428.9 ± 2 keV (corrected for Doppler shift): the ${}^7\text{Li}^*$ - ${}^7\text{Be}^*$ difference is 48.5 ± 1.0 keV (1952TH24). Angular distributions of the n_0 and n_1 groups have been determined at $E_d = 0.56$ to 2.9 MeV (1963BI27), $E_d = 0.6$ to 1.5 MeV (1956NE13), $E_d = 1.8$ to 3.1 MeV (1963CR08) and $E_d = 3.5$ MeV (1952AJ1B). The distributions indicate $l_p = 1$, $J \leq \frac{5}{2}^-$ for both states. Since the n- γ correlations are isotropic (1956NE13) $J = \frac{1}{2}^-$ for the 0.43 MeV excited state is indicated.

Broad maxima are observed in the ratio of low-energy to high-energy neutrons at $E_d = 4.2$ and 5.1 MeV (${}^7\text{Be}^* = 6.5$ MeV and 7.2 MeV, $\Gamma_{\text{c.m.}} = 1.2$ and 0.5 MeV, respectively) (1957SL01). See also (1964PA16, 1965MA1K).

11. ${}^6\text{Li}({}^3\text{He}, d){}^7\text{Be}$ $Q_m = 0.115$

Not reported.

12. ${}^6\text{Li}(\alpha, t){}^7\text{Be}$ $Q_m = -14.206$

Not reported.

13. ${}^6\text{Li}({}^6\text{Li}, n\alpha){}^7\text{Be}$ $Q_m = 1.912$

See (1957NO17, 1962BE16, 1962MC12, 1963BA1Q, 1963CO35) and ${}^{12}\text{C}$.

14. ${}^7\text{Li}(p, n){}^7\text{Be}$ $Q_m = -1.644$

The threshold for this reaction is used as a secondary standard for energy calibrations: the value recommended by (1963MA1R) is 1880.36 ± 0.22 keV (see Table 7.11). See also (1960BR20). Studies of target and shape effects at threshold are reported by (1959WE1A, 1963PA11, 1964BO10).

A determination of Q from the cone angle of neutrons above threshold confirms that the observed threshold corresponds to the true onset of the reaction (1963YO04). A second threshold, corresponding to the first excited state of ${}^7\text{Be}$, yields $E_x = 433 \pm 2$ keV (1960MA1G). Neutrons corresponding to ${}^7\text{Be}^*(4.55)$ are observed for $E_p \gtrsim 7$ MeV (1959AJ81, 1960HI04). At $E_p = 10$ MeV, groups n_0 , n_1 and n_2 account for nearly all the neutrons observed (1963BO06). See also (1952TH1C, 1957BO1F, 1961NI04, 1962AU01). See also (1959GA08, 1960RO21, 1961TO06, 1962BO33, 1963BO1N, 1964BA16, 1964OL1C) and ${}^8\text{Be}$.

15. ${}^7\text{Li}({}^3\text{He}, t){}^7\text{Be}$ $Q_m = -0.880$

Angular distributions have been measured for ground-state tritons (1961WO05: $E({}^3\text{He}) = 4.5$ MeV) and for tritons to both ground and first excited states (1963BO1P: $E({}^3\text{He}) = 3.0$ to 4.0 MeV).

Table 7.11: ${}^7\text{Li}(p, n){}^7\text{Be}$ threshold energies ^a

$E_{\text{thresh.}}$ (keV)	Reference
1879.4 ± 1.0	(1954JO10)
1881.2 ± 0.9	(1959BO14)
1880.8 ± 1.0	b
1880.3 ± 0.5	(1960ST19)
1880.5 ± 0.8	(1961BE13)
1880.48 ± 0.25	c
1879.8 ± 0.6	(1962GA12)
1880.36 ± 0.22	mean: (1963MA1R)
1881.27 ± 0.94	(1964BO10)

^a Revised values as quoted in (1963MA1R), except for the (1964BO10) value.

^b C.P. Browne, as quoted by (1963MA1R).

^c A. Rytz, as quoted by (1963MA1R).

$$16. {}^9\text{Be}(\gamma, 2n){}^7\text{Be} \quad Q_m = -20.561$$

See (1962FO10) and ${}^9\text{Be}$.

$$17. {}^9\text{Be}(p, t){}^7\text{Be} \quad Q_m = -12.079$$

At $E_p = 44$ MeV, triton groups are observed corresponding to the ${}^7\text{Be}$ levels at 0, 0.43 and 4.55 MeV, and to a new level at 10.79 ± 0.04 MeV with $\Gamma = 298 \pm 25$ keV. From the similarity of the angular distribution to that in the $(p, {}^3\text{He})$ reaction to ${}^7\text{Li}^*(11.13)$, it is concluded that the level has $J^\pi = \frac{3}{2}^-$; $T = \frac{3}{2}$ (1965DE08). See also (1954CO02, 1956BE14).

$$18. {}^9\text{Be}(d, tn){}^7\text{Be} \quad Q_m = -14.304$$

See (1955HE83) and ${}^{11}\text{B}$.

$$19. {}^{10}\text{B}(p, \alpha){}^7\text{Be} \quad Q_m = 1.148$$

Alpha groups corresponding to ${}^7\text{Be}(0)$ and ${}^7\text{Be}^*(0.43)$ have been studied by many observers: see (1952AJ38, 1959AJ76). Some reported values for the energy of the first excited state are: 434.4 ± 4 keV (1951BR10), 431 ± 5 keV (1950VA01), 429 ± 3 keV (1952CR30), 428.5 ± 1.8 keV (1952TH24). The mean lifetime of this state is 0.27 ± 0.10 psec (1956BU83: Doppler shift). This value agrees with a shell model calculation by (1955LA1D).

At $E_p = 18$ MeV, α -groups are reported correspondingly to ${}^7\text{Be}^* = 0, 0.49 \pm 0.10, 4.72 \pm 0.08, 6.27 \pm 0.10, 7.21 \pm 0.10$ and 14.6 ± 0.3 MeV. The last group is ten times as intense as any of the others. It is not completely excluded that it may be due to ${}^{10}\text{B}(p, {}^3\text{He}){}^8\text{Be}$ (1955RE16).

Angular distributions of ground and first excited state α -particles have been obtained at 30 energies in the range $E_p = 2.8$ to 7.0 MeV (1964JE01: see also ${}^{11}\text{C}$). See also (1964YA1A).

See also (1960RO21, 1962GR21).

$$20. {}^{12}\text{C}(d, {}^7\text{Li}){}^7\text{Be} \quad Q_m = -17.540$$

See (1965BE1W).

${}^7\text{B}$

(Not illustrated)

From the known location of the ${}^7\text{Be}$ $T = \frac{3}{2}$ state, the mass excess of ${}^7\text{B}$ is calculated as 27.99 ± 0.15 MeV (calculation of the Coulomb energy difference based on the pair ${}^{10}\text{Be}$ - ${}^{10}\text{B}^*$ ($T = 1$)); ${}^7\text{B}$ is then unstable with respect to decay into ${}^5\text{Li} + 2p$ by 1.73 MeV, ${}^6\text{Be} + p$ by 2.33 MeV, and $\alpha + 3p$ by 3.70 MeV (1965DE08). See also (1960GO1B, 1965JA1C).

References

(Closed July 01, 1965)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1948BU31 W.W. Buechner, E.N. Strait, C.G. Stergiopoulos and A. Sperduto, Phys. Rev. 74 (1948) 1569
- 1948EL1A Elliott and Bell, Phys. Rev. 74 (1948) 1869
- 1949EL07 L.G. Elliott and R.E. Bell, Phys. Rev. 76 (1949) 168
- 1949SE20 E. Segre and C.E. Wiegand, Phys. Rev. 75 (1949) 39; Erratum Phys. Rev. 81 (1951) 284
- 1949TU06 C.M. Turner, Phys. Rev. 76 (1949) 148
- 1949TU1B Turner, At. Energy Commission Unclassified Rept. 432 (1949)
- 1949WI13 R.M. Williamson and H.T. Richards, Phys. Rev. 76 (1949) 614
- 1950VA01 D.M. Van Patter, A. Sperduto, E.N. Strait and W.W. Buechner, Phys. Rev. 79 (1950) 900
- 1951BA79 S. Bashkin and H.T. Richards, Phys. Rev. 84 (1951) 1124
- 1951BR10 A.B. Brown, C.W. Snyder, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 82 (1951) 159
- 1951DI12 J.M. Dickson and T.C. Randle, Proc. Phys. Soc. (London) A64 (1951) 902
- 1951GO47 H.E. Gove and J.A. Harvey, Phys. Rev. 82 (1951) 658
- 1952AJ1B Ajzenberg, Phys. Rev. 87 (1952) 205A
- 1952AJ38 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 24 (1952) 321
- 1952CR30 D.S. Craig, D.J. Donahue and K.W. Jones, Phys. Rev. 88 (1952) 808
- 1952GE07 R.W. Gelinas and S.S. Hanna, Phys. Rev. 86 (1952) 253
- 1952TH1C Thomson, Phys. Rev. 88 (1952) 954
- 1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969
- 1953BA04 M.E. Battat and F.L. Ribe, Phys. Rev. 89 (1953) 80
- 1953CO02 E.R. Collins, C.D. McKenzie and C.A. Ramm, Proc. Roy. Soc. A216 (1953) 242
- 1953GE01 R.W. Gelinas and S.S. Hanna, Phys. Rev. 89 (1953) 483
- 1953HO48 J.R. Holt and T.N. Marsham, Prog. Phys. Soc. A66 (1953) 1032
- 1953KR16 J.J. Kraushaar, E.D. Wilson and K.T. Bainbridge, Phys. Rev. 90 (1953) 610

1953PE1A Peaslee and Telegdi, Phys. Rev. 92 (1953) 126
1953ST27 P. Stoll and M. Wachter, Nuovo Cim. 10 (1953) 347
1953TI02 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A66 (1953) 194
1953UE01 R.G. Uebergang and N.W. Tanner, Aust. J. Phys. 6 (1953) 53
1954AL24 D.L. Allan, Nature 174 (1954) 267
1954AL35 K.W. Allen, E. Almqvist, J.T. Dewan and T. Pepper, Phys. Rev. 96 (1954) 684
1954CO02 B.L. Cohen and T.H. Handley, Phys. Rev. 93 (1954) 514
1954CO17 L. Cohen, S.S. Hanna and C.M. Class, Phys. Rev. 94 (1954) 419
1954FR03 G.M. Frye Jr., Phys. Rev. 93 (1954) 1086
1954GO1A Goldemberg and Katz, Can. J. Phys. 32 (1954) 49
1954GO39 J. Goldemberg and L. Katz, Phys. Rev. 95 (1954) 471
1954JO10 K.W. Jones, R.A. Douglas, M.T. McEllistrem and H.T. Richards, Phys. Rev. 94 (1954) 947
1954JO17 C.H. Johnson, H.B. Willard and J.K. Bair, Phys. Rev. 96 (1954) 985
1954MA1D Major and Biedenharn, Rev. Mod. Phys. 26 (1954) 321
1954RU27 R. Rubin and M. Walter, Helv. Phys. Acta 27 (1954) 163
1954ST89 P. Stoll, Helv. Phys. Acta 27 (1954) 395
1954TI16 E.W. Titterton and T.A. Brinkley, Proc. Phys. Soc. (London) A67 (1954) 469
1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77
1955BA59 S. Bashkin and R.R. Carlson, Phys. Rev. 97 (1955) 1245
1955CU17 P. Cuer, D. Magnac-Valette and G. Baumann, Compt. Rend. 240 (1955) 1880
1955HE51 F. Heinrich and R. Rubin, Helv. Phys. Acta 28 (1955) 185
1955HE83 R.E. Heft and W.F. Libby, Phys. Rev. 100 (1955) 799
1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252
1955LA1D Lane, Proc. Phys. Soc. (London) A68 (1955) 189
1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
1955MI55 M. Miwa, J. Phys. Soc. Jpn. 10 (1955) 173
1955RE16 J.B. Reynolds, Phys. Rev. 98 (1955) 1289
1955ST85 J.F. Streib, Phys. Rev. 100 (1955) 1797A
1956BE14 J. Benveniste, R.G. Finke and E.A. Martinelli, Phys. Rev. 101 (1956) 655
1956BO36 P. Bouchez, J. Tobaillem, J. Robert, R. Muxart, R. Mellet and P. Daud, J. Phys. Rad. 17 (1956) 363

- 1956BU83 D.S.T. Bunbury, S. Devons, G. Manning and J.H. Towle, Proc. Phys. Soc. (London) A69 (1956) 165
- 1956CH1B Cheng, Rev. Mod. Phys. 28 (1956) 45
- 1956CO61 H.E. Conzett and R. Silver, Bull. Amer. Phys. Soc. 1 (1956) 388, L7
- 1956DA23 R.B. Day, Phys. Rev. 102 (1956) 767
- 1956HE16 A. Hemmendinger, Bull. Amer. Phys. Soc. 1 (1956) 96, N7
- 1956LI37 J.G. Likely and F.P. Brady, Phys. Rev. 104 (1956) 118
- 1956MA83 E.E. Maslin, J.M. Calvert and A.A. Jaffe, Proc. Phys. Soc. (London) A69 (1956) 754
- 1956MA91 J.B. Marion, G. Weber and F.S. Mozer, Phys. Rev. 104 (1956) 1402
- 1956ME1A Meshkov and Ufford, Phys. Rev. 101 (1956) 734
- 1956NE13 G.C. Neilson and J.B. Warren, Phys. Rev. 103 (1956) 1758
- 1956RE04 J.B. Reynolds and K.G. Standing, Phys. Rev. 101 (1956) 158
- 1956RI34 F.L. Ribe, Phys. Rev. 103 (1956) 741
- 1956WA03 J.B. Warren, T.K. Alexander and G.B. Chadwick, Phys. Rev. 101 (1956) 242
- 1957BA18 G.A. Bartholomew and P.J. Champion, Can. J. Phys. 35 (1957) 1347
- 1957BO1F Bogdanov, Vlasov, Kalinin, Rybakov and Sidorov, Sov. J. At. Energy 3 (1957) 987
- 1957BR97 C.P. Browne, Bull. Amer. Phys. Soc. 2 (1957) 350, N3
- 1957DU37 J.W.M. DuMond, Ann. Phys. 2 (1957) 283
- 1957FR1B French and Fujii, Phys. Rev. 105 (1957) 652
- 1957HO1E Hofstadter, Ann. Rev. Nucl. Sci., Vol. 7 (1957)
- 1957HU1C Huper, Z. Naturforsch. A12 (1957) 295
- 1957JA37 N. Jarmie, J.D. Seagrave et al., LA-2014 (1957)
- 1957LE1E Levinson and Banerjee, Ann. Phys. 2 (1957) 471
- 1957MA04 D.R. Maxson and E.F. Bennett, Bull. Amer. Phys. Soc. 2 (1957) 180, E5
- 1957MA57 J.B. Marion, Nucl. Phys. 4 (1957) 282; Erratum Nucl. Phys. 4 (1957) 492
- 1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
- 1957SC12 C.D. Schrader, J. Benveniste and J.H. Zenger, Bull. Amer. Phys. Soc. 2 (1957) 309, H9
- 1957SI36 R. Silver, Thesis, University of California (1957); UCRL 3887 (1957)
- 1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809
- 1957VA12 S.S. Vasilev, V.V. Komarov and A.M. Popova, Zh. Eksp. Teor. Fiz. 33 (1957) 527; JETP (Sov. Phys.) 6 (1958) 411

- 1957WR37 H.W. Wright, E.I. Wyatt, S.A. Reynolds, W.S. Lyon and T.H. Handle, Nucl. Sci. Eng. 2 (1957) 427
- 1958AL05 R.C. Allen and N. Jarmie, Phys. Rev. 111 (1958) 1129
- 1958BA59 J.B. Ball, A.W. Fairhall and I. Halpern, Bull. Amer. Phys. Soc. 3 (1958) 322, H6
- 1958BE10 O. Beckman and R. Sandstrom, Nucl. Phys. 5 (1958) 595
- 1958BU38 W.E. Burcham, G.P. McCauley, D. Bredin, W.M. Gibson, D.J. Prowse and J. Rotblat, Nucl. Phys. 5 (1958) 141
- 1958EL45 M. El Nadi, Zh. Eksp. Teor. Fiz. 34 (1958) 1207; JETP (Sov. Phys.) 7 (1958) 834
- 1958FO1A Fowler, Astrophys. J. 127 (1958) 551
- 1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
- 1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1
- 1958MI92 P.D. Miller and G.C. Phillips, Phys. Rev. 112 (1958) 2048
- 1958RO49 E.A. Romanovskii and G.F. Timushev, Zh. Eksp. Teor. Fiz. 34 (1958) 1350; JETP (Sov. Phys.) 7 (1958) 932
- 1958RY77 T.W. Rybka and L. Katz, Phys. Rev. 110 (1958) 1123
- 1958WI38 D.H. Wilkinson, Phys. Rev. 109 (1958) 1614
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AJ81 F. Ajzenberg-Selove, C.F. Osgood and C.P. Baker, Phys. Rev. 116 (1959) 1521
- 1959BA1D Baz, Adv. Phys. 8 (1959) 349
- 1959BA1M Balashov, Zh. Eksp. Teor. Fiz. 36 (1959) 1123; JETP (Sov. Phys.) 9 (1959) 798
- 1959BA46 A.J. Bame Jr. and R.L. Cubitt, Phys. Rev. 114 (1959) 1580
- 1959BE84 E.F. Bennett and D.R. Maxson, Phys. Rev. 116 (1959) 131
- 1959BL1C Bloom and Uretsky, Bull. Amer. Phys. Soc. 4 (1959) 78
- 1959BO14 R.O. Bondelid and C.A. Kennedy, Phys. Rev. 115 (1959) 1601
- 1959BO1C Bockelman, Nucl. Phys. 13 (1959) 205
- 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
- 1959CH1B Chen and Hintz, Congres Int. de Phys. Nucl., Paris, 1958 (Dunod, Paris, 1959) 387
- 1959FE1B Feingold, Phys. Rev. 114 (1959) 540
- 1959GA08 F. Gabbard, R.H. Davis and T.W. Bonner, Phys. Rev. 114 (1959) 201
- 1959GR1A Griffiths, Congres Int. de Phys. Nucl., 1958 (Dunod, Paris, 1959) 447
- 1959HA29 E.W. Hamburger, Thesis, Univ. of Pittsburgh (1959)
- 1959HE1C Henley and Jacobsohn, Phys. Rev. 113 (1959) 225

1959HO03 H.D. Holmgren and R.L. Johnston, Phys. Rev. 113 (1959) 1556
 1959JO43 A. Johansson, G. Tibell and P. Hillman, Nucl. Phys. 11 (1959) 540
 1959LE30 R.D. Leamer, Phys. Rev. 114 (1959) 1590
 1959MA1F Maris, Nucl. Phys. 9 (1959) 577
 1959MA1G Mamasakhlisov and Kopaleishvili, Zh. Eksp. Teor. Fiz. 37 (1959) 1134; JETP (Sov. Phys.) 10 (1960) 807
 1959ME1D Meyer-Berkhout, Phys. Rev. 115 (1959) 1300
 1959MU25 R.B. Murray and H.W. Schmitt, Phys. Rev. 115 (1959) 1707
 1959PA02 W.B. Pardo and J.H. Roberts, Bull. Amer. Phys. Soc. 4 (1959) 218, AB6
 1959PH37 G.C. Phillips and P.D. Miller, Phys. Rev. 115 (1959) 1268
 1959PU75 K.H. Purser, Aust. J. Phys. 12 (1959) 231
 1959RO62 T.A. Romanowski and W.H. Voelker, Phys. Rev. 113 (1959) 886
 1959SI1A Sitenko, Usp. Fiz. Nauk 67 (1959) 377
 1959SW63 C.P. Swann, V.K. Rasmussen and F.R. Metzger, Phys. Rev. 114 (1959) 862
 1959VL24 N.A. Vlasov and A.A. Ogloblin, Zh. Eksp. Teor. Fiz. 37 (1959) 54; JETP (Sov. Phys.) 10 (1960) 39
 1959WE1A Weston, Bilpuch and Newson, Bull. Amer. Phys. Soc. 4 (1959) 42
 1960AL18 J. Alster and J. Gonzalez-Vidal, Bull. Amer. Phys. Soc. 5 (1960) 493, C5
 1960AN02 G.O. Andre, Nucl. Phys. 15 (1960) 464
 1960AN14 A.L. Androsenko, D.L. Broder and A.I. Lashuk, At. Energ. 9 (1960) 403; Sov. J. At. Energy 9 (1961) 945
 1960BA51 V.V. Balashov, Atomn. Energ. (USSR) 9 (1960) 48; Sov. J. At. Energy 9 (1961) 544
 1960BL1B Blair, Bull. Amer. Phys. Soc. 5 (1960) 474
 1960BO13 R. Bouchez, C. Delome, J. Fleury, J. Krafft, P. Perrin, L. Goldman, M. Boge and B. Dudek, J. Phys. Rad. 21 (1960) 346
 1960BO23 E.C. Booth, Nucl. Phys. 19 (1960) 426
 1960BR19 D.J. Bredin, J.B.A. England, D. Evans, J.S. McKee, P.V. March, E.M. Mosinger and W.T. Toner, Proc. Roy. Soc. A258 (1960) 202
 1960BR1J Brolley, Stewart and Rosen, Nucl. Forces and Few Nucleon Problem, Vol. 2 (Pergamon, 1960) 455
 1960BR20 C.P. Browne, J.A. Galey, J.R. Erskine and K.L. Warsh, Phys. Rev. 120 (1960) 905
 1960CH1B Chen and Hintz, Nucl. Forces and the Few Nucleon Problem (Pergamon, 1960) 683
 1960FA06 R.W. Fast, P.A. Flournoy, R.S. Tickle and W.D. Whitehead, Phys. Rev. 118 (1960) 535

- 1960GO1B Goldansky, Nucl. Phys. 19 (1960) 482
- 1960HA14 E.W. Hamburger and J.R. Cameron, Phys. Rev. 117 (1960) 781
- 1960HA21 H.J. Hausman, G.F. Dell and H.F. Bowsher, Phys. Rev. 118 (1960) 1237
- 1960HE1F Hetley, Hildebrand, Tompkins and Wooten, Bull. Amer. Phys. Soc. 5 (1960) 454
- 1960HI04 K. Hisatake, Y. Ishizaki, A. Isoya, T. Nakamura, Y. Nakano, B. Saheki, Y. Saji and K. Yuasa, J. Phys. Soc. Jpn. 15 (1960) 741
- 1960HI10 P. Hillman, H. Tyren and Th.A.J. Maris, Phys. Rev. Lett. 5 (1960) 107
- 1960HU08 D.J. Hughes, B.A. Magurno and M.K. Brussel, BNL-325, 2nd Ed., Suppl. 1 (1960)
- 1960JO14 A. Johansson, G. Tibell, K. Parker and P.E. Hodgson, Nucl. Phys. 21 (1960) 383
- 1960KO1C Kotin, Rev. Mex. Fisica 9 (1960) 73
- 1960KU1B Kunz, Ann. Phys. 11 (1960) 275
- 1960KU1C Kulchitskii and Presperin, Zh. Eksp. Teor. Fiz. 39 (1960) 1001; JETP (Sov. Phys.) 12 (1961) 696
- 1960LA1C Lane, Rev. Mod. Phys. 32 (1960) 519
- 1960MA15 K.V. Makariunas and S.V. Starodubtsev, Zh. Eksp. Teor. Fiz. 38 (1960) 372; JETP (Sov. Phys.) 11 (1960) 271
- 1960MA1G Marion, Levesque, Ludemann and Detenbeck, Nucl. Instrum. Meth. 8 (1960) 297
- 1960MA1H Marquez and Redon, Proc. Int. on Nucl. Struct., Kingston (1960)
- 1960MA32 R.D. Macfarlane and J.B. French, Rev. Mod. Phys. 32 (1960) 567
- 1960MC1E McKee, Nucl. Forces and the Few Nucleon Problem (Pergamon, 1960) 609
- 1960MI02 M. Miwa and M. Yamanouchi, J. Phys. Soc. Jpn. 15 (1960) 947
- 1960NE1C Neudachin, Teplov and Tulinov, JETP (Sov. Phys.) 10 (1960) 387
- 1960PE11 L.D. Pearlstein, Y.C. Tang and K. Wildermuth, Nucl. Phys. 18 (1960) 23
- 1960PE1A Perey, Proc. Int. Conf. on Nucl. Struct., Kingston (1960)
- 1960PE24 V.P. Perelygin and K.V. Tolstov, Atomn. Energ. (USSR) 9 (1960) 488; Sov. J. At. Energy 9 (1961) 1028
- 1960PE25 J.M. Peterson, A. Bratenahl and J.P. Stoering, Phys. Rev. 120 (1960) 521
- 1960PH1A Phillips and Tombrello, Nucl. Phys. 19 (1960) 555
- 1960RO21 J.C. Roy, M. Bresesti and J.J. Hawton, Can. J. Phys. 38 (1960) 1428
- 1960SA28 Y. Sakamoto and T. Takemiya, Prog. Theor. Phys. 23 (1960) 172
- 1960SH1A Sheline and Wildermuth, Nucl. Phys. 21 (1960) 196
- 1960ST17 P.H. Stelson and F.K. McGowan, Nucl. Phys. 16 (1960) 92
- 1960ST19 H.H. Staub and H. Winkler, Nucl. Phys. 17 (1960) 271

- 1960TA1C Talmi and Unna, *Ann. Rev. Nucl. Sci.* 10 (1960) 353
- 1960VA1G Vashakidze, Kopaleishvili, Mamasakhlisov and Chilashvili, *Zh. Eksp. Teor. Fiz.* 39 (1960) 666; *JETP (Sov. Phys.)* 12 (1961) 466
- 1961AN07 D.S. Andreev, V.D. Vasilev, G.M. Gusinskii, K.I. Erokhina and I.K. Lemberg, *Izv. Akad. Nauk SSSR Ser. Fiz.* 25 (1961) 832; *Bull. Acad. Sci. USSR* 25 (1962) 842
- 1961BA1D Badalyan and Baz, *Zh. Eksp. Teor. Fiz.* 40 (1961) 549; *JETP (Sov. Phys.)* 13 (1961) 383
- 1961BA1E Balashov, Neudachin and Smirnov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 25 (1961) 170; *Bull. Acad. Sci. USSR Phys.* 25 (1961) 165
- 1961BE13 E.H. Beckner, R.L. Bramblett, G.C. Phillips and T.A. Eastwood, *Phys. Rev.* 123 (1961) 2100
- 1961BE24 A.A. Bergman and F.L. Shapiro, *Zh. Eksp. Teor. Fiz.* 40 (1961) 1270; *JETP (Sov. Phys.)* 13 (1961) 895
- 1961BL1C Blin-Stoyle and Spector, *Phys. Rev.* 124 (1961) 1199
- 1961CH09 R. Chiba, H.E. Conzett, H. Morinaga, N. Mutsuro, K. Shoda and M. Kimura, *J. Phys. Soc. Jpn.* 16 (1961) 1077
- 1961CH1C R.F. Christy and I. Duck, *Nucl. Phys.* 24 (1961) 89
- 1961CL10 A.B. Clegg, *Phil. Mag.* 6 (1961) 1207
- 1961DA04 S.E. Darden, T.R. Donoghue and C.A. Kelsey, *Nucl. Phys.* 22 (1961) 439
- 1961FA02 A.W. Fairhall, *Bull. Amer. Phys. Soc.* 6 (1961) 150, OA5
- 1961GA09 J.P. Garron, J.C. Jacmart, M. Riou, C. Ruhla, J. Teillac, C. Caverzasio and K. Strauch, *Phys. Rev. Lett.* 7 (1961) 261
- 1961GR27 G.M. Griffiths, R.A. Morrow, P.J. Riley and J.B. Warren, *Can. J. Phys.* 39 (1961) 1397
- 1961HA1F Hamburger, Cohen and Price, *Phys. Rev.* 121 (1961) 1143
- 1961HO1F Holmgren and Wolicki, *Proc. Rutherford Jub. Int. Conf., Manchester, England*; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 541
- 1961HO21 H.D. Holmgren and L.M. Cameron, *Proc. Rutherford Jub. Int. Conf., Manchester, England*; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 531
- 1961JA19 von L. Jarczyk, J. Lang, R. Muller and W. Wolffi, *Helv. Phys. Acta* 34 (1961) 483
- 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, *Rev. Mex. Fisica* 10 (1961) 247
- 1961JO18 A. Johansson, U. Svanberg and P.E. Hodgson, *Ark. Fys.* 19 (1961) 541
- 1961KA29 S.L. Kahalas and R.K. Nesbet, *Phys. Rev. Lett.* 6 (1961) 549
- 1961KH01 N.H. Khanh, L. Goldman and R. Bouchez, *J. Phys. Rad.* 22 (1961) 267
- 1961KH03 F.C. Khanna, Y.C. Tang and K. Wildermuth, *Phys. Rev.* 124 (1961) 515

1961KO1J Kowalska, Acta Phys. Pol. 20 (1961) 1019
1961KU1C Kurath, Phys. Rev. 124 (1961) 552
1961LA1A Lane, Langsdorf, Monahan and Elwyn, Ann. Phys. 12 (1961) 135
1961LE1K Lemeille, Saunier and Marquez, Centro Brasileiro de Pesquisas Fisicas 8:5 (1961)
1961NI04 A. Nilsson, Ark. Fys. 19 (1961) 289
1961SL06 R.J. Slobodrian, Bol. Acad. Nac. Cienc. (Cordoba, Rep. Arg.) 42 (1961) 75; Phys. Abs. 67, 1989, Phys. Abs. 22164 (1964)
1961TA05 Y.C. Tang, K. Wildermuth and L.D. Pearlstein, Phys. Rev. 123 (1961) 548
1961TO04 T.A. Tombrello and G.C. Phillips, Phys. Rev. 122 (1961) 224
1961TO06 J.H. Towle and B.E.F. Macefield, Proc. Phys. Soc. (London) 77 (1961) 1217
1961VA03 A.M.K. Van Beek and G.O. Andre, Nucl. Phys. 24 (1961) 102
1961WO05 E.A. Wolicki and A.R. Knudson, Bull. Amer. Phys. Soc. 6 (1961) 415, B2
1961YA04 M. Yamada and Z. Matumoto, J. Phys. Soc. Jpn. 16 (1961) 1497
1962AU01 S.M. Austin, Bull. Amer. Phys. Soc. 7 (1962) 269, CB6
1962BA1D Barber, Ann. Rev. Nucl. Sci. 12 (1962) 1
1962BA1W Batchelor and Towle, Phys. Lett. 2 (1962) 312
1962BA1X Bahcall, Phys. Rev. 128 (1962) 1297
1962BE16 E.H. Berkowitz, Phys. Rev. 126 (1962) 2168
1962BE1J Berggren and Jacob, Phys. Lett. 1 (1962) 258
1962BE1K Berggren, Brown and Jacob, Phys. Lett. 1 (1962) 88
1962BE1N Belousov, Rusakov and Tamm, Zh. Eksp. Teor. Fiz. 43 (1962) 813; JETP (Sov. Phys.) 16 (1963) 576
1962BE1P Beets, Donneaux, Gierts and Winand, Reactor Sci. Tech. 16 (1962) 383; Phys. Abs. 12695 (1963)
1962BI11 J.A. Biggerstaff, R.F. Hood, H. Scott and M.T. McEllistrem, Nucl. Phys. 36 (1962) 631
1962BO33 P.H. Bowen, G.C. Cox, G.B. Huxtable, J.P. Scanlon, J.J. Thresher and A. Langsford, Nucl. Phys. 30 (1962) 475
1962CA1F Casanova, An. Real. Soc. Espan. Fis. y Quim. A58 (1962) 107; Phys. Abs. 18331 (1962)
1962CH26 V.P. Chizhov, et al., Nucl. Phys. 34 (1962) 562
1962CL1F Clegg, Nucl. Phys. 33 (1962) 194
1962CR09 C.L. Critchfield, LA-2714 (1962)

1962DI1A Dietrich, Phys. Lett. 2 (1962) 139
1962EL01 A.J. Elwyn and R.O. Lane, Nucl. Phys. 31 (1962) 78
1962EL1E Elton, Phys. Lett. 2 (1962) 41; Erratum Phys. Lett. 5 (1963) 96
1962FO10 M.S. Foster and A.F. Voigt, J. Inorg. Nucl. Chem. 24 (1962) 343
1962GA09 J.P. Garron, J.C. Jacmart, M. Riou, C. Ruhla, J. Teillac and K. Strauch, Nucl. Phys. 37 (1962) 126
1962GA12 B.R. Gasten, Bull. Amer. Phys. Soc. 7 (1962) 549, T1
1962GL1A Gleason, Gruverman and Need, Int. J. Appl. Rad. Isotopes 13 (1962) 223
1962GO1P Gottschalk, Unpublished Thesis, Harvard Univ. (1962)
1962GR08 A.G. Gregory, T.R. Sherwood and E.W. Titterton, Nucl. Phys. 32 (1962) 543
1962GR21 M.A. Grace, P.P. Singh and R.E. Segel, Bull. Amer. Phys. Soc. 7 (1962) 549, T2
1962HE03 N.P. Heydenburg and I.G. Han, Bull. Amer. Phys. Soc. 7 (1962) 58, R1
1962HE1C Henley, Bull. Amer. Phys. Soc. 7 (1962) 18
1962IN02 D.R. Inglis, Nucl. Phys. 30 (1962) 1
1962IN1A Inglis, Rev. Mod. Phys. 34 (1962) 165
1962IV1A Ivanytskii, Ukr. Fiz. Zh. 7 (1962) 1160; Phys. Abs. 22425 (1963)
1962LA07 J. Laune, J. Phys. Rad. 23 (1962) 238
1962LI13 F.F. Liu, F.J. Loeffler, T.R. Palfrey and Y.S. Kim, Phys. Rev. 128 (1962) 2784
1962MA1R Marshak, Bull. Amer. Phys. Soc. 7 (1962) 305
1962MA40 D.R. Maxson, Phys. Rev. 128 (1962) 1321
1962MA59 K.V. Makariunas, E.K. Makariuniene and V.J. Dienys, Litov. Fiz. Sbornik (USSR) 2 (1962) 351; ; Nucl. Sci. Abstr. 18, 2032, Abs.15242 (1964); Phys. Abs. 3703 (1964)
1962MC12 R.L. McGrath, Phys. Rev. 127 (1962) 2138
1962MO17 W.L. Mouton, J.P.F. Sellschop and R.J. Keddy, Phys. Rev. 128 (1962) 2745
1962NE12 D. Newton, A.B. Clegg, G.L. Salmon, P.S. Fisher and K.J. Foley, Proc. Phys. Soc. (London) 79 (1962) 27
1962PH1B Phillips et al., Phys. Rev. Lett. 9 (1962) 502
1962RI09 R.C. Ritter, P.H. Stelson, F.K. McGowan and R.L. Robinson, Phys. Rev. 128 (1962) 2320
1962RO12 L. Rosen and L. Stewart, Phys. Rev. 126 (1962) 1150
1962RO1F Robson, Nucl. Phys. 30 (1962) 316
1962RO20 L. Rosen and W.T. Leland, Phys. Rev. Lett. 8 (1962) 379
1962RO23 V.M. Rout and W.M. Jones, Phys. Rev. 128 (1962) 263

- 1962RU04 C. Ruhla, M. Riou, J.P. Garron, J.C. Jacmart and L. Massonnet, Phys. Lett. 2 (1962) 44
- 1962SH24 A.K. Shardanov and V.G. Shevchenko, Zh. Eksp. Teor. Fiz. 42 (1962) 1438; JETP (Sov. Phys.) 15 (1962) 996
- 1962SL02 R.J. Slobodrian, Phys. Rev. 125 (1962) 1003
- 1962SL04 R.J. Slobodrian, Phys. Rev. 126 (1962) 1059
- 1962ST1E Strnad, Phys. Rev. 125 (1962) 1639
- 1962ST1F Strnad, Nucl. Phys. 35 (1962) 451
- 1962TA11 J.G.V. Taylor and J.S. Merritt, Can. J. Phys. 40 (1962) 926
- 1962TE1D Temmer, Bull. Amer. Phys. Soc. 7 (1962) 569
- 1962TI01 G. Tibell, O. Sundberg and U. Miklavzic, Phys. Lett. 1 (1962) 172
- 1962VO1C Volkov and Kulchitskii, Zh. Eksp. Teor. Fiz. 42 (1962) 108; JETP (Sov. Phys.) 15 (1962) 77
- 1962WE04 J.J. Wesolowski, J.M. Fowler, J.B. Reynolds and R.J. Wilson, Bull. Amer. Phys. Soc. 7 (1962) 269, CB8
- 1962WO07 C. Wong, J.D. Anderson and J.W. McClure, Nucl. Phys. 33 (1962) 680
- 1963AL1J Alter and Lorenzi, NAA-SR-MEMO-9028 (1963)
- 1963AS01 V.J. Ashby, H.C. Catron, M.D. Goldberg, R.W. Hill, J.M. Le Blanc, L.L. Newkirk, J.P. Stoering, C.J. Taylor and M.A. Williamson, Phys. Rev. 129 (1963) 1771
- 1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527
- 1963BA1Q Ballini and Saunier, J. Physique 24 (1963) 904
- 1963BA1V Batchelor and Towle, AWRE NR 6/63 (1963)
- 1963BA2A Baudinet-robinet, Donneaux, Robaye and Winand, J. Phys. 24 (1963) 803
- 1963BA2B Balashov, Zh. Eksp. Teor. Fiz. 45 (1963) 541; JETP (Sov. Phys.) 18 (1964) 371
- 1963BA50 R. Batchelor and J.H. Towle, Nucl. Phys. 47 (1963) 385
- 1963BA56 J.F. Barry, J. Nucl. Energy 17 (1963) 273
- 1963BE08 H. Beaumevieille, N. Longequeue and J.P. Longequeue, Compt. Rend. 256 (1963) 1494
- 1963BE1A Benioff, Phys. Rev. 129 (1963) 1355
- 1963BE1R Bennett and Grant, 3rd Conf. on Reaction between Complex Nuclei (1963) 50
- 1963BE26 M. Bernheim and G.R. Bishop, Phys. Lett. 5 (1963) 294
- 1963BE42 T. Berggren and G. Jacob, Nucl. Phys. 47 (1963) 481
- 1963BE53 M. Bernheim and G.R. Bishop, J. Phys. 24 (1963) 970

- 1963BI27 M. Birk, G. Goldring, P. Hillman and R. Moreh, Nucl. Phys. 41 (1963) 58
- 1963BO06 R.R. Borchers and C.H. Poppe, Phys. Rev. 129 (1963) 2679
- 1963BO1N Bowen, Cox, Huxtable, Scanlon, Thrasher, Langsford and Appel, Nucl. Phys. 41 (1963) 177
- 1963BO1P Bogart, Devons and Tatcher, Padua (1963) 960
- 1963CH08 C.M. Chesterfield and B.M. Spicer, Nucl. Phys. 41 (1963) 675
- 1963CH20 A. Chatterjee, Nucl. Phys. 47 (1963) 511
- 1963CL03 F.M. Clikeman, A.J. Bureau, J.R. McConnell, M.G. Stewart and D.A. Tripp, Bull. Amer. Phys. Soc. 8 (1963) 290
- 1963CL1B Clegg, CERN 63-28 (1963)
- 1963CO1D Costa et al., Phys. Lett. 6 (1963) 226
- 1963CO35 M. Coste, L. Marquez and S. Shafroth, J. Phys. 24 (1963) 906
- 1963CR08 L. Cranberg, A. Jacquot and H. Liskien, Nucl. Phys. 42 (1963) 608
- 1963DE01 D.W. Devins and H.H. Forster, Bull. Amer. Phys. Soc. 8 (1963) 12, BA13
- 1963DE1F Dearnalay, Prog. in Fast Neutron Phys. (1963) 173
- 1963DR1B Drisko, Satchler and Dassel, 3rd Conf. on Reactions between Complex Nuclei (1963) 85
- 1963EL1C Elton, Padua (1963) 1093
- 1963FU1D Fujii, Prog. Theor. Phys. 29 (1963) 374
- 1963GL1F Glazkov, Atomn. Energ. (USSR) 15 (1963) 416
- 1963GO04 J. Goldemberg and Y. Torizuka, Phys. Rev. 129 (1963) 312
- 1963GO1M Goldberg, May and Stehn, BNL-400, 2nd Ed., Vol. 1 (1963)
- 1963HA49 W.D. Harrison, Bull. Amer. Phys. Soc. 8 (1963) 597, F2
- 1963HA53 W.D. Harrison and A.B. Whitehead, Phys. Rev. 132 (1963) 2607
- 1963HW01 C.F. Hwang, G. Clausnitzer, D.H. Nordby, S. Suwa and J.H. Williams, Phys. Rev. 131 (1963) 2602
- 1963JE03 J.M.F. Jeronimo, G.S. Mani and A. Sadeghi, Nucl. Phys. 43 (1963) 424
- 1963JO07 A. Johansson and Y. Sakamoto, Nucl. Phys. 42 (1963) 625
- 1963KI1C Kim, Liu, Loeffler and Palfrey, Phys. Rev. 129 (1963) 1362
- 1963KI1D Kim, Nucl. Phys. 49 (1963) 651
- 1963KL1A Klapisch, Rameau, Epherre and Gradsztajn, J. Phys. 24 (1963) 839
- 1963KU16 L.A. Kulchitskii and Y.M. Volkov, Zh. Eksp. Teor. Fiz. 44 (1963) 1153; JETP (Sov. Phys.) 17 (1963) 780

1963KU25 L.A. Kulchitskii, Y.M. Volkov, V.P. Denisov and V.I. Ogurtsov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 27 (1963) 1412; *Bull. Acad. Sci. USSR Phys. Ser.* 27 (1964) 1387

1963MA1R Marion, Univ. of Maryland TR 320 (1963)

1963MA61 M.H. MacGregor, R. Booth and W.P. Ball, *Phys. Rev.* 130 (1963) 1471

1963MC09 J.A. McCray, *Phys. Rev.* 130 (1963) 2034

1963ME01 J.B. Mead, L.B. Geesaman and H.B. Knowles, *Bull. Amer. Phys. Soc.* 8 (1963) 292, A15

1963ME09 V. Meyer, W. Pfeifer and H.H. Staub, *Helv. Phys. Acta* 36 (1963) 465

1963MO02 W.L. Mouton, J.P.F. Sellschop and G. Wiechers, *Phys. Rev.* 129 (1963) 361

1963NE02 B.M.K. Nefkens, *Phys. Rev. Lett.* 10 (1963) 55

1963NO02 E. Norbeck, S.A. Coon, R.R. Carlson and E. Berkowitz, *Phys. Rev.* 130 (1963) 1971

1963OP1A Oparin, Saukov and Shuvalov, *Atomn. Energ. (USSR)* 15 (1963) 411; *J. Nucl. Energ.* 18 (1964) 596

1963PA11 D.W. Palmer, *Bull. Amer. Phys. Soc.* 8 (1963) 526, J9

1963PA12 P.D. Parker and R.W. Kavanagh, *Phys. Rev.* 131 (1963) 2578

1963RI1B Riou, Padua (1963) 18

1963SC1M Schmid, Wildermuth and Tang, 3rd Conf. on Reactions between Complex Nuclei (1963) 59

1963SC30 R.G. Schlecht, UCRL-11047 (1963)

1963SH1A Shapiro and Kolybasov, *Nucl. Phys.* 49 (1963) 515

1963SQ1A Squires, Forest and Hodgson, *Nucl. Phys.* 42 (1963) 490

1963TA10 Y.C. Tang, E. Schmid and K. Wildermuth, *Phys. Rev.* 131 (1963) 2631

1963TA1D Takemiya, *Prrog. Theor. Phys.* 30 (1963) 191

1963TE1B Temmer, Pauda (1963) 1013

1963TO04 T.A. Tombrello and P.D. Parker, *Phys. Rev.* 130 (1963) 1112

1963TO06 T.A. Tombrello and P.D. Parker, *Phys. Rev.* 131 (1963) 2582

1963VA19 J.H. van der Merwe, *Phys. Rev.* 131 (1963) 2181

1963WA07 M.H. Wachter, *Phys. Lett.* 4 (1963) 353

1963WA1K Waffler and Walch, Padua (1963) 633

1963WI1B Willey, *Nucl. Phys.* 40 (1963) 529

1963YO04 F.C. Young and J.B. Marion, *Nucl. Phys.* 41 (1963) 561

1964AL08 F.R. Allum, G.M. Crawley and B.M. Spicer, *Nucl. Phys.* 51 (1964) 177

1964AL1N Allison and Smythe, *Bull. Amer. Phys. Soc.* 9 (1964) 544

- 1964AR22 H. Artus, G. Fricke and D.E. Von Stein, Z. Phys. 178 (1964) 109
- 1964AR25 A.H. Armstrong, J. Gammel, L. Rosen and G.M. Frye, Jr., Nucl. Phys. 52 (1964) 505
- 1964BA09 A.C.L. Barnard, C.M. Jones and G.C. Phillips, Nucl. Phys. 50 (1964) 629
- 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209
- 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
- 1964BA1P Balashov and Boyarkina, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1964) 359
- 1964BA2A Bahcall, Phys. Rev. Lett. 12 (1964) 300
- 1964BE1N Beregi, Selenskaja, Neudatchin and Smirnov, Congres Int. de Phys. Nucl., Paris (1964)
- 1964BE37 H. Beaumevieille, J.P. Longequeue, N. Longequeue and R. Bouchez, J. Phys. (Paris) 25 (1964) 933
- 1964BI03 G.R. Bishop and M. Bernheim, Phys. Lett. 8 (1964) 48
- 1964BO10 R.O. Bondelid and E.E.D. Whiting, Phys. Rev. 134 (1964) B591
- 1964BO22 E.C. Booth, B. Chasan and K.A. Wright, Nucl. Phys. 57 (1964) 403
- 1964FA03 U. Fasoli, D. Toniolo and G. Zago, Phys. Lett. 8 (1964) 127
- 1964FA1D Fasoli, Silverstein, Toniolo and Zago, Nuovo Cim. 34 (1964) 1832
- 1964GI1F Gierts, in Photographie Corpusculaire III, Ed. Demers (Press Universitaires de Montreal, 1964) 459
- 1964GR1J Green, Nucl. Phys. 54 (1964) 505
- 1964GR40 L. Green and D.J. Donahue, Phys. Rev. 135 (1964) B701
- 1964GU1B Guth, Congres Int. de Phys. Nucl., Paris (1964)
- 1964HA37 W.D. Harrison, Bull. Amer. Phys. Soc. 9 (1964) 703, B1
- 1964HO31 M. Honda and D. Lal, Nucl. Phys. 51 (1964) 363
- 1964HOZZ J.L. Honsaker, Unpublished Thesis, Cal Tech (1964)
- 1964JA03 J.C. Jacmart, J.P. Garron, M. Riou and C. Ruhla, Phys. Lett. 8 (1964) 269
- 1964JE01 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Nucl. Phys. 50 (1964) 516
- 1964LA19 R.O. Lane, A.J. Elwyn and A. Langsdorf Jr., Phys. Rev. 136 (1964) B1710
- 1964LE1E Lejeune and Lebon, Congres Int. de Phys. Nucl., Paris (1964)
- 1964LI1D Lim and McCarthy, Phys. Rev. 133 (1964) B1006
- 1964LI1F Ligonniere, Vassent and Bernas, Compt. Rend. 259 (1964) 1406
- 1964MA04 S. Mayo, J. Testoni and O.M. Bilaniuk, Phys. Rev. 133 (1964) B350
- 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550

- 1964MA1Y Marty, Geoffrion, Rolland, Tatischeff and Willis, *Congres Int. de Phys. Nucl.*, Paris (1964)
- 1964MA2B Manuzio, Ricco and Sanzone, *Congres Int. de Phys. Nucl.*, Paris (1964)
- 1964MA2C Maximon and Isabelle, *Phys. Rev.* 136 (1964) B674
- 1964MO1K Moscati, Nefkens and Todoroff, *Phys. Lett.* 12 (1964) 45; *Bull. Amer. Phys. Soc.* 9 (1964) 407
- 1964NE1E Neudachin, Shevchenko and Yudin, *Phys. Lett.* 10 (1964) 180
- 1964NE1H Nefkens, Moscati and Todoroff, *Congres Int. de Phys. Nucl.*, Paris (1964)
- 1964OL1A Ollerhead, Chasman and Bromley, *Phys. Rev.* 134 (1964) B74
- 1964OL1C Olness, Seth and Lewis, *Nucl. Phys.* 52 (1964) 529
- 1964PA16 P. Paul, S.S. Hanna and G.D. Sprouse, *Bull. Amer. Phys. Soc.* 9 (1964) 704, B7
- 1964PA1A Parker, Bahcall and Fowler, *Astrophys. J.* 139 (1964) 602
- 1964PA1K Pape, UCRL 11598 (1964)
- 1964PE1E Pendlebury, AWRE O 60/64 (1964)
- 1964PH1A Phillips, *Rev. Mod. Phys.* 36 (1964) 1085
- 1964RA1B Rayudu, *Can. J. Chem.* 42 (1964) 1149; *Phys. Abs.* 25393 (1964)
- 1964RI1E Rickards and Calvillo, *Rev. Mex. de Fisica* 13 (1964) 133, and Private Communication (1964)
- 1964SA1F Sandulescu and Dumitrescu, *Phys. Lett.* 11 (1964) 420
- 1964SA1H Sakamoto, *Phys. Rev.* 134 (1964) B1211
- 1964SA1L Satchler and Haybron, *Phys. Lett.* 11 (1964) 313
- 1964SC1J Schuler and Schopper, *Congres Int. de Phys. Nucl.*, Paris (1964)
- 1964SH1C Shapiro, *Congres Int. de Phys. Nucl.*, Paris, 1964 (CRNS, Paris, 1964) 313
- 1964SP04 R.J. Spiger and T.A. Tombrello, *Bull. Amer. Phys. Soc.* 9 (1964) 703, B2
- 1964ST15 T. Stovall and N.M. Hintz, *Phys. Rev.* 135 (1964) B330
- 1964ST1B Stovall, *Phys. Rev.* 133 (1964) B268
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964TA02 B. Tatischeff, N. Marty, X. De Bouard, J.G. Fox, B. Geoffrion and C. Rolland, *Phys. Lett.* 8 (1964) 54
- 1964TI02 G. Tibell, O. Sundberg and P.U. Renberg, *Ark. Fys.* 25 (1964) 433
- 1964VA19 V. Valkovic, *Nucl. Phys.* 60 (1964) 581
- 1964VE1A Venter and Frahn, *Ann. Phys.* 27 (1964) 385, 401

1964WH01 L. Wharton, L.P. Gold and W. Klemperer, Phys. Rev. 133 (1964) B270
 1964YA1A Yanabu et al., J. Phys. Soc. Jpn. 19 (1964) 1818
 1965BA2C Bahcall, Nucl. Phys. (1965)
 1965BE1R Bernas, Epherre, Gradsztajn, Klapisch and Yiou, Phys. Lett. 15 (1965) 147
 1965BE1W Bender and Newman, Bull. Amer. Phys. Soc. 10 (1965) 602
 1965CH11 B.T. Chertok and E.C. Booth, Nucl. Phys. 66 (1965) 230
 1965DE08 C. Detraz, J. Cerny and R.H. Pehl, Phys. Rev. Lett. 14 (1965) 708
 1965FA1B Fabre de La Ripelle, Prog. Theor. Phys. 33 (1965) 38
 1965FU1G Fuller and Cohen, Appendix I, Nucl. Data Sheets 6-5 (1965)
 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81
 1965HE1B Henkel, Bull. Amer. Phys. Soc. 10 (1965) 601
 1965JA1C Janecke, Nucl. Phys. 61 (1965) 383
 1965JA1L Jackson and Elton, Proc. Phys. Soc. (London) 85 (1965) 659
 1965JU1A Juna and Konecny, Czech. J. Phys. 15 (1965) 95
 1965LA03 H. Langevin-Joliot, C. Stephan, G. Johnson and J. Vernotte, Phys. Lett. 14 (1965) 208
 1965MA1K Marion, Nucl. Phys. 68 (1965) 463
 1965MA1Y C. Mahaux and G. Robaye, Nucl. Phys. 74 (1965) 161
 1965NE1C Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 25
 1965PR1F Present, Bull. Amer. Phys. Soc. 10 (1965) 253
 1965RI1A Riou, Rev. Mod. Phys. 37 (1965) 375
 1965SA1L Sahai, Bull. Amer. Phys. Soc. 10 (1965) 443
 1965SC07 S. Schwarz, L.G. Stromberg and A. Bergstrom, Nucl. Phys. 63 (1965) 593
 1965WE1E Weidenmuller, Nucl. Phys. (1965)
 1965ZH1A Zhdanov, Kuzmin and Yakovlev, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 239
 1966HA09 E.T. Hazzard, F. Ajzenberg-Selove and P.V. Hewka, Nucl. Phys. 75 (1966) 592; Erratum Nucl. Phys. 89 (1966) 706
 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692

