

Energy Levels of Light Nuclei $A = 8$

T. Lauritsen ^a and F. Ajzenberg-Selove ^b

^a *California Institute of Technology, Pasadena, California*

^b *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics* 78 (1966), p. 1. This version of $A = 8$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed July 01, 1965)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 8$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^8\text{He}\$](#) , [\$^8\text{Li}\$](#) , [\$^8\text{Be}\$](#) , [\$^8\text{B}\$](#)

B. Tables of Recommended Level Energies:

[Table 8.1](#): Energy levels of ^8Li

[Table 8.5](#): Energy levels of ^8Be

[Table 8.18](#): Energy levels of ^8B

C. [References](#)

D. Figures: [\$^8\text{Li}\$](#) , [\$^8\text{Be}\$](#) , [\$^8\text{B}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

⁸He
(Not illustrated)

GENERAL: See (1960GO1B, 1960ZE03, 1961BA1C, 1961LO07, 1961YA04, 1964GO25, 1965BA1A).

Mass of ⁸He : From the systematics of neutron binding energies (1960GO36) and from the calculated mass excess of ⁷He, the mass excess of ⁸He is estimated as 31.6 to 32.4 MeV. It is thus stable against ⁶He + 2n by 1.3 to 2.1 MeV and should exhibit β^- -decay to ⁸Li*(0.98) with $E_\beta(\text{max}) = 9.7$ to 10.5 MeV (1965DE08).

1. ⁸He(β^-)⁸Li* $Q_m \approx 10$

An activity with half-life 30 ± 20 msec, $E_\beta(\text{max}) = 13 \pm 2$ MeV, observed in ¹¹B + γ (320 MeV), is ascribed to ⁸He (1963NE07). See also (1965PO03).

2. ¹¹B(γ , 3p)⁸He $Q_m \approx -45$

At $E_\gamma(\text{max}) = 320$ MeV, the production cross section is $> 6 \mu\text{b}$ (1963NE07).

⁸Li
(Figs. 11 and 14)

GENERAL: See (1955LA1D, 1956KU1A, 1957FR1B, 1960TA1C, 1962IN02, 1963LO1E, 1963NA1E, 1964BE1N, 1964GR1J, 1964LO1F, 1964ST1B).

Ground state :

$$\mu = +1.653 \pm 0.0008 \text{ nm: see } ^7\text{Li}(n, \gamma)^8\text{Li, (1965FU1G).}$$

1. ⁸Li(β^-)⁸Be $Q_m = 16.002$

The beta decay leads mainly to ⁸Be*(2.9): see ⁸Be. Reported half-lives are listed in Table 8.2; taking $\tau_{1/2} = 0.849$ sec and $Q = 16.002 - 2.9$, $ft = 4.14 \times 10^5$ (1966BA1A). The distribution of recoil momenta indicates $J^\pi = 2^+$ (see ⁸Be).

The asymmetry of the β -decay has been exploited as an indicator of ⁸Li nuclear polarization: see ⁷Li(n, γ)⁸Li.

Table 8.1: Energy levels of ${}^8\text{Li}$

E_x (MeV \pm keV)	$J^\pi; T$	Γ (keV)	Decay	Reactions
g.s.	$2^+; 1$	$\tau_{1/2} = 849 \pm 4$ msec	β^-	1, 2, 3, 9, 10, 12, 14, 19, 20, 21
0.975 ± 12	$1^+, 2^+; 1$	$\tau < 0.2$ psec	γ	2, 9, 12
2.258 ± 3	$3^+; 1$	31 ± 5	γ, n	3, 4, 9
3.21	$(1^\pm, 2^+); 1$	≈ 1000	n, n_1	5
(5.0)	$(2^-); 1$	broad	n	4
(6.4)		≈ 2000	n	4
6.53 ± 20	$T = 1$	< 40		16
(8.9 ± 400)	$(1^-, 2^-)$	≈ 4000		14

Table 8.2: The half-life of ${}^8\text{Li}$

$\tau_{1/2}$ (sec)	Reference
0.89 ± 0.02	(1947HU06)
0.825 ± 0.02	(1951RA12)
0.89 ± 0.01	(1953BU35)
0.875 ± 0.02	(1953BR1B)
0.85 ± 0.016	(1952SH44)
0.84 ± 0.04	(1954WI25)
0.841 ± 0.004	(1954KL36)
0.84 ± 0.02	(1959FA02)
0.88 ± 0.03	(1959IM04)
0.82 ± 0.02	(1962NE14)
0.87 ± 0.02	(1965BE1P)
0.849 ± 0.004	weighted mean

2. ${}^6\text{Li}(t, p){}^8\text{Li}$ $Q_m = 0.803$

Ground-state protons have been observed with E_t up to 3.3 MeV: see (1952MO19, 1952PE02, 1954AL35, 1955CU1A).

At $E({}^6\text{Li}) = 4.5$ MeV, the 0.98 MeV γ -ray has been observed. Doppler shift measurements give $\tau < 0.2$ psec for ${}^8\text{Li}^*(0.98)$. The transition is thus dipole in character, with a maximum E2 admixture of 2×10^{-3} in intensity (1965MO1P).

3. ${}^7\text{Li}(n, \gamma){}^8\text{Li}$ $Q_m = 2.033$

The thermal capture cross section is 37 ± 4 mb (1964ST25); capture γ -rays of energy $E_\gamma = 2.02, 1.06$ and 0.96 MeV are reported with intensity ratios $0.8/0.2/0.3$ (1961JA19; prelim. values). See also (1959BO1C).

The cross section for capture radiation has been measured for $E_n = 40$ to 1000 keV: it decreases from $50 \mu\text{b}$ at $E_n = 40$ keV to $5 \mu\text{b}$ at 1000 keV. A maximum in the cross section appears at $E_n = 250$ keV corresponding to the known resonance in ${}^7\text{Li}(n, n)$; $\Gamma_\gamma = 0.07 \pm 0.03$ eV (1959IM04).

Measurement of the asymmetry of β -decay of ${}^8\text{Li}$ produced by capture of polarized neutrons indicate that the thermal capture takes place $> 80\%$ in the $J_c = 2$ channel (1959CO68, 1961WA03, 1962AB01). An NMR determination yields $g = 0.8265 \pm 0.0004$ nm/ \hbar , or with $J = 2$, $\mu = +1.653 \pm 0.0008$ nm (1959CO68, 1962CO08). Shell model calculations give this value with $a/K = 2.1$ (1959KU1E).

4. ${}^7\text{Li}(n, n){}^7\text{Li}$ $E_b = 2.033$

Total cross sections for Li and for ${}^7\text{Li}$ are reported in (1960HU1A, 1964ST25). Recent measurements of the total cross section have been reported for $E_n = 1$ to 390 keV (1964HI04), 10 to 155 keV (1959BI1B), 0.2 to 2.2 MeV (1961LA1A, 1964LA19), 1.5 to 7.5 MeV (1963BA50), 14 MeV (1964AR1C), and 18 to 28 MeV (1960PE1B). Angular distribution information is summarized in (1963GO1M). See also (1964LA19). A measurement at $E_n = 96$ MeV (1960SA25) has been analyzed in terms of the optical model by (1960HO14). See also (1962BA1W, 1963LU10). The thermal cross section is 1.07 ± 0.04 b (1960HU1A); the coherent scattering length (thermal, bound) is $a = -2.1$ fm (1964ST25). See also (1961ME02). A pronounced resonance is observed at $E_n = 256$ keV with $J = 3^+$, formed by p-waves (Table 8.3). Polarization measurements confirm the assignment and indicate that the background s-wave scattering is contributed mainly by channel spin $J_c = 2$ (1961DA04, 1962EL01: see also (1961LA1A)) in agreement with earlier results of (1956TH06) but in conflict with (1956WI04). In this case, the coherent scattering length in the $J_c = 1$ channel is $a_1 = [+1.2]$, $a_2 = [-3.6]$ fm, suggesting an s-wave, $J^\pi = 2^-$, resonance at positive neutron energy (1956TH06, 1960LA1C). A good account of the polarization is given by

Table 8.3: ${}^7\text{Li}(n, n){}^7\text{Li}$ resonance parameters ^a

	(1956WI04)	(1956TH06, 1960HU1A)
E_{res} (keV)	256	258 ± 3 ^c
Γ (keV)	32	32 ^d
$\Gamma_n(E_r)$ (keV)	35.8	
Γ_γ (eV)(cm)	0.07 ± 0.03 ^b	
γ_n^2 (keV)	351	307
E_λ (keV)	-49	-43
radius (fm)	4.08	4.0
σ_{max} (b)	11.2 ^e	12.0

^a Energies in laboratory system, except for Γ_γ .

^b (1959IM04).

^c $E_{\text{res}} = 275$ keV, $\sigma_{\text{max}} = 7.0 \pm 0.2$ b (1956GO62).

^d 35 ± 5 keV (1960HU1A).

^e (1961LA1A).

the assumption of levels at $E_n = 0.25$ and 3.4 MeV, with $J^\pi = 3^+$ and 2^- , together with a broad $J^\pi = 3^-$ level at higher energy (1964LA19). It is noted that the polarization near the $J = 3^+$ resonance is similar in shape to that of ${}^7\text{Li}(p, n){}^7\text{Be}$ at the presumptive mirror level, but opposite in sign (1962EL01). A broad maximum centering at $E_n \approx 5$ MeV may indicate a level at $E_x \approx 6.4$ MeV (1956GO62, 1960HU1A, 1963BA50, 1964ST25).

See also (1960PE02, 1962FO1E, 1962OT01, 1963AL1J, 1963GL1E, 1963HA1G, 1963WE1F).

5. ${}^7\text{Li}(n, n'){}^7\text{Li}^*$

$$E_b = 2.033$$

The excitation function for 0.48 MeV γ -rays shows an abrupt rise from threshold (indicating s-wave formation and emission) and a broad maximum ($\Gamma \approx 1$ MeV) at $E_n = 1.35$ MeV. The rise above threshold indicates the existence of a $J = 1^-$ level, which may be identified with the 1.35 MeV resonance, if a strong d-wave contribution is included. Other odd parity assignments to this resonance lead to excessive reduced widths. A good fit is obtained with either $J = 1^-$ or 1^+ , $\Gamma_{\text{lab}} = 1.14$ MeV, $J^\pi = 2^+$ is not excluded (1955FR10). In the range $E_n = 1.5$ to 4 MeV, the cross section for emission of 0.48 MeV γ -rays remains constant at first and then increases to 280 mb at $E_n = 4$ MeV (1963BA50). The cross section for emission of 0.48 MeV γ -rays is ≈ 80 mb ($\pm 12\%$) over the range $E_n = 13.6$ to 14.3 MeV (1962BE35). See also (1963GO1M, 1963MA1Q).

6. ${}^7\text{Li}(n, 2n){}^6\text{Li}$

$$Q_m = -7.253$$

$$E_b = 2.033$$

The cross section is 27 ± 15 mb at $E_n = 10.2$ MeV and 56 ± 5 mb at 14.1 MeV (1963AS01). See also (1964ST25).

7. ${}^7\text{Li}(n, d){}^6\text{He}$

$$Q_m = -7.755$$

$$E_b = 2.033$$

At $E_n = 14$ MeV, the cross section is 9.8 ± 1.1 mb (1953BA04), 14 ± 1.5 mb (1963BA56).

8. ${}^7\text{Li}(n, t){}^4\text{He} + n$

$$Q_m = -2.467$$

$$E_b = 2.033$$

The cross section rises to 450 mb at $E_n \approx 8$ MeV and thereafter decreases slowly to 300 mb at 15 MeV (1964ST25). The large cross section, comparable to the geometric value, supports the hypothesis that ${}^7\text{Li}$ may be described as an $(\alpha + t)$ cluster (1962RO12). See also (1963AL1J, 1963BA50, 1963BR28, 1964VA19).

9. ${}^7\text{Li}(d, p){}^8\text{Li}$

$$Q_m = -0.192$$

Observed proton groups indicate levels at 974 ± 15 (1955LE24), 977 ± 20 keV (1955KH35) and at 2.28 MeV (1955LE24, 1960HA14). At $E_d = 3.5$ MeV, no excited states are observed below the 0.97 MeV level with an intensity $> \frac{1}{160}$ of the ground-state group (1961ER01). At $E_d = 15$ MeV, no states observed in the region $2.28 < E_x < 8$ MeV: a limit of 0.6 mb/sr is placed on groups with widths $\lesssim 100$ keV ($\theta = 10^\circ, 14^\circ$ and 25°) (1960HA14). See also (1965WA12).

Angular distributions at $E_d = 14$ to 15 MeV have been analyzed by PWBA: see Table 8.4. At low bombarding energies, $E_d = 0.5$ to 4 MeV, angular distributions show striking agreement with simple PWBA theory. The applicability of the theory in this range is ascribed to the proximity of a pole in the stripping cross section when Q and E_d are small (1960SE08, 1960WA1G, 1963SE1F). Analysis in both PWBA and DWBA is reported by (1961HA1G): the latter leads to nearly ten times larger θ_n^2 .

The cross section for ${}^7\text{Li}(d, p){}^8\text{Li}^*(\gamma){}^8\text{Li}$ has been measured for $E_d = 1.9$ to 3.3 MeV: no resonances appear. The γ -ray angular distribution, very nearly isotropic for $E_d > 2.3$ MeV, indicates that stripping dominates. If the transition is M1 (see ${}^6\text{Li}(t, p){}^8\text{Li}$), the 0.97 MeV state has $J = 1^+$ or 2^+ : $E_\gamma = 980 \pm 10$ keV (1962CH14).

The polarization of recoil ${}^8\text{Li}$ nuclei has been studied at $E_d = 10$ MeV (1963LE1K).

See also (1959BO1C, 1959HA29, 1959SE1A, 1960PR1D, 1960SE13, 1963AL19, 1963GL1C, 1963SE1G).

Table 8.4: Reduced widths from ${}^7\text{Li}(d, p){}^8\text{Li}$

E_x in ${}^8\text{Li}$ (MeV)	J^π	l	r_0^a (fm)	$\theta_n^2{}^a$	r_0^b (fm)	$\theta_n^2{}^b$	$\theta_n^2{}^c$	$\theta_n^2{}^d$
0	2^+	1	3.8	0.066	4.2	0.053	0.038 – 0.07	0.49 – 0.60
0.97	(1^+)	1	3.5	0.040	4.2	0.028		
2.28	3^+	1	3.0	0.012	4.0	0.015		

^a (1960HA14): PWBA.

^b (1960MA32): PWBA.

^c (1961HA1G): PWBA.

^d (1961HA1G): DWBA.

10. ${}^7\text{Li}(t, d){}^8\text{Li}$ $Q_m = -4.225$

See (1965AJ01).

11. ${}^7\text{Li}(\alpha, {}^3\text{He}){}^8\text{Li}$ $Q_m = -18.546$

Not reported.

12. ${}^9\text{Be}(\gamma, p){}^8\text{Li}$ $Q_m = -16.885$

At $E_{\text{brems.}} = 32$ MeV, transitions are observed to the ground and first excited states of ${}^8\text{Li}$ (1962CU05). See also (1958CH31, 1962NE14, 1963NE02, 1965AL02, 1965KO1B).

13. ${}^9\text{Be}(n, d){}^8\text{Li}$ $Q_m = -14.660$

Not reported.

14. ${}^9\text{Be}(p, 2p){}^8\text{Li}$ $Q_m = -16.885$

At $E_p = 138$ MeV, the cross section of production of 0.98 MeV γ -rays is 2.4 ± 0.4 mb (1961CL09). The summed proton spectrum shows two peaks, corresponding to pickup of protons with binding energies of $\approx 16.7 \pm 0.3$ MeV and 25.8 ± 0.4 MeV ($1p_{\frac{3}{2}}$ and $1s_{\frac{1}{2}}$, respectively) (1958MA1B, 1958TY49, 1962GA23, 1964TI02, 1966TY01). See also ${}^9\text{Be}$.

$$15. {}^9\text{Be}(d, {}^3\text{He}){}^8\text{Li} \quad Q_m = -11.391$$

See (1954WI25).

$$16. {}^9\text{Be}(t, \alpha){}^8\text{Li} \quad Q_m = 2.930$$

At $E_t = 12.98$ MeV, α -particle groups are observed to the ${}^8\text{Li}$ ground state, to the states at 0.98 and 2.26 MeV and to a state at 6.530 ± 0.020 MeV. The width of the 6.53 MeV state is < 40 keV (1965WA12).

$$17. {}^9\text{Be}({}^7\text{Li}, {}^8\text{Be}){}^8\text{Li} \quad Q_m = 0.367$$

See ${}^9\text{Be}$.

$$18. {}^{10}\text{B}(n, {}^3\text{He}){}^8\text{Li} \quad Q_m = -15.754$$

Not reported.

$$19. {}^{10}\text{B}(p, 3p){}^8\text{Li} \quad Q_m = -23.472$$

See (1965BE1P).

$$20. {}^{11}\text{B}(\gamma, {}^3\text{He}){}^8\text{Li} \quad Q_m = -27.210$$

See (1963NE07).

$$21. {}^{11}\text{B}(n, \alpha){}^8\text{Li} \quad Q_m = -6.632$$

See (1959SA04) and ${}^{12}\text{B}$.

^8Be
(Figs. 12 and 14)

GENERAL: See (1956KU1A, 1957FR1B, 1958WI1E, 1959BA1F, 1959BA1D, 1959BR1E, 1959WI1B, 1960BI1E, 1960KU05, 1960PE11, 1960PH1A, 1960PH1C, 1960TA1C, 1961BA1E, 1961CL10, 1961VA17, 1962IN02, 1962IN1A, 1962IW1A, 1963BR1N, 1963BU1C, 1963DA1C, 1963FR1G, 1963KU03, 1963MA1E, 1963MO1H, 1963NA1E, 1963SH1G, 1964AM1D, 1964BA1Y, 1964BE1N, 1964BE1M, 1964BR1H, 1964DA1G, 1964DU1D, 1964GR1J, 1964MA1G, 1964VO1B, 1965BA2H, 1965BE1H, 1965MA2B, 1965MA1G, 1965NE1C, 1965TR1B, 1965YU1D).

Table 8.5: Energy levels of ^8Be


E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	Decay	Tables	Reactions
g.s.	6.8 ± 0.6 eV	$0^+; 0$	α	6, 8	1, 4, 12, 13, 14, 15, 16, 23, 24, 25, 26, 28, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 42, 45, 46, 50, 51
2.90 ± 30	1450 ± 60	$2^+; 0$	α	6, 8, 9	4, 13, 14, 15, 16, 23, 24, 26, 28, 29, 30, 31, 32, 33, 34, 37, 39, 42, 44, 45, 50
11.4 ± 300	≈ 7000	$4^+; 0$	α	8	4, 13, 14, 23, 32, 34
16.628 ± 5	97 ± 11	2^+	α	6, 8, 11, 16	4, 13, 15, 23, 29, 30, 32, 34, 39, 45
16.923 ± 6	83 ± 10	2^+	α	6, 8, 11	4, 13, 15, 23, 34, 39, 45
17.638 ± 2	10.7 ± 0.5	1^+	p, γ	6, 11, 12, 14, 16	13, 15, 16, 18, 23, 24, 34, 45
18.153 ± 5	147	1^+	p, γ	6, 12, 14, 16	15, 16, 18, 19, 23
18.9	> 500	2^-	p, n	13	17, 32
19.05 ± 20	270 ± 20	(3)	p, γ	6, 12, 14	16, 18, 27
19.22	190	3^+	p, n	13, 14	17, 18
19.9	≈ 1000	$2^+; (0)$	p, α	15	22
(19.9)	> 1000	(1^+)	p, n	13	17
20.36	1160	(1^-)	p, n	13	17
20.9 ± 200	1570 ± 180	4^-	p	14	19
21.5	1000	(3^+)	p, n	13	17
21.6	≈ 5000	(1^-)	p, γ	6, 12	16
22.5	(800)	$2^+; (0)$	p, d, α , (n), (γ)	10, 12, 14, 15	6, 7, 11, 15, 16, 18, 22

Table 8.5: Energy levels of ^8Be (continued)


E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	Decay	Tables	Reactions
23.		(4 ⁺)	p, α	15	22, 32
23.6	≈ 5000	(1 ⁻ , 2 ⁻)	p, γ	6, 12	16
(24.0)	(1360)	(0 ⁻ ; 0)	d, α	10	11
25.2	(270)	2 ⁺	d, α , p	10, 15	7, 11, 22
(28.5)	≈ 20 MeV	6 ⁺ ; (0)	α	8	3
(57)	≈ 73 MeV	8 ⁺ ; (0)	α	8	4



The weighted mean of direct Q determinations to 1957 is $Q = 94.1 \pm 0.7$ keV (1957VA11). Reported widths are 4.5 ± 3 eV (1956RU41), 6.8 ± 0.6 eV (1962BA1C), ≤ 3.5 eV (1956HE57), > 0.1 eV (1955TR03): see Table 8.8. (1965BE1B) find $Q = 92.12 \pm 0.05$ keV, $\Gamma = 6.8 \pm 1.7$ eV.



At $E_\alpha = 39$ MeV, $\sigma < 0.7$ mb (1952WA31).



See ^7Li .



Reported differential cross section measurements are cited in Table 8.7. The course of the derived phase shifts with increasing energy from $E_\alpha = 0.15$ to 120 MeV is exhibited in (1956RU41, 1958NI05, 1960JO03, 1963TO02, 1965DA1A).

The s-wave phase shift, δ_0 , decreases smoothly from 180° at 0.3 MeV. The absence of measurable effects in the range $E_\alpha = 0.15$ to 0.2 MeV yields an upper limit of 3.5 eV on the width of the ground state (1956HE57). Analysis of the 0 – 6 MeV data leads to $\Gamma(\text{g.s.}) = 4.5 \pm 3$ eV (1956RU41), 6.8 ± 0.6 eV (1962BA1C): see Table 8.8.

Table 8.6: Electromagnetic transitions in ${}^8\text{Be}$

Transition	Remarks	References
16.6 \rightarrow 2.9 } 16.9 \rightarrow 2.9 }	$\Gamma(\text{M1}) = 1.9 \pm 0.6 \text{ eV}$	(1962NO02: (${}^8\text{Li}$, ${}^8\text{B}$))
17.6 \rightarrow g.s.	$\Gamma(\text{M1}) = 16.7 \text{ eV}$	(1949FO18, 1961ME10)
17.6 \rightarrow 2.9	$\Gamma(\text{M1}) = 8.2 \text{ eV}$, $\Gamma(\text{E2}) = 0.15 \text{ eV}$	(1961ME10)
17.6 \rightarrow 16.6	$\sigma(\text{p}, \gamma) = 6.5 \mu\text{b}$	(1965WI07, 1965WI1K)
17.6 \rightarrow 16.9	$\sigma(\text{p}, \gamma) = 0.4 \mu\text{b}$	(1965WI07, 1965WI1K)
18.15 \rightarrow g.s.	$\Gamma \approx 1.8 \text{ eV}$	(1954KR06, 1960MA23)
18.15 \rightarrow 2.9	$\Gamma \approx 3.6 \text{ eV}$	(1954KR06, 1960MA23)
19.05 \rightarrow 2.9		(1957NE1A, 1963PE15)
21.6 \rightarrow g.s.	E1 giant res.: $\int \sigma dE = 22 \text{ MeV} \cdot \text{mb}$	(1963MI08, 1963RE09)
23.6 \rightarrow 2.9	E1 giant res.	(1963MI08, 1963RE09)

 Table 8.7: Measurements of $d\sigma/d\Omega$ in ${}^4\text{He}(\alpha, \alpha){}^4\text{He}$

E_α (MeV)	References	E_α (MeV)	References
0.15 – 3.0	(1956HE57)	23 – 38	(1959BR71)
3.0 – 6.0	(1956RU41)	23 – 51	(1964SH19)
4.0 – 12.0	(1963TO02)	30	(1951GR45, 1952GR1A)
5.0 – 9.0	(1960JO03)	33.5, 35.5	(1961CH09)
6.4 – 7.8	(1960BE05, 1961DU01)	38.5	(1957BU13)
10 – 20	(1964WE1C)	37 – 47	(1960CO04, 1964CO1D, 1964CO27)
12.3 – 22.9	(1956NI20)	51	(1964VA1J)
12.9 – 21.6	(1953ST52)	53 – 120	(1965DA1A)
20, 20.4	(1951BR92, 1951MA1B)		

The d-wave phase shift becomes appreciable for $E_\alpha > 2.5$ MeV and passes through resonance at $E_\alpha = 6$ MeV ($J = 2^+$, $E_x = 3.18$ MeV, $\Gamma = 1.5$ MeV). The g-wave shift rises from $E_\alpha \approx 11$ MeV and indicates a broad $J^\pi = 4^+$ level at $E_x = 11.4$ MeV. The $l = 6$ and $l = 8$ phase shifts become active above $E_\alpha \approx 30$ and 50 MeV respectively. Dispersion formula fits are somewhat unsatisfactory, but indicate $J^\pi = 6^+$ and 8^+ levels at $E_x \approx 28$ and ≈ 57 MeV respectively. Both the excitation energies and the reduced widths for all five levels are approximately proportional to $J(J + 1)$ (1965DA1A).

Sharp oscillations in the excitation functions at 15° and 45° are observed corresponding to ${}^8\text{Be}^*(16.6)$ and (16.9) . Since no oscillations were observed at 27.8° , these states are assigned $J^\pi = 2^+$ (1964SH19).

Optical model analysis of α - α scattering is discussed by (1960IG02, 1965DA1A); analysis in terms of two-nucleon forces is given by (1959VA1F, 1961SC1B). See also (1959BU07, 1959WI1F, 1960BI1E, 1961GO1T, 1961SH1F, 1962IG1B, 1962SH1F, 1963WI1H, 1964EN1C, 1965OK1B).

5. ${}^6\text{Li}(d, \gamma){}^8\text{Be}$ $Q_m = 22.280$

Not observed: (1953SA1A, 1954SI07).

6. (a) ${}^6\text{Li}(d, n){}^7\text{Be}$ $Q_m = 3.384$ $E_b = 22.280$
 (b) ${}^6\text{Li}(d, n){}^4\text{He} + {}^3\text{He}$ $Q_m = 1.797$

The yield curve has been measured for $E_d = 0.06$ to 5.5 MeV (1952BA1A, 1954HI34, 1956NE13, 1957SL01). A broad s-wave resonance is indicated at $E_d = 0.41$ MeV, $\Gamma = 0.45$ MeV (1952BA1A, 1956NE13). The forward cross section rises from ≈ 22 mb/sr at $E_d = 1.1$ MeV to ≈ 57 mb/sr at 5.5 MeV without sharp resonances (1957SL01). Above $E_d = 0.6$ MeV, angular distributions indicate a strong admixture of stripping process (1956NE13).

Comparison of the yields n_0/n_1 (${}^7\text{Be}(0)$ and ${}^7\text{Be}^*(0.43)$) and p_0/p_1 (${}^7\text{Li}(0)$ and ${}^7\text{Li}^*(0.48)$) over the energy range $E_d = 0.4$ to 3.2 MeV shows that the angular distributions are closely similar for n and p. The yield ratios are also closely equal over this range, consistent with the assumption of charge symmetry. The ratio n_0/n_1 increases rapidly as E_d falls below 0.8 MeV, suggesting a change in the reaction mechanism there (1957WI24, 1963BI1B, 1963CR08).

7. (a) ${}^6\text{Li}(d, p){}^7\text{Li}$ $Q_m = 5.028$ $E_b = 22.280$
 (b) ${}^6\text{Li}(d, p){}^4\text{He} + {}^3\text{H}$ $Q_m = 2.561$

Cross sections and angular distributions have been measured for $E_d = 30$ keV to 5.4 MeV (1959AJ76, 1963BI1B, 1963ME09, 1964PA06). A broad maximum near $E_d = 1.0$ MeV is interpreted as indicating a level at $E_d = 0.4$ MeV (1950WH02). In the range $E_d = 1$ to 5 MeV there

Table 8.8: ${}^8\text{Be}$ parameters from ${}^4\text{He}(\alpha, \alpha){}^4\text{He}$

E_x (MeV)	J^π	$\Gamma_{\text{c.m.}}$	R (fm)	θ^2	References
g.s.	0^+	$4.5 \pm 3 \text{ eV}^{\text{a}}$	5.7	0.15	(1956RU41)
			4.4	[0.40]	(1958NI05)
		$6.8 \pm 0.6 \text{ eV}$	3.5	1.7	(1962BA1C)
		$\leq 3.5 \text{ eV}$			(1956HE57)
		$6.8 \pm 1.7 \text{ eV}$			(1965BE1B)
2.9	2^+	2 MeV	5.0	0.7	(1956RU41)
3.1			3.5	1.32	(1960JO03)
3.18		1.5 MeV	3.5	1.27	(1963TO02)
3.13		1.8 MeV	3.5	2.46	(1962BA1C)
11.8	4^+		4.4	1.3	(1958NI05)
11.4 ± 0.3					(1959BR71)
11.3 ± 0.4					(1962CE01) ^b
≈ 14					(1960IG02)
16.6	2^+				(1964SH19)
16.9	2^+				(1964SH19)
≈ 28.5	6^+	$\approx 20 \text{ MeV}$	≈ 4.5	≈ 1.8	(1964DA12, 1965DA1A)
≈ 57	8^+	$\approx 73 \text{ MeV}$	4.5	≈ 4.1	(1964DA12, 1965DA1A)

^a See also (1962BA1C).

^b From ${}^6\text{Li}(\alpha, d){}^8\text{Be}$.

is evidence for both direct interaction and compound nucleus formation (1963BI1B, 1963ME09, 1964PA06): at back angles the (d, p_1) data show evidence of the $E_d = 3.7 \text{ MeV}$ resonance (see ${}^6\text{Li}(d, \alpha){}^4\text{He}$). See also ${}^6\text{Li}(d, n){}^7\text{Be}$ and (1964FE01).

8. ${}^6\text{Li}(d, d){}^6\text{Li}$

$$E_b = 22.280$$

Excitation functions have been measured for $E_d = 2$ to 4.8 MeV ; they do not show any clear resonance behavior (1964PA06). See also ${}^6\text{Li}$.

9. ${}^6\text{Li}(d, t){}^5\text{Li}$

$$Q_m = 0.595$$

$$E_b = 22.280$$

Table 8.9: Parameters of ${}^8\text{Be}^*(2.9)$ ^a

E_x (MeV)	$\Gamma_{\text{c.m.}}$ (MeV)	R (fm)	γ^2 (MeV)	θ^2 ^b	Reaction	Reference
(3.18)	1.5	3.5	3.36	1.27	${}^4\text{He}(\alpha, \alpha)$ ^c	(1963TO02)
		4.48	2.65	3.4	${}^7\text{Li}(\text{p}, \gamma\alpha)$	(1960GE07)
2.6 ± 0.15	0.7 ± 0.1				${}^7\text{Li}(\text{p}, \gamma)$	(1963BA58)
3.1 ± 0.1	1.75 ± 0.1				${}^7\text{Li}(\text{d}, \text{n})$	(1964JO04)
	1.6 ± 0.4				${}^7\text{Be}(\text{d}, \text{p})$	(1959SP1A)
2.90 ± 0.06	1.53 ± 0.04	5.75	0.60	0.64	${}^7\text{Be}(\text{d}, \text{p})$	(1960KA17)
2.8 ± 0.1	0.8				${}^9\text{Be}(\text{d}, \text{t})$	(1955CU16)
	1.35 ± 0.15	5		0.082	${}^9\text{Be}(\text{d}, \text{t})$	(1959VL24)
2.90 ± 0.04	1.35 ± 0.15				${}^9\text{Be}({}^3\text{He}, \alpha)$	(1963DO08)
3.06		5.5	0.62	0.60	${}^{10}\text{B}(\text{d}, \alpha)$	(1963PU02)
2.88 ± 0.08		4.48	3.0	≈ 2	${}^{10}\text{B}(\text{d}, \alpha)$	(1953TR04)
2.895 ± 0.028	1.45 ± 0.06				mean	

^a Spectra from several reactions – e.g. ${}^8\text{Li}(\beta^-, \alpha)$, ${}^8\text{B}(\beta^+, \alpha)$, ${}^{11}\text{B}(\text{p}, \alpha)$, ${}^9\text{Be}(\text{p}, \text{d})$ – are reported “consistent with phase shifts of ${}^4\text{He}(\alpha, \alpha)$ scattering” and so support the parameters of Table 8.8.

^b Units of $\frac{3}{2}(\hbar^2/\mu R^2)$.

^c See Table 8.8.

The cross section for tritium production rises rapidly to 190 mb at 1 MeV, then more slowly to 290 mb near 4 MeV. There is evidence of deviation from isotropy near 0.4 MeV (1955MA20). See also ${}^5\text{Li}$.

$$10. \quad {}^6\text{Li}(\text{d}, {}^3\text{He}){}^5\text{He} \qquad Q_{\text{m}} = 0.839 \qquad E_{\text{b}} = 22.280$$

See ${}^5\text{He}$.

$$11. \quad {}^6\text{Li}(\text{d}, \alpha){}^4\text{He} \qquad Q_{\text{m}} = 22.375 \qquad E_{\text{b}} = 22.280$$

Cross sections and angular distributions have been measured for $E_{\text{d}} = 0.03$ to 12 MeV (1948HE1A, 1950WH02, 1962HA15, 1962JE02, 1963AN10, 1963ME09, 1964MA2D, 1964PA06, 1965MA13). Maxima are observed at $E_{\text{d}} = 0.8$ MeV, $\Gamma_{\text{lab}} \approx 0.8$ MeV, and $E_{\text{d}} = 3.75$ MeV,

Table 8.10: Levels of ^8Be from $^6\text{Li}(d, \alpha)^4\text{He}$ ^a

E_{res} (MeV)	$E_{\lambda}(\text{c.m.})$ (MeV)	E_x (MeV)	$\Gamma_{\text{c.m.}}$ ^b (MeV)	J^π	θ_d^2	θ_α^2
0.8	0.26	22.54	0.80	2^+	0.23	0.003
	1.74	24.02	1.36	0^+	0.75	0.09
3.75	2.95	25.23	0.27	2^+	0.025	0.022

^a (1965FR02).

^b Partial width, $\Gamma_{\lambda\lambda}$.

$\Gamma_{\text{lab}} \approx 1.4$ MeV. Analysis of the angular distributions in terms of two-level interference favors $J^\pi = 2^+$ for the upper level, and $J^\pi = 0^+$ or 2^+ for the lower (1963ME09, 1964PA06). The upper level appears to have a large deuteron width, $\theta_d^2 \approx 0.2$, and small $\theta_\alpha^2 \approx 0.025$ and $\theta_p^2 < 0.005$ (1964PA06). A more elaborate analysis appears to require that both levels, $E_x = 22.54$ and 25.23 MeV, have $J^\pi = 2^+$ and that a third broad $J^\pi = 0^+$ level exist at $E_x = 24.02$ MeV (1965FR02): see Table 8.10. The same parameters give a good account of the data up to $E_d = 9.5$ MeV. There is no evidence of further resonances (1965MA13). See also (1959HA29, 1960HA14, 1963EL1D, 1963FR1G, 1964FE01).

12. $^6\text{Li}(t, n)^8\text{Be}$ $Q_m = 16.023$

See (1959AJ1C, 1962SE1A).

13. $^6\text{Li}(^3\text{He}, p)^8\text{Be}$ $Q_m = 16.787$

At $E(^3\text{He}) = 1.2$ to 2 MeV, proton groups are reported corresponding to $^8\text{Be}(0)$, $^8\text{Be}^*(2.9)$ and possibly $^8\text{Be}^*(12.3)$ (1956MO19, 1956SC01). At $E(^3\text{He}) = 3.5$ to 4.2 MeV, groups with $Q = 0.163$, -0.143 and -0.854 MeV (± 10 keV) are observed, corresponding to the 16.6 , 16.9 and 17.6 MeV states (1961ER01): see Table 8.11.

The differential cross sections for ground-state protons show backward peaking at $E(^3\text{He}) = 5$, 6 and 7 MeV, forward peaking at $E(^3\text{He}) = 13$ and 17 MeV, and both forwards and backward peaks in the intermediate region (1963MA02). Analysis of the angular distribution of protons leading to the $E_x = 16.6$ MeV level indicates stripping, with $L = 0$ and a large deuteron width (1963MO1K, 1964MO1L). See also (1963WE1B, 1965KA1F, 1965RO1R, 1965YO1D).

14. $^6\text{Li}(\alpha, d)^8\text{Be}$ $Q_m = -1.567$

Table 8.11: ^8Be levels near 16 MeV

E_x^a (MeV \pm keV)	Γ (keV)	Reaction	Reference
16.624 \pm 10	95 \pm 20	$^6\text{Li}(^3\text{He}, \text{p})^8\text{Be}$	(1961ER01)
16.627 \pm 15	105 \pm 30	$^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	(1961ER01)
16.635 \pm 15	96 \pm 20	$^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	(1963DO08)
16.623 \pm 10	95 \pm 20	$^{10}\text{B}(\text{d}, \alpha)^8\text{Be}$	(1961ER01)
16.64 \pm 15		$^7\text{Li}(\text{d}, \text{n})^8\text{Be}$	(1960DI02)
16.67	190	$^7\text{Li}(\text{d}, \text{n})^8\text{Be}$	(1957SL01)
16.4 \pm 200	< 400	$^{12}\text{C}(\gamma, \alpha)^8\text{Be}$	(1955GO59)
16.628 \pm 5	97 \pm 11	mean	
16.930 \pm 10	85 \pm 20	$^6\text{Li}(^3\text{He}, \text{p})^8\text{Be}$	(1961ER01)
16.914 \pm 12	88 \pm 25	$^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	(1961ER01)
16.930 \pm 15	80 \pm 15	$^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	(1963DO08)
16.919 \pm 10	85 \pm 20	$^{10}\text{B}(\text{d}, \alpha)^8\text{Be}$	(1961ER01)
16.9 \pm 50		$^7\text{Li}(\text{d}, \text{n})^8\text{Be}$	(1960DI02)
16.8 \pm 200	< 300	$^{12}\text{C}(\gamma, \alpha)^8\text{Be}$	(1955GO59)
16.923 \pm 6	83 \pm 10	mean	
17.641 \pm 10	< 20	$^6\text{Li}(^3\text{He}, \text{p})^8\text{Be}$	(1961ER01)
17.636 \pm 10	< 15	$^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	(1961ER01)
17.641 \pm 10		$^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	(1963DO08)
17.61	< 20	$^7\text{Li}(\text{d}, \text{n})^8\text{Be}$	(1957SL01)
17.639 \pm 6		mean	

^a Based on listed Q_m .

At $E_\alpha = 43$ MeV, deuteron groups corresponding to ${}^8\text{Be}(0)$ and ${}^8\text{Be}^*(2.9)$ are reported (1959ZE1A); at $E_\alpha = 48$ MeV, an additional group ascribed to ${}^8\text{Be}^*(11.3 \pm 0.4)$ is observed (1962CE01). Angular distributions of d_0 and d_1 show forward peaking, with that of d_0 exhibiting pronounced oscillatory characteristics. Attempts to fit the observations with direct interaction theory are only moderately successful (1959ZE1A, 1962CE01). See also (1959ST30, 1962KO13, 1963BL20, 1963DE1G, 1963DE29, 1963WE1B, 1964DE1K).

15. ${}^6\text{Li}({}^6\text{Li}, \alpha){}^8\text{Be}$ $Q_m = 20.808$

In addition to α -groups to the ground and 2.9 MeV states, there is evidence for a cluster reaction mechanism in which ${}^6\text{Li}$ combines with a deuteron cluster to produce a state of ${}^8\text{Be}$ at 20.95 ± 0.3 MeV, $\Gamma = 3.4$ MeV. At $E({}^6\text{Li}) = 1.9$ MeV, the cross section is 1.3 mb, or 23 times that for ground-state alpha particles (1963KA20). On the other hand (1964QU01) find that the α -particle spectrum is consistent with the involvement of a ${}^8\text{Be}$ level near 22 MeV, with $\Gamma \approx 0.4$ MeV. See also (1962CO05). The ${}^8\text{Be}$ state decays by proton and α emission: $\Gamma_p/\Gamma_\alpha = 1.64$ (1962GA21). At $E({}^6\text{Li}) = 3.2$ MeV, alpha groups corresponding to ${}^8\text{Be}^* = 16.6, 16.9, (17.6), 18.15$ MeV are reported (1964CA1G). See also (1960SH01, 1962CO21, 1963LE19, 1963TA1B, 1964GA1E, 1964SH05, 1965MA2A, 1965NO1A).

16. ${}^7\text{Li}(p, \gamma){}^8\text{Be}$ $E_b = 17.252$

Cross sections and angular distributions have been reported from $E_p = 30$ keV to 14.5 MeV. Two γ -rays are observed, γ_0 to the ground state and γ_1 to the broad, 2^+ , excited state at 2.9 MeV: $E_\gamma = (17.2, 14.3) + \frac{7}{8}E_p$. Resonances for both γ_0 and γ_1 occur at $E_p = 0.44$ and 1.03 MeV, and for γ_1 alone at 2 MeV (see, however, (1963PE15)): see Table 8.12. In the range $E_p = 2.5$ to 9 MeV, broad resonances are reported at $E_p \approx 5$ MeV (γ_0), $\Gamma \approx 5$ MeV (1959GE33, 1963MI08, 1963PE15 ($\Gamma = 5.5$ MeV)), at $E_p \approx 7.3$ MeV (γ_1), $\Gamma \approx 8$ MeV, and possibly at $E_p \approx 6$ MeV (γ_1). The $E_p \approx 5$ MeV resonance ($E_x \approx 21.6$ MeV) is presumed to represent the giant dipole resonance based on ${}^8\text{Be}(0)$, while the γ_1 resonance, 2.0 ± 0.5 (1963MI08), 3 ± 0.5 MeV (1963RE09) higher is similarly related to ${}^8\text{Be}^*(2.9)$. Angular distributions from $E_p = 2.0$ to 7.5 MeV are approximately isotropic (except at 2.59 MeV), suggesting $l = 0$ capture (1963MI08, 1963PE15). The integrated (γ, p) cross section, determined by reciprocity is 22 ± 4 MeV \cdot mb (1963RE09), 20 MeV \cdot mb (1964TA05). See also (1965TA1E).

Angular distributions near the $E_p = 0.44$ MeV resonance show strong $\cos \theta$ interference terms which vanish at resonance. At resonance the radiation is nearly isotropic, consistent with p-wave formation, $J^\pi = 1^+$, with channel spin ratio $\sigma(J_c = 2)/\sigma(J_c = 1) = 5$. A detailed study of the angular distributions of γ_0 and γ_1 yields a small $\cos^2 \theta$ term for γ_0 : $A_2 = 0.067 \pm 0.025$ (1961ME10), $A_2 = 0.028 \pm 0.003$ (1958NE17, 1961BR02), while γ_1 is isotropic: $A_2 = -0.004 \pm 0.008$ (1961ME10); see, however, (1958NE17, 1961BR02). Analysis of these data together with γ - α correlations leads to $\Gamma(E2)/\Gamma(M1)$ for $\gamma_1 = 0.018 \pm 0.009$ (1961ME10), 0.044 ± 0.014 (1960GR21).

Table 8.12: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, \gamma){}^8\text{Be}$

E_{res} (keV)	Γ_{lab} (keV)	${}^8\text{Be}^*$	l_p	J^π	σ_{max} (mb)	γ_0/γ_1	$w\Gamma_\gamma$ (eV)	References
441.4 ± 0.5 ^a	12.2 ± 0.5	17.64	1	1^+	6.0	1.9	9.4	(1949FO18, 1956BU27)
1030 ± 5	168	18.15	1	1^+		0.54	2	(1954KR06, 1960MA33, 1963RI09)
2060 ± 20	310 ± 20	19.05		(-) ^d	γ_1 only			(1957NE1A, 1963PE15, 1963RI09)
5000	≈ 5000	21.6	(0)	(1^-)	0.05 ^b	0.5	^c	(1959GE33, 1963MI08, 1963PE15)
(6000)		(22.5)			(γ_1)	0.3		(1963MI08)
7300	≈ 5500	23.6	(0)	$(1^-, 2^-)$	(γ_1)	0.2		(1963MI08, 1963PE15)

^a See (1959AJ76).

^b γ_0 only (1964TA05).

^c $\int \sigma(p\gamma_0) = 0.37 \pm 0.15 \text{ MeV} \cdot \text{mb}$ (1963RE09).

^d (1964SC19).

With $\Gamma_\gamma = 25 \text{ eV}$, one obtains $\Gamma(\gamma_0) = 16.7 \text{ eV}$, $\Gamma(\gamma_1, \text{M1}) = 8.15 \pm 0.07 \text{ eV}$, $\Gamma(\gamma_1, \text{E2}) = 0.15 \pm 0.07 \text{ eV}$. The observed E2 strength is about 15 times smaller than predicted by intermediate coupling shell model (1961ME10). See (1957KU58). The channel spin mixture $\sigma(2)/\sigma(1)$ is 3.2 ± 0.5 (1961ME10), 4.1 ± 0.1 (1958NE17).

The relative intensity of γ_0/γ_1 at resonance is 2.0 (1961ME10), 2.3 ± 0.04 (1960MA23: $\theta = 90^\circ$), 1.8 ± 0.2 (1963BA58); the ratio falls rapidly above resonance and is approximately constant at 0.54 ± 0.08 for $E_p = 0.8$ to 1.0 MeV (1960MA23: see also (1961BR02)). There is evidence for an increase in the ratio at $E_p = 1.03 \text{ MeV}$ (1964SC19). At $E_p = 5.4, 6.5, 7.5$ and 8.6 MeV , the ratios are $0.5 \pm 0.07, 0.3 \pm 0.04, 0.2 \pm 0.04$ and 0.2 ± 0.04 , respectively (1964TA05). Angular distributions in the range $E_p = 0.2$ to 1.1 MeV indicate interference with s-wave and d-wave non-resonant radiation (1960MA33: see also (1961BR02)); at $E_p = 1.03 \text{ MeV}$ the resonant radiation is isotropic, suggesting p-wave, $J = 1^+$, $\sigma(2)/\sigma(1) = 5$ (1960MA33).

Angular distributions from $E_p = 0.8$ to 1.7 MeV appear to require interference of the $E_p = 0.44$ and $1.03 (1^+)$ MeV levels with at least two odd-parity levels, possibly those at $E_p = 2.1$ and 5.8 MeV (1964SC19).

A pair-spectrometer measurement yields $E_\gamma = 17.647 \pm 0.015 \text{ MeV}$ for γ_0 at the $E_p = 440 \text{ keV}$ resonance: from the spectrum of γ_1 an excitation energy of $E_x = 2.6 \pm 0.15$, $\Gamma = 0.7 \pm 0.1 \text{ MeV}$ is obtained for the first excited state (1963BA58). See Table 8.9.

At $E_p = 440 \text{ keV}$, α -particles from ${}^8\text{Be}^*(16.63)$ are observed, indicating the existence of the γ -transition ${}^8\text{Be}^*(17.64 \rightarrow 16.63)$. The total resonant cross section is $6.45 \pm 0.7 \mu\text{b}$ (1965WI1K). The coincidence spectrum at resonance shows the 1.02 MeV γ -ray ($17.64 \rightarrow 16.63$), and a weak branch $(5.9 \pm 2)\%$ (1965WI07), $(7.5 \pm 2)\%$ (1965KO05), to the 16.92 MeV state. There is no indication of resonance in the yield of α -particles from ${}^8\text{Be}^*(16.6)$ at $E_p = 1.03 \text{ MeV}$ (1965KO05, 1965WI07). Evidence from this and other reactions involving ${}^8\text{Be}^*(16.6)$ and (16.9) indicates that neither has a pure isospin character (1965MA1G). See also (1964PA13).

A careful study of γ - α coincidence spectra at $E_p = 441$ keV reveals only the 2.9 MeV level in the range $E_x = 2$ to 7 MeV; the α -spectra can be matched with a single-level dispersion formula with $E_\lambda = 5.95$ MeV, $\gamma_\alpha^2 = 11.9$ MeV \cdot fm, $\theta^2 = 1.7$, $R = 4.48$ fm (1960GE07). A report of other γ - α groups by (1962CA13) is ascribed to contamination (1964MA25, 1964MI10, 1964PR04, 1964WE1C).

For a review of earlier work, see (1959AJ76). See also (1960SI1D, 1960SI10, 1963FE03, 1963SC1N, 1963TR08)

17. ${}^7\text{Li}(p, n){}^7\text{Be}$

$$Q_m = -1.644$$

$$E_b = 17.252$$

This reaction is widely used as a source of monochromatic neutrons: see summary by (1960GI1A). The threshold is $E_p = 1880.36 \pm 0.22$ keV (1963MA1R: see ${}^7\text{Be}$); a second threshold, for neutrons leading to ${}^7\text{Be}^*(0.43)$, occurs at $E_p = 2376 \pm 2$ keV (1960MA1G), and above $E_p = 7$ MeV, neutrons corresponding to ${}^7\text{Be}^*(4.55)$ may be produced (1963BO06).

Reaction cross sections and angular distributions have been reported by (1948TA16: to 2.55 MeV), (1964BU08: 2.4 to 3.0 MeV; n_1), (1957NE1A, 1958MA07: to 5.5 MeV), (1959GA08: to 3.25 MeV), (1961BE05: 2.6 to 4.2 MeV; n_0 and n_1 groups separately), (1961NI04: 2.7 to 3.5 MeV), (1963BO06: 3 to 13 MeV; n_0, n_1, n_2), (1964BA16: 4 to 14 MeV) and by (1960HI04: 8 to 14 MeV; n_0, n_1, n_2). See also (1959AJ81, 1960GI1A, 1962AU01, 1963PA1K). The yield of ground state neutrons (n_0) rises steeply from threshold and shows pronounced resonances at $E_p = 2.25$ ($\sigma \approx 0.57$ b) and 4.9 MeV ($\sigma \approx 0.36$ b) (1963BO06). The yield of n_1 also rises steeply from threshold (1964BU08) and exhibits a broad maximum near $E_p = 3.5$ MeV (1961BE05).

The behavior of the cross section near the n_0 threshold indicates a broad resonance with $J^\pi = 2^-, T = (0), (l = 0)$ at $E_p = 1.9$ MeV with $\gamma_n^2/\gamma_p^2 \approx 5.0$ (Table 8.13). The structure at $E_p = 2.25$ MeV is ascribed to a $3^+, T = (1), l = 1$ resonance with $\Gamma_n \sim \Gamma_p$ and $\gamma_n^2/\gamma_p^2 = 3$ to 10 (1957NE1A, 1958MA07, 1959WE1A, 1960GI1A, 1961BE05). The broad peak at 4.9 MeV can be fitted by $J^\pi = 3^{(+)}$ with $\Gamma = 1.1$ MeV, $\gamma_n^2 \sim \gamma_p^2$ (1963BO06). An additional resonance at $E_p = 3.0$ MeV may be indicated (1957NE1A, 1961BE05). The behavior of the n_1 cross section can be fitted in terms of a single level with $J^\pi = 1^-$ at $E_p = 3.55$ MeV (1964BU08).

Polarization of the neutrons has been measured by (1961DA04, 1962EL01: 2.0 to 2.4 MeV), (1959CR84: 3.5 to 5 MeV), (1961AU02: 2 to 3 MeV), (1965MO1L: 3.1 to 3.3 MeV), (1965AN09: 3.0 to 4.0 MeV), (1959BA34, 1960BA27: 3 to 6 MeV), (1963MI01, 1963MI20, 1963MI06, 1964MI14: 4 to 5 MeV), (1962BE11: 4 to 10 MeV) and others (1959AJ76, 1963HA1G). Dominant effects are ascribed to the narrow 3^+ level at $E_p = 2.25$ MeV and a broad 2^- background level. At least one level is needed near threshold, and at the higher energies ($E_p > 2.6$ MeV) two additional levels are indicated (1961AU02).

Neutron spectra and polarizations have been measured at $E_p = 143$ MeV by (1962BO1F, 1963BO1N). Some contribution from quasi-free scattering is indicated: see (1963VA1C, 1965VA1E).

18. ${}^7\text{Li}(p, p){}^7\text{Li}$

$$E_b = 17.252$$

Table 8.13: ^8Be levels from $^7\text{Li}(p, n)^7\text{Be}$

E_p (keV)	Γ_{lab} (keV)	$^8\text{Be}^*$	l_p	J^π	σ_{max} (b)	γ_n^2/γ_p^2	References
1900	> 500	18.91	0	2^-		≈ 5.0	(1957NE1A, 1961BE05)
2250	220	19.22	1	3^+	0.57	5.2	(1957NE1A, 1961BE05)
(3000)	> 1000	(19.9)	(1)	(1^+)	(n_0)	(5)	(1957NE1A, 1961BE05)
3550	1300	20.36	(0)	(1^-)	(n_1)	5.2	(1961BE05, 1964BU08)
4900	1100	21.5		$3^{(+)}$	0.36	≈ 1	(1959GI47, 1963BO06)

Absolute differential cross sections are reported for $E_p = 0.4$ to 12 MeV (1953WA27, 1956MA12, 1965GL03) and for $E_p = 14.5$, 20.0 and 31.5 MeV by (1956KI54). Anomalies appear at $E_p = 0.44$, 1.03, 1.88, 2.1, 2.5, 4.2 and 6.0 MeV (see Table 8.14). Both the 0.44 and 1.03 MeV resonances are ascribed to p-waves, $J = 1^+$, with channel spins 1 and 2 in a ratio of 1 to 5 (1953CH1A, 1955LI1B: compare $^7\text{Li}(p, \gamma)^8\text{Be}$). The structure at $E_p = 2.1$ MeV may indicate a state with $J^\pi \leq 3^+$ (1956MA12), 3^- (1957NE1A). The $E_p = 4.2$ MeV resonance is ascribed to an almost pure single-particle level in ^8Be at 20.9 MeV (1965GL03).

The polarization of 14.5 and 40 MeV protons scattered from ^7Li has been studied by (1962RO20) and (1963HW01). See also (1961RO20).

See also (1961JO17, 1961JO18, 1965AN12) and (1959AJ76).

 Table 8.14: ^8Be levels from $^7\text{Li}(p, p)^7\text{Li}$ and $^7\text{Li}(p, p')^7\text{Li}^*$

E_p (keV)	Γ_{lab} (keV)	$^8\text{Be}^*$	l_p	J^π	$\Gamma_{p'}$ (keV)	References
441	12.2 ^b	17.64	1	1^+		(1953CH1A, 1953WA27)
1030 \pm 5	168	18.15	1	1^+	≈ 6	(1954MO04, 1955LI1B)
1880 ^a						(1956MA12, 1957NE1A)
2100	≈ 0.4	19.0		(3) ^c	small	(1956MA12, 1957NE1A)
2250		19.22		3^+	small	(1956MA12, 1957NE1A)
4200 \pm 200	1800 \pm 200	20.9		4^-	(res)	(1965GL03)
5600		22.2				(1965GL03)

^a (p, n) threshold.

^b $\theta_p^2 = 0.064$.

^c (1957NE1A: 3^-), (1956MA12: ≤ 3).

19. ${}^7\text{Li}(p, p'){}^7\text{Li}^*$ $E_b = 17.252$

A pronounced resonance appears in the yield of inelastically scattered protons (1951BR10, 1954MO04) and 0.48 MeV γ -rays (1954KR06) at $E_p = 1.03$ MeV (see Table 8.14); it is an s- or p-wave resonance interfering with a non-resonant wave of opposite parity (1954MO04: see also (1955LI1B)). Excitation functions for the proton group to the 0.48 MeV state have been measured for $E_p = 2.3$ to 12 MeV: a peak is observed at 5.6 MeV (1965GL03). See also (1964FA08). The yield of 0.48 MeV γ -rays rises smoothly from $E_p = 1.5$ to 3.0 MeV except for a pronounced cusp at 1.881 MeV (1955HA34, 1957NE1A). See also (1962CA13).

20. ${}^7\text{Li}(p, d){}^6\text{Li}$ $Q_m = -5.028$ $E_b = 17.252$ See ${}^6\text{Li}$.21. ${}^7\text{Li}(p, t){}^5\text{Li}$ $Q_m = -4.433$ $E_b = 17.252$ See ${}^5\text{Li}$.22. ${}^7\text{Li}(p, \alpha){}^4\text{He}$ $Q_m = 17.347$ $E_b = 17.252$

Excitation functions and angular distributions have been reported by (1962LU01: 0.4 to 3.3 MeV), (1948HE1A: 0.5 to 3.8 MeV), (1958CO62: 0.1 to 0.6 MeV), (1963SA10: 3.0 to 5.5 MeV), (1962CA12: 1.5 to 4.8 MeV), (1960AL18: 12 MeV), (1961HA27: 2.8 to 12 MeV), (1962TE04, 1962TE07: 3.3 to 6.6 MeV), (1964MA51: 4 to 12 MeV), (1962MA40: 15, 18.6 MeV) and (1962CA13, 1963CA1K: 0.5 to 2.5 MeV; see, however, (1964MA25, 1964MI10)). Polarization effects have been studied by (1964AN08: 0.5 to 2.0 MeV), (1962BE14: 1 to 3.2 MeV), (1964AS04: 2 to 3.5 MeV), (1965BO07: 3.2 to 5.3 MeV).

A broad resonance occurs at $E_p = 3.0$ MeV, $\Gamma \approx 1$ MeV, $\sigma_{\max} \approx 90$ mb (1948HE1A): see Table 8.15. A second prominent peak appears at $E_p = 5.6$ MeV, $\Gamma \approx 1$ MeV (1961HA27, 1962TE04, 1964MA51). Some structure is reported near $E_p = 6.0$ to 6.5 MeV, and a further peak occurs at $E_p = 9.0$ MeV (1964MA51).

An earlier suggestion that a $J = 0$ resonance near $E_p \approx 0$ is involved appears to be contradicted by polarization data (1962BE14). The states at $E_x = 19.9$ MeV and 22.1 MeV ($E_p = 3.0$ and 5.5 MeV resonances) appear to have $J = 2^+$ (1959AJ76, 1961HA27, 1962TE04, 1962TE07, 1965FR02) and to have ${}^6\text{Li} + d$ parentage (1962TE01, 1965FR02). Levels at $E_x = 23$ MeV ($E_p = 6.5$ MeV) and 25.2 MeV ($E_p = 9.0$ MeV) are assigned $J^\pi = 4^+$ and 2^+ , respectively (G. Temmer, private communication).

See also (1960BO13, 1962PE20, 1962WE13, 1963AN09, 1964MA2D).

Table 8.15: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, \alpha){}^4\text{He}$

E_{res} (MeV)	Γ (MeV)	${}^8\text{Be}^*$	$J^\pi; T$	References
3.0	≈ 1	19.9	$2^+; 0$	(1948HE1A)
5.6	≈ 1	22.1 ^a	(2^+)	(1961HA27, 1962TE04, 1964MA51)
6.5		23	(4^+)	(1961HA27, 1964MA51)
9.0		25.1	(2^+)	(1964MA51)

^a Possibly same as $E_x = 22.5$ seen in ${}^6\text{Li}(d, \alpha)$: see (1965FR02).

 Table 8.16: Slow neutron thresholds in ${}^7\text{Li}(d, n){}^8\text{Be}$ (1957SL01)

E_d (MeV)	$\Gamma_{\text{c.m.}}$ (MeV)	${}^8\text{Be}^*$ (MeV)
1.35	0.31	(16.08)
2.10	0.19	16.66
3.32	< 0.02	17.61
4.08	0.23	18.20

23. ${}^7\text{Li}(d, n){}^8\text{Be}$

$$Q_m = 15.028$$

For $E_d = 3.6$ to 7.3 MeV, neutron time-of-flight spectra indicate states in ${}^8\text{Be}$ at 0, 2.9, 16.64 ± 0.015 , 16.9 ± 0.05 , 17.64 and 18.15 MeV. Angular distribution of neutrons corresponding to the 16.6, 17.6 and 18.15 MeV states are strongly forward; $l_n = 1$ for the 16.6 MeV state, consistent with $J = 2^+$ (1960DI02). At $E_d = 2$ MeV, recoil proton spectra show only the ground state and $E_x = 2.9$ MeV groups; in the range $E_x < 9$ MeV, no others appear with intensity $> 10\%$ of the ground-state group. The spectrum yields $E_x = 3.1 \pm 0.1$, $\Gamma = 1.75 \pm 0.1$ MeV; comparison with the shape calculated from known α - α phase shifts suggest a considerable contribution of three-body processes (1964JO04). Reported slow neutron thresholds are listed in Table 8.16. See also (1964JO1F, 1964JO1D, 1965DI1F).

Alpha-particle spectra indicate groups corresponding to the break up of ${}^8\text{Be}^*(16.6)$ and (16.9) (1963BI22, 1963PA04, 1964GL03). Alpha particles attributed to ${}^8\text{Be}^*(11.4)$ are also reported (1963PA04). In the range $E_x = 15.0$ to 18.5 MeV, no α -emitting states other than the 16.6 and 16.9 MeV levels are observed (1964GL03). See also (1965BI1F, 1965JO19, 1965MA1G, 1965MA1K).

A large number of additional states with $E_x < 16.1$ MeV has been suggested in addition to ${}^8\text{Be}^*(0, 2.9, 11.4)$: for a listing of early references, see (1959AJ76, 1964JO04). See also (1958CA1E, 1960AG03, 1960JU04, 1963MI1M, 1964MI1J).

24. ${}^7\text{Li}({}^3\text{He}, \text{d}){}^8\text{Be}$ $Q_m = 11.759$

At $E({}^3\text{He}) = 24.3$ MeV, angular distributions of deuterons corresponding to the ground and first excited states of ${}^8\text{Be}$ have been measured (1963WE1B). At $E({}^3\text{He}) = 9$ to 10 MeV, excitation of ${}^8\text{Be}^*(17.6)$ is observed (1965GR08). See also (1964DU1E).

25. ${}^7\text{Li}(\alpha, \text{t}){}^8\text{Be}$ $Q_m = -2.562$

The angular distributions of the tritons to the ground state of ${}^8\text{Be}$ have been determined at a number of energies up to 48 MeV; the distributions indicate strong contribution of direct interaction (1959ST1B, 1960GO04, 1960MA15, 1960VL03, 1962MA59, 1963WE1B). At $E_\alpha = 40$ MeV, higher states are also observed (1960VL03).

26. ${}^7\text{Li}({}^7\text{Li}, {}^6\text{He}){}^8\text{Be}$ $Q_m = 7.272$

See (1957NO17, 1964CA16).

27. (a) ${}^7\text{Be}(\text{n}, \text{p}){}^7\text{Li}$ $Q_m = 1.644$ $E_b = 18.896$
 (b) ${}^7\text{Be}(\text{n}, \alpha){}^4\text{He}$ $Q_m = 18.991$
 (c) ${}^7\text{Be}(\text{n}, \gamma\alpha){}^4\text{He}$

At thermal energies, the (n, p) cross section is $(5.1 \pm 0.6) \times 10^4$ b (1955HA34), the (n, α) cross section is ≤ 0.1 mb (1962BA1B, 1963BA34) and the (n, $\gamma\alpha$) cross section is 155 mb (1963BA34). These observations are consistent with odd parity of ${}^7\text{Be}$; the small value of $\sigma(\text{n}, \alpha)$ leads to an estimated $F^2 < 4 \times 10^{-10}$ for the intensity of a positive parity admixture (1963BA34). Comparison of the thermal cross section for reaction (a) with the (p, n) cross section observed in the inverse reaction supports the assignment $J = \frac{3}{2}$ for ${}^7\text{Be}(0)$ (1957NE1A). See also (1959AJ76).

28. ${}^7\text{Be}(\text{d}, \text{p}){}^8\text{Be}$ $Q_m = 16.672$

For $E_d = 0.8$ to 1.7 MeV, proton groups are observed corresponding to the ground state and the 2.9 MeV level: $\Gamma(2.9) = 1.6 \pm 0.4$ (1959SP1A), 1.53 ± 0.04 MeV (1960KA17). A dispersion formula fit yields parameters $E_k = 2.90 \pm 0.06$ MeV, $\gamma^2 = 0.60$ MeV, $\theta^2 = 0.64$, $R = 5.75$ fm (1960KA17): see Table 8.9.

Table 8.17: α - β angular correlation coefficients^a in ${}^8\text{Li}$, ${}^8\text{B}$

Nuclide	A/W_β	B/W_β	W_β (MeV)	Reference
${}^8\text{Li}$		$(5.7^{+2.9}_{-1.9}) \times 10^{-3}$	7.0	(1960KR03)
${}^8\text{Li}$	$(-8.7 \pm 0.7) \times 10^{-3}$	$(+3.16 \pm 0.6) \times 10^{-3}$	11	(1962NO02)
${}^8\text{Li}$	$(-8.3 \pm 1.1) \times 10^{-3}$	$(+3.7 \pm 1.0) \times 10^{-3}$	7.5	(1963GR11)
${}^8\text{B}$	$(-8.7 \pm 0.9) \times 10^{-3}$	$(-3.86 \pm 1.0) \times 10^{-3}$	11	(1962NO02)

^a $W(\theta) = 1 + A \cos \theta + B \cos^2 \theta$.

 29. ${}^8\text{Li}(\beta^-){}^8\text{Be}$

$Q_m = 16.002$

${}^8\text{Li}$ decays mainly to the broad 2.9 MeV, 2^+ level of ${}^8\text{Be}$, which decays into two α -particles. The β -spectrum, which extends to $\gtrsim 14$ MeV, has been studied by (1950HO01), (1960FA02) and others, while the α -spectrum, extending to ≈ 10 MeV, has been reported by (1938RU01, 1948BO20, 1955FR29, 1960BI10, 1960BI06, 1960FA04, 1961DE1E) and others: see (1955AJ61). The two spectra correspond well, except perhaps for low β -energies (1960GR10). For $E_\alpha \lesssim 5$ MeV, the spectrum is closely matched by a density-of-states derived from the experimental d-wave phase shift of α - α scattering; at higher energies, an increasing excess of α -particles appears which may reflect transition into the tail of the $J^\pi = 2^+$ level at $E_x = 16.67$ MeV (1960GR10, 1963AL19, 1963DA05). See also ${}^8\text{B}(\beta^+)$.

Studies of the distribution of recoil momenta and neutrino-recoil correlation indicate that the decay is at least 90% GT, and that the GT portion is at least 90% axial vector. The observations indicate $J = 2^+$ for ${}^8\text{Li}$ (1958BA1E, 1958LA08, 1959LA11, 1959LA05).

The angular correlations $W(\theta_{\beta\alpha})$ have been measured for the decays of ${}^8\text{Li}$ and ${}^8\text{B}$ as a test of the conserved vector current theory of β -decay. The observed correlation is of the form $W(\theta) = 1 + A \cos \theta + B \cos^2 \theta$; reported values of A and B are listed in Table 8.17. With shell model calculations of the matrix element $\Gamma_\gamma(\text{M1})$ for ${}^8\text{Be}^*(T = 1) \rightarrow {}^8\text{Be}^*(2.9 \text{ MeV})$, the CVC theory predicts $\delta \equiv B({}^8\text{Li}) - B({}^8\text{B}) = (7 \pm 2) \times 10^{-3} W_\beta$ (1960KU05, 1960WE1A); conversely, if CVC is assumed correct, the experiment yields $\Gamma(\text{M1}) = 1.9 \pm 0.6 \text{ eV}$ (1962NO02). See also (1963IS04).

See also (1958KE1B, 1959JA1B, 1960NO01, 1960NO05, 1960ST23, 1961FO1C, 1963DA05, 1963HU1G, 1963LE1K, 1964TO1D).

 30. ${}^8\text{B}(\beta^+){}^8\text{Be}$

$Q_m = 17.979$

The decay proceeds mainly to ${}^8\text{Be}^*(2.9)$. The α and β spectra correspond closely to those of ${}^8\text{Li}$, when account is taken of the different Q -values (1960FA04). The ratio $ft({}^8\text{B})/ft({}^8\text{Li}) = 1.096$

(1964TO1D). Detailed study of the high energy portion of the α -spectrum reveals a maximum near $E_\alpha = 8.3$ MeV, corresponding to transitions to ${}^8\text{Be}^*(16.62)$. The observed shape is reproduced with a Breit-Wigner density-of-states function with parameters $E_x = 16.67$ MeV, $\Gamma = 150$ to 190 keV; parameters $E_x = 16.62$, $\Gamma = 95$ keV are less good, but acceptable. With the former set, $\log ft = 2.9$ (using $\tau_{1/2} = 0.774 \pm 0.005$ sec); with the latter, $\log ft = 3.03$. In either case, the low ft -value supports the identification $T = 1$, $J = 2^+$ for ${}^8\text{Be}^*(16.63)$ (1964MA35). See, however, (1965MA1G).

See also (1960FA02, 1960GR10, 1963DI17) and see ${}^8\text{Li}(\beta^-){}^8\text{Be}$ for a discussion of the work on β - α angular correlations.

31. (a) ${}^9\text{Be}(\gamma, n){}^8\text{Be}$ $Q_m = -1.665$
 (b) ${}^9\text{Be}(n, 2n){}^8\text{Be}$ $Q_m = -1.665$
 (c) ${}^9\text{Be}(\alpha, \alpha n){}^8\text{Be}$ $Q_m = -1.665$

For reaction (a), see ${}^9\text{Be}$. The (n, 2n) reaction appears to proceed largely via excited states of ${}^9\text{Be}$, with subsequent decay to ${}^8\text{Be}$, mainly ${}^8\text{Be}^*(2.9)$ (1959CH1E, 1961MY01, 1963JE05, 1964BO31): see ${}^9\text{Be}$ and ${}^{10}\text{Be}$. For reaction (c), see (1962ST12) and ${}^9\text{Be}$. See also (1959WI41, 1960KO1G, 1961JE01, 1962CU05, 1965LO1K).

32. ${}^9\text{Be}(p, d){}^8\text{Be}$ $Q_m = 0.559$

At $E_p = 5.2$ MeV, the α -spectrum shows a sharp ground-state group α_0 , and a broad group, α_1 , corresponding to ${}^8\text{Be}^*(2.9)$. A pronounced anomaly appears at $E_x \approx 0.75$ MeV, on the side of the α_1 group, which is apparently a “ghost” of α_0 . A quantitative account is given with a density-of-states function derived from α - α scattering phase shifts (1961BE1E: see also (1962BA1C, 1962HA23, 1963AL19)).

At $E_p = 95$ to 155 MeV, ${}^8\text{Be}$ states at 0, 2.9, 11.4, 16.6, 18.8 and (23) MeV are excited (1956SE1A, 1963BA1R, 1963RA01). See also (1959AJ76, 1959FI1B, 1960PH1B, 1964SH07, 1964TO1D, 1964YA1A), ${}^9\text{Be}$ and ${}^{10}\text{B}$.

33. (a) ${}^9\text{Be}(d, t){}^8\text{Be}$ $Q_m = 4.592$
 (b) ${}^9\text{Be}(d, t){}^4\text{He} + {}^4\text{He}$ $Q_m = 4.687$

Angular distributions of ground-state tritons have been measured at a number of energies up to 20 MeV: see (1959AJ76) and (1959VL24, 1960NE09, 1960NE11, 1961RE03, 1962BI11, 1965JA07). The width of the 2.9 MeV state is 0.8 MeV (1955CU16), 1.35 ± 0.15 MeV (1959VL24). See also (1959FI1B, 1959KU1C, 1959ZA01, 1962HA23, 1963OG1A).

34. ${}^9\text{Be}({}^3\text{He}, \alpha){}^8\text{Be}$ $Q_m = 18.913$

At $E({}^3\text{He}) = 3.0$ and 4.0 MeV, angular distributions of the α -particles to the ground-state of ${}^8\text{Be}$ and to the levels at 2.9, 16.6, 16.9 and 17.6 MeV have been measured. The excitation energy and width of the first excited state are found to be 2.90 ± 0.04 MeV and 1.35 ± 0.15 MeV (1963DO08): see Table 8.9. The parameters of the higher states (from (1961ER01) and (1963DO08)) are given in Table 8.11. The angular distributions are amenable to analysis by direct interaction except in the case of the 16.6 MeV state which appears to involve compound nucleus formation (1963DO08) (compare ${}^7\text{Li}(d, n){}^8\text{Be}$). For angular distribution data at lower energies, see (1955AJ61) and (1963WE08).

The α -particles corresponding to the 2.9 MeV excited state are superposed on a broad, intense continuum extending to $E_x > 16$ MeV. Some part of this spectrum may be ascribed to a broad level at $E_x = 12.5$ MeV, with $\Gamma \approx 5$ MeV, but the main contribution appears to be due to a direct 3α breakup (1963DO11: see also (1963WE08)). (1964MO19) find, on the other hand, that the sequential decay, via ${}^8\text{Be}^*(2.9)$ and (16.92) dominates the reaction. See, however, (1965MO1M). Angular correlation studies confirm $J^\pi = 2^+$ for ${}^8\text{Be}^*(16.9)$ (1964MO19, 1965MO1N).

See also (1962WE1C, 1964BL12, 1965MA1G).

35. (a) ${}^9\text{Be}({}^6\text{Li}, {}^7\text{Li}){}^8\text{Be}$ $Q_m = 5.587$
 (b) ${}^9\text{Be}({}^7\text{Li}, {}^8\text{Li}){}^8\text{Be}$ $Q_m = 0.367$

For reaction (a) see (1960MA1H, 1961LE1K). For reaction (b) see ${}^9\text{Be}$.

36. ${}^{10}\text{B}(\gamma, d){}^8\text{Be}$ $Q_m = -6.028$

See ${}^{10}\text{B}$ and (1955AJ61).

37. ${}^{10}\text{B}(n, t){}^8\text{Be}$ $Q_m = 0.229$

Angular distributions of the tritons to the ground and 2.9 MeV states of ${}^8\text{Be}$ have been measured at $E_n = 14.4$ MeV (1964VA14). See also (1959AJ76, 1964VA1E) and ${}^{11}\text{B}$.

38. (a) ${}^{10}\text{B}(p, {}^3\text{He}){}^8\text{Be}$ $Q_m = -0.534$
 (b) ${}^{10}\text{B}(p, pd){}^8\text{Be}$ $Q_m = -6.028$

See ^{11}C and (1964BA1C).

39. (a) $^{10}\text{B}(\text{d}, \alpha)^8\text{Be}$ $Q_{\text{m}} = 17.819$
 (b) $^{10}\text{B}(\text{d}, \alpha)^4\text{He} + ^4\text{He}$ $Q_{\text{m}} = 17.914$

Alpha groups are reported corresponding to ^8Be states at 0, 2.9 (1960BE15, 1960BI11, 1961CI02, 1961LE10, 1963PU02, 1964YA1A; and see (1959AJ76)), 16.6 and 16.8 MeV (1961ER01: see Table 8.11). A coincidence study of the α_1 group yields the following parameters for the level shape: $E_{\text{res}} = 3.15$ MeV, $\Gamma = 2.04$ MeV, $E_{\lambda} = 4.00$ MeV, $\gamma^2 = 3.4$ MeV \cdot fm, $\theta^2 = 0.60$, $R = 5.5$ fm (1963PU02: E relative to $E(2\alpha)$). (1953TR04) reports $E_{\lambda} = 5.29$ MeV (relative to 2α), $\gamma^2 = 13.4$ MeV \cdot fm, $\theta^2 \approx 2$, $R = 4.48$ fm: see Table 8.9.

Arguments are presented for the assignments $J = 2^+$, $T = 1$ for $E_{\text{x}} = 16.63$ and $J^{\pi} = 0^+$ or 2^+ , $T = 0$ for $E_{\text{x}} = 16.93$ MeV. Comparison of yields with $^6\text{Li}(^3\text{He}, \text{p})^8\text{Be}^*$ indicates gross violation of the isospin selection rule, possibly to be ascribed to intermixture in the compound nucleus stage (1961ER01). See, however, (1965MA1G). There is no evidence for an earlier reported level at $E_{\text{x}} = 16.08$ MeV (1961ER01). See also (1961TE02, 1962TE02, 1965LE09).

40. $^{10}\text{B}(^3\text{He}, \text{p}\alpha)^8\text{Be}$ $Q_{\text{m}} = 12.326$

See (1964ET02, 1965AL1B, 1965ET1A, 1965WA1M) and ^{12}C .

41. (a) $^{11}\text{B}(\gamma, \text{t})^8\text{Be}$ $Q_{\text{m}} = -11.226$
 (b) $^{11}\text{B}(\gamma, \text{t})^4\text{He} + ^4\text{He}$ $Q_{\text{m}} = -11.132$

See ^{11}B .

42. $^{11}\text{B}(\text{p}, \alpha)^8\text{Be}$ $Q_{\text{m}} = 8.588$

Alpha groups are reported corresponding to the ground and 2.9 MeV state: see (1959AJ76) and (1962AL20). At $E_{\text{p}} = 3$ to 5 MeV, the α -spectrum shows an anomaly at $E_{\text{x}} \approx 0.75$ MeV, ascribed to successive two-body decays with a final density-of-states related to α - α scattering phase shifts (1961BE1E: see $^9\text{Be}(\text{p}, \text{d})^8\text{Be}$ and (1963BR03, 1964BR05, 1964DE1H)). See also ^{12}C and (1959KA12, 1959KA13, 1961KO08, 1962BE21, 1963PH1A, 1964BA1C, 1964YA1A, 1965DU1C, 1965LE09, 1965PH1A, 1965SW1B).

43. $^{11}\text{B}(\text{d}, \text{n}\alpha)^8\text{Be}$ $Q_{\text{m}} = 6.363$

See (1965OL01).

44. $^{11}\text{B}(^3\text{He}, ^6\text{Li})^8\text{Be}$ $Q_{\text{m}} = 4.566$

At $E(^3\text{He}) = 3.0$ MeV, $^8\text{Be}^*(2.9)$ is produced with an intensity 20 times that of the ground state (1964YO06).

45. (a) $^{12}\text{C}(\gamma, \alpha)^8\text{Be}$ $Q_{\text{m}} = -7.369$

(b) $^{12}\text{C}(\alpha, 2\alpha)^8\text{Be}$

(c) $^{12}\text{C}(\text{p}, \text{p}\alpha)^8\text{Be}$

(d) $^{12}\text{C}(\text{n}, \text{n}\alpha)^8\text{Be}$

Reaction (a) involves states of ^8Be at 0, 2.9, 16.6, 16.9 and (17.6) MeV: see (1955GO59, 1959AJ76, 1964TO1A). See also (1961SE13, 1963SH04, 1964LE1C).

At $E_{\alpha} = 25.4$ MeV, the four-body breakup ($^{12}\text{C} + \alpha \rightarrow 4\alpha$) is greatly favored over reaction (b) (1962BR14: see also (1961VA38, 1962VA25)). See also (1961GO1T, 1962IG1B, 1963LA02).

Reaction (c) at $E_{\text{p}} = 15$ to 29 MeV proceeds predominantly through the ground state and the 2.9 MeV level. It is not clear whether higher levels of ^8Be are involved (1955NE18, 1960VA10, 1962VA1A, 1963VA04). See also (1963JA07, 1964BA1C, 1964KE1F, 1964SY02, 1965YU1C, 1965YU1D).

For reaction (d) see (1955FR35, 1960VA10, 1964BR25).

See also ^{12}C .

46. $^{12}\text{C}(\text{d}, ^6\text{Li})^8\text{Be}$ $Q_{\text{m}} = -5.897$

See (1963DR1B, 1964BL1C, 1964DA1B, 1965BE1W, 1965DE1V, 1965SL1C).

47. $^{12}\text{C}(^3\text{He}, ^7\text{Be})^8\text{Be}$ $Q_{\text{m}} = -5.782$

See (1964PA1K, 1965MA1V).

48. $^{12}\text{C}(^{12}\text{C}, ^6\text{He})$ $Q_{\text{m}} = -14.549$

See (1962BE43, 1963BE1U).

$$49. \text{}^{12}\text{C}(\text{}^{14}\text{N}, \text{}^{18}\text{F})\text{}^8\text{Be} \quad Q_m = -2.953$$

See (1965WI1A).

$$50. (a) \text{}^{16}\text{O}(\gamma, \alpha)\text{}^{12}\text{C}^* \rightarrow \text{}^8\text{Be} + \text{}^4\text{He} \quad Q_m = -14.530$$

$$(b) \text{}^{16}\text{O}(\alpha, \text{}^8\text{Be})\text{}^{12}\text{C} \quad Q_m = -7.256$$

For reaction (a) see (1955AJ61, 1959AJ76, 1964TO1E) and ^{12}C .

For reaction (b) see (1964BR1U) and ^{12}C .

$$51. \text{}^{16}\text{O}(\text{p}, \text{p}')\text{}^4\text{He}\text{}^4\text{He}\text{}^4\text{He}\text{}^4\text{He} \quad Q_m = -14.436$$

See (1962VA1A) and ^{16}O .

$$52. \text{}^{20}\text{Ne}(\alpha, \text{}^{16}\text{O})\text{}^8\text{Be} \quad Q_m = -4.824$$

This reaction has not been observed: see (1962LA15).

Table 8.18: Energy levels of ${}^8\text{B}$

E_x (MeV \pm keV)	$J^\pi; T$	Γ (keV)	Decay	Reactions
g.s.	$(2^+); 1$	$\tau_{1/2} = 774 \pm 4$ msec	β^+	1, 2
0.778 ± 7		42 ± 10		2
2.17 ± 50				2

${}^8\text{B}$

(Figs. 13 and 14)

GENERAL: See (1958FO1C, 1960TA1C, 1962IN02, 1963NA1E, 1964BA2A, 1964BA1X, 1964GR1J, 1964ST1B).

1. ${}^8\text{B}(\beta^+){}^8\text{Be}$ $Q_m = 17.979$

The beta decay leads mainly to ${}^8\text{Be}^*(2.9)$. Reported half lives are listed in Table 8.19; taking $\tau_{1/2} = 0.774$ sec and $Q = 17.979 - 2.90$, $ft = 4.41 \times 10^5$ (1966BA1A). There is also observed a branch to a ${}^8\text{Be}$ state at ≈ 16.6 MeV; $\log ft = 2.9$ (1964MA35). See also (1964BA1X) and ${}^8\text{Be}$.

2. ${}^6\text{Li}({}^3\text{He}, n){}^8\text{B}$ $Q_m = -1.975$

Thresholds for production of slow neutrons are reported at $E({}^3\text{He}) = 2.9661 \pm 0.0017$ MeV (1958DU78), 2.974 ± 0.010 MeV (1959FA02), corresponding to the ground state of ${}^8\text{B}$, and at $E({}^3\text{He}) = 4.16 \pm 0.05$ (1958DU78), 4.159 ± 0.030 MeV (1959FA02), corresponding to an excited state at $E_x = 0.789 \pm 0.030$ MeV with $\Gamma_{\text{c.m.}} = 70 \pm 40$ keV. A threshold at 3.9 MeV reported by (1958DU78) is not confirmed by (1959FA02).

Neutron time-of-flight spectra at $E({}^3\text{He}) = 4$ to 9 MeV show groups corresponding to ${}^8\text{B}(0)$ and ${}^8\text{B}^*(0.8)$, with forward-peaked angular distributions, and to a new state at $E_x = 2.17 \pm 0.05$ MeV (1963DI02). At $E({}^3\text{He}) = 4.5$ to 5.3 MeV, time-of-flight spectra locate the first excited state at 0.767 ± 0.012 (1965FA03), 0.783 ± 0.010 MeV (1965MC06: $\Gamma = 40 \pm 10$ keV). See also (1963MO1J).

3. ${}^7\text{Be}(p, \gamma){}^8\text{B}$ $Q_m = 0.135$

Table 8.19: Half life of ^8B

$\tau_{1/2}$ (sec)	Reference
0.65 ± 0.1	(1950AL57)
0.61 ± 0.11	(1952SH44)
0.78 ± 0.01	(1958DU78)
0.75 ± 0.02	(1958VE20)
0.79 ± 0.02	(1959FA02)
0.774 ± 0.005	(1964MA35)
0.774 ± 0.004	mean

The cross section for proton capture is $0.48 \pm 0.18 \mu\text{b}$ at $E_p = 0.80 \text{ MeV}$ and $0.50 \pm 0.20 \mu\text{b}$ at $E_p = 1.40 \text{ MeV}$ (1960KA05). The relevance of this reaction in stellar energy production is discussed by (1958FO1C, 1960KA05, 1963BA2D, 1964BA2A, 1964PA1A). See also (1961CH1C).

4. (a) $^{10}\text{B}(p, t)^8\text{B}$ $Q_m = -18.532$
 (b) $^9\text{Be}(p, 2n)^8\text{B}$ $Q_m = -20.426$
 (c) $^{12}\text{C}(p, n\alpha)^8\text{B}$ $Q_m = -26.130$

See (1950AL57).

5. (a) $^{10}\text{B}(\gamma, 2n)^8\text{B}$ $Q_m = -27.014$
 (b) $^{11}\text{B}(\gamma, 3n)^8\text{B}$ $Q_m = -38.470$
 (c) $^{12}\text{C}(\gamma, p3n)^8\text{B}$ $Q_m = -54.426$

See (1952SH44).

References

(Closed July 01, 1965)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1938RU01 L.H. Rumbaugh, R.B. Roberts and L.R. Hafstad, Phys. Rev. 54 (1938) 657
1947HU06 D.J. Hughes, D. Hall, C. Eggler and E. Goldfarb, Phys. Rev. 72 (1947) 646
1948BO20 T.W. Bonner, J.E. Evans, C.W. Malich and J.R. Risser, Phys. Rev. 73 (1948) 885
1948HE1A Heydenburg, Hudson, Inglis and Whitehead, Phys. Rev. 74 (1948) 405
1948TA16 R. Taschek and A. Hemmendinger, Phys. Rev. 74 (1948) 373
1949FO18 W.A. Fowler and C.C. Lauritsen, Phys. Rev. 76 (1949) 314
1950AL57 L.W. Alvarez, Phys. Rev. 80 (1950) 519
1950HO01 W.F. Hornyak and T. Lauritsen, Phys. Rev. 77 (1950) 160
1950WH02 W. Whaling and T.W. Bonner, Phys. Rev. 79 (1950) 258
1951BR10 A.B. Brown, C.W. Snyder, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 82 (1951) 159
1951BR92 C.H. Braden, S.M. Carter and A.G. Ford, Phys. Rev. 84 (1951) 837
1951GR45 E. Graves, Phys. Rev. 84 (1951) 1250
1951MA1B Mather, Phys. Rev. 82 (1951) 126
1951RA12 W. Rall and K.G. McNeill, Phys. Rev. 83 (1951) 1244
1952BA1A Baggett and Bame, Phys. Rev. 85 (1952) 434
1952GR1A Graves, Phys. Rev. 85 (1952) 703A
1952MO19 C.D. Moak, W.M. Good and W.E. Kunz, Phys. Rev. 85 (1952) 928
1952PE02 T.P. Pepper, K.W. Allen, E. Almqvist and J.T. Dewan, Phys. Rev. 85 (1952) 155
1952SH44 R.K. Sheline, Phys. Rev. 87 (1952) 557
1952WA31 D. Walker, W.T. Link and W.I.B. Smith, Proc. Phys. Soc. (London) A65 (1952) 861
1953BA04 M.E. Battat and F.L. Ribe, Phys. Rev. 89 (1953) 80
1953BR1B Bretonneau, Compt. Rend. 236 (1953) 913
1953BU35 D.S. Bunbury, Phys. Rev. 90 (1953) 1121; Erratum Phys. Rev. 91 (1953) 1580
1953CH1A Christy, Phys. Rev. 89 (1953) 839
1953SA1A Sawyer and Phillips, Los Alamos Rept.1578 (1953)

1953ST52 F.E. Steigert and M.B. Sampson, Phys. Rev. 92 (1953) 660
 1953TR04 P.B. Treacy, Phil. Mag. 44 (1953) 325
 1953WA27 W.D. Warters, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 91 (1953) 917
 1954AL35 K.W. Allen, E. Almqvist, J.T. Dewan and T. Pepper, Phys. Rev. 96 (1954) 684
 1954HI34 F. Hirst, I. Johnstone and M.J. Poole, Phil. Mag. 45 (1954) 762
 1954KL36 R.M. Kline and D.J. Zaffarano, Phys. Rev. 96 (1954) 1620
 1954KR06 A.A. Kraus, Jr., Phys. Rev. 93 (1954) 1308
 1954MO04 F. Mozer, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 93 (1954) 829
 1954SI07 R.M. Sinclair, Phys. Rev. 93 (1954) 1082
 1954WI25 M.M. Winn, Proc. Phys. Soc. (London) A67 (1954) 946
 1955AJ61 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 27 (1955) 77
 1955CU16 P. Cuer and J. Jung, J. Phys. Rad. 16 (1955) 385
 1955CU1A Cuer, Magnac-Valette and Baumann, Compt. Rend. 240 (1955) 1880
 1955FR10 J.M. Freeman, A.M. Lanem and B. Rose, Phil. Mag. 46 (1955) 17
 1955FR29 R.T. Frost and S.S. Hanna, Phys. Rev. 99 (1955) 8
 1955FR35 G.M. Frye, Jr., L. Rosen and L. Stewart, Phys. Rev. 99 (1955) 1375
 1955GO59 F.K. Goward and J.J. Wilkins, Proc. Roy. Soc. 228 (1955) 376
 1955HA34 R.C. Hanna, Phil. Mag. 46 (1955) 381
 1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252
 1955LA1D Lane, Proc. Phys. Soc. (London) A68 (1955) 189
 1955LE24 S.H. Levine, R.S. Bender and J.N. McGruer, Phys. Rev. 97 (1955) 1249
 1955LI1B Liberman, Thesis, CalTech (1955)
 1955MA20 R.L. Macklin and H.E. Banta, Phys. Rev. 97 (1955) 753
 1955NE18 J.L. Need, Phys. Rev. 99 (1955) 1356
 1955TR03 P.B. Treacy, Proc. Phys. Soc. (London) A68 (1955) 204
 1956BU27 F. Bumiller, H.H. Staub and H.E. Weaver, Helv. Phys. Acta 29 (1956) 83
 1956GO62 G.V. Gorlov, B.M. Gokhberg, V.M. Morozov and G.A. Otroschenko, Dokl. Akad. Nauk SSSR 110 (1956) 963; Sov. Phys. (Doklady) 1 (1956) 598
 1956HE57 N.P. Heydenburg and G.M. Temmer, Phys. Rev. 104 (1956) 123
 1956KI54 B.B. Kinsey and T. Stone, Phys. Rev. 103 (1956) 975
 1956KU1A Kurath, Phys. Rev. 101 (1956) 216

1956MA12 P.R. Malmberg, Phys. Rev. 101 (1956) 114
 1956MO19 C.D. Moak and W.R. Wisseman, Phys. Rev. 101 (1956) 1326
 1956NE13 G.C. Neilson and J.B. Warren, Phys. Rev. 103 (1956) 1758
 1956NI20 R. Nilson, R.O. Kerman, G.R. Briggs and W. Jentschke, Phys. Rev. 104 (1956) 1673
 1956RU41 J.L. Russell, Jr., G.C. Phillips and C.W. Reich, Phys. Rev. 104 (1956) 135
 1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser, Jr., Phys. Rev. 104 (1956) 1064
 1956SE1A Selove, Phys. Rev. 101 (1956) 231
 1956TH06 R.G. Thomas, M. Walt, R.B. Walton and R.C. Allen, Phys. Rev. 101 (1956) 759
 1956WI04 H.B. Willard, J.K. Bair, J.D. Kington and H.O. Cohn, Phys. Rev. 101 (1956) 765
 1957BU13 W.E. Burcham, W.M. Gibson, D.J. Prowse and J. Rotblat, Nucl. Phys. 3 (1957) 217
 1957FR1B French and Fujii, Phys. Rev. 105 (1957) 652
 1957KU58 D. Kurath, Phys. Rev. 106 (1957) 975
 1957NE1A Newson et al, Phys. Rev. 108 (1957) 1294
 1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
 1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809
 1957VA11 D.M. Van Patter and W. Whaling, Revs. Mod. Phys. 29 (1957) 757
 1957WI24 D.H. Wilkinson, Phil. Mag. 2 (1957) 83
 1958BA1E Barnes, Fowler, Greenstein, Lauritsen and Nordberg, Bull. Amer. Phys. Soc. 3 (1958) 328
 1958CA1E Catala, Font, Senent, Aguilar and de La Cuadra, Nuovo Cim. 9, Suppl. 2 (1958) 377
 1958CH31 I.V. Chuvilo and V.G. Shevchenko, Zh. Eksp. Teor. Fiz. 34 (1958) 593; JETP (Sov. Phys.) 7 (1958) 410
 1958CO62 B. Conrad, V. Konig and U. Timm, Naturwiss. 45 (1958) 204
 1958DU78 K.L. Dunning, J.W. Butler and R.O. Bondelid, Phys. Rev. 110 (1958) 1076
 1958FO1C Fowler, Mem. Soc. Roy. Sc. Liege III (1958) 207
 1958KE1B Keszthelyi and Zimanyi, Nuovo Cim. 10 (1958) 909; Repts. Central Research Inst. Phys. Hung. Acad. Sci. 6 (1958) 353
 1958LA08 T. Lauritsen, C.A. Barnes, W.A. Fowler and C.C. Lauritsen, Phys. Rev. Lett. 1 (1958) 326
 1958MA07 R.L. Macklin and J.H. Gibbons, Phys. Rev. 109 (1958) 105
 1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1
 1958NE17 H. Neuert and T. Retz-Schmidt, Z. Naturforsch. A13 (1958) 829

- 1958NI05 R. Nilson, W.K. Jentschke, G.R. Briggs, R.O. Kerman and J.N. Snyder, Phys. Rev. 109 (1958) 850
- 1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10
- 1958VE20 J.F. Vedder, Thesis, Univ. of California (1958); UCRL-8324 (1958)
- 1958WI1E Wildermuth and Kanellopoulos, Nucl. Phys. 7 (1958) 150
- 1959AJ1C Ajzenberg-Selove, Jarmie and haddad, Bull. Amer. Phys. Soc. 4 (1959) 258
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AJ81 F. Ajzenberg-Selove, C.F. Osgood and C.P. Baker, Phys. Rev. 116 (1959) 1521
- 1959BA1D Baz, Adv. Phys. 8 (1959) 349
- 1959BA1F Balashov and Tulinov, Zh. Eksp. Teor. Fiz. 36 (1959) 615; JETP (Sov. Phys.) 9 (1959) 426
- 1959BA34 J.A. Baicker, Thesis, Columbia Univ. (1959); CU-183 (1959)
- 1959BI1B Bilpuch, Weston, Bowman and Newson, Bull. Amer. Phys. Soc. 4 (1959) 42
- 1959BO1C Bockelman, Nucl. Phys. 13 (1959) 205
- 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
- 1959BR71 D.J. Bredin, W.E. Burcham, D. Evans, W.M. Gibson, M.S.C. McKee, D.J. Prowse, J. Rotblat and J.N. Snyder, Proc. Roy. Soc. 251 (1959) 143
- 1959BU07 A.C. Butcher and J.M. McNamee, Proc. Phys. Soc. (London) 74 (1959) 529
- 1959CH1E Chuan, J. Phys. Rad. 20 (1959) 621
- 1959CO68 D. Connor, Phys. Rev. Lett. 3 (1959) 429
- 1959CR84 L. Cranberg, Phys. Rev. 114 (1959) 174
- 1959FA02 B.J. Farmer and C.M. Class, Bull. Amer. Phys. Soc. 4 (1959) 95, C3
- 1959FI1B Filho, An. Acad. Brasil. Cienc. 31 (1959) 1
- 1959GA08 F. Gabbard, R.H. Davis and T.W. Bonner, Phys. Rev. 114 (1959) 201
- 1959GE33 D.S. Gemmell, A.H. Morton and E.W. Titterton, Nucl. Phys. 10 (1959) 33
- 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
- 1959HA29 E.W. Hamburger, Thesis, Univ. of Pittsburgh (1959)
- 1959IM04 W.L. Imhof, R.G. Johnson, F.J. Vaughn and M. Walt, Phys. Rev. 114 (1959) 1037
- 1959JA1B Jancovici, Ann. Phys. 4 (1959) 689
- 1959KA12 D. Kamke, Z. Phys. 156 (1959) 603
- 1959KA13 D. Kamke, Z. Phys. 156 (1959) 621
- 1959KU1C Kurepin and Neudachin, Zh. Eksp. Teor. Fiz. 36 (1959) 1725; JETP (Sov. Phys.) 9 (1959) 1229

- 1959KU1E Kurath, Phys. Rev. Lett. 3 (1959) 431
- 1959LA05 K.H. Lauterjung, B. Schimmer, U. Schmidt-Rohr and H. Maier-Leibnitz, Z. Phys. 155 (1959) 547
- 1959LA11 K.H. Lauterjung, B. Schimmer, U. Schmidt-Rohr and H. Maier-Leibnitz, Z. Phys. 155 (1959) 560
- 1959SA04 M. Sakisaka, J. Phys. Soc. Jpn. 14 (1959) 554
- 1959SE1A Sellschop, Phys. Rev. Lett. 3 (1959) 346
- 1959SP1A Spear, Aust. J. Phys. 12 (1959) 99
- 1959ST1B Starodubtsev and makaryunas, Zh. Eksp. Teor. Fiz. 36 (1959) 1594; JETP (Sov. Phys.) 9 (1959) 1133
- 1959ST30 S.V. Starodubtsev and K.V. Makaryunas, Dokl. Akad. Nauk SSSR 129 (1959) 547; Sov. Phys. Dokl. 4 (1960) 1292
- 1959VA1F Van der Spuy, Nucl. Phys. 11 (1959) 615
- 1959VL24 N.A. Vlasov and A.A. Ogloblin, Zh. Eksp. Teor. Fiz. 37 (1959) 54; JETP (Sov. Phys.) 10 (1960) 39
- 1959WE1A Weston, Bilpuch and Newson, Bull. Amer. Phys. Soc. 4 (1959) 42
- 1959WI1B Wildermuth and Kanellopoulous, Congres. Int. de Phys. Nucl., 1958(Dunod, Paris, 1959) 793
- 1959WI1F Wittern, Naturwiss. 46 (1959) 443
- 1959WI41 K. Winter, B. Torki and E. Remy, Nuovo Cim. 11 (1959) 1
- 1959ZA01 N.I. Zaika and O.F. Nemets, Izv. Akad. Nauk SSSR Ser. Fiz. 23 (1950) 1460
- 1959ZE1A Zeidman and Yntema, Nucl. Phys. 12 (1959) 298
- 1960AG03 J. Aguilar, M. De la Cuadra and R. Font, An. Real Soc. Espan. Fis. Y Quim. (Madrid) 56 (1960) 71; Phys. Abs. 16796 (1961)
- 1960AL18 J. Alster and J. Gonzalez-Vidal, Bull. Amer. Phys. Soc. 5 (1960) 493, C5
- 1960BA27 J.A. Baicker and K.W. Jones, Nucl. Phys. 17 (1960) 424
- 1960BE05 N. Berk, F.E. Steigert and G.L. Salinger, Phys. Rev. 117 (1960) 531
- 1960BE15 R.L. Becker, Phys. Rev. 119 (1960) 1076
- 1960BI06 R. Bilwes, R. Seltz, M. Suffert, C. Gerardin, J. Linck, D. Magnac-Valette and P. Cuer, J. Phys. Rad. 21 (1960) 376
- 1960BI10 R. Bilwes, R. Seltz, C. Gerardin and D. Magnac-Valette, Compt. Rend. 251 (1960) 1771
- 1960BI11 R. Bilwes, C. Gerardin and D. Magnac-Valette, Compt. Rend. 251 (1960) 2157

- 1960BI1E Bifl, Butcher and McNamee, Nucl. Forces and the Few Nucleon Problem (Pergamon, 1960) 641
- 1960BO13 R. Bouchez, C. Delome, J. Fleury, J. Krafft, P. Perrin, L. Goldman, M. Boge and B. Dudek, J. Phys. Rad. 21 (1960) 346
- 1960CO04 H.E. Conzett, G. Igo, H.C. Shaw and R.J. Slobodrian, Phys. Rev. 117 (1960) 1075
- 1960DI02 F.S. Dietrich and L. Cranberg, Bull. Amer. Phys. Soc. 5 (1960) 493, C1
- 1960FA02 B.J. Farmer, J.G. Cramer, Jr. and C.M. Class, Bull. Amer. Phys. Soc. 5 (1960) 101, C1
- 1960FA04 B.J. Farmer and C.M. Class, Nucl. Phys. 15 (1960) 626
- 1960GE07 D.S. Gemmell, Aust. J. Phys. 13 (1960) 116
- 1960GI1A Gibbons and Newson, Fast Neutron Phys.; Eds. J.B. Marion and J.L. Fowler (Academic Press, 1960) 133
- 1960GO04 J. Gonzalez-Vidal, H.C. Conzett and W.H. Wade, Bull. Amer. Phys. Soc. 5 (1960) 230, C9
- 1960GO1B Goldansky, Nucl. Phys. 19 (1960) 482
- 1960GO36 V.I. Goldansky, Zh. Eksp. Teor. Fiz. 38 (1960) 1637; JETP (Sov. Phys.) 11 (1960) 1179
- 1960GR10 T.A. Griffy and L.C. Biedenharn, Nucl. Phys. 15 (1960) 636
- 1960GR21 I.S. Grant, Proc. Phys. Soc. (London) 76 (1960) 737
- 1960HA14 E.W. Hamburger and J.R. Cameron, Phys. Rev. 117 (1960) 781
- 1960HI04 K. Hisatake, Y. Ishizaki, A. Isoya, T. Nakamura, Y. Nakano, B. Saheki, Y. Saji and K. Yuasa, J. Phys. Soc. Jpn. 15 (1960) 741
- 1960HO14 P.E. Hodgson, Nucl. Phys. 21 (1960) 21
- 1960HU1A Hughes, Magurno and Brussel, BNL-325, 2nd Ed., Suppl. 1 (1960)
- 1960IG02 G. Igo, Phys. Rev. 117 (1960) 1079
- 1960JO03 C.M. Jones, G.C. Phillips and P.D. Miller, Phys. Rev. 117 (1960) 525
- 1960JU04 J. Juna, P. Horvath and K. Konecny, Czech. J. Phys. 10 (1960) 715
- 1960KA05 R.W. Kavanagh, Nucl. Phys. 15 (1960) 411
- 1960KA17 R.W. Kavanagh, Nucl. Phys. 18 (1960) 492
- 1960KO1G Komarov, Kurepin and Popova, Zh. Eksp. Teor. Fiz. 38 (1960) 1824; JETP (Sov. Phys.) 11 (1960) 1312; Izv. Akad. Nauk SSSR 24 (1960) 1145; Bul. Acad. Sci. USSR Phys. 24 (1960) 1147
- 1960KR03 K. Krebs, H. Rieseberg and V. Soergel, Z. Phys. 159 (1960) 232
- 1960KU05 D. Kurath, Phys. Rev. Lett. 4 (1960) 180

1960LA1C Lane, *Revs. Mod. Phys.* 32 (1960) 519
 1960MA15 K.V. Makariunas and S.V. Starodubtsev, *Zh. Eksp. Teor. Fiz.* 38 (1960) 372; *JETP (Sov. Phys.)* 11 (1960) 271
 1960MA1G Marion, Levesque, Ludemann and Detenbeck, *Nucl. Instrum. Meth.* 8 (1960) 297
 1960MA1H Marquez and Redon, *Proc. Int. on Nucl. Struct., Kingston* (1960)
 1960MA23 B. Mainsbridge, *Aust. J. Phys.* 13 (1960) 204
 1960MA32 R.D. Macfarlane and J.B. French, *Revs. Mod. Phys.* 32 (1960) 567
 1960MA33 B. Mainsbridge, *Nucl. Phys.* 21 (1960) 1
 1960NO01 M.E. Nordberg, B. Povh and C.A. Barnes, *Phys. Rev. Lett.* 4 (1960) 23
 1960NO05 M.E. Nordberg, F.B. Morinigo and C.A. Barnes, *Phys. Rev. Lett.* 5 (1960) 321
 1960PE02 F.G.J. Perey and P. Lorrain, *Bull. Amer. Phys. Soc.* 5 (1960) 18, E6
 1960PE11 L.D. Pearlstein, Y.C. Tang and K. Wildermuth, *Nucl. Phys.* 18 (1960) 23
 1960PE1B Peterson, Bratenel and Stoering, *Phys. Rev.* 120 (1960) 521
 1960PH1A Phillips and Tombrello, *Nucl. Phys.* 19 (1960) 555
 1960PH1B Phillips and Biedenharn, *Bull. Amer. Phys. Soc.* 5 (1960) 44
 1960PH1C Phillips, Griffy and Biedenharn, *Nucl. Phys.* 21 (1960) 327
 1960PR1D Price, Hamburger and Cohen, *Bull. Amer. Phys. Soc.* 5 (1960) 270
 1960SA25 G.L. Salmon, *Nucl. Phys.* 21 (1960) 15
 1960SE08 J.P.F. Sellschop, *Phys. Rev.* 119 (1960) 251
 1960SE13 J.P.F. Sellschop and D.W. Mingay, *Proc. Int. Conf. on Nucl. Struct., Kingston* (1960) 396
 1960SH01 S.M. Shafroth, *Bull. Amer. Phys. Soc.* 5 (1960) 55, RA1
 1960SI10 L. Simons, *Soc. Sci. Fenn. Comm. Phys. Math* 24 (1960) 13
 1960SI1D Simons, *Proc. Int. on Nucl. Struct., Kingston* (1960)
 1960ST23 R. Stein, H. Braun and P. Cuer, *J. Phys. Rad.* 21 (1960) 374
 1960TA1C Talmi and Unna, *Ann. Revs. Nucl. Sci.* 10 (1960) 353
 1960VA10 S.S. Vasilev, V.V. Komarov and A.M. Popova, *Izv. Akad. Nauk SSSR Ser. Fiz.* 24 (1960) 1149; *Bull. Acad. Sci. Phys.* 24 (1960) 1151
 1960VL03 N.A. Vlasov, S.P. Kalinin, A.A. Ogloblin and V.I. Chuev, *Zh. Eksp. Teor. Fiz.* 39 (1960) 1468; *JETP (Sov. Phys.)* 12 (1961) 1020
 1960WA1G Warburton and Chase, *Phys. Rev.* 120 (1960) 2095
 1960WE1A Weidenmuller, *Phys. Rev. Lett.* 4 (1960) 299

- 1960ZE03 Ya.B. Zeldovich, Zh. Eksp. Teor. Fiz. 38 (1960) 1123; JETP (Sov. Phys.) 11 (1960) 812
- 1961AU02 S.M. Austin, S.E. Darden, A. Okazaki and Z. Wilhelmi, Nucl. Phys. 22 (1961) 451
- 1961BA1C Baz, Goldanskii and Zeldovich, Sov. Phys. Uspekhi 3 (1961) 729
- 1961BA1E Balashov, Neudachin and Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 170; Bull. Acad. Sci. USSR Phys. 25 (1961) 165
- 1961BE05 P.R. Bevington, W.W. Rolland and H.W. Lewis, Phys. Rev. 121 (1961) 871
- 1961BE1E Beckner, Jones and Philips, Phys. Rev. 123 (1961) 255
- 1961BR02 G. Breuer, V. Riech, E. Thormann and H. Neuert, Z. Phys. 161 (1961) 500
- 1961CH09 R. Chiba, H.E. Conzett, H. Morinaga, N. Mutsuro, K. Shoda and M. Kimura, J. Phys. Soc. Jpn. 16 (1961) 1077
- 1961CH1C Christy and Duck, Nucl. Phys. 24 (1961) 89
- 1961CI02 N. Cindro, M. Cerineo and A. Strzalkowski, Nucl. Phys. 24 (1961) 107
- 1961CL09 A.B. Clegg, K.J. Foley, G.L. Salmon and R.E. Segel, Proc. Phys. Soc. (London) 78 (1961) 681
- 1961CL10 A.B. Clegg, Phil. Mag. 6 (1961) 1207
- 1961DA04 S.E. Darden, T.R. Donoghue and C.A. Kelsey, Nucl. Phys. 22 (1961) 439
- 1961DE1E Dekka, Evans, Prowse and Baldo-Ceolin, Nucl. Phys. 23 (1961) 657
- 1961DU01 J.R. Dunning, A.M. Smith and F.E. Steigert, Phys. Rev. 121 (1961) 580
- 1961ER01 J.R. Erskine and C.P. Browne, Phys. Rev. 123 (1961) 958
- 1961FO1C Fowler and Preist, Nucl. Phys. 23 (1961) 667
- 1961GO1T Gooding and Igo, Phys. Rev. Lett. 7 (1961) 28
- 1961HA1G Halton and Hodgson, Proc. Rutherford Jub. Int. Conf. (1961) 593
- 1961HA27 I.G. Han and N.P. Heydenburg, Bull. Amer. Phys. Soc. 6 (1961) 415, B1, and Private Communication (1961)
- 1961JA19 von L. Jarczyk, J. Lang, R. Muller and W. Wolfli, Helv. Phys. Acta 34 (1961) 483
- 1961JE01 H. Jeremie, Compt. Rend. 252 (1961) 403
- 1961JO17 A. Johansson, U. Svanberg and O. Sundberg, Ark. Fys. 19 (1961) 527
- 1961JO18 A. Johansson, U. Svanberg and P.E. Hodgson, Ark. Fys. 19 (1961) 541
- 1961KO08 J. Kokame, J. Phys. Soc. Jpn. 16 (1961) 2101
- 1961LA1A Lane, Langsdorf, Monahan and Elwyn, Ann. Phys. 12 (1961) 135
- 1961LE10 G.J.F. Legge, Nucl. Phys. 26 (1961) 608
- 1961LE1K Lemeille, Saunier and Marquez, Centro Brasileiro de Pesquisas Fisicas 8:5 (1961)

- 1961LO07 O.V. Lozhkin and A.A. Rimskii-Korsakov, Zh. Eksp. Teor. Fiz. 40 (1961) 1519; JETP (Sov. Phys.) 13 (1961) 1064
- 1961ME02 J.W. Meadows and J.F. Whalen, Nucl. Sci. Eng. 9 (1961) 132
- 1961ME10 V. Meyer, H. Muller, H.H. Staub and R. Zurmuhle, Nucl. Phys. 27 (1961) 284
- 1961MY01 S.A. Myachkova and V.P. Pereygin, Zh. Eksp. Teor. Fiz. 40 (1961) 1244; JETP (Sov. Phys.) 13 (1961) 876
- 1961NI04 A. Nilsson, Ark. Fys. 19 (1961) 289
- 1961RE03 F.H. Read and J.M. Calvert, Proc. Phys. Soc. (London) 77 (1961) 65
- 1961RO20 J.R. Rook, Nucl. Phys. 27 (1961) 606
- 1961SC1B Schmid and Wildermuth, Nucl. Phys. 26 (1961) 463
- 1961SE13 W. Sebaoun and H. Gauvin, J. Phys. Rad. 22 (1961) 771
- 1961SH1F Shimodaya, Tamagaki and Tanaka, Prog. Theor. Phys. 25 (1961) 853
- 1961TE02 A. Tejera, M. Mazari, A. Jaidar and G. Lopez, Rev. Mex. Fis. 10 (1961) 229
- 1961VA17 S.S. Vasilev, V.M. Komarov and A.M. Popova, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 1117; Bull. Acad. Sci. Phys. 25 (1960) 1124
- 1961VA38 S.S. Vasilev, V.V. Komarov and A.M. Popova, Zh. Skep. Teor. Fiz. 41 (1961) 1757; JETP (Sov. Phys.) 14 (1962) 1249
- 1961WA03 A.H. Wapstra and D.W. Connor, Nucl. Phys. 22 (1961) 336
- 1961YA04 M. Yamada and Z. Matumoto, J. Phys. Soc. Jpn. 16 (1961) 1497
- 1962AB01 Y.G. Abov, O.N. Yermakov, A.D. Gulko, P.A. Krupchitsky and S.S. Trostin, Nucl. Phys. 34 (1962) 505
- 1962AL20 R.G. Allas, S.S. Hanna and R.E. Segel, Bull. Amer. Phys. Soc. 7 (1962) 453, GA3
- 1962AU01 S.M. Austin, Bull. Amer. Phys. Soc. 7 (1962) 269, CB6
- 1962BA1B Bassi et al, Nuovo Cim. 24 (1962) 560
- 1962BA1C Barker and Treacy, Nucl. Phys. 38 (1962) 33
- 1962BA1W Batchelor and Towle, Phys. Lett. 2 (1962) 312
- 1962BE11 W. Benenson, T.H. May and R.L. Walter, Nucl. Phys. 32 (1962) 510
- 1962BE14 K. Bearpark, I.Hall, R.E. Segel, S.M. Shafroth and N.W. Tanner, Nucl. Phys. 33 (1962) 648
- 1962BE21 H. Beaumevieille, N. Longequeue and J.P. Longequeue, Compt. Rend. 255 (1962) 1340
- 1962BE35 J. Benveniste, A.C. Mitchell, C.D. Schrader and J.H. Zenger, Nucl. Phys. 38 (1962) 300

- 1962BE43 Y.A. Bereznoi, A.P. Klyucharev, Y.N. Ranyuk and N.Y. Rutkevich, Zh. Skep. Teor. Fiz. 43 (1962) 1248; JETP (Sov. Phys.) 16 (1963) 883
- 1962BI11 J.A. Biggerstaff, R.F. Hood, H. Scott and M.T. McEllistrem, Nucl. Phys. 36 (1962) 631
- 1962BO1F Bowen et al, Nucl. Phys. 30 (1962) 475
- 1962BR14 L.B. Brown and H.B. Knowles, Phys. Rev. 125 (1962) 1339
- 1962CA12 Y. Cassagnou, J.M.F. Jeronymo, G.S. Mani, A. Sadeghi and P.D. Forsyth, Nucl. Phys. 33 (1962) 449
- 1962CA13 S. Cavallaro, R. Potenza and A. Rubbino, Nucl. Phys. 36 (1962) 597
- 1962CE01 J. Cerny, B.G. Harvey and R.H. Pehl, Nucl. Phys. 29 (1962) 120
- 1962CH14 L.F. Chase, Jr., R.G. Johnson, F.J. Vaughn and E.K. Warburton, Phys. Rev. 127 (1962) 859
- 1962CO05 M. Coste and L. Marquez, Compt. Rend 254 (1962) 1768
- 1962CO08 D. Connor and T. Tsang, Phys. Rev. 126 (1962) 1506
- 1962CO21 M. Coste, Compt. Rend. 255 (1962) 2750
- 1962CU05 B. Cujec, Nucl. Phys. 37 (1962) 396
- 1962EL01 A.J. Elwyn and R.O. Lane, Nucl. Phys. 31 (1962) 78
- 1962FO1E Foster and Glasgow, Bull. Amer. Phys. Soc. 7 (1962) 334
- 1962GA21 A. Garin and L. Marquez, J. Phys. Rad. 23 (1962) 970
- 1962GA23 J.P. Garron, Ann. Phys. (Paris) 7 (1962) 301
- 1962HA15 I.G. Han and N.P. Heydenburg, Bull. Amer. Phys. Soc. 7 (1962) 570, D10
- 1962HA23 S.S. Hanna, Bull. Amer. Phys. Soc. 7 (1962) 470, LA4
- 1962IG1B Igo, Hansen and Gooding, UCRL 10574 (1962)
- 1962IN02 D.R. Inglis, Nucl. Phys. 30 (1962) 1
- 1962IN1A Inglis, Revs. Mod. Phys. 34 (1962) 165
- 1962IW1A Iwao and Viano, Phys. Lett. 3 (1962) 9
- 1962JE02 J.M.F. Jeronymo, G.S. Mani, F. Picard and A. Sadeghi, Nucl. Phys. 38 (1962) 11
- 1962KO13 M.P. Konstantinova, E.V. Myakinin, A.M. Petrov and A.N. Ronsnov, Zh. Eksp. Teor. Fiz. 43 (1962) 388; JETP (Sov. Phys.) 16 (1963) 278
- 1962LA15 N.O. Lassen, Nucl. Phys. 38 (1962) 442
- 1962LU01 R.A. Lux, R.A. Blue and W. Haeberli, Bull. Amer. Phys. Soc. 7 (1962) 268, CB5
- 1962MA40 D.R. Maxson, Phys. Rev. 128 (1962) 1321

- 1962MA59 K.V. Makariunas, E.K. Makariuniene and V.J. Dienys, Litov. Fiz. Sbornik (USSR) 2 (1962) 351; ; Nucl. Sci. Abstr. 18, 2032, Abs.15242 (1964); Phys. Abs. 3703 (1964)
- 1962NE14 B.M.K. Nefkens, Bull. Amer. Phys. Soc. 7 (1962) 550, T6
- 1962NO02 M.E. Nordberg, Jr., F.B. Morinigo and C.A. Barnes, Phys. Rev. 125 (1962) 321
- 1962OT01 P.S. Otstavnov and V.I. Popov, Zh. Eksp. Teor. Fiz. 43 (1962) 385; JETP (Sov. Phys.) 16 (1963) 276
- 1962PE20 A.M. Petrov, Zh. Eksp. Teor. Fiz. 43 (1962) 66; JETP (Sov. Phys.) 16 (1963) 47
- 1962RO12 L. Rosen and L. Stewart, Phys. Rev. 126 (1962) 1150
- 1962RO20 L. Rosen and W.T. Leland, Phys. Rev. Lett. 8 (1962) 379
- 1962SE1A Serov and Guzhovskii, At. Energ. 12 (1962) 5
- 1962SH1F Shimodaya, Tamagaki and Tanaka, Prog. Theor. Phys. 27 (1962) 793
- 1962ST12 F.A. St. Romain, T.W. Bonner, R.L. Bramblett and J. Hanna, Phys. Rev. 126 (1962) 1794
- 1962TE01 G.M. Temmer, Bull. Amer. Phys. Soc. 7 (1962) 59, R4
- 1962TE02 A.R. Tejera, Bull. Amer. Phys. Soc. 7 (1962) 570, D6
- 1962TE04 I.B. Teplov, I.S. Dmitriev, Y.A. Teplova and O.P. Shevchenko, Zh. Eksp. Teor. Fiz. 42 (1962) 353; JETP (Sov. Phys.) 15 (1962) 243
- 1962TE07 I.B. Teplov, I.S. Dmitriev, Y.A. Teplova and O.P. Shevchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1962) 1150; Bull. Acad. Sci. USSR Phys. Ser. 26 (1962) 1162
- 1962VA1A Vasilyev, Komarcv and Popova, Zh. Skep. Teor. Fiz. 43 (1962) 737; JETP (Sov. Phys.) 16 (1963) 521
- 1962VA25 S.S. Vasilyev, V.V. Komarov and A.M. Popova, Nucl. Phys. 38 (1962) 344
- 1962WE13 C. Weddigen, Z. Phys. 170 (1962) 448
- 1962WE1C Wegner, Rev. Sci. Instrum. 33 (1962) 271
- 1963AL19 D.E. Alburger, P.F. Donovan and D.H. Wilkinson, Phys. Rev. 132 (1963) 334
- 1963AL1J Alter and Lorenzi, NAA-SR-MEMO-9028 (1963)
- 1963AN09 Y. Antoufi ev, I. Bondouk, A. Fikri, F. Machali and P.V. Sorokin, Nucl. Phys. 47 (1963) 426
- 1963AN10 Y.P. Antoufi ev, M.M. El-Shesheni, H.R. Saad, Z.A. Saleh and P.V. Sorokin, Nucl. Phys. 48 (1963) 299; Izv. Akad. Nauk SSSR Ser. Fiz. 27 (1963) 1451
- 1963AS01 V.J. Ashby, H.C. Catron, M.D. Goldberg, R.W. Hill, J.M. Le Blanc, L.L. Newkirk, J.P. Stoering, C.J. Taylor and M.A. Williamson, Phys. Rev. 129 (1963) 1771
- 1963BA1R Bachelier et al, Padua (1963) 1141
- 1963BA2D Bahcall, Fowler, Iben and Sears, Astrophys. J. 137 (1963) 344

- 1963BA34 P. Bassi, B. Ferretti, G. Venturini, G.C. Bertolini, F. Cappellani, V. Mandl, G.B. Restelli and A. Rota, *Nuovo Cim.* 28 (1963) 1049
- 1963BA50 R. Batchelor and J.H. Towle, *Nucl. Phys.* 47 (1963) 385
- 1963BA56 J.F. Barry, *J. Nucl. Energy* 17 (1963) 273
- 1963BA58 R. Balzer, D. Bharucha, F. Heinrich and A. Hofmann, *Helv. Phys. Acta* 36 (1963) 495
- 1963BE1U Y.A. Bereznoi, A.P. Klyucharev, Y.N. Ranyuk and N.Y. Rutkevich, *Zh. Skep. Teor. Fiz.* 45 (1962) 1030; *JETP (Sov. Phys.)* 18 (1963) 713
- 1963BI1B Birk, Goldring, Hillman and Moreh, *Nucl. Phys.* 41 (1963) 58
- 1963BI22 R. Bilwes, B. Bourotte and D. Magnac-Valette, *J. Phys.* 24 (1963) 948
- 1963BL20 H.R. Blieden, G.M. Temmer and K.L. Warsh, *Nucl. Phys.* 49 (1963) 209
- 1963BO06 R.R. Borchers and C.H. Poppe, *Phys. Rev.* 129 (1963) 2679
- 1963BO1N Bowen, Cox, Huxtable, Scanlon, Thresher, Langsford and Appel, *Nucl. Phys.* 41 (1963) 177
- 1963BR03 J.D. Bronson, W.D. Simpson and G.C. Phillips, *Bull. Amer. Phys. Soc.* 8 (1963) 125, U8
- 1963BR1N Brandus, Iosifescu, Sandulescu and Stihi, *Rev. Phys. (Rumania)* 8 (1963) 431
- 1963BR28 F. Brown, R.H. James, J.L. Perkin and J. Barry, *J. Nucl. Energy (Reactor Sci. Tech.)* A17 (1963) 137
- 1963BU1C Bunakov, *Phys. Lett.* 7 (1963) 14
- 1963CA1K Calvi, Cavallaro, Potenza and Rubbino, *Padua* (1963) 1068
- 1963CR08 L. Cranberg, A. Jacquot and H. Liskien, *Nucl. Phys.* 42 (1963) 608
- 1963DA05 J. Darvas, *Z. Naturforsch.* A18 (1963) 569
- 1963DA1C Davis, *Bull. Amer. Phys. Soc.* 8 (1963) 26
- 1963DE1G Demortier, Dossin, Savariraj and Deconninck, *Ann. Soc. Sci. Bruxelles* 77 (1963) 81
- 1963DE29 G. Deconninck and G. Demortier, *Phys. Lett.* 7 (1963) 260
- 1963DI02 F.S. Dietrich, J.L. Honsaker and J.W. Davies, *Bull. Amer. Phys. Soc.* 8 (1963) 120, M5
- 1963DI17 L. Dick, L. Feuvrais and M. Spighel, *Phys. Lett.* 7 (1963) 150
- 1963DO08 W.E. Dorenbusch and C.P. Browne, *Phys. Rev.* 131 (1963) 1212
- 1963DO11 W.E. Dorenbusch and C.P. Browne, *Phys. Rev.* 132 (1963) 1759
- 1963DR1B Drisko, Satchler and Dassel, *3rd Conf. on Reactions between Complex Nuclei* (1963) 85
- 1963EL1D El-Nadi, Rabie and Sherif, *Nucl. Phys.* 48 (1963) 569
- 1963FE03 L. Feldman, B.B. Baliga and M. Nessin, *Bull. Amer. Phys. Soc.* 8 (1963) 290, A6

- 1963FR1G Freeman and Mani, *J. Phys.* 24 (1963) 951
- 1963GL1C Glendenning, *Ann. Rev. Nucl. Sci.* 13 (1963) 191
- 1963GL1E Glasgow and Burge, *Nucl. Phys.* 50 (1963) 327
- 1963GO1M Goldberg, May and Stehn, *BNL-400, 2nd Ed., Vol. 1* (1963)
- 1963GR11 W. Gruhle, K.H. Lauterjung, B. Schimmer, U. Schimmer and U. Schmidt-Rohr, *Nucl. Phys.* 42 (1963) 321
- 1963HA1G Haerberli, in *Fast Neutron Phys.*, Eds. Marion and Fowler (Academic Press, 1963) 1379
- 1963HU1G Huffaker and Greuling, *Bull. Amer. Phys. Soc.* 8 (1963) 315
- 1963HW01 C.F. Hwang, G. Clausnitzer, D.H. Nordby, S. Suwa and J.H. Williams, *Phys. Rev.* 131 (1963) 2602
- 1963IS04 G. Ischenko, D. Seyboth and H. Wegener, *Z. Phy.* 175 (1963) 422
- 1963JA07 A.N. James and H.G. Pugh, *Nucl. Phys.* 42 (1963) 441
- 1963JE05 H. Jeremie, *Nucl. Phys.* 47 (1963) 225
- 1963KA20 M. Kamegai, *Phys. Rev.* 131 (1963) 1701
- 1963KU03 D. Kurath, *Phys. Rev.* 130 (1963) 1525
- 1963LA02 R.A. Lasalle, R.D. Benta and J.G. Cramer, Jr., *Bull. Amer. Phys. Soc.* 8 (1963) 303, D8
- 1963LE19 C. Lemeille, D. Manesse, L. Marquez, N. Saunier and J.L. Quebert, *J. Phys.* 24 (1963) 908
- 1963LE1K Levintov and Pavlovskii, *Zh. Eksp. Teor. Fiz.* 44 (1963) 1442; *JETP (Sov. Phys.)* 17 (1963) 969
- 1963LO1E Lozhkin, Perfi lov and Iakovlev, *Dokl. Akad. Nauk SSSR* 151 (1963) 826; *Sov. Phys. Dokl.* 8 (1964) 791
- 1963LU10 H.F. Lutz, J.B. Mason and M.D. Karvelis, *Nucl. Phys.* 47 (1963) 521
- 1963MA02 J.D. Marshall, N.R. Fletcher and R.H. Davis, *Bull. Amer. Phys. Soc.* 8 (1963) 10, BA1
- 1963MA1E Matkhiz, Neudachin and Smirnov, *Izv. Akad. Nauk. SSSR Ser. Fiz.* 27 (1963) 1273
- 1963MA1Q MacGregor, Booth and Ball, *Phys. Rev.* 130 (1963) 1471
- 1963MA1R Marion, *Univ. of Maryland TR* 320 (1963)
- 1963ME09 V. Meyer, W. Pfeifer and H.H. Staub, *Helv. Phys. Acta* 36 (1963) 465
- 1963MI06 I. Minzatu, A. Calboreanu, N. Martalogu, R. Dumitrescu and M. Molea, *Phys. Lett.* 4 (1963) 357
- 1963MI08 I.V. Mitchell and R.B. Taylor, *Nucl. Phys.* 44 (1963) 664

1963MI1M Milone, Padua (1963) 536
1963MO1H Mottelson, Nucl. Spectroscopy; Ed. Racah (Academis Press, 1962)
1963MO1J Morinigo, Private Communication (1963)
1963MO1K Morinigo, Unpublished Thesis, CalTech (1963)
1963NA1E Nash, Nuovo Cim. 30 (1963) 1064
1963NE02 B.M.K. Nefkens, Phys. Rev. Lett. 10 (1963) 55
1963NE07 B.M.K. Nefkens, Phys. Rev. Lett. 10 (1963) 243
1963OG1A Ogloblin, Nucl. Phys. 47 (1963) 408
1963PA04 P. Paul and D. Kohler, Phys. Rev. 129 (1963) 2698
1963PA1K Palmer, Skofronick and Herb, Bull. Amer. Phys. Soc. 8 (1963) 320
1963PE15 R.R. Perry, B. Mainsbridge and J. Rickards, Nucl. Phys. 45 (1963) 586
1963PH1A Phillips, Bronson, Pitts, Barnard, Weil and Belote, Padua (1963) 1070
1963PU02 K.H. Purser and B.H. Wildenthal, Nucl. Phys. 44 (1963) 22; Erratum Nucl. Phys. 50 (1964) 694
1963RA01 P. Radvanyi, D. Bachelier, M. Bernas, C. Detraz, J. Genin and J.N. Haag, Bull. Amer. Phys. Soc. 8 (1963) 10, BA2
1963RE09 N.W. Reay, N.M. Hintz and L.L. Lee, Jr., Nucl. Phys. 44 (1963) 338
1963RI09 V. Riech, Phys. Lett. 6 (1963) 267
1963SA10 N. Sarma, K.S. Jayaraman and C.K. Kumar, Nucl. Phys. 44 (1963) 205
1963SC1N Schlueter, Univ. of Kansas Rept. (1963)
1963SE1F Sellschop and Mingay, Padua (1963) 425
1963SE1G Sellschop and Mingay, Padua (1963) 545
1963SH04 N.K. Sherman, Phys. Lett. 4 (1963) 113
1963SH1G Shumilov, Zh. Eksp. Teor. Fiz. 44 (1963) 1199; JETP (Sov. Phys.) 17 (1963) 812
1963TA1B Takeda and Nakasima, 3rd Conf. on Reactions between Complex Nuclei (1963) 159
1963TO02 T.A. Tombrello and L.S. Senhouse, Phys. Rev. 129 (1963) 2252
1963TR08 P. Truol and V. Meyer, Helv. Phys. Acta 36 (1963) 494
1963VA04 S.S. Vasilyev, V.V. Komarov and A.M. Popova, Nucl. Phys. 40 (1963) 443
1963VA1C Valentin, Albouy, Cohen and Gusakow, Phys. Lett. 7 (1963) 163
1963WE08 J.A. Weinman and R.K. Smither, Nucl. Phys. 45 (1963) 260
1963WE1B Wegner, Hall and Miller, Padua (1963) 1104
1963WE1F Wells, Tucker and Meyerhof, Phys. Rev. 131 (1963) 1644

1963WI1H Wittern, Z. Naturforsch. A18 (1963) 1254
 1964AM1D Amit and Latz, Nucl. Phys. 58 (1964) 297
 1964AN08 Y.P. Antufev, T. Bunduk, A. Fikri, F. Machali and N.V. Sorokin, Zh. Eksp. Teor. Fiz. 46 (1964) 409; JETP (Sov. Phys.) 19 (1964) 276
 1964AR1C Armstrong, Gammel, Rosen and Frye, Nucl. Phys. 52 (1964) 505
 1964AS04 F. Asfour, I. Bondouk, V.J. Gontchar, V.A. Lutsik, F. Machali and I.I. Zaloubovsky, Nucl. Phys. 58 (1964) 453
 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209
 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
 1964BA1X Bahcall, Phys. Rev. 135 (1964) B137
 1964BA1Y Barker, Proc. Phys. Soc. (London) 84 (1964) 681
 1964BA2A Bahcall, Phys. Rev. Lett. 12 (1964) 300
 1964BE1M Berkowitz, Nucl. Phys. 60 (1964) 555
 1964BE1N Beregi, Selenskaja, Neudatchin and Smirnov, Congres Int. de Phys. Nucl., Paris (1964)
 1964BL12 J.L. Black, G.A. Jones and P.B. Treacy, Nucl. Phys. 54 (1964) 689
 1964BL1C Blieden, Phys. Lett. 9 (1964) 176
 1964BO31 R. Bouchez, J.-C. Gondrand, P. Perrin, C. Perrin, A. Giorni, R. Quivy and M. Dubus, Compt. Rend. 259 (1964) 3501
 1964BR05 J.D. Bronson, W.D. Simpson, W.R. Jackson and G.C. Phillips, Bull. Amer. Phys. Soc. 9 (1964) 406, CA4
 1964BR1H Brown, Paris (1964) 129
 1964BR1U Brown, Blair, Bodansky, Cue and Kavaloski, Bull. Amer. Phys. Soc. 9 (1964) 545
 1964BR25 T.A. Brinkley, B.A. Robson and E.W. Titterton, Proc. Phys. Soc. (London) 84 (1964) 201
 1964BU08 S.G. Buccino, C.E. Hollandsworth and P.R. Bevington, Nucl. Phys. 53 (1964) 375
 1964CA16 R.R. Carlson, R.L. McGrath and E. Norbeck, Phys. Rev. 136 (1964) B1687
 1964CA1G Carlson and Kibler, SUI 64-21 (1964)
 1964CO1D Conzett, Slobodrian, Yamabe and Shield, Congres Int. de Phys. Nucl., Paris (1964)
 1964CO27 H.E. Conzett, E. Shield, R.J. Slobodrian and S. Yamabe, Bull. Amer. Phys. Soc. 9 (1964) 704, B5
 1964DA12 P. Darriulat, G. Igo, H.G. Pugh and H.D. Holmgren, Phys. Lett. 11 (1964) 326
 1964DA1B Daehnick and Denes, Phys. Rev. 136 (1964) B1325
 1964DA1G Davis and Mayer-Boricke, Congres Int. de Phys. Nucl., Paris (1964)

- 1964DE1H Dehnhard, Kamke and Kramer, Phys. Lett. 12 (1964) 113
- 1964DE1K Demortier and Deconninck, Congres. Int. de Phys. Nucl., Paris (1964)
- 1964DU1D Duck, Nucl. Phys. 57 (1964) 643
- 1964DU1E Duncan, Duggan and Purrington, Phys. Rev. 134 (1964) B164
- 1964EN1C Endo, Shimodaya and Miura, Prog. Theor. Phys. 31 (1964) 1157
- 1964ET02 J.E. Etter, M.A. Waggoner, H.D. Holmgren, C. Moazed and A.A. Jaffe, Phys. Lett. 12 (1964) 42
- 1964FA08 U. Fasoli, D. Toniolo and G. Zago, Nuovo Cim. 34 (1964) 542
- 1964FE01 P. Fessenden and D.R. Maxson, Phys. Rev. 133 (1964) B71
- 1964GA1E Garin et al, J. Phys. 25 (1964) 768
- 1964GL03 R. Gleyvod, I.M. Naqib and N.P. Heydenburg, Bull. Amer. Phys. Soc. 9 (1964) 349, E11
- 1964GO25 V.I. Goldansky, Phys. Lett. 9 (1964) 184
- 1964GR1J Green, Nucl. Phys. 54 (1964) 505
- 1964HI04 C.T. Hibdon, Phys. Rev. 133 (1964) B353
- 1964JO04 C.H. Johnson and C.C. Trail, Phys. Rev. 133 (1964) B1183
- 1964JO1D Jones, Bair, Johnson and Willard, Congres Int. de Phys. Nucl., Paris (1964)
- 1964JO1F Jones, Bair, Johnson and Willard, Bull. Amer. Phys. Soc. 9 (1964) 390
- 1964KE1F Kermode, Proc. Phys. Soc. (London) 84 (1964) 557
- 1964LA19 R.O. Lane, A.J. Elwyn and A. Langsdorf, Jr., Phys. Rev. 136 (1964) B1710
- 1964LE1C Letessier, Phys. Lett. 10 (1964) 102
- 1964LO1F Lozhkin and Yakovlev, Zh. Eksp. Teor. Fiz. 47 (1964) 7; JETP (Sov. Phys.) 20 (1965) 4
- 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
- 1964MA25 B.S. Madsen and M. Vedelsby, Nucl. Phys. 55 (1964) 477
- 1964MA2D Mani, Freeman, Redon, Sadeghi and Picard, Congres Int. de Phys. Nucl., Paris (1964)
- 1964MA35 E. Matt, H. Pfander, H. Rieseberg and V. Soergel, Phys. Lett. 9 (1964) 174
- 1964MA51 G.S. Mani, R. Freeman, F. Picard, A. Sadeghi and D. Redon, Nucl. Phys. 60 (1964) 588
- 1964MI10 P.D. Miller, J.L. Duggan and M.M. Duncan, Nucl. Phys. 54 (1964) 155
- 1964MI14 I. Minzatu, A. Calboreanu, N. Martalogu, M. Molea and R. Dumitrescu, Rev. Roum. Phys. 9 (1964) 763
- 1964MI1J Milone and Potenza, Congres Int. de Phys. Nucl., Paris (1964)

1964MO19 C. Moazed, J.E. Etter, H.D. Holmgren and M.A. Waggoner, Phys. Lett. 12 (1964) 45
 1964MO1L Morinigo, Nucl. Phys. 50 (1964) 136
 1964PA06 P. Paul and K.P. Lieb, Nucl. Phys. 53 (1964) 465
 1964PA13 P. Paul, S.L. Blatt and D. Kohler, Phys. Lett. 10 (1964) 201
 1964PA1A Parker, Bahcall and Fowler, Astrophys. J. 139 (1964) 602
 1964PA1K Pape, UCRL 11598 (1964)
 1964PR04 F.W. Prosser, Jr. and F.L. Wilson, Nucl. Phys. 54 (1964) 148
 1964QU01 J.L. Quebert, M. Coste and L. Marquez, Nucl. Phys. 54 (1964) 257
 1964SC19 D.J. Schlueter, R.W. Krone and F.W. Prosser, Jr., Nucl. Phys. 58 (1964) 254
 1964SH05 J.H. Shafer, Phys. Rev. 133 (1964) B920
 1964SH07 T.H. Short and N.M. Hintz, Bull. Amer. Phys. Soc. 9 (1964) 391, BA16
 1964SH19 E. Shield, H.E. Conzett, P. Darriulat, H. Pugh and R.J. Slobodrian, Bull. Amer. Phys. Soc. 9 (1964) 703, B4
 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
 1964SY02 G.D. Symons, Phys. Lett. 10 (1964) 89
 1964TA05 N.W. Tanner, G.C. Thomas and E.D. Earle, Nucl. Phys. 52 (1964) 29
 1964TI02 G. Tibell, O. Sundberg and P.U. Renberg, Ark. Fys. 25 (1964) 433
 1964TO1A Toms, Nucl. Phys. 50 (1964) 561
 1964TO1D Tombrello, Bull. Amer. Phys. Soc. 9 (1964) 704
 1964TO1E Toms, Congre Int. de Phys. Nucl., Paris (1964)
 1964VA14 V. Valkovic, Nucl. Phys. 54 (1964) 465
 1964VA19 V. Valkovic, Nucl. Phys. 60 (1964) 581
 1964VA1E Valkovic and Tomas, Congres Int. de Phys. Nucl., Paris (1964)
 1964VA1J Van Niftrik, Brockman and Van Oers, Congres Int. de Phys. Nucl., Paris (1964)
 1964VO1B Volkov, Phys. Lett. 12 (1964) 118
 1964WE1C Werner and Zimmerer, Congres. Int. de Phys. Nucl., Paris (1964)
 1964YA1A Yanabu et al, J. Phys. Soc. Jpn. 19 (1964) 1818
 1964YO06 F.C. Young, P.D. Forsyth, M.L. Roush, W.F. Hornyak and J.B. Marion, Phys. Lett. 13 (1964) 50
 1965AJ01 F. Ajzenberg-Selove, J.W. Watson and R. Middleton, Phys. Rev. 139 (1965) B592

- 1965AL02 W.J. Alston III, J.E. Draper and R.H. Hilberg, Bull. Amer. Phys. Soc. 10 (1965) 94, HD3
- 1965AL1B Alburger and Donovan, Revs. Mod. Phys. 37 (1965) 448
- 1965AN09 W.D. Andress, Jr., F.O. Purser, J.R. Sawers, Jr. and R.L. Walter, Nucl. Phys. 70 (1965) 313
- 1965AN12 G.B. Andreev, A.S. Deineko and I.Y. Malakhov, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 210; Bull. Acad. Sci. USSR Phys. Ser. 29 (1966) 211
- 1965BA1A A.I. Baz, V.I. Goldanskii and Ya.B. Zeldovich, Usp. Fiz. Nauk 85 (1965) 445; Sov. Phys. Usp. 8 (1965) 177
- 1965BA2H Barker, Bull. Amer. Phys. Soc. 10 (1965) 439
- 1965BE1B Becker and McIntyre, Phys. Rev. 138 (1965) B339
- 1965BE1H Beregi, Zelenskaja, Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 513
- 1965BE1P Bernstein, Ginaven, Chubinsky and Kossler (1965), Unknown Source
- 1965BE1W Bender and Newman, Bull. Amer. Phys. Soc. 10 (1965) 602
- 1965BI1F Bilwes and Bourotte, Revs. Mod. Phys. 37 (1965) 458
- 1965BO07 I. Boca, M. Borsaru, M. Cenja and E. Iliescu, Nucl. Phys. 62 (1965) 75
- 1965DA1A Darriulat, Igo, Pugh and Holmgren, Phys. Rev. 137 (1965) B315
- 1965DE08 C. Detraz, J. Cerny and R.H. Pehl, Phys. Rev. Lett. 14 (1965) 708
- 1965DE1V Dehnhard, Gemmel and Vager, Bull. Amer. Phys. Soc. 10 (1965) 462
- 1965DI1F Dietrich and Zafiratos, Bull. Amer. Phys. Soc. 10 (1965) 439
- 1965DU1C Duck, Revs. Mod. Phys. 37 (1965) 418
- 1965ET1A Etter, Waggoner, Moazed, Holmgren and Han, Revs. Mod. Phys. 37 (1965) 444
- 1965FA03 U. Fasoli, D. Toniolo and G. Zago, Nuovo Cim. 37 (1965) 349
- 1965FR02 R.M. Freeman and G.S. Mani, Proc. Phys. Soc. (London) 85 (1965) 267
- 1965FU1G Fuller and Cohen, Appendix I, Nucl. Data Sheets 6-5 (1965)
- 1965GL03 R. Gleyvod, N.P. Heydenburg and I.M. Naqib, Nucl. Phys. 63 (1965) 650
- 1965GR08 G.M. Griffiths, Nucl. Phys. 65 (1965) 647
- 1965JA07 R. Jahr and R. Santo, Nucl. Phys. 67 (1965) 401
- 1965JO19 C.M. Jones, J.K. Bair, C.H. Johnson, H.B. Willard and M. Reeves III, Revs. Mod. Phys. 37 (1965) 437
- 1965KA1F Kacser and Aitchison, Revs. Mod. Phys. 37 (1965) 350
- 1965KO05 D. Kohler and P. Paul, Phys. Lett. 15 (1965) 157

1965KO1B Komar and Makhnovskii, Dokl. Akad. Nauk SSSR 160 (1965) 1300; Sov. Phys. Dokl. 10 (1965) 150

1965LE09 C. Lemeille, D. Manesse, L. Marquez, N. Saunier and J. Steyaert, J. Phys. (Paris) 26 (1965) 1

1965LO1K Lomakin and Nechaev, Atomnaya Energ. 18 (1965) 33

1965MA13 G.S. Mani, R.M. Freeman, F. Picard, D. Redon and A. Sadeghi, Proc. Phys. Soc. (London) 85 (1965) 281

1965MA1G Marion, Phys. Lett. 14 (1965) 315

1965MA1K Marion, Nucl. Phys. 68 (1965) 463

1965MA1V Markowitz and Pape, Bull. Amer. Phys. Soc. 10 (1965) 602

1965MA2A Mancusi et al, Bull. Amer. Phys. Soc. 10 (1965) 628

1965MA2B Marion, Bull. Amer. Phys. Soc. 10 (1965) 9

1965MC06 W.R. McMurray, P. Van Der Merwe and I.J. Van Heerden, Phys. Lett. 18 (1965) 319

1965MO1L Morgan, Hollandsworth and Walter, Bull. Amer. Phys. Soc. 10 (1965) 440

1965MO1M Moazed, Etter, Holmgren and Waggoner, Revs. Mod. Phys. 37 (1965) 354

1965MO1N Moazed, Etter, Holmgren and Waggoner, Revs. Mod. Phys. 37 (1965) 441

1965MO1P Morrison, Evans, Gale, Ollerhead and Warburton, Private Communication (1965)

1965NE1C Neudatchin and Smirnov, Nucl. Phys. 66 (1965) 25

1965NO1A Norbeck, Mancusi, Carlson and Berkowitz, Revs. Mod. Phys. 37 (1965) 455

1965OK1B Okai and Park, Bull. Amer. Phys. Soc. 10 (1965) 469

1965OL01 W.C. Olsen, W.K. Dawson, G.C. Neilson and J.T. Sample, Nucl. Phys. 61 (1965) 625

1965PH1A Phillips, Revs. Mod. Phys. 37 (1965) 409

1965PO03 A.M. Poskanzer, P.L. Reeder and I. Dostrovsky, Phys. Rev. 138 (1965) B18

1965RO1R Rogers, Bull. Amer. Phys. Soc. 10 (1965) 500

1965SL1C Slee, Bull. Amer. Phys. Soc. 10 (1965) 461

1965SW1B Swan, Revs. Mod. Phys. 37 (1965) 336

1965TA1E Tanner, Nucl. Phys. 63 (1965) 383

1965TR1B Treacy, Phys. Lett. 14 (1965) 125

1965VA1E Valentin, Nucl. Phys. 62 (1965) 81

1965WA12 J.W. Watson, F. Ajzenberg-Selove and R. Middleton, Phys. Lett. 18 (1965) 302

1965WA1M Waggoner, Etter, Holmgren and Moazed, Revs. Mod. Phys. 37 (1965) 358

1965WI07 M. Wilson and J.B. Marion, Phys. Lett. 14 (1965) 313

1965WI1A Williams, Gaedke and Toth, Bull. Amer. Phys. Soc. 10 (1965) 443

- 1965WI1K Wilson, Unpublished Thesis, Univ. of Maryland (1965)
- 1965YO1D Young, Jayaramna, Etter, Holmgren and Waggoner, Revs. Mod. Phys. 37 (1965) 362
- 1965YU1C Yuasa and Hourani, Revs. Mod. Phys. 37 (1965) 399
- 1965YU1D Yuasa and Hourani, Compt. Rend. 260 (1965) 3609
- 1966BA1A J.N. Bahcall, Nucl. Phys. 75 (1966) 10
- 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692