

Energy Levels of Light Nuclei $A = 8$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A490* (1988), p. 1. This version of $A = 8$ differs from the published version in that we have corrected some errors discovered after the article went to press. The introduction and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed June 1, 1988)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 8$

Below is a list of links for items found within the PDF document. The introductory [Table 2](#) is available on this website via the link.

A. Nuclides: [\${}^8\text{n}\$](#) , [\${}^8\text{He}\$](#) , [\${}^8\text{Li}\$](#) , [\${}^8\text{Be}\$](#) , [\${}^8\text{B}\$](#) , [\${}^8\text{C}\$](#)

B. Tables of Recommended Level Energies:

[Table 8.1](#): Energy levels of ${}^8\text{He}$

[Table 8.2](#): Energy levels of ${}^8\text{Li}$

[Table 8.4](#): Energy levels of ${}^8\text{Be}$

[Table 8.9](#): Energy levels of ${}^8\text{B}$

C. [References](#)

D. Figures: [\${}^8\text{Li}\$](#) , [\${}^8\text{Be}\$](#) , [\${}^8\text{B}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

${}^8\text{n}$

(Not illustrated)

${}^8\text{n}$ has not been observed in the interaction of 700 MeV or of 400 GeV protons with uranium: see (1979AJ01). See also (1987FL1A) and (1987SIZX; theor.).

${}^8\text{He}$

(Figs. 1 and 4)

GENERAL: See also (1984AJ01).

Model calculations: (1984VA06, 1985PO10, 1987BL18).

Complex reactions involving ${}^8\text{He}$: (1982AL33, 1983AN13, 1985MA13, 1985TA1D, 1986SA30, 1987AR1G, 1987BO40, 1987KO1Z, 1987PE1C, 1987TAZU, 1988GA10, 1988ST06, 1988TA1A).

Hypernuclei: (1982KA1D, 1983DO1B, 1984BO1H, 1985AH1A, 1985IK1A, 1986BA1W, 1986DA1B, 1987MI38, 1987PO1H).

Other topics: (1983GL1B, 1985AN28, 1987AJ1A, 1988AJ1B).

Ground-state properties of ${}^8\text{He}$: (1983AN1C, 1984FR13, 1985SA32, 1986HE26, 1987BL18, 1987HA30, 1987SA15, 1988JO1C).

Mass of ${}^8\text{He}$: The atomic mass excess of ${}^8\text{He}$ adopted by us and by (1988WA18) is 31598 ± 7 keV. ${}^8\text{He}$ is then stable with respect to decay into ${}^6\text{He} + 2\text{n}$ by 2.137 MeV. See (1979AJ01, 1984AJ01).

The interaction nuclear radius of ${}^8\text{He}$ is 2.48 ± 0.03 fm (1985TA18, 1985TA13) [see also for derived nuclear matter, charge and neutron matter r.m.s. radii].

1. ${}^8\text{He}(\beta^-){}^8\text{Li}$ $Q_m = 10.652$

The half-life of ${}^8\text{He}$ is 119.0 ± 1.5 msec. The decay takes place $(84 \pm 1)\%$ to ${}^8\text{Li}^*(0.98)$ [$\log ft = 4.20$] and $(16 \pm 1)\%$ via the neutron unstable states ${}^8\text{Li}^*(3.21, 5.4)$. $(32 \pm 3)\%$ of the emitted neutrons then populate ${}^7\text{Li}^*(0.48)$. The decay to ${}^8\text{Li}^*(3.21, 5.4)$ suggest $\pi = +$ for ${}^8\text{Li}^*(3.21)$ and 0^+ or 1^+ for ${}^8\text{Li}^*(5.4)$ (1981BJ03). [(1986BO41) suggest $\log ft = 5.0$ for the transition to ${}^8\text{Li}^*(3.21)$]. (1986BO41) report β -delayed tritons with a branching ratio of $(0.9 \pm 0.1)\%$. This decay appears to require a 1^+ state in ${}^8\text{Li}$ at 8.8 MeV with a width, $\Gamma_{\text{c.m.}} \approx 1$ MeV; $\log ft$ is then 4.3 (1986BO41). See also (1988JO1C).

2. ${}^9\text{Be}({}^7\text{Li}, {}^8\text{B}){}^8\text{He}$ $Q_m = -28.264$

Table 8.1: Energy levels of ${}^8\text{He}$

E_x (MeV)	$J^\pi; T$	$\tau_{1/2}$ (msec)	Decay	Reactions
g.s.	$0^+; 2$	119 ± 1.5	β^-	1, 2, 3
2.8 ± 0.4 ^a	$(2^+); 2$			2, 3

^a Excited states are calculated at $E_x = 5.83, 7.92$ and 8.18 MeV, with $J^\pi = 2^+, 1^-$ and 2^- [$(0 + 1)\hbar\omega$ model space]. In the $(0 + 2)\hbar\omega$ model space the excited states are at $5.69, 9.51$ and 11.59 MeV, with $J^\pi = 2^+, 1^+$ and 0^+ (1985PO10). See reaction 3 for possible evidence of other states in ${}^8\text{He}$ (1987BEYI; prelim.).

At $E({}^7\text{Li}) = 83$ MeV, $\theta = 10^\circ$, the population of ${}^8\text{He}_{\text{g.s.}}$, an excited state at 2.8 ± 0.4 MeV (presumably $J^\pi = 2^+$) and a structure near $E_x \approx 7$ MeV are reported by (1985AL1G). See also (1985AL29, 1985ALZL, 1985ALZM).

3. (a) ${}^9\text{Be}({}^9\text{Be}, {}^{10}\text{C}){}^8\text{He}$ $Q_m = -24.602$
 (b) ${}^{11}\text{B}({}^7\text{Li}, {}^{10}\text{C}){}^8\text{He}$ $Q_m = -23.722$

At $E({}^9\text{Be}) = 106.7$ MeV and at $E({}^{11}\text{B}) = 87$ MeV the ground state of ${}^8\text{He}$ is populated. In reaction (a) there is some evidence of a group corresponding to $E_x = 2.6 \pm 0.3$ MeV, $\Gamma = 1.0 \pm 0.5$ MeV, while in reaction (b) excited states are reported at $E_x = 1.3, 2.6$ and 4.0 MeV (± 0.3 MeV). The width of the latter is 0.5 ± 0.3 MeV (1987BEYI). See also (1988BEYJ).

${}^8\text{Li}$
(Figs. 1 and 4)

GENERAL: See also (1984AJ01).

Nuclear models: (1983KU17, 1983SH38, 1984MO1H, 1984REZZ, 1984VA06, 1988WO04).

Special states: (1982PO12, 1983KU17, 1984REZZ, 1984VA06, 1986XU02).

Electromagnetic transitions: (1983KU17).

Astrophysics: (1987MA2C).

Complex reactions involving ${}^8\text{Li}$: (1983FR1A, 1983GU1A, 1983OL1A, 1983WI1A, 1984GR08, 1984HI1A, 1984LA27, 1985JA18, 1985MA02, 1985MA13, 1985MO17, 1985MO24, 1986AV1B, 1986CS1A, 1986GO1G, 1986HA1B, 1986MA19, 1986NAZV, 1986SA30, 1986SIZS, 1986WE1C, 1986XU02, 1987BA38, 1987BE58, 1987BL13, 1987CH26, 1987DE37, 1987GR11, 1987JA06, 1987LY04, 1987TAZU, 1987WA09, 1988AU1A, 1988BL09, 1988CA06, 1988KI05, 1988LI1A, 1988RU01, 1988ST06, 1988TA1A).

Polarization of ${}^8\text{Li}$: (1984KO25, 1986HA1P, 1986NO1C, 1986NO1D, 1987NO04).

Applications: (1985HA40, 1986NO1C, 1987NO04).

Reactions involving pions and other mesons: (1983HA45, 1986CE04, 1986GO1G).

Hypernuclei: (1982KA1D, 1982MO1B, 1983MO1C, 1983SH1E, 1984AS1D, 1984CH1G, 1984MI1C, 1984MI1E, 1984SH1J, 1984ZH1B, 1985MO1F, 1986AN1R, 1986DA1B, 1986YA1F, 1987MI38, 1987PO1H, 1987YA1M, 1988TA29).

Other topics: (1985AN28).

Ground state of ${}^8\text{Li}$: (1983ANZQ, 1985AN28, 1985SA32, 1986GL1A, 1987AR22, 1987HA30, 1987VA26, 1988JO1C, 1988POZS, 1988VA03, 1988WO04).

Fig. 1: Energy levels of ${}^8\text{Li}$. In these diagrams, energy values are plotted vertically in MeV, based on the ground state as zero. Uncertain levels or transitions are indicated by dashed lines; levels which are known to be particularly broad are cross-hatched. Values of total angular momentum J , parity, and isobaric spin T which appear to be reasonably well established are indicated on the levels; less certain assignments are enclosed in parentheses. For reactions in which ${}^8\text{Li}$ is the compound nucleus, some typical thin-target excitation functions are shown schematically, with the yield plotted horizontally and the bombarding energy vertically. Bombarding energies are indicated in laboratory coordinates and plotted to scale in cm coordinates. Excited states of the residual nuclei involved in these reactions have generally not been shown; where transitions to such excited states are known to occur, a brace is sometimes used to suggest reference to another diagram. For reactions in which the present nucleus occurs as a residual product, excitation functions have not been shown; a vertical arrow with a number indicating some bombarding energy, usually the highest, at which the reaction has been studied, is used instead. Further information on the levels illustrated, including a listing of the reactions in which each has been observed, is contained in the master table, entitled "Energy levels of ${}^8\text{Li}$ ".




Table 8.2: Energy levels of ${}^8\text{Li}$ ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$2^+; 1$	$\tau_{1/2} = 838 \pm 6$ msec	β^-	1, 2, 3, 7, 8, 9, 11, 12, 13, 14, 15, 17, 18
0.9808 ± 0.1	$1^+; 1$	$\tau_m = 12 \pm 4$ fsec	γ	2, 7, 8, 10, 11, 12, 13, 17, 18
2.255 ± 3	$3^+; 1$	$\Gamma = 33 \pm 6$ keV	γ, n	2, 3, 4, 7, 11, 12, 13
3.21	$1^+; 1$	≈ 1000	n	5, 10
5.4	$(0, 1)^+; 1$	≈ 650	n	5, 10
6.1 ± 100	$(3); 1$	≈ 1000	n	4
6.53 ± 20	$4^+; 1$	35 ± 15	n	2, 4, 7, 12, 13
7.1 ± 100		≈ 400	n	4
(8)	(1^+)	≈ 1000	t	10
(9)		≈ 6000		11
10.8222 ± 5.5	$0^+; 2$	< 12		16

^a For additional states see reaction 4.

$$\begin{aligned}
 J = 2: & \text{ see (1974AJ01)} \\
 \mu = +1.65335 \pm 0.00035 \text{ nm:} & \text{ see (1978LEZA)} \\
 Q = 24 \pm 2 \text{ mb:} & \text{ see (1979AJ01)}.
 \end{aligned}$$

The interaction nuclear radius of ${}^8\text{Li}$ is 2.36 ± 0.02 fm (1985TA18) [see also for derived nuclear matter, charge and neutron matter r.m.s. radii].

$$1. \text{ } {}^8\text{Li}(\beta^-){}^8\text{Be} \quad Q_m = 16.0039$$

The β^- decay is to the broad 2^+ first-excited state of ${}^8\text{Be}$, which then breaks up into 2α [see reaction 24 in ${}^8\text{Be}$]. The half-life is 838 ± 6 msec [see (1984AJ01)]; $\log ft = 5.4$ (1986WA01).

$$2. \text{ } {}^6\text{Li}(t, p){}^8\text{Li} \quad Q_m = 0.801$$

Angular distributions have been obtained at $E_t = 23$ MeV for the proton groups to ${}^8\text{Li}^*(0, 0.98, 2.26, 6.54 \pm 0.03)$; $\Gamma_{\text{c.m.}}$ for ${}^8\text{Li}^*(2.26, 6.54)$ are 35 ± 10 and 35 ± 15 keV, respectively. J for the latter is ≥ 4 : see (1979AJ01).

Table 8.3: Resonance parameters for ${}^8\text{Li}^*(2.26)$ ^a

E_{res} (keV)	254 ± 3
E_x (MeV) ^b	2.261
Γ (keV)	35 ± 5
$\Gamma_n(E_r)$ (keV)	31 ± 7
Γ_γ (eV) ^b	0.07 ± 0.03
γ_n^2 (keV)	594
θ^2	0.091
radius (fm)	3.30
σ_{max}	12.0
J^π	3^+
l_n	1

^a Energies in lab system except for those labeled ^b. For references see (1974AJ01, 1979AJ01).

^b Energies in c.m. system.

3. ${}^7\text{Li}(n, \gamma){}^8\text{Li}$ $Q_m = 2.033$

The cross section for capture radiation has been measured for $E_n = 40$ to 1000 keV; it decreases from $50 \mu\text{b}$ to $5 \mu\text{b}$ over that interval. The cross section shows the resonance corresponding to ${}^8\text{Li}^*(2.26)$: $E_{\text{res}} = 254 \pm 3$ keV, $\Gamma_n = 31 \pm 7$ keV, $\Gamma_\gamma = 0.07 \pm 0.03$ eV: see Table 8.3 and (1974AJ01). See also (1985SM1B), (1981MUZQ, 1984SH1N, 1986AB1E). The decay of ${}^8\text{Li}^*(2.26) \rightarrow {}^7\text{Li}_{\text{g.s.}} + n$ in the interaction of 35 MeV/A ${}^{14}\text{N}$ ions on Ag is reported by (1987BL13).


4. ${}^7\text{Li}(n, n){}^7\text{Li}$ $E_b = 2.033$

The thermal cross section is 0.97 ± 0.04 b [see (1981MUZQ)], $\sigma_{\text{free}} = 1.07 \pm 0.03$ b (1983KO17). The real coherent scattering length is -2.22 ± 0.01 fm. The complex scattering lengths are $b_+ = -4.15 \pm 0.06$ fm and $b_- = 1.00 \pm 0.08$ fm (1983KO17); see also (1979GL12). See (1984AJ01) for earlier references.

Total and elastic cross sections have been reported for $E_n = 5$ eV to 49.6 MeV: see (1974AJ01, 1979AJ01, 1984AJ01). Cross sections have also been reported for n_0 , n_{0+1} and n_2 at $E_n = 6.82$, 8.90 and 9.80 MeV. (1987SC08; n_2 at the two higher energies).

A pronounced resonance is observed at $E_n = 254$ keV with $J^\pi = 3^+$, formed by p-waves: see Table 8.3. A good account of the polarization is given by the assumption of levels at $E_n = 0.25$ and


3.4 MeV, with $J^\pi = 3^+$ and 2^- , together with a broad $J^\pi = 3^-$ level at higher energy. Broad peaks are reported at $E_n = 4.6$ and 5.8 MeV (± 0.1 MeV) [$^8\text{Li}^*(6.1, 7.1)$] with $\Gamma \approx 1.0$ and 0.4 MeV, respectively, and there is indication of a narrow peak at $E_n = 5.1$ MeV [$^8\text{Li}^*(6.5)$] with $\Gamma \ll 80$ keV and of a weak, broad peak at $E_n = 3.7$ MeV: see (1974AJ01, 1984AJ01). A multi-level, multi-channel R -matrix calculation is reported by (1987KN04). This analysis leads to predictions for the cross section for elastic scattering, for (n, n') to $^7\text{Li}^*(0.48, 4.68, 6.68)$ and for triton production. A number of additional (broad) states of ^8Li , unobserved directly in this and in other reactions, derive from this analysis (1987KN04). See also (1984FE1A, 1984MO1J), (1983DA22, 1983GO1H, 1984SH1N, 1984SH1B, 1986BOZG, 1987LE33, 1988MA1H) and (1983FA17, 1986BA09, 1986FIZX, 1987VE02; theor.).



The excitation function for 0.48 MeV γ -rays shows an abrupt rise from threshold (indicating s-wave formation and emission) and a broad maximum ($\Gamma \approx 1$ MeV) at $E_n = 1.35$ MeV. A good fit is obtained with either $J^\pi = 1^-$ or 1^+ (2^+ not excluded), $\Gamma_{\text{lab}} = 1.14$ MeV. A prominent peak is observed at $E_n = 3.8$ MeV ($\Gamma_{\text{lab}} = 0.75$ MeV) and there is some indication of a broad resonance ($\Gamma_{\text{lab}} = 1.30$ MeV) at $E_n = 5.0$ MeV. At higher energies there is some evidence for structure at $E_n = 6.8$ and 8 MeV followed by a decrease in the cross section to 20 MeV: see (1979AJ01, 1984AJ01). The total cross section for $(n_0 + n_1)$ and n_2 have been reported at $E_n = 8.9$ MeV (1984FE1A; prelim.). For R -matrix analyses see (1987KN04) in reaction 4 and (1984AJ01).

The cross section for reaction (b) rises from threshold to ≈ 360 mb at $E_n \approx 6$ MeV and then decreases slowly to ≈ 250 mb at $E_n \approx 16$ MeV: see (1985SW01, 1987QA01). Cross sections for tritium production have been reported recently from threshold to $E_n = 16$ MeV (1983LI1C; prelim.), 4.57 to 14.1 MeV (1985SW01), 7.9 to 10.5 MeV (1987QA01), 14.74 MeV (1984SMZX; prelim.) and at 14.94 MeV (1985GO18: 302 ± 18 mb). At $E_n = 14.95$ MeV the total α production cross section [which includes to $(n, 2n\text{d})$ process] is 336 ± 16 mb (1986KN06). Spectra at 14.6 MeV may indicate the involvement of states of ^4H (1986MI11). The half-life of ^3H has recently been measured to be 12.38 ± 0.03 mean solar years (1987OL04). See also (1987TI07).

See also (1986DE20, 1986DR10), (1984SH1N, 1985BO1D, 1986BOZG, 1986LI15) and (1986CH1S, 1986FA13, 1986GO1K, 1986IG1A, 1986KO32, 1986MA1R, 1986SE1D, 1986SH1T, 1986SH33, 1986TA1H, 1986VE1A, 1986YA1K, 1986YO1D; applications).



See (1985CH37, 1986CH24). See also (1984SH1B, 1986BOZG, 1988MA1H).

7. ${}^7\text{Li}(p, \pi^+){}^8\text{Li}$ $Q_m = -138.318$

Angular distributions and analyzing powers for the transitions to ${}^8\text{Li}^*(0, 0.98, 2.26)$ have been studied at $E_{\bar{p}} = 200.4$ MeV. [The (p, π^-) reaction to the analog states in ${}^8\text{B}$ is discussed there.] The (p, π^+) cross sections are an order of magnitude greater than the (p, π^-) cross sections and show a much stronger angular dependence (1987CA06). Angular distributions and A_y have also been measured at $E_{\bar{p}} = 250, 354$ and 489 MeV to the first three states of ${}^8\text{Li}$. Those to ${}^8\text{Li}^*(0, 2.26)$ have differential cross sections which exhibit a maximum near the invariant mass of the Δ_{1232} and A_y which are similar to each other and to those of the $\bar{p}p \rightarrow d\pi^+$ reaction. ${}^8\text{Li}^*(6.53)$ is clearly populated (1987HU12).

8. ${}^7\text{Li}(d, p){}^8\text{Li}$ $Q_m = -0.192$

Angular distributions of the p_0 and p_1 groups [$l_n = 1$] at $E_d = 12$ MeV have been analyzed by DWBA: $S_{\text{exp}} = 0.87$ and 0.48 respectively for ${}^8\text{Li}^*(0, 0.98)$. Angular distributions have also been measured at several energies in the range of $E_d = 0.49 \rightarrow 3.44$ MeV (p_0) and 0.95 to 2.94 MeV (p_1). The lifetime of ${}^8\text{Li}^*(0.98)$ is 10.1 ± 4.5 fsec: see (1979AJ01). See also (1985FI1D; astrophysics).

9. (a) ${}^7\text{Li}({}^6\text{Li}, {}^5\text{Li}){}^8\text{Li}$ $Q_m = -3.63$
 (b) ${}^7\text{Li}({}^7\text{Li}, {}^6\text{Li}){}^8\text{Li}$ $Q_m = -5.217$

See (1984KO25).

10. ${}^8\text{He}(\beta^-){}^8\text{Li}$ $Q_m = 10.652$

See ${}^8\text{He}$.

11. (a) ${}^9\text{Be}(e, ep){}^8\text{Li}$ $Q_m = -16.887$
 (b) ${}^9\text{Be}(p, 2p){}^8\text{Li}$ $Q_m = -16.887$

For reaction (a) see (1984AJ01) and (1985KI1A). The summed proton spectrum (reaction (b)) at $E_p = 156$ MeV shows peaks corresponding to ${}^8\text{Li}(0)$ and ${}^8\text{Li}^*(0.98 + 2.26)$ [unresolved]. In addition s-states [$J^\pi = 1^-, 2^-$] are suggested at $E_x = 9$ and 16 MeV, with $\Gamma_{\text{c.m.}} \approx 6$ and 8 MeV; the latter may actually be due to continuum protons: see (1974AJ01). At $E_p = 1$ GeV the separation energy between 5 and 8 MeV broad $1p_{3/2}$ and $1s_{1/2}$ groups is reported to be 10.7 ± 0.5 MeV (1985BE30, 1985DO16). See also (1987GAZM).

12. ${}^9\text{Be}(d, {}^3\text{He}){}^8\text{Li}$ $Q_m = -11.393$

Angular distributions have been reported for the ${}^3\text{He}$ ions to ${}^8\text{Li}^*(0, 0.98, 2.26, 6.53)$ at $E_d = 28$ MeV [C^2S (abs.) = 1.63, 0.61, 0.48, 0.092] and 52 MeV. The distributions to ${}^8\text{Li}^*(6.53)[\Gamma < 100$ keV] are featureless: see (1979AJ01).

13. ${}^9\text{Be}(t, \alpha){}^8\text{Li}$ $Q_m = 2.927$

At $E_t = 12.98$ MeV, angular distributions of the α -particles to ${}^8\text{Li}^*(0, 0.98, 2.26, 6.53 \pm 0.02$ [$\Gamma_{\text{c.m.}} < 40$ keV]) have been measured: see (1974AJ01). At $E_t = 17$ MeV angular distributions to these four states have been analyzed by ZRDWBA and C^2S have been derived (1988LI27). At $E_t = 17$ MeV, $\sigma(\theta)$ and A_y measurements, analyzed by CCBA, lead to $J^\pi = 4^+$ for ${}^8\text{Li}^*(6.53)$: see (1984AJ01). For ${}^8\text{Li}^*(0.98)$, $\tau_m = 14 \pm 5$ fsec, $E_x = 980.80 \pm 0.10$ keV: see (1974AJ01).

14. ${}^9\text{Be}({}^7\text{Li}, {}^8\text{Be}){}^8\text{Li}$ $Q_m = 0.367$

See (1984KO25).

15. ${}^9\text{Be}({}^{11}\text{B}, {}^{12}\text{C}){}^8\text{Li}$ $Q_m = -0.930$

See (1986BE1Q).

16. ${}^{10}\text{Be}(p, {}^3\text{He}){}^8\text{Li}$ $Q_m = -15.981$

At $E_p = 45$ MeV, ${}^3\text{He}$ ions are observed to a state at $E_x = 10.8222 \pm 0.0055$ MeV ($\Gamma_{\text{c.m.}} < 12$ keV): the angular distributions for the transition to this state, and to its analog (${}^8\text{Be}^*(27.49)$), measured in the analog reaction [${}^{10}\text{Be}(p, t){}^8\text{Be}$] are very similar. They are both consistent with $L = 0$ using a DWBA (LZR) analysis: see (1979AJ01).

17. ${}^{11}\text{B}(n, \alpha){}^8\text{Li}$ $Q_m = -6.631$

Angular distributions of the α_0 and α_1 groups have been measured at $E_n = 14.1$ and 14.4 MeV: see (1974AJ01, 1984AJ01).

18. $^{11}\text{B}(^7\text{Li}, ^{10}\text{B})^8\text{Li}$ $Q_m = -9.421$

At $E(^7\text{Li}) = 34$ MeV angular distributions have been studied involving $^8\text{Li}(\text{g.s.}, 0.98)$ and $^{10}\text{B}_{\text{g.s.}}$ (1987CO16).

19. $^{13}\text{C}(\text{d}, ^7\text{Be})^8\text{Li}$ $Q_m = -20.454$

See (1984NE1A).

${}^8\text{Be}$
(Figs. 2 and 4)

GENERAL: See also (1984AJ01).

Shell model: (1984PA04, 1984VA06, 1984ZW1A, 1985FI1E, 1987BL18, 1987KI1C, 1988WO04).

Collective, rotational and deformed models: (1984PA04, 1985RO1G).

Cluster and α -particle models: (1981PL1A, 1983CA12, 1983DR09, 1983FU1D, 1983HA41, 1983JA09, 1983SH38, 1984DE24, 1984DU17, 1984LU1A, 1984LU1B, 1985FI1E, 1986GU1F, 1986KR12, 1986SU06, 1988KR01).

Special states: (1981PL1A, 1983AD1B, 1983BI1C, 1983FE07, 1983FI11, 1983HA41, 1984DE24, 1984DU17, 1984LU1A, 1984LU1B, 1984VA06, 1984VAZS, 1984ZW1A, 1985FI1E, 1985GO1A, 1985PO19, 1985PU03, 1985RO1G, 1986AN10, 1987KA18, 1987KI1C, 1987SV1A, 1987WA36, 1988BA75, 1988KR01, 1988KW02, 1988KH03).

Electromagnetic transitions, giant resonances: (1983FI11, 1984VA1B, 1985FI1E, 1985GO1A, 1985GR1A, 1986AN10, 1986QUZZ, 1987KI1C).

Astrophysical questions: (1985BO1E, 1987FU04).

Complex reactions involving ${}^8\text{Be}$: (1982GU21, 1983DEZW, 1983EL1A, 1983SI1A, 1983WA1F, 1983XU1A, 1984AB1C, 1984PA13, 1985BU16, 1985HA1N, 1985KA1E, 1985KA1G, 1985KAZQ, 1985KW03, 1985PO11, 1985PO19, 1985WA22, 1986BA2D, 1986BL12, 1986BR26, 1986GA24, 1986GU1F, 1986IR01, 1986MA1O, 1986PO06, 1986PO12, 1986TA1M, 1987AR19, 1987BL16, 1987CH26, 1987CH33, 1987CH32, 1987DE1O, 1987DU07, 1987GE1B, 1987GL05, 1987HA45, 1987PE1B, 1987PO23, 1987RUZX, 1988AR05, 1988POZZ, 1988RU01, 1988SA09, 1988VA1E).

Reactions involving pions and other mesons: (1981MC09, 1983SP06, 1985BE1C, 1987HU12).

Hypernuclei: (1982KA1D, 1983SH38, 1983SH1E, 1984ZH1B, 1985AH1A, 1985IK1A, 1985MO1F, 1986BA1W, 1986DA1B, 1987BA2K, 1987MI38, 1987PO1H, 1987YA1M, 1988TA29).

Other topics: (1983AD1B, 1983BI1C, 1983FU1D, 1983GR26, 1983MI1E, 1985AN28, 1986BL1D, 1986GL1E, 1986MA1W, 1987AB21, 1987SV1A, 1988AJ1B, 1988BO04, 1988KW02, 1988RU1B, 1988RU1D, 1988WA1E).

Ground-state properties of ${}^8\text{Be}$: (1983ANZQ, 1983DR09, 1984DU17, 1984LU1A, 1984LU1B, 1985AN28, 1985FI1E, 1985GO1A, 1985SH1A, 1987BL18, 1987BO42, 1987KI1C, 1987KO1U, 1987SA15, 1987SV1A, 1988AR05, 1988WO04).

1. ${}^8\text{Be} \rightarrow 2 {}^4\text{He}$ $Q_m = 0.09189$

$\Gamma_{\text{c.m.}}$ for ${}^8\text{Be}_{\text{g.s.}} = 6.8 \pm 1.7$ eV: see (1974AJ01). See also (1987WE1C, 1988BA86; astrophysics) and (1983DR09; theor.).




Fig. 2: Energy levels of ^8Be . For notation see Fig. 1.

Table 8.4: Energy levels of ${}^8\text{Be}$ ^a

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$0^+; 0$	6.8 ± 1.7 eV	α	1, 2, 4, 10, 11, 12, 13, 14, 19, 20, 21, 22, 23, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52
3.04 ± 30	$2^+; 0$	1500 ± 20	α	2, 4, 10, 11, 12, 13, 14, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 32, 34, 35, 36, 37, 38, 41, 42, 44, 45
11.4 ± 300	$4^+; 0$	≈ 3500 ^b	α	4, 12, 13, 19, 21, 27, 28, 29, 42, 44, 45
16.626 ± 3	$2^+; 0 + 1$	108.1 ± 0.5	γ, α	2, 4, 10, 11, 13, 14, 19, 20, 21, 25, 28, 29, 34, 35, 38, 42, 44
16.922 ± 3	$2^+; 0 + 1$	74.0 ± 0.4	γ, α	2, 4, 10, 11, 13, 14, 19, 20, 21, 27, 28, 29, 34, 35, 38, 42, 44
17.640 ± 1.0	$1^+; 1$	10.7 ± 0.5	γ, p	5, 11, 14, 16, 19, 20, 27, 28, 35, 44
18.150 ± 4	$1^+; 0$	138 ± 6	γ, p	11, 14, 16, 19, 20, 27, 28, 35, 38
18.91	2^-	122 ^e	$\gamma, \text{n}, \text{p}$	11, 14, 15, 16, 19, 23
19.07 ± 30	$3^+; (1)$	270 ± 20	γ, p	11, 14, 16, 19, 27, 28
19.24 ± 25	$3^+; (0)$	230 ± 30	n, p	15, 16, 19, 27, 28, 29, 35
19.4	1^-	≈ 650	n, p	11, 15, 16
19.86 ± 50	$4^+; 0$	700 ± 100	p, α	4, 11, 18, 21, 22, 28, 29, 35
20.1	$2^+; 0$	≈ 1100	$\text{n}, \text{p}, \alpha$	4, 15, 16, 18, 22, 35
20.2	$0^+; 0$	< 1000	α	4, 35
20.9	4^-	1600 ± 200	p	16
21.5	$3^{(+)}$	1000	$\gamma, \text{n}, \text{p}$	14, 15

Table 8.4: Energy Levels of ${}^8\text{Be}$ ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
22.0 ^c	$1^-; 1$	≈ 4000	γ, p	14
22.05 ± 100		270 ± 70		29
22.2	$2^+; 0$	≈ 800	n, p, d, α	4, 9, 13, 15, 16, 18
22.63 ± 100		100 ± 50		29
22.98 ± 100		230 ± 50		29
24.0 ^c	$(1, 2)^-; 1$	≈ 7000	γ, p, α	14, 18
25.2	$2^+; 0$		p, d, α	4, 9, 18
25.5	$4^+; 0$	broad	d, α	9
27.4941 ± 1.8 ^d	$0^+; 2$	5.5 ± 2.0	$\gamma, n, p, d, t, {}^3\text{He}, \alpha$	5, 7, 9, 31
(28.6)		broad	γ, p	14

^a See also Table 8.5 and reaction 4.

^b See, however, reaction 27.

^c Giant resonance: see reaction 14.

^d For the parameters of this state please see Table 8.5 in (1984AJ01).

^e See reaction 23.

2. ${}^4\text{He}(\alpha, \gamma){}^8\text{Be}$ $Q_m = -0.09189$

The yield of γ_1 has been measured for $E_\alpha = 32$ to 36 MeV. The yield of γ_0 for $E_\alpha = 33$ to 38 MeV is twenty times lower than for γ_1 , consistent with E2 decay. An angular correlation measurement at the resonances corresponding to ${}^8\text{Be}^*(16.6 + 16.9)[2^+; T = 0 + 1]$ gives $\delta = 0.19 \pm 0.03$, $\Gamma_\gamma(\text{M1}) = 6.4 \pm 0.5$ eV [weighted mean of the two published measurements listed in (1979AJ01)]. The E_x of ${}^8\text{Be}^*(3.0)$ is determined in this reaction to be 3.18 ± 0.05 MeV [see also Table 8.4 in (1974AJ01)].

The E2 bremsstrahlung cross section to ${}^8\text{Be}_{\text{g.s.}}$ has been calculated as a function of E_x over the 3-MeV state: the total Γ_γ for this transition is 8.3 meV, corresponding to 75 W.u. (1986LA05). A calculation of the Γ_γ from the decay of the 4^+ 11.4-MeV state to the 2^+ state yields 0.46 eV (19 W.u.). The maximum cross section for the intrastate γ -ray transition within the 2^+ resonance is calculated to be ≤ 2.5 nb at $E_x \approx 3.3$ MeV (1986LA19). See also (1985BA45; theor.).

- | | | |
|--|------------------|------------------|
| 3. (a) ${}^4\text{He}(\alpha, n){}^7\text{Be}$ | $Q_m = -18.990$ | $E_b = -0.09189$ |
| (b) ${}^4\text{He}(\alpha, p){}^7\text{Li}$ | $Q_m = -17.3462$ | |
| (c) ${}^4\text{He}(\alpha, d){}^6\text{Li}$ | $Q_m = -22.3716$ | |

Table 8.5: Electromagnetic transitions in ${}^8\text{Be}$ ^a

Transition	Γ_γ (eV)	$ M ^2$ (W.u.)
17.6 \rightarrow 0	16.7	0.15
17.6 \rightarrow 3.0	8.15 ± 0.07 (M1) ^b	0.12
	0.15 ± 0.07 (E2)	
17.6 \rightarrow 16.6	0.032 ± 0.003 ^c	1.48 ± 0.15 (M1)
17.6 \rightarrow 16.9	0.0013 ± 0.0003	0.15 ± 0.04 (M1)
18.15 \rightarrow 0	3.0	
18.15 \rightarrow 3.0	3.8	
18.15 \rightarrow 16.6	0.077 ± 0.019	1.04 ± 0.26 (M1)
18.15 \rightarrow 16.9	0.062 ± 0.007	1.51 ± 0.17 (M1)
18.9 \rightarrow 16.6	0.168	0.053 (E1)
18.9 \rightarrow 16.9	0.099	0.045 (E1)
19.07 \rightarrow 3.0	10.5	

^a See Table 8.7 in (1979AJ01) for the references. See also reaction 2 here.

^b $\delta(\text{E2/M1}) = 0.21 \pm 0.04$, averaged over the energy of the final state.

^c Nearly pure M1: $\delta(\text{E2/M1}) = -0.014 \pm 0.013$.

The cross sections for formation of ${}^7\text{Li}^*(0, 0.48)$ [$E_\alpha = 39$ to 49.5 MeV] and ${}^7\text{Be}^*(0, 0.43)$ [39.4 to 47.4 MeV] both show structures at $E_\alpha \approx 40.0$ and ≈ 44.5 MeV: they are due predominantly to the 2^+ states ${}^8\text{Be}^*(20.1, 22.2)$: see (1979AJ01). The excitation functions for p_0, p_2, d_0, d_1 for $E_\alpha = 54.96$ to 55.54 MeV have been measured in order to study the decay of the first $T = 2$ state in ${}^8\text{Be}$: see Table 8.5 in (1984AJ01). Cross sections for p_{0+1} are also reported at $E_\alpha = 37.5$ to 140.0 MeV: see (1979AJ01, 1984AJ01). The cross sections for reaction (c) has been measured at three energies in the range $E_\alpha = 46.7$ to 49.5 MeV: see (1979AJ01) and below.

The production of ${}^6\text{Li}$, ${}^7\text{Li}$ and ${}^7\text{Be}$ [and ${}^6\text{He}$] has been studied for $E_\alpha = 61.5$ to 158.2 MeV by (1982GL01) and at 198.4 MeV by (1985WO11). The production of ${}^7\text{Li}$ (via reactions (a) and (b)) and of ${}^6\text{Li}$ is discussed. At energies beyond $E_\alpha \approx 250$ MeV the $\alpha + \alpha$ reaction does not contribute to the natural abundance of lithium, reinforcing theories which produce ${}^6\text{Li}$ in cosmic-ray processes and the “missing” ${}^7\text{Li}$ in the Big Bang: thus the universe is open (1985WO11, 1982GL01).

The inclusive cross section for production of ${}^3\text{He}$ has been measured at $E_\alpha = 218$ MeV (1984AL03). For a fragmentation study at 125 GeV see (1985BE1E). See also (1984AJ01, 1984PA1E, 1984RE14).

4. ${}^4\text{He}(\alpha, \alpha){}^4\text{He}$

$E_b = -0.09189$

The α - α scattering reveals the ground state as a resonance with $Q_0 = 92.12 \pm 0.05$ keV, $\Gamma_{\text{c.m.}} = 6.8 \pm 1.7$ eV [$\tau = (0.97 \pm 0.24) \times 10^{-16}$ sec]. For $E_\alpha = 30$ to 70 MeV the $l = 0$ phase shift shows resonant behavior at $E_\alpha = 40.7$ MeV, corresponding to a 0^+ state at $E_x = 20.2$ MeV, $\Gamma < 1$ MeV, $\Gamma_\alpha/\Gamma < 0.5$. No evidence for other 0^+ states is seen above $E_\alpha = 43$ MeV.

The d-wave phase shift becomes appreciable for $E_\alpha > 2.5$ MeV and passes through resonance at $E_\alpha = 6$ MeV ($E_x = 3.18$ MeV, $\Gamma = 1.5$ MeV, $J^\pi = 2^+$): see Table 8.4 in (1974AJ01). Five 2^+ levels are observed from $l = 2$ phase shifts measured from $E_\alpha = 30$ to 70 MeV: ${}^8\text{Be}^*(16.6, 16.9)$ with $\Gamma_\alpha = \Gamma$ [see Table 8.6], and states with $E_x = 20.1, 22.2$ and 25.2 MeV. The latter has a small Γ_α . The $l = 2$ $\alpha - \alpha$ phase shifts have been analyzed by (1986WA01) up to $E_\alpha = 34$ MeV: intruder states below $E_x = 26$ MeV need not be introduced.

The $l = 4$ phase shift rises from $E_\alpha \approx 11$ MeV and indicates a broad 4^+ level at $E_x = 11.5 \pm 0.3$ MeV [$\Gamma = 4.0 \pm 0.4$ MeV]. A rapid rise of δ_4 at $E_\alpha = 40$ MeV corresponds to a 4^+ state at 19.9 MeV with $\Gamma_\alpha/\Gamma \approx 0.96$; $\Gamma < 1$ MeV and therefore $\Gamma_\alpha < 1$ MeV, which is $< 5\%$ of the Wigner limit. A broad 4^+ state is also observed near $E_\alpha = 51.3$ MeV ($E_x = 25.5$ MeV).

Over the range $E_\alpha = 30$ to 70 MeV a gradual increase in δ_6 is observed. Some indications of a 6^+ state at $E_x \approx 28$ MeV and of an 8^+ state at ≈ 57 MeV have been reported; $\Gamma_{\text{c.m.}} \approx 20$ and ≈ 73 MeV, respectively. A resonance is not observed at the first $T = 2$ state, ${}^8\text{Be}^*(27.49)$. See (1979AJ01) for references.

The elastic scattering has also been studied at $E_\alpha = 56.3$ to 95.5 MeV (1987NE1C; pre-lim.), 158.2 MeV, 650 and 850 MeV and at 4.32 and 5.07 GeV/ c [see (1979AJ01, 1984AJ01)] as well as at 198.4 MeV (1985WO11). For α - α correlations involving ${}^8\text{Be}^*(0, 3.0)$ see (1987CH33, 1987PO03). See also (1985KR21, 1986FO04, 1986GOZY, 1986UC1A, 1987FO08) and the ‘‘General’’ section here. For inclusive cross sections see (1984AJ01) and (1984AL03; 218 MeV).

For studies at very high energies see reaction 3 and (1982AB1B, 1984SA39, 1984TA1D, 1984TA1G, 1985AB1A, 1985AK1A, 1985CA1C, 1986BE2N, 1986BE1S, 1986BE1T, 1986TA1N, 1986TA1P, 1987BA13). See also (1986CH1M), (1982NA1B, 1983FA1A, 1984FA1B, 1984FR1C, 1985CA41, 1985FA1A, 1985FR1E, 1985WI1B, 1986AN1F, 1986CH1J, 1986ST1D, 1987HE1B, 1987OT1D), (1985NO1B, 1986LA16, 1987FU04, 1987MU1B; astrophysics) and (1982WE15, 1983AL1C, 1983BU15, 1983FI11, 1983GO25, 1983KO41, 1983MA73, 1983OK06, 1983PR09, 1983SA16, 1984DE24, 1984FI11, 1984FI13, 1984FR10, 1984HE1D, 1984KR10, 1984LI28, 1984MA16, 1984MA1H, 1984MA68, 1984NA11, 1984OK03, 1984TA1E, 1984VAZS, 1984ZA1B, 1985BA45, 1985FI1E, 1985FR1F, 1985HO1B, 1985KI11, 1985PR1A, 1985PR1B, 1985SP05, 1985TH08, 1985YA05, 1985YI1B, 1985YI1C, 1986CR1B, 1986FR12, 1986HO33, 1986LA12, 1986MA03, 1986OC1A, 1986SA30, 1986SU06, 1986WI04, 1986YU1A, 1987BA35, 1987FR1D, 1987KA1W, 1987KR03, 1987OC1B, 1987PR01, 1987SA37, 1987SH1M, 1987WA07, 1988BA75, 1988KR01, 1988MO05; theor.).

5. ${}^6\text{Li}(d, \gamma){}^8\text{Be}$

$$Q_m = 22.2798$$

The yield of γ -rays to ${}^8\text{Be}^*(17.64)$ [1^+ ; $T = 1$] has been measured for $E_d = 6.85$ to 7.10 MeV. A resonance is observed at $E_d = 6965$ keV [$E_x = 27495.8 \pm 2.4$ keV, $\Gamma_{\text{c.m.}} = 5.5 \pm 2.0$ keV];

Table 8.6: Some ${}^8\text{Be}$ states with $16.6 < E_x < 23.0$ MeV ^a

E_x (MeV \pm keV)	$\Gamma_{c.m.}$ (keV)	Reaction
16.627 \pm 5	113 \pm 3 90 \pm 5	${}^7\text{Li}({}^3\text{He}, d)$ ${}^{10}\text{B}(d, \alpha)$
16.623 \pm 3	107.7 \pm 0.5	${}^4\text{He}(\alpha, \alpha)$ ^b
16.630 \pm 3	108.5 \pm 0.5	${}^4\text{He}(\alpha, \alpha)$ ^c
16.626 \pm 3	108.1 \pm 0.5	“best” value
16.901 \pm 5	77 \pm 3 70 \pm 5	${}^7\text{Li}({}^3\text{He}, d)$ ${}^{10}\text{B}(d, \alpha)$
16.925 \pm 3	74.4 \pm 0.4	${}^4\text{He}(\alpha, \alpha)$ ^b
16.918 \pm 3	73.6 \pm 0.4	${}^4\text{He}(\alpha, \alpha)$ ^c
16.922 \pm 3	74.0 \pm 0.4	“best” value
17.640 \pm 1.0	10.7 \pm 0.5	${}^7\text{Li}(p, \gamma)$
18.155 \pm 5	147	${}^7\text{Li}(p, \gamma)$
18.150 \pm 5	138 \pm 6	${}^{10}\text{B}(d, \alpha)$
18.144 \pm 5		${}^9\text{Be}(d, t)$
18.150 \pm 4	138 \pm 6	“best” value
19.06 \pm 20	270 \pm 20	${}^7\text{Li}(p, \gamma)$
19.071 \pm 10	270 \pm 30	${}^9\text{Be}(d, t)$
19.07 \pm 30	270 \pm 20	“best” value
19.21	208 \pm 30	${}^9\text{Be}(p, d)$
19.22 \pm 30	265 \pm 30	${}^9\text{Be}({}^3\text{He}, \alpha)$
19.26 \pm 30	220 \pm 30	${}^9\text{Be}(d, t)$
19.24 \pm 25	230 \pm 30	“best” value
19.86 \pm 50	700 \pm 100	${}^9\text{Be}(d, t)$
22.05 \pm 100	270 \pm 70	${}^9\text{Be}({}^3\text{He}, \alpha)$
22.63 \pm 100	100 \pm 50	${}^9\text{Be}({}^3\text{He}, \alpha)$
22.98 \pm 100	230 \pm 50	${}^9\text{Be}({}^3\text{He}, \alpha)$

^a See Table 8.5 in (1979AJ01) for references. See also Tables 8.7 and 8.8 here.

^b R -matrix theory.

^c Complex eigenvalue theory.

$\Gamma_\gamma = 23 \pm 4$ eV [1.14 ± 0.20 W.u.] for this M1 transition from the first 0^+ ; $T = 2$ state in ${}^8\text{Be}$, in good agreement with the intermediate coupling model: see Table 8.5 in (1984AJ01). See also (1979AJ01).

6. ${}^6\text{Li}(d, n){}^7\text{Be}$ $Q_m = 3.381$ $E_b = 22.2798$

Yield curves and cross sections have been measured for $E_d = 48$ keV to 17 MeV: see (1979AJ01, 1984AJ01). See also (1983SZZY). Polarization measurements are reported at $E_d = 0.27$ to 3.7 MeV. Comparisons of the populations of ${}^7\text{Be}^*(0, 0.43)$ and of ${}^7\text{Li}^*(0, 0.48)$ have been made at many energies, to $E_d = 7.2$ MeV. The n/p ratios are closely equal for analog states, as expected for charge symmetry: see (1979AJ01). However, the n_1/p_1 yield ratio decreases from 1.05 at $E_d = 160$ keV to 0.94 at 60 keV: it is suggested that this is due to polarization of the deuteron (1985CE12). See also ${}^7\text{Be}$, (1985WA1C) and (1984KU15; theor.).

7. ${}^6\text{Li}(d, p){}^7\text{Li}$ $Q_m = 5.0255$ $E_b = 22.2798$

Excitation functions have been measured for $E_d = 30$ keV to 5.4 MeV: see (1979AJ01, 1984AJ01). The thick target yield of 0.48-MeV γ -rays is reported from ≈ 50 to ≈ 170 keV (1985CE12). See also (1983SZZY). An anomaly is observed in the p_1/p_0 intensity ratio at $E_d = 6.945$ MeV, corresponding to the first 0^+ ; $T = 2$ state, $\Gamma = 10 \pm 3$ keV, $\Gamma_{p_0} \ll \Gamma_{p_1}$, $\Gamma_{p_0} < \Gamma_d$. Polarization measurements have been reported at $E_d = 0.6$ to 10.9 MeV: see (1979AJ01). See also ${}^7\text{Li}$ and (1984KU15; theor.).

8. (a) ${}^6\text{Li}(d, d){}^6\text{Li}$ $E_b = 22.2798$
 (b) ${}^6\text{Li}(d, t){}^5\text{Li}$ $Q_m = 0.59$

The yield of elastically scattered deuterons has been measured for $E_d = 2$ to 7.14 MeV. No resonances are observed: see (1974AJ01). See also (1983HA1D, 1985LI1C; theor.). The cross section for tritium production rises rapidly to 190 mb at 1 MeV, then more slowly to 290 mb near 4 MeV: see (1974AJ01). For VAP and TAP measurements at $E_d^- = 191$ and 395 MeV see (1986GA18).

9. (a) ${}^6\text{Li}(d, \alpha){}^4\text{He}$ $Q_m = 22.3716$ $E_b = 22.2798$
 (b) ${}^6\text{Li}(d, \alpha p){}^3\text{H}$ $Q_m = 2.5576$

Table 8.7: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, \gamma){}^8\text{Be}$ ^a

E_{res} (keV)	Γ_{lab} (keV)	${}^8\text{Be}^*$ (MeV)	l_p	J^π	Res. ^d
441.4 ± 0.5 ^b	12.2 ± 0.5	17.640	1	1^+	$\gamma_0, \gamma_1, \gamma_3, \gamma_4$
1030 ± 5	168	18.155	1	1^+	$\gamma_0, \gamma_1, \gamma_3, \gamma_4$
1890	150 ± 50	18.91		(2^-)	γ_3, γ_4
2060 ± 20	310 ± 20	19.06		$J = 1, 2, 3,$ $\pi = (-)$ ^c	γ_1
(3100)		(20.0)			γ_1
4900		21.5			γ_1
5000	≈ 4500	21.6	0	$1^-; T = 1$	γ_0
6000		22.5			γ_1
7500	≈ 8000	23.8	(0)	$(1^-, 2^-); T = 1$	γ_1
(11100)		(27.0)			γ_1
13000	broad	28.6			

^a See Tables 8.6 in (1974AJ01, 1979AJ01) for the references.

^b See (1959AJ76). See also (1983FI13, 1984JE1B).

^c See, however, reaction 16.

^d $\gamma_0, \gamma_1, \gamma_3, \gamma_4$ represent transitions to ${}^8\text{Be}^*(0, 3.0, 16.6, 16.9)$, respectively.

Cross sections and angular distributions (reaction (a)) have been measured at $E_d = 30$ keV to 31 MeV: see (1979AJ01, 1984AJ01). See also (1983SZZY). A critical analysis of the low-energy data has led to a calculation of the reaction rate parameters for thermonuclear reactions for plasma temperatures of 2 keV to 1 MeV: see (1984AJ01). Polarization measurements are reported in the range 0.4 to 11 MeV: see (1979AJ01, 1984AJ01) and see below.

Pronounced variations are observed in the cross sections and in the analyzing powers. Maxima are seen at $E_d = 0.8$ MeV, $\Gamma_{\text{lab}} \approx 0.8$ MeV and $E_d = 3.75$ MeV, $\Gamma_{\text{lab}} \approx 1.4$ MeV. The 4 MeV peak is also observed in the tensor component coefficients with $L = 0, 4$ and 8 and in the vector component coefficients: two overlapping resonances are suggested. At higher energies all coefficients show a fairly smooth behavior which suggests that only broad resonances can exist. The results are in agreement with those from reaction 4, that is with two 2^+ states at $E_x = 22.2$ and 25.2 MeV and a 4^+ state at 25.5 MeV. A strong resonance is seen in the α^* channel [to ${}^4\text{He}(20.1)$, $J^\pi = 0^+$] presumably due to ${}^8\text{Be}^*(25.2, 25.5)$. In addition the ratio of the α^*/α differential cross sections at 30° shows a broad peak centered at $E_x \approx 26.5$ MeV (which may be due to interference effects) and suggests a resonance-like anomaly at $E_x \approx 28$ MeV. $A_{yy} = 1$ points are reported at $E_d = 5.55 \pm 0.12$ ($\theta_{\text{c.m.}} = 29.7 \pm 1.0^\circ$) and 8.80 ± 0.25 MeV ($\theta_{\text{c.m.}} = 90.0 \pm 1.0^\circ$) [corresponds to $E_x = 26.44$ and 28.87 MeV]. For references see (1974AJ01, 1979AJ01).

At $E_d = 6.945$ MeV, the α_0 yield shows an anomaly corresponding to ${}^8\text{Be}^*(27.49)$, the 0^+ ; $T = 2$ analog of ${}^8\text{He}_{\text{g.s.}}$. This $T = 2$ state has recently been studied using both polarized deuterons and ${}^6\text{Li}$ ions. The ratio of the partial widths for decay into ${}^6\text{Li} + \text{d}$ states with channel spin 2 and 0, $\Gamma_2/\Gamma_1 = 0.322 \pm 0.091$ (1986SO07).

A kinematically complete study of reaction (b) has been reported at $E_d = 1.2$ to 8.0 MeV: the transition matrix element squared plotted as a function of $E_{\alpha\alpha^*}$ (the relative energy in the channel ${}^4\text{He}_{\text{g.s.}} + {}^4\text{He}^*(20.1) [0^+]$) shows a broad maximum at $E_x \approx 25$ MeV. Analysis of these results, and of a study of ${}^7\text{Li}(p, \alpha)\alpha^*$ [see reaction 18] which shows a peak of different shape at $E_x \approx 24$ MeV, indicate the formation and decay of overlapping states of high spatial symmetry, if the observed structures are interpreted in terms of ${}^8\text{Be}$ resonances: see (1984AJ01). For other work see (1984AJ01). See also ${}^6\text{Li}$, (1986ST1E), (1984VO1A, 1988KU1E; applications) and (1983HA1D, 1984KR1B, 1984KU15; theor.).

Table 8.8: ${}^8\text{Be}$ levels from ${}^7\text{Li}(p, p_0){}^7\text{Li}$ and ${}^7\text{Li}(p, p_1){}^7\text{Li}^*$ ^a

E_p (MeV)	Γ_{lab} (keV)	${}^8\text{Be}^*$ (MeV)	J^π	$\Gamma_{p'}$ (keV)
0.441	12.2 ^c	17.640 ^h	1^+	
1.030 ± 0.005	168	18.155	1^+	≈ 6
1.88 ^b	55 ± 20	18.90	2^-	
2.05	≈ 400	19.05	3^+	small
2.25		19.22	3^+	small
2.5 ^d	≈ 750	19.4	1^-	res
^e				
4.2 ± 0.2 ^f	1800 ± 200	20.9	4^-	(res)
5.6	broad	22.2	^g	res

^a See references in Table 8.9 (1979AJ01).

^b (p, n) threshold: see reaction 15.

^c $\theta_p^2 = 0.064$.

^d See also Table 8.7, γ_{n1}^2 and $\gamma_{p1}^2 \approx 1\%$ of Wigner limit.

^e A 2^+ state at $E_x \approx 20$ MeV appears to be necessary to account for the cross sections: see Table 8.3 and reaction 4.

^f Reduced width is 70% of the Wigner limit.

^g May be due to two 2^+ states. See also reaction 15.

^h See also (1981BA36; theor.).

10. ${}^6\text{Li}(t, n){}^8\text{Be}$

$Q_m = 16.0225$

At $E_t = 2$ to 4.5 MeV ${}^8\text{Be}^*(0, 3.0, 16.6, 16.9)$ are populated (1984LIZY; prelim.). See also (1966LA04, 1974AJ01).

11. (a) ${}^6\text{Li}({}^3\text{He}, p){}^8\text{Be}$ $Q_m = 16.7863$
 (b) ${}^6\text{Li}({}^3\text{He}, p)2\ {}^4\text{He}$ $Q_m = 16.8782$

Angular distributions have been studied in the range $E({}^3\text{He}) = 0.46$ to 17 MeV and at $E({}^6\text{Li}) = 21$ MeV. ${}^8\text{Be}^*(0, 3.0, 16.63, 16.92, 17.64, 18.15, 19.0, 19.4, 19.9)$ are populated in this reaction: see (1974AJ01, 1979AJ01, 1984AJ01). For reaction (b) see (1974AJ01) and (1987ZA07). See also ${}^9\text{B}$.

12. (a) ${}^6\text{Li}(\alpha, d){}^8\text{Be}$ $Q_m = -1.5669$
 (b) ${}^6\text{Li}(\alpha, 2\alpha){}^2\text{H}$ $Q_m = -1.4750$

Deuteron groups have been observed to ${}^8\text{Be}^*(0, 3.0, 11.3 \pm 0.4)$. Angular distributions have been measured at $E_\alpha = 15.8$ to 48 MeV: see (1974AJ01, 1979AJ01). A study of reaction (b) shows that the peak due to ${}^8\text{Be}^*(3.0)$ is best fitted by using $\Gamma = 1.2 \pm 0.3$ MeV. At $E_\alpha = 42$ MeV the α - α FSI is dominated by ${}^8\text{Be}^*(0, 3.0)$. See also Table 8.4 in (1974AJ01) and (1983BE51; theor.).

13. (a) ${}^6\text{Li}({}^6\text{Li}, \alpha){}^8\text{Be}$ $Q_m = 20.805$
 (b) ${}^6\text{Li}({}^6\text{Li}, \alpha)2\ {}^4\text{He}$ $Q_m = 20.897$
 (c) ${}^6\text{Li}({}^6\text{Li}, 2d)2\ {}^4\text{He}$ $Q_m = -2.950$

At $E_{\text{max}}({}^6\text{Li}) = 13$ MeV reaction (a) proceeds via ${}^8\text{Be}^*(0, 3.0, 16.6, 16.9, 22.5)$. The involvement of a state at $E_x = 19.9$ MeV ($\Gamma = 1.3$ MeV) is suggested. Good agreement with the shapes of the peaks corresponding to ${}^8\text{Be}^*(16.6, 16.9)$ is obtained by using a simple two-level formula with interference, corrected for the effect of final-state Coulomb interaction, assuming $\Gamma(16.6) = 90$ keV and $\Gamma(16.9) = 70$ keV: see also Table 8.6. The ratio of the intensities of the groups corresponding to ${}^8\text{Be}^*(16.6, 16.9)$ remains constant for $E({}^6\text{Li}) = 4.3$ to 5.5 MeV: $I(16.6)/I(16.9) = 1.22 \pm 0.08$. Partial angular distributions for the α_0 group have been measured at fourteen energies for $E({}^6\text{Li}) = 4$ to 24 MeV. See (1979AJ01) for the references.

At $E({}^6\text{Li}) = 36$ to 46 MeV sequential decay (reaction (b)) via ${}^8\text{Be}$ states at $E_x = 3.0, 11.4, 16.9$ and 19.65 MeV is reported: see (1984AJ01). (1987LA25) report the possible involvement of the 2^+ state ${}^8\text{Be}^*(22.2)$.

For reaction (c) see (1983WA09) and ${}^{12}\text{C}$ in (1985AJ01). See also (1983MI10) and (1982LA19, 1985NO1A; theor.).

14. ${}^7\text{Li}(p, \gamma){}^8\text{Be}$

$$Q_m = 17.2543$$

Cross sections and angular distributions have been reported from $E_p = 30$ keV to 18 MeV. Gamma rays are observed to the ground (γ_0) and to the broad, 2^+ , excited state at 3.0 MeV (γ_1) and to ${}^8\text{Be}^*(16.6, 16.9)$ (γ_3, γ_4). Resonances for both γ_0 and γ_1 occur at $E_p = 0.44$ and 1.03 MeV, and for γ_1 alone at 2, 4.9, 6.0, 7.3, and possibly at 3.1 and 11.1 MeV. In addition broad resonances are reported at $E_p \approx 5$ MeV (γ_0), $\Gamma \approx 4 - 5$ MeV, and at $E_p \approx 7.3$ MeV (γ_1), $\Gamma \approx 8$ MeV: see Table 8.7. The $E_p \approx 5$ MeV resonance ($E_x \approx 22$ MeV) represents the giant dipole resonance based on ${}^8\text{Be}(0)$ while the γ_1 resonance, ≈ 2.2 MeV higher, is based on ${}^8\text{Be}^*(3.0)$. The γ_0 and γ_1 giant resonance peaks each contain about 10% of the dipole sum strength. The main trend between $E_p = 8$ and 17.5 MeV is a decreasing cross section.

At the $E_p = 0.44$ MeV resonance ($E_x = 17.64$ MeV) the radiation is nearly isotropic consistent with p-wave formation, $J^\pi = 1^+$, with channel spin ratio $\sigma(J_c = 2)/\sigma(J_c = 1) = 3.2 \pm 0.5$. Radiative widths for the γ_0 and γ_1 decay are displayed in Table 8.5. A careful study of the α -breakup of ${}^8\text{Be}^*(16.63, 16.92)$ [both $J^\pi = 2^+$] for $E_p = 0.44$ to 2.45 MeV shows that the non-resonant part of the cross section for production of ${}^8\text{Be}^*(16.63)$ is accounted for by an extranuclear direct-capture process. Resonances for production of ${}^8\text{Be}^*(16.63, 16.92)$ are observed at $E_p = 0.44, 1.03$ and 1.89 MeV [${}^8\text{Be}^*(17.64, 18.15, 18.9)$]. The results are consistent with the hypothesis of nearly maximal isospin mixing for ${}^8\text{Be}^*(16.63, 16.92)$: decay to these states is not observed from the 3^+ states at $E_x = 19$ MeV, but rather from the 2^- state at $E_x = 18.9$ MeV. Squared $T = 1$ components calculated for ${}^8\text{Be}^*(16.6, 16.9)$ are 40 and 60%, and 95 and 5% for ${}^8\text{Be}^*(17.6, 18.2)$. The cross section for $(\gamma_3 + \gamma_4)$ has also been measured for $E_p = 11.5$ to 30 MeV ($\theta = 90^\circ$) by detecting the γ -rays and for $E_p = 4$ to 13 MeV (at five energies) by detecting the two α -particles from the decay of ${}^8\text{Be}^*(16.6, 16.9)$: a broad bump is observed at $E_p = 8 \pm 2$ MeV (1981MA33). The angle and energy integrated yield only exhausts 8.6% of the classical dipole sum for $E_p = 4$ to 30 MeV, suggesting that this structure does not represent the GDR built on ${}^8\text{Be}^*(16.6, 16.9)$. A weak, very broad [$\Gamma \geq 20$ MeV] peak may also be present at $E_x = 20-30$ MeV. A direct capture calculation adequately describes the observed cross section (1981MA33). A study of the γ -decay of ${}^8\text{Be}^*(17.64, 18.15)$ shows no evidence for a pseudoscalar particle postulated to account for narrow peaks in e^+ spectra in heavy-ion reactions (1988SA2A). For the earlier references see (1979AJ01). See also (1983CH1C), (1986WE1D), (1984DA1H; astrophysics), (1988KI1C; applied) and (1983GO1B, 1984SE16, 1985GO1B, 1987KI1C; theor.).

15. ${}^7\text{Li}(p, n){}^7\text{Be}$

$$Q_m = -1.644$$

$$E_b = 17.2543$$

Measurements of cross sections have been reported for $E_p = 1.9$ to 199.1 MeV [see (1974AJ01, 1979AJ01, 1984AJ01)] and in the range 60.1 to 480.0 MeV (1984DA22; activation σ). Polarization measurements have been reported at $E_p = 2.05$ to 5.5 MeV, 30 and 50 MeV [see (1974AJ01)] and at $E_p = 52.8$ MeV (1988HE08) [$K_z^{z'} = 0.07 \pm 0.02$]. See also below.

The yield of ground state neutrons (n_0) rises steeply from threshold and shows pronounced resonances at $E_p = 2.25$ and 4.9 MeV. The yield of n_1 also rises steeply from threshold and

exhibits a broad maximum near $E_p = 3.2$ MeV and a broad dip at $E_p \approx 5.5$ MeV, also observed in the p_1 yield. Multi-channel scattering length approximation analysis of the 2^- partial wave near the n_0 threshold indicates that the 2^- state at $E_x = 18.9$ MeV is virtual relative to the threshold and that its width $\Gamma = 50 \pm 20$ keV. The ratio of the cross section for ${}^7\text{Li}(p, \gamma){}^8\text{Be}^*(18.9) \rightarrow {}^8\text{Be}^*(16.6 + 16.9) + \gamma$ to the thermal neutron capture cross section ${}^7\text{Be}(n, \gamma){}^8\text{Be}^*(18.9) \rightarrow {}^8\text{Be}^*(16.6 + 16.9) + \gamma$, provides a rough estimate of the isospin impurity of ${}^8\text{Be}^*(18.9)$: $\sigma_{p,\gamma}/\sigma_{n,\gamma} \approx 1.5 \times 10^{-5}$. The $T = 1$ isospin impurity is $\leq 10\%$ in intensity. See also reaction 23. See (1979AJ01, 1984AJ01).

The structure at $E_p = 2.25$ MeV is ascribed to a 3^+ , $T = (1)$, $l = 1$ resonance with $\Gamma_n \approx \Gamma_p$ and $\gamma_n^2/\gamma_p^2 = 3$ to 10: see (1966LA04). At higher energies the broad peak in the n_0 yield at $E_p = 4.9$ MeV can be fitted by $J^\pi = 3^{(+)}$ with $\Gamma = 1.1$ MeV, $\gamma_n^2 \approx \gamma_p^2$. The behavior of the n_1 cross section can be fitted by assuming a 1^- state at $E_x = 19.5$ MeV and a $J = 0, 1, 2$, positive-parity state at 19.9 MeV [presumably the 20.1–20.2 MeV states reported in reaction 4]. In addition the broad dip at $E_p \approx 5.5$ MeV may be accounted for by the interference of two 2^+ states. See Table 8.8 in (1979AJ01). The 0° differential cross section increases rapidly to ≈ 35 mb/sr at 30 MeV and then remains constant to 100 MeV: see (1985BO1C). The total reaction cross section [${}^7\text{Be}^*(0, 0.43)$] decreases inversely with E_p in the range 60.1 to 480.0 MeV (1984DA22) [note: the values of σ_t supersede those reported earlier]. The transverse polarization transfer, $D_{NN}(0^\circ)$, for the g.s. transition has been measured at $E_p = 160$ MeV (1984TA07). See also (1986MC09; $E_p = 800$ MeV), (1987WAZT), (1984BA1U), (1985CA41; astrophysics), (1983LO12; applications), (1986RA21, 1987TA22) and (1988GU1F; theor.).

16. (a) ${}^7\text{Li}(p, p){}^7\text{Li}$

$E_b = 17.2543$

(b) ${}^7\text{Li}(p, p'){}^7\text{Li}^*$

Absolute differential cross sections for elastic scattering have been reported for $E_p = 0.4$ to 12 MeV and at 14.5, 20.0 and 31.5 MeV. The yields of inelastically scattered protons (to ${}^7\text{Li}^*(0.48)$) and of 0.48 MeV γ -rays have been measured for $E_p = 0.8$ to 12 MeV: see (1974AJ01). Polarization measurements have been reported at a number of energies in the range $E_p = 0.67$ MeV to 2.1 GeV/ c [see (1974AJ01, 1979AJ01, 1984AJ01)], at $E_p = 1.89$ to 2.59 MeV (1986SA1P; p_0 ; prelim.) and at 65 MeV (1987TO06; continuum; prelim.). See also (1983GLZZ).

Anomalies in the elastic scattering appear at $E_p = 0.44, 1.03, 1.88, 2.1, 2.5, 4.2$ and 5.6 MeV. Resonances at $E_p = 1.03, 3$ and 5.5 MeV and an anomaly at $E_p = 1.88$ MeV appear in the inelastic channel. A phase-shift analysis and a review of the cross-section data show that the 0.44 and 1.03 MeV resonances are due to 1^+ states which are a mixture of 5P_1 and 3P_1 with a mixing parameter of $+25^\circ$; that the 2^- state at the neutron threshold ($E_p = 1.88$ MeV) has a width of about 50 keV [see also reaction 14]; and that the $E_p = 2.05$ MeV resonance corresponds to a 3^+ state. The anomalous behavior of the 5P_3 phase around $E_p = 2.2$ MeV appears to result from the coupling of the two 3^+ states [resonances at $E_p = 2.05$ and 2.25 MeV]. The 3S_1 phase begins to turn positive after 2.2 MeV suggesting a 1^- state at $E_p = 2.5$ MeV: see Table 8.8. The polarization data show structures at $E_p = 1.9$ and 2.3 MeV. A phase-shift analysis of the (p, p) data finds

no indication of a possible 1^- state with $17.4 < E_x < 18.5$ MeV [see, however, reaction 15 in (1979AJ01)].

An attempt has been made to observe the $T = 2$ state [$^8\text{Be}^*(27.47)$] in the p_0 , p_1 and p_2 yields. None of these shows the effect of the $T = 2$ state. Table 8.5 in (1984AJ01) displays the upper limit for Γ_{p_0}/Γ .

The proton total reaction has been reported for $E_p = 25.1$ to 48.1 MeV by (1985CA36). (1987CH33, 1987PO03) have studied p - ^7Li correlations involving $^8\text{Be}^*(17.64, 18.15, 18.9 + 19.1 + 19.2)$. See also ^7Li (1984BA1U), (1986BA88), (1986RA1D; applications) and (1986HA1K, 1988GUIF; theor.) and the “General” section here.

$$17. \text{}^7\text{Li}(p, d)^6\text{Li} \qquad Q_m = -5.025 \qquad E_b = 17.2543$$

The excitation function for d_0 measured for $E_p = 11.64$ to 11.76 MeV does not show any effect from the $T = 2$ state [$^8\text{Be}^*(27.47)$]: see (1979AJ01). See also (1984BA1T).

$$18. \text{}^7\text{Li}(p, \alpha)^4\text{He} \qquad Q_m = 17.3462 \qquad E_b = 17.2543$$

The cross section increases from $(4.3 \pm 0.9) \times 10^{-5}$ mb at $E_p = 28.1$ keV to 6.33 mb at 998 keV. Astrophysical S -factors have been calculated over that range: $S(0) = 52 \pm 8$ keV \cdot b (1986RO13). For the earlier work see (1984AJ01).

Excitation functions and angular distributions have been measured at many energies in the range $E_p = 23$ keV to 62.5 MeV: see (1979AJ01, 1984AJ01). Polarization measurements have been carried out for $E_p = 0.8$ to 10.6 MeV [see (1974AJ01)]: in the range $E_p = 3$ to 10 MeV the asymmetry has one broad peak in the angular distribution at all energies except near 5 MeV; the peak value is 0.98 ± 0.04 at 6 MeV and is essentially 1.0 for $E_p = 8.5$ to 10 MeV.

Broad resonances are reported to occur at $E_p = 3.0$ MeV [$\Gamma \approx 1$ MeV] and at ≈ 5.7 MeV [$\Gamma \approx 1$ MeV]. Structures are also reported at $E_p = 6.8$ MeV and at $E_p = 9.0$ MeV: see (1979AJ01). The 9.0 MeV resonance is also reflected in the behavior of the A_2 coefficient. The experimental data on yields and on polarization appear to require including two 0^+ states [at $E_x \approx 19.7$ and 21.8 MeV] with very small α -particle widths, and four 2^+ states [at $E_x \approx 15.9, 20.1, 22.2$ and 25 MeV]. See, however, reaction 4. A 4^+ state near 20 MeV was also introduced in the calculation but its contribution was negligible. The observed discrepancies are said to be probably due to the assumption of pure $T = 0$ for these states. At $E_p = 11.64$ to 11.76 MeV the excitation function does not show any effect due to the $T = 2$ state at $E_x = 27.47$ MeV. See (1979AJ01) for references.

A study of the $^7\text{Li}(p, \alpha)^4\text{He}^*$ reaction to $^4\text{He}^*(20.1)$ [0^+] at $E_p = 4.5$ to 12.0 MeV shows a broad maximum at $E_x \approx 24$ MeV: see reaction 9 and (1984AJ01). See also (1986ZA09), (1984HA1M, 1984YA1A, 1985BO1K, 1985CA41, 1985DE1K, 1986BO1H, 1987AS05, 1987KA1R, 1987RO25, 1988BA86, 1988FO1A; astrophysics), (1986RA1D, 1986ST1E, 1986TU1B; applications) and (1984BL21, 1984KR1B; theor.).

19. (a) ${}^7\text{Li}(\text{d}, \text{n}){}^8\text{Be}$ $Q_{\text{m}} = 15.0297$
 (b) ${}^7\text{Li}(\text{d}, \text{n})2\text{ }{}^4\text{He}$ $Q_{\text{m}} = 15.1216$

The population of ${}^8\text{Be}^*(0, 3.0, 16.6, 16.9, 17.6, 18.2, 18.9, 19.1, 19.2)$ has been reported in reaction (a). For the parameters of ${}^8\text{Be}^*(3.0)$ see Table 8.4 in (1974AJ01). Angular distributions of n_0 and n_1 have been reported at $E_{\text{d}} = 0.7$ to 3.0 MeV and at $E_{\text{d}} = 15.25$ MeV [see (1974AJ01, 1979AJ01)] and at 0.19 MeV (1983DA32, 1987DA25) and 0.40 and 0.46 MeV (1984GA07; n_0 only). The angular distributions of the neutrons to ${}^8\text{Be}^*(16.6, 17.6, 18.2)$ are fit by $l_{\text{p}} = 1$: see (1974AJ01).

Reaction (b) at $E_{\text{d}} = 2.85$ to 14.97 MeV proceeds almost entirely through the excitation and sequential decay of ${}^8\text{Be}^*(16.6, 16.9)$ (1987WA21). ${}^8\text{Be}^*(11.4)$ may also be involved [$E_{\text{x}} = 11.4 \pm 0.05$ MeV, $\Gamma_{\text{c.m.}} = 2.8 \pm 0.2$ MeV] as may state(s) at $E_{\text{x}} \approx 20$ MeV: see (1979AJ01). See also ${}^9\text{Be}$, (1983BL17, 1986BA40), (1986LE1E; applications) and (1983MU13, 1984BL21; theor.).

20. (a) ${}^7\text{Li}({}^3\text{He}, \text{d}){}^8\text{Be}$ $Q_{\text{m}} = 11.7608$
 (b) ${}^7\text{Li}({}^3\text{He}, \alpha\text{d}){}^4\text{He}$ $Q_{\text{m}} = 11.8527$

Deuteron groups are observed to ${}^8\text{Be}^*(0, 3.0, 16.6, 16.9, 17.6, 18.2)$. For the parameters of ${}^8\text{Be}^*(3.0)$ see Table 8.4 in (1974AJ01). For the $J^{\pi} = 2^+$ mixed isospin states see Table 8.6. Angular distributions have been measured for $E({}^3\text{He}) = 0.9$ to 24.3 MeV and at $E({}^3\vec{\text{He}}) = 33.3$ MeV: see (1974AJ01, 1979AJ01, 1984AJ01). Reaction (b) has been studied at $E({}^3\text{He}) = 5.0$ MeV (1985DA29) and at $9, 11$ and 12 MeV (1986ZA09). ${}^8\text{Be}^*(0, 3.0)$ are reported to be involved (1985DA29). See also ${}^{10}\text{B}$ and (1983KU17; theor.).

21. (a) ${}^7\text{Li}(\alpha, \text{t}){}^8\text{Be}$ $Q_{\text{m}} = -2.5597$
 (b) ${}^7\text{Li}(\alpha, \alpha\text{t}){}^4\text{He}$ $Q_{\text{m}} = -2.4678$

Angular distributions have been measured to $E_{\alpha} = 50$ MeV: see (1966LA04, 1974AJ01, 1979AJ01). The ground state of ${}^8\text{Be}$ decays isotropically in the c.m. system: $J^{\pi} = 0^+$. Sequential decay (reaction (b)) is reported at $E_{\alpha} = 50$ MeV via ${}^8\text{Be}^*(0, 3.0, 11.4, 16.6, 16.9, 19.9)$: see (1974AJ01). See also (1983BE51, 1985PU03; theor.).

22. ${}^7\text{Li}({}^7\text{Li}, {}^6\text{He}){}^8\text{Be}$ $Q_{\text{m}} = 7.280$

${}^8\text{Be}^*(0, 3.0)$ have been populated in this reaction (1987BO1M; $E({}^7\text{Li}) = 22$ MeV). See also (1988AL1G).

23. (a) ${}^7\text{Be}(n, p){}^7\text{Li}$	$Q_m = 1.644$	$E_b = 18.8985$
(b) ${}^7\text{Be}(n, \alpha){}^4\text{He}$	$Q_m = 18.9905$	
(c) ${}^7\text{Be}(n, \gamma\alpha){}^4\text{He}$	$Q_m = 18.9905$	

The total (n, p) cross section has been measured from 25×10^{-3} eV to 13.5 keV. For thermal neutrons the cross sections to ${}^7\text{Li}^*(0, 0.48)$ are $38\,400 \pm 800$ and 420 ± 120 b, respectively. A departure from a $1/v$ shape in σ_t is observed for $E_n > 100$ eV. The astrophysical reaction rate is $\approx \frac{1}{3}$ lower than that previously used: this could lead to an increase in the calculated rate of production of ${}^7\text{Li}$ in the Big Bang by as much as 20%. A multi-level R -matrix analysis of the data indicates $\Gamma = 122$ keV for the 2^- state ${}^8\text{Be}^*(18.9)$, and a $T = 1$ impurity of $\approx 24\%$ (1988KO03). At thermal energies the (n, α) cross section is ≤ 0.1 mb and the (n, $\gamma\alpha$) cross section is 155 mb: see (1974AJ01). See also (1987GLZZ, 1987GL1D), (1979AJ01, 1988BO15) and (1984YA1A, 1985BO1K, 1985DE1K; astrophysics).

24. ${}^8\text{Li}(\beta^-){}^8\text{Be}$	$Q_m = 16.0039$
---	-----------------

${}^8\text{Li}$ decays to the broad 3.0 MeV, 2^+ level of ${}^8\text{Be}$, which decays into two α -particles. Both the β -spectrum and the resulting α -spectrum have been extensively studied: see (1955AJ61, 1966LA04). See also ${}^8\text{B}(\beta^+)$. Studies of the distribution of recoil momenta and neutrino recoil correlations indicate that the decay is overwhelmingly GT, axial vector [see reaction 1 in ${}^8\text{Li}$] and that the ground state of ${}^8\text{Li}$ has $J^\pi = 2^+$: see (1980MC07).

(1986WA01) has performed a many-level one-channel approximation R -matrix analysis of the β -delayed α -particle spectra in the decay of both ${}^8\text{Li}$ and ${}^8\text{B}$, obtained by (1971WI05) [as well as of the $L = 2$ α - α phase shifts]. Warburton finds that there is no need to introduce “intruder” states below $E_x \approx 26$ MeV [see, e.g., (1974AJ01)]. He extracts the GT matrix elements for the decay to ${}^8\text{Be}^*(3.0)$ and the doublet near 16 MeV; and he points out the difficulties in extracting meaningful E_x and Γ values from the β^\pm decay for ${}^8\text{Be}^*(3.0)$, as well as the $\log ft$ values for the transitions to that state (1986WA01).

Beta- α angular correlations have been measured for the decays of ${}^8\text{Li}$ and ${}^8\text{B}$ for the entire final-state distribution: see Table 8.10 in (1979AJ01). (1980MC07) have measured the β - ν - α correlations as a function of E_x in the decay of ${}^8\text{Li}$ and ${}^8\text{B}$, detecting both α -particles involved in the ${}^8\text{Be}$ decay. They find that the decay is GT for $2 < E_x < 8$ MeV. The absence of Fermi decay strength is expected because the isovector contributions from the tails of ${}^8\text{Be}^*(16.6, 16.9)$ interfere destructively in this energy region: see (1980MC07). The measurement of the β -decay asymmetry as a function of E_β is reported by (1986BI1D, 1985BIZZ; prelim.). (1986NAZZ; prelim.) have measured the β -spectrum and compared it with the spectrum predicted from the α -breakup data. See also (1984KO25, 1985GR1A), (1986HA1P, 1988WA1E), (1986MA1T, 1986NAZZ; astrophysics) and (1983KU17, 1984BA25, 1986QUZZ, 1987LY05, 1988BA75; theor.).

25. ${}^8\text{B}(\beta^+){}^8\text{Be}$

$$Q_m = 17.979$$

The decay [see reaction 1 in ${}^8\text{B}$] proceeds mainly to ${}^8\text{Be}^*(3.0)$ [see Table 8.4 in (1974AJ01) for its parameters]. Detailed study of the high-energy portion of the α -spectrum reveals a maximum near $E_\alpha = 8.3$ MeV, corresponding to transitions to ${}^8\text{Be}^*(16.63)$, for which parameters $E_x = 16.67$ MeV, $\Gamma = 150$ to 190 keV or $E_x = 16.62$ MeV, $\Gamma = 95$ keV are derived: see (1974AJ01). Log ft for the transition to ${}^8\text{Be}^*(16.6)$ is 3.3. An analysis by (1986WA01) of the β^+ delayed α -spectrum is described in reaction 24. See also (1988WA1E) and (1988BA75; theor.). The β^+ spectrum has been measured by (1987NA08) for momenta greater than 9 MeV/c. Then using the α spectra from (1986WA01) the ${}^8\text{B}$ neutrino spectrum is calculated. The average cross section for the “solar neutrino” ${}^{37}\text{Cl}(\nu_e, e^-){}^{37}\text{Ar}$ reaction is then $(1.07 \pm 0.02) \times 10^{-42}$ cm² [certain corrections may increase this value by as much as 4%] (1987NA08). See also (1982BA1J, 1983CO1D, 1983FO1A, 1983HA1B, 1983VO1C, 1984DA1H, 1984HA1M, 1985BA1N, 1985BA1M, 1985CH1B, 1986BA21, 1986BE1K, 1986DE1H, 1986GR04, 1986HA1I, 1986MA1T, 1986RO1N, 1986WO1B, 1987BA1X, 1987BA89, 1987CH1G, 1987FR1C, 1987FU1G, 1987KR10, 1987RI1E, 1987WE1C, 1988BA86, 1988EW1A, 1988HA1M; astrophysics).

26. (a) ${}^9\text{Be}(\gamma, n){}^8\text{Be}$	$Q_m = -1.6654$
(b) ${}^9\text{Be}(n, 2n){}^8\text{Be}$	$Q_m = -1.6654$
(c) ${}^9\text{Be}(p, pn){}^8\text{Be}$	$Q_m = -1.6654$
(d) ${}^9\text{Be}(t, tn){}^8\text{Be}$	$Q_m = -1.6654$
(e) ${}^9\text{Be}(\alpha, \alpha n){}^8\text{Be}$	$Q_m = -1.6654$

Neutron groups to ${}^8\text{Be}^*(0, 3.0)$ have been studied for $E_\gamma = 18$ to 26 MeV: see (1974AJ01, 1979AJ01) and ${}^9\text{Be}$. Reaction (b) appears to proceed largely via excited states of ${}^9\text{Be}$ with subsequent decay mainly to ${}^8\text{Be}^*(3.0)$: see (1966LA04, 1974AJ01), ${}^9\text{Be}$ and ${}^{10}\text{Be}$. Reaction (c) has been studied at $E_p = 45$ and 47 MeV: the reaction primarily populates ${}^8\text{Be}^*(0, 3.0)$: see (1979AJ01), ${}^9\text{Be}$ and ${}^9\text{B}$. For work at $E_p = 1$ GeV see (1985BE30, 1985DO16). For reactions (d) and (e) see (1974AJ01) and ${}^9\text{Be}$. For reaction (e) see (1979AJ01).

27. (a) ${}^9\text{Be}(p, d){}^8\text{Be}$	$Q_m = 0.5592$
(b) ${}^9\text{Be}(p, pn){}^8\text{Be}$	$Q_m = -1.6654$
(c) ${}^9\text{Be}(p, d)2\text{ }^4\text{He}$	$Q_m = 0.6511$

Angular distributions of deuteron groups have been reported at $E_p = 0.11$ to 185 MeV [see (1974AJ01, 1979AJ01, 1984AJ01)] and at 18.6 MeV (1986GO23, 1987GO27; d_0 and d_1) and 50 and 72 MeV (1984ZA07; to ${}^8\text{Be}^*(0, 3.0, 16.9, 19.2)$). For spectroscopic factors see (1979AJ01,

1984ZA07). The angular distributions to ${}^8\text{Be}^*(0, 3.0, 16.9, 17.6, 18.2, 19.1)$ are consistent with $l_n = 1$: see (1974AJ01).

An anomalous group is reported in the deuteron spectra between the d_0 and the d_1 groups. At $E_p = 26.2$ MeV, its (constant with θ) $E_x = 0.6 \pm 0.1$ MeV. Analyses of the spectral shape and transfer cross sections are consistent with this “ghost” feature being part of the Breit-Wigner tail of the $J^\pi = 0^+ {}^8\text{Be}_{g.s.}$: it contains $< 10\%$ of the g.s. transfer strength. An analysis of reported $\Gamma_{c.m.}$ for ${}^8\text{Be}^*(3.0)$ in this reaction shows that there is no E_p dependence. The average $\Gamma_{c.m.}$ at $E_p = 14.3$ and 26.2 MeV is 1.47 ± 0.04 MeV. $\Gamma_{c.m.} = 5.5 \pm 1.3$ eV for ${}^8\text{Be}_{g.s.}$ and 5.2 ± 0.1 MeV for ${}^8\text{Be}^*(11.4)$. Spectroscopic factors for ${}^8\text{Be}_{g.s.}$ (including the “ghost” anomaly) and ${}^8\text{Be}^*(3.0)$ are 1.23 and 0.22 respectively at $E_p = 14.3$ MeV, and 1.53 and 1.02 respectively at $E_p = 26.2$ MeV. The width of ${}^8\text{Be}^*(3.0)$ is not appreciably ($< 10\%$) reaction dependent but the nearness of the decay threshold indicates that care must be taken in comparing decay widths from reaction and from scattering data: $E_R = 3130 \pm 25$ keV (resonance energy in the $\alpha + \alpha$ c.m. system) [$E_x = 3038 \pm 25$ keV] and $\Gamma_{c.m.} = 1.50 \pm 0.02$ MeV for ${}^8\text{Be}^*(3.0)$: the corresponding observed and formal reaction widths and channel radii are $\gamma_R^2 = 580 \pm 50$ keV, $\gamma_\lambda^2 = 680 \pm 100$ keV and $s = 4.8$ fm. See (1979AJ01) for the earlier work. A study of the continuum part of the inclusive deuteron spectra is reported at $E_p = 60$ MeV (1987KA25). For reaction (b) see (1988BO47). For reaction (c) [FSI through ${}^8\text{Be}^*(0, 3.0)$] see (1974AJ01, 1984AJ01). See also (1985PU03; theor.) and ${}^{10}\text{B}$.

$$\begin{aligned} 28. \text{ (a) } & {}^9\text{Be}(d, t){}^8\text{Be} & Q_m &= 4.5919 \\ & \text{(b) } & {}^9\text{Be}(d, t)2\text{ }^4\text{He} & Q_m &= 4.6838 \end{aligned}$$

Angular distributions have been measured for $E_d = 0.3$ to 28 MeV [see (1979AJ01)], at $E_d = 18$ MeV (1988GO02; t_0, t_1) and at $E_d = 2.0$ to 2.8 MeV (1984AN16; t_0). At $E_d = 28$ MeV angular distributions of triton groups to ${}^8\text{Be}^*(16.6, 16.9, 17.6, 18.2, 19.1, 19.2, 19.8)$ have been analyzed using DWUCK: absolute C^2S are 0.074, 1.56, 0.22, 0.17, 0.41, 0.48, 0.40 respectively. See also Table 8.6. An isospin amplitude impurity of 0.21 ± 0.03 is found for ${}^8\text{Be}^*(17.6, 18.2)$: see (1979AJ01).

A kinematically complete study of reaction (b) at $E_d = 26.3$ MeV indicates the involvement of ${}^8\text{Be}^*(0, 3.0, 11.4, 16.9, 19.9 + 20.1)$: see (1974AJ01). (1986PAZN; prelim.) report $E_x = 3.10 \pm 0.15$ MeV, $\Gamma \approx 0.9 - 1.3$ MeV. See also (1988NE1A; theor.).

$$\begin{aligned} 29. \text{ (a) } & {}^9\text{Be}({}^3\text{He}, \alpha){}^8\text{Be} & Q_m &= 18.9124 \\ & \text{(b) } & {}^9\text{Be}({}^3\text{He}, \alpha)2\text{ }^4\text{He} & Q_m &= 19.0043 \end{aligned}$$

Angular distributions have been measured in the range $E({}^3\text{He}) = 3.0$ to 26.7 MeV and at $E({}^3\text{He}) = 33.3$ MeV (to ${}^8\text{Be}^*(16.9, 17.6, 19.2)$) [$S = 1.74, 0.72, 1.17$, assuming mixed isospin for ${}^8\text{Be}^*(16.9)$]. The possibility of a broad state at $E_x \approx 25$ MeV is also suggested: see (1979AJ01). See also (1987VA1I).

Reaction (b) has been studied at $E(^3\text{He}) = 1.0$ to 10 MeV [see (1979AJ01, 1984AJ01)], at $E(^3\text{He}) = 3$ to 12 MeV (1986LA26) and at 11.9 to 24.0 MeV (1987WA25). The reaction is reported to proceed via $^8\text{Be}^*(0, 3.0, 11.4, 16.6, 16.9, 19.9, 22.5)$: see (1979AJ01) and (1986LA26, 1987WA25). For a discussion of the width of $^8\text{Be}^*(11.4)$ see (1987WA25). See also ^9Be , and ^{12}C in (1980AJ01), (1985MC1C, applications) and (1985PU03; theor.).

30. (a) $^9\text{Be}(^6\text{Li}, ^7\text{Li})^8\text{Be}$ $Q_m = 5.585$
 (b) $^9\text{Be}(^7\text{Li}, ^8\text{Li})^8\text{Be}$ $Q_m = 0.367$
 (c) $^9\text{Be}(^9\text{Be}, ^{10}\text{Be})^8\text{Be}$ $Q_m = 5.1466$

Angular distributions have been studied at $E(^6\text{Li}) = 32$ MeV involving $^8\text{Be}^*(0, 3.0)$ and $^7\text{Li}^*(0, 0.48)$ (1985CO09). For reaction (b) see (1984KO25). For reaction (c) see ^{10}Be (1985JA09). For the earlier work see (1979AJ01).

31. $^{10}\text{Be}(p, t)^8\text{Be}$ $Q_m = 0.0045$

Angular distributions for the transition to the first $T = 2$ state $^8\text{Be}^*(27.49)$, and to $^8\text{Li}^*(10.82)$ reached in the $(p, ^3\text{He})$ reaction, are very similar. They are both consistent with $L = 0$ using a DWBA (LZR) analysis: see (1979AJ01, 1984AJ01) and Table 8.5 in (1984AJ01).

32. $^{10}\text{B}(\pi^+, 2p)^8\text{Be}$ $Q_m = 132.100$

See (1988RIZZ; prelim).

33. $^{10}\text{B}(n, t)^8\text{Be}$ $Q_m = 0.2307$

The breakup of ^{10}B by 14.4 MeV neutrons involves, among others, $^8\text{Be}_{g.s.}$ (1984TU02). See also (1979AJ01) and ^{11}B in (1990AJ01).

34. $^{10}\text{B}(p, ^3\text{He})^8\text{Be}$ $Q_m = -0.5332$

Angular distributions of the ^3He ions to $^8\text{Be}^*(0, 3.0, 16.6, 16.9)$ have been studied at $E_p = 39.4$ MeV [see (1974AJ01)] and at $E_p = 51.9$ MeV (1983YA05; see for a discussion of isospin mixing of the 16.8 MeV states).

35. (a) $^{10}\text{B}(\text{d}, \alpha)^8\text{Be}$ $Q_{\text{m}} = 17.8202$
 (b) $^{10}\text{B}(\text{d}, \alpha)^2\ ^4\text{He}$ $Q_{\text{m}} = 17.9121$

Angular distributions have been reported at $E_{\text{d}} = 0.5$ to 7.5 MeV: see (1974AJ01, 1979AJ01). At $E_{\text{d}} = 7.5$ MeV the population of $^8\text{Be}^*(16.63, 16.92)$ is closely the same consistent with their mixed isospin character while $^8\text{Be}^*(17.64)$ is relatively weak consistent with its nearly pure $T = 1$ character. $^8\text{Be}^*(16.63, 16.92, 17.64, 18.15)$ have been studied for $E_{\text{d}} = 4.0$ to 12.0 MeV. Interference between the 2^+ states [$^8\text{Be}^*(16.63, 16.92)$] varies as a function of energy. The cross-section ratios for formation of $^8\text{Be}^*(17.64, 18.15)$ vary in a way consistent with a change in the population of the $T = 1$ part of the wave function over the energy range: at the higher energies, there is very little isospin violation. At higher E_{x} only the 3^+ state at $E_{\text{x}} = 19.2$ MeV is observed, the neighboring 3^+ state at $E_{\text{x}} = 19.07$ MeV is not seen. $\Gamma_{16.6} = 90 \pm 5$ keV, $\Gamma_{16.9} = 70 \pm 5$ keV, $\Delta Q = 290 \pm 7$ keV: see Table 8.6 and (1979AJ01).

Reaction (b) [$E_{\text{d}} < 5$ MeV] takes place mainly by a sequential process involving $^8\text{Be}^*(0, 2.9, 11.4, 16.6, 16.9)$: see (1979AJ01). See also (1983DA11). [The work quoted in (1984AJ01) has not been published.] At $E_{\text{d}} = 13.6$ MeV in addition to $^8\text{Be}^*(16.6, 16.9)$, states with $E_{\text{x}} \approx 19.9$ – 20.2 MeV with $\Gamma \approx 0.7$ – 1.1 MeV are involved (1988KA1K; prelim.). See also (1984SH1D, 1984SHZJ) and (1985PU03, 1988BA75, 1988KA1M; theor.).

36. $^{10}\text{B}(\alpha, ^6\text{Li})^8\text{Be}$ $Q_{\text{m}} = -4.552$

See ^6Li here and reaction 40 in (1984AJ01). See also (1984SH1D, 1988SH1E).

37. (a) $^{11}\text{B}(\text{p}, \alpha)^8\text{Be}$ $Q_{\text{m}} = 8.5906$
 (b) $^{11}\text{B}(\text{p}, \alpha)^2\ ^4\text{He}$ $Q_{\text{m}} = 8.6825$

Angular distributions have been measured at $E_{\text{p}} = 0.78$ to 45 MeV [see (1974AJ01, 1979AJ01, 1984AJ01)], at $E = 0.12$ to 1.10 MeV (1987BE17; ^{11}B and p ; α_0, α_1) and at $E_{\text{p}} = 4.5$ to 7.5 MeV (1983BO19; α_0). Reaction (b) has been studied for $E_{\text{p}} = 0.15$ to 20 MeV: see (1974AJ01, 1984AJ01). The reaction proceeds predominantly by sequential two-body decay via $^8\text{Be}^*(0, 3.0)$. See also ^{12}C in (1990AJ01) and (1983CO1A, 1985MAZG, 1985PU03; theor.).

38. $^{11}\text{B}(^3\text{He}, ^6\text{Li})^8\text{Be}$ $Q_{\text{m}} = 4.5721$

At $E(^3\text{He}) = 71.8$ MeV angular distributions of the ^6Li ions to $^8\text{Be}^*(0, 3.0, 16.6, 16.9, 17.6, 18.2)$ are reported (1986JA14). For the earlier work at 25.6 MeV see (1979AJ01). See also (1986JA02).

39. $^{11}\text{B}(\alpha, ^7\text{Li})^8\text{Be}$ $Q_m = -8.7556$

The work reported in (1984AJ01) has not been published. See also ^7Li here and (1984SH1D, 1988SH1E).

40. $^{11}\text{B}(^9\text{Be}, ^{12}\text{B})^8\text{Be}$ $Q_m = 1.705$

See (1984DA17) and ^{12}B in (1990AJ01).

41. (a) $^{12}\text{C}(\text{n}, \text{n}\alpha)^8\text{Be}$ $Q_m = -7.3666$
 (b) $^{12}\text{C}(\text{p}, \text{p}\alpha)^8\text{Be}$ $Q_m = -7.3666$
 (c) $^{12}\text{C}(\text{p}, \text{d}^3\text{He})^8\text{Be}$ $Q_m = -25.7198$

The first two of these reactions involve $^8\text{Be}^*(0, 3.0)$: see (1974AJ01, 1979AJ01, 1984AJ01, 1985AJ01). See also (1986AN22) [reaction (a)] and (1982ZH06, 1985GA1B, 1986VD01; theor.). For reaction (c) see (1983LI18; theor.).

42. (a) $^{12}\text{C}(\text{d}, ^6\text{Li})^8\text{Be}$ $Q_m = -5.8916$
 (b) $^{12}\text{C}(\text{d}, \text{d}\alpha)^8\text{Be}$ $Q_m = -7.3666$

Angular distributions have been studied at $E_d = 12.7$ to 54.3 MeV [see (1974AJ01, 1979AJ01, 1984AJ01)] and at $E_d = 18$ and 22 MeV (1986YA12; to $^8\text{Be}_{\text{g.s.}}$; also VAP, TAP) and 51.7 MeV (1986YA12; to $^8\text{Be}^*(0, 3.0, 11.4)$; also VAP) as well as at $E_d = 50$ MeV (1987GO1S), 54.2 MeV (1984UM04; FRDWBA) [$S_\alpha = 0.48, 0.51$ and 0.82 for $^8\text{Be}^*(0, 3.0, 11.4)$] and 78.0 MeV (1986JA14; to $^8\text{Be}^*(0, 3.0, 16.6, 16.9)$). See also (1985GO1G; $E_d = 50$ MeV). For reaction (b) see (1984AJ01). See also (1984NE1A) and (1983GA14, 1983SH39, 1985GA1B, 1987KA1L; theor.).

43. $^{12}\text{C}(\text{t}, ^7\text{Li})^8\text{Be}$ $Q_m = -4.8988$

See ^7Li .

44. $^{12}\text{C}(^3\text{He}, ^7\text{Be})^8\text{Be}$ $Q_m = -5.7793$

Angular distributions have been obtained at $E(^3\text{He}) = 25.5$ to 70 MeV [see (1979AJ01, 1984AJ01)] and at $E(^3\vec{\text{He}}) = 33.4$ MeV (1986CL1B; $^8\text{Be}_{\text{g.s.}}$; also A_y ; prelim.). $^8\text{Be}^*(0, 3.0, 11.4, 16.6, 16.9, 17.6)$ have been populated. See also (1986RA15; theor.).

$$45. \text{ (a) } ^{12}\text{C}(\alpha, 2\alpha)^8\text{Be} \quad Q_{\text{m}} = -7.3666$$

$$\text{ (b) } ^{12}\text{C}(\alpha, ^8\text{Be})^8\text{Be} \quad Q_{\text{m}} = -7.4585$$

These reactions have been studied at E_{α} to 104 MeV [see (1979AJ01, 1984AJ01, and ^{12}C in 1985AJ01)] and at 31.2 MeV (1986XI1A; reaction (a)): $^8\text{Be}^*(0, 3.0, 11.4)$ are populated. See also (1984ZE1A, 1985GA1B, 1987KO1E; theor.).

$$46. \text{ (a) } ^{12}\text{C}(^9\text{Be}, ^{13}\text{C})^8\text{Be} \quad Q_{\text{m}} = 3.2810$$

$$\text{ (b) } ^{12}\text{C}(^{11}\text{B}, ^{15}\text{N})^8\text{Be} \quad Q_{\text{m}} = 3.6250$$

Angular distributions involving $^8\text{Be}_{\text{g.s.}} + ^{13}\text{C}_{\text{g.s.}}$ (reaction (a)) have been reported at $E(^9\text{Be}) = 20$ to 22.9 MeV and $E(^{12}\text{C}) = 10.5$ to 13.5 MeV: see (1984AJ01). For both reactions see also (1983DEZW).

$$47. \text{ (a) } ^{12}\text{C}(^{12}\text{C}, ^{16}\text{O})^8\text{Be} \quad Q_{\text{m}} = -0.2047$$

$$\text{ (b) } ^{12}\text{C}(^{16}\text{O}, ^{20}\text{Ne})^8\text{Be} \quad Q_{\text{m}} = -2.631$$

$$\text{ (c) } ^{12}\text{C}(^{20}\text{Ne}, \alpha^{20}\text{Ne})^8\text{Be} \quad Q_{\text{m}} = -7.3666$$

For reaction (a) see ^{16}O in (1986AJ04), (1983DEZW, 1984HU1E, 1984SP1C, 1986ALZN, 1986SH10) and (1984DA1B; theor.). For reaction (b) see reaction 18 in ^{20}Ne (1987AJ02), (1985MU14) and (1988AL07; location of a 10^+ state in ^{20}Ne at $E_x \approx 27.5$ MeV). For reaction (c) see (1987SI06).

$$48. ^{13}\text{C}(\text{d}, ^7\text{Li})^8\text{Be} \quad Q_{\text{m}} = -3.5879$$

See ^7Li .

$$49. ^{13}\text{C}(\alpha, ^9\text{Be})^8\text{Be} \quad Q_{\text{m}} = -10.7395$$

See (1984SH1D, 1988SH1F; prelim.; $E_{\alpha} = 27.2$ MeV) and ^9Be in (1979AJ01).

50. $^{13}\text{C}(^9\text{Be}, ^{14}\text{C})^8\text{Be}$ $Q_m = 6.511$

See ^{14}C in (1986AJ01).

51. $^{14}\text{N}(n, ^7\text{Li})^8\text{Be}$ $Q_m = -8.9139$

See ^7Li .

52. $^{16}\text{O}(p, p2\alpha)^8\text{Be}$ $Q_m = -14.5286$

See (1986VD04; $E_p = 50$ MeV).

53. $^{16}\text{O}(^{16}\text{O}, ^{24}\text{Mg})^8\text{Be}$ $Q_m = -0.483$

See (1987CZ02).

⁸B
(Figs. 3 and 4)

GENERAL: See also (1984AJ01).

Model calculations: (1983SH38).

Special states: (1982PO12, 1988KH03).

Complex reactions involving ⁸B: (1982AL08, 1983OL1A, 1984GR08, 1986HA1B, 1987TAZU, 1988AR05, 1988KI05).

Astrophysical questions: (1984HA1B, 1985BO1E, 1985GI1C, 1985KL1A, 1985LA1C, 1988BA86).

Reactions involving pions: (1983SP06).

Hypernuclei: (1983SH38).

Other topics: (1985AN28).

Ground state of ⁸B: (1983ANZQ, 1985AN28, 1986GL1A, 1987VA26, 1988AR05, 1988VA03)

$$\mu = 1.0355 \pm 0.0003 \text{ nm: see (1978LEZA).}$$

1. ⁸B(β^+)⁸Be $Q_m = 17.979$

The β^+ decay leads mainly to ⁸Be*(3.0). The mean of half-lives listed in (1974AJ01) is 770 ± 3 msec; $\log ft = 5.6$. There is also a branch to ⁸Be*(16.63): see (1986WA01) and reactions 24 and 25 in ⁸Be. $\log ft = 3.3$. See also (1985GR1A) and (1986QUZZ; theor.).

2. ⁶Li(d, π^-)⁸B $Q_m = -135.267$

At $E_d = 300$ and 600 MeV ⁸B*(0, 0.77, 2.32) are populated: see (1984AJ01).

3. ⁶Li(³He, n)⁸B $Q_m = -1.975$

Angular distributions for the n_0 group have been reported at $E(^3\text{He}) = 4.8$ to 5.7 MeV: $L = 0$. Two measurements for the E_x of ⁸B*(0.77) are 767 ± 12 and 783 ± 10 keV [$\Gamma = 40 \pm 10$ keV]: see (1974AJ01) and ⁹B.




Fig. 3: Energy levels of ${}^8\text{B}$. For notation see Fig. 1.

Table 8.9: Energy levels of ${}^8\text{B}$

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$2^+; 1$	$\tau_{1/2} = 770 \pm 3$ msec	β^+	1, 2, 3, 4, 5, 6, 7, 8, 9
0.774 ± 6		$\Gamma = 37 \pm 5$	γ, p	2, 3, 4, 6, 8, 9
2.32 ± 30	$3^+; 1$	350 ± 40		4, 8, 9
10.619 ± 9	$0^+; 2$	< 60		9

4. ${}^7\text{Li}(\text{p}, \pi^-){}^8\text{B}$ $Q_{\text{m}} = -140.293$

Angular distributions and analyzing powers have been measured for the transitions to ${}^8\text{B}^*(0, 0.77, 2.32)$ at $E_{\text{p}} = 199.2$ MeV: the A_y to ${}^8\text{B}^*(2.32)$ is characteristic of that to a stretched high-spin, two-particle one-hole final state [J^π of ${}^8\text{B}^*(2.32)$ is 3^+] (1987CA06). See also (1987CA05).

5. ${}^7\text{Li}({}^7\text{Li}, {}^6\text{H}){}^8\text{B}$ $Q_{\text{m}} = -35.01$

See ${}^6\text{H}$.

6. ${}^7\text{Be}(\text{p}, \gamma){}^8\text{B}$ $Q_{\text{m}} = -0.138$

Absolute cross sections have been measured for $E_{\text{p}} = 134$ keV to 10.0 MeV. A resonance is observed at $E_{\text{p}} = 723$ keV [$E_{\text{R}}(\text{c.m.}) = 632 \pm 10$ keV; $E_{\text{x}} = 770 \pm 10$ keV], $\Gamma_{\text{c.m.}} = 37 \pm 5$ keV [assuming $\Gamma_{\text{p}} \gg \Gamma_{\gamma}$] and $\sigma_{\text{peak}} = 1.18 \pm 0.12$ μb . Γ_{γ} is then 25 ± 4 meV. The zero-energy cross-section factor $S_{17}(0) = 0.0238 \pm 0.0023$ keV \cdot b (1983FI13). See (1979AJ01) for the earlier work, and the discussion in (1986BA38). See also (1984AJ01), (1984HA1F, 1987SA1L) and (1982BA1J, 1982KA1E, 1983BA45, 1983FO1A, 1983HA1B, 1984DA1H, 1984HA1M, 1984YA1A, 1985BA1Q, 1985CA41, 1985FI1D, 1986FI15, 1987KI01, 1987RO25, 1988BA86, 1988BA29, 1988FO1A; astrophysics).

7. ${}^7\text{Be}(\text{d}, \text{n}){}^8\text{B}$ $Q_{\text{m}} = -2.087$

See (1983HA17, 1985HA40).

8. ${}^{10}\text{B}(\text{p}, \text{t}){}^8\text{B}$ $Q_{\text{m}} = -18.530$

At $E_p = 49.5$ MeV [see (1974AJ01)] and 51.9 MeV (1983YA05) angular distributions have been measured for the tritons to ${}^8\text{B}^*(0, 2.32)$: $L = 2$ and $L = 0 + 2$ leading to $J^\pi = 2^+$ and 3^+ , respectively. Measurements of E_x for ${}^8\text{B}^*(2.32)$ yield 2.29 ± 0.05 MeV, 2.34 ± 0.04 MeV [$\Gamma_{\text{lab}} = 0.39 \pm 0.04$ MeV]. ${}^8\text{B}^*(0.77)$ is also observed: see (1974AJ01).

9. ${}^{11}\text{B}({}^3\text{He}, {}^6\text{He}){}^8\text{B}$ $Q_m = -16.913$

At $E({}^3\text{He}) = 72$ MeV the first $T = 2$ state is observed at $E_x = 10.619 \pm 0.009$ MeV, $\Gamma < 60$ keV: $d\sigma/d\Omega$ (lab) = 190 nb/sr at $\theta_{\text{lab}} = 9^\circ$. No other states are observed within 2.4 MeV of this state. ${}^8\text{B}^*(0, 0.77, 2.32)$ have also been populated: see (1979AJ01).

${}^8\text{C}$
(Fig. 4)

Mass of ${}^8\text{C}$: The atomic mass excess of ${}^8\text{C}$ is 35095 ± 24 keV (1985WA02); $\Gamma_{\text{c.m.}} = 230 \pm 50$ keV: see (1979AJ01). ${}^8\text{C}$ is stable with respect to ${}^7\text{B} + \text{p}$ ($Q = -0.13$ MeV) and unstable with respect to ${}^6\text{Be} + 2\text{p}$ ($Q = 21.4$), ${}^5\text{Li} + 3\text{p}$ ($Q = 1.55$), ${}^4\text{He} + 4\text{p}$ ($Q = 3.51$). At $E({}^3\text{He}) = 76$ MeV the differential cross section for formation of ${}^8\text{C}_{\text{g.s.}}$ in the ${}^{14}\text{N}({}^3\text{He}, {}^9\text{Li})$ reaction is ≈ 5 nb/sr at $\theta_{\text{lab}} = 10^\circ$. The ${}^{12}\text{C}(\alpha, {}^8\text{He}){}^8\text{C}$ reaction has been studied at $E_\alpha = 156$ MeV: $d\sigma/d\Omega \approx 20$ nb/sr at $\theta_{\text{lab}} = 20^\circ$: see (1979AJ01). See also (1985AN28) and (1983ANZQ, 1986HE26, 1987BL18, 1987SA15; theor.).




Fig. 4: Isobar diagram, $A = 8$. The diagrams for individual isobars have been shifted vertically to eliminate the neutron-proton mass difference and the Coulomb energy, taken as $E_C = 0.60Z(Z-1)/A^{1/3}$. Energies in square brackets represent the (approximate) nuclear energy, $E_N = M(Z, A) - ZM(\text{H}) - NM(\text{n}) - E_C$, minus the corresponding quantity for ${}^8\text{Be}$: here M represents the atomic mass excess in MeV. Levels which are presumed to be isospin multiplets are connected by dashed lines.

References

(Closed 1 June 1988)

- 1955AJ61 F. Ajzenberg and T. Lauritsen, *Revs. Mod. Phys.* 27 (1955) 77
- 1959AJ76 F. Ajzenberg and T. Lauritsen, *Nucl. Phys.* 11 (1959) 1
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, *Nucl. Phys.* 78 (1966) 1
- 1971WI05 D.H. Wilkinson and D.E. Alburger, *Phys. Rev. Lett.* 26 (1971) 1127
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, *Nucl. Phys.* A227 (1974) 1
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, *Table of Isotopes 7th Ed.* (1978)
- 1979AJ01 F. Ajzenberg-Selove, *Nucl. Phys.* A320 (1979) 1
- 1979GL12 H. Glattli, G.L. Bacchella, M. Fourmond, A. Malinovski, P. Meriel, M. Pinot, P. Roubeau and A. Abragam, *J. Phys. (Paris)* 40 (1979) 629
- 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, *Nucl. Phys.* A336 (1980) 1
- 1980MC07 R.D. McKeown, G.T. Garvey and C.A. Gagliardi, *Phys. Rev. C* 22 (1980) 738; Erratum *Phys. Rev. C* 26 (1982) 2336
- 1981BA36 F.C. Barker and N. Ferdous, *J. Phys. (London)* G7 (1981) 1239
- 1981BJ03 T. Bjornstad, H.A. Gustafsson, B. Jonson, P.O. Larsson, V. Lindfors, S. Mattsson, G. Nyman, A.M. Poskanzer, H.L. Ravn and D. Schardt, *Nucl. Phys.* A366 (1981) 461
- 1981MA33 S. Manglos, N.R. Roberson, H.R. Weller and D.R. Tilley, *Phys. Rev. C* 24 (1981) 2378
- 1981MC09 R.D. McKeown, S.J. Sanders, J.P. Schiffer, H.E. Jackson, M. Paul, J.R. Specht, E.J. Stephenson, R.P. Redwine and R.E. Segel, *Phys. Rev. C* 24 (1981) 211
- 1981MUZQ S.F. Mughabghab, M. Divadeenam and N.E. Holden, *Neutron Cross Sections Part A, Z=1-60* (1981)
- 1981PL1A G.R. Plattner, *Nukleonika* 26 (1981) 1005
- 1982AB1B V.G. Ableev, V.A. Bodyagin, G.G. Vorob'ev, R. Dymazh, S.A. Zaporozhets, V.I. Inozemtsev, A.A. Nomofilov, N.M. Piskunov, I.M. Sitnik, E.A. Stokovskii et al., *Yad. Fiz.* 36 (1982) 1434; *Sov. J. Nucl. Phys.* 36 (1982) 834
- 1982AL08 D.V. Aleksandrov, Yu.A. Glukhov, A.S. Demyanova, V.I. Dukhanov, I.B. Mazurov, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta and D.N. Stepanov, *Yad. Fiz.* 35 (1982) 277; *Sov. J. Nucl. Phys.* 35 (1982) 158
- 1982AL33 D.V. Aleksandrov, Yu.A. Glukhov, A.S. Demyanova, V.I. Dukhanov, I.B. Mazurov, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta and D.N. Stepanov, *Yad. Fiz.* 36 (1982) 1351; *Sov. J. Nucl. Phys.* 36 (1982) 783

- 1982BA1J Bahcall and Davis, *Essays in Nucl. Astrophys.* (1982) 243
- 1982GL01 B.G. Glagola, V.E. Viola, Jr., H. Breuer, N.S. Chant, A. Nadasen, P.G. Roos, S.M. Austin and G.J. Mathews, *Phys. Rev. C*25 (1982) 34
- 1982GU21 I.S. Gurbanovich and N.S. Zelenskaya, *Yad. Fiz.* 36 (1982) 1180; *Sov. J. Nucl. Phys.* 36 (1982) 688
- 1982KA1D K. Kar and J.C. Parikh, *Pramana* 19 (1982) 555
- 1982KA1E Kavanagh, *Essays in Nucl. Astrophys.* (1982) 159
- 1982LA19 M. Lattuada, F. Riggi, C. Spitaleri, D. Vinciguerra and C.M. Sutura, *Phys. Rev. C*26 (1982) 1330
- 1982MO1B T. Motoba, *Proc. Workshop on Hypernucl. Phys.*, Japan (1982) 36
- 1982NA1B D.S. Narayan, *Proc. VI High Energy Phys. Symp.*, 6-11 Dec. 1982, Mysore, India (1982) 57
- 1982PO12 V.S. Popov, A.E. Kudryavtsev, V.I. Lisin and V.D. Mur, *Pisma Zh. Eksp. Teor. Fiz.* 36 (1982) 207; *JETP Lett. (USSR)* 36 (1982) 257
- 1982WE15 K. Wen, P. Zhang and Z. Sun, *Chin. J. Nucl. Phys.* 4 (1982) 289
- 1982ZH06 M.A. Zhusupov and Yu.N. Uzikov, *Yad. Fiz.* 36 (1982) 1396; *Sov. J. Nucl. Phys.* 36 (1982) 810
- 1983AD1B E.G. Adelberger, *Proc. NATO Advanced Study Institute on Symmetries in Nucl. Struct.*, 16-28 Aug. 1982, Dronen, Netherlands; Eds., Abrahams, Allaart and Dieperink (1983) 55
- 1983AL1C G. Alberi, A. Malecki and V. Roberto, *Lett. Nuovo Cim.* 36 (1983) 409
- 1983AN13 L. Anderson, W. Bruckner, E. Moeller, S. Nagamiya, S. Nissen-Meyer, L. Schroeder, G. Shapiro and H. Steiner, *Phys. Rev. C*28 (1983) 1224; *Erratum Phys. Rev. C*28 (1983) 1246
- 1983AN1C Antonov and Petkov, in *Florence* (1983) 334
- 1983ANZQ Y. Ando, M. Uno and M. Yamada, *JAERI-M-83-025* (1983)
- 1983BA45 F.C. Barker, *Phys. Rev. C*28 (1983) 1407
- 1983BE51 T.L. Belyaeva, N.S. Zelenskaya and I.B. Teplov, *Yad. Fiz.* 38 (1983) 901; *Sov. J. Nucl. Phys.* 38 (1983) 540
- 1983BI1C P.G. Bizzeti, *Riv. Nuovo Cim.* 6 (1983) 1
- 1983BL17 S. Blagus, D. Miljanic and D. Rendic, *Nucl. Instrum. Meth.* 216 (1983) 415
- 1983BO19 F. Borchers, H. De Jong, J. Krug and E. Kuhlmann, *Nucl. Phys.* A405 (1983) 141
- 1983BU15 V.V. Burov, O.M. Knyazkov, A.A. Shirokova and K.V. Shitikova, *Z. Phys.* A313 (1983) 319
- 1983CA12 E. Caurier and B. Grammaticos, *J. Phys. (London)* G9 (1983) L125

- 1983CH1C W.H. Chung, J.C. Kim and B.N. Sung, *J. Korean Phys. Soc.* 16 (1983) 105
- 1983CO1A Compani-Tabrizi and Malik, in *Florence* (1983) 666
- 1983CO1D Cowan, *Bull. Amer. Phys. Soc.* 28 (1983) 994
- 1983DA11 V. D'Amico, G. Fazio, G. Giardina and F. Mezzanares, *Can. J. Phys.* 61 (1983) 923
- 1983DA22 J.H. Dave and C.R. Gould, *Phys. Rev. C* 28 (1983) 2212
- 1983DA32 N.X. Dai, Y. Chen, H.J. Lu, B.J. Qi, F. Zhuang and H.C. Sun, *Chin. J. Nucl. Phys.* 5 (1983) 19
- 1983DEZW L.C. Dennis, A.D. Frawley and J.F. Mateja, *Bull. Amer. Phys. Soc.* 28 (1983) 669, BF1
- 1983DO1B C.B. Dover and A. Gal, *Ann. Phys.* 146 (1983) 309
- 1983DR09 S. Drozd, J. Okolowicz and M. Ploszajczak, *Phys. Lett.* B128 (1983) 5
- 1983EL1A A. El-Naghy and M.T. Hussein, 18th Int. Cosmic Ray Conf., 22 Aug.-3 Sept. 1983, Bomay, India, Vol. 5 (1983) 207
- 1983FA17 M.A. Fawzy, A.A. Azzam and A.M. Abd El-Monim, *Indian J. Phys.* A57 (1983) 378
- 1983FA1A M.A. Faessler, *Nucl. Phys.* A400 (1983) 525
- 1983FE07 V.N. Fetisov, L. Majling, J. Zofka and R.A. Eramzhyan, *Z. Phys.* A314 (1983) 239
- 1983FI11 G.F. Filippov, V.S. Vasilevsky and A.V. Nesterov, *Yad. Fiz.* 38 (1983) 584; *Sov. J. Nucl. Phys.* 38 (1983) 347
- 1983FI13 B.W. Filippone, A.J. Elwyn, C.N. Davids and D.D. Koetke, *Phys. Rev. C* 28 (1983) 2222
- 1983FO1A Fowler, *AIP Conf. Proc.* 96 (1983) 80
- 1983FR1A W.A. Friedman and W.G. Lynch, *Phys. Rev. C* 28 (1983) 950
- 1983FU1D Y. Fujiwara, Y.C. Tang and H. Horiuchi, *Prog. Theor. Phys.* 70 (1983) 809
- 1983GA14 L.I. Galanina and N.S. Zelenskaya, *Izv. Akad. Nauk. SSSR Ser. Fiz.* 47 (1983) 1018; *Bull. Acad. Sci. USSR Phys. Ser.* 47 (1983) 191
- 1983GL1B P.W.M. Glaudemans, *Proc. NATO Advanced Study Institute on Symmetries in Nucl. Struct.*, 16-28 Aug. 1982, Dronen, Netherlands; Eds., Abrahams, Allaart and Dieperink (1983) 119
- 1983GLZZ C.W. Glover, C.C. Foster, P. Schwandt, J.R. Comfort, J. Rapaport, T.N. Taddeucci, D. Wong, J. Seubert and G. Wagner, *Bull. Amer. Phys. Soc.* 28 (1983) 996, EC3
- 1983GO1B Goncharova, Kissener and Eramzhyan, in *Orsay* (1983) 61
- 1983GO1H C.R. Gould, J. Dave and R.L. Walter, *Proc. Int. Conf. on Nucl. Data for Sci. and Tech.*, Antwerp, Belgium, 6-10 Sept. 1982 (1983) 766
- 1983GO25 K. Goeke, F. Grummer and P.-G. Reinhard, *Ann. Phys. (New York)* 150 (1983) 504

- 1983GR26 D.H.E. Gross and M.C. Nemes, Phys. Lett. B130 (1983) 131
- 1983GU1A C. Guet, Nucl. Phys. A400 (1983) 191
- 1983HA17 R.C. Haight, G.J. Mathews, R.M. White, L.A. Aviles and S.E. Woodard, Nucl. Instrum. Meth. 212 (1983) 245
- 1983HA1B M.J. Harris, W.A. Fowler, G.R. Caughlin and B.A. Zimmerman, Ann. Rev. Astron. Astrophys. 21 (1983) 165
- 1983HA1D W.S. Han and H.J. Lu, in Florence (1983) 314
- 1983HA41 W.S. Han and H.J. Lu, Chin. J. Nucl. Phys. 5 (1983) 105
- 1983HA45 S. Hayashi and S. Iwata, Ann. Rept. Res. Reactor Inst., Kyoto Univ., 16 (1983) 131
- 1983JA09 L. Jarczyk, B. Kamys, Z. Rudy, A. Strzalkowski, H. Witala, M. Hugi, J. Lang, R. Muller, J. Sromicki and H.H. Wolter. Phys. Rev. C28 (1983) 700
- 1983KO17 L. Koester, K. Knopf and W. Waschkowski, Z. Phys. A312 (1983) 81
- 1983KO41 A. Kolawa, Acta Phys. Pol. B14 (1983) 743
- 1983KU17 N. Kumar, Nucl. Phys. A410 (1983) 50
- 1983LI18 X.H. Liu and X.J. Chen, Chin. J. Nucl. Phys. 5 (1983) 249
- 1983LI1C H. Liskien, R. Wolfle and S.M. Qaim, Proc. Int. Conf. on Nucl. Data for Sci. and Tech., Antwerp, Belgium, 6-10 Sept. 1982 (1983) 349
- 1983LO12 M.A. Lone, R.T. Jones, A. Okazaki, B.M. Townes, D.C. Santry, E.D. Earle, J.K.P. Lee, J.M. Robson, R.B. Moore, L. Nikkinen et al., Nucl. Instrum. Meth. 214 (1983) 333
- 1983MA73 L. Marquez, Phys. Rev. C28 (1983) 2525
- 1983MI10 D. Miljanic, E. Kossionides, G. Vourvopoulos and P. Assimakopoulos, Z. Phys. A312 (1983) 267
- 1983MI1E K. Miyahara, K. Ikeda and H. Bando, Prog. Theor. Phys. 69 (1983) 1717
- 1983MO1C T. Motoba, H. Bando and K. Ikeda, Prog. Theor. Phys. 70 (1983) 189
- 1983MU13 A.M. Mukhamedzhanov, R. Yarmukhamedov and P.O. Dzhamalov, Yad. Fiz. 37 (1983) 1405; Sov. J. Nucl. Phys. 37 (1983) 838
- 1983OK06 S. Okabe, Nucl. Phys. A404 (1983) 179
- 1983OL1A D.L. Olson, B.L. Berman, D.E. Greiner, H.H. Heckman, P.J. Lindstrom and H.J. Crawford, Phys. Rev. C28 (1983) 1602
- 1983PR09 J. Proriol and B. Jargeaix, Nuovo Cim. A77 (1983) 289
- 1983SA16 M. Saraceno, P. Kramer and F. Fernandez, Nucl. Phys. A405 (1983) 88
- 1983SH1E Y.-J. Shi and F. Zhuang, Phys. Energ. Fortis Phys. Nucl. 7 (1983) 605
- 1983SH38 Y.-J. Shi, Phys. Rev. C28 (1983) 2452

- 1983SH39 A.A. Shvedov, V.N. Dobrikov and O.F. Nemets, *Pisma Zh. Eksp. Teor. Fiz.* 38 (1983) 32; *JETP Lett. (USSR)* 38 (1983) 36
- 1983SI1A R.H. Siemssen, *Nucl. Phys. A*400 (1983) 245
- 1983SP06 Ts.P. Spasov, Ch.M. Chernev, Yu.A. Batusov and R.A. Eramzhyan, *Blug. J. Phys.* 10 (1983) 581
- 1983SZZY J. Szabo, M. Varnagy, Z.T. Body and J. Csikai, *Proc. Int. Conf. Nucl. Data for Sci. and Tech., Geel, Belgium, 1982; Ed., K.H. Bockhoff (1983)* 956
- 1983VO1C Vogel, in *Symmetries in Nucl. Struct., Eds., Abrahams, Allaart and Dieperink (1983)* 203
- 1983WA09 R.E. Warner, K. Blum, D. Friesel, P. Schwandt, P.P. Singh and A. Galonsky, *Nucl. Phys. A*401 (1983) 521
- 1983WA1F Wang et al., in *Florence (1983)* 591
- 1983WI1A Wilczynski, *Proc. Int. Conf. on Nucl. Phys., Florence, Aug.-Sept. 1983, Eds., P. Blasi and R.A. Ricci, Vol. 2 (1983)* 305
- 1983XU1A S.-W. Xu, G.-H. Wu, R.-Z. Miao and F. Han, *Chin. Phys.* 3 (1983) 646
- 1983YA05 M. Yasue, H. Yokomizo, T. Hasegawa, F. Soga, M.H. Tanaka, K. Koyama and K. Iwatani, *Nucl. Phys. A*408 (1983) 205
- 1984AB1C U.A. Abdurazakov, R.A. Bondarenko, U.G. Gulyamov and G.M. Chernov, *Yad. Fiz.* 39 (1984) 272; *Sov. J. Nucl. Phys.* 39 (1984) 169
- 1984AJ01 F. Ajzenberg-Selove, *Nucl. Phys. A*413 (1984) 1
- 1984AL03 J.P. Alard, J. Arnold, M.M. Avome Nze, J.P. Costilhes, L. Fraysse, W. Raja and J.C. Tamain, *Lett. Nuovo Cim.* 39 (1984) 139
- 1984AN16 Yu.P. Antufyev, A.S. Deineko, I.I. Zalyubovsky, V.D. Sarana, V.E. Storizhko, A.I. Tutubalin, C. Hategan and N.A. Schliakhov, *Yad. Fiz.* 40 (1984) 53; *Sov. J. Nucl. Phys.* 40 (1984) 35
- 1984AS1D F. Asai, H. Bando and M. Sano, *Phys. Lett. B*145 (1984) 19
- 1984BA1T Bayukov et al., in *Panic (1984)* I16
- 1984BA1U Bayukov et al., in *Panic (1984)* I25
- 1984BA25 L.A. Bandurina, I.Yu. Krivsky, A.I. Lendel and S.Yu. Medvedev, *Yad. Fiz.* 39 (1984) 296; *Sov. J. Nucl. Phys.* 39 (1984) 187
- 1984BL21 L.D. Blokhintsev, A.M. Mukjamedzhanov and A.N. Safronov, *Fiz. Elem. Chastits At. Yadra* 15 (1984) 1296; *Sov. J. Part. Nucl.* 15 (1984) 580
- 1984BO1H L.N. Bogdanova and V.E. Markushin, *Fiz. Elem. Chastits At. Yadra* 15 (1984) 808; *Sov. J. Part. Nucl.* 15 (1984) 361
- 1984CH1G H.Z. Chen, F. Zhuang, X.J. Shi and X.N. Jin, *Chin. J. Nucl. Phys.* 6 (1984) 303

- 1984DA17 B. Dasmahapatra, B. Cujec and F. Lahlou, Nucl. Phys. A427 (1984) 186
- 1984DA1B Danilin and Zhukov, in Alma Ata (1984) 460
- 1984DA1H R. Davis, Jr., B.T. Cleveland and J.K. Rowley, AIP Conf. Proc. 123 (1984) 1037
- 1984DA22 J. D'Auria, M. Dombsky, L. Moritz, T. Ruth, G. Sheffer, T.E. Ward, C.C. Foster, J.W. Watson, B.D. Anderson and J. Rapaport, Phys. Rev. C30 (1984) 1999
- 1984DE24 E. Deumens, Nucl. Phys. A423 (1984) 52
- 1984DU17 S.B. Dubovichenko and M.A. Zhusupov, Izv. Akad. Nauk. SSSR Ser. Fiz. 48 (1984) 935; Bull. Acad. Sci. USSR Phys. Ser. 48 (1984) 95
- 1984FA1B M.A. Faessler, Phys. Rept. 115 (1984) 1
- 1984FE1A Ferch et al., INDC (CCP)-221/L (1984) 18
- 1984FI11 H. Fiedeldey, R. Lipperheide, K. Naidoo and S.A. Sofianos, Phys. Rev. C30 (1984) 434
- 1984FI13 G.F. Filippov, V.S. Vasilevsky and A.V. Nesterov, Nucl. Phys. A426 (1984) 327
- 1984FR10 H. Friedrich, Phys. Rev. C30 (1984) 1102
- 1984FR13 H. Friedrich, Phys. Lett. B146 (1984) 135
- 1984FR1C W. Frati, Nucl. Phys. A418 (1984) 177
- 1984GA07 R.B. Galloway and A.M. Ghazarian, Phys. Rev. C29 (1984) 2349
- 1984GR08 R.E.L. Green, R.G. Korteling and K.P. Jackson, Phys. Rev. C29 (1984) 1806
- 1984HA1B Hampel, Nature 308 (1984) 312
- 1984HA1F Haight, Bull. Amer. Phys. Soc. 29 (1984) 1086
- 1984HA1M W.C. Haxton, AIP Conf. Proc. 123 (1984) 1026
- 1984HE1D E.F. Hefter and K.A. Gridnev, Prog. Theor. Phys. 72 (1984) 549
- 1984HI1A A.S. Hirsch, A. Bujak, J.E. Finn, L.J. Gutay, R.W. Minich, N.T. Porile, R.P. Scharenberg and B.C. Stringfellow, Phys. Rev. C29 (1984) 508
- 1984HU1E I. Hunyadi, I.M. Szoghy and B. Cujec, Nucl. Tracks Radiat. Meas. 8 (1984) 525
- 1984JE1B S.C. Jeong, J.C. Kim, C.W. Nahm, B.N. Sung and S.W. Cho, New Phys. (Korea) 24 (1984) 404
- 1984KO25 I. Koenig, D. Fick, S. Kossionides, P. Egelhof, K.-H. Mobius and E. Steffens, Z. Phys. A318 (1984) 135
- 1984KR10 V.M. Krasnopolskii, V.I. Kukul'in and V.G. Neudachin, Izv. Akad. Nauk. SSSR Ser. Fiz. 48 (1984) 84; Bull. Acad. Sci. USSR Phys. Ser. 48 (1984) 82
- 1984KR1B A.V. Kravtsov, N.P. Popov and G.E. Solvakin, JETP Lett. 40 (1984) 875
- 1984KU15 V.I. Kukul'in, M. Kamal, V.T. Voronchev and V.M. Krasnopolsky, J. Phys. (London) G10 (1984) L213

- 1984LA27 M. Langevin, C. Detraz, M. Epherre, D. Guillemaud-Mueller, B. Jonson, C. Thibault and the ISOLDE Collaboration, Phys. Lett. B146 (1984) 176
- 1984LI28 Y.-G. Li and S.-Y. Lo, Aust. J. Phys. 37 (1984) 255
- 1984LIZY P.W. Lisowski, R.E. Brown, J.C. Gursky, S.D. Howe, N. Jarmie and G.L. Morgan, Bull. Amer. Phys. Soc. 29 (1984) 748, JH13
- 1984LU1A Z.Q. Lu, W.S. Han, X.Z. Zhang and Z.X. Li, Chin. J. Nucl. Phys. 6 (1984) 34
- 1984LU1B Z.Q. Lu, W.S. Han, X.Z. Zhang and Z.X. Li, Chin. Phys. 4 (1984) 654
- 1984MA16 A. Malecki, P. Picozza and L. Satta, Phys. Lett. B136 (1984) 319
- 1984MA1H L. Marquez, Phys. Rev. C30 (1984) 1104
- 1984MA68 D. Majumdar, A.R. Chowdhury and T. Roy, Ann. Phys. (Leipzig) 41 (1984) 218
- 1984MI1C Millener, Gal, Dover and Dalitz, in Panic (1984) M7
- 1984MI1E D.J. Millener, AIP Conf. Proc. 123 (1984) 850
- 1984MO1H T. Motoba, H. Bando and K. Ikeda, Proc. Int. Summer School, Changchun, China, 1983 (1984) 702
- 1984MO1J T. Mori, H. Nishihara, I. Kimura, S.A. Hayashi, K. Kobayashi, S. Yamamoto and M. Nakagawa, Ann. Rept. Res. Reactor Inst., Kyoto Univ. 17 (1984) 22
- 1984NA11 K. Naidoo, H. Fiedeldey, S.A. Sofianos and R. Lipperheide, Nucl. Phys. A419 (1984) 13
- 1984NE1A Nemets, Rudchik and Chuvilski, in Alma Ata (1984) 334
- 1984OK03 I.P. Okhrimenko, Yad. Fiz. 40 (1984) 412; Sov. J. Nucl. Phys. 40 (1984) 262
- 1984PA04 P. Park, J. Carvalho, M. Vassanji, D.J. Rowe and G. Rosensteel, Nucl. Phys. A414 (1984) 93
- 1984PA13 D.J. Parker, J. Asher, T.W. Conlon and I. Naqib, Phys. Rev. C30 (1984) 143
- 1984PA1E G. Paic, B. Antolkovic and K. Kadija, Fizika 16 (1984) 37
- 1984RE14 S.M. Read and V.E. Viola, Jr., At. Data Nucl. Data Tables 31 (1984) 359
- 1984REZZ D.S. Resler, H.D. Knox, R.O. Lane and S.M. Grimes, Bull. Amer. Phys. Soc. 29 (1984) 629, AG3
- 1984SA39 L. Satta, J. Duflo, F. Plouin, P. Picozza, L. Goldzahl, J. Banaigs, R. Frascaria, F.L. Fabbri, A. Codino, J. Berge et al., Phys. Lett. B139 (1984) 263
- 1984SE16 R.G. Seyler and H.R. Weller, Phys. Rev. C30 (1984) 1146
- 1984SH1B K. Shibata, JAERI-M-84-204 (1984)
- 1984SH1D Shvedov, Dobrikov and Nemets, in Alma Ata (1984) 331
- 1984SH1J M. Shoeb and M.Z. Khan, J. Phys. (London) G10 (1984) 1047
- 1984SH1N K. Shibata, JAERI-M-84-163 (1984)

- 1984SHZJ A.A. Shvedov, V.N. Dobrikov and O.F. Nemets, in Alma-Ata (1984) 332
- 1984SMZX D.L. Smith, J.W. Meadows, M.M. Bretscher and S.A. Cox, Bull. Amer. Phys. Soc. 29 (1984) 1036, BD4
- 1984SP1C D. Sprengel, H.V. Buttler, J. Drevermann, W. Hoppe, R. Isenbugel, E.U. Klaus and N. Marquardt, Fizika 16 (1984) 25
- 1984TA07 T.N. Taddeucci, T.A. Carey, C. Gaarde, J. Larsen, C.D. Goodman, D.J. Horen, T. Masterson, J. Rapaport, T.P. Welch and E. Sugarbaker, Phys. Rev. Lett. 52 (1984) 1960
- 1984TA1D M. Tanaka, A.L.S. Anyelis, G. Basinizt, H.-J. Besch, R.E. Breedon, L. Camilleri, T.J. Chapin, C. Chasman, R.L. Cool, P.T. Cox et al., BCMOR Collaboration, Nucl. Phys. A418 (1984) 321
- 1984TA1E F. Takagi, Prog. Theor. Phys. 71 (1984) 585
- 1984TA1G M. Tanaka, AIP Conf. Proc. 123 (1984) 762
- 1984TU02 M. Turk and B. Antolkovic, Nucl. Phys. A431 (1984) 381
- 1984UM04 K. Umeda, T. Yamaya, T. Suehiro, K. Takimoto, R. Wada, E. Takada, S. Shimoura, A. Sakaguchi, S. Murakami, M. Fukada et al., Nucl. Phys. A429 (1984) 88
- 1984VA06 A.G.M. van Hees and P.W.M. Glaudemans, Z. Phys. A315 (1984) 223
- 1984VA1B Vasilevsky, Chopovski and Fillipov, in Alma Ata (1984) 197
- 1984VAZS V.S. Vasilevsky, S.P. Kruchinin, G.F. Filippov and L.L. Chopovsky, in Alma Ata (1984) 463
- 1984VO1A V.T. Voronchev, V.I. Kukulkin and V.M. Krasnopol'skii, Nucl. Fusion 24 (1984) 1117
- 1984YA1A J. Yang, M.S. Turner, G. Steigman, D.N. Schramm and K.A. Olive, Astrophys. J. 281 (1984) 493
- 1984ZA07 N.I. Zaika, A.V. Mokhnach, A.T. Rudchik, V.S. Semenov, V.A. Shityuk and P.L. Shmarin, Yad. Fiz. 39 (1984) 1081; Sov. J. Nucl. Phys. 39 (1984) 682
- 1984ZA1B A.M. Zadorozhnyi, V.V. Uzhinskii and S.Yu. Shmakov, Yad. Fiz. 39 (1984) 1155; Sov. J. Nucl. Phys. 39 (1984) 729
- 1984ZE1A Zelenskaya and Morzabaev, in Alma Ata (1984) 466
- 1984ZH1B F. Zhuang, H.-Z. Chen and X.-N. Jin, Phys. Energ. Fortis Phys. Nucl. 8 (1984) 215
- 1984ZW1A Zwartz, Unpublished Ph.D. Thesis, Utrecht Univ. (1984)
- 1985AB1A V.G. Ableev, V.A. Bodyagin, R. Dymarz, A. Filipkowski, V.I. Innozemtsev, A.A. Nomofilov, N.M. Piskunov, V.I. Sharov, I.M. Sitnik, E.A. Strokovsky et al., Acta Phys. Pol. B16 (1985) 913
- 1985AH1A I. Ahmad, M. Mian and M.Z. Rahman Khan, Phys. Rev. C31 (1985) 1590
- 1985AJ01 F. Ajzenberg-Selove, Nucl. Phys. A433 (1985) 1; Erratum Nucl. Phys. A449 (1986) 155

- 1985AK1A T. Akesson, M.G. Albrow, S. Almeded, R. Batley, O. Benary, H. Boggild, O. Botner, H. Breuker, H. Brody, A.A. Carter et al., Phys. Lett. B152 (1985) 140
- 1985AL1G Aleksandrov et al., Questions in At. Phys. and in Tech., USSR (1985) 3
- 1985AL29 D.V. Aleksandrov, Yu.A. Glukhov, B.G. Novatsky, E.Yu. Nikolsky, A.A. Ogloblin and D.N. Stepanov, Izv. Akad. Nauk SSSR, Ser. Fiz. 49 (1985) 2115; Bull. Acad. Sci. USSR, Phys. Ser. 49 (1985) 30
- 1985ALZL D.V. Aleksandrov, E.A. Ganza, Yu.A. Glukhov, E.Yu. Nikolsky, B.G. Novatsky, A.A. Ogloblin, D.N. Stepanov, in Leningrad (1985) 355
- 1985ALZM D.V. Aleksandrov, Yu.A. Glukhov, E.Yu. Nikolsky, B.G. Novatsky, A.A. Ogloblin, D.N. Stepanov, in Leningrad (1985) 354
- 1985AN28 M.S. Antony, J. Britz, J.B. Bueb and A. Pape, At. Data Nucl. Data Tables 33 (1985) 447
- 1985BA1M J.N. Bahcall, Solar Phys. 100 (1985) 53
- 1985BA1N J.N. Bahcall, AIP Conf. Proc. 126 (1985) 60
- 1985BA1Q D. Baye and P. Descouvemont, Ann. Phys. 165 (1985) 115
- 1985BA45 D. Baye and P. Descouvemont, Nucl. Phys. A443 (1985) 302
- 1985BE1C Berdnikov et al., in Leningrad (1985) 302
- 1985BE1E W. Bell, K. Braune, G. Claesson, D. Drijard, M. A. Faessler, H. G. Fischer, H. Frehse, R. W. Frey, S. Garpman, W. Geist et al; CERN-Heidelberg-Lund Collaboration, Nucl. Phys. B254 (1985) 475
- 1985BE30 S.L. Belostotsky, S.S. Volkov, A.A. Vorobyev, Yu.V. Dotsenko, L.G. Kudin, N.P. Kuropatkin, O.V. Miklukho, V.N. Nikulin and O.E. Prokofyev, Yad. Fiz. 41 (1985) 1425; Sov. J. Nucl. Phys. 41 (1985) 903
- 1985BIZZ R.A. Bigelow, P.A. Quin, S.J. Freedman and J. Napolitano, Bull. Amer. Phys. Soc. 30 (1985) 701, AE8
- 1985BO1C C.D. Bowman, S.A. Wender and G.F. Auchampaugh, AIP Conf. Proc. 124 (1985) 259
- 1985BO1D Body and Mihaly, INDC (HUN)-22/L (1985)
- 1985BO1E R.N. Boyd, R.E. Turner, B. Sur, L. Rybaryk and C. Joseph, AIP Conf. Proc. 126 (1985) 145
- 1985BO1K Boesgaard and Steigman, Ann. Rev. Astron. Astrophys. 23 (1985) 319
- 1985BU16 A. Budzanowski, H. Dabrowski, Y. Chan, R.G. Stokstad, I. Tserruya and S. Wald, Phys. Rev. C32 (1985) 1534
- 1985CA1C V. Cavasinni, T. Del Prete, M. Morganti, F. Schiavo, D.L. Owen, G. Paternoster, S. Patricelli and G. Anzivino, Z. Phys. C28 (1985) 487
- 1985CA36 R.F. Carlson, A.J. Cox, T.N. Nasr, M.S. De Jong, D.L. Ginther, D.K. Hasell, A.M. Sourkes, W.T.H. Van Oers and D.J. Margaziotis, Nucl. Phys. A445 (1985) 57

- 1985CA41 G.R. Caughlan, W. A. Fowler, M.J. Harris and B.A. Zimmerman, *At. Data Nucl. Data Tables* 32 (1985) 197
- 1985CE12 F.E. Cecil, R.J. Peterson and P.D. Kunz, *Nucl. Phys. A*441 (1985) 477
- 1985CH1B H.H. Chen, *AIP Conf. Proc.* 126 (1985) 249
- 1985CH37 S. Chiba, M. Baba, H. Nakashima, M. Ono, N. Yabuta, S. Yukinori and N. Hirakawa, *J. Nucl. Sci. Tech. (Tokyo)* 22 (1985) 771
- 1985CO09 J. Cook and K.W. Kemper, *Phys. Rev. C*31 (1985) 1745
- 1985DA29 V. D'Amico, G. Fazio, S. Femino, G. Giardina, A. Italiano and F. Mezzanares, *Can. J. Phys.* 63 (1985) 1438
- 1985DE1K P. Delbourgo-Salvador, C. Gry, G. Malinie and J. Audouze, *Astron. Astrophys.* 150 (1985) 53
- 1985DO16 Yu.V. Dotsenko and V.E. Starodubsky, *Yad. Fiz.* 42 (1985) 107; *Sov. J. Nucl. Phys.* 42 (1985) 66
- 1985FA1A M.A. Faessler, *Nucl. Phys. A*434 (1985) 563
- 1985FI1D B.W. Filippone, *AIP Conf. Proc.* 126 (1985) 100
- 1985FI1E G.F. Filippov, V.S. Vasilevskii and L.L. Chopovskii, *Fiz. Elem. Chastits At. Yadra* 16 (1985) 349; *Sov. J. Part. Nucl.* 16 (1985) 153
- 1985FR1E S. Frankel, *Comments Nucl. Part. Phys.* 14 (1985) 1
- 1985FR1F V. Franco and Y. Yin, *Phys. Rev. Lett.* 55 (1985) 1059
- 1985GA1B L.I. Galanina and N.S. Zelenskaya, *Nucl. Phys. A*445 (1985) 625
- 1985GI1C R.L. Gilliland, *Astrophys. J.* 290 (1985) 344
- 1985GO18 E. Goldberg, R.L. Barber, P.E. Barry, N.A. Bonner, J.E. Fontanilla, C.M. Griffith, R.C. Haight, D.R. Nethaway and G.B. Hudson, *Nucl. Sci. Eng.* 91 (1985) 173
- 1985GO1A N.G. Goncharova, H.-R. Kissener and R.A. Eramzhyan, *Fiz. Elem. Chastits At. Yadra* 16 (1985) 773; *Sov. J. Part. Nucl.* 16 (1985) 337
- 1985GO1B N.G. Goncharova, Kh.R. Kissener and R.A. Eramzhyan, *Izv. Akad. Nauk. SSSR Ser. Fiz.* 49 (1985) 1032; *Bull. Acad. Sci. USSR Phys. Ser.* 49 (1985) 184
- 1985GO1G Gorionov et al., in *Leningrad* (1985) 310
- 1985GR1A L. Grenacs, *Ann. Rev. Nucl. Part. Sci.* 35 (1985) 455
- 1985HA1N B.G. Harvey, *Nucl. Phys. A*444 (1985) 498
- 1985HA40 R.C. Haight, G.J. Mathews and R.W. Bauer, *Nucl. Instrum. Meth. Phys. Res. B*10-11 (1985) 361
- 1985HO1B H. Horiuchi, *Prog. Theor. Phys.* 74 (1985) 66
- 1985IK1A K. Ikeda, H. Bando and T. Motoba, *Prog. Theor. Phys. Suppl.* 81 (1985) 147

- 1985JA09 L. Jarczyk, B. Kamys, A. Magiera, A. Strzalkowski, A. Szczurek, K. Bodek, M. Hugi, J. Lang, R. Muller and E. Ungricht, *J. Phys. (London)* G11 (1985) 843
- 1985JA18 B.V. Jacak, D. Fox and G.D. Westfall, *Phys. Rev.* C31 (1985) 704
- 1985KA1E Kadenskii and Chuvilskii, in *Leningrad* (1985) 437, 440
- 1985KA1G Kadenskii, Furman and Chuvilskii, in *Leningrad* (1985) 439
- 1985KAZQ S.G. Kadenskii, S.D. Kurgalin and Yu.M. Chuvilskii, in *Leningrad* (1985) 438
- 1985KI11 R. Kircher, H. Kamada and S. Oryu, *Prog. Theor. Phys. (Kyoto)* 73 (1985) 1442
- 1985KI1A Kitching, McDonald, Maris and Vasconcellos, *Adv. Nucl. Phys.* 15 (1985) 43
- 1985KL1A H.V. Klapdor, *Fortschr. Phys.* 33 (1985) 1
- 1985KR21 P. Kristiansson, J.P. Bondorf, L. Carlen, H.-A. Gustafsson, B. Jakobsson, A. Kristiansson, G. Lovhoiden, H. Nifenecker, O.-B. Nielsen, A. Oskarsson et al., *Nucl. Phys.* A447 (1986) 577c
- 1985KW03 E. Kwasniewicz, J. Kisiel and L. Jarczyk, *Acta Phys. Pol.* B16 (1985) 947
- 1985LA1C P.O. Lagage, *Nature* 316 (1985) 420
- 1985LI1C Y.-G. Ling, X. Zhao and F.-A. Zeng, *Phys. Energ. Fortis Phys. Nucl.* 9 (1985) 236
- 1985MA02 H. Machner, D. Protic, G. Riepe, H.G. Bohlen and H. Fuchs, *Phys. Rev.* C31 (1985) 443
- 1985MA13 M.T. Magda, A. Pop and A. Sandulescu, *J. Phys. (London)* G11 (1985) L75
- 1985MAZG B.S. Mazitov and E.N. Rasulov, in *Leningrad* (1985) 298
- 1985MC1C J.R. McNally Jr., *Fusion Tech.* 7 (1985) 331
- 1985MO17 D.J. Morrissey, W. Benenson, E. Kashy, C. Bloch, M. Lowe, R.A. Blue, R.M. Ronningen, B. Sherrill, H. Utsunomiya and I. Kelson, *Phys. Rev.* C32 (1985) 877
- 1985MO1F T. Motoba, H. Bando, K. Ikeda and T. Yamada, *Suppl. Prog. Theor. Phys.* 81 (1985) 42
- 1985MO24 D.J. Morrissey, W. Benenson, E. Kashy, C. Bloch, M. Lowe, B. Sherrill, R.A. Blue, R.M. Ronningen and H. Utsunomiya, *Nucl. Phys.* A447 (1986) 603c
- 1985MU14 T. Murakami, E. Ungricht, Y.-W. Lui, Y. Mihara, E. Takada and R.E. Tribble, *Phys. Rev.* C32 (1985) 1558
- 1985NO1A Norbeck and Lin, *Bull. Amer. Phys. Soc.* 30 (1985) 1248
- 1985NO1B K. Nomoto, F.-K. Thielemann and S. Miyaji, *Astron. Astrophys.* 149 (1985) 239
- 1985PO10 N.A.F.M. Poppelier, L.D. Wood and P.W.M. Glaudemans, *Phys. Lett.* B157 (1985) 120
- 1985PO11 D.N. Poenaru, M. Ivascu, A. Sandulescu and W. Greiner, *Phys. Rev.* C32 (1985) 572

- 1985PO19 J. Pochodzalla, W.A. Friedman, C.K. Gelbke, W.G. Lynch, M. Maier, D. Ardouin, H. Delagrange, H. Doubre, C. Gregoire, A. Kyanowski et al., Phys. Lett. B161 (1985) 275
- 1985PR1A J. Proriot, Lett. Nuovo Cim. 42 (1985) 28
- 1985PR1B J. Proriot, Nouvo Cim. A85 (1985) 310
- 1985PU03 V.M. Pugach, Yu.E. Kozyr, V.I. Medvedev, Yu.N. Pavlenko, V.V. Komarov, A.M. Popova and V.L. Shablov, Izv. Akad. Nauk SSSR Ser. Fiz. 49 (1985) 905; Bull. Acad. Sci. USSR Phys. Ser. 49 (1985) 65
- 1985RO1G D.J. Rowe, Rept. Prog. Phys. 48 (1985) 1419
- 1985SA32 H. Sato and Y. Okuhara, Phys. Lett. B162 (1985) 217
- 1985SH1A K.V. Shitikova, Fiz. Elem. Chastits At. Yadra 16 (1985) 824; Sov. J. Part. Nucl. 16 (1985) 364
- 1985SM1B Smith and Meadows, Bull. Amer. Phys. Soc. 30 (1985) 1252
- 1985SP05 G. Spitz, H. Klar and E.W. Schmid, Z. Phys. A322 (1985) 49
- 1985SW01 M.T. Swinhoe and C.A. Uttley, Nucl. Sci. Eng. 89 (1985) 261
- 1985TA13 I. Tanihata, H. Hamagaki, O. Hashimoto, S. Nagamiya, Y. Shida, N. Yoshikawa, O. Yamakawa, K. Sugimoto, T. Kobayashi, D.E. Greiner et al., Phys. Lett. B160 (1985) 380
- 1985TA18 I. Tanihata, H. Hamagaki, O. Hashimoto, Y. Shida, N. Yoshikawa, K. Sugimoto, O. Yamakawa, T. Kobayashi and N. Takahashi, Phys. Rev. Lett. 55 (1985) 2676
- 1985TA1D I. Tanihata, Hyperfine Interactions 21 (1985) 251
- 1985TH08 J. Thaler, Phys. Rev. C32 (1985) 2189
- 1985WA02 A.H. Wapstra and G. Audi, Nucl. Phys. A432 (1985) 1
- 1985WA1C Walter, Bull. Amer. Phys. Soc. 30 (1985) 113
- 1985WA22 S. Wald, S.B. Gazes, C.R. Albiston, Y. Chan, B.G. Harvey, M.J. Murphy, I. Tserruya, R.G. Stokstad, P.J. Countryman, K. Van Bibber et al., Phys. Rev. C32 (1985) 894
- 1985WI1B D. Wilkinson, Nucl. Phys. A434 (1985) 573
- 1985WO11 L.W. Woo, K. Kwiatkowski, S.H. Zhou and V.E. Viola, Phys. Rev. C32 (1985) 706
- 1985YA05 T. Yamada, K. Ikeda, H. Bando and T. Motoba, Prog. Theor. Phys. 73 (1985) 397
- 1985YI1B Y.-C. Yin, Z.-Q. Tan and K.-Z. Chen, Nucl. Phys. A440 (1985) 685
- 1985YI1C Y.-C. Yin, Z.-Q. Tan and K.-Z. Chen, Phys. Energ. Fortis Phys. Nucl. 9 (1985) 569
- 1986AB1E K. Abrahams, J. Phys. Soc. Jpn. Suppl. 55 (1986) 572
- 1986AJ01 F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1
- 1986AJ04 F. Ajzenberg-Selove, Nucl. Phys. A460 (1986) 1

- 1986ALZN S.C. Allcock, W.D.M. Rae, P.R. Keeling, S. Marsh, A.E. Smith, B.R. Fulton and D.W. Banes, in Harrogate (1986) 46, B20
- 1986AN10 M.S. Antony, J. Britz and J.B. Bueb, Nuovo Cim. A91 (1986) 283
- 1986AN1F B. Andersson, S. Garpman, G. Gustafson, H.-A. Gustafsson, B. Nilsson-Almqvist, I. Otterlund and E. Stenlund, Phys. Scripta 34 (1986) 451
- 1986AN1R H.H. Ansari, M. Shoeb and M.Z.R. Rahman Khan, J. Phys. (London) G12 (1986) 1369
- 1986AN22 B. Antolkovic, M. Turk and K. Kadija, Radiat. Eff. 92 (1986) 183; in Santa Fe 85 (1986) 183
- 1986AV1B Avdeichikov, in Dubna (1986) 122
- 1986BA09 S.A. Badalov and G.F. Filippov, Yad. Fiz. 43 (1986) 71; Sov. J. Nucl. Phys. 43 (1986) 45
- 1986BA1W H. Bando, Nucl. Phys. A450 (1986) 217c
- 1986BA21 J.N. Bahcall and B.R. Holstein, Phys. Rev. C33 (1986) 2121
- 1986BA2D R. Babinet, Ann. Phys. (France) 11 (1986) 113
- 1986BA38 F.C. Barker and R.H. Spear, Astrophys. J. 307 (1986) 847
- 1986BA40 F.M. Baumann, G. Domogala, H. Freiesleben, H.J. Paul, S. Puhlfers and H. Sohlbach, Nucl. Instrum. Meth. Phys. Res. A247 (1986) 359
- 1986BA88 W. Bauhoff, At. Data Nucl. Data Tables 35 (1986) 429
- 1986BE1K H.A. Bethe, Phys. Rev. Lett. 56 (1986) 1305
- 1986BE1Q Belozarov et al., in P7-86-322, Dubna (1986) 53
- 1986BE1S W. Bell, K. Braune, G. Claesson, D. Drijard, M.A. Faessler, H.G. Fischer, H. Frehse, R.W. Frey, S. Garpman, W. Geist et al., Z. Phys. C30 (1986) 513
- 1986BE1T W. Bell, A. Breakstone, G. Claesson, H.B. Crawley, G.M. Dallavalle, K. Doroba, D. Drijard, F. Fabbri, A. Firestone, M.A. Faessler et al., Z. Phys. A325 (1986) 7
- 1986BE2N W. Bell, A. Breakstone, G. Claesson, H.B. Crawley, G.M. Dallavalle, K. Doroba, D. Drijard, F. Fabbri, A. Firestone, M.A. Faessler et al., Z. Phys. C32 (1986) 335
- 1986BI1D R.A. Bigelow, S.J. Freedman, J. Napolitano and P.A. Quin, J. Phys. Soc. Jpn. Suppl. 55 (1986) 1030
- 1986BL12 C. Bloch, W. Benenson, E. Kashy, D.J. Morrissey, R.A. Blue, R.M. Ronningen and H. Utsunomiya, Phys. Rev. C34 (1986) 850
- 1986BL1D Blokhintsev, Razikov, Ubaidullaeva and Yarmukhamedov, in Kharkov (1986) 449
- 1986BO1H A.M. Boesgaard and R.J. Lavery, Astrophys. J. 309 (1986) 762

- 1986BO41 M.J.G. Borge, M. Epherre-Rey-Campagnolle, D. Guillemaud-Mueller, B. Jonson, M. Langevin, G. Nyman, C. Thibault and the ISOLDE Collaboration, Nucl. Phys. A460 (1986) 373
- 1986BOZG I.M. Bondarenko and E.E. Petrov, INDC(CCP)-265/L (1986)
- 1986BR26 M.E. Brandan, A. Menchaca-Rocha, A. Szanto de Toledo, N. Carlin Filho, E.M. Szanto and M.M. Coimbra, J. Phys. (London) G12 (1986) 391
- 1986CE04 C. Cernigoi, N. Grion, G. Pauli, R. Rui and R. Cherubini, Nucl. Phys. A456 (1986) 599
- 1986CH1J N.S. Chant, AIP Conf. Proc. 142 (1986) 246
- 1986CH1M Choi et al., Bull. Amer. Phys. Soc. 31 (1986) 877
- 1986CH1S E.T. Cheng and J.H. Huang, Radiat. Eff. 92 (1985) 231; in Santa Fe 85 (1986) 231
- 1986CH24 S. Chiba, M. Baba, H. Nakashima, M. Ono, N. Yabuta, S. Yukinori and N. Hirakawa, Radiat. Eff. 92 (1986) 227; in Santa Fe 85 (1986) 227
- 1986CL1B N.M. Clarke, P.J. Simmons, K.I. Pearce, S. Roman, A. Farooq and G. Rai, J. Phys. Soc. Jpn. Suppl. 55 (1986) 756
- 1986CR1B E. Cravo and A.C. Fonseca, Few-Body Syst. Suppl. (Austria) 1 (1986) 221
- 1986CS1A L.P. Csernai and J.I. Kapusta, Phys. Rept. 131 (1986) 223
- 1986DA1B D.H. Davis and J. Pniewski, Contemp. Phys. 27 (1986) 91
- 1986DE1H A. De Rujula, S.L. Glashow and L. Hall, Nature 320 (1986) 38
- 1986DE20 E. Dekempeneer, H. Liskien, L. Mewissen and F. Poortmans, Radiat. Eff. 92 (1986) 133; in Santa Fe 85 (1986) 133
- 1986DR10 M. Drosig, P.W. Lisowski, R.A. Hardekopf, D.M. Drake and K. Treitl, Radiat. Eff. 92 (1986) 145; in Santa Fe 85 (1986) 145
- 1986FA13 L.R. Fawcett, Jr., Radiat. Eff. 96 (1986) 239; in Santa Fe 85 (1986) 1581
- 1986FI15 B.W. Filippone, Ann. Rev. Nucl. Part. Sci. 36 (1986) 717
- 1986FIZX G.F. Filippov and S.A. Badalov, in Kharkov (1986) 149
- 1986FO04 D. Fox, D.A. Cebra, Z.M. Koenig, P. Ugorowski and G.D. Westfall, Phys. Rev. C33 (1986) 1540
- 1986FR12 V. Franco and Y. Yin, Phys. Rev. C34 (1986) 608
- 1986GA18 M. Garcon, B. Bonin, G. Bruge, J.C. Duchazeaubeneix, M. Rouger, J. Saudinos, B.H. Silverman, D.M. Sheppard, J. Arvieux, G. Gaillard et al., Nucl. Phys. A458 (1986) 287
- 1986GA24 D.L. Gay, L.C. Dennis and N.R. Fletcher, Phys. Rev. C34 (1986) 2144
- 1986GL1A Glaudemans, AIP Conf. Proc. 142 (1986) 316
- 1986GL1E Glukhov et al., in Kharkov (1986) 377, 378

- 1986GO1G K. Goswami, T.M. Singh and T.D. Goswami, *Indian J. Pure Appl. Phys.* 24 (1986) 7
- 1986GO1K Y. Gohar, *Radiat. Eff.* 92 (1986) 15; in *Sante Fe* 85 (1986) 15
- 1986GO23 S.A. Goncharov, Yu.I. Denisov, A.M. Mukhamedzhanov, E.A. Romanovsky, G.E. Valiev, I.R. Gulamov, T. Iskhakov, G. Ni, N.K. Timofeyuk, R. Yarmukhamedov et al., *Yad. Fiz.* 44 (1986) 303; *Sov. J. Nucl. Phys.* 44 (1986) 191
- 1986GOZY M. Gordon, R.L. McGrath, P. DeYoung, J. Alexander, X. Lu and D.M. De Castro-Rizzo, *Bull. Amer. Phys. Soc.* 31 (1986) 784, BH1
- 1986GR04 K. Grotz, H.V. Klapdor and J. Metzinger, *Phys. Rev. C* 33 (1986) 1263
- 1986GU1F Gupta and Malik, in *Harrogate* (1986) C23
- 1986HA1B B.G. Harvey, *J. Phys. Colloq. (Paris)* 47 (1986) C4-29
- 1986HA1I W.C. Haxton, *AIP Conf. Proc.* 150 (1986) 738
- 1986HA1K Y. Haneishi and T. Fujita, *Phys. Rev. C* 33 (1986) 260
- 1986HA1P Hanna, *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 528
- 1986HE26 E.F. Hefter and I.A. Mitropolsky, *Nuovo Cim.* A95 (1986) 63
- 1986HO33 H. Horiuchi, T. Wada and K. Yabana, *Prog. Theor. Phys. (Kyoto)* 76 (1986) 837
- 1986IG1A T. Iguchi, M. Nakazawa, H. Hashikura, J. Yamamoto, A. Takahashi, K. Sumita, K. Sugiyama, K. Kanda, S. Itagaki, S. Itoh et al., *Radiat. Eff.* 92 (1986) 203; in *Santa Fe* 85 (1986) 203
- 1986IR01 M. Iriondo, D. Jerrestam and R.J. Liotta, *Nucl. Phys.* A454 (1986) 252
- 1986JA02 L. Jarczyk, B. Kamys, Z. Rudy, A. Strzalkowski, B. Styczen, G.P.A. Berg, A. Magiera, J. Meissburger, W. Oelert, P. Von Rossen et al., *Nucl. Phys.* A448 (1986) 1
- 1986JA14 L. Jarczyk, B. Kamys, Z. Rudy, A. Strzalkowski, B. Styczen, G.P.A. Berg, A. Magiera, J. Meissburger, W. Oelert, P. von Rossen et al., *Nucl. Phys.* A459 (1986) 52
- 1986KN06 D.W. Kneff, B.M. Oliver, E. Goldberg and R.C. Haight, *Nucl. Sci. Eng.* 94 (1986) 136
- 1986KO32 A.L. Kochetkov and E.A. Seregina, *At. Energ.* 60 (1986) 271; *Sov. At. Energy* 60 (1986) 322
- 1986KR12 A.T. Kruppa, R.G. Lovas, R. Beck and F. Dickmann, *Phys. Lett.* B179 (1986) 317
- 1986LA05 K. Langanke and C. Rolfs, *Phys. Rev. C* 33 (1986) 790
- 1986LA12 K. Langanke, *Phys. Lett.* B174 (1986) 27
- 1986LA16 K. Langanke, M. Wiescher and F.-K. Thielemann, *Z. Phys.* A324 (1986) 147
- 1986LA19 K. Langanke and C. Rolfs, *Z. Phys.* A324 (1986) 307

- 1986LA26 M. Lattuada, F. Riggi, C. Spitaleri, D. Vinciguerra, D. Miljanic, M. Zadro and J. Yao, Nucl. Phys. A458 (1986) 493
- 1986LE1E Leeper et al., Bull. Amer. Phys. Soc. 31 (1986) 1286
- 1986LI15 H. Liskien, Radiat. Eff. 95 (1986) 271; in Santa Fe 85 (1986) 1277
- 1986MA03 A. Malecki and L. Satta, Phys. Rev. C33 (1986) 382
- 1986MA19 J.F. Mateja, A.D. Frawley, L.C. Dennis and K. Sartor, Phys. Rev. C33 (1986) 1649
- 1986MA1O MacDonald et al., in Harrogate (1986) C214
- 1986MA1R H. Maekawa, M. Nakagawa, Y. Ikeda, Y. Oyama, S. Yamaguchi, K. Tsuda, T. Fukumoto, A. Hasegawa, T. Mori, K. Kosako et al., Radiat. Eff. 92 (1986) 101; in Santa Fe 85 (1986) 101
- 1986MA1T G.J. Mathews, R.W. Bauer, S.D. Bloom, R.C. Haight, W.M. Howard, K. Takahashi and R.A. Ward, Radiat. Eff. 94 (1986) 161; in Santa Fe 85 (1986) 835
- 1986MA1W M. May, Nucl. Phys. A450 (1986) 179c
- 1986MC09 M.W. McNaughton, H. Spinka and H. Shimizu, Nucl. Instrum. Meth. Phys. Res. A243 (1986) 137
- 1986MI11 D. Miljanic, S. Blagus and M. Zadro, Phys. Rev. C33 (1986) 2204
- 1986NAZV S. Nakayama, T. Yamagata, M. Tanaka, K. Yuasa, T. Itahashi and H. Ogata, in Harrogate (1986) 138, B112
- 1986NAZZ J. Napolitano, J. Camp and S.J. Freedom, Bull. Amer. Phys. Soc. 31 (1986) 780, BF9
- 1986NO1C Y. Nojiri, K. Takeyama, K. Matsuta, K. Asahi and T. Minamisono, J. Phys. Soc. Jpn. Suppl. 55 (1986) 391
- 1986NO1D Y. Nojiri, K. Takeyama, K. Matsuta, K. Asahi and T. Minamisono, J. Phys. Soc. Jpn. Suppl. 55 (1986) 1086
- 1986OC1A T. Ochiai, Prog. Theor. Phys. 75 (1986) 1184
- 1986PAZN Yu.N. Pavlenko, Yu.O. Vasilev, A.S. Gass, V.A. Kiva, V.I. Medvedev, V.M. Pugach, V.A. Usik and A.M. Shidlyk, in Kharkov (1986) 320
- 1986PO06 D.N. Poenaru, W. Greiner, K. Depta, M. Ivascu, D. Mazilu and A. Sandulescu, At. Data Nucl. Data Tables 34 (1986) 423
- 1986PO12 J. Pochodzalla, R. Butsch, B. Heck and G. Rosner, Phys. Lett. B181 (1986) 33
- 1986QUZZ P.A. Quin and R.A. Bigelow, Bull. Amer. Phys. Soc. 31 (1986) 1219, CB5
- 1986RA15 Md.A. Rahman and H.M. Sen Gupta, Nuovo Cim. A93 (1986) 236
- 1986RA1D J. Raisanen and R. Lappalainen, Nucl. Instrum. Meth. Phys. Res. B15 (1986) 546
- 1986RA21 J. Rapaport, Radiat. Eff. 95 (1986) 223; in Santa Fe 85 (1986) 1229
- 1986RO13 C. Rolfs and R.W. Kavanagh, Nucl. Phys. A455 (1986) 179

- 1986RO1N S.P. Rosen and J.M. Gelb, in Harrogate (1986) F2
- 1986SA1P K.E. Sale, T.R. Donoghue, A. Abduljalil, J. Cummings, C. Rolfs, P. Schmalbrock, R.G. Seyler, T. Clegg, E.J. Ludwig, R. Fauber et al., J. Phys. Soc. Jpn. Suppl. 55 (1986) 992
- 1986SA30 H. Sato and Y. Okuhara, Phys. Rev. C34 (1986) 2171
- 1986SE1D H. Sekimoto, K. Oishi, K. Hojo, T. Hojo, D.W. Lee, S. Yukinori, N. Yamamuro, T. Hyodo and J. Yamamoto, Radiat. Eff. 92 (1986) 107; in Santa Fe 85 (1986) 107
- 1986SH10 S. Shimoura, A. Sakaguchi, T. Shimoda, T. Fukuda, K. Ogura, K. Katori and H. Ogata, Nucl. Phys. A452 (1986) 123
- 1986SH1T K. Shin, Y. Oyama and M. Abdou, Radiat. Eff. 92 (1986) 239; in Santa Fe 85 (1986) 239
- 1986SH33 K. Shibata and Y. Kikuchi, Radiat. Eff. 96 (1986) 243; in Santa Fe 85 (1986) 1585
- 1986SIZS P.J. Simmonds, N.M. Clarke, K.I. Pearce, R.J. Griffiths, C.A. Ogilvie and M. Mannion, in Harrogate (1986) C128, 315
- 1986SO07 J.P. Soderstrum and L.D. Knutson, Phys. Rev. C34 (1986) 401
- 1986ST1D E. Stenlund, Nucl. Phys. A447 (1986) 181c
- 1986ST1E E. Steffens, J. Phys. Soc. Jpn. Suppl. 55 (1986) 459
- 1986SU06 Y. Suzuki and K.T. Hecht, Nucl. Phys. A455 (1986) 315
- 1986TA1H A. Takahashi, Radiat. Eff. 92 (1986) 59; in Santa Fe 85 (1986) 59
- 1986TA1M B. Tamain, Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., No. 68, Vol. 2 (1986) 247
- 1986TA1N M.J. Tannenbaum, AIP Conf. Proc. 150 (1986) 858
- 1986TA1P M. Tanaka, AIP Conf. Proc. 150 (1986) 872
- 1986TU1B G. Tungate, K. Becker, K. Blatt, R. Butsch, P. Egelhof, D. Fick, B. Heck, H. Jansch, D. Kramer, H. Leucker et al., J. Phys. Soc. Jpn. Suppl. 55 (1986) 1116
- 1986UC1A J. Uckert, M. Burgel, H. Fuchs, H. Homeyer and A. Budzanowski, J. Phys. Colloq. (Paris) 47 (1986) 95
- 1986VD01 A.I. Vdovin, A.V. Golovin and I.I. Loshchakov, Yad. Fiz. 43 (1986) 1443; Sov. J. Nucl. Phys. 43 (1986) 930
- 1986VD04 A.I. Vdovin, I.G. Golikov, A.V. Golovin, M.N. Zhukov and I.I. Loshchakov, Izv. Akad. Nauk SSSR, Ser. Fiz. 50 (1986) 936; Bull. Acad. Sci. USSR, Phys. Ser. 50 (1986) 103
- 1986VE1A G. Velarde, J.M. Aragonés, J.J. Honrubia, J.M. Martínez-Val, E. Minguez, J.M. Perlado and J.M. Santolaya, Radiat. Eff. 92 (1986) 195; in Santa Fe 85 (1986) 195
- 1986WA01 E.K. Warburton, Phys. Rev. C33 (1986) 303
- 1986WE1C G.D. Westfall, Nucl. Phys. A447 (1986) 591c

- 1986WE1D H.R. Weller, *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 113
- 1986WI04 D.H. Wilkinson, *Nucl. Phys.* A452 (1986) 296
- 1986WO1B Wolfenstein, *Nature* 323 (1986) 579
- 1986XI1A X.-L. Xie, X.-J. Kong, C.-W. Sun and S.-M Wang, *Chin. Phys.* 6 (1986) 52
- 1986XU02 H.M. Xu, D.J. Fields, W.G. Lynch, M.B. Tsang, C.K. Gelbke, M.R. Maier, D.J. Morrissey, J. Pochodzalla, D.G. Sarantites, L.G. Sobotka et al., *Phys. Lett.* 182B (1986) 155
- 1986YA12 T. Yamaya, J.I. Hirota, K. Takimoto, S. Shimoura, A. Sakaguchi, S. Kubono, M. Sugitani, S. Kato, T. Suehiro and M. Fukada, *Phys. Rev.* C34 (1986) 2369
- 1986YA1F Y. Yamamoto, *Prog. Theor. Phys.* 75 (1986) 639
- 1986YA1K J. Yamamoto, A. Takahashi, K. Kanda, S. Itoh, K. Yugami, K. Sugiyama and K. Sumita, *Radiat. Eff.* 92 (1986) 171; in *Santa Fe* 85 (1986) 171
- 1986YO1D M. Youssef, J. Jung and M. Sawan, *Radiat. Eff.* 92 (1986) 117; in *Santa Fe* 85 (1986) 117
- 1986YU1A Y.-W. Yu, T. Motoba and H. Bando, *Prog. Theor. Phys.* 76 (1986) 861
- 1986ZA09 M. Zadro, S. Blagus, D. Miljanic, M. Lattuada, F. Riggi and C. Spitaleri, *Z. Phys.* A325 (1986) 119
- 1987AB21 A. Abzouzi, M.S. Antony and V.B. Ndocko Ndongue, *Nuovo Cim.* A97 (1987) 753
- 1987AJ02 F. Ajzenberg-Selove, *Nucl. Phys.* A475 (1987) 1
- 1987AJ1A F. Ajzenberg-Selove, in *Dubna* (1987) 341
- 1987AR19 S.E. Arnell, S. Mattsson, H.A. Roth, M. Rydehell, O. Skeppstedt, A. Johnson, J. Nyberg, A. Kerek and A. Nilsson, *Phys. Scr.* 36 (1987) 214
- 1987AR1G Artemov et al., in *Yurmala* (1987) 386
- 1987AR22 E. Arnold, J. Bonn, R. Gegenwarth, W. Neu, R. Neugart, E.-W. Otten, G. Ulm, K. Wendt and ISOLDE Collaboration, *Phys. Lett.* B197 (1987) 311
- 1987AS05 H.J. Assenbaum, K. Langanke and C. Rolfs, *Z. Phys.* A327 (1987) 461
- 1987BA13 J. Banaigs, J. Berger, P. Berthet, G. Bizard, M. Boivin, M. De Sanctis, J. Dufflo, F.L. Fabbri, R. Frascaria, L. Goldzahl et al., *Phys. Rev.* C35 (1987) 1416
- 1987BA1X J.N. Bahcall, A. Dar and T. Piran, *Nature* 326 (1987) 135
- 1987BA2K H. Bando, T. Yamada and J. Zofka, *Phys. Rev.* C36 (1987) 1640
- 1987BA35 D. Baye, *Phys. Rev. Lett.* 58 (1987) 2738
- 1987BA38 G.J. Balster, P.C.N. Crouzen, P.B. Goldhoorn, R.H. Siemssen and H.W. Wilschut, *Nucl. Phys.* A468 (1987) 93
- 1987BA89 J.N. Bahcall, *Rev. Mod. Phys.* 59 (1987) 505

- 1987BE17 H.W. Becker, C. Rolfs and H.P. Trautvetter, *Z. Phys.* A327 (1987) 341
- 1987BE58 B. Berthier, R. Boisgard, J. Julien, J.M. Hisleur, R. Lucas, C. Mazur, C. Ngo, M. Ribrag and C. Cerruti, *Phys. Lett.* B193 (1987) 417
- 1987BEYI A.V. Belozyorov, C. Borcea, Z. Dlouhy, A.M. Kalinin, N.H. Chau and Yu.E. Penionzhkevich, *JINR-E15-87-733* (1987)
- 1987BL13 C. Bloch, W. Bensenson, A.I. Galonsky, E. Kashy, J. Heltsley, L. Heilbronn, M.Lowe, B. Remington, D.J. Morrissey and J. Kasagi, *Phys. Rev.* C36 (1987) 203
- 1987BL16 C. Bloch and E. Kashy, *Phys. Rev.* C36 (1987) 855
- 1987BL18 R. Blumel and K. Dietrich, *Nucl. Phys.* A471 (1987) 453
- 1987BO1M Bochkarev et al., in *Yurmala* (1987) 384
- 1987BO40 C. Borcea, A.V. Belozyorov, Z. Dlouhy, A.M. Kalinin, N.H. Chau and Yu.E. Penionzhkevich, *Rev. Roum. Phys.* 32 (1987) 497
- 1987BO42 M.C. Bosca, E. Buendia and R. Guardiola, *Phys. Lett.* B198 (1987) 312
- 1987CA05 Z.-J. Cao, R.D. Bent, H. Nann and T.E. Ward, *Phys. Rev.* C35 (1987) 625
- 1987CA06 Z.-J. Cao, T.E. Ward, R.D. Bent, J.D. Brown, E. Korkmaz, H. Nann and T.G. Throwe, *Phys. Rev.* C35 (1987) 825
- 1987CH1G Cherry, Davis and Lande, *Telemark IV: Neutrino Masses and Neutrino Astrophys.*, Ashland (1987)
- 1987CH26 Z. Chen, C.K. Gelbke, J. Pochodzalla, C.B. Chitwood, D.J. Fields, W.G. Gong, W.G. Lynch and M.B. Tsang, *Nucl. Phys.* A473 (1987) 564
- 1987CH32 Z. Chen, C.K. Gelbke, W.G. Gong, Y.D. Kim, W.G. Lynch, M.R. Maier, J. Pochodzalla, M.B. Tsang, F. Saint-Laurent, D. Ardouin et al., *Phys. Lett.* B199 (1987) 171
- 1987CH33 Z. Chen, C.K. Gelbke, W.G. Gong, Y.D. Kim, W.G. Lynch, M.R. Maier, J. Pochodzalla, M.B. Tsang, F. Saint-Laurent, D. Ardouin et al., *Phys. Rev.* C36 (1987) 2297
- 1987CO16 J. Cook, M.N. Stephens and K.W. Kemper, *Nucl. Phys.* A466 (1987) 168
- 1987CZ02 J. Czakanski, W. Zipper, W. Dunnweber, W. Hering, D. Konnerth, W. Trombik, K.G. Bernhardt, H. Bohn, K.A. Eberhard and R. Vandenbosch, *Phys. Lett.* 199 (1987) 166
- 1987DA25 N. Dai, B. Qi, Y. Mao, F. Zhuang, J. Yao and X. Wang, *Chin. J. Nucl. Phys.* 9 (1987) 103
- 1987DE1O Detraz, in *Dubna* (1987) 42
- 1987DE37 F. Deak, A. Kiss, Z. Seres, G. Caskey, A. Galonsky and B. Remington, *Nucl. Instrum. Meth. Phys. Res.* A258 (1987) 67
- 1987DU07 J. Duflo, *Phys. Rev.* C36 (1987) 1425
- 1987FL1A Flerov, in *Dubna* (1987) 9

- 1987FO08 D. Fox, D.A. Cebra, J. Karn, C. Parks, G.D. Westfall and W.K. Wilson, Phys. Rev. C36 (1987) 640
- 1987FR1C G. Friedlander and J. Weneser, Science 235 (1987) 760
- 1987FR1D V. Franco, Phys. Rev. C35 (1987) 1328
- 1987FU04 I. Fushiki and D.Q. Lamb, Astrophys. J. 317 (1987) 368
- 1987FU1G G.M. Fuller, R.W. Mayle, J.R. Wilson and D.N. Schramm, Astrophys. J. 322 (1987) 795
- 1987GAZM S.A. Gaidenko, E.P. Kadkin, M.V. Pasechnik, L.S. Saltykov and A.D. Fursa, in Yurmala (1987) 299
- 1987GE1B C.-K. Gelbke and D.H. Boal, Prog. Part. Nucl. Phys. 19 (1987) 33
- 1987GL05 Yu.A. Glukhov, A.S. Demyanova, A.A. Ogloblin and S.B. Sakuta, Yad. Fiz. 45 (1987) 1236; Sov. J. Nucl. Phys. 45 (1987) 767
- 1987GL1D Gledenov et al., in Yurmala (1987) 404
- 1987GLZZ Yu.M. Gledenov, T.S. Zvarova, I. Kvitek, M.P. Mitrikov, R.S. Mitrikova, Yu.P. Popov, V.I. Salatsky and V.Z. Fung, in Yurmala (1987) 298
- 1987GO1S Goryunov et al., in Yurmala (1987) 474
- 1987GO27 S.A. Goncharov, A.M. Mukhamedzhanov, E.A. Romanovsky, G.E. Valiev, I.R. Gulamov, T. Iskhakov, G. Nie, N.K. Tomopheyuk, R. Yarmukhamedov, V. Kroha et al., Czech. J. Phys. B37 (1987) 168
- 1987GR11 R.E.L. Green, R.G. Korteling, J.M. D'Auria, K.P. Jackson and R.L. Helmer, Phys. Rev. C35 (1987) 1341
- 1987HA30 P.G. Hansen and B. Jonson, Europhys. Lett. 4 (1987) 409
- 1987HA45 D. Hahn and H. Stocker, Phys. Rev. C35 (1987) 1311
- 1987HE1B U. Heinz, Rept. Prog. Phys. 50 (1987) 145
- 1987HU12 G.M. Huber, G.J. Lolos, Z. Papandreou, K.H. Hicks, P.L. Walden, S. Yen, E.G. Auld, F.A. Duncan, R.D. Bent and W.R. Falk, Phys. Rev. C36 (1987) 2683
- 1987JA06 B.V. Jacak, G.D. Westfall, G.M. Crawley, D. Fox, C.K. Gelbke, L.H. Harwood, B.E. Hasselquist, W.G. Lynch, D.K. Scott, H. Stocker et al., Phys. Rev. C35 (1987) 1751
- 1987KA18 K. Kato, H. Kazama and H. Tanaka, Prog. Theor. Phys. 77 (1987) 185
- 1987KA1L Kadmenskii et al., in Yurmala (1987) 473
- 1987KA1R T. Kajino, H. Toki and S.M. Austin, Astrophys. J. 319 (1987) 531
- 1987KA1W M. Kamran and E. Qureshi, Phys. Lett. B188 (1987) 122
- 1987KA25 K. Kadija, G. Paic, D. Vranic, A. Berdoz, F. Foroughi, Ch. Nussbaum, H.H. Muller, P. Schober and St. Steiner, Nucl. Phys. A469 (1987) 183
- 1987KI01 K.H. Kim, M.H. Park and B.T. Kim, Phys. Rev. C35 (1987) 363

- 1987KI1C H.R. Kissener, I. Rotter and N.G. Goncharova, Fortschr. Phys. 35 (1987) 277
- 1987KN04 H.D. Knox, D.A. Resler and R.O. Lane, Nucl. Phys. A466 (1987) 245
- 1987KO1E Kozmyr, in Yurmala (1987) 332
- 1987KO1U L. Kocbach and T.P. Walderhaug, XV ICPEAC Brighton '87 (1987) 623
- 1987KO1Z Kobayashi et al., in Panic (1987) 478
- 1987KR03 D. Krolle, H.J. Assenbaum, C. Funck and K. Langanke, Phys. Rev. C35 (1987) 1631
- 1987KR10 D. Krofcheck, E. Sugarbaker, A.J. Wagner, J. Rapaport, D. Wang, J.N. Bahcall, R.C. Byrd, C.C. Foster, C.D. Goodman, C. Gaarde et al., Phys. Lett. B189 (1987) 299
- 1987LA25 M. Lattuada, F. Riggi, D. Vinciguerra, C. Spitaleri and D. Miljanic, Z. Phys. A328 (1987) 497
- 1987LE33 F.S. Levin, Nucl. Phys. A463 (1987) 487c
- 1987LY04 W.G. Lynch, Nucl. Phys. A471 (1987) 309c
- 1987LY05 Yu.S. Lyutostansky and V.N. Tikhonov, Yad. Fiz. 46 (1987) 66; Sov. J. Nucl. Phys. 46 (1987) 42
- 1987MA2C Malaney and Fowler, OAP-680, To be published in Origin and Distribution of the Elements (1987)
- 1987MI38 M. Mian, Phys. Rev. C35 (1987) 1463
- 1987MU1B R.J. Murphy, C.D. Dermer and R. Ramaty, Astrophys. J. Suppl. 63 (1987) 721
- 1987NA08 J. Napolitano, S.J. Freedman and J. Camp, Phys. Rev. C36 (1987) 298
- 1987NE1C Nemets et al., in Yurmala (1987) 323
- 1987NO04 Y. Nojiri, K. Takeyama, K. Matsuta and T. Minamisono, Hyperfine Interactions 35 (1987) 1015
- 1987OC1B T. Ochiai, Z. Phys. C35 (1987) 209
- 1987OL04 B.M. Oliver, H. Farrar IV and M.M. Bretscher, Int. J. Appl. Radiat. Isotop. 38 (1987) 959
- 1987OT1D I. Otterlund, Nucl. Phys. A461 (1987) 113c
- 1987PE1B Peter, in Dubna (1987) 562
- 1987PE1C Penionshkevich, in Dubna (1987) 364
- 1987PO03 J. Pochodzalla, C.K. Gelbke, W.G. Lynch, M. Maier, D. Ardouin, H. Delagrang, H. Doubre, C. Gregoire, A. Kyanowski, W. Mittag et al., Phys. Rev. C35 (1987) 1695
- 1987PO1H B. Povh, Prog. Part. Nucl. Phys. 18 (1987) 183
- 1987PO23 J. Pochodzalla, Nucl. Phys. A471 (1987) 289c
- 1987PR01 D. Provoost, F. Grummer and K. Goeke, Ann. Phys. (New York) 174 (1987) 202
- 1987QA01 S.M. Qaim and R. Wolfle, Nucl. Sci. Eng. 96 (1987) 52

- 1987RI1E J. Rich, D.L. Owen and M. Spiro, Phys. Rept. 151 (1987) 239
- 1987RO25 C. Rolfs, H.P. Trautvetter and W.S. Rodney, Rept. Prog. Phys. 50 (1987) 233
- 1987RUZX N.Ya. Rutkevich, V.Ya. Golovnya, S.N. Oleinik and E.V. Sheptulenko, in Yurmala (1987) 320
- 1987SA15 H. Sagawa and H. Toki, J. Phys. (London) G13 (1987) 453
- 1987SA1L K.E. Sale, R.W. Bauer, R.N. Boyd, G.J. Mathews, R.C. Haight and P.B. Corn, Nucl. Instrum. Meth. Phys. Res. B24-25 (1987) 490
- 1987SA37 S. Saito, Nucl. Phys. A472 (1987) 161
- 1987SC08 D. Schmidt, D. Seeliger, G.N. Lovchikova and A.M. Trufanov, Nucl. Sci. Eng. 96 (1987) 159
- 1987SH1M S.Yu. Shmakov and V.V. Uzhinskii, Z. Phys. C36 (1987) 77
- 1987SI06 K. Siwek-Wilczynska, J. Wilczynski, C.R. Albiston, Y. Chan, E. Chavez, S.B. Gazes, H.R. Schmidt and R.G. Stokstad, Phys. Rev. C35 (1987) 1316
- 1987SIZX I.V. Simenog, in Yurmala, (1987) 167
- 1987SV1A V.A. Sviciulis and R.K. Kalinauskas, Sov. Phys.-Collect. 27 (1987) 10
- 1987TA22 T.N. Taddeucci, Can. J. Phys. 65 (1987) 557
- 1987TAZU I. Tanihata, H. Hamagaki, O. Hashimoto, Y. Shida, O. Yamakawa, T. Kobayashi, K. Sugimoto and N. Takahashi, in Panic (1987) 474; LBL-22820 (1987) 82
- 1987TI07 D.R. Tilley, H.R. Weller and H.H. Hasan, Nucl. Phys. A474 (1987) 1
- 1987TO06 M. Tosaki, M. Fujiwara, K. Hosono, T. Noro, H. Ito, T. Yamazaki and H. Ikegami, Nucl. Phys. A463 (1987) 429c
- 1987VA1I Valiev et al., in Yurmala (1987) 346
- 1987VA26 A.G.M. van Hees, A.A. Wolters and P.W.M. Glaudemans, Phys. Lett. B196 (1987) 19
- 1987VE02 A.N. Verma, M. Goswami, F. Ahmed and L.S. Kothari, Ann. Nucl. Energy 14 (1987) 159
- 1987WA07 H. Walliser and S. Nakaichi-Maeda, Nucl. Phys. A464 (1987) 366
- 1987WA09 R. Wada, K.D. Hildenbrand, U. Lynen, W.F.J. Muller, H.J. Rabe, H. Sann, H. Stelzer, W. Trautmann, R. Trockel, N. Brummund et al., Phys. Rev. Lett. 58 (1987) 1829
- 1987WA21 R.E. Warner, B.A. Vaughan, J.A. Ditusa, J.W. Rovine, R.S. Wakeland, C.P. Browne, S.E. Darden, S. Sen, A. Nadasen, A. Basak et al., Nucl. Phys. A470 (1987) 339
- 1987WA25 R.E. Warner, F.G. Johnson, C.P. Browne, A. Rollefson, A. Galonsky and A. Nadasen, Nucl. Phys. A472 (1987) 522
- 1987WA36 X.-C. Wang, H. Bando and H. Takaki, Z. Phys. A 327 (1987) 59

- 1987WAZT J.W. Watson, W.P. Alford, R. Helmer, C. Zafiratos, R. Abegg, A. Celler, S. El-Kateb, D. Frekers, O. Hausser, R. Henderson et al., Bull. Amer. Phys. Soc. 32 (1987) 1578, EC1
- 1987WE1C J. Weneser and G. Friedlander, Science 235 (1987) 755
- 1987YA1M Y. Yamamoto, Phys. Rev. C36 (1987) 2166
- 1987ZA07 M. Zadro, D. Miljanic, M. Lattuada, F. Riggi and C. Spitaleri, Nucl. Phys. A474 (1987) 373
- 1988AJ1B F. Ajzenberg-Selove, Interact. and Struct. in Nucl., Proc. in Honor of D.H. Wilkinson, Sussex, 7-9 Sept. 1987 (1988) 181
- 1988AL07 S.C. Allcock, W.D.M. Rae, P.R. Keeling, A.E. Smith, B.R. Fulton and D.W. Banes, Phys. Lett. B201 (1988) 201
- 1988AL1G Aleksandrov et al., in Baku (1988) 377
- 1988AR05 N. Arena, Seb. Cavallaro, G. Fazio, G. Giardina, A. Italiano and F. Mezzanares, Europhys. Lett. 5 (1988) 517
- 1988AU1A Aushev et al., in Baku (1988) 369
- 1988BA29 F.C. Barker, Phys. Rev. C37 (1988) 2920
- 1988BA75 F.C. Barker, Aust. J. Phys. 41 (1988) 743
- 1988BA86 J.N. Bahcall and R.K. Ulrich, Rev. Mod. Phys. 60 (1988) 297
- 1988BEYJ A.V. Belozarov, K. Borchia, I. Vintsour, Z. Dlougy, N.K. Tyau and Yu.Eh. Penionzhkevich, in Baku (1988) 380
- 1988BL09 C. Bloch, W. Benenson, A.I. Galonsky, E. Kashy, J. Heltsley, L. Heilbronn, M. Lowe, R.J. Radtke, B. Remington, J. Kassagi et al., Phys. Rev. C37 (1988) 2469
- 1988BO04 M.C. Bosca and R. Guardiola, Nucl. Phys. A476 (1988) 471
- 1988BO15 C.D. Bowman, J. Phys. (London) G14 Suppl. (1988) S399
- 1988BO47 K. Bodek, B. Bruggemann, J. Krug, P. Lekkas, W. Lubcke, H. Ruhl, M. Steinke, M. Stephan, A. Szczurek and D. Kamke, Few-Body Syst. 3 (1988) 135
- 1988CA06 G. Caskey, L. Heilbronn, B. Remington, A. Galonsky, F. Deak, A. Kiss and Z. Seres, Phys. Rev. C37 (1988) 696
- 1988EW1A G.T. Ewan, AIP Conf. Proc. 164 (1988) 585
- 1988FO1A W.A. Fowler, Interact. and Struct. in Nuclei, Proc. in Honor of D.H. Wilkinson, Sussex, 7-9 Sept. 1987 (1988) 119
- 1988GA10 C.A. Gagliardi, D.R. Semon, E. Takada, D.M. Tanner and R.E. Tribble, Phys. Rev. C37 (1988) 2889
- 1988GO02 S.A. Goncharov, A.M. Mukhamedzhanov, E.A. Romanovsky, G.E. Valiev, I.R. Gulamov, Yu.I. Denisov, T. Iskhakov, A. Karakhodzhaev, G.K. Nie, R. Yarmukhamedov et al., Czech. J. Phys. B38 (1988) 12

- 1988GU1F Guzhovsky, Lazarev and Ershov, in Baku (1988) 482
- 1988HA1M W.C. Haxton and C.W. Johnson, Nature 333 (1988) 325
- 1988HE08 R. Henneck, C. Gysin, P. Haffter, M. Hammans, W. Lorenzon, M.A. Pickar, I. Sick and S. Burzynski, Phys. Rev. C37 (1988) 2224
- 1988JO1C B. Jonson, S. Mattsson, G. Nyman, O. Tengblad, M.J.G. Borge, P.G. Hansen and K. Riisager, AIP Conf. Proc. 164 (1988) 223
- 1988KA1K Karmanov et al., in Baku (1988) 326
- 1988KA1M Karmanov et al., in Baku (1988) 488
- 1988KH03 K.S. Myint and Y. Akaishi, Prog. Theor. Phys. 79 (1988) 454
- 1988KI05 J.M. Kidd, P.J. Lindstrom, H.J. Crawford and G. Woods, Phys. Rev. C37 (1988) 2613
- 1988KI1C Kiptily, in Baku (1988) 534
- 1988KO03 P.E. Koehler, C.D. Bowman, F.J. Steinkruger, D.C. Moody, G.M. Hale, J.W. Starner, S.A. Wender, R.C. Haight, P.W. Lisowski and W.L. Talbert, Phys. Rev. C37 (1988) 917
- 1988KR01 A.T. Kruppa, R.G. Lovas and B. Gyarmati, Phys. Rev. C37 (1988) 383
- 1988KU1E Kukulin, in Baku (1988) 396
- 1988KW02 E. Kwasniewicz and J. Kisiel, Acta Phys. Pol. B19 (1988) 141
- 1988LI1A Liu et al., Bull. Amer. Phys. Soc. 33 (1988) 903
- 1988LI27 G.-B. Liu and H.T. Fortune, Phys. Rev. C38 (1988) 1985
- 1988MA1H Manokhin, INDC(CCP)-283 (1988)
- 1988MO05 K. Mohring, T. Srokowski, D.H.E. Gross and H. Homeyer, Phys. Lett. B203 (1988) 210
- 1988NE1A Nemets, Pavlenko and Pugach, in Baku (1988) 325
- 1988POZS N.A.F.M. Poppelier, J.H. de Vries, A.A. Wolters and P.W.M. Glaudemans, AIP Conf. Proc. 164 (1988) 334
- 1988POZZ J. Pouliot, Y.D. Chan, A. Dacal, B.A. Harmon, R. Knop, M.E. Ortiz, E. Plagnol, R.G. Stokstad, C. Moisan, L. Potvin et al., Bull. Amer. Phys. Soc. 33 (1988) 1179, AG1
- 1988RIZZ B.G. Ritchie, D. Rothenberger, J.R. Comfort, R.A. Giannelli, J. Tinsley, N.S. Chant, D. Mack, P.G. Roos, J.D. Silk, G.S. Kyle et al., Bull. Amer. Phys. Soc. 33 (1988) 903, AI11
- 1988RU01 V.A. Rubchenya and S.G. Yavshits, Z. Phys. A329 (1988) 217
- 1988RU1B Ruza et al., in Baku (1988) 146
- 1988RU1D Rubchenya and Yavchits, in Baku (1988) 341

- 1988SA09 F. Saint-Laurent, A. Kyanowski, D. Ardouin, H. Delagrange, H. Doubre, C. Gregoire, W. Mittig, A. Peghaire, J. Peter, G. Bizard et al., Phys. Lett. B202 (1988) 190
- 1988SA2A M.J. Savage, B.W. Filippone and L.W. Mitchell, Phys. Rev. D37 (1988) 1134
- 1988SH1E Shvedov, Nemets and Rudchik, in Baku (1988) 351
- 1988SH1F Shvedov, Nemets and Rudchik, in Baku (1988) 352
- 1988ST06 J. Stevenson, B.A. Brown, Y. Chen, J. Clayton, E. Kashy, D. Mikolas, J. Nolen, M. Samuel, B. Sherrill, J.S. Winfield et al., Phys. Rev. C37 (1988) 2220
- 1988TA1A I. Tanihata, Nucl. Phys. A478 (1988) 795c
- 1988TA29 H. Tamura, W. Bruckner, H. Dobbeling, R.S. Hayano, T. Ishikawa, M. Iwasaki, T. Motoki, H. Outa, S. Paul, B. Povh et al., Nucl. Phys. A479 (1988) 161c
- 1988VA03 A.G.M. van Hees, A.A. Wolters and P.W.M. Glaudemans, Nucl. Phys. A476 (1988) 61
- 1988VA1E Vagner et al., in Baku (1988) 383
- 1988WA18 A.H. Wapstra, G. Audi and R. Hoekstra, At. Data Nucl. Data Tables 39 (1988) 281
- 1988WA1E E.K. Warburton, Interact. and Struct. in Nucl., Proc. in Honor of D.H. Wilkinson, Sussex, 7-9 Sept. 1987 (1988) 81
- 1988WO04 A.A. Wolters, A.G.M. van Hees and P.W.M. Glaudemans, Europhys. Lett. 5 (1988) 7
- 1990AJ01 F. Ajzenberg-Selove, Nucl. Phys. A506 (1990) 1

