

# Energy Levels of Light Nuclei $A = 9$

F. Ajzenberg-Selove<sup>a</sup> and T. Lauritsen<sup>b</sup>

<sup>a</sup> *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

<sup>b</sup> *California Institute of Technology, Pasadena, California*

**Abstract:** An evaluation of  $A = 5-24$  was published in *Nuclear Physics* 11 (1959), p. 1. This version of  $A = 9$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 01, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 9$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides: [<sup>9</sup>Li](#), [<sup>9</sup>Be](#), [<sup>9</sup>B](#), [<sup>9</sup>C](#)

B. Tables of Recommended Level Energies:

[Table 9.1](#): Energy levels of <sup>9</sup>Be

[Table 9.4](#): Energy levels of <sup>9</sup>B


C. [References](#)

D. Figures: [<sup>9</sup>Be](#), [<sup>9</sup>B](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

**<sup>9</sup>Li**  
(Not illustrated)

*Mass of <sup>9</sup>Li:* From the threshold for <sup>9</sup>Be(d, 2p)<sup>9</sup>Li,  $E_d = 19 \pm 1$  MeV (1951GA30), the mass excess of <sup>9</sup>Li is determined as  $M - A = 28.1 \pm 1$  MeV.



<sup>9</sup>Li decays to excited states of <sup>9</sup>Be which decay by neutron emission. The mean of the reported half-lives is  $0.169 \pm 0.003$  sec (1951GA30, 1952HO25). See also (1952SH44, 1953FR03, 1955BE1E, 1956FL1A, 1958TA04).



The threshold is  $19 \pm 1$  MeV (1951GA30).



See (1952SH44, 1958TA04).



See (1953RE19, 1958TA04).

The following reactions are not reported: <sup>7</sup>Li(t, p)<sup>9</sup>Li ( $Q_m = -2.9$ ), <sup>9</sup>Be(n, p)<sup>9</sup>Li ( $Q_m = -13.3$ ), <sup>9</sup>Be(t, <sup>3</sup>He)<sup>9</sup>Li ( $Q_m = -14.1$ ) and <sup>11</sup>B(n, <sup>3</sup>He)<sup>9</sup>Li ( $Q_m = -23.6$ ).

**<sup>9</sup>Be**  
(Fig. 10)

GENERAL:

*Theory:* See (1955DA1E, 1955FR1F, 1956BL1B, 1956DE1C, 1956KU1A, 1957BA1H, 1957PA1A, 1958KU1B).

Table 9.1: Energy levels of  ${}^9\text{Be}$

$E_x$ in ${}^9\text{Be}$ (MeV)	$J^\pi; T$	$\Gamma$ (keV)	Decay	Reactions
0	$\frac{3}{2}$		stable	2, 7, 13, 14, 15, 16, 17, 18, 19, 20, 22, 25, 26, 27
$1.75 \pm 0.02$	$\frac{1}{2}^{(+)}$	150	n, $\gamma$	7, 10, 14, 15, 16, 17, 25
$2.430 \pm 0.002$	$(\frac{5}{2}^-)$	< 1	(n), $\alpha$	7, 13, 14, 15, 16, 17, 20, 22, 25, 26
$3.04 \pm 0.016$	$(\leq \frac{3}{2})$	$161 \pm 15$		7, 15, 16, 17, 22, 25
$(4.74 \pm 0.08)$		1250		7, 15, 16
$6.76 \pm 0.06$				13, 15, 16, 17
$(7.94 \pm 0.08)$				15
$(9.1 \pm 0.2)$		1200		7
$(11.3 \pm 0.2)$			n, $\gamma$	10, 15, 17
(13.3)			n, $\gamma$	10
(14.5)				15
17.28		200	(d, p, n)	3, 4
17.48		50	(d, p, n)	3, 4, 15
(18.1)		(300)	(d, p, n)	3, 4
(18.6)			(d, p)	3
19.6			(d, p)	3, 15
$(21.7 \pm 0.1)$			p, $\gamma$ , n	11, 15

1. (a) ${}^6\text{Li}(t, d){}^7\text{Li}$	$Q_m = 0.994$	$E_b = 17.687$
(b) ${}^6\text{Li}(t, p){}^8\text{Li}$	$Q_m = 0.803$	
(c) ${}^6\text{Li}(t, n){}^8\text{Be}$	$Q_m = 16.021$	
(d) ${}^6\text{Li}(t, \alpha){}^5\text{He}$	$Q_m = 15.158$	
(e) ${}^6\text{Li}(t, n){}^4\text{He} + {}^4\text{He}$	$Q_m = 16.115$	

The differential cross section at  $90^\circ$  for reaction (a) rises steeply from 8.8 mb/sr at  $E_t = 0.72$  MeV to 19 mb at 0.90 MeV, and then more slowly to 21 mb at  $E_t = 1.15$  MeV. For reaction (d), the differential cross section at  $90^\circ$  rises from 0.75 mb/sr at 0.62 MeV to 5.7 mb/sr at 1.8 MeV and then decreases slowly to 5.3 mb/sr at 2.2 MeV (R.W. Crews, quoted in (1957JA37)).

See also (1952PE02),  ${}^5\text{He}$ ,  ${}^7\text{Li}$ ,  ${}^8\text{Li}$  and  ${}^8\text{Be}$ .

2. ${}^6\text{Li}(\alpha, p){}^9\text{Be}$	$Q_m = -2.125$
--	----------------

See (1956WA29).

3. ${}^7\text{Li}(d, p){}^8\text{Li}$	$Q_m = -0.192$	$E_b = 16.693$
---------------------------------------	----------------	----------------

Cross sections have been measured for  $E_d = 0.4$  to 1.8 MeV by (1952BA1A), and for  $E_d = 0.7$  to 3.3 MeV by (1954BA46); see (1957JA37). The yield for  $E_d = 1.1$  to 4.0 MeV has been measured by (1956BE1A). Resonances are observed at 0.80 and 1.04 MeV (1952BA1A, 1954BA46), 1.4 MeV (1952BA1A: see, however, (1954BA46)), 2.0, 2.5 and 3.7 MeV (1956BE1A: see, however, (1954BA46)). See also (1955AJ61).

4. (a) ${}^7\text{Li}(d, n){}^8\text{Be}$	$Q_m = 15.026$	$E_b = 16.693$
(b) ${}^7\text{Li}(d, \alpha){}^5\text{He}$	$Q_m = 14.163$	
(c) ${}^7\text{Li}(d, n){}^4\text{He} + {}^4\text{He}$	$Q_m = 15.121$	

The cross section for (a) has been measured for  $E_d = 70$  to 110 keV by (1955RA14) and that for (b) has been measured for  $E_d = 30$  to 250 keV by (1953SA1A). Resonances for neutron production occur, in the range  $E_d = 0.2$  to 4.8 MeV, at  $E_d = 0.68$  ( $\Gamma = 250$  keV), 0.98 ( $\Gamma = 60$  keV) and 1.8 MeV ( $\Gamma = 400$  keV) (1952BA1A, 1957SL01). At  $E_d = 0.90$  MeV, the  $\alpha$ -particles in reaction (b) are isotropic to within 2%, consistent with formation by s-wave deuterons. No evidence is found for direct interaction effects at this energy (1957RI39). The angular correlation of ground-state  $\alpha$ -particles with those resulting from breakup of  ${}^5\text{He}$  indicate  $J = \frac{5}{2}^-$  (1956RI37),  $J = \frac{3}{2}^-$  (1957FA10), for the  ${}^9\text{Be}$  level mainly responsible for this reaction at  $E_d = 0.9$  MeV.

Reactions (a) and (c) account for less than 10% of the disintegrations at this energy (1956RI37). See also  $^5\text{He}$ ,  $^8\text{Be}$  and (1956CA1B).

5. (a)  $^7\text{Li}(d, d)^7\text{Li}$   $E_b = 16.693$   
 (b)  $^7\text{Li}(d, t)^6\text{Li}$   $Q_m = -0.994$   
 (c)  $^7\text{Li}(d, ^3\text{He})^6\text{He}$   $Q_m = -4.512$

The cross section for reaction (b) rises steeply from threshold to 95 mb at  $E_d = 2.4$  MeV, and then more slowly to about 165 mb at  $E_d = 4.1$  MeV (1955MA20). See also  $^6\text{He}$  and  $^7\text{Li}$ .

6.  $^7\text{Li}(t, n)^9\text{Be}$   $Q_m = 10.435$

Not observed.

7.  $^7\text{Li}(^3\text{He}, p)^9\text{Be}$   $Q_m = 11.200$

At  $E_d = 0.7$  and 1 MeV, proton groups are observed corresponding to excited states of  $^9\text{Be}$  at  $1.83 \pm 0.04$  MeV ( $\Gamma < 0.4$  MeV),  $2.39 \pm 0.08$  MeV ( $\Gamma < 0.2$  MeV),  $3.10 \pm 0.04$  MeV ( $\Gamma < 0.3$  MeV),  $4.74 \pm 0.08$  MeV ( $\Gamma = 1.25 \pm 0.25$  MeV) and  $(9.1 \pm 0.2)$  MeV ( $\Gamma = 1.2$  MeV) (1954MO92, 1955AL57, 1958MO99). The 1.8-MeV level admits a description in terms of an s-wave n- $^8\text{Be}$  interaction with a scattering length  $a = 23 \times 10^{-13}$  cm (1958MO99: see  $^{11}\text{B}(d, \alpha)^9\text{Be}$ ).

8.  $^7\text{Li}(\alpha, d)^9\text{Be}$   $Q_m = -7.151$

Not observed.

9.  $^9\text{Li}(\beta^-)^9\text{Be}^* \rightarrow ^8\text{Be} + n$   $Q_m = 12.4$

See  $^9\text{Li}$ .

10. (a)  $^9\text{Be}(\gamma, n)^8\text{Be}$   $Q_m = -1.667$   
 (b)  $^9\text{Be}(\gamma, \alpha)^5\text{He}$   $Q_m = -2.529$   
 (c)  $^9\text{Be}(\gamma, n)^4\text{He} + ^4\text{He}$   $Q_m = -1.572$   
 (d)  $^9\text{Be}(\gamma, 2n)^7\text{Be}$   $Q_m = -20.565$

The photo-neutron cross section has been studied from threshold ( $E_\gamma = 1.664 \pm 0.004$  MeV: (1956CO56)) to 320 MeV: see Table 9.2. A sharp peak,  $\Gamma < 50$  keV,  $\sigma \approx 100$  mb, occurs at  $E_\gamma = 1.70$  MeV (1956CO56). A further broad maximum appears at  $E_\gamma \approx 10$  MeV, followed by the giant resonance at 20 – 22 MeV (1953JO1B, 1953NA1A: see (1956CO59)). Measurements from 6 to 18 MeV show pronounced peaks at 11.3 and 13.3 MeV and a broad, low peak at 16 MeV, immediately preceding the giant resonance (1959SP1B). At  $E_\gamma = 6.1$  MeV the main processes appear to involve  ${}^9\text{Be}(\gamma, n){}^8\text{Be}^*(2.9)$  and  ${}^9\text{Be}(\gamma, \alpha){}^5\text{He}$  (1954CA1A). Angular distributions for  $E_\gamma = 1.7$  and 1.8 MeV are spherically symmetric; at  $E_\gamma = 2.76$  MeV,  $W(\theta) = 1.2 + \sin^2\theta$  (1949HA1A: see also (1954NI1B, 1956FA1B)).

Calculations of (1949GU1A) and (1956MA1M) envisage E1 transitions to  $p^4s$  and  $p^4d$  configurations, with levels at  $\approx 1.8$  and  $\approx 5$  MeV (assuming a fixed well depth). The angular distributions and general trend of the cross section are well accounted for, as is the ratio of  $\sigma(\gamma, n)/\sigma(e, e'n)$  (1958BA60: 6 to 17 MeV). (1956CO56) state, on the other hand, that the 1.70-MeV peak is to be attributed to an M1 transition.

See also (1954ER1B, 1955JO1B, 1958BA60, 1958CH31) and (1956CZ1B, 1956CZ1C, 1956DE1C, 1957KO1B; theor.).

The cross section for reaction (c) is  $< 1$  mb at  $E_\gamma = 1.63$  MeV (1952AL26, 1952AL30). For reaction (d) see (1957LO1A).

11. (a)  ${}^9\text{Be}(\gamma, p){}^8\text{Li}$   $Q_m = -16.885$   
 (b)  ${}^9\text{Be}(\gamma, np){}^7\text{Li}$   $Q_m = -18.919$

The yield has a broad maximum,  $\Gamma = 4.7$  MeV, at  $E_\gamma = 22.2$  MeV where  $\sigma = 2.72$  mb (1953HA1A). The angular and energy distributions of photoprotons produced by bremsstrahlung with  $E_{\text{max}} = 25$  to 80 MeV has been studied by (1956CO59, 1956KL19). The angular distributions can be accounted for by the direct interaction of  $\gamma$ -rays with individual nucleons (1956KL19). The energy distributions indicate that transitions to levels in the residual nucleus play an important part in the direct photo effect (1956KL19). (1956CO59) suggest, on the other hand, that the relatively large proportion of low-energy protons indicates predominance of a  $(\gamma, n)$  process, followed by proton emission from  ${}^8\text{Be}$  levels at high excitation. See also (1953HE1B, 1953NA1A, 1957CH24, 1958CH31, 1958PA1B, 1958ST1A, 1958WH1A).

12. (a)  ${}^9\text{Be}(\gamma, d){}^7\text{Li}$   $Q_m = -16.693$   
 (b)  ${}^9\text{Be}(\gamma, t){}^6\text{Li}$   $Q_m = -17.687$   
 (c)  ${}^9\text{Be}(\gamma, \alpha){}^5\text{He}$   $Q_m = -2.529$

See (1955AJ61), (1956CO59, 1958WH1A) and  ${}^9\text{Be}(\gamma, n){}^8\text{Be}$ .

Table 9.2: Photoneutron cross section of  ${}^9\text{Be}$

$E_\gamma$ (MeV)	$\sigma$ (mb)	Refs.
1.69 ( ${}^{124}\text{Sb}$ )	$1.262 \pm 0.069$	(1959GI48)
1.70 (brems.) <sup>a</sup>	100	(1956CO56)
1.77 ( ${}^{56}\text{Mn}$ )	0.6	(1948RU1A: see (1953HA1B, 1956ED1B))
1.85 ( ${}^{88}\text{Y}$ )	$0.654 \pm 0.031$	(1959GI48)
2.185 ( ${}^{144}\text{Pr}$ )	0.39	(1953HA1B)
2.5 ( ${}^{140}\text{La}$ )	0.5	(1948RU1A: see (1953HA1B))
2.61 ( $\text{ThC}''$ )	$0.39 \pm 0.02$	(1956ED1B)
2.76 ( ${}^{24}\text{Na}$ )	0.7	(1948RU1A: see (1953HA1B))
	$0.674 \pm 0.05$	(1950SN67)
4.4 ( ${}^{15}\text{N}(\text{p}, \alpha){}^{12}\text{C}$ )	$0.186 \pm 0.32$	(1956ED1B)
6.2 ( ${}^{19}\text{F}(\text{p}, \alpha){}^{16}\text{O}$ )	$1.13 \pm 0.16$	(1956ED1B)
8.1 ( ${}^{13}\text{C}(\text{p}, \gamma){}^{14}\text{N}$ )	$1.38 \pm 1.16$	(1956ED1B)
10 (brems.) <sup>b</sup>	1.6	(1953NA1A)
11.3 (brems.)	4.2	(1959SP1B)
12 (brems.)	0.1	(1959SP1B)
13.3 (brems.)	3.0	(1959SP1B)
14.2 (brems.)	0.1	(1959SP1B)
15 (brems.)	0.9	(1959SP1B)
17 (brems.)	0.8	(1953NA1A)
22 (brems.) <sup>c</sup>	3.0	(1953NA1A)
25 – 320 (brems.)	3 – 2	(1953JO1B)

<sup>a</sup> Resonance width < 50 keV.

<sup>b</sup> Broad resonance:  $\Gamma \approx 8$  MeV.

<sup>c</sup> Giant resonance.

### 13. ${}^9\text{Be}(e, e'){}^9\text{Be}^*$

Elastic scattering has been studied at  $E_e = 125, 150$  and  $190$  MeV (1953HO79, 1954MC45). An r.m.s. radius of  $2.2 \pm 0.2 \times 10^{-13}$  cm is obtained (1957HO1E: see also (1956HO93)). Inelastic peaks corresponding to levels at 2.54 and 6.96 MeV are reported by (1954MC45). Comparison of  $\sigma(\gamma, n)$  and  $\sigma(e, e'n)$  for  $E_e = 6$  to 17 MeV indicates that the transitions are mainly E1 (1958BA60). See also (1957EH1A, 1958EH1A).

### 14. (a) ${}^9\text{Be}(n, n){}^9\text{Be}$

(b)  ${}^9\text{Be}(n, 2n){}^8\text{Be}$   $Q_m = -1.667$

The neutron spectrum observed when  ${}^9\text{Be}$  is bombarded with 3.7-MeV neutrons exhibits a structure which is consistent with the excitation of the known states at 0, 1.5, 2.4 and 3.1 MeV, with subsequent neutron emission from the latter two. It is concluded that the (n, 2n) process at this energy proceeds mainly via discrete states of  ${}^9\text{Be}$  (1957HU14, 1958WA05). At the same bombarding energy (1955FO1B) observe two discrete groups in coincidence, suggesting the process  ${}^9\text{Be}(n, n'){}^9\text{Be}^*(2.43) \rightarrow n + {}^8\text{Be}$ : see, however, (1957BO83:  ${}^9\text{Be}(\alpha, \alpha'){}^9\text{Be}^*$ ). Using monochromatic neutrons with  $E_n = 2.6$  to 3.25 MeV, (1957FI1B) finds a sharp threshold for (n, 2n) at  $E_n = 2.7$  MeV, corresponding to excitation of  ${}^9\text{Be}^*(2.43)$ . On the other hand, (1958MA22) find evidence for (n, 2n) at  $E_n = 2.6$  MeV, below the threshold for (n, n'); at  $E_n = 5$  to 6 MeV, the spectra show neutrons from the direct (n, 2n) process. At  $E_n = 14$  MeV, the cross section for production of  ${}^9\text{Be}^*$  is  $170 \pm 30$  mb; comparison with  $\sigma(n, 2n) = 530 \pm 40$  mb indicates that about  $\frac{1}{3}$  of the (n, 2n) processes proceed via  ${}^9\text{Be}^*(2.4)$  (1958AN32). See also (1957RO1D) and (1958BE1E; theor.).

A search for  $\gamma$ -transitions from the 1.8 and 2.4-MeV levels yields upper limits of 0.3 and 0.2 mb, respectively, for  $E_n = 2.56$  MeV and 1.8 and 0.3 mb for  $E_n = 2.74$  MeV (1956DA23).

Elastic and inelastic neutron angular distributions show forward peaking at  $E_n = 14$  MeV (1958AN32, 1958NA09): see  ${}^{10}\text{Be}$ , (1958RE1A, 1958TO1A).

### 15. ${}^9\text{Be}(p, p){}^9\text{Be}$

Elastic scattering has been studied at  $E_p = 14.5, 20$  and  $31.5$  MeV by (1956KI54), at 10 MeV by (1956RA32), at 12 MeV by (1958SU14), at 17 MeV by (1956DA03), at 31 MeV by (1953WR1A, 1954FI1A, 1956BE14): see also  ${}^{10}\text{B}$ . All angular distributions show pronounced diffraction maxima characteristic of the optical model. Analysis in terms of the diffuse-surface optical model is discussed by (1957ME21). See also (1956KL55, 1956SH1C).

Inelastic scattering is observed corresponding to levels at (1.8), 2.43, 3.1, 5.0, 6.8, 7.9, 11.3, (14.5), (17.5), (19.9) and (21.7) MeV: see Table 9.3. It is not clear whether the structure observed near 1.8 MeV is properly to be attributed to a level at this energy or to a three-body breakup,

Table 9.3: Levels of  ${}^9\text{Be}$  from  ${}^9\text{Be}(p, p')$ ,  $(d, d')$  and  $(\alpha, \alpha')$ 

$E_x$ (MeV)	$\Gamma$ (keV)	$J^\pi$
0		
(1.8)	150 <sup>a</sup>	$(\frac{1}{2}^+)$ <sup>a</sup>
$2.433 \pm 0.003$	$\leq 1$ <sup>a</sup>	$\frac{1}{2}^-, \frac{5}{2}^-$
3.03	$> 280$ <sup>e</sup>	$\leq \frac{3}{2}$
$5.0 \pm 0.3$ <sup>b</sup>		
$6.76 \pm 0.06$ <sup>c</sup>		$\frac{1}{2}^-, \frac{5}{2}^-, \frac{7}{2}^-$
$7.94 \pm 0.08$ <sup>b</sup>		$\frac{3}{2}^+, \frac{5}{2}^+, \frac{9}{2}^+, \frac{11}{2}^+$
$11.3 \pm 0.2$ <sup>d</sup>		
(14.5) <sup>b</sup>		
(17.5) <sup>b</sup>		
$19.9 \pm 0.1$ <sup>b</sup>		$\frac{1}{2}^-, \frac{3}{2}^-, \frac{5}{2}^-$
$21.7 \pm 0.1$ <sup>b</sup>		$\frac{1}{2}^-, \frac{3}{2}^-, \frac{5}{2}^-$ or $\frac{1}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$

<sup>a</sup> (1955GO48, 1957BO83, 1958SU14).

<sup>b</sup> (1956BE14).

<sup>c</sup> This level may be double: see (1952BR52, 1956BE14, 1956ST30, 1958SU14, 1958TY46).

<sup>d</sup> (1952BR52, 1956BE14, 1956ST30).

<sup>e</sup> See  ${}^{11}\text{B}(d, \alpha){}^9\text{Be}$  (1958KA31).

modified by the proximity of the neutron threshold. If the former is correct, the level width is 0.15 MeV and  $J = \frac{1}{2}^+$  (1955GO48, 1958SU14). If a three-body breakup is involved, some interaction between the neutron and the  ${}^8\text{Be}$  must be involved: (1956BO18) find a satisfactory fit to the observed shape with an s-wave scattering length of  $20 \times 10^{-13}$  cm; see also (1958MI1C) and  ${}^{11}\text{B}(d, \alpha){}^9\text{Be}$ . It is suggested by (1956SU67, 1958SU14) that the observed structure is to be understood in terms of a heavy particle stripping process and it is conjectured that the 4.8-MeV level may reflect the same effect, involving  ${}^8\text{Be}^*(2.9)$ ; see, however, (1958MI1C).

A continuous distribution of neutrons, observed for  $E_p > 4.5$  MeV is attributed to formation of  ${}^9\text{Be}^*(2.4)$  with subsequent breakup into  $n + {}^8\text{Be}^*(2.9)$  (1959MA20). Breakup into  $({}^8\text{Be} + n)$  occurs in  $12 \pm 5$  % of transitions (J.B. Marion, private communication).

The energy of the 2.4-MeV level is given as  $2433 \pm 5$  (1951BR72),  $2434 \pm 5$  (1956BO18),  $2432 \pm 4$  keV (1955GO48); the width is  $\leq 1$  keV (1955GO48). Analysis of angular distributions at  $E_p = 12$  MeV in terms of direct interaction plus compound nucleus formation indicate  $J = \frac{1}{2}^-$  or  $\frac{5}{2}^-$  (1958SU14). At  $E_p = 31$  MeV, analysis indicates  $J = \frac{1}{2}^+, \frac{3}{2}^+$  or  $\frac{5}{2}^+$  (1956BE14: see, however, (1958SU14)). Results at  $E_p = 10$  MeV are consistent with odd parity (1956RA32). See

also  $^{10}\text{B}(n, d)^9\text{Be}$  and (1952DA1B).

The  $3.03 \pm 0.03$  MeV level has a width  $> 0.28$  MeV; the Wigner limit then restricts  $l_n$  to 0 or 1,  $J \leq \frac{3}{2}$  (1956BO18: see, however,  $^{11}\text{B}(d, \alpha)^9\text{Be}$ ). Angular distributions for the higher levels have been studied by (1956BE14); the resulting spin assignments are given in Table 9.3. See also (1954FI1A, 1955GR12, 1958TY46) and (1956BL1B; theor.).

#### 16. $^9\text{Be}(d, d')^9\text{Be}^*$

Elastic scattering angular distributions have been studied at  $E_d = 24$  MeV by (1958SU14). See also (1947GU1A, 1952EL01) and  $^{11}\text{B}$ .

Inelastic groups are reported corresponding to levels at (1.7), 2.4,  $3.01 \pm 0.1$  ( $\Gamma \approx 0.3$ ), 4.8 and 6.8 MeV (1955RA41, 1956GR37, 1958MI1C, 1958SU14). The structure at  $Q = -1.7$  MeV may be accounted for as a three-body breakup with an s-wave ( $n$ - $^8\text{Be}$ ) interaction characterized by a scattering length of  $20 \times 10^{-13}$  cm, or by a resonance of about single-particle width. In either case a  $J = \frac{1}{2}^+$  level of  $^9\text{Be}$  is indicated near the binding energy (1955RA41, 1958MI1C: see also  $^{11}\text{B}(d, \alpha)^9\text{Be}$ ). The angular distribution of the  $Q = -2.43$  MeV group has been studied at  $E_d = 15$  MeV (1956HA90) and  $E_d = 24$  MeV (1958SU14). Analysis by direct interaction theory yields  $l = 2$ ,  $J = \frac{1}{2}^-$ ,  $\frac{5}{2}^-$  or  $\frac{7}{2}^-$  (1958SU14: see, however, (1956HA90)). See also (1956SU1A).

#### 17. $^9\text{Be}(\alpha, \alpha')^9\text{Be}^*$

Elastic scattering has been studied at  $E_\alpha = 48$  MeV by (1958SU14).

Inelastic groups are observed corresponding to  $^9\text{Be}^* = (1.8), 2.4, (3.1), 6.8$  and 11.3 MeV (1955RA41, 1956FA02, 1958SU14). The group corresponding to the 1.8-MeV “level” has a peak at  $Q = -1.83 \pm 0.03$  MeV,  $\Gamma \approx 200$  keV. It is suggested, however, that just such a structure would be expected from a heavy particle stripping process in which the proton also escapes, preferentially with the maximum possible energy. In this case, the 4.8-MeV “level” might arise from a similar process, in which the final nucleus is now  $^8\text{Be}^*(2.9)$ . The angular distribution of the  $Q = -2.4$  MeV group at  $E_\alpha = 48$  MeV indicates  $l = 2$ ,  $J = \frac{1}{2}^-$ ,  $\frac{5}{2}^-$  or  $\frac{7}{2}^-$  (1958SU14). Measurement of the momentum and angular distributions of  $\alpha$ -particles from the breakup of  $^9\text{Be}^*(2.4)$  indicate that the decay proceeds mainly via  $^4\text{He} + ^5\text{He}$ , or by direct three-body breakup. Gamma decay is  $< 1\%$ , neutron emission to  $^8\text{Be}(0)$  is  $< 10\%$  (1957BO83). See also (1956BL1B, 1958PI1B; theor.).

18. (a)  $^9\text{Be}(p, d)^8\text{Be}$   $Q_m = 0.560$

(b)  $^9\text{Be}(d, t)^8\text{Be}$   $Q_m = 4.592$

At  $E_p = 95$  MeV, two groups of deuterons are observed from reaction (a), corresponding to  $^8\text{Be}_{g.s.}$  and to states near 17 MeV. It is suggested that these reflect the “snatching” of the loosely

bound neutron or of one of the tightly bound “alpha-particle” neutrons, respectively. Angular distributions lend some support to the  $\alpha$ -particle model of  ${}^9\text{Be}$ ; the occurrence of high momenta for the tightly bound neutrons indicates the operation of strong short-range two-body forces (1956SE1A).

Angular distributions at  $E_p = 6.5$  and  $22$  MeV, and at  $E_d = 7.7$  MeV are analyzed in terms of pickup theory, using a square-well ( $n$ - ${}^8\text{Be}$ ) interaction to represent the ground state of  ${}^9\text{Be}$  (1955DA1D, 1955DA1E). See also  ${}^{10}\text{B}$ ,  ${}^{11}\text{B}$ .

$$19. \text{ (a) } {}^{10}\text{B}(\gamma, p){}^9\text{Be} \quad Q_m = -6.585$$

$$\text{ (b) } {}^{10}\text{B}(\gamma, p){}^8\text{Be} + n \quad Q_m = -8.251$$

For reaction (a) see (1956GO1F, 1956GO1G). For reaction (b) see  ${}^8\text{Be}$  and  ${}^{10}\text{B}$ .

$$20. {}^{10}\text{B}(n, d){}^9\text{Be} \quad Q_m = -4.358$$

At  $E_n = 14$  MeV, the ground state and the level at  $2.43$  MeV are observed. No other deuteron groups were detected below  $E_x \approx 5.5$  MeV. The angular distribution of the deuterons, analyzed by pickup theory, indicate odd parity,  $\frac{1}{2} < J \leq \frac{9}{2}$  for both states (1954RI15). See also (1956FR18), (1955FR1F; theor.) and  ${}^{11}\text{B}$ .

$$21. {}^{10}\text{B}(d, {}^3\text{He}){}^9\text{Be} \quad Q_m = -1.091$$

Not observed.

$$22. {}^{10}\text{B}(t, \alpha){}^9\text{Be} \quad Q_m = 13.228$$

At  $E_t = 1$  MeV,  $\alpha$ -groups are observed corresponding to  ${}^9\text{Be}$  levels at  $2.39$  and  $3.06$  MeV (1955AL57).

$$23. {}^{11}\text{B}(n, t){}^9\text{Be} \quad Q_m = -9.564$$

Not observed.

$$24. {}^{11}\text{B}(p, {}^3\text{He}){}^9\text{Be} \quad Q_m = -10.329$$

Not observed.

$$25. \text{}^{11}\text{B}(\text{d}, \alpha)^9\text{Be} \quad Q_m = 8.022$$
$$Q_0 = 8.015 \pm 0.010 \text{ MeV (1956BO18)}.$$

Alpha-particle groups are reported to states at (1.75), 2.4 and 3.0 MeV. The structure corresponding to the 1.75-MeV state can be explained in terms of an ( $n$ - $^8\text{Be}$ ) interaction with a scattering length  $a$  of  $20 \times 10^{-13}$  cm or in terms of a resonance near threshold, with  $\theta^2 \approx 1$  (1956BO18, 1958MI1C). (1958KA31) find, on the other hand, that a value of  $a \geq 80 \times 10^{-13}$  cm is required to fit their distribution and suggest that the ( $n$ - $^8\text{Be}$ ) interaction must be very near resonance on this model. The energy of the 2.4-MeV state is given as  $2422 \pm 5$  keV by (1951VA08),  $2431 \pm 6$  keV (1954EL10),  $2424 \pm 5$  keV (1956BO18). The next state is located at  $3.02 \pm 0.03$  (1955LE36),  $3.05 \pm 0.03$  MeV (1956BO18). The width is  $\approx 0.3$  MeV (1956BO18),  $\Gamma_{\text{c.m.}} = 161 \pm 15$  keV (1958KA31). See also (1955HO48).

$$26. \text{}^{12}\text{C}(\text{n}, \alpha)^9\text{Be} \quad Q_m = -5.709$$

At  $E_n = 14$  MeV, the cross section to the ground state of  $^9\text{Be}$  is  $80 \pm 20$  mb; that to the 2.43-MeV level,  $\sigma_{2.43}$ , is  $10 \leq \sigma_{2.43} \leq 210$  mb (1955GR21).

$$27. \text{}^{13}\text{C}(\gamma, \alpha)^9\text{Be} \quad Q_m = -10.654$$

See (1953MI31).

**<sup>9</sup>B**  
(Fig. 11)

1.  ${}^6\text{Li}({}^3\text{He}, \text{p}){}^8\text{Be}$   $Q_m = 16.786$   $E_b = 16.598$

The excitation functions for protons leading to the ground and 2.9-MeV excited states of  ${}^8\text{Be}$  have been measured for  $E({}^3\text{He}) = 0.9$  to  $5.1$  MeV ( $\theta = 0^\circ$  and  $150^\circ$ , lab.). Resonances are observed at  $E({}^3\text{He}) = 1.6$  MeV ( $\Gamma = 0.25$  MeV,  ${}^9\text{B}^* = 17.6$  MeV) and  $3.0$  MeV ( $\Gamma = 1.5$  MeV,  ${}^9\text{B}^* = 18.6$  MeV) (1956SC01). However, J.W. Butler (private communication) suggests that the decrease in the yield for  $E({}^3\text{He}) \gtrsim 3$  MeV can be accounted for by competition from the  ${}^6\text{Li}({}^3\text{He}, \text{n}){}^8\text{B}$  reaction with a threshold at  $2.966$  MeV. Angular distributions, taken at 6 energies, are isotropic at  $\approx 1$  MeV and become increasingly asymmetric at higher energies. In particular, the ground state protons exhibit strong backward peaking for  $E({}^3\text{He}) \gtrsim 3$  MeV (1956SC01).

2.  ${}^6\text{Li}(\alpha, \text{n}){}^9\text{B}$   $Q_m = -3.979$

See (1956RO06).

3.  ${}^7\text{Li}({}^3\text{He}, \text{n}){}^9\text{B}$   $Q_m = 9.346$

Not reported.

4.  ${}^9\text{Be}(\text{p}, \text{n}){}^9\text{B}$   $Q_m = -1.854$

The width of the ground state is  $< 2$  keV (1951ST76). At  $E_p = 6.59$  MeV, a neutron group is observed corresponding to a level at  $2.37 \pm 0.04$  MeV. A continuous distribution of neutrons, attributed to the  $(\text{p}, \text{p}'\text{n})$  reaction, is also observed (1953AJ09, 1959MA20: see also (1950JO1B) and  ${}^9\text{Be}$ ). For  $E_p = 2$  to  $5.8$  MeV, two neutron thresholds have been observed at  $E_p = 2.060 \pm 0.003$  and  $4.645 \pm 0.005$  MeV ( ${}^9\text{B}^* = 2.326 \pm 0.006$  MeV +  $\frac{1}{2}\Gamma$ ). The continued slow neutron yield above threshold suggests  $l = 1$  neutron emission (1955MA84). A broad threshold reported at  $E_p = 3.6$  MeV appears to have been due to variation in counter sensitivity. It is possible, however, that part of the continuum neutron spectrum arises from a  ${}^9\text{B}$  level expected near  $1.4$  MeV (1959MA20). See also (1957KI1B, 1958BO63).

Table 9.4: Energy levels of  ${}^9\text{B}$

$E_x$ in ${}^9\text{B}$ (MeV)	$J^\pi$	$\Gamma$ (keV)	Decay	Reactions
0	$> \frac{1}{2}^-$	$< 2$	(p, $\alpha$ )	4, 5, 6, 7, 8, 9, 11
$2.37 \pm 0.02$	$> \frac{1}{2}^-$	80	(p, $\alpha$ )	4, 7, 9, 11
$2.83 \pm 0.03$		300		9
(7.0)				11
17.6		170	p, ${}^3\text{He}$ , $\alpha$	1
(18.6)		1000	p, ${}^3\text{He}$ , $\alpha$	1

5.  ${}^9\text{Be}({}^3\text{He}, \text{t}){}^9\text{B}$   $Q_m = -1.089$

The ground state angular distributions are peaked in the forward direction at  $E({}^3\text{He}) = 5.7$  MeV (S. Hinds and R. Middleton) and 21 MeV (1958WE1E).

6.  ${}^{10}\text{B}(\gamma, \text{n}){}^9\text{B}$   $Q_m = -8.439$

See (1951SH63).

7.  ${}^{10}\text{B}(\text{p}, \text{d}){}^9\text{B}$   $Q_m = -6.212$

At  $E_p = 18.9$  MeV, deuteron groups are observed corresponding to the ground state of  ${}^9\text{B}$  and to an excited state at  $2.40 \pm 0.15$  MeV. The angular distributions to both states, analyzed by pickup theory, show an  $l = 1$  transfer, indicating odd parity,  $\frac{1}{2} < J \leq \frac{9}{2}$ , for these states (1956RE04). The ratio of cross sections is in good agreement with an intermediate-coupling calculation (1955FR1F).

8.  ${}^{10}\text{B}(\text{d}, \text{t}){}^9\text{B}$   $Q_m = -2.180$   
 $Q_0 = -2.187 \pm 0.010$  (1956BO18).

9.  ${}^{10}\text{B}({}^3\text{He}, \alpha){}^9\text{B}$   $Q_m = 12.139$

At  $E(^3\text{He}) = 1$  MeV, alpha-particle groups have been observed corresponding to the ground state of  $^9\text{B}$  and to an excited state at  $2.58 \pm 0.13$  MeV (1955AL57). At  $E(^3\text{He}) = 2$  to 3 MeV, groups are observed corresponding to the ground state and to levels at  $E_x = 2.37 \pm 0.02$  MeV and  $2.83 \pm 0.03$  MeV. The widths are 80 keV and 300 keV for  $^9\text{B}^*(2.37)$  and  $^9\text{B}^*(2.83)$ , respectively. There is no evidence of a well-defined state corresponding to  $^9\text{Be}^*(1.75)$  (1958PO61, 1959PO61). See also (1959SP17).

10.  $^{11}\text{B}(\text{p}, \text{t})^9\text{B}$   $Q_m = -11.418$

Not observed.

11.  $^{12}\text{C}(\text{p}, \alpha)^9\text{B}$   $Q_m = -7.563$   
 $Q_0 = -7.58 \pm 0.1$  (1955RE16).

At  $E_p = 18.8$  MeV, alpha-particle groups have been observed to the ground state of  $^9\text{B}$  and to an excited state at  $2.39 \pm 0.08$  MeV. The two  $\alpha$ -groups are superimposed on a background possibly due to  $^{12}\text{C}(\text{p}, \text{p}')^{12}\text{C}^* \rightarrow 3\alpha$ . The average differential cross section for the ground state group at  $E_p \approx 18$  MeV is  $3 \pm 1$  mb/sr. No other groups were observed up to  $E_x \approx 7.9$  MeV (1955RE16). The angular distribution of the ground state group at  $E_p = 15.6$  to 18.6 MeV is consistent with triton pickup or knock-on theories, except for discrepancies at low angles (1958MA40). At  $E_p = 29$  MeV, the reaction appears to go to the ground state and to a level at  $3.2 \pm 1.0$  MeV (1955NE18: stars in a cloud chamber;  $^{12}\text{C} + \text{p} \rightarrow \alpha + ^9\text{B} \rightarrow 3\alpha + \text{p}$ ). At  $E_p = 32$  MeV, groups corresponding to  $^9\text{B}(0)$ ,  $^9\text{B}^*(2.4)$  and a new level at 7.0 MeV are observed (1958KN52).

### $^9\text{C}$

(Not illustrated)

Comparison with the mass of  $^9\text{Li}$  leads to an estimated mass excess of  $32.3 \pm 2$  MeV (1955AJ61). Analysis of a single star attributed to  $\beta$ -decay of  $^9\text{C}$  and subsequent breakup into  $\text{p} + 2\alpha$  yields  $Q > 15.4$  MeV, mass excess  $> 30.2$  MeV (1956SW77). Stability against  $^8\text{B} + \text{p}$  requires a mass excess  $< 32.9$  MeV. Two reactions leading to  $^9\text{C}$  which have not been reported are  $^7\text{Be}(^3\text{He}, \text{n})^9\text{C}$  ( $Q_m = -7$ ) and  $^{12}\text{C}(\gamma, 3\text{n})^9\text{C}$  ( $Q_m = -54$ ).

## References

(Closed December 01, 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1947GU1A Guggenheimer, Heitler and Powell, Proc. Roy. Soc. A190 (1947) 196
- 1948RU1A Russell, Sachs, Wattenberg and Fields, Phys. Rev. 73 (1948) 545
- 1949GU1A Guth and Mullin, Phys. Rev. 76 (1949) 234
- 1949HA1A Hamermesh, Hamermesh and Wattenberg, Phys. Rev. 76 (1949) 611
- 1950JO1B Johnson, Ajzenberg and Laubenstein, Phys. Rev. 79 (1950) 187
- 1950SN67 A.H. Snell, E.C. Barker and R.L. Sternberg, Phys. Rev. 80 (1950) 637
- 1951BR72 C.P. Browne, R.M. Williamson, D.S. Craig and D.J. Donahue, Phys. Rev. 83 (1951) 179
- 1951GA30 W.L. Gardner, N. Knable and B.J. Moyer, Phys. Rev. 83 (1951) 1054
- 1951SH63 R. Sher, J. Halpern and A.K. Mann, Phys. Rev. 84 (1951) 387
- 1951ST76 P.H. Stelson and W.M. Preston, Phys. Rev. 83 (1951) 469
- 1951VA08 D.M. Van Patter, A. Sperduto, K. Huang, E.N. Strait and W.W. Buechner, Phys. Rev. 81 (1951) 233
- 1952AL26 D.E. Alburger, Phys. Rev. 88 (1952) 1257
- 1952AL30 D.E. Alburger, Physica 18 (1952) 1034
- 1952BA1A Baggett and Bame, Phys. Rev. 85 (1952) 434
- 1952BR52 R. Britten, Phys. Rev. 88 (1952) 283
- 1952DA1B Davis, Phys. Rev. 86 (1952) 976
- 1952EL01 F.A. El-Bedewi, Proc. Phys. Soc. (London) A65 (1952) 64
- 1952HO25 R.B. Holt, R.N. Thorn and R.W. Waniek, Phys. Rev. 87 (1952) 378
- 1952PE02 T.P. Pepper, K.W. Allen, E. Almqvist and J.T. Dewan, Phys. Rev. 85 (1952) 155
- 1952SH44 R.K. Sheline, Phys. Rev. 87 (1952) 557
- 1953AJ09 F. Ajzenberg and W.W. Buechner, Phys. Rev. 91 (1953) 674
- 1953FR03 W.F. Fry, Phys. Rev. 89 (1953) 325
- 1953HA1A Haslam, Katz, Crosby, Summers-Gill and Cameron, Can. J. Phys. 31 (1953) 210

1953HA1B Hamermesh and Kimball, Phys. Rev. 90 (1953) 1063  
1953HE1B Hendel, Z. F. Phys. 135 (1953) 168  
1953HO79 R. Hofstadter, H.R. Fechter and J.A. McIntyre, Phys. Rev. 92 (1953) 978  
1953JO1B Jones and Terwilliger, Phys. Rev. 91 (1953) 699  
1953MI31 C.H. Millar and A.G.W. Cameron, Can. J. Phys. 31 (1953) 723  
1953NA1A Nathans and Halpern, Phys. Rev. 92 (1953) 940  
1953RE19 D. Reagan, Phys. Rev. 92 (1953) 651  
1953SA1A Sawyer and Phillips, Los Alamos Rept.1578 (1953)  
1953WR1A Wright, UCRL 2422 (1953)  
1954BA46 S. Bashkin, Phys. Rev. 95 (1954) 1012  
1954CA1A Carver, Knodaiah and McDaniel, Phil. Mag. 45 (1954) 948  
1954EL10 R.B. Elliott and D.J. Livesey, Proc. Roy. Soc. A224 (1954) 129  
1954ER1B Erikson and Zaleski, J. Rad. Phys. 15 (1954) 492  
1954FI1A Finke, UCRL 2789 (1954)  
1954MC45 J.A. McIntyre, B. Hahn and R. Hofstadter, Phys. Rev. 94 (1954) 1084  
1954MO92 C.D. Moak, W.M. Good and W.E. Kunz, Phys. Rev. 96 (1954) 1363  
1954NI1B Niewodniczanski and Wielowiejska, Bull. Acad. Polon. Sci. 2 (1954) 25  
1954RI15 F.L. Ribe and J.D. Seagrave, Phys. Rev. 94 (1954) 934  
1955AJ61 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 27 (1955) 77  
1955AL57 E. Almqvist, K.W. Allen and C.B. Bigham, Phys. Rev. 99 (1955) 631A  
1955BE1E Bendt and Scott, Phys. Rev. 97 (1955) 744  
1955DA1D Dabrowski and Sawicki, Acta Phys. Pol. 14 (1955) 143  
1955DA1E Dabrowski and Sawicki, Acta Phys. Pol. 14 (1955) 407; Erratum Acta Phys. Pol. 15 (1956) 431  
1955FO1B Fowler, Hanna and Owen, Phys. Rev. 98 (1955) 249A  
1955FR1F French, Halbert and Pandya, Phys. Rev. 99 (1955) 1387  
1955GO48 C.R. Gossett, G.C. Phillips, J.P. Schiffer and P.M. Windham, Phys. Rev. 100 (1955) 203  
1955GR12 G.W. Greenlees, Proc. Phys. Soc. (London) A68 (1955) 97  
1955GR21 E.R. Graves and R.W. Davis, Phys. Rev. 97 (1955) 1205  
1955HO48 R.E. Holland, D.R. Inglis, R.E. Malm and F.P. Mooring, Phys. Rev. 99 (1955) 92  
1955JO1B Johansson, Phys. Rev. 97 (1955) 434

1955LE36 L.L. Lee, Jr. and D.R. Inglis, Phys. Rev. 99 (1955) 96  
 1955MA20 R.L. Macklin and H.E. Banta, Phys. Rev. 97 (1955) 753  
 1955MA84 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 91  
 1955NE18 J.L. Need, Phys. Rev. 99 (1955) 1356  
 1955RA14 D.C. Ralph and F.E. Dunnam, Phys. Rev. 98 (1955) 249A  
 1955RA41 V.K. Rasmussen, D.W. Miller, M.B. Sampson and U.C. Gupta, Phys. Rev. 100 (1955) 851  
 1955RE16 J.B. Reynolds, Phys. Rev. 98 (1955) 1289  
 1956BE14 J. Benveniste, R.G. Finke and E.A. Martinelli, Phys. Rev. 101 (1956) 655  
 1956BE1A Bezrukov, Panov and Timoshuk, Sov. J. Nucl. Energy 4 (1956) 609  
 1956BL1B Blair and Henley, Bull. Amer. Phys. Soc. 1 (1956) 20; Physica 22 (1956) 1126A  
 1956BO18 C.K. Bockelman, A. Leveque and W.W. Buechner, Phys. Rev. 104 (1956) 456  
 1956CA1B Catala, Aguilar, Senent and Font, An. Real. Soc. Espan. Fis. Y. Quim. A52 (1956) 33  
 1956CO56 D.R. Connors and W.C. Miller, Bull. Amer. Phys. Soc. 1 (1956) 340, P12  
 1956CO59 L. Cohen, A.K. Mann, B.J. Patton, K. Reibel, W.E. Stephens and E.J. Winhold, Phys. Rev. 104 (1956) 108  
 1956CZ1B Czyz, Phys. Rev. 102 (1956) 1185  
 1956CZ1C Czyz and Sawicki, Nuovo Cim. 3 (1956) 864; Physica 22 (1956) 1182A  
 1956DA03 I.E. Dayton and G. Schrank, Phys. Rev. 101 (1956) 1358  
 1956DA23 R.B. Day, Phys. Rev. 102 (1956) 767  
 1956DE1C De Saussure and Guth, Bull. Amer. Phys. Soc. 1 (1956) 21  
 1956ED1B Edge, Nucl. Phys. 2 (1956) 485  
 1956FA02 G.W. Farwell and D.D. Kerlee, Bull. Amer. Phys. Soc. 1 (1956) 20, DA5  
 1956FA1B Fabricand, Allison and Halpern, Phys. Rev. 103 (1956) 1755  
 1956FL1A Flynn, Glendenin and Steinberg, Phys. Rev. 101 (1956) 1492  
 1956FR18 G.M. Frye, Jr. and J.H. Gammel, Phys. Rev. 103 (1956) 328  
 1956GO1F Goldanskii, Physica 22 (1956) 1143A  
 1956GO1G Goldanskii, Zh. Eksp. Teor. Fiz. 30 (1956) 969; JETP (Sov. Phys.) 3 (1956) 791  
 1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28  
 1956HA90 J.W. Haffner, Phys. Rev. 103 (1956) 1398  
 1956HO93 R. Hofstadter, Revs. Mod. Phys. 28 (1956) 214  
 1956KI54 B.B. Kinsey and T. Stone, Phys. Rev. 103 (1956) 975

- 1956KL19 G.K. Kliger, V.I. Riabinkin, I.V. Chuvilo and V.S. Shevchenko, *Physica* 22 (1956) 1142A
- 1956KL55 A.P. Kliucharev, L.I. Bolotin and V.A. Lutsik, *Zh. Eksp. Teor. Fiz.* 30 (1956) 573; *JETP (Sov. Phys.)* 3 (1956) 463
- 1956KU1A Kurath, *Phys. Rev.* 101 (1956) 216
- 1956MA1M Mast and Mullin, *Bull. Amer. Phys. Soc.* 1 (1956) 180
- 1956RA32 S.W. Rasmussen, *Phys. Rev.* 103 (1956) 186
- 1956RE04 J.B. Reynolds and K.G. Standing, *Phys. Rev.* 101 (1956) 158
- 1956RI37 A.C. Riviere, *Nucl. Phys.* 2 (1956) 81
- 1956RO06 A.B. Robbins, *Phys. Rev.* 101 (1956) 1373
- 1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser, Jr., *Phys. Rev.* 104 (1956) 1064
- 1956SE1A Selove, *Phys. Rev.* 101 (1956) 231
- 1956SH1C Shaw, Conzett, Slobodrian and Summers-Gill, *Bull. Amer. Phys. Soc.* 1 (1956) 253
- 1956ST30 K. Strauch and F. Titus, *Phys. Rev.* 104 (1956) 191
- 1956SU1A Summers-Gill, UCRL 3388 (1956)
- 1956SU67 R.G. Summers-Gill, *Bull. Amer. Phys. Soc.* 1 (1956) 253, F11
- 1956SW77 M.S. Swami, J. Schneps and W.F. Fry, *Phys. Rev.* 103 (1956) 1134
- 1956WA29 H.J. Watters, *Phys. Rev.* 103 (1956) 1763
- 1957BA1H Barker, *Phil. Mag.* 2 (1957) 780
- 1957BO83 D. Bodansky, S.F. Eccles and I. Halpern, *Phys. Rev.* 108 (1957) 1019
- 1957CH24 I.V. Chuvilo and V.G. Shevchenko, *Zh. Eksp. Teor. Fiz.* 32 (1957) 1335; *JETP (Sov. Phys.)* 5 (1957) 1090
- 1957EH1A Ehrenberg, Meyer-Berkhout, Hofstadter, Ravenhall and Sobottka, *Bull. Amer. Phys. Soc.* 2 (1957) 390
- 1957FA10 F.J.M. Farley and R.E. White, *Nucl. Phys.* 3 (1957) 561; *Erratum Nucl. Phys.* 3 (1957) 692
- 1957FI1B Fischer, *Phys. Rev.* 108 (1957) 99
- 1957HO1E Hofstadter, *Ann. Rev. Nucl. Sci.*, Vol. 7 (1957)
- 1957HU14 P. Huber and R. Wagner, *Helv. Phys. Acta* 30 (1957) 257
- 1957JA37 N. Jarmie, J.D. Seagrave et al, LA-2014 (1957)
- 1957KI1B Kiselev and Fliagin, *Zh. Eksp. Teor. Fiz.* 32 (1957) 962; *JETP (Sov. Phys.)* 5 (1957) 786
- 1957KO1B Kopaleishvili, *Zh. Eksp. Teor. Fiz.* 32 (1957) 1249; *JETP (Sov. Phys.)* 5 (1958) 1018

1957LO1A Lokan, Proc. Phys. Soc. (London) A70 (1957) 836  
1957ME21 M.A. Melkanoff, J.S. Nodvik, D.S. Saxon and R.D. Woods, Phys. Rev. 106 (1957) 793  
1957PA1A Pal and Mukherjee, Phys. Rev. 106 (1957) 811  
1957RI39 A.C. Riviere and P.B. Treacy, Aust. J. Phys. 10 (1957) 209  
1957RO1D Rosen and Stewart, Phys. Rev. 107 (1957) 824  
1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809  
1958AN32 J.D. Anderson, C.C. Gardner, J.W. McClure, M.P. Nakada and C. Wong, Phys. Rev. 111 (1958) 572  
1958BA60 W.C. Barber, Phys. Rev. 111 (1958) 1642  
1958BE1E Berlin and Owen, Nucl. Phys. 5 (1958) 669  
1958BO63 H.A. Bostick, H.E. Adelson, B.J. Moyer and C.N. Waddel, Bull. Amer. Phys. Soc. 3 (1958) 189, L2  
1958CH31 I.V. Chuvilo and V.G. Shevchenko, Zh. Eksp. Teor. Fiz. 34 (1958) 593; JETP (Sov. Phys.) 7 (1958) 410  
1958EH1A Ehrenberg and Hofstadter, HEPL 129 (1958)  
1958KA31 R.W. Kavanagh and C.A. Barnes, Phys. Rev. 112 (1958) 503  
1958KN52 H.B. Knowles, Bull. Amer. Phys. Soc. 3 (1958) 330, N2  
1958KU1B Kunz and Henley, Bull. Amer. Phys. Soc. 3 (1958) 323  
1958MA22 J.B. Marion, J.S. Levin and L. Cranberg, Bull. Amer. Phys. Soc. 3 (1958) 165, B5  
1958MA40 D.R. Maxson and J.H. Terrell, Bull. Amer. Phys. Soc. 3 (1958) 381, T4  
1958MI1C Miller, Phys. Rev. 109 (1958) 1669  
1958MO99 C.D. Moak, A. Galonsky, R.L. Traugber and C.M. Jones, Phys. Rev. 110 (1958) 1369  
1958NA09 M.P. Nakada, J.D. Anderson, C.C. Gardner and C. Wong, Phys. Rev. 110 (1958) 1439  
1958PA1B Palfrey and Tautfest, Bull. Amer. Phys. Soc. 3 (1958) 173  
1958PI1B Pinkston, Bull. Amer. Phys. Soc. II 3 (1958) 223  
1958PO61 B. Povh, Bull. Amer. Phys. Soc. 3 (1958) 406, F4  
1958RE1A Remy and Winter, Nuovo Cim. 9 (1958) 664  
1958ST1A Stewart, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)  
1958SU14 R.G. Summers-Gill, Phys. Rev. 109 (1958) 1591  
1958TA04 G.W. Tautfest, Phys. Rev. 110 (1958) 708  
1958TO1A Toriki, Winter and Remy, Compt. Rend. 246 (1958) 3047

- 1958TY46 H. Tyren and T.A.J. Maris, Nucl. Phys. 6 (1958) 82
- 1958WA05 R. Wagner and P. Huber, Helv. Phys. Acta 31 (1958) 89
- 1958WE1E Wegner and Hall, Bull. Amer. Phys. Soc. 3 (1958) 338
- 1958WH1A Whitehead, Murray, Aitken, Middlemas and Collie, Phys. Rev. 110 (1958) 941
- 1959GI48 J.H. Gibbons, R.L. Macklin, J.B. Marion and H.W. Schmitt, Phys. Rev. 114 (1959) 1319
- 1959MA20 J.B. Marion and J.S. Levin, Phys. Rev. 115 (1959) 144
- 1959PO61 B. Povh, Phys. Rev. 114 (1959) 1114
- 1959SP17 R.R. Spencer and G.C. Phillips, Bull. Amer. Phys. Soc. 4 (1959) 95, C4 (1959); Oral Report
- 1959SP1B Spicer, (1959)