

# Energy Levels of Light Nuclei $A = 9$

F. Ajzenberg-Selove

*University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

**Abstract:** An evaluation of  $A = 5-10$  was published in *Nuclear Physics A320* (1979), p. 1. This version of  $A = 9$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed 1978)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 9$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides:  [\${}^9\text{n}\$](#) ,  [\${}^9\text{He}\$](#) ,  [\${}^9\text{Li}\$](#) ,  [\${}^9\text{Be}\$](#) ,  [\${}^9\text{B}\$](#) ,  [\${}^9\text{C}\$](#) ,  [\${}^9\text{N}\$](#)

B. Tables of Recommended Level Energies:

[Table 9.1](#): Energy levels of  ${}^9\text{Li}$

[Table 9.2](#): Energy levels of  ${}^9\text{Be}$

[Table 9.9](#): Energy levels of  ${}^9\text{B}$

[Table 9.12](#): Energy levels of  ${}^9\text{C}$

C. [References](#)

D. Figures:  [\${}^9\text{Li}\$](#) ,  [\${}^9\text{Be}\$](#) ,  [\${}^9\text{B}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

**<sup>9</sup>n**  
(Not illustrated)

See (1977DE08).

**<sup>9</sup>He**  
(Not illustrated)

<sup>9</sup>He has not been observed: see (1974AJ01). It is predicted to be particle unstable. Particle instability with respect to <sup>8</sup>He + n, <sup>7</sup>He + 2n and <sup>6</sup>He + 3n implies atomic mass excesses greater than 39.667, 42.253 and 41.808 MeV, respectively. The calculated mass excess of <sup>9</sup>He is 43.49 MeV based on the modified form of the mass equation (1975JE02). See also (1974TH01) and (1974IR04, 1975BE31, 1976IR1B; theor.).

**<sup>9</sup>Li**  
(Figs. 15 and 18)

GENERAL: (See also (1974AJ01).)

*Model calculations:* (1974IR04, 1976IR1B, 1977JA14).

*Special reactions:* (1975AB1D, 1975ZE01, 1976AL1F, 1976BE67, 1976BU16, 1977YA1B).

*Pion and kaon reactions (See also reaction 3.):* (1973CA1C, 1976TR1A, 1977BA1Q, 1977DO06, 1977SH1C).

*Other topics:* (1970KA1A, 1973TO16, 1974IR04, 1975BE56, 1976IR1B).

*Ground state properties:* (1975BE31).

$$\mu = 3.4359 \pm 0.0010 \text{ nm (1976CO1L; prelim.)}$$

*Mass of <sup>9</sup>Li:* The atomic mass excess of <sup>9</sup>Li is determined to be  $24955.4 \pm 2.0$  keV from the *Q*-values of the <sup>7</sup>Li(t, p) and <sup>10</sup>Be(d, <sup>3</sup>He) reactions. This value leads to a cubic coefficient in the IMME for *A* = 9 of  $d = 5.8 \pm 1.5$  keV for the ground state;  $d = 2.3 \pm 2.9$  keV for the second  $T = \frac{3}{2}$  state in *A* = 9 (1975KA18).

1. <sup>9</sup>Li( $\beta^-$ )<sup>9</sup>Be  $Q_m = 13.607$

Table 9.1: Energy levels of  ${}^9\text{Li}$

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions <sup>a</sup>
g.s.	$\frac{3}{2}^-; \frac{3}{2}$	$\tau_{1/2} = 178.3 \pm 0.4$ msec	$\beta^-$	1, 2, 3, 4, 5, 6, 7, 8
$2.691 \pm 5$	$(\frac{1}{2}^-)$		$(\gamma)$	2, 3, 7
$4.31 \pm 20$		$\Gamma = 100 \pm 30$		2, 7
$5.38 \pm 60$		$600 \pm 100$		2
$6.43 \pm 15$	$\geq \frac{9}{2}$	$40 \pm 20$		2, 7

<sup>a</sup> See also (1974AJ01).

The half-life of  ${}^9\text{Li}$  is  $178.3 \pm 0.4$  msec (1976AL02). Other recent values are  $175 \pm 1$  msec (1973AU1A),  $175 \pm 1$  msec (1974RO31). In (1974AJ01) the adopted value was  $176 \pm 2$  msec based on earlier measurements.  ${}^9\text{Li}$  decays to  ${}^9\text{Be}^*(0, 2.43, 2.78)$ : see  ${}^9\text{Be}$  and Table 9.7 (1970CH07).  $\log ft$  values are listed in Table 9.6: the nature of the transitions to  ${}^9\text{Be}^*(0, 2.43, 2.78)$  with  $J^\pi = \frac{3}{2}^-, \frac{5}{2}^-$  and  $(\frac{1}{2}^-)$  is evidence for  $J^\pi = \frac{3}{2}^-$  for  ${}^9\text{Li}_{\text{g.s.}}$  (1970CH07). Delayed neutrons are observed due to the decay of the two excited states of  ${}^9\text{Be}$  involved in the  $\beta^-$  decay: see  ${}^9\text{Be}$ . See also (1976FI03), (1973TA30) and (1973KU1D, 1974CH46, 1974KU06, 1975KR14, 1975WI1E, 1976KU07; theor.).

2.  ${}^7\text{Li}(t, p){}^9\text{Li}$   $Q_m = -2.386$

Protons are observed to excited states at  $E_x = 2.691 \pm 0.005$  MeV (1964MI04),  $4.31 \pm 0.03$ ,  $5.38 \pm 0.06$  and  $6.41 \pm 0.02$  MeV (1971YO04) [ $\Gamma_{\text{c.m.}} = 250 \pm 30, 600 \pm 100, < 100$  keV],  $4.31 \pm 0.02$  and  $6.435 \pm 0.020$  MeV (1978AJ02) [ $\Gamma_{\text{c.m.}} = 100 \pm 30, 40 \pm 20$  keV]. Angular distributions are reported at  $E_t = 11.3$  (1964MI04:  $t_0$ ),  $15$  MeV (1971YO04:  $t_0, t_2, t_4$ ) and at  $23$  MeV (1978AJ02:  $t_0, t_1, t_2, t_4$ ). They are consistent with  $J^\pi = \frac{3}{2}^-, (\frac{1}{2}^-)$  and  $\geq \frac{9}{2}$  for  ${}^9\text{Li}^*(0, 2.69, 6.43)$  (1978AJ02). See also (1974AJ01). For the mass of  ${}^9\text{Li}$  see the ‘‘GENERAL’’ section here (1975KA18). See also (1976HA1M).

3.  ${}^9\text{Be}(\pi^-, \gamma){}^9\text{Li}$   $Q_m = 125.960$

The excitation of  ${}^9\text{Li}^*(0, 2.69)$  is observed: see (1977BA1Q).

4.  ${}^9\text{Be}(n, p){}^9\text{Li}$   $Q_m = -12.825$

See (1977RO01) and (1978NE1A).

5.  $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$   $Q_m = -14.143$

See mass of  $^9\text{Li}$ , the “GENERAL” section here (1975KA18).

6.  $^{10}\text{Be}(^{11}\text{B}, ^{12}\text{C})^9\text{Li}$   $Q_m = -3.680$

See (1976FI03).

7.  $^{11}\text{B}(^6\text{Li}, ^8\text{B})^9\text{Li}$   $Q_m = -25.122$

At  $E(^6\text{Li}) = 80$  MeV the angular distribution to  $^9\text{Li}_{\text{g.s.}}$  has been measured. States at  $E_x = 2.59 \pm 0.10$ ,  $4.36 \pm 0.10$  and  $6.38 \pm 0.12$  MeV are also populated (1977WE03, 1977WE1B). See also (1974CE1A).

8.  $^{18}\text{O}(^7\text{Li}, ^{16}\text{O})^9\text{Li}$   $Q_m = -6.093$

See (1974AJ01).

<sup>9</sup>Be  
(Figs. 16 and 18)

GENERAL: (See also (1974AJ01).)

*Shell model:* (1975KU27, 1975SC1K, 1977CA08, 1977JA14, 1978BO31).

*α and cluster models:* (1974CH19, 1974GR42, 1974PA1B, 1975AB1E, 1975CH28, 1975KR1D, 1975RO1B, 1975SC1K, 1976ZA06, 1977DE1K, 1977HO1F, 1977OK01, 1977SE1D, 1978FO1J, 1978FO1G, 1978HO1E, 1978OK01, 1978OS01).

*Collective and deformed models:* (1974BO25, 1976BR26, 1977CA08).

*Special levels:* (1974IR04, 1974KU06, 1975CH28, 1975LI20, 1975WI1E, 1976IR1B, 1976ZA06, 1977CA08, 1977JA14, 1978BO31, 1978FO1G, 1978HO1E, 1978OK01, 1978OK1B).

*Electromagnetic transitions:* (1974CH46, 1974HA1C, 1974KU06, 1974MU13, 1976KU07, 1977DO06, 1978KI08).

*Astrophysical questions:* (1973CA1B, 1973SC1C, 1973SC1D, 1973TI1A, 1973TR1B, 1973WE1D, 1974AU1A, 1974BO1J, 1974BO1K, 1974RE1A, 1974RO1G, 1975CH1F, 1975LA1E, 1975ME1E, 1975PR1B, 1976AU1B, 1976AU1C, 1976BO1E, 1976BO1H, 1976BO1J, 1976CO1B, 1976EP1A, 1976FU1B, 1976GI1C, 1976HA1F, 1976PE1A, 1976RA1C, 1976RO1J, 1976RO12, 1976SI1C, 1976SI1D, 1976VI1A, 1977AU1B, 1977BO1F, 1977DR1C, 1977DW1A, 1977GA1C, 1977HA1L, 1977MA1H, 1977PR1D, 1977SC1D, 1977WE1D, 1978AU1C, 1978BY1A, 1978DW1A).

*Special reactions:* (1975AR14, 1975FE1A, 1975KU01, 1975RA12, 1975RA14, 1975RA21, 1975ZE01, 1976BE1K, 1976BU16, 1976CH28, 1976FR05, 1976HI05, 1976LE1F, 1976MI13, 1976NA11, 1976OS04, 1976RA1C, 1976RO12, 1977AR06, 1977FE1B, 1977GE08, 1977GO07, 1977GR1D, 1977KU1D, 1977PR05, 1977ST34, 1977ST1G, 1977YA1B, 1978BA1J, 1978BI08, 1978DI04, 1978FA1D, 1978GE1C, 1978GR1F, 1978UN01, 1978WE1D).

*Muon and neutrino capture and reactions:* (1974EN10, 1975CA1H, 1975CH22, 1975FE1B, 1975KN1C, 1976SH1H, 1977CA1E, 1977DO06, 1977MU1A, 1978DE15, 1978LE04).

*Pion capture and reactions:* (1973AL1A, 1973AR1B, 1973BA1G, 1973GO41, 1973WI1A, 1974AM01, 1974CL04, 1974DI20, 1974GO04, 1974HU14, 1974KA07, 1974MI06, 1974SP1A, 1974ST1G, 1974TA18, 1975BA1L, 1975BA1G, 1975BU1A, 1975ER03, 1975HU1D, 1975NI1B, 1975RO1G, 1975TA1C, 1975VO1D, 1975YA02, 1976AB1B, 1976AL1F, 1976AS1B, 1976BA1P, 1976DO1D, 1976DU1B, 1976ED1A, 1976EN02, 1976HE1G, 1976KI1E, 1976PI1B, 1976RA1F, 1976TR1A, 1977AB09, 1977AL1C, 1977AN1D, 1977AP1A, 1977AR1F, 1977AS1D, 1977BA51, 1977BA1Q, 1977BA2D, 1977BE1V, 1977DE1F, 1977DE1G, 1977DE1H, 1977DE1J, 1977DO06, 1977FY1A, 1977HO1B, 1977MA35, 1977MC1E, 1977PI1C, 1977SC1F, 1977SH1C, 1977SP1B, 1977TE1A, 1978BA1L, 1978BO01, 1978ER1C, 1978GE1E, 1978KI08, 1978MI1D, 1978MO01, 1978OL1A, 1978SE1D, 1978TH1B, 1978ZE01).

*Kaon reactions and reactions involving other mesons: (1973GO41, 1975PO1C, 1976AB1B, 1976BO1K, 1976BR1G, 1976KI1E, 1977AN1D, 1977BA27, 1977BE1U, 1977PO1A, 1978AT01, 1978BR1G).*

Table 9.2: Energy levels of  ${}^9\text{Be}$

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$		stable	2, 3, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 45, 46, 48, 49, 50, 51, 52, 53, 54, 55, 56
$1.680 \pm 15$	$\frac{1}{2}^+$	$200 \pm 20$	$\gamma, n$	3, 9, 10, 11, 14, 18, 19, 20, 22, 24, 25, 32, 37, 43, 45
$2.4294 \pm 1.3$	$\frac{5}{2}^-$	$0.77 \pm 0.15$	$\gamma, n, \alpha$	3, 9, 10, 11, 13, 18, 19, 20, 22, 23, 24, 25, 32, 34, 36, 37, 42, 43, 45, 52, 53
$2.78 \pm 120$	$\frac{1}{2}^-$	$1080 \pm 110$	n	3, 9, 13, 52
$3.049 \pm 9$	$\frac{5}{2}^+$	$282 \pm 11$	$\gamma, n$	9, 11, 14, 18, 19, 20, 21, 22, 24, 25, 32, 37, 42, 43, 45
$4.704 \pm 25$	$(\frac{3}{2})^+$	$743 \pm 55$	$\gamma, n$	3, 9, 18, 20, 21, 22, 24, 35, 43, 52
$6.76 \pm 60$	$\frac{7}{2}^-$	$1540 \pm 200$	$\gamma, n$	9, 18, 19, 20, 22, 24, 25, 34
$7.94 \pm 80$		$\approx 1000$		20
$11.283 \pm 24$		$575 \pm 50$	n	9, 20, 25, 34, 37
$11.81 \pm 20$	$T = \frac{1}{2}$	$400 \pm 30$	$\gamma, n$	9, 11, 14, 42, 52
$13.79 \pm 30$	$T = \frac{1}{2}$	$590 \pm 60$	$\gamma, n$	9, 11, 18, 42
$14.3926 \pm 1.8$	$\frac{3}{2}^-; \frac{3}{2}$	$0.329 \pm 0.067$	$\gamma, n, \alpha$	9, 18, 24, 37, 42
$14.4 \pm 300$		$\approx 800$		20, 37
$15.10 \pm 50$			$\gamma$	11, 18, 42

Table 9.2: Energy levels of  ${}^9\text{Be}$  (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
15.97 $\pm$ 30	$T = \frac{1}{2}$	$\approx 300$	$\gamma$	18, 42
16.671 $\pm$ 8		41 $\pm$ 4	$\gamma$	9, 18, 37
16.976 $\pm$ 2	$\frac{1}{2}^-; \frac{3}{2}$	< 0.47	$\gamma, n, p, d$	3, 4, 5, 14, 18
17.297 $\pm$ 10	$(\frac{5}{2})^-$	200	$\gamma, n, p, d, \alpha$	4, 5, 18
17.490 $\pm$ 9	$\leq \frac{7}{2}^+$	47	$\gamma, n, p, d, \alpha$	4, 5, 6, 18
18.02 $\pm$ 50			$\gamma, n$	18
18.58 $\pm$ 40			$\gamma, n, p, d, \alpha$	4, 5, 14, 18
19.20 $\pm$ 50		310 $\pm$ 80	p, d	5
19.51 $\pm$ 50			$\gamma, n$	14, 18
(19.9 $\pm$ 200)			$\gamma, n$	14
(20.47 $\pm$ 40)			$\gamma, p, d$	5, 14
20.74 $\pm$ 30		$\approx 1000$	$\gamma, n, p, t$	1, 14, 18, 20
(21.4 $\pm$ 200)			$\gamma, n$	14, 18, 20
(22.4 $\pm$ 200)		broad	$\gamma, n$	14, 18, 20
(23.8 $\pm$ 200)			$\gamma, n$	14, 35
(27.0 $\pm$ 500)		broad	$\gamma, n$	14

*Other topics:* (1973DZ1A, 1974CH19, 1974IR04, 1974KU06, 1974MU13, 1975CH1E, 1975CH28, 1975KU27, 1975LI20, 1975SC1M, 1976BR26, 1976DA1D, 1976DA1E, 1976ES1A, 1976GA1C, 1976IR1B, 1976KU07, 1976MA04, 1977KO1K, 1977PO1A, 1978OS01).

*Ground state properties:* (1974DE1E, 1974EN10, 1974KU06, 1974MU13, 1974PA1B, 1974SHYR, 1975BE31, 1975CH28, 1976BR26, 1976FU06, 1977BA2E, 1977MA35, 1977OK01, 1978AN07).

$$\mu = -1.1778 \pm 0.0009 \text{ nm (1976WE17);}$$

$$Q = +53 \pm 3 \text{ mb (1978SH1E).}$$

The neutron spectroscopic factor of  ${}^9\text{Be}$  is  $0.42 \pm 0.04$  (1977LA11).

1. (a)  ${}^6\text{Li}(t, n){}^8\text{Be}$   $Q_m = 16.0240$   $E_b = 17.6892$
- (b)  ${}^6\text{Li}(t, p){}^8\text{Li}$   $Q_m = 0.801$
- (c)  ${}^6\text{Li}(t, d){}^7\text{Li}$   $Q_m = 0.993$

- (d)  ${}^6\text{Li}(t, \alpha){}^5\text{He}$   $Q_m = 15.22$   
(e)  ${}^6\text{Li}(t, n){}^4\text{He}{}^4\text{He}$   $Q_m = 16.1159$

The  $0^\circ$  differential cross section for reaction (a) increases monotonically between  $E_t = 0.10$  and 2.4 MeV (1960SE12, 1962SE1A, 1961VA43) except for a resonance at  $E_t = 1.875$  MeV ( ${}^9\text{Be}^* = 18.938$ ). The excitation function for  ${}^8\text{Li}$  (reaction (b)) increases monotonically for  $E_t = 0.275$  to 1.000 MeV (1972CI05). In the range  $E_t = 2.0$  to 6.8 MeV, a broad peak [ $\Gamma \approx 1.3$  MeV] is observed at  $E_t = 4.57$  MeV [ $E_x = 20.73$  MeV] (1973AB10). For reaction (c), (d) and (e) see (1966LA04). See also  ${}^8\text{Li}$  and (1977SU1A).

2.  ${}^6\text{Li}(\alpha, p){}^9\text{Be}$   $Q_m = -2.1248$

Angular distributions of  $p_0$  have been measured at  $E_\alpha = 10.2$  to 14.7 MeV and at 30 MeV: see (1974AJ01). In a study of the inverse reaction [ ${}^9\text{Be}(p, \alpha){}^6\text{Li}$ ] at 45 MeV, the spectroscopic factors for ( ${}^6\text{Li} + t$ ) and ( ${}^5\text{He} + \alpha$ ) cluster structures in  ${}^9\text{Be}$  are calculated to be 0.003 and 0.102, respectively (1972DE01, 1972DE02).

3.  ${}^7\text{Li}(d, \gamma){}^9\text{Be}$   $Q_m = 16.6960$

For  $E_d = 0.1$  to 1.1 MeV, a resonance in the yield of capture  $\gamma$ -rays is observed at  $E_d = 362 \pm 3$  keV (1965WO01),  $361 \pm 2$  keV (1965IM01), corresponding to  ${}^9\text{Be}^*(16.976)$  with  $\Gamma_{c.m.} < 0.47$  keV. The small width of this state and its energy correspondence with  ${}^9\text{Li}^*(2.69)$  argue for  $T = \frac{3}{2}$  (1965WO01). The angular distribution of the  $\gamma$ -rays to  ${}^9\text{Be}(0)$  is isotropic to within 7% (1965IM01). The branching ratios to  ${}^9\text{Be}^*(0, 1.7, 2.4, 2.8, 3.1, 4.7)$  are  $100 / 11.8 \pm 0.6 / 3.3 \pm 0.7 / 13.3 \pm 4.2 / - / 12.9 \pm 1.3$  (1971SC19). The  $E_x$  and  $\Gamma$  of  ${}^9\text{Be}^*(2.8, 4.7)$  are 2.82 and 1.7 MeV, and 4.64 and 0.95 MeV, respectively. The character of the decay suggests  $(\frac{1}{2})^-$  for the second  $T = \frac{3}{2}$  state [ ${}^9\text{Be}^*(16.98)$ ] and is consistent with  $J^\pi = (\frac{1}{2})^-$  for  ${}^9\text{Be}^*(2.8)$  (1971SC19).

4. (a)  ${}^7\text{Li}(d, n){}^8\text{Be}$   $Q_m = 15.031$   $E_b = 16.6960$   
(b)  ${}^7\text{Li}(d, \alpha){}^5\text{He}$   $Q_m = 14.23$   
(c)  ${}^7\text{Li}(d, n){}^4\text{He}{}^4\text{He}$   $Q_m = 15.123$

The yield of neutrons has been measured for  $E_d = 0.2$  to 19 MeV [see (1974AJ01)] and for 0.4 to 3.0 MeV (1975MC02;  $\sigma_t$ ), 2.5 MeV (1974JO1C) and 14.8, 18 and 23 MeV (1977LO10). Resonances in the yield of neutrons are reported at  $E_d = 0.36, 0.68, 0.98$  and (1.8) MeV: see Table 9.3. Polarization measurements have been reported at  $E_d = 0.64$  MeV and at 2.5 to 3.7 MeV: see (1974AJ01). See also (1975SE07, 1977SE09).

Table 9.3: Resonances in  ${}^7\text{Li} + \text{d}$ 

${}^7\text{Li}(\text{d}, \text{n}){}^8\text{Be}$ $E_{\text{res}}$ (keV)	${}^7\text{Li}(\text{d}, \alpha){}^5\text{He}$ $E_{\text{res}}$ (keV)	${}^7\text{Li}(\text{d}, \text{p}){}^8\text{Li}$		$E_{\text{x}}$ (MeV)	$J^{\pi}$
		$E_{\text{res}}$ (keV)	$\Gamma_{\text{lab}}$ (keV)		
360 <sup>a,b</sup>		$360 \pm 3$ <sup>a</sup>	$< 2$	16.976	
680 <sup>c</sup>	750 <sup>d</sup>	$773 \pm 10$ <sup>f</sup>	250	17.297	$(\frac{3}{2}, \frac{5}{2})^{+}$ <sup>i</sup>
980 <sup>c</sup>	1000 <sup>d</sup>	$1025 \pm 10$ <sup>f</sup>	60	17.492	$(\frac{3}{2}, \frac{5}{2}, \frac{7}{2})^{-}$ <sup>i</sup>
(1800) <sup>c</sup>	2500 <sup>e</sup>	2000 <sup>g</sup>		(18.5)	
		$2375 \pm 50$ <sup>g,h</sup>		18.54	
		$3220 \pm 50$ <sup>h</sup>	$400 \pm 100$	19.20	
		$\approx 4800$ <sup>h</sup>		(20.4)	

<sup>a</sup> In  ${}^7\text{Li}(\text{d}, \gamma)$ ,  $E_{\text{res}} = 361 \pm 2$  keV (1965IM01, 1965WO01);  $\Gamma_{\text{no}}/\Gamma_{\gamma} \approx 1.5$ ,  $\Gamma_{\text{d}_0}/\Gamma_{\gamma} < 400$ ,  $\Gamma_{\alpha_0}/\Gamma_{\gamma} < 20$  (1965IM01) for  ${}^9\text{Be}^*(16.98)$ .

<sup>b</sup> (1965IM01).

<sup>c</sup> (1952BA1A, 1957SL01).

<sup>d</sup> (1971FR04, 1969DE31, 1963PA04).

<sup>e</sup> (1963PA04): broad structure.

<sup>f</sup> (1976SC14). See also Table 9.3 in (1974AJ01).

<sup>g</sup> (1956BE1A).

<sup>h</sup> (1973AB10).

<sup>i</sup> (1973RO42). See, however, (1972DE44).

The yield of  $\alpha$ -particles has been studied for  $E_{\text{d}} = 0.2$  to  $3.0$  MeV [see (1974AJ01)] and at  $E_{\text{d}} = 0.60$  to  $1.25$  MeV (1973RO42:  $\alpha_1$ ). Resonances are observed at  $E_{\text{d}} = 0.75$ ,  $1.00$  and  $2.5$  MeV: see Table 9.3.

At  $E_{\text{d}} = 1.0$  MeV reaction (c) is dominated by sequential decay via  ${}^8\text{Be}^*(2.9)$  and  ${}^5\text{He}_{\text{g.s.}}$ . There is evidence also for the involvement of  ${}^9\text{Be}^*(17.50)$ :  $J = \frac{3}{2}$  is suggested (1973HE26). See also (1975DA15, 1978SP03). See also  ${}^5\text{He}$ ,  ${}^8\text{Be}$ , (1975FO19; astrophysics) and (1974JO1C, 1975GO1H, 1977GR1F, 1977LO10, 1977SA1K; applications).

5.  ${}^7\text{Li}(\text{d}, \text{p}){}^8\text{Li}$

$$Q_{\text{m}} = -0.192$$

$$E_{\text{b}} = 16.6960$$

The yield of  $\text{p}_0$  has been measured for  $E_{\text{d}} = 0.29$  to  $0.78$  MeV and the  ${}^8\text{Li}$  yield for  $E_{\text{d}} = 0.4$  to  $7.0$  MeV [see (1974AJ01)] and from threshold to  $3.6$  MeV (1975MC02, 1974MCZS:  $\sigma_{\text{t}}$ ; also cross sections for  $\text{p}_0$  and  $\text{p}_1$ , from  $0.5$  MeV). The yield of  ${}^8\text{Li}$  has also been measured for  $E_{\text{d}} = 0.61$  to  $1.95$  MeV (1976SC14): resonances are displayed in Table 9.3. The total cross section at the  $E_{\text{d}} = 0.77$  MeV resonance is  $181 \pm 8$  mb (1976SC14). For earlier values see (1974AJ01).

The cross section at the 0.77 MeV resonance serves as normalization for the  ${}^7\text{Be}(p, \gamma){}^8\text{B}$  data and to calculations of the  $S$ -factor, of interest in relation to the solar neutrino problem. See also (1976HA1J).

6.  ${}^7\text{Li}(d, d){}^7\text{Li}$

$$E_b = 16.6960$$

The elastic scattering [ $E_d = 0.4$  to  $1.8$  MeV] shows a marked increase in cross section for  $E_d = 0.8$  to  $1.0$  MeV (perhaps related to  ${}^9\text{Be}^*(17.30)$ ) and a conspicuous anomaly at  $E_d = 1.0$  MeV, due to p-wave deuterons [ ${}^9\text{Be}^*(17.50)$ ]. The elastic scattering has also been studied for  $E_d = 1.0$  to  $2.6$  MeV (1974LO10) and  $E_d = 10.0$  to  $12.0$  MeV (1971BI11). See also  ${}^7\text{Li}$ .

7. (a)  ${}^7\text{Li}(d, t){}^6\text{Li}$

$$Q_m = -0.993$$

$$E_b = 16.6960$$

(b)  ${}^7\text{Li}(d, {}^3\text{He}){}^6\text{He}$

$$Q_m = -4.482$$

The cross section for reaction (a) rises steeply from threshold to 95 mb at  $E_d = 2.4$  MeV and then more slowly to  $\approx 165$  mb at  $E_d = 4.1$  MeV (1955MA20). The  $t_0$  yield curve ( $\theta_{\text{lab}} = 155^\circ$ ) decreases monotonically for  $E_d = 10.0$  to  $12.0$  MeV (1971ZA07). See also  ${}^6\text{Li}$ . For reaction (b) see  ${}^6\text{He}$ .

8.  ${}^7\text{Li}(t, n){}^9\text{Be}$

$$Q_m = 10.4387$$

See (1974AJ01) and  ${}^{10}\text{Be}$ .

9. (a)  ${}^7\text{Li}({}^3\text{He}, p){}^9\text{Be}$

$$Q_m = 11.2025$$

(b)  ${}^7\text{Li}({}^3\text{He}, np){}^8\text{Be}$

$$Q_m = 9.537$$

Observed proton groups are listed in Table 9.4. The parameters for the particle and  $\gamma$ -decay of observed states are displayed in Table 9.5, and in Table 9.6 (1976MC10). Angular distributions have been reported in the range  $E({}^3\text{He}) = 0.9$  to  $10$  MeV [see (1974AJ01)], at  $1.0$  to  $2.5$  MeV (1975BO55:  $p_0, p_2$ ) and at  $14$  MeV (1976IR02:  $p_0$ ). See also  ${}^{10}\text{B}$  and (1973WE15, 1975ST16; theor.).

10.  ${}^7\text{Li}(\alpha, d){}^9\text{Be}$

$$Q_m = -7.1507$$

Table 9.4: Excited states of  ${}^9\text{Be}$  from  ${}^7\text{Li}({}^3\text{He}, p){}^9\text{Be}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Refs.
1.64		(1968CO07)
$2.4292 \pm 1.7$	$< 8$	(1968KR02)
$2.9 \pm 250$	$1000 \pm 250$ <sup>e</sup>	(1965LY01, 1971AD01)
$3.076 \pm 15$ <sup>b</sup>	$289 \pm 22$	(1968KR02)
$4.704 \pm 25$ <sup>c</sup>	$743 \pm 55$	(1968KR02)
$6.7 \pm 100$	$2000 \pm 200$	(1968CO07, 1968CO08)
$11.29 \pm 30$	$620 \pm 70$	(1968CO07)
$11.81 \pm 20$	$400 \pm 30$	(1968CO07)
$13.78 \pm 30$	$590 \pm 60$	(1968CO07)
$14.396 \pm 5$ <sup>d</sup>	$< 5$ <sup>d</sup>	(1965LY01, 1971AD01)
$16.671 \pm 8$	$41 \pm 4$	(1968CO07)

<sup>a</sup> See also Table 9.4 in (1974AJ01).

<sup>b</sup> See also (1968CO07).

<sup>c</sup> See also (1968CO07, 1968CO08).

<sup>d</sup> See also Table 9.6.

<sup>e</sup> From  $\gamma$ -decay of  ${}^9\text{Be}^*(14.39)$ .

At  $E_\alpha = 30$  MeV  $d_0$ ,  $d_1$  and  $d_2$  angular distributions have been measured (1972ME07). See also (1976LE1K).

11.  ${}^7\text{Li}({}^6\text{Li}, \alpha){}^9\text{Be}$   $Q_m = 15.223$

Angular distributions of the  $\alpha_0$  group have been measured at  $E({}^7\text{Li}) = 3.78$  to  $5.95$  MeV (1967KI03). See also (1974AJ01).

12.  ${}^7\text{Li}({}^{11}\text{B}, {}^8\text{Be}){}^9\text{Be}$   $Q_m = 7.286$

Not reported: see (1970LK1A; theor.).

13.  ${}^9\text{Li}(\beta^-){}^9\text{Be}$   $Q_m = 13.607$

Table 9.5: Neutron decay of  ${}^9\text{Be}$  states

${}^9\text{Be}$ state (MeV)	$l_n$	Decay (in %) to		$\theta^2$ <sup>a</sup> (%)	Refs.
		${}^8\text{Be}(0)$	${}^8\text{Be}^*(2.9)$		
2.43	3	$7.5 \pm 1$		$2.1 \pm 0.6$	(1966CH20)
		$6.4 \pm 1.2$ <sup>c</sup>			(1970CH07)
2.78	1	mainly		$0.48 \pm 0.06$	(1970CH07)
3.05	2	$87 \pm 13$		$81 \pm 13$	(1966CH20, 1968CO08)
4.70	2	$13 \pm 4$		$6.0 \pm 0.4$	(1968CO08)
6.76	3	$\leq 2$		$\leq 6$	(1968CO08)
	1		$55 \pm 14$	$37 \pm 10$	(1968CO08)
11.28	1	$\leq 2$		$\leq 0.1$	(1968CO08)
	1		$14 \pm 4$	$0.93 \pm 0.28$	(1968CO08)
	3			$4.0 \pm 1.2$	(1968CO08)
11.81	1	$\leq 3$		$\leq 0.1$	(1968CO08)
	1		$12 \pm 4$	$0.48 \pm 0.16$	(1968CO08)
	3			$1.8 \pm 0.6$	(1968CO08)
14.39 <sup>b</sup>					

<sup>a</sup> Expressed in units of  $\hbar^2/mR^2 = 2.47$  MeV (1968CO08, 1970CH07).

<sup>b</sup> See Table 9.6.

<sup>c</sup> (1978CH07) report  $(11 \pm 2)\%$ .

${}^9\text{Li}$  decays by  $\beta^-$  emission with  $\tau_{1/2} = 178.3 \pm 0.4$  msec to  ${}^9\text{Be}^*(0, 2.43, 2.78)$ : see  ${}^9\text{Li}$ , reaction 1, and Table 9.7.  ${}^9\text{Be}(2.43, 2.78)$  are neutron unstable. The probability that  ${}^9\text{Li}$  decays to these two states is 0.35. The branching ratio for the  ${}^9\text{Be}^*(2.43) \rightarrow {}^8\text{Be}(0) + n$  decay is  $(6.4 \pm 1.2)\%$ .  ${}^9\text{Be}^*(2.78)$  decays mainly to  ${}^8\text{Be}(0) + n$ . The assignment  $J^\pi = \frac{1}{2}^-$  to  ${}^9\text{Be}^*(2.78)$  derives from the nature of the  ${}^9\text{Li}$  decay, and the large value of  $\theta_p^2[0.48 \pm 0.06]$  which is in agreement with the shell model prediction that the  $\frac{1}{2}^-$  state should decay mainly by p-wave neutron emission to  ${}^8\text{Be}(0)$ : if  $J^\pi = \frac{3}{2}^-$  this decay branch should be small (1970CH07).

14. (a)  ${}^9\text{Be}(\gamma, n){}^8\text{Be}$   $Q_m = -1.6652$   
 (b)  ${}^9\text{Be}(\gamma, \alpha){}^5\text{He}$   $Q_m = -2.47$   
 (c)  ${}^9\text{Be}(\gamma, n){}^4\text{He}{}^4\text{He}$   $Q_m = -1.5733$   
 (d)  ${}^9\text{Be}(\gamma, 2n){}^7\text{Be}$   $Q_m = -20.565$

Table 9.6: Parameters <sup>a</sup> of the first  $T = \frac{3}{2}$  states in <sup>9</sup>Be and <sup>9</sup>B,  $J^\pi = \frac{3}{2}^-$ 

<sup>9</sup> Be			<sup>9</sup> B		
$E_x$ (MeV)	$14.3926 \pm 0.0018$	(1974KA15)		$14.6550 \pm 0.0025$	(1974KA15)
$\Gamma_{\gamma_0}$ (eV)	$6.9 \pm 0.5$	(1973BE19)		$(6.9 \pm 0.5)^b$	
$\Gamma$ (eV)	$329 \pm 67$	(1971AD01, 1973BE19)		$260^{+90}_{-120}$	(1971AD01, 1973BE19)
				$290 \pm 93$	(1976MC10) <sup>c</sup>
				$275 \pm 93$	(1976MC10)
$\Gamma_{\gamma_1}/\Gamma_{\gamma_0}$	$1.19 \pm 0.16$	(1971AD01)			d
$\Gamma_{\gamma_2}/\Gamma_{\gamma_1}$	$0.30 \pm 0.04$	(1971AD01)			
$\Gamma_{n_0}/\Gamma$	$0.028 \pm 0.021$	(1976MC10)	$\Gamma_{p_0}/\Gamma$	$0.11 \pm 0.04$	(1976MC10)
$\Gamma_{n_1}/\Gamma$	$0.50 \pm 0.11$	(1976MC10)	$\Gamma_{p_1}/\Gamma$	$0.33 \pm 0.09$	(1976MC10)
$\Gamma_{n_0}$ (eV)	$9 \pm 8$	(1976MC10) <sup>e</sup>	$\Gamma_{p_0}$ (eV)	$30 \pm 17$	(1976MC10)
$\Gamma_{n_1}$ (eV)	$141 \pm 34$	(1972AD04, 1973BE19)	$\Gamma_{p_1}$ (eV)	$96 \pm 16$	(1972AD04, 1973BE19)
	$156 \pm 47$	(1976MC10)		$91 \pm 38$	(1976MC10)
	$147 \pm 28$	mean		$95 \pm 15$	mean
$\Gamma_{n_1}/\Gamma_{n_0}$	$18 \pm 14$	(1976MC10)	$\Gamma_{p_1}/\Gamma_{p_0}$	$3.2 \pm 1.9$	(1976MC10)
$\gamma_{n_1}^2/\gamma_{n_0}^2$	$22 \pm 17$	(1976MC10)	$\gamma_{p_1}^2/\gamma_{p_0}^2$	$3.7 \pm 2.2$	(1976MC10)
$\Gamma_{\alpha_0}/\Gamma_{\gamma_0}$	$31.2 \pm 9.8$	(1972AD04, 1973BE19)			

<sup>a</sup> See Table 1 in (1976MC10) and Table 9.6 in (1974AJ01).

<sup>b</sup> Assumed identical to <sup>9</sup>Be.

<sup>c</sup> Calculated from  $\Gamma_{p_1}/\Gamma_{\gamma_0}$ ,  $\Gamma_{p_1}/\Gamma$  and  $\Gamma_{\gamma_0}$ .

<sup>d</sup> Similar to branching ratios in <sup>9</sup>Be (1971AD01):  $\Gamma_{\gamma_1}$  is transition to <sup>9</sup>Be\*(2.4) or <sup>9</sup>B\*(2.4);  $\Gamma_{\gamma_2}$  is transition to <sup>9</sup>Be\*(2.9) or <sup>9</sup>B\*(2.8).

<sup>e</sup> See also (1978MC04; theor.).

 Table 9.7: Branching parameters in <sup>9</sup>Li  $\beta$ -decay (1970CH07)

$E_x$ in <sup>9</sup> Be (MeV)	$J^\pi; T$	Branching ratio (%)	$\log ft^b$
0	$\frac{3}{2}^-; \frac{1}{2}$	$65.0^{+2.7}_{-2.4}$	$5.20^{+0.01}_{-0.02}$
2.43	$\frac{5}{2}^-; \frac{1}{2}$	$32.0^{+2.7}_{-3.7}$	$5.10^{+0.04}_{-0.05}$
$2.78 \pm 0.12^a$	$\frac{1}{2}^-; \frac{1}{2}$	$3.0^{+2.7}_{-0.3}$	$6.06^{+0.05}_{-0.28}$

<sup>a</sup>  $\Gamma_{c.m.} = 1.10 \pm 0.12$  MeV;  $\theta_p^2 = 0.48 \pm 0.06$  (1970CH07).

<sup>b</sup> M. J. Martin, Nucl. Data Project, private communication.

The photoneutron cross section has been measured from threshold to 320 MeV: see Table 9.6 in (1966LA04) and (1975KN03, 1974KN08: 17 to 37 MeV; monoenergetic  $\gamma$ ), (1973AN29, 1975HU03, 1977BU02, 1978BU01). A sharp peak occurs 6 keV above threshold (1967BE49) [but see discussion in (1968BA1C)] with  $\sigma_{\max} = 1.6$  mb. The cross section then decreases slowly to 1.2 mb at  $E_\gamma = 40$  keV (1967BE49). From bremsstrahlung studies, peaks in the  $(\gamma, \text{Tn})$  cross section are observed corresponding to  $E_x = 1.80$  and 3.03 MeV (1975HU03) [see both (1973AN29, 1975HU03) for additional structures]. At higher energies, using monoenergetic photons, (1975KN03) [who also report  $(\gamma, n) + (\gamma, \text{pn})$  and  $(\gamma, 2n)$ ] find the  $(\gamma, \text{Tn})$  cross section to be relatively smooth from  $E_\gamma = 17$  to 37 MeV with weak structures which correspond to  $E_x = 17.1, 18.8, 19.9, 21.4, 22.4, 23.8 [\pm 0.2]$  MeV and  $27 \pm 0.5$  MeV (broad). See also (1976BE1H).

In the range  $E_\gamma = 18$  to 26 MeV the integrated  $(\gamma, n_0)$  cross section is  $< 0.1$  MeV  $\cdot$  mb, that for  $(\gamma, n_1) = 2.4 \pm 0.4$  MeV  $\cdot$  mb (1977BU02) and the combined integrated cross section for  $(\gamma, n)$  to  ${}^8\text{Be}^*(16.6)$  and  $(\gamma, \alpha_0)$  to  ${}^5\text{He}_{\text{g.s.}}$  is  $13.1 \pm 2$  MeV  $\cdot$  mb (1978BU01).

The total absorption cross section has been measured for  $E_\gamma = 10$  to 210 MeV: it rises to  $\approx 5$  mb at  $\approx 21$  MeV, decreases to about 0 at 160 MeV and then increases to  $\approx 1.5$  mb at 210 MeV (1975AH06, 1974AH03). (1969DO09) report an integrated cross section of  $156 \pm 15$  MeV  $\cdot$  mb for  $E_\gamma = 10$  to 29 MeV and resonant structure at  $E_\gamma = 11.8, (13.5), 14.8, (17.3), (19.5), 21.0, (23.0)$  and  $(25.0)$  MeV (1969DO09). Fine structure is reported at  $E_\gamma = 20.47 \pm 0.04$  and  $20.73 \pm 0.04$  MeV (1964TE04). The absorption coefficient of 6.418 MeV  $\gamma$ -rays is reported by (1975MO27). Double pair production by 6.6 MeV  $\gamma$ -rays has been observed by (1976RO1N):  $\sigma_{\pi\pi}/\sigma_\pi = (1.8 \pm 0.6) \times 10^{-5}$  [average of values from interaction with several different nuclei]. See also (1973VE12, 1975SC05), (1974BU1A, 1975AB1F, 1975BR1F, 1977DA1B), (1975FO19; astrophysics) and (1976KH1C, 1978OC1B, 1978OK1B; theor.).

$$\begin{aligned} 15. \text{ (a) } & {}^9\text{Be}(\gamma, \text{p}){}^8\text{Li} & Q_{\text{m}} &= -16.888 \\ & \text{(b) } & {}^9\text{Be}(\gamma, \text{np}){}^7\text{Li} & Q_{\text{m}} &= -18.921 \end{aligned}$$

The yield shows structure in the energy region corresponding to the  ${}^9\text{Be}$  levels at 17 – 19 MeV (1962CL06) followed by the giant resonance at  $E_\gamma \approx 23$  MeV (1962CL06;  $\sigma = 2.64 \pm 0.30$  mb). (1966DE13) report structure attributed to eleven states of  ${}^9\text{Be}$  with  $18.2 < E_x < 32.2$  MeV. Integrated cross sections have been obtained for each of these resonances, and over different energy intervals for protons leading to  ${}^8\text{Li}^*(0 + 0.98, 2.26 + 3.21, 9.0, 17.0)$  (1966DE13). Angular and energy distributions of photoprotons in various energy intervals have been studied by many groups: see (1974AJ01) and (1974GO04). See also (1974DO07, 1975MA1K, 1978DO02) and (1975AN1H; theor.).

$$\begin{aligned} 16. \text{ (a) } & {}^9\text{Be}(\gamma, \text{d}){}^7\text{Li} & Q_{\text{m}} &= -16.6960 \\ & \text{(b) } & {}^9\text{Be}(\gamma, \text{t}){}^6\text{Li} & Q_{\text{m}} &= -17.6892 \end{aligned}$$

The integrated cross sections are reported to be  $1.0 \pm 0.5 \text{ MeV} \cdot \text{mb}$  ( $E_\gamma = 21 \rightarrow 33 \text{ MeV}$ ) for reaction (a) to  ${}^7\text{Li}^*(0 + 0.4)$  and  $0.6 \pm 0.3 \text{ MeV} \cdot \text{mb}$  ( $E_\gamma = 25 \rightarrow 33 \text{ MeV}$ ) for reaction (b) to  ${}^6\text{Li}(0)$ . The total integrated cross section for  $[(\gamma, p) + (\gamma, pn) + (\gamma, d) + (\gamma, t)]$  is given as  $33 \pm 3 \text{ MeV} \cdot \text{mb}$  by (1966DE13), who also report resonances in the  $(\gamma, d)$  and  $(\gamma, t)$  cross sections corresponding to  ${}^9\text{Be}^*(26.0 \pm 0.2)$  and  ${}^9\text{Be}^*(32.2 \pm 0.3)$ , respectively. See also (1974AJ01).

17.  ${}^9\text{Be}(\gamma, \gamma){}^9\text{Be}$

See (1977CR1C) and (1974AJ01).

18. (a)  ${}^9\text{Be}(e, e){}^9\text{Be}$

(b)  ${}^9\text{Be}(e, en){}^8\text{Be}$   $Q_m = -1.6652$

(c)  ${}^9\text{Be}(e, ep){}^8\text{Li}$   $Q_m = -16.888$

(d)  ${}^9\text{Be}(e, e\alpha){}^5\text{He}$   $Q_m = -2.47$

(e)  ${}^9\text{Be}(e, e\pi^+){}^9\text{Li}$   $Q_m = -153.174$

$\langle r^2 \rangle^{1/2} = 2.46 \pm 0.11 \text{ fm}$ ,  $Q = 6.5_{-0.6}^{+0.9} \text{ fm}^2$ ,  $b = 1.5_{-0.2}^{+0.3} \text{ fm}$  [ $b =$  oscillator parameter] (1973BE19);

$\langle r^2 \rangle^{1/2} = 2.519 \pm 0.012 \text{ fm}$ ,  $Q = 6.4 \pm 2.4 \text{ fm}^2$  ((1972JA10) and private communication to J.C. Bergstrom);

$\langle r^2 \rangle_M^{1/2} = 3.2 \pm 0.3 \text{ fm}$ ;  $\Omega = 6 \pm 2 \mu_N \cdot \text{fm}^2$  [this value of the magnetic octupole moment implies a deformation of the average nuclear potential] (1975LA23; see also for calculations of (b)).

The elastic scattering of electrons has been studied for  $E_e$  up to 700 MeV: see (1974AJ01). Magnetic elastic scattering gives indications of both M1 and M3 contributions (1966RA29, 1965GR18). See also (1973FE13).

Inelastic scattering populates a number of levels: Table 9.8 displays the parameters of the states. See also (1974TI05, 1975DI05). Electron bremsstrahlung has been measured at  $E_e = 1.0$  and  $2.0 \text{ MeV}$  by (1968RE11). For reaction (b) see (1974AJ01). At  $E_e = 700 \text{ MeV}$  the proton separation spectra are similar to those observed in  $(p, 2p)$  except for an upward shift in the excitation (1978NA05). See also (1977TU1C, 1977BE1Q:  $E_e = 500 \text{ MeV}$ ; observe a peak at  $E_x \approx 25 \text{ MeV}$ ), (1973KU19:  $1.184 \text{ GeV}$ ) and (1974AJ01). For reaction (d) see (1975GE12). For reaction (e) see (1977SH1C). See also (1974DE1E, 1975FA1A) and (1973GA19, 1973KE1B, 1974BE10, 1974KU06, 1975AB1E, 1976DU05, 1977KU12, 1977OK01, 1977OK1A, 1978OK01; theor.).

19. (a)  ${}^9\text{Be}(n, n){}^9\text{Be}$

Table 9.8: Levels of  ${}^9\text{Be}$  from  ${}^9\text{Be}(e, e'){}^9\text{Be}^*$  <sup>a</sup>

$E_x$ in ${}^9\text{Be}$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Transition	$J^\pi$	$\Gamma_{\gamma_0}$ (eV)	Refs.
$1.78 \pm 30$ <sup>b</sup>	$150 \pm 50$	E1	$\frac{1}{2}^+$	$0.30 \pm 0.12$	(1968CL08)
$2.44 \pm 20$ <sup>c</sup>	$< 30$	M1	$\frac{5}{2}^-$	$0.089 \pm 0.010$	(1968CL08)
		E2		$(1.89 \pm 0.14) \times 10^{-3}$ <sup>d</sup>	(1968CL08)
$3.04 \pm 20$	$450 \pm 150$	E1 <sup>e</sup>	$\frac{5}{2}^+$ <sup>e</sup>	$0.30 \pm 0.25$ <sup>n</sup>	(1968CL08, 1975FA1A)
$4.7 \pm 200$	$700 \pm 300$	E(1)		$2.4 \pm 1.2$ <sup>f</sup>	(1968CL08)
$6.4 \pm 100$	$2000 \pm 500$	E2	$\frac{7}{2}^-$	$0.109 \pm 0.005$	(1963NG01, 1965NG1A)
	$1100 \pm 300$			$0.082 \pm 0.035$	(1968CL08)
<sup>g</sup>					
$13.84 \pm 50$ <sup>h</sup>					(1973BE19)
$14.388 \pm 15$	$< 70$	M1 <sup>i</sup>	$\frac{3}{2}^-$	$6.9 \pm 0.5$ <sup>j</sup>	(1973BE19)
$15.10 \pm 50$ <sup>h</sup>					(1973BE19)
$15.97 \pm 30$ <sup>h</sup>	$\approx 300$	M1		$3.7 \pm 0.8$ <sup>f</sup>	(1973BE19)
$16.631 \pm 15$	$< 70$	M2 <sup>k</sup>	$\leq \frac{7}{2}^+$	$0.26 \pm 0.02$ <sup>f</sup>	(1973BE19)
		M1	$\leq \frac{5}{2}^-$	$2.0 \pm 0.5$ <sup>f</sup>	(1973BE19)
$16.961 \pm 15$	$< 70$	M1	$\leq \frac{1}{2}^-$	$11.5 \pm 1.4$ <sup>l</sup>	(1973BE19)
$17.28$		M1	$\leq \frac{3}{2}^-$	$7.3 \pm 1.3$ <sup>f</sup>	(1973BE19)
$17.480 \pm 20$	$\approx 100$	M2 <sup>i,k</sup>	$\leq \frac{7}{2}^+$	$0.40 \pm 0.03$ <sup>f</sup>	(1973BE19)
$18.02 \pm 50$ <sup>h</sup>					(1973BE19)
$18.62 \pm 50$ <sup>h</sup>					(1973BE19)
$19.51 \pm 50$ <sup>h</sup>					(1973BE19)
$20.76 \pm 50$ <sup>h</sup>					(1973BE19)
<sup>m</sup>					

<sup>a</sup> See also Table 9.8 in (1974AJ01).

<sup>b</sup>  $E_x = 1.79 \pm 0.06$  (1973SL02).

<sup>c</sup> See also (1975FA1A).

<sup>d</sup>  $B(\text{C}2, \omega)^\dagger = 45.7 \pm 3.5 e^2 \cdot \text{fm}^4$ . Form factor measured for  $E_e = 60.7$  to  $120.0$  MeV (1974EN01).

<sup>e</sup> Assumed.

<sup>f</sup>  $g\Gamma_{\gamma_0}$ , where  $g = (2J_f + 1)/(2J_i + 1)$ .

<sup>g</sup> For additional states reported by (1968VA05) see Table 9.8 in (1974AJ01) and footnote <sup>h</sup> in that table.

<sup>h</sup> Weak transition.

<sup>i</sup> See also (1974NA25).

<sup>j</sup> Best value, calculated by (1973BE19): see footnote <sup>f</sup> in Table 9.8 in (1974AJ01).

<sup>k</sup> Or pure spin-flip E1.

<sup>l</sup> See also footnote <sup>j</sup> in (1974AJ01) and (1975FA1A).

<sup>m</sup> Higher states reached by M1 transitions are reported at  $21.6 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $24.4 \pm 0.2$  and  $25.7 \pm 0.2$  MeV (1968VA05).

<sup>n</sup> The group reported by (1968CL08) may consist of two unresolved states, the second one reached by an M1 transition [ $J^\pi = (\frac{1}{2})^-$ ] with  $\Gamma_{\gamma_0} = 0.18 \pm 0.09$  eV. I am indebted to Dr. L.W. Fagg for his help in understanding this point.

- (b)  ${}^9\text{Be}(n, 2n){}^8\text{Be}$   $Q_m = -1.6652$   
(c)  ${}^9\text{Be}(n, nd){}^7\text{Li}$   $Q_m = -16.6960$

Population of  ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1)$  and possibly  ${}^9\text{Be}^*(6.8)$  has been reported in this reaction: see (1974AJ01). For the neutron decay of these states see Table 9.5. Angular distributions have been reported at  $E_n = 14$  MeV [see (1974AJ01)], at 5.9, 10.1 and 14.2 MeV (1977DR09:  $n_0, n_{1+2+3+4}$ ), and 14.1 MeV (1974HY01:  $n_0, n_2$ ). See also (1974HO1E, 1975HO1F, 1976BI1B). For reactions (a) and (b) see also  ${}^{10}\text{Be}$ . For reaction (c) see (1978RI02). See also (1976MC1E) and (1978BY1B, 1978BY1C; theor.).

## 20. ${}^9\text{Be}(p, p){}^9\text{Be}$

Elastic and inelastic scattering angular distributions have been studied at many energies for  $E_p = 5$  to 1000 MeV [see (1974AJ01)] and (1974YA1C:  $p_0(5.0, 6.0$  MeV),  $p_1(5.2, 6.0$  MeV),  $p_2(4.2 \rightarrow 6.0$  MeV)), (1977CO1G:  $p_0(35.2$  MeV) and (1976AL1G:  $p_0; 1$  GeV). The elastic angular distributions show pronounced diffraction maxima characteristic of the optical model. See (1973MO01) for a discussion of optical model parameters. (1973VO02), in a coupled channels analysis, find that a quadrupole-deformed optical model potential with a deformation parameter  $\beta = 1.1$  provides an improved description of the elastic data and a good fit to the  $p_2$  data.

The structure corresponding to  ${}^9\text{Be}^*(1.7)$  is asymmetric: the line shape peaks  $25_{-11}^{+15}$  keV above the threshold for  ${}^8\text{Be} + n$  (1970TU06). The population of  ${}^9\text{Be}^*(1.7)$  has been studied for  $E_p = 4.2$  to 5.5 MeV: it is consistent with a scattering length of about  $-20$  fm between a neutron and  ${}^8\text{Be}$ . An anomaly at  $E_p = 5.1$  MeV is attributed to FSI between  $n, p$  and  ${}^8\text{Be}$  (1974YA05).

The weighted mean of the values of  $E_x$  listed in (1974AJ01) is  $2432 \pm 3$  keV.  ${}^9\text{Be}^*(3.1)$  has a width  $250 \pm 50$  keV (1960SP08),  $J^\pi = \frac{3}{2}^+, \frac{5}{2}^+$ ,  $E_x = 3.03 \pm 0.03$  MeV (1956BO18),  $3.04 \pm 0.05$  MeV (1960SP08). Higher states are observed at  $E_x = 4.8 \pm 0.2, 6.76 \pm 0.06$  [ $J^\pi = \frac{1}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$  (but see below),  $\Gamma = 1.2 \pm 0.2$  MeV],  $7.94 \pm 0.08$  ( $\Gamma \approx 1$  MeV),  $11.3 \pm 0.2$  MeV ( $\Gamma \approx 1$  MeV),  $14.4 \pm 0.3$  ( $\Gamma \approx 1$  MeV),  $16.7 \pm 0.3, 17.4 \pm 0.3, 19.0 \pm 0.4, 21.1 \pm 0.5$  and  $22.4 \pm 0.7$  MeV [the five highest states are all broad] (1965HA17). (1965JA1A) reports for  ${}^9\text{Be}^*(2.4, 6.8)$   $B(E2\uparrow) = 49 \pm 6$  and  $24 \pm 4$  fm<sup>4</sup> and  $\Gamma(E2\downarrow) = 0.0025$  and  $0.10$  eV, respectively. The strong population of  ${}^9\text{Be}^*(2.4, 6.8)$  is consistent with the assumption that they have  $J^\pi = \frac{5}{2}^-$  and  $\frac{7}{2}^-$ , respectively, and are members of the ground state  $K = \frac{3}{2}^-$  band. See also (1966LA04).

(1974BI14) have studied the elastic scattering at  $E_p = 25$  MeV: a non-zero  $D$  (depolarization) is reported; its existence constitutes evidence of a nucleon-nucleus spin-spin interaction. On the other hand (1976MO1B:  $E_p = 32$  MeV) find no evidence for a non-zero value of  $D$  in the transition to  ${}^9\text{Be}^*(2.4)$ . See also (1977KI04, 1978FR1D), (1974LO1B), (1973WE15, 1974GU13, 1974GU1D, 1974JA1F, 1978BY1B, 1978BY1C, 1978WO13; theor.) and  ${}^{10}\text{B}$ .

21. (a)  ${}^9\text{Be}(p, 2p){}^8\text{Li}$   $Q_m = -16.888$

(b) ${}^9\text{Be}(p, pd){}^7\text{Li}$	$Q_m = -16.6960$
(c) ${}^9\text{Be}(p, p\alpha){}^5\text{He}$	$Q_m = -2.47$
(d) ${}^9\text{Be}(p, pn){}^8\text{Be}$	$Q_m = -1.6652$
(e) ${}^9\text{Be}(p, p{}^3\text{He}){}^6\text{He}$	$Q_m = -21.177$
(f) ${}^9\text{Be}(p, d{}^3\text{He}){}^5\text{He}$	$Q_m = -20.82$

The summed proton spectrum (reaction (a)) shows two peaks with  $Q = -16.4 \pm 0.3$  and  $-25.4 \pm 0.5$  MeV, corresponding to removal of a p-proton and an s-proton respectively, and a third peak of uncertain assignment (perhaps due to unresolved states) with  $Q = -32.3 \pm 0.6$  MeV (1966TY01:  $E_p = 460$  MeV). See also  ${}^8\text{Li}$  and (1973MA67; theor.). For reaction (b) see (1974AJ01). Quasi-free scattering of the incident protons by zero-momentum  $\alpha$ -particle clusters is consistent with the data obtained at  $E_p = 26.0, 35.0, 46.8$  and  $57$  MeV (1972QU01, 1968RO19). At  $E_p = 100$  MeV (1977RO02, 1977CO07) find  $\langle S_\alpha \rangle = 0.45 \pm 0.02$  using DWIA analysis: this value is consistent with that predicted by a simple  $LS$  coupling shell model calculation. See also (1975RO1B, 1978CH1H) and (1975SA01; theor.).

At  $E_p = 12.7$  MeV,  ${}^9\text{Be}^*(3.05)$  is strongly populated and  ${}^9\text{Be}^*(4.70)$  is also observed in reaction (d) (1977JE01). See also Table 9.5 (1978CH07),  ${}^8\text{Be}$  (1977WA05;  $E_p = 45$  and  $47$  MeV), (1974MI05, 1974YA05) and (1974AV02; theor.). See also (1974AJ01). In reaction (f), at  $E_p = 100$  MeV, (1977CO07) find  $\langle S_\alpha \rangle = 0.47 \pm 0.04$ .

## 22. ${}^9\text{Be}(d, d){}^9\text{Be}$

Elastic angular distributions have been studied in the range  $1.1$  to  $410$  MeV [see (1974AJ01)] and at  $E_d = 1.0$  to  $2.2$  MeV (1972LO05),  $E_d = 15$  MeV (1976DA15; also  $d_2, d_4$ ) and  $E_d = 27.98$  MeV (1977OO01). Angular distributions to  ${}^9\text{Be}^*(2.4)$  have also been reported at  $E_d = 11.8, 14.35$  and  $15.8$  MeV: see (1974AJ01).

Inelastic groups have been reported to  ${}^9\text{Be}^*(1.7, 2.4, 3.1, 4.7, 6.8)$ : see (1966LA04). (1968KR02) report  $E_x = 2431.9 \pm 7.0$  keV and  $3040 \pm 15$  keV ( $\Gamma = 294 \pm 20$  keV). See also (1978LO1C), (1977TE1A) and (1974BO39, 1974CH58, 1974GUID, 1977DM1A; theor.).

## 23. (a) ${}^9\text{Be}(t, t){}^9\text{Be}$

(b)  ${}^9\text{Be}(t, nt){}^8\text{Be}$   $Q_m = -1.6652$

The angular distribution of elastically scattered tritons has been measured at  $E_t = 2.10$  MeV (1969HE08, 1970CO04). Reaction (b) at  $E_t = 4.2$  and  $4.6$  MeV proceeds via  ${}^9\text{Be}^*(2.4)$  (1967SE11).

24. (a)  ${}^9\text{Be}({}^3\text{He}, {}^3\text{He}){}^9\text{Be}$   
 (b)  ${}^9\text{Be}({}^3\text{He}, 2\alpha){}^4\text{He}$   $Q_m = 19.0045$

Angular distributions of elastically scattered  ${}^3\text{He}$  particles have been obtained at  $E({}^3\text{He}) = 4$  to 32.3 MeV [see (1974AJ01)], at  $E({}^3\text{He}) = 1.6$  to 2.4 MeV (1974BO38) and 32.6 MeV (1975BU11; polarized  ${}^3\text{He}$ ). See also (1975DA1A). At  $E({}^3\text{He}) = 217$  MeV (1973WI07, 1976WI05) angular distributions have been studied to  ${}^9\text{Be}^*(0, 2.4)$ . For optical model parameters see discussions in (1972BU30, 1973WI07, 1976WI05). At  $E({}^3\text{He}) = 39.8$  MeV inelastic  ${}^3\text{He}$  groups are observed to  ${}^9\text{Be}^*(1.7, 2.4, 3.1, 4.7, 6.8, 14.4)$  (1969BA06). See also (1976RO1L). In a study of reaction (b) the  $({}^5\text{He} + \alpha)$  system is found to be mainly in an S-state: the probability of finding such a cluster in  ${}^9\text{Be}$  is  $N_{\text{eff}} = 0.05$  (1975KA05). See also  ${}^8\text{Be}$ .

25. (a)  ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$   
 (b)  ${}^9\text{Be}(\alpha, 2\alpha){}^5\text{He}$   $Q_m = -2.47$   
 (c)  ${}^9\text{Be}(\alpha, \alpha n){}^8\text{Be}$   $Q_m = -1.6652$

Elastic scattering has been studied at many energies for  $E_\alpha = 8.76$  to 104 MeV [see (1974AJ01)] and at 5.00 to 5.50 MeV (1973GO15). See (1972DE01, 1972DE02) for a discussion of optical model parameters. The structure  $({}^5\text{He} + \alpha)$  for  ${}^9\text{Be}$  is found to be much more probable than  $({}^6\text{Li} + t)$ : the ratio of spectroscopic factors is about 30 (1972DE01, 1972DE02).

Inelastic groups have been observed to  ${}^9\text{Be}^*(1.7, 2.4, 3.1, 6.8, 11.3)$ : see (1966LA04). The angular distribution of the  $\alpha_1$  group to  ${}^9\text{Be}^*(1.7)$  is consistent with  $J^\pi = \frac{1}{2}^+$  (1964LU02:  $E_\alpha = 18.4$  MeV). The angular distributions for  $\alpha_2$  are consistent with  $l = 2$  [ $J^\pi = (\frac{1}{2}, \frac{5}{2}, \frac{7}{2})^-$ ] (1958SU14, 1967FU08). Analysis based on the rotational model leads to a deformation coefficient  $\beta_2 = 0.34 \pm 0.01$  (1964GR39).

The summed  $\alpha$ -spectra from reaction (b) show a peak corresponding to quasi-elastic scattering leaving  ${}^5\text{He}$  in its ground state. The angular distribution peaks at the angle corresponding to zero-momentum  $\alpha$ -cluster. The probability of formation of such clusters is  $(7_{-5}^{+13})\%$  (1969PI11). See also (1977WA1M). (1976WO11) find  $S_\alpha = 0.96$ .

A study of continuum neutrons at  $E_\alpha = 4.9$  to 6.4 MeV (reaction (c)) indicates that sequential decay takes place via  ${}^9\text{Be}^*(1.7, 2.4, 3.1)$  (1972OB01). See also (1974AJ01, 1975RO1B) and (1974CH58, 1974PI11, 1974LO09, 1975KU1K, 1976AV05, 1977DE1K, 1977DM1A, 1977MO11; theor.).

26. (a)  ${}^9\text{Be}({}^6\text{Li}, {}^6\text{Li}){}^9\text{Be}$   
 (b)  ${}^9\text{Be}({}^7\text{Li}, {}^7\text{Li}){}^9\text{Be}$

Elastic angular distributions have been measured at  $E({}^6\text{Li}) = 4$  and 6 MeV (1974VO06) and 24 MeV (1968DA20) and at  $E({}^7\text{Li}) = 24$  MeV (1972WE08) and 34 MeV (1977KE09).

27.  ${}^9\text{Be}({}^9\text{Be}, {}^9\text{Be}){}^9\text{Be}$

Elastic angular distributions have been obtained at  $E = 5, 9, 12$  and  $16$  MeV (1977YO02).

28.  ${}^9\text{Be}({}^{12}\text{C}, {}^{12}\text{C}){}^9\text{Be}$

Elastic angular distributions are reported at  $E({}^{12}\text{C}) = 12, 15, 18$  and  $21$  MeV (1970BA49) and  $E({}^9\text{Be}) = 14, 20$  and  $26$  MeV (1978LA1G) and  $43$  MeV (1977MA1L). The neutron spectroscopic factor for  ${}^9\text{Be}$  is  $0.84$  (1970BA49). For spallation studies see (1977IG1D, 1978CH02). See also (1976EC1A, 1977PE1B, 1978MA1L), (1978EN06) and (1978PA1B; theor.).

29.  ${}^9\text{Be}({}^{14}\text{N}, {}^{14}\text{N}){}^9\text{Be}$

Elastic angular distributions have been measured at  $E({}^{14}\text{N}) = 25$  (1966OE1A) and  $27.3$  MeV (1959HA28).

30. (a)  ${}^9\text{Be}({}^{16}\text{O}, {}^{16}\text{O}){}^9\text{Be}$   
(b)  ${}^9\text{Be}({}^{18}\text{O}, {}^{18}\text{O}){}^9\text{Be}$

Elastic angular distributions have been reported at  $E({}^{16}\text{O}) = 15, 18, 21.5$  and  $25$  MeV (1970BA49),  $25.7$  to  $29.5$  MeV (1975FU05) and  $30$  MeV (1969KR03), and at  $E({}^{18}\text{O}) = 12.1, 16$  and  $20$  MeV (1971KN05). See also (1974TA1C, 1978CH02), (1978EN06) and (1978PA1B; theor.).

31. (a)  ${}^9\text{Be}({}^{24}\text{Mg}, {}^{24}\text{Mg}){}^9\text{Be}$   
(b)  ${}^9\text{Be}({}^{27}\text{Al}, {}^{27}\text{Al}){}^9\text{Be}$   
(c)  ${}^9\text{Be}({}^{28}\text{Si}, {}^{28}\text{Si}){}^9\text{Be}$

Elastic angular distributions have been measured at  $E({}^9\text{Be}) = 14, 20$  and  $26$  MeV for reactions (a), (b) and (c) (1977BA71) and at  $14$  and  $17$  MeV for reaction (c) (1977EC04). See also (1978DE1M) and (1978EN06).

32.  ${}^{10}\text{Be}(\text{d}, \text{t}){}^9\text{Be}$

$$Q_m = -0.5545$$

Forward angular distributions have been obtained at  $E_d = 15.0$  MeV for the tritons to  ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1)$ . The ground state transition is well fitted by  $l = 1$ . The transition to  ${}^9\text{Be}^*(1.7)$  [ $\Gamma \approx 165 \pm 25$  keV] is consistent with  $J^\pi = \frac{1}{2}^+$ , that to  ${}^9\text{Be}^*(2.4)$  is quite well fitted with  $l = 3$  [ $J^\pi = \frac{5}{2}^-$ ], and that to  ${}^9\text{Be}^*(3.1)$  [ $\Gamma = 280 \pm 25$  keV] is consistent with  $l = 2$ . No other narrow states are seen up to  $E_x = 5.5$  MeV (1970AU02).

$$\begin{aligned} 33. \text{ (a) } {}^{10}\text{B}(\gamma, \text{p}){}^9\text{Be} & \quad Q_m = -6.5853 \\ \text{ (b) } {}^{10}\text{B}(\text{e}, \text{ep}){}^9\text{Be} & \quad Q_m = -6.5853 \end{aligned}$$

See  ${}^{10}\text{B}$ .

$$34. {}^{10}\text{B}(\text{n}, \text{d}){}^9\text{Be} \quad Q_m = -4.3607$$

A good fit to the angular distributions of the deuterons to  ${}^9\text{Be}^*(0, 2.4)$  has been obtained at  $E_n = 14.4$  MeV, using DWBA. The spectroscopic factors are in close agreement with shell model predictions (1965VA05). See also  ${}^{11}\text{B}$  in (1980AJ01).

$$35. {}^{10}\text{B}(\text{p}, 2\text{p}){}^9\text{Be} \quad Q_m = -6.5863$$

The summed proton spectrum at  $E_p = 460$  MeV yields  $Q = -6.7 \pm 0.5, -11.9 \pm 0.5, -17.1 \pm 0.6$  (all  $l \neq 0$ ) and  $Q = -30.5 \pm 0.6$  MeV ( $l = 0$ ) (1966TY01). See also  ${}^{10}\text{B}$ .

$$36. {}^{10}\text{B}(\text{d}, {}^3\text{He}){}^9\text{Be} \quad Q_m = -1.0918$$

Angular distributions of the  ${}^3\text{He}$  groups corresponding to  ${}^9\text{Be}^*(0, 2.4)$  have previously been measured at  $E_d = 11.8$  and 28 MeV: see (1974AJ01). Recent angular distribution measurements at  $E_d = 15$  MeV (1974LU06; to  ${}^9\text{Be}^*(0, 2.4)$ ) and at  $E_d = 52$  MeV (1975SC41; to  ${}^9\text{Be}^*(0, 2.4, 6.7)$ ) lead to  $S = 0.72$  and 0.82 (1974LU06) and relative  $S = 1.00, 0.98$  and 0.58 (1975SC41). In the latter experiment  ${}^9\text{Be}^*(11.3)$  appears also to be strongly populated. The data for  ${}^9\text{Be}^*(0, 2.4)$  are very similar, as predicted, to those obtained in the mirror reaction: see reaction 12 in  ${}^9\text{B}$ . See also (1974DA1D; theor.) and  ${}^{12}\text{C}$  in (1980AJ01).

$$37. {}^{10}\text{B}(\text{t}, \alpha){}^9\text{Be} \quad Q_m = 13.2287$$

At  $E_t = 12.9$  MeV  $\alpha$ -groups are observed to the ground state of  ${}^9\text{Be}$  and to excited states at  $E_x = 1.75 \pm 0.03, 2.43, 3.02 \pm 0.04$  ( $\Gamma = 320 \pm 60$  keV),  $11.27 \pm 0.04$  ( $\Gamma = 530 \pm 70$  keV), (14.4) [ $\Gamma \approx 800$  keV], 14.39 and 16.67 MeV. The  $T = \frac{3}{2}$  state  ${}^9\text{Be}^*(14.39)$  is very weakly populated [ $\approx 5\%$  of intensity of  $\alpha_2$ ]. The angular distribution of the  $\alpha_2$  group shows sharp forward and backward peaking. The  $\alpha_0$  group is not peaked in the backward direction (1968AJ01). A study at  $E_t = 1.0$  to 3.2 MeV finds  $E_x = 1.750 \pm 0.025$  MeV,  $\Gamma = 220 \pm 8$  keV. The angular distribution of the corresponding  $\alpha$ -group has been determined at  $E_t = 2.5$  MeV (1971GE09). See also (1977CI1A) and  ${}^{13}\text{C}$  in (1981AJ01).

$$38. {}^{10}\text{B}({}^7\text{Li}, {}^8\text{Be}){}^9\text{Be} \quad Q_m = 10.670$$

See (1977KO27).

$$39. {}^{10}\text{B}({}^{14}\text{N}, {}^{15}\text{O}){}^9\text{Be} \quad Q_m = 0.712$$

The ground state angular distribution has been measured at  $E({}^{14}\text{N}) = 27.5$  MeV (1962NE01).

$$40. {}^{10}\text{B}({}^{16}\text{O}, {}^{17}\text{F}){}^9\text{Be} \quad Q_m = -5.985$$

See (1968OK06).

$$41. {}^{11}\text{B}(\text{n}, \text{t}){}^9\text{Be} \quad Q_m = -9.5586$$

The angular distribution of  $t_0$  has been measured at  $E_n = 14.4$  MeV (1970MI14).

$$42. \text{(a) } {}^{11}\text{B}(\text{p}, {}^3\text{He}){}^9\text{Be} \quad Q_m = -10.3224$$

$$\text{(b) } {}^{11}\text{B}(\text{p}, \text{pd}){}^9\text{Be} \quad Q_m = -15.8159$$

At  $E_p = 45$  MeV angular distributions are reported for the  ${}^3\text{He}$  ions corresponding to  ${}^9\text{Be}^*(0, 2.4, 11.8, 13.8, 14.39 [T = \frac{3}{2}], 15.96 \pm 0.04 [T = \frac{1}{2}])$ . In addition one or more states may be located at  ${}^9\text{Be}^*(15.13)$ . It is suggested that  ${}^9\text{Be}^*(11.8, 13.8, 15.96)$  are the  $J^\pi = \frac{3}{2}^-, T = \frac{1}{2}$  analogs to  ${}^9\text{B}^*(12.06, 14.01, 16.02)$  (1971HA10). Angular distributions are also reported at  $E_p = 40$  MeV (1971KA21:  $\alpha_0, \alpha_2$ ). The intensity of the group to  ${}^9\text{Be}^*(3.1)$  is  $\approx 1\%$  of the ground state group (1971KA21). The excitation energy of the first  $T = \frac{3}{2}$  state is  $E_x = 14392.6 \pm 1.8$  keV (1974KA15) using the (1977WA08) mass for  ${}^9\text{Be}_{\text{g.s.}}$ . For reaction (b) see (1964BA1C).

43. (a)  $^{11}\text{B}(\text{d}, \alpha)^9\text{Be}$   $Q_{\text{m}} = 8.0308$   
 (b)  $^{11}\text{B}(\text{d}, \text{n}\alpha)^4\text{He}^4\text{He}$   $Q_{\text{m}} = 6.4575$

Alpha groups are reported corresponding to  $^9\text{Be}^*(0, 1.7, 2.4, 3.1)$ . The width of  $^9\text{Be}^*(1.7)$  [ $E_{\text{x}} = 1.70 \pm 0.01$  MeV (1975FO13)] is  $\Gamma_{\text{c.m.}} = 224 \pm 25$  (1958KA31, 1966PU02),  $200 \pm 10$  keV (1975FO13). The weighted mean of the values of  $E_{\text{x}}$  of  $^9\text{Be}^*(2.4)$ , reported in (1974AJ01), is  $2425 \pm 3$  keV. The  $\frac{5}{2}^+$  state is at  $E_{\text{x}} = 3.02 \pm 0.03$  MeV (1955LE36),  $3.05 \pm 0.03$  MeV (1956BO18):  $\Gamma_{\text{c.m.}} = 257 \pm 25$  keV (1958KA31, 1966PU02). The ratio  $\Gamma_{\gamma}/\Gamma$  of  $^9\text{Be}^*(1.7) \leq 2.4 \times 10^{-5}$ , that for  $^9\text{Be}^*(2.4)$  is  $(1.16 \pm 0.14) \times 10^{-4}$  (1966PU02). Since  $\Gamma_{\gamma}$  is known from (e, e') [see Table 9.8:  $0.089 \pm 0.010$  eV],  $\Gamma = 0.77 \pm 0.15$  keV.

Angular distributions for  $\alpha_0$  and  $\alpha_2$  are reported at  $E_{\text{d}} = 0.39$  to  $2.5$  MeV and at  $12$  MeV [see (1974AJ01)] and at  $3.0$  to  $3.9$  MeV (1974CA34, 1975CA1J, 1975CA1K). Reaction (b), at  $E_{\text{d}} = 10.4$  and  $12.0$  MeV, proceeds via  $^9\text{Be}^*(2.4)$  and to some extent via  $^9\text{Be}^*(3.1, 4.7)$  and possibly some higher excited states. The dominant decay of  $^9\text{Be}^*(2.4)$  is to  $^5\text{He}(0) + \alpha$  while  $^9\text{Be}^*(3.1, 4.7)$  decay to  $^8\text{Be}(0) + \text{n}$  (1971RE19). It should be noted, however, that the peaks corresponding to  $^9\text{Be}^*(3.1)$  have a FWHM of  $\approx 1$  MeV, which may imply that  $^9\text{Be}^*(2.8)$  is involved (1971RE19). See also (1978GR07) and  $^{13}\text{C}$  in (1981AJ01).

44.  $^{11}\text{B}(^{16}\text{O}, ^{18}\text{F})^9\text{Be}$   $Q_{\text{m}} = -8.290$

See (1968OK06).

45. (a)  $^{12}\text{C}(\text{n}, \alpha)^9\text{Be}$   $Q_{\text{m}} = -5.7015$   
 (b)  $^{12}\text{C}(\text{n}, \text{n}\alpha)^4\text{He}^4\text{He}$   $Q_{\text{m}} = -7.2748$

Angular distributions of the  $\alpha_0$  group have been measured at  $E_{\text{n}} = 13.9$  to  $18.8$  MeV;  $^9\text{Be}^*(1.7, 2.4, 3.1)$  are also populated: see (1974AJ01). Reaction (b) at  $E_{\text{n}} = 13$  to  $18$  MeV involves  $^9\text{Be}^*(2.4)$  (1966MO05). See also (1975AN01, 1976AN1A), (1973PE19; theor.) and  $^{12}\text{C}$  and  $^{13}\text{C}$  in (1980AJ01, 1981AJ01).

46.  $^{12}\text{C}(\alpha, ^7\text{Be})^9\text{Be}$   $Q_{\text{m}} = -24.693$

Angular distributions have been obtained for the transition to  $^7\text{Be}^*(0, 0.43)^+ ^9\text{Be}_{\text{g.s.}}$  (1972RU03:  $42$  MeV).

47.  $^{12}\text{C}(^{10}\text{B}, ^{13}\text{N})^9\text{Be}$   $Q_{\text{m}} = -4.642$

See (1975RA13; theor.).

$$48. {}^{12}\text{C}({}^{18}\text{O}, {}^{21}\text{Ne}){}^9\text{Be} \quad Q_m = -6.398$$

See (1974BA15).

$$49. {}^{13}\text{C}(\gamma, \alpha){}^9\text{Be} \quad Q_m = -10.6479$$

See  ${}^{13}\text{C}$  in (1981AJ01).

$$50. {}^{13}\text{C}(\text{p}, \text{p}\alpha){}^9\text{Be} \quad Q_m = -10.6479$$

See (1975OB01) and  ${}^{14}\text{N}$  in (1981AJ01).

$$51. {}^{13}\text{C}(\text{d}, {}^6\text{Li}){}^9\text{Be} \quad Q_m = -9.174$$

See (1966DE09).

$$52. {}^{13}\text{C}({}^3\text{He}, {}^7\text{Be}){}^9\text{Be} \quad Q_m = -9.062$$

Angular distributions have been obtained at  $E({}^3\text{He}) = 70$  MeV for the transitions to  ${}^9\text{Be}^*(0, 2.4)$  and  ${}^7\text{Be}^*(0, 0.43)$ . Broad states at  $2.9$ ,  $4.8 \pm 0.2$ ,  $7.3 \pm 0.2$  and  $11.9 \pm 0.4$  MeV are also populated (1976ST11).

$$53. {}^{13}\text{C}(\alpha, {}^8\text{Be}){}^9\text{Be} \quad Q_m = -10.740$$

At  $E_\alpha = 65$  MeV angular distributions have been measured for the transitions to  ${}^9\text{Be}^*(0, 2.4)$  (1976WO11).

$$54. \text{(a) } {}^{13}\text{C}({}^{17}\text{O}, {}^{21}\text{Ne}){}^9\text{Be} \quad Q_m = -3.300$$

$$\text{(b) } {}^{13}\text{C}({}^{18}\text{O}, {}^{22}\text{Ne}){}^9\text{Be} \quad Q_m = -0.980$$

See (1974CH1Q) and (1978EN06).

$$55. \text{}^{14}\text{C}(\text{}^{18}\text{O}, \text{}^{23}\text{Ne})\text{}^9\text{Be} \quad Q_{\text{m}} = -3.956$$

See (1974BA15) and (1978EN06).

$$56. \text{}^{16}\text{O}(\alpha, \text{}^{11}\text{C})\text{}^9\text{Be} \quad Q_{\text{m}} = -24.310$$

See (1972RU03).

## <sup>9</sup>B

(Figs. 17 and 18)

GENERAL: (See also (1974AJ01).)

*Model calculations:* (1977HO1F, 1977OK01, 1978HO1E).

*Special levels:* (1974IR04, 1975WI1E, 1976IR1B, 1978HO1E).

*Astrophysical questions:* (1977SI1D).

*Pion reactions:* (1974KA07).

*Other topics:* (1974HA1C, 1974IR04, 1976IR1B).

*Ground state properties:* (1975BE31, 1977OK01).

1. (a) ${}^6\text{Li}({}^3\text{He}, n){}^8\text{B}$	$Q_m = -1.975$	$E_b = 16.603$
(b) ${}^6\text{Li}({}^3\text{He}, p){}^8\text{Be}$	$Q_m = 16.7878$	
(c) ${}^6\text{Li}({}^3\text{He}, d){}^7\text{Be}$	$Q_m = 0.113$	
(d) ${}^6\text{Li}({}^3\text{He}, t){}^6\text{Be}$	$Q_m = -4.306$	
(e) ${}^6\text{Li}({}^3\text{He}, {}^3\text{He}){}^6\text{Li}$		
(f) ${}^6\text{Li}({}^3\text{He}, p){}^4\text{He}{}^4\text{He}$	$Q_m = 16.8797$	

The total cross section for reaction (a) increases monotonically from threshold to  $\approx 7$  mb at 3.8 MeV. It then decreases monotonically from  $\approx 20$  mb at  $E({}^3\text{He}) = 5.5$  MeV to 7.6 MeV (1975MC02, 1974MCZS) and the decrease continues in the range  $E({}^3\text{He}) = 8.9$  to 26.5 MeV (1973MA24). See also <sup>8</sup>B.

The excitation functions for protons leading to  ${}^8\text{Be}^*(0, 2.9)$  [ $p_0, p_1$ ] have been measured for  $E({}^3\text{He}) = 0.9$  to 17 MeV (reaction (b)): see (1974AJ01). Resonances are reported at  $E({}^3\text{He}) = 1.6$  MeV [ $J^\pi = \frac{3}{2}^-$  or  $\frac{5}{2}^-$  (1969VI05),  $\Gamma = 0.25$  MeV] and 3.0 MeV ( $\Gamma = 1.5$  MeV) (1956SC01). Total cross sections have been measured for the  $p_1$  and  $p_2$  groups for  $E({}^3\text{He}) = 3 - 6$  MeV (1976GO1F). Polarization measurements are reported at  $E({}^3\text{He}) = 14$  MeV (1977AS04, 1977IR01:  $p_0, p_1$ ). See also <sup>8</sup>Be.

The yields of 0.43 and 0.48 MeV  $\gamma$ -rays (reaction (c)) for  $E({}^3\text{He}) = 0.5$  to 1.3 MeV are reported to show the excitation of  ${}^9\text{B}^*(17.20 \pm 0.02)$  with  $\Gamma = 110 \pm 30$  keV,  $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+$ ;  $T = \frac{1}{2}$  (1970AL25). See also <sup>7</sup>Be and (1976KA1D).

Excitation functions for  $t_0$  (reaction (d)) have been measured for  $E({}^3\text{He}) = 10$  to 16 and 23.3 to 25.4 MeV: see (1974AJ01). See also <sup>6</sup>Be. For reaction (e) see (1972GI07). See also <sup>6</sup>Li. The total cross section for reaction (f) has been measured for  $E({}^3\text{He}) = 3$  to 6 MeV (1976GO1F). The yield of  $\alpha$ - $\alpha$  coincidences [to  ${}^5\text{Li}_{g.s.}$ ] has been measured for  $E({}^3\text{He}) = 1.4$  to 1.8 MeV: a resonance is observed at  $1.57 \pm 0.02$  MeV [ ${}^9\text{B}^*(17.63)$ ],  $\Gamma = 70 \pm 20$  keV (1978GU15).

Table 9.9: Energy levels of  ${}^9\text{B}$ 

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	$0.54 \pm 0.21$	p, $\alpha$	2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17
(1.6) $2.361 \pm 5$	$\frac{5}{2}^-; \frac{1}{2}$	$\approx 700$ $81 \pm 5$	(p, $\alpha$ ) $\alpha$	13 2, 4, 6, 7, 8, 11, 12, 13, 15, 16
$2.788 \pm 30$ ( $4.8 \pm 100$ )	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	$550 \pm 40$ $1000 \pm 200$	p	4, 6, 11, 13, 16 4, 9
$6.97 \pm 60$	$\frac{7}{2}^-; \frac{1}{2}$	$2000 \pm 200$	p	4, 6, 9, 11, 15, 16
$11.70 \pm 70$	$(\frac{7}{2})^-; \frac{1}{2}$	$800 \pm 50$	p	9, 11, 13
$12.06 \pm 60$	$;$ $\frac{1}{2}$	$800 \pm 200$	p	4, 9, 15
$14.01 \pm 70$	$;$ $\frac{1}{2}$	$390 \pm 110$		4, 15
$14.6550 \pm 2.5$	$\frac{3}{2}^-; \frac{3}{2}$	$0.275 \pm 0.093$	$\gamma$ , p	4, 7, 15
$14.7 \pm 180$	$(\frac{5}{2})^-; \frac{1}{2}$	$1350 \pm 200$		11
$15.29 \pm 40$	$;$ $\frac{1}{2}$			15
$15.58 \pm 40$	$;$ $\frac{1}{2}$			15
$16.024 \pm 25$	$;$ $(\frac{1}{2})$	$180 \pm 16$		4, 15
$17.076 \pm 4$	$T = \frac{3}{2}$	$22 \pm 5$		15
$17.190 \pm 25$		$120 \pm 40$	p, d, ${}^3\text{He}$	1, 4, 5, 15
$17.637 \pm 10$		$71 \pm 8$	p, d, ${}^3\text{He}$ , $\alpha$	1, 4, 5, 15
(18.6)		1000	p, ${}^3\text{He}$	1, 11

 2.  ${}^6\text{Li}(\alpha, n){}^9\text{B}$ 

$$Q_m = -3.975$$

Angular distributions at  $E_\alpha = 8.0, 10.0, 12.0$  and  $14.0$  MeV all display strong forward peaking (1963ME08). At  $E_\alpha = 14.4$  MeV, neutron groups are observed to  ${}^9\text{B}^*(0, 2.4)$ : the upper limit of the cross section to a state at  $\approx 1.7$  MeV is  $100 \mu\text{b/sr}$  or  $< 0.1$  of the ground state group (1964BA29). See also  ${}^{10}\text{B}$ .

 3.  ${}^6\text{Li}({}^6\text{Li}, t){}^9\text{B}$ 

$$Q_m = 0.809$$

Angular distributions of the  $t_0$  group have been measured for  $E({}^6\text{Li}) = 4.0$  to  $5.5$  MeV and at  $7.35$  and  $9.0$  MeV. No evidence was observed for a group corresponding to  ${}^9\text{B}^*(1.6)$  (1966KI09).

4. (a)  ${}^7\text{Li}({}^3\text{He}, n){}^9\text{B}$   $Q_m = 9.352$   
 (b)  ${}^7\text{Li}({}^3\text{He}, np){}^8\text{Be}$   $Q_m = 9.537$

For  $E({}^3\text{He})$  to 12.5 MeV this reaction populates  ${}^9\text{B}^*(0, 2.4, 2.8, (7.0))$  (1963DU12) and levels at  $12.06 \pm 0.06$  [ $0.8 \pm 0.2$ ],  $14.01 \pm 0.07$  [ $0.39 \pm 0.11$ ],  $14.670 \pm 0.016$  [ $< 0.045$ ],  $16.024 \pm 0.025$  [ $0.180 \pm 0.016$ ], 17.19 and 17.63 MeV (1965DI03) [widths in brackets].

${}^9\text{B}^*(14.66)$  is the first  $T = \frac{3}{2}$  state in  ${}^9\text{B}$ . Its decay properties are displayed in Table 9.6 and compared with those of  ${}^9\text{Be}^*(14.40)$  (1971AD01, 1972AD04, 1976MC10).

The  $n_0$  angular distribution has been measured at  $E({}^3\text{He}) = 3.1$  MeV. A state at  $E_x = 4.8 \pm 0.1$  MeV,  $\Gamma = 1.0 \pm 0.2$  MeV is also reported (1970GU08). See also (1976RA1G, 1976SL1A) and  ${}^{10}\text{B}$ .

5.  ${}^7\text{Be}(d, p){}^8\text{Be}$   $Q_m = 16.675$   $E_b = 16.490$

For  $E_d = 0.75$  to 1.70 MeV, resonances in the yields of protons are observed at  $E_d = 0.900 \pm 0.025$  MeV ( $p_0, p_1$ ) and  $1.475 \pm 0.010$  MeV ( $p_1$  only) with  $\Gamma_{\text{c.m.}} = 120 \pm 40$  and  $71 \pm 8$  keV, respectively [ ${}^9\text{B}^* = 17.19$  and 17.64 MeV] (1960KA17).

6. (a)  ${}^9\text{Be}(p, n){}^9\text{B}$   $Q_m = -1.850$   
 (b)  ${}^9\text{Be}(p, pn){}^8\text{Be}$   $Q_m = -1.6652$

At  $E_p = 20$  MeV,  ${}^9\text{B}^*(0, 2.4)$  are populated. The results are consistent with the excitation of  ${}^9\text{B}^*(2.8)$  [ $\Gamma \approx 0.3$  MeV] and  ${}^9\text{B}^*(7.0)$  [ $\Gamma > 1$  MeV]. No other states are excited for  $E_x < 7.1$  MeV (1970AN07). Additional states have been reported by a number of groups: see (1974AJ01) and (1978CH07). The width of the ground state is  $540 \pm 210$  eV (1964TE01).

Angular distributions have been reported at many energies in the range  $E_p = 3.5$  to 49.3 MeV [see (1974AJ01)] and at 22.8 MeV (1975CA18: quasi elastic). See also (1978WA1D). A study of reaction (b) at  $E_p = 5.5$  MeV is said to suggest a  $(19 \pm 5)\%$  decay of  ${}^9\text{B}^*(2.4)$  to  ${}^8\text{Be}_{\text{g.s.}}$  (1978CH07): see, however, reaction 13. See also (1974BO1H, 1976AN1B, 1976CA17, 1977JE01, 1977LO10, 1978BY1B, 1978BY1C, 1978GO1G, 1978LO1C) and  ${}^{10}\text{B}$ .

7.  ${}^9\text{Be}({}^3\text{He}, t){}^9\text{B}$   $Q_m = -1.0860$

Angular distributions have been measured in the range  $E({}^3\text{He}) = 3.0$  to 25 MeV [see (1974AJ01)] and at 217 MeV (1976WI05:  $t_0, t_1$ ). At  $E({}^3\text{He}) = 39.8$  MeV,  ${}^9\text{B}_{\text{g.s.}}$  is strongly populated and  ${}^9\text{B}^*(2.4, 14.7)$  are also observed. There is some indication that other known  ${}^9\text{B}$  states are also populated (1969BA06). See also (1976UE01), (1974LO1B) and  ${}^{12}\text{C}$  in (1975AJ02, 1980AJ01).

Table 9.10: Delayed protons following the  $\beta^+$  decay of  ${}^9\text{C}$  <sup>a</sup>

$E_p$ (c.m.) (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	Corresponding state in ${}^9\text{B}$ (MeV)	
		if decay is to ${}^8\text{Be}(\text{g.s.})$	if decay is to ${}^8\text{Be}^*(2.9)$
$3.45 \pm 0.25$	$200 \pm 100$	$3.26 \pm 0.25$ <sup>d</sup>	c
$(4.2 \pm 0.3)$	$1000 \pm 200$	$4.0 \pm 0.3$	$6.9 \pm 0.3$
$(5.0 \pm 0.2)$	$400 \pm 200$	$4.8 \pm 0.2$	c
$6.10 \pm 0.10$	$400 \pm 100$	$5.91 \pm 0.10$	c
$9.28 \pm 0.24$ <sup>b</sup>	$1800 \pm 200$	$9.09 \pm 0.24$	$11.99 \pm 0.24$
$12.30 \pm 0.10$ <sup>b</sup>	$450 \pm 100$	$12.11 \pm 0.10$	c

<sup>a</sup> (1972ES05).


<sup>b</sup> Ratio of the intensities  $I_{9.28}/I_{12.30} = 1.2 \pm 0.2$ .

<sup>c</sup> The relatively narrow width of the proton group does not permit this option.

<sup>d</sup> By analogy with the  ${}^9\text{Li}$  decay, this decay may involve a  $J^\pi = \frac{1}{2}^-$  analog of  ${}^9\text{Be}^*(2.78)$ . Such a state in  ${}^9\text{B}$  has not been reported in any other reaction.



At  $E({}^6\text{Li}) = 30.8$  and  $31.8$  MeV the ground state of  ${}^9\text{B}$  is strongly excited.  ${}^9\text{B}^*(2.4)$  is also observed (1970CH19, 1971CH1B). A partial angular distribution for the ground state transition is reported at the higher energy by (1971CH1B).



Several groups of delayed protons are observed indicating the involvement of a number of  ${}^9\text{B}$  states: see Table 9.10 (1972ES05). It is not possible to determine  $ft$  values since some of the  ${}^9\text{B}$  states involved in the  ${}^9\text{C}$  decay via  ${}^5\text{Li} + \alpha$ : see (1972ES05).



See  ${}^{10}\text{B}$ .




Table 9.11: Levels of  ${}^9\text{B}$  from  ${}^{10}\text{B}(\text{p}, \text{d}){}^9\text{B}$

$E_x$ <sup>a</sup> (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ <sup>a</sup> (MeV)	$l_n$ <sup>b</sup>	$J\pi$ <sup>c</sup>
0		1	$\frac{3}{2}^-$
$2.35 \pm 20$ (2.8) <sup>d</sup>		1	$\frac{5}{2}^-$
$7.1 \pm 140$	$2.15 \pm 0.15$	1	$\frac{7}{2}^-$
$11.70 \pm 70$ <sup>b,e</sup>	$0.80 \pm 0.05$	1	$(\frac{7}{2})^-$
$14.7 \pm 180$ (18.4) <sup>f</sup>	$1.35 \pm 0.2$	1	$(\frac{5}{2})^-$

<sup>a</sup> (1968KU04):  $E_p = 33.6$  MeV.

<sup>b</sup> (1968KU04) and (1969BA05):  $E_p = 155.6$  MeV.

<sup>c</sup>  $J$  from best fit to theoretical spectroscopic factor (1968KU04).

<sup>d</sup> Weak group.

<sup>e</sup> (1970SQ01):  $E_p = 49.5$  MeV.

<sup>f</sup> (1969BA05).

Observed groups are displayed in Table 9.11 (1968KU04, 1969BA05, 1970SQ01). See also (1977GU14;  $\alpha$  to  ${}^9\text{B}^*(0, 2.36)$ ;  $E_p = 16.7$  and  $17.7$  MeV).

12.  ${}^{10}\text{B}(\text{d}, \text{t}){}^9\text{B}$   $Q_m = -2.179$

Angular distributions have been measured at  $E_d = 11.8, 13.5$  and  $28$  MeV [see (1974AJ01)], at  $13.6$  MeV (1973ZA06:  $t_0$ ) and at  $E_d = 15$  MeV (1974LU06:  $t_0, t_1$ ).  $S$  for  ${}^9\text{B}^*(0, 2.4)$  are  $0.66$  and  $0.58$ , respectively (1974LU06). See also (1975ZA06), (1974DA1D; theor.) and  ${}^{12}\text{C}$  in (1975AJ02, 1980AJ01).

13. (a)  ${}^{10}\text{B}({}^3\text{He}, \alpha){}^9\text{B}$   $Q_m = 12.142$

(b)  ${}^{10}\text{B}({}^3\text{He}, \alpha\text{p}){}^8\text{Be}$   $Q_m = 12.3273$

Alpha-particle spectra show the excitation of  ${}^9\text{B}^*(0, 2.4, 2.8, 11.8)$ : see (1966LA04). Measurements by (1968KR02) determine  $E_x = 2.361 \pm 0.005$  and  $2.788 \pm 0.030$  MeV,  $\Gamma = 81 \pm 5$  and  $548 \pm 40$  keV, respectively [see Table 9.11 in (1966LA04) for other values]. There is some evidence for a state with  $E_x \approx 1.6$  MeV,  $\Gamma \approx 0.7$  MeV, but it is not conclusive, in agreement with the older work. No evidence is found for any narrow levels in  ${}^9\text{B}$  with  $\Gamma \leq 100$  keV and

$4 < E_x < 7$  MeV: the upper limit to the intensity of the corresponding  $\alpha$ -group is 1% of the intensity of the group to  ${}^9\text{B}^*(2.4)$  (1968KR02). Angular distributions have been determined at  $E({}^3\text{He}) = 5.5$  MeV (1966CA02:  $\alpha_0$ ) and 33.7 MeV (1971SQ03:  $\alpha_0, \alpha_{2.4}, \alpha_{11.8}$ ). DWBA does not seem to give a good description of the transition to  ${}^9\text{B}^*(11.8)$  (1971SQ03).

In reaction (b) study of the decays of  ${}^9\text{B}^*(2.4, 2.8)$  shows that  ${}^9\text{B}^*(2.4)$  decays  $< 0.5\%$  by proton emission to  ${}^8\text{Be}(0)$  [ $\theta_f^2 < 5.1 \times 10^{-3}$ ] [it decays to  ${}^5\text{Li}(0)$  by  $\alpha$ -emission] while the second state,  $E_x = 2.71 \pm 0.03$  MeV [ $\Gamma = 0.71 \pm 0.06$  MeV] decays almost 100% by that channel [ $\theta_d^2 = 0.74$ ] (1966WI08). No other excited states of  ${}^9\text{B}$  with  $3.5 < E_x < 9.5$  MeV decay by proton emission to  ${}^8\text{Be}(0)$  (1968KR02). A number of additional states of  ${}^9\text{B}$  are reported by (1974FO08). See also (1974LO1B) and (1974AJ01).

$$14. {}^{10}\text{B}({}^{16}\text{O}, {}^{17}\text{O}){}^9\text{B} \quad Q_m = -4.292$$

See (1968OK06).

$$15. {}^{11}\text{B}(\text{p}, \text{t}){}^9\text{B} \quad Q_m = -11.409$$

At  $E_p = 45$  MeV angular distributions have been obtained for the triton groups to  ${}^9\text{B}^*(0, 2.36, 12.06, 14.01, 14.66, 16.02)$ . In addition the spectra show some indication of the groups corresponding to  ${}^9\text{B}^*(7.0, 17.19, 17.64)$ .  $T = \frac{1}{2}$  states are reported at  $E_x = 15.29 \pm 0.04$  and  $15.58 \pm 0.04$  MeV (1971HA10). The first two  $T = \frac{3}{2}$  states have been observed at  $E_x = 14.6550 \pm 0.0025$  (1974KA15) and  $17.076 \pm 0.004$  MeV [ $\Gamma = 22 \pm 5$  keV] (1974BE66). See also (1975BE56) and reaction 42 in  ${}^9\text{Be}$ .

$$16. \text{(a) } {}^{12}\text{C}(\text{p}, \alpha){}^9\text{B} \quad Q_m = -7.552$$

$$\text{(b) } {}^{12}\text{C}(\text{p}, \text{p}){}^4\text{He}{}^4\text{He} \quad Q_m = -7.2748$$

Angular distributions have been measured for the  $\alpha_0$  group at  $E_p = 14.0$  to 54.1 MeV. Alpha groups are also observed to  ${}^9\text{B}^*(2.3, 2.9 \pm 0.2, 6.97 \pm 0.06)$ : see (1974AJ01). The angular distribution to  ${}^9\text{B}^*(6.97)$  is consistent with  $J^\pi = \frac{7}{2}^-$ ,  $\Gamma \approx 2$  MeV (1972MA21:  $E_p = 54.1$  MeV). Angular distributions involving the  $\alpha_0$  and  $\alpha^*$  groups [to  ${}^4\text{He}^*(20.1), 0^+$ ] to  ${}^9\text{B}_{g.s.}$  have been studied at  $E_p = 45.2$  MeV (1978DA1C). For reaction (b) see (1974AJ01). See also  ${}^{12}\text{C}$  in (1975AJ02, 1980AJ01),  ${}^{13}\text{N}$  in (1976AJ04) and (1978HA04; theor.).

$$17. {}^{12}\text{C}({}^3\text{He}, {}^6\text{Li}){}^9\text{B} \quad Q_m = -11.572$$

Angular distributions of  ${}^6\text{Li}$  ions have been obtained at  $E({}^3\text{He}) = 30.0$  and  $40.7$  MeV (1972OH01). See also (1974AJ01).

${}^9\text{C}$   
(Figs. 17 and 18)

GENERAL: (See also (1974AJ01).)


*Model calculations:* (1974IR04, 1976IR1B).

*Pion reactions:* (1974KA07, 1976HE1G, 1978SE1D).

*Other topics:* (1974IR04, 1975BE56, 1976IR1B, 1977CE05).

*Ground state properties:* (1975BE31).


*Mass of  ${}^9\text{C}$ :* The atomic mass excess of  ${}^9\text{C}$  is  $28912 \pm 3$  keV (1975KA18) based on the  $Q$ -value of the  ${}^{12}\text{C}({}^3\text{He}, {}^6\text{He}){}^9\text{C}$  reaction (1971TR03) and the threshold energy of the  ${}^7\text{Be}({}^3\text{He}, n){}^9\text{C}$  reaction (1971MO01).



The half-life of  ${}^9\text{C}$  is  $126.5 \pm 1.0$  msec (1971HA05, 1972ES05),  $126.5 \pm 2$  msec (1971MO01). Several groups of delayed protons are observed indicating the involvement of a number of  ${}^9\text{B}$  states: see Table 9.10 (1972ES05). See also (1974KU06, 1975WI1E; theor.).



$E_{\text{thresh.}} = 8980 \pm 5$  keV (1971MO01).



At  $E({}^3\text{He}) = 74.1$  MeV a  ${}^6\text{He}$  group is observed to the first excited state whose atomic mass excess is measured to be  $31.131 \pm 0.011$  MeV,  $\Gamma = 100 \pm 20$  keV. Based on the adopted AME for the ground state of  ${}^9\text{C}$  of  $28.912 \pm 0.003$  MeV,  $E_{\text{x}} = 2.219 \pm 0.011$  MeV (1974BE66).

Table 9.12: Energy levels of  ${}^9\text{C}$

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma$	Decay	Reactions
g.s. $2.219 \pm 11$	$(\frac{3}{2}^-); \frac{3}{2}$	$126.5 \pm 0.9$ msec $\Gamma = 100 \pm 20$ keV	$\beta^+$	1, 2, 3, 4 <sup>a</sup> 3

<sup>a</sup> See also (1974AJ01).

4.  ${}^{14}\text{N}({}^3\text{He}, {}^8\text{Li}){}^9\text{C}$   $Q_m = -32.064$

See  ${}^8\text{Li}$  (1976RO04).

${}^9\text{N}$   
(Not illustrated)

Not observed: see (1974IR04, 1975BE31, 1976IR1B; theor.).

## References

(Closed 1978)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1952BA1A Baggett and Bame, Phys. Rev. 85 (1952) 434
- 1955LE36 L.L. Lee, Jr. and D.R. Inglis, Phys. Rev. 99 (1955) 96
- 1955MA20 R.L. Macklin and H.E. Banta, Phys. Rev. 97 (1955) 753
- 1956BE1A Bezrukov, Panov and Timoshuk, Sov. J. Nucl. Energy 4 (1956) 609
- 1956BO18 C.K. Bockelman, A. Leveque and W.W. Buechner, Phys. Rev. 104 (1956) 456
- 1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser, Jr., Phys. Rev. 104 (1956) 1064
- 1957SL01 J.C. Slattery, R.A. Chapman and T.W. Bonner, Phys. Rev. 108 (1957) 809
- 1958KA31 R.W. Kavanagh and C.A. Barnes, Phys. Rev. 112 (1958) 503
- 1958SU14 R.G. Summers-Gill, Phys. Rev. 109 (1958) 1591
- 1959HA28 M.L. Halbert and A. Zucker, Phys. Rev. 115 (1959) 1635
- 1960KA17 R.W. Kavanagh, Nucl. Phys. 18 (1960) 492
- 1960SE12 R. Seltz and D. Magnac-Valette, Compt. Rend. 251 (1960) 2006
- 1960SP08 R.R. Spencer, G.C. Phillips and T.E. Young, Nucl. Phys. 21 (1960) 310
- 1961VA43 A.K. Valter, P.I. Vatsset, L.Y. Kolesnikov, S.G. Tonapetyan, K.K. Chernyavskii and A.I. Shpetnyi, Atomnaya Energ. 10 (1961) 577; At. Energy (USSR) 10 (1962) 574
- 1962CL06 F.M. Clikeman, A.J. Bureau and M.G. Stewart, Phys. Rev. 126 (1962) 1822
- 1962NE01 E. Newman, Phys. Rev. 125 (1962) 600
- 1962SE1A Serov and Guzhovskii, At. Energ. 12 (1962) 5
- 1963DU12 J.L. Duggan, P.D. Miller and R.F. Gabbard, Nucl. Phys. 46 (1963) 336
- 1963ME08 M.K. Mehta, W.E. Hunt, H.S. Plendl and R.H. Davis, Nucl. Phys. 48 (1963) 90
- 1963NG01 H. Nguyen Ngoc, N. Hors and J. Perez y Jorba, Nucl. Phys. 42 (1963) 62
- 1963PA04 P. Paul and D. Kohler, Phys. Rev. 129 (1963) 2698
- 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
- 1964BA29 R.W. Bauer, J.D. Anderson and C. Wong, Nucl. Phys. 56 (1964) 117

1964GR39 G. Gregoire and P.C. Macq, Phys. Lett. 8 (1964) 328  
 1964LU02 B.T. Lucas, S.W. Cosper and O.E. Johnson, Phys. Rev. 133 (1964) B963  
 1964MI04 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 50  
 1964TE01 E. Teranishi and B. Furubayashi, Phys. Lett. 9 (1964) 157  
 1964TE04 G. Tessler and W.E. Stephens, Phys. Rev. 135 (1964) B129  
 1965DI03 F.S. Dietrich, Nucl. Phys. 69 (1965) 49  
 1965GR18 T.A. Griffy and D.U.L. Yu, Phys. Rev. 139 (1965) B880  
 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81  
 1965IM01 W.L. Imhof, L.F. Chase, Jr. and D.B. Fossan, Phys. Rev. 139 (1965) B904  
 1965JA1A Jacmart, Cahiers Phys. (France) 19 (1965) 1  
 1965LY01 B. Lynch, G.M. Griffiths and T. Lauritsen, Nucl. Phys. 65 (1965) 641  
 1965NG1A Nguyen Ngoc, Ann. Phys. (France) 10 (1965) 315  
 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331  
 1965WO01 J.B. Woods and D.H. Wilkinson, Nucl. Phys. 61 (1965) 661  
 1966CA02 R.T. Carpenter and H.R. Hiddleston, Phys. Lett. 20 (1966) 286  
 1966CH20 P.R. Christensen and C.L. Cocke, Nucl. Phys. 89 (1966) 656  
 1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097  
 1966DE13 V.P. Denisov and L.A. Kulchitskii, Yad. Fiz. 3 (1966) 268; Sov. J. Nucl. Phys. 3 (1966) 192  
 1966KI09 K.G. Kibler, Phys. Rev. 152 (1966) 932  
 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1  
 1966MO05 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. 75 (1966) 113  
 1966OE1A Oertzen, Bock and Grosse-Schulte, Z. Naturforsch. 21A (1966) 946  
 1966PU02 P. Purdom, Jr., P.A. Seeger and R.W. Kavanagh, Nucl. Phys. 83 (1966) 513  
 1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, Phys. Rev. 144 (1966) 859; Erratum Phys. Rev. 148 (1966) 1246  
 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692  
 1966WI08 D.H. Wilkinson, J.T. Sample and D.E. Alburger, Phys. Rev. 146 (1966) 662  
 1967BE49 B.L. Berman, R.L. Van Hemert and C.D. Bowman, Phys. Rev. 163 (1967) 958; Erratum Phys. Rev. 168 (1968) 1414  
 1967FU08 K. Fukunaga, H. Nakamura and N. Fujiwara, J. Phys. Soc. Jpn. 23 (1967) 911

- 1967KI03 K.G. Kibler, Phys. Rev. 155 (1967) 1110
- 1967SE11 V.I. Serov, S.N. Abramovich, V.A. Pereshivkin and L.A. Morkin, Izv. Akad. Nauk SSSR Ser. Fiz. 31 (1967) 333; Bull. Acad. Nauk USSR Phys. Ser. 31 (1968) 317
- 1968AJ01 F. Ajzenberg-Selove, R.D. Wardaski and R. Middleton, Nucl. Phys. A116 (1968) 481
- 1968BA1C Barker and Fitzpatrick, Aust. J. Phys. 21 (1968) 415
- 1968CL08 H.-G. Clerc, K.J. Wetzel and E. Spamer, Nucl. Phys. A120 (1968) 441
- 1968CO07 C.L. Cocke, Nucl. Phys. A110 (1968) 321
- 1968CO08 C.L. Cocke and P.R. Christensen, Nucl. Phys. A111 (1968) 623
- 1968DA20 V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Yad. Fiz. 7 (1968) 758; Sov. J. Nucl. Phys. 7 (1968) 463
- 1968KR02 J.J. Kroepfl and C.P. Browne, Nucl. Phys. A108 (1968) 289
- 1968KU04 L.A. Kull and E. Kashy, Phys. Rev. 167 (1968) 963
- 1968OK06 Y. Okuma, J. Phys. Soc. Jpn. 25 (1968) 1
- 1968RE11 D.H. Rester, Nucl. Phys. A118 (1968) 129
- 1968RO19 P.G. Roos, H.G. Pugh, M. Jain, H.D. Holmgren, M. Epstein and C.A. Ludemann, Phys. Rev. 176 (1968) 1246
- 1968VA05 G.J. Vanpraet and W.C. Barber, Z. Phys. 211 (1968) 213
- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1969DE31 J.M. Delbrouck-Habaru, P.D. Dumont, M. Huez, G. Robaye and L. Winand, Bull. Soc. Roy. Sci. Liege 38 (1969) 240
- 1969DO09 B.S. Dolbilkin, A.I. Isakov, V.I. Korin, L.E. Lazareva, N.V. Linkova and F.A. Nikolaev, Yad. Fiz. 9 (1969) 914; Sov. J. Nucl. Phys. 9 (1969) 534
- 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, Phys. Rev. 178 (1969) 1551
- 1969KR03 E. Krubasik, H. Voit, E. Blatt, H.-D. Helb and G. Ischenko, Z. Phys. 219 (1969) 185
- 1969PI11 J.R. Pizzi, M. Gaillard, P. Gaillard, A. Guichard, M. Gusakow, G. Reboulet and C. Ruhla, Nucl. Phys. A136 (1969) 496
- 1969VI05 B. Vignon, E. Ligeon and J.P. Longequeue, J. Phys. (Paris) 30 (1969) 913
- 1970AL25 M.R. Aleksic, R.V. Popic, D.M. Stanojevic and B.Z. Stepanic, Fizika (Yugoslavia) 2 (1970) 113; Phys. Abs. 60823 (1970)
- 1970AN07 J.D. Anderson, C. Wong, B.A. Pohl and J.W. McClure, Phys. Rev. C2 (1970) 319
- 1970AU02 D.L. Auton, Nucl. Phys. A157 (1970) 305

- 1970BA49 P.H. Barker, A. Huber, H. Knoth, U. Matter, A. Gobbi and P. Marmier, Nucl. Phys. A155 (1970) 401
- 1970CH07 Y.S. Chen, T.A. Tombrello and R.W. Kavanagh, Nucl. Phys. A146 (1970) 136
- 1970CH19 V.I. Chuev, V.V. Davydov, V.I. Manko, B.G. Novatsky, S.B. Sakuta and D.N. Stepanov, Phys. Lett. B31 (1970) 624
- 1970CO04 L. Cohen and G.H. Herling, Nucl. Phys. A141 (1970) 595
- 1970GU08 K. Gul, B.H. Armitage and B.W. Hooton, Nucl. Phys. A153 (1970) 390
- 1970KA1A Kasim et al, Sci. Res. (Bangladesh) 7 (1970) 140; Phys. Abs. 74520 (1973)
- 1970LK1A Lkhagva and Rotter, Sov. J. Nucl. Phys. 11 (1970) 576
- 1970MI14 D. Miljanic, V. Valkovic, D. Rendic and M. Furic, Nucl. Phys. A156 (1970) 193
- 1970SQ01 G.T.A. Squier, A.R. Johnston, E.W. Spiers, S.A. Harbison and N.M. Stewart, Nucl. Phys. A141 (1970) 158
- 1970TU06 S.N. Tucker, P.B. Treacy and V.V. Komarov, Aust. J. Phys. 23 (1970) 651
- 1971AD01 J.C. Adloff, K.H. Souw and C.L. Cocke, Phys. Rev. C3 (1971) 1808
- 1971BI11 H.G. Bingham, A.R. Zander, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A173 (1971) 265
- 1971CH1B Chuev et al, Suppl. J. Phys. 32 (1971) C6-167
- 1971FR04 E. Friedland and I. Venter, Z. Phys. 243 (1971) 126
- 1971GE09 C. Gerardin, R. Seltz and D. Magnac-Valette, Nucl. Phys. A169 (1971) 521
- 1971HA05 J.C. Hardy, J.E. Esterl, R.G. Sextro and J. Cerny, Phys. Rev. C3 (1971) 700
- 1971HA10 J.C. Hardy, J.M. Loiseaux, J. Cerny and G.T. Garvey, Nucl. Phys. A162 (1971) 552
- 1971KA21 D.G. Kamke and C.D. Goodman, Nucl. Phys. A172 (1971) 555
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971MO01 J.M. Mosher, R.W. Kavanagh and T.A. Tombrello, Phys. Rev. C3 (1971) 438
- 1971RE19 D. Rendic, N.D. Gabitzsch, V. Valkovic, W. von Witsch and G.C. Phillips, Nucl. Phys. A178 (1971) 49
- 1971SC19 M. Schaeffer, M. Suffert and D. Magnac-Valette, Nucl. Phys. A175 (1971) 217; Erratum Nucl. Phys. A196 (1972) 639
- 1971SQ03 G.T.A. Squier, A.M. Hanna, J.B.A. England, D.L. Clough and L.H. Watson, Nucl. Phys. A170 (1971) 571
- 1971TR03 G.F. Trentelman, B.M. Preedom and E. Kashy, Phys. Rev. C3 (1971) 2205
- 1971YO04 P.G. Young and R.H. Stokes, Phys. Rev. C4 (1971) 1597
- 1971ZA07 A.R. Zander, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A173 (1971) 273

- 1972AD04 J.C. Adloff, W.K. Lin, K.H. Souw and P. Chevallier, Phys. Rev. C5 (1972) 664
- 1972BU30 A.J. Buffa, Jr. and M.K. Brussel, Nucl. Phys. A195 (1972) 545
- 1972CI05 D. Ciric, B. Stepancic, R. Popic, D. Stanojevic and M. Aleksic, Fizika 4 (1972) 40, 193
- 1972DE01 R.M. Devries, J.W. Sunier, J.-L. Perrenoud, M. Singh, G. Paic and I. Slaus, Nucl. Phys. A178 (1972) 417
- 1972DE02 R.M. Devries, J.-L. Perrenoud, I. Slaus and J.W. Sunier, Nucl. Phys. A178 (1972) 424
- 1972DE44 J. Decharge, G. Surget, G. Bruno and M.Y. Decharge, J. Phys. (Paris) 33 (1972) 485
- 1972ES05 J.E. Esterl, D. Allred, J.C. Hardy, R.G. Sextro and J. Cerny, Phys. Rev. C6 (1972) 373
- 1972GI07 R.W. Givens, M.K. Brussel and A.I. Yavin, Nucl. Phys. A187 (1972) 490
- 1972JA10 J.A. Jansen, R.T. Peerdeman and C. de Vries, Nucl. Phys. A188 (1972) 337
- 1972LO05 J.M. Lombaard and E. Friedland, Z. Phys. 249 (1972) 349
- 1972MA21 C. Maples and J. Cerny, Phys. Lett. B38 (1972) 504
- 1972ME07 F. Merchez, J. Arvieux, T. Honda and S. Morita, Nucl. Phys. A183 (1972) 417
- 1972OB01 A.W. Obst, T.B. Grandy and J.L. Weil, Phys. Rev. C5 (1972) 738
- 1972OH01 I.K. Oh, C.S. Zaidins, C.D. Zafiratos and S.I. Hayakawa, Nucl. Phys. A178 (1972) 497
- 1972QU01 J.R. Quinn, M.B. Epstein, S.N. Bunker, J.W. Verba and J.R. Richardson, Nucl. Phys. A181 (1972) 440
- 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430
- 1972WE08 K.A. Weber, K. Meier-Ewert, H. Schmidt-Bocking and K. Bethge, Nucl. Phys. A186 (1972) 145
- 1973AB10 S.N. Abramovich, B.Y. Guzhovskii, A.G. Zvenigorodskii and S.V. Trusillo, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1967; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 144
- 1973AL1A Allardyce et al, Report RL-73-038, Rutherford High Energy Lab. (1973); Phys. Abs. 74609 (1973)
- 1973AN29 G.P. Antropov, V.A. Vakhlamov, I.E. Mitrofanov and V.S. Russkikh, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 2657; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 172
- 1973AR1B Arthur, Thesis LA 5230 T (1973)
- 1973AU1A Augustson and Menlove, Trans. Amer. Nucl. Soc. 17 (1973) 494
- 1973BA1G Backenstoss et al, Nucl. Phys. B66 (1973) 125
- 1973BE19 J.C. Bergstrom, I.P. Auer, M. Ahmad, F.J. Kline, J.H. Hough, H.S. Caplan and J.L. Groh, Phys. Rev. C7 (1973) 2228

- 1973CA1B Cameron, Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 3
- 1973CA1C Catala et al, An. de Fisica 69 (1973) 267
- 1973DZ1A Dzhibuti, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1562
- 1973FE13 G. Fey, H. Frank, W. Schutz and H. Theissen, Z. Phys. 265 (1973) 401
- 1973GA19 S. Gamba, G. Ricco and G. Rottigni, Nucl. Phys. A213 (1973) 383
- 1973GO15 J.D. Goss, S.L. Blatt, D.R. Parsignault, C.D. Porterfeld and F.L. Riffle, Phys. Rev. C7 (1973) 1837
- 1973GO41 Y.P. Gorin, S.P. Denisov, S.V. Donskov, R.N. Krasnokutskii, A.I. Petrukhin, Y.D. Prokoshkin and D.A. Stoyanova, Yad. Fiz. 18 (1973) 336; Sov. J. Nucl. Phys. 18 (1974) 173
- 1973HE26 J.C.P. Heggie and P.W. Martin, Nucl. Phys. A212 (1973) 78
- 1973KE1B Kerimov, Aruri and Safin, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1768
- 1973KU19 E.L. Kuplennikov, N.G. Afanasev, V.A. Goldshtein, V.I. Ogurtsov and V.G. Vlasenko, Yad. Fiz. 18 (1973) 20; Sov. J. Nucl. Phys. 18 (1974) 10
- 1973KU1D Kubodera, Delorme and Rho, Nucl. Phys. B66 (1973) 253
- 1973MA24 R.E. Marrs, D. Bodansky and E.G. Adelberger, Phys. Rev. C8 (1973) 427
- 1973MA67 Z. Maric and V.C. Suslenko, Fizika (Yugoslavia) 5 (1973) 43
- 1973MO01 D.G. Montague, R.K. Cole, P.S. Lewis, C.N. Waddell and D.L. Hendrie, Nucl. Phys. A199 (1973) 433
- 1973PE19 C.-C. Perng, J.-L. Hwang and Y.-C. Hsu, Chin. J. Phys. (Taiwan) 11 (1973) 113; Erratum Chin. J. Phys. (Taiwan) 12 (1974) 52; Phys. Abs. 39997 (1974)
- 1973RO42 G. Robaye, J.M. Delbrouck-Habaru, H.P. Garnir and M. Huez, Bull. Soc. Roy. Sci. Liege (Belgium) 42 (1973) 598; Phys. Abs. 51834 (1974)
- 1973SC1C Schramm, Astrophys. J. 185 (1973) 293
- 1973SC1D Schramm, Explosive Nucleosynthesis, (Univ. Texas Press, 1973) p. 84
- 1973SL02 A.G. Slight, T.E. Drake and G.R. Bishop, Nucl. Phys. A208 (1973) 157
- 1973TA30 K. Takahashi, M. Yamada and T. Kondoh, At. Data Nucl. Data Tables 12 (1973) 101
- 1973TI1A Tinsley, Explosive Nucleosynth. (Univ. Texas Press, 1973) 22
- 1973TO16 L. Tomlinson, At. Data Nucl. Data Tables 12 (1973) 179
- 1973TR1B Truran, in Cosmochem., Ed. Cameron (Reidel Pub. Co., 1973) 23
- 1973VE12 F. Vesely and F. Bensch, J. Nucl. Energy 27 (1973) 828
- 1973VO02 H.J. Votava, T.B. Clegg, E.J. Ludwig and W.J. Thompson, Nucl. Phys. A204 (1973) 529; Erratum Nucl. Phys. A217 (1973) 628
- 1973WE15 M.F. Werby and S. Edwards, Nucl. Phys. A213 (1973) 294

- 1973WE1D Webber, Lezniak, Kish and Damle, *Astrophys. Space Sci.* 24 (1973) 17
- 1973WI07 N. Willis, I. Brissaud, Y. Le Bornec, B. Tatischeff and G. Duhamel, *Nucl. Phys.* A204 (1973) 454
- 1973WI1A Wilkin et al, *Nucl. Phys.* B62 (1973) 61
- 1973ZA06 N.I. Zaika, Y.V. Kibkalo, A.V. Mokhnach, O.F. Nemets, V.S. Semenov, V.P. Tokarev, E.D. Chesnokov and P.L. Shmarin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 1073; *Bull. Acad. Sci. USSR Phys. Ser.* 37 (1974) 140
- 1974AH03 J. Ahrens, H.B. Eppler, H. Gimm, M. Kroning, P. Riehn, A. Zieger and B. Ziegler, *Phys. Lett.* B52 (1974) 43
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, *Nucl. Phys.* A227 (1974) 1
- 1974AM01 J. Amato, R.L. Burman, R. Macek, J. Oostens, W. Schlaer, E. Arthur, S. Sobottka and W.C. Lam, *Phys. Rev.* C9 (1974) 501
- 1974AU1A Audouze and Tinsley, *Astrophys. J.* 192 (1974) 487
- 1974AV02 G.V. Avakov, E.S. Belyaeva and E.I. Dolinsky, *Nucl. Phys.* A220 (1974) 205
- 1974BA15 G.C. Ball, J.G. Costa, W.G. Davies, J.S. Forster, J.C. Hardy and A.B. McDonald, *Phys. Lett.* B49 (1974) 33
- 1974BE10 J. Bernabeu and J. Ros, *Nucl. Phys.* A220 (1974) 1
- 1974BE66 W. Benenson and E. Kashy, *Phys. Rev.* C10 (1974) 2633
- 1974BI14 J. Birchall, H.E. Conzett, J. Arvieux, W. Dahme and R.M. Larimer, *Phys. Lett.* B53 (1974) 165
- 1974BO1H Bonner et al, *Bull. Amer. Phys. Soc.* 19 (1974) 594
- 1974BO1J Boesgaard, *Astron. Astrophys.* 34 (1974) 9
- 1974BO1K Boesgaard et al, *Astrophys. J.* 194 (1974) L143
- 1974BO25 M. Bouten and M.-C. Bouten, *Lett. Nuovo Cim.* 10 (1974) 161
- 1974BO38 I. Bondouk, Z.A. Saleh, F. Machali and D.A. Darwish, *Rev. Roum. Phys.* 19 (1974) 653
- 1974BO39 I.I. Bondouk, *Rev. Roum. Phys.* 19 (1974) 843
- 1974BU1A Bulow and Forkman, in IAEA, STI/DOC/10/156 (1974) 475
- 1974CA34 A. Castellanos, J. Casanova, B. Leroux and R. Chastel, *An. Fis.* 70 (1974) 50
- 1974CE1A Cerny, in Nashville Vol. 2 (1974) 483
- 1974CH19 D.S. Chuu, C.S. Han and D.L. Lin, *Phys. Rev.* C9 (1974) 2086
- 1974CH1Q Chechik, Stocker, Eyal and Fraenkel, in Proc. Conf. Rxns Between Complex Nuclei, Vol. 1 (North-Holland, 1974) p. 6
- 1974CH46 M. Chemtob and S. Furui, *Nucl. Phys.* A233 (1974) 435

- 1974CH58 I.P. Chernov, B.I. Kuznetsov, V.V. Kozyr and V.A. Matusevich, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2530; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 58
- 1974CL04 A.S. Clough, G.K. Turner, B.W. Allardyce, C.J. Batty, D.J. Baugh, W.J. McDonald, R.A.J. Riddle, L.H. Watson, M.E. Cage, G.J. Pyle et al, *Nucl. Phys.* B76 (1974) 15
- 1974DA1D Datta and Thompson, *Bull. Amer. Phys. Soc.* 19 (1974) 552
- 1974DE1E De Jager, De Vries and De Vries, *At. Data Nucl. Data Tables* 14 (1974) 479
- 1974DI20 M. Dillig and M.G. Huber, *Lett. Nuovo Cim.* 11 (1974) 728
- 1974DO07 P. Dougan and W. Stiefler, *Z. Phys.* 269 (1974) 97
- 1974EN01 N. Ensslin, W. Bertozzi, S. Kowalski, C.P. Sargent, W. Turchinets, C.F. Williamson, S.P. Fivozinsky, J.W. Lightbody, Jr. and S. Penner, *Phys. Rev. C* 9 (1974) 1705
- 1974EN10 R. Engfer, H. Schneuwly, J.L. Vuilleumier, H.K. Walter and A. Zehnder, *At. Data Nucl. Data Tables* 14 (1974) 509
- 1974FO08 C.M. Fou and R.M. Gagne, *Z. Phys.* 267 (1974) 331
- 1974GO04 N.V. Goncharov, S.G. Tonapetyan and V.M. Khvorostyan, *Yad. Fiz.* 19 (1974) 21; *Sov. J. Nucl. Phys.* 19 (1974) 10
- 1974GR42 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.M. Pugach, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2543; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 69
- 1974GU13 V.B. Gubin and E.A. Romanovskii, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 752; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 76
- 1974GU1D Gurbanovich et al, *Ann. Phys.* 87 (1974) 263
- 1974HA1C Hanna, *Int. Conf. Nucl. Struct. Spectrosc., Pt. II, Amsterdam, 1974* (Amsterdam, Scholars Press 1974) 249
- 1974HO1E Hollandsworth, Bucher and Youngblood, *Bull. Amer. Phys. Soc.* 19 (1974) 574
- 1974HU14 J. Hufner, L. Tauscher and C. Wilkin, *Nucl. Phys.* A231 (1974) 455
- 1974HY01 M. Hyakutake, M. Sonoda, A. Katase, Y. Wakuta, M. Matoba, H. Tawara and I. Fujita, *J. Nucl. Sci. Tech. (Japan)* 11 (1974) 407
- 1974IR04 J.M. Irvine, G.S. Mani and M. Vallieres, *Czech. J. Phys.* B24 (1974) 1269
- 1974JA1F Jackson, *Prog. in Phys.* 37 (1974) 55
- 1974JO1C Jones and Bartle, *Nucl. Instrum. Meth.* 118 (1974) 525
- 1974KA07 W.B. Kaufmann, J.C. Jackson and W.R. Gibbs, *Phys. Rev. C* 9 (1974) 1340
- 1974KA15 E. Kashy, W. Benenson and J.A. Nolen, Jr., *Phys. Rev. C* 9 (1974) 2102
- 1974KN08 U. Kneissl, G. Kuhl, K.H. Leister and A. Weller, *Z. Naturf.* A29 (1974) 1688
- 1974KU06 N. Kumar, *Nucl. Phys.* A225 (1974) 221

1974LO09 W.G. Love, Nucl. Phys. A226 (1974) 319  
 1974LO10 J.M. Lombaard and E. Friedland, Z. Phys. 268 (1974) 413  
 1974LO1B Lorenzen and Brune, IAEA, STI/DOC/10/156 (1974) 325  
 1974LU06 E.J. Ludwig, C.E. Busch, T.B. Clegg, S.K. Datta and A.C. Watkins, Nucl. Phys. A230 (1974) 271  
 1974MCZS C.R. McClenahan, Thesis, Northwestern Univ. (1974); ANL-8088 (1974)  
 1974MI05 C.A. Miller, J.W. Watson, D.I. Bonbright, F.J.S. Wilson and D.O. Wells, Phys. Rev. Lett. 32 (1974) 684  
 1974MI06 G.A. Miller, Nucl. Phys. A224 (1974) 269  
 1974MU13 N.C. Mukhopadhyay and F. Cannata, Phys. Lett. B51 (1974) 225  
 1974NA25 I.C. Nascimento, C.F. Wong and H.S. Caplan, Rev. Brasil. Fis. 4 (1974) 241; Phys. Abs. 84312 (1975)  
 1974PA1B Park, Cap, Leung and Rickett, Bull. Amer. Phys. Soc. 19 (1974) 666  
 1974PI11 C.F. Pineda, M.L. de Toledo and C. Ruiz Bauza, An. Fis. 70 (1974) 30  
 1974RE1A Reeves, Ann. Rev. Astron. Astrophys. 12 (1974) 437  
 1974RO1G Ross and Aller, Solar Phys. 36 (1974) 11  
 1974RO31 E. Roeckl, P.F. Dittner, C. Detraz, R. Klapisch, C. Thibault and C. Rigaud, Phys. Rev. C10 (1974) 1181  
 1974SHYR V.S. Shirley and C.M. Lederer, LBL-3450 (1974)  
 1974SP1A Spence, Thesis, College Williams & Mary (1974); Phys. Abs. 80252 (1975)  
 1974ST1G G.L. Strobel, Nucl. Phys. A232 (1974) 502  
 1974TA18 L. Tauscher and W. Schneider, Z. Phys. 271 (1974) 409  
 1974TA1C Takimoto et al, Bull. Inst. Chem. Res. Kyoto Univ. 52 (1974) 240  
 1974TH01 C. Thibault and R. Klapisch, Phys. Rev. C9 (1974) 793  
 1974TI05 Y.I. Titov, E.V. Stepula, N.G. Afanasev, R.V. Akhmerov and N.F. Severin, Yad. Fiz. 19 (1974) 479; Sov. J. Nucl. Phys. 19 (1974) 240  
 1974VO06 P.L. Von Behren, E. Norbeck and G.L. Payne, Phys. Rev. C10 (1974) 550  
 1974YA05 M. Yasue, J. Phys. Soc. Jpn. 36 (1974) 1254  
 1974YA1C Yasue et al, Bull. Inst. Chem. Res. Kyoto Univ. 52 (1974) 177  
 1975AB1D Abdul-Magd, Prog. Theor. Phys. 53 (1975) 997  
 1975AB1E Abe, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 500  
 1975AB1F Abramov, INDC(CCP)-52/LN, Vienna (1975)

- 1975AH06 J. Ahrens, H. Borchert, K.H. Czock, H.B. Eppler, H. Gimm, H. Gundrum, M. Kroning, P. Riehn, G. Sita Ram, A. Zieger et al, Nucl. Phys. A251 (1975) 479
- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
- 1975AN01 B. Antolkovic and Z. Dolenc, Nucl. Phys. A237 (1975) 235
- 1975AN1H Andersson, Dougan and Stiefler, Z. Phys. A272 (1975) 263
- 1975AR14 A.G. Artyukh, V.V. Volkov, G.F. Gridnev and V.L. Mikheev, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2; Bull. Acad. Sci. USSR Phys. Ser. 39 (1975) 1
- 1975BA1G Baldin, in LASL, AIP Conf. Proc. 26 (1975) 621
- 1975BA1L Batty, Czech. J. Phys. 25 (1975) 286
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BE56 W. Benenson, Nukleonika (Poland) 20 (1975) 775
- 1975BO55 I.I. Bondouk, F. Asfour, Z. Saleh and F. Machali, Rev. Roum. Phys. 20 (1975) 1095
- 1975BR1F Brown, At. Data Nucl. Data Tables 15 (1975) 111
- 1975BU11 W.E. Burcham, J.B.A. England, R.G. Harris, O. Karban and S. Roman, Nucl. Phys. A246 (1975) 269
- 1975BU1A Bugg, Few Body Prob., Quebec, 1974 (Univ. Laval, 1975) 267
- 1975CA18 J.D. Carlson, C.D. Zafiratos and D.A. Lind, Nucl. Phys. A249 (1975) 29
- 1975CA1H Cannata, Lett. Nuovo Cim. 13 (1975) 319
- 1975CA1J Casanova, Villamanan and Casanova, An. Fis. 71 (1975) 150
- 1975CA1K Castellanos, An. Fis. 71 (1975) 197
- 1975CH1E Christillin and Rosa-Clot, Nuovo Cim. A28 (1975) 29
- 1975CH1F Chmielewski, Muller and Brault, Astron. Astrophys. 42 (1975) 37
- 1975CH22 P. Christillin, A. Dellafiore and M. Rosa-Clot, Phys. Rev. C12 (1975) 691
- 1975CH28 D.S. Chuu and C.S. Han, Chin. J. Phys. (Taiwan) 13 (1975) 59
- 1975DA15 V. D'Amico, G. Fazio, S. Jannelli, F. Mezzanares and R. Potenza, Lett. Nuovo Cim. 13 (1975) 489
- 1975DA1A Dam et al, Bull. Amer. Phys. Soc. 20 (1975) 1156
- 1975DI05 W.R. Ditzler, M. Breidenbach, J.I. Friedman, H.W. Kendall, J.S. Poucher, E.D. Bloom, R.L.A. Cottrell, D.H. Coward, H. DeStaebler, C.L. Jordan et al, Phys. Lett. B57 (1975) 201
- 1975ER03 M. Ericson and M. Krell, Nucl. Phys. A241 (1975) 487
- 1975FA1A Fagg, Revs. Mod. Phys. 47 (1975) 683
- 1975FE1A Feather, Commun. R. Soc. Edinb. (Phys. Sci.) 2 (1975) 19

- 1975FE1B Feinberg, in LASL, AIP Conf. Proc. 26 (1975) 468
- 1975FO13 C.M. Fou, R.M. Gagne and P.T. Wu, Can. J. Phys. 53 (1975) 2445
- 1975FO19 W.A. Fowler, G.R. Caughlan and B.A. Zimmerman, Ann. Rev. Astron. Astrophys. 13 (1975) 69
- 1975FU05 K. Fujii, J. Schimizu, K. Takimoto, I. Yamane and J. Muto, J. Phys. Soc. Jpn. 38 (1975) 1563
- 1975GE12 J.P. Genin, J. Julien, M. Rambaut, C. Samour, A. Palmeri and D. Vinciguerra, Lett. Nuovo Cim. 13 (1975) 693
- 1975GO1H Goland, Snead, Parkin and Theus, Nucl. Sci. NS-22 (1975) 1776
- 1975HO1F Hollandsworth, Bucher, Youngblood and Niiler, Rept. BRL-1764 (1975); Phys. Abs. 982 (1976)
- 1975HU03 R.J. Hughes, R.H. Sambell, E.G. Muirhead and B.M. Spicer, Nucl. Phys. A238 (1975) 189
- 1975HU1D Hufner, Phys. Rept. C21 (1975) 1
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, Phys. Rev. C11 (1975) 2049
- 1975KA05 J. Kasagi, T. Nakagawa, N. Sekine, T. Tohei and H. Ueno, Nucl. Phys. A239 (1975) 233
- 1975KA18 E. Kashy, W. Benenson, D. Mueller, R.G.H. Robertson and D.R. Goosman, Phys. Rev. C11 (1975) 1959
- 1975KN03 U. Kneissl, G. Kuhl, K.-H. Leister and A. Weller, Nucl. Phys. A247 (1975) 91
- 1975KN1C B. Knapp, W. Lee, P. Leung, S.D. Smith, A. Wijangco, J. Knauer, D. Yount, D. Nease, J. Bronstein, R. Coleman et al, Phys. Rev. Lett. 34 (1975) 1044
- 1975KR14 S. Krol and S. Szpikowski, Acta Phys. Pol. B6 (1975) 409
- 1975KR1D Krasnopol'sky and Kukjlin, in Clustering Phenomena in Nuclei, II, ORO-4856-26 (1975) p. 157
- 1975KU01 D. Kurath and D.J. Millener, Nucl. Phys. A238 (1975) 269
- 1975KU1K Kuznetsov and Chernov, Sov. J. Nucl. Phys. 20 (1975) 340
- 1975KU27 N. Kumar and S.K. Soni, Indian J. Pure Appl. Phys. 13 (1975) 826
- 1975LA1E Lavrukhina, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 395
- 1975LA23 L. Lapikas, G. Box and H. De Vries, Nucl. Phys. A253 (1975) 324
- 1975LI20 D.L. Lin and M.Y. Chen, Nucl. Phys. A251 (1975) 457
- 1975MA1K Matthews, Proc. Int. Symp. Interactions, Studies in Nuclei, Germany, 1975 (Amsterdam, North-Holland, 1975) p. 429
- 1975MC02 C.R. McClenahan and R.E. Segel, Phys. Rev. C11 (1975) 370

- 1975ME1E Meneguizzi and Reeves, *Astron. Astrophys.* 40 (1975) 91, 99
- 1975MO27 R. Moreh and Y. Wand, *Nucl. Phys.* A252 (1975) 423
- 1975NI1B Nichitiu, *Few Body Problems, Quebec, 1974 (Univ. Laval, 1975)* p. 90
- 1975OB01 D.L. Oberg, D. Bodansky, D. Chamberlin and W.W. Jacobs, *Phys. Rev.* C11 (1975) 410
- 1975PO1C Povh, in *Lasl, AIP Conf. Proc.* 26 (1975) 173
- 1975PR1B Preszler, Kish, Iezniak and Webber, *Bull. Amer. Phys. Soc.* 20 (1975) 711
- 1975RA12 G.M. Raisbeck and F. Yiou, *Phys. Rev. Lett.* 35 (1975) 155
- 1975RA13 L. Ray, W.R. Coker and T. Udagawa, *Phys. Lett.* B56 (1975) 318
- 1975RA14 G.M. Raisbeck, J. Lestringuez and F. Yiou, *Phys. Lett.* B57 (1975) 186
- 1975RA21 G.M. Raisbeck, P. Boerstling, R. Klapisch and T.D. Thomas, *Phys. Rev.* C12 (1975) 527
- 1975RO1B Roos and Chant, in *Clustering Phenom. in Nucl., II, ORO-4856-26 (1975)* 242, 265
- 1975RO1G Rosen, in *Lasl, AIP Conf. Proc.* 26 (1975) 265
- 1975SA01 Y. Sakamoto, P. Cuer and F. Takeuchi, *Phys. Rev.* C11 (1975) 668
- 1975SC05 H. Schier and B. Schoch, *Lett. Nuovo Cim.* 12 (1975) 334
- 1975SC1K Schenzle, in *Clustering Phenom. in Nucl., II, ORO-4856-26 (1975)* 85
- 1975SC1M J.P. Schiffer and H.J. Lipkin, *Phys. Rev. Lett.* 35 (1975) 708
- 1975SC41 U. Schwinn, G. Mairle, G.J. Wagner and C. Ramer, *Z. Phys.* A275 (1975) 241
- 1975SE07 F. Seiler, *Nucl. Phys.* A244 (1975) 236
- 1975ST16 D.M. Stanojevic and R.V. Popic, *Fizika (Yugoslavia)* 7 (1975) 143; *Phys. Abs.* 25126 (1976)
- 1975TA1C Tauscher, in *LASL, AIP Conf. Proc.* 26 (1975) 541
- 1975VO1D K.G. Vosburgh, W. Schimmerling, K. Koepke and W.D. Wales, *Phys. Rev.* D11 (1975) 1743
- 1975WI1E Wilkinson, *Proc. Int. Symp. on Inter. Studies in Nucl., Germany, 1975 (Amsterdam, North-Holland, 1975)* 147
- 1975YA02 Y.-Y. Yam, *Phys. Rev.* C11 (1975) 73
- 1975ZA06 N.I. Zaika and P.L. Shmarin, *Ukr. Fiz. Zh.* 20 (1975) 238; *Phys. Abs.* 45405 (1975)
- 1975ZE01 A.M. Zebelman, A.M. Poskanzer, J.D. Bowman, R.G. Sextro and V.E. Viola, Jr., *Phys. Rev.* C11 (1975) 1280
- 1976AB1B Abramov et al, *Sov. J. Nucl. Phys.* 23 (1976) 636
- 1976AJ04 F. Ajzenberg-Selove, *Nucl. Phys.* A268 (1976) 1

- 1976AL02 D.E. Alburger and D.H. Wilkinson, Phys. Rev. C13 (1976) 835
- 1976AL1F Alder et al, in Meson-Nucler Phys. 1976 (AIP, 1976) 158
- 1976AL1G Alkhazov et al, in Contributions to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) p. 158
- 1976AN1A Antokovic, Fiz. (Yugoslavia) 8 (1976) 163; Phys. Abs. 6364 (1977)
- 1976AN1B Anderson et al, Bull. Amer. Phys. Soc. 21 (1976) 978
- 1976AS1B Aslanides, Meson-Nucl. Phys., 1976 (AIP, 1976) 204
- 1976AU1B Audouze and Lequeux, Astron. Astrophys. 49 (1976) 133
- 1976AU1C Audouze, Meneguzzi and Reeves, in Spallation Nucl. Rxns. & Their Applic., Shen/Marker Eds. (D. Reidel, Holland 1976) 113
- 1976AV05 G.V. Avakov and E.I. Dolinsky, Nucl. Phys. A270 (1976) 74
- 1976BA1P Barabash et al, Sov. J. Nucl. Phys. 24 (1976) 188
- 1976BE1H Berman, Preprint UCRL-78482 (1976)
- 1976BE1K Bertini, in Spallation Nucl. Rxns. and Their Appl., Eds., Shen and Merker (D. Reidel, The Netherlands, 1976) p. 27
- 1976BE67 B.N. Belyaev, V.D. Domkin, Y.G. Korobulin and L.M. Krizhanskii, Yad. Fiz. 24 (1976) 1177; Sov. J. Nucl. Phys. 24 (1976) 617
- 1976BI1B Bilpuch et al, in Lowell Conf. 760715-P2 (1976) 1309
- 1976BO1E Boesgaard, Publ. Astron. Soc. Pac. 88 (1976) 353
- 1976BO1H Boesgaard, Astrophys. J. 210 (1976) 466
- 1976BO1J Boesgaard and Chesley, Astrophys. J. 210 (1976) 475
- 1976BO1K T. Bowen, D.A. DeLise and A.E. Pifer, Phys. Rev. C14 (1976) 195
- 1976BR1G W. Bruckner, B. Granz, D. Ingham, K. Kilian, U. Lynen, J. Niewisch, B. Pietrzyk, B. Povh, H.G. Ritter and H. Schroder, Phys. Lett. B62 (1976) 481
- 1976BR26 F. Brut and S. Jang, Phys. Rev. C14 (1976) 1638
- 1976BU16 M. Buenerd, C.K. Gelbke, B.G. Harvey, D.L. Hendrie, J. Mahoney, A. Menchaca-Rocha, C. Olmer and D.K. Scott, Phys. Rev. Lett. 37 (1976) 1191
- 1976CA17 C.G. Cassapakis, H.C. Bryant, B.D. Dieterle, C.P. Leavitt, D.M. Wolfe, B.E. Bonner, J.E. Simmons, C.W. Bjork, P.J. Riley, M.L. Evans et al, Phys. Lett. B63 (1976) 35
- 1976CH28 A. Chevarier, N. Chevarier, A. Demeyer, P. Martin and P. de Saintignon, Nucl. Phys. A274 (1976) 223
- 1976CO1B Colgate, Audouze and Fowler, OAP-458 (1976)
- 1976CO1L Correll et al, Bull. Amer. Phys. Soc. 21 (1976) 513
- 1976DA15 S.E. Darden, G. Murillo and S. Sen, Nucl. Phys. A266 (1976) 29

- 1976DA1D R.H. Dalitz and A. Gal, Phys. Rev. Lett. 36 (1976) 362; Erratum Phys. Rev. Lett. 36 (1976) 628
- 1976DA1E R.H. Dalitz and A. Gal, Phys. Lett. B64 (1976) 154
- 1976DO1D Dover, Meson-Nucl. Phys.-1976 (AIP, 1976) p. 249
- 1976DU05 J. Dubach, J.H. Koch and T.W. Donnelly, Nucl. Phys. A271 (1976) 279
- 1976DU1B Dumbrajs, Meson-Nucl. Phys. 1976 (AIP, 1976) 192
- 1976EC1A Eck and Gray, Bull. Amer. Phys. Soc. 21 (1976) 555
- 1976ED1A Edge, Tompkins and Glenn, Nuovo Cim. A31 (1976) 641
- 1976EN02 H.D. Engelhardt, C.W. Lewis and H. Ullrich, Nucl. Phys. A258 (1976) 480
- 1976EP1A Epsteain, Arnett and Schramm, Astrophys. J. Suppl. 31 (1976) 111
- 1976ES1A Esmael, Nuovo Cim. A35 (1976) 419
- 1976FI03 P. Fintz, G. Guillaume, F. Jundt, I. Ordonez, A. Gallmann and D.E. Alburger, Nucl. Phys. A259 (1976) 493
- 1976FR05 S. Frankel, W. Frati, V. Highland, O. Van Dyck and R. Werbeck, Phys. Rev. C13 (1976) 737
- 1976FU06 G.H. Fuller, J. Phys. Chem. Ref. Data 5 (1976) 835
- 1976FU1B Fukui, Enge and Beaujean, Z. Phys. A277 (1976) 99
- 1976GA1C Gal, Meson-Nucl. Phys.-1976 (AIP,1976) p. 694
- 1976GI1C Ginzburg and Ptjskin, Revs. Mod. Phys. 48 (1976) 161
- 1976GO1F Gould and Boyce, Nucl. Sci. Eng. 60 (1976) 477
- 1976HA1F Hainebach, Schramm and Blake, Astrophys. J. 205 (1976) 920
- 1976HA1J Hanna, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 407
- 1976HA1M Hardekopf, Bull. Amer. Phys. Soc. 21 (1976) 551
- 1976HE1G Hess et al, Bull. Amer. Phys. Soc. 21 (1976) 518
- 1976HI05 M.D. High and B. Cujec, Nucl. Phys. A259 (1976) 513
- 1976IR02 M. Irshad, J. Asai, S. Sen, R. Pigeon and R.J. Slobodrian, Nucl. Phys. A265 (1976) 349
- 1976IR1B Irvine et al, Ann. Phys. 102 (1976) 129
- 1976KA1D Karban et al, in Contributions to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) p. 95
- 1976KH1C Khandelwal, Pritchard and Singh, Nucl. Phys. Sci. Eng. 60 (1976) 481
- 1976KI1E Kilian, Meson-Nucl. Phys.-1976 (AIP,1976) p. 497
- 1976KU07 N. Kumar and O. Prakash, Z. Phys. A278 (1976) 105

- 1976LE1F Lefort, Rep. Prog. 39 (1976) 129
- 1976LE1K Lebedev et al, in Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) 98
- 1976MA04 F. Malaguti and P.E. Hodgson, Nucl. Phys. A257 (1976) 37
- 1976MC10 A.B. McDonald, T.K. Alexander, O. Hausser, D. Disdier, E.G. Adelberger, H.B. Mak, A.P. Shukla and A.V. Nero, Nucl. Phys. A273 (1976) 451
- 1976MC1E McEllistrem, in Lowell Conf., CONF-760715-P1 (1976) p. 171
- 1976MI13 T. Mikumo, I. Kohno, K. Katori, T. Motobayashi, S. Nakajima, M. Yoshie and H. Kamitsubo, Phys. Rev. C14 (1976) 1458
- 1976MO1B Moss, Brown and Cornelius, Bull. Amer. Phys. Soc. 21 (1976) 990
- 1976NA11 M.N. Namboodiri, E.T. Chulick and J.B. Natowitz, Nucl. Phys. A263 (1976) 491
- 1976OS04 R. Ost, N.E. Sanderson, S. Mordechai, J.B.A. England, B.R. Fulton, J.M. Nelson and G.C. Morrison, Nucl. Phys. A265 (1976) 142
- 1976PE1A C. Perron, Phys. Rev. C14 (1976) 1108
- 1976PI1B Pilkuhn, Schlaile and Zoyko, Meson-Nucl. Phys., 1976 (AIP, 1976) p. 194
- 1976RA1C Raisbeck and Yiou, Spallation Nucl. Rxns. & Their Appl., Eds., Shen and Merker (D. Reidel, Netherlands, 1976) 83
- 1976RA1F J. Ranft and G. Ranft, Phys. Rev. Lett. 36 (1976) 988
- 1976RA1G Ramavataram et al, Proc. 7th Int. Conf. on Few Body Problems in Nucl. and Particle Phys., Delhi, 1975 (Amsterdam, North-Holland, 1976) p. 273; Phys. Abs. 18590 (1977)
- 1976RO04 R.G.H. Robertson, W. Benenson, E. Kashy and D. Mueller, Phys. Rev. C13 (1976) 1018
- 1976RO12 C.T. Roche, R.G. Clark, G.J. Mathews and V.E. Viola, Jr., Phys. Rev. C14 (1976) 410
- 1976RO1J Ross and Aller, Science 191 (1976) 1223
- 1976RO1L Roman, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 255
- 1976RO1N A. Robertson, T.J. Kennett and W.V. Prestwich, Phys. Rev. C13 (1976) 1552
- 1976SC14 A.E. Schilling, N.F. Mangelson, K.K. Nielsen, D.R. Dixon, M.W. Hill, G.L. Jensen and V.C. Rogers, Nucl. Phys. A263 (1976) 389
- 1976SH1H Sharma and Mukherjee, Pramana 7 (1976) 181
- 1976SI1C Simpson, Bull. Amer. Phys. Soc. 21 (1976) 602
- 1976SI1D Silberberg, Tsao and Shapiro, in Spallation Nucl. Rxns. & Their Appl., Eds., Shen and Mercer (D. Reidel, Netherlands, 1976) p. 49
- 1976SL1A Slobodrian et al, Proc. 7th Int. Conf. on Few Body Prob. in Nucl. & Part. Phys., Delhi, 1975 (Amsterdam, North-Holland, 1976) 263; Phys. Abs. 18589 (1977)

- 1976ST11 W.F. Steele, P.A. Smith, J.E. Finck and G.M. Crawley, Nucl. Phys. A266 (1976) 424
- 1976TR1A Truol, Meson-Nucl. Phys., 1976 (AIP, 1976) p. 581
- 1976UE01 H. Ueno, T. Nakagawa, M. Baba, J. Kasagi, H. Orihara and T. Tohei, J. Phys. Soc. Jpn. 40 (1976) 1537
- 1976VI1A Vidal-Quadras et al, Nuovo Cim. B34 (1976) 380
- 1976WE17 E.W. Weber and J. Vetter, Phys. Lett. A56 (1976) 446
- 1976WI05 N. Willis, I. Brissaud, L. Bimbot, Y. Le Bornec and B. Tatischeff, Nucl. Phys. A261 (1976) 45
- 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, Phys. Rev. C14 (1976) 815
- 1976ZA06 W. Zahn, Nucl. Phys. A269 (1976) 138
- 1977AB09 B.M. Abramov, I.A. Dukhovskoi, V.V. Kishkurno, L.A. Kondratyuk, A.P. Krutenkova, V.V. Kulikov, M.A. Matsyuk, I.A. Radkevich, E.N. Turdakina, V.S. Fedorets et al, Zh. Eksp. Teor. Fiz. Pisma 25 (1977) 123; JETP Lett. 25 (1977) 111
- 1977AL1C Alder et al, Helv. Phys. Acta 50 (1977) 197
- 1977AN1D Antipov et al, JETP Lett. 26 (1977) 414
- 1977AP1A Apokin et al, Yad. Fiz. 25 (1977) 555
- 1977AR06 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A283 (1977) 350
- 1977AR1F Arefyev et al, in Proc., Zurich-Sin (1977) p. 130
- 1977AS04 J. Asai, C. Rioux, M. Irshad, R. Pigeon and R.J. Slobodrian, Phys. Rev. C16 (1977) 1762
- 1977AS1D Aslanides et al, in Tokyo (1977), Contrib. Papers p. 844
- 1977AU1B Austin, MSUCL-254 (1977)
- 1977BA1Q Baer, in High Energy Phys. & Nucl. Struct., Zurich 1977 (Birkhauser Verlag, 1977) 245
- 1977BA27 C.J. Batty, S.F. Biagi, M. Blecher, R.A.J. Riddle, B.L. Roberts, J.D. Davies, G.J. Pyle, G.T.A. Squier and D.M. Asbury, Nucl. Phys. A282 (1977) 487
- 1977BA2D Bayukov et al, in Proc., Zurich-Sin (1977) p. 150
- 1977BA2E Bansal and Kumar, Pramana 9 (1977) 273
- 1977BA51 B. Bassalleck, W.-D. Klotz, F. Takeutchi, H. Ullrich and M. Furic, Phys. Rev. C16 (1977) 1526
- 1977BA71 R. Balzer, M. Hugi, B. Kamys, J. Lang, R. Muller, E. Ungricht, J. Unternahrer, L. Jarczyk and A. Strzalkowski, Nucl. Phys. A293 (1977) 518
- 1977BE1Q Bernheim et al, in Proc. Zurich-Sin (1977) 267

- 1977BE1U Bejidian et al, in Proc., Zurich-Sin (1977) 308
- 1977BE1V Belikov et al, in Proc., Zurich-Sin (1977) 152
- 1977BO1F Boesgaard, Heacox and Conti, *Astrophys. J.* 214 (1977) 124
- 1977BU02 A. Buchnea, R.G. Johnson and K.G. McNeill, *Can. J. Phys.* 55 (1977) 364
- 1977CA08 E. Caurier and B. Grammaticos, *Nucl. Phys. A279* (1977) 333
- 1977CA1E Carter et al, in Proc., Zurich-Sin (1977) p. 318
- 1977CE05 J. Cerny and J.C. Hardy, *Ann. Rev. Nucl. Sci.* 27 (1977) 333
- 1977CI1A Ciric et al, *Fizika (Yugoslavia)* 9, Suppl. 1 (1977) 39; *Phys. Abs.* 17216 (1978)
- 1977CO07 A.A. Cowley, P.G. Roos, N.S. Chant, R. Woody, III, H.D. Holmgren and D.A. Goldberg, *Phys. Rev. C15* (1977) 1650
- 1977CO1G Colombo et al, in Tokyo (1977), *Contrib. Papers* p. 490
- 1977CR1C L. Criegee, G. Franke, A. Giese, Th. Kahl, G. Poelz, U. Timm, H. Werner and W. Zimmermann, *Nucl. Phys. B121* (1977) 38
- 1977DA1B Davydov, Naumov and Shcherbachenko, *INDC(CCP)-104/LN* (1977)
- 1977DE08 C. Detraz, *Phys. Lett. B66* (1977) 333
- 1977DE1F Delorme and Figureau, in Tokyo (1977), *Contrib. Papers* p. 821
- 1977DE1G Deutsch et al, in Proc., Zurich-Sin (1977) p. 46
- 1977DE1H Delorme and Figureau, in Proc., Zurich-Sin (1977) p. 120
- 1977DE1J Demianov, Murzin and Sarycheva, in Proc., Zurich-Sin (1977) 143
- 1977DE1K Delbar, Gregoire and Paic, *Fizika (Yugoslavia)* 9, Suppl. 2 (1977) 1; *Phys. Abs.* 615 (1978)
- 1977DM1A Dmitrieva, Lebedev and Matusевич, *Izv. Akad. Nauk SSSR Ser. Fiz.* 41 (1977) 143
- 1977DO06 G.E. Dogotar, R.A. Eramzhyan, H.R. Kissener and R.A. Sakaev, *Nucl. Phys. A282* (1977) 474
- 1977DR09 D.M. Drake, G.F. Auchampaugh, E.D. Arthur, C.E. Ragan and P.G. Young, *Nucl. Sci. Eng.* 63 (1977) 401
- 1977DR1C Dravins and Hultqvist, *Astron. Astrophys.* 55 (1977) 463
- 1977DW1A Dwyer and Meyer, *Bull. Amer. Phys. Soc.* 22 (1977) 600
- 1977EC04 J.S. Eck, T.J. Gray and R.K. Gardner, *Phys. Rev. C16* (1977) 1873
- 1977FE1B Feshbach, in *High Energy Phys. & Nucl. Struct., Zurich 1977* (Birkhauser Verlag, 1977) 291
- 1977FY1A Fyodorov et al, in Proc., Zurich-Sin (1977) 157
- 1977GA1C Garcia-Munoz, Mason and Simpson, *Astrophys. J.* 217 (1977) 859

- 1977GE08 C.K. Gelbke, D.K. Scott, M. Bini, D.L. Hendrie, J.L. Laville, J. Mahoney, M.C. Mer-  
maz and C. Olmer, Phys. Lett. B70 (1977) 415
- 1977GO07 J. Gosset, H.H. Gutbrod, W.G. Meyer, A.M. Poskanzer, A. Sandoval, R. Stock and  
G.D. Westfall, Phys. Rev. C16 (1977) 629
- 1977GR1D Grabisch, Beaujean, Scherzer and Enge, Nucl. Instrum. Meth. 147 (1977) 263
- 1977GR1F Grand and Goland, Nucl. Instrum. Meth. 145 (1977) 49
- 1977GU14 M.G. Gulyamov, B.S. Mazitov, G.A. Radyuk, E.I. Rasulov and R. Yarmukhamedov,  
Izv. Akad. Nauk SSSR Ser. Fiz. 41 (1977) 2214; Bull. Acad. Sci. USSR Phys. Ser. 41  
(1977) 189
- 1977HA1L Hagen, Fisher and Ormes, Astrophys. J. 212 (1977) 262
- 1977HO1B Hoistad, in High Energy Phys. Nucl. Struct., Zurich 1977 (Birkhauser Verlag, 1977)  
215
- 1977HO1F Horiuchi, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44  
(1978) 85
- 1977IG1D Igo et al, in Tokyo (1977), Contrib. Papers p. 768
- 1977IR01 M. Irshad, C. Rioux, J. Asai, R. Pigeon and R.J. Slobodrian, Nucl. Phys. A286 (1977)  
483
- 1977JA14 H.U. Jager and M. Kirchbach, Nucl. Phys. A291 (1977) 52
- 1977JE01 H. Jeremie and G. Kennedy, Can. J. Phys. 55 (1977) 261
- 1977KE09 K.W. Kemper, G.E. Moore, R.J. Puigh and R.L. White, Phys. Rev. C15 (1977) 1726
- 1977KI04 A. Kiss, E. Koltay, G. Szabo and L. Vegh, Nucl. Phys. A282 (1977) 44
- 1977KO1K Kolesnikov, Krokhin, Kopylov and Chernov, Izv. Vuz Fiz. 6 (1977) 75; Phys. Abs.  
74791 (1977)
- 1977KO27 W. Kohler, G. Gruber, A. Steinhauser and K. Bethge, Nucl. Phys. A290 (1977) 233
- 1977KU12 E.L. Kuplennikov, V.A. Goldshtein, N.G. Afanasev and V.G. Vlasenko, Yad. Fiz. 25  
(1977) 1129; Sov. J. Nucl. Phys. 25 (1977) 599
- 1977KU1D R. Kullberg, K. Kristiansson, B. Lindkvist and I. Otterlund, Nucl. Phys. A280 (1977)  
491
- 1977LA11 J. Lang, R. Muller, J. Unternahrer, L. Jarczyk, B. Kamys and A. Strzalkowski, Phys.  
Rev. C16 (1977) 1448
- 1977LO10 M.A. Lone, C.B. Bigham, J.S. Fraser, H.R. Schneider, T.K. Alexander, A.J. Ferguson  
and A.B. McDonald, Nucl. Instrum. Meth. 143 (1977) 331
- 1977MA1H Mason, Garcia-Munoz and Simpson, Bull. Amer. Phys. Soc. 22 (1977) 567
- 1977MA1L Mateja, Parks, Frawley and Nagel, Bull. Amer. Phys. Soc. 22 (1977) 1002
- 1977MA35 V.B. Mandelzweig, Nucl. Phys. A292 (1977) 333

- 1977MC1E K.W. McVoy, Nucl. Phys. A276 (1977) 491
- 1977MO11 P.J. Moffa, Phys. Rev. C16 (1977) 1431
- 1977MU1A Mukhopadhyay, Phys. Rept. C30 (1977) 1
- 1977OK01 S. Okabe, Y. Abe and H. Tanaka, Prog. Theor. Phys. 57 (1977) 866
- 1977OK1A Oka and Kubodera, in Proc., Zurich-Sin. (1977) 329
- 1977OO01 M.A. Oothoudt and G.T. Garvey, Nucl. Phys. A284 (1977) 41
- 1977PE1B Perez-Mendez et al, in Proc., Zurich-Sin (1977) 183
- 1977PI1C Pile, Bent, Debevec and Pollock, Bull. Amer. Phys. Soc. 22 (1977) 590
- 1977PO1A Povh, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 775
- 1977PR05 P.B. Price, J. Stevenson and K. Frankel, Phys. Rev. Lett. 39 (1977) 177
- 1977PR1D Praderie, Boesgaard, Milliard and Pitois, Astrophys. J. 214 (1977) 130
- 1977RO01 E. Rosario-Garcia and R.E. Benenson, Nucl. Phys. A275 (1977) 453
- 1977RO02 P.G. Roos, N.S. Chant, A.A. Cowley, D.A. Goldberg, H.D. Holmgren and R. Woody, III, Phys. Rev. C15 (1977) 69
- 1977SA1K Saltmarsh, Ludemann, Fulmer and Styles, Nucl. Instrum. Meth. 145 (1977) 81
- 1977SC1D Schramm and Wagoner, Ann. Rev. Nucl. Sci. 27 (1977) 37
- 1977SC1F Schoch et al, Bull. Amer. Phys. Soc. 22 (1977) 591
- 1977SE09 F. Seiler, Nucl. Phys. A286 (1977) 1
- 1977SE1D Seya et al, in Tokyo (1977), Contrib. Papers p. 149
- 1977SH1C Shoda et al, Proc. Int. Conf. Nucl, Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 482
- 1977SI1D Silberberg and Tsao, Astrophys. J. Suppl. 35 (1977) 129
- 1977SP1B Spencer, in High Energy Phys. Nucl. Struct., Zurich (Birkhauser Verlag, 1977) p. 153
- 1977ST1G Steiner, in High Energy Phys. Nucl. Struct., Zurich (Birkhauser Verlag, 1977) p. 261
- 1977ST34 R.G. Stokstad, M.N. Namboodiri, E.T. Chulick, J.B. Natowitz and D.L. Hanson, Phys. Rev. C16 (1977) 2249
- 1977SU1A Subotic, Lalovic and Stepancic, Fizika (Yugoslavia) 9, Suppl. 1 (1977) 43; Phys. Abs. 17217 (1978)
- 1977TE1A Terrien, in Tokyo (1977), Contrib. Papers p. 843
- 1977TU1C Turck et al, in Tokyo (1977), Contrib. Papers p. 152
- 1977WA05 C.N. Waddell, E.M. Diener, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, A.G. Pieper, R.B. Theus, C.C. Chang and N.S. Chant, Nucl. Phys. A281 (1977) 418

- 1977WA08 A.H. Wapstra and K. Bos, *At. Data Nucl. Data Tables* 19 (1977) 175; Erratum *At. Data Nucl. Data Tables* 20 (1977) 1
- 1977WA1M Wang, Chant, Cowley and Roos, in Tokyo (1977), *Contrib. Papers* p. 568
- 1977WE03 R.B. Weisenmiller, N.A. Jelley, D. Ashery, K.H. Wilcox, G.J. Wozniak, M.S. Zisman and J. Cerny, *Nucl. Phys. A280* (1977) 217
- 1977WE1B Weisenmiller, Thesis, Univ. California, Berkeley (1976) - LBL-5077 (1977)
- 1977WE1D Weller, Furst, Tombrello and Burnett, *Astrophys. J.* 214 (1977) L39
- 1977YA1B Yakovlev, *Fiz. Elem. Chastits At. Yadra* 8 (1977) 255
- 1977YO02 R.C. York and R.T. Carpenter, *Nucl. Phys. A282* (1977) 351
- 1978AJ02 F. Ajzenberg-Selove, E.R. Flynn and O. Hansen, *Phys. Rev. C17* (1978) 1283
- 1978AN07 I. Angeli and M. Csatlos, *Atomki Kozlem.* 20 (1978) 1
- 1978AT01 M. Atarashi, K. Hira and H. Narumi, *Prog. Theor. Phys.* 60 (1978) 209
- 1978AU1C Austin, *Bull. Amer. Phys. Soc.* 23 (1978) 609
- 1978BA1J Back et al, *Bull. Amer. Phys. Soc.* 23 (1978) 958
- 1978BA1L Baker et al, *Bull. Amer. Phys. Soc.* 23 (1978) 592
- 1978BI08 R. Billerey, C. Cerruti, A. Chevarier, N. Chevarier and A. Demeyer, *Z. Phys. A285* (1978) 389
- 1978BO01 T. Bowles, D.F. Geesaman, R.J. Holt, H.E. Jackson, R.M. Laszewski, J.R. Specht, L.L. Rutledge, Jr., R.E. Segel, R.P. Redwine and M.A. Yates-Williams, *Phys. Rev. Lett.* 40 (1978) 97
- 1978BO31 M. Bouten, E. Flerackers and M.C. Bouten, *Nucl. Phys. A307* (1978) 413
- 1978BR1G T.J. Brodbeck, P. Coddington, J.V. Morris, D. Newton and T. Sloan, *Nucl. Phys. B136* (1978) 95
- 1978BU01 A. Buchnea, R.G. Johnson and K.G. McNeill, *Can. J. Phys.* 56 (1978) 47
- 1978BY1A Byrnak, *Nuovo Cim.* A44 (1978) 1
- 1978BY1B Byrd, Walter and Cotanch, *Bull. Amer. Phys. Soc.* 23 (1978) 526
- 1978BY1C Byrd, Walter and Cotanch, *Bull. Amer. Phys. Soc.* 23 (1978) 926
- 1978CH02 H.C. Cheung, M.D. High and B. Cujec, *Nucl. Phys. A296* (1978) 333
- 1978CH07 J.-C. Chou, C.-M. Fou, C.-S. Lin, P.-S. Song, M. Wen and Y.-C. Liu, *J. Phys. Soc. Jpn.* 44 (1978) 1
- 1978CH1H Chrien et al, in *Proc. Cluster, Winnipeg* (1978) D5
- 1978DA1C Datta et al, in *Proc., Cluster, Winnipeg* (1978) F6
- 1978DE15 P. Desgrolard, P.A.M. Guichon and J. Joseph, *Nuovo Cim.* A43 (1978) 475
- 1978DE1M de Vries, *Bull. Amer. Phys. Soc.* 23 (1978) 624

- 1978DI04 S. Di Liberto, F. Meddi, G. Romano, G. Rosa and C. Sgarbi, Nucl. Phys. A296 (1978) 519
- 1978DO02 P. Dougan, Z. Phys. A284 (1978) 165
- 1978DW1A Dwek, Astrophys. J. 225 (1978) L149
- 1978EN06 P.M. Endt and C. van der Leun, Nucl. Phys. A310 (1978) 96
- 1978ER1C Ericson, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 27
- 1978FA1D Fallica, Riggi, Spitaleri and Sutera, Lett. Nuovo Cim. 22 (1978) 547
- 1978FO1G Fonseca, Revai and Matveencko, in Proc. Cluster, Winnipeg (1978) B1
- 1978FO1J Fonseca and Revai, Bull. Amer. Phys. Soc. 23 (1978) 498
- 1978FR1D S. Frankel, W. Frati, G. Blanpied, G.W. Hoffmann, T. Kozlowski, C. Morris, H.A. Thiessen, O. Van Dyck, R. Ridge and C. Whitten, Phys. Rev. C18 (1978) 1375
- 1978GE1C Gelbke et al, Phys. Rept. C42 (1978) 311
- 1978GE1E Geesaman et al, Bull. Amer. Phys. Soc. 23 (1978) 593
- 1978GO1G Goodman et al, Bull. Amer. Phys. Soc. 23 (1978) 526
- 1978GR07 V.I. Grantsev, I.P. Dryapachenko, V.A. Kornilov, O.F. Nemets, V.V. Ostashko, V.M. Pugach, V.A. Pilipchenko, B.A. Rudenko, M.V. Sokolov and B.G. Struzhko, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 147; Bull. Acad. Sci. USSR Phys. Ser. 42 (1978) 126
- 1978GR1F Green and Korteling, in Proc., Cluster, Winnipeg (1978) 89
- 1978GU15 O.P. Gupta, A. Bond and P.M.S. Lesser, Phys. Rev. C18 (1978) 1075
- 1978HA04 M.Y.M. Hassan, E.H. Ismail and A. Rabie, Z. Phys. A285 (1978) 37
- 1978HO1E Horiuchi, 3rd Int. Conf. on Clustering Aspects of Nucl. Struct. & Nucl. Rxns, Winnipeg (1978)
- 1978KI08 H.R. Kissener, G.E. Dogotar, R.A. Eramzhyan and R.A. Sakaev, Nucl. Phys. A302 (1978) 523
- 1978LA1G Lang et al, in Proc., Cluster, Winnipeg (1978) F2
- 1978LE04 H.C. Lee, Nucl. Phys. A294 (1978) 473
- 1978LO1C Lone, Ferguson and Earle, Bull. Amer. Phys. Soc. 23 (1978) 500
- 1978MA1L Mateja et al, Bull. Amer. Phys. Soc. 23 (1978) 616
- 1978MC04 A.B. McDonald and E.G. Adelberger, Phys. Rev. Lett. 40 (1978) 1692
- 1978MI1D Milder et al, Bull. Amer. Phys. Soc. 23 (1978) 611
- 1978MO01 C.L. Morris, R.L. Boudrie, J.J. Kraushaar, R.J. Peterson, R.A. Ristinen, G.R. Smith, J.E. Bolger, W.J. Braithwaite, C.F. Moore and L.E. Smith, Phys. Rev. C17 (1978) 227

- 1978NA05 K. Nakamura, S. Hiramatsu, T. Kamae, H. Muramatsu, N. Izutsu and Y. Watase, Nucl. Phys. A296 (1978) 431
- 1978NE1A Needham et al, Bull. Amer. Phys. Soc. 23 (1978) 943
- 1978OC1B O'Connell, Dressler and MacDonald, Bull. Amer. Phys. Soc. 23 (1978) 507
- 1978OK01 S. Okabe and Y. Abe, Prog. Theor. Phys. 59 (1978) 315
- 1978OK1B Okabe and Abe, in Proc., Cluster, Winnipeg (1978) B10
- 1978OL1A Olmer et al, Bull. Amer. Phys. Soc. 23 (1978) 592
- 1978OS01 A. Osman, Phys. Rev. C17 (1978) 341
- 1978PA1B Park, Greiner and Scheid, Bull. Amer. Phys. Soc. 23 (1978) 538
- 1978RI02 P.J. Riley, C.W. Bjork, C.R. Newsom, R.A. Kenefick, M.L. Evans, G. Glass, J. Hiebert, M. Jain, L.C. Northcliffe, B.E. Bonner et al, Phys. Rev. C17 (1978) 1881
- 1978SE1D Seya et al, in Tokyo (1977) Contrib. Papers p. 149
- 1978SH1E Shirley and Lederer, in Table of Isotopes, 7th Edition, John Wiley (1978)
- 1978SP03 C. Spitaleri, M. Lattuada, F. Riggi, N. Arena and D. Vinciguerra, Lett. Nuovo Cim. 21 (1978) 345
- 1978TH1B Thiessen, Bull. Amer. Phys. Soc. 23 (1978) 940
- 1978UN01 J. Unternahrer, J. Lang and R. Muller, Phys. Rev. Lett. 40 (1978) 1077
- 1978WA1D Walter et al, Bull. Amer. Phys. Soc. 23 (1978) 926
- 1978WE1D Wessels et al, Nucl. Instrum. Meth. 141 (1977) 157
- 1978WO13 R.M. Woloshyn, Nucl. Phys. A306 (1978) 333
- 1978ZE01 B. Zeidman, C. Olmer, D.F. Geesaman, R.L. Boudrie, R.H. Siemssen, J.F. Amann, C.L. Morris, H.A. Thiessen, G.R. Burlison, M.J. Devereux et al, Phys. Rev. Lett. 40 (1978) 1316; Erratum Phys. Rev. Lett. 40 (1978) 1539
- 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, Nucl. Phys. A336 (1980) 1
- 1981AJ01 F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1