

Energy Levels of Light Nuclei $A = 9$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A*413 (1984), p. 1. This version of $A = 9$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and Introductory tables have been omitted from this manuscript. Also, [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed June 1, 1983)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 9$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\${}^9\text{n}\$](#) , [\${}^9\text{He}\$](#) , [\${}^9\text{Li}\$](#) , [\${}^9\text{Be}\$](#) , [\${}^9\text{B}\$](#) , [\${}^9\text{C}\$](#) , [\${}^9\text{N}\$](#)

B. Tables of Recommended Level Energies:

[Table 9.1](#): Energy levels of ${}^9\text{Li}$

[Table 9.2](#): Energy levels of ${}^9\text{Be}$

[Table 9.9](#): Energy levels of ${}^9\text{B}$

[Table 9.12](#): Energy levels of ${}^9\text{C}$

C. [References](#)

D. Figures: [\${}^9\text{Li}\$](#) , [\${}^9\text{Be}\$](#) , [\${}^9\text{B}\$](#) , [Isobar diagram](#)

⁹n
(Not illustrated)

Not observed: see (1977DE08).

⁹He
(Not illustrated)

⁹He has been observed in the ⁹Be(π^- , π^+)⁹He reaction at $E_{\pi^-} = 194$ MeV; the atomic mass excess is 40.81 ± 0.12 MeV. ⁹He is then unstable with respect to decay into ⁸He + n by 1.14 MeV (1981SE1B, 1980NA1D, 1980SE1C, 1980SE1F). See also (1979AJ01) and (1982PO1C; hypernuclei) and (1982NG01; theor.).

⁹Li
(Figs. 15 and 18)

GENERAL: (See also (1979AJ01).)

Model calculations: (1979LA06).

Complex reactions involving ⁹Li: (1978DU1B, 1979AL22, 1979BO22, 1979JA1C, 1980BO31, 1980WI1L, 1981BO1X, 1981MO20, 1982BO1Y).

Muon and neutrino capture and reactions: (1980MU1B).

Reactions involving pions and other mesons (See also reaction 3.): (1978FU09, 1979BO21, 1979PE1C, 1979WI1E, 1980NI03, 1980ST15, 1981YA1A).

Hypernuclei: (1978DA1A, 1978PO1A, 1980MA1Z, 1981ST1G, 1981WA1J, 1982KO11).

Other topics: (1979BE1H, 1979LA06, 1982NG01).

Ground-state properties of ⁹Li: (1981AV02, 1982NG01).

$\mu = 3.4391 \pm 0.0006$ nm: see (1983CO1Q).

$Q = 0.88 \pm 0.18$ Q(⁷Li): see (1983CO1Q).

1. ⁹Li(β^-)⁹Be

$Q_m = 13.606$

Table 9.1: Energy levels of ${}^9\text{Li}$

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions ^a
g.s.	$\frac{3}{2}^-; \frac{3}{2}$	$\tau_{1/2} = 178.3 \pm 0.4$ msec	β^-	1, 2, 3, 4, 5, 6
2.691 ± 5	$(\frac{1}{2}^-)$		(γ)	2, 6
4.31 ± 20		$\Gamma = 100 \pm 30$		2, 6
5.38 ± 60		600 ± 100		2
6.43 ± 15	$\geq \frac{9}{2}$	40 ± 20		2, 6

^a For other reactions on which no new work is reported, see (1979AJ01).

The half-life of ${}^9\text{Li}$ is 178.3 ± 0.4 msec (1976AL02). See also (1979AJ01). ${}^9\text{Li}$ decays to a number of states in ${}^9\text{Be}$: see reaction 12 in ${}^9\text{Be}$ and Table 9.7. The nature of the decay to ${}^9\text{Be}^*(0, 2.43)$ with $J^\pi = \frac{3}{2}^-, \frac{5}{2}^-$ is evidence for $J^\pi = \frac{3}{2}^-$ for ${}^9\text{Li}_{\text{g.s.}}$. The probability for delayed neutron decay, P_n , is $(49.5 \pm 5)\%$ (1981LA11): see Table 9.7.

2. ${}^7\text{Li}(t, p){}^9\text{Li}$ $Q_m = -2.386$

Protons are observed to excited states at $E_x = 2.691 \pm 0.005$ MeV (1964MI04), 4.31 ± 0.03 , 5.38 ± 0.06 and 6.41 ± 0.02 MeV (1971YO04) [$\Gamma_{\text{c.m.}} = 250 \pm 30, 600 \pm 100, < 100$ keV], 4.31 ± 0.02 and 6.435 ± 0.020 MeV (1978AJ02) [$\Gamma_{\text{c.m.}} = 100 \pm 30, 40 \pm 20$ keV]. Angular distributions are reported at $E_t = 11.3$ (1964MI04; t_0), 15 MeV (1971YO04: t_0, t_2, t_4) and at 23 MeV (1978AJ02: t_0, t_1, t_2, t_4). They are consistent with $J^\pi = \frac{3}{2}^-, (\frac{1}{2}^-)$ and $\geq \frac{9}{2}$ for ${}^9\text{Li}^*(0, 2.69, 6.43)$ (1978AJ02). See also (1979AB11, 1980AB16, 1983AB1A, 1983CO11) and ${}^{10}\text{Be}$.

3. ${}^9\text{Be}(\gamma, \pi^+){}^9\text{Li}$ $Q_m = -153.174$

At $E_\gamma = 175, 187$ and 200 MeV, the cross section for production of ${}^9\text{Li}_{\text{g.s.}}$ at $\theta_\pi = 90^\circ$ has been measured by (1982TE01). See also (1979BO21, 1980NI03).

4. ${}^9\text{Be}(n, p){}^9\text{Li}$ $Q_m = -12.824$

See (1981BR1K) and (1979AJ01).

5. ${}^9\text{Be}({}^7\text{Li}, {}^7\text{Be}){}^9\text{Li}$ $Q_m = -14.468$

See (1982AL1G: $E(^7\text{Li}) = 78 \text{ MeV}$).

$$6. \text{}^{11}\text{B}(\text{}^6\text{Li}, \text{}^8\text{B})\text{}^9\text{Li} \quad Q_m = -25.121$$

At $E(^6\text{Li}) = 80 \text{ MeV}$ the angular distribution to $^9\text{Li}_{\text{g.s.}}$ has been measured. States at $E_x = 2.59 \pm 0.10, 4.36 \pm 0.10$ and $6.38 \pm 0.12 \text{ MeV}$ are also populated (1977WE03, 1977WE1B).

⁹Be
(Figs. 16 and 18)

GENERAL: (See also (1979AJ01).)

Shell model: (1978AR1H, 1979LA06, 1981BO1Y, 1982OR03).

Cluster and α -particle models: (1978AR1H, 1978RE1A, 1979CH1D, 1979FO16, 1979LU1A, 1979OK02, 1980FU1G, 1980RE02, 1981BA2N, 1981KR1J, 1982LA11, 1982OR03, 1982PO1B).

Collective and rotational models: (1979MA1J, 1981BA2N).

Special states: (1979CH1D, 1979KI10, 1979OK02, 1980FU1G, 1981BO1Y, 1981SE06, 1982OR03, 1983GO1R).

Electromagnetic transitions and giant resonance: (1979DO17, 1980MI1G, 1981BO1Y, 1981KO1Q, 1982AW02).

Astrophysical questions: (1978BU1B, 1978OR1A, 1979MC1A, 1980CO1R, 1980RE1B, 1980WI1M, 1981AU1D, 1981AU1G, 1981GA1C, 1981SC1M).

Applied work: (1979FO1F, 1979GR1E, 1979JU1B, 1980KI1B, 1981FA1E, 1981SA1G, 1981SC1D, 1982KL1A, 1982KU1G).

Complex reactions involving ⁹Be: (1978BE1G, 1978BH03, 1978DU1B, 1978HE1C, 1979AL22, 1979BA10, 1979BA34, 1979BO22, 1979BU1G, 1979DY01, 1979FR12, 1979GA04, 1979GE1A, 1979JA1C, 1979LO11, 1979SA17, 1979SA27, 1979SA1E, 1979SC08, 1979SC1D, 1979VI05, 1979WE06, 1980BR1P, 1980GR10, 1980ME1F, 1980MI01, 1980MO28, 1980WI1L, 1981BH02, 1981BL1G, 1981CI03, 1981HN02, 1981LE03, 1981ME13, 1981MO20, 1981TA16, 1981TA22, 1982BO1J, 1982BO35, 1982GU1H, 1982JA1C, 1982LU01, 1982LY1A, 1982MO1N, 1982WU1B, 1983FU04, 1983SA06).

Muon and neutrino capture and reactions: (1977GR1C, 1980MU1B, 1980SC18, 1981MU1E).

Reactions involving pions (See also reaction I7): (1978AR1J, 1978ER1A, 1978FU09, 1978GE11, 1978KI13, 1978VL1B, 1978WA1B, 1979AL1J, 1979BA16, 1979BO21, 1979BO1N, 1979BU1D, 1979CO1D, 1979DE06, 1979DE1G, 1979FR1F, 1979FU1G, 1979GI1B, 1979GI1C, 1979HO1C, 1979HO13, 1979IN1A, 1979KI1G, 1979MA02, 1979MA1P, 1979NA1E, 1979OH1A, 1979PE1D, 1979PI1C, 1979SC1E, 1979ST1G, 1979TI1A, 1979WI1E, 1980BU15, 1980CR03, 1980ER01, 1980HA29, 1980JE1C, 1980LE02, 1980MA1F, 1980NA1D, 1980NA1L, 1980NI03, 1980ST15, 1980TH1C, 1980TR1A, 1981AS1H, 1981BA2M, 1981BA2R, 1981BE13, 1981BE1X, 1981BE63, 1981BO06, 1981DU1H, 1981FE2A, 1981GE03, 1981IO01, 1981IS11, 1981KU15, 1981LE14, 1981MC09, 1981NA05, 1981NI03, 1981SE1B, 1981WA1H, 1981YA1A, 1982AP1A, 1982AS01, 1982GR1F, 1982MU1C, 1983AS02).

Reactions involving kaons and other mesons: (1978AN1B, 1978DA1A, 1978SO1A, 1979GA1E, 1979LI1A, 1980AU1C, 1980BE30, 1980BE12, 1980DE11, 1980DO1E, 1980GA1C, 1980PO1A,

1981BA2N, 1981BE17, 1981BE45, 1981BO09, 1981DA1C, 1981HU1C, 1981KA1U, 1981PO1D, 1981WA1H, 1981ZH1C, 1982AB07, 1982BA17, 1982BO1U, 1982DEZQ, 1982DO1C, 1982ER1E, 1982PI1J, 1982PO1C).

Table 9.2: Energy levels of ${}^9\text{Be}$

E_x ^a (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$		stable	2, 3, 4, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45
1.685 ± 15	$\frac{1}{2}^+$	≈ 150	γ, n	9, 10, 13, 16, 18, 19, 21, 23, 24, 32, 33, 36, 40, 41
2.4294 ± 1.3	$\frac{5}{2}^-$	0.77 ± 0.15	γ, n, α	9, 10, 12, 16, 17, 18, 19, 21, 22, 23, 24, 26, 32, 35, 36, 39, 40, 41, 42, 43
2.78 ± 120	$\frac{1}{2}^-$	1080 ± 110	n	9, 12, 42
3.049 ± 9	$\frac{5}{2}^+$	282 ± 11	γ, n	9, 13, 16, 18, 19, 20, 21, 23, 24, 32, 36, 39, 40, 41
4.704 ± 25	$(\frac{3}{2})^+$	743 ± 55	γ, n	9, 16, 19, 20, 21, 23, 24, 34, 40, 42
6.76 ± 60	$\frac{7}{2}^-$	1540 ± 200	γ, n	9, 16, 17, 18, 19, 20, 21, 23, 24, 35, 41
7.94 ± 80	$(\frac{1}{2}^-)$	≈ 1000		12, 19
11.283 ± 24	$(\frac{3}{2}^-)$	575 ± 50	n	9, 12, 19, 24, 35, 36
11.81 ± 20	$T = \frac{1}{2}$	400 ± 30	γ, n	9, 13, 39, 42
13.79 ± 30	$T = \frac{1}{2}$	590 ± 60	γ, n	9, 16, 39
14.3929 ± 1.8	$\frac{3}{2}^-; \frac{3}{2}$	0.381 ± 0.033	γ, n, α	9, 16, 23, 36, 39
14.4 ± 300		≈ 800		19, 36
15.10 ± 50			γ	16, 39
15.97 ± 30	$T = \frac{1}{2}$	≈ 300	γ	16, 39
16.671 ± 8		41 ± 4	γ	9, 16, 19, 36

Table 9.2: Energy levels of ${}^9\text{Be}$ (continued)

E_x ^a (MeV \pm keV)	J^π, T	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
16.975 \pm 2	$\frac{1}{2}^-; \frac{3}{2}$	< 0.47	γ, n, p, d	4, 5, 6, 16
17.298 \pm 7	$(\frac{5}{2})^-$	200	γ, n, p, d, α	5, 6, 7, 13, 16, 19
17.493 \pm 7	$\leq \frac{7}{2}^+$	47	γ, n, p, d, α	5, 6, 7, 16, 19
18.02 \pm 50			γ	16
18.58 \pm 40			γ, n, p, d, α	6, 16
19.20 \pm 50		310 \pm 80	n, p, d, t	1, 6, 19
19.51 \pm 50			γ	13, 16
(19.9 \pm 200)			γ, n	13, 19
(20.47 \pm 40)			γ, p, d	13, 19
20.74 \pm 30		\approx 1000	γ, n, p, t	1, 13, 16
(21.4 \pm 200)			γ, n	13, 19
(22.4 \pm 200)		broad	γ, n	13, 19
(23.8 \pm 200)			γ, n	13, 34
(27.0 \pm 500)		broad	γ, n	13
b				

^a See also reactions 14 and 16.

^b See footnote ⁱ in Table 9.8.

Hypernuclei: (1977KO1D, 1978DA1A, 1978PO1A, 1978SO1A, 1979GA1E, 1979VE1B, 1980AU1C, 1980BE30, 1980BE12, 1980DO1E, 1980GA1P, 1980GI1C, 1980PO1A, 1980ZH1C, 1980ZO1A, 1981BA2N, 1981BE17, 1981BE45, 1981BO09, 1981DA1C, 1981DA1D, 1981PO1D, 1981PO1F, 1981RE1B, 1981ST1G, 1981SU1D, 1981WA1J, 1982BA1P, 1982BA17, 1982BO1U, 1982DA1R, 1982DEZQ, 1982DO1C, 1982DO1M, 1982ER1B, 1982ER1E, 1982GA1J, 1982IK1A, 1982JO1C, 1982KO11, 1982MO1Q, 1982PO1C, 1982SH1P, 1982ZH1F, 1983BA2K, 1983JO1E).

Other topics: (1978OS1B, 1979BE1H, 1979LA06, 1979LI1A, 1979OS02, 1980BO1N, 1980GI1C, 1981SE06, 1982AW02, 1982BA2G, 1982NG01, 1983BO1E, 1983GO1R).

Ground-state properties of ${}^9\text{Be}$: (1978HE1D, 1979AL26, 1979OK02, 1979SA27, 1980BO1N, 1981AV02, 1981BO1Y, 1981HN02, 1981SE06, 1982BA2G, 1982DE35, 1982LO13, 1982NG01, 1982OR03, 1983BO1E).

$\mu = -1.1778 \pm 0.0009$ nm: see (1976WE17).

$Q = +53 \pm 3$ mb: see (1978LEZA).

1. (a) ${}^6\text{Li}(t, n){}^8\text{Be}$	$Q_m = 16.0223$	$E_b = 17.6878$
(b) ${}^6\text{Li}(t, p){}^8\text{Li}$	$Q_m = 0.801$	
(c) ${}^6\text{Li}(t, d){}^7\text{Li}$	$Q_m = 0.9929$	
(d) ${}^6\text{Li}(t, \alpha){}^5\text{He}$	$Q_m = 15.22$	
(e) ${}^6\text{Li}(t, n){}^4\text{He}{}^4\text{He}$	$Q_m = 16.1141$	
(f) ${}^6\text{Li}(t, 2n){}^7\text{Be}$	$Q_m = -2.876$	

The 0° differential cross section for reaction (a) increases monotonically between $E_t = 0.10$ and 2.4 MeV except for a resonance at $E_t = 1.875$ MeV (${}^9\text{Be}^* = 18.937$). The excitation function for ${}^8\text{Li}$ (reaction (b)) increases monotonically for $E_t = 0.275$ to 1.000 MeV. In the range $E_t = 2.0$ to 6.8 MeV, a broad peak [$\Gamma \approx 1.3$ MeV] is observed at $E_t = 4.57$ MeV [$E_x = 20.73$ MeV]: see (1974AJ01) for references. For reaction (d) see (1982CE1C). Cross sections for ${}^7\text{Be}$ production [reaction (f)] ($E_t = 4.4$ to 11.99 MeV) are reported by (1980GU26). For reactions (c) and (e) see (1966LA04).

2. ${}^6\text{Li}({}^3\text{He}, \pi^+){}^9\text{Be}$	$Q_m = -121.898$
---	------------------

See (1981BI1F, 1982LE1L).

3. ${}^6\text{Li}(\alpha, p){}^9\text{Be}$	$Q_m = -2.1262$
--	-----------------

Angular distributions of p_0 have been measured at $E_\alpha = 10.2$ to 14.7 MeV and at 30 MeV: see (1974AJ01). See also (1983BE1P; theor.).

4. ${}^7\text{Li}(d, \gamma){}^9\text{Be}$	$Q_m = 16.6950$
--	-----------------

For $E_d = 0.1$ to 1.1 MeV, a resonance in the yield of capture γ -rays is observed at $E_d = 361 \pm 2$ keV, corresponding to ${}^9\text{Be}^*(16.975)$ with $\Gamma_{c.m.} < 0.47$ keV. The small width of this state and its energy correspondence with ${}^9\text{Li}^*(2.69)$ argue for $T = \frac{3}{2}$. The angular distribution of the γ -rays to ${}^9\text{Be}(0)$ is isotropic to within 7%: see (1974AJ01) for references.

5. (a) ${}^7\text{Li}(d, n){}^8\text{Be}$	$Q_m = 15.0296$	$E_b = 16.6950$
(b) ${}^7\text{Li}(d, \alpha){}^5\text{He}$	$Q_m = 14.23$	
(c) ${}^7\text{Li}(d, n){}^4\text{He}{}^4\text{He}$	$Q_m = 15.1213$	
(d) ${}^7\text{Li}(d, 2n){}^7\text{Be}$	$Q_m = -3.869$	

Table 9.3: Resonances in ${}^7\text{Li} + \text{d}$ ^a

${}^7\text{Li}(\text{d}, \text{n})$ E_{res} (keV)	${}^7\text{Li}(\text{d}, \alpha)$ E_{res} (keV)	${}^7\text{Li}(\text{d}, \text{p})$		E_x (MeV)
		E_{res} (keV)	Γ_{lab} (keV)	
361 ± 2 ^b		360 ± 3	< 0.5 ^b	16.975
680	750	776 ± 7 ^c	250	17.298
980	1000	1027 ± 7 ^c	60	17.493
		2000	broad	(18.5)
		2375 ± 50		18.54
		3220 ± 50	400 ± 100	19.20
		≈ 4800		20.4

^a For references and additional information see Tables 9.3 in (1974AJ01, 1979AJ01).

^b E_{res} from ${}^7\text{Li}(\text{d}, \gamma)$: $\Gamma_{\text{n}\alpha}/\Gamma_{\gamma} \approx 1.5$, $\Gamma_{\text{d}\alpha}/\Gamma_{\gamma} < 400$, $\Gamma_{\alpha\alpha}/\Gamma_{\gamma} < 20$.

^c (1976SC14, 1979MI1E).

The yield of neutrons has been measured for $E_{\text{d}} = 0.2$ to 23 MeV: observed resonances are displayed in Table 9.3. For polarization measurements [$E_{\text{d}} = 0.64$ MeV and 2.5 to 3.7 MeV] see (1974AJ01). See also (1979SE04; theor.).

The yields of α -particles have been measured for $E_{\text{d}} = 0.25$ to 0.75 MeV (1977RO1C; α_1) and 0.2 to 3.0 MeV: see Table 9.3. For reaction (c) see (1974AJ01) and ${}^5\text{He}$. The ${}^7\text{Be}$ production cross section (reaction (d)) has been studied for $E_{\text{d}} = 5.38$ to 11.94 MeV (1980GU26). See also (1978BA1F, 1979JO1B; applications).

6. ${}^7\text{Li}(\text{d}, \text{p}){}^8\text{Li}$

$$Q_{\text{m}} = -0.192$$

$$E_{\text{b}} = 16.6950$$

Excitation functions have been measured for $E_{\text{d}} = 0.29$ to 7 MeV [see (1974AJ01, 1979AJ01)] and 0.60 to 1.2 MeV (1982EL03, 1982FI03; σ_{t}) and 0.7 to 3.4 MeV (1979MI1E; σ_{t}): observed resonances are displayed in Table 9.3. Recent measurements of the total cross section at the $E_{\text{d}} = 0.78$ MeV resonance are 181 ± 8 mb (1976SC14), 174 ± 16 mb (1979MI1E), 146 ± 13 mb (1982EL03) and 148 ± 12 mb (1982FI03): we adopt 147 ± 9 mb. See also (1983FI01). For earlier values see (1974AJ01). This cross section serves as normalization for the ${}^7\text{Be}(\text{p}, \gamma){}^8\text{B}$ data and to calculations of the S -factor. The new value predicts that the neutrino-capture rate in ${}^{37}\text{Cl}$ should be decreased by about 15%.

Polarization measurements are reported at $E_{\text{d}} = 4$ MeV (1982WAZS). See also (1980HU1D, 1982AB1D) and ${}^8\text{Li}$.

Table 9.4: Excited states of ${}^9\text{Be}$ from ${}^7\text{Li}({}^3\text{He}, \text{p}){}^9\text{Be}$ ^a

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)
1.64	
2.4292 ± 1.7	< 8
2.9 ± 250	1000 ± 250
3.076 ± 15	289 ± 22
4.704 ± 25	743 ± 55
6.7 ± 100	2000 ± 200
11.29 ± 30	620 ± 70
11.81 ± 20	400 ± 30
13.78 ± 30	590 ± 60
14.396 ± 5 ^b	0.38 ± 0.03
16.671 ± 8	41 ± 4

^a See also Tables 9.4 in (1974AJ01, 1979AJ01) for references.

^b See also Table 9.6.

7. ${}^7\text{Li}(\text{d}, \text{d}){}^7\text{Li}$

$$E_b = 16.6950$$

The elastic scattering [$E_d = 0.4$ to 1.8 MeV] shows a marked increase in cross section for $E_d = 0.8$ to 1.0 MeV (perhaps related to ${}^9\text{Be}^*(17.30)$) and a conspicuous anomaly at $E_d = 1.0$ MeV, due to p-wave deuterons [${}^9\text{Be}^*(17.50)$]. The elastic scattering has also been studied for $E_d = 1.0$ to 2.6 MeV and 10.0 to 12.0 MeV: see (1979AJ01). See also (1982AB1D) and ${}^7\text{Li}$.

8. (a) ${}^7\text{Li}(\text{d}, \text{t}){}^6\text{Li}$

$$Q_m = -0.993$$

$$E_b = 16.6950$$

(b) ${}^7\text{Li}(\text{d}, {}^3\text{He}){}^6\text{He}$

$$Q_m = -4.481$$

The cross section for reaction (a) rises steeply from threshold to 95 mb at $E_d = 2.4$ MeV and then more slowly to ≈ 165 mb at $E_d = 4.1$ MeV. The t_0 yield curve ($\theta_{\text{lab}} = 155^\circ$) decreases monotonically for $E_d = 10.0$ to 12.0 MeV: see (1974AJ01). See also ${}^6\text{Li}$. For reaction (b) see ${}^6\text{He}$.

9. ${}^7\text{Li}({}^3\text{He}, \text{p}){}^9\text{Be}$

$$Q_m = 11.2015$$

Table 9.5: Neutron decay of ${}^9\text{Be}$ states ^a

${}^9\text{Be}$ state (MeV)	l_n	Decay (in %) to		θ^2 (%) ^b
		${}^8\text{Be}(0)$	${}^8\text{Be}^*(2.9)$	
2.43	3	7.0 ± 1.0 ^a		2.1 ± 0.6
2.78	1	mainly		0.48 ± 0.06
3.05	2	87 ± 13		81 ± 13
4.70	2	13 ± 4		6.0 ± 0.4
6.76	3	≤ 2		≤ 6
	1		55 ± 14	37 ± 10
11.28	1	≤ 2		≤ 0.1
	1		14 ± 4	0.93 ± 0.28
	3			4.0 ± 1.2
11.81	1	≤ 3		≤ 0.1
	1		12 ± 4	0.48 ± 0.16
	3			1.8 ± 0.6
14.39 ^c				

^a For references see Table 9.5 in (1979AJ01).

^b Expressed in units of $\hbar^2/mR^2 = 2.47$ MeV (1968CO08, 1970CH07).

^c See Table 9.6.

Observed proton groups are listed in Table 9.4. The parameters for the particle and γ -decay of observed states are displayed in Tables 9.5 and 9.6 Angular distributions have been reported in the range $E({}^3\text{He}) = 0.9$ to 14 MeV: see (1974AJ01, 1979AJ01). See also (1981SL03), (1981VI1B) and ${}^{10}\text{B}$.

The γ -decay of ${}^9\text{Be}^*(14.39)$ [$J^\pi = \frac{3}{2}^-$; $T = \frac{3}{2}$] has been measured by (1978DI08) and compared with the decay from the mirror state in ${}^9\text{B}$: see Table 9.6. No evidence is found for violation of the selection rule for $\Delta T = 1$ mirror γ transitions (1978DI08).

$$10. {}^7\text{Li}(\alpha, d){}^9\text{Be} \quad Q_m = -7.1517$$

Angular distributions of d_0 , d_1 and d_2 are reported at $E_\alpha = 30$ MeV: see (1974AJ01). See also (1980ZE05, 1983BE1P; theor.).

$$11. {}^7\text{Li}({}^6\text{Li}, \alpha){}^9\text{Be} \quad Q_m = 15.220$$

Table 9.6: Parameters ^a of the first $T = \frac{3}{2}$ states in ⁹Be and ⁹B, $J^\pi = \frac{3}{2}^-$

	⁹ Be		⁹ B
E_x (keV)	14392.9 ± 1.8		14655.0 ± 2.5
Γ_{γ_0} (eV)	6.9 ± 0.5		$(6.9 \pm 0.5)^b$
Γ (eV)	381 ± 33^c		395 ± 42^d
$\Gamma_{\gamma_0}(\text{to } \frac{3}{2}^-)/\Gamma(\%)$	1.81 ± 0.09^d		1.85 ± 0.15^d
$\Gamma_{\gamma_1}(\text{to } \frac{1}{2}^+)/\Gamma(\%)$	0.03 ± 0.04^d		0.00 ± 0.08^d
$\Gamma_{\gamma_2}(\text{to } \frac{5}{2}^-)/\Gamma(\%)$	2.05 ± 0.11^d		1.93 ± 0.22^d
$\Gamma_{\gamma_3}(\text{to } \frac{1}{2}^-)/\Gamma(\%)$	$< 0.2^d$		} 0.31 ± 0.18^d
$\Gamma_{\gamma_4}(\text{to } \frac{5}{2}^+)/\Gamma(\%)$	0.33 ± 0.07^d		
$\Gamma_{\gamma_5}(\text{to } \frac{3}{2}^+)/\Gamma(\%)$	0.23 ± 0.05^d		
$\Gamma_{\gamma_2}/\Gamma_{\gamma_0}$	1.13 ± 0.05^d		1.03 ± 0.11^d
Γ_{n_0}/Γ	0.028 ± 0.021	Γ_{p_0}/Γ	0.11 ± 0.04
Γ_{n_1}/Γ	0.50 ± 0.11	Γ_{p_1}/Γ	0.33 ± 0.09
Γ_{n_0} (eV)	9 ± 8	Γ_{p_0} (eV)	30 ± 17
Γ_{n_1} (eV)	147 ± 28	Γ_{p_1} (eV)	95 ± 15
$\Gamma_{n_1}/\Gamma_{n_0}$	18 ± 14	$\Gamma_{p_1}/\Gamma_{p_0}$	3.2 ± 1.9
$\gamma_{n_1}^2/\Gamma_{n_0}^2$	22 ± 17	$\gamma_{p_1}^2/\gamma_{p_0}^2$	3.2 ± 2.2
$\Gamma_{\alpha_0}/\Gamma_{\gamma_0}$	31.2 ± 9.8		

^a See Table 9.6 in (1979AJ01) for references.

^b Assumed identical to ⁹Be.

^c Calculated from Γ_{γ_0}/Γ and Γ_{γ_0} (1978DI08).

^d (1978DI08): $\Gamma(^9\text{B})/\Gamma(^9\text{Be}) = 1.04 \pm 0.10$.

Table 9.7: Branching parameters in ${}^9\text{Li}$ β -decay (1981LA11)

E_x in ${}^9\text{Be}$ (MeV)	$J^\pi; T$	Branching ratio (%)	$\log ft$ ^a
0	$\frac{3}{2}^-; \frac{1}{2}$	50.5 ± 5 ^c	5.31
2.43	$\frac{5}{2}^-; \frac{1}{2}$	34 ± 4	5.07
2.78 ^b	$\frac{1}{2}^-; \frac{1}{2}$	10 ± 2	5.54
7.94	$(\frac{1}{2}^-)$ ^d ; $\frac{1}{2}$	1.5 ± 0.5	5.04
11.28	$(\frac{3}{2}^-)$ ^d ; $\frac{1}{2}$	4 ± 0.5	2.87 ^e
11.81		< 0.1	> 4.0

^a M.J. Martin, private communication.

^b 2.78 ± 0.12 MeV, $\Gamma_{\text{c.m.}} = 1.10 \pm 0.12$ MeV; $\theta_p^2 = 0.48 \pm 0.06$: see Table 9.7 in (1979AJ01).

^c $P_n = (50 \pm 4)\%$ (1981BJ01); $P_n = (49.5 \pm 5)\%$ (1981LA11).

^d Suggested by (1981LA11) on the basis of the branching ratios.

^e (1981LA11) suggest that the very low $\log ft$ value implies that ${}^9\text{Be}^*(11.28)$ is the anti-analog to ${}^9\text{Li}_{\text{g.s.}}$.

Angular distributions of α_0 have been studied at $E({}^7\text{Li}) = 3.78$ to 5.95 MeV: see (1974AJ01).

$$12. {}^9\text{Li}(\beta^-){}^9\text{Be} \quad Q_m = 13.606$$

${}^9\text{Li}$ decays by β^- emission with $\tau_{1/2} = 178.3 \pm 0.4$ msec to several ${}^9\text{Be}$ states: see ${}^9\text{Li}$, reaction 1 and Table 9.7. Measurement of $\beta - \alpha$ coincidences involving ${}^9\text{Be}^*(11.28)$ show contributions from the direct $n + 2\alpha$ breakup process as well as the sequential n-emission to ${}^8\text{Be}^*(2.9)$ [$J^\pi = 2^+$], followed by breakup into 2α (1981LA11, 1981NY1A). The branching ratio for the ${}^9\text{Be}^*(2.43) \rightarrow {}^8\text{Be}_{\text{g.s.}} + n$ decay is $(6.4 \pm 1.2)\%$. ${}^9\text{Be}^*(2.78)$ [$J^\pi = \frac{1}{2}^-$] decays mainly to ${}^8\text{Be}_{\text{g.s.}} + n$ presumably by p-wave neutron emission: see reaction 13 in (1979AJ01). See also (1981JOZV) and (1979DE15; theor.).

$$13. \begin{aligned} \text{(a)} \quad & {}^9\text{Be}(\gamma, n){}^8\text{Be} & Q_m &= -1.6655 \\ \text{(b)} \quad & {}^9\text{Be}(\gamma, \alpha){}^5\text{He} & Q_m &= -2.46 \\ \text{(c)} \quad & {}^9\text{Be}(\gamma, n){}^4\text{He}{}^4\text{He} & Q_m &= -1.5737 \end{aligned}$$

The photoneutron cross section has been measured from threshold to 320 MeV: see Table 9.6 in (1966LA04) and (1979AJ01). A pronounced peak occurs ≈ 29 keV above threshold with

$\sigma_{\max} = 1.33 \pm 0.24$ mb. The shape of the resonance has been measured very accurately for $E_\gamma = 1675$ to 2168 keV. The FWHM of the peak is estimated to be 100 keV (1982FU11). See also (1983BA52; theor.). The cross section then decreases slowly to 1.2 mb at 40 keV above threshold. From bremsstrahlung studies, peaks in the (γ, Tn) cross section are observed corresponding to $E_x = 1.80$ and 3.03 MeV. At higher energies, using monoenergetic photons, the (γ, Tn) cross section is found to be relatively smooth from $E_\gamma = 17$ to 37 MeV with weak structures which correspond to $E_x = 17.1, 18.8, 19.9, 21.4, 22.4, 23.8 [\pm 0.2]$ MeV and 27 ± 0.5 MeV (broad). In the range $E_\gamma = 18$ to 26 MeV the integrated (γ, n_0) cross section is < 0.1 MeV \cdot mb, that for $(\gamma, n_1) = 2.4 \pm 0.4$ MeV \cdot mb and the combined integrated cross section for (γ, n) to ${}^8\text{Be}^*(16.6)$ and (γ, α_0) to ${}^5\text{He}_{\text{g.s.}}$ is 13.1 ± 2 MeV \cdot mb.

The total absorption cross section has been measured for $E_\gamma = 10$ to 210 MeV: it rises to ≈ 5 mb at ≈ 21 MeV, decreases to about 0 at 160 MeV and then increases to ≈ 1.5 mb at 210 MeV. An integrated cross section of 156 ± 15 MeV \cdot mb is reported for $E_\gamma = 10$ to 29 MeV as is resonant structure at $E_\gamma = 11.8, (13.5), 14.8, 917.3, (19.5), 21.0, (23.0)$ and (25.0) MeV. Fine structure is also reported at $E_\gamma = 20.47 \pm 0.04$ and 20.73 ± 0.04 MeV. See (1979AJ01) for references. See also (1978CH1E, 1979ZI1A, 1980AH1A, 1982DE1H), (1982BA1T; applications) and (1979BA60, 1979KI1D, 1980TA1D, 1981DE18, 1981DE33, 1981IS11; theor.).

14. (a) ${}^9\text{Be}(\gamma, p){}^8\text{Li}$	$Q_m = -16.887$
(b) ${}^9\text{Be}(\gamma, np){}^7\text{Li}$	$Q_m = -18.920$
(c) ${}^9\text{Be}(\gamma, d){}^7\text{Li}$	$Q_m = -16.6950$
(d) ${}^9\text{Be}(\gamma, t){}^6\text{Li}$	$Q_m = -17.6878$

The yield shows structure in the energy region corresponding to the ${}^9\text{Be}$ levels at 17–19 MeV followed by the giant resonance at $E_\gamma \approx 23$ MeV ($\sigma = 2.64 \pm 0.30$ mb). Structure attributed to eleven states of ${}^9\text{Be}$ with $18.2 < E_x < 32.2$ MeV has also been reported. Integrated cross sections have been obtained for each of these resonances, and over different energy intervals for protons leading to ${}^8\text{Li}^*(0 + 0.98, 2.26 + 3.21, 9.0, 17.0)$. Angular and energy distributions of photoprotons in various energy intervals have been studied by many groups: see (1974AJ01) for references. See also (1976MA34). For momentum spectra using tagged photons with $E_\gamma = 180 - 420$ MeV and 360–600 MeV, see (1980HO22) and (1982BA32), respectively. See also (1981AR04, 1981TA1T, 1983ST1G).

The integrated cross sections are reported to be 1.0 ± 0.5 MeV \cdot mb ($E_\gamma = 21 \rightarrow 33$ MeV) for reaction (c) to ${}^7\text{Li}^*(0 + 0.4)$ and 0.6 ± 0.3 MeV \cdot mb ($E_\gamma = 25 \rightarrow 33$ MeV) for reaction (d) to ${}^6\text{Li}(0)$. The total integrated cross section for $[(\gamma, p) + (\gamma, pn) + (\gamma, d) + (\gamma, t)]$ is given as 33 ± 3 MeV \cdot mb by (1966DE13), who also report resonances in the (γ, d) and (γ, t) cross sections corresponding to ${}^9\text{Be}^*(26.0 \pm 0.2)$ and ${}^9\text{Be}^*(32.2 \pm 0.3)$, respectively. See also (1974AJ01), (1980IS1E, 1982AV06, 1983HO01, 1983LI1J) and (1981IS11, 1982SU09; theor.).

15. ${}^9\text{Be}(\gamma, \gamma){}^9\text{Be}$

See (1980IS1E, 1982NO1C). See also (1979AJ01) and (1983ZH1D; theor.).

16. (a) ${}^9\text{Be}(e, e){}^9\text{Be}$
 (b) ${}^9\text{Be}(e, en){}^8\text{Be}$ $Q_m = -1.6655$
 (c) ${}^9\text{Be}(e, ep){}^8\text{Li}$ $Q_m = -16.887$
 (d) ${}^9\text{Be}(e, e\alpha){}^5\text{He}$ $Q_m = -2.46$
 (e) ${}^9\text{Be}(e, e\pi^+){}^9\text{Li}$ $Q_m = -153.174$

$$\langle r^2 \rangle^{1/2} = 2.519 \pm 0.012 \text{ fm}, Q = 6.5_{-0.6}^{+0.9} \text{ fm}^2,$$

$$b = 1.5 \pm_{-0.2}^{+0.3} \text{ fm} [b = \text{oscillator parameter}]$$

$\langle r^2 \rangle_M^{1/2} = 3.2 \pm 0.3 \text{ fm}, \Omega = 6 \pm 2 \mu_N \cdot \text{fm}^2$ [this value of the magnetic octupole moment implies a deformation of the average nuclear potential].

The elastic scattering of electrons has been studied for E_e up to 700 MeV. Magnetic elastic scattering gives indications of both M1 and M3 contributions. See also (1978DE32). Inelastic scattering populates a number of levels: Table 9.8 displays the parameters of the states. See (1974AJ01, 1979AJ01) for other work and for references.

At $E_e = 134.7$ and 237 MeV (1979BU11) have determined transverse and longitudinal form factors for electrodisintegration in the region of the quasielastic scattering peak. They report the observation of E1 and E2 giant resonance peaks: see Table 9.8. The quasielastic process has been studied in the range 250–580 MeV/c by (1978KU06). For reaction (c) see ${}^8\text{Li}$. For reaction (e) see (1980ST15). (1979JE04) have measured the ratio π^-/π^+ in the electroproduction of 10 MeV pions. See also (1979SC1E) and the “General” section here.

See also (1979FR1B, 1979FR1F, 1979FR1C, 1983LI1K), (1979MO1G, 1979TI1A, 1980MO1C, 1981MO1H, 1982DE1J) and (1978KE13, 1978TA1D, 1979OK02, 1980ER01, 1980FU1G, 1980SA1E, 1980ST15, 1981BO1T, 1981CH30, 1981DE1T, 1981IS11, 1981KE15, 1981LA1E, 1982BO1G, 1982KE1C, 1983BO1E, 1983OC01; theor.).

17. ${}^9\text{Be}(\pi^\pm, \pi^\pm){}^9\text{Be}$ (See also the “General” section here.)

The elastic scattering, and inelastic scattering to ${}^9\text{Be}^*(2.43, 6.76)$, have been studied at $E_{\pi^\pm} = 162$ MeV (1978GE11) and 291 MeV (1981GE03). Quadrupole contributions appear to be quite important for the elastic scattering at 162 MeV (1978GE11), but are much less so at higher energy (1981GE03). The inelastic scattering suggest the importance of radial localization effects (1981GE03). For a study of the (π^+, π^0) reaction see (1981BO06). For measurements of inclusive charged pion scattering yields see (1981MC09).

Table 9.8: Levels of ${}^9\text{Be}$ from ${}^9\text{Be}(e, e'){}^9\text{Be}^*$ ^a

E_x in ${}^9\text{Be}$ (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Transition	J^π	Γ_{γ_0} (eV)
1.78 \pm 30	150 \pm 50	E1	$\frac{1}{2}^+$	0.30 \pm 0.12
2.44 \pm 20	< 30	M1	$\frac{5}{2}^-$	0.089 \pm 0.010
		E2		(1.89 \pm 0.14) $\times 10^{-3}$ ^b
3.04 \pm 20	450 \pm 150	E1 ^c	$\frac{5}{2}^+$ ^c	0.30 \pm 0.25 ^d
4.7 \pm 200	700 \pm 300	E(1)		2.4 \pm 1.2 ^e
6.4 \pm 100	1000 \pm 300	E2	$\frac{7}{2}^-$	0.082 \pm 0.035
		f		
13.84 \pm 50 ^g				
14.388 \pm 15 ^h	< 70	M1	$\frac{3}{2}^-$	6.9 \pm 0.5 ^f
15.10 \pm 50 ^g				
15.97 \pm 30 ^g	\approx 300	M1		3.7 \pm 0.8 ^e
16.631 \pm 15 ^h	< 70	M2 ⁱ	$\leq \frac{7}{2}^+$	0.26 \pm 0.02 ^e
		M1	$\leq \frac{5}{2}^-$	2.0 \pm 0.5 ^e
16.961 \pm 15 ^h	< 70	M1	$\frac{1}{2}^-$	11.5 \pm 1.4 ^f
17.28		M1	$\leq \frac{5}{2}^-$	7.3 \pm 1.3 ^e
17.480 \pm 20 ^h	\approx 100	M2 ⁱ	$\leq \frac{7}{2}^+$	0.40 \pm 0.03 ^e
18.02 \pm 50 ^g				
18.62 \pm 50 ^g				
19.51 \pm 50 ^g				
20.76 \pm 50 ^g				
		j		

^a For references see Table 9.8 in (1979AJ01).

^b $B(\text{C}2, \omega) \uparrow = 45.7 \pm 3.5 e^2 \cdot \text{fm}^4$.

^c Assumed.

^d The group may consist of two unresolved states, the second one reached by an M1 transition [$J^\pi = (\frac{1}{2})^-$] with $\Gamma_{\gamma_0} = 0.18 \pm 0.09$ eV. I am indebted to Dr. L.W. Fagg for his help in understanding this point.

^e $g\Gamma_{\gamma_0}$; where $g = (2J_f + 1)/(2J_i + 1)$.

^f See (1974AJ01) and Table 9.6 here.

^g Weak transition.

^h (1983LO1G, 1983LOZZ) report transverse form factors for this state ($0.9 \leq q \leq 2.5 \text{ fm}^{-1}$).

ⁱ Or pure spin-flip E1.

^j States belonging to the E1 and E2 giant resonances are reported at $E_x = 19.2 \pm 0.2$ [4.0 \pm 0.6], 28.7 \pm 0.5 [8.9 \pm 1.7], 36.3 \pm 0.5 [2.8 \pm 1.3], 41 \pm 1 [7.2 \pm 2.0] and 58.0 \pm 0.4 MeV [3.1 \pm 0.5] MeV (1979BU11) [Γ in brackets]. See also (1974AJ01) for other reported states.

18. (a) ${}^9\text{Be}(n, n){}^9\text{Be}$

(b) ${}^9\text{Be}(n, 2n){}^8\text{Be}$ $Q_m = -1.6655$

The population of ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1, (6.8))$ has been reported in this reaction: see (1974AJ01). For the neutron decay of these states see Table 9.5. Angular distributions have been measured at $E_n = 5.9$ to 14.2 MeV [see (1974AJ01, 1979AJ01)] and at 3.5 to 7.1 MeV (1980CO1U; n_0, n_2), 6.97 to 14.93 MeV (1978HO23; n_0, n_2) and at 14.6 MeV (1982HAZK; n_0). See also (1981DA1K, 1983ANZY), (1979BY01, 1980AK01; theor.) and ${}^{10}\text{Be}$.

19. ${}^9\text{Be}(p, p){}^9\text{Be}$

Elastic and inelastic scattering angular distributions have been studied at many energies in the range $E_p = 5$ to 1000 MeV [see (1974AJ01, 1979AJ01)] and at $E_p = 2.31$ to 2.73 MeV (1980BO1L; p_0), 35.2 MeV (1980FA07; p_0), 72 MeV (1982AU1B; p_0), 225 MeV (1981RO1F; p_0) and 1 GeV (1979AL26; p_0). The elastic distributions show pronounced diffraction maxima characteristic of the optical model. A quadrupole-deformed optical model potential is necessary to obtain a good fit to the p_0 and p_2 angular distributions: see (1974AJ01).

The structure corresponding to ${}^9\text{Be}^*(1.7)$ is asymmetric: the line shape peaks 25_{-11}^{+15} keV above the threshold for ${}^8\text{Be} + n$. The population of ${}^9\text{Be}^*(1.7)$ has been studied for $E_p = 4.2$ to 5.5 MeV: it is consistent with a scattering length of about -20 fm between a neutron and ${}^8\text{Be}$. The weighted mean of the values of E_x for ${}^9\text{Be}^*(2.4)$ listed in (1974AJ01) is 2432 ± 3 keV. ${}^9\text{Be}^*(3.1)$ has $E_x = 3.03 \pm 0.03$ MeV, $\Gamma = 250 \pm 50$ keV, $J^\pi = \frac{3}{2}^+, \frac{5}{2}^+$. Higher states are observed at $E_x = 4.8 \pm 0.2, 6.76 \pm 0.06$ [$J^\pi = \frac{1}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$ (but see below), $\Gamma = 1.2 \pm 0.2$ MeV], 7.94 ± 0.08 ($\Gamma \approx 1$ MeV), 11.3 ± 0.2 MeV ($\Gamma \approx 1$ MeV), 14.4 ± 0.3 ($\Gamma \approx 1$ MeV), $16.7 \pm 0.3, 17.4 \pm 0.3, 19.0 \pm 0.4, 21.1 \pm 0.5$ and 22.4 ± 0.7 MeV [the five highest states are all broad]. For ${}^9\text{Be}^*(2.4, 6.8)$ $B(E2\uparrow) = 49 \pm 6$ and 24 ± 4 fm⁴ and $\Gamma(E2\downarrow) = 0.0025$ and 0.10 eV, respectively. The strong population of ${}^9\text{Be}^*(2.4, 6.8)$ is consistent with the assumption that they have $J^\pi = \frac{5}{2}^-$ and $\frac{7}{2}^-$, respectively, and are members of the ground-state $K = \frac{3}{2}^-$ band. See (1966LA04, 1974AJ01) for references.

The elastic scattering studied at $E_p = 25$ MeV (1974BI14) and 225 MeV (1981RO1F) shows that the depolarization parameter, D , deviates from unity and thus that a spin-spin term must be included in the optical parameter. The spin-flip probability at $E_{p\text{-bar}} = 31$ MeV is ≈ 0 for the p_2 group, which is expected in view of the collective nature of the transition (1981CO08). See also (1978AL1G) and (1979BY01, 1980KO1V, 1981BA50, 1981CO08, 1981GU1B, 1981GU1F, 1981KR17, 1982BA1X, 1982BA14, 1982GU1E; theor.).

20. (a) ${}^9\text{Be}(p, 2p){}^8\text{Li}$

$Q_m = -16.887$

(b) ${}^9\text{Be}(p, pd){}^7\text{Li}$

$Q_m = -16.6950$

(c) ${}^9\text{Be}(p, p\alpha){}^5\text{He}$	$Q_m = -2.46$
(d) ${}^9\text{Be}(p, pn){}^8\text{Be}$	$Q_m = -1.6655$
(e) ${}^9\text{Be}(p, d{}^3\text{He}){}^5\text{He}$	$Q_m = -20.82$

The summed proton spectrum (reaction (a)) shows two peaks with $Q = -16.4 \pm 0.3$ and -25.4 ± 0.5 MeV, corresponding to removal of a p-proton and an s-proton respectively, and a third peak of uncertain assignment (perhaps due to unresolved states) with $Q = -32.3 \pm 0.6$ MeV. See (1974AJ01). See also (1979KO21; also for (p, pX)), (1979KR1A, 1981FR24). For reaction (b) see (1974AJ01). Reactions (c) and (f) are dominated by quasifree processes at $E_p = 101.5$ MeV: $S_\alpha(L = 0) = 0.45$, $S_\alpha(L = 2) = 0.55$ (1980NA09). See also (1982NA1N). Reaction (d) at $E_p = 10 - 24$ MeV involves the $\frac{5}{2}^+$ state [3.01 ± 0.02 MeV]. ${}^9\text{Be}^*(4.7)$ is also involved (1978JE01). See also (1982PE1F). For pion production see (1979BU1D). See also (1979BA28) and (1978WO1A, 1979BO27, 1979DO03, 1979KI10, 1979MA1M, 1980BO12, 1981IS11; theor.).

21. ${}^9\text{Be}(d, d){}^9\text{Be}$

Angular distributions have been reported in the range 1.0 to 410 MeV [see (1974AJ01, 1979AJ01)] and at $E_{d\text{-bar}} = 2.0$ to 2.8 MeV (1982DE1P; d_0) and $E_d = 12.17$ to 14.43 MeV (1978TA12; d_0, d_1, d_2).

Inelastic groups have been reported to ${}^9\text{Be}^*(1.7)$, to states with $E_x = 2431.9 \pm 7.0$ keV and 3040 ± 15 keV ($\Gamma = 294 \pm 20$ keV), and to ${}^9\text{B}^*(4.7, 6.8)$: see (1974AJ01). See also (1978ZE03, 1979DO03; theor.) and ${}^{11}\text{B}$ in (1985AJ01).

22. (a) ${}^9\text{Be}(t, t){}^9\text{Be}$

(b) ${}^9\text{Be}(t, nt){}^8\text{Be}$ $Q_m = -1.6655$

Angular distributions of elastically scattered tritons have been measured at $E_t = 2.10$ MeV [see (1974AJ01)] and at $E_t = 15$ and 17 MeV (1978SC02). Reaction (b) at $E_t = 4.2$ and 4.6 MeV proceeds via ${}^9\text{Be}^*(2.4)$: see (1974AJ01). See also (1981ME1D; theor.).

23. (a) ${}^9\text{Be}({}^3\text{He}, {}^3\text{He}){}^9\text{Be}$

(b) ${}^9\text{Be}({}^3\text{He}, 2\alpha){}^4\text{He}$ $Q_m = 19.0041$

Angular distributions have been studied for $E({}^3\text{He}) = 1.6$ to 32.6 MeV and at 217 MeV [see (1974AJ01, 1979AJ01)] and at $E({}^3\text{He}) = 46.1$ MeV (1979GO07; elastic). For optical model parameters see references in (1979AJ01). At $E({}^3\text{He}) = 39.8$ MeV, ${}^9\text{Be}^*(1.7, 2.4, 3.1, 4.7, 6.8,$

14.4) are populated. In a study of reaction (b) the (${}^5\text{He} + \alpha$) system is found to be mainly in an S-state: the probability of finding such a cluster in ${}^9\text{Be}$ is $N_{\text{eff}} = 0.05$ (1975KA05). At $E({}^3\text{He}) = 2.1$ to 2.7 MeV quasifree processes are reported by (1978AR21): see ${}^8\text{Be}$. See also (1981DE1X) and (1979KA1G).

24. (a) ${}^9\text{Be}(\alpha, \alpha){}^9\text{Be}$

(b) ${}^9\text{Be}(\alpha, 2\alpha){}^5\text{He}$ $Q_m = -2.46$

(c) ${}^9\text{Be}(\alpha, \alpha n){}^8\text{Be}$ $Q_m = -1.6655$

Angular distributions have been studied at many energies in the range $E_\alpha = 5.0$ to 104 MeV [see (1974AJ01)] and at $E_\alpha = 35.5$ MeV (1982PE03; $\alpha_0, \alpha_1, \alpha_2, \alpha_4, \alpha_5$ and α to ${}^9\text{Be}^*(11.28)$) and 65 MeV (1980HA33; $\alpha_0, \alpha_2, \alpha_5$). See (1972DE01) for a discussion of optical model parameters. The structure (${}^5\text{He} + \alpha$) for ${}^9\text{Be}$ is found to be much more probable than (${}^6\text{Li} + t$): the ratio of spectroscopic factors is about 30 (1972DE01).

At $E_\alpha = 35.5$ MeV, states belonging to the $K = \frac{3}{2}^-$ ground-state band are strongly excited [${}^9\text{Be}^*(0, 2.43, 6.76, 11.28)$]; it is suggested that the latter has $J^\pi = (\frac{9}{2}^-)$; see, however, reaction 12]. The first three states belonging to the $K = \frac{1}{2}^+$ band are also excited [${}^9\text{Be}^*(1.68, 3.05, 4.70)$] (1982PE03; coupled channels analysis). For reaction (b) see (1974AJ01) and (1980ZH1A; 18 MeV) and (1980WA07; 140 MeV). $S_\alpha = 0.96$ (1976WO11). For reaction (c) see (1974AJ01). See also (1978ZE03, 1981BA20, 1982BU1D; theor.).

25. (a) ${}^9\text{Be}({}^6\text{Li}, {}^6\text{Li}){}^9\text{Be}$

(b) ${}^9\text{Be}({}^7\text{Li}, {}^7\text{Li}){}^9\text{Be}$

Elastic angular distributions have been measured at $E({}^6\text{Li}) = 4, 6$ and 24 MeV and at $E({}^7\text{Li}) = 24$ and 34 MeV: see (1979AJ01).

26. ${}^9\text{Be}({}^9\text{Be}, {}^9\text{Be}){}^9\text{Be}$

Elastic angular distributions have been obtained at $E({}^9\text{Be}) = 5, 9, 12$ and 16 MeV [see (1979AJ01)], at 14, 20 and 26 MeV (1979UN01) and at 40 to 55 MeV (1983OMZZ; also to ${}^9\text{Be}^*(2.43)$; σ (fusion)).

27. ${}^9\text{Be}({}^{10}\text{B}, {}^{10}\text{B}){}^9\text{Be}$

Elastic angular distributions are reported at $E(^{10}\text{B}) = 20.1$ and 30 MeV ([1980BO14](#)).

28. (a) ${}^9\text{Be}(^{12}\text{C}, ^{12}\text{C})^9\text{Be}$
(b) ${}^9\text{Be}(^{13}\text{C}, ^{13}\text{C})^9\text{Be}$

Elastic angular distributions have been measured for reaction (a) at $E(^{12}\text{C}) = 12, 15, 18$ and 21 MeV and $E(^9\text{Be}) = 14, 20, 26$ and 43 MeV [see ([1979AJ01](#))] and at $E(^9\text{Be}) = 14, 20$ and 26 MeV ([1979UN01](#)), 20 MeV ([1979BO1K](#)), 20 and 26.1 MeV ([1979JA04](#)), 50 MeV ([1977ST20](#)), 69.7 and 76.6 MeV ([1979MA21](#)) and 158.4 MeV ([1983FUZY](#)). For yield and fusion cross-section measurements see ([1978MA44](#), [1979MA21](#), [1981JA09](#), [1982DEZL](#), [1982HU06](#)). Elastic angular distributions for reaction (b) are reported at $E(^9\text{Be}) = 14, 20$ and 26 MeV ([1979UN01](#)) and 19.5 and 25.0 MeV ([1979JA04](#)). See also ([1978TA1B](#), [1982JA1E](#)) and ([1978GR22](#), [1980BR05](#), [1980OH1B](#), [1981GR17](#), [1981HU07](#), [1982LO13](#), [1983DE1U](#); theor.).

29. ${}^9\text{Be}(^{14}\text{N}, ^{14}\text{N})^9\text{Be}$

Elastic angular distributions have been measured at $E(^{14}\text{N}) = 25$ and 27.3 MeV: see ([1974AJ01](#)). For fusion studies see ([1981MA1V](#), [1982MAZA](#)).

30. (a) ${}^9\text{Be}(^{16}\text{O}, ^{16}\text{O})^9\text{Be}$
(b) ${}^9\text{Be}(^{18}\text{O}, ^{18}\text{O})^9\text{Be}$

Elastic angular distributions have been reported in the range $E(^{16}\text{O}) = 15$ to 30 MeV [see ([1979AJ01](#))] and at $E(^9\text{Be}) = 14, 20$ and 26 MeV ([1979UN01](#)) and 158.4 MeV ([1983FUZY](#)), as well as at $E(^{18}\text{O}) = 12.1, 16$ and 20 MeV [see ([1974AJ01](#))]. See also ([1981ST1P](#), [1982BE54](#)), ([1979HU1B](#), [1979MA30](#), [1979PA1B](#), [1981GR17](#), [1981GU1B](#); theor.).

31. (a) ${}^9\text{Be}(^{24}\text{Mg}, ^{24}\text{Mg})^9\text{Be}$
(b) ${}^9\text{Be}(^{26}\text{Mg}, ^{26}\text{Mg})^9\text{Be}$
(c) ${}^9\text{Be}(^{27}\text{Al}, ^{27}\text{Al})^9\text{Be}$
(d) ${}^9\text{Be}(^{28}\text{Si}, ^{28}\text{Si})^9\text{Be}$
(e) ${}^9\text{Be}(^{39}\text{K}, ^{39}\text{K})^9\text{Be}$
(f) ${}^9\text{Be}(^{40}\text{Ca}, ^{40}\text{Ca})^9\text{Be}$
(g) ${}^9\text{Be}(^{44}\text{Ca}, ^{44}\text{Ca})^9\text{Be}$

Elastic angular distributions have been measured at $E(^9\text{Be}) = 14, 20$ and 26 MeV for reactions (a), (c), (d) and at 14 and 17 MeV for reaction (d): see (1979AJ01). Recent measurements are at $E(^9\text{Be}) = 12$ to 30 MeV (1980BO11, 1981HU08; reaction (d)), $14, 20$ and 26 MeV (1978UN02, 1979UN01; reactions (a) (nat. Mg), (c), (d), (f)), 20 MeV (1978JA07; reaction (d)), 27 and 40 MeV (1981GL03; reaction (d)), 40 MeV (1981HN04; reactions (e), (f), (g)), 45 and 60 MeV (1980EC01; reactions (d), (f)), 50 MeV (1977ST20; reactions (d), (f)), 158.4 MeV (1983FUZY; reactions (b), (c)) and 121 and 201.6 MeV (1980ZI02; reaction (d)). For fusion studies see (1980EC03, 1983EC01). See also (1980EC04) and (1979SA1E, 1980HU09; theor.).

$$32. \text{}^{10}\text{Be}(d, t)^9\text{Be} \quad Q_m = -0.5548$$

Forward angular distributions have been obtained at $E_d = 15.0$ MeV for the tritons to $^9\text{Be}^*(0, 1.7, 2.4, 3.1)$. The ground-state transition is well fitted by $l = 1$. The transition to $^9\text{Be}^*(1.7)$ [$\approx 165 \pm 25$ keV] is consistent with $J^\pi = \frac{1}{2}^+$, that to $^9\text{Be}^*(2.4)$ is quite well fitted with $l = 3$ [$J^\pi = \frac{5}{2}^-$], and that to $^9\text{Be}^*(3.1)$ [$\Gamma = 280 \pm 25$ keV] is consistent with $l = 2$. No other narrow states are seen up to $E_x = 5.5$ MeV (1970AU02).

$$33. \text{}^{10}\text{B}(n, d)^9\text{Be} \quad Q_m = -4.3620$$

Angular distributions of the d_0 and d_1 groups have been measured at $E_n = 14.4$ MeV: see (1974AJ01). See also ^{11}B in (1980AJ01).

$$34. \text{}^{10}\text{B}(p, 2p)^9\text{Be} \quad Q_m = -6.5865$$

The summed proton spectrum at $E_p = 460$ MeV yields $Q = -6.7 \pm 0.5, -11.9 \pm 0.5, -17.1 \pm 0.6$ (all $l \neq 0$) and $Q = -30.5 \pm 0.6$ MeV ($l = 0$): see (1974AJ01).

$$35. \text{}^{10}\text{B}(d, ^3\text{He})^9\text{Be} \quad Q_m = -1.0930$$

Angular distributions of the ^3He groups corresponding to $^9\text{Be}^*(0, 2.4)$ ahev been studied at $E_d = 11.8, 28$ and 52 MeV [the latter also to $^9\text{Be}^*(6.7)$], and at $E_d = 15$ MeV: $S = 0.72$ and 0.82 for $^9\text{Be}^*(0, 2.4)$. At $E_{d\text{-bar}} = 52$ MeV $^9\text{Be}^*(11.3)$ appears to be strongly populated: see (1979AJ01).

$$36. \text{}^{10}\text{B}(t, \alpha)^9\text{Be} \quad Q_m = 13.2275$$

At $E_t = 12.9$ MeV α -groups are observed to the ground state of ${}^9\text{Be}$ and to excited states at $E_x = 1.75 \pm 0.03$, 2.43 , 3.02 ± 0.04 ($\Gamma = 320 \pm 60$ keV), 11.27 ± 0.04 ($\Gamma = 530 \pm 70$ keV), (14.4) [$\Gamma \approx 800$ keV], 14.39 and 16.67 MeV. The $T = \frac{3}{2}$ state ${}^9\text{Be}^*(14.39)$ is very weakly populated [$\approx 5\%$ of intensity of α_2]. The angular distribution of the α_2 group shows sharp forward and backward peaking. The α_0 group is not peaked in the backward direction: see (1979AJ01). Angular distributions are also reported at $E_t = 0.9$ and 1.1 MeV to ${}^9\text{Be}^*(2.4)$ (1980CI1B). A study at $E_t = 1.0$ to 3.2 MeV finds $E_x = 1.750 \pm 0.025$ MeV, $\Gamma = 220 \pm 8$ keV: see (1974AJ01). See also (1978KR16; theor.) and ${}^{13}\text{C}$ in (1981AJ01).

$$37. {}^{10}\text{B}(\alpha, {}^5\text{Li}){}^9\text{Be} \quad Q_m = -8.551$$

See ${}^5\text{Li}$.

$$38. {}^{11}\text{B}(\text{n}, \text{t}){}^9\text{Be} \quad Q_m = -9.5589$$

The angular distribution of t_0 has been measured at $E_n = 14.4$ MeV: see (1974AJ01).

$$39. {}^{11}\text{B}(\text{p}, {}^3\text{He}){}^9\text{Be} \quad Q_m = -10.3227$$

At $E_p = 45$ MeV angular distributions are reported for the ${}^3\text{He}$ ions corresponding to ${}^9\text{Be}^*(0, 2.4, 11.8, 13.8, 14.39)$ [$T = \frac{3}{2}$], 15.96 ± 0.04 [$T = \frac{1}{2}$]). In addition one or more states may be located at ${}^9\text{Be}^*(15.13)$. It is suggested that ${}^9\text{Be}^*(11.8, 13.8, 15.96)$ are the $J = \frac{3}{2}^-$, $T = \frac{1}{2}$ analogs to ${}^9\text{B}^*(12.06, 14.01, 16.02)$. Angular distributions are also reported at $E_p = 40$ MeV. The intensity of the group to ${}^9\text{Be}^*(3.1)$ is $\approx 1\%$ of the ground-state group at that energy: see (1974AJ01). The excitation energy of the first $T = \frac{3}{2}$ state is $E_x = 14392.9 \pm 1.8$ keV (1974KA15), using Q_m . See also (1981CO2D) and ${}^{12}\text{C}$ in (1985AJ01).

$$40. \text{(a) } {}^{11}\text{B}(\text{d}, \alpha){}^9\text{Be} \quad Q_m = 8.031$$

$$\text{(b) } {}^{11}\text{B}(\text{d}, \text{n}\alpha){}^4\text{He}{}^4\text{He} \quad Q_m = 6.457$$

Alpha groups reported corresponding to ${}^9\text{Be}^*(0, 1.7, 2.4, 3.1)$. The width of ${}^9\text{Be}^*(1.7)$ [$E_x = 1.70 \pm 0.01$ MeV] is $\Gamma_{\text{c.m.}} = 220 \pm 20$ keV. The weighted mean of the values of E_x of ${}^9\text{Be}^*(2.4)$, reported in (1974AJ01), is 2425 ± 3 keV. The $\frac{5}{2}^+$ state is at $E_x = 3.035 \pm 0.025$ MeV: $\Gamma_{\text{c.m.}} = 257 \pm 25$ keV. The ratio Γ_γ/Γ of ${}^9\text{Be}^*(1.7) \leq 2.4 \times 10^{-5}$, that for ${}^9\text{Be}^*(2.4)$ is reported to be $(1.16 \pm 0.14) \times 10^{-4}$. Since Γ_γ is known from (e, e') [see Table 9.8: 0.089 ± 0.010 eV], $\Gamma = 0.77 \pm 0.15$ keV. See (1974AJ01, 1979AJ01) for references.

Angular distributions for α_0 and α_2 are reported at $E_d = 0.39$ to 3.9 MeV and at 12 MeV [see (1974AJ01, 1979AJ01)]. Reaction (b), at $E_d = 10.4$ and 12.0 MeV, proceeds via ${}^9\text{Be}^*(2.4)$ and to some extent via ${}^9\text{Be}^*(3.1, 4.7)$ and possibly some higher excited states. The dominant decay of ${}^9\text{Be}^*(2.4)$ is to ${}^5\text{He}(0) + \alpha$ while ${}^9\text{Be}^*(3.1, 4.7)$ decay to ${}^8\text{Be}(0) + n$. It should be noted, however, that the peaks corresponding to ${}^9\text{Be}^*(3.1)$ have a FWHM of ≈ 1 MeV, which may imply that ${}^9\text{Be}^*(2.8)$ is involved.

$$41. \text{ (a) } {}^{12}\text{C}(n, \alpha){}^9\text{Be} \quad Q_m = -5.7010$$

$$\text{ (b) } {}^{12}\text{C}(n, n\alpha){}^4\text{He}{}^4\text{He} \quad Q_m = -7.2747$$

Angular distributions of the α_0 group have been measured at $E_n = 13.9$ to 18.8 MeV [see (1974AJ01)] and at 14 MeV (1981HA1K). ${}^9\text{Be}^*(1.7, 2.4, 3.1)$ are also populated. At $E_n = 14.7$ MeV (1981GU12) report the involvement of ${}^9\text{Be}^*(6.76)$ which subsequently decays by n-emission. Reaction (b) at $E_n = 13$ to 18 MeV populates ${}^9\text{Be}^*(2.4)$. See also (1979SU1C) and ${}^{12}\text{C}$, ${}^{13}\text{C}$ in (1980AJ01, 1981AJ01).

$$42. {}^{13}\text{C}({}^3\text{He}, {}^7\text{Be}){}^9\text{Be} \quad Q_m = -9.060$$

Angular distributions have been obtained at $E({}^3\text{He}) = 70$ MeV for the transitions to ${}^9\text{Be}^*(0, 2.4)$ and ${}^7\text{Be}^*(0, 0.43)$. Broad states at 2.9 , 4.8 ± 0.2 , 7.3 ± 0.2 and 11.9 ± 0.4 MeV are also populated (1976ST11).

$$43. {}^{13}\text{C}(\alpha, {}^8\text{Be}){}^9\text{Be} \quad Q_m = -10.7392$$

At $E_\alpha = 65$ MeV angular distributions have been measured for the transitions to ${}^9\text{Be}^*(0, 2.4)$ (1976WO11). See also (1979BE1J).

$$44. {}^{19}\text{F}(d, {}^{12}\text{C}){}^9\text{Be} \quad Q_m = 0.3009$$

See (1981GO16) in ${}^{12}\text{C}$ (1985AJ01).

$$45. {}^{23}\text{Na}(d, {}^{16}\text{O}){}^9\text{Be} \quad Q_m = -3.0059$$

See (1981RU09; $E_d = 13.6$ MeV).

⁹B
(Figs. 17 and 18)

GENERAL: (See also (1979AJ01).)

Model calculations: (1978AR1H, 1979LA06, 1979MA1J, 1981KO1Q).

Special states: (1981KO1Q).

Reactions involving pions: (1978WA1B, 1979AL1J, 1982EL07, 1982HI02, 1983HU02).

Hypernuclei: (1978PO1A, 1978SO1A, 1979MA1L, 1981WA1J, 1982KO11).

Other topics: (1979BE1H, 1979LA06, 1982NG01).

Ground state of ⁹B: (1982NG01).

1. (a) ${}^6\text{Li}({}^3\text{He}, \gamma){}^9\text{B}$	$Q_m = 16.601$	
(b) ${}^6\text{Li}({}^3\text{He}, n){}^8\text{B}$	$Q_m = -1.975$	$E_b = 16.601$
(c) ${}^6\text{Li}({}^3\text{He}, p){}^8\text{Be}$	$Q_m = 16.7861$	
(d) ${}^6\text{Li}({}^3\text{He}, d){}^7\text{Be}$	$Q_m = 0.112$	
(e) ${}^6\text{Li}({}^3\text{He}, t){}^6\text{Be}$	$Q_m = -4.307$	
(f) ${}^6\text{Li}({}^3\text{He}, {}^3\text{He}){}^6\text{Li}$		
(g) ${}^6\text{Li}({}^3\text{He}, \alpha){}^5\text{Li}$	$Q_m = 14.91$	
(h) ${}^6\text{Li}({}^3\text{He}, p){}^4\text{He}{}^4\text{He}$	$Q_m = 16.8779$	

The 90° yields of γ_0 and of γ to ${}^9\text{Be}^*(2.36)$ (reaction (a)) has been measured for $E({}^3\text{He}) = 0.6$ to 1.2 MeV [as have the 2 α -particles from the decay of ${}^8\text{Be}^*(16.6)$ (reaction (c))]: they are reported to show a resonance at $E({}^3\text{He}) = 765 \pm 5$ keV [${}^9\text{B}^*(17.111)$], attributed to ${}^9\text{B}^*(17.076)$ [$T = \frac{3}{2}$] (1978AL37). The total cross section for reaction (b) increases monotonically from threshold to ≈ 7 mb at 3.8 MeV. It then decreases monotonically from ≈ 20 mb at $E({}^3\text{He}) = 5.5$ to 7.6 MeV and also from 8.9 to 26.5 MeV: see (1979AJ01), and ${}^8\text{B}$.

Absolute cross sections for protons (reaction (c)) to ${}^8\text{Be}^*(0, 2.9, 16.6, 16.9)$ as well as for the continuum protons (reaction (h)) have been measured for $E({}^3\text{He}) = 0.5$ to 1.85 MeV. Reaction rate parameters, $\langle\sigma\nu\rangle$, have been calculated for $kT = 0.01$ to 10.0 MeV (1980EL02). Excitation functions for p_0 and p_1 have been measured for $E({}^3\text{He}) = 0.9$ to 17 MeV, and polarization measurements are reported at $E({}^3\text{He}) = 14$ MeV. Resonances are reported at $E({}^3\text{He}) = 1.6$ and 3.0 MeV [$\Gamma = 0.25$ and 1.5 MeV]: see (1974AJ01, 1979AJ01), and ${}^8\text{Be}$. See also (1979SL01). Polarization measurements are also reported at $E({}^6\text{Li}) = 21$ MeV (1983KO04; VAP; p_0). A resonance in the excitation function for deuterons (reaction (d)) is observed corresponding to ${}^9\text{B}^*(17.6)$ (1979BU1E) [$E({}^3\text{He}) = 0.7$ to 2.0 MeV]. Polarization measurements at $E({}^3\text{He}) = 33.3$ MeV for

Table 9.9: Energy levels of ${}^9\text{B}$

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	0.54 ± 0.21	p, α	1, 2, 3, 4, 6, 7, 8, 10, 11, 12, 13, 14, 16, 17, 18, 19
(1.6)		≈ 700	(p, α)	4, 13
2.361 ± 5	$\frac{5}{2}^-; \frac{1}{2}$	81 ± 5	α	1, 2, 4, 6, 7, 8, 11, 12, 13, 14, 16, 18
2.788 ± 30	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	550 ± 40	p	4, 6, 11, 13, 16, 18
(4.8 \pm 100)		1000 ± 200		4, 9
6.97 ± 60	$\frac{7}{2}^-; \frac{1}{2}$	2000 ± 200	p	4, 6, 9, 11, 14, 16
11.70 ± 70	$(\frac{7}{2})^-; \frac{1}{2}$	800 ± 50	p	9, 11, 13
12.06 ± 60	$;$ $\frac{1}{2}$	800 ± 200	p	4, 9, 14
14.01 ± 70	$;$ $\frac{1}{2}$	390 ± 110		4, 14
14.6550 ± 2.5	$\frac{3}{2}^-; \frac{3}{2}$	0.395 ± 0.042	γ , p	4, 7, 14
14.7 ± 180	$(\frac{5}{2})^-; \frac{1}{2}$	1350 ± 200		11
15.29 ± 40	$;$ $\frac{1}{2}$			14
15.58 ± 40	$;$ $\frac{1}{2}$			14
16.024 ± 25	$;$ $(\frac{1}{2})$	180 ± 16		4, 14
17.076 ± 4	$T = \frac{3}{2}$	22 ± 5	$(\gamma, {}^3\text{He})$	1, 14
17.190 ± 25		120 ± 40	p, d, ${}^3\text{He}$	4, 5, 14
17.637 ± 10		71 ± 8	p, d, ${}^3\text{He}$, α	1, 4, 5, 14
(18.6)		1000	p, ${}^3\text{He}$	1, 11

the d_0 and d_1 groups are reported by (1981BA38). See also (1979AJ01), and ${}^7\text{Be}$. Excitation functions for t_0 (reaction (e)) have been measured for $E({}^3\text{He}) = 10$ to 16 and 23.3 to 25.4 MeV: see (1974AJ01). Polarization measurements are reported at $E({}^3\vec{\text{He}}) = 33.3$ MeV for the t_0 group as well as for the ${}^3\text{He}$ ions to ${}^6\text{Li}^*(0, 2.19)$ (1981BA37) (reaction (f)). The elastic scattering has also been studied for $E({}^3\text{He}) = 0.7$ to 2.0 MeV (1979BU1E). See also ${}^6\text{Li}$ and ${}^6\text{Be}$.

The $\alpha - \alpha$ coincidences from the ${}^5\text{Li}_{\text{g.s.}}$ decay (reaction (g)) have been measured for $E({}^3\text{He}) = 1.4$ to 1.8 MeV: a resonance is observed at 1.57 ± 0.02 MeV [${}^9\text{B}^*(17.63)$], $\Gamma = 70 \pm 20$ keV (1978GU15). Polarization measurements of the α -particles to ${}^5\text{Li}^*(0, 16.7)$ are reported at $E({}^3\vec{\text{He}}) = 33.3$ MeV (1981BA38). For reaction (h) see also (1979AJ01). See also (1982LA20), (1979KA1G, 1981HO1E) and (1980LE06, 1982HI02, 1982LE10; theor.).

2. ${}^6\text{Li}(\alpha, n){}^9\text{B}$ $Q_m = -3.977$

At $E_\alpha = 14.4$ MeV, neutron groups are observed to ${}^9\text{B}^*(0, 2.4)$: the upper limit of the cross section to a state at ≈ 1.7 MeV is $100 \mu\text{b}/\text{sr}$ or < 0.1 of the n_0 group: see (1974AJ01). See also ${}^{10}\text{B}$.

3. ${}^6\text{Li}({}^6\text{Li}, t){}^9\text{B}$ $Q_m = 0.805$

Angular distributions of the t_0 group have been measured for $E({}^6\text{Li}) = 4.0$ to 5.5 MeV and at 7.35 and 9.0 MeV. No evidence was observed for a group, corresponding to ${}^9\text{B}^*(1.6)$: see (1974AJ01).

4. (a) ${}^7\text{Li}({}^3\text{He}, n){}^9\text{B}$ $Q_m = 9.351$
 (b) ${}^7\text{Li}({}^3\text{He}, np){}^8\text{Be}$ $Q_m = 9.5360$

For $E({}^3\text{He})$ to 12.5 MeV this reaction populates ${}^9\text{B}^*(0, (1.6), 2.4, 2.8, (7.0))$, and states at $E_x = 4.8 \pm 0.1$ MeV [1.0 ± 0.2 MeV], 12.06 ± 0.06 [0.8 ± 0.2], 14.01 ± 0.07 [0.39 ± 0.11], 14.657 ± 0.005 (based on Q_m) [< 0.045], 16.024 ± 0.025 [0.180 ± 0.016], 17.19 and 17.63 MeV [Γ in brackets]: see (1974AJ01). ${}^9\text{B}^*(14.66)$ is the first $T = \frac{3}{2}$ state in ${}^9\text{B}$. Its decay properties are displayed in Table 9.6 and compared with those of ${}^9\text{Be}^*(14.40)$: see reaction 9 in ${}^9\text{Be}$ and (1974AJ01). Angular distributions have been measured at $E({}^3\text{He}) = 1.56$ to 5.27 MeV: see (1974AJ01).

5. ${}^7\text{Be}(d, p){}^8\text{Be}$ $Q_m = 16.6738$ $E_b = 16.489$

For $E_d = 0.75$ to 1.70 MeV, resonances in the yields of protons are observed at $E_d = 0.900 \pm 0.025$ MeV (p_0, p_1) and 1.475 ± 0.010 MeV (p_1 only) with $\Gamma_{c.m.} = 120 \pm 40$ and 71 ± 8 keV, respectively [${}^9\text{B}^* = 17.19$ and 17.64 MeV] (1960KA17).

6. (a) ${}^9\text{Be}(p, n){}^9\text{B}$ $Q_m = -1.851$
 (b) ${}^9\text{Be}(p, pn){}^8\text{Be}$ $Q_m = -1.6655$

At $E_p = 20$ MeV, ${}^9\text{B}^*(0, 2.4)$ are populated. The results are consistent with the excitation of ${}^9\text{B}^*(2.8)$ [$\Gamma \approx 0.3$ MeV] and ${}^9\text{B}^*(7.0)$ [$\Gamma > 1$ MeV]. No other states are excited for $E_x < 7.1$ MeV (1970AN07). Additional states have been reported: see (1979AJ01) and the comments in (1978JE01). See also (1983BY01). The width of the ground state is 540 ± 210 eV (1964TE01).

Angular distributions have been reported at many energies in the range $E_p = 3.5$ to 49.3 MeV [see (1979AJ01)] and at 8.15 to 16.98 MeV (1983BY01; n₀). At $E_p = 10$ to 24 MeV reaction (b) does not appear to involve states of ${}^9\text{B}$ (1978JE01) [see ${}^9\text{Be}$]: see, however, (1979AJ01). For polarization and yield measurements see ${}^{10}\text{B}$. See also (1978BA1F, 1979WA1E, 1979WA1F; applications) and (1979BY01, 1980BA2H, 1983GU1G; theor.).

$$7. {}^9\text{Be}({}^3\text{He}, t){}^9\text{B} \quad Q_m = -1.087$$

Angular distributions have been measured for $E({}^3\text{He}) = 3.0$ to 25 MeV and at 217 MeV: see (1974AJ01, 1979AJ01). At $E({}^3\text{He}) = 39.8$ MeV, ${}^9\text{B}_{\text{g.s.}}$ is strongly populated and ${}^9\text{B}^*(2.4, 14.7)$ are also observed (1969BA06). See also ${}^{12}\text{C}$ in (1980AJ01).

$$8. {}^9\text{Be}({}^6\text{Li}, {}^6\text{He}){}^9\text{B} \quad Q_m = -4.575$$

At $E({}^6\text{Li}) \approx 31$ MeV ${}^9\text{B}_{\text{g.s.}}$ is strongly excited. ${}^9\text{B}^*(2.4)$ is also populated: see (1974AJ01).

$$9. {}^9\text{C}(\beta^+){}^9\text{B} \rightarrow {}^8\text{Be} + \text{p} \quad Q_m = 16.682$$

$$\quad \quad \quad \rightarrow {}^5\text{Li} + \alpha \quad Q_m = 14.81$$

Several groups of delayed protons are observed indicating the involvement of a number of ${}^9\text{B}$ states: see Table 9.10 (1972ES05). It is not possible to determine ft values since some of the ${}^9\text{B}$ states involved decay via ${}^5\text{Li} + \alpha$: (1972ES05).

$$10. {}^{10}\text{B}(\gamma, n){}^9\text{B} \quad Q_m = -8.437$$

See ${}^{10}\text{B}$.

$$11. {}^{10}\text{B}(\text{p}, \text{d}){}^9\text{B} \quad Q_m = -6.212$$

See Table 9.11. See also (1979AJ01).

$$12. {}^{10}\text{B}(\text{d}, \text{t}){}^9\text{B} \quad Q_m = -2.180$$

Table 9.10: Delayed protons following the β^+ decay of ${}^9\text{C}$ ^a

E_p (c.m.) (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	Corresponding state in ${}^9\text{B}$ (MeV)	
		if decay is to ${}^8\text{Be}_{\text{g.s.}}$	if decay is to ${}^8\text{Be}^*(2.9)$
3.45 ± 0.25	200 ± 100	3.26 ± 0.25 ^d	c
(4.2 ± 0.3)	1000 ± 200	4.0 ± 0.3	6.9 ± 0.3
(5.0 ± 0.2)	400 ± 200	4.8 ± 0.2	c
6.10 ± 0.10	400 ± 100	5.91 ± 0.10	c
9.28 ± 0.24 ^b	1800 ± 200	9.09 ± 0.24	11.99 ± 0.24
12.30 ± 0.10 ^b	450 ± 100	12.11 ± 0.10	c

^a (1972ES05).

^b Ratio of the intensities $I_{9.28}/I_{12.30} = 1.2 \pm 0.2$.

^c The relatively narrow width of the proton group does not permit this option.

^d By analogy with the ${}^9\text{Li}$ decay, this decay may involve a $J^\pi = \frac{1}{2}^-$ analog of ${}^9\text{Be}^*(2.78)$.

Such a state in ${}^9\text{B}$ has not been reported in any other reaction.

Table 9.11: Levels of ${}^9\text{B}$ from ${}^{10}\text{B}(p, d){}^9\text{B}$ ^a

E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (MeV)	l_n	J^π ^b
0		1	$\frac{3}{2}^-$
2.35 ± 20		1	$\frac{5}{2}^-$
(2.8) ^c			
7.1 ± 140	2.15 ± 0.15	1	$\frac{7}{2}^-$
11.70 ± 70	0.80 ± 0.05	1	$(\frac{7}{2})^-$
14.7 ± 180	1.35 ± 0.2	1	$(\frac{5}{2})^-$
(18.4)			

^a For references see Table 9.11 in (1974AJ01).

^b J from best fit to theoretical spectroscopic factor (1968KU04).

^c Weak group.

Angular distributions have been measured at $E_d = 11.8$ to 28 MeV. S for ${}^9\text{B}^*(0, 2.4)$ are 0.66 and 0.58 , respectively. See ([1974AJ01](#), [1979AJ01](#)).

13. (a) ${}^{10}\text{B}({}^3\text{He}, \alpha){}^9\text{B}$ $Q_m = 12.141$
 (b) ${}^{10}\text{B}({}^3\text{He}, \alpha\text{p}){}^8\text{Be}$ $Q_m = 12.3258$

Alpha-particle spectra show the excitation of ${}^9\text{B}^*(0, 2.4, 2.8, 11.8)$: see ([1966LA04](#)). Measurements by ([1968KR02](#)) determine $E_x = 2.361 \pm 0.005$ and 2.788 ± 0.030 MeV, $\Gamma = 81 \pm 5$ and 548 ± 40 keV, respectively. There is some evidence for a state with $E_x \approx 1.6$ MeV, $\Gamma \approx 0.7$ MeV, but it is not conclusive, in agreement with the older work. No evidence is found for any narrow levels in ${}^9\text{B}$ with $\Gamma \leq 100$ keV and $4 < E_x < 7$ MeV: the upper limit to the intensity of the corresponding α -group is 1% of the intensity of the group to ${}^9\text{B}^*(2.4)$ ([1968KR02](#)). Angular distributions have been determined at $E({}^3\text{He}) = 5.5$ and 33.7 MeV.

In reaction (b) study of the decays of ${}^9\text{B}^*(2.4, 2.8)$ shows that ${}^9\text{B}^*(2.4)$ decays $< 0.5\%$ by proton emission to ${}^8\text{Be}(0)$ [$\theta_f^2 < 5.1 \times 10^{-3}$] [it decays to ${}^5\text{Li}(0)$ by α -emission] while the second state, $E_x = 2.71 \pm 0.03$ MeV [$\Gamma = 0.71 \pm 0.06$ MeV] decays almost 100% by that channel [$\theta_d^2 = 0.74$] ([1966WI08](#)). No other excited states of ${}^9\text{B}$ with $3.5 < E_x < 9.5$ MeV decay by proton emission to ${}^8\text{Be}(0)$ ([1968KR02](#)). See also ([1974AJ01](#)).

14. ${}^{11}\text{B}(\text{p}, \text{t}){}^9\text{B}$ $Q_m = -11.409$

At $E_p = 45$ MeV angular distributions have been obtained for the triton groups to ${}^9\text{B}^*(0, 2.36, 12.06, 14.01, 14.66, 16.02)$. In addition the spectra show some indication of the groups corresponding to ${}^9\text{B}^*(7.0, 17.19, 17.64)$. $T = \frac{1}{2}$ states are reported at $E_x = 15.29 \pm 0.04$ and 15.58 ± 0.04 MeV ([1971HA10](#)). The first two $T = \frac{3}{2}$ states have been observed at $E_x = 14.6550 \pm 0.0025$ ([1974KA15](#)) and 17.076 ± 0.004 MeV [$\Gamma = 22 \pm 5$ keV] ([1974BE66](#)).

15. ${}^{12}\text{C}(\pi^+, \pi^+\text{t}){}^9\text{B}$ $Q_m = -27.366$

See ([1979EL12](#)).

16. (a) ${}^{12}\text{C}(\text{p}, \alpha){}^9\text{B}$ $Q_m = -7.552$
 (b) ${}^{12}\text{C}(\text{p}, \text{p}){}^4\text{He}{}^4\text{He}$ $Q_m = -7.2747$

Angular distributions have been measured for the α_0 group at $E_p = 14.0$ to 54.1 MeV. Alpha groups are also observed to ${}^9\text{B}^*(2.3, 2.9 \pm 0.2, 6.97 \pm 0.06)$. The angular distribution to ${}^9\text{B}^*(6.97)$ at $E_p = 54.1$ MeV is consistent with $J^\pi = \frac{7}{2}^-$, $\Gamma \approx 2$ MeV: see (1974AJ01). Angular distributions involving the α_0 and α^* groups [to ${}^4\text{He}^*(20.1), 0^+$] to ${}^9\text{B}_{\text{g.s.}}$ have been studied at $E_p = 45.2$ MeV (1980DA07). At $E_p = 72.4$ MeV, a study of the inclusive (p, dX) reaction shows a structure which, if attributed to ${}^9\text{B}$, would correspond to $E_x \approx 20.7$ MeV (1981KO28). See also ${}^{13}\text{N}$ (1981AJ01) and (1978GO14, 1981KA04; theor.).

17. ${}^{12}\text{C}({}^3\text{He}, {}^6\text{Li}){}^9\text{B}$ $Q_m = -11.570$

Angular distributions of ${}^6\text{Li}$ ions have been obtained at $E({}^3\text{He}) = 30.0$ and 40.7 MeV: see (1974AJ01).

18. ${}^{12}\text{C}(\alpha, {}^7\text{Li}){}^9\text{B}$ $Q_m = -24.898$

At $E_\alpha = 65$ MeV, ${}^9\text{B}_{\text{g.s.}}$ and possibly ${}^9\text{B}^*(2.36, 2.79)$ are populated (1978SA26).

19. ${}^{12}\text{C}({}^{16}\text{O}, {}^{19}\text{F}){}^9\text{B}$ $Q_m = -15.665$

See (1982LE1N).

${}^9\text{C}$
(Figs. 17 and 18)

GENERAL: (See also (1979AJ01) for other references in this category and for some reactions on which no new work has been done.)


Model calculations: (1979LA06).

Complex reactions involving ${}^9\text{C}$: (1981MO20).

Reactions involving pions: (1979AS01, 1979NA1E, 1980BU15, 1983HU02).

Other topics: (1979BE1H, 1979LA06, 1982NG01).

Mass of ${}^9\text{C}$: The recent Q_0 value for the ${}^{12}\text{C}({}^3\text{He}, {}^6\text{He}){}^9\text{C}$ reaction (see reaction 3) has not been published. We therefore adopt the Wapstra mass excess: 28913.1 ± 2.5 keV.



The half-life of ${}^9\text{C}$ is 126.5 ± 0.9 msec: see (1974AJ01). Several groups of delayed protons are observed indicating the involvement of a number of ${}^9\text{B}$ states: see Table 9.10 (1972ES05). See also (1978RA2A).

Table 9.12: Energy levels of ${}^9\text{C}$

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or Γ	Decay	Reactions
g.s. 2.218 ± 11	$(\frac{3}{2}^-); \frac{3}{2}$	$\tau_{1/2} = 126.5 \pm 0.9$ msec $\Gamma = 100 \pm 20$ keV	β^+	1, 2, 3 ^a 3

^a See also (AJ74, AJ79).



${}^9\text{C}_{\text{g.s.}}$ has been observed at $E_{\pi^+} = 180$ MeV (1980BU15). See also (1979NA1E).



$$Q_0 = -31.5762 \pm 0.0032: \text{ see (1980KA1J).}$$

At $E(\text{}^3\text{He}) = 74.1$ MeV a ${}^6\text{He}$ group is observed to the first excited state of ${}^9\text{C}$ (1974BE66). Based on $Q_m, E_x = 2218 \pm 11$ keV and $\Gamma = 100 \pm 20$ keV.

${}^9\text{N}$
(Not illustrated)

Not observed: see (1979AJ01). See also (1982NG01; theor.).

References

(Closed 1 June 1983)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1960KA17 R.W. Kavanagh, Nucl. Phys. 18 (1960) 492
- 1964MI04 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 50
- 1964TE01 E. Teranishi and B. Furubayashi, Phys. Lett. 9 (1964) 157
- 1966DE13 V.P. Denisov and L.A. Kulchitskii, Yad. Fiz. 3 (1966) 268; Sov. J. Nucl. Phys. 3 (1966) 192
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966WI08 D.H. Wilkinson, J.T. Sample and D.E. Alburger, Phys. Rev. 146 (1966) 662
- 1968CO08 C.L. Cocke and P.R. Christensen, Nucl. Phys. A111 (1968) 623
- 1968KR02 J.J. Kroepfl and C.P. Browne, Nucl. Phys. A108 (1968) 289
- 1968KU04 L.A. Kull and E. Kashy, Phys. Rev. 167 (1968) 963
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1970AN07 J.D. Anderson, C. Wong, B.A. Pohl and J.W. McClure, Phys. Rev. C2 (1970) 319
- 1970AU02 D.L. Auton, Nucl. Phys. A157 (1970) 305
- 1970CH07 Y.S. Chen, T.A. Tombrello and R.W. Kavanagh, Nucl. Phys. A146 (1970) 136
- 1971HA10 J.C. Hardy, J.M. Loiseaux, J. Cerny and G.T. Garvey, Nucl. Phys. A162 (1971) 552
- 1971YO04 P.G. Young and R.H. Stokes, Phys. Rev. C4 (1971) 1597
- 1972DE01 R.M. Devries, J.W. Sunier, J.-L. Perrenoud, M. Singh, G. Paic and I. Slaus, Nucl. Phys. A178 (1972) 417
- 1972ES05 J.E. Esterl, D. Allred, J.C. Hardy, R.G. Sextro and J. Cerny, Phys. Rev. C6 (1972) 373
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974BE66 W. Benenson and E. Kashy, Phys. Rev. C10 (1974) 2633
- 1974BI14 J. Birchall, H.E. Conzett, J. Arvieux, W. Dahme and R.M. Larimer, Phys. Lett. B53 (1974) 165
- 1974KA15 E. Kashy, W. Benenson and J.A. Nolen, Jr., Phys. Rev. C9 (1974) 2102
- 1975KA05 J. Kasagi, T. Nakagawa, N. Sekine, T. Tohei and H. Ueno, Nucl. Phys. A239 (1975) 233

- 1976AL02 D.E. Alburger and D.H. Wilkinson, Phys. Rev. C13 (1976) 835
- 1976MA34 J.L. Matthews, D.J.S. Findlay, S.N. Gardiner and R.O. Owens, Nucl. Phys. A267 (1976) 51
- 1976SC14 A.E. Schilling, N.F. Mangelson, K.K. Nielsen, D.R. Dixon, M.W. Hill, G.L. Jensen and V.C. Rogers, Nucl. Phys. A263 (1976) 389
- 1976ST11 W.F. Steele, P.A. Smith, J.E. Finck and G.M. Crawley, Nucl. Phys. A266 (1976) 424
- 1976WE17 E.W. Weber and J. Vetter, Phys. Lett. A56 (1976) 446
- 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, Phys. Rev. C14 (1976) 815
- 1977DE08 C. Detraz, Phys. Lett. B66 (1977) 333
- 1977GR1C Grechukhin and Soldatov, Zh. Eksp. Teor. Fiz. 72 (1977) 390
- 1977KO1D Kolesnikov, Khrokhin and Kopylov, Bull. Acad. Sci. USSR Phys. Ser. 41 (1977) 85
- 1977RO1C Robaye and Delbrouck-Habaru, Bull. Soc. R. Sci. Liege (Belgium) 46 (1977) 375
- 1977ST20 D.P. Stahel, G.J. Wozniak, M.S. Zisman, B.D. Jeltema and J. Cerny, Phys. Rev. C16 (1977) 1456
- 1977WE03 R.B. Weisenmiller, N.A. Jelley, D. Ashery, K.H. Wilcox, G.J. Wozniak, M.S. Zisman and J. Cerny, Nucl. Phys. A280 (1977) 217
- 1977WE1B Weisenmiller, Thesis, Univ. California, Berkeley (1976) - LBL-5077 (1977)
- 1978AJ02 F. Ajzenberg-Selove, E.R. Flynn and O. Hansen, Phys. Rev. C17 (1978) 1283
- 1978AL1G Alkhazov, Belostotsky and Vorobyov, Phys. Rep. C42 (1978) 89
- 1978AL37 M.R. Aleksic and R.V. Popic, Fizika (Yugoslavia) 10 (1978) 273
- 1978AN1B Antipov et al, Yad. Fiz. 28 (1978) 1299
- 1978AR1H Arima, in Clustering, Winnipeg 1978, AIP Conf. Proc. 47 (1978) 1
- 1978AR1J Arefev et al, Yad. Fiz. 28 (1978) 1534
- 1978AR21 N. Arena, D. Vinciguerra, M. Lattuada, F. Riggi and C. Spitaleri, Nuovo Cim. A45 (1978) 405
- 1978BA1F Barschall, Ann. Rev. Nucl. Part. Sci. 28 (1978) 207
- 1978BE1G Berezhnoi, Klyucharev and Rutkevich, Ukr. Fiz. Zh. 23 (1978) 1841
- 1978BH03 R.K. Bhowmik, E.C. Pollacco, N.E. Sanderson, J.B.A. England and G.C. Morrison, Phys. Lett. B80 (1978) 41
- 1978BU1B Buffington, Orth and Mast, Astrophys. J. 226 (1978) 355
- 1978CH1E Cherkasov, Yad. Fiz. 28 (1978) 639
- 1978DA1A Dalitz and Gal, Ann. Phys. 116 (1978) 167
- 1978DE32 S.V. Dementii, V.I. Ogurtsov and N.G. Afanasev, Yad. Fiz. 28 (1978) 865; Sov. J. Nucl. Phys. 28 (1978) 443

- 1978DI08 P.A. Dickey, P.L. Dyer, K.A. Snover and E.G. Adelberger, *Phys. Rev. C* 18 (1978) 1973
- 1978DU1B Dubna-Warsaw-Leningrad Collaboration, *Yad. Fiz.* 27 (1978) 1246
- 1978ER1A Ericson, *Prog. Part. Nucl. Phys.* 1 (1978) 67
- 1978FU09 S. Furui, *Nucl. Phys.* A312 (1978) 311
- 1978GE11 D.F. Geesaman, C. Olmer, B. Zeidman, R.L. Boudrie, R.H. Siemssen, J.F. Amann, C.L. Morris, H.A. Thiessen, G.R. Burleson, M.J. Devereux et al, *Phys. Rev. C* 18 (1978) 2223
- 1978GO14 N.F. Golovanova and I.M. Ilin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 1528; *Bull. Acad. Sci. USSR Phys. Ser.* 42 (1978) 154
- 1978GR22 K.A. Gridnev, N.Z. Darvish, A.S. Demyanova, V.M. Semenov, V.B. Subbotin and E.F. Khefter, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 2361; *Bull. Acad. Sci. USSR Phys. Ser.* 42 (1978) 124
- 1978GU15 O.P. Gupta, A. Bond and P.M.S. Lesser, *Phys. Rev. C* 18 (1978) 1075
- 1978HE1C Hendrie, in *Clustering, Winnipeg 1978*, AIP Conf. Proc. 47 (1978) 402
- 1978HE1D Heitzmann, *Atomkernenergie* 31 (1978) 262
- 1978HO23 H.H. Hogue, P.L. Von Behren, D.H. Epperson, S.G. Glendinning, P.W. Lisowski, C.E. Nelson, H.W. Newson, F.O. Purser, W. Tornow, C.R. Gould et al, *Nucl. Sci. Eng.* 68 (1978) 38
- 1978JA07 L. Jarczyk, J. Okolowicz, A. Strzalkowski, H. Witala, W. Zipper, J. Lang, R. Muller, E. Ungricht and J. Unternahrer, *Phys. Rev. C* 17 (1978) 2106
- 1978JE01 H. Jeremie, L. Lemay, M. Irshad and G. Kennedy, *Nucl. Phys.* A312 (1978) 43
- 1978KE13 B.K. Kerimov, M.Y. Safin and I.M. Al-Hamis, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 1976; *Bull. Acad. Sci. USSR Phys. Ser.* 42 (1978) 153
- 1978KI13 H.R. Kissener, G.E. Dogotar, R.A. Eramzhyan and R.A. Sakaev, *Nucl. Phys.* A312 (1978) 394
- 1978KR16 S.P. Krekoten and A.M. Mukhamedzhanov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 2405; *Bull. Acad. Sci. USSR Phys. Ser.* 42 (1978) 162
- 1978KU06 E.L. Kuplennikov, V.A. Goldshtein and N.G. Afanasev, *Sov. J. Nucl. Phys.* 27 (1978) 313
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, *Table of Isotopes 7th ed.* (New York: John Wiley & Sons, 1978)
- 1978MA44 J.F. Mateja, A.D. Frawley, A. Roy, J.R. Hurd and N.R. Fletcher, *Phys. Rev. C* 18 (1978) 2622
- 1978OR1A Orth, Buffington, Smoot and Mast, *Astrophys. J.* 226 (1978) 1147

- 1978OS1B Osman. *Nuovo Cim.* A46 (1978) 477
- 1978PO1A B. Povh, *Ann. Rev. Nucl. Part. Sci.* 28 (1978) 1
- 1978RA2A S. Raman, C.A. Houser, T.A. Walkiewicz and I.S. Towner, *At. Data Nucl. Data Tables* 21 (1978) 567; *IBID* 22 (1978) 369
- 1978RE1A Redish, in *Few Body Syst. and Nucl. Forces, Graz 1978*, Springer Lecture Notes 87 (1978) 427
- 1978SA26 A. Saha, R. Kamermans, J. van Driel and H.P. Morsch, *Phys. Lett.* B79 (1978) 363
- 1978SC02 P.A. Schmelzbach, R.A. Hardekopf, R.F. Haglund, Jr. and G.G. Ohlsen, *Phys. Rev.* C17 (1978) 16
- 1978SO1A Sokol, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 1829
- 1978TA12 S. Tanaka, K. Fukunaga, S. Kakigi, T. Ohsawa, N. Fujiwara and T. Yanabu, *J. Phys. Soc. Jpn.* 45 (1978) 733
- 1978TA1B Taras, in *Clustering, Winnipeg*, AIP Conf. Proc. 47 (1978) 234
- 1978TA1D Tartakovskii, Kozlovskii and Malyarzh, *Ukr. Fiz. Zh. USSR* 23 (1978) 368
- 1978UN02 J. Unternahrer, M. Hugi, J. Lang, R. Muller, E. Ungricht, L. Jarczyk, B. Kamys and A. Strzalkowski, *Helv. Phys. Acta* 51 (1978) 115
- 1978VL1B Vlasov et al, *Yad. Fiz.* 27 (1978) 413
- 1978WA1B Walter, in *Clustering, Winnipeg 1978*, AIP Conf. Proc. 47 (1978) 444
- 1978WO1A Woloshyn, in *Clustering, Winnipeg 1978*, AIP Conf. Proc. 47 (1978) 457
- 1978ZE03 N.S. Zelenskaya, V.M. Lebedev and T.A. Yushchenko, *Yad. Fiz.* 28 (1978) 90; *Sov. J. Nucl. Phys.* 28 (1978) 44
- 1979AB11 S.N. Abramovich, B.Y. Guzhovsky, A.G. Zvenigorodsky and S.V. Trusillo, *Yad. Fiz.* 30 (1979) 1276; *Sov. J. Nucl. Phys.* 30 (1979) 665
- 1979AJ01 F. Ajzenberg-Selove, *Nucl. Phys.* A320 (1979) 1
- 1979AL1J Alster and Warszawski, *Phys. Rept.* 52 (1979) 87
- 1979AL22 Y. Alhassid, R.D. Levine, J.S. Karp and S.G. Steadman, *Phys. Rev.* C20 (1979) 1789
- 1979AL26 G.D. Alkhazov, S.L. Belostotskii, A.A. Vorobev, O.A. Domchenkov, Y.V. Dotsenko, N.P. Kuropatkin, V.N. Nikulin and M.A. Shuvaev, *Pisma Zh. Eksp. Teor. Fiz.* 29 (1979) 88; *JETP Lett.* 29 (1979) 80
- 1979AS01 E. Aslanides, P. Fassnacht, F. Hibou, E. Chiavassa, G. Dellacasa, M. Gallio, A. Musso, T. Bressani and G. Puddu, *Phys. Rev. Lett.* 43 (1979) 1466
- 1979BA10 G. Baur, M. Pauli, F. Rosel and D. Trautmann, *Nucl. Phys.* A315 (1979) 241
- 1979BA16 B. Bassalleck, H.-D. Engelhardt, W.D. Klotz, F. Takeutchi, H. Ullrich and M. Furic, *Nucl. Phys.* A319 (1979) 397

- 1979BA28 Y.D. Bayukov, V.I. Efremenko, S. Frankel, W. Frati, M. Gazzaly, G.A. Leksin, N.A. Nikiforov, C.F. Perdrisat, V.I. Tchistilin and Y.M. Zaitsev, Phys. Rev. C20 (1979) 764
- 1979BA34 Y.D. Bayukov, V.I. Efremenko, S. Frankel, W. Frati, M. Gazzaly, G.A. Leksin, N.A. Nikiforov, C.F. Perdrisat, V.I. Tchistilin and Y.M. Zaitsev, Phys. Lett. B85 (1979) 315
- 1979BA60 H.W. Barz, Nucl. Phys. A330 (1979) 273
- 1979BE1H Benenson and Kashy, Revs. Mod. Phys. 51 (1979) 527
- 1979BE1J Bennett and Kouzes, Bull. Amer. Phys. Soc. 24 (1979) 43
- 1979BO1K Bodek et al, Helv. Phys. Acta 52 (1979) 88
- 1979BO1N Bobchenko et al, Yad. Fiz. 30 (1979) 1553
- 1979BO21 E.C. Booth, B. Chasan, F.L. Milder and B.L. Roberts, Phys. Rev. C20 (1979) 1603
- 1979BO22 V.I. Bogatin, O.V. Lozhkin and Y.P. Yakovlev, Nucl. Phys. A326 (1979) 508
- 1979BO27 D.H. Boal and R.M. Woloshyn, Phys. Rev. C20 (1979) 1878
- 1979BU11 A.Y. Buki, N.G. Shevchenko, V.N. Polishchuk, A.A. Khomich and I.I. Chkalov, Yad. Fiz. 30 (1979) 5; Sov. J. Nucl. Phys. 30 (1979) 2
- 1979BU1D Burgov et al, in Conf. Proc. TRIUMF, Vancouver (1979) papers 1E4, 1E6
- 1979BU1E Buzhinski et al, Izv. Akad. Nauk SSSR Ser. Fiz. 43 (1979) 838
- 1979BU1G Burgov et al, Yad. Fiz. 30 (1979) 720
- 1979BY01 R.C. Byrd, R.L. Walter and S.R. Cotanch, Phys. Rev. Lett. 43 (1979) 260
- 1979CH1D Chen, Zheng and Lu, Phys. Energ. Fortis Phys. Nucl. 3 (1979) 494; Phys. Abs. 93518 (1979)
- 1979CO1D Cooper, in NATO Institute Vol. 45 (Plenum Press, 1979) 667
- 1979DE06 J. Deutsch, D. Favart, M. Lebrun, P. Lipnik, P. Macq and R. Prieels, Phys. Lett. B80 (1979) 347
- 1979DE15 P. Desgrolard and P.A.M. Guichon, Z. Phys. A290 (179) 373
- 1979DE1G Deutsch, in NATO Institute Vol. 45 (Plenum Press, 1979) 165
- 1979DO03 E.I. Dolinskii, S.P. Krekoten, A.M. Mukhamedzhanov and E.A. Romanovskii, Yad. Fiz. 29 (1979) 71; Sov. J. Nucl. Phys. 29 (1979) 34
- 1979DO17 G.E. Dogotar, R.A. Eramzhyan, M. Gmitro, H.R. Kissener and E. Tinkova, J. Phys. G5 (1979) L221
- 1979DY01 P. Dyer, T.C. Awes, C.K. Gelbke, B.B. Back, A. Mignerey, K.L. Wolf, H. Breuer, V.E. Viola, Jr. and W.G. Meyer, Phys. Rev. Lett. 42 (1979) 560
- 1979EL12 R.J. Ellis, H.J. Ziock, K.O.H. Ziock, J. Bolger, E. Boschitz, J. Arvieux, R. Corfu and J. Piffaretti, Phys. Lett. B88 (1979) 253
- 1979FO16 A.C. Fonseca, J. Revai and A. Matveenko, Nucl. Phys. A326 (1979) 182

- 1979FO1F Fou, Rasmussen, Swann and Van Patter, IEEE Trans. on Nucl. Sci. 26 (1979) 1378
- 1979FR12 S. Frankel, W. Frati, M. Gazzaly, Y.D. Bayukov, V.I. Efremenko, G.A. Leksin, N.A. Nikiforov, V.I. Tchistilin, Y.M. Zaitsev and C.F. Perdrisat, Phys. Rev. C20 (1979) 2257
- 1979FR1B Franz et al, Int. Conf. Nucl. Phys. with Electromagnetic interact., Mainz (1979) 11
- 1979FR1C Franz et al, in Conf. Proc. TRIUMF, Vancouver (1979) Papers 5C5, 5C6
- 1979FR1F Franz et al, Int. Conf. Nucl. Phys. with Electromagnetic interact., Mainz (1979) 12
- 1979FU1G Furui, in 1978 RPI Symp. "Photopion Nucl. Phys.," (Plenum Press, New York, 1979) p. 199
- 1979GA04 J.A. Gaidos, L.J. Gutay, A.S. Hirsch, R. Mitchell, T.V. Ragland, R.P. Scharenberg, F. Turkot, R.B. Willmann and C.L. Wilson, Phys. Rev. Lett. 42 (1979) 82
- 1979GA1E Gal, in NATO Inst. Vol. 45 (Plenum Press 1979) 485
- 1979GE1A Gelbke, in BNL-51115 (1979) 1
- 1979GI1B Gibbs et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper 1C14
- 1979GI1C Gibbs, in NATO Institute Vol. 45 (Plenum Press, 1979) p. 595
- 1979GO07 R. Gorgen, F. Hinterberger, R. Jahn, P. von Rossen and B. Schuller, Nucl. Phys. A320 (1979) 296
- 1979GR1E Greenwood, in Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York 1979) 441
- 1979HO13 B. Hoistad, G.S. Adams, M. Gazzaly, G. Igo, F. Irom and H. Nann, Phys. Rev. Lett. 43 (1979) 487
- 1979HO1C Holt et al, Bull. Amer. Phys. Soc. 24 (1979) 687
- 1979HU1B M.S. Hussein, Phys. Rev. C19 (1979) 807
- 1979IN1A Ingram, in Houston, AIP Conf. Proc. 54 (1979) 455
- 1979JA04 L. Jarczyk, J. Okolowicz, A. Strzalkowski, K. Bodek, M. Hugi, J. Lang, R. Muller and E. Ungricht, Nucl. Phys. A316 (1979) 139
- 1979JA1C Jackson et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper A417
- 1979JE04 P. Jennewein and B. Schoch, Z. Phys. A292 (1979) 27
- 1979JO1B Johnson et al, Bull. Amer. Phys. Soc. 24 (1979) 886
- 1979JU1B Journey, in Proc. Third Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York, 1979) p. 461
- 1979KA1G Karban, Microscopic Optical Potentials. Hamburg, Germany (Springer-Verlag 1979) p. 264; Phys. Abs. 97712 (1979)
- 1979KI10 M. Kirchbach and H.-U. Jager, Yad. Fiz. 29 (1979) 1191; Sov. J. Nucl. Phys. 29 (1979) 614

- 1979KI1D Kissener and Eramzhyan, Int. Conf. Nucl. Phys. With Electromag. Inter., MAINZ (1979) L.20
- 1979KI1G Kim and Primakoff, in "Mesons in Nucl.", Eds. Rho and Wilkenson (North-Holland, Amsterdam, 1979) 69
- 1979KO21 V.I. Komarov, G.E. Kosarev, H. Muller, D. Netzband, V.D. Toneev, T. Stiehler, S. Tesch, K.K. Gudima and S.G. Mashnik, Nucl. Phys. A326 (1979) 297
- 1979KR1A Krieger et al, Bull. Amer. Phys. Soc. 24 (1979) 62
- 1979LA06 R.D. Lawson, Phys. Rev. C19 (1979) 2359
- 1979LI1A Li and Shen, Phys. Energ. Fortis & Phys. Nucl. 3 (1979) 327; Phys. Abs. 80956 (1979)
- 1979LO11 H. Lohner, B. Ludewigt, D. Frekers, G. Gaul and R. Santo, Z. Phys. A292 (1979) 35
- 1979LU1A Lu et al, Scientia Sinica 22 (1979) 1248
- 1979MA02 W.M. MacDonald, E.T. Dressler and J.S. O'Connell, Phys. Rev. C19 (1979) 455
- 1979MA1J MacGregor, Phys, Rev. Lett. 42 (1979) 1724
- 1979MA1L Mandal, Bhattacharya, Saha and Sen, Z. Naturforschung 34 (1979) 242
- 1979MA1M Mahaux, in NATO Institute Vol. 45 (Plenum Press, 1979) p. 265
- 1979MA1P Mazzucato et al, in Houston, AIP Conf. Proc. 54 (1979) 418
- 1979MA21 J.F. Mateja, A.D. Frawley, P.B. Nagel and L.A. Parks, Phys. Rev. C20 (1979) 176
- 1979MA30 R.A. Malfliet and Y. Karant, Phys. Lett. B86 (1979) 251
- 1979MC1A McGuire, Von Roseninge and McDonald, Bull. Amer. Phys. Soc. 24 (1979) 693
- 1979MI1E Mingay, S. Afr. J. Phys. 2 (1979) 107
- 1979MO1G Mougey, Proc. Mainz, 1979, Springer Lecture Notes 108 (1979) 124
- 1979NA1E Nann et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper 1E27
- 1979OH1A Ohashi et al, 1978 RPI Symp. "Photopion Nucl. Phys., (Plenum Press, New York, 1979) p. 193
- 1979OK02 S. Okabe and Y. Abe, Prog. Theor. Phys. 61 (1979) 1049
- 1979OS02 A. Osman, Phys. Rev. C19 (1979) 1127
- 1979PA1B Park, Scheid and Greiner, Bull. Amer. Phys. Soc. 24 (1979) 665
- 1979PE1C Perroud, in 1978 RPI Symp. "Photopion Nucl. Phys., (Plenum Press, New York, 1979) p. 69
- 1979PE1D Petrov and Shabelskii, Sov. J. Nucl. Phys. 30 (1979) 66
- 1979PI1C Pickar et al, Bull. Amer. Phys. Soc. 24 (1979) 819
- 1979SA17 G.R. Satchler, Phys. Lett. B83 (1979) 284
- 1979SA1E Satchler and Love, Phys. Rept. 55 (1979) 183

- 1979SA27 G.R. Satchler, Nucl. Phys. A329 (1979) 233
- 1979SC08 T.C. Schweizer, R.R. Doering, S.T. Thornton, L.C. Dennis, K.R. Cordell and R.L. Parks, Phys. Rev. C19 (1979) 1408
- 1979SC1D Scott, Prog. Nucl. Phys. 4 (1979) 5
- 1979SC1E Schmitt, Borkowski, Simon and Walther, Int. Conf. Nucl. Phys. with Electromagnetic Interat., Mainz (1979) 6.9
- 1979SE04 F. Seiler and H.W. Roser, Nucl. Phys. A315 (1979) 45
- 1979SL01 R.J. Slobodrian, M. Irshad, R. Labrie, C. Rioux, R. Roy and R. Pigeon, Nucl. Instrum. Meth. 159 (1979) 413
- 1979ST1G Stephenson, Gibbs, Gibson and Devereux, Bull. Amer. Phys. Soc. 24 (1979) 574
- 1979SU1C Subramanian et al, Bull. Amer. Phys. Soc. 24 (1979) 593
- 1979TI1A Titov, Int. Conf. Nucl. Phys. with Electromagnetic Interact., Mainz (1979) 6.14
- 1979UN01 E. Ungricht, D. Balzer, M. Hugi, J. Lang, R. Muller, L. Jarczyk, B. Kamys and A. Strzalkowski, Nucl. Phys. A313 (1979) 376
- 1979VE1B S.P. Verma and D.P. Sural, Phys. Rev. C20 (1979) 781
- 1979VI05 A. Vidal-Quadras and M. Ortega, Nuovo Cim. A49 (1979) 235
- 1979WA1E Watermann et al, Med. Phys. (USA) 6 (1979) 160
- 1979WA1F Watermann et al, Med. Phys. (USA) 6 (1979) 432
- 1979WE06 J.P. Wefel and J.M. Kidd, Phys. Rev. C19 (1979) 1380
- 1979WI1E Winhold et al, Int. Conf. Nucl. Phys. with Electromagnetic Interat., Mainz (1979) 6.16
- 1979ZI1A Ziegler, Proc. Mainz, 1979, Springer Lecture Notes 108 (1979) 148
- 1980AB16 S.N. Abramovich, A.I. Baz and B.Ya. Guzhovsky, Yad. Fiz. 32 (1980) 402; Sov. J. Nucl. Phys. 32 (1980) 208
- 1980AH1A J. Ahrens, Nucl. Phys. A335 (1980) 67
- 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, Nucl. Phys. A336 (1980) 1
- 1980AK01 J.M. Akkermans, H. Gruppelaar and G. Reffo, Phys. Rev. C22 (1980) 73
- 1980AU1C N. Auerbach and N. Van Giai, Phys. Lett. B90 (1980) 354
- 1980BA2H Baryshnikov, Gulyamov and Krekoten, IAN (1980) 2421
- 1980BE12 R. Bertini, O. Bing, P. Birien, W. Bruckner, H. Catz, A. Chaumeaux, J.M. Durand, M.A. Faessler, T.J. Ketel, K. Kilian et al, Phys. Lett. B90 (1980) 375
- 1980BE30 M. Bedjidian, E. Descroix, J.Y. Grossiord, A. Guichard, M. Gusakow, M. Jacquin, M.J. Kudla, H. Piekarcz, J. Piekarcz, J.R. Pizzi et al, Phys. Lett. B94 (1980) 480
- 1980BO11 K. Bodek, M. Hugi, J. Lang, R. Muller, E. Ungricht, K. Jankowski, W. Kipper, L. Jarczyk, A. Strzalkowski, G. Willim et al, Nucl. Phys. A339 (1980) 353

- 1980BO12 D.H. Boal, Phys. Rev. C21 (1980) 1913
- 1980BO14 K. Bodek, M. Hugi, J. Lang, R. Muller, A. Schiltz, J. Sromicki, E. Ungricht, L. Jarczyk and A. Strzalkowski, Phys. Lett. B92 (1980) 79
- 1980BO1L Boucenna, Allab and Hadda, Bull. Amer. Phys. Soc. 25 (1980) 785
- 1980BO1N Bohigas, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 116
- 1980BO31 V.I. Bogatin, E.A. Ganza, O.V. Lozhkin, Yu.A. Murin and V.S. Oplavin, Yad. Fiz. 32 (1980) 27; Sov. J. Nucl. Phys. 32 (1980) 14
- 1980BR05 V.N. Bragin, M.V. Zhukov, D.N. Stepanov, L.V. Chulkov and J. Jakiel, Yad. Fiz. 31 (1980) 29; Sov. J. Nucl. Phys. 31 (1980) 14
- 1980BR1P Braun and Vechernin, Sov. J. Nucl. Phys. 31 (1980) 312
- 1980BU15 G.R. Burlison, G.S. Blanpied, G.H. Daw, A.J. Viescas, C.L. Morris, H.A. Thiessen, S.J. Greene, W.J. Braithwaite, W.B. Cottingham, D.B. Holtkamp et al, Phys. Rev. C22 (1980) 1180
- 1980CI1B Ciric, Zb. Rad. Prir-Mat. Fak. Ser. Fiz. 10 (1980) 257
- 1980CO1R Cook, Stone and Vogl, Astrophys. J. 238 (1980) L97
- 1980CO1U Corcalciuc, Stud. Cercet. Fiz. 32 (1980) 1135, 1166
- 1980CR03 J.F. Crawford, M. Daum, G.H. Eaton, R. Frosch, H. Hirschmann, R. Horisberger, J.W. McCulloch, E. Steiner, R. Hausammann, R. Hess et al, Phys. Rev. C22 (1980) 1184
- 1980DA07 S.K. Datta, W.R. Falk, S.P. Kwan, R. Abegg and O.A. Abou-Zeid, Phys. Rev. C21 (1980) 1687
- 1980DE11 A. Deloff, Phys. Rev. C21 (1980) 1516
- 1980DO1E Dover and Gal, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 795
- 1980EC01 J.S. Eck, T.R. Ophel, P.D. Clark and D.C. Weissner, Nucl. Phys. A334 (1980) 519
- 1980EC03 J.S. Eck, J.R. Leigh, T.R. Ophel and P.D. Clark, Phys. Rev. C21 (1980) 2352
- 1980EC04 J.S. Eck, T.R. Ophel, P.D. Clark, D.C. Weissner and G.R. Satchler, Nucl. Phys. A341 (1980) 178
- 1980EL02 A.J. Elwyn, R.E. Holland, C.N. Davids, J.E. Monahan, F.P. Mooring and W. Ray, Jr., Phys. Rev. C22 (1980) 1406
- 1980ER01 R.A. Eramzhyan, M. Gmitro and H.R. Kissener, Nucl. Phys. A338 (1980) 436
- 1980FA07 E. Fabrici, S. Micheletti, M. Pignanelli, F.G. Resmini, R. De Leo, G. D'Erasmo and A. Pantaleo, Phys. Rev. C21 (1980) 844
- 1980FU1G Furutani et al, Suppl. Prog. Theor. Phys. 68 (1980) 193
- 1980GA1C A. Gal and C.B. Dover, Phys. Rev. Lett. 44 (1980) 379
- 1980GA1P Gal, Nukelonika 25 (1980) 447

- 1980GI1C Gai and Auerbach, *Nukleonika* 25 (1980) 596
- 1980GR10 R.E.L. Green and R.G. Korteling, *Phys. Rev. C* 22 (1980) 1594
- 1980GU26 B.Ya. Guzhovsky, S.N. Abramovich, A.G. Zvenigorodsky, V.P. Protopopov, G.N. Sleptsov and S.V. Trusillo, *Izv. Akad. Nauk. SSSR Ser. Fiz.* 44 (1980) 1983
- 1980HA29 S. Hayashi and S. Iwata, *Phys. Lett.* B95 (1980) 39
- 1980HA33 M.N. Harakeh, J. Van Popta, A. Saha and R.H. Siemssen, *Nucl. Phys.* A344 (1980) 15
- 1980HO22 S. Homma, M. Kanazawa, K. Maruyama, Y. Murata, H. Okuno, A. Sasaki and T. Taniguchi, *Phys. Rev. Lett.* 45 (1980) 706
- 1980HU09 M. Hugi, L. Jarczyk, B. Kamys, J. Lang, R. Muller, A. Strzalkowski, E. Ungricht and W. Zipper, *J. Phys.* G6 (1980) 1257
- 1980HU1D Hugg and Hanna, *Proc. Int. Conf. on Nucl. Phys. (Berkeley)* (1980) 891
- 1980IS1E Ishkanov, Morkeev, Novikov and Piskarev, *Yad. Fiz.* 32 (1980) 11
- 1980JE1C Jenkins et al, *Proc. Int. Conf. on Nucl. Phys. (Berkeley)* (1980) 685
- 1980KA1J Kashy, Beneson, Nolen and Robertson, *Proc. Int. Conf. in At. Masses and Fund. Constants, Eds., Nolen and Benenson, Plenum Publ.* (1980) 39
- 1980KI1B Kilius et al, *Nucl. Instrum. Meth. Phys. Res.* 171 (1980) 355
- 1980KO1V Koptev, Maev, Makarov and Khanzadeev, *Yad. Fiz.* 31 (1980) 1501
- 1980LE02 T.-S.H. Lee and D. Kurath, *Phys. Rev. C* 21 (1980) 293
- 1980LE06 M. LeMere and Y.C. Tang, *Phys. Rev. C* 21 (1980) 1170
- 1980MA1F Maleki, Rao and Levinger, *Bull. Amer. Phys. Soc.* 25 (1980) 606
- 1980MA1Z Mandal and Saha, *Can. J. Phys.* 58 (1980) 300
- 1980ME1F Meyer, *Proc. Int. Conf. on Nucl. Phys. (Berkeley)* (1980) 683
- 1980MI01 T. Mikumo, M. Sasagase, M. Sato, T. Ooi, Y. Higashi, Y. Nagashima and M. Yamanoichi, *Phys. Rev. C* 21 (1980) 620
- 1980MI1G D.J. Millener, *Phys. Rev. C* 22 (1980) 1355
- 1980MO1C J. Mougey, *Nucl. Phys.* A335 (1980) 35
- 1980MO28 G.C. Morrison, R.K. Bhowmik, E.C. Pollacco, J.B.A. England and N.E. Sanderson, *J. Phys. (Paris) Colloq.* C10 (1980) 243
- 1980MU1B N.C. Mukhopadhyay, *Nucl. Phys.* A335 (1980) 111
- 1980NA09 A. Nadasen, N.S. Chant, P.G. Roos, T.A. Carey, R. Cowen, C. Samanta and J. Wesick, *Phys. Rev. C* 22 (1980) 1394
- 1980NA1D Nann, *Proc. Int. Conf. in At. Masses and Fund. Constants, Eds., Nolen and Benenson, Plenum Publ.* (1980) 51

- 1980NA1L Nakamura, J. Sci. Hiroshima Univ. A44 (1980) 215; Phys. Abs. 50656 (1981)
- 1980NI03 M. Nilsson, B. Schroder, B. Bulow, J. Grintals, G.G. Jonsson, B. Lindner, K.S. Rao and S. Susila, Z. Phys. A294 (1980) 253
- 1980OH1B Ohkubo, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 477
- 1980PO1A B. Povh, Nucl. Phys. A335 (1980) 233
- 1980RE02 J. Revai and A.V. Matveenko, Nucl. Phys. A339 (1980) 448
- 1980RE1B Read et al, Bull. Amer. Phys. Soc. 25 (1980) 592
- 1980SA1E R. Sartor and C. Mahaux, Phys. Rev. C21 (1980) 2613
- 1980SC18 L.A. Schaller, L. Schellenberg, A. Ruetschi and H. Schneuwly, Nucl. Phys. A343 (1980) 333
- 1980SE1C Seth et al, Bull. Amer. Phys. Soc. 25 (1980) 505
- 1980SE1F Seth et al, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 164
- 1980ST15 P. Stoler, E.J. Winhold, F. O'Brien, P.F. Yergin, D. Rowley, K. Min, J. LeRose, A.M. Bernstein, K.I. Blomqvist, G. Franklin et al, Phys. Rev. C22 (1980) 911
- 1980TA1D Tavares et al, Lett. Nuovo Cim. 27 (1980) 358
- 1980TH1C Thomas and Landau, Phys. Rept. 58 (1980) 121
- 1980TR1A P. Truol, Nucl. Phys. A335 (1980) 55
- 1980WA07 C.W. Wang, N.S. Chant, P.G. Roos, A. Nadasen and T.A. Carey, Phys. Rev. C21 (1980) 1705
- 1980WI1L L. Winsberg, Phys. Rev. C22 (1980) 2123
- 1980WI1M Wiedenbeck and Greiner, Astrophys. J. 239 (1980) L139
- 1980ZE05 N.S. Zelenskaya, V.M. Lebedev and A.G. Serebryakov, Izv. Akad. Nauk SSSR Ser. Fiz. 44 (1980) 189; Bull. Acad. Sci. USSR Phys. Ser. 44 (1980) 158
- 1980ZH1A Zhang et al, Proc. Int. Conf. on Nucl. Phys., Berkeley (1980) p. 920
- 1980ZH1C Zheng, Li and Shen, Transactions of the N.Y. Acad. of Sci. 40 (1980) 274
- 1980ZI02 M.S. Zisman, J.G. Cramer, D.A. Goldberg, J.W. Watson and R.M. DeVries, Phys. Rev. C21 (1980) 2398
- 1980ZO1A Zofka, Czech. J. Phys. B30 (1980) 95
- 1981AJ01 F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1
- 1981AR04 J. Arends, J. Eyink, A. Hegerath, K.G. Hilger, B. Mecking, G. Noldeke and H. Rost, Nucl. Phys. A358 (1981) 367c
- 1981AS1H Aslanides et al, Proc. Versailles Conf. (1981) p. 203
- 1981AU1D Audouze, Prog. Part. Nucl. Phys. 6 (1981) 125
- 1981AU1G Austin, Prog. Part. Nucl. Phys. 7 (1981) 1

- 1981AV02 I.K. Averyanov, A.I. Golubev and A.A. Sadovoy, *Yad. Fiz.* 33 (1981) 66
- 1981BA20 O.L. Bartaya and J.V. Mebonia, *Yad. Fiz.* 33 (1981) 987
- 1981BA2M Bayukov et al, *Yad. Fiz.* 34 (1981) 785
- 1981BA2N Bando, Seki and Shono, *Prog. Theor. Phys.* 66 (1981) 2118; *Prog. Theor. Phys.* 68 (1982) 346
- 1981BA2R Bayukov et al, *Yad. Fiz.* 34 (1981) 1511
- 1981BA37 A.K. Basak, O. Karban, S. Roman, G.C. Morrison, C.O. Blyth and J.M. Nelson, *Nucl. Phys. A368* (1981) 74
- 1981BA38 A.K. Basak, O. Karban, S. Roman, G.C. Morrison, C.O. Blyth and J.M. Nelson, *Nucl. Phys. A368* (1981) 93
- 1981BA50 A.G. Baryshnikov, E.I. Dolinsky and S.P. Krekoten, *Izv. Akad. Nauk SSSR Ser. Fiz.* 45 (1981) 169
- 1981BE13 B. Bellinghausen, A. Christ, H.J. Gassen, G. Goerigk, R. Muller, G. Noldeke, T. Reichelt, H. Stanek and P. Stipp, *Nucl. Phys. A358* (1981) 373c
- 1981BE17 R. Bertini, O. Bing, P. Birien, K. Braune, W. Bruckner, H. Catz, A. Chaumeaux, M.A. Faessler, R.-W. Frey, D. Garreta et al, *Nucl. Phys. A360* (1981) 315
- 1981BE1X G. Berlad and A. Dar, *Phys. Lett. B102* (1981) 385
- 1981BE45 R. Bertini, O. Bing, P. Birien, K. Braune, W. Bruckner, A. Chaumeaux, M.A. Faessler, R.W. Frey, D. Garreta, T.J. Ketel et al, *Nucl. Phys. A368* (1981) 365
- 1981BE63 V.B. Belyaev and O.P. Solovtsova, *Yad. Fiz.* 33 (1981) 699; *Sov. J. Nucl. Phys.* 33 (1981) 363
- 1981BH02 R.K. Bhowmik, E.C. Pollacco, J.B.A. England, G.C. Morrison and N.E. Sanderson, *Nucl. Phys. A363* (1981) 516
- 1981BI1F Bimbot et al, *Proc. Versailles Conf.* (1981) p. 201
- 1981BJ01 T. Bjornstad, H.A. Gustafsson, P.G. Hansen, B. Jonson, V. Lindfors, S. Mattsson, A.M. Poskanzer, H.L. Ravn, the ISOLDE Collaboration, *Nucl. Phys. A359* (1981) 1
- 1981BL1G M. Blann and T.T. Komoto, *Phys. Rev. C24* (1981) 426
- 1981BO06 T.J. Bowles, D.F. Geesaman, R.J. Holt, H.E. Jackson, J. Julien, R.M. Laszewski, J.R. Specht, E.J. Stephenson, R.P. Redwine, L.L. Rutledge, Jr. et al, *Phys. Rev. C23* (1981) 439
- 1981BO09 A. Bouyssy, *Phys. Lett. B99* (1981) 373
- 1981BO1T Bogdanov, Kerimov and Safi n, *Proc. Samarkand Conf.* (1981) p. 516
- 1981BO1X Bogatin et al, *Yad. Fiz.* 34 (1981) 104
- 1981BO1Y Bouten and Bouten, *Prog. Part. Nucl. Phys.* 5 (1981) 55
- 1981BR1K Brady et al, *Bull. Amer. Phys. Soc.* 26 (1981) 1128

- 1981CH30 A.S. Cherkasov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 45 (1981) 783
- 1981CI03 M.A. Cirit and F. Yazici, *Phys. Rev. C*23 (1981) 2627
- 1981CO08 W.D. Cornelius, J.M. Moss and T. Yamaya, *Phys. Rev. C*23 (1981) 1364
- 1981CO2D Conzett, in *Polarization, Santa Fe 1980*, AIP Conf. Proc. 69 (1981) 1422
- 1981DA1C R.H. Dalitz, *Nucl. Phys.* A354 (1981) 101
- 1981DA1D Dalitz and Gal, *Ann. Phys.* 131 (1981) 314
- 1981DA1K Dave et al, *Bull. Amer. Phys. Soc.* 26 (1981) 551
- 1981DE18 E.S. de Almeida and V. di Napoli, *Lett. Nuovo Cim.* 31 (1981) 311
- 1981DE1T Dementij, Fartushnyj and Afanas'ev, *Proc. Versailles Conf.* (1981) p.127
- 1981DE1X Delbar, IPC-N-8102 (1981)
- 1981DE33 E.S. de Almeida, V. di Napoli and F. Salvetti, *Lett. Nuovo Cim.* 32 (1981) 109
- 1981DU1H Dumbrajs, *Fortschr. Phys.* 29 (1981) 487
- 1981FA1E Farwell, Grootes and Schmidt, *Bull. Amer. Phys. Soc.* 26 (1981) 1159
- 1981FE2A H.W. Fearing, *Prog. Part. Nucl. Phys.* 7 (1981) 113
- 1981FR24 S. Frankel, W. Frati, C.F. Perdrisat and O.B. Van Dyck, *Phys. Rev. C*24 (1981) 2684
- 1981GA1C Garcia-Munoz, Guzik, Simpson and Wefel, *Bull. Amer. Phys. Soc.* 26 (1981) 557
- 1981GE03 D.F. Geesaman, C. Olmer, B. Zeidman, R.L. Boudrie, G.S. Blanpied, M.J. Devereux, G.R. Burlson, R.E. Segel, L.W. Swenson and H.A. Thiessen, *Phys. Rev. C*23 (1981) 2635
- 1981GL03 Yu.A. Glukhov, A.S. Demyanova, S.I. Drozdov, M.V. Zhukov, V.I. Manko, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov and L.V. Chulkov, *Yad. Fiz.* 34 (1981) 312; *Sov. J. Nucl. Phys.* 34 (1981) 177
- 1981GO16 O.Yu. Goryunov, V.N. Dobrikov, O.F. Nemets, A.t. Rudchik, A.S. Gass and V.A. Stepanenko, *Ukr. Fiz. Zh.* 26 (1981) 1273
- 1981GR17 K.A. Gridnev, K. Mikulas, V.M. Semenov and E.F. Khefter, *Izv. Akad. Nauk SSSR Ser. Fiz.* 45 (1981) 134
- 1981GU12 K. Gul, M. Anwar, M. Ahmad, S.M. Saleem and N.A. Khan, *Phys. Rev. C*24 (1981) 2458
- 1981GU1B Gurbanovich and Zelenskaya, *Proc. Versailles Conf.* (1981) p. 142
- 1981GU1F S.A. Gurvitz, *Phys. Rev. Lett.* 47 (1981) 560
- 1981HA1K Haight, Grimes and Johnson, *Bull. Amer. Phys. Soc.* 26 (1981) 551
- 1981HN02 V. Hnizdo, K.W. Kemper and J. Szymakowski, *Phys. Rev. Lett.* 46 (1981) 590
- 1981HN04 V. Hnizdo, J. Szymakowski, K.W. Kemper and J.D. Fox, *Phys. Rev. C*24 (1981) 1495
- 1981HO1E Holland et al, *Nucl. Sci.* 28 (1981) 1344

- 1981HU07 M. Hugi, L. Jarczyk, B. Kamys, J. Lang, R. Muller, J. Sromicki, A. Strzalkowski, E. Ungricht, G. Willim, Z. Wrobel et al, J. Phys. G7 (1981) 1395
- 1981HU08 M. Hugi, J. Lang, R. Muller, E. Ungricht, K. Bodek, L. Jarczyk, B. Kamys, A. Magiera, A. Strzalkowski and G. Willim, Nucl. Phys. A368 (1981) 173
- 1981HU1C Ed V. Hungerford, Nucl. Phys. A353 (1981) 75
- 1981IO01 D.B. Ion and R. Ion-Mihai, Nucl. Phys. A360 (1981) 400
- 1981IS11 B.S. Ishkhanov, I.M. Kapitonov, V.G. Neudachin and R.A. Eramzhyan, Fiz. Elem. Chastits At. Yadra 12 (1981) 905; Sov. J. Part. Nucl. 12 (1981) 362
- 1981JA09 L. Jarczyk, B. Kamys, A. Magiera, J. Sromicki, A. Strzalkowski, G. Willim, Z. Wrobel, D. Balzer, K. Bodek, M. Hugi et al, Nucl. Phys. A369 (1981) 191
- 1981JOZV B. Jonson, H.A. Gustafsson, P.G. Hansen, P. Hoff, P.O. Larsson, S. Mattsson, G. Nyman, H.L. Ravn and D. Schardt, Proc. Int. Conf. Nuclei Far from Stability, Helsingor, Denmark, 1 (1981) 265; CERN 81-09 (1981)
- 1981KA04 C. Kalbach and F.M. Mann, Phys. Rev. C23 (1981) 112
- 1981KA1U Kanzaki et al, J. Phys. Soc. Jpn. 50 (1981) 3849
- 1981KE15 B.K. Kerimov, M.Ya. Safin and I.M. Al-Khamisi, Yad. Fiz. 34 (1981) 996; Sov. J. Nucl. Phys. 34 (1981) 554
- 1981KO1Q Kostin and Migaleny, Ukr. Fiz. Zh. SSSR 26 (1981) 803
- 1981KO28 KM. Kozlowski, H.H. Muller, P. Schober, Z. Lewandowski, E. Loeffler and R. Wagner, Lett. Nuovo Cim. 31 (1981) 565
- 1981KR17 S.P. Krekoten, A.M. Mukhamedzhanov and E.A. Romanovsky, Izv. Akad. Nauk SSSR Ser. Fiz. 45 (1981) 790
- 1981KR1J Kramer, John and Schenzle, Clustering Phenomena in Nuclei 2 (1981)
- 1981KU15 K. Kume, Phys. Lett. B106 (1981) 271
- 1981LA11 M. Langevin, C. Detraz, D. Guillemaud, F. Naulin, M. Epherre, R. Klapisch, S.K.T. Mark, M. De Saint Simon, C. Thibault and F. Touchard, Nucl. Phys. A366 (1981) 449
- 1981LA1E Laget and Chretien-Marquet, Proc. Versailles Conf. (1981) p. 418
- 1981LE03 D. Lebrun, M. Buenerd, M. Bini, B.G. Harvey, R. Legrain, J. Mahoney, T.J.M. Symons and K. Van Bibber, Phys. Rev. C23 (1981) 552
- 1981LE14 M. Lebrun, J. Deutsch, D. Favart, P. Lipnik, P. Macq and R. Prieels, Phys. Rev. C24 (1981) 754
- 1981MA1V Mateja et al, Bull. Amer. Phys. Soc. 26 (1981) 1132
- 1981MC09 R.D. McKeown, S.J. Sanders, J.P. Schiffer, H.E. Jackson, M. Paul, J.R. Specht, E.J. Stephenson, R.P. Redwine and R.E. Segel, Phys. Rev. C24 (1981) 211
- 1981ME13 M.C. Mermaz, J. Barrette and H.E. Wegner, Phys. Rev. C24 (1981) 2148

- 1981ME1D Meyer and Elbaz, Santa Fe 1980, AIP Conf. Proc. 69 (1981) 499
- 1981MO1H J. Mougey, Nucl. Phys. A358 (1981) 293
- 1981MO20 J. Mougey, R. Ost, M. Buenerd, A.J. Cole, C. Guet, D. Lebrun, J.M. Loiseaux, P. Martin, M. Maurel, E. Monnard et al, Phys. Lett. B105 (1981) 25
- 1981MU1E Mukhopadhyay and Hintermann, Santa Fe 1980, AIP Conf. Proc. 69 (1981) 1068
- 1981NA05 T. Nakamura, Y. Chiba, M. Aoki and S. Kaneko, Nucl. Phys. A365 (1981) 457
- 1981NI03 T. Nishi, I. Fujiwara, N. Imanishi, H. Moriyama, K. Otozai, R. Arakawa, S. Saito, T. Tsuneyoshi, N. Takahashi, S. Iwata et al, Nucl. Phys. A352 (1981) 461
- 1981NY1A Nyman et al, Proc. Conf., in Helsingor, June 1981, CERN 81-09 (1981) p. 312
- 1981PO1D B. Povh, Nucl. Phys. A358 (1981) 113
- 1981PO1F Povh, Prog. Part. Nucl. Phys. 5 (1981) 245
- 1981RE1B J. Revai and J. Zofka, Phys. Lett. B101 (1981) 228
- 1981RO1F Roy et al, Santa Fe 1980, AIP Conf. Proc.69 (1981) 1317
- 1981RU09 A.T. Rudchik, V.N. Dobrikov, O.Yu. Goryunov, O.F. Nemets, A.S. Gass and V.A. Stepanenko, Yad. Fiz. 34 (1981) 306; Sov. J. Nucl. Phys. 34 (1981) 173
- 1981SA1G Savin, Kruse and Moniot, Bull. Amer. Phys. Soc. 26 (1981) 543
- 1981SC1D Schmidt, Farwell and Grootes, Bull. Amer. Phys. Soc. 26 (1981) 593
- 1981SC1M Schatzman, CERN 81-11
- 1981SE06 M. Seya, M. Kohno and S. Nagata, Prog. Theor. Phys. 65 (1981) 204
- 1981SE1B Seth, 4th Int. Conf. on Nucl. Far From Stability, Helsingor, CERN Report 81-09 (1981) 655
- 1981SL03 R.J. Slobodrian, C. Rioux, R. Roy, H.E. Conzett, P. von Rossen and F. Hinterberger, Phys. Rev. Lett. 47 (1981) 1803
- 1981ST1G W. Stepien-Rudzka and S. Wycech, Nucl. Phys. A362 (1981) 349
- 1981ST1P Stokstad, Nukleonika 26 (1981) 373
- 1981SU1D Sunami and Narumi, Prog. Theor. Phys. 66 (1981) 355
- 1981TA16 S.L. Tabor, L.C. Dennis, K.W. Kemper, J.D. Fox, K. Abdo, G. Neuschaefer, D.G. Kovar and H. Ernst, Phys. Rev. C24 (1981) 960
- 1981TA1T Takeshita, J. Sci. Hiroshima Univ. A45 (1981) 257; Phys. Abs. 74755 (1982)
- 1981TA22 S.L. Tabor, L.C. Dennis and K. Abdo, Phys. Rev. C24 (1981) 2552
- 1981VI1B Vigdor, Santa Fe 1980, AIP Conf. proc. 69 (1981) 1429
- 1981WA1H M. Wakamatsu, Phys. Rev. C23 (1981) 2203
- 1981WA1J Wang, Zhang, Li and Ruan, Proc. Versailles Conf. (1981) 374

- 1981YA1A Yamazaki et al, Proc. Versailles Conf. (1981) 115
- 1981ZH1C Zhang, Li, Wang and Ruan, Proc. Versailles Conf. (1981) 369
- 1982AB07 N.K. Abrosimov, V.A. Volchenkov, V.A. Gordeev, V.A. Eliseev, E.M. Ivanov, V.P. Koptev, S.P. Kruglov, Yu.A. Malov, S.M. Mikirtychyants, G.A. Ryabov et al, Pisma Zh. Eksp. Teor. Fiz. 36 (1982) 211; JETP Lett. 36 (1982) 261
- 1982AB1D Abramovich, Gushovskii and Protopopov, in Kiev (1982) 566
- 1982AL1G Aleksandrov et al, in Kiev (1982) 358
- 1982AP1A Apokin et al, Yad. Fiz. 35 (1982) 382
- 1982AS01 E. Aslanides, A.M. Bergdolt, O. Bing, P. Fassnacht, F. Hibou, N. Willis, P. Kitching, Y. Le Bornec, B. Tatischeff, K. Baba et al, Phys. Lett. B108 (1982) 91
- 1982AU1B Auchev et al, in Kiev (1982) 307
- 1982AV06 R.O. Avakyan, A.E. Avetisyan, N.Z. Akopov, S.S. Danagulyan, I.Kh. Kosakov, A.A. Oganessian, Zh.V. Petrosyan, S.P. Taroyan and G.M. Elbakyan, Yad. Fiz. 35 (1982) 1089; Sov. J. Nucl. Phys. 35 (1982) 636
- 1982AW02 A.M. Awin and P.E. Shanley, Nucl. Phys. A386 (1982) 101
- 1982BA14 A.G. Baryshnikov, E.I. Dolinsky and S.P. Krekoten, Izv. Akad. Nauk SSSR Ser. Fiz. 46 (1982) 166
- 1982BA17 H.W. Barz and R. Wunsch, Czech. J. Phys. B32 (1982) 335
- 1982BA1P H. Bando and M. Bando, Phys. Lett. B109 (1982) 164
- 1982BA1T Baldwin, Bull. Amer. Phys. Soc. 27 (1982) 542
- 1982BA1X Barshnikov and Krekoten, in Kiev (1982) 411
- 1982BA2G B.F. Bayman, P.J. Ellis and Y.C. Tang, Phys. Rev. Lett. 49 (1982) 532
- 1982BA32 K. Baba, I. Endo, H. Fukuma, K. Inoue, T. Kawamoto, T. Ohsugi, Y. Sumi, T. Takeshita, S. Uehara, Y. Yano et al, Phys. Lett. B113 (1982) 459
- 1982BE54 Yu.A. Berezhnoi, A.P. Klyucharev and N.Ya. Rutkevich, Ukr. Fiz. Zh. 27 (1982) 1475; Phys. Abs. 11339 (1983)
- 1982BO1G Bogdanov, Kerimov and Safin, in Kiev (1982) 400
- 1982BO1J D.H. Boal, Phys. Rev. C25 (1982) 3068
- 1982BO1U Bouyssy, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 11
- 1982BO1Y Bogatin et al, Yad. Fiz. 36 (1982) 33
- 1982BO35 C. Borcea, E. Gierlik, A.M. Kalinin, R. Kalpakchieva, Yu.Ts. Oganessian, T. Pawlat, Yu.E. Penionzhkevich and A.V. Rykhlyuk, Nucl. Phys. A391 (1982) 520
- 1982BU1D Burtebaev, Duisebaev and Ivanov, in Kiev (1982) 327
- 1982CE1C Cecil, Bull. Amer. Phys. Soc. 27 (1982) 812

- 1982DA1R Daskaloyannis, Grypeos, Koutroulos and Saloupos, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 171
- 1982DE1H N. de Botton, Nucl. Phys. A374 (1982) 143
- 1982DE1J P.K.A. de Witt Huberts, Nucl. Phys. A374 (1982) 165
- 1982DE1P Deineko et al, in Kiev (1982) 320
- 1982DE35 A. Dellafore and E. Lipparini, Nucl. Phys. A388 (1982) 639
- 1982DEZL L.C. Dennis, A.D. Frawley, R.L. Parks and S.T. Thornton, Bull. Amer. Phys. Soc. 27 (1982) 478, BYa7
- 1982DEZQ M. Deutsch, J. Piekarz, R. Chrien, M. Le Vine, M. May, D. Maurizio, P. Pile, S. Bart, R. Hackenburg, E. Hungerford et al, Bull. Amer. Phys. Soc. 27 (1982) 721, DD9
- 1982DO1C C.B. Dover, Nucl. Phys. A374 (1982) 359
- 1982DO1M Dover and Walker, Phys. Rept. 89 (1982) 1
- 1982EL03 A.J. Elwyn, R.E. Holland, C.N. Davids and W.Ray, Jr., Phys. Rev. C25 (1982) 2168
- 1982EL07 R.J. Ellis, H.J. Ziock, K.O.H. Ziock, Y. Tzeng, J. Arvieux, R. Corfu, J. Piffaretti, L.C. Liu and E.R. Siciliano, Phys. Rev. C26 (1982) 1544
- 1982ER1B Eramzhyan et al, Czech. J. Phys. 32 (1982) 326
- 1982ER1E Eramzhyan, Fetisov, Majling and Zofka, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 91
- 1982FI03 B.W. Filippone, A.J. Elwyn, W. Ray, Jr. and D.D. Koetke, Phys. Rev. C25 (1982) 2174
- 1982FU11 M. Fujishiro, T. Tabata, K. Okamoto and T. Tsujimoto, Can. J. Phys. 60 (1982) 1672
- 1982GA1J Gal, Conf. on Hypernucl. Kaon Phys., Heidelberg, June, 1982 (1982) 27
- 1982GR1F S.J. Greene, W.J. Braithwaite, D.B. Holtkamp, W.B. Cottingham, C.F. Moore, G.R. Burleson, G.S. Blanpied, A.J. Viescas, G.H. Daw, C.L. Morris et al, Phys. Rev. C25 (1982) 927
- 1982GU1E Gurbanovich and Zelenskaia, in Kiev (1982) 390
- 1982GU1H H.H. Gutbrod, A.I. Warwick and H. Wieman, Nucl. Phys. A387 (1982) 177
- 1982HAZK L.F. Hansen, B.A. Pohl, C.H. Poppe, C. Wong and F.S. Dietrich, Bull. Amer. Phys. Soc. 27 (1982) 721, DE1
- 1982HI02 B. Hiller and H.J. Pirner, Phys. Lett. B109 (1982) 338
- 1982HU06 M. Hugi, J. Lang, R. Muller, J. Sromicki, E. Ungricht, K. Bodek, L. Jarczyk, B. Kamys, A. Strzalkowski and H. Witala, Phys. Rev. C25 (1982) 2403
- 1982IK1A Ikeda, Miyahara and Bando, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 149
- 1982JA1C Jacak et al, in MSU (1982) 32; Bull. Amer. Phys. Soc. 27 (1982) 715

- 1982JA1E Jarczyk, in Bad Honnef Symp. 1981, Springer-Verlag (1982) 37; Phys. Abs. 101932 (1982)
- 1982JO1C Johnstone and Thomas, J. Phys. G8 (1982) L105
- 1982KE1C Kerimov, Agalarov, Ishankuliev and Safin, in Kiev (1982) 397
- 1982KL1A Klein, Middleton and Tang, Nucl. Instrum. Meth. Phys. Res. 193 (1982) 601
- 1982KO11 N.N. Kolesnikov, D. Amarasingam and V.I. Tarasov, Yad. Fiz. 35 (1982) 32; Sov. J. Nucl. Phys. 35 (1982) 20
- 1982KU1G Ku et al, Nature 299 (1982) 240
- 1982LA11 M. Lattuada, F. Riggi, C. Spitaleri, D. Vinciguerra, C.M. Sutura and A. Pantaleo, Nuovo Cim. A69 (1982) 1
- 1982LA20 M. Lattuada, F. Riggi, C. Spitaleri, D. Vinciguerra, C.M. Sutura and A. Pantaleo, Nuovo Cim. A71 (1982) 429
- 1982LE10 M. LeMere, Y.C. Tang and H. Kanada, Phys. Rev. C25 (1982) 2902
- 1982LE1L Le Bornec and Willis, in "Pion Prod. and Absorp. in Nucl.-Indiana 1981", AIP Conf. Proc. 79 (1982) 155
- 1982LE1N R. Legrain, Nucl. Phys. A387 (1982) 219
- 1982LO13 M. Lozano, J.I. Escudero and G. Madurga, J. Phys. (London) G8 (1982) 1259
- 1982LU01 B. Ludewigt, G. Gaul, R. Glasow, H. Lohner and R. Santo, Phys. Lett. B108 (1982) 15
- 1982LY1A U. Lynen, H. Ho, W. Kuhn, D. Pelte, U. Winkler, W. F.J. Muller, Y.-T. Chu, P. Doll, A. Gobbi, K. Hildenbrand et al, Nucl. Phys. A387 (1982) 129
- 1982MAZA J.F. Mateja, R.L. Kozub, J. Garman, A.D. Frawley, L.C. Dennis and K.W. Kemper, Bull. Amer. Phys. Soc. 27 (1982) 478, BYa3
- 1982MO1N J. Mougey, Nucl. Phys. A387 (1982) 109
- 1982MO1Q Motoba, Ikeda and bando, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 129
- 1982MU1C Murphy et al, Bull. Amer. Phys. Soc. 27 (1982) 32
- 1982NA1N Nadasen et al, Bull. Amer. Phys. Soc. 27 (1982) 730
- 1982NG01 Nguyen Tien Nguyen and I. Ulehla, Czech. J. Phys. B32 (1982) 1040
- 1982NO1C Novikov, in Kiev (1982) p. 279
- 1982OR03 M.C.L. Orlowski, Bao Cheng-Guang and Liu Yuen, Z. Phys. A305 (1982) 249
- 1982PE03 R.J. Peterson, Nucl. Phys. A377 (1982) 41
- 1982PE1F Pella et al, Bull. Amer. Phys. Soc. 27 (1982) 729
- 1982PI1J Piekarczyk, in Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 72

- 1982PO1B Pomerantsev, Kukulín and Klimov, in Kiev (1982) p. 406
- 1982PO1C Povh, Czech. J. Phys. 32 (1982) 183
- 1982SH1P Shi Yi-Jin and Zhuang Fei, Phys. Energ. Fortis Phys. Nucl. 6 (1982) 750; Phys. Abs. 40985 (1983)
- 1982SU09 J.-I. Suzumura and Y. Futami, Prog. Theor. Phys. 68 (1982) 2060
- 1982TE01 P.K. Teng, A. Chaudhury, J.J. LeRose, K. Min, D. Rowley, B.O. Sapp, P. Stoler, E.J. Winhold, P.F. Yergin, A.M. Bernstein et al, Phys. Rev. C26 (1982) 1313
- 1982WAZS Z.-J. Wang and P.A. Quin, Bull. Amer. Phys. Soc. 27 (1982) 700, AE6
- 1982WU1B Wu et al, Chin. Phys. 2 (1982) 726
- 1982ZH1F Zhang, Li, Wang and Ruan, Phys. Energ. Fortis Phys. Nucl. 6 (1982) 64
- 1983AB1A Abramovitch et al, in Moscow (1983) p. 362
- 1983ANZY R.E. Anderson, C.R. Gould, B.C. Karp, K. E.Nash, R.S. Pedroni, H.G. Pfutzner and R.L. Walter, Bull. Amer. Phys. Soc. 28 (1983) 648, AF1
- 1983AS02 E. Aslanides, P. Baumann, G. Bergdolt, P. Engelstein, P. Fassnacht, F. Hibou, E. Chiavassa, G. Dellacasa, M. Gallio, A. Musso et al, Nucl. Phys. A393 (1983) 314
- 1983BA2K Bando, Ikeda and Motoba, Prog. Theor. Phys. 69 (1983) 918
- 1983BA52 F.C. Barker, Can. J. Phys. 61 (1983) 1371
- 1983BE1P Belyaeva, Zelenskaia and Teplov, in Moscow (1983) p. 477
- 1983BO1E Bogdanov and Safin, in Moscow (1983) p. 185
- 1983BY01 R.C. Byrd, C.E. Floyd, P.P. Guss, K. Murphy and R.L. Walter, Nucl. Phys. A399 (1983) 94
- 1983CO11 F.D. Correll, L. Madansky, R.A. Hardekopf and J.W. Sunier, Phys. Rev. C28 (1983) 862
- 1983CO1Q Correll et al, (1983)
- 1983DE1U Dennis, Phys. Rev. C27 (1983) 2641; Erratum Phys. Rev. C28 (1983) 2547
- 1983EC01 J.S. Eck, A.R. Omar, J.R. Leigh and T.R. Ophel, Phys. Rev. C27 (1983) 1807
- 1983FI01 B.W. Filippone, A.J. Elwyn, C.N. Davids and D.D. Koetke, Phys. Rev. Lett. 50 (1983) 412
- 1983FU04 T. Fukuda, M. Ishihara, M. Tanaka, H. Ogata, I. Miura, M. Inoue, T. Shimoda, K. Katori and S. Nakayama, Phys. Rev. C27 (1983) 2029
- 1983FUZY C.B. Fulmer, K.A. Erb, D.C. Hensley, R.L. Auble, J.B. Ball, F.E. Bertrand, E.E. Gross and G.R. Satchler, Bull. Amer. Phys. Soc. 28 (1983) 716, GF9
- 1983GO1R Gontcharova, Kissener and Eramzhian, in Moscow (1983) p. 188
- 1983GU1G Guliamov, Kim, Bekbaev and Islomov, in Moscow (1983) p. 428

- 1983HO01 S. Homma, M. Kanazawa, K. Maruyama, Y. Murata, H. Okuno, A. Sasaki and T. Taniguchi, Phys. Rev. C28 (1983) 31
- 1983HU02 M.G. Huber, K. Klingenberg and R. Hupke, Nucl. Phys. A396 (1983) 191c
- 1983JO1E J.A. Johnstone and A.W. Thomas, Nucl. Phys. A392 (1983) 409
- 1983KO04 H.-G. Korber, R. Beckmann and U. Holm, Nucl. Phys. A394 (1983) 257
- 1983LI1J Likhachev et al, in Moscow (1983) 339
- 1983LI1K Likhachev et al, in Moscow (1983) 338
- 1983LO1G Lourie et al, Unknown Source (1983)
- 1983LOZZ R.W. Lourie, W. Bertozzi, T.N. Buti, J.M. Finn, F.W. Hersman, C. Hyde, J. Kelly, M.A. Kovash, S. Kowalski, M.V. Hynes et al, Bull. Amer. Phys. Soc. 28 (1983) 673, BH4
- 1983OC01 J.S. O'Connell, E. Hayward, J.W. Lightbody, Jr., X.K. Maruyama, P. Bosted, K.I. Blomqvist, G. Franklin, J-O. Adler, K. Hansen and B. Schroder, Phys. Rev. C27 (1983) 2492
- 1983OMZZ A.R. Omar, J.S. Eck, T.R. Ophel and J.R. Leigh, Bull. Amer. Phys. Soc. 28 (1983) 715, GF1
- 1983SA06 M. Sato, M. Sasagase, Y. Nagashima, J. Schimizu, T. Nakagawa, Y. Fukuchi and T. Mikumo, Phys. Rev. C27 (1983) 2621
- 1983ST1G Stibunov, in Moscow (1983) 352
- 1983ZH1D Zhivopistev and Slivoi, in Moscow (1983) 354
- 1985AJ01 F. Ajzenberg-Selove, Nucl. Phys. A433 (1985) 1; Erratum Nucl. Phys. A449 (1986) 155