

Energy Levels of Light Nuclei $A = 10$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 5-10$ was published in *Nuclear Physics A*413 (1984), p. 1. This version of $A = 10$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and Introductory tables have been omitted from this manuscript. Also, [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed June 1, 1983)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 10$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{10}\text{n}\$](#) , [\$^{10}\text{He}\$](#) , [\$^{10}\text{Li}\$](#) , [\$^{10}\text{Be}\$](#) , [\$^{10}\text{B}\$](#) , [\$^{10}\text{C}\$](#) , [\$^{10}\text{N}\$](#) , [\$^{10}\text{O}\$](#) , [\$^{10}\text{F}\$](#) , [\$^{10}\text{Ne}\$](#)

B. Tables of Recommended Level Energies:

[Table 10.1](#): Energy levels of ^{10}Be

[Table 10.5](#): Energy levels of ^{10}B

[Table 10.19](#): Energy levels of ^{10}C

C. [References](#)

D. Figures: [\$^{10}\text{Be}\$](#) , [\$^{10}\text{B}\$](#) , [\$^{10}\text{C}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

¹⁰n
(Not illustrated)

¹⁰n has not been observed in the interaction of 0.7 and 400 GeV protons with uranium: the cross section is $< 0.7 \times 10^{-5} \mu\text{b}$ (1977TU02) at 0.7 GeV and $< 0.5 \mu\text{b}$ (1977TU1B) at 400 GeV.

¹⁰He
(Not illustrated)

¹⁰He has not been observed in the bombardment of ²³²Th by 4.8 GeV/c deuterons [the production cross section is $\lesssim 2 \mu\text{b}$] (1979BE60) nor in the interaction of ¹⁰B, ¹¹B and ²²Ne [8 to 10 MeV/A] with Ti and ²³²Th targets [the cross section is $< 5 \times 10^{-4} \mu\text{b/sr}$] (1982OG02). See also (1974AJ01). The calculated value of the atomic mass excess of ¹⁰He is 49.40 MeV: ¹⁰He is then unstable with respect to breakup into ⁹He + n and ⁸He + 2n by 0.52 and 1.66 MeV, respectively (1980SE1F). See also (1979AJ01), (1980BO31, 1982BI1C) and (1979BO22, 1981AV02, 1981KI04, 1982AV1A, 1982NG01, 1982VE05; theor.).

¹⁰Li
(Fig. 22)

At $E(^9\text{Be}) = 121 \text{ MeV}$, ¹⁰Li has been observed in the ⁹Be(⁹Be, ⁸B)¹⁰Li reaction with a differential cross section (c.m.) of $\approx 30 \text{ nb/sr}$ at $\theta = 14^\circ$ (lab): $Q_0 = -34.06 \pm 0.25 \text{ MeV}$, and the atomic mass excess of ¹⁰Li is $33.83 \pm 0.25 \text{ MeV}$ if the group observed ($\Gamma \approx 1.2 \pm 0.3 \text{ MeV}$) corresponds to the ground state. ¹⁰Li_{g.s.} would be unbound with respect to breakup into ⁹Li + n by $0.80 \pm 0.25 \text{ MeV}$ (1975WI26). However (1979AB11, 1980AB16), on the basis of the possible location of the first $T = 2$ state in ¹⁰Be (see reaction 2) with $J^\pi = 2^-$, suggest that (1975WI26) have observed an excited state of ¹⁰Li, with $J^\pi = 1^+$. See also (1979AJ01) and (1979BO22, 1980MA1Z, 1981AV02, 1981WA1J, 1982NG01; theor.).

¹⁰Be
(Figs. 19 and 22)

GENERAL: (See also (1979AJ01).)

Model calculations: (1980FU1G, 1981DE2G).

Special states: (1981DE2G, 1981SE06).

Electromagnetic transitions: (1982LA26, 1982RI04).

Astrophysical questions: (1978BU1B, 1979MO04, 1979SI1D, 1980WI1M, 1981GA1B, 1981GA1C, 1981KR1C, 1981PR1B, 1981SA1G, 1982BR1N, 1982KU1J, 1982PA1F).

Applied work: (1978MU1B, 1978RA1C, 1979GO1L, 1979IN1C, 1979LI1C, 1979RA1E, 1980EL1B, 1980EL1C, 1980GO1B, 1980KI1B, 1980LA1B, 1981FA1E, 1981LI1K, 1981RA1F, 1981SA1G, 1981SC1D, 1981TH1C, 1982BO35, 1982BR1N, 1982KL1A, 1982KU1G, 1982KU1J, 1982PA1F, 1983KR1B, 1983SO1B, 1983SU1C).

Complex reactions involving ¹⁰Be: (1978DU1B, 1978HE1C, 1978TU06, 1979AL22, 1979BO22, 1979JA1C, 1979SC1D, 1979VI05, 1980GR10, 1980WI1L, 1981CI03, 1981MO20, 1982BI1C, 1982BO1J, 1982GU1H, 1982LU01, 1982LY1A, 1983SA06).

Muon and neutrino capture and reactions: (1979GO1M, 1980MU1B, 1981GI08).

Reactions involving pions and kaons (See also reactions 10, 16 and 18): (1978DA1A, 1978WA1B, 1979PE1C, 1980BO1B, 1981AU1C, 1981FE2A, 1981SI09, 1981WH01, 1981YA1A, 1982AU02, 1982RO04, 1982WA1G).

Hypernuclei: (1978DA1A, 1978PO1A, 1978SO1A, 1980RO1F, 1980ZH1C, 1981WA1J, 1982IK1A, 1982KO1L, 1982KO1I, 1982RA1L).

Other topics: (1981SE06, 1982DE1N, 1982NG01).

Ground-state properties of ¹⁰Be: (1978SM02, 1981AV02, 1982NG01).

1. $^{10}\text{Be}(\beta^-)^{10}\text{B}$ $Q_m = 0.5568$

The half-life of ¹⁰Be is $(1.6 \pm 0.2) \times 10^6$ y; $\log ft = 13.42$: see (1974AJ01). See also (1979FE1E; theor.) and (1979AJ01).

| | | |
|---|-----------------|-----------------|
| 2. (a) $^7\text{Li}(t, \gamma)^{10}\text{Be}$ | $Q_m = 17.2498$ | |
| (b) $^7\text{Li}(t, n)^9\text{Be}$ | $Q_m = 10.4377$ | $E_b = 17.2498$ |
| (c) $^7\text{Li}(t, p)^9\text{Li}$ | $Q_m = -2.386$ | |
| (d) $^7\text{Li}(t, d)^8\text{Li}$ | $Q_m = -4.225$ | |
| (e) $^7\text{Li}(t, t)^7\text{Li}$ | | |
| (f) $^7\text{Li}(t, \alpha)^6\text{He}$ | $Q_m = 9.839$ | |

Table 10.1: Energy levels of ^{10}Be ^a

| E_x (MeV \pm keV) | $J^\pi; T$ | τ or $\Gamma_{c.m.}$ (keV) | Decay | Reactions |
|-----------------------|-------------|--|----------------|---|
| g.s. | $0^+; 1$ | $\tau_{1/2} = (1.6 \pm 0.2) \times 10^6$ y | β^- | 1, 2, 3, 4, 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 |
| 3.3680 ± 0.2 | $2^+; 1$ | $\tau_m = 180 \pm 17$ fsec | γ | 2, 3, 4, 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29 |
| 5.9583 ± 0.3 | $2^+; 1$ | $\tau_m < 80$ fsec | γ | 5, 10, 11, 13, 16, 17, 18, 20, 21, 25, 29 |
| 5.9599 ± 0.6 | $1^-; 1$ | | γ | 5, 10, 11, 13, 21, 25, 29 |
| 6.1793 ± 0.7 | $0^+; 1$ | $\tau_m = 1.1^{+0.4}_{-0.3}$ psec | π, γ | 5, 10, 11, 21 |
| 6.2633 ± 5 | $2^-; 1$ | | γ | 5, 11, 13 |
| 7.371 ± 1 | $3^-; 1$ | $\Gamma = 15.7 \pm 0.5$ keV | n | 4, 6, 10, 11, 13 |
| 7.542 ± 1 | $2^+; 1$ | 6.3 ± 0.8 | n | 3, 4, 6, 11, 13, 29 |
| 9.27 | $(4^-); 1$ | 150 ± 20 | n | 4, 6, 10, 11, 13 |
| 9.4 | $(2^+); 1$ | 291 ± 20 | n | 4, 6, 10, 11, 13, 20, 25, 29 |
| 10.57 ± 30 | $\geq 1; 1$ | | n | 3, 4, 6, 11 |
| 11.76 ± 20 | | 121 ± 10 | | 3, 4, 10, 11, 13, 29 |
| 17.79 | | 110 ± 35 | γ, n, t | 2, 3, 4 |
| 18.55 | | ≈ 350 | n, t | 2, 3, 4 |
| (21.22) | $(2^-; 2)$ | sharp | n, p, t | 2 |
| (24) | | | | 24 |

^a See also Table 10.4.

The yield of γ_0 and γ_1 has been studied for $E_t = 0.4$ to 1.1 MeV (1978SU02) [$^{10}\text{Be}^*(17.79)$ is said to be involved]. See also (1974AJ01). The neutron yield exhibits a weak structure at $E_t = 0.24$ MeV and broad resonances at $E_t \approx 0.77$ MeV [$\Gamma = 160 \pm 50$ keV] and 1.74 MeV: see (1966LA04) [$^{10}\text{Be}^*(17.79, 18.47)$]. The total cross section for reaction (c), the yield of neutrons (reaction (b) to $^9\text{Be}^*(14.39)$), and the yield of γ -rays from $^7\text{Li}^*(0.48)$ (reaction (e)) all show a sharp anomaly at $E_t = 5.685$ MeV: $J^\pi = 2^-$; $T = 2$ is suggested for a state at $E_x = 21.22$ MeV. The total cross section for α_0 (reaction (e)) and the all-neutrons yield do not show this structure (1979AB11, 1980AB16, 1982AB1D, 1983AB1A). Differential cross sections and S -factors are reported by (1983CE01) for $E_t = 70$ to 110 keV for $^6\text{He}^*(0, 1.80)$. The zero-energy S -factor for $^6\text{He}^*(1.80)$ is 14 ± 2.5 MeV \cdot b. The relevance to an Li-seeded tritium plasma is discussed by (1983CE01). See also (1974AJ01, 1979AJ01).

$$3. \text{}^7\text{Li}(\alpha, p)^{10}\text{Be} \quad Q_m = -2.5642$$

Angular distributions have been measured at $E_\alpha = 30$ and 50 MeV: see (1979AJ01).

$$4. \text{}^7\text{Li}(\text{}^7\text{Li}, \alpha)^{10}\text{Be} \quad Q_m = 14.782$$

Angular distributions have been reported for $E(^7\text{Li}) = 2.1$ to 5.75 MeV and at 30.3 MeV: see (1974AJ01).

$$5. \text{}^9\text{Be}(n, \gamma)^{10}\text{Be} \quad Q_m = 6.8121$$

$$Q_0 = 6812.1 \pm 0.2 \text{ keV: see (1980IS02)}$$

$$Q_0 = 6912.08 \pm 0.10 \text{ keV: see (1981KE02).}$$

The thermal capture cross section is 7.6 ± 0.8 mb (1981MUZQ). Reported γ -ray transitions are displayed in Table 10.2.

$$6. \text{}^9\text{Be}(n, n)^9\text{Be} \quad E_b = 6.8121$$

The scattering amplitude (bound) $a = 7.778 \pm 0.003$ fm, $\sigma_{\text{free}} = 6.151 \pm 0.005$ b (1981MUZQ). Earlier cross-section measurements are listed in (1979AJ01, 1981MUZQ). Recent σ_t measurements have been reported for $E_n = 0.002$ to 0.005 eV (1979AD1B), 23.5 keV (1975BL07; 5.903 ± 0.011 b), 1.0 to 13.5 MeV (1979AU07), 6.97 to 14.94 MeV (1978HO23; total elastic) and 200 to 590 MeV (1980FR1K). See also (1980AD1A).

Observed resonances are displayed in Table 10.3. Analysis of polarization and differential cross-section data leads to the 3^- , 2^+ assignments for $^{10}\text{Be}^*(7.37, 7.55)$. Below $E_n = 0.5$ MeV the scattering cross

Table 10.2: Neutron-capture γ -rays in ^{10}Be ^a

| E_γ ^b (keV) | Transition | Intensities ^c | E_x (keV) ^b |
|-------------------------------|---|--------------------------|--------------------------|
| 6809.4 ± 0.4 | capt. \rightarrow g.s. | 64.6 | |
| 5955.9 ± 0.5 | 5.96 ^d \rightarrow g.s. | 1.5 | 5958.3 ± 0.3 |
| 3443.3 ± 0.3 | capt. \rightarrow 3.37 | 11.3 | |
| 3367.4 ± 0.2 | 3.37 \rightarrow g.s. | 33.0 | 3368.0 ± 0.2 |
| 2896.4 ^e | 6.26 \rightarrow 3.37 | 0.15 | |
| 2811.8 ^e | 6.18 \rightarrow 3.37 | 0.13 | |
| 2589.9 ± 0.25 | 5.96 ^d \rightarrow 3.37 | 22.4 | |
| 853.5 ± 0.3 | capt. \rightarrow 5.96 ^d | 26.4 | |
| 631.8 ^e | capt. \rightarrow 6.18 | 0.24 | |
| 547.4 ^e | capt. \rightarrow 6.26 | 0.16 | |
| 219.3 ^e | 6.18 \rightarrow 5.96 ^f | 0.05 | |

^a See also Tables 10.2 in (1974AJ01, 1979AJ01).

^b (1966GR18).

^c Gamma rays per 100 captures (1979JU1B).

^d This is the 2^+ member of the doublet at $E_x = 5.96$ MeV.

^e (1979JU1B).

^f This is the 1^- member of the doublet.

section reflects the effect of bound 1^- and 2^- states, presumably $^{10}\text{Be}^*(5.960, 6.26)$. There is also indication of interference with s -wave background and with a broad $l = 1$, $J^\pi = 3^+$ state. The structure at $E_n = 2.73$ MeV is ascribed to two levels: a broad state at about 2.85 MeV with $J^\pi = 2^+$, and a narrow one, $\Gamma \approx 100$ keV, at $E_n = 2.73$ MeV with a probable assignment of $J^\pi = 4^-$. The 4^- assignment results from a study of the polarization of the n_0 group at $E_n = 2.60$ to 2.77 MeV. A rapid variation of the polarization over this interval is observed, and the data are consistent with $4^-(l = 2)$ for $^{10}\text{Be}^*(9.27)$. A weak dip at $E_n \approx 4.3$ MeV is ascribed to a level with $J \geq 1$. See (1974AJ01) for references. The analyzing power has been measured for $E_n = 1.6$ to 4.8 MeV (1975HO01) and $E_n = 9$ to 15 MeV (1981FL1A, 1981FL04; n_0, n_2). See also (1981WA1G, 1983ANZY) and (1979BY01, 1983BY01; theor.).

7. (a) $^9\text{Be}(n, n')^9\text{Be}^*$

$$E_b = 6.8121$$

(b) $^9\text{Be}(n, 2n)^8\text{Be}$

$$Q_m = -1.6655$$

The non-elastic and the $(n, 2n)$ cross sections rise rapidly to ≈ 0.6 b (≈ 0.5 b for $(n, 2n)$) at $E_n \approx 3.5$ MeV and then stay approximately constant to $E_n = 15$ MeV: see (1979AJ01). Recent measurements include those of (1978HO23; $E_n = 6.97$ to 14.94 MeV; inelastic cross sections to unresolved states at $E_x = 1.7$ –3.1 MeV). A_y measurements are reported for the n_2 group [to $^9\text{Be}^*(2.4)$] at $E_n = 9$ to 15 MeV (1981FL1A). For reaction (b) see also (1980BA2L; applied work) and (1978HE1F; theor.).

Table 10.3: Resonances in ${}^9\text{Be}(n, n){}^9\text{Be}$ ^a

| E_{res} (MeV \pm keV) | ${}^{10}\text{Be}^*$ (MeV) | $\Gamma_{\text{c.m.}}$ (keV) | J^π | l | θ^2 (%) ^b |
|----------------------------------|----------------------------|------------------------------|----------|-----|-----------------------------|
| 0.6220 ± 0.8 | 7.371 | 15.7 ± 0.5 | 3^- | 2 | 7.5 |
| 0.8118 ± 0.7 | 7.542 | 6.3 ± 0.8 | 2^+ | 1 | 0.28 |
| 2.73 | 9.27 | ≈ 100 | (4^-) | (2) | |
| (2.85) | 9.4 | ≈ 400 | (2^+) | (1) | |
| 4.3 | 10.7 | | ≥ 1 | | |

^a For references see Table 10.3 in (1979AJ01).

^b $R = 5.6$ fm.

8. (a) ${}^9\text{Be}(n, p){}^9\text{Li}$ $Q_m = -12.824$ $E_b = 6.8121$
 (b) ${}^9\text{Be}(n, d){}^8\text{Li}$ $Q_m = -14.662$
 (c) ${}^9\text{Be}(n, t){}^7\text{Li}$ $Q_m = -10.438$

Cross sections have been measured at $E_n = 14.1$ – 14.9 MeV for reaction (a), 16.3 to 18.7 MeV for (b) and 13.3 to 15.0 (t_1) and 22.5 MeV (reaction (c)): see (1979AJ01). See also (1981HAZJ, 1982HA1A).

9. ${}^9\text{Be}(n, \alpha){}^6\text{He}$ $Q_m = -0.598$

The cross section for production of ${}^6\text{He}$ shows a smooth rise to a broad maximum of 104 ± 7 mb at 3.0 MeV, followed by a gradual decrease to 70 mb at 4.4 MeV. From $E_n = 3.9$ to 8.6 MeV, the cross section decreases smoothly from 100 mb to 32 mb. Excitation functions have been measured for α_0 and α_1 for $E_n = 12.2$ to 18.0 MeV: see (1979AJ01) for references. See also (1981HAZJ, 1982HA1A).

10. ${}^9\text{Be}(p, \pi^+){}^{10}\text{Be}$ $Q_m = -133.538$

Angular distributions have been studied at $E_p = 185$ MeV (1973DA09: to ${}^{10}\text{Be}^*(0, 3.37, 6.07 \pm 0.13, 7.39 \pm 0.13, 9.31 \pm 0.24, 11.76)$), 200 MeV (1978AU07: to ${}^{10}\text{Be}^*(0, 3.37)$), 410 and 605 MeV (1980DI02: to ${}^{10}\text{Be}^*(0, 3.37, 6.1)$) and 800 MeV (1979HO13: to ${}^{10}\text{Be}^*(0, 3.37, 11.8)$); the latter may be unresolved). (1982AU02) have studied $A_y(\theta)$ for the π^+ to ${}^{10}\text{Be}^*(0, 3.37)$ for $E_{\vec{p}} = 200$ and 250 MeV: little variation with E is observed. The cross section for π^+ production near threshold has been studied by (1979MA38). See also the ‘‘General’’ section here, and (1981NI1B, 1982HO1C, 1982NA1K).

11. ${}^9\text{Be}(d, p){}^{10}\text{Be}$ $Q_m = 4.5875$

Table 10.4: Radiative transitions in ${}^9\text{Be}(d, p){}^{10}\text{Be}$ ^a

| E_x (keV) | Transition | ΔJ^π | Mult. | Branch (%) | τ_m (psec) | Γ_γ (meV) |
|------------------|--------------------------------|-----------------------|-------|-----------------|---------------------|-----------------------|
| 3368.0 ± 0.2 | $3.37 \rightarrow \text{g.s.}$ | $2^+ \rightarrow 0^+$ | E2 | 100 | 0.189 ± 0.020 | 3.48 ± 0.37 |
| | | | | | 0.160 ± 0.030 | 4.11 ± 0.78 |
| 5958.3 ± 0.3 | $5.96 \rightarrow 3.37$ | $2^+ \rightarrow 2^+$ | M1 | > 90 | < 0.08 | |
| | $5.96 \rightarrow \text{g.s.}$ | $2^+ \rightarrow 0^+$ | E2 | < 10 | | |
| 5959.9 ± 0.6 | $5.96 \rightarrow \text{g.s.}$ | $1^- \rightarrow 0^+$ | E1 | 83_{-6}^{+10} | $1.1_{-0.3}^{+0.4}$ | 0.14 ± 0.05 |
| | $5.96 \rightarrow 3.37$ | $1^- \rightarrow 2^+$ | E1 | 17_{-10}^{+6} | | |
| 6179.3 ± 0.7 | $6.18 \rightarrow 5.96$ | $0^+ \rightarrow 1^-$ | E1 | 24 ± 2 | $1.1_{-0.3}^{+0.4}$ | 0.46 ± 0.28 |
| | $6.18 \rightarrow 3.37$ | $0^+ \rightarrow 2^+$ | E2 | 76 ± 2 | | |
| | $6.18 \rightarrow \text{g.s.}$ | $0^+ \rightarrow 0^+$ | E0 | ^b | | |
| 6263.3 ± 5 | $6.26 \rightarrow 5.96$ | $2^- \rightarrow 1^-$ | M1 | ≤ 1 | | |
| | | $2^- \rightarrow 2^+$ | E1 | | | |
| | $6.26 \rightarrow 3.37$ | $2^- \rightarrow 2^+$ | E1 | 99_{-2}^{+1} | | |
| | $6.26 \rightarrow \text{g.s.}$ | $2^- \rightarrow 0^+$ | M2 | 1 ± 1 | | |

^a See Table 10.4 in (1979AJ01) for references. However, note that there are several typographical errors in the ${}^{10}\text{Be}^*(6.18)$ decay.

^b (1975WA06).

Angular distributions of proton groups have been studied at many energies in the range $E_d = 0.15$ to 17.13 MeV [see (1979AJ01)], at $E_d = 2.0$ to 2.8 MeV (1983DE1P; p_0, p_1) and at 698 MeV (1981BO03; p to ${}^{10}\text{Be}^*(0, 3.37, 6.1, 7.37 + 7.54, 9.27 + 9.4, 11.8)$). At $E_d = 15$ MeV (1976DA15) find $S = 2.1, 0.23$ ($j_n = \frac{3}{2}$) and 0.12 ($j_n = \frac{1}{2}$), $\leq 1.0, 0.065$ ($j_n = \frac{5}{2}$) and 0.132 ($j_n = \frac{1}{2}$), for ${}^{10}\text{Be}^*(0, 3.37, 5.96, 6.62)$ (1976DA15). The angular distributions show $l_n = 1$ transfer for ${}^{10}\text{Be}^*(0, 3.37, 5.958, 7.54)$, $l_n = 0$ transfer for ${}^{10}\text{Be}^*(5.960, 6.26)$, $l_n = 2$ transfer for ${}^{10}\text{Be}^*(7.37)$. ${}^{10}\text{Be}^*(6.18, 9.27, 9.4)$ are also populated, as are two states at $E_x = 10.57 \pm 0.03$ and 11.76 ± 0.02 MeV (1974AN27). ${}^{10}\text{Be}^*(9.27, 9.4, 11.76)$ have $\Gamma_{\text{c.m.}} = 150 \pm 20, 291 \pm 20$ and 121 ± 10 keV (1974AN27).

Attempts to understand the γ -decay of ${}^{10}\text{Be}^*(5.96)$ and its population in ${}^9\text{Be}(n, \gamma){}^{10}\text{Be}$ led to the discovery that it consisted of two states separated by 1.6 ± 0.5 keV. The lower of the two has $J^\pi = 2^+$ and decays primarily by a cascade transition via ${}^{10}\text{Be}^*(3.37)$ [it is the state fed directly in the ${}^9\text{Be}(n, \gamma)$ decay]; the higher state has $J^\pi = 1^-$ and goes mainly by a crossover to ${}^{10}\text{Be}_{\text{g.s.}}$. Angular correlations measured with their γ -ray detector located normal to the reaction plane (\equiv angular distributions) lead to l_n values consistent with the assignments of 2^+ and 1^- for ${}^{10}\text{Be}^*(5.9658, 5.9660)$ obtained from the character of the γ -decay (1969RO12). ${}^{10}\text{Be}^*(6.18)$ decays primarily to ${}^{10}\text{Be}^*(3.37)$: $E_\gamma = 219.4 \pm 0.3$ keV for the $6.18 \rightarrow 5.96$ transition (1969AL17). See Table 10.4 for a listing of the information on radiative transitions obtained in this reaction and lifetime measurements. For (p, γ) correlations through ${}^{10}\text{Be}^*(3.37)$ see (1974AJ01). For polarization measurements see ${}^{11}\text{B}$ in (1980AJ01). See also (1981CE04), (1974FI1D) and (1982GO05; theor.).

12. ${}^9\text{Be}({}^3\text{He}, 2p){}^{10}\text{Be}$ $Q_m = -0.9061$

See (1980CO12; $E({}^3\text{He}) = 13$ MeV) and ${}^{12}\text{C}$ in (1985AJ01).

13. ${}^9\text{Be}(\alpha, {}^3\text{He}){}^{10}\text{Be}$ $Q_m = -13.7657$

Angular distributions have been studied at $E_\alpha = 65$ MeV to ${}^{10}\text{Be}^*(0, 3.37, 5.96, 6.26, 7.37, 7.54, 9.33, 11.88)$. DWBA analyses of these lead to spectroscopic factors which are in poor agreement with those reported in other reactions (1980HA33).

14. ${}^9\text{Be}({}^7\text{Li}, {}^6\text{Li}){}^9\text{Be}$ $Q_m = -0.438$

Angular distributions have been measured at $E({}^7\text{Li}) = 34$ MeV to ${}^{10}\text{Be}^*(0, 3.4)$: $S = 2.07$ and $0.42(p_{1/2}), 0.38(p_{3/2})$ (1977KE09).

15. (a) ${}^{10}\text{Be}(p, p){}^{10}\text{Be}$

(b) ${}^{10}\text{Be}(d, d){}^{10}\text{Be}$

Angular distributions of the p_0 and p_1 groups have been measured at $E_p = 12.0$ to 16.0 MeV. The elastically scattered deuterons have been studied at $E_d = 12.0$ and 15.0 MeV: see (1974AJ01).

16. (a) ${}^{10}\text{Be}(\gamma, \pi^+){}^{10}\text{Be}$ $Q_m = -140.124$

(b) ${}^{10}\text{B}(e, e'\pi^+){}^{10}\text{Be}$ $Q_m = -140.124$

Differential cross sections have been measured for reaction (a) to ${}^{10}\text{Be}^*(0, 3.37)$ for $E_\gamma \approx 230$ to 340 MeV (1980BO24). Measurements are also reported at $E_{\pi^+} = 17, 29$ and 48 MeV to ${}^{10}\text{Be}_{\text{g.s.}}$ and at the two higher energies for the transitions to ${}^{10}\text{Be}^*(3.37)$ (1982RO04). See also (1981YA1A; $E_\gamma \approx 183$ MeV; g.s.). At $E_e = 158.5$ to 165.0 MeV the angular distributions of 12.3 ± 0.7 MeV pions are reported by (1982ZU03). The (e, π^+) angular distributions to ${}^{10}\text{Be}^*(0, 3.37)$ are in good agreement with the M3 form factors. It is suggested that ${}^{10}\text{Be}^*(5.96)$ is also populated through an M1 transition. (1982ZU03) also present a review of the data for the 90° photopion cross section leading to ${}^{10}\text{Be}_{\text{g.s.}}$ as a function of E_{π^+} . See also (1982DE14; theor.).

17. ${}^{10}\text{B}(\mu^-, \nu){}^{10}\text{Be}$ $Q_m = 105.1026$

Partial capture rates leading to the 2^+ states $^{10}\text{Be}^*(3.37, 5.96)$ are reported by (1981GI08).

18. $^{10}\text{B}(\pi^-, \gamma)^{10}\text{Be}$ $Q_m = 139.011$

The total radiative capture branching ratio for stopped pions is $(2.27 \pm 0.22)\%$. The γ -spectrum is dominated by the transition to $^{10}\text{Be}^*(6.0) [J^\pi = 2^+]$ (1975BA52). See also (1979AL1M) and (1979TR1B).

19. $^{10}\text{B}(p, p\pi^+)^{10}\text{Be}$ $Q_m = -140.124$

See (1979AJ01).

20. $^{10}\text{B}(d, 2p)^{10}\text{Be}$ $Q_m = -3.5637$

Angular distributions are reported at $E_d = 55$ MeV to $^{10}\text{Be}^*(0, 3.37, 5.96, 9.4)$ (1979ST15).

21. $^{11}\text{Li}(\beta^-)^{11}\text{Be} \rightarrow ^{10}\text{Be} + n$

^{11}Li populates several states of ^{10}Be , via delayed neutron emission. Gamma rays have been observed for the transitions $6.18 \rightarrow 5.96$, $6.18 \rightarrow 3.37$, 5.96 (unres.) $\rightarrow 3.37$ and $3.37 \rightarrow$ g.s. with $I_\gamma = (0.95 \pm 0.35)$, (1.65 ± 0.70) , (3.5 ± 1.0) and $(21 \pm 6)\%$, respectively (1980DE39). See also ^{11}Be in (1985AJ01).

22. $^{11}\text{B}(\gamma, p)^{10}\text{Be}$ $Q_m = 11.2287$

Transitions to $^{10}\text{Be}^*(3.4)$ from the upper region of the giant resonance in ^{11}B are reported to be about twice as intense as those from the lower region: see ^{11}B in (1975AJ02). Angular distributions of the p_0 and p_1 groups have been measured with $E_{\text{bs}} = 18.5$ MeV and the yield of the 3.37 MeV γ -ray has been reported for $E_{\text{bs}} = 100$ to 800 MeV: see (1979AJ01). See also (1982GO03; theor.).

23. $^{11}\text{B}(n, d)^{10}\text{Be}$ $Q_m = -9.0042$

The angular distribution of the d_0 group has been measured at $E_n = 14.4$ MeV: see (1974AJ01).

24. $^{11}\text{B}(p, 2p)^{10}\text{Be}$ $Q_m = -11.2287$

Structure is observed in the summed proton spectrum corresponding to $Q = -10.9 \pm 0.35, -14.7 \pm 0.4, -21.1 \pm 0.4, -35 \pm 1$ MeV: see (1974AJ01).

25. $^{11}\text{B}(\text{d}, ^3\text{He})^{10}\text{Be}$ $Q_{\text{m}} = -5.7352$

Angular distributions have been measured at $E_{\text{d}} = 11.8$ and 22 MeV to $^{10}\text{Be}_{\text{g.s.}}$ [see (1974AJ01)] and at 52 MeV to $^{10}\text{Be}^*(0, 3.37, 5.96, 9.60)$: $S = 0.65, 2.03, 0.13, 1.19$ (normalized to the theoretical value for the ground state); $\pi = +$ for $^{10}\text{Be}^*(9.6)$ (1975SC41).

26. $^{11}\text{B}(\text{t}, \alpha)^{10}\text{Be}$ $Q_{\text{m}} = 8.5853$

See (1979AJ01).

27. $^{11}\text{B}(^{14}\text{N}, ^{15}\text{O})^{10}\text{Be}$ $Q_{\text{m}} = -3.932$

See (1974AJ01). See also (1978MA1F).

28. $^{12}\text{C}(\text{p}, 3\text{p})^{10}\text{Be}$ $Q_{\text{m}} = -27.1849$

See (1979KO36).

29. $^{12}\text{C}(^6\text{Li}, ^8\text{B})^{10}\text{Be}$ $Q_{\text{m}} = -21.442$

At $E(^6\text{Li}) = 80$ MeV, $^{10}\text{Be}^*(0, 3.37, 5.96, 7.54, (9.4), 11.8)$ are populated and the angular distribution to $^{10}\text{Be}_{\text{g.s.}}$ has been measured: see (1979AJ01). See also (1982AL08).

¹⁰B
(Figs. 20 and 22)

GENERAL: (See also (1979AJ01).)

Shell and deformed models: (1978FU13, 1979FL06, 1979KU05, 1980NIIF, 1981BO1Y, 1981DE2G, 1982BA52).

Cluster and α -particle models: (1979AD1A, 1980FU1G, 1980NIIF, 1980OK1B, 1981KR1J, 1983RO1G).

Special states: (1979FL06, 1980BR21, 1980FU1G, 1980NIIF, 1980OK1B, 1980RI06, 1981BA64, 1981BO1Y, 1981DE2G, 1981KU04, 1981SE06, 1982BA52, 1983GO1R).

Electromagnetic transitions and giant resonances: (1978FU13, 1979FL06, 1979KU05, 1980KO1L, 1980NIIF, 1980RI06, 1981BA64, 1981BO1Y, 1981KN06, 1982BA52, 1982RI04, 1982VE11).

Astrophysical questions: (1978BU1B, 1979MO04, 1979RA1C, 1980RE1B, 1981AU1D, 1981AU1G, 1981GU1D).

Applied work: (1979AT01, 1979FL1A, 1979FO1F, 1979JU1B, 1980MU1D, 1983ST1H).

Complex reactions involving ¹⁰B: (1978BH03, 1978HE1C, 1979AL22, 1979BO22, 1979JA1C, 1979LO11, 1979SA27, 1979ST1D, 1979VI05, 1980GR10, 1980GU1E, 1980MI01, 1980OL1C, 1980RI06, 1980WI1L, 1981BL1G, 1981ME13, 1981MO20, 1981TA22, 1981VA1D, 1982CH1M, 1982FU04, 1982GO1E, 1982LU01, 1982LY1A, 1982MO1K, 1982VE11, 1983BE02, 1983SA06).

Muon and neutrino capture and reactions: (1979BE1N, 1979DE01, 1979GO1M, 1980MU1B, 1981GI08, 1981MU1E, 1981OL01, 1982NA01).

Pion and kaon capture and reactions (See also reactions 23, 24 and 47.): (1978AN20, 1978DA1A, 1978TS1A, 1979AL1J, 1979AL21, 1979BA16, 1979BE1N, 1979BO1P, 1979BO1N, 1979PI06, 1979TI1A, 1979TR1B, 1980CR03, 1980DE11, 1980LE02, 1980MA1F, 1980NA1B, 1980ST25, 1981BE63, 1981FR17, 1981GE1B, 1981GI15, 1981SI1D, 1981ST05, 1981YA1A, 1982RO04).

Hypernuclei: (1978DA1A, 1978PO1A, 1978SO1A, 1980IW1A, 1981WA1J, 1982ER1E, 1982KO11, 1982RA1L).

Other topics: (1980BR21, 1981AV02, 1981KU04, 1981SE06, 1982BA2G, 1982CH1M, 1982DE1N, 1982NG01, 1982VE02, 1983GO1R).

Ground-state properties of ¹⁰B: (1978HE1D, 1979BE1N, 1979SA27, 1980FU1G, 1981AV02, 1981BO1Y, 1981OL01, 1981SE06, 1982BA2G, 1982LO13, 1982NG01).

$$\mu = +1.80065 \pm 0.00001 \text{ nm: see (1978LEZA)}$$

$$Q = +84.72 \pm 0.56 \text{ mb: see (1978LEZA)}.$$

Mass of ¹⁰B: (A.H. Wapstra (private communication) adopts 12050.0 ± 0.39 keV, and we shall too. See also (1983CH08).

¹⁰B(0.72):* $\mu = +0.63 \pm 0.12$ nm: see (1978LEZA). See also (1982VE11).

Table 10.5: Energy levels of ^{10}B ^a

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|------------|---|------------------|--|
| g.s. | $3^+; 0$ | stable | | 1, 4, 5, 6, 11, 12, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 43, 44, 45, 48, 49, 50, 51, 52, 53, 55, 56, 57 |
| 0.71835 ± 0.04 | $1^+; 0$ | $\tau_m = 1.020 \pm 0.005$ nsec $g = +0.63 \pm 0.12$ | γ | 1, 4, 5, 6, 11, 12, 17, 18, 19, 20, 25, 26, 27, 29, 30, 35, 41, 43, 44, 45, 47, 48, 49, 50, 52, 55, 56, 57 |
| 1.74015 ± 0.17 | $0^+; 1$ | 7 ± 3 fsec | γ | 1, 4, 12, 17, 18, 19, 20, 23, 25, 26, 29, 41, 42, 43, 44, 45, 48, 49, 53, 56 |
| 2.1543 ± 0.5 | $1^+; 0$ | 2.13 ± 0.20 psec | γ | 1, 4, 12, 17, 18, 19, 20, 25, 26, 27, 29, 30, 35, 43, 44, 45, 47, 48, 49, 50, 52, 55, 56 |
| 3.5871 ± 0.5 | $2^+; 0$ | 153 ± 12 fsec | γ | 1, 4, 6, 12, 17, 18, 19, 25, 26, 27, 29, 30, 42, 43, 45, 48, 49, 50, 52, 55, 56 |
| 4.7740 ± 0.5 | $3^+; 0$ | $\Gamma = 8.7 \pm 2.2$ eV | γ, α | 1, 4, 6, 17, 18, 19, 26, 27, 30, 45, 48, 49, 50, 55 |
| 5.1103 ± 0.6 | $2^-; 0$ | 0.98 ± 0.07 keV | γ, α | 1, 12, 17, 18, 26, 30, 45, 49 |
| 5.1639 ± 0.6 | $2^+; 1$ | $\tau_m < 6$ fsec | γ, α | 1, 12, 17, 18, 23, 26, 27, 42, 45, 48 |
| 5.180 ± 10 | $1^+; 0$ | $\Gamma = 100 \pm 10$ | γ, α | 1, 3, 12, 17, 18, 27, 30, 45 |
| 5.9195 ± 0.6 | $2^+; 0$ | 6 ± 1 | γ, α | 1, 3, 12, 17, 18, 19, 26, 27, 29, 30, 45, 48, 49, 50 |
| 6.0250 ± 0.6 | 4^+ | 0.05 ± 0.03 | γ, α | 1, 3, 17, 18, 19, 23, 25, 26, 27, 29, 30, 43, 45, 49, 50, 53, 55, 56 |
| 6.1272 ± 0.7 | 3^- | 2.36 ± 0.03 | α | 3, 17, 18, 19, 26, 27, 29, 43, 45, 49, 56 |

Table 10.5: Energy levels of ^{10}B ^a (continued)

| E_x (MeV \pm keV) | $J^\pi; T$ | τ_m or $\Gamma_{c.m.}$ (keV) | Decay | Reactions |
|-----------------------|---|-----------------------------------|---|---|
| 6.561 \pm 1.9 | (4) ⁻ | 25.1 \pm 1.1 | α | 3, 17, 18, 19, 26, 27, 29, 30, 43, 45, 48, 49 |
| 6.873 \pm 5 | 1 ⁻ ; 0 + 1 | 120 \pm 5 | γ , p, d, α | 1, 12, 14, 16, 17 |
| 7.002 \pm 6 | (1, 2) ⁺ ; (0) | 100 \pm 10 | p, d, α | 3, 16, 17, 19, 26, 27, 29, 45, 49, 55 |
| 7.430 \pm 10 | 2 ⁽⁻⁾ ; 0 + 1 | 100 \pm 10 | γ , p, d, α | 1, 12, 16, 23 |
| 7.467 \pm 10 | 1 ⁺ | 65 \pm 10 | p | 14, 45 |
| 7.479 \pm 2 | 2 ⁺ ; 1 | 74 \pm 4 | γ , p | 12, 14, 23, 45 |
| 7.5607 \pm 0.9 | 0 ⁺ ; 1 | 2.65 \pm 0.18 | γ , p | 12, 14, 17, 45 |
| (7.67 \pm 30) | (1 ⁺ ; 0) | 250 \pm 20 | p, d | 14, 16 |
| 7.819 \pm 20 | 1 ⁻ | 260 \pm 30 | p | 14, 17, 19, 45 |
| 8.07 | 2 ⁺ | 800 \pm 200 | γ , p, d | 16, 17, 23 |
| (8.7) | (1 ⁺ , 2 ⁺) | (\approx 200) | p | 14, 16, 55 |
| 8.889 \pm 6 | 3 ⁻ ; 1 | 84 \pm 7 | γ , n, p, α | 13, 14, 16, 19, 23, 48 |
| 8.895 \pm 2 | 2 ⁺ ; 1 | 40 \pm 1 | γ , p, α | 12, 14, 16, 19, 23, 48 |
| (9.7) | ($T = 1$) | (\approx 700) | n, p, α | 13, 16 |
| 10.84 \pm 10 | (2 ⁺ , 3 ⁺ , 4 ⁺) | 300 \pm 100 | γ , n, p | 12, 13, 14, 23, 45 |
| 11.52 \pm 35 | | 500 \pm 100 | (γ) | 23, 43, 45 |
| 12.56 \pm 30 | (0 ⁺ , 1 ⁺ , 2 ⁺) | 100 \pm 30 | γ , p | 12, 14, 23, 45 |
| 13.49 \pm 5 | (0 ⁺ , 1 ⁺ , 2 ⁺) | 300 \pm 50 | γ , p | 12, 23, 45 |
| 14.4 \pm 100 | | 800 \pm 200 | γ , p, α | 3, 12, 43, 45 |
| (18.2 \pm 200) | | (1500 \pm 300) | | 45 |
| 18.43 | 2 ⁻ ; 1 | 340 | γ , ^3He | 6, 8 |
| 18.80 | 2 ⁺ , 1 ⁺ | < 600 | γ , ^3He , α | 6, 10 |
| 19.29 | 2 ⁻ ; 1 | 190 \pm 20 | γ , n, ^3He , α | 6, 7, 8, 10 |
| 20.2 | 1 ⁻ ; 1 | broad | γ , n, t, ^3He , α | 6, 7, 8, 9, 10, 22 |
| (21.1) | | | γ , ^3He | 6 |
| 23.1 | | broad | γ , n | 22 |

^a See also Tables 10.6, 10.7 and 10.11.

Table 10.6: Electromagnetic transitions in ^{10}B ^a

| Initial state | $J^\pi; T$ | Γ_γ (total) (eV) | Branching ratios (%) to final states at: | | | | | Γ_γ/Γ | |
|---------------|--------------------|----------------------------------|--|----------------------------|----------------------------|----------------------------|----------------------------|--------------------------------|-----------------|
| | | | g.s. 3 ⁺ ; 0 | 0.72 1 ⁺ ; 0 | 1.74 0 ⁺ ; 1 | 2.15 1 ⁺ ; 0 | 3.59 2 ⁺ ; 0 | | |
| 0.72 | 1 ⁺ ; 0 | 6.5×10^{-7} | 100 | | | | | $(2.3 \pm 0.3) \times 10^{-3}$ | |
| 1.74 | 0 ⁺ ; 1 | 0.09 ± 0.04 ^b | < 0.2 | 100 | | | | | |
| 2.15 | 1 ⁺ ; 0 | $(3.1 \pm 0.3) \times 10^{-4}$ | 21.1 ± 1.6 | 27.3 ± 0.9 | 51.6 ± 1.6 | | | | |
| 3.59 | 2 ⁺ ; 0 | $(4.31 \pm 0.34) \times 10^{-3}$ | 19 ± 3 | 67 ± 3 | < 0.3 | 14 ± 2 | | | |
| 4.77 | 3 ⁺ ; 0 | 0.020 ± 0.004 | 0.5 ± 0.1 | > 99 | | | | | |
| 5.11 | 2 ⁻ ; 0 | | 64 ± 7 | 31 ± 7 | 5 ± 5 | | | | |
| 5.16 | 2 ⁺ ; 1 | 1.5 ± 0.1 ^c | 4.4 ± 0.4 | 22.4 ± 0.6 | 0.7 ± 0.2 | 64.8 ± 0.9 | 7.7 ± 0.3 | | 0.87 ± 0.04 |
| 5.18 | 1 ⁺ ; 0 | 0.06 ± 0.03 | | | ≈ 100 | | | | |
| 5.92 | 2 ⁺ ; 0 | 0.15 ± 0.04 | 82 ± 5 | 18 ± 5 | | | | | ≤ 0.009 |
| 6.03 | 4 ⁺ | 0.11 ± 0.02 | ≈ 100 | ^d | | | | | ≤ 0.009 |
| 6.13 | 3 ⁻ | ≤ 21 | | | | | | ≤ 0.009 | |
| ^e | | | | | | | | | |

^a For references see Table 10.6 in (1979AJ01).

^b From Table 10.7.

^c See also Table 10.8 here. Branching ratios and Γ_γ/Γ from (1979KE08). The mixing ratios $\delta = 0.12 \pm 0.05, 0.03 \pm 0.03, 0.02 \pm 0.03$ and 0.00 ± 0.02 for the transitions to $^{10}\text{B}^*(0, 0.72, 2.15, 3.59)$, respectively (1979KE08).

^d Other branches < 3%.

^e For γ -decay of higher ^{10}B states see Tables 10.8, 10.10, 10.11 and 10.12. See also Table 10.16.

Table 10.7: Lifetimes of ^{10}B states

| $^{10}\text{B}^*$ (MeV) | τ_m | Reactions | Refs. |
|-------------------------|------------------------|-------------------------------|---------------------|
| 0.72 | 1.020 ± 0.005 nsec | mean | ^a |
| 1.74 | 7 ± 3 fsec | $^6\text{Li}(\alpha, \gamma)$ | (KE79A) |
| 2.15 | 2.30 ± 0.26 psec | mean | (AJ79) ^b |
| | 1.9 ± 0.3 psec | $^6\text{Li}(\alpha, \gamma)$ | (KE79A) |
| 3.59 | 2.13 ± 0.20 psec | mean | all values |
| | 153 ± 13 fsec | mean | (AJ79) |
| | 150 ± 30 fsec | $^6\text{Li}(\alpha, \gamma)$ | (KE79A) |
| 5.16 | 153 ± 12 fsec | mean | all values |
| | < 6 fsec | $^6\text{Li}(\alpha, \gamma)$ | (KE79A) |

^a Weighted mean of last 7 values in Table 10.20 of (1966LA04) and of a new measurement by Vermeer, Bhalla and Polletti, $\tau_m = 1.020 \pm 0.005$ nsec. I am indebted to Prof. F.C. Barker for informing me of this new value.

^b Table 10.9 in (1979AJ01).

1. $^6\text{Li}(\alpha, \gamma)^{10}\text{B}$ $Q_m = 4.4603$

Observed resonances are displayed in Table 10.8. See also (1981GA1A, 1981MUIF).

2. (a) $^6\text{Li}(\alpha, n)^9\text{B}$ $Q_m = -3.977$ $E_b = 4.4603$
 (b) $^6\text{Li}(\alpha, p)^9\text{Be}$ $Q_m = -2.1262$
 (c) $^6\text{Li}(\alpha, d)^8\text{Be}$ $Q_m = -1.5671$

The excitation functions for neutrons [from threshold to $E_\alpha = 15.5$ MeV] and for deuterons [$E_\alpha = 9.5$ to 25 MeV; d_0, d_1 over most of range] do not show resonance structure. See also ^9B , ^9Be and ^8Be . For reaction (a) see also (1979BA48). See (1979AJ01) for references.

3. (a) $^6\text{Li}(\alpha, \alpha)^6\text{Li}$ $E_b = 4.4603$
 (b) $^6\text{Li}(\alpha, 2\alpha)^2\text{H}$ $Q_m = -1.4753$

Table 10.8: Levels of ^{10}B from $^6\text{Li}(\alpha, \gamma)^{10}\text{B}$ ^a

| E_{res} (keV) | E_x (MeV \pm keV) | $J^\pi; T$ | Γ_{lab} (keV) | Decay to E_f | Branch (%) | $\omega\gamma$ (eV) | Γ_γ (eV) |
|------------------------|-----------------------|--------------------------|------------------------------|----------------|----------------|---|--------------------------------|
| 500 \pm 25 | 4.761 | 3 ⁺ ; 0 | | 0 | 0.5 \pm 0.1 | | 0.020 \pm 0.004 ^h |
| | | | | 0.72 | > 99 | | |
| 1085 | 5.112 | 2 ⁻ ; 0 | 1.63 \pm 0.11 ^b | 0 | 64 \pm 7 | 0.059 \pm 0.012 ^c | |
| | | | | 0.72 | 31 \pm 7 | 0.028 \pm 0.008 | |
| | | | | 1.74 | 5 \pm 5 | 0.005 \pm 0.005 | |
| 1175 ^d | 5.166 | 2 ⁺ ; 1 | < 0.5 | 0 | 4.4 \pm 0.4 | 0.018 \pm 0.002 | 0.068 \pm 0.007 |
| | | | | 0.72 | 22.4 \pm 0.6 | 0.090 \pm 0.008 | 0.33 \pm 0.03 |
| | | | | 1.74 | 0.7 \pm 0.2 | (2.8 \pm 0.8) \times 10 ⁻³ | 0.010 \pm 0.003 |
| | | | | 2.15 | 64.8 \pm 0.9 | 0.259 \pm 0.024 | 0.942 \pm 0.090 |
| | | | | 3.59 | 7.7 \pm 0.3 | 0.031 \pm 0.004 | 0.114 \pm 0.015 |
| 1210 \pm 35 | 5.187 | 1 ⁺ ; 0 | 340 \pm 50 | 1.74 | \approx 100 | | 0.06 \pm 0.03 |
| 2435 ^e | 5.922 | 2 ⁺ | 10 \pm 1 | 0 | 82 \pm 5 | 0.19 \pm 0.04 | 0.13 \pm 0.03 |
| | | | | 0.72 | 18 \pm 5 | 0.04 \pm 0.01 | 0.02 \pm 0.01 |
| | | | | 1.74 | | < 0.02 | |
| 2605 ^e | 6.024 | 4 ⁺ | 0.08 \pm 0.05 | 0 | \approx 100 | 0.34 \pm 0.05 | 0.11 \pm 0.02 |
| | | | | 0.72 | | < 0.02 | |
| 4019 ^f | 6.873 \pm 5 | 1 ⁻ ; 0 + 1 | 200 \pm 10 | 0 | 6 \pm 2 | | |
| | | | | 0.72 | 21 \pm 4 | | |
| | | | | 1.74 | 59 \pm 3 | | |
| | | | | 2.15 | 14 \pm 4 | | |
| 4963 ^g | 7.440 \pm 20 | 2 ⁽⁻⁾ ; 0 + 1 | 150 \pm 15 | ^g | | | |

^a For earlier references see Table 10.7 in (1979AJ01).

^b (1983NAZZ) and J. Napolitano, private communication.

^c Absolute error only.

^d Branching ratios from (1979KE08); $\omega_{\gamma_{c.m.}} = 0.40 \pm 0.04$ eV (1979SP01), 0.43 ± 0.07 eV (prelim., J. Napolitano, private communication); $\Gamma_{\alpha}/\Gamma = 0.16 \pm 0.04$ (1979SP01).

^e Values of ω_{γ} (1966FO05) have been multiplied by 0.6 to convert them to the c.m. system. I am indebted to Prof. F.C. Barker for pointing out this problem.

^f Branching ratios calculated from 0° relative intensities; $\Gamma_{\alpha}/\Gamma_p = 1.25 \pm 0.12$.

^g At 0° the branches to $^{10}\text{B}^*(0, 0.72)$ are equally strong ($(50 \pm 12)\%$).

^h $\Gamma_{\gamma}/\Gamma = (2.3 \pm 0.3) \times 10^{-3}$; $\Gamma_{\alpha} = 8.4 \pm 1.8$ eV (E.K. Warburton and D.E. Alburger, private communication).

Table 10.9: ^{10}B levels from $^6\text{Li}(\alpha, \alpha)^6\text{Li}$ ^a

| E_α (keV) | E_x (MeV) | Γ_{lab} (keV) | $J^\pi; T$ |
|-------------------------------|-------------|-----------------------------|--------------|
| 1210 ± 30 | 5.19 | 175 | $1^+; 0$ |
| 2440 ^b | 5.93 | ≈ 30 | $2^+; 0$ |
| 2606.0 ± 1.5 | 6.025 | 0.09 ± 0.04 | 4^+ |
| 2785.5 ± 1.5 ^c | 6.133 | 3.93 ± 0.05 | 3^- |
| 3498.5 ± 1.6 ^d | 6.561 | 41.8 ± 1.9 | $4^-, 2^-$ |
| 4250 ± 15 ^d | 7.012 | 183 ± 25 | $(2)^+; (0)$ |
| 16000 | 14.1 | broad | |

^a For references see Table 10.8 in (1979AJ01).

^b $\Gamma_\alpha = 9.7 \pm 0.1$ keV (1981HE05).

^c $\Gamma_\alpha = 2.45 \pm 0.12$ keV and $\Gamma_d = 0.08 \pm 0.05$ keV (1981HE05).

^d There is evidence of broad structure near these states.

Excitation functions of α_0 and α_1 have been reported for $E_\alpha \leq 18.0$ MeV and 9.5 to 12.5 MeV, respectively: see (1974AJ01). Reported anomalies are displayed in Table 10.9. Elastic scattering and VAP measurements are reported for $E_d^- = 15.1$ to 22.7 MeV (1979EG01). Small anomalies are observed in the excitation function of the magnitude parameter (reaction (b)) corresponding to $^{10}\text{B}^*(8.67, 9.65, 10.32, 11.65)$ (1983GO07). See, however, Table 10.5. See also ^6Li , (1980SK1A) and (1979NO1C, 1979SU09, 1979SU1F, 1980FU1G, 1982LE10, 1983SH04; theor.).

$$4. \ ^6\text{Li}(^6\text{Li}, d)^{10}\text{B} \quad Q_m = 2.985$$

Angular distributions of deuteron groups have been determined at $E(^6\text{Li}) = 2.4$ to 9.0 MeV (d_0, d_1, d_3) and 7.35 and 9.0 MeV (d_4, d_5). The d_2 group is also observed but its intensity is weak: see (1974AJ01) and ^{12}C in (1980AJ01). See also (1979WA13).

$$5. \ ^6\text{Li}(^7\text{Li}, t)^{10}\text{B} \quad Q_m = 1.992$$

Angular distributions of the t_0 and t_1 groups have been measured at $E(^6\text{Li}) = 3.3$ MeV and $E(^7\text{Li}) = 3.78$ to 5.95 MeV: see (1974AJ01).

$$6. \ ^7\text{Li}(^3\text{He}, \gamma)^{10}\text{B} \quad Q_m = 17.7880$$

Table 10.10: Resonances in ${}^7\text{Li} + {}^3\text{He}$ ^a

| E_{res} (MeV) | $\Gamma_{\text{c.m.}}$ (keV) | E_x (MeV) | $J^\pi; T$ | Γ_γ (eV) for transition to | | | | Γ_α (keV) | Γ_n | Γ_p | Γ_t |
|------------------------|------------------------------|-------------|------------|--|-----------|------------------------|-----------|-----------------------|------------|------------|------------|
| | | | | g.s. | 0.72 | 3.59 | 4.77 | | | | |
| 0.92 | 340 | 18.43 | $2^-; 1$ | ≥ 3 | | | ≥ 17 | | | | |
| 1.45 ^b | < 600 | 18.80 | $2^+; 1^+$ | | ≥ 20 | ≥ 20 ^c | | res < 80 | | | |
| 2.15 ^b | 280 ^d | 19.29 | $2^-; 1$ | ≥ 12 | | | ≥ 49 | res < 20 | res n_0 | (p) | |
| 3.4 | 910 ^d | 20.2 | $1^-; 1$ | | | ≥ 350 | | res α_2 | res n_0 | (p) | res t_0 |
| (4.7) | | (21.1) | | | res | | | | | | |

^a See references listed in Table 10.10 in (1974AJ01, 1979AJ01).

^b See (1979LI1B, 1980LI1F; abstracts).

^c Assumes isotropy of angular distribution.

^d $\Gamma_{\text{c.m.}} = 190 \pm 20$ and 350 ± 70 keV, respectively, from the n_0 yield.

Capture γ -rays have been observed for $E({}^3\text{He}) = 0.8$ to 6.0 MeV. The γ_0 and γ_5 yields [to ${}^{10}\text{B}^*(0, 4.77)$] show resonances at $E({}^3\text{He}) = 1.1$ and 2.2 MeV [$E_{\text{res}} = 0.92$ and 2.1 MeV], the γ_1 and γ_4 yields [to ${}^{10}\text{B}^*(0.72, 3.59)$] at 1.4 MeV and the γ_4 yield at 3.4 MeV: see Table 10.10. Both the 1.1 and 2.2 MeV resonances [${}^{10}\text{B}^*(18.4, 19.3)$] appear to result from s-wave capture; the subsequent decay is to two 3^+ states [${}^{10}\text{B}^*(0, 4.77)$]. Therefore the most likely assignment is $2^-, T = 1$ for both [there appears to be no decay of these states via α_2 to ${}^6\text{Li}^*(3.56)$ which has $J^\pi = 0^+, T = 1$: see reaction 10]. The assignment for ${}^{10}\text{B}^*(18.8)$ [1.4 MeV resonance] is 1^+ or 2^+ but there appears to be α_2 decay and therefore $J^\pi = 2^+$. ${}^{10}\text{B}^*(20.2)$ [3.4 MeV resonance] has an isotropic angular distribution of γ_4 and therefore $J^\pi = 0^+, 1^-, 2^-$. The γ_2 group resonates at this energy which eliminates 2^- , and 0^+ is eliminated on the basis of the strength of the transition which is too large for E2. See (1974AJ01) for references.

7. ${}^7\text{Li}({}^3\text{He}, n){}^9\text{Be}$

$$Q_m = 9.351$$

$$E_b = 17.7880$$

The excitation curve is smooth up to $E({}^3\text{He}) = 1.8$ MeV and the n_0 yield shows resonance behavior at $E({}^3\text{He}) = 2.2$ and 3.25 MeV, $\Gamma_{\text{lab}} = 270 \pm 30$ and 500 ± 100 keV. No other resonances are observed up to $E({}^3\text{He}) = 5.5$ MeV. See Table 10.10 and (1974AJ01) for references.

8. ${}^7\text{Li}({}^3\text{He}, p){}^9\text{Be}$

$$Q_m = 11.2015$$

$$E_b = 17.7880$$

The yield of protons has been measured for $E({}^3\text{He}) = 0.60$ to 4.8 MeV: there is some indication of weak maxima at $1.1, 2.3$ and 3.3 MeV. Polarization measurements are reported at $E({}^3\text{He}) = 14$ MeV, and more recently at $E({}^3\text{He}) = 13.6 - 13.7$ MeV (1981SL03) and 14.01 MeV (1983RI01). See also (1981VI1B). For the earlier references see (1974AJ01, 1979AJ01).

9. (a) ${}^7\text{Li}({}^3\text{He}, \text{d}){}^8\text{Be}$ $Q_{\text{m}} = 11.7606$ $E_{\text{b}} = 17.7880$
 (b) ${}^7\text{Li}({}^3\text{He}, \text{t}){}^7\text{Be}$ $Q_{\text{m}} = -0.881$
 (c) ${}^7\text{Li}({}^3\text{He}, {}^3\text{He}){}^7\text{Li}$

Yields of deuterons have been measured for $E({}^3\text{He}) = 1.0$ to 2.5 MeV (d_0) and yields of tritons are reported for 2.0 to 4.2 MeV (t_0): a broad peak is reported at $E({}^3\text{He}) \approx 3.5$ MeV in the t_0 yield. See (1979AJ01) for references. Polarization measurements are reported at $E({}^3\vec{\text{He}}) = 33.3$ MeV for the deuteron groups to ${}^8\text{Be}^*(16.63, 17.64, 18.15)$ (1981BA38) and for the triton and ${}^3\text{He}$ groups to ${}^7\text{Be}^*(0, 0.43)$ and ${}^7\text{Li}^*(0, 0.48, 4.63)$ (1981BA37). See also (1979KA1G).

10. ${}^7\text{Li}({}^3\text{He}, \alpha){}^6\text{Li}$ $Q_{\text{m}} = 13.328$ $E_{\text{b}} = 17.7880$

Excitation functions have been measured for $E({}^3\text{He}) = 1.3$ to 18.0 MeV: see (1974AJ01). The α_0 group (at 8°) shows a broad maximum at ≈ 2 MeV, a minimum at 3 MeV, followed by a steep rise which flattens off between $E({}^3\text{He}) = 4.5$ and 5.5 MeV. Integrated α_0 and α_1 yields rise monotonically to 4 MeV and then tend to decrease. Angular distributions give evidence of the resonances at $E({}^3\text{He}) = 1.4$ and 2.1 MeV seen in ${}^7\text{Li}({}^3\text{He}, \gamma){}^{10}\text{B}$: $J^\pi = 2^+$ or 1^- , $T = (1)$ for both [see, however, reaction 6]: Γ_α is small. The α_2 yield [to ${}^6\text{Li}^*(3.56)$, $J^\pi = 0^+$, $T = 1$] shows some structure at $E({}^3\text{He}) = 1.4$ MeV and a broad maximum at ≈ 3.3 MeV: see Table 10.10. Recent (unpublished) excitation studies have been reported at $E({}^3\text{He}) = 0.6$ to 2.5 MeV for the α_0 , α_1 , α_2 and α_4 groups: the α_4 group shows possible resonances at $E({}^3\text{He}) = 1.45$ and 2.15 MeV. $J^\pi = 2^+$; $T = 1$ is suggested for the lower structure (1979LI1B, 1980LI1F). Polarization measurements are reported at $E({}^3\vec{\text{He}}) = 33.3$ MeV to ${}^6\text{Li}^*(0, 2.19, 3.56)$ (1981BA38).

11. ${}^7\text{Li}(\alpha, \text{n}){}^{10}\text{B}$ $Q_{\text{m}} = -2.7898$

Angular distributions have been measured for $E_\alpha = 4.78$ to 13.9 MeV [see (1974AJ01)] and more recently for $E_\alpha = 4.4$ to 5.1 MeV (1981SE04). See also (1979AJ01).

12. ${}^9\text{Be}(\text{p}, \gamma){}^{10}\text{B}$ $Q_{\text{m}} = 6.5865$

Parameters of observed resonances are listed in Tables 10.11 and 10.12. Table 10.6 summarizes the γ -transitions from this and other reactions. For references to the discussion below, see (1974AJ01, 1979AJ01).

The $E_{\text{p}} = 0.32$ MeV resonance (${}^{10}\text{B}^* = 6.87$ MeV) is ascribed to s-wave protons because of its comparatively large proton width [see ${}^9\text{Be}(\text{p}, \text{p})$] and because of the isotropy of the γ -radiation. The strong transition to ${}^{10}\text{B}^*(1.74)$ requires E1 and hence $J^\pi = 1^-$, $T = 0$. $T = 0$ is also indicated by the large deuteron width. On the other hand, the strength of E1 transitions to ${}^{10}\text{B}^*(0.7, 2.1)$ indicates $T = 1$. The amplitudes for the $T = 0$ and $T = 1$ parts of the wave function for ${}^{10}\text{B}^*(6.87)$ are 0.92 and 0.39 , respectively. See (1982RI04) for the $5.16 \rightarrow 1.74$ decay (Table 10.6).

Table 10.11: Resonances in ${}^9\text{Be}(p, \gamma){}^{10}\text{B}$ ^a

| E_p (MeV \pm keV) | E_x (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | $J^\pi; T$ | Γ_p/Γ | Γ_γ (eV) |
|-----------------------|-----------------------|------------------------------|-----------------------|-------------------|----------------------|
| 0.319 | 6.873 ± 5 | 120 ± 5 | $1^-; 0 + 1$ | 0.30 | 4.8 |
| 0.938 ± 10 | 7.430 | 140 ± 30 | $2^{(-)}; 0 + 1$ | 0.7 | 2.4 |
| (0.98) | (7.47) | | (2^+) | | |
| 0.992 ± 2 | 7.479 | 72 ± 4 | $2^-; 1$ ^b | ≈ 0.65 | 25.8 |
| 1.0832 ± 0.4 | 7.5607 | 2.65 ± 0.18 | $0^+; 1$ | 1.0 | 8.5 |
| 1.29 | 7.75 | 210 ± 60 | $2^-; (1)$ | ≈ 0.65 | 8.5 |
| 2.567 ± 2 | 8.895 | 36 ± 2 | $2^+; 1$ | | |
| 4.72 ^c | 10.83 | ≈ 500 | $2^+, 3^+, 4^+$ | | |
| 6.7 ^c | 12.6 | < 200 | $0^+, 1^+, 2^+$ | | |
| (7.0) ^c | (12.9) | (≈ 100) | ($\pi = ^+$) | | |
| 7.5 ^c | 13.3 | ≈ 300 | $0^+, 1^+, 2^+$ | | |
| 8.4 ^c | 14.1 | ≈ 250 | $0^+, 1^+, 2^+$ | | |
| 8.9 ^c | 14.6 | ≈ 150 | $2^+, 3^+, 4^+$ | | |
| 10.0 ^c | 15.6 | ≈ 400 | $2^+, 3^+, 4^+$ | | |
| 14.6 ^c | 19.7 | ≈ 500 | $2^-, 3^-, 4^-$ | | |

^a For references and for additional comments see table 10.11 in (1979AJ01). See Table 10.12 for decay schemes.

^b See (1974AJ01). This state is assigned $J^\pi = 2^+$ on the basis of the (e, e') work (see Table 10.16). I am indebted to Dr. D. Kurath for his comments.

^c Unpublished Ph.D. thesis.

Table 10.12: Radiative transitions in ${}^9\text{Be}(p, \gamma){}^{10}\text{B}$ ^a

| Initial state (MeV) | $J^\pi; T$ | Γ_γ (tot) (eV) | Relative intensities to final states | | | | | | | | |
|-----------------------------------|--------------------------|----------------------------|--------------------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| | | | ground 3 ⁺ ; 0 | 0.72 1 ⁺ ; 0 | 1.74 0 ⁺ ; 1 | 2.15 1 ⁺ ; 0 | 3.59 2 ⁺ ; 0 | 5.11 2 ⁻ ; 0 | 5.16 2 ⁺ ; 1 | 5.18 1 ⁺ ; 0 | 5.92 2 ⁺ ; 1 |
| 6.87 $E_p = 0.32$ | 1 ⁻ ; 0 + 1 | 4.8 ^b | < 0.05 | 0.20 ± 0.02 | 0.53 ± 0.02 | 0.13 ± 0.01 | < 0.01 | 0.04 ± 0.01 | 0.03 ± 0.01 | < 0.01 | 0.035 ± 0.01 |
| 7.43 $E_p = 0.94$ | 2 ⁽⁻⁾ ; 0 + 1 | [2.4] | < 2 | 1.3 | [< 0.14] | 0.62 | 0.5 | | | [< 1] | |
| 7.48 $E_p = 0.99$ | 2 ⁺ ; 1 | [25.8] | 25 | 0.3 | [< 0.14] | 0.49 | 0 | | | [< 1] | |
| 7.56 $E_p = 1.08$ | 0 ⁺ ; 1 | [8.5] | < 0.2 | 6.7 | < 0.3 | 0.8 | < 0.2 | | | 1.0 | |
| 7.75 ^c $E_p = 1.29$ | (2 ⁻ ; 1) | [8.5] | 6.6 | 0.9 | < 0.08 | 0.3 | 0.3 | | | 0.4 | |

^a For references and other values see Table 10.12 in (1979AJ01).

^b See, however, Table 10.8, footnote ^f.

^c See, however, Table 10.14.

The proton-capture data near $E_p = 1$ MeV appears to require at least five resonant states, at $E_p = 938$, (980), 992, 1083 and 1290 keV. The narrow $E_p = 1083$ keV level ($^{10}\text{B}^* = 7.56$ MeV) is formed by p-wave protons, $J^\pi = 0^+$ [see $^9\text{Be}(p, p)$, $^9\text{Be}(p, \alpha)$]. The isotropy of the γ -rays supports this assignment. The strong M1 transitions to $J^\pi = 1^+$; $T = 0$ levels at 0.72, 2.15 and 5.18 MeV (Table 10.12) indicate $T = 1$. The width of $^{10}\text{B}^*(5.18)$ observed in the decay is 100 ± 10 keV.

The excitation function for the ground-state radiation shows resonance at $E_p = 992$ ($\Gamma = 80$ keV) and 1290 keV ($\Gamma = 230$ keV). Elastic scattering studies indicate s-wave formation and $J^\pi = 2^-$ for both. For the lower level ($E_x = 7.48$ MeV) the intensity of the g.s. capture radiation, $\Gamma_\gamma = 25$ eV indicates E1 and $T = 1$. The angular distribution of γ -rays, $1 + 0.1 \sin^2 \theta$, is consistent with s-wave formation with some d-wave admixture or with some contribution from a nearby p-wave resonance; possibly a $J^\pi = 2^+$ level at $E_p = 980$ keV.

The angular distribution of ground-state radiation at $E_p = 1330$ keV is isotropic and $\Gamma_\gamma = 8.5$ eV, supporting E1, $T = 1$ for this level ($E_x = 7.75$ MeV).

Transitions to $^{10}\text{B}^*(0.7)$ [γ_1] show resonances at $E_p = 992$, 1290 and 938 keV, $\Gamma = 155$ keV. The latter is presumably also a resonance for (p, d) and (p, α). An assignment of $J^\pi = 2^-$, $T = 0$ is consistent with the data, although the E1 radiation then seems somewhat too strong for a $\Delta T = 0$ transition.

A resonance for capture radiation at $E_p = 2.567 \pm 0.003$ ($E_x = 8.895$ MeV) has a width of 40 ± 2 keV and decays mainly via $^{10}\text{B}^*(0.7)$ (unpublished Ph.D. thesis). It appears from the width that this resonance corresponds to that observed in $^9\text{Be}(p, \alpha)$, $J^\pi = 2^+$, $T = 1$ and not to the $^9\text{Be}(p, n)$ resonance at the same energy. A further resonance is reported at $E_p = 4.72 \pm 0.01$ MeV, $\Gamma \approx 0.5$ MeV.

In the range $E_p = 4$ to 18 MeV, the γ_0 yield at 90° shows the resonance at $E_p = 4.7$ MeV ($E_x = 10.7$ MeV) and shows fluctuations suggesting states at $E_x \approx 14.6$, 15.6 and 19.7 MeV. It is suggested that $^{10}\text{B}^*(19.7)$ decays via E1 and therefore $J^\pi = 2^-, 3^-, 4^-$. The other three states presumably decay by M1 and therefore $J^\pi = 2^+, 3^+, 4^+$. These fluctuations appear on a nearly constant γ_0 yield with a 90° differential cross section $\approx 1.5 \mu\text{b}/\text{sr}$. The average yield of γ_1 is $\approx \frac{2}{3}$ of the γ_0 yield. The broad giant resonance peak is centered at $E_x \approx 14.5$ MeV. Fluctuations in the γ_1 yield are reported at $E_x \approx 12.6$, 13.3 and 14.1 MeV. These states presumably decay by M1 to $^{10}\text{B}^*(0.7)$ [$J^\pi = 1^+$] and therefore $J_1^\pi = 0^+, 1^+, 2^+$. The weak γ_2 yield (to $^{10}\text{B}^*(1.74)$ [$J^\pi; T = 0^+; 1$]) seems to exhibit a broad peak centered near $E_x = 15$ MeV (maximum 90° differential cross section $\approx 0.5 \mu\text{b}/\text{sr}$) and possibly some structure near $E_x = 20$ MeV. The γ_3 yield (to $^{10}\text{B}^*(2.15)$ [$J^\pi = 1^+$]) increases to $\approx 0.4 \mu\text{b}/\text{sr}$ at $E_x \approx 16$ MeV and seems to remain constant beyond that energy, with some suggestion of a fluctuation corresponding to $E_x \approx 12.9$ MeV. $^{10}\text{B}^*(12.9)$ appears to have positive parity. Angular distributions of γ_0 , γ_1 , γ_2 and γ_3 are also reported (unpublished Ph.D. thesis).

The magnetic moment of $^{10}\text{B}^*(0.72)$ has been studied via $\gamma - \gamma$ correlations from $^{10}\text{B}^*(7.56)$: $g = +0.63 \pm 0.12$.

13. $^9\text{Be}(p, n)^9\text{B}$

$$Q_m = -1.851$$

$$E_b = 6.5865$$

Resonances in the neutron yield occur at $E_p = 2562 \pm 6$, 4720 ± 10 and, possibly, at 3500 keV with $\Gamma_{\text{c.m.}} = 84 \pm 7$, ≈ 500 and ≈ 700 keV. These three resonances correspond to $^{10}\text{B}^*(8.891, 10.83, 9.7)$: see Table 10.13 in (1974AJ01). Cross-section measurements for the (p, n) and (p, n_0) reactions have been obtained by (1983BY01; $E_p = 8.15$ to 15.68 MeV) [see also for a review of earlier work]. They indicate possible structure in ^{10}B near 13–14 MeV (1983BY01).

The $E_p = 2.56$ MeV resonance is considerably broader than that observed at the same energy in ${}^9\text{Be}(p, \alpha)$ and ${}^9\text{Be}(p, \gamma)$ and the two resonances are believed to be distinct. The shape of the resonance and the magnitude of the cross section can be accounted for with $J^\pi = 3^-$ or 3^+ ; the former assignment is in better accord with ${}^{10}\text{Be}^*(7.37)$. For $J^\pi = 3^-$, $\theta_n^2 = 0.135$, $\theta_p^2 = 0.115$ ($R = 4.47$ fm): see (1974AJ01).

The analyzing power for n_0 has been measured for $E_{\bar{p}} = 2.7$ to 17 MeV (1980MA33, 1981BY1B, 1981BY1C, 1981MU1D), as has P^y near 8.1 MeV. See also (1983BY01). The polarization transfer coefficient has been studied for $E_{\bar{p}} = 3.9$ to 15.1 MeV by (1976LI08): negative values of $K_y^{y'}$ (0) are reported near $E_p = 7$ MeV in a region where several states are known to exist in ${}^{10}\text{B}$; a spin-flip mechanism may also be involved. Polarization measurements are also reported at $E_{\bar{p}} = 135$ MeV (1981MA1J) and 800 MeV (1981RI06). See also ${}^9\text{B}$, (1979AJ01), (1979BA68), (1980WA1K, 1981WA1G, 1982BY1A) and (1978BA1F, 1981UL1B; applications).

14. (a) ${}^9\text{Be}(p, p){}^9\text{Be}$

$$E_b = 6.5865$$

(b) ${}^9\text{Be}(p, pn){}^8\text{Be}$

$$Q_m = -1.6655$$

The elastic scattering has been studied for $E_p = 0.2$ to 9.5 MeV [see (1974AJ01, 1979AJ01)] and (1980BO1L; $E_p = 2.31 \rightarrow 2.73$ MeV; p_0). Below $E_p = 0.7$ MeV only s-waves are present exhibiting resonance at $E_p = 330$ keV [${}^{10}\text{B}^*(6.88)$], $J^\pi = 1^-$. Between $E_p = 0.8$ to 1.6 MeV polarization and cross-section measurements are well fitted by a phase-shift analysis, using only the ${}^3\text{S}_1$, ${}^5\text{S}_2$, ${}^5\text{P}_1$ and ${}^5\text{P}_2$ phases. Four levels satisfy the data, 1^+ and 2^- states at $E_x = 7.48$ MeV, a sharp 0^+ state at $E_x = 7.56$ MeV, and a 1^- state at 7.82 MeV: see Table 10.13. Pronounced minima at $E_p = 2.48$ and 2.55 are observed in the polarization (p_0): these are ascribed to $T = 1$ analogs of the 3^- and 2^+ states ${}^{10}\text{Be}^*(7.37, 7.52)$. A strong anomaly is observed at $E_p = 6.7$ MeV: see Table 10.13.

Polarization measurements have been reported at $E_p = 0.9$ to 49.8 MeV, at 138.2 and 145 MeV, and at 990 MeV [see (1974AJ01, 1979AJ01)] and at $E_{\bar{p}} = 780$ MeV (1982RA20) as well as at 1 GeV (1983BE16). See also ${}^9\text{Be}$. A_y and K_y^y have been measured at $E_{\bar{p}} = 225$ MeV (1981RO1M; p_0). Total reaction cross sections have been measured at eight energies in the range $E_p = 225$ to 557 MeV (1979SC07). Inclusive cross sections have been measured by (1979FR12, 1979KO21, 1980NI09). Polarization transfer parameters have been measured at $E_p = 800$ MeV (1981RI06). Hadron multiple production has been studied by (1978AR1J). For pion and kaon production at 400 GeV see (1980NI09). For reaction (b) see ${}^9\text{Be}$ and (1982PE1F). See also (1981CO1D, 1981BA1R, 1981NA05, 1981WA1G, 1982BA2T, 1983SEZW), (1983LE1J; astrophysics) and (1978BH1B, 1979BY01, 1979WE1C, 1981BO1C, 1981KR17, 1983BY01; theor.).

15. (a) ${}^9\text{Be}(p, t){}^7\text{Be}$

$$Q_m = -12.082$$

$$E_b = 6.5865$$

(b) ${}^9\text{Be}(p, {}^3\text{He}){}^7\text{Li}$

$$Q_m = -11.201$$

Polarization measurements (reaction (b)) are reported at $E_{\bar{p}} = 23.06$ MeV (1983RI01). See also (1979AR04, 1981BA1R, 1981SL03, 1983SEZW).

Table 10.13: Resonances in ${}^9\text{Be}(p, p){}^9\text{Be}$ ^a

| E_{res} (keV) | E_x (MeV) | $\Gamma_{\text{c.m.}}$ (keV) | J^π | Γ_p/Γ |
|------------------------|-------------|------------------------------|--------------------|-------------------|
| 330 | 6.88 | 145 | 1^- | 0.30 |
| 980 ± 10 | 7.468 | 65 ± 10 | 1^+ | 1.0 |
| 980 ± 10 | 7.468 | 80 ± 8 | 2^- ^b | 0.90 ± 0.05 |
| 1084 ± 2 | 7.561 | 2.7 | 0^+ | 1.0 |
| (1200 ± 30) | (7.66) | 250 ± 20 | (1^+) | 0.30 ± 0.10 |
| 1370 ± 20 | 7.819 | 265 ± 30 | 1^- | 0.90 ± 0.05 |
| (2070 ± 10) | (8.4) | 70 ± 10 | $(1^-, 2^-)$ | 0.43 |
| (2300) | (8.65) | ≈ 300 | $(1^+, 2^+)$ | |
| (2480) | (8.82) | | $(3^-; 1)$ | |
| 2560 | 8.89 | | $\geq 2; (1)^c$ | large |
| (4600) | (10.7) | | | |
| (5100) | (11.2) | | | |
| 6700^d | 12.6 | broad | | |

^a For references and for a listing of other reported resonances see Table 10.13 in (1979AJ01).

^b See, however, Table 10.16 and footnote ^a in Table 10.13 of (1979AJ01).

^c Resonance shape shows $l_p = 2$ formation with a large Γ_p/Γ : the contribution from the 2^+ state appears small (1977KI04).

^d Weak resonance near $E_p = 6.5$ MeV in p_0 .

16. (a) ${}^9\text{Be}(p, d){}^8\text{Be}$

$$Q_m = 0.5591$$

$$E_b = 6.5865$$

(b) ${}^9\text{Be}(p, \alpha){}^6\text{Li}$

$$Q_m = 2.126$$

Knowledge of the cross sections of these two reactions at low energies is of importance for power generation and astrophysical considerations. Absolute cross sections for the d_0 and α_0 groups have been measured for $E_p = 28$ to 697 keV with $\pm 5 - 6\%$ uncertainty. The value of $S_{\text{c.m.}}(E = 0)$ for the combined cross sections is estimated to be 35_{-15}^{+45} MeV \cdot b. At the 0.33 MeV resonance ($J^\pi = 1^-$), $\sigma_{\alpha_0} = 360 \pm 20$ mb and $\sigma_{d_0} = 470 \pm 30$ mb. The data (including angular distributions), analyzed by an R -matrix compound nucleus model, were fitted by assuming three states at $E_p(\text{c.m.}) = -20$ keV ($J^\pi = 2^+$; 3^+ possible) [$E_x = 6.57$ MeV] [see, however, Table 10.9], 310 keV (1^-) and 410 keV (1^+ ; 2^+ or 3^+ possible) (1973SI27).

Measurements of excitation functions for deuterons and α -particles have been reported at a number of energies to $E_p = 15$ MeV: see (1974AJ01, 1979AJ01). Observed resonances are displayed in Table 10.14.

Polarization measurements have been made in the range $E_p = 0.30$ to 15 MeV and at 185 MeV: see (1974AJ01, 1979AJ01). See also ${}^6\text{Li}$, ${}^8\text{Be}$ and (1979AR04, 1983SEZW).

Table 10.14: Resonances in ${}^9\text{Be}(p, d){}^8\text{Be}$ and ${}^9\text{Be}(p, \alpha){}^6\text{Li}$ ^a

| E_p (MeV) | E_x (MeV) | $\Gamma_{\text{c.m.}}$ (keV) | $J^\pi; T$ | Γ_p/Γ | θ_p^2 | θ_d^2 | θ_α^2 |
|-------------------|-------------|------------------------------|------------------------------|-------------------|---------------|--------------|-------------------|
| 0.34 | 6.89 | | $1^-; 0$ | 0.30 | 0.34 | 0.15 | 0.055 |
| 0.46 | 7.00 | | $1^+(2^+, 3^+)$ ^b | | 0.3 | 0.3 | 0.1 |
| (0.68) | (7.20) | | | | | | |
| 0.94 | 7.43 | 140 | $(2^-; 0)$ | 0.7 | 0.04 | 0.02 | |
| 1.15 | 7.62 | 225 ± 50 | $(1^+; 0)$ | ≈ 0.4 | ≈ 0.1 | | |
| 1.65 | 8.07 | 800 ± 200 | $(2^-; 0)$ | ≈ 0.07 | 0.18 | 0.21 | |
| (2.3) | (8.7) | (≈ 220) | | | | | |
| 2.56 ^c | 8.89 | 36 ± 2 | $2^+; 1$ | | | | |
| 3.5 ^d | 9.7 | | $T = 1$ | | | | |
| 4.49 ^d | 10.62 | | $T = 1$ | | | | |

^a For references and for a listing of other reported references see Table 10.14 in (1979AJ01).

^b See, however, Table 10.9.

^c (1977K104) have analyzed the $(\alpha_2\gamma)$ and p_0 yields with an R -matrix formalism and find the following parameters:

$$\left. \begin{array}{cc} 2.566 \pm 0.001 & 2^+ \\ 2.561_{-2}^{+10} & 3^- \end{array} \right\} \Gamma_{\text{c.m.}} = \left\{ \begin{array}{l} 40 \pm 1 \text{ keV} \\ 100 \pm 20 \text{ keV} \end{array} \right.$$

^d Resonance for α_2 to ${}^6\text{Li}^*(3.56)$, $J^\pi = 0^+$, $T = 1$.

17. ${}^9\text{Be}(d, n){}^{10}\text{B}$

$$Q_m = 4.3620$$

Neutron groups are observed corresponding to the ${}^{10}\text{B}$ states listed in Table 10.15. Angular distributions have been measured for $E_d = 0.5$ to 16 MeV: see (1974AJ01, 1979AJ01). Observed γ -transitions are listed in Table 10.16 of (1979AJ01). See Tables 10.6 and 10.7 here for the parameters of radiative transitions and for τ_m .

From all the various experiments the following picture emerges: the first five states of ${}^{10}\text{B}$ have even parity [from l_p]. The ground state is known to have $J = 3$, by direct measurement, and ${}^{10}\text{B}^*(1.74)$ has $J^\pi = 0^+$ and is the $T = 1$ analog of the ${}^{10}\text{C}_{\text{g.s.}}$ [from the β^+ decay of ${}^{10}\text{C}$]. Then looking at the branching ratios and lifetimes of the other states, the sequence for ${}^{10}\text{B}^*(0, 0.72, 1.74, 2.15, 3.59)$ is $J^\pi = 3^+, 1^+, 0^+, 1^+, 2^+$ [see discussion in (1966LA04, 1966WA10)].

For polarization measurements see (1981BR1E) and ${}^{11}\text{B}$ in (1980AJ01, 1985AJ01). See also (1977BA1L, 1978PL1B) and (1978BA1F, 1979WA1F, 1982OV1A, 1982SM1F; applications).

18. ${}^9\text{Be}({}^3\text{He}, d){}^{10}\text{B}$

$$Q_m = 1.0930$$

Table 10.15: Levels of ^{10}B from $^9\text{Be}(\text{d}, \text{n})$ and $^9\text{Be}({}^3\text{He}, \text{d})$ ^a

| E_x (MeV) ^a | $^9\text{Be}(\text{d}, \text{n})$ ^b | | $^9\text{Be}({}^3\text{He}, \text{d})$ ^c | | $J^\pi; T$ |
|--------------------------|--|------------------|---|----------------|-----------------|
| | l_p | S_{rel} | l_p | $(2J + 1)C^2S$ | |
| 0 | 1 | 1.0 | 1 | 3.30 | $3^+; 0$ |
| 0.72 | 1 | 1.97 | 1 | 2.76 | $1^+; 0$ |
| 1.74 | 1 | 1.42 | 1 | 1.20 | $0^+; 1$ |
| 2.15 | 1 | 0.41 | 1 | 0.82 | $1^+; 0$ |
| 3.59 | 1 | 0.10 | 1 | 0.29 | $2^+; 0$ |
| 4.77 | (≥ 2) | | 1^+ | 0.10 | $3^+; 0$ |
| | | | (3) ^d | ≤ 0.82 | |
| 5.11 | 0 | 0.14 | $0 + 2$ | 0.34, 0.14 | $2^-; 0$ |
| 5.16 | } | 1 | 1 | 0.86 | $2^+; 1$ |
| 5.18 | | | | | $1^+; 0$ |
| 5.92 | 1 | 0.49 | 1 | 2.05 | $2^+; 0$ |
| 6.03 | | | (3) ^d | ≤ 0.20 | 4^+ |
| 6.13 | (2) | | (2) ^e | 3.04 | 3^- |
| 6.56 | (3) | | (2) ^e | 2.01 | $(4)^-$ |
| 6.89 ± 15 | (1) | | | | $1^-; 0 + 1$ |
| 7.00 ± 15 | (1) | | | | $(1, 2)^+; (0)$ |
| 7.48 ± 15 | f | | | | ^g |
| 7.56 ± 25 | f | | | | $0^+; 1$ |
| (7.85 ± 50) | f | | | | 1^- |
| (8.07 ± 50) | f | | | | $(2^-; 0)$ |
| (8.12 ± 50) | f | | | | |

^a Values without uncertainties are from Table 10.5; others are from Table 10.15 in (1979AJ01). See that table for additional information and for references.

^b S_{rel} from experiment at $E_d = 12.0 - 16.0$ MeV. (1974KE06) have reanalyzed the results for $^{10}\text{B}^*(0, 1.74)$ and find S_{rel} (ave.) = 1.0 and 1.36. For values at other energies see Table 10.15 in (1979AJ01).

^c (1980BL02; $E({}^3\text{He}) = 18$ MeV); DWBA analysis; values shown are those obtained with one of the two optical model potentials used in the analysis. For earlier (${}^3\text{He}, \text{d}$) results see Table 10.17 in (1979AJ01).

^d Angular distribution poorly fitted by DWBA (1980BL02).

^e See (1980BL02) for a discussion of these two states, including a comparison with the (d, n) data: $l_p = 2$ is slightly preferred to $l_p = 1$ on the basis of observed strengths. Neither $l_p = 2$ or 1 gives a good DWBA fit.

^f State observed in (d, n) reaction; l_p not determined.

^g Group shown corresponds to unresolved states in ^{10}B .

Deuteron groups have been observed to a number of states of ^{10}B : see Table 10.15. Angular distributions have been reported at $E(^3\text{He}) = 10$ to 25 MeV, at $E(^3\text{He}) = 33.3$ MeV [see (1974AJ01, 1979AJ01)] and at $E(^3\text{He}) = 18$ MeV (1980BL02). Spectroscopic factors obtained in the (d, n) and (^3He , d) reactions are not in good agreement: see the discussions in (1974KE06, 1980BL02).

$$19. \ ^9\text{Be}(\alpha, t)^{10}\text{B} \quad Q_m = -13.2275$$

Angular distributions have been studied at $E_\alpha = 27, 28.3$ and 43 MeV [see (1979AJ01)], at $E_\alpha = 30.1$ MeV (1983VA1H; $\alpha_0, \alpha_1, \alpha_3, \alpha_4$) and at 65 MeV (1980HA33). In the latter experiment DWBA analyses have been made of the distributions to $^{10}\text{B}^*(0, 0.72, 1.74, 2.15, 3.59, 5.2, 5.92, 6.13, 6.56, 7.00, 7.5, 7.82, 8.9)$ and spectroscopic factors were derived. The distributions to $^{10}\text{B}^*(4.77, 6.03)$ could not be fitted by either DWBA or coupled-channel analyses. In general, coupled-channels calculations give a better fit to the 65 MeV data than does DWBA (1980HA33; see also for a comparison with the (d, n) and (^3He , d) results). See also (1978ZE03; theor.).

$$20. \ ^9\text{Be}(^7\text{Li}, ^6\text{He})^{10}\text{B} \quad Q_m = -3.388$$

At $E(^7\text{Li}) = 34$ MeV angular distributions have been obtained for the ^6He ions to the first four states of ^{10}B . Absolute values of the spectroscopic factors are $S = 0.88, 1.38$ ($p_{1/2}$ or $p_{3/2}$), 1.40 and 0.46 ($p_{1/2}$), 0.54 ($p_{3/2}$) for $^{10}\text{B}^*(0, 0.74, 1.74, 2.15)$ (FRDWBA analysis) (1977KE09).

$$21. \ ^{10}\text{Be}(\beta^-)^{10}\text{B} \quad Q_m = 0.5568$$

See ^{10}Be .

$$\begin{aligned} 22. \text{ (a) } & \ ^{10}\text{B}(\gamma, n)^9\text{B} & Q_m &= -8.437 \\ \text{ (b) } & \ ^{10}\text{B}(\gamma, p)^9\text{Be} & Q_m &= -6.5865 \\ \text{ (c) } & \ ^{10}\text{B}(\gamma, d)^8\text{Be} & Q_m &= -6.0274 \\ \text{ (d) } & \ ^{10}\text{B}(\gamma, \alpha)^6\text{Li} & Q_m &= -4.4603 \end{aligned}$$

Absolute measurements have been made of the $^{10}\text{B}(\gamma, \text{Tn})$ cross section from threshold to 35 MeV with quasimonoenergetic photons; the integrated cross section is 0.54 in units of the classical dipole sum ($60 NZ/A$ MeV \cdot mb). The $(\gamma, 2n) + (\gamma, 2np)$ cross section is zero, within statistics, for $E_\gamma = 16$ to 35 MeV (1976KN04). The giant resonance is broad with the major structure contained in two peaks at $E_x = 20.1 \pm 0.1$ and 23.1 ± 0.1 MeV ($\sigma_{\text{max}} \approx 5.5$ mb for each of the two maxima). For reaction (b) see (1974AJ01); for reactions (c) and (d) see (1959AJ76, 1966LA04). See also (1978DI1A, 1979TA1C) and (1981KE16, 1983GO1T; theor.).

Table 10.16: Radiative widths for $^{10}\text{B}(e, e')$ ^a

| E_x in ^{10}B (MeV) | $J^\pi; T$ | Mult. | Γ_{γ_0} (eV) |
|--------------------------------|-----------------------|----------|----------------------------------|
| 1.74 | $0^+; 1$ | M3 | $(1.05 \pm 0.25) \times 10^{-7}$ |
| 5.16 ± 0.04 ^b | $2^+; 1$ | { M1 | 0.05 ± 0.05 |
| | | { M3 | $(1.1 \pm 0.1) \times 10^{-6}$ |
| 6.03 | 4^+ | { C2 | 0.106 ± 0.005 |
| | | { C4 | $(3.3 \pm 0.8) \times 10^{-7}$ |
| 7.48 | $2^+; 1$ ^c | M1 | 11.75 ± 0.75 |
| 8.07 | 2^+ ^d | C2 | 0.19 ± 0.02 |
| 8.9 | $2^+; 1$ | { M1 | 0.3 ± 0.1 |
| | | { M3 | $(1.0 \pm 0.1) \times 10^{-5}$ |
| | $3^-; 1$ | M2 | $(1.2 \pm 0.1) \times 10^{-3}$ |
| 10.79 ^c | | M1 or C2 | |
| 11.56 ^c | | (M1) | 11.4 ± 2.3 ^c |
| 12.6 | | | |
| 13.3 | | | |

^a (1979AN08; $E_e = 67$ to 194 MeV). See also Table 10.18 in (AJ79) and (SH78N).

^b Assumed to correspond to 2^+ state at 5.16 MeV. $\Gamma_{\gamma_0} = (3.5 \pm 0.3) \times 10^{-4}$ eV for M2 if the transition were to the 2^- state at 5.11 MeV: see also footnote ^g in Table 10.18 (1979AJ01).

^c (1976FA13, 1979AN08).

^d Determined by (1979AN08); $\Gamma \approx 760$ keV.

23. (a) $^{10}\text{B}(e, e)^{10}\text{B}$

(b) $^{10}\text{B}(e, en)^9\text{B}$ $Q_m = -8.437$

(c) $^{10}\text{B}(e, ep)^9\text{Be}$ $Q_m = -6.5865$

(d) $^{10}\text{B}(e, e\pi^+)^{10}\text{Be}$ $Q_m = -140.124$

Inelastic electron groups are displayed in Table 10.16 (1979AN08). For reactions (b) and (c) see (1978SH14); for reaction (c) see (1978NA05); for reaction (d) see ^{10}Be (1982ZU03). See also (1979AJ01, 1979TI1A, 1979TR1B) and (1978BO09, 1978FU13, 1981KE15; theor.).

24. $^{10}\text{B}(\pi^\pm, \pi^\pm)^{10}\text{B}$

See (1981GE1B).

25. $^{10}\text{B}(n, n')^{10}\text{B}$

Angular distributions have been studied for $E_n = 1.5$ to 14.1 MeV: see (1974AJ01, 1979AJ01). See also ^{11}B in (1985AJ01), (1979GL1E, 1981DA1K) and (1979GL1D; theor.).

26. (a) $^{10}\text{B}(p, p')^{10}\text{B}$

(b) $^{10}\text{B}(p, 2p)^9\text{Be}$ $Q_m = -6.5865$

Angular distributions have been measured for $E_p = 3.0$ to 49.5 MeV [see (1974AJ01, 1979AJ01)] and at 6.0 (1977SA1B) and 800 MeV (1979MO1E). Table 10.17 displays the states observed in this reaction. The earlier γ -decay results are presented in (1979AJ01) and in Table 10.6 here. See also ^{11}C in (1985AJ01), (1980FA07, 1981HO13) and (1979GL1D, 1980KO1V; theor.). For reaction (b) see ^9Be and (1974AJ01).

27. $^{10}\text{B}(d, d')^{10}\text{B}$

Angular distributions have been reported at $E_d = 4$ to 28 MeV: see (1974AJ01, 1979AJ01). Observed deuteron groups are displayed in Table 10.17. The very low intensity of the group to $^{10}\text{B}^*(1.74)$ and the absence of the group to $^{10}\text{B}^*(5.16)$ is good evidence of their $T = 1$ character: see (1974AJ01).

28. $^{10}\text{B}(t, t)^{10}\text{B}$

Angular distributions of elastically scattered tritons have been measured at $E_t = 1.5$ to 3.3 MeV: see (1974AJ01).

29. $^{10}\text{B}(^3\text{He}, ^3\text{He}')^{10}\text{B}$

Angular distributions have been measured at $E(^3\text{He}) = 4$ to 32.5 MeV [see (1974AJ01, 1979AJ01)] and at 41 MeV (1980TR02; elastic) and 46.1 MeV (1979GO07). $L = 2$ gives a good fit to the distributions of ^3He ions to $^{10}\text{B}^*(0.72, 2.15, 3.59, 6.03)$: derived β_L are shown in Table 10.19 of (1979AJ01). See also Table 10.17 here.

30. (a) $^{10}\text{B}(\alpha, \alpha')^{10}\text{B}$

(b) $^{10}\text{B}(\alpha, 2\alpha)^6\text{Li}$ $Q_m = -4.4603$

Table 10.17: ^{10}B levels from $^{10}\text{B}(\text{p}, \text{p}')$, $^{10}\text{B}(\text{d}, \text{d}')$ and $^{10}\text{B}({}^3\text{He}, {}^3\text{He}')$ ^a

| E_x (MeV \pm keV) ^b | $\Gamma_{\text{c.m.}}$ (keV) | L | β_{I} ^{b,c} |
|------------------------------------|------------------------------|-----|-----------------------------------|
| 0 ^d | | | |
| 0.7183 \pm 0.4 ^{d,e,f} | | 2 | 0.67 \pm 0.05 |
| \equiv 1.7402 ^{f,g} | | (3) | |
| 2.1541 \pm 0.5 ^d | | 2 | 0.49 \pm 0.04 |
| 3.5870 \pm 0.5 ^d | | 2 | 0.45 \pm 0.04 |
| 4.7740 \pm 0.5 ^h | | | |
| 5.1103 \pm 0.6 | | 3 | 0.45 \pm 0.04 |
| 5.1639 \pm 0.6 | | | |
| 5.18 \pm 10 ^{h,i} | 110 \pm 10 | | |
| 5.9195 \pm 0.6 ^d | < 5 | | 0.28 \pm 0.03 |
| 6.0250 \pm 0.6 ^d | < 5 | 2 | 0.95 \pm 0.04 |
| 6.1272 \pm 0.7 ^d | < 5 | 3 | 0.58 \pm 0.03 |
| 6.55 \pm 10 ^d | 25 \pm 5 | 3 | 0.46 \pm 0.04 ^j |
| 7.00 \pm 10 ^d | 95 \pm 10 | | |
| 7.48 \pm 10 | 90 \pm 15 | | |

^a For references and a more complete presentation see Table 10.19 in (1979AJ01).

^b From (p, p').

^c See results obtained from (${}^3\text{He}$, ${}^3\text{He}'$) in Table 10.19 of (1979AJ01).

^d Also observed in (d, d') and (${}^3\text{He}$, ${}^3\text{He}'$).

^e $E_x = 718.35 \pm 0.04$ (1979RI12; from E_γ).

^f $E_x = 718.5 \pm 0.2$ and 1740.0 ± 0.6 keV (1966FR09; from E_γ).

^g Also observed in (${}^3\text{He}$, ${}^3\text{He}'$).

^h Also observed in (d, d').

ⁱ Not reported in (p, p') at $E_p = 10$ MeV.

^j If $J^\pi = 4^-$; $\beta_{\text{L}} = 0.59 \pm 0.03$ if $J^\pi = 2^-$.

Angular distributions have been measured for $E_\alpha = 5$ to 56 MeV [see (1974AJ01, 1979AJ01)] and at $E_\alpha = 31.2$ MeV (1981KO1U; α_0). $^{10}\text{B}^*(1.74)$ is not observed. S_α for $^{10}\text{B}_{\text{g.s.}} = 0.16$ (1976WO11).

Reaction (b) has been studied at $E_\alpha = 24$ MeV [see (1979AJ01)] and at 700 MeV (1979DO04). Using a width parameter of 141 MeV/c, (1979DO04) find that the effective number of $\alpha + \text{d}$ clusters for $^{10}\text{B}_{\text{g.s.}}$, $n_{\text{eff}} = 1.19 \pm 0.23$; the results are very model dependent. See also (1978ZE03, 1981LA13; theor.).

31. (a) $^{10}\text{B}(^6\text{Li}, ^6\text{Li})^{10}\text{B}$
 (b) $^{10}\text{B}(^7\text{Li}, ^7\text{Li})^{10}\text{B}$

Elastic scattering angular distributions have been studied at $E(^6\text{Li}) = 5.8$ and 30 MeV and at $E(^7\text{Li}) = 24$ MeV: see (1979AJ01).

32. $^{10}\text{B}(^9\text{Be}, ^9\text{Be})^{10}\text{B}$

Elastic scattering angular distributions have been reported at $E(^{10}\text{B}) = 20.1$ and 30.0 MeV (1980BO14).

33. (a) $^{10}\text{B}(^{10}\text{B}, ^{10}\text{B})^{10}\text{B}$
 (b) $^{10}\text{B}(^{11}\text{B}, ^{11}\text{B})^{10}\text{B}$

Elastic angular distributions (reaction (a)) have been studied at $E(^{10}\text{B}) = 8, 13$ and 21 MeV. For yields and reaction (b) see (1979AJ01). See also (1979SH22) and (1978TA1B; theor.).

34. (a) $^{10}\text{B}(^{12}\text{C}, ^{12}\text{C})^{10}\text{B}$
 (b) $^{10}\text{B}(^{13}\text{C}, ^{13}\text{C})^{10}\text{B}$

Elastic angular distributions have been measured at $E(^{10}\text{B}) = 18$ and 100 MeV [see (1979AJ01)] for reaction (a) and at 18, 25, 32, 39 and 46 MeV for reaction (b) (1982MA20). For fusion measurements see (1979AJ01) and (1981MA18, 1982MA20). See also (1978VA1A, 1981DE13, 1982HA42; theor.).

35. $^{10}\text{B}(^{14}\text{N}, ^{14}\text{N})^{10}\text{B}$

Angular distributions are reported at $E(^{10}\text{B}) = 100$ MeV [see (1979AJ01)] and at $E(^{14}\text{N}) = 73.9$ and 93.6 MeV (1979MO14; to $^{10}\text{B}^*(0, 0.72, 2.15)$). For fusion cross-section measurements see (1979AJ01) and (1980PA19, 1982BE54, HO82F, 1982HO1F, 1982OR02). See also (1978TA1B) and (1981AB1A; theor.).

36. (a) $^{10}\text{B}(^{16}\text{O}, ^{16}\text{O})^{10}\text{B}$
 (b) $^{10}\text{B}(^{17}\text{O}, ^{17}\text{O})^{10}\text{B}$
 (c) $^{10}\text{B}(^{18}\text{O}, ^{18}\text{O})^{10}\text{B}$

Elastic angular distribution (reaction (a)) have been studied at $E(^{16}\text{O}) = 15.0$ to 32.5 MeV and at $E(^{10}\text{B}) = 100$ MeV [see (1979AJ01)] as well as at $E(^{10}\text{B}) = 33.7, 41.6, 49.5$ MeV (1980PA01) and 65.8 MeV (1977MO1A, 1979MO14). The ground-state quadrupole moment of ^{10}B is observed to influence the scattering (1980PA01). The elastic scattering for reaction (c) has been studied at $E(^{18}\text{O}) = 20, 24$ and 30.5 MeV: see (1974AJ01). For fusion cross-section measurements and excitation functions see (1979GO09, 1981TH1A) for reaction (a) and (1980WI09, 1982CH07) for reaction (b). See also (1981ST1P) and (1979HU1B, 1980VA03, 1981VA1E, 1983CI08; theor.).

37. (a) $^{10}\text{B}(^{19}\text{F}, ^{19}\text{F})^{10}\text{B}$
 (b) $^{10}\text{B}(^{20}\text{Ne}, ^{20}\text{Ne})^{10}\text{B}$

The elastic scattering has been investigated for $E(^{19}\text{F}) = 20$ and 24 MeV [see (1974AJ01)] (reaction (a)) and for $E(^{10}\text{B}) = 65.9$ MeV (1979MO14) (reaction (b)).

38. (a) $^{10}\text{B}(^{24}\text{Mg}, ^{24}\text{Mg})^{10}\text{B}$
 (b) $^{10}\text{B}(^{25}\text{Mg}, ^{25}\text{Mg})^{10}\text{B}$

The elastic scattering for both reactions has been studied at $E(^{10}\text{B}) = 87.4$ MeV (1982FU09).

39. (a) $^{10}\text{B}(^{27}\text{Al}, ^{27}\text{Al})^{10}\text{B}$
 (b) $^{10}\text{B}(^{28}\text{Si}, ^{28}\text{Si})^{10}\text{B}$
 (c) $^{10}\text{B}(^{30}\text{Si}, ^{30}\text{Si})^{10}\text{B}$

The elastic scattering for all three reactions has been studied at $E(^{10}\text{B}) = 41.6$ and ≈ 50 MeV (1979PA09; also 33.7 MeV for reaction (b)). See also (1980GL03).

40. $^{10}\text{B}(^{40}\text{Ca}, ^{40}\text{Ca})^{10}\text{B}$

The elastic scattering angular distribution has been measured for $E(^{10}\text{B}) = 46.6$ MeV (1980GL03).

41. $^{10}\text{C}(\beta^+)^{10}\text{B}$ $Q_m = 3.6488$

The half-life of ^{10}C is 19.255 ± 0.53 sec (1975HA45) [and see (1974AJ01, 1979AJ01)]: the decay is to $^{10}\text{B}^*(0.72, 1.74)$ with branching ratios of $(98.53 \pm 0.02)\%$ and $(1.465 \pm 0.014)\%$ and $\log ft = 3.047$ for the transition to $^{10}\text{B}^*(0.72)$ and 3.492 ± 0.005 for that to the analog state, $^{10}\text{B}^*(1.74)$: see Table 10.20 in (1979AJ01). The excitation energies of the two states are 718.32 ± 0.09 and 1740.16 ± 0.17 keV [$E_\gamma = 718.29 \pm 0.09$ and 1021.78 ± 0.14 keV] (1969FR02). See (1979AJ01) for a further discussion of the decay. See also (1979DE15, 1979FE02, 1979KU05; theor.).

$$42. \quad ^{11}\text{B}(\gamma, n)^{10}\text{B} \qquad Q_m = -11.454$$

The intensities of the transitions to $^{10}\text{B}^*(3.59, 5.16)$ [$T = 0$ and 1 , respectively] depend on the region of the giant dipole resonances in ^{11}B from which the decay takes place: it is suggested that the lower-energy region consists mainly of $T = \frac{1}{2}$ states and the higher-energy region of $T = \frac{3}{2}$ states: see ^{11}B in (1980AJ01). (1979BR1D) report observation of the 1.02 MeV line from the decay of $^{10}\text{B}^*(1.74)$. See also ^{11}B in (1985AJ01) and (1982GO03, 1983GO1T; theor.).

$$43. \quad ^{11}\text{B}(p, d)^{10}\text{B} \qquad Q_m = -9.230$$

Angular distributions of deuteron groups have been measured at several energies in the range $E_p = 17.7$ to 154.8 MeV: see (1979AJ01). The population of the first five states of ^{10}B and $^{10}\text{B}^*(5.18(u), 6.04(u), 6.56, 7.5, 11.4 \pm 0.2, 14.1 \pm 0.2)$ is reported. For VAP measurements see (1982BU03) in ^{12}C (1985AJ01).

$$44. \quad ^{11}\text{B}(d, t)^{10}\text{B} \qquad Q_m = -5.197$$

Angular distributions have been measured at $E_d = 11.8$ MeV ($t_0 \rightarrow t_3$; $l = 1$; $S = 1.88, 0.94, 1.35, 1.35$, respectively): see (1974AJ01).

$$45. \quad \begin{aligned} \text{(a)} \quad ^{11}\text{B}(^3\text{He}, \alpha)^{10}\text{B} & \qquad Q_m = 9.124 \\ \text{(b)} \quad ^{11}\text{B}(^3\text{He}, 2\alpha)^6\text{Li} & \qquad Q_m = 4.663 \end{aligned}$$

Reported levels are displayed in Table 10.18. Angular distributions have been measured at a number of energies between $E(^3\text{He}) = 1.0$ and 33 MeV: see (1974AJ01). For the decay of observed states see Table 10.6.

The $\alpha\alpha$ angular correlations (reaction (b)) have been measured for the transitions via $^{10}\text{B}^*(5.92, 6.03, 6.13, 6.56, 7.00)$. The results are consistent with $J^\pi = 2^+$ and 4^+ for $^{10}\text{B}^*(5.92, 6.03)$ and require $J^\pi = 3^-$ for $^{10}\text{B}^*(6.13)$. There is substantial interference between levels of opposite parity for the α -particles due to $^{10}\text{B}^*(6.56, 7.00)$: the data are fitted by $J^\pi = 3^+$ for $^{10}\text{B}^*(7.00)$ and $(3, 4)^-$ for $^{10}\text{B}^*(6.56)$ [the $^6\text{Li}(\alpha, \alpha)$ results then require $J^\pi = 4^-$]. See however reaction 16, and see (1974AJ01) for the references.

Table 10.18: ^{10}B levels from $^{11}\text{B}(^3\text{He}, \alpha)^{10}\text{B}$ ^a

| E_x (MeV \pm keV) | $\Gamma_{\text{c.m.}}$ (keV) | l | S_{rel} |
|-----------------------|------------------------------|-----|------------------|
| 0 | | 1 | 1.0 |
| 0.718 \pm 7 | | 1 | 0.22 |
| 1.744 \pm 7 | | 1 | 0.73 |
| 2.157 \pm 6 | | 1 | 0.44 |
| 3.587 \pm 6 | | 1 | 0.09 |
| 4.777 \pm 5 | | 1 | 0.09 |
| 5.114 \pm 5 | | | |
| 5.166 \pm 5 | | 1 | 1.81 |
| 5.923 \pm 5 | | | |
| 6.028 \pm 5 | | | |
| 6.131 \pm 5 | | | |
| 6.570 \pm 7 | 30 \pm 10 | | |
| 7.002 \pm 10 | 95 \pm 10 | | |
| 7.475 \pm 10 | | | |
| 7.567 \pm 10 | | | |
| 7.87 \pm 40 | 240 \pm 50 | | |
| 10.85 \pm 100 | 300 \pm 100 | | |
| 11.52 \pm 35 | 500 \pm 100 | | |
| 12.56 \pm 30 | 100 \pm 30 | | |
| 13.49 \pm 50 | 300 \pm 50 | | |
| 14.4 \pm 100 | 800 \pm 200 | | |
| (18.2 \pm 200) | (1500 \pm 300) | | |

^a See Table 10.21 in (1979AJ01) for references.

46. $^{12}\text{C}(\gamma, \text{pn})^{10}\text{B}$ $Q_{\text{m}} = -27.4104$

See (1982DO08), ^{12}C in (1985AJ01) and (1981KH08; theor.).

47. $^{12}\text{C}(\pi^{\pm}, \pi^{\pm}\text{d})^{10}\text{B}$ $Q_{\text{m}} = -25.1858$

At $E_{\pi^+} = 180$ MeV and $E_{\pi^-} = 220$ MeV, $^{10}\text{B}^*(0.72, 2.15)$ are populated (1981ST05). See also (1979EL12, 1982EL07).

48. (a) $^{12}\text{C}(\text{p}, ^3\text{He})^{10}\text{B}$ $Q_{\text{m}} = -19.6923$

(b) $^{12}\text{C}(\text{p}, \text{pd})^{10}\text{B}$ $Q_{\text{m}} = -25.1858$

Angular distributions of ^3He ions have been measured for $E_{\text{p}} = 39.8, 51.9$ and 185 MeV: see (1979AJ01). $^{10}\text{B}^*(0, 0.72, 1.74, 2.15, 3.59, 4.77, 5.16, 5.92, 6.56, 7.50, 8.90)$ are populated. For reaction (b) see (1981ER10; 670 MeV) and (1979AJ01). See also (1978GO14; theor.).

49. $^{12}\text{C}(\text{d}, \alpha)^{10}\text{B}$ $Q_{\text{m}} = -1.3391$

Alpha groups have been observed to most of the known states of ^{10}B below $E_{\text{x}} = 7.1$ MeV: see Table 10.23 in (1974AJ01). Angular distributions have been measured for $E_{\text{d}} = 5.0$ to 40 MeV: see (1979AJ01). Single-particle S -values are $1.5, 0.5, 0.1, 0.1$ and 0.3 for $^{10}\text{B}^*(0, 0.72, 2.15, 3.59, 4.77)$ (1976VA07; ZRD-WBA). A study of the $m_{\text{s}} = 0$ yield at $E_{\text{d}} = 14.5$ MeV ($\theta = 0^\circ$) leads to assignments of $3^+, 2^-$ and $(3^+, 4^-)$ for $^{10}\text{B}^*(4.77, 5.11, 6.56)$ (1975KU15). VAP measurements are reported at $E_{\text{d}} = 52$ MeV (1982MA25): see ^{14}N in (1986AJ01).

The population of the isospin forbidden group to $^{10}\text{B}^*(1.74) [\alpha_2]$ has been studied with E_{d} up to 30 MeV: see ^{14}N in (1976AJ04). See also (1981JO02, 1982BA24; theor.).

50. $^{12}\text{C}(\alpha, ^6\text{Li})^{10}\text{B}$ $Q_{\text{m}} = -23.7105$

Angular distributions have been reported at $E_{\alpha} = 42$ and 46 MeV: see (1979AJ01). At $E_{\alpha} = 65$ MeV, an investigation of the ^6Li breakup shows that $^{10}\text{B}^*(0, 0.72, 2.16, 3.57, 4.77, 5.2, 5.9, 6.0)$ are involved (1978SA26).

51. $^{12}\text{C}(^{10}\text{B}, 3\alpha)^{10}\text{B}$ $Q_{\text{m}} = -7.2747$

See (1978BE1G).

52. (a) $^{12}\text{C}(^{12}\text{C}, ^{14}\text{N})^{10}\text{B}$ $Q_m = -14.9134$
(b) $^{12}\text{C}(^{14}\text{N}, ^{16}\text{O})^{10}\text{B}$ $Q_m = -4.4495$

Angular distributions (reaction (a)) involving $^{10}\text{B}^*(0, 0.7)$ have been studied at $E(^{12}\text{C}) = 49.0$ to 75.5 MeV (1980CO10, 1979CL06) and 93.8 MeV (1979FU04). Angular distributions (reaction (b)) involving $^{10}\text{B}^*(0, 0.72, 2.15, 3.59)$ have been measured at $E(^{14}\text{N}) = 53$ MeV [see (1979AJ01)] and 78.8 MeV (1979MO14; not to $^{10}\text{B}^*(3.59)$).

53. $^{13}\text{C}(\text{p}, \alpha)^{10}\text{B}$ $Q_m = -4.0609$

Angular distributions have been measured for $E_p = 5.8$ to 18 MeV and 43.7 and 50.5 MeV: see (1979AJ01). Polarization measurements are reported at $E_p = 65$ MeV (1980KA03; p_0): see ^{14}N in (1986AJ01).

54. $^{14}\text{N}(\text{p}, \text{p}\alpha)^{10}\text{B}$ $Q_m = -11.6115$

See (1978GO14; theor.).

55. $^{14}\text{N}(\text{d}, ^6\text{Li})^{10}\text{B}$ $Q_m = -10.136$

At $E_d = 80$ MeV angular distributions are reported to $^{10}\text{B}^*(0, 0.72, 2.15, 3.59, 4.8, 6.04, 7.05, 8.68)$ (1979OE01; see for S_α).

56. $^{14}\text{N}(^3\text{He}, ^7\text{Be})^{10}\text{B}$ $Q_m = -10.024$

At $E(^3\text{He}) = 41$ MeV groups to $^{10}\text{B}^*(0, 0.72, 2.15, 3.59, 6.1)$ have been observed. The transition to $^{10}\text{B}^*(1.74)$ is very weak: see (1979AJ01).

57. $^{16}\text{O}(^{14}\text{N}, ^{20}\text{Ne})^{10}\text{B}$ $Q_m = -6.877$

At $E(^{14}\text{N}) = 76.2$ MeV angular distributions involving $^{10}\text{B}^*(0, 0.7)$ are presented by (1979MO14).

¹⁰C
(Figs. 21 and 22)

GENERAL: (See also (1979AJ01).)

Model calculations: (1981DE2G, 1982SA1U).

Electromagnetic transitions: (1982RI04).

Astrophysical questions: (1979MO04, 1979RA1C).

Complex reactions involving ¹⁰C: (1979BO22, 1981MO20).

Reactions involving pions (See also reactions 2 and 6.): (1979AL1J, 1979LI1D, 1980LE02, 1981AU1C, 1982COZV, 1982RO04).

Other topics: (1979NO1B, 1980NO1A, 1982DE1N, 1982NG01).

Mass of ¹⁰C: Based on Q_0 for ¹⁰B(p, n)¹⁰C and ¹²C(p, t)¹⁰C and the Wapstra masses for n, t and ¹⁰B, the atomic mass excess of ¹⁰C is 15698.8 ± 0.5 keV.

1. ¹⁰C(β^+)¹⁰B $Q_m = 3.6488$

¹⁰C decays with a half-life of 19.255 ± 0.053 sec to ¹⁰B*(0.7, 1.7): the branching ratios are $(98.53 \pm 0.02)\%$ and $(1.465 \pm 0.014)\%$, respectively (1972RO03): see reaction 41 in ¹⁰B.

2. ⁹Be(p, π^-)¹⁰C $Q_m = -136.6296$

Angular distributions of π^- groups have been measured at $E_p = 185$ MeV (1973DA09; to ¹⁰C*(0, 3.36, 5.28, 6.63)), at $E_p = 200$ MeV (1980SJ02; ¹⁰C_{g.s.}) and at $E_p = 800$ MeV (1979HO13; to ¹⁰C*(0, 3.35, 5.3, 6.6)). A_y measurements have been reported at $E_p = 200$ MeV (1980SJ02; ¹⁰C_{g.s.}) and 200, 225 and 250 MeV (1982LO02; ¹⁰C*(0, 3.35, 5.28, 6.6)). See also (1979AJ01, 1979JO1C, 1979ME2A, 1981AU1C, 1981BO1D, 1982HO1C, 1982NA1K).

3. ¹⁰B(γ , π^-)¹⁰C $Q_m = -143.2161$

See (1982RO04).

4. ¹⁰B(p, n)¹⁰C $Q_m = -4.411$

Table 10.19: Energy levels of ^{10}C

| E_x (MeV \pm keV) | $J^\pi; T$ | τ or $\Gamma_{\text{c.m.}}$ (keV) | Decay | Reactions |
|-----------------------|--------------|--|-----------|------------------------|
| g.s. | $0^+; 1$ | $\tau_{1/2} = 19.255 \pm 0.053$ sec | β^+ | 1, 2, 3, 4, 5, 6, 7, 8 |
| 3.3536 ± 0.9 | 2^+ | $\tau_m = 155 \pm 25$ fsec | γ | 2, 4, 5, 6, 7, 8 |
| 5.22 ± 40 | ^a | $\Gamma = 225 \pm 45$ keV | | 2, 4, 5, 7 |
| 5.38 ± 70 | ^a | 300 ± 60 | | 2, 4, 5, 7 |
| 6.580 ± 20 | (2^+) | 200 ± 40 | | 2, 5, 7 |

^a One of these two states is presumably a 2^+ state.

$$Q_0 = -4430.17 \pm 0.34 \text{ keV: P.H. Barker (private communication).}$$

The E_x of $^{10}\text{C}^*(3.35) = 3352.7 \pm 1.5$ keV, $\tau_m = 155 \pm 25$ fsec, $\Gamma_\gamma = 4.25 \pm 0.69$ meV. Angular distributions have been measured for n_0 and n_1 groups and for the neutrons to $^{10}\text{C}^*(5.2 \pm 0.3)$ at $E_p = 30$ and 50 MeV: see (1974AJ01, 1979AJ01).

$$5. \ ^{10}\text{B}(^3\text{He}, t)^{10}\text{C} \quad Q_m = -3.6674$$

Angular distributions have been measured at $E(^3\text{He}) = 14$ MeV and 217 MeV: see (1979AJ01). The latter [to $^{10}\text{C}^*(0, 3.35, 5.6)$] have been compared with microscopic calculations using a central + tensor interaction [$J^\pi = 0^+, 2^+, 2^+$] (1976WI05). Structures have been reported at $E_x = 5.22 \pm 0.04$ [$\Gamma = 225 \pm 45$ keV], 5.38 ± 0.07 [300 ± 60 keV] and 6.580 ± 0.020 MeV [190 ± 35 keV] (1975SC27). [It is not clear which of the 5.2–5.4 MeV states is the 2^+ state studied by (1976WI05).]

$$6. \ ^{12}\text{C}(\pi^+, d)^{10}\text{C} \quad Q_m = 110.7327$$

See (1982DO01).

$$7. \ ^{12}\text{C}(p, t)^{10}\text{C} \quad Q_m = -23.3597$$

$$Q_0 = -23360.74 \pm 0.5 \text{ keV: J.A. Nolen (private communication).}$$

Angular distributions have been reported at $E_p = 30.0$ to 54.1 MeV [see (1974AJ01, 1979AJ01)] and at $E_p = 80$ MeV (1979SH09; t to $^{10}\text{C}^*(0, 3.35, 5.28)$. $L = 0, 2$ and 2 thus leading to $0^+, 2^+$ and 2^+ for these states [but note that the “5.28 MeV” states is certainly unresolved]: see reaction 5 and Table 10.19. $^{10}\text{C}^*(6.6)$ is also populated. The excitation energy of $^{10}\text{C}^*(3.4)$ is 3353.3 ± 1.0 keV (1974BE66), 3354.1 ± 1.1 keV (1978RO08) [based on Q_m]. See also (1979NO1B, 1980NO1A) and (1982BE1Z; theor.).

8. $^{13}\text{C}(^3\text{He}, ^6\text{He})^{10}\text{C}$ $Q_m = -15.235$

At $E(^3\text{He}) = 70.3$ MeV the angular distributions of the ^6He ions corresponding to the population of $^{10}\text{C}^*(0, 3.35)$ have been measured. The group to $^{10}\text{C}^*(3.35)$ is much more intense than the ground-state group: multi-step processes may be important (1973KA16). (1976DE27) suggest, on the basis of an FRDWBA analysis, that the process can be interpreted as a direct-cluster transfer to both final states.

$^{10}\text{N}, ^{10}\text{O}, ^{10}\text{F}, ^{10}\text{Ne}$
(Not illustrated)

Not observed: see (1979AJ01). A.H. Wapstra (private communication) suggests 39.5 MeV for the atomic mass excess of ^{10}N . See also (1982NG01; theor.).

References

(Closed 1 June 1983)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1966FO05 P.D. Forsyth, H.T. Tu and W.F. Hornyak, Nucl. Phys. 82 (1966) 33
- 1966FR09 J.M. Freeman, J.G. Jenkin and G. Murray, Phys. Lett. 22 (1966) 177
- 1966GR18 R.C. Greenwood, Phys. Lett. 23 (1966) 482
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966WA10 E.K. Warburton, J.W. Olness, K.W. Jones, C. Chasman, R.A. Ristinen and D.H. Wilkinson, Phys. Rev. 148 (1966) 1072
- 1969AL17 D.E. Alburger, E.K. Warburton, A. Gallmann and D.H. Wilkinson, Phys. Rev. 185 (1969) 1242
- 1969FR02 J.M. Freeman, J.G. Jenkin and G. Murray, Nucl. Phys. A124 (1969) 393
- 1969RO12 M.L. Roush, F.C. Young, P.D. Forsyth and W.F. Hornyak, Nucl. Phys. A128 (1969) 401
- 1972RO03 D.C. Robinson, J.M. Freeman and T.T. Thwaites, Nucl. Phys. A181 (1972) 645
- 1973DA09 S. Dahlgren, P. Grafstrom, B. Hoistad and A. Asberg, Nucl. Phys. A204 (1973) 53
- 1973KA16 E. Kashy, W. Benenson, I.D. Proctor, P. Hauge and G. Bertsch, Phys. Rev. C7 (1973) 2251
- 1973SI27 A.J. Sierk and T.A. Tombrello, Nucl. Phys. A210 (1973) 341
- 1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227 (1974) 1
- 1974AN27 R.E. Anderson, J.J. Kraushaar, M.E. Rickey and W.R. Zimmerman, Nucl. Phys. A236 (1974) 77
- 1974BE66 W. Benenson and E. Kashy, Phys. Rev. C10 (1974) 2633
- 1974FI1D Fick, Nukleonika 19 (1974) 693
- 1974KE06 K.W. Kemper, S. Cotanch, G.E. Moore, A.W. Obst, R.J. Puigh and R.L. White, Nucl. Phys. A222 (1974) 173
- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
- 1975BA52 H.W. Baer, J.A. Bistirlich, N. de Botton, S. Cooper, K.M. Crowe, P. Truol and J.D. Vergados, Phys. Rev. C12 (1975) 921

- 1975BL07 R.C. Block, Y. Fujita, K. Kobayashi and T. Oosaki, *J. Nucl. Sci. Tech.* 12 (1975) 1
- 1975HA45 J.C. Hardy and I.S. Towner, *Nucl. Phys. A254* (1975) 221
- 1975HO01 R.J. Holt, F.W.K. Firk, G.T. Hickey and R. Nath, *Nucl. Phys. A237* (1975) 111
- 1975KU15 J.A. Kuehner, P.W. Green, G.D. Jones and D.T. Petty, *Phys. Rev. Lett.* 35 (1975) 423
- 1975SC27 M.J. Schneider, B.W. Ridley, M.E. Rickey, J.J. Kraushaar and W.R. Zimmerman, *Phys. Rev. C12* (1975) 335
- 1975SC41 U. Schwinn, G. Mairle, G.J. Wagner and C. Ramer, *Z. Phys. A275* (1975) 241
- 1975WA06 P. Wagner, R.M. Freeman, A. Gallman and E.K. Warburton, *Phys. Rev. C11* (1975) 1459
- 1975WI26 K.H. Wilcox, R.B. Weisenmiller, G.J. Wozniak, N.A. Jelley, D. Ashery and J. Cerny, *Phys. Lett. B59* (1975) 142
- 1976AJ04 F. Ajzenberg-Selove, *Nucl. Phys. A268* (1976) 1
- 1976DA15 S.E. Darden, G. Murillo and S. Sen, *Nucl. Phys. A266* (1976) 29
- 1976DE27 G. Delic and D. Kurath, *Phys. Rev. C14* (1976) 619
- 1976FA13 L.W. Fagg, R.A. Lindgren, W.L. Bendel and E.C. Jones, Jr., *Phys. Rev. C14* (1976) 1727
- 1976KN04 U. Kneissl, K.H. Leister, H.O. Neidel and A. Weller, *Nucl. Phys. A264* (1976) 30
- 1976LI08 P.W. Lisowski, R.L. Walter, C.E. Busch and T.B. Clegg, *Nucl. Phys. A264* (1976) 188
- 1976VA07 A. van der Woude and R.J. de Meijer, *Nucl. Phys. A258* (1976) 199
- 1976WI05 N. Willis, I. Brissaud, L. Bimbot, Y. Le Bornec and B. Tatischeff, *Nucl. Phys. A261* (1976) 45
- 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, *Phys. Rev. C14* (1976) 815
- 1977BA1L Badovskii et al, *Sov. At. Energy* 43 (1977) 838
- 1977KE09 K.W. Kemper, G.E. Moore, R.J. Puigh and R.L. White, *Phys. Rev. C15* (1977) 1726
- 1977KI04 A. Kiss, E. Koltay, G. Szabo and L. Vegh, *Nucl. Phys. A282* (1977) 44
- 1977MO1A Motobayashi, *Rep. Inst. Phys. Chem. Res.* 53 (1977) 81
- 1977SA1B Sadkovskii, Feofilov, Denisov and Kolalis, *Yad. Fiz.* 26 (1977) 1200
- 1977TU02 A. Turkevich, J.R. Cadieux, J. Warren, T. Economou, J. La Rosa and H.R. Heydegger, *Phys. Rev. Lett.* 38 (1977) 1129
- 1977TU1B A. Turkevich, J.R. Cadieux, J. Warren, T. Economou and J. LaRosa, *Phys. Lett. B72* (1977) 163
- 1978AN20 M.K. Anikina, G.L. Vardenga, A.I. Golokhvastov, M.S. Zhuravleva, V.L. Ilina, E.S. Kuznetsova, Y. Lukstinsh, E.O. Okonov, T.G. Ostanovich, S.A. Khorozov et al, *Yad. Fiz.* 27 (1978) 724; *Sov. J. Nucl. Phys.* 27 (1978) 387

- 1978AR1J Arefev et al, *Yad. Fiz.* 28 (1978) 1534
- 1978AU07 E.G. Auld, A. Haynes, R.R. Johnson, G. Jones, T. Masterson, E.L. Mathie, D. Ottewell, P. Walden and B. Tatischeff, *Phys. Rev. Lett.* 41 (1978) 462
- 1978BA1F Barschall, *Ann. Rev. Nucl. Part. Sci.* 28 (1978) 207
- 1978BE1G Berezhnoi, Klyucharev and Rutkevich, *Ukr. Fiz. Zh.* 23 (1978) 1841
- 1978BH03 R.K. Bhowmik, E.C. Pollacco, N.E. Sanderson, J.B.A. England and G.C. Morrison, *Phys. Lett.* B80 (1978) 41
- 1978BH1B Bhakar and Van Oers, in *Few Body Syst. Nucl. Forces, Graz 1978, Springer Lecture Notes* 87 (1978) 364
- 1978BO09 M.C. Bouten and M. Bouten, *Nucl. Phys.* A299 (1978) 141
- 1978BU1B Buffington, Orth and Mast, *Astrophys. J.* 226 (1978) 355
- 1978DA1A Dalitz and Gal, *Ann. Phys.* 116 (1978) 167
- 1978DI1A Di Napoli et al, *J. Inorg. Nucl. Chem.* 40 (1978) 1973
- 1978DU1B Dubna-Warsaw-Leningrad Collaboration, *Yad. Fiz.* 27 (1978) 1246
- 1978FU13 A.V. Fursaev, V.K. Tartakovskij and A.A. Pasichny, *Ukr. Fiz. Zh.* 23 (1978) 827
- 1978GO14 N.F. Golovanova and I.M. Ilin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 42 (1978) 1528; *Bull. Acad. Sci. USSR Phys. Ser.* 42 (1978) 154
- 1978HE1C Hendrie, in *Clustering, Winnipeg 1978, AIP Conf. Proc.* 47 (1978) 402
- 1978HE1D Heitzmann, *Atomkernenergie* 31 (1978) 262
- 1978HE1F Hendricks, *Trans. Amer. Nucl. Soc.* 30 (1978) 732
- 1978HO23 H.H. Hogue, P.L. Von Behren, D.H. Epperson, S.G. Glendinning, P.W. Lisowski, C.E. Nelson, H.W. Newson, F.O. Purser, W. Tornow, C.R. Gould et al, *Nucl. Sci. Eng.* 68 (1978) 38
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, *Table of Isotopes 7th ed.* (New York: John Wiley & Sons, 1978)
- 1978MA1F Malecki, Namyslowski, Reale and Minetti, *Riv. Nuovo Cim.* 1 (1978) 1
- 1978MU1B Muller, *Phys. Today*, Feb. (1969) 23
- 1978NA05 K. Nakamura, S. Hiramatsu, T. Kamae, H. Muramatsu, N. Izutsu and Y. Watase, *Nucl. Phys.* A296 (1978) 431
- 1978PL1B Plattard et al, *Conf. Int. Sur La Physique Neutronique Et Les Donnees Nucleaires Pour Les Reacteurs Et Autres Applications-Harwell (G.B.)* (1978)
- 1978PO1A B. Povh, *Ann. Rev. Nucl. Part. Sci.* 28 (1978) 1
- 1978RA1C Raisbeck et al, *Nature* 275 (1978) 731

- 1978RO08 R.G.H. Robertson, T.L. Khoo, G.M. Crawley, A.B. McDonald, E.G. Adelberger and S.J. Freedman, Phys. Rev. C17 (1978) 1535
- 1978SA26 A. Saha, R. Kamermans, J. van Driel and H.P. Morsch, Phys. Lett. B79 (1978) 363
- 1978SH14 N.G. Shevchenko, A.Y. Buki, B.V. Mazanko, V.N. Polishchuk and A.A. Khomich, Yad. Fiz. 28 (1978) 12; Sov. J. Nucl. Phys. 28 (1978) 5
- 1978SM02 Y.F. Smirnov and G.F. Filippov, Yad. Fiz. 27 (1978) 73; Sov. J. Nucl. Phys. 27 (1978) 39
- 1978SO1A Sokol, Izv. Akad. Nauk SSSR Ser. Fiz. 42 (1978) 1829
- 1978SU02 K.M. Subotic, B. Lalovic and B.Z. Stepancic, Nucl. Phys. A296 (1978) 141
- 1978TA1B Taras, in Clustering, Winnipeg, AIP Conf. Proc. 47 (1978) 234
- 1978TS1A Tsangarides, Wills and Bent, Bull. Amer. Phys. Soc. 23 (1978) 952
- 1978TU06 P.P. Tung, K.A. Erb, M.W. Sachs, G.B. Sherwood, R.J. Ascutto and D.A. Bromley, Phys. Rev. C18 (1978) 1663
- 1978VA1A L.C. Vaz and J.M. Alexander, Phys. Rev. C18 (1978) 2152
- 1978WA1B Walter, in Clustering, Winnipeg 1978, AIP Conf. Proc. 47 (1978) 444
- 1978ZE03 N.S. Zelenskaya, V.M. Lebedev and T.A. Yushchenko, Yad. Fiz. 28 (1978) 90; Sov. J. Nucl. Phys. 28 (1978) 44
- 1979AB11 S.N. Abramovich, B.Y. Guzhovsky, A.G. Zvenigorodsky and S.V. Trusillo, Yad. Fiz. 30 (1979) 1276; Sov. J. Nucl. Phys. 30 (1979) 665
- 1979AD1A S.K. Adhikari, Phys. Rev. C19 (1979) 325
- 1979AD1B Adib et al, Bull. Amer. Phys. Soc. 24 (1979) 864
- 1979AJ01 F. Ajzenberg-Selove, Nucl. Phys. A320 (1979) 1
- 1979AL1J Alster and Warszawski, Phys. Rept. 52 (1979) 87
- 1979AL1M Alder et al, 1978 RPI Symp. Photopion Nucl. Phys., (Plenum Press, New York, 1979) 101
- 1979AL21 V.V. Alizade, A.V. Kuptsov, V.P. Kurochkin, L.L. Nemenov, G.I. Smirnov and D.M. Khazins, Yad. Fiz. 30 (1979) 363; Sov. J. Nucl. Phys. 30 (1979) 187
- 1979AL22 Y. Alhassid, R.D. Levine, J.S. Karp and S.G. Steadman, Phys. Rev. C20 (1979) 1789
- 1979AN08 E.J. Ansaldo, J.C. Bergstrom, R. Yen and H.S. Caplan, Nucl. Phys. A322 (1979) 237; Erratum Nucl. Phys. A342 (1980) 532
- 1979AR04 A.V. Arefev, Y.D. Bayukov, A.E. Buklei, V.B. Gavrilov, N.A. Goryainov, L.N. Kondratev, G.A. Leksin, V.S. Pavlov, V.Y. Rusinov, V.B. Fedorov et al, Yad. Fiz. 29 (1979) 410; Sov. J. Nucl. Phys. 29 (1979) 205
- 1979AT01 M. Attrep, Jr., W.B. Ledbetter and D.K. Riddle, J. Inorg. Nucl. Chem. 41 (1979) 1
- 1979AU07 G.F. Auchampaugh, S. Plattard and N.W. Hill, Nucl. Sci. Eng. 69 (1979) 30

- 1979BA16 B. Bassalleck, H.-D. Engelhardt, W.D. Klotz, F. Takeutchi, H. Ullrich and M. Furic, Nucl. Phys. A319 (1979) 397
- 1979BA48 J.K. Bair and J. Gomez del Campo, Nucl. Sci. Eng. 71 (1979) 18
- 1979BA68 V.N. Baturin, V.P. Koptev, E.M. Maev, M.M. Makarov, V.V. Nelyubin, V.V. Sulimov, A.V. Khanzadeev and G.V. Shcherbakov, Pisma Zh. Eksp. Teor. Fiz. 30 (1979) 86; JETP Lett. 30 (1979) 78
- 1979BE1N Beer et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper 1B27
- 1979BE60 G.G. Beznogikh, N.K. Zhidkov, L.F. Kirillova, V.A. Nikitin, P.V. Nomokonov, V.V. Avdeichikov, Yu.A. Murin, V.S. Oplavin, V.D. Maisyukov, Yu.V. Maslennikov et al, Pisma Zh. Eksp. Teor. Fiz. 30 (1979) 130; JETP Lett. 30 (1979) 323
- 1979BO1N Bobchenko et al, Yad. Fiz. 30 (1979) 1553
- 1979BO1P Boudard et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper 4A16
- 1979BO22 V.I. Bogatin, O.V. Lozhkin and Y.P. Yakovlev, Nucl. Phys. A326 (1979) 508
- 1979BR1D Brajnik et al, Int. Conf. Nucl. Phys. with Electromag. Int., Mainz (1979) 46
- 1979BY01 R.C. Byrd, R.L. Walter and S.R. Cotanch, Phys. Rev. Lett. 43 (1979) 260
- 1979CL06 M.R. Clover, T.M. Cormier, B.R. Fulton and B.J. Herman, Phys. Lett. 43 (1979) 256
- 1979DE01 P. Desgrolard and P.A.M. Guichon, Phys. Rev. C19 (1979) 120
- 1979DE15 P. Desgrolard and P.A.M. Guichon, Z. Phys. A290 (1979) 373
- 1979DO04 W.E. Dollhopf, C.F. Perdrisat, P. Kitching and W.C. Olsen, Nucl. Phys. A316 (1979) 350
- 1979EG01 P. Egelhof, J. Barrette, P. Braun-Munzinger, W. Dreves, C.K. Gelbke, D. Kassen, E. Steffens, W. Weiss and D. Fick, Phys. Lett. B84 (1979) 176
- 1979EL12 R.J. Ellis, H.J. Ziock, K.O.H. Ziock, J. Bolger, E. Boschitz, J. Arvieux, R. Corfu and J. Piffaretti, Phys. Lett. B88 (1979) 253
- 1979FE02 J.M. Feagin, E. Merzbacher and W.J. Thompson, Phys. Lett. B81 (1979) 107
- 1979FE1E Feagin, Merzbacher and Thompson, IEEE Trans, on Nucl. Sci. 26 (1979) 1223
- 1979FL06 E. Flerackers, Nucl. Phys. A329 (1979) 45
- 1979FL1A Fleischer, American Scientist 67 (1979) 194
- 1979FO1F Fou, Rasmussen, Swann and Van Patter, IEEE Trans. on Nucl. Sci. 26 (1979) 1378
- 1979FR12 S. Frankel, W. Frati, M. Gazzaly, Y.D. Bayukov, V.I. Efremenko, G.A. Leksin, N.A. Nikiforov, V.I. Tchistilin, Y.M. Zaitsev and C.F. Perdrisat, Phys. Rev. C20 (1979) 2257
- 1979FU04 C.B. Fulmer, R.M. Wieland, D.C. Hensley, S. Raman, G.R. Satchler, A.H. Snell, P.H. Stelson and R.G. Stokstad, Phys. Rev. C20 (1979) 670
- 1979GL1D Glendinning, Purswer and Gould, Bull. Amer. Phys. Soc. 24 (1979) 830

- 1979GL1E Glendinning et al, Bull. Amer. Phys. Soc. 24 (1979) 656
- 1979GO07 R. Gorgen, F. Hinterberger, R. Jahn, P. von Rossen and B. Schuller, Nucl. Phys. A320 (1979) 296
- 1979GO09 J. Gomez del Campo, R.A. Dayras, J.A. Biggerstaff, D. Shapira, A.H. Snell, P.H. Stelson and R.G. Stokstad, Phys. Rev. Lett. 43 (1979) 26
- 1979GO1L Gove et al, IEEE Trans. on Nucl. Sci. 26 (1979) 1414
- 1979GO1M Goncalves et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper 5B3
- 1979HO13 B. Hoistad, G.S. Adams, M. Gazzaly, G. Igo, F. Irom and H. Nann, Phys. Rev. Lett. 43 (1979) 487
- 1979HU1B M.S. Hussein, Phys. Rev. C19 (1979) 807
- 1979IN1C Inoue and Tanaka, Nature 277 (1979) 209
- 1979JA1C Jackson et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper A417
- 1979JO1C Jones, in Houston, AIP Conf. Proc. 54 (1979) 116
- 1979JU1B Journey, in Proc. Third Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York, 1979) p. 461
- 1979KA1G Karban, Microscopic Optical Potentials. Hamburg, Germany (Springer-Verlag 1979) p. 264; Phys. Abs. 97712 (1979)
- 1979KE08 J. Keinonen and A. Anttila, Nucl. Phys. A330 (1979) 397
- 1979KO21 V.I. Komarov, G.E. Kosarev, H. Muller, D. Netzband, V.D. Toneev, T. Stiehler, S. Tesch, K.K. Gudima and S.G. Mashnik, Nucl. Phys. A326 (1979) 297
- 1979KO36 V.I. Komarov, G.E. Kosarev, H. Muller, D. Netzband, T. Stiehler and S. Tesch, J. Phys. G5 (1979) 1717
- 1979KU05 D. Kurath, Nucl. Phys. A317 (1979) 175
- 1979LI1B Lin and Lesser, Bull. Amer. Phys. Soc. 24 (1979) 593
- 1979LI1C Litherland, Bull. Amer. Phys. Soc. 24 (1979) 599
- 1979LI1D Li and Jiang, Phys. Energ. Fortis & Phys. Nucl. 3 (1979) 595; Phys. Abs. 22861 (1980)
- 1979LO11 H. Lohner, B. Ludewigt, D. Frekers, G. Gaul and R. Santo, Z. Phys. A292 (1979) 35
- 1979MA38 R.E. Marrs, R.E. Pollock and W.W. Jacobs, Phys. Rev. C20 (1979) 2308
- 1979ME2A D.F. Measday and G.A. Miller, Ann. Rev. Nucl. Part. Sci. 29 (1979) 121
- 1979MO04 R.A. Moyle, B.G. Glagola, G.J. Mathews and V.E. Viola, Jr., Phys. Rev. C19 (1979) 631
- 1979MO14 T. Motobayashi, I. Kohno, T. Ooi and S. Nakajima, Nucl. Phys. A331 (1979) 193
- 1979MO1E Moss et al, in Conf. Proc. TRIUMF, Vancouver (1979) Paper 4A11

- 1979NO1B Nolen, Barker and Curtin, Bull. Amer. Phys. Soc. 24 (1979) 63
- 1979NO1C Noya and Ohnishi, in Tsukuba Symp. on Polariz. Phenomena, Nov. 1979, Ed. K. Yagi (1979) 217
- 1979OE01 W. Oelert, A. Djaloeis, C. Mayer-Boricke and P. Turek, Phys. Rev. C19 (1979) 1747
- 1979PA09 L.A. Parks, K.W. Kemper, R.I. Cutler and L.H. Harwood, Phys. Rev. C19 (1979) 2206
- 1979PE1C Perroud, in 1978 RPI Symp. 'Photopion Nucl. Phys., (Plenum Press, New York, 1979) p. 69
- 1979PI06 P.H. Pile, R.D. Bent, R.E. Pollock, P.T. Debevec, R.E. Marrs, M.C. Green, T.P. Sjoreen and F. Soga, Phys. Rev. Lett. 42 (1979) 1461
- 1979RA1C Ramaty, Kozlovsky and Lingenfelter, Astrophys. J. Suppl. Series 40 (1979) 487
- 1979RA1E Raisbeck et al, Nature 282 (1979) 279
- 1979RI12 Y. Rihet, G. Costa, C. Gerardin and R. Seltz, Phys. Rev. C20 (1979) 1583
- 1979SA27 G.R. Satchler, Nucl. Phys. A329 (1979) 233
- 1979SC07 P. Schwaller, M. Pepin, B. Favier, C. Richard-Serre, D.F. Measday and P.U. Renberg, Nucl. Phys. A316 (1979) 317
- 1979SC1D Scott, Prog. Nucl. Phys. 4 (1979) 5
- 1979SH09 J.R. Shepard, R.E. Anderson, J.J. Kraushaar, R.A. Ristinen, J.R. Comfort, N.S.P. King, A. Bacher and W.W. Jacobs, Nucl. Phys. A322 (1979) 92
- 1979SH22 D. Shapira, R. Dayras, J.L.C. Ford, Jr., J. Gomez del Campo, A.H. Snell, P.H. Stelson and R.G. Stokstad, Nucl. Instrum. Meth. Phys. Res. 163 (1979) 325
- 1979SI1D Simpson, Bull. Amer. Phys. Soc. 24 (1979) 69
- 1979SP01 R.H. Spear, Z.E. Switkowski, D.L. Kennedy and J.C.P. Heggie, Nucl. Phys. A318 (1979) 21
- 1979ST15 D.P. Stahel, R. Jahn, G.J. Wozniak and J. Cerny, Phys. Rev. C20 (1979) 1680
- 1979ST1D Stanley, Glover and Petrovich, Bull. Amer. Phys. Soc. 24 (1979) 816
- 1979SU09 W. Sunkel and Y.C. Tang, Nucl. Phys. A329 (1979) 10
- 1979SU1F Suzuki and Kubo, in Tsukuba Symp. on Polarization Phenomena, Nov. 1979, Ed. K. Yagi, (1979) p. 211
- 1979TA1C Taneichi, Ueno and Shoda, Nucl. Interact., Canberra, Australia (Springer-Verlag 1979) p. 456; Phys. Abs. 63169 (1980)
- 1979TI1A Titov, Int. Conf. Nucl. Phys. with Electromagnetic Interact., Mainz (1979) 6.14
- 1979TR1B Truol, Proc. Mainz 1979, Springer Lect. Notes 108 (1979) 351
- 1979VI05 A. Vidal-Quadras and M. Ortega, Nuovo Cim. A49 (1979) 235

- 1979WA13 R.E. Warner, D.C. Martin, G.C. Ball, W.G. Davies, A.J. Ferguson and D. Horn, Nucl. Phys. A326 (1979) 209
- 1979WA1F Watermann et al, Med. Phys. (USA) 6 (1979) 432
- 1979WE1C Weber and Eisenberg, in Houston, AIP Conf. Proc. 54 (1979) 190
- 1980AB16 S.N. Abramovich, A.I. Baz and B.Ya. Guzhovsky, Yad. Fiz. 32 (1980) 402; Sov. J. Nucl. Phys. 32 (1980) 208
- 1980AD1A Adib et al, Proc. Int. Conf. on Nucl. Cross Sections for Tech., Knoxville, TN (1980) p. 101; Phys. Abs. 3925 (1982)
- 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, Nucl. Phys. A336 (1980) 1
- 1980BA2L Basu, Subba Rao and Srinivasan, Atomkernenerg Kerntech. 36 (1980) 30, 32
- 1980BL02 L. Bland and H.T. Fortune, Phys. Rev. C21 (1980) 11
- 1980BO14 K. Bodek, M. Hugi, J. Lang, R. Muller, A. Schiltz, J. Sromicki, E. Ungricht, L. Jarczyk and A. Strzalkowski, Phys. Lett. B92 (1980) 79
- 1980BO1B Bosted et al, Bull. Amer. Phys. Soc. 25 (1980) 559
- 1980BO1L Boucenna, Allab and Hadda, Bull. Amer. Phys. Soc. 25 (1980) 785
- 1980BO24 P.E. Bosted, K.I. Blomqvist, A.M. Bernstein, S.A. Dytman and R.A. Miskimen, Phys. Rev. Lett. 45 (1980) 1544
- 1980BO31 V.I. Bogatin, E.A. Ganza, O.V. Lozhkin, Yu.A. Murin and V.S. Oplavin, Yad. Fiz. 32 (1980) 27; Sov. J. Nucl. Phys. 32 (1980) 14
- 1980BR21 B.A. Brown, W.A. Richter and N.S. Godwin, Phys. Rev. Lett. 45 (1980) 1681
- 1980CO10 T.M. Cormier and B.R. Fulton, Phys. Rev. C22 (1980) 565
- 1980CO12 T.V. Congedo, I.S. Lee-Fan and B.L. Cohen, Phys. Rev. C22 (1980) 985
- 1980CR03 J.F. Crawford, M. Daum, G.H. Eaton, R. Frosch, H. Hirschmann, R. Horisberger, J.W. McCulloch, E. Steiner, R. Hausammann, R. Hess et al, Phys. Rev. C22 (1980) 1184
- 1980DE11 A. Deloff, Phys. Rev. C21 (1980) 1516
- 1980DE39 C. Detraz, D. Guillemaud, M. Langevin, F. Naulin, M. Epherre, R. Klapisch, M. de Saint-Simon, C. Thibault and F. Touchard, J. Phys. (Paris) Lett. 41 (1980) L-459
- 1980DI02 M. Dillig, P. Couvert, T.S. Bauer, R. Beurtey, A. Boudard, G. Bruge, H. Catz, A. Chaumeaux, H.H. Duhm, J.L. Escudie et al, Nucl. Phys. A333 (1980) 477
- 1980EL1B Elmore et al, Bull. Amer. Phys. Soc. 25 (1980) 784
- 1980EL1C Elmore et al, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 655
- 1980FA07 E. Fabrici, S. Micheletti, M. Pignanelli, F.G. Resmini, R. De Leo, G. D'Erasmus and A. Pantaleo, Phys. Rev. C21 (1980) 844
- 1980FR1K Franz et al, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 890

- 1980FU1G Furutani et al, Suppl. Prog. Theor. Phys. 68 (1980) 193
- 1980GL03 C.W. Glover, K.W. Kemper, L.A. Parks, F. Petrovich and D.P. Stanley, Nucl. Phys. A337 (1980) 520
- 1980GO1B Gove et al, Nucl. Instrum. Meth. Phys. Res. 168 (1980) 425
- 1980GR10 R.E.L. Green and R.G. Korteling, Phys. Rev. C22 (1980) 1594
- 1980GU1E Guet et al, Proc. Int. Conf. on Nucl. Phys., Berkeley (1980) 471
- 1980HA33 M.N. Harakeh, J. Van Popta, A. Saha and R.H. Siemssen, Nucl. Phys. A344 (1980) 15
- 1980IS02 M.A. Islam, T.J. Kennett, S.A. Kerr and W.V. Prestwich, Can. J. Phys. 58 (1980) 168
- 1980IW1A Iwao, Lett. Nuovo Cim. 29 (1980) 40
- 1980KA03 T. Kammuri, H. Shimaoka, P.D. Kunz, S. Kato, K. Okada, M. Kondo, K. Hosono, T. Saito, N. Matsuoka, S. Nagamachi et al, Phys. Lett. B90 (1980) 197
- 1980KI1B Kilius et al, Nucl. Instrum. Meth. Phys. Res. 171 (1980) 355
- 1980KO1L Kostin, Koval, Kopanets and Tsytko, Ukr. Fiz. Zh. 25 (1980) 881
- 1980KO1V Koptev, Maev, Makarov and Khanzadeev, Yad. Fiz. 31 (1980) 1501
- 1980LA1B Lanford et al, Nucl. Instrum. Meth. Phys. Res. 168 (1980) 505
- 1980LE02 T.-S.H. Lee and D. Kurath, Phys. Rev. C21 (1980) 293
- 1980LI1F Lin and Lesser, Bull. Amer. Phys. Soc. 25 (1980) 577
- 1980MA1F Maleki, Rao and Levinger, Bull. Amer. Phys. Soc. 25 (1980) 606
- 1980MA1Z Mandal and Saha, Can. J. Phys. 58 (1980) 300
- 1980MA33 G. Mack, R.C. Byrd, P.W. Lisowski and R.L. Walter, Nucl. Phys. A345 (1980) 241
- 1980MI01 T. Mikumo, M. Sasagase, M. Sato, T. Ooi, Y. Higashi, Y. Nagashima and M. Yamanoichi, Phys. Rev. C21 (1980) 620
- 1980MU1B N.C. Mukhopadhyay, Nucl. Phys. A335 (1980) 111
- 1980MU1D Muller et al, Nucl. Instrum. Meth. Phys. Res. 170 (1980) 151
- 1980NA1B Nagl and Uberall, Bull. Amer. Phys. Soc. 25 (1980) 606
- 1980NI09 N.A. Nikiforov, Y.D. Bayukov, V.I. Efremenko, G.A. Leskin, V.I. Tchistilin, Y.M. Zaitsev, S. Frankel, W. Frati, M. Gazzaly and C.F. Perdrisat, Phys. Rev. C22 (1980) 700
- 1980NI1F Nishioka, Euro. Symp. on Few Body Prob. in Nucl. and Part. Phys. (1980); Phys. Abs. 28380 (1981)
- 1980NO1A Nolen, Proc. Int. Conf. in At. Masses and Fund. Constants, Nolen and Beneson Eds., Plenum Publ. (1980) 505
- 1980OK1B Okabe, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 9

- 1980OL1C Olson et al, Proc. Int. Conf. on Nucl. Phys., Berkeley (1980) p. 200
- 1980PA01 L.A. Parks, D.P. Stanley, L.H. Courtney and K.W. Kemper, Phys. Rev. C21 (1980) 217
- 1980PA19 R.L. Parks, S.T. Thornton, L.C. Dennis and K.R. Cordell, Nucl. Phys. A348 (1980) 350
- 1980RE1B Read et al, Bull. Amer. Phys. Soc. 25 (1980) 592
- 1980RI06 W.A. Richter and P.R. de Kock, Z. Phys. A297 (1980) 343
- 1980RO1F Roy-Choudhury, Gautam and Surak, Pramana 14 (1980) 57
- 1980SE1F Seth et al, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 164
- 1980SJ02 T.P. Sjoreen, M.C. Green, W.W. Jacobs, R.E. Pollock, F. Soga, R.D. Bent and T.E. Ward, Phys. Rev. Lett. 45 (1980) 1769
- 1980SK1A Skrzypczak et al, Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 575
- 1980ST25 K. Stricker, J.A. Carr and H. McManus, Phys. Rev. C22 (1980) 2043
- 1980TR02 H.-J. Trost, A. Schwarz, U. Feindt, F.H. Heimlich, S. Heinzl, J. Hintze, F. Korber, R. Lekebusch, P. Lezoch, G. Mock et al, Nucl. Phys. A337 (1980) 377
- 1980VA03 R. Vandenbosch, Nucl. Phys. A339 (1980) 167
- 1980WA1K Walter and Byrd, Proc. Int. Conf. in the (p, n) reaction and the Nucleon-Nucleon Force, Plenum (1980) p. 469
- 1980WI09 J.P. Wieleczko, S. Harar, M. Conjeaud and F. Saint-Laurent, Phys. Lett. B93 (1980) 35
- 1980WI1L L. Winsberg, Phys. Rev. C22 (1980) 2123
- 1980WI1M Wiedenbeck and Greiner, Astrophys. J. 239 (1980) L139
- 1980ZH1C Zheng, Li and Shen, Transactions of the N.Y. Acad. of Sci. 40 (1980) 274
- 1981AB1A Abe, RIFP-421 (1981) 428
- 1981AU1C Auld, Santa Fe 1980, AIP Conf. Proc. 69 (1981) 93
- 1981AU1D Audouze, Prog. Part. Nucl. Phys. 6 (1981) 125
- 1981AU1G Austin, Prog. Part. Nucl. Phys. 7 (1981) 1
- 1981AV02 I.K. Averyanov, A.I. Golubev and A.A. Sadovoy, Yad. Fiz. 33 (1981) 66
- 1981BA1R Bayukov et al, Yad. Fiz. 33 (1981) 183
- 1981BA37 A.K. Basak, O. Karban, S. Roman, G.C. Morrison, C.O. Blyth and J.M. Nelson, Nucl. Phys. A368 (1981) 74
- 1981BA38 A.K. Basak, O. Karban, S. Roman, G.C. Morrison, C.O. Blyth and J.M. Nelson, Nucl. Phys. A368 (1981) 93
- 1981BA64 F.C. Barker, Aust. J. Phys. 34 (1981) 7

- 1981BE63 V.B. Belyaev and O.P. Solovtsova, *Yad. Fiz.* 33 (1981) 699; *Sov. J. Nucl. Phys.* 33 (1981) 363
- 1981BL1G M. Blann and T.T. Komoto, *Phys. Rev. C*24 (1981) 426
- 1981BO03 A. Boudard, Y. Terrien, R. Beurtey, L. Bimbot, G. Burge, A. Chaumeaux, P. Couvert, J.M. Fontaine, M. Garcon, Y. Le Bornec et al, *Phys. Rev. Lett.* 46 (1981) 218; *Phys. Rev. Lett.* 47 (1981) 147
- 1981BO1C D.H. Boal and R.M. Woloshyn, *Phys. Rev. C*23 (1981) 1206
- 1981BO1D Boyd, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 373
- 1981BO1Y Bouten and Bouten, *Prog. Part. Nucl. Phys.* 5 (1981) 55
- 1981BR1E Brooks, Lister, Nelson and Dhuga, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 656
- 1981BY1B Byrd and Walter, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 1475
- 1981BY1C Byrd and Walter, *Bull. Amer. Phys. Soc.* 26 (1981) 624
- 1981CE04 F.E. Cecil and R.F. Fahlsing, *Phys. Rev. C*24 (1981) 1769; Erratum *Phys. Rev. C*25 (1982) 2137
- 1981CI03 M.A. Cirit and F. Yazici, *Phys. Rev. C*23 (1981) 2627
- 1981CO1D Conzett, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 1452
- 1981DA1K Dave et al, *Bull. Amer. Phys. Soc.* 26 (1981) 551
- 1981DE13 J.-L. Dethier and Fl. Stancu, *Phys. Rev. C*23 (1981) 1503
- 1981DE2G de Wet, *Found. Phys.* 11 (1981) 155
- 1981ER10 J. Ero, Z. Fodor, P. Koncz, Z. Seres, M. Csatos, B.A. Khomenko, N.N. Khovanskij, Z.V. Krumstein, Yu.P. Merekov and V.I. Petrukhin, *Nucl. Phys. A*372 (1981) 317
- 1981FA1E Farwell, Grootes and Schmidt, *Bull. Amer. Phys. Soc.* 26 (1981) 1159
- 1981FE2A H.W. Fearing, *Prog. Part. Nucl. Phys.* 7 (1981) 113
- 1981FL04 C.E. Floyd, P.P. Guss, K. Murphy, R.C. Byrd, G. Tungate, S.A. Wender, R.L. Walter and T.B. Clegg, *Phys. Rev. Lett.* 47 (1981) 1042
- 1981FL1A Floyd et al, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 407
- 1981FR17 E. Friedman, *Phys. Lett.* B104 (1981) 357
- 1981GA1A Gagliardi et al, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 939
- 1981GA1B Garcia-Munoz, Simpson and Wefel, *Bull. Amer. Phys. Soc.* 26 (1981) 557
- 1981GA1C Garcia-Munoz, Guzik, Simpson and Wefel, *Bull. Amer. Phys. Soc.* 26 (1981) 557
- 1981GE1B Geesaman et al, *Proc. Versailles Conf.* (1981) p. 302
- 1981GI08 M. Giffon, A. Goncalves, P.A.M. Guichon, J. Julien, L. Roussel and C. Samour, *Phys. Rev. C*24 (1981) 241
- 1981GI15 Yu.R. Gismatullin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 45 (1981) 674

- 1981GU1D Guzik, *Astrophys. J.* 244 (1981) 695
- 1981HAZJ R.C. Haight and S.M. Grimes, *Bull. Amer. Phys. Soc.* 26 (1981) 1138, CE1
- 1981HE05 P. Heusi, M. Berta, V. Meyer and R.E. Pixley, *Nucl. Phys.* A357 (1981) 381
- 1981HO13 J. Hohn, J. Kayser, W. Pilz, D. Schmidt and D. Seeliger, *J. Phys.* G7 (1981) 803
- 1981JO02 P.L. Jolivette, *Nucl. Phys.* A370 (1981) 256
- 1981KE02 T.J. Kennett, M.A. Islam and W.V. Prestwich, *Can. J. Phys.* 59 (1981) 93
- 1981KE15 B.K. Kerimov, M.Ya. Safin and I.M. Al-Khamisi, *Yad. Fiz.* 34 (1981) 996; *Sov. J. Nucl. Phys.* 34 (1981) 554
- 1981KE16 B.K. Kerimov, A.I. Elgavkhari and A.G. Ganiev, *Izv. Akad. Nauk SSSR Ser. Fiz.* 45 (1981) 2189; *Bull. Acad. Sci. USSR Phys. Ser.* 45 (1981) 145
- 1981KH08 A.F. Khodyachikh, P.I. Vatset, V.N. Gurev, I.V. Dogyust and V.V. Kirichenko, *Yad. Fiz.* 34 (1981) 1425; *Sov. J. Nucl. Phys.* 34 (1981) 789
- 1981KI04 Yu.V. Kirichenko, V.Yu. Gonchar, E.V. Inopin and V.N. Tarasov, *Ukr. Fiz. Zh.* 26 (1981) 904
- 1981KN06 W. Knupfer and A. Richter, *Phys. Lett.* B101 (1981) 375
- 1981KO1U Kong et al, *Phys. Energ. Fortis Phys. Nucl.* 5 (1981) 600
- 1981KR17 S.P. Krekoten, A.M. Mukhamedzhanov and E.A. Romanovsky, *Izv. Akad. Nauk SSSR Ser. Fiz.* 45 (1981) 790
- 1981KR1C Kruse et al, *Bull. Amer. Phys. Soc.* 26 (1981) 543
- 1981KR1J Kramer, John and Schenzle, *Clustering Phenomena in Nuclei* 2 (1981)
- 1981KU04 D. Kurath and W. Teeters, *Phys. Lett.* B101 (1981) 5
- 1981LA13 M. Lattuada, F. Riggi, C. Spitaleri, D. Vinciguerra and C.M. Sutura, *Nuovo Cim.* A63 (1981) 530
- 1981LI1K Litherland et al, *Nucl. Sci.* 28 (1981) 1469
- 1981MA18 J.F. Mateja, A.D. Frawley, L.C. Dennis, K. Abdo and K.W. Kemper, *Phys. Rev. Lett.* 47 (1981) 311
- 1981MA1J Maday et al, *Proc. Versailles Conf.* (198) p. 538
- 1981ME13 M.C. Mermaz, J. Barrette and H.E. Wegner, *Phys. Rev.* C24 (1981) 2148
- 1981MO20 J. Mougey, R. Ost, M. Buenerd, A.J. Cole, C. Guet, D. Lebrun, J.M. Loiseaux, P. Martin, M. Maurel, E. Monnard et al, *Phys. Lett.* B105 (1981) 25
- 1981MU1D Murphy et al, *Santa Fe 1980, AIP Conf. Proc.* 69 (1981) 1478
- 1981MU1E Mukhopadhyay and Hintermann, *Santa Fe 1980, AIP Conf. Proc.* 69 (1981) 1068
- 1981MU1F Murnick, Chung, Niv and McGlashan-Powell, *Bull. Amer. Phys. Soc.* 26 (1981) 623

- 1981MUZQ S.F. Mughabghab, M. Divadeenam and N.E. Holden, Neutron Cross Sections 1A (1981) Z=1–60
- 1981NA05 T. Nakamura, Y. Chiba, M. Aoki and S. Kaneko, Nucl. Phys. A365 (1981) 457
- 1981NI1B Niskanen, Santa Fe 1980. AIP Conf. Proc. 69 (1981) 62
- 1981OL01 A. Olin, P.R. Poffenberger, G.A. Beer, J.A. Macdonald, G.R. Mason, R.M. Pearce and W.C. Sperry, Nucl. Phys. A360 (1981) 426
- 1981PR1B Protheroe, Bull. Amer. Phys. Soc. 26 (1981) 556
- 1981RA1F Raisbeck et al, Nature 293 (1981) 825
- 1981RI06 P.J. Riley, C.L. Hollas, C.R. Newsom, R.D. Ransome, B.E. Bonner, J.E. Simmons, T.S. Bhatia, G. Glass, J.C. Hiebert, L.C. Northcliffe et al, Phys. Lett. B103 (1981) 313
- 1981RO1M Roy et al, Proc. Versailles Conf. (1981) p. 133
- 1981SA1G Savin, Kruse and Moniot, Bull. Amer. Phys. Soc. 26 (1981) 543
- 1981SC1D Schmidt, Farwell and Grootes, Bull. Amer. Phys. Soc. 26 (1981) 593
- 1981SE04 R.M. Sealock, H.-Y. Wu and J.C. Overley, Nucl. Phys. A357 (1981) 279
- 1981SE06 M. Seya, M. Kohno and S. Nagata, Prog. Theor. Phys. 65 (1981) 204
- 1981SI09 M.K. Singham and F. Tabakin, Ann. Phys. 135 (1981) 71
- 1981SI1D Singham and Tabakin, Proc. Versailles Conf. (1981) 129
- 1981SL03 R.J. Slobodrian, C. Rioux, R. Roy, H.E. Conzett, P. von Rossen and F. Hinterberger, Phys. Rev. Lett. 47 (1981) 1803
- 1981ST05 C.E. Stronach, B.J. Lieb, H.O. Funsten, W.J. Kossler, H.S. Plendl and V.G. Lind, Phys. Rev. C23 (1981) 2150
- 1981ST1P Stokstad, Nukleonika 26 (1981) 373
- 1981TA22 S.L. Tabor, L.C. Dennis and K. Abdo, Phys. Rev. C24 (1981) 2552
- 1981TH1A Thornton, Parks, Cordell and Woedner, Bull. Amer. Phys. Soc. 26 (1981) 611
- 1981TH1C Thomas, Mangini and Parker, Nucl. Sci. 28 (1981) 1478
- 1981UL1B Ullmann et al, Med. Phys. 8 (1981) 396
- 1981VA1D van Driel et al, Kernfysisch Versneller Inst. (KVI) 235 (1980) 1
- 1981VA1E R. Vandenbosch and A.J. Lazzarini, Phys. Rev. C23 (1981) 1074
- 1981VI1B Vigdor, Santa Fe 1980, AIP Conf. proc. 69 (1981) 1429
- 1981WA1G Walter, Santa Fe 1980, AIP Conf. proc. 69 (1981) 344
- 1981WA1J Wang, Zhang, Li and Ruan, Proc. Versailles Conf. (1981) 374
- 1981WH01 W.R. Wharton and B.D. Keister, Phys. Rev. C23 (1981) 1141

- 1981YA1A Yamazaki et al, Proc. Versailles Conf. (1981) 115
- 1982AB1D Abramovich, Gushovskii and Protopopov, in Kiev (1982) 566
- 1982AL08 D.V. Aleksandrov, Yu.A. Glukhov, A.S. Demyanova, V.I. Dukhanov, I.B. Mazurov, B.G. Novatsky, A.A. Ogloblin, S.B. Sakuta and D.N. Stepanov, Yad. Fiz. 35 (1982) 277; Sov. J. Nucl. Phys. 35 (1982) 158
- 1982AU02 E.G. Auld, G. Jones, G.J. Lolos, E.L. Mathie, P.L. Walden and R.B. Taylor, Phys. Rev. C25 (1982) 2222
- 1982AV1A Averyanoy, Golubev and Sadovi, Sov. J. Nucl. Phys. 35 (1982) 484
- 1982BA24 M.T. Barriuso and F. Fernandez, Lett. Nuovo Cim. 33 (1982) 305
- 1982BA2G B.F. Bayman, P.J. Ellis and Y.C. Tang, Phys. Rev. Lett. 49 (1982) 532
- 1982BA2T Barabash et al, Yad. Fiz. 36 (1982) 155
- 1982BA52 F.C. Barker, Aust. J. Phys. 35 (1982) 291
- 1982BE1Z Bevelacqua, Rev. Mex. Fis. 28 (1982) 573
- 1982BE54 Yu.A. Berezhnoi, A.P. Klyucharev and N.Ya. Rutkevich, Ukr. Fiz. Zh. 27 (1982) 1475; Phys. Abs. 11339 (1983)
- 1982BI1C Bikov et al, in Kiev (1982) 559
- 1982BO1J D.H. Boal, Phys. Rev. C25 (1982) 3068
- 1982BO35 C. Borcea, E. Gierlik, A.M. Kalinin, R. Kalpakchieva, Yu.Ts. Oganessian, T. Pawlat, Yu.E. Penionzhkevich and A.V. Rykhlyuk, Nucl. Phys. A391 (1982) 520
- 1982BR1N Brown et al, Nature 299 (1982) 718
- 1982BU03 B.L. Burks, R.E. Anderson, T.B. Clegg, H. Paetz gen.Schieck, E.J. Ludwig, R.L. Varner and J.F. Wilkerson, Phys. Rev. C25 (1982) 1168
- 1982BY1A Byrd, Murphy and Walter, Bull. Amer. Phys. Soc. 27 (1982) 460
- 1982CH07 Y.-D. Chan, D.E. DiGregorio, J.L.C. Ford, Jr., J. Gomez del Campo, M.E. Ortiz and D. Shapira, Phys. Rev. C25 (1982) 1410
- 1982CH1M Choudhury et al, in MSU (1982) 88
- 1982COZV B. Cottman, A.M. Bernstein, K.I. Blomqvist, S.A. Dytman and J. Nelson, Bull. Amer. Phys. Soc. 27 (1982) 709, BE7
- 1982DE14 V. DeCarlo and N. Freed, Phys. Rev. C25 (1982) 2162
- 1982DE1N de Wet, Found. Phys. 12 (1982) 285
- 1982DO01 K.G.R. Doss, P.D. Barnes, N. Colella, S.A. Dytman, R.A. Eisenstein, C. Ellegaard, F. Takeuchi, W.R. Wharton, J.F. Amann, R.H. Pehl et al, Phys. Rev. C25 (1982) 962
- 1982DO08 I.V. Dogyust, P.I. Vatsset, V.V. Kirichenko and A.F. Khodyachikh, Yad. Fiz. 35 (1982) 810

- 1982EL07 R.J. Ellis, H.J. Ziock, K.O.H. Ziock, Y. Tzeng, J. Arvieux, R. Corfu, J. Piffaretti, L.C. Liu and E.R. Siciliano, Phys. Rev. C26 (1982) 1544
- 1982ER1E Eramzhyan, Fetisovm Majling and Zofka, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 91
- 1982FU04 T. Fukuda, M. Ishihara, M. Tanaka, I. Miura, H. Ogata and H. Kamitsubo, Phys. Rev. C25 (1982) 2464
- 1982FU09 C.B. Fulmer, S. Mukhopadhyay, G.R. Satchler, R.L. Auble, J.B. Ball, F.E. Bertrand, E.E. Gross and D.C. Hensley, Nucl. Phys. A385 (1982) 83
- 1982GO03 N.G. Goncharova, B.S. Ishkhanov and V.I. Mokeev, Yad. Fiz. 35 (1982) 43; Sov. J. Nucl. Phys. 35 (1982) 26
- 1982GO05 S.A. Goncharov, Ya. Dobesh, E.I. Dolinsky, A.M. Mukhamedzhanov and Ya. Tseipek, Yad. Fiz. 35 (1982) 662
- 1982GO1E Gogitidze et al, in Kiev (1982) 338
- 1982GU1H H.H. Gutbrod, A.I. Warwick and H. Wieman, Nucl. Phys. A387 (1982) 177
- 1982HA1A Haight, Proc. 4th Int. Symp. Grenoble 1981 (IOP 1982) 510
- 1982HA42 Q. Haider and F.B. Malik, Phys. Rev. C26 (1982) 989
- 1982HO1C Hoistad, in "Pion Prod. & Absorp. in Nucl.-Indiana 1981", AIP Conf. Proc. 79 (1982) 105
- 1982HO1F Hoppe et al, in Bad Honnef Symp. 1981, Springer-Verlag (1982) 135; Phys. Abs. 101808 (1982)
- 1982IK1A Ikeda, Miyahara and Bando, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 149
- 1982KL1A Klein, Middleton and Tang, Nucl. Instrum. Meth. Phys. Res. 193 (1982) 601
- 1982KO11 N.N. Kolesnikov, D. Amarasingam and V.I. Tarasov, Yad. Fiz. 35 (1982) 32; Sov. J. Nucl. Phys. 35 (1982) 20
- 1982KO1L Kolesnikov, Kopilov andf Kolesov, in Kiev (1982) 187
- 1982KU1G Ku et al, Nature 299 (1982) 240
- 1982KU1J Kusakabe et al, Nature 299 (1982) 712
- 1982LA26 J. Lange, K. Kumar and J.H. Hamilton, Revs. Mod. Phys. 54 (1982) 119
- 1982LE10 M. LeMere, Y.C. Tang and H. Kanada, Phys. Rev. C25 (1982) 2902
- 1982LO02 G.J. Lolos, E.L. Mathie, P.L. Walden, G. Jones, E.G. Auld and R.B. Taylor, Phys. Rev. C25 (1982) 1082
- 1982LO13 M. Lozano, J.I. Escudero and G. Madurga J. Phys. (London) G8 (1982) 1259
- 1982LU01 B. Ludewigt, G. Gaul, R. Glasow, H. Lohner and R. Santo, Phys. Lett. B108 (1982) 15

- 1982LY1A U. Lynen, H. Ho, W. Kuhn, D. Pelte, U. Winkler, W. F.J. Muller, Y. -T. Chu, P. Doll, A. Gobbi, K. Hildenbrand et al, Nucl. Phys. A387 (1982) 129
- 1982MA20 J.F. Mateja, A.D. Frawley, L.C. Dennis, K. Abdo and K.W. Kemper, Phys. Rev. C25 (1982) 2963
- 1982MA25 G. Mairle, G.J. Wagner, P. Grabmayr, K.T. Knopff, Liu Ken Pao, H. Riedesel, K. Schindler, V. Bechtold, L. Friedrich and P. Ziegler, Nucl. Phys. A382 (1982) 173
- 1982MO1K Mohr, Aust. J. Phys. 35 (1982) 1
- 1982NA01 J. Navarro, J. Bernabeu, J.M.G. Gomez and J. Martorell, Nucl. Phys. A375 (1982) 361
- 1982NA1K Nann, in ‘Pion Prod. and Absorp. in Nucl.-Indiana 1981’, AIP Conf. Proc. 79 (1982) 219
- 1982NG01 Nguyen Tien Nguyen and I. Ulehla, Czech. J. Phys. B32 (1982) 1040
- 1982OG02 Yu.Ts. Oganessian, Yu.E. Penionzhkevich, E. Gierlik, R. Kalpakchieva, T. Pawlat, C. Borcea, A.V. Belozerov, Yu.P. Kharitonov, S.P. Tretyakova, V.G. Subbotin et al, Pisma Zh. Eksp. Teor. Fiz. 36 (1982) 104; JETP Lett. 36 (1982) 129
- 1982OR02 M.E. Ortiz, J. Gomez del Campo, Y.D. Chan, D.E. DiGregorio, J.L.C. Ford, D. Shapira, R.G. Stokstad, J.P.F. Sellschop, R.L. Parks and D. Weiser, Phys. Rev. C25 (1982) 1436
- 1982OV1A Overley, Bull. Amer. Phys. Soc. 27 (1982) 770
- 1982PA1F D.K. Pal, C. Tuniz, R.K. Moniot, T.H. Kruse and G.F. Herzog, Science 218 (1982) 787
- 1982PE1F Pella et al, Bull. Amer. Phys. Soc. 27 (1982) 729
- 1982RA1L Rahman Khan, Conf. on Hypernucl. Kaon Phys., Heidelberg, June 1982 (1982) 115
- 1982RA20 R.D. Ransome, C.L. Hollas, P.J. Riley, B.E. Bonner, W.D. Cornelius, O.B. van Dyck, E.W. Hoffman, S.A. Wood and K. Toshioka, Nucl. Instrum. Meth. Phys. Res. 201 (1982) 315
- 1982RI04 L. Ricken, D. Bohle, G. Domogala, K. Glasner and E. Kuhlmann, Z. Phys. A306 (1982) 67
- 1982RO04 D. Rowley, J. LeRose, K. Min, B.O. Sapp, P. Stoler, E.J. Winhold, P.F. Yergin, A.M. Bernstein, K.I. Blomqvist, H.S. Caplan et al, Phys. Rev. C25 (1982) 2652
- 1982SA1U Sakai, Inst. for Nucl. Study (Japan) (1982)
- 1982SM1F Smith and Liskien, Eur. Appl. Res. Nucl. Sci. Technol. Sect. 4 (1982) 969; Phys. Abs. 31396 (1983)
- 1982VE02 J.J.M. Verbaarschot, G.A. Timmer and P.J. Brussaard, Nucl. Phys. A378 (1982) 280
- 1982VE05 W.J. Vermeer and A.R. Poletti, J. Phys. G8 (1985) 851

- 1982VE11 W.J. Vermeer, T.H. Zabel, M.T. Esat, J.A. Kuehner, R.H. Spear and A.M. Baxter, Aust. J. Phys. 35 (1982) 283
- 1982WA1G P.L. Walden, Nucl. Phys. A374 (1982) 277
- 1982ZU03 B.W. Zulkoskey, R.M. Sealock, H.S. Caplan and J.C. Bergstrom, Phys. Rev. C26 (1982) 1610
- 1983AB1A Abramovitch et al, in Moscow (1983) p. 362
- 1983ANZY R.E. Anderson, C.R. Gould, B.C. Karp, K. E.Nash, R.S. Pedroni, H.G. Pfurtner and R.L. Walter, Bull. Amer. Phys. Soc. 28 (1983) 648, AF1
- 1983BE02 E. Betak and V.D. Toneev, J. Phys. G9 (1983) L47
- 1983BE16 S.L. Belostotsky, Yu.V. Dotsenko, J. Ero, Z. Fodor, L.G. Kudin, N.P. Kuropatkin, A.A. Lobodenko, O.V. Miklukho, V.N. Nikulin, O.E. Prokofev et al, Phys. Lett. B124 (1983) 469
- 1983BY01 R.C. Byrd, C.E. Floyd, P.P. Guss, K. Murphy and R.L. Walter, Nucl. Phys. A399 (1983) 94
- 1983CE01 F.E. Cecil, R.F. Fahlsing, N. Jarmie, R.A. Hardekopf and R. Martinez, Phys. Rev. C27 (1983) 6
- 1983CH08 A. Chalupka, H. Vonach, E. Huenges and H.J. Scheerer, Z. Phys. A310 (1983) 135
- 1983CI08 O. Civitarese, B.V. Carlson, M.S. Hussein and A. Szanto de Toledo, Phys. Lett. B125 (1983) 22
- 1983DE1P Deineko et al, in Moscow (1983) p. 318
- 1983GO07 D. Gola, W. Bretfeld, W. Burgmer, H. Eichner, Ch. Heinrich, H.J. Helten, H. Kretzer, K. Prescher, H. Oswald, W. Schnorrenberg et al, Phys. Rev. C27 (1983) 1394
- 1983GO1R Gontcharova, Kissener and Eramzhian, in Moscow (1983) p. 188
- 1983GO1T Goltsov, Goncharova and Matveev, in Moscow (1983) p. 211
- 1983KR1B Kruse, Bull. Amer. Phys. Soc. 28 (1983) 44
- 1983LE1J Letaw, Silberberg and Tsao, Astrophys. J. 51 (1983) 271
- 1983NAZZ J. Napolitano and S.J. Freedman, Bull. Amer. Phys. Soc. 28 (1983) 650, AG4
- 1983RI01 C. Rioux, R. Roy, R.J. Slobodrian and H.E. Conzett, Nucl. Phys. A394 (1983) 428
- 1983RO1G Rotter, Phys. Rev. C27 (1983) 2261
- 1983SA06 M. Sato, M. Sasagase, Y. Nagashima, J. Schimizu, T. Nakagawa, Y. Fukuchi and T. Mikumo, Phys. Rev. C27 (1983) 2621
- 1983SEZW R.E. Segel, A. Hassan, S.M. Levenson, S. Mukhopadhyay, P. Zupranski, J.V. Maher and P.P. Singh, Bull. Amer. Phys. Soc. 28 (1983) 691, DG10
- 1983SH04 V. Shkolnik and Y.C. Tang, Nucl. Phys. A397 (1983) 132
- 1983SO1B Southon et al, Nucl. Instrum. Meth. Phys. Res. 205 (1983) 251

1983ST1H Stepanenko et al, in Moscow (1983) 387
1983SU1C Suter et al, IEEE Trans. Nucl. Sci. 30 (1983) 1528
1983VA1H Vasilieva et al, in Moscow (1983) 331
1985AJ01 F. Ajzenberg-Selove, Nucl. Phys. A433 (1985) 1; Erratum Nucl. Phys. A449 (1986)
155
1986AJ01 F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1
HO82F Unknown Source