

Energy Levels of Light Nuclei $A = 11$

F. Ajzenberg-Selove^a and T. Lauritsen^b

^a *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

^b *California Institute of Technology, Pasadena, California*

Abstract: An evaluation of $A = 11$ –12 was published in *Nuclear Physics A114* (1968), p. 1. This version of $A = 11$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 15, 1967)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 11$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{11}\text{Li}\$](#) , [\$^{11}\text{Be}\$](#) , [\$^{11}\text{B}\$](#) , [\$^{11}\text{C}\$](#) , [\$^{11}\text{N}\$](#)

B. Tables of Recommended Level Energies:

[Table 11.1](#): Energy levels of ^{11}Be

[Table 11.3](#): Energy levels of ^{11}B

[Table 11.14](#): Energy levels of ^{11}C

C. [References](#)

D. Figures: [\$^{11}\text{Be}\$](#) , [\$^{11}\text{B}\$](#) , [\$^{11}\text{C}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

¹¹Li
(Fig. 4)

¹¹Li has been identified in the 5.3 GeV proton bombardment of uranium. It is particle stable (1966PO09). See also (1966GA25, 1967CO1K).

¹¹Be
(Figs. 1 and 4)

GENERAL:

Mass of ¹¹Be: The Q -value of the ${}^9\text{Be}(t, p){}^{11}\text{Be}$ reaction is given as $Q = -1.164 \pm 0.015$ MeV by (1962PU01) (based on ${}^{12}\text{C}(t, p){}^{14}\text{C}^*$). This value has been adjusted by (1965RY01) to -1.170 MeV, leading to $M - A$ for ${}^{11}\text{Be} = 20.181 \pm 0.015$ MeV (relative to ${}^{12}\text{C}$) (1965MA54).

See (1960TA07, 1960TA1C, 1961DO03, 1966RO11, 1967DE1V). The ground state of ${}^{11}\text{Be}$ has even parity (1964AL22).

1. ${}^{11}\text{Be}(\beta^-){}^{11}\text{B}$ $Q_m = 11.513$

The decay proceeds to the ${}^{11}\text{B}$ states at 0, 2.12, 6.79 and 8.00 MeV ($J^\pi = \frac{3}{2}^-, \frac{1}{2}^-, \frac{3}{2}^+, \frac{3}{2}^+$, respectively); the half-life is 13.68 ± 0.14 sec. The nature of the decay indicates even parity, $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+$ or $\frac{5}{2}^+$ for ${}^{11}\text{Be}_{\text{g.s.}}$ (1961DO03, 1964AL22): see ${}^{11}\text{B}$.

2. ${}^9\text{Be}(t, p){}^{11}\text{Be}$ $Q_m = -1.170$

Proton groups have been observed corresponding to seven states of ${}^{11}\text{Be}$: see Table 11.2 (1962PU01). Angular distributions measured at $E_t = 6, 10$ and 14 MeV do not show a clear stripping pattern for the ground-state protons. At $E_t = 14$ MeV, the distribution of protons corresponding to ${}^{11}\text{Be}^*(0.319)$ exhibits an $L = 2$ pattern indicating $J^\pi = \frac{1}{2}^-, \frac{5}{2}^-$ or $\frac{7}{2}^-$ (1962PU01, 1964HI08). (1960TA07) suggest $J^\pi = \frac{1}{2}^+$ and $\frac{1}{2}^-$ for the ground and first excited states, respectively. The dimensionless reduced widths θ_n^2 for $E_x = 1.8, 2.7$ and 3.4 MeV indicate $l = 0, 1$ or 2 decay to ${}^{10}\text{Be} + n$, $J^\pi = \frac{1}{2}^\pm, \frac{3}{2}^\pm$ or $\frac{5}{2}^+$ (1962PU01). See also (1968HO1F).

3. ${}^9\text{Be}({}^6\text{He}, \alpha){}^{11}\text{Be}$ $Q_m = 6.343$

Table 11.1: Energy levels of ^{11}Be

E_x (MeV \pm keV)	J^π	Γ_{lab} (keV)	Decay	Reactions
0	$\frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+$	$\tau_{1/2} = 13.81 \pm 0.08$ sec	β^-	1, 2, 3, 4
0.319 ± 10	$\frac{1}{2}^-, \frac{5}{2}^-, \frac{7}{2}^-$	< 10	(γ)	2
1.780 ± 20	$\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+$	110 ± 15	(n)	2
2.700 ± 25	$\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+$	250 ± 20	(n)	2
3.410 ± 25	$\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+$	150 ± 20	(n)	2
3.890 ± 20		< 10	(n)	2
3.960 ± 20		< 10	(n)	2

 Table 11.2: Levels of ^{11}Be from $^9\text{Be}(t, p)^{11}\text{Be}$ ^a

E_x (MeV \pm keV)	Γ_{lab} (keV)	J^π	θ_n^2 ^b			
			s	p	d	f
0		$\frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+$				
0.319 ± 10	< 10	$\frac{1}{2}^-, \frac{5}{2}^-, \frac{7}{2}^-$				
1.780 ± 20	110 ± 15	$\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+$	0.028	0.051	0.40	10
2.700 ± 25	250 ± 20	$\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+$	0.046	0.080	0.31	3.6
3.410 ± 25	150 ± 20	$\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+$	0.025	0.038	0.073	0.89
3.890 ± 20	< 10					
3.960 ± 20	< 10					

^a (1962PU01, 1964HI08).

^b Units of $\hbar^2/\mu R^2$.

See (1967ST21) and ^{15}C .

4. $^{11}\text{B}(\text{n}, \text{p})^{11}\text{Be}$ $Q_{\text{m}} = -10.731$

At $E_{\text{n}} = 14.8$ MeV, $\sigma(\text{n}\alpha)/\sigma(\text{np}) = 10.0 \pm 1.5$ (1958NU40).

5. $^{11}\text{B}(\gamma, \pi^+)^{11}\text{Be}$ $Q_{\text{m}} = -151.09$

See (1962DY1A, 1963NE07).

^{11}B
(Figs. 2 and 4)

GENERAL:

Shell model: (1956KU1A, 1957KU58, 1960BI07, 1960TA1C, 1961BA1E, 1961BA1F, 1961KO1A, 1961KU1C, 1961TR1B, 1961UM1A, 1964AM1D, 1964NE1E, 1965CO25, 1965FA1B, 1965FA1C, 1966HA18, 1966MA1P, 1967CO32, 1967FA1A, 1968KU1D).

Collective model: (1959BR1E, 1961CL10, 1962CL13, 1964MA1G, 1965NE1B, 1966EL08, 1966MI1F, 1967RI1B, 1968GO01).

Ground state properties: (1962BE1D, 1963BE36, 1964LI14, 1964LI1B, 1964ST1B, 1965HU13, 1966RI12, 1966WI1E, 1967BA78, 1967RH1C, 1967SH05, 1968BA2G).

Other: (1963SE1K, 1964OL1A, 1964TH03, 1966WI1F, 1967BA1E, 1967PO1J).

$$\mu = +2.6885 \text{ nm (1965FU1G);}$$

$$Q = +0.04 \text{ b (1965FU1G), } +0.0372 \text{ b (1966ST12).}$$

1. $^6\text{Li}(^6\text{Li}, \text{p})^{11}\text{B}$ $Q_{\text{m}} = 12.220$

Angular distributions of the protons to ^{11}B states below $E_{\text{x}} = 9.3$ MeV have been measured for $E(^6\text{Li}) = 2.4$ to 9.0 MeV (1966KI09). Gamma spectra have been studied at $E(^6\text{Li}) = 2.6$ MeV. Relative populations to ^{11}B levels are reported (1962BE24). See also (1960SH01, 1962BE16, 1963LE09).

Table 11.3: Energy levels of ^{11}B

E_x (MeV \pm keV)	J^π	τ_m or Γ (sec) (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-$	stable		1, 2, 7, 8, 9, 13, 14, 15, 16, 17, 18, 25, 26, 28, 29, 30, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 48, 50, 51, 52, 53, 54
2.1244 ± 0.7^a	$\frac{1}{2}^-$	0.122 ± 0.005 eV	γ	1, 7, 8, 9, 14, 15, 16, 25, 26, 29, 30, 34, 35, 36, 39, 42, 43, 45, 46, 52, 53
4.4444 ± 1.4^a	$\frac{5}{2}^-$	0.54 ± 0.05 eV	γ	1, 2, 7, 8, 9, 14, 15, 16, 25, 26, 30, 34, 36, 42, 43, 45, 46, 52, 53
5.0189 ± 1.7^a	$\frac{3}{2}^- (\frac{1}{2}^-)$	1.73 ± 0.14 eV	γ	1, 7, 8, 9, 14, 15, 16, 25, 26, 30, 34, 36, 42, 43, 45, 46, 52, 53
6.7429 ± 1.8^a	$\frac{7}{2}^-$	$\tau_m < 3 \times 10^{-13}$	γ	1, 2, 7, 14, 15, 25, 26, 34, 36, 42, 43, 45
6.7928 ± 1.8^a	$(\frac{1}{2}, \frac{3}{2})^+$	$\tau_m < 5 \times 10^{-14}$	γ	1, 7, 14, 16, 25, 26, 29, 34, 43
7.296 ± 4	$(\frac{3}{2}, \frac{5}{2})^+$	1.0 ± 0.5 eV	γ	1, 7, 14, 16, 25, 26, 30, 34, 36
7.996 ± 6	$\frac{3}{2}^+$	$\tau_m < 5 \times 10^{-14}$	γ	1, 7, 14, 16, 25, 29, 34
8.566 ± 4	$\leq \frac{5}{2}^-$	2.0 ± 0.7 eV	γ	1, 7, 14, 25, 34
8.925 ± 4	$\frac{5}{2}^-$	4.0 ± 0.6 eV	γ, α	1, 2, 14, 25, 30, 34, 36
9.185 ± 1	$\frac{7}{2}^+$	3 eV	γ, α	1, 2, 14, 25
9.274 ± 1	$\frac{5}{2}^+$	7 eV	γ, α	1, 2, 14, 25
9.87 ± 20	$\frac{3}{2}^+$	250	α	6, 14, 36
10.25 ± 35	$(\frac{3}{2}^-)$	200	α	6, 14
10.33 ± 15	$\frac{5}{2}^- (\frac{7}{2}^-)$	54 ± 17	α	6, 14, 25
10.595 ± 6	$\frac{7}{2}^+ (\frac{5}{2}^+)$	90	α	6, 14
11.0	$\frac{5}{2}^- (\frac{3}{2}^-)$	4500	α	6
11.266 ± 7	$\frac{7}{2}^+, \frac{9}{2}^+$	100	α	6, 14
11.462 ± 10		70 ± 30	α	6, 14
11.60 ± 35		150 ± 50	n, α	3, 6
11.884 ± 12	$(\frac{5}{2}^+)$	150 ± 50	n, α	3, 6, 14, 24
12.0 ± 130		≈ 1000	α	6, 14
12.565 ± 12	$(T = \frac{3}{2})$	145 ± 30	α	6, 14, 16
13.02 ± 80	$\frac{1}{2}^-; T = \frac{3}{2}$	358 ± 60		51
13.15 ± 100	$\geq \frac{11}{2}^+$	270 ± 50	n, α	3, 6, 14, 19, 20, 24, 34, 36
14.04 ± 100		500 ± 200	n, α	3, 6, 19, 20, 24

Table 11.3: Energy levels of ^{11}B (continued)

E_x (MeV \pm keV)	J^π	τ_m or Γ (sec) (keV)	Decay	Reactions
14.563 \pm 15		< 120	n, α	3, 6, 14, 20, 24, 36
15.12 \pm 100		750	n, α	3, 20, 24
15.80 \pm 130		180 \pm 100	n, α	3, 6, 20
16.7 \pm 300			n, α , d, p, (t)	3, 10, 11, 24
(16.9)			d, p, t	11, 14
17.53 \pm 30			n, α , d, p	3, 11, 24
18.3 \pm 800	$\frac{1}{2}^+$			41

^a (1966BR18).

2. $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$

$$Q_m = 8.664$$

Three resonances for capture radiation are reported at $E_\alpha = 401, 819$ and 958 keV, corresponding to ^{11}B levels at $8.92, 9.19$ and 9.27 MeV: see Table 11.4. No others appear for $E_\alpha < 2.5$ MeV (1954HE22). See also (1960SI02). Angular distributions and correlations have been studied by (1959JO25, 1962GR07) with the following conclusions:

$E_x = 9.27$ MeV. The angular distribution of the ground-state transition requires $J^\pi = \frac{3}{2}^\pm$ or $\frac{5}{2}^\pm$. Correlations in the cascades eliminate all but $J^\pi = \frac{5}{2}^+$, formed by p-wave. $\theta_\alpha^2 \approx 0.1$. The g.s. radiation is E1, $\Gamma_\gamma = 0.6$ eV [cm].

$E_x = 4.44$ MeV. The angular distribution in the cascade $9.28 \rightarrow 4.44 \rightarrow$ g.s. determines $J = \frac{5}{2}$; stripping results give odd parity. The E2/M1 amplitude ratio in the transition to the ground state is $x = -0.2 \pm 0.02$ ¹.

$E_x = 6.74$ MeV. Angular distributions and branching ratios indicate $J^\pi = \frac{7}{2}^\pm$, with $\frac{7}{2}^-$ favored (and indicated by stripping). The transition $6.74 \rightarrow 4.44$ is either pure M1 or pure E2. The branching ratio $6.74 \rightarrow$ g.s./ $6.74 \rightarrow 4.44$ is 1.75 ± 0.1 (1962GR07), 4.9 (1958FE70). See also $^9\text{Be}(^3\text{He}, p)^{11}\text{B}$.

$E_x = 9.19$ MeV. Angular correlations in the cascade transitions require $J^\pi = \frac{7}{2}^\pm$; the ground-state transition fixes $J^\pi = \frac{7}{2}^+$ formed by f-waves, $\theta^2 = 0.04$, with an M2/E3 amplitude of 0.9 and an E3 strength of 6 to 17 Weisskopf units. See, however, $^9\text{Be}(^3\text{He}, p)^{11}\text{B}$.

$E_x = 8.93$ MeV. The angular distribution of the ground-state transition yields $J^\pi = \frac{3}{2}^+$ or $\frac{5}{2}^\pm$. Consideration of the transition strength favors $\frac{5}{2}^-$.

¹ The sign and magnitude are in good accord with IPM, if collective enhancement is included (PO66E).

Table 11.4: Resonances in ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ ^a

E_r^b (keV)	$\Gamma_{\text{c.m.}}^b$ (keV)	${}^{11}\text{B}^*$ (MeV)	J^π	$\omega\Gamma_s^c$ (eV)	Percentage decay ^d to states of ${}^{11}\text{B}^*$ at			
					0	4.44	6.74	6.79
401	< 1	8.919	$(\frac{5}{2})^-$	0.04	93 ± 5	2.3 ± 1		
819 ± 1^e	$\approx 4^f$	9.185	$\frac{7}{2}^+$	0.55	0.9 ± 0.3	82.8 ± 2.0	12.8 ± 0.4	< 1.3
958 ± 1^e	7	9.274	$\frac{5}{2}^+$	3.5	19.7 ± 1.0	67.5 ± 2.0	12.8 ± 0.7	< 0.6

^a See also Tables 11.9 and 11.13.

^b (1951BE13). See also (1954HE22).

^c $\omega\Gamma_\gamma\Gamma_\alpha/\Gamma$, in c.m. (1951BE13, 1959JO25). See also (1965OL03).

^d (1962GR07) $\theta = 55^\circ$; see also (1958FE70, 1959JO25), and Table 11.9.

^e (1962GR07) find 815, 951 keV; (1957BR18) find 957 ± 2 keV.

^f $\Gamma \approx 3$ eV (1965OL03).

3. ${}^7\text{Li}(\alpha, n){}^{10}\text{B}$

$$Q_m = -2.792$$

$$E_b = 8.664$$

Observed resonances are listed in Table 11.5. Comparison with ${}^{10}\text{B}(n, \alpha){}^7\text{Li}$ indicates that the resonances at $E_\alpha = 5.15$ MeV ($E_x = 12.0$ MeV) has $J^\pi = \frac{3}{2}^-$ or $\frac{5}{2}^+$, $\Gamma_n \approx 20$ keV, $\Gamma_\alpha \approx 300$ keV formed by $l_n = 0$ or 1 (1959GI47).

4. ${}^7\text{Li}(\alpha, p){}^{10}\text{Be}$

$$Q_m = -2.564$$

$$E_b = 8.664$$

See ${}^{10}\text{Be}$ in (1966LA04).

5. ${}^7\text{Li}(\alpha, t){}^8\text{Be}$

$$Q_m = -2.562$$

$$E_b = 8.664$$

See ${}^8\text{Be}$ on (1966LA04).

6. ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$

$$E_b = 8.664$$

Elastic scattering has been studied for $E_\alpha = 1.6$ to 12 MeV by (1966CU02). The inelastic scattering, leading to ${}^7\text{Li}^*(0.48)$ has been studied by (1954HE22: $E_\alpha = 1.5$ to 3.5 MeV), (1954LI48: $E_\alpha = 1.2$ to 2.8 MeV), (1957BI84: $E_\alpha = 1.5$ to 6.0 MeV), (1966CU02: $E_\alpha = 1.6$ to 12 MeV); see Table 11.6.

Table 11.5: Thresholds and resonances in ${}^7\text{Li}(\alpha, n){}^{10}\text{B}$

(1957BI84)		(1959GI47)	(1963ME08)	
E_α (MeV \pm keV)	Γ_{lab} (keV)	E_α (MeV \pm keV)	E_α (MeV \pm keV)	E_x (MeV \pm keV)
4.379 \pm 6 (thresh.)	220	5.15 \pm 70 (5.64) 7.15 ^a	4.380 \pm 20	thresh.
4.7			[4.72]	11.67 \pm 100
5.15 \pm 80			[5.22]	11.99 \pm 100
			5.5	thresh.
			7.05	13.15 \pm 100
			[8.44]	14.04 \pm 100
			[9.21]	14.53 \pm 50
			10.14	15.12 \pm 100
			[11.33]	(15.88 \pm 200)
			11.90	thresh.
			12.56	(16.7 \pm 300)
			13.92	17.53 \pm 30
			14.53	thresh.

^a The width of this resonance is large.

A detailed fit to the elastic and inelastic data below the first (α, n) threshold, $E_\alpha = 4.38$ MeV, requires at least seven resonances (Table 11.7) (1966CU02). For the 9.87 MeV level, the weakness of the elastic effect eliminates $J = \frac{1}{2}$, while the reduced width in the inelastic channel eliminates $J^\pi = \frac{3}{2}^-$ and $J \geq \frac{5}{2}$; thus $J^\pi = \frac{3}{2}^+$. Of the next two levels, $E_x = 10.25$ and 10.32 MeV, the first influences mainly the inelastic yield, while the second (seen also in ${}^{10}\text{B}(\text{d}, \text{p}){}^{11}\text{B}$) appears only in the elastic scattering. For the lower, possible assignments are $J = \frac{1}{2}^\pm, \frac{3}{2}^\pm$ or $\frac{5}{2}^\pm$, with $\frac{3}{2}^-$ preferred; the upper has $J^\pi = \frac{5}{2}^-$ or $\frac{7}{2}^-$. The narrow level at $E_x = 10.60$ MeV, also seen in ${}^9\text{Be}({}^3\text{He}, \text{p}){}^{11}\text{B}$ shows no effect on elastic scattering at $\theta_{\text{cm}} = 141^\circ$, indicating $l = 3$; $J^\pi = \frac{7}{2}^+$ is preferred, with $J = \frac{5}{2}^+$ possible. A broad maximum in the inelastic cross section at $E_\alpha \approx 3.6$ MeV may indicate a level at $E_x = 11.0$ MeV. An underlying general rise is ascribed to a $J^\pi = \frac{1}{2}^+$ level, with $\Gamma \approx 4.4$ MeV. With this assumption, the elastic yields require $J^\pi = \frac{3}{2}^-$ or $\frac{5}{2}^-$ for the 11.0 MeV level, with $\frac{5}{2}^-$ preferred. The narrow level at $E_\alpha = 4.1$ MeV, $E_x = 11.27$ MeV, appears only weakly in the inelastic cross section; it is seen also in ${}^9\text{Be}({}^3\text{He}, \text{p}){}^{11}\text{B}$. The elastic scattering indicates $J^\pi = \frac{7}{2}^+$ or $\frac{9}{2}^+$.

Table 11.6: Structure in ${}^7\text{Li}(\alpha, \alpha'){}^7\text{Li}^*$ and ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$

A		B	C		D			
E_α (keV)	Γ_{cm} (keV)	E_α (keV)	E_α (keV)	Γ_{cm} (keV)	E_α^{a} (keV)	E_α^{b} (keV)	Γ_{cm} (keV)	E_x (MeV \pm keV)
1889 ± 10	125 ± 10	1910	1910 ± 20	160	1920 ± 30		130 ± 30	9.89 ± 20
2500 ± 30	≈ 155	2460	2490 ± 50	220	2480 ± 50		150 ± 40	10.24 ± 35
						2630 ± 30	80 ± 30	10.34 ± 20
		3060	3060 ± 30	100	3032 ± 10	3032	70 ± 10	10.595 ± 6
			3600 ± 100	670	3600 ± 50		≥ 900	10.96 ± 35
						4120 ± 30	90 ± 50	11.29 ± 20
					4430 ± 50	4430		$(11.49)^{\text{e}}$
					4600 ± 50		150 ± 50	11.59 ± 35
					5050 ± 30		150 ± 50	11.88 ± 20
						5300 ± 200	≈ 1000	12.04 ± 130
						5500 ± 100	60 ± 50	$(12.17)^{\text{e}}$
					6100 ± 30		150 ± 50	12.55 ± 20
					6850 ± 60		270 ± 50	13.03 ± 45
					$(7200 \pm 50)^{\text{c}}$		50 ± 50	$(13.25)^{\text{e}}$
						7800 ± 100	500 ± 200	$(13.63)^{\text{e}}$
					$(8450 \pm 200)^{\text{d}}$		500 ± 200	(14.04 ± 130)
					$(9450 \pm 200)^{\text{d}}$		≤ 250	(14.68 ± 130)
						9950 ± 200	500 ± 200	$(15.00)^{\text{e}}$
					$(11200 \pm 200)^{\text{d}}$		180 ± 100	(15.80 ± 130)

 A: (1954LI48): ${}^7\text{Li}(\alpha, \alpha'\gamma){}^7\text{Li}$.

 B: (1954HE22): ${}^7\text{Li}(\alpha, \alpha'\gamma){}^7\text{Li}$.

 C: (1957BI84): ${}^7\text{Li}(\alpha, \alpha'\gamma){}^7\text{Li}$.

D: (1966CU02): see also Table 11.7.

^a ${}^7\text{Li}(\alpha, \alpha'\gamma){}^7\text{Li}$: $\sigma(\text{total})$.

^b ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$.

^c Anomaly in angular distribution.

^d ${}^7\text{Li}(\alpha, \alpha'){}^7\text{Li}$: $\theta = 60^\circ$.

^e ${}^7\text{Li}(\alpha, \text{n}){}^{10}\text{B}$ threshold.

Table 11.7: Resonance parameters in ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$ ^a

E_r (MeV)	E_x (MeV)	J^π	$\Gamma_{\text{c.m.}}$ (keV)	Γ_i/Γ_e ^b
1.88	9.86	$\frac{3}{2}^+$	250	4.0
2.50	10.25	$\frac{3}{2}^-$	200	0.04
2.60	10.32	$\frac{5}{2}^-$	100	0
3.03	10.60	$\frac{7}{2}^+$	90	1.0
3.54	11.0	$\frac{5}{2}^-$	4500	1.5
^c		$\frac{1}{2}^+$	4000	4.6
4.10	11.27	$\frac{9}{2}^+$	100	0

^a (1966CU02); assignments not unique: see text.

^b Width ratio: inelastic/elastic.

^c Broad level in background: $E_\lambda(\text{c.m.}) = 1.71$.

Analysis of the structure above $E_\alpha = 4.38$ MeV is rendered difficult by the increasing number of open channels. Anomalies observed at $E_\alpha = 4.4, 5.5, 7.2, 7.8$ and 9.95 MeV coincide with thresholds in ${}^7\text{Li}(\alpha, n){}^{10}\text{B}^*$ (1966CU02).

7. ${}^7\text{Li}({}^6\text{Li}, d){}^{11}\text{B}$ $Q_m = 7.192$

The lifetimes of ${}^{11}\text{B}^*(6.79, 7.30, 8.00)$ are less than 50 fsec (1967THZX). Gamma spectra indicate that all bound levels are populated at $E({}^6\text{Li}) = 2.6$ MeV (1962BE24). Angular distributions are not symmetric about 90° ; an α -transfer process is suggested (1961MO02). Angular distributions have also been obtained at $E({}^7\text{Li}) = 3.78$ to 5.95 MeV for the deuterons to ${}^{11}\text{B}^*(0, 2.12, 4.44, 5.02, 6.74 + 6.79, 7.30)$ (1967KI03), and at $E({}^7\text{Li}) = 3.3$ MeV (${}^{11}\text{B}^* = 0, 2.12, 4.44, 5.02$) (1967GA06). See also (1966RO1E, 1966RO1F).

8. ${}^7\text{Li}({}^7\text{Li}, t){}^{11}\text{B}$ $Q_m = 6.197$

Triton-gamma coincidence studies yield the following branching ratios: ${}^{11}\text{B}^*(5.02) \rightarrow \text{g.s.}$ ($85 \pm 5\%$), $5.02 \rightarrow 2.12$ ($15 \pm 5\%$), $5.02 \rightarrow 4.44$ ($< 5\%$), $4.44 \rightarrow \text{g.s.}$ (100%), $4.44 \rightarrow 2.12$ ($< 5\%$) (1963CA09): see Table 11.9. The lifetimes of ${}^{11}\text{B}^*(6.79, 7.30, 8.00)$ are less than 0.5 msec (1967THZX). See also (1961MO02, 1962BE24, 1966RO1E, 1966RO1F, 1967WY1B).

Table 11.8: Energy levels of ^{11}B from $^9\text{Be}(^3\text{He}, p)^{11}\text{B}$

(1959HI69, 1963GR20)		(1965MA1E)	(1966BR18)	L^c
E_x (MeV \pm keV)	Γ (keV)	E_x (MeV \pm keV)	E_x (MeV \pm keV)	
0		0	0	0
2.144 ± 10		2.142 ± 10	2.1243 ± 0.9	0
4.459 ^a		4.454 ± 10	4.4434 ± 1.8	0
5.037 ± 10		5.033 ± 10	5.0187 ± 2.3	0
6.753 ± 10		6.743 ± 10	6.7411 ± 3.0	
6.805 ± 10		6.793 ± 10	6.7909 ± 3.1	1
7.299 ± 10		7.285 ± 10		
7.989 ± 10		8.018 ± 10		
8.567 ± 10		8.560 ± 10		0
8.923 ± 10		8.920 ± 10		0
9.189 ± 10		9.182 ± 10		
9.278 ± 10		9.268 ± 10		
9.87 ± 20	≈ 150			
(10.26)				
10.337 ± 15				
10.594 ± 12				
11.266 ± 7	< 30			
11.462 ± 10				
11.884 ± 12				
11.97				
12.565 ± 12^b	145 ± 30			
13.3 ± 100				
14.563 ± 15				

^a Values up to $E_x = 9.87$ MeV normalized to this group (1959HI69): see (1966BR18).

^b Possibly $T = \frac{3}{2}$: see text.

^c Angular momentum transfer (1960HI08).

Table 11.9: Gamma decay of ^{11}B levels ^a

E_i (MeV)	J_i^π	τ_m or Γ_γ (sec) (eV)	E_f (MeV)	Branch %	Mult.	χ^2 ^b
2.12	$\frac{1}{2}^-$	0.122 eV ^c	0	100		
4.44	$\frac{5}{2}^-$	0.54 eV ^c	0	100	M1	0.04 ^d
5.02	$(\frac{1}{2}^-) \frac{3}{2}^-$	1.73 eV ^c	2.12	< 0.5		
			0	85 ± 2 ^j	M1 ^c	
			2.12	15 ± 2		
6.74	$\frac{7}{2}^-$	< 3×10^{-13} ^e	4.44	< 0.3		
			0	70 ± 2	E2	0.4
			2.12	< 3		
			4.44	30 ± 2	M1, E2 ^g	
6.79	$(\frac{1}{2}, \frac{3}{2})^+$	< 5×10^{-14} ^f	5.02	< 1		
			0	71 ± 5 ^k		
			2.12	29 ± 5		
			4.44	< 8		
			5.02	< 8		
7.30	$\frac{3}{2}^+, \frac{5}{2}^+$	1.0 eV ^c	0	87 ± 2	E1	0.3
			2.12	< 1		
			4.44	5.5 ± 1		
			5.02	7.5 ± 1		
7.99	$\frac{3}{2}^+$	< 5×10^{-13} < 5×10^{-14} ^f	0	47 ± 2	E1	0.2
			2.12	53 ± 2	E1	0.4
			4.44	< 1		
			5.02	< 1		
8.57	$\leq \frac{5}{2}^{(-)}$	2.0 eV ^c	0	56 ± 2	M1, E2 ^c	
			2.12	30 ± 2		
			4.44	5 ± 1		
			5.02	9 ± 1		
8.93	$\frac{5}{2}^-$	4.0 eV ^c	0	95 ± 1	M1	0.7 ^h
			2.12	< 1		
			4.44	4.5 ± 0.5		
			5.02	< 1		

Table 11.9: Gamma decay of ^{11}B levels ^a (continued)

E_i (MeV)	J_i^π	τ_m or Γ_γ (sec) (eV)	E_f (MeV)	Branch %	Mult.	χ^2 ^b
9.19	$\frac{7}{2}^+$	$\Gamma_\gamma = 0.3 \text{ eV}$	6.74	< 1	M2, E3 ⁱ	
			6.79	< 1		
			0	0.9 ± 0.3		
			4.44	82.8 ± 2		
			6.74	12.8 ± 0.4		
9.27	$\frac{5}{2}^+$	$\Gamma_\gamma = 2.3 \text{ eV}^g$	6.79	< 1.3	E1	
			0	19.7 ± 1.0		
			4.44	67.5 ± 2.0		
			6.74	12.8 ± 0.7		
			6.79	< 0.6		

^a From $^9\text{Be}(^3\text{He}, p)^{11}\text{B}$ and $^{10}\text{B}(d, p)^{11}\text{B}$ (1965OL03 and references therein).

^b χ^2 = maximum intensity of quadrupole radiation in dipole transitions or of M1 in E2 transitions.

^c See Table 11.13.

^d Amplitude E2/M1 = -0.2 ± 0.02 (1962GR07): see (PO66E).

^e (1966WA10).

^f (1967THZX).

^g From $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$ (1962GR07). See Table 11.4.

^h 0.6 % E2 (1962GR07). See also (1966GO12).

ⁱ Amplitude M2/E3 = 0.9 (1962GR07).

^j $88 \pm 2, 12 \pm 2$ (1968EA03).

^k $67 \pm 3, 33 \pm 3$ (1968EA03).

9. $^9\text{Be}(d, \gamma)^{11}\text{B}$

$$Q_m = 15.819$$

Radiative transitions have been observed to $^{11}\text{B}^*(0, 2.12, 4.44 + 5.02)$. For $E_d = 0.5$ to 1.4 MeV, the intensity ratios are $1:(0.30 \pm 0.08):(0.78 \pm 0.15)$. At $E_d = 1.4$ MeV, $\sigma(\text{g.s.}) = 4.3 \pm 0.8 \mu\text{b}$: the g.s. radiation is isotropic at $E_d = 0.7$ and 1.2 MeV (1966ZI01). The 90° differential cross section has been measured for $E_d = 0.5$ to 5.5 MeV for ground state γ -rays: no resonance structure is observed. The peak cross section is about $8 \mu\text{b}$. Angular distributions are anisotropic for the higher bombarding energies (1963SU09, 1966SU05, 1966SU1C). See also (1960SU09, 1961SU17).

10. ${}^9\text{Be}(d, n){}^{10}\text{B}$

$$Q_m = 4.363$$

$$E_b = 15.819$$

The cross section follows the Gamow function for $E_d = 70$ to 110 keV (1955RA14). The fast neutron and γ -ray yield rise smoothly to $E_d = 1.8$ MeV except for a possible “resonance” at $E_d \approx 0.94$ MeV (the fast neutron yield then remains approximately constant to 3 MeV) (1949EV1A, 1955BO1A, 1957SH65, 1960BA46, 1963KO15, 1965SI12). This resonance is

observed in the total neutron yield and in the yield of the fast neutrons to each of the first five states of ${}^{10}\text{B}$ (1957SH65; see however, (1959NE1A)). The polarization angular distributions of the neutrons to ${}^{10}\text{B}^*(0, 0.72, 1.74, 2.15)$ have been measured at $E_d = 1.57, 2.06$ and 2.48 MeV (1967MI1F). See also ${}^{10}\text{B}$ in (1966LA04) and (1965BU10, 1967SC43).

11. (a) ${}^9\text{Be}(d, p){}^{10}\text{Be}$

$$Q_m = 4.590$$

$$E_b = 15.819$$

(b) ${}^9\text{Be}(d, \alpha){}^7\text{Li}$

$$Q_m = 7.154$$

(c) ${}^9\text{Be}(d, t){}^8\text{Be}$

$$Q_m = 4.592$$

(d) ${}^9\text{Be}(d, 2\alpha){}^3\text{H}$

$$Q_m = 4.687$$

(e) ${}^9\text{Be}(d, tn){}^7\text{Be}$

$$Q_m = -14.304$$

(1952CA19) reports broad maxima in the 90° yield of the ground-state protons at $E_d \approx 0.9, (1.3)$ and 2.1 MeV (see also (1961IS01)). (1957MC35) observe broad resonances at 1.3 and possibly at 1.8 MeV in the yield of 3.37 MeV γ -rays (from ${}^{10}\text{Be}^*$) in the range $E_d = 1.0$ to 5.6 MeV. No resonances are observed in the yield of 6 MeV γ -rays for $E_d = 2.0$ to 5.6 MeV. The yields of protons (at 40°) to the ground state and to the first excited state both decrease slowly and monotonically with energy in the range $E_d = 3.8$ to 6.3 MeV (1961RE03, 1961RE04). See also (1962BI11, 1966AM1C, 1967FA03) and (1959AJ76). Polarization of the protons has been studied at $E_d = 1$ to 6 (1967BL02), 1.6 (1961VA03), 6 (1960HI09), 7.8 (1962GR14), $7.8, 8.9, 10$ and 13.8 (1967SA07), 8.9 (1959HI1E), 10 (1962AL10), 11 (1964PA1E), 13.8 (1962PA12), 15 (1964RE04) and 21 MeV (1963BO1J). See also (1960LU04, 1966MI1E).

The yields of α -particles (reaction (b)) to the ground and first excited states have been measured in the range $E_d = 0.5$ to 2.3 MeV. There is no clear indication of resonance structure. The cross sections for the two groups are quite similar but their angular distributions are different over the entire energy range (1962BI11). Yields have also been measured for $E_d = 8$ to 12.4 MeV (α_0, α_1) and 9 to 12.4 MeV (α_2) (1966DO1A). See also ${}^7\text{Li}$ in (1966LA04) and (1966AM1C).

The cross section for reaction (c) has been measured for $E_d = 0.15$ to 0.62 MeV by (1952DE24), for $E_d = 0.6$ to 1.5 MeV by (1955JU10, 1955JU1B), for $E_d = 3.8$ to 6.3 MeV by (1961RE03, 1961RE04) and at several energies in the range $E_d = 3$ to 19 MeV by (1955HE83). See also (1962BI11, 1966AM1C). The forward yield of tritons shows a peak at $E_d = 1.38$ MeV, and possibly, at 0.87 MeV (1955JU10, 1955JU1B, 1958JU38). In the range $E_d = 3.8$ to 6.3 MeV,

the 40° differential cross section for ground state protons is approximately constant (1961RE03, 1961RE04).

The direct three-body reaction [reaction (d)] does not appear to occur (1953GE01). For reaction (e) see (1955HE83). See also (1962WI15, 1965MA57).

12. ${}^9\text{Be}(d, d){}^9\text{Be}$

$$E_b = 15.819$$

The differential cross section for elastic scattering has been measured for $E_d = 0.4$ to 1.8 MeV ($\theta = 90^\circ, 126^\circ, 163.5^\circ$). No anomalies were observed (1963RE16: see, however, (1956JU17)). Asymmetries have been measured with polarized deuterons at $E_d = 11.7$ MeV (1966DO1B). See also (1960BU25, 1963NE1H) and ${}^9\text{Be}$ in (1966LA04).

13. ${}^9\text{Be}(t, n){}^{11}\text{B}$

$$Q_m = 9.561$$

See (1962SE1A).

14. ${}^9\text{Be}({}^3\text{He}, p){}^{11}\text{B}$

$$Q_m = 10.325$$

$$Q_0 = 10.344 \pm 0.013 \text{ (1964MA57);}$$

$$Q_0 = 10.3221 \pm 0.0023 \text{ (1967OD01).}$$

Proton groups have been observed to 23 states of ${}^{11}\text{B}$: see Table 11.8 (1959HI69, 1963GR20, 1966BR18). Angular distributions of many of these proton groups have been observed at $E({}^3\text{He}) = 1.0$ to 3.0 MeV (1967CO03), 2 MeV (1963WE08), 4.5 MeV (1959WO53), 5.7 MeV (1959HI69) and 8.8 and 10.2 MeV (1960HI08). Of the first ten groups, seven show stripping patterns: the exceptions are ${}^{11}\text{B}^*(6.74, 7.30, 8.00)$ (1960HI08). The narrow level at $E_x = 12.565$ MeV does not appear in ${}^{10}\text{B}(d, p){}^{11}\text{B}$ and may therefore have $T = \frac{3}{2}$ (1963GR20: see, however, ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$). See also (1959AJ76) and ${}^{12}\text{C}$.

Gamma-ray branching ratios and multiplicities for ${}^{11}\text{B}$ levels up to $E_x = 9.19$ MeV have been extensively studied by (1958FE70, 1961DO03, 1964AL22, 1965OL03): see Table 11.9. The following remarks on individual levels derive largely from (1965OL03): see also ${}^{10}\text{B}(d, p){}^{11}\text{B}$.

$\underline{E_x = 9.19 \text{ MeV. } \Gamma_\gamma/\Gamma = 0.1_{-0.05}^{+0.2}}$. From ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$, $\Gamma_s = 0.275 \text{ eV}$, $J^\pi = \frac{7}{2}^+$. Then $\Gamma_\alpha = 2.8_{-1.8}^{+2.8} \text{ eV}$, $\Gamma_\gamma = 0.3_{-0.01}^{+0.1} \text{ eV}$. The ground-state transition of 0.9% is 0.45 E3 and 0.55 M2 leading to M2 and E3 strengths of 0.3 ± 0.1 and 78 ± 26 Weisskopf units, respectively.

$\underline{E_x = 8.93 \text{ MeV. } \Gamma_\gamma/\Gamma = 1.08 \pm 0.12}$. From ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$, $J^\pi = \frac{3}{2}^+, \frac{5}{2}^+, \frac{5}{2}^-$. The internal pair correlation excludes $J^\pi = \frac{3}{2}^+, \frac{5}{2}^+$, and confirms $\frac{5}{2}^-$. The ground-state transition is then M1 with

0.6% E2. With $\Gamma_s = 0.025$ eV, $0.025 \leq \Gamma_\alpha \leq 0.03$ eV, $\Gamma_\gamma \geq 0.15$ eV. See (1966GO12) and $^{11}\text{B}(e, e')^{11}\text{B}^*$.

$E_x = 8.57$ MeV. Correlation of internal pairs indicate that the g.s. transition is M1 + E2 or E1 + M2, $J^\pi \leq \frac{5}{2}^+$ or $\leq \frac{7}{2}^-$; the lifetime to $^{11}\text{B}^*(2.12)$ excludes $\frac{7}{2}^-$. If the level has even parity, the required M2 admixture is excessive. $J^\pi \leq \frac{5}{2}^-$ is favored. The odd parity assignment agrees with stripping results in the present reaction, but disagrees with the results from $^{10}\text{B}(d, p)^{11}\text{B}$. See $^{11}\text{B}(e, e')^{11}\text{B}^*$.

$E_x = 8.00$ MeV. Transitions to $^{11}\text{B}_{\text{g.s.}}$ and (2.12) are predominantly E1; thus $E_x = 8.00$ MeV has even parity, and the odd parity of $^{11}\text{B}^*(2.12)$ is confirmed. The transition to $^{11}\text{B}^*(2.12)$ is not isotropic, so $J^\pi = \frac{3}{2}^+$. Comparison of branching ratios with analogous case in ^{11}C suggests some deviation from charge symmetry.

$E_x = 7.30$ MeV. The g.s. transition is mainly E1, so $J^\pi \leq \frac{5}{2}^+$. The assignment $\frac{1}{2}^+$ is excluded by the strength of (7.30 \rightarrow 4.44).

$E_x = 6.79$ MeV. The allowed β -decay from ^{11}Be indicates $J^\pi \leq \frac{7}{2}^+$. The relatively strong γ -branch to $^{11}\text{B}^*(2.12)$ favors $\frac{1}{2}^+$, $\frac{3}{2}^+$. (1968EA03) finds that all γ 's from this level are isotropic, suggesting $J^\pi = \frac{1}{2}^+$, but not excluding $\frac{3}{2}^+$.

$E_x = 6.74$ MeV. From $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$, $J = \frac{7}{2}$. Internal pairs indicate practically pure E2 g.s. radiation: $J^\pi = \frac{7}{2}^-$.

$E_x = 5.02$ MeV. Stripping work fixes the parity as odd. The internal pair correlations permit M1, E2 for the g.s. transition, $J^\pi \leq \frac{7}{2}^-$; the lifetime $\tau_m < 0.5$ psec excludes $J = \frac{7}{2}$. The comparatively strong branch to $^{11}\text{B}^*(2.12)$ argues against $J = \frac{5}{2}$, therefore $J^\pi \frac{1}{2}^-$ or $\frac{3}{2}^-$ (1961DO03). Angular correlation studies indicate a $P_2(\cos\theta)$ term of $(10 \pm 5)\%$; if this term is real, $J \neq \frac{1}{2}$ (1968EA03).

$E_x = 4.44$ MeV. The g.s. transition is predominantly M1, consistent with $J^\pi = \frac{5}{2}^-$ (see $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$).

$E_x = 2.12$ MeV. The lifetime (4.6 fsec) demands dipole radiation. The E1 transition from $^{11}\text{B}^*(8.00)$ requires odd parity; the weakness of the 4.44 ($\frac{5}{2}^-$) to 2.12 transition leaves only $J^\pi = \frac{1}{2}^-$ (1961DO03).

The properties of the first five levels of ^{11}B are consistent with ascription to p^7 ; there is evidence that $^{11}\text{B}^*(8.93)$ also belongs to p^7 , while $^{11}\text{B}^*(8.57)$ appears more likely to belong to a higher configuration.

See also (1960EL1C, 1961EL1A, 1962WE1C, 1963VA16, 1966WA1C).

15. $^9\text{Be}(\alpha, d)^{11}\text{B}$

$$Q_m = -8.028$$

At $E_\alpha = 28.3$ MeV, differential cross sections have been measured ($\theta = 15^\circ$ to 47°) for the deuteron groups to $^{11}\text{B}^*(0, 2.12, 4.44, 5.02)$ (1965KA14). See also (1955RA41, 1962WE1C, 1967ZE1A).

$$16. \text{}^9\text{Be}(\text{}^6\text{Li}, \alpha)\text{}^{11}\text{B} \quad Q_m = 14.347$$

Angular distributions and total cross sections have been determined for seven α -groups. In the range $E(^6\text{Li}) = 3$ to 4 MeV, the distributions change only slowly. A direct interaction mechanism is suggested (1961EL1A, 1961HO19, 1961LE01, 1961LE09, 1961RU1B, 1966SA04). Gamma radiation has been studied by (1963NO02, 1964CA18). Evidence of participation of a level near $E_x = 12.5$ MeV is reported by (1961CO33, 1961LE08). See also (1963BA1T, 1966BA1T, 1966RO1E, 1966RO1F, 1966RO1H).

$$17. (a) \text{}^9\text{Be}(\text{}^7\text{Li}, \alpha n)\text{}^{11}\text{B} \quad Q_m = 7.094$$

$$(b) \text{}^9\text{Be}(\text{}^{14}\text{N}, \text{}^{12}\text{C})\text{}^{11}\text{B} \quad Q_m = 5.547$$

For reaction (a) see (1964CA18). For reaction (b) see (1964BO1M).

$$18. \text{}^{10}\text{B}(n, \gamma)\text{}^{11}\text{B} \quad Q_m = 11.456$$

The thermal neutron capture cross section is 0.5 ± 0.2 b. The direct ground-state decay has $\Gamma_\gamma = 0.01$ eV (1957BA18). See also (1959AJ76).

$$19. \text{}^{10}\text{B}(n, n)\text{}^{10}\text{B} \quad E_b = 11.456$$

The thermal scattering cross section (bound) for B is 4.4 ± 0.2 bn (1958HU18). The coherent scattering amplitude (bound) for ^{10}B is $a = +1.4 \pm 1.5$ fm. Combined with the potential scattering amplitude of 5.0 fm (derived from (1955WI25)), this low value indicates a resonance level above the neutron binding energy (1965DO14, 1966DO11). According to (1960BI04) the total cross section $\sigma_s + \sigma_{n,\alpha}$ fits the expression $\sigma = 2.43 + 642/E^{1/2}$ up to 70 keV; the $1/\nu$ term is ascribed to $\sigma_{n,\alpha}$ and σ_s is taken to be 2.43 b: see also (1964ST25, 1965MO1J). Elastic scattering differential cross sections at $E_n = 0.55, 1.0$ and 1.5 MeV are analyzed in terms of phase shifts by (1955WI25): only s-waves appear at $E_n = 0.55$ MeV.

The total cross section (including $\sigma_{n,\alpha}$) shows broad maxima at $E_n = 1.9$ and 2.8 MeV (1951BO45) and at 4.3 MeV (1961FO07); an additional peak at $E_n = 0.2$ MeV may be indicated (1951BO45). Above $E_n = 5.5$ MeV, σ_{tot} is constant at 1.5 b to 16 MeV (1961FO07, 1964ST25). Polarization and differential cross sections have been measured for 70 energies in the range $E_n = 0.075$ to 2.25 MeV. The polarization is nearly constant for $E_n = 0.5$ to 2.25 MeV and shows no appreciable effect of the resonance at $E_n = 1.9$ MeV. Most of the width of this resonance appears to be in the α -channel (1967LA1N). See also (1959MA1C, 1960AN14, 1966AG1A, 1967CI1B) and (1959AJ76).

Table 11.10: Resonances in $^{10}\text{B} + \text{n}$ ^a

$^{10}\text{B}(\text{n}, \text{n}'\gamma)^{10}\text{B}$ (1960DA08)		$^{10}\text{B}(\text{n}, \alpha)^7\text{Li}$ (1961DA16)			$^{11}\text{B}^*$ (MeV)
E_{res} (MeV)	Γ (keV)	E_{res} (MeV)	Γ (keV)	Yield of	
					11.67 ^a
		0.53	140	α_0, α_1	11.94
1.93	260	1.86	570	$\alpha_0, \alpha_1, \text{t}$	13.18
(2.6)	broad	2.79	530	α_0, α_1	13.99
3.31	370	3.43	< 120	α_0, t	14.52
4.1		4.1	800	α_0, α_1	15.2
4.73					15.75
		5.7	broad	α_0, t	16.7
		6.4	broad	α_0, t	17.3

^a $E_{\text{res}} = 0.230$ MeV (1966MO09).

20. $^{10}\text{B}(\text{n}, \text{n}')^{10}\text{B}^*$

$$E_b = 11.456$$

The yield of 0.7 MeV γ -rays has been studied from threshold to $E_n = 5.2$ MeV: resonances are observed at $E_n = 1.93, (2.6), 3.31, 4.1$ and 4.73 MeV (see Table 11.10) (1960DA08). See also (1964ST25) and (1963GL1F, 1963GO1M).

21. $^{10}\text{B}(\text{n}, \text{p})^{10}\text{Be}$

$$Q_m = 0.228$$

$$E_b = 11.456$$

The thermal cross section is < 0.2 b (1958HU18); the cross section for fast pile neutrons is 3 mb (1948EG1A).

22. $^{10}\text{B}(\text{n}, \text{d})^9\text{Be}$

$$Q_m = -4.363$$

$$E_b = 11.456$$

See (1954RI15). See also (1963GO1M).

23. $^{10}\text{B}(\text{n}, \text{t})^8\text{Be}$

$$Q_m = 0.229$$

$$E_b = 11.456$$

The $^{10}\text{B}(\text{n}, \text{t})^4\text{He}^4\text{He}$ cross section has been measured for $E_n = 1.4$ to 8.2 MeV by (1961DA16). Fluctuations in the cross section are observed at some of the resonant energies in the $^{10}\text{B}(\text{n}, \alpha)$ reaction: see Table 11.10. See also (1964ST25, 1966JE1B).

24. $^{10}\text{B}(\text{n}, \alpha)^7\text{Li}$

$$Q_m = 2.792$$

$$E_b = 11.456$$

The “recommended” value of the thermal isotropic absorption cross section is 3837 ± 10 b (1964ST25), which includes recent determinations by (1963PR11: 3837 ± 9 b; (1961ME02): 3843 ± 17 b; (1960SA13, 1961SA02): 3838 ± 11 b; (1960SC15): 3848 ± 38 b). See also (1964AL05, 1965WY01). The cross section follows the $1/\nu$ law from 4 meV to 500 keV (1964ST25, 1965MO1J, 1966CO1K, 1967CO1N) except for a weak resonance at $E_n = 230$ keV (1966MO09). There appears to be some discrepancy between reported values of σ_{abs} and $\sigma(\text{n}, \alpha)$ (1966MO09).

Observed resonances are listed in Table 11.10 (1951PE18, 1957BI84, 1961DA16). Comparison of $\sigma(\text{n}, \alpha)$ and $\sigma(\text{n}, \text{n})$, at the $E_n = 0.53$ MeV resonance indicates $J^\pi = \frac{1}{2}^+$, $\frac{3}{2}^\pm$, or $\frac{5}{2}^\pm$. For the $E_n = 1.86$ MeV resonance, the minimum spin assignment consistent with the observed cross section is $J^\pi = \frac{11}{2}^+$ (1961DA16). The ratio of ground-state to excited state transitions varies monotonically with energy although the 0.53 MeV resonance appears to influence the cross-section ratio to some extent: see (1959AJ76, 1960YO02, 1963MA39, 1965MA65). Recent determinations of the branching ratio for thermal neutrons yield $(6.50 \pm 0.03)\%$ (see (1966TO01)), $(6.70 \pm 0.2)\%$ (1966MA2P) for the ground state function. At $E_n = 0.16$ MeV, the ratio is $(8.4 \pm 0.5)\%$ (1966MA2P).

See also (1957SE1B, 1961BE24, 1961MA1F).

25. $^{10}\text{B}(\text{d}, \text{p})^{11}\text{B}$

$$Q_m = 9.231$$

Proton groups reported by (1951VA1A, 1953EL12, 1961JA23, 1966BR18) are listed in Table 11.11. Angular distributions have been studied at many energies: see (1959AJ76) and (1960HA08: 0.17 to 0.25 MeV), (1957SJ66: 0.8 MeV), (1961RE01: 1.2 MeV), (1959GO69, 1960GO14: 1.25 MeV), (1967PO01: 1.75 to 3.0 MeV), (1965LE1B: 2 to 12 MeV), (1961PU1B: 3.0 MeV), (1960GO11, 1960GO25: 4.6 MeV), (1963MO07: 6 MeV), (1960BI08: 7.8 MeV), (1958ZE01: 8.1, 9.2, 10 MeV), (1962HI07: 10.1 MeV), (1960TA27: 11.4 MeV), (1967MO1Q: 12 MeV), (1965BA31: 13.5 MeV), (1961ZE02: 12.5, 15.5, 18.5, 21.5 MeV) and (1962SL04: 28 MeV).

The lowest five levels are formed by $l_n = 1$, except for $^{11}\text{B}^*(2.12)$ which appears to involve a spin-flip process (1957WI26, 1958EV01, 1958HE47, 1961LE1F). They are presumed to comprise the set $\frac{3}{2}^-, \frac{1}{2}^-, \frac{5}{2}^-, \frac{3}{2}^-, \frac{7}{2}^-$ expected as the lowest p^7 levels ($a/K \approx 4.0$) (1956KU1A, 1960BI08, 1965OL03). There is some disagreement about the parities of $^{11}\text{B}^*(8.57, 8.93)$: see $^9\text{Be}(^3\text{He}, \text{p})^{11}\text{B}$ and (1966GO12, 1967PO01). The levels at 9.19 and 9.28 MeV, $J^\pi = \frac{7}{2}^+$ and $\frac{5}{2}^+$, respectively, show strong $l = 0$ stripping and are ascribed to capture of a 2s neutron by ^{10}B (1960BI07, 1960BI08, 1967PO01). See also (1960MA32, 1964BA1G, 1967BA2J, 1967MO1N, 1967ST1H).

Table 11.11: ^{11}B levels from $^{10}\text{B}(\text{d}, \text{p})^{11}\text{B}$

E_x (MeV \pm keV)		E_x (MeV \pm keV)		$(2J + 1)\theta^2$	l_n	$(2J + 1)\theta^2$
(1951VA1A, 1953EL12)	(1961JA23)	(1966BR18)	l_n	(1962HI07)		(1960BI08)
0	0	0	1	0.120	1	1.00
$2.140 \pm 14^{\text{d}}$	2.128 ± 10	2.1246 ± 1.1				0.09
$4.464 \pm 14^{\text{d}}$	4.449 ± 8	4.4458 ± 2.1	1	0.048		0.46
$5.039 \pm 14^{\text{d}}$	5.023 ± 8	5.0192 ± 2.4	(1)	(0.010)		0.11
$6.765 \pm 13^{\text{a,d}}$		6.7439 ± 2.2	1	0.210^{c}	1	1.72
$6.815 \pm 13^{\text{a,d}}$		6.7938 ± 2.2				
7.298 ± 6			(2?)	(0.022)		
7.987 ± 9					isotropic	
8.568 ± 5	8.565 ± 10		(2?)		2	
8.927 ± 5	8.926 ± 10		1	0.186	0, 2	
9.191 ± 5	9.190 ± 10		0	0.242	0	
9.276 ± 5	9.278 ± 10		0	0.175	0	
$10.32 \pm 20^{\text{b}}$						

^a $6.752 \pm 6, 6.804 \pm 6$: see (1964AL22).

^b $\Gamma = 54 \pm 17$ keV (1953EL12).

^c See also (1967PO01).

^d Corrected by (1966BR18).

Studies of (p, γ) correlations are reported by (1959CR1A, 1959GA05, 1959GO69, 1960CR1A, 1960GO11, 1960GO14, 1960GO25, 1961RE01, 1965OL03, 1966GO1N: see also (1959AJ76)). The principal results are exhibited in Table 11.9 and are discussed in reaction 14.

(1963GR20: $E_d = 10.0$ MeV) report that the 11.27 MeV state is populated in this reaction while the narrow 12.57 MeV state observed in $^9\text{Be}(\text{}^3\text{He}, \text{p})^{11}\text{B}$ is not seen. This suggests $T = \frac{1}{2}$ and $\frac{3}{2}$, respectively, for these two states. See, however, $^7\text{Li}(\alpha, \alpha)^7\text{Li}$.

See also (1959BO1C, 1959BU1F, 1959HO1D, 1959LE1B, 1959TO1A, 1960BI1B, 1960NE1C, 1961BU16, 1961TE02, 1961ZE03, 1962ME1B, 1963GL1C, 1963MO1E, 1963SM05, 1963TA1A, 1963TA1C, 1964MA57, 1965BA31, 1965ST1E, 1966BA1X, 1966BA2R, 1967SC1K) and ^{12}C .

Polarization studies are reported by (1958HE47, 1959HI1E, 1960ER1A, 1960TA27, 1961ZI02, 1962PA12, 1962TA13, 1962ZI01, 1963BO1J, 1963ER1A, 1964BE08, 1964PA1E, 1965SZ01, 1965ZI1A, 1966BA2V).

26. $^{10}\text{B}(\text{t}, \text{d})^{11}\text{B}$

$Q_m = 5.199$

At $E_d = 5.5$ MeV, deuteron groups are observed to the ground state of ^{11}B and to states at $E_x = 2.126, 4.449, 5.027, 6.769, 6.806$ and 7.301 MeV (± 10 keV). All the angular distributions appear to be characteristic of $l_n = 1$ (1961BA10). See also (1963HO19, 1963KN02, 1967BI1E) and ^{13}C in (1970AJ04).

$$27. \ ^{10}\text{B}(\alpha, \ ^3\text{He})^{11}\text{B} \quad Q_m = -9.122$$

Not reported.

$$28. \ ^{10}\text{B}(^{14}\text{N}, \ ^{13}\text{N})^{11}\text{B} \quad Q_m = 0.903$$

See (1963TO1D, 1963TO1E, 1966CO27, 1966GA04, 1967CO1T).

$$29. \ ^{11}\text{Be}(\beta^-)^{11}\text{B} \quad Q_m = 11.513$$

The decay properties of ^{11}Be are exhibited in Table 11.12. The transition energy to the ground state is $E_{\beta(\text{max})} = 11.48 \pm 0.15$ MeV; $\tau_{1/2} = 13.57 \pm 0.15$ sec, $\log ft = 6.77$ (1958AL96, 1959WI49), $\tau_{1/2} = 14.1 \pm 0.3$ sec (1958NU40). The small ft values of the transitions to the $\frac{3}{2}^+$ states at 6.79 and 8.00 MeV indicate even parity [$J = \frac{1}{2}, \frac{3}{2}$ or $\frac{5}{2}$] for the ground state of ^{11}Be (1961DO03, 1964AL22).

$$30. \ ^{11}\text{B}(\gamma, \gamma)^{11}\text{B}$$

Mean gamma widths of low-lying levels obtained by resonance scattering and transmission studies are listed in Table 11.13. M1 transition strengths calculated in intermediate coupling are given by (1965CO25).

From a threshold determination, (1965SC04) finds $E_x = 2124 \pm 3$ keV for the first excited state. (1959HA1J) report $(2J + 1) \Gamma_\gamma = 0.6 \pm 0.1$ eV for $^{11}\text{B}^*(9.19)$: see, however, $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$. Resonance scattering from levels at 4.4, 5.0, 7.3 and 8.8 MeV is reported by (1962SE02). See also (1959FA1A, 1960BO23, 1960RE05, 1962BO17, 1964LO1C, 1967LO1B).

$$31. \ ^{11}\text{B}(\gamma, n)^{10}\text{B} \quad Q_m = -11.456$$

The cross section shows many peaks in the range $E_\gamma = 12$ to 28 MeV. The integrated cross section to 29 MeV is 68.6 ± 4 MeV \cdot mb (1965HA19). See also (1961BA1D, 1963CO1D).

Table 11.12: Beta decay of ^{11}Be (1958AL96, 1959WI49, 1964AL22)

$^{11}\text{B}^*$ (MeV)	$J\pi^a$	% betas	$\log ft$	E_γ (MeV \pm keV)	% gammas ^b
0	$\frac{3}{2}^-$	61	6.78		
2.12	$\frac{1}{2}^-$	29	6.68	2.121 ± 10	32
4.44	$\frac{5}{2}^-$	≤ 0.2	≥ 8.2		
5.02	$\frac{3}{2}^-, (\frac{1}{2}^-)$	≤ 0.2	≥ 8.2		
6.74	$\frac{7}{2}^-$		≥ 6.9		
6.79	$(\frac{1}{2}, \frac{3}{2})^+$	6.5	5.92	6.792 ± 6	4.4
				4.64 ± 20	2.1
7.30	$(\frac{3}{2}, \frac{5}{2})^+$	≤ 1.5	≥ 6.4		
8.00	$\frac{3}{2}^+$	4.1	5.55	7.97 ± 30	1.7
				5.86 ± 40	2.4
8.57	$\leq \frac{5}{2}^-$	≤ 0.3	≥ 6.3		
8.93	$\frac{5}{2}^-$	≤ 0.15	≥ 6.3		

^a Assignments from summary Table 11.3.

^b Relative to all β -transitions.

32. (a) $^{11}\text{B}(\gamma, p)^{10}\text{Be}$ $Q_m = -11.228$
 (b) $^{11}\text{B}(\gamma, d)^9\text{Be}$ $Q_m = -15.819$

See (1961DO08, 1962CH26, 1962LI13, 1962VO1C, 1963KI1C) for reaction (a), and (1962CH26, 1962VO1C, 1963BA1K) for reaction (b).

33. (a) $^{11}\text{B}(\gamma, t)^4\text{He} + ^4\text{He}$ $Q_m = -11.132$
 (b) $^{11}\text{B}(\gamma, \alpha)^7\text{Li}$ $Q_m = -8.664$

See (1955AJ61, 1959AJ76, 1962VO1C).

34. $^{11}\text{B}(e, e)^{11}\text{B}$

The charge-scattering radius is 1.55 fm (1959ME24). Magnetic elastic scattering at $\theta = 180^\circ$ shows strong M3 effects: the derived ratio of static M3/M1, $2.9 \pm 0.2 \text{ fm}^2$, suggests a $j - j$

Table 11.13: Gamma widths from $^{11}\text{B}(\gamma, \gamma)^{11}\text{B}$ and $^{11}\text{B}(e, e')^{11}\text{B}^*$ ^e

E_x (MeV)	J^π	Γ_γ to g.s. (eV)	Reaction	Ref.
2.12	$\frac{1}{2}^-$	0.14 ± 0.018	$\gamma\gamma$	(1958ME79)
		0.22 ± 0.06^c	$\gamma\gamma$	(1963VA10)
		0.11 ± 0.04	$\gamma\gamma$	(1964BO22)
		0.137 ± 0.020	$\gamma\gamma$	(1965KE05)
		0.114 ± 0.005	$\gamma\gamma$	(1965SC04)
		$0.130 \pm 0.031^{a,c}$	$\gamma\gamma$	(1965HA1K)
		0.16 ± 0.02	ee	(1966KO08)
		0.17 ± 0.034	ee	(1962ED02)
4.44	$\frac{5}{2}^-$	0.122 ± 0.005		mean
		0.56 ± 0.08	$\gamma\gamma$	(1958RA14)
		0.43 ± 0.095	$\gamma\gamma$	(1959CO95)
		$0.500 \pm 0.200^{a,c}$	$\gamma\gamma$	(1965HA1K)
		0.60 ± 0.2^c	ee	(1966KO08)
		1.1 ± 0.4^c	ee	(1962ED02)
		0.60 ± 0.09 (M1) + 0.016 ± 0.002 (E2)	ee	(1967SP02)
5.02	$\frac{3}{2}^-$	0.54 ± 0.05		mean
		1.1 ± 0.2^d	$\gamma\gamma$	(1959CO95)
		0.500 ± 0.200^a	$\gamma\gamma$	(1965HA1K)
		2.4 ± 0.8	ee	(1966KO08)
		3.7 ± 1.5	ee	(1962ED02)
		1.73 ± 0.14 (M1) < 0.0034 (E2) } 	ee	(1967SP02)
7.3	$(\frac{5}{2}^-)$	1.0 ± 0.5	ee	(1962ED02)
8.56	$(\frac{3}{2}^-)$	0.72 ± 0.30 (M1) 0.40 ± 0.10 (E2) }	ee	(1966SP02)
8.93	$\frac{5}{2}^-$	4.0 ± 0.6 (M1) ^b	ee	(1966SP02)

^a Preliminary values.

^b See also (1962ED02, 1966KO08) for higher levels.

^c Omitted from mean.

^d See (1967SP02).

^e See also Table 11.9.

coupling scheme for $^{11}\text{B}_{\text{g.s.}}$ (1966RA29). The quadrupole contribution to the elastic form factor is best accounted for by the undeformed shell model, $Q = 3.72 (\pm 20\%) \text{ b}$, $r(\text{r.m.s.}) = 2.42 \text{ fm}$ (1966IS1A, 1966ST12). See also (1963GO04, 1963GU1A, 1965GO1K, 1965GR18, 1965RA1C, 1965VA1G).

Ground-state transition widths for various excited states determined by inelastic scattering are listed in Table 11.13 (1962ED02, 1966KO08, 1966SP02, 1967SP02). For $^{11}\text{B}^*(8.57)$, the dependence of the transition probability on q^2 indicates an E2-M1 mixture: hence the parity is odd, $J \leq \frac{5}{2}$. The observed γ -width if $^{11}\text{B}^*(8.93)$ implies $\Gamma_\alpha = 0.03 \text{ eV}$ (compare $^9\text{Be}(^3\text{He}, \text{p})^{11}\text{B}$ (1966SP02)). See also (1962BA1D, 1964BR1N, 1966RI1G, 1967KA1A, 1967LE1E).

35. $^{11}\text{B}(\text{n}, \text{n}')^{11}\text{B}^*$

See (1955GR18, 1961LI04, 1962TE05, 1963OP1A, 1966FR18).

36. (a) $^{11}\text{B}(\text{p}, \text{p}')^{11}\text{B}^*$

(b) $^{11}\text{B}(\text{p}, 2\text{p})^{10}\text{Be}$ $Q_{\text{m}} = -11.228$

Study of the pair line spectrum of the 2.12 MeV γ -ray establishes that there is no parity change between the ground and first excited states: therefore the parity of the excited state is odd (1963WI01; see also (1962GO09, 1963SC33)). The 2.12 MeV excited state decays by emission of a $2.134 \pm 0.005 \text{ MeV}$ γ -ray (1957MC35): it exhibits $< 2.0 \times 10^{-3}$ part of circular polarization; this observation places an upper limit of $F^2 \lesssim 1 \times 10^{-7}$ for the intensity of the parity non-conserving part of the wave function (1958WI41).

At $E_{\text{p}} = 185 \text{ MeV}$, proton groups are observed to ^{11}B levels at 2.2 ± 0.1 , 4.5 ± 0.1 , 5.1 ± 0.15 , 6.8 ± 0.1 (probably corresponds to $^{11}\text{B}^*(6.74)$), 7.4 ± 0.3 , 8.95 ± 0.15 , ≈ 9.8 , 10.5 ± 0.2 , 11.9 ± 0.2 , 13.0 ± 0.2 ($\Gamma \approx 0.4 \text{ MeV}$) and $14.5 \pm 0.4 \text{ MeV}$ (1965HA17). The state at $E_{\text{x}} = 6.79 \text{ MeV}$ is not reported by (1965HA17) or by (1964NE06). Assuming that the (p, p') transitions involve mainly E2, $\Gamma(\text{E2}\downarrow)$ is $22 \pm 9 \text{ meV}$ for $^{11}\text{B}^*(4.44)$ and $90 \pm 40 \text{ meV}$, for $^{11}\text{B}^*(6.74)$ (1964JA03). See also (1965JA1A).

See also (1959EG1C, 1959JO43) for polarization studies; (1961KO08, 1962KI02, 1965HU10) for angular distributions at $E_{\text{p}} = 5.8$ to 7.5 MeV and (1959EG20, 1959TO1A, 1961CL09, 1962CL13).

For reaction (b) see (1964LI1D, 1964TI02, 1966JA1A, 1967JA1E, 1968JA1G) and ^{10}Be in (1966LA04).

37. $^{11}\text{B}(\text{d}, \text{d})^{11}\text{B}$

The angular distribution of elastically scattered deuterons has been determined at $E_{\text{d}} = 11.8 \text{ MeV}$ and analyzed by the optical model (1967FI07). See (1955KH35, 1962SL03, 1965GA02).

38. $^{11}\text{B}(t, t)^{11}\text{B}$

Angular distributions of elastically scattered tritons have been measured at $E_t = 1.75$ and 2.10 MeV. Optical model fits have been made (1968HE1N).

39. $^{11}\text{B}(^3\text{He}, ^6\text{Li})^8\text{Be}$ $Q_m = 4.566$

Angular distributions observed at $E(^3\text{He}) = 3.0$ and 5.2 MeV suggest a cluster pick-up mechanism, indicating a substantial ($^8\text{Be} + t$) configuration in ^{11}B (1967YO02).

40. $^{11}\text{B}(\alpha, \alpha)^{11}\text{B}$

At $E_\alpha = 28.5$ MeV, angular distributions are obtained for the elastic α -particles and the inelastic α 's corresponding to $^{11}\text{B}^*(2.12, 4.44, 5.02, 6.74 + 6.79, 7.30, 8.57)$ (1967NA06). See also (1964ST1K, 1966GE12).

41. (a) $^{11}\text{B}(^{11}\text{B}, ^{11}\text{B})^{11}\text{B}$
 (b) $^{11}\text{B}(^{16}\text{O}, ^{16}\text{O})^{11}\text{B}$

For reaction (a), see (1967GU1A). For reaction (b) see (1966OE1A).

42. $^{11}\text{C}(\beta^+)^{11}\text{B}$ $Q_m = 1.981$

See ^{11}C .

43. (a) $^{12}\text{C}(\gamma, p)^{11}\text{B}$ $Q_m = -15.957$
 (b) $^{12}\text{C}(e, e'p)^{11}\text{B}$ $Q_m = -15.957$

For reaction (a) see (1962PA08, 1963WA18, 1965AL02, 1966PA05). For reaction (b) see (1962DO1A, 1962PA08, 1965AM1B, 1965CO1E). See also ^{12}C .

44. $^{12}\text{C}(p, 2p)^{11}\text{B}$ $Q_m = -15.957$

In the summed proton spectrum, gross structure is observed corresponding to $Q = -15.6 \pm 0.6$ and -34.3 ± 0.8 MeV corresponding to $^{11}\text{B}_{\text{g.s.}}$ and an excited state with $J^\pi = \frac{1}{2}^+$ at $E_x = 18.3$ MeV (ejection of p- and s-protons, respectively) (1965RI1A, 1966TY01). See also (1958MA1B, 1958TY49, 1960GO1M, 1961PU1A, 1962GA09, 1962GA23, 1962GO1P, 1965BE1E, 1965CO1E, 1966BE1B, 1966WA12). A high resolution experiment shows groups corresponding to $^{11}\text{B}^*(0, 2.12, 4.44, 5.02, 6.79)$ (1965PU02, 1967PU01). The angular correlation and energy distribution of the protons emitted at $E_p = 45.5$ MeV have been measured by (1967RI08) and compared with distorted-wave t -matrix approximation theory.

See also (1960GO16, 1961CL09, 1961GA14, 1962CO17, 1962VA1G, 1963CL1B, 1963GR1G, 1963RI1B, 1964BO1L, 1966YU1A, 1967YU02; expt.) and (1960RI1B, 1962BA1J, 1962CL13, 1962DI1A, 1962IN1A, 1963BE42, 1963CL07, 1963YU1A, 1964BA1C, 1965YU1B, 1966LI1E, 1966NG1B, 1967JA1E; theor.).

$$45. \ ^{12}\text{C}(n, d)^{11}\text{B} \quad Q_m = -13.732$$

See (1962AU1A, 1966ME03, 1967ME11).

$$46. \ ^{12}\text{C}(d, \ ^3\text{He})^{11}\text{B} \quad Q_m = -10.463$$

Angular distributions with DWBA indicate $l = 1$ pickup for $^{11}\text{B}^*(0, 2.0, 4.8)$. No $l = 3$ fit is possible for the $\frac{5}{2}^-$ and $\frac{7}{2}^-$ states [$^{11}\text{B}^*(4.3, 6.5)$], suggesting a double process involving ^{12}C (2^+) as an intermediate stage (1966CH1K). The ground-state angular distribution has also been determined at $E_d = 82$ MeV (1967AR21). At $E_d = 28.5$ MeV, the cross sections for the (d, ^3He) and (d, t) reactions to the mirror ground states ^{11}B and ^{11}C are the same (1966DE1C). See also (1965PE17, 1967FI1G).

$$47. \ ^{12}\text{C}(t, \alpha)^{11}\text{B} \quad Q_m = 3.858$$

Angular distributions of the α -particles to the ground and first excited states of ^{11}B have been measured at $E_t = 1$ to 2 MeV (1962GU01), 10.1 MeV (1962PU01) and 13 MeV (1965AJ01). At the two higher energies, the angular distributions of the α -particles to the 4.44 and 5.02 MeV states have also been determined. Ground state angular distributions have been measured at 18 energies between $E_t = 1.5$ and 3.2 MeV (1966SE1D, 1968SE1F). See also (1960MU07, 1961HO1F, 1962NE1D, 1963NI04, 1964BE1K, 1965HO1C, 1965NE1B).

$$48. \ ^{12}\text{C}(^{12}\text{C}, \ ^{13}\text{N})^{11}\text{B} \quad Q_m = -14.013$$

See (1967VO1D, 1967WI04).

$$49. {}^{12}\text{C}({}^{14}\text{N}, {}^{15}\text{O}){}^{11}\text{B} \quad Q_{\text{m}} = -8.664$$

See (1966PO1B, 1967BI06).

$$50. {}^{13}\text{C}(\text{n}, \text{t}){}^{11}\text{B} \quad Q_{\text{m}} = -12.422$$

This reaction has not been reported.

$$51. {}^{13}\text{C}(\text{p}, {}^3\text{He}){}^{11}\text{B} \quad Q_{\text{m}} = -13.185$$

At $E_{\text{p}} = 43.7$ MeV, ${}^3\text{He}$ particles are observed corresponding to a $T = \frac{3}{2}$ state in ${}^{11}\text{B}$: $E_{\text{x}} = 13.02 \pm 0.08$ MeV ($\Gamma = 358 \pm 60$ keV); $L = 0$ ($J = \frac{1}{2}^-$) (1966MA2N). See also ${}^{11}\text{C}$.

$$52. {}^{13}\text{C}(\text{d}, \alpha){}^{11}\text{B} \quad Q_{\text{m}} = 5.168$$

Angular distributions of ground-state α -particles have been determined at $E_{\text{d}} = 3.3$ to 4.2 MeV (1963MA24), 10 to 12 MeV (1967CU1A). Alpha groups corresponding to the states at 2.12, 4.44 and 6.79 MeV have been observed by (1951LI29, 1953SP1A). The γ -decay of the 4.44 (1955BE62, 1958RA13) and 5.02 MeV states (1963AL21) is also reported. See also (1959EL43, 1965NE10, 1966KL06, 1966KL1E, 1966KL1F) and ${}^{15}\text{N}$ in (1970AJ04).

$$53. {}^{14}\text{C}(\text{p}, \alpha){}^{11}\text{B} \quad Q_{\text{m}} = -0.784$$

Angular distributions of the α -particles to the first four states of ${}^{11}\text{B}$ have been determined at $E_{\text{p}} = 18$ MeV (1962BR34). See also (1967EL1D).

$$54. {}^{14}\text{N}(\text{n}, \alpha){}^{11}\text{B} \quad Q_{\text{m}} = -0.157$$

See (1958DO63, 1959GA14, 1959HA13, 1963MO04, 1963SE08, 1964MO1D, 1966CS1B, 1966MA2M, 1967MO21) and ${}^{15}\text{N}$ in (1970AJ04).

¹¹C
(Figs. 3 and 4)

GENERAL:

Shell model: (1956KU1A, 1957KU58, 1960TA1C, 1961KU1C, 1965FA1C).

Collective model: (1966EL08).

Ground-state properties: (1963BE36, 1964ST1B, 1967SH05).

Other: (1962DY1A, 1964RE1B, 1967AU1B, 1967DE1V, 1967FA1A, 1967NE1D, 1967PO1J, 1968DI1B).

$$Q = (+)0.0308 \pm 0.0006 \text{ b (1963KO1G, 1964HA46);}$$

$$\mu = (-)1.027 \pm 0.010 \text{ nm (1963KO1G, 1964HA46);}$$

$$J = \frac{3}{2} \text{ (1961SN01).}$$

1. ¹¹C(β^+)¹¹B $Q_m = 1.981$

The mean value of reported half-lives is 20.34 ± 0.04 min (see Table 11.15); $ft = 3882$. The ratio of K-capture of positron emission is $(0.19 \pm 0.03)\%$ (1957SC29), $(0.230_{-0.011}^{+0.014})\%$ (1967CA09). See also (1965GA1D, 1966MI1F, 1967AM1H).

2. (a) ⁶Li(⁶Li, n)¹¹C $Q_m = 9.457$

(b) ⁷Li(⁶Li, 2n)¹¹C $Q_m = 2.204$

At $E(^6\text{Li}) = 4.1$ MeV, neutron groups have been observed to the first seven excited states and to the 8.42 MeV state. Associated γ -rays were used for the stop triggering of the time-of-flight system. The group to the 8.4 MeV state had an intensity $\approx 10\%$ of that to the highest bound state at $E_x = 7.5$ MeV suggesting a small Γ_α for the 8.42 MeV state (compare ¹²C(³He, α)¹¹C). No evidence for the γ -decay of the 8.11 MeV state was found (1966BA2U). See also (1957NO17, 1960NO1A, 1962BE16, 1962BE24).

3. ⁹Be(³He, n)¹¹C $Q_m = 7.562$

Reported neutron groups are listed in Table 11.16. Angular distributions and excitation func-

Table 11.14: Energy levels of ^{11}C

E_x in ^{11}C (MeV \pm keV)	$J^\pi; T$	τ (psec) or Γ (keV) (keV)	Decay	Reactions
0	$\frac{3}{2}^-$	$\tau_{1/2} = 20.39 \pm 0.06$ min	β^-	1, 2, 3, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 26
1.995 ± 3	$\frac{1}{2}^-$	$\tau_m < 0.5$ psec < 0.5 < 0.11 < 0.25 < 0.16 < 0.5	γ	2, 3, 11, 13, 16, 20, 22, 25
4.305 ± 6	$\frac{5}{2}^-$		γ	2, 3, 11, 13, 16, 22, 25
4.794 ± 6	$\frac{3}{2}^-$		γ	2, 3, 11, 13, 16, 20, 22, 25
6.339 ± 5	$\frac{1}{2}^+$		γ	2, 3, 11, 13, 16, 22
6.480 ± 6	$\frac{7}{2}^-$		γ	2, 3, 13, 16, 22
6.906 ± 4	$\frac{5}{2}^+$		γ	2, 3, 11, 13, 20, 22
7.509 ± 6	$\frac{3}{2}^+$		γ	2, 3, 11, 13, 22
8.107 ± 3	$\leq \frac{5}{2}^-$	Γ_α small	(γ)	3, 11, 13, 22
8.420 ± 4	$\frac{5}{2}^-$		γ	2, 11, 13, 22
8.652 ± 4	+	$\Gamma = 450 \pm 50$ keV ≈ 200 200 ± 30		11
8.694 ± 4	(+)			11
9.731 ± 5	$(\frac{5}{2}^+)$		γ, p, α_0	4, 10, 11
10.086 ± 5	$\frac{7}{2}^+$		p, α_0, α_1	6, 10, 11
10.682 ± 5	$\frac{9}{2}^+$		p, α_0, α_1	6, 10, 11
(10.801 \pm 5)			p, α_1	10
(11.032 \pm 5)			p, α_1	10
11.45 ± 10			370 p, α_1	10, 11
(11.957 \pm 7)			p, α_1	10, 11
12.45 ± 80	$\frac{1}{2}^-; T = \frac{3}{2}$		560 ± 60	
12.65 ± 20	$(\frac{7}{2}^+)$	370	$\gamma, p, \alpha_0, \alpha_1$	4, 7, 10
13.33			p, α_0, α_1	10
13.90 ± 20		≈ 400	p	7
14.07 ± 20		broad	$n, p, (\alpha_0, \alpha_1)$	5, 10
14.76 ± 40		broad	n, p, α_1	5, 10
15.35 ± 50		broad	n, p, α_1	5, 10
15.60 ± 50		broad	n, p	5
≈ 16		≈ 3200	(n), p	7

Table 11.15: Half-life of ^{11}C

$\tau_{1/2}$ (min)	Ref.
20.35 ± 0.08	(1941SM11)
20.50 ± 0.6	(1941SO01)
20.0 ± 0.1	(1951DI12)
20.74 ± 0.10	(1953KU08)
20.26 ± 0.1	(1955BA63)
20.8 ± 0.2	(1957PR53)
20.11 ± 0.13	(1958AR15)
20.34 ± 0.04	Weighted mean

tions have been studied in the range $E(^3\text{He}) = 1.3$ to 10 MeV by (1961DU1B, 1963DU12, 1963VA16, 1965DI06, 1965TO06, 1966AD1D: see (1966WE1B, 1968OK1D)). In the middle range of these energies, the reaction is dominated by direct interaction, but it is not clear whether two-particle stripping or knock-on processes are involved. Large back angle cross sections are unexplained by DWBA (1965TO06). At 10 MeV, the distributions are consistent with a predominant role for the two-nucleon stripping process. The contribution of the knock-on process appears to be small (1968OK1D).

Gamma branching ratios and multiplicities for ^{11}C levels up to $E_x = 7.5$ MeV have been studied by (1965OL03, 1965RO07): see Table 11.17, $^{10}\text{B}(d, n)^{11}\text{C}$ and $^{12}\text{C}(^3\text{He}, \alpha\gamma)^{11}\text{C}$. The following remarks on individual levels derive largely from (1965OL03, 1965RO07):

Transitions to $^{11}\text{C}_{g.s.}$ and (2.0) are predominantly E1, fixing the parity of $^{11}\text{C}^*(7.51)$ as even and that of $^{11}\text{C}^*(2.0)$ as odd. The transition to $^{11}\text{C}^*(2.0)$ is not isotropic, so $J \neq \frac{1}{2}$; $J^\pi = \frac{3}{2}^+$ or $\frac{5}{2}^+$ are possible, but if $^{11}\text{C}^*(2.0)$ is $\frac{1}{2}^-$, $J^\pi = \frac{5}{2}^+$ is eliminated (see also $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$). The relatively large strength of the cascade transition is difficult to understand, and comparison with $^{11}\text{B}^*(8.00)$ suggests some deviation from charge symmetry (1965OL03, 1965RO07).

$E_x = 6.91$ MeV. The g.s. transition is predominantly E1 so $J^\pi \leq \frac{5}{2}^+$ (1965OL03). Branching ratios favor $J = \frac{3}{2}$ or $\frac{5}{2}$ (1965RO07): $J = \frac{5}{2}$ from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$. The analogue is presumably $^{11}\text{B}^*(7.30)$ (1965OL03). $\tau_m < 0.16$ psec (1966WA10).

$E_x = 6.48$ MeV. The E2 character of the g.s. transition indicates $J^\pi \leq \frac{7}{2}^-$; $J = \frac{7}{2}$ from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$. This level is weakly populated in $^9\text{Be}(^3\text{He}, n)^{11}\text{C}$. The analogue is presumably $^{11}\text{B}^*(6.74)$. $\tau_m < 0.25$ psec (1966WA10). See also (1966GO12).

$E_x = 6.34$ MeV. The g.s. transition is predominantly E1 so $J^\pi \leq \frac{5}{2}^+$ (1965OL03). Relative transition strengths are consistent with $J = \frac{1}{2}$ or $\frac{3}{2}$ with the assumption that $^{11}\text{B}^*(6.79)$ is the mirror level (1965RO07). $\tau_m < 0.11$ psec (1966WA10). $J^\pi = \frac{1}{2}^+$ from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$.

Table 11.16: ^{11}C levels from $^9\text{Be}(^3\text{He}, n)^{11}\text{C}$

(1963VA16)	(1965TO06)			(1965RO07)	(1968MC1C)
E_x^a	E_x^a	L	J^π	E_x^a	E_x^a
0	0	0 + 2	$\leq \frac{5}{2}^-$		-0.010 ± 18
2.01 ± 30	1.97 ± 30	(0) + 2			1.968 ± 13
4.29 ± 20	4.25 ± 30	(0) + 2			4.304 ± 10
4.79 ± 20	4.76 ± 30	0 + 2	$\leq \frac{5}{2}^-$		4.797 ± 9
6.39 ± 20	6.37 ± 30^b	1 + 3		6.37 ± 100^b	6.338 ± 6
					6.480 ± 6
6.88 ± 20	6.89 ± 30	weak		6.93 ± 80	6.909 ± 5
	7.48 ± 30	weak		7.50 ± 50	7.516 ± 6
	8.10 ± 30	0	$\leq \frac{5}{2}^-$		8.108 ± 4
					8.418 ± 5
					8.651 ± 4
					8.694 ± 4

^a E_x given in MeV \pm keV.

^b Probably unresolved doublet: the higher component is only weakly populated if at all.

$E_x = 4.79$ MeV. Stripping results indicate odd parity, $J \leq \frac{5}{2}^-$ (1965TO06), consistent with the observed M1, E2 nature of the g.s. transition. The strength of the transition to $^{11}\text{C}^*(2.0)$ argues against $J > \frac{5}{2}$. $J = \frac{3}{2}$ from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$.

$E_x = 4.31$ MeV. The M1 g.s. transition indicates $J^\pi \leq \frac{5}{2}^-$. $J = \frac{5}{2}$ from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$.

$E_x = 2.00$ MeV. The E1 transition from $^{11}\text{C}^*(7.50)$ fixes the parity as odd, $J^\pi \leq \frac{7}{2}^-$. $J^\pi = \frac{1}{2}^-$ is fixed from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$.

See also (1959LI1D, 1962SE1A, 1963MA1K, 1964BR13, 1966WA1C, 1967BA1E, 1967BO1D).

4. $^{10}\text{B}(p, \gamma)^{11}\text{C}$

$$Q_m = 8.693$$

In the range $E_p = 0.7$ to 4.7 MeV, two broad resonances are reported at 1.14 and ≈ 4.4 MeV: see Table 11.18 (1954DA20, 1955BA22, 1956CH20, 1957HU79, 1962OP03). At the $E_p = 1.14$ MeV resonance ($E_x = 9.73$ MeV), capture γ -rays are observed corresponding to the direct ground-state transition (relative intensity 1.0 ± 0.2) and to cascades via the states at 6.48 (0.3), 4.31 (0.18 ± 0.02) and 2.00 MeV (0.05), with $(2J + 1)\Gamma_\gamma = 10, 1.8$ and 3 eV for the ground state transition and the cascades through the 6.48 and 4.31 MeV states, respectively (1961JA11). The

6.48 MeV state decays $\approx \frac{1}{3}$ of the time via a cascade through the 4.31 MeV state (1961JA11). The 4.31 MeV state goes by direct ground-state decay ($> 98\%$) while the 4.79 MeV state decays in $(15 \pm 5\%)$ of the cases via the 2.0 MeV state ($E_\gamma = (2.72 \pm 0.08) + (2.05 \pm 0.08)$ MeV) (1961DO03). See Table 11.17.

Table 11.17: Gamma decay of ^{11}C levels

E_i (MeV)	J_i^π	τ_m^b (psec)	E_f (MeV)	Branch ^a (%)	Mult. ^a	Branch ^c (%)	X^c
2.00	$\frac{1}{2}^-$		0	100			
4.31	$\frac{5}{2}^-$	< 0.5	0	100	M1	100	$+0.17 \pm 0.03^g$
			2.0	< 2		< 2	
4.79	$\frac{3}{2}^-$	< 0.5	0	83 ± 4	M1	86 ± 2^f	e
			2.0	17 ± 4		14 ± 2^f	e
6.34	$\frac{1}{2}^+$	< 0.11	0	65 ± 3	E1	68 ± 3	
			2.0	35 ± 3		32 ± 3^d	
			4.3			< 7	
			4.8	< 4		< 3	
6.48	$\frac{7}{2}^-$	< 0.25	0	89 ± 2	E2	88 ± 2	-0.01 ± 0.06
			2.0	< 2		< 4	
			4.3	11 ± 2		12 ± 2	
			4.8			< 2	
6.91	$\frac{5}{2}^+$	< 0.16	0	89 ± 3	E1	91 ± 2	0.02 ± 0.03
			2.0	< 2		< 1	
			4.3	11 ± 3		4.5 ± 1	
			4.8	< 3		4.5 ± 1	
			6.3	< 5			
			6.5	< 5			
7.51	$\frac{3}{2}^+$	< 0.5	0	36 ± 2	E1	37 ± 3	-0.04 ± 0.04
			2.0	64 ± 2	E1	63 ± 8	0 ± 0.03
			4.3	< 3		< 1	
			4.8	< 3		< 1	
			6.3	< 3			
			6.5	< 3			
			6.9	< 4			

^a From ${}^9\text{Be}({}^3\text{He}, n){}^{11}\text{C}$ and ${}^{10}\text{B}(d, n){}^{11}\text{C}$ (1965OL03): includes earlier measurements.

^b (1966WA10).

^c (1968EA03): ${}^{12}\text{C}({}^3\text{He}, \alpha){}^{11}\text{C}$; $X \equiv$ amplitude ratio of $(L + 1)/L$.

^d The cascade is through ${}^{11}\text{C}^*(2.0)$ and not ${}^{11}\text{C}^*(4.3)$ (1968EA03).

^e See ${}^{12}\text{C}({}^3\text{He}, \alpha){}^{11}\text{C}$.

^f 86 ± 3 , $14 \pm 3\%$ (1966GA19); 84 ± 3 , $16 \pm 3\%$ (1967BL22).

^g $+0.16$ (-0.02 , $+0.06$) (1966GA19); 0.13 ± 0.04 (1967BL22).

5. ${}^{10}\text{B}(p, n){}^{10}\text{C}$

$$Q_m = -4.432$$

$$E_b = 8.693$$

$E_{\text{thresh.}} = 4880.1 \pm 2.0$ keV; $Q_0 = -4432.8 \pm 1.8$ keV (1966FR09). This result is 44 keV lower than the value used by (1965MA54), and considerably more accurate. We adopt $(M - A)$ ${}^{10}\text{C} = 15702.6 \pm 2.0$ keV for the present review.

The total (p, n) cross section has been measured to $E_p = 10.6$ MeV by (1963EA01): broad maxima are observed at $E_p = 5.92 \pm 0.02$, 6.68 ± 0.04 , 7.33 ± 0.05 and 7.60 ± 0.05 MeV (see Table 11.18). The cross section for formation of ${}^{10}\text{C}_{\text{g.s.}}$ measured up to $E_p = 12$ MeV, shows similar behavior to 8 MeV. At $E_p \approx 8$ MeV, a sharp maximum is observed. The cross section for production of 3.35 MeV γ -rays (from ${}^{10}\text{C}^*$) does not appear to show structure for $E_p = 8.5$ to 12 MeV (1966SE03). See also (1959AJ76, 1963VA1C, 1965VA23).

6. ${}^{10}\text{B}(p, p){}^{10}\text{B}$

$$E_b = 8.693$$

The elastic scattering shows two conspicuous anomalies, at $E_p = 1.50 \pm 0.02$ MeV and at 2.18 MeV, corresponding to states at $E_x = 10.06$ and 10.68 MeV with $J^\pi = \frac{7}{2}^+$ and $\frac{9}{2}^+$ (see Tables 11.18 and 11.19). Below $E_p = 0.7$ MeV, the scattering can be explained in terms of pure s-wave potential scattering but the possibility of a state near $E_p = 0.270$ MeV ($E_x = 8.95$ MeV) cannot be excluded (1960OV1A, 1962OV02: $E_p = 0.15$ to 3.0 MeV). (1962OV02) also report evidence for a state formed by s-wave protons near $E_p = 2.8$ MeV. On the other hand (1962AN11) who have studied scattering to $E_p = 3.5$ MeV interpret the data as indicating a resonance above $E_p = 3.5$ MeV. See also (1951BR10, 1961RO05, 1966JA1F).

7. ${}^{10}\text{B}(p, p'){}^{10}\text{B}^*$

$$E_b = 8.693$$

The yield of 0.71 MeV radiation, from ${}^{10}\text{B}^*$, rises monotonically from $E_p = 1.5$ to 4.1 MeV (1952DA05, 1954DA20, 1957HU79, 1964BE31), and then shows resonance behavior at $E_p = 4.35 \pm 0.02$ MeV and 5.73 ± 0.02 MeV (1962OP03: see Table 11.18). For $E_p = 6$ to 12 MeV, the

Table 11.18: Resonances in $^{10}\text{B} + \text{p}$ ^a

E_{res} (MeV \pm keV)	E_x (MeV)	J^π	Γ_{lab} (keV)	Decay	Refs.
1.145 \pm 5	9.731	$(\frac{5}{2}^+)$	500 \pm 50	γ , p, α_0	(1951BR10, 1955HA01, 1956CH20, 1956CR07, 1957HU79)
1.533 \pm 5	10.086	$\frac{7}{2}^+$	\approx 250	p, α_0 , α_1	(1951BR10, 1956AL23, 1956CH20, 1956CR07, 1957HU79, 1962OV02)
2.189 \pm 5	10.682	$\frac{9}{2}^+$	220 \pm 30	p, α_0 , α_1	(1962OV02, 1964BE31, 1964JE01)
2.320 \pm 5	10.801			p, α_1	(1964BE31)
2.574 \pm 5	11.032			p, α_1	(1964BE31)
3.03 \pm 10	11.45		400	p, α_0 , α_1	(1962OP03, 1964JE01)
3.592 \pm 7	11.957			p, α_1	(1964BE31)
4.36 \pm 20	12.65	$(\frac{7}{2}^+)$	400	γ , p, α_0 , α_1 , ^3He	(1962OP03, 1964JE01, 1966SE03)
5.10	13.33			p, α_0 , α_1	(1962OP03, 1964JE01, 1966SE03)
5.73 \pm 20	13.90		\approx 500	p	(1962OP03)
5.92 \pm 20	14.07		broad	n, p ₀ , (α_0 , α_1)	(1963EA01, 1964JE01, 1966SE03)
6.68 \pm 40	14.76		broad	n, p, (^3He), α_1	(1963EA01, 1964JE01, 1966SE03)
7.33 \pm 50	15.35		\approx 2500	n, p, (α_1)	(1963EA01, 1966SE03)
7.60 \pm 50	15.60		broad	n, p	(1963EA01)
8 ^b	16		\approx 3500	(n), p	(1966SE03)

^a See also Table 11.19.^b See reaction 7. This broad state may be a giant resonance in the ($\alpha + \alpha + \text{d} + \text{p}$) system.

Table 11.19: Level parameters for $^{10}\text{B}(p, p)^{10}\text{B}$ and $^{10}\text{B}(p, \alpha)^7\text{Be}$ ^a

E_r (MeV)	E_x (MeV)	J^π	Γ_p (MeV)	Γ_{α_0} (MeV)	Γ_{α_1} (MeV)	θ_p^2	$\theta_{\alpha_0}^2$	$\theta_{\alpha_1}^2$
1.17 ^b	9.76	$(\frac{5}{2}^+)$	0.045	0.255				
1.50 ^c	10.06	$\frac{7}{2}^+$	0.090	0.100	0.060	0.02	0.26	0.35
2.18 ^c	10.67	$\frac{9}{2}^+$	0.100	0.100		0.17	0.11	
4.36 ^d	12.65	$(\frac{7}{2}^+)$	0.20	0.15	0.05	0.02	0.29	0.08

^a See also Table 11.18.

^b (1956CR07, 1962OV02).

^c (1962OV02).

^d (1964JE01).

cross section shows several sharp maxima superposed on a broad maximum ($\Gamma \approx 2.5$ MeV) at $E_p \approx 7.2$ MeV (1966SE03). (1966SE03) have also measured the yields of the 1.02 ($^{10}\text{B}^*(1.7) \rightarrow ^{10}\text{B}^*(0.7)$), 1.43 ($^{10}\text{B}^*(2.1) \rightarrow ^{10}\text{B}^*(0.7)$) and $^{10}\text{B}^*(3.6) \rightarrow ^{10}\text{B}^*(2.2)$), 2.86 ($^{10}\text{B}^*(3.6) \rightarrow ^{10}\text{B}^*(0.7)$) and 4.44 ($^{10}\text{B}^*(5.16) \rightarrow ^{10}\text{B}^*(0.7)$) MeV gamma rays for $E_p = 4$ to 12 MeV. The yields of both the 2.15 and 3.58 MeV states of ^{10}B show a broad resonance, about 4 MeV wide, centered at $E_p \approx 8$ MeV, with some fine structure superimposed. The yields of the 1.74 and 5.16 MeV states show this resonance weakly or not at all (this is also true of the neutrons in reaction 6). For the groups that show the broad resonance, the maximum cross sections vary from 20 to 160 mb and tend to increase with increasing Q . For the groups that do not show the broad resonance, the cross sections are all below 5 mb and reach their maxima in the 7–12 MeV region. The weak reactions are those in which the residual nucleus is left in a $T = 1$ state. This is explained in terms of an $\alpha + \alpha + d + p$ character for the broad resonance. This suggests a significant $(\alpha + \alpha + d)$ cluster structure for the ^{10}B ground state as well as for the first few low-lying states of ^{10}B (1966SE03, 1967WA1L).

$$8. \ ^{10}\text{B}(p, d)^9\text{B} \qquad Q_m = -6.213 \qquad E_b = 8.693$$

See ^9B IN (1966LA04).

$$9. \ ^{10}\text{B}(p, ^3\text{He})^8\text{Be} \qquad Q_m = -0.534 \qquad E_b = 8.693$$

The ground-state yield has been obtained for $E_p = 4$ to 10 MeV ($\theta = 50^\circ, 90^\circ$). There are slight maxima at similar energies to those in the (p, α) yield. However, the angular distributions do

not vary strongly over the region covered and it is suggested that a direct interaction mechanism dominates (1963JE01). (1966SE03) report two strong maxima at $E_p \approx 4.5$ and 6.5 MeV.

10. $^{10}\text{B}(p, \alpha)^7\text{Be}$

$$Q_m = 1.148$$

$$E_b = 8.693$$

The parameters of observed resonances are displayed in Tables 11.18 and 11.19. The ground state (α_0) α -particles exhibit broad resonances at $E_p = 1.17, 1.53, 2.18, 3.0, 4.4, 5.1$ and 6.3 MeV (1959AJ76, 1962OV02, 1964JE01). Alpha particles to the 0.43 MeV ^7Be state (α_1) and 0.43 MeV γ -rays exhibit all but the 1.2 MeV resonance (1959AJ76, 1962OP03, 1964BE31, 1964JE01, 1966SE03). Weak resonances are also reported at 2.32, 2.57 and 3.59 MeV (1964BE31). A broad maximum dominates the region from $E_p = 4$ MeV to about 7.5 MeV (1966SE03). See also discussion under reaction 6, and (1963VA1C).

11. $^{10}\text{B}(d, n)^{11}\text{C}$

$$Q_m = 6.468$$

Neutron spectra have been studied by (1959AJ76), (1952JO10, 1956CE1B, 1956CE73, 1956GR54, 1956MA83, 1957GR50, 1959NE1A, 1960MC1C, 1963OV02, 1965SI13) and others. Angular distributions, γ spectra and (n- γ) correlations are reported by (1960FE13, 1960MC1C, 1960NE17, 1961JA12, 1962FR06, 1963BR1H, 1965OL03, 1966AD1D), (1966GO12, 1966RU01, 1966WE1B, 1967DI01) and (1959AJ76). The principal results are exhibited in Table 11.20.

DWBA fits to the ground-state distribution are reported by (1967DI01: $E_d = 5$ to 7.6 MeV). At $E_d = 7.6$ and 9.0 MeV, the angular distributions corresponding to the first four levels match the PWBA $l = 1$ patterns moderately well, although the $^{11}\text{C}^*(2.0)$ level is relatively weak (1956CE1B, 1956CE73: 7.6 MeV) and (1956MA83: 9 MeV). For $E_d = 1$ to 4 MeV, levels at $E_x = 6.48, 8.43$ and 8.66 MeV show good patterns; $^{11}\text{C}^*(2.0, 4.3, 4.8, 6.34, 6.91)$ are quite weak (1960FE13, 1960NE17, 1965SI13: $E_d = 5$ to 8 MeV) and (1961JA12). The $^{11}\text{C}^*(2.0)$ is believed to have $J^\pi = \frac{1}{2}^-$ and is presumably formed by an exchange or spin-flip mechanism (1963AU1A). In $j - j$ coupling, it is expected that only the levels $(\frac{3}{2})_I, (\frac{7}{2})_I$ and $(\frac{5}{2})_{II}$ will show large stripping widths: these are presumably $^{11}\text{C}^*(0, 6.48, 8.42)$ (1960BI08, 1960MA32, 1961JA12).

Information on the gamma decay of levels up to $^{11}\text{C}^*(7.50)$ has been summarized by (1965OL03) and is incorporated in Table 11.17. See also (1966GO12).

Neutron threshold measurements indicate levels at 8.103, 8.426 and 8.656 MeV (1955MA76: ± 8 keV; based on Q_m). (1963OV02) has investigated this region with time-of-flight techniques and finds that two levels exist near $E_x = 8.7$ MeV, at $E_x = 8.657$ and 8.702 MeV. A broad level is located at $E_x = 10.69$ MeV ($\Gamma \approx 200$ keV) but no evidence is found for other levels reported above $E_x = 8.7$ MeV. The doublet appears to correspond to the s-particle doublet in $^{11}\text{B}^*(9.19, 9.28)$ (1963OV02).

See also (1959BR75, 1959BU1F, 1960FE01, 1963MO07, 1965MA1K, 1966GO1N, 1966RO1X, 1967SC1K) and ^{12}C .

Table 11.20: Levels of ^{11}C from $^{10}\text{B}(\text{d}, \text{n})^{11}\text{C}$

E_x (MeV \pm keV)	J^π ^f	l	$(2J + 1)\theta^2$ ^a	$(2J + 1)\theta^2$ ^b	Refs. ^c
0	$\frac{3}{2}^-$	1	(4)	4	(1956MA83, 1967DI01)
1.94 \pm 50	$\frac{1}{2}^-$	(1)	weak		(1956MA83, 1960NE17)
4.260 \pm 21 ^d	$\frac{5}{2}^-$	1	(2.8)	1.2	(1960MC1C, 1960NE17)
4.749 \pm 17 ^d	$\frac{3}{2}^-$	1	(0.9)	0.54	(1955SA1B, 1960NE17, 1962FR06, 1966RU01)
6.34	$(\frac{1}{2})^+$				(1961JA12)
6.487 \pm 14 ^d	$\frac{7}{2}^-$	1	7.5		(1955BE81, 1955SA1B, 1960FE13, 1960MC1C, 1960NE17, 1962FR06, 1963BR1H, 1966RU01)
6.91 \pm 20 ^e	$\frac{5}{2}^+$	(1)	(0.9)		(1962FR06)
7.40 \pm 40	$\frac{3}{2}^+$				(1955SA1B)
8.103 \pm 8	$(\leq \frac{5}{2}^-)$				(1955MA76)
8.426 \pm 8	-	1	2.9		(1955MA76)
8.656 \pm 8	+	0	10.3		(1955MA76, 1963OV02)
8.702 \pm 20	(+)	(0)			(1963OV02)
(8.97 \pm 20)					(1957GR50)
(9.13 \pm 20)					(1952JO10)
(9.28 \pm 30)					(1956CE1B, 1956CE73)
(9.69 \pm 30)					(1956CE1B, 1956CE73)
(10.09 \pm 20)					(1957GR50)
10.69	(+)	(0, 2)			(1963OV02)
(10.89 \pm 20)					(1957GR50)
(11.26 \pm 20)					(1956CE1B, 1956CE73)
(11.52 \pm 20)					(1956CE1B, 1956CE73)

^a (1961JA12); $E_d = 1$ to 4 MeV: relative values.

^b (1956CE1B, 1956CE73, 1960MA32); $E_d = 7.55$ MeV: relative values.

^c (1952JO10) lists 12 levels, (1956CE1B, 1956CE73) lists 17, (1961JA12) lists 11: see also (1959AJ76).

^d (1959NE1A) and references therein.

^e (1961JA12).

^f J^π from Table 11.17.

12. $^{10}\text{B}(t, 2n)^{11}\text{C}$ $Q_m = 0.211$

See (1961RO21).

13. $^{10}\text{B}(^3\text{He}, d)^{11}\text{C}$ $Q_m = 3.199$
 $Q_0 = 3.226 \pm 0.010$ (1964MA57).

Angular distributions of the neutrons to the ground state of ^{11}C and to the first five excited states have been analyzed by stripping theory [(1961HI08: $E(^3\text{He}) = 9.84$ MeV) and (1965PA10: $E(^3\text{He}) = 3.5$ to 10 MeV)] and the excitation energies of nine states have been determined (1961HI08): see Table 11.21.

Table 11.21: Energy levels of ^{11}C from $^{10}\text{B}(^3\text{He}, d)^{11}\text{C}$ and $^{11}\text{B}(p, n)^{11}\text{C}$

(1961HI08) ^a			(1965OV01) ^d
E_x (MeV \pm keV)	l	θ_p^2 ^b	E_x (MeV \pm keV)
0	1	0.019	0
2.002 ± 10 ^c	^c		2.008 ± 20
4.322 ± 10	1	0.009	4.320 ± 20
4.808 ± 10	(1)	(0.006)	4.806 ± 20
6.345 ± 10			6.330 ± 20
6.476 ± 10	1	0.033	6.481 ± 20
6.903 ± 10			
7.498 ± 10			
8.107 ± 10			
8.42			

^a $^{10}\text{B}(^3\text{He}, d)$: $E(^3\text{He}) = 9.84$ MeV.

^b Values based on known J^π of states.

^c (1960FO01) find $E_x = 1.990 \pm 0.005$ MeV, $l = 1$.

^d $^{11}\text{B}(p, n)^{11}\text{C}$; time-of-flight.

See also (1959AL96, 1960FO01, 1960SP08, 1962BR10).

14. $^{10}\text{B}(\alpha, t)^{11}\text{C}$ $Q_m = -11.121$

The tritons corresponding to the ground state have been observed at $E_\alpha = 43$ MeV (1967DE1K).

15. (a) $^{10}\text{B}(^6\text{Li}, ^5\text{He})^{11}\text{C}$ $Q_m = 4.038$
 (b) $^{10}\text{B}(^{14}\text{N}, ^{13}\text{C})^{11}\text{C}$ $Q_m = 1.143$

For reaction (a) see (1957NO17). For reaction (b) see (1962NE01, 1963TO1E, 1966GA04).

16. $^{11}\text{B}(\text{p}, \text{n})^{11}\text{C}$ $Q_m = -2.763$

$$E_{\text{thresh.}} = 3.0164 \pm 0.0015 \text{ (1961BE13);}$$

$$E_{\text{thresh.}} = 3.019 \pm 0.004 \text{ (1959BR06).}$$

Neutron groups have been observed to $^{11}\text{C}_{\text{g.s.}}$ and to the first excited states: see Table 11.21 (1965OV01). Angular distributions of the ground-state neutrons have been studied at many energies up to $E_p = 18.5$ MeV (1960HI04, 1961AL07, 1964AN1B, 1964SA1D, 1964ST1C, 1965WA04). See also (1956AJ22, 1961GO13, 1961TA12, 1963PA1E, 1964BA16, 1965VA23, 1966UN1A) and ^{12}C .

17. $^{11}\text{B}(^3\text{He}, \text{t})^{11}\text{C}$ $Q_m = -1.999$
 $Q_0 = -2.0023 \pm 0.0012 \text{ (1965GO05).}$

The ground-state tritons have been observed at $E(^3\text{He}) = 4.8$ to 5.0 MeV, $d\sigma/d\Omega \approx 0.26$ mb/sr (1965GO05). Angular distributions of t_0 and t_1 have been obtained at $E(^3\text{He}) = 10$ MeV (1967CR04).

18. $^{12}\text{C}(\gamma, \text{n})^{11}\text{C}$ $Q_m = -18.720$

See (1963FU05, 1967FI1E) and ^{12}C .

19. (a) $^{12}\text{C}(\text{n}, 2\text{n})^{11}\text{C}$ $Q_m = -18.720$
 (b) $^{12}\text{C}(\text{p}, \text{pn})^{11}\text{C}$ $Q_m = -18.720$

For reaction (a) see ^{13}C in (1970AJ04). For reaction (b) see ^{13}N in (1970AJ04) and (1962AU1A, 1962BA1A, 1962GU10, 1966PA08).

$$20. \ ^{12}\text{C}(\text{p}, \text{d})^{11}\text{C} \quad Q_{\text{m}} = -16.495$$

At $E_{\text{p}} = 155$ MeV, deuteron groups have been observed to the ground state and to excited states of ^{11}C at 2.0 ± 0.1 , 4.9 ± 0.1 and (7.0 ± 0.2) MeV (1962RA01, 1963BA1R, 1963RA01, 1967BA2L). See also (1965DE1A, 1965KE02, 1965PU02, 1965VE1B). Angular distributions have been measured at $E_{\text{p}} = 19 - 20$ MeV (1962WA31), 27.5 MeV (1967GL01), 30 MeV (1967CH15), 31.1 and 33.5 MeV (1967GR1M), 36 MeV (1965KE02), 40 MeV (1963KA26, 1966SH1A), 45 MeV (1966MA2B), 54.9 MeV (1968TA1P), and 60 MeV (1965IS06). For spectroscopic factors, see (1967BA2L). See also (1956SE1A, 1959GR1B, 1960RA12, 1961CL09, 1962BE1H, 1963CL07, 1963IS1B, 1963MA1J, 1964JA1C, 1964SH07, 1965GL1E, 1967HO1M, 1967TO1D).

$$21. \ ^{12}\text{C}(\text{d}, \text{t})^{11}\text{C} \quad Q_{\text{m}} = -12.462$$

Ten states of ^{11}C have been observed at $E_{\text{d}} = 50$ MeV. The angular distributions are identical with those seen in the mirror reaction $^{12}\text{C}(\text{d}, \ ^3\text{He})^{11}\text{B}$ to analogue states (1966CH1K): see ^{11}B . At $E_{\text{d}} = 28.5$ MeV the (d, t) and (d, ^3He) cross sections to the ground states of the mirror nuclei ^{11}C and ^{11}B are equal (1966DE1C). See also (1959VL23, 1967FI1G), and ^{14}N in (1970AJ04).

$$22. \ ^{12}\text{C}(\ ^3\text{He}, \alpha)^{11}\text{C} \quad Q_{\text{m}} = 1.858$$

$$Q_0 = 1.856 \pm 0.012 \text{ (1960HI07).}$$

Angular distributions of alpha particles to the ground state have been observed at $E(^3\text{He}) = 1.8$ to 5.4 MeV (1964KU05), 2.4 to 4.5 MeV (1963LU05), 4.8 to 5.9 MeV (1966BL01), 8.5 to 10.0 MeV (1966SC22), 6.0, 8.8, 9.4 and 10.1 MeV (1960HI07), 16 to 18 MeV (1967GR1L, 1967GR1N), 24.8 MeV (1967HA21), 26 to 33 MeV (1963PA1G), and 26 MeV (1966DA1H). The α_1 distributions (to the 2 MeV state) have been investigated at $E(^3\text{He}) = 4.2, 4.9, 5.6, 6.0, 8.8, 9.4, 10.1, 16$ to 18 MeV (1960HI07, 1964KU05, 1966BL01, 1967GR1L, 1968GR1G). Angular distributions at the higher energies are consistent with $l_{\text{n}} = 1$ for both α_0 and α_1 (1960HI07). See also (1966SC22). The energy of the first excited state is $E_{\text{x}} = 1.990 \pm 0.010$ (1959PO61), 2.000 ± 0.010 (1960HI07), 1.999 ± 0.004 MeV (1968EA03). See also (1961CA1D, 1962AG01, 1962CA29, 1962GA17, 1962WE1C, 1963PA15, 1964BE1K, 1965GR1R). See also ^{15}O in (1970AJ04) and (1966HA1Q, 1967BR1N).

Studies of $\alpha - \gamma$ correlations have been carried out by (1965WA06: $E(^3\text{He}) = 4.7$ MeV), (1965NE06, 1965SC1D: $E(^3\text{He}) = 5.1$ MeV), (1967BL22: $E(^3\text{He}) = 9$ MeV), (1966GA19:

$E(^3\text{He}) = 10$ MeV), and (1968EA03: $E(^3\text{He}) = 6$ to 12 MeV) with the following results (1968EA03) (see also $^9\text{Be}(^3\text{He}, \text{n})^{11}\text{C}$ and Table 11.17):

$E_x = 8.42$ MeV. This level is unbound $\Gamma_\gamma/\Gamma = 0.2 \pm 0.1$.

$E_x = 8.11$ MeV. $\Gamma_\gamma/\Gamma < 0.04$; $J^\pi \leq (\frac{5}{2}^-)$ from $^9\text{Be}(^3\text{He}, \text{n})^{11}\text{C}$ (Table 11.16).

$E_x = 7.51$ MeV. For the ground-state transition, the $\alpha - \gamma$ correlation permits $J = \frac{3}{2}$ or $\frac{5}{2}$. With the $J = \frac{5}{2}$ assignment, an unreasonable M2 enhancement is required, therefore $J^\pi = \frac{3}{2}^+$ and $x(\equiv (L+1)/L$ amplitude) = -0.04 ± 0.04 (1968EA03). The correlation in the cascade transition excludes $J^\pi = \frac{3}{2}^-$ for $^{11}\text{C}^*(2.0)$ and indicates $J^\pi = \frac{1}{2}^-$, $x = 0.0 \pm 0.03$ (1968EA03).

$E_x = 6.91$ MeV. The correlation analysis gives $J = \frac{5}{2}$, $x = 0.02 \pm 0.03$; $J = \frac{1}{2}$ and $\frac{3}{2}$ are eliminated.

$E_x = 6.48$ MeV. Analysis of $\alpha - \gamma$ correlation gives $J = \frac{7}{2}$, $x = -0.01 \pm 0.06$.

$E_x = 6.34$ MeV. The cascade transition is via $^{11}\text{C}^*(2.0)$ and not $^{11}\text{C}^*(4.31)$. From $^9\text{Be}(^3\text{He}, \text{n})^{11}\text{C}$, $J^\pi = \frac{1}{2}^+$ or $\frac{3}{2}^+$. For $J^\pi = \frac{3}{2}^+$ correlation analysis requires an excessive M2 admixture. The angular distribution of α 's at $E(^3\text{He}) = 11.0$ MeV gives an $l = 0$ pattern: $J^\pi = \frac{1}{2}^+$.

$E_x = 4.79$ MeV. The decay is to $^{11}\text{C}_{\text{g.s.}}$ and (2.0). With $J^\pi = \frac{1}{2}^-$ for $^{11}\text{C}^*(2.0)$, the correlation analysis eliminates $J^\pi = \frac{5}{2}^-$ and $\frac{1}{2}^-$ and gives for $J^\pi = \frac{3}{2}^-$, $x(\text{g.s.}) = -(0.04 \pm 0.04)$ or $-(3.28 \pm 0.49)$, and $x(4.8 \rightarrow 2.0) = +(0.03 \pm 0.05)$ or $+(1.60 \pm 0.12)$. See also (1966GA19, 1967BL22).

$E_x = 4.31$ MeV. Analysis of angular correlations gives $J = \frac{5}{2}$, $x = 0.17 \pm 0.03$. See also (1966GA19, 1967BL22).

$E_x = 2.00$ MeV. Stripping results give $l_n = 1$, $J^\pi = \frac{1}{2}^-$ or $\frac{3}{2}^-$. The $\alpha - \gamma$ correlation is isotropic (1965NE06, 1965WA06). The cascade from $^{11}\text{C}^*(7.51)$ excludes $J^\pi = \frac{3}{2}^-$. See also (1967BL22).

$$23. \ ^{12}\text{C}(^{14}\text{N}, ^{15}\text{N})^{11}\text{C} \quad Q_m = -7.885$$

See (1966PO1B, 1967BI06).

$$24. \ ^{13}\text{C}(\text{p}, \text{t})^{11}\text{C} \quad Q_m = -15.185$$

At $E_p = 43.7$ MeV, a triton group was observed corresponding to a $T = \frac{3}{2}$ state at $E_x = 12.45 \pm 0.08$ MeV, $\Gamma = 566 \pm 60$ keV, $J^\pi = \frac{1}{2}^-$ (1966MA2N). See also (1968TA1V).

$$25. \ ^{14}\text{N}(\text{p}, \alpha)^{11}\text{C} \quad Q_m = -2.920$$

At $E_p = 38$ MeV angular distributions have been obtained for $\alpha_0, \alpha_1, \alpha_2 + \alpha_3$ (1968GA1N).
See also (1961CL09, 1962MA1L, 1963BR14) and ^{15}O in (1970AJ04).

26. $^{16}\text{O}(\gamma, n\alpha)^{11}\text{C}$ $Q_m = -25.881$

See (1955AJ61) and (1962BI09).

^{11}N
(Not illustrated)

See (1960GO1B, 1966KE16).

References

(Closed 15 December 1967)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1941SM11 J.H.C. Smith and D.B. Cowie, *J. Appl. Phys.* 12 (1941) 78
- 1941SO01 A.K. Solomon, *Phys. Rev.* 60 (1941) 279
- 1948EG1A Egger, Hughes and Huddleston, *Phys. Rev.* 74 (1948) 1238
- 1949EV1A Evans, Malich and Risser, *Phys. Rev.* 75 (1949) 1161
- 1951BE13 W.E. Bennett, P.A. Roys and B.J. Toppel, *Phys. Rev.* 82 (1951) 20
- 1951BO45 C.K. Bockelman, D.W. Miller, R.K. Adair and H.H. Barschall, *Phys. Rev.* 84 (1951) 69
- 1951BR10 A.B. Brown, C.W. Snyder, W.A. Fowler and C.C. Lauritsen, *Phys. Rev.* 82 (1951) 159
- 1951DI12 J.M. Dickson and T.C. Randle, *Proc. Phys. Soc. (London)* A64 (1951) 902
- 1951LI29 C.W. Li and W. Whaling, *Phys. Rev.* 82 (1951) 122
- 1951PE18 B. Petree, C.H. Johnson and D.W. Miller, *Phys. Rev.* 83 (1951) 1148
- 1951VA1A Van Patter, Buechner and Sperduto, *Phys. Rev.* 82 (1951) 248
- 1952CA19 F.L. Canavan, *Phys. Rev.* 87 (1952) 136
- 1952DA05 R.B. Day and R.L. Walker, *Phys. Rev.* 85 (1952) 582
- 1952DE24 D. deJong, P.M. Endt and L.J.G. Simons, *Physica* 18 (1952) 676
- 1952JO10 V.R. Johnson, *Phys. Rev.* 86 (1952) 302
- 1953EL12 M.M. Elkind, *Phys. Rev.* 92 (1953) 127
- 1953GE01 R.W. Gelinas and S.S. Hanna, *Phys. Rev.* 89 (1953) 483
- 1953KU08 D.N. Kundu, T.W. Donaven, M.L. Pool and J.K. Long, *Phys. Rev.* 89 (1953) 1200
- 1953SP1A Sperduto and Fader, *M.I.T. Prog. Rept.* (May, 1953)
- 1954DA20 R.B. Day and T. Huus, *Phys. Rev.* 95 (1954) 1003
- 1954HE22 N.P. Heydenberg and G.M. Temmer, *Phys. Rev.* 94 (1954) 1252
- 1954LI48 C.W. Li and R. Sherr, *Phys. Rev.* 96 (1954) 389
- 1954RI15 F.L. Ribe and J.D. Seagrave, *Phys. Rev.* 94 (1954) 934
- 1955AJ61 F. Ajzenberg and T. Lauritsen, *Rev. Mod. Phys.* 27 (1955) 77

1955BA22 J.K. Bair, J.D. Kington and H.B. Willard, Phys. Rev. 100 (1955) 21.
1955BA63 W.C. Barber, W.D. George and D.D. Reagan, Phys. Rev. 98 (1955) 73
1955BE62 R.D. Bent, T.W. Bonner and R.F. Sippel, Phys. Rev. 98 (1955) 1237
1955BE81 R.D. Bent, T.W. Bonner, J.H. McCrary, W.A. Ranken and R.F. Sippel, Phys. Rev. 99 (1955) 710
1955BO1A Borelli and Lalovic, Nature 176 (1955) 1021
1955GR18 G.L. Griffith, Phys. Rev. 98 (1955) 579
1955HA01 T.M. Hahn Jr., B.D. Kern and G.K. Farney, Phys. Rev. 98 (1955) 1183A
1955HE83 R.E. Heft and W.F. Libby, Phys. Rev. 100 (1955) 799
1955JU10 M.K. Juric, Phys. Rev. 98 (1955) 85
1955JU1B Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 5 (1955) 7
1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252
1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847.
1955RA14 D.C. Ralph and F.E. Dunnam, Phys. Rev. 98 (1955) 249A
1955RA41 V.K. Rasmussen, D.W. Miller, M.B. Sampson and U.C. Gupta, Phys. Rev. 100 (1955) 851
1955SA1B Sample, Neilson, Chadwick and Warren, Can. J. Phys. 33 (1955) 828
1955WI25 H.B. Willard, J.K. Bair and J.D. Kington, Phys. Rev. 98 (1955) 669
1956AJ22 F. Ajzenberg-Selove, G.D. Johnson, A. Rubin and M. Mazari, Phys. Rev. 103 (1956) 356
1956AL23 H.R. Allan, M. Govindjee and N. Sarma, Proc. Phys. Soc. (London) A69 (1956) 350
1956CE1B M. Cerineo, Physica 22 (1956) 1154A
1956CE73 M. Cerineo, Nucl. Phys. 2 (1956) 113
1956CH20 G.B. Chadwick, T.K. Alexander and J.B. Warren, Can. J. Phys. 34 (1956) 381
1956CR07 J.W. Cronin, Phys. Rev. 101 (1956) 298
1956GR54 A. Graue, Phil. Mag. 1 (1956) 1027
1956JU17 M.J. Juric and S.D. Cirilov, Bull. Inst. Nucl. Sci. Boris Kidrich 6 (1956) 45
1956KU1A Kurath, Phys. Rev. 101 (1956) 216
1956MA83 E.E. Maslin, J.M. Calvert and A.A. Jaffe, Proc. Phys. Soc. (London) A69 (1956) 754
1956SE1A Selove, Phys. Rev. 101 (1956) 231
1957BA18 G.A. Bartholomew and P.J. Campion, Can. J. Phys. 35 (1957) 1347
1957BI84 H. Bichsel and T.W. Bonner, Phys. Rev. 108 (1957) 1025

- 1957BR18 D.A. Bromley, E. Almqvist, H.E. Gove, A.E. Litherland, E.B. Paul and A.J. Ferguson, Phys. Rev. 105 (1957) 957
- 1957GR50 A. Graue and B. Trumpy, Phil. Mag. 2 (1957) 138
- 1957HU79 S.E. Hunt, R.A. Pope and W.W. Evans, Phys. Rev. 106 (1957) 1012
- 1957KU58 D. Kurath, Phys. Rev. 106 (1957) 975
- 1957MC35 J.H. McCrary, T.W. Bonner and W.A. Ranken, Phys. Rev. 108 (1957) 392
- 1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
- 1957PR53 Y.D. Prokoshkin and A.A. Tyapkin, Zh. Eksp. Teor. Fiz. 32 (1957) 177; JETP (Sov. Phys.) 5 (1957) 148
- 1957SC29 J. Scobie and G.M. Lewis, Phil. Mag. 2 (1957) 1089
- 1957SE1B Sehgal, Indian J. Phys. 31 (1957) 329
- 1957SH65 A.I. Shpetni, Zh. Eksp. Teor. Fiz. 32 (1957) 423; JETP (Sov. Phys.) 5 (1957) 357
- 1957SJ66 B. Sjogren, Ark. Fys. 12 (1957) 537
- 1957WI26 D.H. Wilkinson, Phys. Rev. 105 (1957) 666
- 1958AL96 D.E. Alburger and D.H. Wilkinson, Phil. Mag. 3 (1958) 1332
- 1958AR15 S.E. Arnell, J. Dubois and O. Almen, Nucl. Phys. 6 (1958) 196
- 1958DO63 T. Doke, S. Suzuki and I. Ogawa, J. Phys. Soc. Jpn. 13 (1958) 656
- 1958EV01 N.T.S. Evans and A.P. French, Phys. Rev. 109 (1958) 1272
- 1958FE70 A.J. Ferguson, H.E. Gove, J.A. Kuehner, A.E. Litherland, E. Almqvist and D.A. Bromley, Phys. Rev. Lett. 1 (1958) 414
- 1958HE47 J.C. Hensel and W.C. Parkinson, Phys. Rev. 110 (1958) 128
- 1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
- 1958JU38 M.K. Juric and N.D. Zarubica, Bull. Inst. Nucl. Sci. Boris Kidrich 8 (1958) 17
- 1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1
- 1958ME79 F.R. Metzger, C.P. Swann and V.K. Rasmussen, Phys. Rev. 110 (1958) 906
- 1958NU40 M.J. Nurmia and R.W. Fink, Phys. Rev. Lett. 1 (1958) 23
- 1958RA13 W.A. Ranken, T.W. Bonner, J.M. McCrary and T.A. Rabson, Phys. Rev. 109 (1958) 917
- 1958RA14 V.K. Rasmussen, F.R. Metzger and C.P. Swann, Phys. Rev. 110 (1958) 154
- 1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10
- 1958WI41 D.H. Wilkinson, Phys. Rev. 109 (1958) 1610
- 1958ZE01 B. Zeidman and J.M. Fowler, Phys. Rev. 112 (1958) 2020

- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AL96 E. Almqvist, D. A. Bromley, A. J. Ferguson, H. E. Gove, and A. E. Litherland, Phys. Rev. 114 (1959) 1040
- 1959BO1C Bockelman, Nucl. Phys. 13 (1959) 205
- 1959BR06 D.A. Bromley, A.J. Ferguson, H.E. Gove, J.A. Kuehner, A.E. Litherland, E. Almqvist and R. Batchelor, Can. J. Phys. 37 (1959) 1514
- 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
- 1959BR75 O.D. Brill and L.V. Sumin, Atomnaya Energ. 7 (1959) 377
- 1959BU1F Buck and Satchler, Congres Int. de Phys. Nucleaire (1958), Dundod, Paris (1959) 363
- 1959CO95 L. Cohen, R.A. Tobin and J. McElhinney, Phys. Rev. 114 (1959) 590
- 1959CR1A Croissiaux, Thesis, Univ. of Starsbourg (1959)
- 1959EG1C Egardt, Nucl. Phys. 12 (1959) 84
- 1959EG20 L. Egardt, Nucl. Phys. 11 (1959) 349
- 1959EL43 M. El Nadi and M. El Khishin, Proc. Phys. Soc. (London) A73 (1959) 705
- 1959FA1A Fagg and Hanna, Rev. Mod. Phys. 31 (1959) 711
- 1959GA05 A. Gallmann, Ann. Phys. 4 (1959) 185
- 1959GA14 F. Gabbard, H. Bischel and T.W. Bonner, Nucl. Phys. 14 (1959) 277
- 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
- 1959GO69 S. Gorodetzky, M. Croissiaux, A. Gallmann, P. Fintz, J. Samuel and G. Frick, Compt. Rend. 248 (1959) 550
- 1959GR1B Grieder, Phys. Rev. 114 (1959) 786
- 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295
- 1959HA1J Hanna and Meyer-Schutzmeister, Congres Int. de Phys. Nucl., 1958 (Dunod, Paris, 1959) 650
- 1959HI1E Hird, Cookson and Bokhari, Congres Int. de Phys. Nucl., 1958 (Dunod, Paris, 1959) 470
- 1959HI69 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 196
- 1959HO1D Honda and Nagasaki, Proc. Phys. Soc. (London) A74 (1959) 517
- 1959JO25 G.A. Jones, C.M.P. Johnson and D.H. Wilkinson, Phil. Mag. 4 (1959) 796
- 1959JO43 A. Johansson, G. Tibell and P. Hillman, Nucl. Phys. 11 (1959) 540
- 1959LE1B Levin and MacDonald, Bull. Amer. Phys. Soc. 4 (1959) 58
- 1959LI1D Litherland, Almqvist, Batchelor and Gove, Phys. Rev. Lett. 2 (1959) 104
- 1959MA1C MacGregor, Congr. Int. De. Phys. Nucl., 1958 (Dunod, Paris, 1959) 612

- 1959ME24 U. Meyer-Berkhout, K.W. Ford and A.E.S. Green, *Ann. Phys. (N.Y.)* 8 (1959) 119
- 1959NE1A Neilson, Dawson and Johnson, *Rev. Sci. Instrum.* 30 (1959) 963
- 1959PO61 B. Povh, *Phys. Rev.* 114 (1959) 1114
- 1959TO1A Tobocman, *Phys. Rev.* 115 (1959) 98
- 1959VL23 N.A. Vlasov, S.P. Kalinin, A.A. Ogloblin and V.J. Chuev, *Zh. Eksp. Teor. Fiz.* 37 (1959) 1187; *JETP (Sov. Phys.)* 10 (1960) 844
- 1959WI49 D.H. Wilkinson and D.E. Alburger, *Phys. Rev.* 113 (1959) 563
- 1959WO53 E.A. Wolicki, H.D. Holmgren, R.L. Holmgren, R.L. Johnston and E.G. Illsley, *Phys. Rev.* 116 (1959) 1585
- 1960AN14 A.L. Androsenko, D.L. Broder and A.I. Lashuk, *At. Energ.* 9 (1960) 403; *Sov. J. At. Energy* 9 (1961) 945
- 1960BA46 R. Bardes and G.E. Owen, *Phys. Rev.* 120 (1960) 1369
- 1960BI04 E.G. Bilpuch, L.W. Weston and H.W. Newson, *Ann. Phys. (N.Y.)* 10 (1960) 455
- 1960BI07 O.M. Bilaniuk and J.B. French, *Nucl. Phys.* 17 (1960) 435
- 1960BI08 O.M. Bilaniuk and J.C. Hensel, *Phys. Rev.* 120 (1960) 211
- 1960BI1B Biedenharn and Satchler, *Helv. Phys. Acta Suppl.* 6 (1960) 372
- 1960BO23 E.C. Booth, *Nucl. Phys.* 19 (1960) 426
- 1960BU25 J. Button and R. Mermod, *Phys. Rev.* 118 (1960) 1333
- 1960CR1A Croissiaux, *Ann. Phys.* 5 (1960) 409
- 1960DA08 R.B. Day and M. Walt, *Phys. Rev.* 117 (1960) 1330
- 1960EL1C El-Nadi, *Phys. Rev.* 120 (1960) 1360
- 1960ER1A Ero, Pocs, Szentpetery and Zimanyi, *Rept. Central Research Inst. Phys. Hung. Acad. Sci* 8 (1960) 263
- 1960FE01 A.J. Ferguson, H.E. Gove, A.E. Litherland and R. Batchelor, *Bull. Amer. Phys. Soc.* 5 (1960) 45, MA7
- 1960FE13 A.J. Ferguson, H.E. Gove, A.E. Litherland and R. Batchelor, *Proc. Int. Conf. on Nucl. Struct., Kingston, Canada*; Eds. D.A. Bromley and E.W. Vogt (Univ. of Toronto Press, 1960) 364
- 1960FO01 P.D. Forsyth, F. de S. Barros, A.A. Jaffe, I.J. Taylor and S. Ramavataram, *Proc. Phys. Soc. (London)* A75 (1960) 291
- 1960GO11 S. Gorodetzky, M. Croissiaux, P. Fintz, J. Samuel, G. Bassompierre, R. Armbruster and P. Chevallier, *Compt. Rend.* 250 (1960) 2874
- 1960GO14 S. Gorodetzky, M. Croissiaux, A. Gallmann, P. Fintz, J. Samuel and G. Bassompierre, *Compt. Rend.* 250 (1960) 3153

- 1960GO16 T.J. Gooding and H.G. Pugh, Nucl. Phys. 18 (1960) 46
- 1960GO1B Goldansky, Nucl. Phys. 19 (1960) 482
- 1960GO1M Gottschalk and Strauch, Phys. Rev. 120 (160) 1005
- 1960GO25 S. Gorodetzky, M. Croissiaux, A. Gallmann, P. Fintz, J. Samuel and G. Bassompierre, Nucl. Phys. 18 (1960) 286
- 1960HA08 G.R. Harrison, G.D. Schmidt and C.D. Curtis, Phys. Rev. 117 (1960) 532
- 1960HI04 K. Hisatake, Y. Ishizaki, A. Isoya, T. Nakamura, Y. Nakano, B. Saheki, Y. Saji and K. Yuasa, J. Phys. Soc. Jpn. 15 (1960) 741
- 1960HI07 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) 75 (1960) 745
- 1960HI08 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) 75 (1960) 754
- 1960HI09 B. Hird and A. Strzalkowski, Proc. Phys. Soc. (London) 75 (1960) 868
- 1960LU04 C.R. Lubitz, Bull. Amer. Phys. Soc. 5 (1960) 346, F2
- 1960MA32 R.D. Macfarlane and J.B. French, Rev. Mod. Phys. 32 (1960) 567
- 1960MC1C McDonnell et al., Aust. J. Phys. 13 (1960) 99
- 1960MU07 J. Muto, F. de S. Barros and A.A. Jaffe, Proc. Phys. Soc. (London) 75 (1960) 929
- 1960NE17 G.C. Neilson, W.K. Dawson and J.T. Sample, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds. D.A. Bromley and E.W. Vogt (Univ. of Toronto Press, 1960) 505
- 1960NE1C Neudachin, Teplov and Tulinov, JETP (Sov. Phys.) 10 (1960) 387
- 1960NO1A Norbeck, Bull. Amer. Phys. Soc. 5 (1960) 476
- 1960OV1A Overley, Thesis, CalTech (1960)
- 1960RA12 P. Radvanyi and J. Genin, J. Phys. Rad. 21 (1960) 322
- 1960RE05 K. Reibel and A.K. Mann, Phys. Rev. 118 (1960) 701
- 1960RI1B Riley, Pugh and Gooding, Nucl. Phys. 18 (1960) 65
- 1960SA13 G.J. Safford, T.I. Taylor, B.M. Rustad and W.W. Havens Jr., Phys. Rev. 119 (1960) 1291
- 1960SC15 H.W. Schmitt, R.C. Block and R.L. Bailey, Nucl. Phys. 17 (1960) 109
- 1960SH01 S.M. Shafroth, Bull. Amer. Phys. Soc. 5 (1960) 55, RA1
- 1960SI02 P.P. Singh and G.M. Griffiths, Bull. Amer. Phys. Soc. 5 (1960) 346, F3
- 1960SP08 R.R. Spencer, G.C. Phillips and T.E. Young, Nucl. Phys. 21 (1960) 310
- 1960SU09 M. Suffert, D. Magnac-Valette and J. Yoccoz, Compt. Rend. 251 (1960) 2335
- 1960TA07 I. Talmi and I. Unna, Phys. Rev. Lett. 4 (1960) 469
- 1960TA1C Talmi and Unna, Ann. Rev. Nucl. Sci. 10 (1960) 353

- 1960TA27 M. Takeda, S. Kato, C. Hu and N. Takahashi, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds. D.A. Bromley and E.W. Vogt (Univ. of Toronto Press, 1960) 400
- 1960YO02 J.C. Young, C.W. Cook and R.B. Walton, Bull. Amer. Phys. Soc. 5 (1960) 230, C12
- 1961AL07 R.D. Albert, S.D. Bloom and N.K. Glendenning, Phys. Rev. 122 (1961) 862
- 1961BA10 F.De S. Barros, P.D. Forsyth, A.A. Jaffe and I.J. Taylor, Proc. Phys. Soc. (London) 77 (1961) 853
- 1961BA1D Badalyan and Baz, Zh. Eksp. Teor. Fiz. 40 (1961) 549; JETP (Sov. Phys.) 13 (1961) 383
- 1961BA1E Balashov, Neudachin and Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 170; Bull. Acad. Sci. USSR Phys. 25 (1961) 165
- 1961BA1F Barker, Phys. Rev. 122 (1961) 572
- 1961BE13 E.H. Beckner, R.L. Bramblett, G.C. Phillips and T.A. Eastwood, Phys. Rev. 123 (1961) 2100
- 1961BE24 A.A. Bergman and F.L. Shapiro, Zh. Eksp. Teor. Fiz. 40 (1961) 1270; JETP (Sov. Phys.) 13 (1961) 895
- 1961BU16 B. Buck and P.E. Hodgson, Phil. Mag. 6 (1961) 1371
- 1961CA1D Catala, Lleo, Garcia and Aguilar, An. Real. Soc. Espan. Fis. y Quim. A57 (1961) 51; Phys. Abs. 3612 (1962)
- 1961CL09 A.B. Clegg, K.J. Foley, G.L. Salmon and R.E. Segel, Proc. Phys. Soc. (London) 78 (1961) 681
- 1961CL10 A.B. Clegg, Phil. Mag. 6 (1961) 1207
- 1961CO33 M. Coste, C. Lemeille, N. Saunier and L. Marquez, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 583; Centro Brasileiro De Pesquisas Fisicas 8 (1961)
- 1961DA16 E.A. Davis, F. Gabbard, T.W. Bonner and R. Bass, Nucl. Phys. 27 (1961) 448
- 1961DO03 P.F. Donovan, J.V. Kane, R.E. Pixley and D.H. Wilkinson, Phys. Rev. 123 (1961) 589.
- 1961DO08 W.R. Dodge, Thesis, Stanford Univ. (1961); HEPL-246 (1961); Nucl. Sci. Abstr. 16, 750, Abstr. 5921 (1962)
- 1961DU1B Duggan, Thesis, Louisiana State Univ. (1961)
- 1961EL1A El Nadi and Sherif, Nucl. Phys. 28 (1961) 331
- 1961FO07 D.B. Fossan, R.L. Walter, W.E. Wilson and H.H. Barschall, Phys. Rev. 123 (1961) 209
- 1961GA14 J.P. Garron, J.C. Jacmart, M. Riou and Ch. Ruhla, J. Phys. Rad. 22 (1961) 622
- 1961GO13 H.E. Gove, A.E. Litherland and R. Batchelor, Nucl. Phys. 26 (1961) 480

- 1961HI08 J. Hinds and R. Middleton, Proc. Phys. Soc. (London) 78 (1961) 81
- 1961HO19 R.K. Hobbie, C.W. Lewis and J.M. Blair, Phys. Rev. 124 (1961) 1506
- 1961HO1F Holmgren and Wolicki, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 541
- 1961IS01 T. Ishimatsu, N. Takano, Y. Hachiya and T. Nakashima, J. Phys. Soc. Jpn. 16 (1961) 367
- 1961JA11 A.N. James, Nucl. Phys. 24 (1961) 675
- 1961JA12 A.N. James, A.T.G. Ferguson and C.M.P. Johnson, Nucl. Phys. 25 (1961) 282
- 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, Rev. Mex. Fisica 10 (1961) 247
- 1961KO08 J. Kokame, J. Phys. Soc. Jpn. 16 (1961) 2101
- 1961KO1A Koltun, Phys. Rev. 124 (1961) 1162
- 1961KU1C Kurath, Phys. Rev. 124 (1961) 552
- 1961LE01 J.J. Leigh and J.M. Blair, Phys. Rev. 121 (1961) 246
- 1961LE08 O.C. Lemeille, L. Marquez, N. Saunier and M. Coste, J. Phys. Rad. 22 (1961) 349; IBID 21 (1960) 355; Centro Brasileiro De Pesquisas Fisicas 8 (1961)
- 1961LE09 J.J. Leigh, Phys. Rev. 123 (1961) 2145
- 1961LE1F Levin, Bull. Amer. Phys. Soc. 6 (1961) 67
- 1961LI04 D.A. Lind and R.B. Day, Ann. Phys. 12 (1961) 485
- 1961MA1F Maluszynska, Natanson and Yuan-Han-Yung, Bull. Acad. Pol. Sci. Ser. Sci. Math. Astron. Phys. 9 (1961) 119
- 1961ME02 J.W. Meadows and J.F. Whalen, Nucl. Sci. Eng. 9 (1961) 132
- 1961MO02 G.C. Morrison, Phys. Rev. 121 (1961) 182
- 1961PU1A Pugh and Riley, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 195
- 1961PU1B Pullen, Wilkinson and Whitehead, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 565
- 1961RE01 W. Reichelt and G.E. Owen, Phys. Rev. 121 (1961) 547
- 1961RE03 F.H. Read and J.M. Calvert, Proc. Phys. Soc. (London) 77 (1961) 65
- 1961RE04 F.H. Read, J.M. Calvert and G. Schork, Nucl. Phys. 23 (1961) 386
- 1961RO05 L. Rosen, J.E. Brolley Jr. and L. Stewart, Phys. Rev. 121 (1961) 1423
- 1961RO21 J.C. Roy and J.J. Hawton, Can. J. Phys. 39 (1961) 1528
- 1961RU1B Rustgi, Nucl. Phys. 27 (1961) 58
- 1961SA02 G.J. Safford, T.I. Taylor, B.M. Rustad and W.W. Havens Jr., Nucl. Sci. Eng. 9 (1961) 99

- 1961SN01 J.L. Snider, M. Posner, A.M. Bernstein and D.R. Hamilton, *Bull. Amer. Phys. Soc.* 6 (1961) 224, A1
- 1961SU17 M. Suffert, D. Magnac-Valette and J. Yoccoz, *J. Phys. Rad.* 22 (1961) 565
- 1961TA12 S. Takayanagi, N.H. Gale, J.B. Garg and J.M. Calvert, *Nucl. Phys.* 28 (1961) 494
- 1961TE02 A. Tejera, M. Mazari, A. Jaidar and G. Lopez, *Rev. Mex. Fis.* 10 (1961) 229
- 1961TR1B True and Warburton, *Nucl. Phys.* 22 (1961) 426
- 1961UM1A Umezawa, *Nucl. Phys.* 25 (1961) 661
- 1961VA03 A.M.K. Van Beek and G.O. Andre, *Nucl. Phys.* 24 (1961) 102
- 1961ZE02 B. Zeidman and J.L. Yntema, *Bull. Amer. Phys. Soc.* 6 (1961) 225, AA3
- 1961ZE03 B. Zeidman, J.L. Yntema and G.R. Satchler, *Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 515*
- 1961ZI02 J. Zimanyi, J. Ero, L. Pocs and J. Szentpeteryi, *Zh. Eksp. Teor. Fiz.* 40 (1961) 709; *JETP (Sov. Phys.)* 13 (1961) 496
- 1962AG01 J. Aguilar, J.B.A. England and A. Garcia, *An. Real. Soc. Espan. Fis. y Quim. (Madrid)* A58 (1962) 319; *Phys. Abs.* 12699 (1963)
- 1962AL10 R.G. Allas, R.W. Bercaw and F.B. Shull, *Phys. Rev.* 127 (1962) 1252
- 1962AN11 G.B. Andreev, A.S. Deineko and I.Y. Malakhov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 26 (1962) 1134; *Bull. Acad. Sci. USSR Phys. Ser.* 26 (1963) 1147
- 1962AU1A Austin, Slamon, Clegg, Foley and Newton, *Proc. Phys. Soc. (London)* 80 (1962) 383
- 1962BA1A Balashov and Boyarkina, *Nucl. Phys.* 38 (1962) 629
- 1962BA1D Barber, *Ann. Rev. Nucl. Sci.* 12 (1962) 1
- 1962BA1J Balashov and Boyarkina, *Izv. Akad. Nauk SSSR Ser. Fiz.* 26 (1962) 1196
- 1962BE16 E.H. Berkowitz, *Phys. Rev.* 126 (1962) 2168
- 1962BE1D Bessis, Lefebvre-Brion and Moser, *Phys. Rev.* 128 (1962) 213
- 1962BE1H Benioff, *Phys. Rev.* 128 (1962) 740
- 1962BE24 E. Berkowitz, S. Bashkin, R.R. Carlson, S.A. Coon and E. Norbeck, *Phys. Rev.* 128 (1962) 247
- 1962BI09 G.R. Bishop, B. Grossetete and J.C. Risset, *J. Phys. Rad.* 23 (1962) 31
- 1962BI11 J.A. Biggerstaff, R.F. Hood, H. Scott and M.T. McEllistrem, *Nucl. Phys.* 36 (1962) 631
- 1962BO17 E.C. Booth and K.A. Wright, *Nucl. Phys.* 35 (1962) 472
- 1962BR10 C.P. Browne, W.E. Dorenbusch and J.R. Erskine, *Phys. Rev.* 125 (1962) 992
- 1962BR34 F.P. Brady and E.K. Warburton, *Phys. Rev.* 128 (1962) 790

- 1962CA29 J. Catala, A. Garcia and J. de Rubia, An. Real. Soc. Espan. Fis. y Quim. A58 (1962) 227
- 1962CH26 V.P. Chizhov, et al., Nucl. Phys. 34 (1962) 562
- 1962CL13 A.B. Clegg, Nucl. Phys. 38 (1962) 353
- 1962CO17 C.F. Coleman, P.E. Cavanagh, B.W. Ridley and J.F. Turner, Conf. Low Energy Nucl. Phys. Harwell, AERE-R-4131, 8, 3.3; Sept. 1962 (1962)
- 1962DI1A Dietrich, Phys. Lett. 2 (1962) 139
- 1962DO1A Dodge and Barber, Phys. Rev. 127 (1962) 1746
- 1962DY1A Dyal and Hummel, Phys. Rev. 127 (1962) 2217
- 1962ED02 R.D. Edge and G.A. Peterson, Phys. Rev. 128 (1962) 2750
- 1962FR06 R.M. Freeman, Nucl. Phys. 37 (1962) 215.
- 1962GA09 J.P. Garron, J.C. Jacmart, M. Riou, C. Ruhla, J. Teillac and K. Strauch, Nucl. Phys. 37 (1962) 126
- 1962GA17 A. Garcia, J. Kirk, J.B.A. England and P.E. Hodgson, Nucl. Phys. 38 (1962) 372
- 1962GA23 J.P. Garron, Ann. Phys. (Paris) 7 (1962) 301
- 1962GO09 S. Gorodetzky, F. Scheibling, P. Chevallier, P. Mennrath and G. Sutter, Phys. Lett. 1 (1962) 24
- 1962GO1P Gottschalk, Unpublished Thesis, Harvard Univ. (1962)
- 1962GR07 L.L. Green, G.A. Stephens and J.C. Willmott, Proc. Phys. Soc. (London) 79 (1962) 1017
- 1962GR14 J.A. Green and W.C. Parkinson, Phys. Rev. 127 (1962) 926
- 1962GU01 G.D. Gutsche, H.D. Holmgren, L.M. Cameron and R.L. Johnston, Phys. Rev. 125 (1962) 648
- 1962GU10 M. Gusakow, Ann. Phys. 7 (1962) 67
- 1962HI07 S. Hinds and R. Middleton, Nucl. Phys. 38 (1962) 114
- 1962IN1A Inglis, Rev. Mod. Phys. 34 (1962) 165
- 1962KI02 K. Kimura, Y. Uemura, R. Ishiwari, J. Kokame, K. Fukunaga, A. Katase, J. Muto, I. Kumabe, H. Ogata, T. Oasme et al., J. Phys. Soc. Jpn. 17 (1962) 9
- 1962LI13 F.F. Liu, F.J. Loeffler, T.R. Palfrey and Y.S. Kim, Phys. Rev. 128 (1962) 2784
- 1962MA1L MacLeod and Reid, AERE-R-4131 (1962)
- 1962ME1B Menyhard and Zimanyi, Nucl. Phys. 29 (1962) 687
- 1962NE01 E. Newman, Phys. Rev. 125 (1962) 600
- 1962NE1D Nedvedyk, Joint Inst. Nucl. Res. Lab. Neutron Phys. USSR, Rept. No. P-1098 (1962)
- 1962OP03 T.R. Ophel, R.N. Glover and E.W. Titterton, Nucl. Phys. 33 (1962) 198

1962OV02 J.C. Overley and W. Whaling, Phys. Rev. 128 (1962) 315
 1962PA08 Parikh, Nucl. Phys. 38 (1962) 529
 1962PA12 M.V. Pasechnik, L.S. Saltykov and D.I. Tambovtsev, Zh. Eksp. Teor. Fiz. 43 (1962) 1575; JETP (Sov. Phys.) 16 (1963) 1111
 1962PU01 D.J. Pullen, A.E. Litherland, S. Hinds and R. Middleton, Nucl. Phys. 36 (1962) 1
 1962RA01 P. Radvanyi, J. Genin and C. Detraz, Phys. Rev. 125 (1962) 295
 1962SE02 F.D. Seward, Phys. Rev. 125 (1962) 335
 1962SE1A Serov and Guzhovskii, Atomn. Energ. (USSR) 12 (1962) 5
 1962SL03 R.J. Slobodrian, Nucl. Phys. 32 (1962) 684
 1962SL04 R.J. Slobodrian, Phys. Rev. 126 (1962) 1059
 1962TA13 D.I. Tambovtsev, Ukr. Fiz. Zh. 7 (1962) 244; Nucl. Sci. Abstr. 16, 3442, Abstr. 26347 (1962); Phys. Abs. 20672 (1962)
 1962TE05 K. Tesch, Nucl. Phys. 37 (1962) 412
 1962VA1G Varfolomeev, Zh. Eksp. Teor. Fiz. 42 (1962) 713; JETP (Sov. Phys.) 15 (1962) 496
 1962VO1C Volkov and Kulchitskii, Zh. Eksp. Teor. Fiz. 42 (1962) 108; JETP (Sov. Phys.) 15 (1962) 77
 1962WA31 E.K. Warburton and H.O. Funsten, Phys. Rev. 128 (1962) 1810
 1962WE1C Wegner, Rev. Sci. Instrum. 33 (1962) 271
 1962WI15 B. Wilkins and G. Igo, Phys. Lett. 3 (1962) 48
 1962ZI01 J. Zimanyi, J. Ero, L. Pocs and I. Szentpetery, Nucl. Phys. 33 (1962) 524
 1963AL21 D.E. Alburger and E.K. Warburton, Phys. Rev. 132 (1963) 790
 1963AU1A Austern, in Fast Neutron Phys., Interscience, Ed. Marion and Fowler (1963)
 1963BA1K Balashov and Fetisov, Zh. Eksp. Teor. Fiz. 45 (1963) 532; JETP (Sov. Phys.) 18 (1964) 365
 1963BA1R Bachelier et al., Padua (1963) 1141
 1963BA1T Ballini, Lemeille, Marquez and Saunier, Padua (1963) 1008A
 1963BE36 N. Bessis, H. Lefebvre-Brion and C.M. Moser, Rev. Mod. Phys. 35 (1963) 548
 1963BE42 T. Berggren and G. Jacob, Nucl. Phys. 47 (1963) 481
 1963BO1J Boschitz, Padua (1963) 640
 1963BR14 R.E. Brown, Astrophys. J. 137 (1963) 338
 1963BR1H D.W. Braben, P.J. Riley and G.C. Neilson, Can. J. Phys. 41 (1963) 784.
 1963CA09 R.R. Carlson and E. Norbeck, Phys. Rev. 131 (1963) 1204
 1963CL07 A.B. Clegg, Phys. Lett. 7 (1963) 132

1963CL1B Clegg, CERN 63-28 (1963)
 1963CO1D Costa et al., Phys. Lett. 6 (1963) 226
 1963DU12 J.L. Duggan, P.D. Miller and R.F. Gabbard, Nucl. Phys. 46 (1963) 336
 1963EA01 L.G. Earwaker, J.G. Jenkin and E.W. Titterton, Nucl. Phys. 42 (1963) 521
 1963ER1A Ero, Pocs, Szentpetery and Zimanyi, Padua (1963) 558A
 1963FU05 H. Fuchs and D. Haag, Z. Physik 171 (1963) 403
 1963GL1C Glendenning, Ann. Rev. Nucl. Sci. 13 (1963) 191
 1963GL1F Glazkov, Atomn. Energ. (USSR) 15 (1963) 416
 1963GO04 J. Goldemberg and Y. Torizuka, Phys. Rev. 129 (1963) 312
 1963GO1M Goldberg, May and Stehn, BNL-400, 2nd Ed., Vol. 1 (1963)
 1963GR1G Griffiths, Knight, Gibson and Johnston, Padua (1963) 58
 1963GR20 D.E. Groce, J.H. McNally and W. Whaling, Bull. Amer. Phys. Soc. 8 (1963) 486, T6
 1963GU1A Gupta and Waghmare, Nucl. Phys. 48 (1963) 321
 1963HO19 H.D. Holmgren, L.M. Cameron and R.L. Johnston, Nucl. Phys. 48 (1963) 1
 1963IS1B Ishizaki et al., Padua (1963) 560A
 1963JE01 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Phys. Lett. 4 (1963) 142
 1963KA26 C.D. Kavaloski, G. Bassani and N.M. Hintz, Phys. Rev. 132 (1963) 813
 1963KI1C Kim, Liu, Loeffler and Palfrey, Phys. Rev. 129 (1963) 1362
 1963KN02 A.R. Knudson, Bull. Amer. Phys. Soc. 8 (1963) 292, A14
 1963KO15 E. Koltay, Acta Phys. Acad. Sci. Hung. 16 (1963) 93
 1963KO1G Kossler, PUC-1963-126 (1963)
 1963LE09 C. Lemeille, L. Marquez and N. Saunier, Phys. Lett. 5 (1963) 267
 1963LU05 C.A. Ludemann, H.D. Holmgren and J.E. Etter, Bull. Amer. Phys. Soc. 8 (1963) 303, D6
 1963MA1J Matsuda, Padua (1963) 499
 1963MA1K Manley and Stein, Bull. Amer. Phys. Soc. 8 (1963) 611
 1963MA24 B.B. Marsh and O.M. Bilaniuk, Phys. Rev. 130 (1963) 2373
 1963MA39 S. Malmskog, Physica 29 (1963) 987
 1963ME08 M.K. Mehta, W.E. Hunt, H.S. Plendl and R.H. Davis, Nucl. Phys. 48 (1963) 90
 1963MO04 I.L. Morgan, D.O. Nellis, R. Benjamin and J.B. Ashe, Bull. Amer. Phys. Soc. 8 (1963) 120, M6
 1963MO07 W.H. Moore and H.A. Enge, Bull. Amer. Phys. Soc. 8 (1963) 291, A9

- 1963MO1E Morinigo, Phys. Lett. 5 (1963) 215
- 1963NE07 B.M.K. Nefkens, Phys. Rev. Lett. 10 (1963) 243
- 1963NE1H Nemilov and Pobedonotsev, Zh. Eksp. Teor. Fiz. 45 (1963) 103; JETP (Sov. Phys.) 18 (1964) 76
- 1963NI04 K. Niedzwiedziuk, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teor. Fiz. 44 (1963) 1450; JETP (Sov. Phys.) 17 (1963) 974
- 1963NO02 E. Norbeck, S.A. Coon, R.R. Carlson and E. Berkowitz, Phys. Rev. 130 (1963) 1971
- 1963OP1A Oparin, Saukov and Shuvalov, Atomn. Energ. (USSR) 15 (1963) 411; J. Nucl. Energ. 18 (1964) 596
- 1963OV02 J.C. Overley, Nucl. Phys. 49 (1963) 537
- 1963PA15 V.M. Pankratov and I.N. Serikov, Zh. Eksp. Teor. Fiz. 44 (1963) 887; JETP (Sov. Phys.) 17 (1963) 604
- 1963PA1E Parks, Beard, Bilpuch and Newson, Bull. Amer. Phys. Soc. 8 (1963) 48
- 1963PA1G Pankratov and Serikov, Zh. Eksp. Teor. Fiz. 45 (1963) 910; JETP (Sov. Phys.) 18 (1964) 627
- 1963PR11 A. Prosdocimi and A.J. Deruytter, J. Nucl. Energy, Pt. A Reactor Sci. Tech. 17 (1963) 83
- 1963RA01 P. Radvanyi, D. Bachelier, M. Bernas, C. Detraz, J. Genin and J.N. Haag, Bull. Amer. Phys. Soc. 8 (1963) 10, BA2
- 1963RE16 J.H. Renken, Phys. Rev. 132 (1963) 2627
- 1963RI1B Riou, Padua (1963) 18
- 1963SC33 F. Scheibling, Ann. Phys. 8 (1963) 353; Thesis, Univ. of Strasbourg (1962)
- 1963SE08 B. Sen, Nucl. Phys. 41 (1963) 435
- 1963SE1K Serov and Zhmailo, Zh. Eksp. Teor. Fiz. 44 (1963) 332; JETP (Sov. Phys.) 17 (1963) 227
- 1963SM05 W.R. Smith and E.V. Ivash, Phys. Rev. 131 (1963) 304
- 1963SU09 M. Suffert, G. Costa and D. Magnac-Valette, J. Phys. 24 (1963) 1029
- 1963TA1A Tanifuji, Nucl. Phys. 40 (1963) 357
- 1963TA1C Tanifuji, Phys. Lett. 6 (1963) 276
- 1963TO1D Toth and Newman, 3rd Conf. on Reactions between Complex Nuclei (1963) 114
- 1963TO1E Toth, Phys. Rev. 131 (1963) 379
- 1963VA10 V.J. Vanhuyse and G.J. Vanpraet, Nucl. Phys. 43 (1963) 344
- 1963VA16 L. van der Zwan, A.T. Stewart, J.Y. Park and E. Merzbacher, Can. J. Phys. 41 (1963) 1036

1963VA1C Valentin, Albouy, Cohen and Gusakow, Phys. Lett. 7 (1963) 163
 1963WA18 H.D. Warren and A.P. Batson, Nucl. Phys. 48 (1963) 361
 1963WE08 J.A. Weinman and R.K. Smither, Nucl. Phys. 45 (1963) 260
 1963WI01 D.H. Wilkinson, D.E. Alburger, E.K. Warburton and R.E. Pixley, Phys. Rev. 129 (1963) 1643
 1963YU1A Yuasa and Poulet, J. Phys. 24 (1963) 1062
 1964AL05 J. Als-Nielsen and O. Dietrich, Phys. Rev. 133 (1964) B925
 1964AL22 D.E. Alburger, C. Chasman, K.W. Jones, J.W. Olness and R.A. Ristinen, Phys. Rev. 136 (1964) B916
 1964AM1D Amit and Latz, Nucl. Phys. 58 (1964) 297
 1964AN1B Anderson, Wong, McClure and Walker, Phys. Rev. 136 (1964) B118
 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209
 1964BA1C Balashov, Boyarkina and Rotter, Nucl. Phys. 59 (1964) 417
 1964BA1G Barz, Ann. Phys. 13 (1964) 164
 1964BE08 R.W. Bercaw and F.B. Shull, Phys. Rev. 133 (1964) B632
 1964BE1K Beregi, Neudatchin and Smirnov, Nucl. Phys. 60 (1964) 305
 1964BE31 E.M. Bernstein, Nucl. Phys. 59 (1964) 525
 1964BO1L Bowden, Bowman and Yuasa, Paris (1964) 136
 1964BO1M Bock, Duhm, Hortig and Rudel, Paris (1964) 136
 1964BO22 E.C. Booth, B. Chasan and K.A. Wright, Nucl. Phys. 57 (1964) 403
 1964BR13 H.C. Bryant, J.G. Beery, E.R. Flynn and W.T. Leland, Nucl. Phys. 53 (1964) 97
 1964BR1N Brix et al., Paris (1964) 41
 1964CA18 R.R. Carlson and M. Throop, Phys. Rev. 136 (1964) B630
 1964HA46 R.A. Haberstroh, W.J. Kossler, O. Ames and D.R. Hamilton, Phys. Rev. 136 (1964) B932
 1964HI08 S. Hinds and R. Middleton, Phys. Lett. 9 (1964) 149
 1964JA03 J.C. Jacmart, J.P. Garron, M. Riou and C. Ruhla, Phys. Lett. 8 (1964) 269
 1964JA1C Jackson, Nucl. Phys. 54 (1964) 561
 1964JE01 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Nucl. Phys. 50 (1964) 516
 1964KU05 H.-M. Kuan, T.W. Bonner and J.R. Risser, Nucl. Phys. 51 (1964) 481
 1964LI14 I. Lindgren, Perturbed Angular Correlations; Eds. E. Karlsson, E. Matthias and K. Siegbahn (Amsterdam, North-Holland Publ. Co., 1964) 379

- 1964LI1B Lindskog, Sundstrom and Sparrman, Perturbed Angular Correlations (Amsterdam, North-Holland Publ. Company, 1964) 411
- 1964LI1D Lim and McCarthy, Phys. Rev. 133 (1964) B1006
- 1964LO1C Loiseaux, Langevin and Maison, Congres Int. de Phys. Nucl., Paris (1964)
- 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
- 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed. W.H. Johnson Jr. (Springer Verlag, Vienna, 1964) 305
- 1964MO1D Morgan, Bull. Amer. Phys. Soc. 9 (1964) 653
- 1964NE06 D. Newton, A.B. Clegg, G.L. Salmon and D.J. Rowe, Nucl. Phys. 53 (1964) 433
- 1964NE1E Neudachin, Shevchenko and Yudin, Phys. Lett. 10 (1964) 180
- 1964OL1A Ollerhead, Chasman and Bromley, Phys. Rev. 134 (1964) B74
- 1964PA1E Pasechnik, Pucherov, Saltykov and Tambovtsev, Congres Int. de Phys. Nucl., Paris, 1964 (CNRS, Paris, 1964)
- 1964RE04 L.H. Reber and J.X. Saladin, Phys. Rev. 133 (1964) B1155
- 1964RE1B Reeder and Markowitz, Congres Int. de Phys. Nucl., Paris, 1964 (CRNS, Paris, 1964) 213
- 1964SA1D Satchler, Drisko and Bassel, Phys. Rev. 136 (1964) B637
- 1964SH07 T.H. Short and N.M. Hintz, Bull. Amer. Phys. Soc. 9 (1964) 391, BA16
- 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
- 1964ST1C Stromberg, Wiedling and Holmqvist, AE 139 (1964)
- 1964ST1K Starodubtsev and Khrushchev, Atomnaya Energ. 17 (1964) 59
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964TH03 M.J. Throop and R.R. Carlson, Bull. Amer. Phys. Soc. 9 (1964) 406, CA3
- 1964TI02 G. Tibell, O. Sundberg and P.U. Renberg, Ark. Fys. 25 (1964) 433
- 1965AJ01 F. Ajzenberg-Selove, J.W. Watson and R. Middleton, Phys. Rev. 139 (1965) B592
- 1965AL02 W.J. Alston III, J.E. Draper and R.H. Hilberg, Bull. Amer. Phys. Soc. 10 (1965) 94, HD3
- 1965AM1B Amaldi et al., Suppl. Nuovo Cim. 3 (1965) 1149
- 1965BA31 H.W. Barz, R. Fulle, D. Netzband, R. Reif, K. Schlott and J. Slotta, Nucl. Phys. 73 (1965) 473
- 1965BE1E Berggren, Ark. Fys. 30 (1965) 508
- 1965BU10 S.G. Buccino and A.B. Smith, Phys. Lett. 19 (1965) 234

1965CO1E Cortellessa, Suppl. Nuovo Cim. 3 (1965) 820
 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707
 1965DE1A Detraz, Cahiers Phys. 19 (1965) 359
 1965DI06 G. Din and J.L. Weil, Nucl. Phys. 71 (1965) 641
 1965DO14 R.E. Donaldson, L. Passell, W. Bartolini and D. Groves, Phys. Rev. 138 (1965) B1116
 1965FA1B Fabre de La Ripelle, Prog. Theor. Phys. 33 (1965) 38
 1965FA1C Wang, Chen and Sze, Acta Phys. Sin. 21 (1965) 140
 1965FU1G Fuller and Cohen, Appendix I, Nucl. Data Sheets 6-5 (1965)
 1965GA02 A. Gallman, F. Hibou, P. Fintz, P.E. Hodgson and E.K. Warburton, Phys. Rev. 138 (1965) B560
 1965GA1D Gaponov, Yad. Fiz. 2 (1965) 1002
 1965GL1E Glashausser, Thesis, Princeton Univ. (1965)
 1965GO05 S. Gorodetzky, A. Gallmann and R. Rebmeister, Phys. Rev. 137 (1965) B1466
 1965GO1K Goldemberg et al., Phys. Lett. 16 (1965) 141
 1965GR18 T.A. Griffy and D.U.L. Yu, Phys. Rev. 139 (1965) B880
 1965GR1R Gromov et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P-2184 (1965)
 1965HA17 D. Hasselgren, P.U. Renberg, O. Sundberg and G. Tibell, Nucl. Phys. 69 (1965) 81
 1965HA19 E. Hayward and T. Stovall, Nucl. Phys. 69 (1965) 241
 1965HA1K Hans, Thomas and Bollinger, Bull. Amer. Phys. Soc. 10 (1965) 513
 1965HO1C Honda, Horie, Kudo and Ui, Nucl. Phys. 62 (1965) 561
 1965HU10 H. Hulubei, M. Ivascu, A. Berinde, I. Neamu, N. Scintei, I. Francz, N. Martalogu and E. Marincu, Rev. Roum. Phys. 10 (1965) 403
 1965HU13 R.W. Huggins and J.H. Sanders, Proc. Phys. Soc. (London) 86 (1965) 53
 1965IS06 Y. Ishizaki, K. Kikuchi, K. Matsuda, T. Mikumo, Y. Nakajima, I. Nonaka, Y. Saji, A. Suzuki, H. Yamaguchi and R. Eisberg, J. Phys. Soc. Jpn. 20 (1965) 2118
 1965JA1A Jacmart, Cahiers Phys. (France) 19 (1965) 1
 1965KA14 S. Kakigi, J. Phys. Soc. Jpn. 20 (1965) 1967
 1965KE02 W.H. Kelly, C.A. Ludemann and C.D. Goodman, Bull. Amer. Phys. Soc. 10 (1965) 121, KD1
 1965KE05 E.F. Kennedy and W.C. Miller, Nucl. Phys. 63 (1965) 634
 1965LE1B Lee and Siemssen, Bull. Amer. Phys. Soc. 10 (1965) 510
 1965MA1E Mazari, Private Communication (1965)

- 1965MA1K Marion, Nucl. Phys. 68 (1965) 463
- 1965MA54 J.H.E. Mattauch, W. Thiele and A.H. Wapstra, Nucl. Phys. 67 (1965) 1
- 1965MA57 S. Mayo, W. Schimmerling, M.J. Sametband and R.M. Eisberg, Nucl. Phys. 62 (1965) 393
- 1965MA65 R.L. Macklin and J.H. Gibbons, Phys. Rev. 140 (1965) B324
- 1965MO1J Mooring and Monohan, Bull. Amer. Phys. Soc. 10 (1965) 513
- 1965NE06 J.O. Newton, R.S. Blake, D.J. Jacobs and J.P. Schapira, Nucl. Phys. 71 (1965) 113
- 1965NE10 Y.A. Nemilov, V.S. Sadkovskii, E.D. Teterin, A.E. Denisov and R.P. Kolalis, Izv. Akad. Nauk SSSR Ser. Fiz. 29 (1965) 1197; Bull. Acad. Sci. USSR Phys.Ser. 29 (1966) 1202
- 1965NE1B Neudachin and Smirnov, At. Energy Rev. 3 (1965) 157
- 1965OL03 J.W. Olness, E.K. Warburton, D.E. Alburger and J.A. Becker, Phys. Rev. 139 (1965) B512
- 1965OV01 J.C. Overley, R.R. Borchers, Nucl. Phys. 65 (1965) 156.
- 1965PA10 J.R. Patterson, J.M. Poate, E.W. Titterton and B.A. Robson, Proc. Phys. Soc. (London) 86 (1965) 1297
- 1965PE17 R.H. Pehl, J. Cerny, E. Rivet and B.G. Harvey, Phys. Rev. 140 (1965) B605
- 1965PU02 H.G. Pugh, D.L. Hendrie, M. Chabre and E. Boschitz, Phys. Rev. Lett. 14 (1965) 434.
- 1965RA1C Rand, Frosch and Yearian, Phys. Rev. Lett. 14 (1965) 234
- 1965RI1A Riou, Rev. Mod. Phys. 37 (1965) 375
- 1965RO07 M.L. Roush, A.A. Jaffe, A.S. Figuera and W.F. Hornyak, Nucl. Phys. 67 (1965) 577.
- 1965RY01 A. Rytz, Nucl. Phys. 70 (1965) 369
- 1965SC04 L.A. Schaller and W.C. Miller, Bull. Amer. Phys. Soc. 10 (1965) 120, KC10
- 1965SC1D Schapira, Ann. Physique 10 (1965) 743
- 1965SI12 R.H. Siemssen, M. Cosack and R. Felst, Nucl. Phys. 69 (1965) 209
- 1965SI13 R.H. Siemssen, M. Cosack and R. Felst, Nucl. Phys. 69 (1965) 227
- 1965ST1E Strobel, Thesis, Univ. of Southern California (1965)
- 1965SZ01 I. Szentpetery, J. Ero, I. Fodor and J. Zimanyi, Yad. Fiz. 1 (1965) 1002; Sov. J. Nucl. Phys. 1 (1965) 714
- 1965TO06 J.H. Towle and B.E.F. Macefield, Nucl. Phys. 66 (1965) 65
- 1965VA1G Van Praet and Kossanyi-Demay, Nuovo Cim. 39 (1965) 388
- 1965VA23 L. Valentin, Nucl. Phys. 62 (1965) 81
- 1965VE1B Verba et al., Bull. Amer. Phys. Soc. 10 (1965) 1194

- 1965WA04 B.D. Walker, C. Wong, J.D. Anderson and J.W. McClure, Phys. Rev. 137 (1964) B1504
- 1965WA06 E.K. Warburton, J.S. Lopes, R.W. Ollerhead, A.R. Poletti and M.F. Thomas, Phys. Rev. 138 (1965) B104
- 1965WY01 S.A.R. Wynchank, A.E. Cox and C.H. Collie, Nucl. Phys. 62 (1965) 491
- 1965YU1B Yuasa and Hourani, Phys. Lett. 18 (1965) 146
- 1965ZI1A Zimanyi, Magyar Fiz. Folyoirat 13 (1965) 461; Phys. Abs. 10804 (1966)
- 1966AD1D Adams et al., Antwerp 1965 Neutron Conf. (North Holland, 1966) 566
- 1966AG1A Agee and Rosen, LA-3538-MS (1966)
- 1966AM1C Ambrosino et al., J. Phys. C1-62 (1966)
- 1966BA1T Balashov and Rotter, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 479
- 1966BA1X Barz et al., Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 222
- 1966BA2R Barz, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK 122 (196) 207
- 1966BA2U Bahnsen, Carpenter and Mendelson, Bull. Amer. Phys. Soc. 11 (1966) 301
- 1966BA2V Barrett, Bull. Amer. Phys. Soc. 11 (1966) 904
- 1966BE1B Berggren and Tyren, Ann. Rev. Nucl. Sci. 16 (1966) 153
- 1966BL01 R.S. Blake, D.J. Jacobs, J.O. Newton and J.P. Schapira, Nucl. Phys. 77 (1966) 254
- 1966BR18 C.P. Browne and F.H. O'Donnell, Phys. Rev. 149 (1966) 767
- 1966CH1K Chabre, Hendrie, Pugh and Detraz, Bull. Amer. Phys. Soc. 11 (1966) 317
- 1966CO1K Cox, Bull. Amer. Phys. Soc. 11 (1966) 654
- 1966CO27 J.G. Couch, J.A. McIntyre and J.C. Hiebert, Phys. Rev. 152 (1966) 883
- 1966CS1B Csikai and Nagy, Acta Phys. Acad. Sci. Hung. 21 (1966) 303
- 1966CU02 R.Y. Cusson, Nucl. Phys. 86 (1966) 481
- 1966DA1H Das, Ali and Sen Gupta, Sci. Res. (Pakistan) 3 (1966) 73; Phys. Abs. 13609 (1967)
- 1966DE1C Detraz et al., Phys. Lett. 22 (1966) 638
- 1966DO11 R.E. Donaldson, D.J. Groves and R.K. Pearson, Phys. Rev. 146 (1966) 660
- 1966DO1A Dolinov and Melikov, Vest. Mosk. Univ. Fiz. Astron. (March-April, 1966) 116
- 1966DO1B Dore, Griffith and Roman, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 60
- 1966EL08 F. El-Batanoni and A.A. Kresnin, Nucl. Phys. 89 (1966) 577
- 1966FR09 J.M. Freeman, J.G. Jenkin and G. Murray, Phys. Lett. 22 (1966) 177

- 1966FR18 A.J. Frasca, R.W. Finlay, R.D. Koshel and R.L. Cassola, Phys. Rev. 144 (1966) 854
- 1966GA04 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 141 (1966) 996
- 1966GA19 A. Gallmann, F. Haas and N. Balaux, Phys. Rev. 151 (1966) 735
- 1966GA25 G.T. Garvey and I. Kelson, Phys. Rev. Lett. 16 (1966) 197
- 1966GE12 L.B. Geesaman, H.B. Knowles and J.B. Mead, Nucl. Phys. 88 (1966) 177
- 1966GO12 S. Gorodetzky, J.P. Coffin, A. Gallmann, G. Frick and F. Jundt, Nucl. Phys. 82 (1966) 113
- 1966GO1N Gorodetzky et al., J. Phys. C1-118 (1966)
- 1966HA18 E.C. Halbert, Y.E. Kim and T.T.S. Kuo, Phys. Lett. 20 (1966) 657
- 1966HA1Q Hahn and Ricci, Bull. Amer. Phys. Soc. 11 (1966) 724
- 1966IS1A Isabelle, J. Phys. C1-17 (1966)
- 1966JA1A Jacob and Maris, Rev. Mod. Phys. 38 (1966) 121
- 1966JA1F O.N. Jarvis, B. Rose and A.E. Taylor, Nucl. Phys. 79 (1966) 275
- 1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KI09 K.G. Kibler, Phys. Rev. 152 (1966) 932
- 1966KL06 A.P. Klyucharev and Y.I. Titov, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 435; Bull. Acad. Sci. USSR Phys. 30 (1966) 443
- 1966KL1E Klyucharev and Titov, Bull. Acad. Sci. USSR 30 (1966) 234
- 1966KL1F Klyucharev, Titov and Vipirailenko, Izv. Vys. Uch. Zaved. Fiz. 3 (1966) 104
- 1966KO08 P. Kossanyi-Demay and G.J. Vanpraet, Nucl. Phys. 81 (1966) 529
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966LI1E K.L. Lim and I.E. McCarthy, Nucl. Phys. 88 (1966) 433
- 1966MA1P MacFarlane, Nucl. Spin-Parity Assignments, Ed. Gove (Academic Press, 1966) 411
- 1966MA2B Mark, Standing and Wu, Bull. Amer. Phys. Soc. 11 (1966) 471
- 1966MA2M Maxson, Murphy and Zatzick, Bull. Amer. Phys. Soc. 11 (1966) 43
- 1966MA2N Maples, Cerny, Fleming and Cospers, Bull. Amer. Phys. Soc. 11 (1966) 903
- 1966MA2P Macklin and Gibbons, Antwerp 1965 Neutron Conf. (North Holland, 1966) 498
- 1966ME03 D.F. Measday, Phys. Lett. 21 (1966) 62
- 1966MI1E Miller, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 410
- 1966MI1F Migdal, Proc. Int. School Enrico Fermi, Course 36; Ed. C. Bloch (Academic Press, 1966) 171

- 1966MO09 F.P. Mooring, J.E. Monahan and C.M. Huddleston, Nucl. Phys. 82 (1966) 16
- 1966NG1B Nguyen-Trung and Sakamoto, Prog. Theor. Phys. 35 (1966) 423
- 1966OE1A Oertzen, Bock and Grosse-Schulte, Z. Naturforsch. A21 (1966) 946
- 1966PA05 V. Parikh, Nucl. Phys. 77 (1966) 157
- 1966PA08 J.A. Panontin, L.L. Schwartz, A.F. Stehney, E.P. Steinberg and L. Winsberg, Phys. Rev. 145 (1966) 754
- 1966PO09 A.M. Poskanzer, S.W. Cospers, E.K. Hyde and J. Cerny, Phys. Rev. Lett. 17 (1966) 1271
- 1966PO1B J.E. Poth, J. Birnbaum and D.A. Bromley, Phys. Rev. Lett. 17 (1966) 200
- 1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, Phys. Rev. 144 (1966) 859; Erratum Phys. Rev. 148 (1966) 1246
- 1966RI12 G. Ripka and L. Zamick, Phys. Lett. 23 (1966) 347
- 1966RI1G Riskalla and Vinciguerra, J. Phys. C1-106 (1966)
- 1966RO11 H.P.C. Rood, Nucl. Phys. 87 (1966) 367
- 1966RO1E Rotter and Zhusupov, Ann. Phys. 17 (1966) 57
- 1966RO1F Rotter, Ann. Phys. 17 (1966) 247
- 1966RO1H Rotter, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf; Ed. J. Schintlmeister, ZFK-122 (1966) 244
- 1966RO1X Rosier, Deschamps and Avignon, Antwerp 1965 Neutron Conf. (North Holland, 1966) 501
- 1966RU01 A.S. Rupaal, B.L. White and J.R. Prescott, Can. J. Phys. 44 (1966) 1213
- 1966SA04 B. Sahai, Phys. Rev. 142 (1966) 612
- 1966SC22 J.J. Schwartz, W.P. Alford, L.M. Blau and D. Cline, Nucl. Phys. 88 (1966) 539
- 1966SE03 R.E. Segel, P.P. Singh, S.S. Hanna and M.A. Grace, Phys. Rev. 145 (1966) 736.
- 1966SE1D Seltz et al., J. Phys. C1-148 (1966)
- 1966SH1A Shute and Brown, Bull. Amer. Phys. Soc. 11 (1966) 471
- 1966SP02 E. Spamer, Z. Physik 191 (1966) 24
- 1966ST12 T. Stovall, J. Goldemberg and D.B. Isabelle, Nucl. Phys. 86 (1966) 225
- 1966SU05 M. Suffert, Nucl. Phys. 75 (1966) 226
- 1966SU1C Suffert, Ann. Phys. (Paris) 1 (1966) 547
- 1966TO01 W.M. Toney and A.W. Waltner, Nucl. Phys. 80 (1966) 237
- 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692

- 1966UN1A Une, Yamaji and Yoshida, Prog. Theor. Phys. 35 (1966) 1010
- 1966WA10 E.K. Warburton, J.W. Olness, K.W. Jones, C. Chasman, R.A. Ristinen and D.H. Wilkinson, Phys. Rev. 148 (1966) 1072
- 1966WA12 N.S. Wall and P.R. Roos, Phys. Rev. 150 (1966) 811; Erratum Phys. Rev. 159 (1967) 1062
- 1966WA1C Warburton and Alburger, Nucl. Spin-Parity Assign., Ed. Gove (Academic Press, 1966) 114
- 1966WE1B Weinberg, Antwerp 1965 Neutron Conf. (Noth Holland, 1966) 37
- 1966WI1E D.H. Wilkinson and M.E. Mafethe, Nucl. Phys. 85 (1966) 97
- 1966WI1F D.H. Wilkinson, Nucl. Phys. 85 (1966) 114
- 1966YU1A Yuasa and Hourany, J. Phys. C1-76 (1966)
- 1966ZI01 B. Ziegler, W. Buss and H. Waffler, Nucl. Phys. 83 (1966) 145
- 1967AM1H Amiet, Ebenhoh and Huguenin, Helv. Phys. Acta 40 (1967) 283
- 1967AR21 M. Arditì, H. Doubre, A. Guichard, M. Riou, D. Royer and C. Ruhla, Contrib. 1967 Int. Conf. Nucl. Struct., Tokyo, Japan (1968) p. 692
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
- 1967BA1E Bang, Zelemskaya, Magzumov and Neudachin, Sov. J. Nucl. Phys. 4 (1967) 688
- 1967BA2J H.W. Barz, Nucl. Phys. A91 (1967) 262
- 1967BA2L Bachelier et al., Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 865
- 1967BA78 G. Backenstoss, S. Charalambus, H. Daniel, H. Koch, G. Poelz, H. Schmitt and L. Tauscher, Phys. Lett. B25 (1967) 547
- 1967BI06 J. Birnbaum, J.C. Overley and D.A. Bromley, Phys. Rev. 157 (1967) 787
- 1967BI1E Bing, Seltz, Gerardin and Magnac-Valette, Private Communication (1967)
- 1967BL02 R.A. Blue, K.J. Stout and G. Marr, Nucl. Phys. A90 (1967) 601
- 1967BL22 R.S. Blake, E.B. Paul, C.H. Sinex and S.T. Emerson, Nucl. Phys. A102 (1967) 305
- 1967BO1D Bodart and Deconninck, Ann. Soc. Sci. Bruxelles 81 (1967) 137
- 1967BR1N Broadhurst and Pyle, Nucl. Instrum. Meth. 48 (1967) 117
- 1967CA09 J.L. Campbell, W. Leiper, K.W.D. Ledingham and R.W.P. Drever, Nucl. Phys. A96 (1967) 279
- 1967CH15 N.S. Chant, P.S. Fisher and D.K. Scott, Nucl. Phys. A99 (1967) 669
- 1967CI1B Cinda (1967)
- 1967CO03 W.R. Coker, M.M. Duncan, J.L. Duggan and P.D. Miller, Nucl. Phys. A91 (1967) 97
- 1967CO1K S.W. Cospèr, J. Cerny and R.C. Gatti, Phys. Rev. 154 (1967) 1193

1967CO1N Cox and Pontet, J. Nucl. Energy 21 (1967) 271

1967CO1T Couch, McIntyre and Hiebert, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 174

1967CO32 S. Cohen and D. Kurath, Nucl. Phys. 101 (1967) 1

1967CR04 M.A. Crosby and J.C. Legg, Nucl. Phys. A95 (1967) 639

1967CU1A Curry, Coker, Lock and Riley, Bull. Amer. Phys. Soc. 12 (1967) 1198

1967DE1K Dehnhard and Siemssen, Bull. Amer. Phys. Soc. 12 (1967) 17

1967DE1V V. Devanathan, K.S. Rao and R. Sridhar, Phys. Lett. B25 (1967) 456

1967DI01 G.U. Din, M.A. Nagarajan and R. Pollard, Nucl. Phys. A93 (1967) 190

1967EL1D El-Nadi, Ismail and Rihan, Nuovo Cim. B50 (1967) 224

1967FA03 M.A. Farouk, M.N.H. Comsan, A.Z. El-Behay, A.A. El-Kamhawy, A.N. Lvov and A.I. Popov, Z. Phys. 201 (1967) 52

1967FA1A W.M. Fairbairn, Nucl. Phys. A90 (1967) 135

1967FI07 W. Fitz, R. Jahr and R. Santo, Nucl. Phys. A101 (1967) 449

1967FI1E Firk, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 352

1967FI1G Fitz, Jahr and Santo, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 170

1967GA06 D.S. Gale and J.M. Blair, Phys. Rev. 161 (1967) 1061

1967GL01 C. Glashauser and M.E. Rickey, Phys. Rev. 154 (1967) 1033

1967GR1L Gray, Fortune, Trost and Fletcher, Bull. Amer. Phys. Soc. 12 (1967) 34

1967GR1M Gruhn and Kelly, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 167

1967GR1N Gray, Fortune, Trost and Fletcher, Bull. Amer. Phys. Soc. 12 (1967) 1197

1967GU1A G. Gunther, K. Bethge and K.R. Schubert, Nucl. Phys. A101 (1967) 288

1967HA21 R.L. Hahn, Nucl. Phys. A101 (1967) 545

1967HO1M Horowitz and Sherman, Bull. Amer. Phys. Soc. 12 (1967) 632

1967JA1E B.K. Jain and D.F. Jackson, Nucl. Phys. A99 (1967) 113

1967KA1A Kabachnik and Grishanova, Sov. J. Nucl. Phys. 4 (1967) 583

1967KI03 K.G. Kibler, Phys. Rev. 155 (1967) 1110

1967LA1N Lane, Elwyn, Mooring and Langsdorf, Bull. Amer. Phys. Soc. 12 (1967) 87

1967LE1E Lehmann, Proc. Int. Conf. Electromag. Int. Dubna, Feb. 1967, Vol. 3 (Moscow, 1967) 66

1967LO1B Loiseaux, Maison and Langevin, J. Phys. 28 (1967) 11

1967ME11 D.F. Measday and J.N. Palmieri, Phys. Rev. 161 (1967) 1071
1967MI1F Miller and Biggerstaff, Bull. Amer. Phys. Soc. 12 (1967) 633
1967MO1N F.B. Morinigo, Nucl. Phys. A90 (1967) 113
1967MO1Q Morrison, Schiffer, Siemssen and Zeidman, Bull. Amer. Phys. Soc. 12 (1967) 501
1967MO21 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. A103 (1967) 238
1967NA06 H. Nakamura, J. Phys. Soc. Jpn. 22 (1967) 685
1967NE1D Nemirovskii, Sov. J. Nucl. Phys. 4 (1967) 334
1967OD01 F.H. O'Donnell and C.P. Browne, Phys. Rev. 158 (1967) 957
1967PO01 R.V. Poore, P.E. Shearin, D.R. Tilley and R.M. Williamson, Nucl. Phys. A92 (1967) 97
1967PO1J A.R. Poletti, E.K. Warburton and D. Kurath, Phys. Rev. 155 (1967) 1096
1967PU01 H.G. Pugh, D.L. Hendrie, M. Chabre, E. Boschitz and I.E. McCarthy, Phys. Rev. 155 (1967) 1054
1967RH1C Rhee and Bray, Bull. Amer. Phys. Soc. 12 (1967) 641
1967RI08 K.E. Richie and B.T. Wright, Phys. Rev. 159 (1967) 839
1967RI1B Ripka, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 833
1967SA07 L.S. Saltykov, Izv. Akad. Nauk SSSR Ser. Fiz. 31 (1967) 323; Bull. Acad. Sci. USSR Phys. Ser. 31 (1968) 307
1967SC1K Schier, Anderson, Bernard and Plattner, Bull. Amer. Phys. Soc. 12 (1967) 33
1967SC43 G.W. Schweimer, Nucl. Phys. A100 (1967) 537
1967SH05 Y.Y. Sharon, Nucl. Phys. A99 (1967) 321
1967SP02 E. Spamer and H. Artus, Z. Phys. 198 (1967) 445
1967ST1H G.L. Strobel, Phys. Rev. 154 (1967) 941
1967ST21 B. Stepanic, R. Popic and M. Aleksic, Phys. Rev. Lett. 19 (1967) 1137
1967THZX M.J. Throop, Bull. Amer. Phys. Soc. 12 (1967) 484, BD13
1967TO1D I.S. Towner, Nucl. Phys. A93 (1967) 145
1967VO1D Volkov, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 85
1967WA1L Watson, Kroepel and Singh, Bull. Amer. Phys. Soc. 12 (1967) 484
1967WI04 J. Wilczynski and V.V. Volkov, Nucl. Phys. A93 (1967) 133
1967WY1B Wyborny and Carlson, Bull. Amer. Phys. Soc. 12 (1967) 1177
1967YO02 F.C. Young, P.D. Forsyth and J.B. Marion, Nucl. Phys. A91 (1967) 209
1967YU02 T. Yuasa and E. Hourany, Nucl. Phys. A103 (1967) 577

- 1967ZE1A Zelenskaya, El-Nagar and Yarba, Joint Inst. Nucl. Res., Lab. Nucl. Problems, Rept. P1 (1967) 3598
- 1968BA2G Backenstoss et al., Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 500
- 1968DI1B Di Napoli et al., Nuovo Cim. B55 (1968) 95
- 1968EA03 L.G. Earwaker and J.H. Montague, Nucl. Phys. A109 (1968) 507
- 1968GA1N Gambarini et al., Proc. 1967 Int. Conf. on Nucl. Struct., Tokyo, Japan (1968)
- 1968GO01 P. Goldhammer, J.R. Hill and J. Nachamkin, Nucl. Phys. A106 (1968) 62
- 1968GR1G Greene, Bull. Amer. Phys. Soc. 13 (1968) 606
- 1968HE1N Herling, Cohen and Silverstein, Proc. 1967 Int. Conf. on Nucl. Struct., Tokyo, Japan (1968)
- 1968HO1F Honda, Mori and Yoshida, Tokyo (1968) 260
- 1968JA1G Jackson, Tokyo (1968) 105
- 1968KU1D Kurath, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan (1967); Suppl. J. Phys. Soc. Jpn. 24 (1968) 393
- 1968MC1C Mc Donald, To be published (1968)
- 1968OK1D Okano, Kikuchi, Nishimura and Harada, Proc. 1967 Int. Conf. on Nucl. Struct., Tokyo, Japan (1968)
- 1968SE1F Seltz et al., Proc. 1967 Int. Conf. on Nucl. Struct., Tokyo, Japan (1968)
- 1968TA1P Taketani, Yamaguchi, Kokame and Muto, Tokyo (1968) 270
- 1968TA1V Taketani, Muto, Yamaguchi and Kokame, Tokyo (1968) 277
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- PO66E Unknown Source

