

Energy Levels of Light Nuclei $A = 11$

F. Ajzenberg-Selove and C.L. Busch

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 11$ –12 was published in *Nuclear Physics A336* (1980), p. 1. This version of $A = 11$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed July 1, 1979)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [EY-76-S-02-2785.*000]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 11$

Below is a list of links for items found within the PDF document. Figures from this evaluation are not included in this PDF document but have been scanned in and are available on this website or via the link below.

A. Nuclides: [¹¹He](#), [¹¹Li](#), [¹¹Be](#), [¹¹B](#), [¹¹C](#), [¹¹N](#), [¹¹O](#), [¹¹F](#), [¹¹Ne](#)

B. Tables of Recommended Level Energies:

[Table 11.1](#): Energy levels of ¹¹Be

[Table 11.3](#): Energy levels of ¹¹B

[Table 11.19](#): Energy levels of ¹¹C

C. [References](#)

D. Figures: [¹¹Be](#), [¹¹B](#), [¹¹C](#), [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

¹¹He
(Not illustrated)

¹¹He has not been observed: see (1976IR1B; theor.).

¹¹Li
(Figs. 1 and 4)

¹¹Li has been observed in the bombardment of iridium by 24 GeV protons. Its mass excess is 40.94 ± 0.08 MeV (1975TH08). The cross section for its formation is ≈ 50 μ b (1976TH1A). ¹¹Li is bound: E_b for break up into ⁹Li + 2n and ¹⁰Li + n are 158 ± 80 and 960 ± 250 keV, respectively [see (1979AJ01) for discussions of the masses of ⁹Li and ¹⁰Li]. The half-life of ¹¹Li is 8.5 ± 0.2 msec (1974RO31): it decays to neutron unstable states of ¹¹Be [$P_n = (60.8 \pm 7.2)\%$: (1974RO31)]. Since systematics suggest $J^\pi = \frac{1}{2}^-$ for ¹¹Li_{g.s.}, the remainder of the decay is presumably to ¹¹Be*(0.32) ($J^\pi = \frac{1}{2}^-$). $\log ft = 5.0$ (1975TH08). (1977BA11; theor.) suggest that the ground state of ¹¹Li has large admixtures of higher configurations and that the mass of ¹¹Li should be used with caution in mass equations. These calculations predict a large fraction of high energy neutrons suggesting that the actual P_n is greater than 61% (1977BA11). See also (1973TO16, 1974HA1H, 1975AJ02) and (1973TA30, 1975BE31, 1975JE02, 1976IR1B; theor.).

¹¹Be
(Figs. 1 and 4)

GENERAL: (See also (1975AJ02).)

Model calculations: (1974TA1E, 1975MI12, 1976IR1B, 1977SE1D, 1978BO31, 1979BE1H).

Special reactions: (1975FE1A, 1976OS04, 1977AR06, 1977YA1B).

Muon capture (See also reaction 2.): (1978DE15).

Pion reactions: (1975CO06, 1976CO1M, 1977DO06, 1977GE1D).

Ground state of ¹¹Be: (1975BE31, 1978BO31).

1. ¹¹Be(β^-)¹¹B $Q_m = 11.508$

The decay proceeds to ¹¹B*(0, 2.12, 5.02, 6.79, 7.98, 9.88) with $J^\pi = \frac{3}{2}^-, \frac{1}{2}^-, \frac{3}{2}^-, \frac{1}{2}^+, \frac{3}{2}^+, \frac{3}{2}^+$, respectively: see Table 11.15 (1971AL07). The half-life is 13.81 ± 0.08 sec (1970AL21). The nature of the decay indicates $J = \frac{1}{2}$ or $\frac{3}{2}$, even parity for the ground state of ¹¹Be: see (1971AL07). See also (1977TE01; theor.).

Table 11.1: Energy Levels of ^{11}Be

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^+; \frac{3}{2}$	$\tau_{1/2} = 13.81 \pm 0.08$ sec	β^-	1, 2, 4
0.3198 ± 0.2	$\frac{1}{2}^-$	$\tau_m = 0.18 \pm 0.06$ psec	γ	2, 4
1.778 ± 12	$(\frac{5}{2}, \frac{3}{2})^+$	$\Gamma = 100 \pm 20$	(n)	2, 4
2.69 ± 20	$(\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+)$	200 ± 20	(n)	2
3.41 ± 20	$(\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+)$	125 ± 20	(n)	2
3.887 ± 15	$\geq \frac{7}{2}$	< 10	(n)	2
3.956 ± 15	$\frac{3}{2}^-$	15 ± 5	(n)	2
5.240 ± 21		45 ± 10	(n)	2
(5.86)		≈ 300	(n)	2
6.51 ± 50		120 ± 50	(n)	2
6.705 ± 21		40 ± 20	(n)	2
7.03 ± 50		300 ± 100	(n)	2
8.816 ± 32		200 ± 50	(n)	2
10.59 ± 50		210 ± 40	(n)	2



Proton groups have been observed to many states of ^{11}Be : see Table 11.2 (1962PU01, 1972AJ01, 1978AJ02). The first excited state has an excitation energy of 319.8 ± 0.2 keV and $\tau_m = 0.18 \pm 0.06$ psec corresponding to a very strong E1 transition of 0.33 W.u. (1971HA25). The J^π of $^{11}\text{Be}^*(0.32)$ is $\frac{1}{2}^-$, as determined by a study of the yield of 320 keV γ -rays as a function of time in μ^- capture by ^{11}B (1968DE20). The strength of the E1 transition fixes J^π of $^{11}\text{Be}_{g.s.}$ to be $\frac{1}{2}^+$ or $\frac{3}{2}^+$, using the parity information obtained from the nature of the β^- decay of the ground state [see reaction 1]. $^{11}\text{Be}^*(5.24, 6.71, 8.82)$ are strongly populated at $E_t = 20$ MeV indicating that these states have a large overlap with $^9\text{Be}_{g.s.} +$ two neutrons (1972AJ01).



See (1975AJ02).



Table 11.2: Levels of ^{11}Be from $^9\text{Be}(t, p)^{11}\text{Be}$

E_x (keV) ^a	E_x (keV) ^b	E_x (keV) ^c	Γ_{lab} (keV) ^a	$\Gamma_{\text{c.m.}}$ (keV) ^b	J^π
0	0	0			$\frac{1}{2}^+$ ^d
319.8 ± 0.2 ^g	322 ± 10	318 ± 10			$\frac{1}{2}^-$ ^d
1780 ± 20	1790 ± 20	1764 ± 20	110 ± 15	130 ± 25	$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$ ^{c,e}
2700 ± 25	2680 ± 30	f	250 ± 20	250 ± 50	$(\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+)$ ^e
3410 ± 25	3410 ± 30	f	150 ± 20	145 ± 30	$(\frac{1}{2}^\pm, \frac{3}{2}^\pm, \frac{5}{2}^+)$ ^e
3890 ± 20	3890 ± 30	3877 ± 30	< 10	≤ 10	$\geq \frac{7}{2}$ ^c
3960 ± 20	3960 ± 30	3943 ± 30	< 10	15 ± 5	$\frac{3}{2}^-$ ^c
	5250 ± 30	5231 ± 30		45 ± 5	
	(5860)	f		≈ 300	
	6510 ± 50	f		120 ± 50	
	6720 ± 30	6690 ± 30		40 ± 20	
	7030 ± 50	f		300 ± 100	
	8840 ± 50	8800 ± 40		200 ± 50	
		10590 ± 50		210 ± 40 ^c	

^a (1962PU01): $E_t = 14$ MeV.

^b (1972AJ01): $E_t = 20$ MeV.

^c (1978AJ02): $E_t = 23$ MeV; angular distributions to $^{11}\text{Be}^*(0, 0.32, 1.78, 3.89, 3.96, 5.24, 6.71, 8.82)$.

^d See text.

^e From angular distributions: see (1962PU01).

^f This E_x region not studied (1978AJ02).

^g (1971HA25).

Angular distributions of the p_0 and p_1 groups have been measured at $E_d = 6$ MeV (1970GO11: p_1 only) and 12 MeV (1970AU02): $l_n = 0$ [and therefore $J^\pi = \frac{1}{2}^+$ for $^{11}\text{Be}(0)$] and 1, $S = 0.73 \pm 0.06$ and 0.63 ± 0.15 , respectively (1970AU02). At $E_d = 25$ MeV $^{11}\text{Be}^*(0, 0.32, 1.78)$ are strongly populated: $S = 0.77, 0.96$ and 0.50 , respectively, $J^\pi = (\frac{5}{2}, \frac{3}{2})^+$ for $^{11}\text{Be}^*(1.78)$ [$l_n = 2$] (1979ZW01).

5. $^{11}\text{Li}(\beta^-)^{11}\text{Be}$

$$Q_m = 20.76$$

See ^{11}Li .

6. $^{11}\text{B}(n, p)^{11}\text{Be}$ $Q_m = -10.726$

See ^{12}B .

7. $^{13}\text{C}(^6\text{Li}, ^8\text{B})^{11}\text{Be}$ $Q_m = -25.886$

At $E(^6\text{Li}) = 80$ MeV, $^{11}\text{Be}^*(0.32)$ is strongly populated and the angular distribution to this state has been measured (1977WE03). $^{11}\text{Be}^*(2.69, 4.0)$ are also observed. It is suggested that these states have odd parity (1976WE09) [see, however, reaction 2 for evidence concerning the positive parity of $^{11}\text{Be}^*(2.69)$]. See also (1977WE1B).

8. $^{14}\text{C}(^{18}\text{O}, ^{21}\text{Ne})^{11}\text{Be}$ $Q_m = -12.206$

See (1974BA15).

¹¹B
(Figs. 2 and 4)

GENERAL: (See also (1975AJ02).)

Shell model: (1977BO07, 1977JA14, 1977TE01, 1978BO31).

Cluster, collective and rotational models: (1976BR26, 1977BO07, 1977NI1A, 1977OK1C, 1979NI06).

Special levels: (1976IR1B, 1977BO07, 1977JA14, 1977TE01, 1977TE02, 1978BO31, 1979NI06).

Electromagnetic transitions: (1974HA1C, 1977BO07, 1977DO06, 1977OK1C, 1977TE01, 1977TE02, 1978KI08, 1979NI06).

Astrophysical questions: (1975AR1E, 1975HA1N, 1975ME1E, 1975PR1B, 1975TR1A, 1976AU1B, 1976AU1C, 1976BO1E, 1976CO1B, 1976EP1A, 1976GI1C, 1976HA1F, 1976RO1J, 1976SI1C, 1976VIIA, 1977AU1B, 1977DW1A, 1977HA1L, 1977KO1L, 1977MA1H, 1977SC1D, 1978AU1C, 1978BO1K, 1978BU1B, 1978BY1A, 1978DI04, 1978DI1D, 1978DW1A, 1978DW1B, 1978ME1D, 1979MC1A).

Special reactions: (1975AB1D, 1975AR14, 1975HU14, 1975RA21, 1976BE1K, 1976BU16, 1976HE1H, 1976LE1F, 1976NA11, 1976OS04, 1976RA1C, 1976RO12, 1976YO1A, 1977AR06, 1977FO04, 1977GE08, 1977GO07, 1977KU1D, 1977RE08, 1977SH1D, 1977ST1G, 1977UD1A, 1977YA1B, 1978BH03, 1978BI08, 1978GE1C, 1978GR1F, 1978HE1C, 1978KO01, 1978LE15, 1978PE1E, 1978TU06, 1978VO1D, 1978WE1D, 1978ZE02, 1979CR03, 1979DY01, 1979GA04, 1979GO11, 1979PA09, 1979SC08, 1979VI05, 1979WE06).

Muon and neutrino capture and reactions: (1974EN10, 1974WA1C, 1975DO1F, 1975DO1D, 1976DO1G, 1977CA1E, 1977DO06, 1977MU1A, 1977WA1G, 1978DE15, 1978LE04, 1979BE1N, 1979DE01).

Pion capture and reactions (See also reactions 27, 38 and 54.): (1975HU1D, 1975ZI1B, 1976BA1R, 1976CA1H, 1976KA1F, 1976MI15, 1976NO1D, 1977BA2H, 1977BA1Q, 1977BA2G, 1977BE1L, 1977DO06, 1977HO1B, 1977KO25, 1977MA35, 1977PI1D, 1977SI01, 1978CO02, 1978KI08, 1978KI13, 1978MO01, 1978SI1D, 1978TS1A, 1979BE1N, 1979BO1U, 1979EL1E, 1979HU02, 1979TI1A, 1979ZI1C).

Reactions and kaons and other mesons: (1975TA1C, 1976DE1D, 1978AT01, 1978DA1A).

Applied topics: (1975AU1C, 1979AN1L, 1979GR1E, 1979JA1F, 1979JU1B).

Ground state of ¹¹B: (1974DE1E, 1974EN10, 1974SHYR, 1975AL19, 1975BE31, 1975EP02, 1976BR26, 1976DO1G, 1976FU06, 1976PH01, 1977AN21, 1977MA35, 1977PA25, 1977YO1D, 1978AN07, 1978HE1D, 1978ZA1D, 1979NI06).

$$\mu = +2.68837 (2) (1978LEZA);$$

$Q = 40.65 (26)$ mb [(1970NE05; theor.): adopted by (1978LEZA); see also (1968SC18, 1969SC34)].

Table 11.3: Energy Levels of ^{11}B ^a

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	stable		1, 2, 8, 9, 10, 14, 15, 16, 17, 18, 19, 20, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74
2.1248 ± 0.4	$\frac{1}{2}^-$	$\Gamma = 0.128 \pm 0.010$ eV	γ	1, 8, 9, 10, 14, 15, 16, 17, 19, 20, 27, 28, 29, 36, 37, 39, 40, 41, 44, 45, 49, 53, 55, 57, 58, 59, 65, 66, 67, 69, 70, 72, 73
4.4451 ± 0.5	$\frac{5}{2}^-$	0.61 ± 0.04 eV	γ	1, 2, 8, 9, 10, 14, 15, 16, 20, 27, 28, 29, 31, 32, 37, 39, 40, 41, 44, 45, 49, 53, 54, 55, 57, 58, 59, 66, 67, 69, 70, 72, 73
5.0208 ± 0.5	$\frac{3}{2}^-$	2.16 ± 0.10 eV	γ	1, 8, 9, 10, 15, 16, 27, 28, 29, 31, 36, 37, 39, 40, 41, 44, 45, 53, 57, 58, 59, 66, 67, 69, 70, 73
6.7429 ± 0.6	$\frac{7}{2}^-$	0.043 ± 0.007 eV	γ	1, 2, 8, 15, 20, 28, 29, 31, 32, 40, 41, 44, 45, 53, 58, 66, 67, 70, 72
6.7929 ± 1.0	$\frac{1}{2}^+$	0.47 ± 0.06 eV	γ	1, 8, 15, 28, 29, 36, 40, 41, 45, 49, 57, 58, 67, 70, 73

Table 11.3: Energy Levels of ^{11}B ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
7.2856 ± 1.5	$(\frac{3}{2}, \frac{5}{2})^+$	1.46 ± 0.20 eV ^b	γ	1, 8, 14, 15, 28, 29, 39, 41, 45, 58, 73
7.9780 ± 1.3	$\frac{3}{2}^+$	1.46 ± 0.22 eV	γ	1, 15, 28, 36, 41
8.5594 ± 1.9	$\leq \frac{5}{2}^-$	1.04 ± 0.16 eV ^c	γ	1, 14, 15, 28, 39, 41, 45, 67
8.920 ± 1.5	$\frac{5}{2}^-$	4.78 ± 0.29 eV	γ, α	1, 2, 14, 15, 20, 28, 31, 39, 41, 66, 67
9.185 ± 1.5	$\frac{7}{2}^+$	0.3 ± 0.1 eV	γ, α	1, 2, 12, 28, 41, 70
9.274 ± 1.5	$\frac{5}{2}^+$	7 keV	γ, α	1, 2, 15, 28, 41, 70
9.875 ± 10	$\frac{3}{2}^+$	130 ± 30	α	7, 15, 36
10.26 ± 20	$\frac{3}{2}^{(-)}, \frac{1}{2}$	150 ± 40	γ, α	2, 7
10.33 ± 20	$(\frac{5}{2}, \frac{7}{2})^-$	70 ± 20	γ, α	2, 7, 28
(10.45 ± 50)		≈ 140	γ, α	2, 41
10.601 ± 10	$\frac{7}{2}^+$	70 ± 10	γ, α	2, 7
10.96 ± 50	$\frac{5}{2}^-$	4500	α	7
11.29 ± 30	$\frac{9}{2}^+$	90 ± 50	α	7
11.49 ± 50			α	7
11.61 ± 50	$\frac{5}{2}^+$	150 ± 30	n, α	3, 7, 21, 26, 27, 41
11.8	$\frac{7}{2}^+$	880	n, α	26
11.90 ± 20	$\frac{5}{2}^-$	150 ± 30	n, α	3, 7, 26
(12.18 ± 40)			γ, p	19
12.56 ± 30	$\frac{1}{2}^+(\frac{3}{2}^+); T = \frac{3}{2}$	240 ± 50	γ, p, α	7, 17, 19, 44
12.91 ± 20	$\frac{1}{2}^-; T = \frac{3}{2}$	240 ± 30	γ, p	19, 41, 44, 66
13.03 ± 60		270 ± 50	α	7
13.12	$\frac{9}{2}^-$	425	n, α	3, 21, 26
13.16	$\frac{5}{2}^+, \frac{7}{2}^+$	430	n, α	3, 21, 26
14.04 ± 100	$\frac{11}{2}^+$	500 ± 200	n, α	3, 7, 21, 26
14.33 ± 20	$\frac{5}{2}^{(+)}, (\frac{3}{2}^-); T = \frac{3}{2}$	250 ± 40	γ, p	19, 44
14.53 ± 50		< 120	$\text{n}, \text{t}, \alpha$	3, 25, 26, 44
15.32 ± 100	$(\frac{3}{2}, \frac{5}{2}, \frac{7}{2})^+; T = \frac{3}{2}$	635 ± 180	γ, p	19, 21, 26, 41
15.8 ± 200			n, α	3, 7, 22
(16.43 ± 15)		≈ 40	$\text{p}, \text{d}, \alpha$	12, 41

Table 11.3: Energy Levels of ^{11}B ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
(16.7 \pm 300)		broad	n, t, α	3, 25, 26, 41
17.33		\approx 1000	n, p, d, t, α	12, 25, 26
17.50 \pm 50		180 \pm 40	γ , n, p, d, α	3, 10, 12
18.37 \pm 50	$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$	260 \pm 80	γ , d	10
(19.6)	$(\frac{1}{2}^+)$	broad	γ , d	10, 38, 57
23.7	$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$		γ , d	10
26.5		broad	γ , n	38

^a See also Tables 11.4, 11.5 and 11.16.

^b Assuming $J^\pi = \frac{5}{2}^+$: see Table 11.16.

^c Assuming $J^\pi = \frac{3}{2}^-$: see Table 11.16.

1. $^6\text{Li}(^6\text{Li}, \text{p})^{11}\text{B}$ $Q_m = 12.218$

Angular distributions have been measured for proton groups to states with $E_x < 9.3$ MeV at $E(^6\text{Li}) = 2.4$ to 9.0 MeV (1966KI09). For γ -spectra and for τ_m see Tables 11.4 and 11.5. See also (1975AJ02) and (1976EP1A; astrophys.).

2. $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$ $Q_m = 8.665$

Resonances for capture radiation are displayed in Table 11.6. Results of studies of angular correlations, distributions and branching ratios are displayed in Tables 11.6 and 11.4: see (1968AJ02) for a discussion of the results. See also (1978SC1D; applied).

3. $^7\text{Li}(\alpha, \text{n})^{10}\text{B}$ $Q_m = -2.790$ $E_b = 8.665$

Table 11.7 displays the thresholds and resonances observed in this reaction. See also (1976SE1E), (1974LO1B, 1975AJ02, 1977LI19).

4. $^7\text{Li}(\alpha, \text{p})^{10}\text{Be}$ $Q_m = -2.5635$ $E_b = 8.665$

See ^{10}Be in (1979AJ01).

Table 11.4: Electromagnetic transitions in ^{11}B ^a

Initial state	$J^\pi; T$	Γ_γ (total) ^b (eV)	Branching ratios (%) to final states					Refs. ^a	
			g.s. $\frac{3^-}{2}; \frac{1}{2}$	2.12 $\frac{1^-}{2}; \frac{1}{2}$	4.45 $\frac{5^-}{2}; \frac{1}{2}$	5.02 $\frac{3^-}{2}; \frac{1}{2}$	6.74 $\frac{7^-}{2}; \frac{1}{2}$		6.79 $(\frac{1}{2}, \frac{3}{2})^+; \frac{1}{2}$
2.12	$\frac{1^-}{2}; \frac{1}{2}$	0.128 ± 0.010	100						
4.45	$\frac{5^-}{2}; \frac{1}{2}$	0.61 ± 0.04	100 ^f	< 0.5					
5.02	$\frac{3^-}{2}; \frac{1}{2}$	2.16 ± 0.10	85 ± 2	15 ± 2	0.3				
			88 ± 2	12 ± 2					(1968EA03)
			88 ± 2.5 ^g	12 ± 2.5 ^h					(1968BE30)
			83 ± 5	17 ± 5					(1971AL07)
6.74	$\frac{7^-}{2}; \frac{1}{2}$	0.043 ± 0.007	70 ± 2 ^c	< 3	30 ± 2	< 1			
6.79	$\frac{1^+}{2}; \frac{1}{2}$	0.47 ± 0.06	66.0 ± 1.5	30.0 ± 1.5	< 0.5	4.1 ± 0.9			(1971AL07)
7.29	$(\frac{3}{2}, \frac{5}{2})^+; \frac{1}{2}$	1.46 ± 0.20 ⁱ	87 ± 2	< 1	5.5 ± 1	7.5 ± 1			
7.98	$\frac{3^+}{2}; \frac{1}{2}$	1.46 ± 0.22	47 ± 2	53 ± 2	< 1	< 1			
			45 ± 2	55 ± 2	< 1.6	< 1.6	< 1.5	< 1.4	(1971AL07)
8.56	$\frac{5^-}{2}; \frac{1}{2}$	^b	56 ± 2	30 ± 2	5 ± 1	9 ± 1			
8.92	$\frac{5^-}{2}; \frac{1}{2}$	4.78 ± 0.29	95 ± 1 ^d	< 1	4.5 ± 0.5	< 1	< 1	< 1	
9.19	$\frac{7^+}{2}; \frac{1}{2}$	0.3 ± 0.1 ^a	0.9 ± 0.3		82.8 ± 2.0		12.8 ± 0.4	< 1.3	(1962GR07)
9.27	$\frac{5^+}{2}; \frac{1}{2}$		19.7 ± 1.0		67.5 ± 2.0		12.8 ± 0.7	< 0.6	(1962GR07)
^e									

^a (1965OL03) except where shown. See also Table 11.4 in (1975AJ02).

^b From Table 11.16, corrected for branching to other states.

^c $\delta = -0.45 \pm 0.18$ (1968CO09). This value leads to too large a value of Γ_γ for an M3 transition (P.M. Endt, private communication).

^d $\delta = -0.11 \pm 0.04$ (1968CO09).

^e See Tables 11.6, 11.11 and 11.16 for higher states.

^f $\delta = -0.19 \pm 0.03$ (1968BE30).

^g $\delta = 0.03 \pm 0.05$ (1968BE30).

^h $\delta = -0.05 \pm 0.2$ (1968BE30).

ⁱ Assuming $J^\pi = \frac{5}{2}^+$.

Comments [mainly from (1962GR07, 1965OL03)]:

(1) 4.45 MeV. $9.28 \rightarrow 4.45 \rightarrow 0$ angular distribution fixes $J = \frac{5}{2}$. Odd parity determined from direct interaction assignments.

(2) 5.02 MeV. Internal pair correlation permit M1, E2 for the g.s. transition: $J^\pi \leq \frac{7}{2}^-$ (parity from l -assignments). τ_m excludes $\frac{7}{2}$, branch to 2.12, $\frac{5}{2}$. Angular correlation (1968BE30) fixes $\frac{5}{2}^-$.

(3) 6.74 MeV. Internal pairs indicate practically pure E2 g.s. radiation. Angular distributions and branching ratios (and l -assignments) all lead to $\frac{7}{2}^-$.

(4) 6.79 MeV. The allowed β -decay from ^{11}Be indicates $J^\pi \leq \frac{7}{2}^+$. The relatively strong γ -branch to $^{11}\text{B}^*(2.12)$ favors $\frac{1}{2}^+$, $\frac{3}{2}^+$. (1968EA03) finds that all γ 's from this level are isotropic, suggesting $J^\pi = \frac{1}{2}^+$, but not excluding $\frac{3}{2}^+$.

(5) 7.29 MeV. The g.s. transition is mainly E1, so $J^\pi \leq \frac{5}{2}^+$. The assignment $\frac{1}{2}^+$ is excluded by the strength of (7.29 \rightarrow 4.45).

(6) 7.98 MeV. Transitions to $^{11}\text{B}_{(\text{g.s.})}$ and (2.12) are predominantly E1; thus $^{11}\text{B}^*(7.98)$ has even parity, and odd parity of $^{11}\text{B}^*(2.12)$ is confirmed. The transition to $^{11}\text{B}^*(2.12)$ is not isotropic, so $J^\pi = \frac{3}{2}^+$.

(7) 8.56 MeV. Correlation of internal pairs indicate the the g.s. transition is M1 + E2 or E1 + M2, $J^\pi \leq \frac{5}{2}^+$ or $\leq \frac{7}{2}^-$; the lifetime to $^{11}\text{B}^*(2.12)$ excludes $\frac{7}{2}^-$. If the level has even parity, the required M2 admixture is excessive. $J^\pi \leq \frac{5}{2}^-$ is favored.

(8) 8.92 MeV. From $^7\text{Li}(\alpha, \gamma)^{11}\text{B}$, $J^\pi = \frac{3}{2}^+$, $\frac{5}{2}^+$, $\frac{5}{2}^-$. The internal pair correlation confirms $\frac{5}{2}^-$. For higher states see comments under individual reactions and (1968AJ02).

Table 11.5: Lifetimes of some ^{11}B states ^a

^{11}B state	τ_m (fsec)	Refs.
2.12	5.1 ± 0.4	Table 11.4
4.45	1.08 ± 0.07	Table 11.4
5.02	0.31 ± 0.02	Table 11.4
6.74	15 ± 3	(1979MO1M)
6.79	1.4 ± 0.2	(1979MO1M)
7.29	0.45 ± 0.06	Table 11.4
7.98	0.45 ± 0.09	(1979MO1M)
8.56	1.04 ± 0.16	(1979MO1M)
8.92	0.14 ± 0.01	Table 11.4
9.19	0.3 ± 0.1	Table 11.4, (1965OL03)

^a See also Table 11.5 in (1975AJ02).

5. ${}^7\text{Li}(\alpha, d){}^9\text{Be}$ $Q_m = -7.1507$ $E_b = 8.665$

See (1976LE1K) and ${}^9\text{Be}$ in (1979AJ01).

6. ${}^7\text{Li}(\alpha, t){}^8\text{Be}$ $Q_m = -2.559$ $E_b = 8.665$

Excitation functions have been measured for $E_\alpha = 14$ to 25 MeV (t_0) and 18 to 25 MeV (t_1) (1973VA1A, 1974DM01). See also (1976LE1K) and ${}^8\text{Be}$ in (1979AJ01).

7. ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$ $E_b = 8.665$

The elastic scattering and the scattering to ${}^7\text{Li}^*(0.48)$ have been studied at many energies to $E_\alpha = 22.5$ MeV: see (1968AJ02, 1975AJ02). Observed resonances are displayed in Table 11.8. See also (1968AJ02, 1975AJ02, 1975VO1B), (1977BE51, 1977NI1A; theor.) and ${}^7\text{Li}$ in (1979AJ01).

8. ${}^7\text{Li}({}^6\text{Li}, d){}^{11}\text{B}$ $Q_m = 7.192$

Table 11.6: Resonances in ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ ^a

E_r (keV)	$\Gamma_{\text{c.m.}}$ (keV)	${}^{11}\text{B}^*$ (MeV)	J^π	$\omega\gamma_\alpha$ ^d (eV)	Γ_{γ_0} ^c (eV)	Percentage decay to ${}^{11}\text{B}^*$ ^f			
						0	4.45	6.74	6.79
401 ^b	< 1 ^b	8.920	$(\frac{5}{2})^-$	0.04	i	93 ± 5	2.3 ± 1		
819 ± 1 ^b	3×10^{-3} ^g	9.186	$\frac{7}{2}^+$	0.55		0.9 ± 0.3	82.8 ± 2.0	12.8 ± 0.4	< 1.3
958 ± 1 ^b	7 ^b	9.275	$\frac{5}{2}^+$	3.5		19.7 ± 1.0	67.5 ± 2.0	12.8 ± 0.7	< 0.6 ^k
2500 ± 20 ^c	433	10.26			17	h			
2620 ± 20 ^c	100	10.33			1.0	h			
2800 ± 50 ^c	≈ 140 ⁱ	10.45			$10/2J + 1$				
(3040) ^c	90	(10.60)			< 0.2	h			

^a See also Tables 11.4, 11.8, 11.9 and 11.16 and Table 11.4 in (1968AJ02).

^b (1951BE13).

^c (1967PA19).

^d $\omega\Gamma_\gamma\Gamma_\alpha/\Gamma$, in c.m. (1951BE13, 1959JO25).

^e (1967PA19); based on analysis using $R = 4.9$ fm, $\gamma_w^2 = 1.0$ MeV.

^f (1962GR07). See also Table 11.4.

^g (1965OL03).

^h $< 10\%$ to ${}^{11}\text{B}^*(2.12)$ (1967PA19).

ⁱ Observed width (1967PA19).

^j See Table 11.16.

^k The decay to ${}^{11}\text{B}^*(7.29, 7.98)$ [$J^\pi = \frac{5}{2}^+, \frac{3}{2}^+$] is also observed: $\approx 1\%$ and $\approx 0.03\%$ respectively (1977MO1K).

Table 11.7: Thresholds and resonances in ${}^7\text{Li}(\alpha, n){}^{10}\text{B}$

(1959GI47)	(1963ME08)	
E_α (MeV \pm keV)	E_α (MeV \pm keV)	E_x (MeV \pm keV)
	4.380 \pm 20	thresh.
	[4.72]	11.67 \pm 100
5.15 \pm 70 ^a	[5.22] ^c	11.99 \pm 100
(5.64)	5.5	thresh.
7.15 ^b	7.05 ^c	13.15 \pm 100
	[8.44]	14.04 \pm 100
	[9.21]	14.53 \pm 50
	10.14	15.12 \pm 100
	[11.33]	(15.88 \pm 200)
	11.90	thresh.
	12.56	(16.7 \pm 300)
	13.92	17.52 \pm 30
	14.53	thresh.

^a $J^\pi = \frac{3}{2}^-$ or $\frac{5}{2}^+$, $\Gamma_n \approx 20$ keV, $\Gamma_\alpha \approx 300$ keV formed by $l_n = 0$ or 1 (1959GI47) [comparison with ${}^{10}\text{B}(n, \alpha)$], $\Gamma_{\text{lab}} = 220$ keV (1957BI84).

^b The width of this resonance is large.

^c The n_0 yield shows the resonances at $E_\alpha \approx 5.2$ and 7.05 MeV: no others are seen in the interval $4.5 < E_\alpha < 8$ MeV (1972VA02).

Angular distributions have been measured for $E({}^7\text{Li}) = 3.3$ to 5.95 MeV: see (1975AJ02). For γ -spectra and τ_m see Tables 11.4 and 11.5. See also ${}^{13}\text{C}$ in (1981AJ01).

9. ${}^7\text{Li}({}^7\text{Li}, t){}^{11}\text{B}$ $Q_m = 6.199$

Angular distributions have been measured at $E({}^7\text{Li}) = 2.10$ to 5.75 MeV (1969CA1A: $t_0 \rightarrow t_3$). See also (1976CA1J). At $E({}^7\text{Li}) = 79.6$ MeV transitions are observed to several ${}^{11}\text{B}$ states. ${}^{11}\text{B}_{\text{g.s.}}$ is particularly strongly populated (1974CE06).

10. ${}^9\text{Be}(d, \gamma){}^{11}\text{B}$ $Q_m = 15.8159$

Table 11.8: Structure in ${}^7\text{Li}(\alpha, \alpha){}^7\text{Li}$ and ${}^7\text{Li}(\alpha, \alpha'){}^7\text{Li}$ ^a

E_α ^b (keV)	E_α ^c (keV)	$\Gamma_{\text{c.m.}}$ (keV)	E_x (MeV \pm keV)	J^π
1900 \pm 10 ^d		130 \pm 30	9.875 \pm 10	$\frac{3}{2}^+$
2480 \pm 50		150 \pm 40	10.24 \pm 50	$\frac{3}{2}^{(-)}, \frac{1}{2}$
	2630 \pm 30	80 \pm 30	10.34 \pm 30	$\frac{5}{2}^-, \frac{7}{2}^-$
3040 \pm 10 ^d	3040	70 \pm 10	10.601 \pm 10	$\frac{7}{2}^+$
3600 \pm 50		4500	10.96 \pm 50	$\frac{5}{2}^-$
	4120 \pm 30	90 \pm 50	11.29 \pm 30	$\frac{9}{2}^+$
4430 \pm 50	4430		11.49 \pm 50	
4600 \pm 50		150 \pm 50	11.59 \pm 50	
5050 \pm 30		150 \pm 50	11.88 \pm 30	
	5300 \pm 200	\approx 1000	12.0 \pm 200	
	5500 \pm 100	60 \pm 50	(12.17 \pm 100) ^e	
6100 \pm 30		150 \pm 50	12.55 \pm 30	
6850 \pm 60		270 \pm 50	13.03 \pm 60	
(7200 \pm 50) ^f		50 \pm 50	(13.25 \pm 50) ^e	
	7800 \pm 100	500 \pm 200	(13.63 \pm 100) ^e	
(8450 \pm 200) ^g		500 \pm 200	(14.0 \pm 200)	
(9450 \pm 200) ^g		\leq 250	(14.7 \pm 200)	
	9950 \pm 20	500 \pm 200	(15.00 \pm 20) ^e	
(11200 \pm 200) ^g			(15.8 \pm 200)	
h				

^a (1966CU02), except where shown. See also Table 11.9 in (1975AJ02).

^b ${}^7\text{Li}(\alpha, \alpha'\gamma){}^7\text{Li}$: $\sigma(\text{total})$.

^c ${}^7\text{Li}(\alpha, \alpha_0){}^7\text{Li}$.

^d (1967PA19). Other values are listed in Table 11.8 of (1975AJ02).

^e ${}^7\text{Li}(\alpha, n){}^{10}\text{B}$ threshold.

^f Anomaly in angular distribution.

^g Observed at $\theta = 60^\circ$.

^h For possible higher structures see (1971BI12, 1973KE13).

The 90° differential cross section has measured for $E_d = 0.5$ to 11.9 MeV: see (1975AJ02). The behavior of the γ_0 , γ_1 and γ_{2+3} total cross sections of the angular distributions of these γ -rays indicate two resonances at $E_d = 1.98 \pm 0.05$ and 3.12 ± 0.05 MeV with $\Gamma_{\text{lab}} = 225 \pm 50$ and 320 ± 100 keV, corresponding to $^{11}\text{B}^*(17.43, 18.37)$. The higher resonance was not observable in the $\gamma_2 + \gamma_3$ cross section which was not measured beyond $E_d = 2.5$ MeV. The maximum γ_0 cross section observed is $10.1 \pm 3.5 \mu\text{b}$ at $E_d \approx 0.96$ MeV (1971BA72). (1974DE39) observe resonant behavior in the $90^\circ \gamma_0$ cross section at $E_d \approx 3.4$ and 9.65 MeV ($^{11}\text{B}^*(18.6, 23.7)$) in addition to a wide structure at 4.7 MeV ($^{11}\text{B}^*(19.7)$). The angular distributions of γ_0 from $^{11}\text{B}^*(18.6, 23.7)$ are typical of E1 transitions. The (d, γ_0) reaction appears to proceed via excitation of the $T = \frac{1}{2}$ component of the giant dipole resonance in ^{11}B . See also (1974LO1B).

11. $^9\text{Be}(d, n)^{10}\text{B}$

$$Q_m = 4.3607$$

$$E_b = 15.8159$$

The cross section follows the Gamow function for $E_d = 70$ to 110 keV (1955RA14). The fast neutron and γ -yield rise smoothly to $E_d = 1.8$ MeV except for a possible “resonance” at $E_d \approx 0.94$ MeV. The fast neutron yield then remains approximately constant to 3 MeV: see (1968AJ02) for references. The excitation functions for $n_0 \rightarrow n_4$, and n to $^{10}\text{B}^*(5.1, 6.57)$ have been measured for $E_d = 14$ to 16 MeV: no strong fluctuations are observed (1973PA14). Thick target yields are reported at $E_d = 14.8, 18.0$ and 23.0 MeV (1977LO10). Polarization measurements have been carried out at $E_d = 0.9$ to 5.5 MeV [see (1975AJ02)] and at 0.4 MeV (1977BA2K; n_0, n_1). See also (1975SA1C), (1977GR1F, 1977SA1K, 1978BA1F, 1978HA1M, 1979CH11; applied) and ^{10}B in (1979AJ01).

12. (a) $^9\text{Be}(d, p)^{10}\text{Be}$

$$Q_m = 4.5872$$

$$E_b = 15.8159$$

(b) $^9\text{Be}(d, \alpha)^7\text{Li}$

$$Q_m = 7.151$$

(c) $^9\text{Be}(d, t)^8\text{Be}$

$$Q_m = 4.5921$$

Measurements of proton yields have been carried out up to $E_d = 6.0$ MeV for p_0 and p_1 [see (1975AJ02)] and at 0.9 to 3.1 MeV (1975ZW01: σ_t, p_0, p_1). The p_0 and p_1 yields show a resonance at $E_d = 750 \pm 15$ keV [$^{11}\text{B}^*(16.43)$], $\Gamma \approx 40$ keV (1974AN01) and the p_1 yield resonates at 1.85 MeV [$^{11}\text{B}^*(17.33)$], $\Gamma_{\text{c.m.}} \approx 1.0$ MeV (1974FR02, 1975ZW01) and 2.3 MeV (1974FR02) [sharp state]. See also (1975AJ02) for other possible structures. Polarization of the protons have been measured at a number of energies for $E_d = 1$ to 21 MeV [see (1975AJ02)] and at $E_d = 12$ MeV (1977BA39: TAP; p_0, p_1) and 15 MeV (1976DA15: VAP; $p_0, p_1, p_{2+3}, p_{4+5}, p_6, p_7, p_9$). See also ^{10}Be in (1979AJ01) and (1974BO35, 1974BO37, 1975BO20, 1975BO21; theor.).

The yields of α -particles [α_0, α_1], reaction (b), have been measured for $E_d = 0.3$ to 12.4 MeV [see (1975AJ02)] and at $E_d = 2.25$ to 3.10 MeV (1977SL02: α_0, α_1) and 12.17 to 14.43 MeV (1978TA04: $\alpha_0, \alpha_1, \alpha_2, \alpha_4$). (1974AN01) report a weak indication of the 0.75 MeV resonance, observed in the proton yield, in the α_0 yield. See also ^7Li in (1979AJ01).

The cross section for reaction (c) has been measured for $E_d = 0.15$ to 19 MeV [see (1968AJ02, 1975AJ02)] and at 0.9 to 2.5 MeV (1975BO1C; t_0), 0.9 to 3.1 MeV (1975ZW01; t_0) and 12.17 to 14.43 MeV (1978TA04; t_0, t_1). Polarization studies have been carried out at $E_d = 12$ MeV (1977BA39: TAP; t_0, t_1) and 15 MeV (1976DA15; VAP; t_0, t_1). There is no clear evidence of resonance structure. See also ^8Be in (1979AJ01). See also (1974CH1L).

13. $^9\text{Be}(d, d)^9\text{Be}$

$$E_b = 15.8159$$

Excitation functions for elastically scattered deuterons have been measured for $E_d = 0.4$ to 7.0 MeV [see (1975AJ02)] and for 12.17 to 14.43 MeV (1978TA12; also d_1, d_2). Polarization measurements involving the d_0 group have been reported at $E_d = 6.3$ to 12.6 MeV [see (1975AJ02)] and at $E_d = 15$ MeV (1976DA15: TAP; d to $^9\text{Be}^*(0, 2.43, 3.06)$). See also ^9Be in (1979AJ01).

14. $^9\text{Be}(t, n)^{11}\text{B}$

$$Q_m = 9.5586$$

Angular distributions have been measured at $E_t = 1.1$ to 1.7 MeV (1977MA1N; $n_0, n_1, n_2, n_6, n_8, n_9$ [at 1.3 MeV]).

15. $^9\text{Be}(^3\text{He}, p)^{11}\text{B}$

$$Q_m = 10.3224$$

Proton groups have been observed to a number of ^{11}B states: see Table 11.9. Angular distributions of many of these proton groups have been studied at $E(^3\text{He}) = 1.0$ to 10.2 MeV [see (1968AJ02, 1975AJ02)] and at 3 to 6 MeV (1977LI1F) and 14 MeV (1977IR01; p_0, p_1). Gamma-ray branching ratios and multiplicities are displayed in Table 11.4: see the discussion in (1968AJ02). See also (1966YO1A) and (1976EP1A; astrophys.).

16. $^9\text{Be}(\alpha, d)^{11}\text{B}$

$$Q_m = -8.0308$$

Angular distributions have been measured at $E_\alpha = 23.4, 24.7, 25.2$ MeV (1975VA19; d_0, d_1 (at two higher energies only)), 27.0 MeV (1975PU01; $d_0 \rightarrow d_6$) and 28.3 MeV (1965KA14; $d_0 \rightarrow d_3$). The predominant L -transfers are $L = 0, 2; 0; 0$ for $^{11}\text{B}^*(0, 2.12, 5.02)$. The angular distribution for $^{11}\text{B}^*(4.45)$ is flat (1975PU01). See also (1976LE1K) and (1978ZE03; theor.).

17. (a) $^9\text{Be}(^6\text{Li}, \alpha)^{11}\text{B}$

$$Q_m = 14.342$$

(b) $^9\text{Be}(^6\text{Li}, 2\alpha)^7\text{Li}$

$$Q_m = 5.677$$

(c) $^9\text{Be}(^6\text{Li}, \alpha p)^{10}\text{Be}$

$$Q_m = 3.114$$

(d) $^9\text{Be}(^6\text{Li}, \alpha t)^8\text{Be}$

$$Q_m = 3.119$$

Table 11.9: Energy levels of ^{11}B from $^9\text{Be}(^3\text{He}, \text{p})^{11}\text{B}$

(1959HI69) ^a	(1966BR18)	(1960HI08)
E_x (MeV \pm keV)		L
0	0	0
2.130 ± 10	2.1243 ± 0.9	0
4.445^b	4.4434 ± 1.8	0
5.023 ± 10	5.0187 ± 2.3	0
6.739 ± 10	6.7411 ± 3.0	
6.791 ± 10	6.7909 ± 3.1	1
7.285 ± 10		
7.975 ± 10		
8.553 ± 10		0
8.909 ± 10		0
9.175 ± 10		
9.264 ± 10		
9.86 ± 20^c		

^a The original results were normalized to the second excited state taken to be at 4.459 MeV. Here they are shown normalized to 4.445 MeV.

^b (1966BR18).

^c $\Gamma \approx 150$ keV.

Angular distributions have been determined for seven α -groups at $E(^6\text{Li}) = 3$ to 4 MeV: see (1968AJ02). Angular distributions have also been obtained at $E(^6\text{Li}) = 24$ MeV to $^{11}\text{B}^*(0, 2.12)$ and to a number of unresolved levels with $E_x \leq 13.2$ MeV (1968DA20, 1967CH34). See also (1977EL10; theor.).

The breakup reactions have been studied at $E(^6\text{Li}) = 3.5$ MeV by (1968JA08): reaction (b) goes mainly via a sequential process involving $^{11}\text{B}^*(10.3, 11.4, 12.6, 13.16, 13.5)$. The results for reaction (c) are not conclusive. $^{11}\text{B}^*(12.6, 13.16)$ may possibly contribute to reaction (d) (1968JA08).

18. (a) $^9\text{Be}(^7\text{Li}, \alpha\text{n})^{11}\text{B}$ $Q_m = 7.092$
 (b) $^9\text{Be}(^9\text{Be}, ^7\text{Li})^{11}\text{B}$ $Q_m = -0.880$

See (1975AJ02).

Table 11.10: Levels of ^{11}B from the $^{10}\text{Be}(p, \gamma)^{11}\text{B}$ reaction (1970GO04)

E_p (MeV \pm keV)	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	$(J + \frac{1}{2})(\Gamma_p/\Gamma)\Gamma_{\gamma_0}$ ^e (eV)	Γ_{γ_0} ^{a,e} (eV)	$\Gamma_{\gamma_1}/\Gamma_{\gamma_0}$	J^π
(1.05 ± 40) ^b	(12.18)	230 ± 90	$3.1^{+2.9}_{-2.0}$			
1.46 ± 30	12.56	230 ± 65	10^{+7}_{-5}	10^{+7}_{-5}	0.25 ± 0.08	$\frac{1}{2}^+(\frac{3}{2}^+)$
1.85 ± 20	12.91	235 ± 27	29 ± 9	29 ± 9 ^f	≤ 0.06	$\frac{1}{2}^-$ ^c
3.41 ± 20	14.33	255 ± 36	29 ± 9	14.5 ± 4.3	≤ 0.1	$\frac{5}{2}^+(\frac{3}{2}^-)$
4.5 ± 100	15.32	635 ± 180	53^{+34}_{-26} ^d			

^a These values assume that $J \neq \frac{3}{2}$: see (1970GO04).

^b May be due to $^{10}\text{B}^*(0.7) + n$ threshold.

^c See Table 11.3.

^d Assumes that $\sigma_{\text{total}} = 4\pi d\sigma/d\Omega$ (90°).

^e Values reported in (1970GO04) are here shown multiplied by 1.7: see (1973GO09).

^f In the (e, e') work of (1975KA02) a strong group is observed at $E_x = 13.0 \pm 0.1$ MeV. If it corresponds to the excitation of $^{11}\text{B}^*(12.91)$ with $J^\pi = \frac{1}{2}^-$; $T = \frac{3}{2}$, $\Gamma_{\gamma_0} = 36 \pm 7$ eV (1975KA02).

19. $^{10}\text{Be}(p, \gamma)^{11}\text{B}$

$$Q_m = 11.2287$$

The yield of ground state γ -rays has been measured at 90° for $E_p = 0.6$ to 6.3 MeV. Observed resonances are displayed in Table 11.10. The four resonances correspond to states in ^{11}B whose energies match well with those of the first four states in ^{11}Be : $T = \frac{3}{2}$. Several known $T = \frac{1}{2}$ states in ^{11}B are not observed in this reaction: see Table 11.3 (1970GO04, 1973GO09). See also (1974LO1B).

20. $^{10}\text{B}(n, \gamma)^{11}\text{B}$

$$Q_m = 11.4552$$

The thermal capture cross section is 0.5 ± 0.2 b (1973MU14). For a listing of the observed capture γ -rays see Table 11.12 (1967TH05). For τ_m see Table 11.4. The yield of γ -rays at $E_n = 14$ MeV shows a smaller fore-aft anisotropy than is observed in proton capture, suggesting that the forward peaking in the (p, γ) reaction is due mainly to direct rather than collective capture amplitudes (1975AR19). See also (1976MA1J).

21. $^{10}\text{B}(n, n)^{10}\text{B}$

$$E_b = 11.4552$$

Table 11.11: Neutron capture γ -rays from $^{10}\text{B} + \text{n}$ ^a

E_γ (keV)	ΔE_x (keV)	I_γ ^b	Assignment
11447 ± 2 ^c	11453 ± 2	6 ± 1	capt. \rightarrow g.s.
8916 ± 2	8920 ± 2	15 ± 2	$8.92 \rightarrow$ g.s.
7006 ± 2	7008 ± 2	54 ± 3	capt. \rightarrow 4.45
6739 ± 2	6741 ± 2	19 ± 1	$6.74 \rightarrow$ g.s.
	5020	< 2	$5.02 \rightarrow$ g.s.
4711 ± 2	4712 ± 2	25 ± 1	capt. \rightarrow 6.74
4444 ± 2	4445 ± 2	65 ± 3	$4.45 \rightarrow$ g.s.
2534 ± 2	2534 ± 2	15 ± 2	capt. \rightarrow 8.92
2295 ± 2	2296 ± 2	10 ± 3	$6.74 \rightarrow$ 4.45
	2120	< 3	$2.12 \rightarrow$ g.s.

^a (1967TH05).

^b Photons/100 captures.

^c $\Gamma_\gamma = 0.01$ eV (1957BA18).

The “free” neutron scattering cross section, $\bar{\sigma}_s = 2.23 \pm 0.06$ b. The coherent scattering amplitude (bound) is $a = +1.4 \pm 0.5$ fm (1973MU14). The total scattering cross section is constant at 2.23 ± 0.06 b for $E_n = 0.7$ to 10 keV and then rises to 2.97 b at $E_n = 127$ keV (1970AS10).

Total cross section measurements in the range $E_n = 10$ to 500 keV confirm the broad maximum near $E_n = 0.23$ MeV, originally suggested by (1951BO45) and also observed in the (n, α) cross section (1966MO09). At higher energies the total cross section shows broad maxima at $E_n = 1.9$ and 2.8 MeV (1951BO45) and at 4.3 MeV (1961FO07): see Table 11.12. In the range $E_n = 5.5$ to 16 MeV σ_{tot} is constant at 1.5 b (1961FO07, 1979AU07). (1976GAYV) displays reported σ_{tot} measurements.

Polarization measurements (0.075 to 2.2 MeV and 2.63 MeV) and measurements of differential cross sections (0.075 to 4.4 MeV) by (1971LA10, 1973CO05, 1973HA64) have been analyzed using R -matrix calculations: the results are shown in Table 11.13. They are consistent with results from $^{10}\text{B}(\text{n}, \text{n}'\gamma)$ and $^7\text{Li}(\alpha, \text{n})$.

Elastic differential cross sections have also been reported at $E_n = 7.02$ to 9.72 MeV [see (1975AJ02)] and at 4 to 8 MeV (1978KN01; n_0, n_1) and 14.1 MeV (1974HY01). See also ^{10}B in (1979AJ01).

22. $^{10}\text{B}(\text{n}, \text{n}')^{10}\text{B}^*$

$E_b = 11.4552$

Table 11.12: Resonances in $^{10}\text{B} + \text{n}$ ^a

$^{10}\text{B}(\text{n}, \text{n}'\gamma)^{10}\text{B}$ (1960DA08)		$^{10}\text{B}(\text{n}, \alpha)^7\text{Li}$ (1961DA16)		Yield of	$^{11}\text{B}^*$ (MeV)
E_{res} (MeV)	Γ (keV)	E_{res} (MeV)	Γ (keV)		
		0.23 ^b		$\sigma_{\text{t}}, \alpha$	11.66
		0.53 ^f	140	α_0, α_1	11.94
1.93	260	1.86 ^e	570	$\sigma_{\text{t}}^{\text{c}}, \alpha_0, \alpha_1, \text{t}, \text{n}'$	13.2
(2.6)	broad	2.79	530	$\sigma_{\text{t}}^{\text{c}}, \alpha_0, \alpha_1, \text{n}'$	14.0
3.31	370	3.43	< 120	$\alpha_0, \text{t}, \text{n}'$	14.57
4.1		4.1	800	$\sigma_{\text{t}}^{\text{c}}, \alpha_0, \alpha_1, \text{n}'$	15.2
4.73		^d		n'	15.75
		5.7	broad	α_0, t	16.6
		6.4	broad	α_0, t	17.3

^a See also Table 11.13.

^b (1951BO45, 1966MO09, 1970NE03).

^c (1951BO45). See also (1979AU07).

^d (1961FO07).

^e $J^\pi \geq \frac{11}{2}^+$ (1961DA16). See, however, Table 11.13.

^f See footnote ^c in Table 11.13.

The yield of 0.7 MeV γ -rays has been studied from threshold to $E_{\text{n}} = 5.2$ MeV: observed resonances are displayed in Table 11.12 (1960DA08). Inelastic scattering cross sections for formation of various ^{10}B states have been measured at a number of energies in the range $E_{\text{n}} = 1.45$ to 14.8 MeV: see (1975AJ02). See also ^{10}B in (1979AJ01).

$$23. \ ^{10}\text{B}(\text{n}, \text{p})^{10}\text{Be} \qquad Q_{\text{m}} = 0.2265 \qquad E_{\text{b}} = 11.4552$$

The thermal cross section is < 0.2 b (1973MU14). See also (1976SL2A).

$$24. \ ^{10}\text{B}(\text{n}, \text{d})^9\text{Be} \qquad Q_{\text{m}} = -4.3607 \qquad E_{\text{b}} = 11.4552$$

See ^9Be in (1979AJ01) and (1976SL2A).

$$25. \ (a) \ ^{10}\text{B}(\text{n}, \text{t})^8\text{Be} \qquad Q_{\text{m}} = 0.2314 \qquad E_{\text{b}} = 11.4552$$

$$(b) \ ^{10}\text{B}(\text{n}, \text{t})^4\text{He}^4\text{He} \qquad Q_{\text{m}} = 0.3233$$

Table 11.13: R -matrix analysis of resonant states in $^{10}\text{B} + \text{n}$ ^a

E_n (MeV)	E_x (MeV)	J^π	l_n	Γ_n	Γ_{α_0}	Γ_{α_1}	$\Gamma_{\text{c.m.}}$ (keV)
	10.60	$\frac{7}{2}^+$	0		0.030	0.070	100
0.17	11.61	$\frac{5}{2}^+$	0	0.004	0.296	0.0	300
0.37 ^b	11.79	$\frac{7}{2}^+$	0	0.770	0.001	0.113	884
0.53 ^c	11.94	$\frac{5}{2}^-$	1	0.031	0.080	0.090	201
1.83	13.12	$\frac{9}{2}^-$	1	0.100	0.275	0.050	425
1.88	13.16	$\frac{5}{2}^+, \frac{7}{2}^+$	2	0.080	0.200	0.150	430
2.82	14.02	$\frac{11}{2}^+$	2	0.800	0.045	0.010	855
4.2	15.3	$(\frac{3}{2}, \frac{3}{2}, \frac{7}{2})^+$	2	0.500	0.100	0.100	700

^a (1971LA10, 1972HA04, 1973CO05, 1973HA64): analysis based on polarization and differential cross-section measurements of the elastic scattering, and on results from $^{10}\text{B}(n, \alpha_0)$ and (n, α_1) . The analysis used a two-level, four-channel R -matrix formalism with a non-diagonal background R^0 matrix: see (1973HA64). This analysis does not include $^{11}\text{B}^*(14.53)$ because the resonance is weak, narrow and almost entirely in the α -channel (1973CO05). See also Table 11.12.

^b See also (1976SE06).

^c (1978SC31) report $E_{\text{res}} = 495 \pm 5$ keV, $\Gamma = 140 \pm 15$ keV, $\sigma_{\text{max}} [\text{in } (n, \alpha_1\gamma)] = 94 \pm 6$ mb.

The cross section for reaction (b) has been measured for $E_n = 1.4$ to 8.2 MeV by (1961DA16). Fluctuations are observed at some of the resonant energies in the $^{10}\text{B}(n, \alpha)$ reaction: see Table 11.12. For a display of the cross section see (1976GAYV). See also (1975BI07) and (1976LI1H, 1976SL2A).

26. $^{10}\text{B}(n, \alpha)^7\text{Li}$

$$Q_m = 2.790$$

$$E_b = 11.4552$$

The “recommended” value of the thermal isotropic absorption cross section is 3837 ± 9 b (1973MU14). The ground state branching for thermal neutrons is $(6.308 \pm 0.006)\%$ (1967DE15), $(6.699 \pm 0.012)\%$ (1979ST03). At $E_n = 2$ and 24 keV the values are $(7.05 \pm 0.16)\%$ and $(7.13 \pm 0.15)\%$, respectively (1979ST03). The cross section for $E_n = 10$ eV to 200 keV is given by the expression

$$13.837/\sqrt{E} = 0.312 - 1.014 \times 10^{-2}\sqrt{E} + \frac{2.809 \times 10^5}{\sqrt{E}[(170.3-E)^2 + 2.243 \times 10^4]} \text{ b (1970SO1A)}.$$

See also (1974ST1F).

The total cross section for the (n, α) reaction is displayed in (1976GAYV). The cross section has been measured for $E_n = 1.03$ to 300 keV (1974CO1C; $\alpha_1\gamma$; no evidence for resonances), 4.70 to 630.64 keV (1978SC31; $\alpha_1\gamma$), 0.15 to 1.25 MeV (1975SE1H, 1976SE06; α_0, α_1 ; $0^\circ, 120^\circ$; evidence for broad resonance at $E_n \approx 0.4$ MeV in the $\alpha_1(0^\circ)$ cross section) and at higher energies up to 14.8 MeV: see (1975AJ02) and Tables 11.12 and 11.13. At $E_n = 0.79$ MeV, the ratio for the $\alpha_1/\alpha_{\text{tot}}$ yield is 0.66 ± 0.03 (1975LA08). See ${}^7\text{Li}$ in (1979AJ01), (1975AJ02, 1976LI1H, 1976SL2A), (1975FR1C), (1976BO1L, 1976DA1F; applied) and (1975HA1L, 1977LO1C; theor.).

$$27. {}^{10}\text{B}(p, \pi^+){}^{11}\text{B} \quad Q_m = -128.894$$

Angular distributions for proton capture to ${}^{11}\text{B}^*(0, 2.15 \pm 0.10)$ have been measured at $E_p = 168$ to 186 MeV. The population of ${}^{11}\text{B}^*(4.45 + 5.02)$ and of unresolved higher states is also observed at 186 MeV but an attempt to observe $T = \frac{3}{2}$ analogue states in ${}^{11}\text{B}$ and ${}^{11}\text{N}$ (the latter populated via (p, π^-)) was unsuccessful (1974DA27, 1979HO09). Polarization measurements are reported by (1979PI1E). See also (1977BE1X, 1979BE1W, 1979JO1C, 1979PI06, 1979SO1C) and (1979BO1V; theor.).

$$28. {}^{10}\text{B}(d, p){}^{11}\text{B} \quad Q_m = 9.2306$$

Reported proton groups are displayed in Table 11.14. Angular distributions have been studied at many energies in the range $E_d = 0.17$ to 28 MeV [see (1968AJ02, 1975AJ02)] and at 0.75 to 2.5 MeV (1977AR12; $p_0 \rightarrow p_3$). The lowest five levels are formed by $l_n = 1$ except for ${}^{11}\text{B}^*(2.12)$ which appears to involve a spin-flip process. They are presumed to comprise the set $\frac{3}{2}^-, \frac{1}{2}^-, \frac{5}{2}^-, \frac{3}{2}^-, \frac{7}{2}^-$ expected as the lowest p^7 levels ($a/K \approx 4.0$). ${}^{11}\text{B}^*(9.19, 9.27)$ [$J^\pi = \frac{7}{2}^+, \frac{5}{2}^+$] show strong $l = 0$ stripping and are ascribed to capture of a 2s neutron by ${}^{10}\text{B}$: see (1968AJ02) for a listing of all the relevant references. Studies of $p\gamma$ correlations are discussed in reaction 14 of (1968AJ02) and displayed in Table 11.4 of this paper. See also ${}^{12}\text{C}$ and (1975ZA06).

$$29. {}^{10}\text{B}(t, d){}^{11}\text{B} \quad Q_m = 5.1979$$

At $E_t = 5.5$ MeV, deuteron groups are observed to the ground state of ${}^{11}\text{B}$ and to states at $E_x = 2.126, 4.449, 5.027, 6.769, 6.806$ and 7.301 MeV (± 10 keV). All the angular distributions appear to be characteristics of $l_n = 1$ (1961BA10). See also ${}^{13}\text{C}$ in (1981AJ01).

$$30. {}^{10}\text{B}(\alpha, {}^3\text{He}){}^{11}\text{B} \quad Q_m = -9.1226$$

Table 11.14: ^{11}B levels from $^{10}\text{B}(\text{d}, \text{p})^{11}\text{B}$

E_x (MeV \pm keV) ^a	(1962HI07)		(1960BI08)	
	l_n	$(2J + 1)\theta^2$	l_n	$(2J + 1)\theta^2$
0	1	0.120	1	1.00
2.1246 \pm 1.1				0.09
4.4458 \pm 2.1	1	0.048		0.46
5.0192 \pm 2.4	(1)	(0.010)		0.11
6.7439 \pm 2.2	1	0.210	1	1.72
6.7938 \pm 2.2				
7.298 \pm 6	(2?)	(0.022)		
7.987 \pm 9			isotropic	
8.568 \pm 5	(2?)		2	
8.927 \pm 5	1	0.186	0, 2	
9.191 \pm 5	0	0.242	0	
9.276 \pm 5	0	0.175	0	
10.32 \pm 20 ^b				

^a (1951VA1A, 1953EL12, 1966BR18). See also Table 11.15 in (1975AJ02) and (1977AR12).

^b $\Gamma = 54 \pm 17$ keV (1953EL12).

Angular distributions have been measured at $E_\alpha = 56$ MeV for the ground state transitions in this, and in the analog (α , t) reactions: the average ratio of the (α , ^3He) to the (α , t) differential cross sections is 1.2 ± 0.1 (1969GA11).

31. (a) $^{10}\text{B}(^6\text{Li}, ^5\text{Li})^{11}\text{B}$ $Q_m = 5.79$
 (b) $^{10}\text{B}(^7\text{Li}, ^6\text{Li})^{11}\text{B}$ $Q_m = 4.205$

At $E(^7\text{Li}) = 24$ MeV angular distributions are reported to $^{11}\text{B}^*(0, 6.74, 8.92)$. $^{11}\text{B}^*(4.45, 5.02)$ are also populated. The (^7Li , ^6He) reaction [see reaction 16 in ^{11}C] has also been studied (1977KO27). See also (1975AJ02).

32. $^{10}\text{B}(^{13}\text{C}, ^{12}\text{C})^{11}\text{B}$ $Q_m = 6.5088$

Total cross section measurements involving $^{11}\text{B}^*(4.45, 6.74)$ have been carried out for $E(^{13}\text{C}) = 8.0$ to 12.7 MeV: $S_1S_2 = 0.6$ and 1.7 , respectively (1975SE03). τ_m for $^{11}\text{B}^*(6.74) < 50$ fsec (1975SE04): see Table 11.5.

$$33. \ ^{10}\text{B}(^{14}\text{N}, ^{13}\text{N})^{11}\text{B} \quad Q_m = 0.902$$

See (1975NA15, 1977MO1A). See also ^{13}N in (1976AJ04, 1981AJ01), (1975AJ02) and (1975OS01; theor.).

$$34. \ ^{10}\text{B}(^{16}\text{O}, ^{15}\text{O})^{11}\text{B} \quad Q_m = -4.209$$

See (1975NA15), ^{15}O in (1976AJ04) and (1977OK1D; theor.).

$$35. \ (a) \ ^{10}\text{B}(^{18}\text{O}, ^{17}\text{O})^{11}\text{B} \quad Q_m = 3.411$$

$$(b) \ ^{10}\text{B}(^{19}\text{F}, ^{18}\text{F})^{11}\text{B} \quad Q_m = 1.024$$

Angular distributions have been measured at $E(^{18}\text{O})$ and $E(^{19}\text{F}) = 20$ and 24 MeV (1971KN05). See also (1975AJ02).

$$36. \ ^{11}\text{Be}(\beta^-)^{11}\text{B} \quad Q_m = 11.508$$

^{11}Be decays to $^{11}\text{B}^*(0, 2.12, 5.02, 6.79, 7.98, 9.88)$: see Table 11.15 for the parameters of the observed β and γ transitions (1971AL07). Delayed α -particles are also observed with a total I_α of 3.0%/decay. These α -particles are not observed to be in coincidence with 478 keV γ -rays [upper limit = 5% for $E_\alpha > 0.3$ MeV], suggesting that they result from the decay of $^{11}\text{B}^*(9.88)$ to $^7\text{Li}_{g.s.}$ (1971AL07).

$$37. \ ^{11}\text{B}(\gamma, \gamma)^{11}\text{B}$$

Mean gamma widths of low-lying states obtained by resonance scattering and transmission studies are listed in Table 11.16 (1978KU12, 1979MO1M). The excitation energies of $^{11}\text{B}^*(2.13, 4.45, 5.02)$ are 2126.5 ± 1.5 , 4446.4 ± 1.8 and $5.021.1 \pm 1.8$ keV (1977WE1C). See also (1975BR1F, 1979MO1L).

Table 11.15: Beta decay of ^{11}Be (1971AL07) ^a

$^{11}\text{B}^*$ ^b (keV)	J^π ^c	I_β (%)	$\log ft$	E_γ (keV)	I_γ ^d (%)	Transition to $^{11}\text{B}^*$
g.s.	$\frac{3}{2}^-$	57 ± 3	6.81 ± 0.02			
2125.0 ± 0.7	$\frac{1}{2}^-$	29 ± 3	6.68 ± 0.04	2124.8 ± 0.7	33 ± 3	g.s.
4445	$\frac{5}{2}^-$	< 0.06	> 10.9 ^e			
5020.1 ± 1.7	$\frac{3}{2}^-$	0.28 ± 0.11	7.94 ± 0.14	5019.3 ± 1.7	0.47 ± 0.09	g.s.
				2893.1 ± 0.8	0.093 ± 0.028	2125
6742.7 ± 1.8 ^g	$\frac{7}{2}^-$	< 0.08				
6792.6 ± 1.8	$\frac{1}{2}^+$	6.8 ± 0.8	5.91 ± 0.05	6790.5 ± 1.8	4.51 ± 0.69	g.s.
				4666.3 ± 1.8	2.00 ± 0.28	2125
				1772.2 ± 0.7	0.28 ± 0.06	5020
7286	$(\frac{3}{2}, \frac{5}{2})^+$	< 0.16				
7978.1 ± 1.9	$\frac{3}{2}^+$	3.9 ± 0.5	5.58 ± 0.05	7974.7 ± 1.9	1.74 ± 0.30	g.s.
				5851.8 ± 1.9	2.13 ± 0.34	2125
8559	$\leq \frac{5}{2}^-$	< 0.06	> 7.0			
8920	$\frac{5}{2}^-$	< 0.02	> 8.5 ^e			
9875	$\frac{3}{2}^+$	3.0 ± 0.7 ^f	4.03 ± 0.15 ^f			

^a See also Table 11.12 in (1968AJ02).

^b When errors are indicated the excitation energies are determined in this experiment from the measured E_γ .

^c From Table 11.3.

^d Intensity in % per β -decay, normalized to $(33 \pm 3)\%$ for the 2.13 MeV γ -intensity.

^e $\log f_1 t$. Q_0 assumed to be 11.506 ± 0.007 MeV.

^f Assuming that the breakup of $^{11}\text{B}^*(9.88)$ is solely to $^7\text{Li}_{\text{g.s.}}$. If the inelasticity for the breakup of $^{11}\text{B}^*(9.88)$ is that suggested by (1966CU02), then the β -branch is $15 \pm 3.5\%$, $\log ft = 3.33 \pm 0.15$, and the β branches to the other ^{11}B states have to be recalculated: see (1971AL07).

^g Energy derived from E_x of $^{11}\text{B}^*(6.79)$ and known ΔE of 4th and 5th excited states (1970BR23) [49.9 keV].

38. (a) $^{11}\text{B}(\gamma, \pi^-)^{11}\text{C}$	$Q_m = -141.549$
(b) $^{11}\text{B}(\gamma, n)^{10}\text{B}$	$Q_m = -11.4552$
(c) $^{11}\text{B}(\gamma, p)^{10}\text{Be}$	$Q_m = -11.2287$
(d) $^{11}\text{B}(\gamma, d)^9\text{Be}$	$Q_m = -15.8159$
(e) $^{11}\text{B}(\gamma, t)^4\text{He} + ^4\text{He}$	$Q_m = -11.1319$
(f) $^{11}\text{B}(\gamma, \alpha)^7\text{Li}$	$Q_m = -8.665$

For reaction (a) see (1975YE1A) and reaction 20 in ^{11}C . The giant dipole resonance is shown to consist mainly of $T = \frac{1}{2}$ states in the lower energy region and of $T = \frac{3}{2}$ states in the higher energy region by observing the decay to states in ^{10}B and ^{10}Be [reactions (b) and (c)] (1971PA10). See also (1979BR1D). Absolute measurements of the $^{11}\text{B}(\gamma, \text{all n})$ cross section from threshold to 35 MeV have been carried out by (1973HU09, 1976KN04): the cross section exhibits a main peak

at $E_\gamma = 25$ to 28 MeV and weak shoulders at 13 and 16 MeV. The integrated cross section to 35 MeV is 69.1 ± 0.8 MeV · mb (1976KN04). See also (1975AD04) and (1974BU1A, 1976BE1H). For other structures reported in the (γ, n) and (γ, p) cross sections see (1975AJ02). The (γ, d_0) cross section peaks at ≈ 19 MeV, lower than it would if $T = \frac{3}{2}$ states were involved (1974DE01). The yield of 3.37 MeV γ -rays [from $^{10}\text{Be}^*(3.37)$, reaction (c)] has been measured for $E_{\text{bs}} = 100$ to 800 MeV (1975AD04). See also (1977BR1J). Double positron-electron pair production by 6.6 MeV γ -rays in boron has been studied by (1976RO1N). See (1959AJ76) for reactions (e) and (f). See also (1975AJ02) and (1978DI1A).

Table 11.16: Gamma widths ^a from $^{11}\text{B}(\gamma, \gamma)^{11}\text{B}$ and $^{11}\text{B}(e, e)^{11}\text{B}$

E_x (MeV)	J^π	Γ_{γ_0} (eV)	Reaction	Refs.
2.12	$\frac{1}{2}^-$	0.137 ± 0.020	$\gamma\gamma$	(1965KE05)
		0.12 ± 0.02	$\gamma\gamma$	(1968CR07)
		0.11 ± 0.02	$\gamma\gamma$	(1978KU12)
		0.125 ± 0.04	$\gamma\gamma$	(1979MO1M)
		0.14 ± 0.04	ee	(1975KA02)
		0.17 ± 0.034	ee	(1962ED02)
4.45	$\frac{5}{2}^-$	0.128 ± 0.010		mean
		0.73 ± 0.07 (M1)	ee	(1975KA02)
		$+0.020 \pm 0.002$ (E2)		
		0.60 ± 0.09 (M1)	ee	(1967SP02)
		$+0.016 \pm 0.002$ (E2)		
		0.58 ± 0.04	$\gamma\gamma$	(1978KU12)
5.02	$\frac{3}{2}^-$	0.58 ± 0.04	$\gamma\gamma$	(1979MO1M)
		0.61 ± 0.04		mean
		1.73 ± 0.14 (M1)	ee	(1967SP02)
		< 0.0034 (E2)		
		2.12 ± 0.21	ee	(1975KA02)
		1.80 ± 0.13	$\gamma\gamma$	(1978KU12)
6.74	$\frac{7}{2}^-$	1.88 ± 0.17	$\gamma\gamma$	(1979MO1M)
		1.84 ± 0.07		mean
		0.030 ± 0.005	$\gamma\gamma$	(1979MO1M)
		0.31 ± 0.04	$\gamma\gamma$	(1979MO1M)
		1.17 ± 0.26	$\gamma\gamma$	(1978KU12)
		1.34 ± 0.22 ^b	$\gamma\gamma$	(1979MO1M)
6.79	$\frac{1}{2}^+$			
7.29	$(\frac{3}{2}, \frac{5}{2})^+$			

Table 11.16: Gamma widths ^a from ¹¹B(γ, γ)¹¹B and ¹¹B(e, e)¹¹B (continued)

E_x (MeV)	J^π	Γ_{γ_0} (eV)	Reaction	Refs.
7.98	$\frac{3}{2}^+$	1.27 ± 0.17	$\gamma\gamma$	mean
		0.67 ± 0.10		(1979MO1M)
8.56	$\leq \frac{5}{2}^-$	0.73 ± 0.07 (M1)	ee	(1975KA02)
		$+0.23 \pm 0.03$ (E2)		
8.92	$\frac{5}{2}^-$	0.58 ± 0.09 ^c	$\gamma\gamma$	(1979MO1M)
		4.0 ± 0.6 (M1)	ee	(1966SP02)
		4.93 ± 0.50	ee	(1975KA02)
		4.20 ± 0.52	$\gamma\gamma$	(1978KU12)
		5.00 ± 0.60	$\gamma\gamma$	(1979MO1M)
		4.54 ± 0.28		mean

^a See also Tables 11.4 and 11.5, and Table 11.17 in (1975AJ02).

^b Assuming $J^\pi = \frac{5}{2}^+$.

^c Assuming $J^\pi = \frac{3}{2}^-$.

39. (a) ¹¹B(e, e)¹¹B

(b) ¹¹B(e, ep)¹⁰Be $Q_m = -11.2287$

The charge-scattering radius leads to an oscillator length parameter of 1.55 fm (1959ME24). Magnetic elastic scattering at $\theta = 180^\circ$ shows strong M3 effects: the derived ratio of static M3/M1, 2.9 ± 0.2 fm², suggests a $j - j$ coupling scheme for ¹¹B_{g.s.} (1966RA29). The quadrupole contribution to the elastic form factor is best accounted for by the undeformed shell model, $Q = 3.72 (\pm 20\%)$ fm², $r(\text{r.m.s.}) = 2.42$ fm (1966IS1A, 1966ST12). See also (1979BE1N, muonic X-rays): $\langle r^2 \rangle^{1/2} = 2.47 \pm 0.04$ fm. The excitation of ¹¹B*(2.1, 4.4, 5.0, 8.6, 8.9) has been studied by (1975KA02: $E_e = 52$ to 90 MeV). The giant resonance region, centered at ≈ 18 MeV, is characterized by a lack of prominent features except for a pronounced peak at $E_x = 13.0 \pm 0.1$ MeV (mixed M1–E2) and a broad transverse group at $E_x = 15.5$ MeV (1975KA02).

Ground state transition widths are listed in Table 11.16. For reaction (b) see (1975AJ02). See also (1974DE1E) and (1974WA1C, 1975DO1D, 1976DO1G, 1977TE02, 1977WA1G, 1978BO09, 1978TA1D; theor.).

40. ¹¹B(n, n')¹¹B*

Angular distributions of neutrons have been measured for $E_n = 7.55, 9.72$ and 14.1 MeV [see (1975AJ02)] and for $E_n = 4$ to 8 MeV (1976LA1C, 1976WH1A, 1977WH1A, 1977WH1B; n_0), $5.46, 5.62$ and 6.00 MeV (1979KO1Q; n_1), 8 to 14 MeV (1978GL1D; $n_0 \rightarrow n_3$) and 14.1 MeV (1974HY01; $n_0 \rightarrow n_3, n_{4+5}$). See also ^{12}B .

41. (a) $^{11}\text{B}(p, p')^{11}\text{B}^*$

(b) $^{11}\text{B}(p, 2p)^{10}\text{Be}$ $Q_m = -11.2287$

(c) $^{11}\text{B}(p, pn)^{10}\text{B}$ $Q_m = -11.4552$

Observed proton groups are displayed in Table 11.17. Angular distributions have been measured for $E_p = 12$ to 185 MeV [see (1975AJ02)] and at $E_p = 6$ MeV (1977SA1B; p_0) and 30 MeV (1975CA08: to states shown in column 1 of Table 11.17, except $^{11}\text{B}^*(8.92)$). See also (1976AL1G, 1977CO1G). The analysis of the $\frac{3}{2}^- \rightarrow \frac{1}{2}^-$ transition [$^{11}\text{B}^*(0 \rightarrow 2.12)$] at $E_p = 32$ MeV shows less spin flip than predicted by DWBA (1977MO09). For reaction (b) see ^{10}Be in (1979AJ01). For reaction (c) see (1977WA05). See also (1978AL1G) and (1976PH01, 1977PH02; theor.).

42. $^{11}\text{B}(d, d)^{11}\text{B}$

The elastic scattering has been studied at $E_d = 5.5$ MeV (1971HIZF) and 11.8 MeV (1967FI07). See also (1974CH58; theor.).

43. $^{11}\text{B}(t, t)^{11}\text{B}$

The elastic scattering has been studied at $E_t = 1.8$ and 2.1 MeV (1969HE08, 1969SI12).

44. $^{11}\text{B}(^3\text{He}, ^3\text{He})^{11}\text{B}$

The elastic scattering has been studied has been studied at $E(^3\text{He}) = 8$ to 74 MeV [see (1975AJ02)], at 17.5 and 40.0 MeV (1977SH09) and at 46.1 MeV (1979GO07). In the recent work angular distributions have also been studied for the ^3He ions to $^{11}\text{B}^*(2.12, 4.45, 5.02, 6.74)$ (1977SH09). $T = \frac{3}{2}$ states populated in this reaction are displayed in Table 11.18. There is a weak indication also of a state at $E_x = 14.51$ MeV (1971WA21). See also (1975AJ02).

45. $^{11}\text{B}(\alpha, \alpha)^{11}\text{B}$

Angular distributions have been reported at $E_\alpha = 28.3$ to 29.0 MeV [see (1975AJ02)] and at 24 MeV (1976DE34: $\alpha_0 \rightarrow \alpha_3, \alpha_{4+5}$). See also (1978ZE03; theor.).

Table 11.17: States of ^{11}B from $^{11}\text{B}(\text{p}, \text{p}')^{11}\text{B}^*$ and $^{13}\text{C}(\text{d}, \alpha)^{11}\text{B}$ ^a

(1971BR41, 1974KA15) ^b	(1970BR23) ^c
E_x (keV)	
0	0
2124.7 ± 0.5	2125.4 ± 1.4
4445.2 ± 0.5	4444.5 ± 1.6
5021.1 ± 0.6	5020.2 ± 1.9
6743.0 ± 0.7 ^d	6745.8 ± 3.4
6792.6 ± 1.6	6795 ± 3.0
7285.6 ± 1.5	
7978.0 ± 1.7	
8559.4 ± 1.9	8520 ± 70
8920.2 ± 2.0	8910 ± 60
9185.0 ± 2.0	
9274.4 ± 2.0	
10450 ± 150 ^e	
11650 ± 150	
12850 ± 100	
15200 ± 150	
16400 ± 150	

^a See also Table 11.18 in (1975AJ02).

^b $^{11}\text{B}(\text{p}, \text{p}')^{11}\text{B}$.

^c $^{13}\text{C}(\text{d}, \alpha)^{11}\text{B}$.

^d Values below are normalized to $E_x = 4445.3, 5020.0$ and 6743.4 keV.

^e This value and the values below are from (1969SU03).

Table 11.18: Possible $T = \frac{3}{2}$ states in ^{11}B ^a

Reaction	E_x (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	References
$^{10}\text{Be}(p, \gamma)^{11}\text{B}$	12.56 ± 30	230 ± 65	(1970G004)
$^{11}\text{B}(^3\text{He}, ^3\text{He})^{11}\text{B}^*$	12.51 ± 50	260 ± 50	(1971WA21)
	12.56 ± 30	240 ± 50	“best”
$^{10}\text{Be}(p, \gamma)^{11}\text{B}$	12.91 ± 20	235 ± 27	(1970G004)
$^{11}\text{B}(^3\text{He}, ^3\text{He})^{11}\text{B}^*$	12.98 ± 90	390 ± 120	(1971WA21)
$^{13}\text{C}(p, ^3\text{He})^{11}\text{B}$	12.94 ± 50	350 ± 50	(1968CO26)
$^{13}\text{C}(p, ^3\text{He})^{11}\text{B}$	12.91 ± 30	260 ± 50	(1974BE20)
	12.91 ± 20	240 ± 30	“best”
$^{10}\text{Be}(p, \gamma)^{11}\text{B}$	14.33 ± 20	255 ± 36	(1970G004)
$^{11}\text{B}(^3\text{He}, ^3\text{He})^{11}\text{B}^*$	14.40 ± 50	220 ± 50	(1971WA21)
	14.33 ± 20	250 ± 40	“best”
$^{10}\text{Be}(p, \gamma)^{11}\text{B}$	15.32 ± 100	635 ± 180	(1970G004)

^a These states have also been seen in other reactions: see Table 11.3. The parameters shown in that table reflect all the pertinent data. See also Table 11.21 for $T = \frac{3}{2}$ states in ^{11}C .

46. $^{11}\text{B}(^6\text{Li}, ^6\text{Li})^{11}\text{B}$

The elastic scattering has been measured at $E(^6\text{Li}) = 28$ MeV (1972BA52).

47. $^{11}\text{B}(^9\text{Be}, ^9\text{Be})^{11}\text{B}$

See (1975AJ02).

48. (a) $^{11}\text{B}(^{10}\text{B}, ^{10}\text{B})^{11}\text{B}$

(b) $^{11}\text{B}(^{11}\text{B}, ^{11}\text{B})^{11}\text{B}$

See (1976HI05). See also (1975AJ02).

49. $^{11}\text{B}(^{12}\text{C}, ^{12}\text{C})^{11}\text{B}$

The elastic scattering has been studied at $E(^{11}\text{B}) = 18.8$ to 34.1 MeV (1978FR20) and 28 MeV (1969VO07, 1969VO10), at $E(^{12}\text{C}) = 15$ to 24 MeV (1974BO15, 1975DU11) and at $E(^{12}\text{C}) = 87$ MeV (1971LI11). The population of $^{11}\text{B}^*(2.12, 4.45, 6.79)$ and of $^{12}\text{C}^*(0, 4.43)$ are also reported: see (1969VO07). See also (1976ST12), (1978RO1D; astophys.) and (1975RE04, 1978AV1A, 1978VA1A; theor.).

50. $^{11}\text{B}(^{14}\text{N}, ^{14}\text{N})^{11}\text{B}$

The elastic scattering has been investigated at $E(^{14}\text{N}) = 41, 77$ and 133 MeV (1971LI11).

51. $^{11}\text{B}(^{16}\text{O}, ^{16}\text{O})^{11}\text{B}$

The elastic scattering has been studied at $E(^{16}\text{O}) = 14.5$ to 60 MeV [see (1975AJ02)] and at $E(^{11}\text{B}) = 115$ MeV (1979BR03).

52. $^{11}\text{C}(\beta^+)^{11}\text{B}$ $Q_m = 1.982$

See ^{11}C .

53. $^{12}\text{C}(\gamma, p)^{11}\text{B}$ $Q_m = -15.9569$

The fraction of transitions to the ground and to excited states of ^{11}B [and to ^{11}C states reached in the (γ, n) reaction] has been measured at $E_{\text{bs}} = 24.5, 27, 33$ and 42 MeV: the ground state is predominantly populated. The population of analog states in the (γ, n) and (γ, p) reactions are similar: see ^{11}C (1970ME17). See also (1979GO1N: $E_{\text{bs}} = 21.7$ to 31.0 MeV) and (1978KI03: E_{bs} to 120 MeV). Differential cross sections have been measured at $E_\gamma = 60, 80$ and 100 MeV (1976MA34, 1974FI17; $p_0, p_1, p_2 + p_3 + p_4$). See also (1975MA1E, 1979KI1H), ^{12}C and (1975AJ02).

54. $^{12}\text{C}(\pi^+, \pi^+p)^{11}\text{B}$ $Q_m = -15.9569$

At $E_{\pi^+} = 100$ MeV the reaction proceeds primarily to $^{11}\text{B}_{\text{g.s.}}$ (1978CO02). At $E_\pi = 200$ MeV the ratios for σ_n/σ_p for the first excited states in $^{11}\text{C}/^{11}\text{B}$ are 1.4 ± 0.2 for π^- and $1/1.8 \pm 0.2$ for π^+ (1979LI11). At $E_{\pi^+} = 60$ to 300 MeV $^{11}\text{B}^*(4.45)$ [$J^\pi = \frac{5}{2}^-$] is strongly populated, as is the analog state in the mirror reaction (1978MO01). See also reaction 25 in ^{11}C .

55. $^{12}\text{C}(\text{e}, \text{ep})^{11}\text{B}$ $Q_{\text{m}} = -15.9569$

At $E_{\text{e}} = 497$ MeV $^{11}\text{B}^*(0, 2.1, 5.0)$ are populated: $l = 1$ (1976MO17). See also (1978SH14), (1975AJ02), ^{12}C and (1975RO18, 1979BO07; theor.).

56. $^{12}\text{C}(\text{n}, \text{d})^{11}\text{B}$ $Q_{\text{m}} = -13.7323$

See (1974KI1A, 1975MC19, 1976KI1D).

57. $^{12}\text{C}(\text{p}, 2\text{p})^{11}\text{B}$ $Q_{\text{m}} = -15.9569$

Gross structure is seen in the summed proton spectrum corresponding to $^{11}\text{B}_{\text{g.s.}}$ and to an excited state with $J^{\pi} = \frac{1}{2}^{+}$ at $E_{\text{x}} \approx 19.5$ MeV: see (1975AJ02). A high resolution experiment at $E_{\text{p}} = 100$ MeV shows groups corresponding to $^{11}\text{B}^*(0, 2.12, 4.45, 5.02, 6.79)$ (1977DE1M) [see (1975AJ02) for the earlier work]. See also (1977DE1L, 1978DE1P). At $E_{\text{p}} = 100$ MeV DWIA gives reasonably good agreement with angular correlation distributions ($\text{p}_0, \text{p}_1, \text{p}_{2+3}, \text{p}_{4+5}$) (1976BH02). See also ^{12}C , (1976SI1A, 1977MC1F, 1977RO1E) and (1975PI1A, 1977RE1A, 1978HA35, 1979FA1B, 1979MA20; theor.).

58. $^{12}\text{C}(\text{d}, ^3\text{He})^{11}\text{B}$ $Q_{\text{m}} = -10.4634$

Angular distributions of ^3He ions have been measured for $E_{\text{d}} = 20$ to 50 MeV [see (1975AJ02)] and at $E_{\text{d}} = 29$ MeV (1978CO13: to $^{11}\text{B}^*(0, 2.12, 4.45)$) and $E_{\text{d}} = 52$ MeV (1968HI01, 1975MA41: to $^{11}\text{B}^*(0, 2.12, 4.45, 5.02, 6.74 + 6.79, 7.29)$) and 80 MeV [(1974AS04: to $^{11}\text{B}^*(0, 4.45)$), (1976DI05: to $^{11}\text{B}^*(0, 2.12, 5.02)$)]. Reported C^2S for $^{11}\text{B}_{\text{g.s.}}$ are 3.22 or 4.42 depending on the choice of parameters (1978CO13: $E_{\text{d}} = 29$ MeV; see also (d, t) results in reaction 28 of ^{11}C), and $2.98, 0.69, 0.31$ for $^{11}\text{B}^*(0, 2.12, 5.02)$ (1975MA41: $E_{\text{d}} = 52$ MeV, DWBA). See also (1975LU07, 1976CH1E, 1977CH01; theor.).

59. $^{12}\text{C}(\text{t}, \alpha)^{11}\text{B}$ $Q_{\text{m}} = 3.8571$

Angular distributions have been measured for $E_{\text{t}} = 1$ to $3.4, 10.1$ and 13 MeV: see (1975AJ02). For electromagnetic transitions see Table 11.4 (1968BE30).

60. (a) $^{12}\text{C}(\alpha, ^5\text{Li})^{11}\text{B}$ $Q_m = -17.92$
 (b) $^{12}\text{C}(\alpha, \alpha p)^{11}\text{B}$ $Q_m = -15.9569$

At $E_\alpha = 65$ MeV, $^{11}\text{B}^*(0, 2.12, 6.74 + 6.79)$ are strongly populated (1978SA26). For reaction (b) see (1977YA1A, 1979KU08).

61. $^{12}\text{C}(^{10}\text{B}, ^{11}\text{C})^{11}\text{B}$ $Q_m = -7.266$

See reaction 31 in ^{11}C (1975NA15, 1975VO05, 1976NA09).

62. $^{12}\text{C}(^{12}\text{C}, ^{13}\text{N})^{11}\text{B}$ $Q_m = -14.014$

The angular distribution involving the ground state transitions has been measured at $E(^{12}\text{C}) = 114$ MeV (1974AN36). See also (1979HE1E).

63. $^{12}\text{C}(^{14}\text{N}, ^{15}\text{O})^{11}\text{B}$ $Q_m = -8.660$

See (1975AJ02) and ^{15}O in (1981AJ01).

64. $^{12}\text{C}(^{16}\text{O}, ^{17}\text{O})^{11}\text{B}$ $Q_m = -12.595$

See (1977ZE1A; theor.).

65. $^{12}\text{C}(^{19}\text{F}, ^{20}\text{Ne})^{11}\text{B}$ $Q_m = -3.1123$

At $E(^{19}\text{F}) = 40, 60$ and 68.8 MeV angular distributions involving $^{20}\text{Ne}_{\text{g.s.}} + ^{11}\text{B}_{\text{g.s.}}, ^{20}\text{Ne}_{1.63}^* + ^{11}\text{B}_{\text{g.s.}}$ and $^{20}\text{Ne}_{\text{g.s.}} + ^{11}\text{B}_{2.12}^*$ ($E = 68.8$ MeV only) have been measured by (1972SC03). See also (1975PU02).

66. $^{13}\text{C}(p, ^3\text{He})^{11}\text{B}$ $Q_m = -13.1851$

At $E_p = 50.5$ MeV, in addition to $^{11}\text{B}^*(0, 2.12, 4.45, 5.02, 6.74, 8.92)$, a state is observed at $E_x = 12.94 \pm 0.05$ MeV, $\Gamma = 350 \pm 50$ keV. Comparison of the angular distributions of the ^3He and of the tritons [in the analog reaction] at $E_p = 43.7$ and 50.3 MeV lead to the assignments $J^\pi = \frac{1}{2}^-$, $T = \frac{3}{2}$ for this state and for $^{11}\text{C}^*(12.50)$: the strong proton and the weak α -decay are consistent with this assignment (1968CO26). See Table 11.18 and (1974BE20). Angular distributions have been measured to $^{11}\text{B}^*(2.12, 4.45, 5.02)$, at $E_p = 26.9$ to 49.6 MeV [see (1975AJ02)] and at $E_{\bar{p}} = 49.6$ MeV (1974MA12; also to $^{11}\text{B}^*(6.74, 12.94)$). See also reaction 34 in ^{11}C for the mirror reaction, and see ^{14}N in (1981AJ01).

$$67. \ ^{13}\text{C}(d, \alpha)^{11}\text{B} \quad Q_m = 5.1680$$

Observed proton groups are displayed in Table 11.17 (1970BR23). Angular distributions are reported at $E_d = 0.41$ to 14.1 MeV: see (1975AJ02). See also ^{15}N in (1981AJ01).

$$68. \ ^{13}\text{C}(^{17}\text{O}, ^{19}\text{F})^{11}\text{B} \quad Q_m = -4.865$$

See (1977CH22).

$$69. \ ^{14}\text{C}(p, \alpha)^{11}\text{B} \quad Q_m = -0.7839$$

Angular distributions of the $\alpha_0 \rightarrow \alpha_3$ groups have been measured at $E_p = 18$ MeV (1962BR34).

$$70. \ (a) \ ^{14}\text{N}(n, \alpha)^{11}\text{B} \quad Q_m = -0.1580$$

$$(b) \ ^{14}\text{N}(n, 2\alpha)^7\text{Li} \quad Q_m = -8.823$$

Angular distributions have been measured for $E_n = 4.9$ to 18.8 MeV [see (1975AJ02)] and at 13.9 MeV (1978MO09; $\alpha_0 \rightarrow \alpha_3$). At $E_n = 14.1$ and 15.7 MeV various states of ^{11}B with $8.9 < E_x < 14.5$ MeV appear to be involved in the sequential decay to ^7Li . Angular correlation results are consistent with $J^\pi = \frac{7}{2}$ and $\frac{5}{2}$ for $^{11}\text{B}^*(9.19, 9.27)$ respectively (1971SC16). See also (1979SU1C), ^7Li in (1979AJ01), ^{15}N in (1981AJ01), and (1975AJ02).

$$71. \ ^{14}\text{N}(^{10}\text{B}, ^{13}\text{N})^{11}\text{B} \quad Q_m = 0.902$$

See ^{13}N in (1981AJ01).

$$72. \text{}^{15}\text{N}(\alpha, \text{}^8\text{Be})\text{}^{11}\text{B} \quad Q_m = -11.0832$$

At $E_\alpha = 72.5$ MeV angular distributions are reported to $^{11}\text{B}^*(0, 2.10 \pm 0.04, 4.50 \pm 0.07, 6.75 \pm 0.04)$: $S = 0.23, 0.12, 0.29, 0.45$ (unresolved groups) ([1974WO1C](#), [1976WO11](#)).

$$73. \text{}^{16}\text{O}(\text{d}, \text{}^7\text{Be})\text{}^{11}\text{B} \quad Q_m = -16.039$$

At $E_d = 80$ MeV angular distributions have been measured to $^{11}\text{B}^*(0, 2.12, 4.45 + 5.02, 6.74 + 6.79 + 7.29)$ ([1978OE1A](#)).

$$74. \text{}^{19}\text{F}(\text{}^{12}\text{C}, \text{}^{20}\text{Ne})\text{}^{11}\text{B} \quad Q_m = -3.1123$$

See ^{20}Ne in ([1978AJ03](#)).

¹¹C
(Figs. 3 and 4)

GENERAL: (See also (1975AJ02).)

Special levels: (1976IR1B).

Astrophysical questions: (1976VIIA, 1977SC1D, 1977SI1D, 1978BU1B).

Special reactions: (1975HU14, 1976BE1K, 1976BU16, 1976DI01, 1976HE1H, 1976LE1F, 1976SM07, 1977AR06, 1977AS03, 1977SC1G, 1978DI1A, 1978GE1C, 1978HE1C, 1979KA07, 1979VI05).

Muon and neutrino capture and reactions: (1975DO1F, 1976DO1G).

Pion capture and reactions (See also reactions 25 and 26.): (1975BA66, 1975BH03, 1975HU13, 1975KA1G, 1975PA1D, 1975SC1N, 1975SI18, 1975ZI1B, 1976MI15, 1977BA2H, 1977LE1G, 1977MA1M, 1978AM01, 1978HI03, 1978MO01, 1978WA1B, 1979AN1J, 1979HU02).

Applied topics: (1975AL1B, 1975CH1H, 1977MA1P, 1977WO1A, 1978LO1D, 1978TI1A, 1978WI1F, 1978WO1C, 1979AL1Q, 1979WI1G).

Other topics: (1976IR1B, 1979BE1H).

Ground state of ¹¹C: (1969SC34, 1974SHYR, 1975BE31, 1976DO1G, 1977KA07, 1977YO1D).

$$\mu = -0.964 \pm 0.001 \text{ nm (1969WO03);}$$

$$Q = 34.26 \text{ mb (1978LEZA).}$$

1. ¹¹C(β^+)¹¹B $Q_m = 1.982$

The half-life of ¹¹C is 1223.1 ± 1.2 sec [mean of previously reported values (see (1975AJ02)); mainly based on $\tau_{1/2} = 1222.9 \pm 1.2$ sec (1975AZ01)]. See also (1975BE28). $\log ft = 3.599 \pm 0.002$ (1975AZ01) based on $E_{\beta^+}(\text{max}) = 960 \pm 1$ keV (from Q_m). See also (1975AZ01). (1975BE28) measure $E_{\beta^+}(\text{max}) = 960.8 \pm 2.6$ keV. The ratio of K-capture to positron emission is $(0.230_{-0.011}^{+0.014})\%$ (1967CA09). See also (1975AJ02, 1977BA48, 1978RA2A) and (1975KR14, 1975WI1E, 1976BE1E, 1976DO1G, 1977RI08, 1977YO1D, 1977YO1E, 1979DE15; theor.).

2. (a) ⁶Li(⁶Li, n)¹¹C $Q_m = 9.453$
 (b) ⁷Li(⁶Li, 2n)¹¹C $Q_m = 2.203$
 (c) ⁷Li(⁷Li, 3n)¹¹C $Q_m = -5.048$

At $E(^6\text{Li}) = 4.1$ MeV [reaction (a)] angular distributions have been obtained for the neutrons to $^{11}\text{C}^*(2.00, 4.32, 4.80, 6.34 + 6.48, 6.90, 7.50)$. In addition, n- γ coincidences via $^{11}\text{C}^*(8.42)$ [and a 8.42 MeV γ -ray] are reported. $^{11}\text{C}^*(8.10)$ was not observed (1967BA53). The lifetimes, τ_m , for $^{11}\text{C}^*(4.32, 6.90, 7.50)$ are < 140 , < 69 and < 91 fsec, respectively. The upper limits for τ_m of $^{11}\text{C}^*(6.34, 6.48)$ [which were unresolved] are 0.5 psec. The ground state transition from $^{11}\text{C}^*(7.50)$ has $E_\gamma = 7505 \pm 8$ keV (1969TH01). For reaction (b) see (1968AJ02). For reaction (c) see (1974CE06).

Table 11.19: Energy levels of ^{11}C ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
g.s.	$\frac{3}{2}^-; \frac{1}{2}$	$\tau_{1/2} = 20.40 \pm 0.04$ min	β^+	1, 2, 4, 5, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39
2.0000 ± 0.5	$\frac{1}{2}^-$	$\tau_m < 0.5$ psec	γ	2, 4, 5, 12, 13, 14, 16, 21, 22, 25, 26, 27, 28, 29, 30, 31, 34, 35, 38
4.3188 ± 1.2	$\frac{5}{2}^-$	< 0.14 psec	γ	2, 4, 5, 12, 13, 16, 21, 22, 25, 26, 27, 28, 29, 34, 38
4.8042 ± 1.2	$\frac{3}{2}^-$	< 0.5 psec	γ	2, 4, 12, 16, 21, 22, 27, 29, 34, 38
6.3392 ± 1.4	$\frac{1}{2}^+$	< 0.11 psec	γ	2, 4, 13, 21, 26, 27, 29, 38
6.4782 ± 1.3	$\frac{7}{2}^-$	< 0.25 psec	γ	2, 4, 5, 12, 13, 16, 21, 22, 26, 27, 29, 34, 38
6.9048 ± 1.4	$\frac{5}{2}^+$	< 69 fsec	γ	2, 4, 12, 13, 17, 27, 29, 34, 38
7.4997 ± 1.5	$\frac{3}{2}^+$	< 91 fsec	γ	2, 4, 13, 27, 29, 34
8.1045 ± 1.7	$\frac{3}{2}^-$		(γ)	4, 13, 22, 27, 29
8.424 ± 8	$\frac{5}{2}^-$		γ	2, 4, 12, 13, 16, 26, 27, 29
8.655 ± 8	$\frac{7}{2}^+$	$\Gamma \ll 9$ keV	(γ)	12, 13, 16, 26, 27
8.701 ± 20	$\frac{5}{2}^+$	15 ± 1		12, 13, 16, 26, 27
9.732 ± 5	($\frac{5}{2}^+$)	450 ± 50	γ, p, α	5, 7, 11, 27
10.084 ± 5	$\frac{7}{2}^+$	≈ 230	p, α	7, 11, 13, 27
10.680 ± 5	$\frac{9}{2}^+$	200 ± 30	p, α	7, 11, 12, 27

Table 11.19: Energy levels of ^{11}C ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
11.03 \pm 30	$T = \frac{1}{2}$	300 \pm 60		27, 29, 34
11.44 \pm 10		360	p, α	11, 27, 29
(12.16 \pm 40)	$(T = \frac{3}{2})$	270 \pm 50	p	4, 8, 22, 26
12.4	$\pi = -$	1–2 MeV	γ , p	5
12.51 \pm 30	$\frac{1}{2}^-; \frac{3}{2}$	490 \pm 40	p, α	4, 8, 22, 26, 29, 34
12.65 \pm 20	$(\frac{7}{2}^+)$	360	p, ^3He , α	10, 11, 26
(13.01)			γ , p	5
13.33 \pm 60		270 \pm 80		34
13.4		1100 \pm 100	p, α	11
13.90 \pm 20	$(T = \frac{3}{2})$	200 \pm 100	p	4, 8, 22, 34
14.07 \pm 20		135 \pm 50	n, p	6, 34
14.76 \pm 40		\approx 450	n, p, ^3He	4, 6, 8, 10
15.35 \pm 50	$\pi = -$	broad	γ , n, p	5, 6, 8, 29
15.59 \pm 50		\approx 450	n, p	6, 8
16.7	$\pi = -$	800 \pm 100	γ , p	5
(18.2)			γ , p	5
(23.0)				29
(28.0)				29

^a See also Table 11.20.



See (1972PA1C, 1975FO19).



Reported neutron groups are listed in Table 11.16 of (1968AJ02). Angular distributions have been studied in the range $E(^3\text{He}) = 1.3$ to 13 MeV: see (1975AJ02) and (1979OS10). The dominant L -values from the angular distributions reported by (1974FU11) are 0 for $^{11}\text{C}^*(0, 8.10)$, 1 for $^{11}\text{C}^*(6.34, 7.50)$, 2 for $^{11}\text{C}^*(2.00, 4.32, 4.80, 6.48, 8.42)$ and 3 for $^{11}\text{C}^*(6.90)$. Neutron groups to

Table 11.20: Gamma decay of ^{11}C levels

E_i (MeV)	J^π	τ_m^b (psec)	E_f (MeV)	Branch ^a (%)	Mult. ^a	Branch ^c (%)	X^c
2.00	$\frac{1}{2}^-$	< 0.5	0	100			
4.32	$\frac{5}{2}^-$	< 0.5	0	100	M1	100	$+0.17 \pm 0.03^g$
		$< 0.14^h$	2.00	< 2		< 2	
4.80	$\frac{3}{2}^-$	< 0.5	0	83 ± 4	M1	86 ± 2^f	e
			2.00	17 ± 4		14 ± 2^f	e
6.34	$\frac{1}{2}^+$	< 0.11	0	65 ± 3	E1	68 ± 3	
			2.00	35 ± 3		32 ± 3^d	
			4.32			< 7	
			4.80	< 4		< 3	
6.48	$\frac{7}{2}^-$	< 0.25	0	89 ± 2	E2	88 ± 2	-0.01 ± 0.06
			2.00	< 2		< 4	
			4.32	11 ± 2		12 ± 2	
			4.80			< 2	
6.90	$\frac{5}{2}^+$	< 0.16	0	89 ± 3	E1	91 ± 2	0.02 ± 0.03
		$< 0.07^h$	2.00	< 2		< 1	
			4.32	11 ± 3		4.5 ± 1	
			4.80	< 3		4.5 ± 1	
			6.34	< 5			
			6.48	< 5			
7.50	$\frac{3}{2}^+$	< 0.5	0	36 ± 2	E1	37 ± 3	-0.04 ± 0.04
		$< 0.09^h$	2.00	64 ± 2	E1	63 ± 8	0 ± 0.03
			4.32	< 3		< 1	
			4.80	< 3		< 1	
			6.34	< 3			
			6.48	< 3			
			6.90	< 4			
9.73 ⁱ	$(\frac{5}{2}^+)$		0	65 ± 15			
			2.00	3			
			4.32	12 ± 2			
			6.48	20			

^a From ${}^9\text{Be}({}^3\text{He}, n){}^{11}\text{C}$ and ${}^{10}\text{B}(d, n){}^{11}\text{C}$ (1965OL03): includes earlier measurements, except for ${}^{11}\text{C}^*(9.73)$: see footnote ⁱ.

^b (1966WA10).

^c (1968EA03): ${}^{12}\text{C}({}^3\text{He}, \alpha){}^{11}\text{C}$: $X \equiv$ amplitude ratio of $(L + 1)/L$.

^d The cascade id through ${}^{11}\text{C}^*(2.0)$ and not ${}^{11}\text{C}^*(4.3)$ (1968EA03).

^e See ${}^{12}\text{C}({}^3\text{He}, \alpha){}^{11}\text{C}$.

^f (86 ± 3) , $(14 \pm 3)\%$ (1966GA19); (89 ± 3) , $(16 \pm 3)\%$ (1967BL22).

^g $+0.16$ $(-0.02, +0.06)$ (1966GA19); 0.13 ± 0.04 (1967BL22).

^h (1969TH01).

ⁱ (1961JA11): ${}^{10}\text{B}(p, \gamma){}^{11}\text{C}$.

to $T = \frac{3}{2}$ states have been reported by (1971WA21) [$E_x = 12.17 \pm 0.05$ and 12.55 ± 0.05 MeV] and by (1969BR30) [$E_x = 12.5 \pm 0.1$, 13.7 ± 0.1 and 14.7 ± 0.1 MeV]: see Table 11.22.

Gamma branching ratios and multiplicities for ${}^{11}\text{C}$ levels up to $E_x = 7.5$ MeV have been studied by (1965OL03, 1965RO07): see Table 11.20. Together with evidence from reactions 12 and 27 they lead to assignments of $J^\pi = \frac{1}{2}^-, \frac{5}{2}^-, \frac{3}{2}^-, \frac{1}{2}^+, \frac{7}{2}^-, \frac{5}{2}^+, \frac{3}{2}^+$ for ${}^{11}\text{C}^*(2.00, 4.32, 4.80, 6.34, 6.48, 6.90, 7.50)$: see (1965OL03, 1965RO07) and reaction 3 in (1968AJ02) for a summary of the evidence concerning these assignments. See also (1977ZA1B; applied), (1977OS08; theor.) and ${}^{12}\text{C}$.

5. ${}^{10}\text{B}(p, \gamma){}^{11}\text{C}$

$$Q_m = 8.691$$

The thick target yield for this reaction has been measured for $E_p = 0.27$ to 3.0 MeV (1975PE1A): see also reaction 11. A broad resonance is reported at $E_p = 1.15$ MeV: see Table 11.22. Capture γ -rays are observed corresponding to the ground state transition and to cascades via ${}^{11}\text{C}^*(2.00, 4.32, 6.48)$: see Table 11.21 (1961JA11). The 90° yield of γ_0 has been measured for $E_p = 2.6$ to 17 MeV and angular distributions have been obtained for $E_p = 2.8$ to 14 MeV. The excitation function is consistent with the giant resonance centered at $E_x \approx 16$ MeV. In addition to weak structures at $E_p = 4.75$ MeV and 10.5 MeV, there are three major peaks at $E_p = 4.1, 7.0$ and 8.8 MeV ($\Gamma = 1 - 2$ MeV) [$E_x = 12.4, 15.0, 16.7$ MeV]. At ${}^{11}\text{C}^*(12.4)$, the γ_0 angular distribution is essentially isotropic: $\Gamma_p \Gamma_\gamma / \Gamma \approx 200$ eV, $\Gamma_\gamma \approx 5$ keV (assuming $\Gamma_p \approx 10$ keV). The $E_p = 4.1$ MeV resonance is probably part of the E1 giant resonance and is formed by s-wave capture. At the two higher resonances the angular distributions are characteristic of E1 giant resonances in light nuclei. The ${}^{10}\text{B}(p, \gamma_1)$ cross section is small for $E_p = 2.6$ to 17 MeV (1970KU09). See also (1974LO1B, 1975AJ02), (1976RO1Q; astrophysics) and (1977YO1F; applied).

6. ${}^{10}\text{B}(p, n){}^{10}\text{C}$

$$Q_m = -4.434$$

$$E_b = 8.691$$

The total (p, n) cross section has been measured to $E_p = 10.6$ MeV: broad maxima are observed at $E_p = 5.92 \pm 0.02, 6.68 \pm 0.04, 7.33 \pm 0.05$ and 7.60 ± 0.05 MeV (see Table 11.23) (1963EA01).

Table 11.21: Possible $T = \frac{3}{2}$ states in ^{11}C ^a

Reaction	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	Refs.
$^9\text{Be}(^3\text{He}, \text{n})^{11}\text{C}$	12.17 ± 0.05	200 ± 100	(1971WA21)
$^{10}\text{B}(\text{p}, \text{p}_2)^{10}\text{B}^{**}$	12.20 ± 0.10		(1971WA21)
$^{11}\text{B}(^3\text{He}, \text{t})^{11}\text{C}$	12.15 ± 0.05	290 ± 50	(1971WA21)
	12.16 ± 0.04 ^b	270 ± 50	mean
$^9\text{Be}(^3\text{He}, \text{n})^{11}\text{C}$	12.55 ± 0.05	350 ± 100	(1971WA21)
$^{10}\text{B}(\text{p}, \text{p}_2)^{10}\text{B}^{**}$	12.45 ± 0.10	400 ± 100	(1971WA21)
$^{11}\text{B}(^3\text{He}, \text{t})^{11}\text{C}$	12.57 ± 0.07	370 ± 90	(1971WA21)
$^{13}\text{C}(\text{p}, \text{t})^{11}\text{C}$	12.47 ± 0.06	550 ± 50	(1968CO26) ^c
$^{13}\text{C}(\text{p}, \text{t})^{11}\text{C}$	12.48 ± 0.04	540 ± 60	(1974BE20)
	12.51 ± 0.03	490 ± 40	mean
$^9\text{Be}(^3\text{He}, \text{n})^{11}\text{C}$	13.7 ± 0.1		(1969BR30)
$^{11}\text{B}(^3\text{He}, \text{t})^{11}\text{C}$	13.92 ± 0.05	260 ± 50	(1971WA21) ^a

^a See also Table 11.18 for $T = \frac{3}{2}$ states in ^{11}B .

^b See, however, reaction 34 (1974BE20).

^c See also (1974MA12).

The cross section for formation of $^{10}\text{C}_{\text{g.s.}}$ measured up to 12 MeV shows similar behavior to 8 MeV. At $E_p \approx 8$ MeV, a sharp maximum is observed. The cross section for production of 3.35 MeV γ -rays (from $^{10}\text{C}^*$) does not appear to show structure for $E_p = 8.5$ to 12 MeV (1966SE03). See also ^{10}C in (1979AJ01).

7. $^{10}\text{B}(\text{p}, \text{p})^{10}\text{B}$

$$E_b = 8.691$$

Below $E_p = 0.7$ MeV the scattering can be explained in terms of pure s-wave potential scattering but the possibility of a state near $E_p = 0.27$ MeV ($E_x = 8.95$ MeV) cannot be excluded. The elastic scattering then shows two conspicuous anomalies at $E_p = 1.50 \pm 0.02$ MeV and at 2.18 MeV [$E_x = 10.05$ and 10.67 MeV] with $J^\pi = \frac{7}{2}^+$ and $\frac{9}{2}^+$: see Table 11.22 (1962OV02: $E_p = 0.15$ to 3.0 MeV). At higher energies (to $E_p = 10.5$ MeV) a single broad resonance is reported at $E_p \approx 5$ MeV (1969WA11) while (1970BO17) suggest additional structure. Polarization measurements are reported at 30.3 MeV (1976DE15, 1977PH02): optical model parameters are derived. The depolarization parameter D has been measured for polarized protons with $E_p = 26$ MeV (1976BI1E) and 50 MeV (1970BA05). See also (1977SA1B) and (1975BL10; theor.).

Table 11.22: Resonances ^a in ¹⁰B + p

E_{res} (MeV \pm keV)	E_x (MeV)	J^π	Γ_{lab} (keV)	Decay	Refs.
1.145 ± 5	9.732	$(\frac{5}{2}^+)$	500 ± 50	γ, p_0, α_0	A
1.533 ± 5	10.084	$\frac{7}{2}^+$	≈ 250	p_0, α_0, α_1	A
2.189 ± 5	10.680	$\frac{9}{2}^+$	220 ± 30	p_0, α_0, α_1	A
3.03 ± 10	11.44		400	α_0, α_1	(1962OP03, 1964JE01)
3.9 ± 100	12.20	$T = \frac{3}{2}$		p_2	(1971WA21)
4.1 ± 100	12.45	$T = \frac{3}{2}$	440 ± 100	p_2	(1971WA21)
$4.1^{b,c}$	12.4	$\pi = -$	1–2 MeV	γ_0	(1970KU09)
4.36 ± 20	12.65	$(\frac{7}{2}^+)$	400	$\alpha_0, \alpha_1, {}^3\text{He}$	A
(4.75)	(13.01)			γ_0	(1970KU09)
5.2	13.4		1200 ± 100	α_0, α_1	A
5.73 ± 20	13.90		≈ 500	p	A
5.92 ± 20	14.07		broad	n	A
6.68 ± 40	14.76		≈ 500	n, p, ³ He	A
7.33 ± 50^c	15.35	$\pi = -$	broad	γ_0, n, p	A
7.60 ± 50	15.59		≈ 500	n, p	A
8.8^c	16.7	$\pi = -$	900 ± 100	γ_0	(1970KU09)
(10.5)	(18.2)			γ_0	(1970KU09)

A: See references listed for this resonance in Table 11.23 (1975AJ02).

^a See also Table 11.20 here, and Tables 11.23 and 11.24 in (1975AJ02). Table 11.23 displays some other reported resonances; Table 11.24 gives detailed parameters for ¹¹C*(9.73, 10.08, 10.68, 12.65).

^b $\Gamma_p \Gamma_\gamma / \Gamma \approx 20$ eV (1970KU09).

^c Probably part of the E1 giant resonance (1970KU09).

8. $^{10}\text{B}(\text{p}, \text{p}')^{10}\text{B}$

$$E_b = 8.691$$

The yield γ_1 [from $^{10}\text{B}^*(0.72)$] rises monotonically from $E_p = 1.5$ to 4.1 MeV and then shows resonance behavior at $E_p = 4.36$ and 5.73 MeV (1962OP03: see Table 11.22). For $E_p = 6$ to 12 MeV, the cross section for γ_1 shows several sharp maxima superposed on a broad maximum ($\Gamma \approx 2.5$ MeV) at $E_p \approx 7.2$ MeV (1966SE03). [The cross section below $E_p = 10$ MeV appears to be in error: see (1969WA23).] Yields of five other γ -rays involved in the decay of $^{10}\text{B}^*(1.74, 2.16, 3.59, 5.18)$ have also been measured by (1966SE03) in the range $E_p = 4$ to 12 MeV.

Yield curves for inelastically scattered protons have been measured at $E_p = 5.0$ to 16.4 MeV (p_1, p_2, p_3), 6.6 to 16.4 MeV (p_4), 8.9 to 16.4 MeV (p_5) and 10.9 to 16.4 MeV (p to $^{10}\text{B}^*(6.03)$): the principal feature for all groups, except that to $^{10}\text{B}^*(6.03)$, is a structure at $E_p \approx 7.5$ MeV, $\Gamma \approx 4$ MeV. In addition narrower structures are observed, including three at $E_p = 5.75, 6.90$ and 7.80 MeV (± 0.2 MeV) and widths of ≈ 500 keV (1969WA23). It had previously been suggested by (1966SE03) that the formation of $T = 1$ states was relatively suppressed in this reaction. (1969WA23) find that the isotopic spin effect disappears when a correction factor $(2J_f + 1)$ is included. Excitation curves for the p_1, p_2 and p_3 groups have been measured for $E_p = 3.5$ to 5.0 MeV. Possible resonances are observed in the p_2 yield [to the $T = 1$ state $^{10}\text{B}^*(1.74)$] corresponding to the first $T = \frac{3}{2}$ states at $E_x = 12.16$ and 12.50 MeV [see Table 11.21]: these do not occur in the yield of p_1 and p_3 (1971WA21). See also (1976DE15), ^{10}B in (1979AJ01) and (1975AJ02).

9. $^{10}\text{B}(\text{p}, \text{d})^9\text{B}$

$$Q_m = -6.211$$

$$E_b = 8.691$$

Polarization measurements have been carried out at $E_p = 49.6$ MeV for the deuterons to $^9\text{B}^*(0, 2.36)$ (1971SQ02). See also ^9B in (1979AJ01).

10. $^{10}\text{B}(\text{p}, ^3\text{He})^8\text{Be}$

$$Q_m = -0.5323$$

$$E_b = 8.691$$

The ground-state yield shows slight maxima at energies similar to those in the (p, α) yield in the range $E_p = 4$ to 10 MeV. However, the angular distributions do not vary strongly over the region and it is suggested that a direct interaction mechanism dominates (1963JE01). (1966SE03) report two strong maxima at $E_p \approx 4.5$ and 6.5 MeV. See also ^8Be in (1979AJ01).

11. $^{10}\text{B}(\text{p}, \alpha)^7\text{Be}$

$$Q_m = 1.146$$

$$E_b = 8.691$$

The total cross section for this reaction has been measured for $E_p = 60$ to 180 keV by (1972SZ02): the extrapolated cross section at the Gamow energy, taken to be 19.1 keV, is $\approx 10^{-12}$ b. The thick target yield for $E_p = 75$ keV to 3 MeV has been measured by (1975PE1A): the ^7Be yield constitutes a potential problem if natural boron is used as fuel in CTR devices.

The parameters of observed resonances are displayed in Table 11.22. The ground state (α_0) α -particles exhibit broad resonances at $E_p = 1.17, 1.53, 2.18, 3.0, 4.4, 5.1$ and 6.3 MeV. Alpha particles to ${}^7\text{Be}^*(0.43)$ [α_1] and 0.43 -MeV γ -rays exhibit all but the 1.2 MeV resonance: see (1975AJ02). A broad maximum dominates the region from $E_p = 4$ MeV to about 7.5 MeV (1966SE03). For spallation studies see (1976BO08, 1977AV01). See also (1976GR1F, 1976IN04) and (1976RO1Q; astrophysics).

$$12. {}^{10}\text{B}(\text{d}, \text{n}){}^{11}\text{C} \quad Q_m = 6.466$$

Table 11.23 presents the results obtained in this reaction and in the (${}^3\text{He}, \text{d}$) reaction (1955MA76, 1963OV02, 1970BO34). Information on the γ -decay of ${}^{11}\text{C}$ states has been summarized by (1965OL03) and is incorporated in Table 11.20. For the earlier work see (1959AJ76, 1968AJ02 [particularly Table 11.20], 1975AJ02). See also (1972RE1C) and (1974LO1B).

$$13. {}^{10}\text{B}({}^3\text{He}, \text{d}){}^{11}\text{C} \quad Q_m = 3.197$$

Table 11.23 displays the information derived from this reaction and from the (d, n) reaction. The study of the angular distributions of the deuterons to ${}^{11}\text{C}^*(8.66, 8.70)$ shows that these levels are the analogs, respectively, of ${}^{11}\text{B}^*(9.19, 9.27)$ whose J^π are $\frac{7}{2}^+$ and $\frac{5}{2}^+$ [the ${}^{11}\text{B}$ states were studied in the (d, p) reaction]: $\Gamma_{\text{c.m.}}$ are $\ll 9$ keV and 15 ± 1 keV, respectively, for ${}^{11}\text{C}^*(8.66, 8.70)$ (1973FO02). Singlet deuteron emission has been studied at $E({}^3\text{He}) = 8, 10$ and 11 MeV by (1970BO07). For the earlier work see (1975AJ02).

$$14. {}^{10}\text{B}(\alpha, \text{t}){}^{11}\text{C} \quad Q_m = -11.123$$

Angular distributions have been measured at $E_\alpha = 25.1$ MeV (1974DM01; t_1) and at 56 MeV (1969GA11; t_0). See also (1975AJ02).

$$15. {}^{10}\text{B}({}^6\text{Li}, {}^5\text{He}){}^{11}\text{C} \quad Q_m = 4.10$$

See (1957NO17).

$$16. {}^{10}\text{B}({}^7\text{Li}, {}^6\text{He}){}^{11}\text{C} \quad Q_m = -1.284$$

Table 11.23: Energy levels of ^{11}C from $^{10}\text{B}(\text{d}, \text{n})^{11}\text{C}$ and $^{10}\text{B}({}^3\text{He}, \text{d})^{11}\text{C}$

E_x (MeV \pm keV)	J^π	l^f	l^g	$S_{\text{d,n}}^g$	$S_{{}^3\text{He,d}}^g$	l^i	$S_{{}^3\text{He,d}}^i$
0	$\frac{3}{2}^-$	1	1	1.12	0.88	1	1.09
2.0006 \pm 0.9 ^a	$\frac{1}{2}^-$	(1)	(1)	(0.18)	(0.036)		
			(3)		≤ 0.09	(3)	< 0.40
4.322 \pm 10 ^b	$\frac{5}{2}^-$	1	1	0.27	0.20	1	0.17, 0.19
4.808 \pm 10 ^b	$\frac{3}{2}^-$	1	1	< 0.02		(1)	< 0.08
						(3)	< 0.35
6.345 \pm 10 ^b	$\frac{1}{2}^+$		2		0.07	2	0.08
6.476 \pm 10 ^b	$\frac{7}{2}^-$	1	1	0.86	0.56	1	0.73, 0.79
6.903 \pm 10 ^b	$\frac{5}{2}^+$	(1)				2	0.06
						0	< 0.04
7.498 \pm 10 ^b	$\frac{3}{2}^+$					2	0.08
8.107 \pm 10 ^b	$\frac{3}{2}^-$ ^k					1	0.07
8.424 \pm 8 ^c	$\frac{5}{2}^-$	1	1	0.65	0.46	1	0.73, 0.79
8.655 \pm 8 ^{c,d}	$\frac{5}{2}^+$	0	0	<u>0.84</u>	0.45		
			2	0.8	<u>0.32</u>		
	$\frac{7}{2}^+$ ^j		0	<u>0.63</u>	0.33	2	0.41
			2	0.6	<u>0.24</u>	0	< 0.34
8.701 \pm 20 ^d	$\frac{5}{2}^+$ ^j	(0)	0	<u>0.40</u>	0.14	0	< 0.8
			2	≤ 0.2	0.13		
	$\frac{7}{2}^+$		0	<u>0.30</u>	0.11		
			2	≤ 0.15	0.10		
10.08 ^h							
10.68 ^{d,e}			(0, 2)				

^a (${}^3\text{He}, \text{d}$): (1970BR23).

^b (${}^3\text{He}, \text{d}$): (1961HI08). $E({}^3\text{He}) = 9.84$ MeV.

^c (d, n): neutron threshold measurements (1955MA76); based on Q_m .

^d (d, n): observed by time-of-flight technique (1963OV02).

^e $\Gamma \approx 200$ keV (1963OV02).

^f From (d, n) work summarized in Table 11.20 of (1968AJ02).

^g From (1970BO34): $S_{\text{d,n}}$ obtained at $E_d = 5.8$ MeV, $S_{{}^3\text{He,d}}$ obtained at $E({}^3\text{He}) = 11.0$ MeV [both $\pm 30\%$]. When $S_{\text{d,n}}$ and $S_{{}^3\text{He,d}}$ differ appreciably, the more reliable value is underlined.

^h See (1971CO07).

ⁱ From $E({}^3\text{He}) = 21$ MeV work of (1971CO07); when two values are shown for $S_{{}^3\text{He,d}}$, they are in order of descending j .

^j Value determined by (1973FO02).

^k See (1970FO05).

Angular distributions of ${}^6\text{He}$ ions have been measured at $E({}^7\text{Li}) = 3.0$ to 3.8 MeV (1968ST12; to ${}^{11}\text{C}_{\text{g.s.}}$) and at 24 MeV (1977KO27: to ${}^{11}\text{C}^*(0, 4.32, 6.48)$). ${}^{11}\text{C}^*(2.0, 4.80, 8.42, 8.66 + 8.70)$ are also populated (1977KO27).

17. ${}^{10}\text{B}({}^{14}\text{N}, {}^{13}\text{C}){}^{11}\text{C}$ $Q_{\text{m}} = 1.140$

See (1975NA15, 1977MO1A), (1975AJ02) and ${}^{13}\text{C}$ in (1981AJ01).

18. ${}^{10}\text{B}({}^{16}\text{O}, {}^{15}\text{N}){}^{11}\text{C}$ $Q_{\text{m}} = -3.437$

See (1975NA15). See also ${}^{15}\text{N}$ in (1981AJ01) and (1977OK1D; theor.).

19. ${}^{10}\text{B}({}^{19}\text{F}, {}^{18}\text{O}){}^{11}\text{C}$ $Q_{\text{m}} = 0.697$

See (1968GA03).

20. ${}^{11}\text{B}(\gamma, \pi^-){}^{11}\text{C}$ $Q_{\text{m}} = -141.549$

Yields of ${}^{11}\text{C}$ have been measured from threshold to $E_{\gamma} = 169$ MeV (1976MI15; see it also for other references).

21. ${}^{11}\text{B}(\text{p}, \text{n}){}^{11}\text{C}$ $Q_{\text{m}} = -2.764$

Neutron groups have been observed to ${}^{11}\text{C}^*(0, 2.008, 4.320, 4.806, 6.330, 6.481)$ (± 20 keV) (1965OV1A; $E_{\text{p}} = 10.9$ MeV). Angular distributions of the n_0 group have been measured at many energies up to 49.5 MeV [see (1975AJ02)] and at $E_{\text{p}} = 3.15$ to 4.85 MeV (1978VA12). See also (1977ME1C) and ${}^{12}\text{C}$.

22. ${}^{11}\text{B}({}^3\text{He}, \text{t}){}^{11}\text{C}$ $Q_{\text{m}} = -2.001$

Angular distributions of t_0 and t_1 have been measured at $E({}^3\text{He}) = 10$ MeV (1967CR04), 14 MeV (1970NU02) and 217 MeV (1976WI05; also to ${}^{11}\text{C}^*(4.3, 4.8, 6.48, 8.10)$). The latter have been compared with microscopic calculations and calculations of two-step processes (1976WI05). At $E({}^3\text{He}) = 26$ MeV the known states of ${}^{11}\text{C}$ below $E_{\text{x}} = 11$ MeV are populated and triton groups are also observed to states, assumed to be $T = \frac{3}{2}$, at $E_{\text{x}} = 12.15, 12.57$ and 13.92 MeV [see Table 11.21] and, possibly, 14.15 MeV (1971WA21).

$$23. \text{}^{11}\text{B}(\text{}^6\text{Li}, \text{}^6\text{He})\text{}^{11}\text{C} \quad Q_m = -5.489$$

See (1975AJ02).

$$24. \text{}^{12}\text{C}(\gamma, \text{n})\text{}^{11}\text{C} \quad Q_m = -18.721$$

The fraction of transitions to the ground and to excited states of ^{11}C [and to ^{11}B states reached in the (γ, p) reaction] has been measured at $E_{\text{bs}} = 24.5, 27, 33$ and 42 MeV: the ground state is predominantly populated. The population of analog states in the (γ, n) and (γ, p) reactions are similar. And a significant decay strength is found to the positive parity states with $6 < E_x < 8$ MeV. In general the main contribution to the strength of the transitions to the various excited states of ^{11}B , ^{11}C lies in rather localized energy bands in ^{12}C which are a few MeV wide (1970ME17). See also (1975SC05, 1979KI1H). For the (e, n) reaction see (1978KL03).

$$25. \text{(a) } \text{}^{12}\text{C}(\text{e}, \text{en})\text{}^{11}\text{C} \quad Q_m = -18.721$$

$$\text{(b) } \text{}^{12}\text{C}(\pi^+, \pi^+\text{n})\text{}^{11}\text{C} \quad Q_m = -18.721$$

$$\text{}^{12}\text{C}(\pi^-, \pi^-\text{n})\text{}^{11}\text{C}$$

$$\text{(c) } \text{}^{12}\text{C}(\text{n}, 2\text{n})\text{}^{11}\text{C} \quad Q_m = -18.721$$

$$\text{(d) } \text{}^{12}\text{C}(\text{p}, \text{pn})\text{}^{11}\text{C} \quad Q_m = -18.721$$

For reaction (a) see (1977KN04, 1978SH14). (1978MO01) report the strong population of the $\frac{5}{2}^-$ state $^{11}\text{C}^*(4.32)$ (and of the analog state in ^{11}B) for $E_{\pi^+} = 60$ to 300 MeV. See also (1973HO43, 1979LI11) for reactions (b), reaction 54 in ^{11}B and the “GENERAL” section here. For reaction (c) see ^{13}C in (1981AJ01). For reaction (d) see (1976MC1F). See also (1975AJ02) and ^{12}C .

$$26. \text{}^{12}\text{C}(\pi^+, \text{p})\text{}^{11}\text{C} \quad Q_m = 121.628$$

At $E_{\pi^+} = 49.3$ MeV angular distributions have been obtained to $^{11}\text{C}^*(0, 2.0 \pm 0.2, 4.35 \pm 0.2, 6.4 \pm 0.2, 8.5 \pm 0.2, 12.5 \pm 0.3)$. At the same momentum transfer this reaction and the (p, d) reaction give similar intensities to the low-lying energy states of ^{11}C (1978AM01). See also (1979AN1K, 1979AN1M).

$$27. \text{}^{12}\text{C}(\text{p}, \text{d})\text{}^{11}\text{C} \quad Q_m = -16.497$$

Table 11.24: Levels of ^{11}C from $^{12}\text{C}(\text{p}, \text{d})^{11}\text{C}$ ^a

E_x (MeV \pm keV)	J^π ^c	S_{rel}	C^2S ^d
0	$\frac{3}{2}^-$	100	2.5
1.9997 ± 0.5 ^b	$\frac{1}{2}^-$	17.4	0.61
4.30 ± 50	$\frac{5}{2}^-$	< 0.06	(0.08)
4.80 ± 50	$\frac{3}{2}^-$	9.7	0.33
(6.34)	$\frac{1}{2}^+$	< 0.03	
6.49 ± 50	$\frac{7}{2}^-$	0.6	
6.92 ± 50	$\frac{5}{2}^+$	0.7	
7.53 ± 50	$\frac{3}{2}^+$	0.4	
8.13 ± 50	$\frac{3}{2}^-$	0.7	
8.43 ± 50	$\frac{5}{2}^-$	0.08	
8.67 ± 80	$\frac{7}{2}^+ + \frac{5}{2}^+$		
9.3 ± 100			
9.7 ± 100	$(\frac{5}{2}^+)$		
10.1 ± 200	$\frac{7}{2}^+$		
10.7 ± 200	$\frac{9}{2}^+$		
11.0 ± 100			
11.5 ± 200			

^a (1971KA56): $E_p = 185$ MeV.

^b (1974NO07).

^c From Table 11.19.

^d (1975RO27).

Angular distributions have been measured for $E_p = 19$ to 700 MeV [see (1968AJ02, 1975AJ02)] and recently at 65 MeV (1975RO27; see Table 11.24), 121.2 MeV (1976AN1C) and 650 and 800 MeV (1978IG1A, 1978WH1A, 1979BO1A). Observed states of ^{11}C are displayed in Table 11.24. See also (1975AJ02, 1975BE1Q, 1975IG1A, 1977HA1P, 1977MO1F, 1978AZ02, 1978IG1B, 1978SH1F, 1979WH1A), (1977HA1N [in ^{13}N , (1981AJ01)], and (1975GE1H, 1975RO29, 1976GR1G, 1976SH1K, 1976WA15, 1977KA07, 1977LU1A, 1978CO1J, 1978LU1B; theor.).

28. $^{12}\text{C}(\text{d}, \text{t})^{11}\text{C}$

$$Q_m = -12.464$$

At $E_d = 28$ MeV the t_0 angular distribution has been measured and a detailed comparison has been made with the results for the mirror reaction $^{12}\text{C}(\text{d}, ^3\text{He})^{11}\text{B}$ (1968GA13, 1966DE1C). At

$E_{\bar{a}} = 29$ MeV the t_0 angular distribution leads to spectroscopic factors $C^2S = 2.82$ or 3.97 depending on different sets of parameters for $^{11}\text{C}_{\text{g.s.}}$. $^{11}\text{C}^*(2.0, 4.32)$ are also populated (1978CO13). See also ^{14}N in (1981AJ01), and (1975AJ02).

$$29. \ ^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C} \quad Q_{\text{m}} = 1.856$$

Angular distributions have been measured at many energies to $E(^3\text{He}) = 217$ MeV [see (1968AJ02, 1975AJ02) for the earlier results] and at 11.0 MeV (1978CU02: α_0, α_1), 18, 20 and 22 MeV (1977AD07: α_0, α_1), at $E(^3\text{He}) = 33$ MeV (1976KA23: α_0, α_1), and at $E(^3\text{He}) = 35.6$ MeV (1978FO13: see Table 11.25), 82.1 MeV (1976TA12: $\alpha_0, \alpha_1, \alpha_{2+3}, \alpha_{4+5}$) and at 217 MeV (1978VA05: α_0, α_1). See also (1976ST1B: 60 MeV) and (1978CH1P: 132 MeV). At $E(^3\text{He}) \approx 217$ MeV (1975GE16, 1978VA05) also report the population of unresolved low-lying states, as well as the excitation of states with $E_x = 11.2, 12.4, 15.3, 23$ and (28) MeV. See Table 11.25 for a display of the results.

At $E(^3\text{He}) = 35.6$ MeV (1978FO13) find good fits by DWBA for strong $l = 1$ transitions, and reasonable agreement in the forward direction, as well as with $S_{\text{theor.}}$, for weak $l = 1$ transitions. Transitions involving $l = 0$ or 2 (and 3) are weak and the agreement with theory is poor. It is suggested that $^{11}\text{C}^*(8.10) [\frac{3}{2}^-]$ is predominantly a $p_{3/2}$ hole state coupled to $^{12}\text{C}^*(7.65) [0^+]$ (1978FO13).

Alpha- γ correlations have been studied for $E(^3\text{He}) = 4.7$ to 12 MeV: see, in particular, (1968EA03). Their results are summarized in Table 11.20 and are discussed in detail in reaction 22 of (1968AJ02). A measurement of the linear polarization of the 2.00 MeV γ -ray (together with knowledge of the τ_{m}) fixes $J^\pi = \frac{1}{2}^-$ for $^{11}\text{C}^*(2.00)$ (1968BL09). See also (1976RO1L, 1977MO1F) and ^{15}O in (1981AJ01).

$$30. \ ^{12}\text{C}(^6\text{Li}, ^7\text{Li})^{11}\text{C} \quad Q_{\text{m}} = -11.471$$

At $E(^6\text{Li}) = 36$ MeV the angular distributions involving $^7\text{Li}_{\text{g.s.}} + ^{11}\text{C}_{\text{g.s.}}$ and $^7\text{Li}_{0.48}^* + ^{11}\text{C}_{2.00}^*$ have been studied by (1973SC26).

$$31. \ ^{12}\text{C}(^{10}\text{B}, ^{11}\text{B})^{11}\text{C} \quad Q_{\text{m}} = -7.266$$

At $E(^{10}\text{B}) = 100$ MeV, angular distributions have been measured involving $^{11}\text{B}_{\text{g.s.}} + ^{11}\text{C}_{\text{g.s.}}$, $^{11}\text{B}_{\text{g.s.}} + ^{11}\text{C}_{2.00}$ and $(^{11}\text{C}_{\text{g.s.}} + ^{11}\text{B}_{2.12}) + (^{11}\text{B}_{\text{g.s.}} + ^{11}\text{C}_{2.00})$ [unresolved], as well as unresolved groups involving the first few states of ^{11}B and ^{11}C (1975NA15, 1975VO05, 1976NA09). See also (1975AJ02).

Table 11.25: Levels of ^{11}C from $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$ ^a

E_x (MeV \pm keV)	l ^d	S_{rel}				S 33 MeV ⁱ
		$E(^3\text{He}) = 16$ MeV ^f	24 MeV ^g	28 MeV ^g	35.6 MeV ^h	
0	1	1	1	1	1.00	3.06
1.999 ± 4 ^b	1	0.10	≤ 0.6	≤ 0.6	0.19	0.65
4.3188 ± 1.2 ^c	3 ^{d,h}	0.057	(0.04)	(0.06)	(0.031)	
4.8042 ± 1.2 ^c	1	0.11	0.22	0.22	0.13	
6.3392 ± 1.4 ^c	0	0.003 ^j	≤ 0.07	≤ 0.07	($\lesssim 0.2$)	
6.4782 ± 1.4 ^c	3 ^{d,h}	0.11 ^j	0.06	(0.06)	(0.21)	
6.9048 ± 1.4 ^c	2	0.018	(0.15)	(0.17)	(0.054)	
7.4997 ± 1.5 ^c	2	0.006 ^j	(0.07)	(0.09)	(0.046)	
8.1045 ± 1.7 ^c	1 ^e	0.017 ^{j,k}			(0.035)	
8.42	3 ^{e,h}	0.034 ^{j,l}			(0.041)	

^a See Table 11.20 for γ -decay work (1968EA03). Higher excited states are reported by (1978VA05): see text.

^b (1968EA03).

^c (1970BR23).

^d (1970BO34).

^e (1970FO05).

^f (1970GR08).

^g (1973FU02).

^h (1978FO13).

ⁱ (1976KA23). Polarized ^3He .

^j At $E(^3\text{He}) = 18$ MeV.

^k Assuming $J^\pi = \frac{3}{2}^-$.

^l Assuming $J^\pi = \frac{5}{2}^-$.

$$32. {}^{12}\text{C}({}^{12}\text{C}, {}^{13}\text{C}){}^{11}\text{C} \quad Q_{\text{m}} = -13.775$$

The angular distribution involving the ground state transitions has been studied at $E({}^{12}\text{C}) = 114$ MeV (1974AN36). See also (1979HE1E).

$$33. {}^{12}\text{C}({}^{14}\text{N}, {}^{15}\text{N}){}^{11}\text{C} \quad Q_{\text{m}} = -7.888$$

See (1967BI06, 1974AN36) and ${}^{15}\text{N}$ in (1981AJ01).

$$34. {}^{13}\text{C}(\text{p}, \text{t}){}^{11}\text{C} \quad Q_{\text{m}} = -15.186$$

At $E_{\text{p}} = 43.7$ to 50.5 MeV (1968CO26, 1968FL02) and at $E_{\bar{\text{p}}} = 49.6$ MeV (1974MA12) angular distributions of the tritons have been studied to ${}^{11}\text{C}^*(0, 2.00, 4.32, 4.80, 6.48, 6.90, 7.50)$ and to a $T = \frac{3}{2}$ state at $E_{\text{x}} = 12.47$ MeV [see Table 11.21] whose J^{π} is determined to be $\frac{1}{2}^{-}$ [it is thus the analog of ${}^{11}\text{Be}^*(0.32)$]. The state decays primarily by $\text{p} \rightarrow {}^{10}\text{B}^*(1.74)$. Alpha decay to ${}^7\text{Be}_{\text{g.s.}+0.4}^*$ is also observed (1968CO26). Angular distributions have also been measured for $E_{\text{p}} = 26.9$ to 43.1 MeV (1975MI01: $\text{t}_0 \rightarrow \text{t}_3$). At $E_{\text{p}} = 46.7$ MeV the $T = \frac{3}{2}$ state is also observed by (1974BE20) who, in addition, report the population of states with $E_{\text{x}} = 11.03 \pm 0.03$, 13.33 ± 0.06 , 13.90 ± 0.04 and 14.07 ± 0.04 MeV [$\Gamma = 300 \pm 60$, 270 ± 80 , 150 ± 50 and 135 ± 50 keV, respectively]. However, the $T = \frac{3}{2}$ state at $E_{\text{x}} = 12.16$ MeV reported by (1971WA21) in reactions 4, 8 and 22 is not observed by (1974BE20). See also ${}^{14}\text{N}$ in (1981AJ01).

$$35. {}^{14}\text{N}(\text{p}, \alpha){}^{11}\text{C} \quad Q_{\text{m}} = -2.922$$

Angular distributions have been reported at a number of energies in the range $E_{\text{p}} = 5.00$ to 44.3 MeV for the α_0 and α_1 groups: see (1975AJ02) for the earlier work and (1974HU02: 9.0 to 12.0 MeV; α_0). See also ${}^{15}\text{O}$ in (1981AJ01), (1976IN01; astrophys.), (1975CH1H; applied) and (1975KU1L; theor.).

$$36. {}^{14}\text{N}({}^{10}\text{B}, {}^{13}\text{C}){}^{11}\text{C} \quad Q_{\text{m}} = 1.140$$

This reaction has been studied at $E({}^{10}\text{B}) = 100$ MeV: see (1975NA15, 1976NA09) and ${}^{13}\text{C}$ in (1981AJ01).

$$37. {}^{16}\text{O}(\text{p}, {}^6\text{Li}){}^{11}\text{C} \quad Q_{\text{m}} = -22.185$$

See (1975AJ02).

$$38. {}^{16}\text{O}(\text{d}, {}^7\text{Li}){}^{11}\text{C} \quad Q_{\text{m}} = -17.159$$

At $E_{\text{d}} = 80$ MeV angular distributions involving ${}^{11}\text{C}^*(0, 2.0, 4.3 + 4.8, 6.3 + 6.5 + 6.9)$ have been measured (1978OE1A).

$$39. {}^{16}\text{O}(\alpha, {}^9\text{Be}){}^{11}\text{C} \quad Q_{\text{m}} = -24.310$$

At $E_{\alpha} = 42$ MeV, the angular distribution involving the two ground state transitions has been measured (1972RU03).

${}^{11}\text{N}$
(Fig. 4)

The ${}^{14}\text{N}({}^3\text{He}, {}^6\text{He}){}^{11}\text{N}$ reaction has been studied at $E({}^3\text{He}) = 70$ MeV. A ${}^6\text{He}$ group is observed which corresponds to a state in ${}^{11}\text{N}$ with an atomic mass excess of 25.23 ± 0.10 MeV and $\Gamma = 740 \pm 100$ keV. The cross section for forming this state is $0.5 \mu\text{b}/\text{sr}$ at 10° . The observed state is interpreted as being the $J^\pi = \frac{1}{2}^-$ mirror of ${}^{11}\text{Be}^*(0.32)$ because of its width; the $\frac{1}{2}^+$ mirror of ${}^{11}\text{Be}_{\text{g.s.}}$ would be expected to be much broader (1974BE20). The ${}^{11}\text{N}$ state is unbound with respect to decay into ${}^{10}\text{C} + \text{p}$ by 2.24 MeV. See also (1975BE31, 1975HU14, 1976IR1B, 1978GU10, 1979BE1H; theor.) and (1974DA27).

${}^{11}\text{O}, {}^{11}\text{F}, {}^{11}\text{Ne}$
(Not illustrated)

These nuclei have not been observed: see (1975BE31, 1976IR1B; theor.).

References

(Closed 01 July 1979)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1951BE13 W.E. Bennett, P.A. Roys and B.J. Toppel, Phys. Rev. 82 (1951) 20
- 1951BO45 C.K. Bockelman, D.W. Miller, R.K. Adair and H.H. Barschall, Phys. Rev. 84 (1951) 69
- 1951VA1A Van Patter, Buechner and Sperduto, Phys. Rev. 82 (1951) 248
- 1953EL12 M.M. Elkind, Phys. Rev. 92 (1953) 127
- 1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847
- 1955RA14 D.C. Ralph and F.E. Dunnam, Phys. Rev. 98 (1955) 249A
- 1957BA18 G.A. Bartholomew and P.J. Campion, Can. J. Phys. 35 (1957) 1347
- 1957BI84 H. Bichsel and T.W. Bonner, Phys. Rev. 108 (1957) 1025
- 1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
- 1959HI69 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 196
- 1959JO25 G.A. Jones, C.M.P. Johnson and D.H. Wilkinson, Phil. Mag. 4 (1959) 796
- 1959ME24 U. Meyer-Berkhout, K.W. Ford and A.E.S. Green, Ann. Phys. (N.Y.) 8 (1959) 119
- 1960BI08 O.M. Bilaniuk and J.C. Hensel, Phys. Rev. 120 (1960) 211
- 1960DA08 R.B. Day and M. Walt, Phys. Rev. 117 (1960) 1330
- 1960HI08 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) 75 (1960) 754
- 1961BA10 F.De S. Barros, P.D. Forsyth, A.A. Jaffe and I.J. Taylor, Proc. Phys. Soc. (London) 77 (1961) 853
- 1961DA16 E.A. Davis, F. Gabbard, T.W. Bonner and R. Bass, Nucl. Phys. 27 (1961) 448
- 1961FO07 D.B. Fossan, R.L. Walter, W.E. Wilson and H.H. Barschall, Phys. Rev. 123 (1961) 209
- 1961HI08 J. Hinds and R. Middleton, Proc. Phys. Soc. (London) 78 (1961) 81
- 1961JA11 A.N. James, Nucl. Phys. 24 (1961) 675
- 1962BR34 F.P. Brady and E.K. Warburton, Phys. Rev. 128 (1962) 790

1962ED02 R.D. Edge and G.A. Peterson, Phys. Rev. 128 (1962) 2750
1962GR07 L.L. Green, G.A. Stephens and J.C. Willmott, Proc. Phys. Soc. (London) 79 (1962) 1017
1962HI07 S. Hinds and R. Middleton, Nucl. Phys. 38 (1962) 114
1962OP03 T.R. Ophel, R.N. Glover and E.W. Titterton, Nucl. Phys. 33 (1962) 198
1962OV02 J.C. Overley and W. Whaling, Phys. Rev. 128 (1962) 315
1962PU01 D.J. Pullen, A.E. Litherland, S. Hinds and R. Middleton, Nucl. Phys. 36 (1962) 1
1963EA01 L.G. Earwaker, J.G. Jenkin and E.W. Titterton, Nucl. Phys. 42 (1963) 521
1963JE01 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Phys. Lett. 4 (1963) 142
1963ME08 M.K. Mehta, W.E. Hunt, H.S. Plendl and R.H. Davis, Nucl. Phys. 48 (1963) 90
1963OV02 J.C. Overley, Nucl. Phys. 49 (1963) 537
1964JE01 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Nucl. Phys. 50 (1964) 516
1965KA14 S. Kakigi, J. Phys. Soc. Jpn. 20 (1965) 1967
1965KE05 E.F. Kennedy and W.C. Miller, Nucl. Phys. 63 (1965) 634
1965OL03 J.W. Olness, E.K. Warburton, D.E. Alburger and J.A. Becker, Phys. Rev. 139 (1965) B512
1965OV1A Overley and Borchers, Nucl. Phys. 65 (1965) 156
1965RO07 M.L. Roush, A.A. Jaffe, A.S. Figuera and W.F. Hornyak, Nucl. Phys. 67 (1965) 577.
1966BR18 C.P. Browne and F.H. O'Donnell, Phys. Rev. 149 (1966) 767
1966CU02 R.Y. Cusson, Nucl. Phys. 86 (1966) 481
1966DE1C Detraz et al., Phys. Lett. 22 (1966) 638
1966GA19 A. Gallmann, F. Haas and N. Balaux, Phys. Rev. 151 (1966) 735
1966IS1A Isabelle, J. Phys. C1-17 (1966)
1966KI09 K.G. Kibler, Phys. Rev. 152 (1966) 932
1966MO09 F.P. Mooring, J.E. Monahan and C.M. Huddleston, Nucl. Phys. 82 (1966) 16
1966RA29 R.E. Rand, R. Frosch and M.R. Yearian, Phys. Rev. 144 (1966) 859; Erratum Phys. Rev. 148 (1966) 1246
1966SE03 R.E. Segel, P.P. Singh, S.S. Hanna and M.A. Grace, Phys. Rev. 145 (1966) 736.
1966SP02 E. Spamer, Z. Physik 191 (1966) 24
1966ST12 T. Stovall, J. Goldemberg and D.B. Isabelle, Nucl. Phys. 86 (1966) 225
1966WA10 E.K. Warburton, J.W. Olness, K.W. Jones, C. Chasman, R.A. Ristinen and D.H. Wilkinson, Phys. Rev. 148 (1966) 1072
1966YO1A Young et al., Nucl. Instrum. Meth. Phys. Res. 44 (1966) 109

- 1967BA53 R.M. Bahnsen and R.T. Carpenter, Phys. Rev. 164 (1967) 1235
- 1967BI06 J. Birnbaum, J.C. Overley and D.A. Bromley, Phys. Rev. 157 (1967) 787
- 1967BL22 R.S. Blake, E.B. Paul, C.H. Sinex and S.T. Emerson, Nucl. Phys. A102 (1967) 305
- 1967CA09 J.L. Campbell, W. Leiper, K.W.D. Ledingham and R.W.P. Drever, Nucl. Phys. A96 (1967) 279
- 1967CH34 V.I. Chuev, V.V. Davidov, A.A. Ogloblin and S.B. Sakuta, Ark. Fys. 36 (1967) 263
- 1967CR04 M.A. Crosby and J.C. Legg, Nucl. Phys. A95 (1967) 639
- 1967DE15 A.J. Deruytter and P. Pelfer, J. Nucl. Energy 21 (1967) 833
- 1967FI07 W. Fitz, R. Jahr and R. Santo, Nucl. Phys. A101 (1967) 449
- 1967PA19 P. Paul, N.G. Puttaswamy and D. Kohler, Phys. Rev. 164 (1967) 1332
- 1967SP02 E. Spamer and H. Artus, Z. Phys. 198 (1967) 445
- 1967TH05 G.E. Thomas, D.E. Blatchley and L.M. Bollinger, Nucl. Instrum. Meth. 56 (1967) 325
- 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
- 1968BE30 R.A.I. Bell, R.D. Gill, B.C. Robertson, J.S. Lopes and H.J. Rose, Nucl. Phys. A118 (1968) 481
- 1968BL09 R.S. Blake, G. Johnson, H. Laurent, J.P. Schapira and F. Picard, Nucl. Phys. A117 (1968) 561
- 1968CO09 M.N.H. Comsan, M.A. Farouk, A.A. El-Kamhawy, M.S.M. El-Tahawy and A.N. Lvov, Z. Phys. 212 (1968) 71.
- 1968CO26 S.W. Cosper, R.L. McGrath, J. Cerny, C.C. Maples, G.W. Goth and D.G. Fleming, Phys. Rev. 176 (1968) 1113
- 1968CR07 W.L. Creten, R.J. Jacobs and H.M. Ferdinande, Nucl. Phys. A120 (1968) 126
- 1968DA20 V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Yad. Fiz. 7 (1968) 758; Sov. J. Nucl. Phys. 7 (1968) 463
- 1968DE20 J.P. Deutsch, L. Grenacs, J. Lehmann, P. Lipnik and P.C. Macq, Phys. Lett. B28 (1968) 178
- 1968EA03 L.G. Earwaker and J.H. Montague, Nucl. Phys. A109 (1968) 507
- 1968FL02 D.G. Fleming, J. Cerny and N.K. Glendenning, Phys. Rev. 165 (1968) 1153
- 1968GA03 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 167 (1968) 957
- 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161
- 1968HI01 F. Hinterberger, G. Mairle, U. Schmidt-Rohr, P. Turek and G.J. Wagner, Nucl. Phys. A106 (1968) 161
- 1968JA08 R. Jambunathan and R.K. Hobbie, Phys. Rev. 172 (1968) 1065

1968SC18 H.F. Schaefer III, R.A. Klemm and F.E. Harris, Phys. Rev. 176 (1968) 49
 1968ST12 R.A. Stryk and J.M. Blair, Phys. Rev. 169 (1968) 767
 1969BR30 O.D. Brill, A.D. Vongai and A.A. Ogloblin, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 615; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 567
 1969CA1A R.R. Carlson and H.W. Wyborny, Phys. Rev. 178 (1969) 1529
 1969GA11 P. Gaillard, R. Bouche, L. Feuveys, M. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.R. Pizzi, Nucl. Phys. A131 (1969) 353
 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, Phys. Rev. 178 (1969) 1551
 1969SC34 H.F. Schaefer, III, R.A. Klemm and F.E. Harris, Phys. Rev. 181 (1969) 137.
 1969SI12 J.D. Silverstein and G.H. Herling, Phys. Rev. 181 (1969) 1512
 1969SU03 O. Sundberg and G. Tibell, Ark. Fys. 39 (1969) 397
 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
 1969VO07 W. Von Oertzen, H.H. Gutbrod, U.C. Voos and R. Bock, Nucl. Phys. A133 (1969) 101
 1969VO10 U.C. Voos, W. von Oertzen and R. Bock, Nucl. Phys. A135 (1969) 207
 1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977
 1969WA23 B.A. Watson, R.E. Segel, J.J. Kroepfl and P.P. Singh, Phys. Rev. 187 (1969) 1351
 1969WO03 G. Wolber, H. Figger, R.A. Haberstroh and S. Penselin, Phys. Lett. A29 (1969) 461
 1970AL21 D.E. Alburger and G.A.P. Engelbertink, Phys. Rev. C2 (1970) 1594
 1970AS10 A. Asami and M.C. Moxon, J. Nucl. Energy 24 (1970) 85.
 1970AU02 D.L. Auton, Nucl. Phys. A157 (1970) 305
 1970BA05 C.J. Batty and C. Tschalar, Nucl. Phys. A143 (1970) 151
 1970BO07 W. Bohne, M. Hagen, H. Homeyer, H. Lettau, K.H. Maier, H. Morgenstern and J. Scheer, Phys. Rev. Lett. 24 (1970) 1028
 1970BO17 F. Boreli, R. Little and A. Rego, Fiz. (Yugoslavia) 2 (1970) 19
 1970BO34 W. Bohne, J. Bommer, H. Fuchs, K. Grabisch, M. Hagen, H. Homeyer, U. Janetzki, H. Lettau, H. Morgenstern, G. Roschert et al., Nucl. Phys. A157 (1970) 593
 1970BR23 C.P. Browne, G. Maille, R. Tarara and J.R. Duray, Nucl. Phys. A153 (1970) 289
 1970FO05 H.T. Fortune, J.R. Comfort, J.V. Maher and B. Zeidman, Phys. Rev. C2 (1970) 425
 1970GO04 D.R. Goosman, E.G. Adelberger and K.A. Snover, Phys. Rev. C1 (1970) 123
 1970GO11 D.R. Goosman and R.W. Kavanagh, Phys. Rev. C1 (1970) 1939
 1970GR08 T.H. Gray, H.T. Fortune, W. Trost and N.R. Fletcher, Nucl. Phys. A144 (1970) 129

- 1970KU09 H.M. Kuan, M. Hasinoff, W.J. O'Connell and S.S. Hanna, Nucl. Phys. A151 (1970) 129
- 1970ME17 H.A. Medicus, E.M. Bowey, D.B. Gayther, B.H. Patrick and E.J. Winhold, Nucl. Phys. A156 (1970) 257
- 1970NE03 D.O. Nellis, W.E. Tucker and I.L. Morgan, Phys. Rev. C1 (1970) 847
- 1970NE05 R.K. Nesbet, Phys. Rev. Lett. 24 (1970) 1155.
- 1970NU02 F. Nusslin and P. Braun-Munzinger, Z. Phys. 240 (1970) 217
- 1970SO1A Sowerby, Patrick, Uttley and Diment, J. Nucl. Energy 24 (1970) 323
- 1971AL07 D.E. Alburger and D.H. Wilkinson, Phys. Rev. C3 (1971) 1492
- 1971BA72 K. Battleson and D.K. McDaniels, Phys. Rev. C4 (1971) 1601
- 1971BI12 H.G. Bingham, K.W. Kemper and N.R. Fletcher, Nucl. Phys. A175 (1971) 374
- 1971BR41 C.P. Browne and H. Stocker, Phys. Rev. C4 (1971) 1481
- 1971CO07 J.R. Comfort, H.T. Fortune, J.V. Maher and B. Zeidman, Phys. Rev. C3 (1971) 1086
- 1971HA25 S.S. Hanna, K. Nagatani, W.R. Harris and J.W. Olness, Phys. Rev. C3 (1971) 2198
- 1971HIZF F. Hibou, Thesis, Univ. Louis Pasteur, Strasbourg (1971)
- 1971KA56 J. Kallne and E. Hagberg, Phys. Scr. 4 (1971) 151
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971LA10 R.O. Lane, S.L. Hausladen, J.E. Monahan, A.J. Elwyn, F.P. Mooring and A. Langsdorf, Jr., Phys. Rev. C4 (1971) 380
- 1971LI11 M. Liu, W. von Oertzen, J.C. Jacmart, F. Pougheon, M. Riou, J.C. Roynette and C. Stephan, Nucl. Phys. A165 (1971) 118
- 1971PA10 B.H. Patrick, H.A. Medicus, G.K. Mehta, E.M. Bowey and D.B. Gayther, Phys. Lett. B34 (1971) 488
- 1971SC16 G. Schmidt, J. Mosner and J. Schintlmeister, Nucl. Phys. A173 (1971) 449
- 1971SQ02 G.T.A. Squier, A.R. Johnston, J.H.P.C. Megaw, R.J. Griffiths and F.G. Kingston, Nucl. Phys. A167 (1971) 465
- 1971WA21 B.A. Watson, C.C. Chang and M. Hasinoff, Nucl. Phys. A173 (1971) 634
- 1972AJ01 F. Ajzenberg-Selove, R.F. Casten, O. Hansen and T.J. Mulligan, Phys. Lett. B40 (1972) 205
- 1972BA52 G. Bassani, N. Saunier, B.M. Traore, J. Raynal, A. Foti and G. Pappalardo, Nucl. Phys. A189 (1972) 353
- 1972HA04 S.L. Hausladen, R.O. Lane and J.E. Monahan, Phys. Rev. C5 (1972) 277
- 1972PA1C Parker, Astrophys. J. 175 (1972) 261

- 1972RE1C Rendic, Conf. Nucl. Struct. Study with Neutrons, Budapest, 1972 (Budapest Central Res. Inst., 1972) 28; Phys. Abs. 67324 (1972)
- 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430.
- 1972SC03 U.C. Schlotthauer-Voos, H.G. Bohlen, W. von Oertzen and R. Bock, Nucl. Phys. A180 (1972) 385
- 1972SZ02 J. Szabo, J. Csikai and M. Varnagy, Nucl. Phys. A195 (1972) 527
- 1972VA02 L. van der Zwan and K.W. Geiger, Nucl. Phys. A180 (1972) 615
- 1973CO05 J.M. Cox, H.D. Knox, R.O. Lane and R.W. Finlay, Nucl. Phys. A203 (1973) 89
- 1973FO02 H.T. Fortune, H.G. Bingham, J.D. Garrett and R. Middleton, Phys. Rev. C7 (1973) 136
- 1973FU02 H. Fuchs and H. Oeschler, Nucl. Phys. A202 (1973) 396
- 1973GO09 D.R. Goosman and R.W. Kavanagh, Phys. Rev. C7 (1973) 1717
- 1973HA64 S.L. Hausladen, C.E. Nelson and R.O. Lane, Nucl. Phys. A217 (1973) 563
- 1973HO43 K.R. Hogstrom, B.W. Mayes, L.Y. Lee, J.C. Allred, C. Goodman, G.S. Mutchler, C.R. Fletcher and G.C. Phillips, Nucl. Phys. A215 (1973) 598
- 1973HU09 R.J. Hughes and E.G. Muirhead, Nucl. Phys. A215 (1973) 147
- 1973KE13 H. Kelleter, G. Hrehuss and C. Mayer-Boricke, Nucl. Phys. A210 (1973) 502
- 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
- 1973PA14 Y.S. Park, A. Niiler and R.A. Lindgren, Phys. Rev. C8 (1973) 1557; Erratum Phys. Rev. C9 (1974) 1673
- 1973SC26 P. Schumacher, N. Ueta, H.H. Duhm, K.-I. Kubo and W.J. Klages, Nucl. Phys. A212 (1973) 573
- 1973TA30 K. Takahashi, M. Yamada and T. Kondoh, At. Data Nucl. Data Tables 12 (1973) 101
- 1973TO16 L. Tomlinson, At. Data Nucl. Data Tables 12 (1973) 179
- 1973VA1A Vasileva et al., Sov. J. Nucl. Phys. 16 (1973) 257
- 1974AN01 H.J. Annegarn, D.W. Mingay and J.P.F. Sellschop, Phys. Rev. C9 (1974) 419
- 1974AN36 N. Anyas-Weiss, J.C. Cornell, P.S. Fisher, P.N. Hudson, A. Menchaca-Rocha, D.J. Millener, A.D. Panagiotou, D.K. Scott, D. Strottman, D.M. Brink et al., Phys. Rept. 12 (1974) 201
- 1974AS04 O. Aspelund, D. Ingham, A. Djaloeis, H. Kelleter and C. Mayer-Boricke, Phys. Lett. B50 (1974) 441
- 1974BA15 G.C. Ball, J.G. Costa, W.G. Davies, J.S. Forster, J.C. Hardy and A.B. McDonald, Phys. Lett. B49 (1974) 33

- 1974BE20 W. Benenson, E. Kashy, D.H. Kong-A-Siou, A. Moalem and H. Nann, Phys. Rev. C9 (1974) 2130
- 1974BO15 W. Bohne, C.K. Gelbke, P. Braun-Munzinger, W. Grochulski, H.L. Harney and H. Oeschler, Nucl. Phys. A222 (1974) 117
- 1974BO35 I.I. Bondouk, Atomkernenergie 23 (1974) 219
- 1974BO37 I.I. Bondouk, Atomkernenergie 23 (1974) 283
- 1974BU1A Bulow and Forkman, IAEA, STI/DOC/10/156 (1974) 475
- 1974CE06 J. Cerny, R.B. Weisenmiller, N.A. Jelley, K.H. Wilcox and G.J. Wozniak, Phys. Lett. B53 (1974) 247.
- 1974CH1L Chiang et al., Acta Phys. Sin. (China) 23 (1974) 362
- 1974CH58 I.P. Chernov, B.I. Kuznetsov, V.V. Kozyr and V.A. Matusевич, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2530; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 58
- 1974CO1C Coates, Hunt and Uttley, NSRD, IAEA STI/PUB/371 (1974) 129
- 1974DA27 S. Dahlgren and P. Grafstrom, Phys. Scr. 10 (1974) 104
- 1974DE01 W. Del Bianco, P. Boucher, S. Kundu and B. Rouben, Can. J. Phys. 52 (1974) 92.
- 1974DE1E De Jager, De Vries and De Vries, At. Data Nucl. Data Tables 14 (1974) 479
- 1974DE39 W. Del Bianco, S. Kundu and B. Rouben, Nucl. Phys. A232 (1974) 333
- 1974DM01 T.A. Dmitrieva, V.M. Lebedev, V.G. Markman, A.V. Spasskii and I.B. Teplov, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2567; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 91
- 1974EN10 R. Engfer, H. Schneuwly, J.L. Vuilleumier, H.K. Walter and A. Zehnder, At. Data Nucl. Data Tables 14 (1974) 509
- 1974FI17 D.J.S. Findlay, S.N. Gardiner, J.L. Matthews and R.O. Owens, J. Phys. A7 (1974) L157
- 1974FR02 E. Friedland, H.W. Alberts and J.C. van Staden, Z. Phys. 267 (1974) 97
- 1974FU11 H. Fuchs, K. Grabisch, D. Hilscher, U. Jahnke, H. Kluge, T. Masterson and H. Morgenstern, Nucl. Phys. A234 (1974) 61
- 1974HA1C Hanna, Int. Conf. Nucl. Struct. Spectrosc., Pt. II, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 249
- 1974HA1H Hansen, Int. Conf. Nucl. Struct. Spectrosc., Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 662
- 1974HU02 G.A. Huttlin and A.A. Rollefson, Phys. Rev. C9 (1974) 576
- 1974HY01 M. Hyakutake, M. Sonoda, A. Katase, Y. Wakuta, M. Matoba, H. Tawara and I. Fujita, J. Nucl. Sci. Tech. (Japan) 11 (1974) 407
- 1974KA15 E. Kashy, W. Benenson and J.A. Nolen, Jr., Phys. Rev. C9 (1974) 2102

- 1974KI1A King et al., Bull. Amer. Phys. Soc. 19 (1974) 432
- 1974LO1B Lorenzen and Brune, IAEA, STI/DOC/10/156 (1974) 325
- 1974MA12 J.A. Macdonald, J. Cerny, J.C. Hardy, H.L. Harney, A.D. Bacher and G.R. Plattner, Phys. Rev. C9 (1974) 1694
- 1974NO07 J.A. Nolen, Jr., G. Hamilton, E. Kashy and I.D. Proctor, Nucl. Instrum. Meth. 115 (1974) 189
- 1974RO31 E. Roeckl, P.F. Dittner, C. Detraz, R. Klapisch, C. Thibault and C. Rigaud, Phys. Rev. C10 (1974) 1181
- 1974SHYR V.S. Shirley and C.M. Lederer, LBL-3450 (1974)
- 1974ST1F Stewart, NSRD, IAEA STI/PUB/371 (1974) 149
- 1974TA1E Talmi, Proc. Int. Conf. Nucl. Struct. Spectrosc., Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 696
- 1974WA1C Walecka, Int. Conf. Nucl. Struct. Spectrosc., Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 438
- 1974WO1C Wozniak, LBL 2999 (1974)
- 1975AB1D Abdul-Magd, Prog. Theor. Phys. 53 (1975) 997
- 1975AD04 J.-O. Adler, G.G. Jonsson and K. Lindgren, Nucl. Phys. A239 (1975) 440
- 1975AJ02 F. Ajzenberg-Selove, Nucl. Phys. A248 (1975) 1
- 1975AL19 J.M. Alexander, H. Delagrangue and A. Fleury, Phys. Rev. C12 (1975) 149
- 1975AL1B Almond, Phys. Rept. C17 (1975) 1
- 1975AR14 A.G. Artyukh, V.V. Volkov, G.F. Gridnev and V.L. Mikheev, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2; Bull. Acad. Sci. USSR Phys. Ser. 39 (1975) 1
- 1975AR19 E.D. Arthur, D.M. Drake and I. Halpern, Phys. Rev. Lett. 35 (1975) 914
- 1975AR1E Arnould and Norgaard, Astron. Astrophys. 42 (1976) 55
- 1975AU1C August et al., Bull. Amer. Phys. Soc. 20 (1975) 1181
- 1975AZ01 G. Azuelos and J.E. Kitching, Phys. Rev. C12 (1975) 563
- 1975BA66 L.H. Batist, V.D. Vitman, V.P. Koptev, M.M. Makarov, A.A. Naberezhnov, V.V. Nelyubin, G.Z. Obrant, V.V. Sarantsev and G.V. Scherbakov, Nucl. Phys. A254 (1975) 480
- 1975BE1Q Beurty, in LASL, AIP Conf. Proc. 26 (1975) 653
- 1975BE28 H. Behrens, M. Kobelt, L. Szybisz and W.-G. Thies, Nucl. Phys. A246 (1975) 317
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BH03 R.S. Bhalerao and Y.R. Waghmare, Pramana 5 (1975) 154

- 1975BI07 T. Biro, S. Sudar, Z. Miligy, Z. Dezso and J. Csikai, *J. Inorg. Nucl. Chem.* 37 (1975) 1583
- 1975BL10 J.S. Blair, M.P. Baker and H.S. Sherif, *Phys. Lett.* B60 (1975) 25
- 1975BO1C Bondouk, Asfour and Machali, *Ann. Phys.* 32 (1975) 255
- 1975BO20 I.I. Bondouk, *Ann. Phys. (Leipzig)* 32 (1975) 202
- 1975BO21 I.I. Bondouk, *Ann. Phys. (Leipzig)* 32 (1975) 119
- 1975BR1F Brown, *At. Data Nucl. Data Tables* 15 (1975) 111
- 1975CA08 J.F. Cavaignac, S. Jang and D.H. Worledge, *Nucl. Phys.* A243 (1975) 349
- 1975CH1H Christman, Finn, Karlstrom and Wolf, *Int. J. App. Rad. Isotopes* 26 (1975) 435
- 1975CO06 B. Coupat, P.Y. Bertin, D.B. Isabelle, P. Vernin, A. Gerard, J. Miller, J. Morgenstern, J. Picard and B. Saghai, *Phys. Lett.* B55 (1975) 286
- 1975DO1D Donnelly and Walecka, *Ann. Rev. Nucl. Sci* 25 (1975) 329
- 1975DO1F Donnelly, in *LASL, AIP Conf. Proc.* 26 (1975) 454
- 1975DU11 P. Duck, W. Treu, W. Galster, E. Haindl, F. Siller and H. Voit, *Nucl. Phys.* A251 (1975) 344
- 1975EP02 B.W. Epperlein, O. Lutz and A. Schwenk, *Z. Naturforsch.* 30A (1975) 955
- 1975FE1A Feather, *Commun. R. Soc. Edinb. (Phys. Sci.)* 2 (1975) 19
- 1975FO19 W.A. Fowler, G.R. Caughlan and B.A. Zimmerman, *Ann. Rev. Astron. Astrophys.* 13 (1975) 69
- 1975FR1C Friesenhahn et al., *Nucl. Cross Sections and Tech., Pt. I, Washington, D.C., 1975 (Wash., D.C., Nat. Bur. Standards 1975)* p. 232; *Phys. Abs.* 26755 (1977)
- 1975GE16 E. Gerlic, H. Langevin-Joliot, P. Roos, J. Van de Wiele, J.P. Didelez and G. Duhamel, *Phys. Rev.* C12 (1975) 2106
- 1975GE1H Geramb and Hefter, in *Jul-Conf-16* (1975) 93
- 1975HA1L Hale, *Nucl. Cross Sections and Tech., Pt. I, Washington, D.C., 1975 (Wash., D.C., Nat. Bur. Standards 1975)* p. 302; *Phys. Abs.* 23041 (1977)
- 1975HA1N Hall and Engvol, *Astrophys. J.* 197 (1975) 513
- 1975HU13 J. Hufner, H.J. Pirner and M. Thies, *Phys. Lett.* B59 (1975) 215
- 1975HU14 J. Hufner, K. Schafer and B. Schurmann, *Phys. Rev.* C12 (1975) 1888
- 1975HU1D Hufner, *Phys. Rept.* C21 (1975) 1
- 1975IG1A Igo, in *Lasl, AIP Conf. Proc.* 26 (1975) 63
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, *Phys. Rev.* C11 (1975) 2049
- 1975KA02 P.T. Kan, G.A. Peterson, D.V. Webb, S.P. Fivozinsky, J.W. Lightbody, Jr. and S. Penner, *Phys. Rev.* C11 (1975) 323

- 1975KA1G Karol, Yester, Klobuchar and Caretto, Phys. Lett. B58 (1975) 489
- 1975KR14 S. Krol and S. Szpikowski, Acta Phys. Pol. B6 (1975) 409
- 1975KU1L Kurath, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 439
- 1975LA08 G.P. Lamaze, A.D. Carlson and M.M. Meier, Nucl. Sci. Eng. 56 (1975) 94
- 1975LU07 V.K. Lukyanov and A.I. Titov, Yad. Fiz. 22 (1975) 83; Sov. J. Nucl. Phys. 22 (1976) 40
- 1975MA1E Matthews, in LASL, AIP Conf. Proc. 26 (1975) 437
- 1975MA41 G. Mairle and G.J. Wagner, Nucl. Phys. A253 (1975) 253
- 1975MC19 M.W. McNaughton, N.S.P. King, F.P. Brady and J.L. Ullmann, Nucl. Instrum. Meth. Phys. Res. 129 (1975) 241
- 1975ME1E Meneguizzi and Reeves, Astron. Astrophys. 40 (1975) 91, 99
- 1975MI01 S. Micheletti, M. Pignanelli and P. Guazzoni, Phys. Rev. C11 (1975) 64
- 1975MI12 D.J. Millener and D. Kurath, Nucl. Phys. A255 (1975) 315
- 1975NA15 K.G. Nair, H. Voit, C.W. Towsley, M. Hamm, J.D. Bronson and K. Nagatani, Phys. Rev. C12 (1975) 1575
- 1975OS01 A. Osman, Nuovo Cim. A25 (1975) 571
- 1975PA1D Pak, Pocs and Tarasov, Yad. Fiz. 21 (1975) 520
- 1975PE1A R.J. Peterson, C.S. Zaidins, M.J. Fritts, N.A. Roughton and C.J. Hansen, Ann. Nucl. Energy 2 (1975) 503
- 1975PI1A Picklesimer and Walker, Bull. Amer. Phys. Soc. 20 (1975) 690
- 1975PR1B Preszler, Kish, Iezniak and Webber, Bull. Amer. Phys. Soc. 20 (1975) 711
- 1975PU01 R.J. Puigh, K.W. Kemper, G.E. Moore and R.L. White, Nucl. Phys. A237 (1975) 1
- 1975PU02 F. Puhlhofer, W. Pfeffer, B. Kohlmeyer and W.F.W. Schneider, Nucl. Phys. A244 (1975) 329
- 1975RA21 G.M. Raisbeck, P. Boerstling, R. Klapisch and T.D. Thomas, Phys. Rev. C12 (1975) 527
- 1975RE04 W. Reisdorf, Nucl. Phys. A242 (1975) 406
- 1975RO18 D. Royer, J. Mougey, M. Priou, M. Bernheim, A. Bussiere, A. Gillebert and G.J. Wagner, Phys. Rev. C12 (1975) 327
- 1975RO27 P.G. Roos, S.M. Smith, V.K.C. Cheng, G. Tibell, A.A. Cowley and R.A.J. Riddle, Nucl. Phys. A255 (1975) 187
- 1975RO29 E. Rost and J.R. Shepard, Phys. Lett. B59 (1975) 413
- 1975SA1C Saltmarsh and Smith, Bull. Amer. Phys. Soc. 20 (1975) 562
- 1975SC05 H. Schier and B. Schoch, Lett. Nuovo Cim. 12 (1975) 334

- 1975SC1N Schiffer, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 329
- 1975SE03 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 151
- 1975SE04 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 159
- 1975SE1H Sealock, Thesis Univ. Oregon (1975); Phys. Abs. 25118 (1976)
- 1975SI18 R.R. Silbar, Phys. Rev. C12 (1975) 341
- 1975TA1C Tauscher, in LASL, AIP Conf. Proc. 26 (1975) 541
- 1975TH08 C. Thibault, R. Klapisch, C. Rigaud, A.M. Poskanzer, R. Prieels, L. Lessard and W. Reisdorf, Phys. Rev. C12 (1975) 644
- 1975TR1A Trimble, Rev. Mod. Phys. 47 (1975) 877
- 1975VA19 O.I. Vasileva, V.M. Lebedev, A.V. Spasskii, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 22 (1975) 446; Sov. J. Nucl. Phys. 22 (1976) 229
- 1975VO05 H. Voit, M. Hamm, K.G. Nair, C. Towsley, A.D. Bacher, K. Nagatani and F. Siller, Phys. Lett. B58 (1975) 152
- 1975VO1B Von Oertzen and Bohlen, Phys. Rept. 19 (1975) 1
- 1975WI1E Wilkinson, Proc. Int. Symp. on Inter. Studies in Nucl., Germany, 1975 (Amsterdam, North-Holland, 1975) 147
- 1975YE1A Yergin et al., Bull. Amer. Phys. Soc. 20 (1975) 661
- 1975ZA06 N.I. Zaika and P.L. Shmarin, Ukr. Fiz. Zh. 20 (1975) 238; Phys. Abs. 45405 (1975)
- 1975ZI1B D. Zieminska, Nucl. Phys. A242 (1975) 461
- 1975ZW01 Zwieglinski, Saganek, Sledzinska and Wilhelmi, Nucl. Phys. A250 (1975) 93
- 1976AJ04 F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1
- 1976AL1G Alkhazov et al., in Contributions to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) p. 158
- 1976AN1C Anderson, Shepard and Comfort, Bull. Amer. Phys. Soc. 21 (1976) 979
- 1976AU1B Audouze and Lequeux, Astron. Astrophys. 49 (1976) 133
- 1976AU1C Audouze, Meneguzzi and Reeves, in Spallation Nucl. Reactions & Their Applic., Shen/Marker Eds. (D. Reidel, Holland 1976) 113
- 1976BA1R Bacher et al., Bull. Amer. Phys. Soc. 21 (1976) 983
- 1976BE1E Behrens and Lzybisz, Phys. Data (1976) (ZAED)
- 1976BE1H Berman, Preprint UCRL-78482 (1976)
- 1976BE1K Bertini, in Spallation Nucl. Reactions and Their Appl., Eds., Shen and Merker (D. Reidel, The Netherlands, 1976) p. 27

- 1976BH02 R.K. Bhowmik, C.C. Chang, J.-P. Didelez and H.D. Holmgren, *Phys. Rev. C*13 (1976) 2105
- 1976BI1E Birchall et al., in *Polariz.*, Zurich, 1975 (Birkhauser Verlag, 1976) 630
- 1976BO08 V.I. Bogatin, V.F. Litvin, O.V. Lozhkin, N.A. Perfilov and Y.P. Yakovlev, *Nucl. Phys. A*260 (1976) 446
- 1976BO1E Boesgaard, *Publ. Astron. Soc. Pac.* 88 (1976) 353
- 1976BO1L Borza et al., *Rev. Roum. Phys.* 21 (1976) 431
- 1976BR26 F. Brut and S. Jang, *Phys. Rev. C*14 (1976) 1638
- 1976BU16 M. Buenerd, C.K. Gelbke, B.G. Harvey, D.L. Hendrie, J. Mahoney, A. Menchaca-Rocha, C. Olmer and D.K. Scott, *Phys. Rev. Lett.* 37 (1976) 1191
- 1976CA1H Cammarata and Donnelly, *Meson-Nucl. Phys.*, 1976 (AIP, 1976) 618
- 1976CA1J Carlson, *Bull. Amer. Phys. Soc.* 21 (1976) 1004
- 1976CH1E Chang, Didelez and Wu, *Bull. Amer. Phys. Soc.* 21 (1976) 965
- 1976CO1B Colgate, Audouze and Fowler, *OAP-458* (1976)
- 1976CO1M Coupat et al., *Meson-Nucl. Phys.*, 1976 (AIP, 1976) 332
- 1976DA15 S.E. Darden, G. Murillo and S. Sen, *Nucl. Phys. A*266 (1976) 29
- 1976DA1F Davis, in *Spallation Nucl. Reactions & Their Appl.*, Eds., Shen and Merker (D. Reidel Pub. Co., Netherlands, 1976) 207
- 1976DE15 R. de Swiniarski, F.G. Resmini, C. Glashausser and A.D. Bacher, *Helv. Phys. Acta* 49 (1976) 227
- 1976DE1D Deloff and Law, *Meson-Nucl. Phys.*, 1976 (AIP, 1976) 544, 546
- 1976DE34 A.E. Denisov, R.P. Kolalis, V.S. Sadkovskii and G.A. Feofilov, *Yad. Fiz.* 24 (1976) 249; *Sov. J. Nucl. Phys.* 24 (1976) 129
- 1976DI01 V. di Napoli, G. Rosa, F. Salvetti, M.L. Terranova, H.G. de Carvalho, J.B. Martins and O.A.P. Tavares, *J. Inorg. Nucl. Chem.* 38 (1976) 1
- 1976DI05 J.-P. Didelez, C.C. Chang, N.S. Chant, H.D. Holmgren, R.I. Steinberg and J.R. Wu, *Phys. Rev. C*13 (1976) 1388
- 1976DO1G T.W. Donnelly and J.D. Walecka, *Nucl. Phys. A*274 (1976) 368
- 1976EP1A Epsteain, Arnett and Schramm, *Astrophys. J. Suppl.* 31 (1976) 111
- 1976FU06 G.H. Fuller, *J. Phys. Chem. Ref. Data* 5 (1976) 835
- 1976GAYV D.I. Garber and R.R. Kinsey, *BNL 325, Vol. 2* (1976)
- 1976GI1C Ginzburg and Ptjskin, *Rev. Mod. Phys.* 48 (1976) 161
- 1976GR1F Griggs et al., *Bull. Amer. Phys. Soc.* 21 (1976) 514
- 1976GR1G Grabowski, Fleming and Vogt, *Can. J. Phys.* 54 (1976) 870

- 1976HA1F Hainebach, Schramm and Blake, *Astrophys. J.* 205 (1976) 920
- 1976HE1H H.H. Heckman and P.J. Lindstrom, *Phys. Rev. Lett.* 37 (1976) 56
- 1976HI05 M.D. High and B. Cujec, *Nucl. Phys. A*259 (1976) 513
- 1976IN01 P.D. Ingalls, J.S. Schweitzer, B.D. Anderson and M. Rios, *Phys. Rev. C*13 (1976) 524
- 1976IN04 T. Inoue and S. Tanaka, *J. Inorg. Nucl. Chem. (GB)* 38 (1976) 1425
- 1976IR1B Irvine et al., *Ann. Phys.* 102 (1976) 129
- 1976KA1F Karol, *Meson-Nucl. Phys.*, 1976 (AIP, 1976) 305
- 1976KA23 O. Karban, A.K. Basak, J.B.A. England, G.C. Morrison, J.M. Nelson, S. Roman and G.G. Shute, *Nucl. Phys. A*269 (1976) 312
- 1976KI1D King et al., in *Lowell Conf.*, CONF-760715-P2 (1976) 1344
- 1976KN04 U. Kneissl, K.H. Leister, H.O. Neidel and A. Weller, *Nucl. Phys. A*264 (1976) 30
- 1976LA1C Lane, White and Knox, in *Lowell Conf.*, CONF-760715-P2 (1976) 1042
- 1976LE1F Lefort, *Rept. Prog.* 39 (1976) 129
- 1976LE1K Lebedev et al., in *Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct.*, Vol. 1, JINR D-9682 (1976) 98
- 1976LI1H Liskien, in *Lowell Conf.*, CONF-760715-P2 (1976) p. 1110
- 1976MA1J Malik and Nath, in *Lowell Conf.*, Conf-760715-P2 (1976) 1280
- 1976MA34 J.L. Matthews, D.J.S. Findlay, S.N. Gardiner and R.O. Owens, *Nucl. Phys. A*267 (1976) 51
- 1976MC1F McDonald et al., *Bull. Amer. Phys. Soc.* 21 (1976) 775
- 1976MI15 K. Min, P. Stoler, S. Trentalange, E.J. Winhold, P.F. Yergin, A.M. Bernstein, W. Turchinets and K.S.R. Sastry, *Phys. Rev. C*14 (1976) 807
- 1976MO17 J. Mougey, M. Bernheim, A. Bussiere, A. Gillebert, Phan Xuan Ho, M. Priou, D. Royer, I. Sick and G.J. Wagner, *Nucl. Phys. A*262 (1976) 461
- 1976NA09 K.G. Nair, K. Nagatani and H. Voit, *Phys. Rev. Lett.* 36 (1976) 1293
- 1976NA11 M.N. Namboodiri, E.T. Chulick and J.B. Natowitz, *Nucl. Phys. A*263 (1976) 491
- 1976NO1D Noble, *Meson-Nucl. Phys.-1976 (AIP, 1976)* 221
- 1976OS04 R. Ost, N.E. Sanderson, S. Mordechai, J.B.A. England, B.R. Fulton, J.M. Nelson and G.C. Morrison, *Nucl. Phys. A*265 (1976) 142
- 1976PH01 D.-L. Pham, *J. Phys. Lett.* 37 (1976) L-67
- 1976RA1C Raisbeck and Yiou, *Spallation Nucl. Reactions & Their Appl.*, Eds., Shen and Merker (D. Reidel, Netherlands, 1976) 83
- 1976RO12 C.T. Roche, R.G. Clark, G.J. Mathews and V.E. Viola, Jr., *Phys. Rev. C*14 (1976) 410
- 1976RO1J Ross and Aller, *Science* 191 (1976) 1223

- 1976RO1L Roman, in *Polariz.*, Zurich, 1975 (Birkhauser Verlag, 1976) 255
- 1976RO1N A. Robertson, T.J. Kennett and W.V. Prestwich, *Phys. Rev. C*13 (1976) 1552
- 1976RO1Q Roughton et al., *Astrophys. J.* 205 (1976) 302
- 1976SE06 R.M. Sealock and J.C. Overley, *Phys. Rev. C*13 (1976) 2149
- 1976SE1E Sealock, *Bull. Amer. Phys. Soc.* 21 (1976) 538
- 1976SH1K Shepard, Kunz, Sparrow and Rose, *Bull. Amer. Phys. Soc.* 21 (1976) 519
- 1976SI1A Siemssen, *Proc. Dubna Conf. JINR D-992C*, Vol. 2 (1976) 106
- 1976SI1C Simpson, *Bull. Amer. Phys. Soc.* 21 (1976) 602
- 1976SL2A I. Slaus, in *Lowell Conf.*, CONF-760715-P1 (1976) 272
- 1976SM07 A.R. Smith and R.H. Thomas, *Nucl. Instrum. Meth. Phys. Res.* 137 (1976) 459
- 1976ST12 R.G. Stokstad, Z.E. Switkowski, R.A. Dayras and R.M. Wieland, *Phys. Rev. Lett.* 37 (1976) 888
- 1976ST1B Stahel, Jahn, Wozniak and Cerny, *Bull. Amer. Phys. Soc.* 21 (1976) 1295
- 1976TA12 T. Tanabe, K. Koyama, M. Yasue, H. Yokomizo, K. Sato, J. Kokame, N. Koori and S. Tanaka, *J. Phys. Soc. Jpn.* 41 (1976) 361
- 1976TH1A Thibault, in *Cargese Conf.*, CERN 76-14 (1976) 93
- 1976VI1A Vidal-Quadras et al., *Nuovo Cim.* B34 (1976) 380
- 1976WA15 G.L. Wales and R.C. Johnson, *Nucl. Phys.* A274 (1976) 168
- 1976WE09 R.B. Weisenmiller, N.A. Jelley, K.H. Wilcox, G.J. Wozniak and J. Cerny, *Phys. Rev. C*13 (1976) 1330
- 1976WH1A White, Cox, Knox and Lane, *Bull. Amer. Phys. Soc.* 21 (1976) 513
- 1976WI05 N. Willis, I. Brissaud, L. Bimbot, Y. Le Bornec and B. Tatischeff, *Nucl. Phys.* A261 (1976) 45
- 1976WO11 G.J. Wozniak, D.P. Stahel, J. Cerny and N.A. Jelley, *Phys. Rev. C*14 (1976) 815
- 1976YO1A Young et al., *Bull. Amer. Phys. Soc.* 21 (1976) 30
- 1977AD07 G.D. Adeev, A.M. Kazantseva, V.V. Kozyr, V.A. Matusevich and I.P. Chernov, *Yad. Fiz.* 26 (1977) 683; *Sov. J. Nucl. Phys.* 26 (1977) 361
- 1977AN21 I. Angeli and M. Csatlos, *Nucl. Phys.* A288 (1977) 480
- 1977AR06 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, *Nucl. Phys.* A283 (1977) 350
- 1977AR12 N. Arena, G. Calvi, S. Cavallaro, R. Potenza, G. Inghima and M. Sandoli, *Nuovo Cim.* A38 (1977) 101

- 1977AS03 V.M. Asaturyan, E.O. Grigoryan, G.A. Vartapetyan, A.S. Danagulyan, N.A. Demekhina and A.G. Khudaverdyan, *Yad. Fiz.* 25 (1977) 1133; *Sov. J. Nucl. Phys.* 25 (1977) 601
- 1977AU1B Austin, MSUCL-254 (1977)
- 1977AV01 V.V. Avdeychikov, V.I. Bogatin and O.V. Lozhkin, *Yad. Fiz.* 25 (1977) 3; *Sov. J. Nucl. Phys.* 25 (1977) 1
- 1977BA11 F.C. Barker and G.T. Hickey, *J. Phys. (London)* G3 (1977) L23
- 1977BA1Q Baer, in *High Energy Phys. & Nucl. Struct., Zurich 1977* (Birkhauser Verlag, 1977) 245
- 1977BA2G Barnes, *Proc. Int. Conf. Nucl. Struct., Tokyo (1977)*; *J. Phys. Soc. Jpn. Suppl.* 44 (1978) 783
- 1977BA2H Bacher et al., in *Tokyo (1977), Contrib. Papers* p. 812
- 1977BA2K Bains and Galloway, *Nucl. Instrum. Meth. Phys. Res.* 143 (1977) 295
- 1977BA39 A.K. Basak, J.A.R. Griffith, O. Karban, S. Roman and G. Tungate, *Nucl. Phys.* A286 (1977) 420
- 1977BA48 W. Bambynek, H. Behrens, M.H. Chen, B. Crasemann, M.L. Fitzpatrick, K.W.D. Ledingham, H. Genz, M. Mutterer and R.L. Intemann, *Rev. Mod. Phys.* 49 (1977) 77; *Erratum Rev. Mod. Phys.* 49 (1977) 961
- 1977BE1L Bernstein, *Bull. Amer. Phys. Soc.* 22 (1977) 522
- 1977BE1X Bent et al., in *Proc. Zurich-Sin (1977)* 14
- 1977BE51 F. Becker and I. Strykowski, *Nucl. Phys.* A289 (1977) 446
- 1977BO07 M.-C. Bouten and M. Bouten, *Z. Phys.* A280 (1977) 45
- 1977BR1J Brajnik et al., *Fiz. (Yugoslavia)* 9, Suppl. 1 (1977) 48; *Phys. Abs.* 13056 (1978)
- 1977CA1E Carter et al., in *Proc., Zurich-Sin (1977)* p. 318
- 1977CH01 N.S. Chant, N.S. Wall, C.F. Clement, S.M. Perez and J.N. Craig, *Phys. Rev.* C15 (1977) 53
- 1977CH22 R. Chechik, Z. Fraenkel, H. Stocker and Y. Eyal, *Nucl. Phys.* A287 (1977) 353
- 1977CO1G Colombo et al., in *Tokyo (1977), Contrib. Papers* p. 490
- 1977DE1L Devins et al., *Bull. Amer. Phys. Soc.* 22 (1977) 562
- 1977DE1M Devins, in *Momentum Wave Funct., 1976*, Ed. Devins (AIP Conf. Proc. no 36, AIP, New York, 1977) 77
- 1977DO06 G.E. Dogotar, R.A. Eramzhyan, H.R. Kissener and R.A. Sakaev, *Nucl. Phys.* A282 (1977) 474
- 1977DW1A Dwyer and Meyer, *Bull. Amer. Phys. Soc.* 22 (1977) 600
- 1977EL10 M. El-Nadi and A. Osman, *Ann. Phys. (Leipzig)* 34 (1977) 91

- 1977FO04 P. Fontes, Phys. Rev. C15 (1977) 2159
- 1977GE08 C.K. Gelbke, D.K. Scott, M. Bini, D.L. Hendrie, J.L. Laville, J. Mahoney, M.C. Mer-
maz and C. Olmer, Phys. Lett. B70 (1977) 415
- 1977GE1D Gerard, Bull. Inf. Sci. Tech. (France) 224-225 (1977) 88; Phys. Abs. 572 (1979)
- 1977GO07 J. Gosset, H.H. Gutbrod, W.G. Meyer, A.M. Poskanzer, A. Sandoval, R. Stock and
G.D. Westfall, Phys. Rev. C16 (1977) 629
- 1977GR1F Grand and Golland, Nucl. Instrum. Meth. 145 (1977) 49
- 1977HA1L Hagen, Fisher and Ormes, Astrophys. J. 212 (1977) 262
- 1977HA1N Hasegawa et al., in Tokyo (1977), Contrib. Papers p. 539
- 1977HA1P Haerberli, in Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44
(1978) 435
- 1977HO1B Hoistad, in High Energy Phys. Nucl. Struct., Zurich 1977 (Birkhauser Verlag, 1977)
215
- 1977IR01 M. Irshad, C. Rioux, J. Asai, R. Pigeon and R.J. Slobodrian, Nucl. Phys. A286 (1977)
483
- 1977JA14 H.U. Jager and M. Kirchbach, Nucl. Phys. A291 (1977) 52
- 1977KA07 J. Kallne and A.W. Obst, Phys. Rev. C15 (1977) 477
- 1977KN04 U. Kneissl, G. Kuhl and K.H. Leister, Z. Phys. A281 (1977) 35
- 1977KO1L Kohl, Parkison and Withbroe, Astrophys. J. 212 (1977) L101
- 1977KO25 D.S. Koltun and F. Myhrer, Z. Phys. A283 (1977) 397
- 1977KO27 W. Kohler, G. Gruber, A. Steinhauser and K. Bethge, Nucl. Phys. A290 (1977) 233
- 1977KU1D R. Kullberg, K. Kristiansson, B. Lindkvist and I. Otterlund, Nucl. Phys. A280 (1977)
491
- 1977LE1G Lenz, in High Energy Phys. & Nucl. Struct., Zurich, 1977 (Birkhauser Verlag, 1977)
175
- 1977LI19 H. Liskien and A. Paulsen, Atomkernenergie 30 (1977) 59
- 1977LI1F Lin, Hou and Wen, in Tokyo (1977), Contrib. Papers p. 560
- 1977LO10 M.A. Lone, C.B. Bigham, J.S. Fraser, H.R. Schneider, T.K. Alexander, A.J. Ferguson
and A.B. McDonald, Nucl. Instrum. Meth. 143 (1977) 331
- 1977LO1C Lobov and Danilyan, Izv. Akad. Nauk SSSR Ser. Fiz. 41 (1977) 1548
- 1977LU1A Ludeking and Vary, Bull. Amer. Phys. Soc. 22 (1977) 1030
- 1977MA1H Mason, Garcia-Munoz and Simpson, Bull. Amer. Phys. Soc. 22 (1977) 567
- 1977MA1M Mausner et al., Bull. Amer. Phys. Soc. 22 (1977) 547
- 1977MA1N Maluszynska, Niedzwiedziuk, Salackij and Holwek, Acta Phys. Pol. B8 (1977) 309

- 1977MA1P Marazano, Maziere, Berger and Comar, *Int. J. Appl. Rad. & Isotopes* 28 (1977) 49
- 1977MA35 V.B. Mandelzweig, *Nucl. Phys. A*292 (1977) 333
- 1977MC1F McCarthy, in *Momentum Wave Functions-1976*, AIP Conf. Proc. 36 (1977) 1
- 1977ME1C Meadows, ANL/NDM-25 (1977)
- 1977MO09 J.M. Moss, W.D. Cornelius and D.R. Brown, *Phys. Lett. B*69 (1977) 154
- 1977MO1A T. Motobayashi, *Rept. Inst. Phys. Chem. Res.* 53 (1977) 81.
- 1977MO1F Mougey, *Proc. Int. Conf. Nucl. Struct., Tokyo*, (1977); *J. Phys. Soc. Jpn. Suppl.* 44 (1978) 420
- 1977MO1K Morinaga et al., in *Tokyo (1977)*, *Contrib. Papers* p. 56
- 1977MU1A Mukhopadhyay, *Phys. Rept.* C30 (1977) 1
- 1977NI1A Nishioka, Saito and Yasuno, *Proc. Int. Conf. Nucl. Struct., Tokyo*, (1977), *J. Phys. Soc. Jpn. Suppl.* 44 (1978) 239
- 1977OK1C Okabe, in *Tokyo (1977)*, *Contrib. Papers* p. 154
- 1977OK1D Okuma, in *Tokyo (1977)*, *Contrib. Papers* p. 598
- 1977OS08 A. Osman, *Nuovo Cim.* A39 (1977) 138
- 1977PA25 L.A. Parks, K.W. Kemper, A.H. Lumpkin, R.I. Cutler, L.H. Harwood, D. Stanley, P. Nagel and F. Petrovich, *Phys. Lett. B*70 (1977) 27
- 1977PH02 D.-L. Pham and R. de Swiniarski, *Nuovo Cim.* A41 (1977) 543
- 1977PI1D Pickar et al., *Bull. Amer. Phys. Soc.* 22 (1977) 591
- 1977RE08 H. Reiss, H.V. Klapdor, G. Rosner, J.L.C. Ford, Jr. and S.T. Thornton, *Nucl. Phys.* A283 (1977) 149
- 1977RE1A Redish, in *Momentum Wave Functions, 1976*, Ed. Devins, AIP Conf. Proc. 37 (AIP, New York, 1977) 111
- 1977RI08 W.A. Richter and P.R. de Kock, *Z. Phys.* A282 (1977) 207
- 1977RO1E Roos, in *Momentum Wave Functions, 1976*, Ed. Devins, AIP Conf. Proc. 37 (AIP, New York, 1977) 32
- 1977SA1B Sadkovskii, Feofilov, Denisov and Kolalis, *Yad. Fiz.* 26 (1977) 1200
- 1977SA1K Saltmarsh, Ludemann, Fulmer and Styles, *Nucl. Instrum. Meth.* 145 (1977) 81
- 1977SC1D Schramm and Wagoner, *Ann. Rev. Nucl. Sci.* 27 (1977) 37
- 1977SC1G Schroder and Huizenga, *Ann. Rev. Nucl. Sci.* 27 (1977) 465
- 1977SE1D Seya et al., in *Tokyo (1977)*, *Contrib. Papers* p. 149
- 1977SH09 M.A.M. Shahabuddin, C.J. Webb and V.R.W. Edwards, *Nucl. Phys.* A284 (1977) 83
- 1977SH1D Shimoda et al., in *Tokyo (1977)*, *Contrib. Papers* p. 703

- 1977SI01 R.R. Silbar, J.N. Ginocchio and M.M. Sternheim, Phys. Rev. C15 (1977) 371
- 1977SI1D Silberberg and Tsao, Astrophys. J. Suppl. 35 (1977) 129
- 1977SL02 I. Sledzinska, A. Saganek, Z. Wilhelmi and B. Zwieglinski, Acta Phys. Pol. B8 (1977) 227
- 1977ST1G Steiner, in High Energy Phys. Nucl. Struct., Zurich (Birkhauser Verlag, 1977) p. 261
- 1977TE01 W.D. Teeters and D. Kurath, Nucl. Phys. A275 (1977) 61
- 1977TE02 W.D. Teeters and D. Kurath, Nucl. Phys. A283 (1977) 1
- 1977UD1A Udagawa, Proc. Int. Conf. Nucl. Struct., Tokyo (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 667
- 1977WA05 C.N. Waddell, E.M. Diener, R.G. Allas, L.A. Beach, R.O. Bondelid, E.L. Petersen, A.G. Pieper, R.B. Theus, C.C. Chang and N.S. Chant, Nucl. Phys. A281 (1977) 418
- 1977WA1G Walecka, AIP Conf. Proc. no. 37 (Indiana Univ.), Weak Inter. Phys., 1977, Ed. Lichtenberg (AIP, New York, 1977) 125
- 1977WE03 R.B. Weisenmiller, N.A. Jelley, D. Ashery, K.H. Wilcox, G.J. Wozniak, M.S. Zisman and J. Cerny, Nucl. Phys. A280 (1977) 217
- 1977WE1B R.B. Weisenmiller, Thesis, Univ. of California, Berkeley (1976); LBL-5077 (1977)
- 1977WE1C Wessels et al., Nucl. Instrum. Meth. Phys. Res. 141 (1977) 157
- 1977WH1A White et al., Bull. Amer. Phys. Soc. 22 (1977) 631
- 1977WH1B White, Thesis Ohio Univ. (1977) (COO-2490-6)
- 1977WO1A Wolf and Redvanly, Int. J. Appl. Rad. Isotopes 28 (1977) 29
- 1977YA1A Yasue et al., in Tokyo (1977), Contrib. Papers p. 520
- 1977YA1B Yakovlev, Fiz. Elem. Chastits At. Yadra 8 (1977) 255
- 1977YO1D Yoro and Arima, in Tokyo (1977), Contrib. Papers p. 170
- 1977YO1E K. Yoro, Phys. Lett. B70 (1977) 147
- 1977YO1F Young, Golden and Kapetanakos, Rev. Sci. Instrum. 48 (1977) 432
- 1977ZA1B Zatolokin et al., Int. J. Appl. Rad. Isotopes 28 (1977) 459
- 1977ZE1A Zelenskaya and Teplov, Izv. Akad. Nauk SSSR Ser. Fiz. 41 (1977) 1711
- 1978AJ02 F. Ajzenberg-Selove, E.R. Flynn and O. Hansen, Phys. Rev. C17 (1978) 1283.
- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1
- 1978AL1G Alkhazov, Belostotsky and Vorobyov, Phys. Rep. C42 (1978) 89
- 1978AM01 J.F. Amann, P.D. Barnes, K.G.R. Doss, S.A. Dytman, R.A. Eisenstein, J.D. Sherman and W.R. Wharton, Phys. Rev. Lett. 40 (1978) 758
- 1978AN07 I. Angeli and M. Csatlos, Atomki Kozlem. 20 (1978) 1

- 1978AT01 M. Atarashi, K. Hira and H. Narumi, *Prog. Theor. Phys.* 60 (1978) 209
- 1978AU1C Austin, *Bull. Amer. Phys. Soc.* 23 (1978) 609
- 1978AV1A Avishai, *Z. Phys.* A285 (1978) 333
- 1978AZ02 L.S. Azhgirei, I.K. Vzorov, V.N. Zhmyrov, V.V. Ivanov, M.A. Ignatenko, A.S. Kuznetsov, M.G. Meshcheryakov, S.V. Razin and G.D. Stoletov, *Yad. Fiz.* 28 (1978) 1017; *Sov. J. Nucl. Phys.* 28 (1978) 522
- 1978BA1F Barschall, *Ann. Rev. Nucl. Part. Sci.* 28 (1978) 207
- 1978BH03 R.K. Bhowmik, E.C. Pollacco, N.E. Sanderson, J.B.A. England and G.C. Morrison, *Phys. Lett.* B80 (1978) 41
- 1978BI08 R. Billerey, C. Cerruti, A. Chevarier, N. Chevarier and A. Demeyer, *Z. Phys.* A285 (1978) 389
- 1978BO09 M.C. Bouten and M. Bouten, *Nucl. Phys.* A299 (1978) 141
- 1978BO1K Boesgaard and Heacox, *Astrophys. J.* 226 (1978) 888
- 1978BO31 M. Bouten, E. Flerackers and M.C. Bouten, *Nucl. Phys.* A307 (1978) 413
- 1978BU1B Buffington, Orth and Mast, *Astrophys. J.* 226 (1978) 355
- 1978BY1A Byrnak, *Nuovo Cim.* A44 (1978) 1
- 1978CH1P Chen et al., *Bull. Amer. Phys. Soc.* 23 (1978) 953
- 1978CO02 J.C. Comiso, H.J. Ziock, K.O.H. Ziock and K.P. Ziock, *Z. Phys.* A285 (1978) 101
- 1978CO13 J.D. Cossairt, S.B. Talley, D.P. May, R.E. Tribble and R.L. Spross, *Phys. Rev.* C18 (1978) 23
- 1978CO1J J.R. Comfort and B.C. Karp, *Phys. Lett.* B80 (1978) 1
- 1978CU02 G. Cutrona, G. Lanzano, F. Porto and S. Sambataro, *Lett. Nuovo Cim.* 21 (1978) 368
- 1978DA1A Dalitz and Gal, *Ann. Phys.* 116 (1978) 167
- 1978DE15 P. Desgrolard, P.A.M. Guichon and J. Joseph, *Nuovo Cim.* A43 (1978) 475
- 1978DE1P Devins et al., *Bull. Amer. Phys. Soc.* 23 (1978) 97
- 1978DI04 S. Di Liberto, F. Meddi, G. Romano, G. Rosa and C. Sgarbi, *Nucl. Phys.* A296 (1978) 519
- 1978DI1A Di Napoli et al., *J. Inorg. Nucl. Chem.* 40 (1978) 1973
- 1978DI1D Dietrich and Simpson, *Astrophys. J.* 225 (1978) L41
- 1978DW1A Dwek, *Astrophys. J.* 225 (1978) L149
- 1978DW1B Dwyer, *Astrophys. J.* 224 (1978) 691
- 1978FO13 H.T. Fortune, D.J. Crozier, B. Zeidman and M.E. Cobern, *Nucl. Phys.* A303 (1978) 14
- 1978FR20 A.D. Frawley, J.F. Mateja, A. Roy and N.R. Fletcher, *Phys. Rev. Lett.* 41 (1978) 846

- 1978GE1C Gelbke et al., Phys. Rept. C42 (1978) 311
- 1978GL1D Glendinning et al., Bull. Amer. Phys. Soc. 23 (1978) 635
- 1978GR1F Green and Korteling, in Proc., Cluster, Winnipeg (1978) 89
- 1978GU10 K. Gul, J. Phys. Soc. Jpn. 44 (1978) 353
- 1978HA1M Harrison and Kubiczek, Nucl. Instrum. Meth. Phys. Res. 154 (1978) 399
- 1978HA35 H.-B. Hakansson, T. Berggren and R. Bengtsson, Nucl. Phys. A306 (1978) 406
- 1978HE1C Hendrie, in Clustering, Winnipeg 1978, AIP Conf. Proc. 47 (1978) 402
- 1978HE1D Heitzmann, Atomkernenergie 31 (1978) 262
- 1978HI03 M. Hirata, Phys. Rev. Lett. 40 (1978) 704
- 1978IG1A Igo et al., Bull. Amer. Phys. Soc. 23 (1978) 47
- 1978IG1B Igo, Rev. Mod. Phys. 50 (1978) 523
- 1978KI03 V.V. Kirichenko, Y.M. Arkatov, P.I. Vatset, I.V. Dogyust and A.F. Khodyachikh, Sov. J. Nucl. Phys. 27 (1978) 314
- 1978KI08 H.R. Kissener, G.E. Dogotar, R.A. Eramzhyan and R.A. Sakaev, Nucl. Phys. A302 (1978) 523
- 1978KI13 H.R. Kissener, G.E. Dogotar, R.A. Eramzhyan and R.A. Sakaev, Nucl. Phys. A312 (1978) 394
- 1978KL03 F.J. Kline and E. Hayward, Phys. Rev. C17 (1978) 1531
- 1978KN01 H.D. Knox, R.M. White and R.O. Lane, Nucl. Sci. Eng. 65 (1978) 65
- 1978KO01 D.G. Kovar, W. Henning, B. Zeidman, Y. Eisen, J.R. Erskine, H.T. Fortune, T.R. Ophel, P. Sperr and S.E. Vigdor, Phys. Rev. C17 (1978) 83
- 1978KU12 N. Kumagai, G. Isoyama, E. Tanaka, K. Kageyama and T. Ishimatsu, Nucl. Instrum. Meth. Phys. Res. 157 (1978) 423
- 1978LE04 H.C. Lee, Nucl. Phys. A294 (1978) 473
- 1978LE15 R.D. Levine, S.G. Steadman, J.S. Karp and Y. Alhassid, Phys. Rev. Lett. 41 (1978) 1537
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, Table of Isotopes 7th Ed. (John Wiley and Sons, Inc., New York, 1978)
- 1978LO1D Loose, Wagenbach and Clausnitzer, Nucl. Instrum. Meth. Phys. Res. 150 (1978) 345
- 1978LU1B Ludeking and Vary, Bull. Amer. Phys. Soc. 23 (1978) 552
- 1978ME1D Meyer, Nature 272 (1978) 675
- 1978MO01 C.L. Morris, R.L. Boudrie, J.J. Kraushaar, R.J. Peterson, R.A. Ristinen, G.R. Smith, J.E. Bolger, W.J. Braithwaite, C.F. Moore and L.E. Smith, Phys. Rev. C17 (1978) 227

1978MO09 M. Morike, G. Staudt and F. Weng, *Z. Phys.* A287 (1978) 211

1978OE1A W. Oelert, A. Djalois, C. Mayer-Boricke, P. Turek and S. Wiktor, *AIP Conf. Proc.* 47 (1978) 730.

1978PE1E Peterson, Seglie and Ascuitto, *Bull. Amer. Phys. Soc.* 23 (1978) 947

1978RA2A S. Raman, C.A. Houser, T.A. Walkiewicz and I.S. Towner, *At. Data Nucl. Data Tables* 21 (1978) 567; Erratum *At. Data Nucl. Data Tables* 22 (1978) 369

1978RO1D Rolfs and Trautvetter, *Ann. Rev. Nucl. & Part. Sci.* 28 (1978) 115

1978SA26 A. Saha, R. Kamermans, J. van Driel and H.P. Morsch, *Phys. Lett.* B79 (1978) 363

1978SC1D Schulte, *Nucl. Instrum. Meth. Phys. Res.* 149 (1978) 932

1978SC31 R.A. Schrack, G.P. Lamaze and O.A. Wasson, *Nucl. Sci. Eng.* 68 (1978) 189

1978SH14 N.G. Shevchenko, A.Y. Buki, B.V. Mazanko, V.N. Polishchuk and A.A. Khomich, *Yad. Fiz.* 28 (1978) 12; *Sov. J. Nucl. Phys.* 28 (1978) 5

1978SH1F Shepard, *Bull. Amer. Phys. Soc.* 23 (1978) 11

1978SI1D R.R. Silbar, J.N. Ginocchio and M.M. Sternheim, *Phys. Rev.* C18 (1978) 2785

1978TA04 S. Tanaka, *J. Phys. Soc. Jpn.* 44 (1978) 1405

1978TA12 S. Tanaka, K. Fukunaga, S. Kakigi, T. Ohsawa, N. Fujiwara and T. Yanabu, *J. Phys. Soc. Jpn.* 45 (1978) 733

1978TA1D Tartakovskii, Kozlovskii and Malyarzh, *Ukr. Fiz. Zh. USSR* 23 (1978) 368

1978TI1A Tilbury et al., *Bull. Amer. Phys. Soc.* 23 (1978) 1033

1978TS1A Tsangarides, Wills and Bent, *Bull. Amer. Phys. Soc.* 23 (1978) 952

1978TU06 P.P. Tung, K.A. Erb, M.W. Sachs, G.B. Sherwood, R.J. Ascuitto and D.A. Bromley, *Phys. Rev.* C18 (1978) 1663

1978VA05 J. van de Wiele, E. Gerlic, H. Langevin-Joliot and G. Duhamel, *Nucl. Phys.* A297 (1978) 61

1978VA12 L. Van der Zwan and K.W. Geiger, *Nucl. Phys.* A306 (1978) 45

1978VA1A L.C. Vaz and J.M. Alexander, *Phys. Rev.* C18 (1978) 2152

1978VO1D Volkov, *Phys. Rept.* 44 (1978) 93

1978WA1B Walter, in *Clustering, Winnipeg 1978*, *AIP Conf. Proc.* 47 (1978) 444

1978WE1D Wessels et al., *Nucl. Instrum. Meth.* 141 (1977) 157

1978WH1A Whitten et al., *Bull. Amer. Phys. Soc.* 23 (1978) 934

1978WI1F Weiland and Highfall, *Bull. Amer. Phys. Soc.* 23 (1978) 1033

1978WO1C Wolf, *Bull. Amer. Phys. Soc.* 23 (1978) 1033

1978ZA1D Zalm, Van Hienen and Glaudemans, *Z. Phys.* A287 (1978) 255

- 1978ZE02 A.F. Zeller, G.T. Hickey, D.C. Weisser, D.F. Hebbard and B.A. Robson, Nucl. Phys. A301 (1978) 130
- 1978ZE03 N.S. Zelenskaya, V.M. Lebedev and T.A. Yushchenko, Yad. Fiz. 28 (1978) 90; Sov. J. Nucl. Phys. 28 (1978) 44
- 1979AJ01 F. Ajzenberg-Selove, Nucl. Phys. A320 (1979) 1
- 1979AL1Q Alonso, Chatterjee and Tobias, Bull. Amer. Phys. Soc. 24 (1979) 142
- 1979AN1J Aniol et al., in Houston, AIP Conf. Proc. 54 (1979) 276
- 1979AN1K Anderson et al., in Houston, AIP Conf. Proc. 54 (1979) 194
- 1979AN1L Anderson, Failey, Zoller and Gordon, in Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York, 1979) 546
- 1979AN1M Anderson et al., in Conf. Proc. TRIUMF, Vancouver (1979) Paper 4B14
- 1979AU07 G.F. Auchampaugh, S. Plattard and N.W. Hill, Nucl. Sci. Eng. 69 (1979) 30
- 1979BE1H Benenson and Kashy, Rev. Mod. Phys. 51 (1979) 527
- 1979BE1N Beer et al., in Conf. Proc. TRIUMF, Vancouver (1979) Paper 1B27
- 1979BE1W Bent, in Houston, AIP Conf. Proc. 54 (1979) 142
- 1979BO07 S. Boffi, C. Giusti, F.D. Pacati and S. Frullani, Nucl. Phys. A319 (1979) 461
- 1979BO1A Boudrie et al., Conf. Proc. Triumf, Vancouver (1979) Paper 4A3
- 1979BO1U Booth, in 1978 RPI Symp. Photopion Nucl. Phys. (Plenum Press, New York, 1979) 129
- 1979BO1V Boudard, Couvert and Dillig, in Houston, AIP Conf. Proc. 54 (1979) 185
- 1979BR03 H.S. Bradlow, W.D.M. Rae, P.S. Fisher, N.S. Godwin, G. Proudfoot and D. Sinclair, Nucl. Phys. A314 (1979) 171
- 1979BR1D Brajnik et al., Int. Conf. Nucl. Phys. with Electromag. Int., Mainz (1979) 46
- 1979CH11 D. Christenson, M.W. McNaughton and J.A. Jungerman, Nucl. Instrum. Meth. Phys. Res. 160 (1979) 499
- 1979CR03 B.J.B. Crowley, Ann. Phys. 117 (1979) 323
- 1979DE01 P. Desgrolard and P.A.M. Guichon, Phys. Rev. C19 (1979) 120
- 1979DE15 P. Desgrolard and P.A.M. Guichon, Z. Phys. A290 (179) 373
- 1979DY01 P. Dyer, T.C. Awes, C.K. Gelbke, B.B. Back, A. Mignerey, K.L. Wolf, H. Breuer, V.E. Viola, Jr. and W.G. Meyer, Phys. Rev. Lett. 42 (1979) 560
- 1979EL1E Ellis et al., in Houston, AIP Conf. Proc. 54 (1979) 274
- 1979FA1B Famery, J. Phys. G5 (1979) 241
- 1979GA04 J.A. Gaidos, L.J. Gutay, A.S. Hirsch, R. Mitchell, T.V. Ragland, R.P. Scharenberg, F. Turkot, R.B. Willmann and C.L. Wilson, Phys. Rev. Lett. 42 (1979) 82

- 1979GO07 R. Gorgen, F. Hinterberger, R. Jahn, P. von Rossen and B. Schuller, Nucl. Phys. A320 (1979) 296
- 1979GO11 J. Gomez del Campo, R.G. Stokstad, J.A. Biggerstaff, R.A. Dayras, A.H. Snell and P.H. Stelson, Phys. Rev. C19 (1979) 2170
- 1979GO1N Goncharova, Int. Conf. Nucl. Phys. with Electromag. Inter., Mainz (1979) 4.9
- 1979GR1E Greenwood, in Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York 1979) 441
- 1979HE1E Herman, Cormier and Fulton, Bull. Amer. Phys. Soc. 24 (1979) 664
- 1979HO09 B. Hoistad, S. Dahlgren, T. Johansson and O. Jonsson, Nucl. Phys. A319 (1979) 409
- 1979HU02 J. Hufner and M. Thies, Phys. Rev. C20 (1979) 273
- 1979JA1F D.F. Jackson and D.J. Brenner, Prog. Part. Nucl. Phys. 5 (1981) 143
- 1979JO1C Jones, in Houston, AIP Conf. Proc. 54 (1979) 116
- 1979JU1B Journey, in Proc. Third Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York, 1979) p. 461
- 1979KA07 S.B. Kaufman, M.W. Weisfield, E.P. Steinberg, B.D. Wilkins and D.J. Henderson, Phys. Rev. C19 (1979) 962
- 1979KI1H Klingbeck and Huber, Int. Conf. Nucl. Phys. with Electromag. Inter., Mainz (1979) 3.19
- 1979KO1Q Koehler et al., Bull. Amer. Phys. Soc. 24 (1979) 656
- 1979KU08 P.D. Kunz, A. Saha and H.T. Fortune, Phys. Rev. Lett. 43 (1979) 341
- 1979LI11 B.J. Lieb, H.S. Plendl, C.E. Stronach, H.O. Funsten and V.G. Lind, Phys. Rev. C19 (1979) 2405
- 1979MA20 T.A.J. Maris, M.R. Teodoro and C.A.Z. Vasconcellos, Nucl. Phys. A322 (1979) 461
- 1979MC1A McGuire, Von Rosenvinge and McDonald, Bull. Amer. Phys. Soc. 24 (1979) 693
- 1979MO1L Moreh, Vodhanel and Sellyey, Bull. Amer. Phys. Soc. 24 (1979) 648
- 1979MO1M Moreh, Vodhanel and Sellyey, Int. Conf. Nucl. Phys. with Electromag. Inter., Mainz (1979) 4.26
- 1979NI06 H. Nishioka, S. Saito and M. Yasuno, Prog. Theor. Phys. 62 (1979) 424
- 1979OS10 A. Osman and G.U. Din, Ann. Phys. (Leipzig) 36 (1979) 56
- 1979PA09 L.A. Parks, K.W. Kemper, R.I. Cutler and L.H. Harwood, Phys. Rev. C19 (1979) 2206
- 1979PI06 P.H. Pile, R.D. Bent, R.E. Pollock, P.T. Debevec, R.E. Marrs, M.C. Green, T.P. Sjoreen and F. Soga, Phys. Rev. Lett. 42 (1979) 1461
- 1979PI1E Pile et al., Bull. Amer. Phys. Soc. 24 (1979) 614

- 1979SC08 T.C. Schweizer, R.R. Doering, S.T. Thornton, L.C. Dennis, K.R. Cordell and R.L. Parks, Phys. Rev. C19 (1979) 1408
- 1979SO1C Soga et al., Bull. Amer. Phys. Soc. 24 (1979) 614
- 1979ST03 M.L. Stelts, R.E. Chrien, M. Goldhaber, M.J. Kenny and C.M. McCullagh, Phys. Rev. C19 (1979) 1159
- 1979SU1C Subramanian et al., Bull. Amer. Phys. Soc. 24 (1979) 593
- 1979TI1A Titov, Int. Conf. Nucl. Phys. with Electromagnetic Interact., Mainz (1979) 6.14
- 1979VI05 A. Vidal-Quadras and M. Ortega, Nuovo Cim. A49 (1979) 235
- 1979WE06 J.P. Wefel and J.M. Kidd, Phys. Rev. C19 (1979) 1380
- 1979WH1A Whitten et al., Conf. Proc. Triumf, Vancouver (1979) Paper 4A4
- 1979WI1G Wieland, Highfall and King, IEEE Trans. on Nucl. Sci. 26 (1979) 1713
- 1979ZI1C Ziock et al., in Conf. Proc. TRIUMF, Vancouver (1979) Paper IE29
- 1979ZW01 B. Zwioglinski, W. Benenson, R.G.H. Robertson and W.R. Coker, Nucl. Phys. A315 (1979) 124
- 1981AJ01 F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1.