

# Energy Levels of Light Nuclei $A = 12$

F. Ajzenberg-Selove<sup>a</sup> and T. Lauritsen<sup>b</sup>

<sup>a</sup> *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

<sup>b</sup> *California Institute of Technology, Pasadena, California*

**Abstract:** An evaluation of  $A = 5-24$  was published in *Nuclear Physics* 11 (1959), p. 1. This version of  $A = 12$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the TUNL/NNDC format.

(References closed December 1, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 12$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides: <sup>12</sup>B, <sup>12</sup>C, <sup>12</sup>N

B. Tables of Recommended Level Energies:

Table 12.1: Energy levels of <sup>12</sup>B

Table 12.4: Energy levels of <sup>12</sup>C

Table 12.12: Energy levels of <sup>12</sup>N

C. References

D. Figures: <sup>12</sup>B, <sup>12</sup>C, <sup>12</sup>N

**<sup>12</sup>B**  
(Fig. 19)

GENERAL:

*Theory:* See (1956KU1A, 1958FR1C).

1.  $^{12}\text{B}(\beta^-)^{12}\text{C}$   $Q_m = 13.376$

The spectrum is complex: see  $^{12}\text{C}$ . The transition to  $^{12}\text{C}_{\text{g.s.}}$  is allowed; hence  $J(^{12}\text{B}) = 1^+$ .

2.  $^6\text{Li}(^7\text{Li}, \text{p})^{12}\text{B}$   $Q_m = 8.338$

Three groups of protons are reported, corresponding to the ground state and to the excited states at 0.95 and 1.67 MeV. At  $E(^7\text{Li}) = 2.0$  MeV,  $\theta = 90^\circ$  (lab), the relative intensities are 1 : 1.1 : 0.8 (1957NO14). See also (1957NO17).

3.  $^7\text{Li}(^7\text{Li}, \text{d})^{12}\text{B}$   $Q_m = 3.311$

See  $^{14}\text{C}$ .

4.  $^9\text{Be}(\alpha, \text{p})^{12}\text{B}$   $Q_m = -6.884$

At  $E_\alpha = 21.7$  MeV, proton groups are observed corresponding to  $^{12}\text{B}^*(0, 0.95, 1.65, 3.38, 3.82)$  (1951MC57, 1955RA41).

5.  $^9\text{Be}(^7\text{Li}, \alpha)^{12}\text{B}$   $Q_m = 10.463$

See (1957NO17).

6.  $^{10}\text{B}(\text{t}, \text{p})^{12}\text{B}$   $Q_m = 6.344$

Table 12.1: Energy levels of  $^{12}\text{B}$

$E_x$ (MeV $\pm$ keV)	$J^\pi$	$\tau_{1/2}$ or $\Gamma$ (keV)	Decay	Reactions
0	$1^+$	$\tau_{1/2} = 20.34 \pm 0.5$ msec	$\beta^-$	1, 2, 3, 4, 5, 10, 13, 15, 20, 22
$0.947 \pm 5$	$\leq 3^+$	$< 10$	$\gamma$	2, 4, 6, 10
$1.674 \pm 11$	$1^-, 2^-$	$< 10$	$\gamma$	2, 4, 6, 10
$2.618 \pm 11$		$< 10$		6, 10
$2.723 \pm 11$		$< 10$		10
$3.383 \pm 9$	$\leq 3^+$	$< 10$		4, 6, 10
$3.76 \pm 10$	$2^+$	$40 \pm 5$	n	4, 6, 8, 10
$4.54 \pm 20$	$3^-$	$140 \pm 20$	n	6, 8, 10
$5.00 \pm 20$	1	$60 \pm 20$	n	8
$5.61 \pm 20$	2	$120 \pm 40$	n	8
$5.73 \pm 20$	3	$60 \pm 20$	n	8

At  $E_t = 0.90$  MeV, proton groups are observed corresponding to  $^{12}\text{B}^*(0.94, 1.65, 2.61, 3.37, 3.75, 4.46)$  (1955BI26:  $\pm \approx 0.1$  MeV).

7.  $^{11}\text{B}(n, \gamma)^{12}\text{B}$   $Q_m = 3.365$

$\sigma_{\text{th}} < 50$  mb (1958HU18). See also (1953WI1C).

8.  $^{11}\text{B}(n, n)^{11}\text{B}$   $E_b = 3.365$

The parameters of observed resonances at  $E_n = 0.43, 1.28, 1.78, 2.45$  and  $2.58$  MeV are exhibited in Table 12.2. The rise of the cross section at low energies may indicate a broad level formed by s-wave neutrons (1951BO45). The angular distributions of elastically scattered neutrons have been studied for  $E_n = 0.43$  to  $1.50$  MeV by (1955WI25). It is found that the 0.43-MeV resonance has  $J = 2^+$ , formed by p-waves, either all in channel spin 1 or all in channel spin 2. The 1.28 MeV resonance had  $J = 3^-$ , formed by d-waves, with the level width of channel spin 2 about 10 times that of channel spin 1. Potential scattering at  $E_n = 1.5$  MeV is nearly all s-wave:  $\delta_0 = -90^\circ$  (1955WI25).

The total cross section for natural boron shows no sharp discontinuities for  $E_n = 4.4$  to  $5.6$  MeV ( $\sigma \approx 1.6$  b) and for  $E_n = 7.8$  to  $8.6$  MeV ( $\sigma \approx 1.4$  b) (1956BE1D). The total cross section has also been measured for  $E_n = 6$  to  $9.7$  MeV (1954NE1A) and from  $E_n = 14.1$  to  $18.0$  MeV

Table 12.2: Resonances in  $^{11}\text{B}(n, n)^{11}\text{B}$

$E_n^a$ (MeV)	$\sigma_{\max} - \sigma_{\text{tot}}^b$ b	$\Gamma_n^a$ (keV)	$^{12}\text{B}^*$ (MeV)	$l^c$	$\theta_n^2^b$	$J^\pi$
$0.43 \pm 0.01$	2.9	$40 \pm 5$	3.76	1	0.036	$2^+$
$1.28 \pm 0.02$	2	$140 \pm 20$	4.54	2	0.28	3
$1.78 \pm 0.02$	0.4	$60 \pm 20$	5.00	1	0.012	1
				2	0.056	
$2.45 \pm 0.02$	0.5	$120 \pm 40$	5.61	1	0.017	2
				2	0.053	
$2.58 \pm 0.02$	0.6	$60 \pm 20$	5.73	1	0.008	3
				2	0.025	

<sup>a</sup> (1951BO45, 1958HU18).

<sup>b</sup> (1951BO45):  $R = 4.5 \times 10^{-13}$  cm.

<sup>c</sup> (1951BO45, 1955WI25, 1958HU18).

by (1952CO1B, 1954CO16); see (1955HU1B). See also (1955HI1C, 1956LA1C, 1957LA14, 1958HU18). At  $E_n = 14.5$  MeV the non-elastic cross section for natural boron is  $0.64 \pm 0.04$  b (1956FL1B).

9. (a)  $^{11}\text{B}(n, d)^{10}\text{Be}$   $Q_m = -9.010$   $E_b = 3.365$   
 (b)  $^{11}\text{B}(n, t)^9\text{Be}$   $Q_m = -9.564$   
 (c)  $^{11}\text{B}(n, \alpha)^8\text{Li}$   $Q_m = -6.636$   
 (d)  $^{11}\text{B}(n, p)^{11}\text{Be}$   $Q_m = -10.69$

The cross section for reaction (c) decreases from 27 mb at  $E_n = 12.6$  MeV to 16 mb at  $E_n = 20.0$  MeV (1956AR21). At  $E_n = 14.1$  MeV, the cross section for reaction (b) is  $15 \pm 5$  mb (1958WY67). Reaction (a) has not been reported. See also (1954HE1B). For reaction (d) see  $^{11}\text{Be}$ .

10.  $^{11}\text{B}(d, p)^{12}\text{B}$   $Q_m = 1.138$

Proton groups observed at  $E_d = 4.0$  to 8.5 MeV are listed in Table 12.3. No other groups are observed for  $E_x = 0$  to 3.15 MeV with an intensity greater than 4% of the ground-state

Table 12.3:  $^{12}\text{B}$  levels from  $^{11}\text{B}(\text{d}, \text{p})^{12}\text{B}$

$^{12}\text{B}^*$ (MeV)	$l_n$	$J^\pi$	$\frac{\Lambda}{2J+1}$ <sup>e</sup>
0	1	$1^+$ <sup>c</sup>	3.4
$0.947 \pm 0.005$	1	$3^+, 2^+, 1^+, (0^+)$	1.5, 2.1, 3.5, (10)
$1.674 \pm 0.011$	0	$2^-, (1^-)$	3.5, (5.8)
$2.618 \pm 0.011$			
$2.723 \pm 0.011$			
$3.383 \pm 0.009$	1	$3^+, 2^+, 1^+, (0^+)$	1.6, 2.2, 3.7, (11)
$3.76$ <sup>b</sup>			
$4.53$ <sup>b</sup>	2	$3^-$ <sup>d</sup>	3.8

<sup>a</sup> (1950BU1A, 1953EL12).

<sup>b</sup> (1953HO48).

<sup>c</sup> From  $^{12}\text{B}(\beta^-)^{12}\text{C}$ .

<sup>d</sup> From  $^{11}\text{B}(\text{n}, \text{n})^{11}\text{B}$ .

<sup>e</sup> Neutron capture probability; proportional to reduced width; values greater than 4 are regarded as improbable (1953HO48).

group, or from 3.15 to 3.5 MeV excitation with an intensity greater than 8% (1953EL12). See also (1954KH1A, 1955KH35). Angular distributions and absolute cross sections have been determined at  $E_d = 8$  MeV. Analysis by stripping theory leads to the assignments indicated in Table 12.3. Neutron capture probabilities are compared with expected single-particle values to eliminate certain spin possibilities (1953HO48). At  $E_d = 1.05$  MeV,  $\gamma$ -rays with energy 0.94 and 1.64 MeV are observed with an intensity ratio of 2 : 1 (1954TH1B). See also (1955RA41, 1956KA1A, 1956TA07, 1957CH25, 1957CO59).

11.  $^{11}\text{B}(\text{t}, \text{d})^{12}\text{B}$   $Q_m = -2.894$

Not reported.

12.  $^{11}\text{B}(\alpha, ^3\text{He})^{12}\text{B}$   $Q_m = -17.213$

Not reported.

13.  $^{12}\text{C}(\text{n}, \text{p})^{12}\text{B}$   $Q_m = -12.593$

See (1948JE03).

14.  $^{12}\text{C}(t, ^3\text{He})^{12}\text{B}$   $Q_m = -13.358$

Not reported.

15.  $^{13}\text{C}(\gamma, p)^{12}\text{B}$   $Q_m = -17.539$

See  $^{13}\text{C}$ .

16.  $^{13}\text{C}(d, ^3\text{He})^{12}\text{B}$   $Q_m = -12.045$

Not reported.

17.  $^{13}\text{C}(t, \alpha)^{12}\text{B}$   $Q_m = 2.274$

Not reported.

18.  $^{14}\text{C}(n, t)^{12}\text{B}$   $Q_m = -17.228$

Not reported.

19.  $^{14}\text{C}(p, ^3\text{He})^{12}\text{B}$   $Q_m = -17.993$

Not reported.

20.  $^{14}\text{C}(d, \alpha)^{12}\text{B}$   $Q_m = 0.358$   
 $Q_0 = 0.362 \pm 0.0015$  (1956DO41).

See also (1950HU72).

21.  $^{14}\text{N}(n, ^3\text{He})^{12}\text{B}$   $Q_m = -17.365$

Not reported.

22.  $^{15}\text{N}(n, \alpha)^{12}\text{B}$   $Q_m = -7.629$

See (1948JE03).

$^{12}\text{C}$   
(Fig. 20)

GENERAL:

*Theory* : See (1955FE1A, 1955HE1E, 1956CA1F, 1956EL1C, 1956GL1B, 1956HA1C, 1956HA1D, 1956KU1A, 1956MO1D, 1956NA1A, 1956PE1A, 1956RE1C, 1956WI1F, 1957BA1H, 1957BI1C, 1957HE1B, 1957KU58, 1957PA1A, 1957RE1A, 1957SA1B, 1958CA1G, 1958FR1C).

1.  $^7\text{Li}(^6\text{Li}, n)^{12}\text{C}$   $Q_m = 20.931$

See (1957NO17).

2. (a) $^9\text{Be}(^3\text{He}, n)^{11}\text{C}$	$Q_m = 7.565$	$E_b = 26.286$
(b) $^9\text{Be}(^3\text{He}, p)^{11}\text{B}$	$Q_m = 10.329$	
(c) $^9\text{Be}(^3\text{He}, \alpha)^8\text{Be}$	$Q_m = 18.911$	
(d) $^9\text{Be}(^3\text{He}, d)^{10}\text{B}$	$Q_m = 1.093$	

The yields and angular distributions of protons leading to the ground state and several excited states of  $^{11}\text{B}$  have been investigated by (1956AL1E:  $E(^3\text{He})$  up to 2.7 MeV), by (1955HO1D, 1956HO1C:  $E(^3\text{He}) = 2.0$  MeV), by ((1956WO1A, 1956WO1C, 1957JO1B) and E. Wolicki, private communication: 2 to 4.5 MeV) and by (D.R. Sweetman, private communication: 6.05 MeV). The yield rises rapidly to  $E(^3\text{He}) = 1.8$  MeV and remains approximately constant to 4.5 MeV, with no indication of resonance. Angular distributions show fore and aft asymmetry and vary only slowly with energy. At  $E(^3\text{He}) = 2$  MeV, it appears that both direct interaction and compound nucleus formation, involving interfering resonances with  $l \leq 3$ , may be taking place. At higher energies the forward peaking suggestive of direct interaction becomes more obvious. See also  $^{11}\text{B}$ . For reactions (a), (c) and (d), see  $^{11}\text{C}$ ,  $^8\text{Be}$  and  $^{10}\text{B}$ .

Table 12.4: Energy levels of  $^{12}\text{C}$ 

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma$ (keV)	Decay	Reactions
0	$0^+; 0$	$-^a$	stable	1, 3, 10, 11, 16, 17, 19, 20, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 37, 39, 40, 41, 42
$4.433 \pm 5$	$2^+; 0$	$0.01 - 0.02 \text{ eV}^b$	$\gamma$	3, 10, 16, 19, 20, 24, 25, 26, 27, 29, 30, 34, 39, 41
$7.656 \pm 7$	$0^+; 0$	$< 25$	$\alpha, (\gamma)$	3, 10, 16, 19, 24, 25, 26, 29, 30, 39
$9.63 \pm 14$	$(1^-); 0$	$30 \pm 8^a$	$\alpha$	10, 16, 24, 25, 26, 27, 29, 39, 42
$10.1 \pm 200$	$0^+; 0$	$\approx 2000$	$\alpha$	19, 25, 42
$10.84 \pm 50$				10, 16, 26
$11.1 \pm 100$				16
$11.81 \pm 50$				10, 16
$12.73 \pm 50$	$(1^+); 0$		$(\alpha), \gamma$	10, 16, 26, 29, 30
$(13.21 \pm 50)$				16
$13.30 \pm 50$				10, 16
$14.05 \pm 60$				10, 16
$15.11 \pm 10$	$1^+; 1$	0.069	$(\alpha), \gamma$	3, 10, 16, 20, 23, 25, 26, 34, 39
$15.62 \pm 60$				10, 16
$16.11 \pm 2$	$2^+; 1$	6	$\alpha, p, \gamma$	10, 12, 14, 16, 23
$16.58 \pm 15$	$2^-; (1)$	295	$\alpha, p, \gamma$	10, 12, 23
17.23	$1^-; (1)$	1160	$\alpha, p, \gamma$	12, 14, 22, 23
17.77	$(0^+)$	140	$\alpha, p$	12
18.37	$(2^+)$	280	$\alpha, p, \gamma$	12, 23
18.40		46	$p, p'$	14
18.85		90	$n, p, \gamma$	12, 13, 14
19.26		450	$n, p, \gamma$	12, 13, 14, 21
19.42		45	$p$	14
19.67		180	$n, p, (\gamma)$	13, 21
19.88		90	$p, p'$	14
20.27		180	$n, p, (\gamma)$	13, 14, 21
20.49		180	$(n), p, \gamma$	12, 21
20.65		180	$n, p, \gamma$	12, 13, 14, 21, 22, 26
21.34			$n, p, (\gamma)$	12, 13, 21, 22

Table 12.4: Energy levels of  $^{12}\text{C}$  (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\Gamma$ (keV)	Decay	Reactions
21.80		$\approx 4000$	n, ( $\alpha$ ), p, ( $\gamma$ )	12, 13, 21, 22, 23
22.55 $\pm$ 100			n, p, $\gamma$	12, 13, 20, 21, 22, 23
(22.8)			(n), ( $\alpha$ ), (p), ( $\gamma$ )	21, 22
(24.3)			n, ( $\alpha$ ), ( $\gamma$ )	13, 23
(25.4)			n, p, d	6, 13
(26.0)			n, $\alpha$ , (p), d, ( $\gamma$ )	5, 6, 8, 22
(26.4)			( $\alpha$ ), p, d	6, 8
(26.8)			$\alpha$ , p, d	6, 8
(27.0)			p, d	6
(27.2)			p, d	6
(27.4)			p, d	6
(29.4)		( $\alpha$ ), ( $\gamma$ )	23	

<sup>a</sup> For reduced width, see  $^{11}\text{B}(d, n)^{12}\text{C}$  and Table 12.8.

<sup>b</sup> See reactions 3, 20 and 24.

### 3. $^9\text{Be}(\alpha, n)^{12}\text{C}$

$$Q_m = 5.709$$

Neutron groups corresponding to states of  $^{12}\text{C}^*(0, 4.4, 7.6)$  have been observed with  $E_\alpha$  up to 5.3 MeV. At  $E_\alpha = 5.3$  MeV the forward yield of the group leading to the 7.6 MeV state is about  $\frac{1}{8}$  of that leading to the 4.4 MeV state: see (1952GU1A, 1955ST1C, 1956ME1B, 1957RI38); see also (1957DI1B).

The energy of the  $\gamma$ -ray from the first excited state is  $4.425 \pm 0.020$  (1954MI68),  $4.48 \pm 0.06$  MeV (1955BE1G) (both values corrected for Doppler shift). The internal pair conversion, coefficient indicates an E2 transition (1954MI68); the angular correlation of pairs admits M1 or E2, favoring the latter (1954HA07, 1956GO73, 1956GO1K, 1958AR1B). The angular distribution of  $\gamma$ -rays observed at several bombarding energies is consistent with  $J = 2^+$  (1955TA28, 1956PR1A). The n- $\gamma$  correlation at  $E_\alpha = 5.3$  MeV (thick target) is isotropic within 6.5% (1956ST1E: see also (1958TA05)). The mean lifetime of the 4.4 MeV level is  $(2.6 \pm 0.9) \times 10^{-14}$  sec, about one-eighth of the single-particle value for an E2 transition (1956DE22).

The 7.7 MeV state appears to decay predominantly into  $^8\text{Be} + \alpha$  (see  $^{12}\text{B}(\beta^-)^{12}\text{C}$  and  $^{12}\text{C}(\alpha, \alpha')^{12}\text{C}^*$ ). A gamma ray of energy 3.1 MeV has been reported by (1953BE1C, 1954UE1A, 1956ST1E, 1956ST1F, 1957ST1E), but (1955BE1G, 1957GO1C, 1957KR1A) find no evidence of the 7.7

MeV nuclear pairs which should accompany the decay to the ground state. (1955BE1G) estimate that at least 96% of the decays proceed to  ${}^8\text{Be} + \alpha$ ; (1957KR1A, 1958KR70) find  $< 1.6 \times 10^{-5}$  7 MeV pairs per 4.4 MeV  $\gamma$ -ray; assuming a population ratio of 1 : 8, this result yields  $\Gamma_{\pi}/\Gamma < 1.3 \times 10^{-4}$ . An upper limit of  $\frac{1}{600}$  for the ratio of 7.6/4.4 MeV pairs is reported by (1957GO1C). (1954DI1A) find no n- $\gamma$  coincidences other than those associated with the 4.4 MeV level. See also (1957RO1E).

At  $E_{\alpha} = 21.7$  and 175 MeV,  $\gamma$ -radiation from the 15 MeV,  $J = 1^+$ ;  $T = 1$  state (see  ${}^{12}\text{C}(p, p'){}^{12}\text{C}^*$ ) is reported (1954RA35, 1957WA04). At the higher energy, the ratio of 15 MeV to 4.4 MeV radiation is  $1.2 \times 10^{-2}$  (1957WA1F). See also (1954EL1B, 1955BR1A, 1955HA1E, 1956GO1N, 1957BR1J) and (1955MA1J; theor.).

$$4. {}^{10}\text{B}(d, \gamma){}^{12}\text{C} \quad Q_m = 25.195$$

At  $E_d = 0.95$  MeV, the upper limit to the capture cross section is  $0.1 \mu\text{b}$  (1955SA1B).

$$5. {}^{10}\text{B}(d, n){}^{11}\text{C} \quad Q_m = 6.473 \quad E_b = 25.195$$

The thin-target excitation function in the forward direction in the range  $E_d = 0.3$  to 4.6 MeV shows some indication of a broad resonance near  $E_d = 0.9$  MeV. Above  $E_d = 2.4$  MeV, the cross section increases rapidly to 210 mb/sr at 3.8 MeV, and then remains constant to 4.6 MeV (1954BU06, 1955MA76). Angular distributions seem to be dominated by the stripping process: see  ${}^{11}\text{C}$ . The yield of 6.5 MeV  $\gamma$ -rays has been measured at four bombarding energies between 0.8 and 2.2 MeV (1955SA1B). See also  ${}^{11}\text{C}$ .

$$6. {}^{10}\text{B}(d, p){}^{11}\text{B} \quad Q_m = 9.237 \quad E_b = 25.195$$

Absolute yields and angular distributions are reported for various proton groups by (1952EN19, 1954BU06, 1954PA28, 1956MA69, 1956VA17) for  $E_d = 0.18$  to 3.1 MeV. Although the excitation functions show several broad peaks, no clear resonances can be identified, and it must be assumed that many overlapping resonances are involved (1956MA69). Angular distributions indicate both stripping and compound nucleus processes even at low bombarding energies (1954PA1D). However, the  $p_1$  group, leading to  ${}^{11}\text{B}^*(2.1)$ , shows no stripping even at  $E_d = 3$  MeV; it is suggested that an orbital angular momentum selection rule is operative here (1956MA69: see  ${}^{11}\text{B}$ ). Absolute cross sections reported by (1954BU06, 1954PA28, 1956MA69) differ rather greatly.

The yields of 6.8 and 7.3 MeV  $\gamma$ -rays have been measured at four bombarding energies between 0.8 and 2.2 MeV (1955SA1B). At  $E_d = 1.70$  MeV,  $\theta(\text{lab}) = 58^\circ$ , the cross section for protons leading to the 6.76 MeV state is  $6.1 (\pm 15\%) \text{ mb/sr}$  (1956KA1A).

$$7. {}^{10}\text{B}(\text{d}, \text{d}){}^{10}\text{B} \qquad E_{\text{b}} = 25.195$$

See  ${}^{10}\text{B}$ .

$$8. {}^{10}\text{B}(\text{d}, \alpha){}^8\text{Be} \qquad Q_{\text{m}} = 17.819 \qquad E_{\text{b}} = 25.195$$

Excitation curves for ground state  $\alpha$ -particles have been measured for  $E_{\text{d}} = 0.9$  to  $2.6$  MeV at  $45^\circ$ ,  $90^\circ$  and  $150^\circ$ . Broad maxima are observed at  $1.0$ ,  $(1.4)$  and  $2.0$  MeV. At  $E_{\text{d}} = 0.91$  MeV, the angular distribution of  $\alpha_0$  particles shows a peaking in the forward direction ([1956MA69](#)). See also  ${}^8\text{Be}$ .

$$9. {}^{10}\text{B}(\text{t}, \text{n}){}^{12}\text{C} \qquad Q_{\text{m}} = 18.937$$

Not reported.

$$10. {}^{10}\text{B}({}^3\text{He}, \text{p}){}^{12}\text{C} \qquad Q_{\text{m}} = 19.702$$

Proton groups reported by ([1955BI26](#)) and ([1958MO99](#)) are listed in Table [12.5](#). A careful search, at  $E({}^3\text{He}) = 1.25$  MeV, reveals no other level in the range  $E_{\text{x}} = 4.4$  to  $7.7$  MeV ([1958MO99](#): region at  $E_{\text{x}} = 6.4$  MeV obscured). At  $E({}^3\text{He}) = 2.0$  MeV, the proton group leading to the  $15.11$  MeV level was found to be in coincidence with a  $15.10$  MeV  $\gamma$ -ray:  $\Gamma_{\gamma}/\Gamma$  for this level is  $0.77 \pm 0.20$ . The ratio of the width for  $\gamma$ -emission to the  $4.4$  MeV level to the ground state  $\Gamma_{\gamma}$  is  $\approx 0.03$ . The  $12.76$  MeV level also emits  $\gamma$ -rays:  $\Gamma_{\gamma}/\Gamma \approx 0.02$ , suggested  $J^{\pi} = 1^+$  ([1957GO1B](#)) and H.E. Gove, private communication). Coincidence studies by ([1958MO99](#)) lead to  $\Gamma_{\gamma}/\Gamma < 0.9\%$  for the  $7.7$  MeV level,  $\Gamma_{\gamma}/\Gamma = 3 \pm 1\%$  for the  $12.76$  MeV level, and  $50 \pm 25\%$  for the  $15$  MeV level. See also  ${}^{11}\text{B}(\text{d}, \text{n}){}^{12}\text{C}$ , ([1958BR1D](#), [1958SW63](#)) and  ${}^{13}\text{N}$ .

$$11. {}^{10}\text{B}(\alpha, \text{d}){}^{12}\text{C} \qquad Q_{\text{m}} = 1.351$$

$$\qquad Q_0 = 1.341 \pm 0.002 \text{ ([1956DO41](#))}.$$

$$\qquad Q_0 = 1.36 \pm 0.09 \text{ ([1956PI1A](#))}.$$

$$\qquad Q_0 = 1.39 \pm 0.01 \text{ ([1953SH64](#))}.$$

See also  ${}^{14}\text{N}$ .

$$12. \text{(a) } {}^{11}\text{B}(\text{p}, \gamma){}^{12}\text{C} \qquad Q_{\text{m}} = 15.958$$

$$\text{(b) } {}^{11}\text{B}(\text{p}, \alpha){}^8\text{Be} \qquad Q_{\text{m}} = 8.582 \qquad E_{\text{b}} = 15.958$$

Table 12.5:  $^{12}\text{C}$  states from  $^{10}\text{B}(^3\text{He}, \text{p})^{12}\text{C}$ 

A		B	
$E_x$ (MeV)	$\Gamma_\gamma/\Gamma$	$E_x$ (MeV $\pm$ keV)	$\Gamma_\gamma/\Gamma$
4.43	1	4.43	1
7.77		7.65	< 0.009
9.61		9.61	
10.75		10.90 $\pm$ 60	
11.83		11.84 $\pm$ 60	
12.76	0.02	12.69 $\pm$ 60	0.03 $\pm$ 0.01
13.31		13.30 $\pm$ 60	
13.97		14.05 $\pm$ 60	
15.10	0.77 $\pm$ 0.20	14.97 $\pm$ 60	0.50 $\pm$ 0.25
		15.62 $\pm$ 60	
16.04		16.10	
16.57		16.57 $\pm$ 60	

A: (1955BI26, 1957GO1B):  $E(^3\text{He}) = 0.9$  and  $2.0$  MeV, values  $\pm 0.1$  MeV.

B: (1958MO99):  $E(^3\text{He}) = 1.25$  MeV.

In the range  $E_p = 0$  to  $3$  MeV, five principal resonances occur, at  $E_p = 0.16, 0.67, 1.4, 2.0$  and  $2.6$  MeV (see Table 12.6). All except the second and fourth exhibit resonance for  $\alpha_0, \alpha_1, \gamma_0$  and  $\gamma_1$  (to  $^8\text{Be}^*(0, 2.9)$  and  $^{12}\text{C}^*(0, 4.4)$ ); at  $E_p = 0.67$  MeV, only  $\alpha_1, \gamma_1$  are resonant. It follows from angular momentum selection rules that resonances for  $\alpha_0$  must have the character  $J^\pi = 0^+, 1^-, 2^+, 3^- \dots$ ;  $J = 0^+$  is excluded by observation of  $\gamma_0$ .

The  $E_p = 0.16$  MeV resonance ( $^{12}\text{C}^*(16.11)$ ) is well established as  $J = 2^+$ ; probably the  $T = 1$  analogue of  $^{12}\text{B}^*(0.95)$ . The angular distribution of  $\alpha_0$  particle is strongly anisotropic at resonance and shows a  $(\cos \theta)$  term varying with energy near resonance. The assumption  $J = 2^+, l_p = 1$ , with interference from an s-wave state at higher energy gives a good account of the observed angular distributions from  $E_p = 0.13$  to  $0.3$  MeV. The channel spin ratio  $\chi = 0.42 \pm 0.02$ ; the relative amplitude of the interfering  $J = 1^-$  state is  $0.022 \pm 0.002$  (1952TH1B). The angular correlation of  $\alpha_1$  and the subsequent breakup of  $^8\text{Be}^*(2.9)$  also requires  $J = 2^+$ , with the ratio of reduced matrix elements for outgoing d to s-waves,  $B = 0.80$ , phase difference  $\cos \beta = 0.60$  (1955GE1A). The angular distribution of  $\gamma_1$  and of the following  $4.4$  MeV radiation is consistent with the scheme  $2^+(M1)2^+(E2)0^+$  with the channel spin ratio  $\chi = 0.42$  (1954GR1C); (1956CR1C) obtain  $\chi = 0.51 \pm 0.03$ . Angular distributions of the  $16$  MeV radiation,  $\gamma_0$ , require  $J = 2^+$ , with interference from a  $J = 1^-$  level at  $E_p = 1.4$  MeV (1954GR1C, 1956CR1C). ( $\gamma_0$  is not resonant at  $E_p = 0.67$  MeV, so this state cannot be involved here.) The resonant energy is  $163.1 \pm 0.2$  keV;

Table 12.6: Resonances in  $^{11}\text{B}(p, \gamma)^{12}\text{C}$  and  $^{11}\text{B}(p, \alpha)^8\text{Be}$ 

$E_p$ (MeV)	$\Gamma_{\text{lab}}$ (keV)	$\sigma(\gamma_{16})^a$ ( $\mu\text{b}$ )	$\sigma(\gamma_{12})^a$ ( $\mu\text{b}$ )	$\sigma(\alpha_0)^e$ (mb)	$\sigma(\alpha_1)^e$ (mb)	$\Gamma_{\gamma_{16}}^e$ (eV)	$\Gamma_{\gamma_{12}}^e$ (eV)	$\Gamma_{\alpha_0}^e$ (keV)	$\Gamma_{\alpha_1}^e$ (keV)	$\Gamma_p^e$ (keV)	$^{12}\text{C}^*$ (MeV)	$J^\pi; T$
0.163	7	5.5	152	0.2	10	$\lesssim 3$	70	0.1	5	0.005	16.11	$2^+; 1$
0.675 <sup>b</sup>	322	(< 2.3) <sup>g</sup>	48	(< 0.2) <sup>g</sup>	600	$\lesssim 0.5$	15	$\lesssim 0.05$	150	150	16.58	$2^-; (1)$
1.388 <sup>b</sup>	1270	35	18	6	150	40	20	7	200	1000 <sup>h</sup>	17.23	$1^-; (1)$
1.98 <sup>c,d</sup>	150	non-res.	non-res.	[8] <sup>f</sup>	[39] <sup>f</sup>						17.77	$(0^+)$
2.63 <sup>c,d</sup>	300	weak	res.	[16] <sup>f</sup>	[180] <sup>f</sup>						18.37	$(2^+)$
3.13 <sup>d</sup>	100	weak	res.								18.83	
3.55 <sup>d</sup>	500	res.	res.								19.21	
4.94 <sup>d</sup>	200	non-res.	res.								20.49	
5.12 <sup>d</sup>	200	non-res.	res.								20.65	

<sup>a</sup> (1953HU29); ratio  $\sigma(\gamma_{16})/\sigma(\gamma_{12}) = 3.3 \pm 1\%$  at  $E_p = 0.16$  MeV (1956CR1C).

<sup>b</sup> (1953HU29).

<sup>c</sup> (1955HO48).

<sup>d</sup> (1955BA22).

<sup>e</sup> (1953BE61).

<sup>f</sup> (1953PA1B): normalized at  $E_p = 1.4$  MeV. See also (1955HO48).

<sup>g</sup> Non-resonant.

<sup>h</sup> According to (1957DE11),  $\Gamma_p \approx 50$  keV: see  $^{11}\text{B}(p, p)^{11}\text{B}$ . If this value is used,  $\alpha$ -widths should be increased by a factor of 6, and  $\gamma$ -widths by a factor of 20.

$\Gamma_{\text{lab}} = 6.5 \pm 0.6$  keV (see (1955AJ61)). The very small  $\alpha$ -width suggests  $T = 1$  (1953BE61).

For the  $E_p = 1.4$  MeV state ( $^{12}\text{C}^*(17.23)$ ), the possible assignments are  $1^-$  (s-wave),  $2^+$  (p-wave),  $1^-$  or  $3^-$  (d-wave); d-wave formation would seem to be excluded by the observed width. As indicated above,  $J = 1^-$  appears to be required to account for the interference at lower energies in  $\alpha_0$  and  $\gamma_0$ ; known higher resonances are probably too narrow to produce the observed effects. (1957DE11) find that the  $\alpha_0$ -distributions for  $E_p = 0.6$  to 1.4 MeV are well accounted for by the assumption of s-wave formation of  $J = 1^-$  through channel spin ( $\chi = 0$ ) with a relative d-wave amplitude  $A = 0.5$ , and interference by the 2.6 MeV,  $J = 2^+$ , state, with relative amplitude  $C = 0.25$ . A qualitative fit to the behavior of  $\alpha_1$  can be obtained with the same assumptions (1957DE11). Angular distributions of  $\gamma_0$  at  $E_p = 1.4$  MeV admit either  $J = 2^+$  or  $1^-$ ; for the latter, however, formation in channel spin 2 ( $\chi = \infty$ , d-waves) is required (1955GO10). The angular correlation of internal pairs indicates E1 for  $\gamma_0$  (1956GO1K, 1956GO1N, 1958AR1B). The large E1 width suggests  $T = 1$  for this state (1953BE61).

The  $E_p = 0.67$  MeV state ( $^{12}\text{C}^*(16.58)$ ) may be formed by s- or p-waves; d-waves are excluded by the width (1953BE61). The angular distribution of  $\alpha_1$  at  $E_p = 0.64$  and 0.93 MeV indicates s-wave formation: if  $J = 2^-$  is assumed, the d-wave admixture is  $< 10\%$ . The correlation of  $\alpha_1$  with the subsequent  $^8\text{Be}^*(2.9)$  breakup is consistent with  $J = 2^-$  and excludes  $1^-$ ; an appreciable f-wave admixture in outgoing  $\alpha_1$ -particles is indicated (1957DE11). Correla-

tion results at  $E_p = 270$  keV can be accounted for by  $J = 2^-$  with interference from the  $1^-$ ,  $E_p = 1.4$  MeV state (1955GE1A, 1957DE11). The angular distribution of  $\gamma_1$  is reported to require  $J = 2^+$ , with interference from the  $1^-$ ,  $E_p = 1.4$  MeV state (1954GR1B): according to (1957DE11), however, the distributions observed by (1954GR1C, 1955GO10) can equally well be ascribed to  $J = 2^-$ , with interference from a broad, even parity state, possibly at  $E_p = 2.0$  or  $2.6$  MeV (see, however, (1954GI1B)). The angular correlation of internal pairs indicates E1 for the  $\gamma_1$  radiation (1956GO1K, 1956GO1N, AR57). The relatively large E1 width suggest  $T = 1$  for this state (1953BE61).

The  $E_p = 2.0$  MeV level is reported to be resonant for  $\alpha_0$  and  $\alpha_1$ ; the relative weakness of  $\alpha_1$  suggests  $J = 0^+$  (1953PA1B). These seems to be no clear indication of resonance for  $\gamma_0$  or  $\gamma_1$  at this energy (1955GO10: see also (1953HU29)). At  $E_p = 2.65$  MeV, resonance occurs for  $\alpha_0$ ,  $\alpha_1$  (1953PA1B) and, weakly, for  $\gamma_0$ ,  $\gamma_1$  (1955BA22, 1955GO10). A large  $P_2$  coefficient in the angular distribution of  $\gamma_0$  suggests  $J = 2^+$  (1955GO10). (1955HO48) find  $E_p = 1.98$  and  $2.61$  MeV for the resonant energies for  $\alpha_0$ . Additional resonances for  $\gamma_0$  and  $\gamma_1$ , reported by (1955BA22) are listed in Table 12.6. (1959GE33) have examined the excitation function for ground-state transitions from  $E_p = 4$  to  $7.7$  MeV. The experiment locates the maximum of the  $^{12}\text{C}$  giant resonance at  $E_x = 22.55 \pm 0.1$  MeV but does not resolve individual levels. Two additional peaks, at  $E_x = 21.4$  and  $22.1$  MeV are suggested. The maximum value of  $\sigma(\gamma, p)$  is calculated to be  $29 \pm 5$  mb.

An upper limit for the total cross section (average value,  $E_p = 1.7$  to  $4.0$  MeV) for the production of nuclear pairs with  $E_\pi = 6.5$  to  $9.5$  MeV is  $0.03 \mu\text{b}$  (1955BE62). See also (1955AJ61, 1956MA1T), (1957SI1B; theor.) and  $^8\text{Be}$ .

$$13. \text{}^{11}\text{B}(p, n)^{11}\text{C} \qquad Q_m = -2.764 \qquad E_b = 15.958$$

Observed maxima in the (p, n) cross section are listed in Table 12.7 (1951BL1A, 1955BA22, 1957KA1C, 1959GI47). The region covered is characterized by considerable overlapping of resonances (1959GI47). See also (1956KO1D, 1958MA1F, 1958TA03).

$$14. \text{(a) } ^{11}\text{B}(p, p)^{11}\text{B} \qquad E_b = 15.958$$

$$\text{(b) } ^{11}\text{B}(p, p')^{11}\text{B}^*$$

Absolute elastic scattering cross sections are reported for one angle for  $E_p = 0.6$  to  $2.0$  MeV by (1956TA16), for four angles for  $E_p = 0.3$  to  $1.0$  MeV by (1957DE11). A pronounced anomaly is observed near  $E_p = 0.67$  MeV at all angles; the level is therefore formed by s-waves. The  $0.3$  to  $1.0$  MeV results are well accounted for by two resonances:  $E_p = 0.67$  MeV, s-wave,  $J = 2^-$ ,  $\Gamma = 0.33$  MeV,  $\Gamma_p/\Gamma = 0.5$ , d-wave  $< 10\%$ , and  $E_p = 1.4$  MeV, s-wave,  $J = 1^-$ ,  $\Gamma = 1.27$  MeV,  $\Gamma_p/\Gamma = 0.05$  (1957DE11). (The reported  $\Gamma_p/\Gamma$  for the  $1.4$  MeV resonance appears to be inconsistent with the values  $0.8$  or  $0.2$  derived by (1953BE61) from (p,  $\gamma$ ) and (p,  $\alpha$ ) cross sections.) (1956TA16)

Table 12.7: Maxima in yields of  $^{11}\text{B}(p, n)^{11}\text{C}$  and  $^{11}\text{B}(p, p')^{11}\text{B}^*$

A	B		C	D		E		$^{12}\text{C}^*$
$E_p$ (MeV)	$E_p$ (MeV)	$\Gamma$ (keV)	$E_p$ (MeV)	$E_p$ (MeV)	$\sigma$ (mb)	$E_p$ (MeV)	$\Gamma$ (keV)	(MeV)
						2.664	48	18.40
	3.17	100		3.18	25	3.15	100	18.87
3.7	3.65	500		3.67	62	3.4	500	19.31
						3.78	50	19.42
	4.05	200						19.67
	4.70	200	$\approx 4.5$	4.70	55	4.28	100	19.88
5.18				5.10	105	4.68	200	20.27
5.87						5.13	200	20.63
6.37								21.34
			6.6					21.81
			8.8					22.0
			10.1					24.0
								25.3

A: (p, n) (1951BL1A): stacked foils.

B: (p, n) (1955BA22):  $\theta = 0 - 15^\circ$ .

C: (p, n) (1957KA1C): stacked foils.

D: (p, n) (1959GI47): total cross section (estimated from curve).

E: (p, p') (1955BA22).

find no rapid variation in cross section near  $E_p = 2.0$  MeV. The absence of a detectable anomaly near  $E_p = 0.16$  MeV confirms the small value of  $\Gamma_p$  assumed for this resonance;  $\Gamma_p < 200$  eV (J.C. Overley, private communication). See also (1956KI54).

Maxima in the yield of 2.1 MeV  $\gamma$ -radiation from  $^{11}\text{B}^*(2.1)$  are observed at  $E_p = 2.664$  MeV,  $\Gamma = 48$  keV: (1953HU29, 1955BA22) and at  $E_p = 3.15, 3.4, 3.78, 4.28, 4.68$  and 5.13 MeV (1955BA22: see Table 12.7). (Judging from the width, the 2.66 MeV resonance is not that observed, e.g., in  $^{11}\text{B}(p, \gamma)^{12}\text{C}$ .)

15.  $^{11}\text{B}(p, d)^{10}\text{B}$

$$Q_m = -9.237$$

$$E_b = 15.958$$

See  $^{10}\text{B}$ .

16.  $^{11}\text{B}(\text{d}, \text{n})^{12}\text{C}$

$$Q_{\text{m}} = 13.731$$

$$Q_0 = 13.63 \pm 0.05 \text{ (1957BI78)}.$$

Reported neutron groups are listed in Table 12.8. The group corresponding to the 7.6 MeV state is weak, relative to neighboring groups, at all bombarding energies investigated, and the stripping pattern is poorly developed. The relative weakness of the 7.6 MeV state in this reaction and in the  $^{12}\text{C}(\text{e}, \text{e}')^{12}\text{C}^*$  and  $^{12}\text{C}(\text{p}, \text{p}')^{12}\text{C}^*$  reactions is attributed to lack of parentage overlap with the ground state of  $^{12}\text{C}$  (1955LA1C). At  $E_{\text{d}} = 0.92$  MeV, there is no indication of a state in the range  $E_{\text{x}} = 5.1 - 6.6$  MeV: the upper limit of the intensity of the corresponding neutron group is  $\lesssim 1\%$  of the intensity of the group corresponding to the 4.4 MeV state (1957BI78). For  $E_{\text{d}} = 1.1$  to 2.0 MeV only the groups corresponding to  $^{12}\text{C}^*(4.4, 12.76)$  are accompanied by  $\gamma$ -radiation; an upper limit for (n,  $\gamma$ ) coincidences from  $^{12}\text{C}^*(7.6)$  is 0.2% of  $^{12}\text{C}^*(4.4)$  (1958DA11, 1958NE38, 1959NE1A).

Angular distributions of the neutrons to the first four states of  $^{12}\text{C}$  have been reported for a number of energies in the range  $E_{\text{d}} = 0.5$  to 10 MeV. At the higher energies, the distributions are understood in terms of simple stripping theory (except for the 7.6 MeV state). At the lower energies,  $E_{\text{d}} = 0.5$  to 5 MeV, a good account of the angular distributions of ground-state neutrons is obtained with the theory of (1957OW03) which includes not only stripping of the deuteron but also the possibility of stripping a neutron from  $^{11}\text{B}$ , and the interference between the two processes. The relative probability of the exchange stripping increases with energy until the Coulomb barrier is surmounted. The exchange process seems to involve s-wave deuteron capture by a  $^{10}\text{B}$  core with  $J = 1^+$  (1956PR1B, 1957AM48, 1957OW03: see also (1959NE1A)). Angular distributions at  $E_{\text{d}} = 9$  MeV indicate odd parity for the 9.6 MeV state (1956MA83: see also (1954GR53) and Table 12.8). For other work on angular distributions, see (1957AM48, 1958AM13:  $E_{\text{d}} = 0.50$  to 1.15 MeV), (1955WA30:  $E_{\text{d}} = 0.6$  MeV), (1955IH1B:  $E_{\text{d}} = 0.69$  MeV), (1954GR53:  $E_{\text{d}} = 0.85$  MeV), (1957BI78:  $E_{\text{d}} = 0.92$  MeV), (1956PR1B:  $E_{\text{d}} = 1.5$  to 5 MeV), (1953GI05:  $E_{\text{d}} = 8.1$  MeV), and (1957ZE1A:  $E_{\text{d}} = 10$  MeV). See also (1954BU06, 1956BO1F, 1956BO43, 1956KO1E, 1957RA1A) and (1955MA1J, 1958ED1C; theor.).

In the range  $E_{\text{d}} = 1.0$  to 5.5 MeV, two slow neutron thresholds are observed at  $1.627 \pm 0.004$  MeV ( $E_{\text{x}} = 15.11 \pm 0.01$  MeV) and near 4.1 MeV (broad;  $E_{\text{x}} = 17.23$  MeV) (1955MA76). Gamma rays are observed with  $E_{\gamma} = 4.44 \pm 0.05$  (1951RU1A) and  $12.8 \pm 0.3$  MeV (1958KA31). If the latter  $\gamma$ -ray is properly attributed to decay of the 12.8 MeV level, it is not clear why the level should so decay in view of its instability with respect to  $^8\text{Be}$  and  $^8\text{Be}^*(2.9)$  (1958KA31: see also  $^{10}\text{B}(^3\text{He}, \text{p})^{12}\text{C}$ ). A 15.1 MeV  $\gamma$ -ray is observed with a threshold of  $E_{\text{d}} = 1633 \pm 3$  keV, attributed to the first  $T = 1$  state of  $^{12}\text{C}$  at 15.11 MeV. The observed width is  $< 2$  keV. A search for  $\alpha$ -particles to  $^8\text{Be}$  and  $^8\text{Be}^*$  gives  $\Gamma_{\alpha}/\Gamma_{\gamma} < 1.5$ . At  $E_{\text{d}} = 2.96$  MeV the cross section for production of the 16.1 MeV  $T = 1$  state is  $< 1$  mb/sr (1958KA31). See also (1955AJ61) and (1957WA04).

17.  $^{11}\text{B}(^3\text{He}, \text{d})^{12}\text{C}$

$$Q_{\text{m}} = 10.464$$

Table 12.8: Energy levels of  $^{12}\text{C}$  from  $^{11}\text{B}(d, n)^{12}\text{C}$

$^{12}\text{C}^*$ <sup>a</sup> (MeV)	$l_p$	$(2J + 1)\gamma^2$ ( $10^{-13}$ MeV-cm)	$(2J + 1)\theta^2$
0	1 <sup>e</sup>	1.6 <sup>e</sup>	0.11
4.38 ± 0.07	1 <sup>c,e</sup>	2.0 <sup>e</sup>	0.14
7.57 ± 0.11	d		
9.6 ± 0.1	2 <sup>c,e</sup>	1.9 <sup>e</sup>	0.13
10.8 ± 0.1			
11.1 ± 0.1			
11.74 ± 0.08			
12.76 ± 0.08			
13.21 ± 0.05 <sup>b</sup>			
13.36 ± 0.05 <sup>b</sup>			
(14.16 ± 0.05)			
15.09 ± 0.03			
(15.52 ± 0.03)			
16.07 ± 0.03			

<sup>a</sup> (1952JO10, 1954GR53, 1957BI78).

<sup>b</sup> May represent a single level at 13.3 MeV (1952JO10).

<sup>c</sup> (1954GR53).

<sup>d</sup> Angular distribution shows peaking forward and backward: (see (1954GR53, 1957BI78)).

<sup>e</sup> (1956MA83:  $E_d = 9$  MeV).

At  $E(^3\text{He}) = 4.5$  MeV, the ground state deuteron group is strongly peaked in the forward direction (1957HO61).

18.  $^{11}\text{B}(\alpha, t)^{12}\text{C}$   $Q_m = -3.855$

Not reported.

19.  $^{12}\text{B}(\beta^-)^{12}\text{C}$   $Q_m = 13.376$

Table 12.9: Branching in  $^{12}\text{B}(\beta^-)^{12}\text{C}$  (1958CO66)

$^{12}\text{C}^*$ (MeV)	$J^\pi$	Branching fraction (%)	$\log ft^a$
0	$0^+$	97	$4.1 \pm 0.1$
4.43	$2^+$	$1.5 \pm 0.3^b$	$5.1 \pm 0.2$
$7.653 \pm 0.008$	$0^+$	$1.3 \pm 0.4$	$4.2 \pm 0.3$
$10.1 \pm 0.2^c$	$0^+$	$0.13 \pm 0.04$	$4.1 \pm 0.4$

<sup>a</sup> Based on  $\tau_{1/2} = 20.6$  msec.

<sup>b</sup> (1958KA31):  $1.4 \pm 0.4$  %. (1956TA07):  $1.7 \pm 0.4$  %. (1958VE20):  $6 \pm 3$  %.

<sup>c</sup>  $\Gamma = 2.5$  MeV,  $\theta_\alpha^2 = 1.5$ .

The half-life is  $20.34 \pm 0.5$  msec (weighted mean of (1955AJ61, 1956NO1A, 1957CO57, 1958KR65, 1958VE20, 1959KR1B), excluding (1948JE03)).  $E_\beta(\text{max}) = 13.40 \pm 0.05$  MeV (1958VE20). Branches are observed leading to  $^{12}\text{C}^*(0, 4.4, 7.65, 10.1)$ : see Table 12.9. The fact that the transition to the  $0^+$  ground state is allowed establishes  $J = 1^+$  for  $^{12}\text{B}$ . This assignment is confirmed by the allowed character of the transition to  $^{12}\text{C}^*(4.4)$ ,  $J = 2^+$ . The 7.7 MeV level decays mainly by  $\alpha$ -emission to  $^8\text{Be}(0)$  with  $Q = 278 \pm 4$  keV;  $E_x = 7.653 \pm 0.008$  MeV (1957CO59). Gamma transitions with  $E_\gamma > 6$  MeV accompany  $< 10^{-4}$  of all  $\beta$ -decays (1956KA1A, 1958KA31); an upper limit of  $3 \times 10^{-5}$  is obtained in a search for  $\beta$ - $\gamma$  (7.6) coincidences (1958KA14). Upper limits for (3.2 + 4.4) MeV cascades are given as  $(0.4 \pm 2) \times 10^{-3}$  and  $10^{-5}$  of all decays by (1956TA07) and (1958KA14) respectively. It follows that the relative partial width of the 7.6 MeV level is  $< 3 \times 10^{-3}$  for 7.6 MeV  $\gamma$ -rays and  $< 10^{-3}$  for 3.2 MeV  $\gamma$ -rays (see also  $^9\text{Be}(\alpha, n)^{12}\text{C}$ ). Since the  $\beta$ -transition is allowed, the 7.7 MeV state has  $J = 0^+, 1^+$  or  $2^+$ ;  $J = 1^+$  is ruled out by the  $\alpha$ -decay. The preponderance of  $\alpha$ -decay over  $\gamma$ -decay speaks for  $J = 0^+$  (1957CO59).

The 10.1 MeV level decays mainly via  $\alpha$ -emission to  $^8\text{Be}(0)$ ; transitions to  $^8\text{Be}^*(2.9)$  amount to  $< 4\%$ . The c.m. width is about 2.5 MeV (after removing the  $E_\beta^5$  factor): the best account of the observed  $\alpha$ -spectrum is obtained with  $J = 0^+$ ,  $E_\lambda = 10.4$  MeV,  $\theta_\alpha^2 = 1.5$ ,  $R = 5.21 \times 10^{-13}$  cm (1958CO66).

The  $\beta$ -spectrum has been reported by (1950HO01, 1958VE20). (1958GE1C) discusses a possible distortion of high-energy  $\beta$ -spectra in axial-vector coupling due to a “weak-magnetic” interaction.

See also (1957CH25), (1955JA1C, 1957FE1C, 1959SC1B; theor.).

## 20. $^{12}\text{C}(\gamma, \gamma')^{12}\text{C}^*$

Table 12.10: Parameters of the 15.1 MeV  $^{12}\text{C}$  level from  $^{12}\text{C}(\gamma, \gamma')^{12}\text{C}^*$

	(1957HA13)	(1959GA09)
peak absorption, $\sigma_n^0$ (b)	$22.2 \pm 2.2$	$29.7 \pm 1.1$
integrated scattering, $\int \sigma_s dE$ (MeV · mb)	$1.90 \pm 0.27$	$2.33 \pm 0.19$
$\Gamma(\text{total})$ (eV)	$79 \pm 16$	$64.5 \pm 10.4$
$\Gamma_\gamma(15 \rightarrow \text{g.s.})$ (eV)	$54.5 \pm 9.3$	$59.2 \pm 9.7$
$\Gamma_\gamma(15 \rightarrow 4.4)$ (eV)	$< 5.5$	$3.2 \pm 2.5$
$\Gamma_\alpha$ (eV)	$18 - 25^a$	$2.1 \pm 3.2$

<sup>a</sup>  $\Gamma_\alpha/\Gamma < 1.5$  from  $^{11}\text{B}(\text{d}, \text{n})^{12}\text{C}$  (1958KA31).

The lifetime of the 4.4 MeV state has been determined by resonance scattering and resonant absorption (of  $\gamma$ -radiation from  $^{15}\text{N}(\text{p}, \alpha)^{12}\text{C}^*$ ) as  $\tau_{\text{mean}} = (6.5 \pm 1.2) \times 10^{-14}$  sec (1958RA14).

Excitation of the 15.1 MeV level by bremsstrahlung is reported by (GA57B, 1957HA13, 1959GA09). From measurement of the yield of resonance scattered radiation and of the self-absorption coefficient, the integrated scattering cross section  $\int \sigma_s dE$ , and the peak absorption cross section  $\sigma_n^0$  are deduced. The resulting values for partial widths are given in Table 12.10. It is noted the  $\gamma$ -ray width for the ground state transition is near the single-particle M1 value of 65 eV, and that the very small  $\alpha$ -width strongly suggests  $T = 1$  for this level (1957HA13). The scattering angular distribution indicates dipole radiation (1956LE1E, GA57B, 1959GA09). The strength of the M1 radiation also indicates  $T = 1$ : see (1958MO17). Inelastic scattering in the giant-resonance region has been studied by (1959GA09). See also (1957GO1F, 1958AX1A, 1958MC1D).

## 21. $^{12}\text{C}(\gamma, \text{n})^{11}\text{C}$

$$Q_m = -18.722$$

The cross section for production of  $^{11}\text{C}$  exhibits a broad peak at  $E_\gamma = 22.5$  MeV,  $\Gamma \approx 4$  MeV,  $\sigma_{\text{max}} = 8.3$  mb (1955BA63). Other reported values for  $\sigma_{\text{max}}$  are summarized by (1957CO57: note a 10% correction in this work). See also (1957CA1D). At high energies, the cross section exhibits a long tail, falling off approximately as  $E_\gamma^{-3}$ . The integrated cross section to  $E_\gamma = 250$  MeV is 80 MeV·mb, accounting for about  $\frac{1}{3}$  of the sum-rule limit for all absorption processes. It is noted that the relative prominence of the high-energy tail is not a general feature of  $(\gamma, \text{n})$  reactions in heavy elements (1955BA63, 1957CO57). Comparison of  $(\gamma, \text{n})$  and  $(\text{e}, \text{n})$  cross section for 28 to 145 MeV are consistent with the assumption that the transitions are predominantly E1 (1958BA60). The angular distribution of photoneutrons at the giant resonance is  $W(\theta) = 1 + (1.35 \pm 0.88) \sin^2 \theta$ , indicating considerable emission of neutrons with  $l > 0$  (1956FA1B). See  $^{12}\text{C}(\gamma, \text{p})^{11}\text{B}$  and (1954TE1A, 1955MO1B, 1957BA1K; theor.).

Discontinuities in the yield function are reported to indicate levels at 19.3, 19.8, 20.1, 20.5, 20.7, 21.1, 21.6, 22.4, and 22.8 MeV (1954GO39, 1954KA1A). The first two are given as  $19.09 \pm$

0.05 and  $19.55 \pm 0.05$  MeV by (1955SP1A). Eighteen discontinuities observed between  $E_\gamma = 18.90$  and 22.88 MeV are tabulated by (1958KA1D). A search for resonance absorption near  $E_\gamma = 22.8$  MeV indicates a width  $\Gamma > 580$  keV, in apparent contradiction of the activation results (1956TZ1A). (1958WO1B), using monochromatic gamma rays, find no evidence of fine structure in the total cross section from  $E_\gamma = 20.3$  to 20.8 MeV. The upper limit is, however, not in conflict with the recent report of (1958KA1D). See also (1955JO1B, 1955SA1F, 1958BA1K, 1958SM1A).

22.  $^{12}\text{C}(\gamma, \text{p})^{11}\text{B}$   $Q_m = -15.958$

The cross section exhibits a giant resonance at  $E_\gamma = 21.5 \pm 0.5$  MeV,  $\Gamma = 1.7 \pm 0.5$  MeV (1951HA1C). The peak cross section is 22 mb, and the integrated cross section to 24 MeV is 56 MeV-mb (1956CO59): compare  $^{11}\text{B}(\text{p}, \gamma)^{12}\text{C}$  (1959GE33). The photoproton spectrum shows the general features of the inverse reaction,  $^{11}\text{B}(\text{p}, \gamma)^{12}\text{C}$ , and suggests resonances at  $E_\gamma = 17.3$ , (20.8), 22.6, and (23.1) MeV (1956CO59: see, however, (1958WO1B)). (1957LI1A) finds indications of peaks at  $E_\gamma = 21.8$ , 22.6, 23.3 and 25.8 MeV. An absolute measurement, based on  $\sigma(^{12}\text{C}^*(\gamma, 3\alpha))$  for  $E_\gamma = 17.6$  MeV yields  $\sigma(\gamma, \text{p}) = 1.19 \pm 0.21$  mb, in good agreement with the value  $1.09 \pm 0.16$  mb calculated from the inverse reaction (1956MA1T). See also (1956GO1G, 1958CH31, 1958PE1A, 1958WH1A).

Angular distributions of photoprotons show a pronounced  $90^\circ$  peaking, somewhat skewed in the forward direction (1952HA1B, 1953HE1B, 1955JO1B, 1956KL19, 1957DO1A, 1957LI1A, 1957MI1A). Such distributions are inconsistent with s-wave proton emission from  $J = 1^-$ ;  $^{12}\text{C}$  compound states formed by E1 absorption and suggest a direct interaction involving independent-particle states:  $L$ - $S$  coupling seems to be favored (1955MA1H). The angular distributions of  $^{12}\text{C}(\gamma, \text{n})$  and  $(\gamma, \text{p})$  are evidently quite similar, as expected on the assumption of charge independence; the difference of about a factor of 2 in total cross section is ascribed to a 1%  $T = 0$  admixture in the intermediate state (1957BA1K). See also (1953HE1B, 1957CH24, 1958BA1M, 1958BA30, 1958PA1B, 1958PE1B, 1958SM1A) and (1955MO1B, 1957SI1B; theor.).

23.  $^{12}\text{C}(\gamma, 3\alpha)$   $Q_m = -7.281$

Maxima in the yield of 3-prong stars are reported at  $E_\gamma = 17.3$ , 18.3, 21.9, 24.3 and 29.4 MeV; some evidence of fine structure is also found. The integrated cross section is  $1.21 \pm 0.16$  MeV-mb for  $E_\gamma < 20.5$  MeV,  $2.8 \pm 0.4$  MeV-mb for  $20.5 \leq E_\gamma < 42$  MeV, and  $< 0.2$  MeV-mb for  $42 \leq E_\gamma < 60$  MeV (1953GO13, 1955GO59). (1955CA19) summarize cross section measurements for the  $^7\text{Li}(\text{p}, \gamma)$  radiation and find evidence for a resonance near  $E_\gamma = 12.3$  MeV and possibly others at 15 and 16 MeV. According to (1955JO1C), peaks occur at  $E_\gamma = 14.7$ , 15.8, 16.6, 18.3, 24.3, and  $> 29$  MeV, in fair agreement with (1953GO13). Absolute cross sections are reported for  $E_\gamma = 13$  to 30 MeV, and integrated cross sections agree well with (1953GO13, 1955JO1C).

See also (1953DA1A, 1953GU1A, 1953MI31, 1955HA1D). According to (1955GO59), the three-body reaction is not involved for  $E_\gamma < 40$  MeV (see, however, (1953MI31, 1954CH1B)). The reaction  $^{12}\text{C}(\gamma, \alpha)^8\text{Be}^*(\text{p})^7\text{Li}$  is reported by (1956LI05).

Studies of angular distributions indicate that for  $E_\gamma = 12$  to 15.6 MeV, the reaction involves mainly E2 absorption ( $^{12}\text{C}^*$ :  $J = 2^+$ ;  $T = 0$ ); from 15.6 to 20 MeV both E1 ( $J = 1^-$ ;  $T = 1$ ) and E2 ( $J = 2^+$ ;  $T = 1$ ), and for  $E_\gamma > 20$  MeV, mainly E1 ( $J = 1^-$ ;  $T = 1$ ). Significant E2 absorption ( $J = 2^+$ ;  $T = 0$ ) also occurs for  $E_\gamma = 20$  to 25 MeV (1955GO59: see, however, (1951TE1A, 1953GE1B)). See also (1953LI1C, 1954GR1B, 1955CO1A, 1955SO1B, 1955TH1A, 1956MA1T, 1957MU1C).

#### 24. $^{12}\text{C}(\text{e}, \text{e}')^{12}\text{C}$

Both elastic and inelastic scattering angular distributions have been studied at  $E_e = 80, 150$  and 187 MeV by (1955FR1G, 1956FR27) and at 420 MeV by (1958EH1B). The elastic data are well accounted for by a modified Gaussian charge distribution of r.m.s. radius  $2.50 \times 10^{-13}$  cm, derived from a harmonic well with a characteristic length parameter of  $1.68 \times 10^{-13}$  cm (1956FR27, 1958EH1B). See also (1953HO79, 1956FE1B, 1956HO93, 1957HO1E, 1958EH1A, 1958RA43).

Inelastic peaks corresponding to  $^{12}\text{C}^*(4.4, 7.7, 9.6)$  are observed, in addition to some unresolved structure near 11 MeV. There is no indication of the 15.1 MeV level. The observed angular distributions agree well with shell-model calculations of (1955RA1D, 1956MO1E, 1956TA1C, 1957TA1B) with a harmonic well of r.m.s. radius  $2.40 \times 10^{-13}$  cm. Predicted absolute cross sections are low by a factor of 2 in  $L$ - $S$  coupling, 6 in  $j$ - $j$  coupling; it is presumed that some collective modes of excitation are involved.

Excitation of the 4.4 MeV level is electric; a width of  $(12.5 \pm 2.5) \times 10^{-3}$  eV,  $\tau_m = (0.53 \pm 0.11) \times 10^{-13}$  sec, is obtained (1956HE83). The 7.7 and 9.6 MeV levels are also electrically excited; the angular distributions indicate either monopole or quadrupole transitions,  $J = 0^+, 2^+$ . The matrix element for the 7.7 MeV E0 transition is 50 mb, in good agreement with that observed for the  $^{16}\text{O}$  monopole transition (1955SC1B, 1956FR27). A shell model calculation in intermediate coupling indicates that configuration mixing is required to give a non-zero matrix element for the  $0^+ \rightarrow 0^+$  transition and suggests that a semi-collective model is indicated (1955SC1B, 1956SH1F, 1957TA1B: see also (1955LA1C)). According to (1956EL1C), satisfactory agreement is obtained with a 50% admixture of  $1s^3 1p^8 2s$  and  $1s^4 1p^7 2p$ . (1956RE1C) also finds reasonable agreement using the  $1s^{-1} 2s$  configuration and suggests that such a "core" excitation might appear quite generally in the light nuclei (see  $^{16}\text{O}$ ,  $^{14}\text{C}$  and (1954CH1A)). Calculations using an independent-particle approach to a collective description give a good account of the form factors for both the 4.4 and 7.7 MeV excitations. The form factor for the 9.6 MeV level is consistent with  $J = 1^-$  (1956FE1B: see also (1957PA1B)). See also (1958EL48; theor.).

Neutron production with  $E_e = 35$  to 150 MeV has been studied by (1956GE1B). Comparison of  $\sigma(\gamma, n)$  and  $\sigma(\text{e}, \text{e}'n)$  for  $E_e = 24$  to 145 MeV indicates that the transitions are largely E1 (1958BA60).

25. (a)  $^{12}\text{C}(n, n)^{12}\text{C}$   
 (b)  $^{12}\text{C}(n, n')^{12}\text{C}^*$   
 (c)  $^{12}\text{C}(n, n')^4\text{He}^4\text{He}^4\text{He}$   $Q_m = -7.281$

For  $E_n \gtrsim 14$  MeV, elastic scattering angular distributions show pronounced optical-model effects: see (1956BU95, 1956CU1A, 1958NA09) and  $^{13}\text{C}$ .

A gamma ray of energy  $4.42 \pm 0.03$  MeV is observed at  $E_n = 6.58$  MeV (1956DA23: see also (1954TH1A, 1955BA1N, 1955BE1H)). Production of 15.1 MeV  $\gamma$ -rays is observed at  $E_n = 90$  MeV (1957WA04, 1957WA1F). At  $E_n = 14$  MeV, inelastic neutron groups corresponding to  $^{12}\text{C}^*(4.4, 9.6)$  are reported by (1956WO1B: see also (1953WH1A, 1956BE1F, 1956CA1E, 1958AN32)). The angular distributions for neutrons corresponding to  $^{12}\text{C}^*(4.4)$  agree well with (p, p') distributions and with direct interaction theory (1958AN32:  $E_n = 14$  MeV).

Inelastic excitation leading to  $\alpha$ -particle stars has been studied by (1955FR35); levels of  $^{12}\text{C}$  at 7.7 and 9.6 MeV are involved. (1953JA1C) find that for  $E_n < 20$  MeV, most events proceed through a level at  $10 \pm 0.8$  MeV,  $\Gamma_{\text{obs}} = 1.6$  MeV, to  $^8\text{Be}(0)$  (See (1955FR35) and  $^{12}\text{B}(\beta^-)^{12}\text{C}$ ). See also  $^{13}\text{C}$ , (1953LI1C) and (1956LA1D; theor.).

26. (a)  $^{12}\text{C}(p, p')^{12}\text{C}^*$   
 (b)  $^{12}\text{C}(p, p')^4\text{He}^4\text{He}^4\text{He}$   $Q_m = -7.281$

Elastic scattering differential cross sections have been determined by (1953BU72, 1957GI14, 1957GR53:  $E_p = 9.4$  to  $9.6$  MeV), (1954FI1B:  $E_p = 10$  MeV), (1956SH1C:  $E_p = 12$  MeV), (1957PE14:  $E_p = 14$  to  $19.4$  MeV), (1956DA03:  $E_p = 17.0$  MeV, c.m.), (1955KI43, 1956KI54:  $E_p = 14.5, 20$  and  $31.5$  MeV), (1952BR52, 1953WR1A:  $E_p = 30.6$  MeV), (1957HI1C:  $E_p = 40$  MeV) and (1957GE08:  $E_p = 96$  MeV). Calculations based on the diffuse-surface optical model give a good account of the data at higher energies (1957GL58, 1957ME21, 1958GL11). At  $E_p = 9.5$  MeV, some evidence for compound elastic scattering is seen in the marked energy variation at back angles (1957GI14). See also  $^{13}\text{N}$ . Polarization of elastic and inelastic protons has been studied by (1958BR24) for  $E_p = 16$  to  $18$  MeV. Polarization of elastic protons for  $E_p = 5 - 7$  MeV is reported by (1958WA1D, 1958WA1E). See also (1957LE1G, 1958KO03, 1958SQ53; theor.). Polarization studies at high energies are discussed by (1956ER1A, 1956NI1B, 1957AL1B, 1957HE1A, 1957HI98, 1957TY1B, 1958NI1B, 1958NI26).

Excitation of the 4.4 MeV level is commonly observed for all energies above  $E_p \approx 5$  MeV: see (1955AJ61) and (1956RE39:  $E_p = 5$  to  $6$  MeV), (1957GI14, 1957HO1H:  $E_p = 9.5$  MeV), (1957CO53:  $E_p = 12$  MeV), (1957PE14:  $E_p = 14$  to  $19$  MeV), (1952BR52:  $E_p = 31.5$  MeV), (1958CH26:  $E_p = 40$  MeV), (1956ST65:  $E_p = 96$  MeV), (1957DI28:  $E_p = 95$  and  $135$  MeV) and (1957AL39, 1957MA1G, 1957TY36, 1958MA1B, 1958TY49:  $E_p = 182$  MeV).  $E_x = 4431 \pm 8$  keV (1957BU36). The 7.7 and 9.6 MeV levels are reported by (1957CO53, 1957HO1H, 1957MA1G, 1957PE14, 1958CH26). Levels at 12.6, 15.0, and  $\approx 20$  MeV are

also reported by (1957MA1G, 1957TY36); the last appears strongly in the work of (1956ST65) ( $E_x = 20.8$  MeV) and is there associated with the giant resonance seen in  $^{12}\text{C}(\gamma, n)$  and  $^{12}\text{C}(\gamma, p)$ .

For  $E_p \lesssim 30$  MeV, the angular distribution of the proton group corresponding to the 4.4 MeV level exhibits a minimum near  $90^\circ - 100^\circ$  (c.m.) and a definite fore-and-aft asymmetry. Neither the compound nucleus model nor the direct interaction theory appears to give a satisfactory account of these distributions (1957CO53, 1957GI14, 1957PE14). A direct interaction calculation, using distorted waves, does reproduce the general features of the distributions and also fits the observed ( $p'$ - $\gamma$ ) correlation of (1956SH1E) and (1958LE06). See also (1956BE1G, 1957BA1L, 1957BU52, 1957MA58, 1958BR83, 1958CH26, 1958MO98).

The angular distribution of protons corresponding to the 7.7 MeV state depends strongly on energy in the range  $E_p = 14$  to 19 MeV, but consistently shows a strong forward peak, indicative of  $l = 0$  and hence  $J = 0^+$  (1957PE14). See also (1955LA1C). An attempt to observe  $\gamma$ -decay of this level yields an upper limit of 3% for  $\Gamma_\gamma/\Gamma_\alpha$  (1956HO1D). The angular distribution of protons leading to the 9.6 MeV level is consistent with  $l = 1, J = 0^-, 1^-, 2^-$  (1957PE14).

The angular distribution of pick-up deuterons at  $E_p = 95$  MeV indicates significant contributions of high momentum components in the bound,  $1p$  neutron wave function, suggesting strong interactions, ( $\approx 200$  MeV), for close distances,  $R \approx 1 \times 10^{-13}$  cm (1955SE1C, 1956SE1A). See also (1956GR1E).

Emission of  $(15.1 \pm 0.2)$  MeV  $\gamma$ -radiation, ascribed to the first  $T = 1$  level, has been studied in the range  $E_p = 15$  to 340 MeV by (1957WA04, 1957WA1F). The general shape of the excitation function indicates direct nucleon-nucleon interaction for the higher energies; near threshold, emission of s-wave protons is indicated. Estimates of the  $\alpha$ -particle width, and comparison with isobaric spin forbidden reactions (e.g.  $^{12}\text{C}(\alpha, \alpha')^{12}\text{C}^*$ ,  $^{12}\text{C}(d, d')^{12}\text{C}^*$ ) indicate a  $T = 0$  admixture  $a^2 \approx 10^{-3}$ . At 31 MeV,  $\theta_{\text{lab}} = 80^\circ$ ,  $\gamma$ -rays of energy 15.1, 12.8 and 10.7 MeV are observed, with relative intensities  $1/0.090 \pm 0.015/0.095 \pm 0.014$ . The 12.8 MeV radiation is ascribed to excitation of a  $^{12}\text{C}$  level of that energy, while the 10.7 MeV radiation represents a cascade from  $^{12}\text{C}^*(15.1)$  to  $^{12}\text{C}^*(4.4)$  (1957WA1F: see also  $^{10}\text{B}(^3\text{He}, p)^{12}\text{C}$ ).

A study of reaction (c) at an energy of 29 MeV shows no indication of direct 4-body decay of  $^{13}\text{N}^*$ ;  $\frac{1}{4}$  of the events proceed via  $^8\text{Be}(0)$ , and  $> \frac{1}{2}$  via  $^8\text{Be}^*(2.9)$ . Evidence is found for the participation of  $^{12}\text{C}^*(9.6, \approx 12, 16, 20, 25)$  (1955NE18). See also (1955RE16), (1957JA1B) and (1955CU1C, 1956SA1C, 1956SA1D).

See also (1956ST30, 1957AL39, 1957GO1D) and (1957KA1D, 1958EL48; theor.).

## 27. $^{12}\text{C}(d, d')^{12}\text{C}^*$

The angular distribution of elastic scattering has been studied at  $E_d = 19$  MeV by (1954FR24); several diffraction peaks appear. See also (1958WA07).

Inelastic groups corresponding to  $^{12}\text{C}^*(4.4, 9.6)$  are reported by (1951KE02, 1954FR24, 1956GR37, 1956HA90: see also (1956CA65)). The 7.7 MeV level has not been observed. The angular distribution of the  $Q = -4.4$  MeV group at  $E_d = 15$  MeV has been analyzed in terms of direct nuclear

interaction theory and in terms of electric interaction theory by (1956HA90); neither appears to give a satisfactory account of the observations.

A search for 15 MeV  $\gamma$ -radiation at  $E_d = 85$  MeV yielded a negative result; a  $T = 0$  admixture of  $< 4 \times 10^{-2}$  is indicated (1957WA04, 1957WA1F).

See also (1955KH31, 1955KH35).

28.  $^{12}\text{C}(^3\text{He}, ^3\text{He}')^{12}\text{C}^*$

See (1958WE1E).

29. (a)  $^{12}\text{C}(\alpha, \alpha')^{12}\text{C}^*$

(b)  $^{12}\text{C}(\alpha, \alpha n)^{11}\text{C} \quad Q_m = -18.722$

Elastic scattering has been studied at  $E_\alpha = 19$  MeV by (1958PR65), at 31.5 MeV by (1956WA29), at 40 MeV by (1956IG02, 1956WE1C, 1957IG03) and at 48 MeV by (1955VA1A). The angular distributions show strong diffraction effects indicative of a direct interaction. Inelastic groups corresponding to levels at 4.4,  $7.64 \pm 0.07$ , 9.6 and possibly, 12.7 MeV are observed.  $^{12}\text{C}$  recoils corresponding to the ground and 4.4 MeV states are also reported; the absence of recoils corresponding to the 7.7 MeV state is taken to indicate that this state disintegrates primarily ( $> 80\%$ ) by  $\alpha$ -emission (1955RA1B: see  $^{11}\text{B}(d, n)^{12}\text{C}$ ). From a similar experiment, (1958EC12) find that the chance is less than 0.1 for  $\Gamma_\gamma/\Gamma > 10^{-3}$ .

Angular distribution of the  $Q = -4.4$  MeV inelastic group at  $E_\alpha = 31.5$  MeV are consistent with the direct surface interaction theory of (1953AU1A). A similar analysis of the  $Q = -7.7$  MeV group gives good agreement for  $J = 0^+$  (1956WA29). At  $E_\alpha = 42$  MeV, the angular distribution of this group is well matched by the  $j_0^2(kr)$  or  $j_2^2(kr)$  functions, indicating  $J^\pi = 0^+$  or  $2^+$ . The former is preferred in view of the small  $\gamma$ -width (1958EC12). See also (1956WE1C, 1957FI1C, 1958PR65, 1958SH65).

A search for  $\gamma$ -radiation from the de-excitation of the 15 MeV,  $T = 1$  level at  $E_\alpha = 48$  and 175 MeV gives an upper limit of  $\approx 10^{-3}$  for the  $T = 0$  admixture (1957WA04, 1957WA1F).

See also (1954JU1B). For reaction (b), see (1953LI1B).

30.  $^{12}\text{N}(\beta^+)^{12}\text{C} \quad Q_m = 17.46$

The decay is mainly to the ground state via an allowed transition. Transitions to  $^{12}\text{C}^*(4.4, 7.65)$  also are allowed. Branching ratios are 100/15/3;  $\log ft = 4.17, 4.4$  and  $4.4$ , respectively (1958VE20). Delayed  $\alpha$ -particles with a total energy of  $\approx 4$  MeV are also observed, suggesting that a state of  $^{12}\text{C}$  in the region 11 to 12 MeV is involved (1950AL57). See  $^{12}\text{N}$ .

31.  $^{13}\text{C}(\gamma, n)^{12}\text{C}$   $Q_m = -4.946$

See  $^{13}\text{C}$ .

32.  $^{13}\text{C}(\text{p}, \text{d})^{12}\text{C}$   $Q_m = -2.719$   
 $Q_0 = -2.720 \pm 0.007$  (1957BU36).

See (1955NE18) and (1957BE49).

33.  $^{13}\text{C}(\text{d}, \text{t})^{12}\text{C}$   $Q_m = 1.313$

Angular distributions of the ground state tritons have been measured at  $E_d = 2.2$  and  $3.3$  MeV (1954HO48).

34.  $^{13}\text{C}(^3\text{He}, \alpha)^{12}\text{C}$   $Q_m = 15.632$

Angular distributions of the  $\alpha$ -particle groups to the ground and  $4.4$  MeV states have been obtained at  $E(^3\text{He}) = 2$  MeV (1957HO63) and  $4.5$  MeV (1957HO62). Some direct interaction appears to be involved at both energies. At  $E(^3\text{He}) = 2$  MeV, a  $15.1$  MeV  $\gamma$ -ray is observed (1957BR18, 1957GO1B, 1958BR1D).

35.  $^{14}\text{C}(\text{p}, \text{t})^{12}\text{C}$   $Q_m = -4.635$

Not observed.

36.  $^{14}\text{N}(\gamma, \text{d})^{12}\text{C}$   $Q_m = -10.265$

Not observed.

37.  $^{14}\text{N}(\text{n}, \text{t})^{12}\text{C}$   $Q_m = -4.007$

See  $^{15}\text{N}$  and (1952LI1A).

Table 12.11:  $^{12}\text{C}$  states from  $^{14}\text{N}(\text{d}, \alpha)^{12}\text{C}$

A		B	C	D	E	F	G
Q (MeV $\pm$ keV)	$\sigma$ (mb)	Q (MeV $\pm$ keV)	Q (MeV $\pm$ keV)	Q (MeV $\pm$ keV)	Q (MeV $\pm$ keV)	Q (MeV $\pm$ keV)	$^{12}\text{C}^*$ (MeV)
13.39 $\pm$ 80	1	13.575 $\pm$ 12					0
9.02 $\pm$ 70	3	9.137 $\pm$ 6					4.442
5.77 $\pm$ 70	0.3		5.89 $\pm$ 30	5.910 $\pm$ 15	5.912 $\pm$ 13		7.664
		3.955 $\pm$ 3				3.933 $\pm$ 14	9.63

A:  $E_\alpha = 1.01$  MeV,  $\theta = 90^\circ$  (1940HO1A).

B: (1951MA08).

C: Based on  $Q_1 = 9.137$  for the second group (1953DU23). At  $E_\alpha = 0.62$  MeV,  $\theta = 90^\circ$ , this group has 6% the intensity of the first.

D: (1955PA50).

E: (1956AH32).

F: (1956DO41): it is suggested that a systematic error exists in (1951MA08)'s value for this state.

G: Based on  $Q_0 = 13.579$ .

38.  $^{14}\text{N}(\text{p}, ^3\text{He})^{12}\text{C}$

$$Q_m = -4.772$$

Not observed.

39.  $^{14}\text{N}(\text{d}, \alpha)^{12}\text{C}$

$$Q_m = 13.579$$

For  $\alpha$ -groups have been observed corresponding to  $^{12}\text{C}^*(0, 4.4, 7.7, 9.6)$ : see Table 12.11 and (1957HO1H). Alpha-gamma correlations give  $J = 2^+$  for the 4.4 MeV state (1954ST1C) while  $\gamma$ - $\gamma$  correlations give  $J = 0$  or  $> 2$  for the 7.7 MeV state (1955SE1B: see, however,  $^{12}\text{B}(\beta^-)^{12}\text{C}$ ). The width of the 7.7 MeV state is  $< 25$  keV (1953DU23, 1956AH32) and that of the 9.6 MeV state is  $30 \pm 8$  keV (c.m.). The  $J^\pi$  values for the 9.6 MeV state are limited to  $0^+, 1^-, 2^+, 3^-, 4^+$  (1956DO41). Angular distributions of the  $\alpha$ -particles to the ground, 4.4 and 9.6 MeV states have been measured at  $E_d = 20.9$  MeV (1957FI1C). See also (1952GI01, 1958BO18, 1958BO71). A small yield of 15 MeV  $\gamma$ -radiation is observed at  $E_d = 10.8$  MeV, presumably due to excitation of the 15.1 MeV,  $T = 1$  state (1954RA35). See also (1956GR37, 1958RA13) and  $^{16}\text{O}$ .

40.  $^{14}\text{N}(^3\text{He}, \text{p}\alpha)^{12}\text{C}$

$$Q_m = 8.086$$

See  $^{16}\text{O}$ .

41.  $^{15}\text{N}(\text{p}, \alpha)^{12}\text{C}$   $Q_{\text{m}} = 4.964$

Alpha particles have been observed to a state of  $^{12}\text{C}$  at  $4.432 \pm 0.010$  MeV (1952SC1B). The  $\gamma$ -ray energy after Doppler correction of 20 keV is  $4.443 \pm 0.020$  MeV. The necessity for the correction implies a lifetime  $< 3 \times 10^{-13}$  sec (1952TH24); see  $^{12}\text{C}(\gamma, \gamma')^{12}\text{C}^*$ . The angular distributions of short-range alpha particles and 4.4 MeV  $\gamma$ -radiation indicate that the 4.4 MeV state has  $J = 2^+$  or  $> 4$  (1953KR1B; see also (1957GO1E)). See also  $^{16}\text{O}$  and (1958RA14).

42. (a)  $^{16}\text{O}(\gamma, \alpha)^{12}\text{C}$   $Q_{\text{m}} = -7.148$   
 (b)  $^{16}\text{O}(\gamma, 4\alpha)$   $Q_{\text{m}} = -14.429$

There is evidence for the involvement of  $^{12}\text{C}$  states at 9.6 and  $\approx 11$  MeV which decay to the ground state of  $^8\text{Be}$ , a state at 12 – 13 MeV, decaying mainly to the 2.9 MeV state of  $^8\text{Be}$ , and  $T = 1$  state at  $\approx 16$  and 18 – 19 MeV, again leading mainly to the 2.9 MeV  $^8\text{Be}$  state. The 4.4 and 7.7 MeV states of  $^{12}\text{C}$  seem to occur rarely, if at all (see  $^{16}\text{O}$ ).

43.  $^{19}\text{F}(\text{p}, 2\alpha\gamma)^{12}\text{C}$   $Q_{\text{m}} = 0.971$

See (1957ZA1A).

$^{12}\text{N}$   
(Fig. 21)

GENERAL:

*Mass of  $^{12}\text{N}$ :* From the  $Q$  of the  $^{10}\text{B}(^3\text{He}, \text{n})^{12}\text{N}$  reaction,  $1.46 \pm 0.06$  MeV, and the Wapstra (1955WA1A) atomic masses for  $^{10}\text{B}$ ,  $^3\text{He}$  and  $\text{n}$ , the mass excess of  $^{12}\text{N}$  is  $21.00 \pm 0.06$  MeV.

1.  $^{12}\text{N}(\beta^+)^{12}\text{C}$   $Q_{\text{m}} = 17.46$

The half life is  $12.5 \pm 1$  msec;  $E_{\beta}(\text{max}) = 16.6 \pm 0.2$  MeV (1949AL05),  $\tau_{1/2} = 11.43 \pm 0.05$  msec;  $E_{\beta}(\text{max}) = 16.37 \pm 0.06$  MeV (1958VE20). The decay is complex;  $^{12}\text{N}$  decays to the ground state of  $^{12}\text{C}$  and to  $^{12}\text{C}^*(4.4, 7.6)$ : see  $^{12}\text{C}$ .  $\text{Log } ft = 4.17$  for the ground state transition (1958VE20). See also (1959SC1B; theor.).

Table 12.12: Energy levels of  $^{12}\text{N}$

$E_x$ (MeV)	$J^\pi$	$\tau_{1/2}$	Decay	Reactions
0	$1^+$	$(11.43 \pm 0.05) \times 10^{-3}$ sec	$\beta^+$	1, 2, 3, 5
$1.06 \pm 0.08$				2
$1.56 \pm 0.08$				2
$(1.97 \pm 0.10)$				(2)
$2.35 \pm 0.08$				2
$3.18 \pm 0.15$				2
$3.46 \pm 0.15$				2

2.  $^{10}\text{B}(^3\text{He}, \text{n})^{12}\text{N}$

$$Q_m = 1.46$$

$$Q_0 = 1.46 \pm 0.06 \text{ (1957AJ71)}.$$

The neutron spectrum has been studied at  $E(^3\text{He}) = 2.54$  and  $3.60$  MeV. Seven neutron groups are observed, corresponding to the ground state and to excited states at  $1.06 \pm 0.08$ ,  $1.56 \pm 0.08$ ,  $(1.97 \pm 0.10)$ ,  $2.35 \pm 0.08$ ,  $3.18 \pm 0.15$  and  $3.46 \pm 0.15$  MeV, in good agreement with the levels of the mirror nucleus,  $^{12}\text{B}$ . At the lower energy, the (c.m.) angular distribution of the ground state neutrons is isotropic within  $\approx 15\%$  statistics; at the higher energy, the distribution is peaked forward (1957AJ71).

3.  $^{12}\text{C}(\text{p}, \text{n})^{12}\text{N}$

$$Q_m = -18.24$$

$$E_{\text{thresh}} = 20.0 \pm 0.1 \text{ MeV (1949AL05)}. \text{ See also (1957CA1E, 1957GR1F) and } ^{13}\text{N}.$$

4.  $^{12}\text{C}(^3\text{He}, \text{t})^{12}\text{N}$

$$Q_m = -17.48$$

Not reported.

5.  $^{14}\text{N}(\gamma, 2\text{n})^{12}\text{N}$

$$Q_m = -30.73$$

See  $^{14}\text{N}$ .

6.  $^{14}\text{N}(\text{p}, \text{t})^{12}\text{N}$

$$Q_m = -22.25$$

Not reported.

## References

(Closed 1 December 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1940HO1A M.G. Holloway and B.L. Moore, *Phys. Rev.* 58 (1940) 847
- 1948JE03 J.V. Jelley and E.B. Paul, *Proc. Cambridge Phil. Soc.* 44 (1948) 133
- 1949AL05 L.W. Alvarez, *Phys. Rev.* 75 (1949) 1815
- 1950AL57 L.W. Alvarez, *Phys. Rev.* 80 (1950) 519
- 1950BU1A Buechner, van Patter, Strait and Sperduto, *Phys. Rev.* 70 (1950) 262
- 1950HO01 W.F. Hornyak and T. Lauritsen, *Phys. Rev.* 77 (1950) 160
- 1950HU72 E.L. Hudspeth, C.P. Swann and N.P. Heydenburg, *Phys. Rev.* 80 (1950) 643
- 1951BL1A Blaser, Boehm, Marmier and Scherrer, *Helv. Phys. Acta* 24 (1951) 465
- 1951BO45 C.K. Bockelman, D.W. Miller, R.K. Adair and H.H. Barschall, *Phys. Rev.* 84 (1951) 69
- 1951HA1C J. Halpern and A.K. Mann, *Phys. Rev.* 83 (1951) 370
- 1951KE02 K.K. Keller, J.B. Niedner, C.F. Wang and F.B. Shull, *Phys. Rev.* 81 (1951) 481
- 1951MA08 R. Malm and W.W. Buechner, *Phys. Rev.* 81 (1951) 519
- 1951MC57 W.O. McMinn, M.B. Sampson and V.K. Rasmussen, *Phys. Rev.* 84 (1951) 963
- 1951RU1A Rutherglen, Rae and Smith, *Proc. Phys. Soc. (London)* A64 (1951) 906
- 1951TE1A V.L. Telegdi, *Phys. Rev.* 84 (1951) 600
- 1952BR52 R. Britten, *Phys. Rev.* 88 (1952) 283
- 1952CO1B J.H. Coon, E.R. Graves and H.H. Barschall, *Phys. Rev.* 88 (1952) 562
- 1952EN19 P.M. Endt, C.H. Paris, H.M. Jongerius and F.P.G. Valckx, *Physica* 18 (1952) 423
- 1952GI01 W.M. Gibson and E.E. Thomas, *Proc. Roy. Soc.* A210 (1952) 543
- 1952GU1A W.H. Guier, H.W. Bertini and J.H. Roberts, *Phys. Rev.* 85 (1952) 426
- 1952HA1B J. Halpern, A.K. Mann and M. Rothman, *Phys. Rev.* 87 (1952) 164
- 1952JO10 V.R. Johnson, *Phys. Rev.* 86 (1952) 302
- 1952LI1A A.B. Lillie, *Phys. Rev.* 87 (1952) 716
- 1952SC1B Schardt, Fowler and Lauritsen, *Phys. Rev.* 86 (1952) 527

1952TH1B Thomson, Cohen, French and Hutchinson, Proc. Phys. Soc. (London) A65 (1952) 745  
1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969  
1953AU1A Austern, Butler and McManus, Phys. Rev. 92 (1953) 350  
1953BE1C L.E. Beghian, H.H. Halban, T. Husain and L.G. Sanders, Phys. Rev. 90 (1953) 1129  
1953BE61 O. Beckman, T. Huus and C. Zupancic, Phys. Rev. 91 (1953) 606  
1953BU72 W.E. Burcham, W.M. Gibson, A. Hossain and J. Rotblat, Phys. Rev. 92 (1953) 1266  
1953DA1A Dawson and Bigham, Can. J. Phys. 31 (1953) 167  
1953DU23 D.N. Dunbar, R.E. Pixley, W.A. Wenzel and W. Whaling, Phys. Rev. 92 (1953) 649  
1953EL12 M.M. Elkind, Phys. Rev. 92 (1953) 127  
1953GE1B Gell-Mamm and Telegdi, Phys. Rev. 91 (1953) 169  
1953GI05 W.M. Gibson, Phil. Mag. 44 (1953) 297  
1953GO13 F.K. Goward and J.J. Wilkins, Proc. Roy. Soc. A217 (1953) 357  
1953GU1A Gurewitsch, Diss. E.T.H. Zurich, Paper 2205 (1953); Phys. Abs. 3103 (1955)  
1953HE1B Hendel, Z. F. Phys. 135 (1953) 168  
1953HO48 J.R. Holt and T.N. Marsham, Prog. Phys. Soc. A66 (1953) 1032  
1953HO79 R. Hofstadter, H.R. Fechter and J.A. McIntyre, Phys. Rev. 92 (1953) 978  
1953HU29 T. Huus and R.B. Day, Phys. Rev. 91 (1953) 599  
1953JA1C Jackson and Wanklyn, Phys. Rev. 90 (1953) 381, W6  
1953KR1B A.A. Kraus, A.P. French, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 89 (1953) 299  
1953LI1B Lindner and Osborne, Phys. Rev. 91 (1953) 1501  
1953LI1C Livesey and Smith, Proc. Phys. Soc. (London) A66 (1953) 689  
1953MI31 C.H. Millar and A.G.W. Cameron, Can. J. Phys. 31 (1953) 723  
1953PA1B Paul and Clarke, Phys. Rev. 91 (1953) 463, KA14  
1953SH64 E.S. Shire, J.R. Wormald, G. Lindsay-Jones, A. Lundan and A.G. Stanley, Phil. Mag.  
44 (1953) 1197  
1953WH1A B.G. Whitmore, Phys. Rev. 92 (1953) 654  
1953WI1C Wilkinson, Phil. Mag. 44 (1953) 1019  
1953WR1A Wright, UCRL 2422 (1953)  
1954BU06 W.H. Burke, J.R. Risser and G.C. Phillips, Phys. Rev. 93 (1954) 188  
1954CH1A Christy and Fowler, Phys. Rev. 96 (1954) 851, F11  
1954CH1B Chastel, J. Phys. Rad. 15 (1954) 240  
1954CO16 C.F. Cook and T.W. Bonner, Phys. Rev. 94 (1954) 651

1954DI1A Diller and Crouch, Phys. Rev. 93 (1954) 362, 17  
1954EL1B J.O. Elliot, W I. McGarry and W.R. Faust, Phys. Rev. 93 (1954) 1348  
1954FI1B Fischer, UCRL-2546 (1954)  
1954FR24 R.G. Freemantle, W.M. Gibson and J. Rotblat, Phil. Mag. 45 (1954) 1200  
1954GI1B H.H. Givin, G.K. Farney, T.M. Hahn and B.D. Kern, Phys. Rev. 96 (1954) 1337  
1954GO39 J. Goldemberg and L. Katz, Phys. Rev. 95 (1954) 471  
1954GR1B L.H. Greenberg, J.G. Taylor and R.N. Haslam, Phys. Rev. 95 (1954) 1540  
1954GR1C Grant, Flack, Rutherglen and Deuchars, Proc. Phys. Soc. (London) A67 (1954) 751  
1954GR53 A. Graue, Phil. Mag. 45 (1954) 1205  
1954HA07 G. Harries, Proc. Phys. Soc. (London) A67 (1954) 153  
1954HE1B Heiberg, Phys. Rev. 96 (1954) 856, I12  
1954HO48 H.D. Holmgren, J.M. Blair, B.E. Simmons, T.F. Stratton and R.V. Stuart, Phys. Rev. 95 (1954) 1544  
1954JU1B Juric, Atomics and Atomic Tech. 5 (1954) 333  
1954KA1A L. Katz, R.N. Haslam, R.J. Horsley, A.G. Cameron and R. Montalbetti, Phys. Rev. 95 (1954) 464  
1954KH1A Khromchenko, Dokl. Akad. Nauk SSSR 94 (1954) 1037  
1954MI68 W.R. Mills and R.J. Mackin, Phys. Rev. 95 (1954) 1206  
1954NE1A Nereson et al, LA 1655 (1954)  
1954PA1D Pauli, Ark. Fys. 8 (1954) 16A  
1954PA28 C.H. Paris, F.P.G. Valckx and P.M. Endt, Physica 20 (1954) 573  
1954RA35 V.K. Rasmussen, J.R. Rees, M.B. Sampson and N.S. Wall, Phys. Rev. 96 (1954) 812  
1954ST1C Stanley, Phil. Mag. 45 (1954) 807  
1954TE1A Telegdi, Phys. in Canada 10 (1954) 44  
1954TH1A L.C. Thompson and J.R. Risser, Phys. Rev. 94 (1954) 941  
1954TH1B L.C. Thompson, Phys. Rev. 96 (1954) 369  
1954UE1A Uebergang, Aust. J. Phys. 7 (1954) 279  
1955AJ61 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 27 (1955) 77  
1955BA1N M.E. Battat and E.R. Graves, Phys. Rev. 97 (1955) 1266  
1955BA22 J.K. Bair, J.D. Kington and H.B. Willard, Phys. Rev. 100 (1955) 21  
1955BA63 W.C. Barber, W.D. George and D.D. Reagan, Phys. Rev. 98 (1955) 73  
1955BE1G R.D. Bent, T.W. Bonner, J.H. McCrary and W.A. Ranken, Phys. Rev. 100 (1955) 771

1955BE1H Beghian, Hicks and Milman, *Phil. Mag.* 464 (1955) 924  
 1955BE62 R.D. Bent, T.W. Bonner and R.F. Sippel, *Phys. Rev.* 98 (1955) 1237  
 1955BI26 C.B. Bigham, K.W. Allen and E. Almqvist, *Phys. Rev.* 99 (1955) 631A  
 1955BR1A R.J. Breen and M.R. Hertz, *Phys. Rev.* 98 (1955) 599  
 1955CA19 J.H. Carver, H.J. Hay and E.W. Titterton, *Phil. Mag.* 46 (1955) 841  
 1955CO1A D. Cooper and Dan J. Zaffarano, *Phys. Rev.* 98 (1955) 1199, X13  
 1955CU1C Cuer, Samman and Combe, *Compt. Rend.* 240 (1955) 1527  
 1955FE1A Ferrell and Visscher, *Phys. Rev.* 100 (1955) 1796, F8  
 1955FR1G J.H. Fregeau and R. Hofstadter, *Phys. Rev.* 99 (1955) 1503  
 1955FR35 G.M. Frye, Jr., L. Rosen and L. Stewart, *Phys. Rev.* 99 (1955) 1375  
 1955GE1A Geer, Nelson and Wolicki, *Phys. Rev.* 100 (1955) 215  
 1955GO10 H.E. Gove and E.B. Paul, *Phys. Rev.* 97 (1955) 104  
 1955GO59 F.K. Goward and J.J. Wilkins, *Proc. Roy. Soc.* 228 (1955) 376  
 1955HA1D Havlicek and Dobovisek, *Phys. Rev.* 100 (1955) 1355  
 1955HA1E Harding, *Proc. Phys. Soc. (London)* A68 (1955) 52  
 1955HE1E Herzenberg, *Nuovo Cim.* 1 (1955) 986, 1008  
 1955HI1C Hibdon and Langsdorf, *Phys. Rev.* 98 (1955) 223, B3  
 1955HO1D H.D. Holmgren, M.L. Bullock, and W.E. Kunz, *Phys. Rev.* 100 (1955) 436  
 1955HO48 R.E. Holland, D.R. Inglis, R.E. Malm and F.P. Mooring, *Phys. Rev.* 99 (1955) 92  
 1955HU1B Hughes and Harvey, BNL-325 (1955)  
 1955IH1B Ihsan, *Proc. Phys. Soc. (London)* A68 (1955) 393  
 1955JA1C Jancovici, *Nuovo Cim.* 1 (1955) 840  
 1955JO1B Johansson, *Phys. Rev.* 97 (1955) 434  
 1955JO1C Jones, *Proc. Glasgow Conf. (Pergamon Press, 1955)*  
 1955KH31 L.M. Khromchenko and V.A. Blinov, *Zh. Eksp. Teor. Fiz.* 28 (1955) 741; *JETP (Sov. Phys.)* 1 (1955) 596  
 1955KH35 L.M. Khromchenko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 19 (1955) 277; *Columbia Tech. Transl.* 19 (1956) 252  
 1955KI43 B.B. Kinsey, *Phys. Rev.* 99 (1955) 332  
 1955LA1C Lane and Wilkinson, *Phys. Rev.* 97 (1955) 1199  
 1955MA1H A.K. Mann, W.E. Stephens and D.H. Wilkinson, *Phys. Rev.* 97 (1955) 1184  
 1955MA1J L. Madansky and G.E. Owen, *Phys. Rev.* 99 (1955) 1608

1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847  
1955MO1B H. Morinaga, Phys. Rev. 97 (1955) 1185  
1955NE18 J.L. Need, Phys. Rev. 99 (1955) 1356  
1955PA50 R.T. Pauli, Ark. Fys. 9 (1955) 571  
1955RA1B Rasmussen, Miller and Sampson, Phys. Rev. 100 (1955) 181  
1955RA1D Ravenhall, Phys. Rev. 100 (1955) 1797, H6  
1955RA41 V.K. Rasmussen, D.W. Miller, M.B. Sampson and U.C. Gupta, Phys. Rev. 100 (1955) 851  
1955RE16 J.B. Reynolds, Phys. Rev. 98 (1955) 1289  
1955SA1B Sample, Neilson, Chadwick and Warren, Can. J. Phys. 33 (1955) 828  
1955SA1F Santos, Goldemberg, Pieroni, Silva, Borelli, Villaca and Leite Lopes, Anais Acad. Brasil. Ciencias 27 (1955) 437  
1955SC1B L.I. Schiff, Phys. Rev. 98 (1955) 1281  
1955SE1B Seed, Phil. Mag. 46 (1955) 100  
1955SE1C W. Selove, Phys. Rev. 98 (1955) 208  
1955SO1B S.D. Softky, Phys. Rev. 98 (1955) 173  
1955SP1A B.M. Spicer and A.S. Penfold, Phys. Rev. 100 (1955) 1375  
1955ST1C L. Stewart, Phys. Rev. 98 (1955) 740  
1955TA28 N.W. Tanner, Proc. Phys. Soc. (London) A68 (1955) 1195  
1955TI1A Titterton, Prog. Nucl. Phys. 4 (1955) 1  
1955VA1A Vaughn, UCRL-3174 (1955)  
1955WA1A Wapstra, Physica 21 (1955) 367  
1955WA30 A. Ward and P.J. Grant, Proc. Phys. Soc. (London) A68 (1955) 637  
1955WI25 H.B. Willard, J.K. Bair and J.D. Kington, Phys. Rev. 98 (1955) 669  
1956AH32 K. Ahnlund, Ark. Fys. 10 (1956) 369  
1956AL1E Almqvist, Bromley, Gove, Litherland and Paul, Bull. Amer. Phys. Soc. 1 (1956) 195  
1956AR21 A.H. Armstrong and G.M. Frye, Jr., Phys. Rev. 103 (1956) 335  
1956BE1D Becker and Barschall, Phys. Rev. 102 (1956) 1384  
1956BE1F Benveniste, Gardner, Nakada and Wong, Bull. Amer. Phys. Soc. 1 (1956) 175  
1956BE1G Benoist, Marty and Meyer, Physica 22 (1956) 1173A  
1956BO1F Bogdanov, Vlasov, Kalinin, Rybakov and Sidorov, Physica 22 (1956) 1150  
1956BO43 G.F. Bogdanov, N.A. Vlasov, S.P. Kalinin, B.V. Rybakov and V.A. Sidorov, Zh. Eksp. Teor. Fiz. 30 (1956) 981; JETP (Sov. Phys.) 3 (1956) 793

1956BU95 E.J. Burge, Y. Fujimoto and A. Hossain, *Phil. Mag.* 1 (1956) 19  
1956CA1E Casson, *Bull. Amer. Phys. Soc.* 1 (1956) 56  
1956CA1F Cameron, *Bull. Amer. Phys. Soc.* 1 (1956) 191  
1956CA65 J. Catala, E. Villar and A. Bonet, *An. Real. Soc. Espan. Fis. y Quim.* A52 (1956) 93  
1956CO59 L. Cohen, A.K. Mann, B.J. Patton, K. Reibel, W.E. Stephens and E.J. Winhold, *Phys. Rev.* 104 (1956) 108  
1956CR1C D.S. Craig, W.G. Gross and R.G. Jarvis, *Phys. Rev.* 103 (1956) 1414  
1956CU1A G. Culler, S. Fernbach and N. Sherman, *Phys. Rev.* 101 (1956) 1047  
1956DA03 I.E. Dayton and G. Schrank, *Phys. Rev.* 101 (1956) 1358  
1956DA23 R.B. Day, *Phys. Rev.* 102 (1956) 767  
1956DE22 S. Devons, G. Manning and J.H. Towle, *Proc. Phys. Soc. (London)* A69 (1956) 173  
1956DO41 R.A. Douglas, J.W. Broer, R. Chiba, D.F. Herring and E.A. Silverstein, *Phys. Rev.* 104 (1956) 1059  
1956EL1C Elliott, *Phys. Rev.* 101 (1956) 1212  
1956ER1A T. Eriksson, *Nucl. Phys.* 2 (1956) 91  
1956FA1B Fabricand, Allison and Halpern, *Phys. Rev.* 103 (1956) 1755  
1956FE1B R.A. Ferrell and W.M. Visscher, *Phys. Rev.* 104 (1956) 475  
1956FL1B Flerov and Talyzin, *Sov. J. Nucl. Energy* 4 (1956) 617  
1956FR27 J.H. Fregeau, *Phys. Rev.* 104 (1956) 225  
1956GE1B George, *Bull. Amer. Phys. Soc.* 1 (1956) 387  
1956GL1B Glassgold and Galonsky, *Phys. Rev.* 103 (1956) 701  
1956GO1G Goldanskii, *Zh. Eksp. Teor. Fiz.* 30 (1956) 969; *JETP (Sov. Phys.)* 3 (1956) 791  
1956GO1K Gorodetzky, Armbruster, Chevallier and Gallmann, *J. Phys. Rad.* 17 (1956) 548  
1956GO1N Gorodetzky, Armbruster and Chevallier, *Physica* 22 (1956) 1159A  
1956GO73 S. Gorodetzky, R. Armbruster, P. Chevallier and A. Gallman, *Compt. Rend.* 242 (1956) 898  
1956GR1E Greider, *Bull. Amer. Phys. Soc.* 1 (1956) 376  
1956GR37 T.S. Green and R. Middleton, *Proc. Phys. Soc. (London)* A69 (1956) 28  
1956HA1C Hayakawa, Hayashi, Imoto and Kikuchi, *Prog. Theor. Phys.* 16 (1956) 507  
1956HA1D Hayashi and Nishida, *Prog. Theor. Phys.* 16 (1956) 613  
1956HA90 J.W. Haffner, *Phys. Rev.* 103 (1956) 1398  
1956HE83 R.H. Helm, *Phys. Rev.* 104 (1956) 1466  
1956HO1C Holmgren, Bullock and Kunz, *Phys. Rev.* 104 (1956) 1446

1956HO1D Hornyak, Bull. Amer. Phys. Soc. 1 (1956) 197  
1956HO93 R. Hofstadter, Revs. Mod. Phys. 28 (1956) 214  
1956IG02 G. Igo, H.E. Wegner and R.M. Eisberg, Phys. Rev. 101 (1956) 1508  
1956KA1A Kavanagh, Ph.D. Thesis, CalTech (1956)  
1956KI54 B.B. Kinsey and T. Stone, Phys. Rev. 103 (1956) 975  
1956KL19 G.K. Kligler, V.I. Riabinkin, I.V. Chuvilo and V.S. Shevchenko, Physica 22 (1956) 1142A  
1956KO1D Koch and Turkevich, Bull. Amer. Phys. Soc. 1 (1956) 94  
1956KO1E Kondaiah, Proc. Indian Acad. Sci. A43 (1956) 130  
1956KU1A Kurath, Phys. Rev. 101 (1956) 216  
1956LA1C Langsdorf, Lane and Monahan, ANL 5567 (1956)  
1956LA1D J.R. Lamarsh and H. Feshbach, Phys. Rev. 104 (1956) 1633  
1956LE1E Leiss and Wyckoff, Bull. Amer. Phys. Soc. 1 (1956) 197  
1956LI05 D.L. Livesey, Can. J. Phys. 34 (1956) 216  
1956MA1T Mann and Titterton, Proc. Phys. Soc. (London) A69 (1956) 917  
1956MA69 J.B. Marion and G. Weber, Phys. Rev. 103 (1956) 1408  
1956MA83 E.E. Maslin, J.M. Calvert and A.A. Jaffe, Proc. Phys. Soc. (London) A69 (1956) 754  
1956ME1B Medveczky, Acta Phys. Hungar. 6 (1956) 261; Phys. Abs. 4703 (1957)  
1956MO1D H. Morinaga, Phys. Rev. 101 (1956) 254  
1956MO1E Morpurgo, Nuovo Cim. 3 (1956) 430  
1956NA1A Nakagawa, Ohmura, Takebe and Obi, Prog. Theor. Phys. 16 (1956) 389  
1956NI1B Nishimura and Ruderman, Bull. Amer. Phys. Soc. 1 (1956) 383  
1956NO1A Norbeck, Bull. Amer. Phys. Soc. 1 (1956) 329  
1956PE1A Perring and Skyrme, Proc. Phys. Soc. (London) A69 (1956) 600  
1956PI1A G.F. Pieper and G.S. Stanford, Phys. Rev. 101 (1956) 672  
1956PR1A Prosser, Kraus and Schiffer, Bull. Amer. Phys. Soc. 1 (1956) 94  
1956PR1B Price, Class and Risser, Bull. Amer. Phys. Soc. 1 (1956) 94  
1956RE1C P.J. Redmond, Phys. Rev. 101 (1956) 751  
1956RE39 C.W. Reich, G.C. Phillips and J.L. Russell, Jr., Phys. Rev. 104 (1956) 143  
1956SA1C Samman, Compt. Rend. 242 (1956) 2232  
1956SA1D Samman, Compt. Rend. 242 (1956) 3062  
1956SE1A Selove, Phys. Rev. 101 (1956) 231

1956SH1C Shaw, Conzett, Slobodrian and Summers-Gill, Bull. Amer. Phys. Soc. 1 (1956) 253  
 1956SH1E Sherr and Hornyak, Bull. Amer. Phys. Soc. 1 (1956) 197  
 1956SH1F B.F. Sherman and D.G. Ravenhall, Phys. Rev. 103 (1956) 949  
 1956ST1E Steffen, Hinrichs and Neuert, Z. F. Phys. 145 (1956) 156  
 1956ST1F Steffen and Neuert, Z. F. Phys. 147 (1956) 125  
 1956ST30 K. Strauch and F. Titus, Phys. Rev. 104 (1956) 191  
 1956ST65 K. Strauch and F. Titus, Phys. Rev. 103 (1956) 200  
 1956TA07 N.W. Tanner, Phil. Mag. 1 (1956) 47  
 1956TA16 G.W. Tautfest and S. Rubin, Phys. Rev. 103 (1956) 196  
 1956TA1C Tassie, Aust. J. Phys. 9 (1956) 407  
 1956TZ1A Tzara, Compt. Rend. 242 (1956) 2340  
 1956VA17 F.P.G. Valckx, Thesis, Univ. of Utrecht (1956)  
 1956WA29 H.J. Watters, Phys. Rev. 103 (1956) 1763  
 1956WE1C Wegner and Igo, Bull. Amer. Phys. Soc. 1 (1956) 21  
 1956WI1F R. Wilson, Phys. Rev. 104 (1956) 1424  
 1956WO1A Wolicki, Geer, Holmgren and Johnston, Bull. Amer. Phys. Soc. 1 (1956) 196  
 1956WO1B Wood and Singletary, Bull. Amer. Phys. Soc. 1 (1956) 246  
 1956WO1C Wolicki, Holmgren, Johnston and Geer, Bull. Amer. Phys. Soc. 1 (1956) 325  
 1957AJ71 F. Ajzenberg-Selove, M.L. Bullock and E. Almqvist, Phys. Rev. 108 (1957) 1284  
 1957AL1B Alphonse, Johansson and Tobell, Nucl. Phys. 3 (1957) 185  
 1957AL39 R. Alphonse, A. Johansson and G. Tibell, Nucl. Phys. 4 (1957) 672  
 1957AM48 O. Ames, G.E. Owen and C.D. Swartz, Phys. Rev. 106 (1957) 775  
 1957BA1H Barker, Phil. Mag. 2 (1957) 780  
 1957BA1K Barker and Mann, Phil. Mag. 2 (1957) 5  
 1957BA1L Banerjee and Levinson, Ann. Phys. (USA) 2 (1957) 499  
 1957BE49 E.F. Bennett, Bull. Amer. Phys. Soc. 2 (1957) 208, P10  
 1957BI1C Biel, Phil. Mag. A70 (1957) 866  
 1957BI78 J.R. Bird and R.H. Spear, Aust. J. Phys. 10 (1957) 268  
 1957BR18 D.A. Bromley, E. Almqvist, H.E. Gove, A.E. Litherland, E.B. Paul and A.J. Ferguson,  
 Phys. Rev. 105 (1957) 957  
 1957BR1J Breen, Hertz and Wright, Bull. Amer. Phys. Soc. 2 (1957) 357  
 1957BU36 W.W. Buechner and A. Sperduto, Phys. Rev. 106 (1957) 1008

1957BU52 S.T. Butler, Phys. Rev. 106 (1957) 272  
 1957CA1D Carver and Lokan, Aust. J. Phys. 10 (1957) 312  
 1957CA1E Carpenter and Wilson, Bull. Amer. Phys. Soc. 2 (1957) 382  
 1957CH24 I.V. Chuvilo and V.G. Shevchenko, Zh. Eksp. Teor. Fiz. 32 (1957) 1335; JETP (Sov. Phys.) 5 (1957) 1090  
 1957CH25 L.F. Chase, Jr. and G. Igo, Bull. Amer. Phys. Soc. 2 (1957) 381, J3  
 1957CO53 H.E. Conzett, Phys. Rev. 105 (1957) 1324  
 1957CO57 B.C. Cook, Phys. Rev. 106 (1957) 300  
 1957CO59 C.W. Cook, W.A. Fowler, C.C. Lauritsen and T. Lauritsen, Phys. Rev. 107 (1957) 508  
 1957DE11 G. Dearnaley, G.A. Dissanaiké, A.P. French and G.L. Jones, Phys. Rev. 108 (1957) 743  
 1957DI1B Dixon, Bielesch and Geiger, Can. J. Phys. 35 (1957) 699  
 1957DI28 J.M. Dickson and D.C. Salter, Nuovo Cim. 6 (1957) 235  
 1957DO1A Dodge and Barber, Bull. Amer. Phys. Soc. 2 (1957) 377  
 1957FE1C A.M. Feingold, Phys. Rev. 105 (1957) 944  
 1957FI1C Fischer and Fischer, Bull. Amer. Phys. Soc. 2 (1957) 182  
 1957GE08 G. Gerstein, J. Niederer and K. Strauch, Phys. Rev. 108 (1957) 427  
 1957GI14 W.M. Gibson, D.J. Prowse and J. Rotblat, Proc. Roy. Soc. A243 (1957) 237  
 1957GL58 A.E. Glassgold and P.J. Kellogg, Phys. Rev. 107 (1957) 1372  
 1957GO1B Gove, Litherland, Almqvist, Bromley and Ferguson, Bull. Amer. Phys. Soc. 2 (1957) 51  
 1957GO1C G. Goldring, Y. Wolfson and R. Wiener, Phys. Rev. 107 (1957) 1667  
 1957GO1D Gooding, Bull. Amer. Phys. Soc. 2 (1957) 350  
 1957GO1E S. Gorodetzky, A. Gallmann, M. Croissiaux and R. Armbruster, Nucl. Phys. 4 (1957) 112  
 1957GO1F Goldemberg, An. Acad. Bras. Cienc. 29 (1957) 23  
 1957GR1F Gross, Bull. Amer. Phys. Soc. 2 (1957) 14  
 1957GR53 G.W. Greenlees, I.G. Kuo and M. Petracic, Proc. Roy. Soc. A243 (1957) 206  
 1957HA13 E. Hayward and E.G. Fuller, Phys. Rev. 106 (1957) 991  
 1957HE1A Heiberg, Phys. Rev. 106 (1957) 1271  
 1957HE1B A. Herzenberg, Nucl. Phys. 3 (1957) 1  
 1957HI1C Hintz, Bull. Amer. Phys. Soc. 2 (1957) 14  
 1957HI98 P. Hillman, A. Johansson and H. Tyren, Nucl. Phys. 4 (1957) 648

1957HO1E Hofstadter, Ann. Rev. Nucl. Sci., Vol. 7 (1957)  
 1957HO1H Hossain and Prowse, Nuovo Cim. 5 (1957) 973  
 1957HO61 H.D. Holmgren, E.A. Wolicki and R.L. Johnston, Bull. Amer. Phys. Soc. 2 (1957) 181, E7  
 1957HO62 H.D. Holmgren, E.H. Geer, R.L. Johnston and E.A. Wolicki, Phys. Rev. 106 (1957) 102  
 1957HO63 H.D. Holmgren, Phys. Rev. 106 (1957) 100  
 1957IG03 G. Igo and R.M. Thaler, Phys. Rev. 106 (1957) 126  
 1957JA1B Jannelli and Mezzanares, Nuovo Cim. 5 (1957) 1047  
 1957JO1B Johnston, Holmgren and Wolicki, Bull. Amer. Phys. Soc. 2 (1957) 181  
 1957KA1C Kalinin, Ogloblin and Petrov, Sov. J. At. Energy 2 (1957) 193  
 1957KA1D Kajikawa and Watari, Prog. Theor. Phys. 18 (1957) 103  
 1957KR1A Kruse, Bent and Eklund, Bull. Amer. Phys. Soc. 2 (1957) 29  
 1957KU58 D. Kurath, Phys. Rev. 106 (1957) 975  
 1957LA14 A. Langsdorf, Jr., R.O. Lane and J.E. Monahan, Phys. Rev. 107 (1957) 1077  
 1957LE1G Leksin, Joint Inst. for Nucl. Research, Lab. of Nucl. Problems, USSR (1957)  
 1957LI1A Livesey, Can. J. Phys. 35 (1957) 987  
 1957MA1G H. Tyren and Th.A.J. Maris, Nucl. Phys. 3 (1957) 52  
 1957MA58 Th.A.J. Maris and H. Tyren, Nucl. Phys. 4 (1957) 662  
 1957ME21 M.A. Melkanoff, J.S. Nodvik, D.S. Saxon and R.D. Woods, Phys. Rev. 106 (1957) 793  
 1957MI1A Milone, Ricamo and Rubbino, Nuovo Cim. 5 (1957) 528  
 1957MU1C Muto, Takekoshi, Nakamura, Imamura and Tsuneoka, J. Phys. Soc. Jpn. 12 (1957) 109  
 1957NO14 E. Norbeck, Jr., Phys. Rev. 105 (1957) 204  
 1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754  
 1957OW03 G.E. Owen and L. Madansky, Phys. Rev. 105 (1957) 1766; Erratum Phys. Rev. 108 (1957) 1647  
 1957PA1A Pal and Mukherjee, Phys. Rev. 106 (1957) 811  
 1957PA1B M.K. Pal and M.A. Nagarajan, Phys. Rev. 108 (1957) 1577  
 1957PE14 R.W. Peelle, Phys. Rev. 105 (1957) 1311  
 1957RA1A Rask, AMes, Edwards, Madansky and Owen, Bull. Amer. Phys. Soc. 2 (1957) 351  
 1957RE1A Reiner, Physica 23 (1957) 338

1957RI38 J.R. Risser, J.E. Price and C.M. Class, Phys. Rev. 105 (1957) 1288  
 1957RO1E Roy, Goes and Pirson, Compt. Rend. 244 (1957) 3054  
 1957SA1B E.E. Salpeter, Phys. Rev. 107 (1957) 516  
 1957SI1B Simons, Soc. Sci. Fennica, Commentationes Phys.-Math. 20 (1957) No. 3  
 1957ST1E Steffen and Neuert, Z. F. Phys. 147 (1957) 125  
 1957TA1B Tassie, Nuovo Cim. 5 (1957) 1497  
 1957TY1B Tyren, Hillman and Johansson, Nucl. Phys. 3 (1957) 336  
 1957TY36 H. Tyren and Th.A.J. Maris, Nucl. Phys. 4 (1957) 637  
 1957WA04 C.N. Waddell, H.E. Adelson, B.J. Moyer and H.C. Shaw, Bull. Amer. Phys. Soc. 2 (1957) 181, E10  
 1957WA1F Waddell, Thesis, Univ. of California (1957); UCRL-3901 (1957)  
 1957ZA1A Zabel, WASH 192 (1957)  
 1957ZE1A Zeidman and Fowler, Bull. Amer. Phys. Soc. 2 (1957) 351  
 1958AM13 O. Ames and G.E. Owen, Phys. Rev. 109 (1958) 1639  
 1958AN32 J.D. Anderson, C.C. Gardner, J.W. McClure, M.P. Nakada and C. Wong, Phys. Rev. 111 (1958) 572  
 1958AR1B Armbuster, Ann. Phys. (France) 3 (1958) 88  
 1958AX1A Axel, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)  
 1958BA1K P.S. Baranov, V.I. Gol'danskii and V.S. Roganov, Phys. Rev. 109 (1958) 1801  
 1958BA1M W.C. Barber, W.R. Dodge and V.J. Van Huyse, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)  
 1958BA30 W.C. Barber, W.R. Dodge and V.J. Van Huyse, Bull. Amer. Phys. Soc. 3 (1958) 173, E7  
 1958BA60 W.C. Barber, Phys. Rev. 111 (1958) 1642  
 1958BO18 D.L. Booth, F.V. Price, D. Roaf and G.L. Salmon, Proc. Phys. Soc. (London) A71 (1958) 325  
 1958BO71 D. Bodansky, S.F. Eccles, G.W. Farwell, M. Rickey and P.C. Robinson, Bull. Amer. Phys. Soc. 3 (1958) 327, K9  
 1958BR1D Bromley, Proc. Rehovoth Conf. (North-Holland Pub. Co., Amsterdam, 1958)  
 1958BR24 K.W. Brockman, Jr., Phys. Rev. 110 (1958) 163  
 1958BR83 T.H. Braid and J.L. Yntema, Bull. Amer. Phys. Soc. 3 (1958) 188, K11  
 1958CA1G Cameron, Bull. Amer. Phys. Soc. 3 (1958) 269  
 1958CH26 S.W. Chen, Bull. Amer. Phys. Soc. 3 (1958) 189, K14

- 1958CH31 I.V. Chuvilo and V.G. Shevchenko, Zh. Eksp. Teor. Fiz. 34 (1958) 593; JETP (Sov. Phys.) 7 (1958) 410
- 1958CO66 C.W. Cook, W.A. Fowler, C.C. Lauritsen and T. Lauritsen, Phys. Rev. 111 (1957) 567
- 1958DA11 W.K. Dawson, G.C. Neilson and J.T. Sample, Bull. Amer. Phys. Soc. 3 (1958) 323, H12
- 1958EC12 S.F. Eccles and D. Bodansky, Bull. Amer. Phys. Soc. 3 (1958) 188, K12
- 1958ED1C Edwards, Owen and Madansky, Bull. Amer. Phys. Soc. 3 (1958) 268
- 1958EH1A Ehrenberg and Hofstadter, HEPL 129 (1958)
- 1958EH1B Ehrenberg, Hofstadter, Meyer-Berkhout, Ravenhall and Sobottka, HEPL 149 (1958)
- 1958EL48 L.R.B. Elton, Nucl. Phys. 5 (1958) 173; Erratum Nucl. Phys. 8 (1958) 396
- 1958FR1C French, Univ. of Pittsburgh Tech. Rept. 9 (1958)
- 1958GE1C M. Gell-Mann, Phys. Rev. 111 (1958) 362
- 1958GL11 A.E. Glassgold and P.J. Kellogg, Phys. Rev. 109 (1958) 1291
- 1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
- 1958KA14 R.W. Kavanagh, Bull. Amer. Phys. Soc. 3 (1958) 316, B7
- 1958KA1D Katz, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)
- 1958KA31 R.W. Kavanagh and C.A. Barnes, Phys. Rev. 112 (1958) 503
- 1958KO03 E. Kondaiah, K.V.K. Iyengar and C. Badrinathan, Nucl. Phys. 5 (1958) 346
- 1958KR65 W.E. Kreger and B.D. Kern, Bull. Amer. Phys. Soc. 3 (1958) 188, K13
- 1958KR70 T.H. Kruse and R.D. Bent, Phys. Rev. 112 (1958) 931
- 1958LE06 C.A. Levinson and M.K. Banerjee, Ann. Phys. (USA) 3 (1958) 67
- 1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1
- 1958MA1F Macklin and Gibbons, Bull. Amer. Phys. Soc. 3 (1958) 26
- 1958MC1D McElhinney, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)
- 1958MO17 G. Morpurgo, Phys. Rev. 110 (1958) 721
- 1958MO98 C.B.O. Mohr, Proc. Phys. Soc. (London) A71 (1958) 717
- 1958MO99 C.D. Moak, A. Galonsky, R.L. Traughber and C.M. Jones, Phys. Rev. 110 (1958) 1369
- 1958NA09 M.P. Nakada, J.D. Anderson, C.C. Gardner and C. Wong, Phys. Rev. 110 (1958) 1439
- 1958NE38 G.C. Neilson, W.K. Dawson and J.T. Sample, Bull. Amer. Phys. Soc. 3 (1958) 323, H11
- 1958NI1B K. Nishimura, Phys. Rev. 110 (1958) 1166

1958NI26 K. Nishimura, Nucl. Phys. 7 (1958) 425  
1958PA1B Palfrey and Tautfest, Bull. Amer. Phys. Soc. 3 (1958) 173  
1958PE1A Penner and Leiss, Bull. Amer. Phys. Soc. 3 (1958) 56  
1958PE1B Penner and Leiss, See Koch in Conf. on Photonucle. Reactions, National Bureau of Standards (1958)  
1958PR65 J.R. Priest, J.C. Corradi, E. Bleuler and D.J. Tendam, Bull. Amer. Phys. Soc. 3 (1958) 199, P8  
1958RA13 W.A. Ranken, T.W. Bonner, J.M. McCrary and T.A. Rabson, Phys. Rev. 109 (1958) 917  
1958RA14 V.K. Rasmussen, F.R. Metzger and C.P. Swann, Phys. Rev. 110 (1958) 154  
1958RA43 D.G. Ravenhall, Revs. Mod. Phys. 30 (1958) 430  
1958SH65 G.B. Shook, Bull. Amer. Phys. Soc. 3 (1958) 330, N4  
1958SM1A Smith and Baker, Bull. Amer. Phys. Soc. 3 (1958) 172  
1958SQ53 E.J. Squires, Nucl. Phys. 6 (1958) 504  
1958SW63 D.R. Sweetman, Bull. Amer. Phys. Soc. 3 (1958) 186, K1  
1958TA03 Y.-K. Tai, G.P. Millburn, S.N. Kaplan and B.J. Moyer, Phys. Rev. 109 (1958) 2086  
1958TA05 N.W. Tanner, Proc. Phys. Soc. (London) A72 (1958) 457  
1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10  
1958VE20 J.F. Vedder, Thesis, Univ. of California (1958); UCRL-8324 (1958)  
1958WA07 S. Watanabe, Nucl. Phys. 8 (1958) 484  
1958WA1D Warner and Alford, Bull. Amer. Phys. Soc. 3 (1958) 204  
1958WA1E Warner and Alford, Bull. Amer. Phys. Soc. 3 (1958) 364  
1958WE1E Wegner and Hall, Bull. Amer. Phys. Soc. 3 (1958) 338  
1958WH1A Whitehead, Murray, Aitken, Middlemas and Collie, Phys. Rev. 110 (1958) 941  
1958WO1B Wolff, Univ. of Penn., Tech. Rept. 4 (1958); Phys. Rev. Lett. 1 (1958) 354  
1958WY67 M.E. Wyman, E.M. Fryer and M.M. Thorpe, Phys. Rev. 112 (1958) 1264  
1959GA09 E.L. Garwin, Phys. Rev. 114 (1959) 143  
1959GE33 D.S. Gemmell, A.H. Morton and E.W. Titterton, Nucl. Phys. 10 (1959) 33  
1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571  
1959KR1B W.E. Kreger and B.D. Kern, Phys. Rev. 113 (1959) 890  
1959NE1A Neilson, Dawson and Johnson, Rev. Sci. Instrum. 30 (1959) 963  
1959SC1B Schwarzschild, Bull. Amer. Phys. Soc. 4 (1959) 79  
AR57 Unknown Source

GA57B    Unknown Source