

Energy Levels of Light Nuclei $A = 13$

F. Ajzenberg-Selove^a and T. Lauritsen^b

^a *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

^b *California Institute of Technology, Pasadena, California*

Abstract: An evaluation of $A = 5-24$ was published in *Nuclear Physics* 11 (1959), p. 1. This version of $A = 13$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 1, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 13$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{13}\text{B}\$](#) , [\$^{13}\text{C}\$](#) , [\$^{13}\text{N}\$](#)

B. Tables of Recommended Level Energies:

[Table 13.1](#): Energy levels of ^{13}B

[Table 13.2](#): Energy levels of ^{13}C

[Table 13.9](#): Energy levels of ^{13}N

C. [References](#)

D. Figures: [\$^{13}\text{C}\$](#) , [\$^{13}\text{N}\$](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

Table 13.1: Energy levels of ^{13}B

E_x (MeV)	$J^\pi; T$	$\tau_{1/2}$ (sec)	Decay	Reactions
0	$(\frac{3}{2}^-); \frac{3}{2}$	$(35 \pm 15) \times 10^{-3}$	β^-	1, 2

^{13}B
(Not illustrated)

GENERAL:

Mass of ^{13}B : The mass excess of ^{13}B is 20.40 ± 0.05 MeV from the Q of the reaction $^7\text{Li}(^7\text{Li}, \text{p})^{13}\text{B}$ (1957NO14), and Wapstra's masses (1955WA1A) for ^7Li and ^1H . ^{13}B is then stable by 4.88 MeV to decay into $^{12}\text{B} + \text{n}$, by 11.00 MeV to decay into $^{10}\text{Be} + \text{t}$ and by 11.3 MeV to decay into $^9\text{Li} + \alpha$.

1. $^{13}\text{B}(\beta^-)^{13}\text{C}$ $Q_m = 13.44$

The half-life of ^{13}B is $(35 \pm 15) \times 10^{-3}$ sec (1956NO1A). Attempts to observe delayed neutrons from the decay of neutron-unstable states of ^{13}C have been unsuccessful (1953HU1C, 1956NO1A). It is pointed out by (1957NO14) that transitions to such states are unlikely if ^{13}B has the expected $J = \frac{3}{2}^-$. See also (1948SN1A, 1952SH44).

2. $^7\text{Li}(^7\text{Li}, \text{p})^{13}\text{B}$ $Q_m = 5.97$
 $Q_0 = 5.97 \pm 0.05$ (1957NO14).

This reaction has been observed for $E(^7\text{Li}) = 1.4$ to 2.0 MeV (1956AL1F, 1957NO14, 1958LI42). At $E(^7\text{Li}) = 2$ MeV, no proton groups have been observed corresponding to excited states of ^{13}B below $E_x = 2.9$ MeV (1958LI42). See also ^{14}C .

3. $^{11}\text{B}(\text{t}, \text{p})^{13}\text{B}$ $Q_m = 0.24$

Not reported.

¹³C
(Fig. 22)

GENERAL:

Theory: See (1955AU1A, 1955LA1A, 1956DA1G, 1956DE1C, 1956KU1A, 1957BA1H, 1958FR1C, 1958SK1A).

- | | | |
|--|----------------|----------------|
| 1. (a) ${}^6\text{Li}({}^7\text{Li}, \text{p}){}^{12}\text{B}$ | $Q_m = 8.338$ | $E_b = 25.876$ |
| (b) ${}^6\text{Li}({}^7\text{Li}, \text{n}){}^{12}\text{C}$ | $Q_m = 20.931$ | |
| (c) ${}^6\text{Li}({}^7\text{Li}, 2\text{n}){}^{11}\text{C}$ | $Q_m = 2.209$ | |

See (1957NO17).

- | | |
|--|----------------|
| 2. ${}^7\text{Li}({}^7\text{Li}, \text{n}){}^{13}\text{C}$ | $Q_m = 18.624$ |
|--|----------------|

See (1957NO17).

- | | |
|---|----------------|
| 3. ${}^9\text{Be}(\alpha, \gamma){}^{13}\text{C}$ | $Q_m = 10.654$ |
|---|----------------|

At $E_\alpha = 1.60$ MeV, the capture cross section is less than $30 \mu\text{b}$ (1955AL16).

- | | | |
|---|---------------|----------------|
| 4. ${}^9\text{Be}(\alpha, \text{n}){}^{12}\text{C}$ | $Q_m = 5.709$ | $E_b = 10.654$ |
|---|---------------|----------------|

Resonances for neutrons and for γ -rays from ${}^{12}\text{C}^*(4.4)$ are given in Table 13.3. Absolute cross sections for several resonances are reported by (1956BO61, 1959GI47). For the prominent 1.9 MeV resonance, $d\sigma/d\Omega$ (90°) for 4.4 MeV γ -rays is given as 12 mb/sr (lab) by (1956BO61) and as 26 mb/sr by (1955TA28). For $E_\alpha = 2.5$ to 8.2 MeV, absolute neutron yields have been measured by (1958MA1J, 1959GI47).

Separate excitation curves (at 0°) for ground state neutrons (n_0) and for neutrons to the 4.4 MeV state (n_1) are reported by (1957RI38) in the range $E_\alpha = 1.7$ to 4.8 MeV (n_0) and 3.1 to 4.8 MeV (n_1). The n_0 yield curves show broad maxima at $E_\alpha = 1.9, 2.0, 2.6, 4.2$ and 4.5 MeV. The sharp 3.98 MeV resonance is strong for n_1 , but quite weak for n_0 . Angular distributions of ground-state neutrons suggest two broad resonances in the region $E_\alpha = 3.9$ to 4.6 MeV, probably $J = \frac{3}{2}^+$ and $\frac{5}{2}^+$ (1957RI38).

Table 13.2: Energy levels of ^{13}C

E_x (MeV \pm keV)	J^π	τ_m or Γ (keV)	Decay	Reactions
0	$\frac{1}{2}^-$	—	stable	2, 9, 15, 17, 23, 29, 30, 32, 34, 35, 37, 40, 41
3.085 ± 5	$\frac{1}{2}^+$	$\tau_m < 3 \times 10^{-13}$ sec	γ	9, 15, 18, 23, 29, 32, 40
3.680 ± 7	$\frac{3}{2}^-$	$\tau_m < 3 \times 10^{-13}$ sec	γ	9, 15, 17, 23, 29, 32, 34, 40
3.850 ± 10	$\frac{5}{2}^+$	$\tau_m > 3 \times 10^{-13}$ sec	γ	9, 15, 23, 32, 40
5.51 ± 50				15
6.10 ± 50				15
6.86 ± 10	$\frac{5}{2}^+$	$\Gamma = 6$	n	15, 18, 23, 40
7.470 ± 20				23, 40
7.533 ± 20				15, 23, 40
7.641 ± 20	$\frac{3}{2}^+$	55 ± 15	n	18, 23, 40
8.33 ± 100	$\frac{3}{2}^+$	1000 ± 250	n	18, 23, 35
8.82 ± 40			n	15, 18, 40
9.50 ± 20			n	15, 18, 23, 40
9.90 ± 20	> 0		n	15, 18, 23, 40
10.76 ± 20			n	18, 19, 23
10.94 ± 100			n	15, 18, 19
11.02 ± 30	$(\frac{1}{2}^+)$	50	α, n	4, 15
11.08 ± 30		sharp	α, n	4, 15
11.97 ± 15	$(\frac{7}{2}^-)$	70	α, n	4, 15, 18, 19
12.21 ± 30		≈ 140	α, n	4, 15, 18, 19
12.44 ± 30	$(\frac{1}{2}^-)$	≈ 140	α, n	4, 19
12.81 ± 100				15
13.41 ± 30		50	α, n	4
13.77 ± 30		≈ 280	α, n	4
14.1 ± 100		≈ 210	α, n	4
14.64 ± 30			α, n	4
16.1 ± 100			α, n	4
20.52 ± 20		115 ± 10	d, n	10
21.28 ± 20		160 ± 15	d, n	10

Table 13.3: Resonances in ${}^9\text{Be}(\alpha, n){}^{12}\text{C}$

E_α ^a (MeV)	E_α ^b (MeV)	Γ (keV)	J^π	${}^{13}\text{C}^*$ (MeV)	Refs.
0.53	0.53	70	$(\frac{1}{2}^+)$ j	11.02	c
0.61	0.61	sharp		11.08	c
1.9	1.905	180	$(\frac{7}{2}^-)$ j	11.97 ^k	d
2.24		≈ 200		12.21 ^k	e
2.58	2.6	≈ 200	$(\frac{1}{2}^-)$ j	12.44 ^k	f
4.00	3.98	70		13.41	h,i
(4.2)		(≈ 300)		(13.6)	g
4.50	4.4	≈ 400		13.77 ^k	h,i
5.0	5.0	≈ 300		14.1 ^k	h
5.75				14.64	i
7.8				16.1	i

^a Resonances in neutron yield.

^b Resonances for 4.4 MeV γ -rays.

^c (1954BE08).

^d (1953TA06, 1954BE08, 1954TR09, 1955TA28, 1956BO61), $\Gamma_n/\Gamma_\alpha > 12$ (1955TA28).

^e (1954TR09, 1956BO61).

^f (1953TA06, 1954TR09, 1956BO61, 1958MA1J).

^g (1957RI38).

^h (1956BO61).

ⁱ (1958MA1J, 1959GI47).

^j (1956JA28).

^k Not corrected for effects of Coulomb barrier penetration.

Extensive angular distribution studies have been made for $E_\alpha < 2$ MeV by (1955TA28, 1956JA28). According to (1956JA28), the best fit to the distributions in the range $E_\alpha = 0.4$ to 1.3 MeV is obtained from the assignments $J = \frac{1}{2}^+, \frac{7}{2}^-, \frac{1}{2}^-$ for the 0.5, 1.9 and 2.6 (?) MeV resonances (see also (1955TA28)). The angular correlation of neutrons and 4.4 MeV gamma-rays is isotropic at $E_\alpha = 1.2$ and 2.8 MeV, indicating that stripping plays only a minor role at these energies (1958TA05). See also (1955MA1J; theor.) and (1956BE1D).

5. ${}^9\text{Be}(\alpha, p){}^{12}\text{B}$

$$Q_m = -6.884$$

$$E_b = 10.654$$

See ${}^{12}\text{B}$.

$$6. \text{}^9\text{Be}(\alpha, \text{d})^{11}\text{B} \qquad Q_{\text{m}} = -8.022 \qquad E_{\text{b}} = 10.654$$

See ^{11}B .

$$7. \text{(a) } ^9\text{Be}(\alpha, \alpha')^9\text{Be}^* \qquad E_{\text{b}} = 10.654$$

$$\text{(b) } ^9\text{Be}(\alpha, \alpha'\text{n})^8\text{Be} \qquad Q_{\text{m}} = -1.667$$

$$\text{(c) } ^9\text{Be}(\alpha, \text{n})^4\text{He}^4\text{He}^4\text{He} \qquad Q_{\text{m}} = -1.572$$

For reaction (a) see ^9Be and (1955TA28). For reactions (b) and (c), see (1952AJ38).

$$8. \text{}^{10}\text{B}(\text{t}, \alpha)^9\text{Be} \qquad Q_{\text{m}} = 13.210 \qquad E_{\text{b}} = 23.882$$

See ^9Be .

$$9. \text{}^{10}\text{B}(\alpha, \text{p})^{13}\text{C} \qquad Q_{\text{m}} = 4.070$$

$$Q_0 = 4.064 \pm 0.012 \text{ (W.J. Fader, quoted in (1957VA11))}.$$

$$Q_0 = 4.08 \pm 0.03 \text{ (1956PI1A)}.$$

$$Q_0 = 4.10 \pm 0.03 \text{ (1956PA1B)}.$$

Four proton groups are observed, corresponding to the ^{13}C levels at 0, 3.09, 3.68 and 3.85 MeV (1953SH64, 1954FA1A, 1954FA1B, 1956PI1A). Additional groups are reported by (1957RO1F). The relative intensities depend strongly on bombarding energy (see ^{14}N , (1953SH64)). See also (1955AJ61).

A study of gamma rays from this reaction and from $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$ shows three lines with $E_{\gamma} = 0.1695 \pm 0.0004$, 3.844 ± 0.015 and 3.69 ± 0.02 MeV. The 3.85 MeV γ -ray exhibits no Doppler shift and therefore has a lifetime greater than 3×10^{-13} sec; the 3.69 MeV line shows approximately the maximum possible Doppler shift ($\tau < 3 \times 10^{-13}$ sec). The 170 keV line is due to the cascade transition between the 3.84 and 3.68 MeV states; the internal conversion coefficient is consistent with E1, although M1 cannot be excluded. The probability of this cascade decay of the 3.84 MeV state is 0.24 ± 0.05 . Cascade transitions to the 3.1 MeV excited state have not been observed. Their intensities are less than 3% of the ground state transitions (1956MA1Q, 1956MA52): see Fig. 23. (For earlier work see (1953SH64) and (1954ST20)). The angular distributions and p- γ correlations for the 3.8 MeV radiation contain terms in $(\cos^4 \theta)$, indicating $J = \frac{5}{2}^+$ for the 3.84 MeV state (1954ST20: see $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$). If the 170 keV line is due to an E1 transition, the J^{π} of the 3.68 MeV state is then $\frac{3}{2}^-$ ($J^{\pi} = \frac{1}{2}^-$, $\frac{3}{2}^-$ follows from $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$); the angular distribution of the 3.7 MeV radiation is consistent with M1 (1954ST20).

Angular distributions of the ground state protons are reported at $E_\alpha = 4.9, 6.0, 7.0$ and 8.1 MeV (1957VO25) and at $E_\alpha = 30.5$ MeV (1957HU1E): in both cases direct interaction appears to be involved. See also (1957BA1K) and (1955BR1A).

$$10. \text{}^{11}\text{B}(\text{d}, \text{n})\text{}^{12}\text{C} \qquad Q_{\text{m}} = 13.731 \qquad E_{\text{b}} = 18.677$$

The yield of neutrons has been measured for $E_{\text{d}} = 0.2$ to 5.4 MeV. The total cross section for ground state neutrons in the 0.5 to 1.15 MeV range rises from 0.5 to 30 mb; both direct and exchange stripping processes seem to be involved (1956PA23, 1957AM48). The yield of the excited-state group, $^{12}\text{C}^*(4.4)$ rises smoothly from $E_{\text{d}} = 0.5$ to 1.1 MeV and is essentially flat from $E_{\text{d}} = 1.1$ to 2.0 MeV (1959NE1A). At $E_{\text{d}} = 600$ keV, angular distributions indicate that stripping is important for the ground-state group. For the excited-state group ($^{12}\text{C}^*(4.4)$), the interpretation is less clear; the observed distribution can be accounted for by p-wave formation of a $J = \frac{7}{2}^+$ level in ^{13}C (1955WA30). (1959NE1A) find evidence of strong “heavy-particle” stripping for this group in the range $E_{\text{d}} = 0.5$ to 2.0 MeV. The cross section for emission of neutrons in the forward direction is ≈ 270 mb/sr at $E_{\text{d}} = 5.4$ MeV (1955MA76). For $E_{\text{d}} = 1.6$ to 3.2 MeV, the yield of 15.1 MeV γ -rays shows resonances at $E_{\text{d}} = 2.180 \pm 0.010$ and 3.080 ± 0.015 MeV, corresponding to $^{13}\text{C}^*(20.52, 21.28)$ with $\Gamma_{\text{c.m.}} = 115 \pm 10$ and 160 ± 15 keV, respectively; the cross section at $E_{\text{d}} = 2.2$ MeV is 29 ± 7 mb (1958KA31). See also (1954BU06, 1955RI1B) and ^{12}C .

$$11. \text{}^{11}\text{B}(\text{d}, \text{p})\text{}^{12}\text{B} \qquad Q_{\text{m}} = 1.138 \qquad E_{\text{b}} = 18.677$$

The thin-target yield rises smoothly from $E_{\text{d}} = 0.3$ to 3.1 MeV with no evidence of resonances (1949HU41, 1958KA31). At $E_{\text{d}} = 1.5$ MeV, $\sigma \approx 0.38$ b (1958KA31: see, however, (1949HU41)). See also ^{12}B and (1957JA37).

$$12. \text{}^{11}\text{B}(\text{d}, \text{d})\text{}^{11}\text{B} \qquad E_{\text{b}} = 18.677$$

See ^{11}B .

$$13. \text{}^{11}\text{B}(\text{d}, \alpha)\text{}^9\text{Be} \qquad Q_{\text{m}} = 8.022 \qquad E_{\text{b}} = 18.677$$

Some absolute cross sections are given by (1958KA31). See also ^9Be .

$$14. \text{}^{11}\text{B}(\text{t}, \text{n})\text{}^{13}\text{C} \qquad Q_{\text{m}} = 12.419$$

Table 13.4: Levels of ^{13}C from $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$

E_x^a (MeV \pm keV)	E_x^b (MeV)
0	0
3.09	3.08
3.68	3.77
3.86	
5.51 \pm 50	
6.10 \pm 50	
6.87	6.89
7.55 \pm 40	7.63
8.87 \pm 50	8.96
9.52 \pm 60	
9.91 \pm 50	10.00
10.9 \pm 150 ^c	10.99
11.1 \pm 150 ^c	
	11.67
12.08 \pm 100 ^c	
12.81 \pm 100 ^c	

^a (1958MO99): $E(^3\text{He}) = 1.23$ MeV.

^b (1955BI26): $E(^3\text{He}) = 0.9$ MeV; values are ± 100 keV.

^c (1957GA01): $E(^3\text{He}) = 1.25$ MeV.

Not reported.

15. $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$

$$Q_m = 13.184$$

Levels derived from reported proton groups are listed in Table 13.4. The levels at 5.5 and 6.1 MeV have not been observed in any other reaction. From the fact that they do not appear in $^{12}\text{C}(n, n)^{12}\text{C}$, an upper limit of $\Gamma = 10$ keV is estimated; the mirror levels in ^{13}N must be assumed to have $\theta_p^2 < 0.02$ (1958MO99).

Angular distributions have been measured for the p_0 , p_1 and $(p_2 + p_3)$ groups at $E(^3\text{He}) = 4.5$ MeV. The p_0 group appears to be peaked in both the forward and the backward direction. The

other groups do not exhibit a strong angular variation (1957HO61). At $E(^3\text{He}) = 6.05$ MeV the p_0 group is strongly peaked forward ((1958SW63), and D.R. Sweetman, private communication).

$$16. \text{}^{11}\text{B}(\alpha, \text{d})^{13}\text{C} \quad Q_{\text{m}} = -5.167$$

Not reported.

$$17. \text{}^{12}\text{C}(\text{n}, \gamma)^{13}\text{C} \quad Q_{\text{m}} = 4.946$$

The thermal capture cross section is 3.3 ± 0.2 mb (1958HU18). In addition to the 4.95 MeV ground state transition ($E_{\gamma} = 4948 \pm 8$ keV), a γ -ray is reported with an energy of 3.68 ± 0.05 MeV and an intensity of 0.3 γ /capture. If 3.1 and 3.9 MeV γ -rays occur, their intensities are less than 0.10 and 0.06 γ /capture, respectively (1953BA18).

$$18. \text{}^{12}\text{C}(\text{n}, \text{n})^{12}\text{C} \quad E_{\text{b}} = 4.946$$

The cross section is approximately constant to 160 keV, then decreases monotonically to $E_{\text{n}} = 2$ MeV. There follows a region of resonances to 8.5 MeV, followed by a smooth variation of the cross section to $E_{\text{n}} = 100$ MeV (1958HU18). (1958CO07) finds a minimum of 1.3 b at $E_{\text{n}} = 14$ MeV, followed by a rise to 1.5 b at $E_{\text{n}} = 15.5$ MeV: see also $^{12}\text{C}(\text{n}, \text{p})^{12}\text{B}$. The average total cross section in the range $E_{\text{n}} = 14$ to 10000 eV is 4.69 ± 0.10 b (1956BR1E).

The parameters of observed resonances are displayed in Table 13.5. A careful search for resonances in the region $E_{\text{n}} = 20$ to 1360 keV with 10 and 22 keV resolution revealed no deviation $> 5\%$ from a smooth monotonic decrease in σ_{t} (1950MI1A) (see, however, $^{11}\text{B}(^3\text{He}, \text{p})^{13}\text{C}$). The course of the cross section in this region can be accounted for by the broad $s_{1/2}$ state at $^{13}\text{C}^*(3.09)$ (1952TH1D). A similar search, with 5 keV resolution, in the range $E_{\text{n}} = 2.73$ to 2.80 MeV revealed no deviations > 0.2 b from a smooth function (1958WI01: see $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$).

Below $E_{\text{n}} = 4.0$ MeV, three d-wave resonances occur, at 2.08, 2.95, and 3.67 MeV. The total cross section and angular distributions establish the first as $D_{\frac{5}{2}}$ (1958WI36); phase shift analyses of angular distributions yield $D_{\frac{3}{2}}$ for the other two. The s-wave phase shift is everywhere negative and decreases slowly with energy: the behavior for $E_{\text{n}} = 0$ to 4 MeV can be accurately reproduced by a static potential well with a diffuse boundary (1958WI36). See also (1954HU1A). The $p_{1/2}$ and $p_{3/2}$ phase shifts are negative and small, not inconsistent with hard-sphere scattering (1954ME95, 1955BU56, 1958WI36). Polarization of scattered neutrons is discussed in these three papers and in (1956MC70, 1957MC1B). Other angular distribution studies in this range are reported by (1957LA14: $E_{\text{n}} = 0.06$ to 1.8 MeV), (1955WI25: $E_{\text{n}} = 0.55$ to 1.5 MeV), (1956MU1A: $E_{\text{n}} = 1.66$ MeV), (1955LI1C: $E_{\text{n}} = 2.7$ MeV), (1955WA27: $E_{\text{n}} = 4.1$ MeV).

Table 13.5: Resonances in $^{12}\text{C}(n, n)^{12}\text{C}$

E_{res} (MeV)	Γ_{lab} (keV)	$^{13}\text{C}^*$ (MeV)	l_n	J^π	θ^2
2.076 ± 0.008 ^a	7 ^a	6.862	2 ^a	$\frac{5}{2}^+$ ^a	0.006 ^a
2.95 ^b	90 ^g	7.67	2 ^h	$\frac{3}{2}^+$ ^h	0.038 ^a
3.67 ^b	1690 ^g	8.33	2 ^h	$\frac{3}{2}^+$ ^h	0.51 ^a
4.4 ^c		9.0			
4.95 ^d		9.51			
5.40 ^d		9.93	> 0 ^d	$> \frac{1}{2}$ ^d	
6.3 ^e		10.8			
7.5 ^f		11.9			
7.83 ^f		12.17			

^a (1958WI36): see also (1951BO45).

^b See (1950RI1B, 1951BO45, 1951RI1A, 1951RI1B, 1958WI36).

^c (1950FR1A).

^d (1956BE1D).

^e See (1953NE1A, 1957BO13, 1958HU18).

^f (1956HA1E).

^g (1958WI36): (1954ME95) report 60 and 1200 keV.

^h See (1950RI1B, 1951BO45, 1951RI1A, 1951RI1B, 1954ME95, 1955BU56, 1958WI36).

For $E_n = 4$ to 8 MeV, several additional resonances are reported by (1950FR1A, 1956BE1D, 1956HA1E, 1957BO13): see Table 13.5 and (1958HU18). The structure above $E_n = 7$ MeV is undoubtedly quite complex: see $^9\text{Be}(\alpha, n)^{12}\text{C}$ and $^{12}\text{C}(n, n')^{12}\text{C}^*$. Further angular distribution measurements are reported by (1955JE1A: 4.4 MeV), (1958HI68: 5 MeV), (1958BR1F: 5.6 MeV), (1956BE32: 7 MeV), (1956DO1D, 1958NA09: 14 MeV), (1958CO1E: 14.5 MeV) and (1957KH1B: 14.8 MeV): see also (1956HU1A). Optical model effects become apparent at the higher energies; see (1956CU1A; theor.).

In the region beyond the resolved resonances, recent measurements of the total cross section have been made by (1958BR16: $E_n = 7$ to 14 MeV), (1958CO07: $E_n = 13.1$ to 15.6 MeV), (1958VE15, 1958VE21: $E_n = 13.6$ to 14.75 MeV) and (1957KH1A: $E_n = 14.8$ MeV). For a review of the earlier work, see (1955AJ61) and (1957HU1D). Non-elastic cross section measurements are reported by (1956BE32: $E_n = 7$ MeV), (1958BA03: $E_n = 7$ to 14 MeV), (1955TA1B: $E_n = 12.7$ and 14.1 MeV), (1955GR21, 1956FL1B, 1957ST1F: $E_n = 14$ MeV) and (1958MA54: $E_n = 21, 26$ and 29 MeV): see also (1947HU03, 1955MA1G, 1957ZA1A). See also (1956LA1C) and (1956KA1B; theor.).

19. (a) $^{12}\text{C}(n, n')^{12}\text{C}^*$ $E_b = 4.946$
 (b) $^{12}\text{C}(n, n')^4\text{He}^4\text{He}^4\text{He}$ $Q_m = -7.281$

In the range $E_n = 4.4$ to 8 MeV, four resonances are observed in the yield of 4.4 MeV γ -rays, at $E_n = 6.30, 6.49, 7.6, 7.87$ and 8.15 MeV, corresponding to $^{13}\text{C}^*(10.76, 10.94, 12.0, 12.21, 12.47)$. The differential cross section at 90° reaches a maximum of 60 mb/sr at 7.87 MeV ([1956HA1E](#), [1958HU18](#)). At $E_n = 6.58$ MeV, the cross section for production of 4.4 MeV γ -rays is 353 ± 59 mb ([1956DA23](#)); at 14 MeV, it is 245 ± 35 mb ([1955BA1N](#)). ([1955GR21](#)) estimate 160 mb for the inelastic cross section at $E_n = 14$ MeV to ^{12}C levels at 9.6 to ≈ 13 MeV and 100 to 300 mb as the cross section to the ^{12}C states at 4.4 and 7.6 MeV. See $^{12}\text{C}(n, n)^{12}\text{C}$ above, for further references on non-elastic cross sections.

Reaction (b) has been studied for $E_n = 12.3$ to 20.1 MeV. The cross section is 190 ± 50 mb at 12.9 MeV. It goes through a broad maximum of $\approx 300 \pm 60$ mb at ≈ 16.5 MeV and then decreases to 240 ± 50 mb at $E_n = 20.1$ MeV ([1955FR35](#)). See also ([1955BE1D](#), [1955BE1F](#), [1958VA1D](#)), ([1956SA1E](#); theor.) and ([1955AJ61](#)).

20. $^{12}\text{C}(n, 2n)^{11}\text{C}$ $Q_m = -18.722$ $E_b = 4.946$

See ([1952BR61](#), [1958AS63](#)).

21. $^{12}\text{C}(n, p)^{12}\text{B}$ $Q_m = -12.593$ $E_b = 4.946$

The cross section has been measured from threshold to $E_n = 17.5$ MeV. At $E_n = 17.5$ MeV, the cross section is 29.1 ± 4 mb ([1958KR65](#), [1959KR1B](#)). See also ([1956KR1A](#), [1956KR1B](#)).

22. $^{12}\text{C}(n, \alpha)^9\text{Be}$ $Q_m = -5.709$ $E_b = 4.946$

See ([1955GR21](#)) and ^9Be .

23. $^{12}\text{C}(d, p)^{13}\text{C}$ $Q_m = 2.719$
 $Q_0 = 2.721 \pm 0.002$ ([1957VA11](#)).

Measurements on the proton groups are summarized in Table [13.6](#). The level assignments were obtained by analysis of angular distributions (with deuterons of energies up to 24 MeV) in terms of direct interaction theories. A careful search with $E_d = 5$ to 8.5 MeV ($\theta = 90^\circ$) reveals no further

Table 13.6: Levels of ^{13}C from $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$

$^{13}\text{C}^*$ (MeV \pm keV)					l_n	J^π	$\sigma(\theta)^e$ (mb/sr)	$\theta_n^2{}^h$ (%)
(1951ST19, 1951VA1A)	(1954SP01)	(1956DO41)	(1953KH1A, 1955KH35) ^a	(1955MC75)				
0	0		0	0	1 ^c	$\frac{1}{2}^-, \frac{3}{2}^-$ ^c	26	7
3.086 \pm 6	3.090 \pm 10		3.107	3.09 ^b	0 ^c	$\frac{1}{2}^+$ ^c	103	25
3.686 \pm 11	3.684 \pm 10	[3.681 \pm 3]	3.699	3.68 ^b	1 ^c	$\frac{1}{2}^-, \frac{3}{2}^-$ ^c	16	2
	3.855 \pm 7	[3.851 \pm 3]	3.869	3.84 ^b	2 ^c	$\frac{3}{2}^+, \frac{5}{2}^+$ ^c	152	10
				6.87 ^b	(0), (2) ^d	$(\leq \frac{5}{2}^+)$ ^d	36	
				7.470 \pm 20			≈ 0.8	
				7.533 \pm 20			9.6	
				7.641 \pm 20 ^f			7.5	
				8.4 \pm 300 ^g			100	
				9.500 \pm 20			1.6	
				9.897 \pm 20			2.2	
				10.759 \pm 20			4.5	

^a ± 10 to 50 keV.

^b Energies given for identification only.

^c (1951RO1B, 1951RO1C, 1952BL1B, 1953CA1B, 1954FR24, 1955MC75, 1956CA1D, 1956GR37, 1957CO68).

^d (1955MC75).

^e (1955MC75); differential cross sections at the first maximum or in the forward direction; $\pm 25\%$.

^f $\Gamma = 70 \pm 15$ keV.

^g $\Gamma = 1.1 \pm 0.3$ MeV.

^h (1956EL1A, 1956GR37, 1958MC63).

proton proton groups corresponding to levels in the range 0 to 4.9 MeV with intensity greater than 0.5% of the ground state group (1954SP01). At $E_d = 14.8$ MeV, all groups show pronounced stripping distributions except that corresponding to $^{13}\text{C}^*(9.50)$, for which the distribution is roughly isotropic. The proton spectrum exhibits a conspicuous broad structure attributed to a ^{13}C level at $E_x = 8.4$ MeV, $\Gamma = 1.1 \pm 0.3$ MeV. (It seems probable that this level is to be identified with the $D_{\frac{3}{2}}$ level of similar width observed in $^{12}\text{C}(\text{n}, \text{n})^{12}\text{C}$ at $E_x = 8.33$ MeV: see Table 13.5.) Only one other level has a measurable width: $E_x = 7.64$ MeV, $\Gamma_{\text{lab}} = 70 \pm 15$ keV (compare Table 13.5) (1955MC75). It is of interest that the 7.47 and 7.53 MeV levels do not appear in $^{12}\text{C}(\text{n}, \text{n})^{12}\text{C}$ (1958WI01).

Angular distributions at low energies have been studied by (1954TA1A: $E_d = 0.52$ to 0.84 MeV), by (1956JU1F, 1956JU1E, 1957JU1A: $E_d = 0.60$ to 1.45 MeV), by (1955AL1D, 1955AL1E: $E_d = 1.4$ to 2.0 MeV), by (1956BE1H, 1956MC88: $E_d = 1.86$ to 2.86 MeV for the p_0 group - on and off resonances - and $E_d = 2.74$ and 2.89 MeV for the p_1 group), by (1954HO48, 1956BO08: $E_d = 3.2$ to 4.4 MeV) and by (1956KO1A, 1956VA17: $E_d = 0.26$ to 0.59 MeV). In the range $E_d = 1$ to 6 MeV, the ($^{12}\text{C} + \text{d}$) reactions are characterized by numerous strong, overlapping reso-

nances (see ^{14}N); the angular distributions show evidence of both stripping and compound nucleus formation, even below 1 MeV (1956JU1E, 1956KO1A, 1956VA17, 1957JU1A). From $E_d = 2$ to 6 MeV, angular distributions of the p_0 group (to $^{13}\text{C}_{\text{g.s.}}$) generally show a stripping maximum near 25° , as expected for an $l = 1$ transfer; several of the “resonances” appear most conspicuously at the angle. The p_1 -group, $^{13}\text{C}^*(3.09)$, show even stronger stripping effects, with a pronounced forward maximum (1956BO08, 1956MC88, 1958MC63). A detailed comparison of distributions for $^{12}\text{C}(d, p)^{13}\text{C}$ and $^{12}\text{C}(d, n)^{13}\text{N}$ at $E_d = 2.68$ and 3.26 MeV indicates equality of the ground-state reduced widths (1956BE1H, 1958MC63). At $E_d = 9$ MeV, a similar comparison yields $\gamma^2(^{13}\text{C})/\gamma^2(^{13}\text{N}) = 0.86$ (1956CA1D). See also (1955WI43).

Observed gamma rays are listed in Table 13.7. No γ -rays are observed with $E_\gamma = 3.9$ to 5.8 MeV with intensity $> 10\%$ of the 3.85 MeV γ -ray (1955BE62). An upper limit of 3% is placed on the fraction of cascade transitions from the 3.67 and 3.84 MeV levels via the 3.1 MeV level. The internal conversion coefficient of the 170 keV radiation indicates E1, though M1 is not excluded (1956MA1Q, 1956MA52). The internal pair formation coefficient for the 3.09 MeV level indicates an E1 transition (1952TH24); the angular correlation of internal pairs also indicates E1 (1954GO1E, 1956GO1K, 1958AR1B). Polarization of protons accompanying the formation of $^{13}\text{C}_{\text{g.s.}}$ and $^{13}\text{C}^*(3.1)$ has been studied by (1956HI1B, 1958BO67, 1958HE47, 1958HI74, 1958JU39, 1958JU42). The sense of polarization is correlated with the coupling of l_n and s_n : $P = \pm$ when $j = l \mp \frac{1}{2}$ (1958HE47). See (1954CH1C, 1957SA1C; theor.). See also (1952CA1B, 1954CA1B, 1955KH31, 1956CA1H, 1957SE1C), (1957DA1C; theor.) and ^{14}N .

Table 13.7: Gamma radiation from $^{12}\text{C}(d, p)^{13}\text{C}$

E_γ ^a (MeV \pm keV)	E_γ ^b (MeV \pm keV)	Reference
3.86 ± 20	(3.84 ± 30) ^c	(1955BE62)
3.844 ± 15		(1956MA1Q, 1956MA52)
0.1695 ± 0.4 ^f		(1956MA1Q, 1956MA52)
(3.76 ± 20) ^c	3.74 ± 30	(1955BE62)
(3.69 ± 20) ^c	3.675 ± 15 ^d	(1956MA1Q, 1956MA52)
(3.097 ± 5) ^e	3.082 ± 7	(1952TH24)

^a Uncorrected for Doppler shift.

^b Corrected for Doppler shift.

^c Doppler shift correction is not required for the 3.86 MeV radiation, but is required for the 3.67 MeV radiation (1956MA1Q, 1956MA52): see $^{10}\text{B}(\alpha, p)^{13}\text{C}$.

^d Value obtained by subtraction: $3.844 - 0.170$ (1956MA1Q, 1956MA52).

^e Doppler shift required (1952TH24).

^f From the proton groups $\Delta E = 170 \pm 3$ keV (1954SP01) and 170 ± 1.5 keV (1956DO41).

24. $^{12}\text{C}(t, d)^{13}\text{C}$ $Q_m = -1.313$

Not reported.

25. $^{12}\text{C}(\alpha, ^3\text{He})^{13}\text{C}$ $Q_m = -15.632$

Not reported.

26. $^{13}\text{C}(\gamma, n)^{12}\text{C}$ $Q_m = -4.946$

The cross section for neutron production has been determined to 38 MeV. The (γ, n) cross section exhibits two peaks at 13.3 ± 1 MeV ($\Gamma = 5 \pm 1$ MeV, $\sigma = 3.3$ mb) and at ≈ 22 MeV ($\sigma \approx 6$ mb, $\Gamma \approx 7$ MeV). The total absorption cross section, $\sigma(\gamma, xn) + \sigma(\gamma, p)$ shows maxima at $E_\gamma = 13.5$ and 25 MeV. The lower resonance is much too large to be explained on a single-particle model (1957CO57). See also (1949SE1B, 1953GO13, 1956CO72).

27. $^{13}\text{C}(\gamma, p)^{12}\text{B}$ $Q_m = -17.539$

The yield of β -particles from the ^{12}B decay has been determined to 45 MeV. The cross section shows a broad maximum of 8.8 mb near 25.5 MeV (1956CO72, 1957CO57).

28. $^{13}\text{C}(\gamma, \alpha)^9\text{Be}$ $Q_m = -10.654$

See (1953MI31).

29. $^{13}\text{C}(p, p')^{13}\text{C}^*$

Angular distributions of the 3.09 MeV γ -rays are isotropic for $E_p = 3.7$ to 4.2 MeV, consistent with the assignment $J = \frac{1}{2}$ to $^{13}\text{C}^*(3.09)$. Angular distributions of the 3.68 MeV radiation have also been studied near the $E_p = 4.5$ MeV resonance (1957BA29). See also (1952CO1C).

30. $^{13}\text{N}(\beta^+)^{13}\text{C}$ $Q_m = 2.222$

See ^{13}N .

$$31. \ ^{14}\text{C}(\text{p}, \text{d})^{13}\text{C} \quad Q_{\text{m}} = -5.947$$

Not reported.

$$32. \ ^{14}\text{C}(\text{d}, \text{t})^{13}\text{C} \quad Q_{\text{m}} = -1.915$$

At $E_{\text{d}} = 14.8$ MeV, triton groups have been observed leading to the ^{13}C states at 0, 3.09, 3.68 and 3.85 MeV (1958MO97). See ^{14}C .

$$33. \ ^{14}\text{C}(\text{}^3\text{He}, \alpha)^{13}\text{C} \quad Q_{\text{m}} = 12.404$$

Not reported.

$$34. \ ^{14}\text{N}(\text{n}, \text{d})^{13}\text{C} \quad Q_{\text{m}} = -5.319$$

At 14 MeV, deuteron groups to the 0 and 3.68 MeV (but not to the 3.09 and 3.84 MeV) states of ^{13}C have been observed (1957CA07). See also (1952LI1A) and ^{14}N .

$$35. \ ^{14}\text{N}(\text{p}, 2\text{p})^{13}\text{C} \quad Q_{\text{m}} = -7.546$$

At $E_{\text{p}} = 185$ MeV, the summed proton spectrum shows two peaks, corresponding to ejection of $\text{p}_{1/2}$ and $\text{p}_{3/2}$ protons, with binding energies of ≈ 7 and ≈ 15 MeV, $^{13}\text{C}^* = 0$ and ≈ 8 MeV (1958MA1B, 1958TY49).

$$36. \ ^{14}\text{N}(\text{d}, \text{}^3\text{He})^{13}\text{C} \quad Q_{\text{m}} = -2.052$$

Not reported.

$$37. \ ^{14}\text{N}(\text{t}, \alpha)^{13}\text{C} \quad Q_{\text{m}} = 12.267$$

Table 13.8: ^{13}C states from $^{15}\text{N}(d, \alpha)^{13}\text{C}$

(1951MA08)	(1957WA01)	
$^{13}\text{C}^*$ (MeV \pm keV)	$^{13}\text{C}^*$ (MeV \pm keV) ^a	$d\sigma/d\Omega$ ($\theta_{\text{lab}} = 12.6^\circ$) (mb/sr)
0	0	0.85 ± 0.09
3.083 ± 5	3.09	0.18 ± 0.03
3.677 ± 5	3.68	2.4 ± 0.3
	3.85	< 0.3
	6.87	1.3 ± 0.2 ^c
	7.47, 7.53, 7.64 ^b	1.1 ± 0.2
	8.80 ± 40	0.56 ± 0.11
	9.5	0.15 ± 0.04
	9.9	0.26 ± 0.06

^a Level energies for identification purposes only except for $^{13}\text{C}^*(8.80)$.

^b Not resolved.

^c Measured at $\theta = 18^\circ$.

This reaction has been observed at $E_t = 1.9$ MeV (1958JA06).

38. $^{15}\text{N}(n, t)^{13}\text{C}$ $Q_m = -9.903$

Not reported.

39. $^{15}\text{N}(p, ^3\text{He})^{13}\text{C}$ $Q_m = -10.668$

Not reported.

40. $^{15}\text{N}(d, \alpha)^{13}\text{C}$ $Q_m = 7.683$

Observed alpha particle groups are displayed in Table 13.8 (1951MA08, 1957WA01). The broad level at 8.4 MeV observed in $^{12}\text{C}(d, p)^{13}\text{C}$ does not appear in the present reaction; it is suggested that the direct (d, α) transition is forbidden by the nature of the configurations involved

(1957WA01). The angular distribution of the ground-state alpha particles at $E_d = 21$ MeV shows a maximum at 70° (c.m.) (1958FI27).

41. $^{16}\text{O}(n, \alpha)^{13}\text{C}$ $Q_m = -2.203$

See (1951HU1A, 1952LI1A) and ^{17}O .

^{13}N
(Fig. 24)

GENERAL:

Theory: See (1955LA1A, 1956DA1G, 1958FR1C, 1958HA1D, 1958SK1A).

1. $^{13}\text{N}(\beta^+)^{13}\text{C}$ $Q_m = 2.222$

Recent determinations of the half-life give 10.05 ± 0.03 min (1953CH34), 10.08 ± 0.04 min (1955WI43), 10.07 ± 0.06 min (1957NO17), 9.96 ± 0.03 min (1958AR15), 9.96 ± 0.03 min (1958DA09); see also (1957DE22). $E_\beta(\text{max}) = 1.202 \pm 0.005$ MeV (1950HO01), 1.185 ± 0.025 MeV (1954GR66), 1.190 ± 0.003 MeV (1958DA09). The positron spectrum shows no deviation from the allowed shape; it is concluded that the Fierz coefficient in the Fermi interaction is $< 11\%$. $\text{Log } ft = 3.66$ (1957DA08, 1958DA09). The positron polarization has been studied by (1957BO65, 1957HA27). The results indicate that the positrons are completely polarized and hence that Fermi transitions as well as G-T transitions exhibit the maximum effect of parity nonconservation.

2. $^9\text{Be}(^6\text{Li}, 2n)^{13}\text{N}$ $Q_m = 3.952$

See (1957NO17).

3. $^{10}\text{B}(^3\text{He}, n)^{12}\text{N}$ $Q_m = 1.46$ $E_b = 21.642$

At $E(^3\text{He}) = 2.54$ MeV, the cross section for formation of the ground state is $5.2_{-1.6}^{+2.1}$ mb. At $E(^3\text{He}) = 3.60$ MeV, the differential cross section for formation of the ground state at $\theta = 0^\circ$ is 0.73 ± 0.30 mb/sr (1957AJ71).

Table 13.9: Energy levels of ^{13}N

E_x (MeV \pm keV)	J^π	$\tau_{1/2}$ or Γ (keV)	Decay	Reactions
0	$(\frac{1}{2})^-$	$\tau_{1/2} = 10.02 \pm 0.02$ min	β^+	1, 2, 7, 9, 15, 16, 18, 19, 20, 21, 22
2.365 ± 3	$\frac{1}{2}^+$	$\Gamma = 32 \pm 2$	p, γ	7, 9, 12, 15, 22
3.507 ± 7	$\frac{3}{2}^-$	63 ± 6	p, γ	7, 9, 12, 15, 22
3.555 ± 10	$\frac{5}{2}^+$	61	p	12, 15, 22
6.379 ± 10	$\frac{5}{2}^+$	11	p	12
6.908 ± 10	$\frac{3}{2}^+$	115	p	12
7.415 ± 10	$(\frac{5}{2}^+)$	≈ 85	p	12
(8.08)	$(\frac{3}{2}^+)$	(350)	p	12
22.7 ± 300		≈ 1400	p	12
23.2		400	p, ^3He	4
24.5		550	p, ^3He	4
24.8		90	p, ^3He	4
25.2		120	p, ^3He	4

4. $^{10}\text{B}(^3\text{He}, \text{p})^{12}\text{C}$

$$Q_m = 19.702$$

$$E_b = 21.642$$

The yields of the protons to the ground and 4.4 MeV excited states of ^{12}C have been measured for $E(^3\text{He}) = 1.3$ to 5 MeV. Resonances are observed at 2.0, 3.7, 4.1 and 4.6 MeV, with widths of 0.5, 0.7, 0.12 and 0.15 MeV, respectively, corresponding to $^{13}\text{N}^*(23.2, 24.5, 24.8, 25.2)$. Angular distributions taken at six energies in the above range tend to be more asymmetric at the higher energies (1956SC01). See also (1956JO1B).

5. $^{10}\text{B}(^3\text{He}, \text{d})^{11}\text{C}$

$$Q_m = 3.206$$

$$E_b = 21.642$$

See ^{11}C .

6. $^{10}\text{B}(^3\text{He}, \alpha)^9\text{B}$

$$Q_m = 12.139$$

$$E_b = 21.642$$

See ^9B .

Table 13.10: Resonances in $^{12}\text{C}(\text{p}, \gamma)^{13}\text{N}$

E_p (keV)	Γ_{lab} (keV)	σ_{res} (mb)	$\omega\Gamma_\gamma$ (eV)	$^{13}\text{N}^*$ (MeV)	References
456.8 ± 0.5	39.5 ± 1.0			2.363	(1953HU18)
456 ± 2	35				(1949FO18)
450	35	127	0.67		(1951SE1B, 1951SE67)
1697 ± 12	74 ± 9			3.507	(1949VA1A)
1698 ± 5	70 ± 10	35	1.39		(1951SE1B, 1951SE67)

 7. $^{10}\text{B}(\alpha, \text{n})^{13}\text{N}$ $Q_m = 1.065$

Measurements at $E_\alpha = 8$ MeV with a proton recoil telescope and a neutron threshold detector are reported to indicate ^{13}N states at 2.4 ± 0.3 , 3.6 ± 0.3 , (4.3 ± 0.3) and 5.0 ± 0.3 MeV (1956QU1A). See also (1957BA1K) and ^{14}N .

 8. $^{11}\text{B}(^3\text{He}, \text{n})^{13}\text{N}$ $Q_m = 10.179$

Not reported.

 9. (a) $^{12}\text{C}(\text{p}, \gamma)^{13}\text{N}$ $Q_m = 1.941$

 (b) $^{12}\text{C}(\text{p}, \gamma\text{p}')^{12}\text{C}$

Two resonances for capture radiation are reported, at $E_p = 0.46$ and 1.70 MeV (Table 13.10). The resonance at $E_p = 1.75$ MeV observed in $^{12}\text{C}(\text{p}, \text{p})^{12}\text{C}$ does not appear in the γ -excitation curve (1951SE67). The displacement of the lower level ($^{13}\text{N}^*(2.37)$, $J = \frac{1}{2}^+$) from its mirror in $^{13}\text{C}^*(3.09)$ is ascribed to the large reduced width (1951EH1A, 1952TH1D). The angular distribution of the ground-state radiation from the upper resonance ($^{13}\text{N}^*(3.51)$, $J = \frac{3}{2}^-$) has the form $W(\theta) = 1 - 0.52 \cos^2 \theta$ (1951DA1A, 1951DA1B).

The capture cross section at low energy is of interest in connection with stellar energy generation. Measurements have been reported in the range $E_p = 80$ to 360 keV by (1950BA1B, 1950HA1A, 1957DE22, 1957LA1C). At 80 keV, $\sigma = (1.4 \pm 0.4) \times 10^{-5}$ μb ; from 80 to 126 keV, the course of the cross section is reasonably well accounted for by extrapolation of the $E_p = 0.46$ MeV resonance (1957LA1C: see also (1957DE22)).

From $E_p = 5$ to 11 MeV, the cross section for formation of ^{13}N changes only from 2.5 to 1.8 mb; this small change strongly indicates the predominance of direct capture in this region

(1955CO1B). According to (1956RE39), however, the 90° differential cross section for formation of ^{13}N is $< 1 \mu\text{b/sr}$ at $E_p = 4.8 \text{ MeV}$. See also (1956CH1D).

In the range $E_p = 1.2$ to 2.5 MeV , reaction (b) is observed, involving a γ -transition to the 2.37 MeV state. Excitation functions at $\theta = 0^\circ$ and 90° indicate interference between p-wave resonant capture at $E_p = 1.70 \text{ MeV}$, with $\Gamma_\gamma = 0.04 \text{ eV}$, and direct p-wave capture (1954WO09). The angular distributions at $E_p = 1.37$ and 1.58 MeV have the form $W(\theta) = (0.02 \pm 0.02) + \sin^2 \theta$ (1955HE1F).

$$10. \text{}^{12}\text{C}(\text{p}, \text{n})\text{}^{12}\text{N} \qquad Q_m = -18.24 \qquad E_b = 1.941$$

See (1957ST1D, 1958TA03) and ^{12}N .

$$11. \text{}^{12}\text{C}(\text{p}, \text{pn})\text{}^{11}\text{C} \qquad Q_m = -18.722 \qquad E_b = 1.941$$

See (1947CH1A, 1948MC1A, 1958WH34).

$$12. \text{(a) } \text{}^{12}\text{C}(\text{p}, \text{p})\text{}^{12}\text{C} \qquad E_b = 1.941 \\ \text{(b) } \text{}^{12}\text{C}(\text{p}, \text{p}')\text{}^{12}\text{C}^*$$

Elastic scattering studies indicate a number of pronounced resonances in the range $E_p = 0$ to 6 MeV : see Table 13.11. The first five excited states correspond in character and approximately in reduced width to those of ^{13}C : see $^{12}\text{C}(\text{d}, \text{p})\text{}^{13}\text{C}$ and $^{12}\text{C}(\text{n}, \text{n})\text{}^{12}\text{C}$. The relatively large reduced widths of the first and third ($s_{1/2}$ and $d_{5/2}$) excited states indicate a single-particle character (1953JA1B). The small and roughly equal widths of $^{13}\text{N}^*(6.4, 6.9)$ suggest that they may comprise a doublet, built upon $^{12}\text{C}^*(4.4) + \text{p}$ (1956RE39: see also (1953BL1A, 1953MA1D, 1956SC29)). Angular distribution measurements above $E_p = 10 \text{ MeV}$ generally show direct interaction effects: see ^{12}C . Some form of resonance structure may exist near 23 MeV (1955KI43). See also (1956KL55).

The yields of 4.4 MeV gamma rays and inelastic protons from $^{12}\text{C}^*(4.4)$ show resonances at $E_p = 5.39$ and 5.93 MeV (1953MA1D, 1956BR27, 1957LI1B). Angular distributions of inelastic protons at $E_p = 6.1$ to 6.9 MeV do not fit direct interaction theory and suggest the effects of still higher compound nucleus levels (1956BR27).

Polarization studies for $E_p < 6 \text{ MeV}$ are reported by (1956GA66, 1956SO1C, 1958WA1D): see also ^{12}C . See also (1955DE50, 1956ER1A, 1956NI1B, 1957GL58; theor.) and (1957GO1D).

$$13. \text{}^{12}\text{C}(\text{p}, \text{d})\text{}^{11}\text{C} \qquad Q_m = -16.495 \qquad E_b = 1.941$$

Table 13.11: ^{13}N levels from $^{12}\text{C}(p, p)^{12}\text{C}$ and $^{12}\text{C}(p, p')^{12}\text{C}^*$

E_{res} (MeV \pm keV)	$^{13}\text{N}^*$ (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	l_p	J^π	θ_p^2
0.461 ± 3^a	2.367	31	0	$\frac{1}{2}^+$	0.54
1.698^b	3.508	55	1	$\frac{3}{2}^-$	0.031
1.748^b	3.555	61	2	$\frac{5}{2}^+$	0.21
4.808^c	6.379	11	2	$\frac{5}{2}^+$	0.0031
$(5.05)^e$		(75)	(0)	$(\frac{1}{2}^+)$	(0.014)
5.381 ± 7^d	6.908	115	2	$\frac{3}{2}^+$	0.012
5.930 ± 7^f	7.415	≈ 85	2	$\frac{5}{2}^+$	
6.65^g	(8.08)	(350)	(2)	$(\frac{3}{2}^+)$	(0.11)

^a (1953JA1B). (1954MI05) finds $E_{\text{res}} = 0.462$ MeV, $\Gamma = 32$ keV.

^b (1953JA1B).

^c (1956RE39).

^d (1956RE39): parameters estimated from elastic scattering; $\theta_p^2 = 0.2$.

E_{res} from $^{12}\text{C}(p, p')^{12}\text{C}^*$ (1957LI1B). See also (1956SC29).

^e (1956SC29): $E_p = 4.8$ MeV, $J = \frac{5}{2}^+$ level not observed.

^f $^{12}\text{C}(p, p')^{12}\text{C}^*$ (1957LI1B). (1956BR27) finds $E_{\text{res}} = 5.891$ MeV, $\Gamma_{\text{lab}} = 59$ keV. See also (1956SC29).

^g (1956SC29). See, however, (1956BR27).

See ¹¹C.

14. $^{12}\text{C}(p, \alpha)^9\text{Be}$

$$Q_m = -7.563$$

$$E_b = 1.941$$

See ⁹B.

15. $^{12}\text{C}(d, n)^{13}\text{N}$

$$Q_m = -0.286$$

Neutron groups have been observed corresponding to excited states of ^{13}N at 2.29 ± 0.12 (1949GR1A), 2.38 ± 0.05 MeV (1953MI10) and 3.48 ± 0.12 (1949GR1A), 3.74 ± 0.05 (1957GR1A), 3.53 ± 0.05 MeV (1953MI10). The angular distributions of the ground state group and the groups corresponding to the 2.37 and (3.51 ± 3.56) MeV states at $E_d = 9.0$ MeV are consistent with $l_p = 1, 0$ and 2. The dimensionless reduced widths of the ground and (3.56) MeV states are

respectively 0.056 and 0.19 (1957CA02: see also (1953MI10)). (1958MC63) finds that the reduced widths of the ground states of ^{13}C and ^{13}N are the same, 0.09 ± 0.035 (see also (1956BE1H, 1956CA1D, 1958KA16) and $^{12}\text{C}(\text{d}, \text{p})^{13}\text{C}$). In the range $E_{\text{d}} = 2.8$ to 3.7 MeV, a single neutron threshold is observed, at $E_{\text{d}} = 3.09 \pm 0.02$ MeV, corresponding to $^{13}\text{N}^*(2.36 \pm 0.02)$; the slow rise above threshold is attributed to p-wave neutron emission (1955MA76).

Polarization of neutrons has been studied for $E_{\text{d}} = 2.5$ to 3.6 MeV by (1957HA1J). See also (1956BO1F, 1956BO43, 1956DE1D).

16. $^{12}\text{C}(^3\text{He}, \text{d})^{13}\text{N}$ $Q_{\text{m}} = -3.553$

See (1952FR1A, 1958WE1E).

17. $^{12}\text{C}(\alpha, \text{t})^{13}\text{N}$ $Q_{\text{m}} = -17.872$

Not reported.

18. $^{13}\text{C}(\text{p}, \text{n})^{13}\text{N}$ $Q_{\text{m}} = -3.005$

$$E_{\text{thresh.}} = 3.2372 \pm 0.0016 \text{ (1958BO76)}.$$

See also (1950RI59, 1955MA84, 1958BI1B) and ^{14}N .

19. $^{13}\text{C}(^3\text{He}, \text{t})^{13}\text{N}$ $Q_{\text{m}} = -2.240$

See (1952FR1A).

20. $^{14}\text{N}(\gamma, \text{n})^{13}\text{N}$ $Q_{\text{m}} = -10.551$

See ^{14}N .

21. $^{14}\text{N}(\text{p}, \text{d})^{13}\text{N}$ $Q_{\text{m}} = -8.324$

See ^{14}N .

$$22. \text{}^{14}\text{N}(\text{d}, \text{t})^{13}\text{N} \quad Q_{\text{m}} = -4.292$$

At $E_{\text{d}} = 14.8$ MeV, triton groups are observed corresponding to the states at 0, 2.37 and (3.51 ± 3.56) MeV. The cross section for the transition to the 2.37 MeV state is two orders of magnitude smaller than that for the ground state transition. Transitions to $^{13}\text{N}^*(2.37, 3.56)$ are shell-model forbidden (1957WA01).

$$23. \text{}^{14}\text{N}(\text{}^3\text{He}, \alpha)^{13}\text{N} \quad Q_{\text{m}} = 10.027$$

Not reported.

$$24. \text{}^{15}\text{N}(\text{p}, \text{t})^{13}\text{N} \quad Q_{\text{m}} = -12.908$$

Not reported.

$$25. \text{}^{16}\text{O}(\text{p}, \alpha)^{13}\text{N} \quad Q_{\text{m}} = -5.208$$

See (1958WH34).

References

(Closed 1 December 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1947CH1A W.W. Chupp and E.M. McMillan, Phys. Rev. 72 (1947) 873
1947HU03 D.J. Hughes, C. Eggler and C.M. Huddleston, Phys. Rev. 71 (1947) 269
1948MC1A E.M. McMillan and R.D. Miller, Phys. Rev. 73 (1948) 80
1948SN1A Snell, Science 108 (1948) 167
1949FO18 W.A. Fowler and C.C. Lauritsen, Phys. Rev. 76 (1949) 314
1949GR1A J.C. Grosskreutz, Phys. Rev. 76 (1949) 482
1949HU41 E.L. Hudspeth and C.P. Swann, Phys. Rev. 76 (1949) 1150
1949SE1B Sexl, Acta Phys. Aust. 2 (1949) 377
1949VA1A Van Patter, Phys. Rev. 76 (1949) 1264
1950BA1B C.L. Bailey and W.R. Stratton, Phys. Rev. 77 (1950) 194
1950FR1A G. Freier, M. Fulk, E.E. Lampi and J.H. Williams, Phys. Rev. 78 (1950) 508
1950HA1A R.N. Hall and W.A. Fowler, Phys. Rev. 77 (1950) 197
1950HO01 W.F. Hornyak and T. Lauritsen, Phys. Rev. 77 (1950) 160
1950MI1A D.W. Miller, Phys. Rev. 78 (1950) 806
1950RI1B Ricamo, Zunti, Baldinger and Huber, Helv. Phys. Acta 23 (1950) 508
1950RI59 H.T. Richards, R.V. Smith and C.P. Browne, Phys. Rev. 80 (1950) 524
1951BO45 C.K. Bockelman, D.W. Miller, R.K. Adair and H.H. Barschall, Phys. Rev. 84 (1951) 69
1951DA1A R.B. Day and J.E. Perry, Jr., Phys. Rev. 81 (1951) 662; J14
1951DA1B Day, Ph.D. Thesis, CIT (1951)
1951EH1A J.B. Ehrman, Phys. Rev. 81 (1951) 412
1951HU1A Huber, Baldinger and Proctor, Helv. Phys. Acta 24 (1951) 302
1951MA08 R. Malm and W.W. Buechner, Phys. Rev. 81 (1951) 519
1951RI1A Ricamo and Zunti, Helv. Phys. Acta 24 (1951) 419
1951RI1B Ricamo, Nuovo Cim. 8 (1951) 893

1951RO1B Rotblat, Nature 167 (1951) 1027
 1951RO1C J. Rotblat, Phys. Rev. 83 (1951) 1271
 1951SE1B Seagrave, Ph.D. Thesis, CIT (1951)
 1951SE67 J.D. Seagrave, Phys. Rev. 84 (1951) 1219
 1951ST19 E.N. Strait, D.M. Van Patter, W.W. Buechner and A. Sperduto, Phys. Rev. 81 (1951) 747
 1951VA1A Van Patter, Buechner and Sperduto, Phys. Rev. 82 (1951) 248
 1952AJ38 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 24 (1952) 321
 1952BL1B Black, Phys. Rev. 87 (1952) 205; L7
 1952BR61 J.E. Brolley, Jr., J.L. Fowler and L.K. Schlacks, Phys. Rev. 88 (1952) 618
 1952CA1B Catala, Senent and Aguilar, An. Real. Soc. Espan. Fis. y Quim. A48 (1952) 73; Phys. Abs. 57 (1954) 1709
 1952CO1C D.B. Cowie, N.P. Heydenburg and G.C. Phillips, Phys. Rev. 87 (1952) 304
 1952FR1A Fremlin, Proc. Phys. Soc. (London) A65 (1952) 762
 1952LI1A A.B. Lillie, Phys. Rev. 87 (1952) 716
 1952SH44 R.K. Sheline, Phys. Rev. 87 (1952) 557
 1952TH1D R.G. Thomas, Phys. Rev. 88 (1952) 1109
 1952TH24 R.G. Thomas and T. Lauritsen, Phys. Rev. 88 (1952) 969
 1953BA18 G.A. Bartholomew and B.B. Kinsey, Can. J. Phys. 31 (1953) 49
 1953BL1A Bleuler, Helv. Phys. Acta 26 (1953) 597
 1953CA1B Catala, Senent and Casanova, An. Real. Soc. Espan. Fis. y Quim. A49 (1953) 91; Nucl. Sci. Abs. (1953)
 1953CH34 J.L.W. Churchill, W.M. Jones and S.E. Hunt, Nature 172 (1953) 460
 1953GO13 F.K. Goward and J.J. Wilkins, Proc. Roy. Soc. A217 (1953) 357
 1953HU18 S.E. Hunt and W.M. Jones, Phys. Rev. 89 (1953) 1283
 1953HU1C E.L. Hubbard, L. Ruby and W.F. Stubbins, Phys. Rev. 92 (1953) 1494
 1953JA1B Jackson and Galonsky, Phys. Rev. 89 (1953) 370
 1953KH1A Khromchenko, Dokl. Akad. Nauk. SSSR 93 (1953) 451
 1953MA1D Martin, Schneider and Sempert, Helv. Phys. Acta 26 (1953) 595
 1953MI10 R. Middleton, F.A. El Bedewi and C.T. Tai, Proc. Phys. Soc. (London) A66 (1953) 95
 1953MI31 C.H. Millar and A.G.W. Cameron, Can. J. Phys. 31 (1953) 723
 1953NE1A Nereson and Darden, Phys. Rev. 89 (1953) 775

- 1953SH64 E.S. Shire, J.R. Wormald, G. Lindsay-Jones, A. Lundan and A.G. Stanley, *Phil. Mag.* 44 (1953) 1197
- 1953TA06 F.L. Talbott and N.P. Heydenburg, *Phys. Rev.* 90 (1953) 186
- 1954BE08 W.E. Bennett, P.A. Roys and B.J. Toppel, *Phys. Rev.* 93 (1954) 924A
- 1954BU06 W.H. Burke, J.R. Risser and G.C. Phillips, *Phys. Rev.* 93 (1954) 188
- 1954CA1B Catala and Senent, *An. Real. Soc. Espan. Fis. y Quim.* A50 (1954) 55
- 1954CH1C W.B. Cheston, *Phys. Rev.* 96 (1954) 1590
- 1954FA1A Fader and Sperduto, *Phys. Rev.* 94 (1954) 748; C8
- 1954FA1B Fader, MIT Prog. Rept. (May, 1954)
- 1954FR24 R.G. Freemantle, W.M. Gibson and J. Rotblat, *Phil. Mag.* 45 (1954) 1200
- 1954GO1E Gorodetzky, Armbruster, Chevallier and Gallmann, *Compt. Rend.* 239 (1954) 1623
- 1954GR66 T. Grabowski and L. Natanson, *Bull. Acad. Pol. Sci.* 2 (1954) 379
- 1954HO48 H.D. Holmgren, J.M. Blair, B.E. Simmons, T.F. Stratton and R.V. Stuart, *Phys. Rev.* 95 (1954) 1544
- 1954HU1A Huber and Budde, *Helv. Phys. Acta* 27 (1954) 512
- 1954ME95 R.W. Meier, P. Scherrer and G. Trumpy, *Helv. Phys. Acta* 27 (1954) 577
- 1954MI05 E.A. Milne, *Phys. Rev.* 93 (1954) 762
- 1954SP01 A. Sperduto, W.W. Buechner, C.K. Bockelman and C.P. Browne, *Phys. Rev.* 96 (1954) 1316
- 1954ST20 A.G. Stanley, *Phil. Mag.* 45 (1954) 430
- 1954TA1A Takemoto, Dazai, Chiba, Ito, Suganomata and Watanabe, *J. Phys. Soc. Jpn.* 9 (1954) 447
- 1954TR09 R.E. Trumble, Jr., *Phys. Rev.* 94 (1954) 748A
- 1954WO09 H.H. Woodbury, A.V. Tollestrup and R.B. Day, *Phys. Rev.* 93 (1954) 1311
- 1955AJ61 F. Ajzenberg and T. Lauritsen, *Revs. Mod. Phys.* 27 (1955) 77
- 1955AL16 H.R. Allan and N. Sarma, *Proc. Phys. Soc. (London)* A68 (1955) 535
- 1955AL1D Alba, Fernandez, Mazari, Serment and Vazquez, *Rev. Mex. Fis.* 4 (1955) 207
- 1955AL1E Alba, Brody, Fernandez, Mazari, Serment and Vazquez, *An. del Inst. Fis., Univ. Nacional. Autonoma de Mexico* 1 (1955) 1
- 1955AU1A T. Auerbach and J.B. French, *Phys. Rev.* 98 (1955) 1276
- 1955BA1N M.E. Battat and E.R. Graves, *Phys. Rev.* 97 (1955) 1266
- 1955BE1D Beyster, Henkel and Nobles, *Phys. Rev.* 97 (1955) 563
- 1955BE1F Beyster, Henkel, Nobles and Kister, *Phys. Rev.* 98 (1955) 1216

1955BE62 R.D. Bent, T.W. Bonner and R.F. Sippel, Phys. Rev. 98 (1955) 1237
 1955BI26 C.B. Bigham, K.W. Allen and E. Almqvist, Phys. Rev. 99 (1955) 631A
 1955BR1A R.J. Breen and M.R. Hertz, Phys. Rev. 98 (1955) 599
 1955BU56 R. Budde and P. Huber, Helv. Phys. Acta 28 (1955) 49
 1955CO1B B.L. Cohen, Phys. Rev. 100 (1955) 206
 1955DE50 P.S. de Toledo and S. Watanabe, An. Acad. Brasil. Cienc. 27 (1955) 471; Phys. Abs. 644 (1959)
 1955FR35 G.M. Frye, Jr., L. Rosen and L. Stewart, Phys. Rev. 99 (1955) 1375
 1955GR21 E.R. Graves and R.W. Davis, Phys. Rev. 97 (1955) 1205
 1955HE1F Heiberg, James and Alexander, Can. J. Phys. 33 (1955) 34
 1955JE1A B. Jennings, J. Weddell, I. Alexeff and R.L. Hellens, Phys. Rev. 98 (1955) 582
 1955KH31 L.M. Khromchenko and V.A. Blinov, Zh. Eksp. Teor. Fiz. 28 (1955) 741; JETP (Sov. Phys.) 1 (1955) 596
 1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252
 1955KI43 B.B. Kinsey, Phys. Rev. 99 (1955) 332
 1955LA1A Lane, Proc. Phys. Soc. (London) A68 (1955) 197
 1955LI1C R.N. Little, B.P. Leonard, J.T. Prud'homme and L. D. Vincent, Phys. Rev. 98 (1955) 634
 1955MA1G MacGregor, Ball and Booth, Phys. Rev. 100 (1955) 1793A
 1955MA1J L. Madansky and G.E. Owen, Phys. Rev. 99 (1955) 1608
 1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847
 1955MA84 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 91
 1955MC75 J.N. McGruer, E.K. Warburton and R.S. Bender, Phys. Rev. 100 (1955) 235
 1955RI1B Risser, Price and Class, Phys. Rev. 98 (1955) 1183, RA3
 1955TA1B Taylor, Lonsjo and Bonner, Phys. Rev. 100 (1955) 174
 1955TA28 N.W. Tanner, Proc. Phys. Soc. (London) A68 (1955) 1195
 1955WA1A Wapstra, Physica 21 (1955) 367
 1955WA27 M. Walt and J.R. Beyster, Phys. Rev. 98 (1955) 677
 1955WA30 A. Ward and P.J. Grant, Proc. Phys. Soc. (London) A68 (1955) 637
 1955WI25 H.B. Willard, J.K. Bair and J.D. Kington, Phys. Rev. 98 (1955) 669
 1955WI43 D.H. Wilkinson, Phys. Rev. 100 (1955) 32
 1956AL1F S.K. Allison, P.G. Murphy and E. Norbeck, Phys. Rev. 102 (1956) 1182

- 1956BE1D Becker and Barschall, Phys. Rev. 102 (1956) 1384
- 1956BE1H R.E. Benenson, K.W. Jones and M.T. McEllistrem, Phys. Rev. 101 (1956) 308
- 1956BE32 J.R. Beyster, M. Walt and E.W. Salmi, Phys. Rev. 104 (1956) 1319
- 1956BO08 T.W. Bonner, J.T. Eisinger, A.A. Kraus, Jr. and J.B. Marion, Phys. Rev. 101 (1956) 209
- 1956BO1F Bogdanov, Vlasov, Kalinin, Rybakov and Sidorov, Physica 22 (1956) 1150
- 1956BO43 G.F. Bogdanov, N.A. Vlasov, S.P. Kalinin, B.V. Rybakov and V.A. Sidorov, Zh. Eksp. Teor. Fiz. 30 (1956) 981; JETP (Sov. Phys.) 3 (1956) 793
- 1956BO61 T.W. Bonner, A.A. Kraus, Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348
- 1956BR1E R.M. Brugger, J.E. Evans, E.G. Joki and R.S. Shankland, Phys. Rev. 104 (1956) 1054
- 1956BR27 C.P. Browne and J.R. Lamarsh, Phys. Rev. 104 (1956) 1099
- 1956CA1D Calvert, Jaffe and Maslin, Phys. Rev. 101 (1956) 501
- 1956CA1H Catala, Villar, Bonett and Miralles, An. Real. Soc. Espan. Fis. y Quim. A52 (1956) 85
- 1956CH1D Chick, Evans, Hancock, Hunt and Pope, Proc. Phys. Soc. (London) A69 (1956) 624
- 1956CO72 B.C. Cook, A.S. Penfold and V.L. Telegdi, Phys. Rev. 104 (1956) 554
- 1956CU1A G. Culler, S. Fernbach and N. Sherman, Phys. Rev. 101 (1956) 1047
- 1956DA1G Davidson and Giambiagi, Centro Brasil. Pesquisas Fis., Notas Fis. Vol. II, No. 12 (1956)
- 1956DA23 R.B. Day, Phys. Rev. 102 (1956) 767
- 1956DE1C De Saussure and Guth, Bull. Amer. Phys. Soc. 1 (1956) 21
- 1956DE1D De Pangher, HW-47284 (1956)
- 1956DO1D Dolan, Fincher, Kenny, Berko and Whitehead, Bull. Amer. Phys. Soc. 1 (1956) 339
- 1956DO41 R.A. Douglas, J.W. Broer, R. Chiba, D.F. Herring and E.A. Silverstein, Phys. Rev. 104 (1956) 1059
- 1956EL1A El-Bedewi, Proc. Phys. Soc. (London) A69 (1956) 221
- 1956ER1A T. Eriksson, Nucl. Phys. 2 (1956) 91
- 1956FL1B Flerov and Talyzin, Sov. J. Nucl. Energy 4 (1956) 617
- 1956GA66 B.V. Gavrilovskii, V.Y. Golovnya, K.V. Karadzhev, A.P. Kliucharev and V.I. Manko, Dokl. Akad. Nauk SSSR 111 (1956) 59; Sov. Phys. Dokl. 1 (1957) 611
- 1956GO1K Gorodetzky, Armbruster, Chevallier and Gallmann, J. Phys. Rad. 17 (1956) 548
- 1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28
- 1956HA1E Hall and Bonner, Bull. Amer. Phys. Soc. 1 (1956) 96

1956HI1B P. Hillman, Phys. Rev. 104 (1956) 176
1956HU1A Hughes and Carter, BNL-400 (1956)
1956JA28 D.B. James, G.A. Jones and D.H. Wilkinson, Phil. Mag. 1 (1956) 949
1956JO1B Johnston and Holmgren, Bull. Amer. Phys. Soc. 1 (1956) 134
1956JU1E Juric, Physica 22 (1956) 1154A
1956JU1F Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 6 (1956) 35
1956KA1B Kalos and Goldstein, NDA 12-16 (1956)
1956KL55 A.P. Kliucharev, L.I. Bolotin and V.A. Lutsik, Zh. Eksp. Teor. Fiz. 30 (1956) 573;
JETP (Sov. Phys.) 3 (1956) 463
1956KO1A Koudijs, Thesis, Univ. of Utrecht (1956)
1956KR1A Kreger, Bolotin and Edelsack, USNRDL TR 81 (1956)
1956KR1B Kreger, Bolotin and Edelsack, Bull. Amer. Phys. Soc. 1 (1956) 325
1956KU1A Kurath, Phys. Rev. 101 (1956) 216
1956LA1C Langsdorf, Lane and Monahan, ANL 5567 (1956)
1956MA1Q R.J. Mackin, Jr., W.R. Mills, Jr. and J. Thirion, J. Phys. Rad. 17 (1956) 551
1956MA52 R.J. Mackin, Jr., W.R. Mills, Jr. and J. Thirion, Phys. Rev. 102 (1956) 802
1956MC70 B.M. McCormac, M.F. Steuer, C.D. Bond and F.L. Hereford, Phys. Rev. 104 (1956)
718
1956MC88 M.T. McEllistrem, K.W. Jones, R. Chiba, R.A. Douglas, D.F. Herring and E.A. Sil-
verstein, Phys. Rev. 104 (1956) 1008
1956MU1A C.O. Muehlhause, S.D. Bloom, H.E. Wegner and G.N. Glasoe, Phys. Rev. 103 (1956)
720
1956NI1B Nishimura and Ruderman, Bull. Amer. Phys. Soc. 1 (1956) 383
1956NO1A Norbeck, Bull. Amer. Phys. Soc. 1 (1956) 329
1956PA1B Papkow, Z. Naturforsch. A11 (1956) 776
1956PA23 E.B. Paul, Physica 22 (1956) 1140A
1956PI1A G.F. Pieper and G.S. Stanford, Phys. Rev. 101 (1956) 672
1956QU1A A.R. Quinton and W.T. Doyle, Phys. Rev. 101 (1956) 669
1956RE39 C.W. Reich, G.C. Phillips and J.L. Russell, Jr., Phys. Rev. 104 (1956) 143
1956SA1E Sachs, Phys. Rev. 103 (1956) 671
1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser, Jr., Phys. Rev. 104 (1956)
1064
1956SC29 H. Schneider, Helv. Phys. Acta 29 (1956) 55

1956SO1C Sorokin and Taranov, Dokl. Akad. Nauk SSSR 111 (1956) 82
 1956VA17 F.P.G. Valckx, Thesis, Univ. of Utrecht (1956)
 1957AJ71 F. Ajzenberg-Selove, M.L. Bullock and E. Almqvist, Phys. Rev. 108 (1957) 1284
 1957AM48 O. Ames, G.E. Owen and C.D. Swartz, Phys. Rev. 106 (1957) 775
 1957BA1H Barker, Phil. Mag. 2 (1957) 780
 1957BA1K Barker and Mann, Phil. Mag. 2 (1957) 5
 1957BA29 J.K. Bair, H.O. Cohn and H.B. Willard, ORNL-2430 (1957) 26
 1957BO13 R.O. Bondelid, K.L. Dunning and F.L. Talbott, Phys. Rev. 105 (1957) 193
 1957BO65 F. Boehm, T.B. Novy, C.A. Barnes and B. Stech, Phys. Rev. 108 (1957) 1497
 1957CA02 J.M. Calvert, A.A. Jaffe and E.E. Maslin, Proc. Phys. Soc. (London) A70 (1957) 78
 1957CA07 R.R. Carlson, Phys. Rev. 107 (1957) 1094
 1957CO57 B.C. Cook, Phys. Rev. 106 (1957) 300
 1957CO68 H.E. Conzett and D.G. Hoffman, Bull. Amer. Phys. Soc. 2 (1957) 382, J10
 1957DA08 H. Daniel and U. Schmidt-Rohr, Z. Naturforsch. A12 (1957) 750
 1957DA1C Dabrowski and Tulczyjew, Acta Phys. Pol. 16 (1957) 231
 1957DE22 A.S. Deineko, A.I. Taranov and A.K. Valter, Zh. Eksp. Teor. Fiz. 32 (1957) 251; JETP (Sov. Phys.) 5 (1957) 201
 1957GA01 A. Galonsky, C.D. Moak, R.L. Traughber and C.M. Jones, Bull. Amer. Phys. Soc. 2 (1957) 51, RA1
 1957GL58 A.E. Glassgold and P.J. Kellogg, Phys. Rev. 107 (1957) 1372
 1957GO1D Gooding, Bull. Amer. Phys. Soc. 2 (1957) 350
 1957GR1A Grismore and Parkinson, Rev. Sci. Instrum. 28 (1957) 245
 1957HA1J Haerberli and Rolland, Bull. Amer. Phys. Soc. 2 (1957) 234
 1957HA27 S.S. Hanna and R.S. Preston, Phys. Rev. 108 (1957) 160
 1957HO61 H.D. Holmgren, E.A. Wolicki and R.L. Johnston, Bull. Amer. Phys. Soc. 2 (1957) 181, E7
 1957HU1D Hughes and Schwartz, BNL-325, Suppl. 1 (1957)
 1957HU1E Hunting and Wall, Bull. Amer. Phys. Soc. 2 (1957) 181
 1957JA37 N. Jarmie, J.D. Seagrave et al, LA-2014 (1957)
 1957JU1A Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 7 (1957) 1
 1957KH1A Khaletskii, Dokl. Akad. Nauk SSSR 113 (1957) 305; Sov. Phys., Dokl. 2 (1958) 129
 1957KH1B Khaletskii, Dokl. Akad. Nauk SSSR 113 (1957) 553; Sov. Phys. Dokl. 2 (1958) 152
 1957LA14 A. Langsdorf, Jr., R.O. Lane and J.E. Monahan, Phys. Rev. 107 (1957) 1077

1957LA1C W.A. Lamb and R.E. Hester, Phys. Rev. 107 (1957) 550
 1957LI1B Lidofsky, Weil, Bent and Jones, Bull. Amer. Phys. Soc. 2 (1957) 29
 1957MC1B McCormac, Steuer, Bond and Hereford, Phys. Rev. 108 (1957) 116
 1957NO14 E. Norbeck, Jr., Phys. Rev. 105 (1957) 204
 1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
 1957RI38 J.R. Risser, J.E. Price and C.M. Class, Phys. Rev. 105 (1957) 1288
 1957RO1F Roy, Lagasse, Goes, Achari and de Henau, Compt. Rend. 244 (1957) 2907
 1957SA1C J. Sawicki, Phys. Rev. 106 (1957) 172
 1957SE1C Senent, Villar and Bonet, An. Real. Soc. Espan. Fis. y Quim. A53 (1957) 109; Phys. Abs. 8264 (1958)
 1957ST1D Stafford, Tornabene and Whitehead, Phys. Rev. 106 (1957) 831
 1957ST1F Strizhak, Sov. J. At. Energy 2 (1957) 72
 1957VA11 D.M. Van Patter and W. Whaling, Revs. Mod. Phys. 29 (1957) 757
 1957VO25 P. von Hermann and G.F. Pieper, Phys. Rev. 105 (1957) 1556
 1957WA01 E.K. Warburton and J.N. McGruer, Phys. Rev. 105 (1957) 639
 1957ZA1A Zabel, WASH 192 (1957)
 1958AR15 S.E. Arnell, J. Dubois and O. Almen, Nucl. Phys. 6 (1958) 196
 1958AR1B Armbuster, Ann. Phys. (France) 3 (1958) 88
 1958AS63 V.J. Ashby, H.C. Catron, L.L. Newkirk and C.J. Taylor, Phys. Rev. 111 (1958) 616
 1958BA03 W.P. Ball, M. MacGregor and R. Booth, Phys. Rev. 110 (1958) 1392
 1958BI1B Bichsel, Phys. Rev. Lett. 1 (1958) 384
 1958BO67 M.S. Bokhari, J.A. Cookson, B. Hird and B. Weesakul, Proc. Phys. Soc. (London) A72 (1958) 88
 1958BO76 R.O. Bondelid and C.A. Kennedy, NRL Rept. 5083 (1958)
 1958BR16 A. Bratenahl, J.M. Peterson and J.P. Steering, Phys. Rev. 110 (1958) 927
 1958BR1F Braley and Cook, Bull. Amer. Phys. Soc. 3 (1958) 267
 1958CO07 J.P. Conner, Phys. Rev. 109 (1958) 1268
 1958CO1E J.H. Coon, R.W. Davis, H.E. Felthuser and D.B. Nicodemus, Phys. Rev. 111 (1958) 250
 1958DA09 H. Daniel and U. Schmidt-Rohr, Nucl. Phys. 7 (1958) 516
 1958FI27 V.K. Fischer and G.E. Fischer, Bull. Amer. Phys. Soc. 3 (1958) 199, P4
 1958FR1C French, Univ. of Pittsburgh Tech. Rept. 9 (1958)
 1958HA1D Haig, Nucl. Phys. 7 (1958) 429

1958HE47 J.C. Hensel and W.C. Parkinson, Phys. Rev. 110 (1958) 128
1958HI68 R.W. Hill, Phys. Rev. 109 (1958) 2105
1958HI74 B. Hird, J.A. Cookson and M.S. Bokhari, Proc. Phys. Soc. (London) A72 (1958) 489
1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
1958JA06 N. Jarmie and R.C. Allen, Phys. Rev. 111 (1958) 1121
1958JU39 A.C. Juveland and W. Jentschke, Phys. Rev. 110 (1958) 456
1958JU42 M.K. Juric and S.D. Cirilov, Phys. Rev. 112 (1958) 1224
1958KA16 J. Kane, A.J. Elwyn, S. Ofer and D. Wilkinson, Bull. Amer. Phys. Soc. 3 (1958) 381, T6
1958KA31 R.W. Kavanagh and C.A. Barnes, Phys. Rev. 112 (1958) 503
1958KR65 W.E. Kreger and B.D. Kern, Bull. Amer. Phys. Soc. 3 (1958) 188, K13
1958LI42 C.S. Littlejohn and G.G. Morrison, Bull. Amer. Phys. Soc. 3 (1958) 227, XA3; Oral Report
1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1
1958MA1J Macklin and Gibbons, Bull. Amer. Phys. Soc. 3 (1958) 187
1958MA54 M.H. MacGregor, W.P. Ball and R. Booth, Phys. Rev. 111 (1958) 1155
1958MC63 M.T. McEllistrem, Phys. Rev. 111 (1958) 596
1958MO97 W.E. Moore, J.N. McGruer and A.I. Hamburger, Phys. Rev. Lett. 1 (1958) 29
1958MO99 C.D. Moak, A. Galonsky, R.L. Traugher and C.M. Jones, Phys. Rev. 110 (1958) 1369
1958NA09 M.P. Nakada, J.D. Anderson, C.C. Gardner and C. Wong, Phys. Rev. 110 (1958) 1439
1958SK1A Skyrme, Proc. Rehovoth Conf. (North-Holland Publishing Co., Amsterdam, 1958)
1958SW63 D.R. Sweetman, Bull. Amer. Phys. Soc. 3 (1958) 186, K1
1958TA03 Y.-K. Tai, G.P. Millburn, S.N. Kaplan and B.J. Moyer, Phys. Rev. 109 (1958) 2086
1958TA05 N.W. Tanner, Proc. Phys. Soc. (London) A72 (1958) 457
1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10
1958VA1D Vasilev, Komarov and Popova, JETP (Sov. Phys.) 6 (1958) 1016
1958VE15 J.F. Vervier and A. Martegani, Nucl. Phys. 6 (1958) 260
1958VE21 J.F. Vervier and A. Martegani, Phys. Rev. 109 (1958) 947
1958WA1D Warner and Alford, Bull. Amer. Phys. Soc. 3 (1958) 204
1958WE1E Wegner and Hall, Bull. Amer. Phys. Soc. 3 (1958) 338
1958WH34 A.B. Whitehead and J.S. Foster, Can. J. Phys. 36 (1958) 1276

- 1958WI01 H.B. Willard, J.K. Bair and H.O. Cohn, Bull. Amer. Phys. Soc. 3 (1958) 18, F2
1958WI36 J.E. Wills, Jr., J.K. Bair, H.O. Cohn and H.B. Willard, Phys. Rev. 109 (1958) 891
1959GI47 J.H. Gibbons and R.L. Macklin, Phys. Rev. 114 (1959) 571
1959KR1B W.E. Kreger and B.D. Kern, Phys. Rev. 113 (1959) 890
1959NE1A Neilson, Dawson and Johnson, Rev. Sci. Instrum. 30 (1959) 963