

Energy Levels of Light Nuclei $A = 15$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 13-15$ was published in *Nuclear Physics A152* (1970), p. 1. This version of $A = 15$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1969)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 15$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{15}\text{B}\$](#) , [\$^{15}\text{C}\$](#) , [\$^{15}\text{N}\$](#) , [\$^{15}\text{O}\$](#) , [\$^{15}\text{F}\$](#)

B. Tables of Recommended Level Energies:

[Table 15.1](#): Energy levels of ^{15}C

[Table 15.4](#): Energy levels of ^{15}N

[Table 15.18](#): Energy levels of ^{15}O

C. [References](#)

D. Figures: [\$^{15}\text{C}\$](#) , [\$^{15}\text{N}\$](#) , [\$^{15}\text{O}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

¹⁵B

¹⁵B has been identified in the 5.3 GeV proton bombardment of uranium. It is particle stable (1966PO09). See also (1960ZE03, 1961BA1C, 1966GA25).

¹⁵C

(Figs. 9 and 12)

GENERAL:

See (1960TA1C, 1962TA1E, 1964LI1B, 1964ST1B, 1967LO03). See also (1969AR13).

1. ¹⁵C(β^-)¹⁵N $Q_m = 9.773$

The half-life is 2.25 ± 0.05 sec (1956DO37), 2.49 ± 0.07 sec (1964NE09). The β -spectrum is complex. Transitions have been observed both to the ground state and to the upper of the 5.3 MeV levels of ¹⁵N: the latter transition is clearly allowed: see Table 15.2 (1959AL06, 1966AL12, 1969GA05). The ground state transition of ¹⁵N*(5.3) has $E_\gamma = 5.29903 \pm 0.00043$ MeV (1967CH19); J^π for this state is $\frac{1}{2}^+$ (see Table 15.4). Thus $J^\pi(^{15}\text{C}) = \frac{1}{2}^+$ or $\frac{3}{2}^+$ (1959AL06, 1964AL21: see also (1965WA03)). See also (1963KI1B, 1968ZH1B) and ¹⁵N.

2. ⁹Be(⁶He, α)¹¹Be $Q_m = 6.343$ $E_b = 19.076$

⁹Be was irradiated with 14 MeV neutrons and the decay of ¹¹Be was observed [⁹Be(n, α)⁶He, followed by the above reaction]: the cross section (which is determined for a large ⁶He energy spread) is 11 ± 10 mb (1967ST21).

3. ⁹Be(⁷Li, p)¹⁵C $Q_m = 9.095$

Observed proton groups are listed in Table 15.3 (1957MU99, 1964CA05). τ_m for ¹⁵C*(0.75) is (3.77 ± 0.11) nsec (1968ME08: delayed coincidence technique). Angular distributions of the protons to ¹⁵C*(0, 0.75) are reported at $E(^7\text{Li}) = 5.6, 5.8, 6.0, 6.2$ MeV by (1969SN02).

4. ¹⁴C(d, p)¹⁵C $Q_m = -1.006$

Table 15.1: Energy levels of ^{15}C

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
g.s.	$\frac{1}{2}^+; \frac{3}{2}$	$\tau_{1/2} = 2.33 \pm 0.08$ sec	β^-	1, 3, 4
0.747 ± 7	$\frac{5}{2}^+; \frac{3}{2}$	$\tau_m = 3.76 \pm 0.10$ nsec	γ	3, 4
(2.48 ± 50)				3
3.08 ± 40				3, 4
4.21 ± 30				3, 4
(4.60)				3
5.93 ± 30				3, 4
6.38 ± 30				3, 4
6.58 ± 40				3
(6.84)				3
(7.06)				3
7.32 ± 30				3, 4
(7.69)				3
(8.00)				3
8.12 ± 60				3
(8.47)				3

 Table 15.2: Beta decay of ^{15}C (1959AL06, 1966AL12, 1969GA05)

Decay to $^{15}\text{N}^*$ (MeV)	J^π	Branch (%)	$\log ft^a$ (exp)	$\log ft^b$ (theor)
g.s.	$\frac{1}{2}^-$	32 ± 2	5.96 ± 0.03	5.8
5.30	$\frac{1}{2}^+$	68 ± 2	4.04 ± 0.02	4.80
7.30	$\frac{1}{2}^+, \frac{3}{2}^+$	$(0.8 \pm 0.2) \times 10^{-2}$	6.82 ± 0.13	6.49
8.31	$\frac{1}{2}^+, \frac{3}{2}^+$	$(5.0 \pm 0.6) \times 10^{-2}$	5.06 ± 0.06	4.51
8.58	$\frac{1}{2}^+, \frac{3}{2}^+$	$\leq 2.8 \times 10^{-2}$	≥ 5.0	
9.05	$\frac{1}{2}^+, \frac{3}{2}^+$	$(3.5 \pm 0.5) \times 10^{-2}$	3.99 ± 0.07	

^a Using $\tau_m = 2.25 \pm 0.05$ sec.

^b See (1959AL97).

Table 15.3: Proton groups from ${}^9\text{Be}({}^7\text{Li}, \text{p}){}^{15}\text{C}$ and ${}^{14}\text{C}(\text{d}, \text{p}){}^{15}\text{C}$

E_x (MeV \pm keV)		E_x (MeV)
(1957MU99)	(1959MO1B)	(1964CA05) ^c
0	0	0
0.62 ± 60 ^a	0.75 ± 30	0.74
2.48 ± 50 ^b		
3.08 ± 40	3.09	3.08
4.26 ± 40	4.21	4.16
		4.60
5.93 ± 40	5.94	5.81
	6.38	6.39
6.58 ± 40		6.58
		6.84
		7.06
	7.32	7.31
		7.69
		8.00
8.16 ± 60		8.08
		8.47

^a (1957NO14) reports $E_x = 0.70 \pm 0.05$ MeV.

^b Not observed by (1964CA05).

^c ± 30 keV (private communication).

Identification of ${}^{15}\text{C}_{\text{g.s.}}$ with ${}^{15}\text{N}^*(11.62)$, $T = \frac{3}{2}$, $J^\pi = \frac{1}{2}^+$, is suggested by (1956BA16). At $E_d = 14.9$ MeV, proton groups are observed to the ground state of ${}^{15}\text{C}$ and to the levels at 0.75, 3.09, 4.21, 5.94, 6.38 and 7.32 MeV (± 30 keV) (1959MO1B). The angular distribution of ground state protons implies $l_n = 0$, $J^\pi = \frac{1}{2}^+$ (1959MO1B: see also (1961PU1B) and ref. ⁹ there, (1964AL21)); for the first excited state, $l_n = 2$. $\theta^2 = 0.093$ and 0.032 (1959MO1B), 0.16 and 0.063 (1966GL01) for the ground state and the first excited state, respectively. See also (1967NE06).

The 0.75 MeV level has a mean life of 3.73 ± 0.23 nsec (1962LO02); $E_\gamma = 750 \pm 7$ keV. The angular distribution of the γ -rays requires $J \geq \frac{5}{2}$. Since the $l_n = 2$ stripping requires $J^\pi = \frac{3}{2}^+$ or $\frac{5}{2}^+$, $J^\pi = \frac{5}{2}^+$ is established (1962CH14). The observed lifetime excludes $J > \frac{5}{2}$ (1962LO02). See also (1964NE09, 1965WA03, 1966AL12, 1967CH19).

5. $^{14}\text{C}(\text{n}, \gamma)^{15}\text{C}$ $Q_{\text{m}} = 1.218$

The capture cross section is $< 1 \mu\text{b}$ (1951YA1A).

6. $^{18}\text{O}(\text{n}, \alpha)^{15}\text{C}$ $Q_{\text{m}} = -5.009$

See (1966BA1F).

The following reactions leading to ^{15}C have not been reported:

$^{13}\text{C}(\text{t}, \text{p})^{15}\text{C}$ $Q_{\text{m}} = 0.912$

$^{14}\text{C}(\text{t}, \text{d})^{15}\text{C}$ $Q_{\text{m}} = -5.039$

$^{14}\text{C}(\alpha, ^3\text{He})^{15}\text{C}$ $Q_{\text{m}} = -19.360$

$^{15}\text{N}(\text{n}, \text{p})^{15}\text{C}$ $Q_{\text{m}} = -8.990$

^{15}N
(Figs. 10 and 12)

GENERAL:

Model calculations: (1957HA1E, 1959BR1E, 1959FE1B, 1960TA1C, 1961BA1E, 1963BU1C, 1963KU1B, 1964MA1G, 1965CO25, 1965FA1B, 1965GR1H, 1965GU1A, 1965ZA1B, 1966EL08, 1966SO05, 1967CO32, 1967EL03, 1967PA05, 1968EL1A, 1968HO1H, 1968MA2B, 1968SH08, 1968WA04, 1968ZH05, 1969CH1R, 1969EL1B).

General calculations and reviews: (1964EV1A, 1965BE1B, 1966OL1C, 1966WI1E, 1967FA1A, 1967LO03, 1968BI1C, 1968ZH1B, 1969HA1G, 1969IW1B).

Electromagnetic transitions: (1965RO1N, 1966HA31, 1966PO11, 1966RO1P, 1966RO1U, 1966WA1E, 1967KU1E, 1967PO1J, 1967WA1C, 1968BI1C, 1968SH08, 1968YA1E, 1968ZH06, 1968ZH1B, 1969KH1C, 1969ZH1A).

Meson interactions: (1969KA1R).

Other: (1961BA05, 1964VA1D).

Ground state: $\mu = -0.28309$ nm (1962BA63, 1964LI14, 1967CO1D).

See also (1961BR13, 1964ST1B, 1965IC1A, 1965MA1T, 1966MA1V, 1966WI1E, 1967SH14, 1968PE16, 1968RO1E, 1969CH1R, 1969FU11).

1. (a) $^9\text{Be}(^6\text{Li}, \text{p})^{14}\text{B}$	$Q_m = 15.130$	$E_b = 25.339$
(b) $^9\text{Be}(^6\text{Li}, \alpha)^{11}\text{B}$	$Q_m = 14.347$	
(c) $^9\text{Be}(^6\text{Li}, 2\text{n})^{13}\text{N}$	$Q_m = 3.951$	
(d) $^9\text{Be}(^6\text{Li}, ^8\text{Be})^7\text{Li}$	$Q_m = 5.587$	
(e) $^9\text{Be}(^6\text{Li}, ^5\text{He})^{10}\text{B}$	$Q_m = 1.933$	

The yield of p_0 and p_1 (reaction (a)) for $E(^6\text{Li}) = 3.84$ to 6.40 MeV shows some broad structure: analysis in terms of Ericson fluctuation theory gives a value of ≈ 0.4 MeV for the average level width at $E_x = 28$ MeV in ^{15}N (1967SE08). The excitation functions for α_0 , α_1 and α_{2+3} (reaction (b)) [$E(^6\text{Li}) = 2$ to 4 MeV: (1961LE01)] and the yield of ^{13}N (reaction (c)) [$E(^6\text{Li}) = 1.5$ to 3.5 MeV: (1961NO05)] show a smooth increase in the cross section with energy. For reactions (d) and (e) see (1962MC12). See also (1963BA1Q).

2. $^9\text{Be}(^7\text{Li}, \text{n})^{15}\text{N}$	$Q_m = 18.086$
--	----------------

At $E(^7\text{Li}) = 2.9$ MeV, γ -rays are observed which are assigned to the ground state decay of $^{15}\text{N}^*(9.05, 9.83, 10.80)$ (1964CA18). See also (1957NO17).

Table 15.4: Energy levels of ^{15}N ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
0	$\frac{1}{2}^-$		stable	2, 3, 4, 8, 9, 10, 12, 13, 14, 21, 22, 23, 24, 25, 26, 29, 30, 32, 39, 40, 41, 42, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66
5.27055 ± 0.25	$\frac{5}{2}^+$	$\tau_m = 2.9 \pm 0.5$	γ	3, 4, 8, 12, 13, 21, 22, 23, 26, 29, 30, 32, 39, 42, 52, 53, 55, 58, 59, 62
5.29921 ± 0.25	$\frac{1}{2}^+$	< 0.01	γ	3, 4, 8, 12, 13, 21, 22, 23, 26, 29, 30, 32, 39, 46, 52, 53, 55, 58, 59
6.3235 ± 0.4	$\frac{3}{2}^-$	< 0.040	γ	3, 4, 8, 10, 12, 13, 21, 22, 26, 29, 30, 32, 39, 49, 52, 53, 55, 57, 58, 59, 62
7.1550 ± 0.4	$\frac{5}{2}^+$	< 0.018	γ	3, 4, 8, 12, 21, 22, 26, 29, 32, 39, 52, 53, 62
7.3010 ± 0.5	$\frac{3}{2}^+$	< 0.025	γ	3, 4, 8, 12, 13, 21, 26, 29, 32, 39, 46, 52, 53, 55, 62
7.566 ± 3	$\frac{7}{2}^+$	0.06 ± 0.02	γ	3, 4, 8, 12, 13, 21, 22, 29, 39, 52, 53, 62
8.3126 ± 0.7	$\frac{1}{2}^+$	< 0.010	γ	3, 4, 8, 21, 26, 29, 32, 39, 46, 52, 53, 55, 62
8.576 ± 2	$\frac{3}{2}^+$		γ	3, 4, 8, 13, 21, 22, 26, 32, 39, 52, 53, 62
9.053 ± 2	$\frac{1}{2}^+$		γ	2, 3, 4, 8, 21, 26, 29, 32, 39, 46, 55
9.1518 ± 0.5	$\frac{3}{2}^-$		γ	3, 4, 8, 13, 21, 22, 29, 32, 39, 52, 53
9.1549 ± 0.5	$(\frac{5}{2})$	< 0.010	γ	3, 4, 8, 13, 21, 22, 29, 32, 39, 52, 53
9.225 ± 3.5	$\frac{3}{2}$ or $\frac{1}{2}$ ^b	< 0.1	γ	21, 29, 39, 55
9.762 ± 3.5	$\frac{5}{2}^-$		γ	21, 39, 52, 53
9.829 ± 3	$\frac{7}{2}$	< 0.19	γ	2, 3, 4, 8, 21, 22, 29, 52, 53
9.929 ± 4	$(\frac{1}{2}, \frac{3}{2})^+$		γ	21, 39

Table 15.4: Energy levels of ^{15}N ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
10.070 \pm 3	$\frac{3}{2}^+$		γ	13, 21, 39, 52, 53
10.451 \pm 1	$\frac{3}{2} \rightarrow \frac{7}{2}$		γ, p	21, 22, 26, 39
10.536 \pm 1	$\frac{5}{2}^+$		γ, p	21, 26, 39
10.700 \pm 1	$\frac{3}{2}^+$		γ, p	21, 22, 26, 27, 39, 52
10.800 \pm 1	$\frac{3}{2}^-$		γ, p	2, 3, 4, 8, 13, 21, 26, 39
(10.94 \pm 30)				21
11.236 \pm 5	$\geq \frac{3}{2}$	$\Gamma = 3.3$	n	33
11.2943 \pm 1	$\frac{1}{2}^-$	7.9 \pm 0.3	γ, n, p	26, 27, 28, 33, 52
11.438 \pm 1	$\frac{1}{2}^+$	41.4 \pm 1.1	γ, n, p, α	5, 26, 27, 28, 33, 35
11.615 \pm 4	$\frac{1}{2}; \frac{3}{2}$	404.9 \pm 6.3	γ, n, p	26, 27, 28
11.764 \pm 3	$\frac{3}{2}^+$	40 \pm 3	n, p, α	5, 28, 33, 35
11.877 \pm 3	$\frac{3}{2}^-$	21 \pm 4	n, p, α	5, 28, 33, 35, 52
11.943 \pm 6	$(\frac{9}{2}^-)$	≤ 3	n	22, 33, 52
11.965 \pm 3	$\frac{1}{2}^-$	17 \pm 5	n, p, α	5, 28, 33, 35
12.097 \pm 4	$\frac{5}{2}^+$	14 \pm 5	γ, n, p, α	5, 6, 27, 28, 33, 35, 38
12.145 \pm 3	$\frac{3}{2}^-$	47 \pm 7	γ, n, p, α	5, 6, 27, 28, 33, 35, 39
12.326 \pm 4	$\frac{5}{2}^+$	22	n, p	22, 28, 33, 35
12.493 \pm 4	$\frac{5}{2}^+; \frac{1}{2}$	42	γ, n, p, α	5, 6, 28, 33, 35, 38, 52
12.52 \pm 10	$\frac{5}{2}^+; \frac{3}{2}$	80	p	27, 52
12.921 \pm 4	$\frac{3}{2}^-$	67 \pm 8	n, p, α	5, 6, 28, 33, 35, 38
12.93	$\frac{7}{2}^-$	30	p, α	6
13.028 \pm 20	$(\frac{11}{2}^-)$			22
13.14		< 3	n, p, α	5, 6, 38
13.19		6	n, p, α	5, 6, 28, 38
13.36	$\frac{3}{2}^-$	29 \pm 8	n, p, α	5, 6, 28, 38
13.40	$\frac{5}{2}^+$	≈ 60	n, p, α	6, 28, 35
(13.52)			n, p	28
13.60	$(\frac{5}{2}, \frac{7}{2})^-$	15 \pm 4	n, p, α	5, 6, 28, 33, 35, 38
13.67	$\frac{1}{2}^+$	≈ 80	n, p, α	6, 28
13.71		≈ 40	n, p, α	5, 35, 38
13.75			n, α	5
13.84		≈ 40	n, p, α	28, 35, 38
13.89			n, α	5

Table 15.4: Energy levels of ^{15}N ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
14.03	$\frac{3}{2}^{(+)}$		p, α	6
14.06			n, α	5
14.11		105	n, p, α	5, 28, 35, 38, 52
14.17		30 ± 5	n, p, α	5, 28, 35, 38
14.18			n, α	5
14.23			n, α	5
14.4		≈ 2000	n, p, α	33, 35, 38
14.46		≈ 180	p, α	6
14.51		130	n, p	28
14.64		50 ± 3	n, p, α	5, 35, 38
14.7		≈ 280	n, p, α	35, 38
14.81		99	n, p	28
14.90		37	n, p, α	5, 28, 38
15.00			n, p, α	5, 28
15.11			n, p, α	5, 28, 38, 52
15.29			n, α	5
15.37			n, t, α	5, 11
15.52			n, α	38
15.61			n, α	5
15.74			p, t, α	11
15.83		< 3	n, p, t, α	5, 6, 11
15.89 ± 20		< 3	n, t, α	5, 11
15.96 ± 20			n, t, α	5, 11
15.99			n, α	5
16.03			n, p, t, α	5, 6, 11, 38
16.08			n, p, t, α	5, 6, 11, 38
16.17 ± 40		n, p, t, α	5, 6, 11	
16.29		n, p, t, α	5, 6, 11	
16.33 ± 20		n, p, t, α	5, 6, 11, 38	
16.43 ± 20		n, p, t, α	5, 6, 11, 38	
16.49 ± 30		n, p, d, t, α	5, 6, 11, 16	
16.59 ± 25	70	n, p, t, α	5, 6, 11, 38	
16.67 ± 30	100	n, p, d, t, α	5, 6, 11, 15, 16, 33, 38	

Table 15.4: Energy levels of ^{15}N ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or Γ (keV)	Decay	Reactions
16.74			n, p, α	5, 6
16.76 \pm 30			n, p, d, t, α	5, 6, 11, 15, 16
16.85 \pm 30			t, α	11
16.90		\approx 350	n, p, d, t, α	5, 6, 11, 15, 33, 38
16.98			n, p, α	5, 6, 33, 38
17.05			p, t	11
17.10		broad	d, t, α	11, 19
17.16 \pm 50			n, p, t, α	5, 6, 11
17.19			n, p, α	5, 6, 18
17.30		190	n, p, α	5, 6, 18, 38
17.36		350	n, p, d, t, α	5, 6, 11, 16, 18, 19
17.50			n, p, α	5, 6, 38
17.56			n, p, α	5, 6
17.81 \pm 40		\approx 170	n, p, d, t, α	5, 6, 18
17.70 \pm 50		\approx 500	n, d, α	15, 19
17.72 \pm 10		48 \pm 9	(p), d, t, α	16, 18, 19
17.81		170	n, α	33, 38
17.95			n, p, α	5, 6
18.07 \pm 10		19 \pm 4	(n), d, α	15, 19
18.09 \pm 20		\approx 45	(n), p, d, t	15, 16, 18
18.22		160	n, α	38
18.28 \pm 30		230 \pm 60	n, p, d, α	15, 16, 19, 38
19.16 \pm 30		\approx 130	n, d	15
19.5			γ , p	48
20.4			γ , p	48
22.7			γ , p	48
24.5			γ , p	48

^a See also Tables 15.7 and 15.10.

^b See (1967PH03).

3. $^{10}\text{B}(^6\text{Li}, \text{p})^{15}\text{N}$

$$Q_m = 18.751$$

Table 15.5: Resonances in $^{11}\text{B} + \alpha$

E_α (MeV)	Γ_{lab} (keV)	Particle out	E_x (MeV)	J^π	Refs. ^a
0.60		n	11.43		(1954BE08) ^b
1.03		n	11.75		(1954BE08)
1.18		n	11.86		(1954BE08)
1.30		n	11.95		(1954BE08)
1.51		n, p	12.10		(1955SH46)
1.58		n, p	12.15		(1955SH46)
2.06	66	n_0, p_0	12.50	$\frac{5}{2}^+$	(1955SH46, 1956BO61, 1958HA1B, 1959LE28)
2.63	80	n_0, p_0	12.92	$\frac{3}{2}^-$	(1956BO61, 1958HA1B, 1959LE28, 1963MA28, 1966MA04)
2.64	40	p_0	12.93	$\frac{7}{2}^-$	(1959LE28, 1963MA28)
2.94	< 6	n_0, p_0	13.15		(1956BO61, 1958HA1B, 1959LE28)
2.99	8	n_0, p_0	13.18		(1956BO61, 1958HA1B, 1959LE28, 1966MA04)
3.23	29	n_0, p	13.36	$\frac{3}{2}^-$	(1956BO61, 1958HA1B, 1959LE28, 1963MA28, 1966MA04)
3.31	40	p	13.42	$\frac{5}{2}^+$	(1959LE28, 1963MA28)
3.56	20	n_0, p	13.61	$(\frac{5}{2}, \frac{7}{2})^-$	(1956BO61, 1958HA1B, 1959LE28, 1963MA28, 1966MA04)
3.64	≈ 110	p_0	13.66		(1963MA28)
3.71	40	n_0	13.71		(1956BO61, 1958HA1B, 1963MA28, 1966MA04)
3.76		n_0	13.75		(1966MA04)
3.95		n_0	13.89		(1963MA28, 1966MA04)
4.15		p_0	14.03		(1963MA28)
4.19		n_1	14.06		(1963MA28, 1966MA04)
4.24		n_0	14.10		(1963MA28, 1966MA04)
4.31	35	n_1	14.15		(1956BO61, 1958HA1B, 1963MA28, 1966MA04)
4.35		n_0	14.18		(1963MA28, 1966MA04)
4.41		n_0	14.23		(1963MA28, 1966MA04)
4.56		p_0	(14.33)		(1963MA28)

Table 15.5: Energy levels of ^{15}N ^a (continued)

E_α (MeV)	Γ_{lab} (keV)	Particle out	E_x (MeV)	J^π	Refs. ^a
4.73	≈ 250	p ₀	14.46		(1963MA28)
4.96	72	n ₀	14.63		(1956BO61, 1958HA1B)
5.34		n ₀	14.90		(1958HA1B)
5.49		n ₀	15.01		(1958HA1B)
5.58		n ₀	15.08		(1958HA1B)
5.86		n ₀	15.29		(1958HA1B)
5.98		n ₀ , n ₂	15.37		(1958HA1B)
6.30		n ₀ , (n ₂)	15.61		(1958HA1B)
6.60		n, p	15.83		(1963ED01) ^c
6.72		n ₀ , (n ₂)	15.92		(1958HA1B)
6.74		n ₀ , n ₂	15.93		(1958HA1B)
6.82		n ₀ , n ₂	15.99		(1958HA1B)
6.89		n, p	16.04		(1958HA1B, 1963ED01)
6.94		n, p	16.08		(1963ED01)
7.08		n, p	16.18		(1963ED01)
7.26		n, p	16.31		(1963ED01)
7.31		n, p	16.35		(1963ED01)
7.41		n, p	16.42		(1963ED01)
7.56		n, p	16.53		(1963ED01)
7.65		n, p	16.60		(1963ED01)
7.77		n, p	16.69		(1963ED01)
7.84		n, p	16.74		(1963ED01)
7.86		n, p	16.75		(1963ED01)
8.07		n, p	16.91		(1963ED01)
8.17		n, p	16.98		(1963ED01)

Table 15.5: Energy levels of ^{15}N ^a (continued)

E_α (MeV)	Γ_{lab} (keV)	Particle out	E_x (MeV)	J^π	Refs. ^a
8.44		n, p	17.18		(1963ED01)
8.46		n, p	17.19		(1963ED01)
8.59		n, p	17.29		(1963ED01)
8.68		n, p	17.35		(1963ED01)
8.88		n, p	17.50		(1963ED01)
8.96		n, p	17.56		(1963ED01)
8.99		n, p	17.58		(1963ED01)
9.49		n, p	17.95		(1963ED01)

^a It should be noted that (1954BE08), (1958HA1B) and (1963ED01) are unpublished. They are quoted here only because of the dearth of published information on these resonances.

^b And private communication.

^c Resonant energies listed by (1963ED01) are ± 40 keV.

At $E(^6\text{Li}) = 4.9$ MeV, thirty proton groups are observed corresponding to ^{15}N states with $E_x < 16.8$ MeV. Angular distributions have been measured for the proton groups corresponding to $^{15}\text{N}^*(5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.58, 9.05 + 9.15)$ (1966MC05). The ground-state γ -decay of $^{15}\text{N}^*(9.83, 10.80)$ is reported by (1964CA18). See also (1963MO1B).

$$4. \text{}^{10}\text{B}(^7\text{Li}, \text{d})^{15}\text{N} \quad Q_m = 13.723$$

At $E(^7\text{Li}) = 5.2$ MeV, thirty deuteron groups are observed corresponding to ^{15}N states with $E_x < 15.1$ MeV. Angular distributions have been measured for the deuteron groups corresponding to $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.58, 9.05 + 9.15, 9.83)$ (1966MC05). See also (1963MO1B). The ground state γ -decay of $^{15}\text{N}^*(8.31, 9.05, 9.83, 10.80)$ is reported by (1964CA18).

$$5. \text{}^{11}\text{B}(\alpha, \text{n})^{14}\text{N} \quad Q_m = 0.157 \quad E_b = 10.992$$

Reported resonances are displayed in Table 15.5 (1954BE08, 1955SH46, 1956BO61, 1958HA1B, 1963ED01, 1963MA28, 1966MA04). See also (1962GO1J, 1963GO1J, 1965TS1A) and (1966WE1B). See also ^{14}N .

$$6. \text{}^{11}\text{B}(\alpha, \text{p})^{14}\text{C} \quad Q_m = 0.784 \quad E_b = 10.992$$

Reported resonances are listed in Table 15.5 (1955SH46, 1959LE28, 1963ED01, 1963MA28). Partial widths for several resonances are listed by (1959LE28, 1963MA28). See also ^{14}C .

$$7. \text{}^{11}\text{B}(\alpha, \text{d})^{13}\text{C} \quad Q_m = -5.168 \quad E_b = 10.992$$

The yield of ground state deuterons has been measured for $E_\alpha = 17$ to 22 MeV by (1967AL16). See also ^{13}C .

$$8. \text{(a) } ^{11}\text{B}(^6\text{Li}, \text{d})^{15}\text{N} \quad Q_m = 9.520$$

$$\text{(b) } ^{11}\text{B}(^7\text{Li}, \text{t})^{15}\text{N} \quad Q_m = 8.525$$

At $E(^6\text{Li}) = 4.72$ MeV and at $E(^7\text{Li}) = 5.00$ MeV, angular distributions are reported for the deuterons and the tritons corresponding to $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.58, 9.05 + 9.15)$ (1966MC05). See also (1963MO1B). Gamma rays are observed in reaction (a) which are assigned to the ground state transitions of $^{15}\text{N}^*(9.15, 9.83, 10.80)$ (1964CA18). See also (1963HO1E) and (1964HA09).

9. $^{11}\text{B}(^9\text{Be}, ^5\text{He})^{15}\text{N}$ $Q_m = 8.464$

See (1963HO1E).

10. $^{11}\text{B}(^{16}\text{O}, ^{12}\text{C})^{15}\text{N}$ $Q_m = 3.831$

At $E(^{16}\text{O}) = 27, 30, 32.5$ and 35 MeV, angular distributions corresponding to transitions to $^{15}\text{N}^*(0, 6.32)$ have been measured by (1965BO14, 1966BO22): the ground state angular distributions show strong diffraction structure. See also (1969GO1R, 1969BR1D, 1969KA1G).

11. (a) $^{12}\text{C}(t, n)^{14}\text{N}$ $Q_m = 4.015$ $E_b = 14.850$
 (b) $^{12}\text{C}(t, p)^{14}\text{C}$ $Q_m = 4.641$
 (c) $^{12}\text{C}(t, t)^{12}\text{C}$
 (d) $^{12}\text{C}(t, \alpha)^{11}\text{B}$ $Q_m = 3.858$

Reported resonances are listed in Table 15.6 (1961VA13, 1962GU01, 1962KU09, 1963NI04, 1965SE05, 1969ET01). The triton yield has been measured for $E_t = 9$ to 13 MeV by (1965GL04). See also (1962NE1D, 1964GR1H, 1967CH35), ^{11}B and ^{12}C in (1968AJ02), and ^{14}C and ^{14}N .

12. $^{12}\text{C}(\alpha, p)^{15}\text{N}$ $Q_m = -4.965$

Angular distributions of the protons corresponding to the ground state transition have been measured at $E_\alpha = 13.4$ to 16.0 MeV (1967IV1B), 16.1 to 19.0 MeV (1960PR13), 19.7 to 22.1 MeV (1963YA1C), 20.6 to 22.2 MeV (1961KO04), 25 to 39 MeV (1959NO38) and at 42 MeV (1962LI07). Angular distributions are also reported for the groups to $^{15}\text{N}^*(5.27+5.30, 6.32, 7.16+7.30+7.57)$ (1959NO38: 34.6 MeV) and $^{15}\text{N}^*(6.32)$ (1962LI07: 42 MeV). See also (1962EI03, 1969GL1D), (1961KR1A, 1962HO1D, 1962TE1B, 1963DA1B, 1964DA1D, 1964KE1C, 1965NE1D, 1966HI1C, 1967RO1K; theor.) and (1959AJ76).

13. $^{12}\text{C}(^7\text{Li}, \alpha)^{15}\text{N}$ $Q_m = 12.382$

Angular distributions have been measured at $E(^7\text{Li}) = 3.2$ to 4.0 MeV (1962HO06; $\alpha_0, \alpha_{1+2}, \alpha_3$) and 30.3 MeV (1969GL07; $\alpha_0, \alpha_{1+2}, \alpha_3$ and the α 's corresponding to $^{15}\text{N}^*(7.35, 7.6, 8.6, 9.2, 10.1, 10.9, 12.8, 13.5, 15.2)$. (1969BA2U) report the excitation of $^{15}\text{N}^*(12.5, 13.3, 14.9, 15.6, 16.2, 16.8, 18.9, 19.8)$ and (1969GL07) also report α groups corresponding to $^{15}\text{N}^*(11.8, 16.5)$

Table 15.6: Resonances in $^{12}\text{C} + t$

E_t (MeV \pm keV)	E_x (MeV)	Particles out	Refs.
0.66	15.38	α_0	(1963NI04, 1969ET01)
1.11	15.74	p_0, t_0, α_1	(1962GU01, 1962KU09, 1963NI04, 1969ET01)
1.21	15.82	t_0	(1969ET01)
1.30 ± 20	15.89	n, α_0	(1961VA13, 1969ET01)
1.39 ± 20	15.96	n, t_0, α_0	(1961VA13, 1969ET01)
1.46	16.02	p_0	(1969ET01)
1.54	16.08	n, α_0, α_1	(1961VA13, 1969ET01)
1.65 ± 40	16.17	n, α_0	(1961VA13, 1965SE05, 1969ET01)
1.78	16.27	α_0	(1969ET01)
1.85 ± 20	16.33	$n, p_0, \alpha_0, \alpha_1$	(1961VA13, 1965SE05, 1969ET01)
1.98 ± 20	16.43	n, p_0	(1961VA13, 1969ET01)
2.05 ± 30	16.49	p_0, t_0, α_0	(1965SE05, 1969ET01)
2.18 ± 25	16.59	$n, p_0, t_0, \alpha_0, \alpha_1$	(1961VA13, 1969ET01)
2.28 ± 30	16.67	$n, p_0, \alpha_0, \alpha_1$	(1961VA13, 1965SE05, 1969ET01)
2.39 ± 30	16.76	$n, t_0, \alpha_0, \alpha_1$	(1961VA13, 1965SE05, 1969ET01)
2.50 ± 30	16.85	α_0, α_1	(1965SE05, 1969ET01)
2.60	16.93	α_0	(1969ET01)
2.75	17.05	p_0	(1969ET01)
2.82	17.10	t_0, α_0, α_1	(1969ET01)
2.89 ± 50	17.16	α_0	(1965SE05, 1969ET01)
3.14	17.36	α_1	(1969ET01)

and, possibly $^{15}\text{N}^*(17.1, 18.3)$. The mean lifetime of $^{15}\text{N}^*(9.83) < 1.9 \times 10^{-13}$ sec (see Table 15.7): $E_\gamma = 4562.6 \pm 4.0$ keV ($^{15}\text{N}^*(9.83) \rightarrow 5.27$) (1969TH01). See also (1960SH05) and (1969GI1B, 1969RO1G).

14. (a) $^{12}\text{C}(^{14}\text{N}, ^{11}\text{C})^{15}\text{N}$ $Q_m = -7.885$
 (b) $^{12}\text{C}(^{19}\text{F}, ^{16}\text{O})^{15}\text{N}$ $Q_m = 3.150$

For reaction (a) see (1969BR1D); for reaction (b) see (1969RO1G).

15. $^{13}\text{C}(d, n)^{14}\text{N}$ $Q_m = 5.325$ $E_b = 16.160$

Table 15.7: Lifetimes of some ^{15}N states

E_x (MeV)	τ_m (psec)	Reaction	Refs.
5.27	> 1	$^{16}\text{O}(t, \alpha)$	(1965AL19)
	≈ 1	$^9\text{Be}(^{14}\text{N}, ^8\text{Be})$	(1969NI09)
	2.9 ± 0.5	$^{14}\text{N}(d, p)$	(1967BI11)
5.30	< 0.01	$^{14}\text{C}(p, \gamma)$	(1968COZV)
	$(4.3 \pm 1.8) \times 10^{-2}$	$^{14}\text{N}(d, p)$	(1965AL19)
	$(2.2 \pm 0.7) \times 10^{-2}$	$^{14}\text{N}(n, \gamma)$	(1969WE07)
6.32	< 0.045	$^{14}\text{N}(d, p)$	(1968GI11)
	< 0.040	$^{14}\text{N}(n, \gamma)$	(1969WE07)
	< 0.010	$^{14}\text{C}(p, \gamma)$	(1968COZV)
7.16	< 0.018	$^{14}\text{N}(d, p)$	(1968GI11)
	0.115 ± 0.025	$^{13}\text{C}(^3\text{He}, p)$	(1966LI07)
	0.010 ± 0.002	$^{14}\text{C}(p, \gamma)$	(1968COZV)
7.30	< 0.025	$^{14}\text{N}(d, p)$	(1968GI11)
	< 0.030	$^{14}\text{N}(n, \gamma)$	(1969WE07)
	< 0.010	$^{14}\text{C}(p, \gamma)$	(1968COZV)
7.57	0.15 ± 0.05	$^{13}\text{C}(^3\text{He}, p)$	(1966LI07)
	0.06 ± 0.02	$^{14}\text{N}(d, p)$	(1968GI11)
8.31	< 0.021	$^{14}\text{N}(d, p)$	(1968GI11)
	< 0.020	$^{14}\text{N}(n, \gamma)$	(1969WE07)
	< 0.010	$^{14}\text{C}(p, \gamma)$	(1968COZV)
9.155	< 0.010	$^{14}\text{C}(p, \gamma)$	(1968COZV)
	< 0.010	$^{14}\text{N}(n, \gamma)$	(1969WE07)
9.23	< 0.1	$^{16}\text{O}(\gamma, p)$	(1969MU07)
9.83	< 0.19	$^{12}\text{C}(^7\text{Li}, \alpha)$	(1969TH01)

Table 15.8: Resonances in $^{13}\text{C} + \text{d}$

E_d (MeV)	Emitted particles	Γ_{lab} (keV)	$^{15}\text{N}^*$ (MeV)	Refs.
0.37	p		16.47	(1956VA17)
0.64	n, p_0	≈ 100	16.72	(1950CU13, 1950RI57, 1953KO42, 1956MA46, 1956VA17)
0.85	n	≈ 400	16.90	(1950RI57)
1.10	α_0	broad	17.11	(1956MA35, 1966KL06)
1.24 ± 0.04	t_0	≈ 200	17.24	(1956MA35)
1.40 ± 0.04	p_0, t_0, α_0	≈ 400	17.37	(1956MA35, 1956MA46)
1.55 ^a	n	≈ 100		(1941BE1A, 1950RI57, 1966KL06)
1.64 ± 0.04	t_0	≈ 200	17.58	(1956MA35)
1.78 ± 0.05	n, α_0	≈ 600	17.70	(1950RI57, 1955MA76, 1956MA35)
1.80 ± 0.01	(p_0), t_0, α_1	55 ± 10	17.72	(1956MA35, 1956MA46)
2.20 ± 0.01	(n), α_0, α_1	22 ± 4	18.07	(1956MA35, 1963AL21)
2.23 ± 0.02	(n), p_0, t	≈ 50	18.09	(1956MA35, 1956MA46, 1963AL21)
2.45 ± 0.03	n, p_0, α_0	270 ± 70	18.28	(1955MA76, 1956MA35, 1956MA46)
3.46 ± 0.03	n	≈ 150	19.16	(1955MA76, 1963DE19)

^a Possibly to be identified with 1.40 MeV resonance (1956MA35).

Observed resonances are displayed in Table 15.8 (1950RI57, 1955MA76, 1963AL21, 1963DE19). Excitation functions have recently been measured for $E_d = 0.8$ to 1.3 MeV (1961JA09; n_0, n_1, n_2, n_3, n_5), 2.0 to 3.2 MeV (1963AL21: 7.03 MeV γ -ray) and 3.2 to 4.0 MeV (1963DE19; n_0). See also (1960VA11). Polarization measurements are reported for $E_d = 2.5$ to 4.0 MeV (1967ME1N; n_0, n_1, n_2) and 2.8 MeV (1965GA1G; n_3, n_4, n_5). See also ^{14}N .

16. $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$

$$Q_m = 5.952$$

$$E_b = 16.160$$

Observed resonances are displayed in Table 15.8 (1941BE1A, 1950CU13, 1953KO42, 1956MA46). Excitation functions have recently been measured for $E_d = 1$ to 3.4 MeV (1968LI1L; p_0, p_1), 3.0 to 4.0 MeV (1965LA09: $^{14}\text{C}^*(6.09, 6.73, 7.34)$), 3.1 to 4.1 MeV (1963DE19; p_0), and 4.1 to 6.2 MeV (1968CO04; p_1). See also (1963AL21). See also ^{14}C .

17. $^{13}\text{C}(\text{d}, \text{d})^{13}\text{C}$

$$E_b = 16.160$$

Excitation functions for elastically scattered deuterons have been measured for $E_d = 1$ to 3.4 MeV (1968LI1L) and 4.5 to 5.7 MeV (1968CO04).

$$18. \text{}^{13}\text{C}(\text{d}, \text{t})\text{}^{12}\text{C} \qquad Q_m = 1.311 \qquad E_b = 16.160$$

Observed resonances are listed in Table 15.8 (1956MA35). (1968LI1L) report measurement of the t_0 excitation function for $E_d = 1$ to 3.4 MeV. A polarization study has been made at $E_d = 12.3$ MeV (1969DE1H; t_0, t_1). See also ^{12}C .

$$19. \text{}^{13}\text{C}(\text{d}, \alpha)\text{}^{11}\text{B} \qquad Q_m = 5.168 \qquad E_b = 16.160$$

Observed resonances are listed in Table 15.8 (1956MA35). See also (1966KL06, 1966KL1F, 1968CO04, 1968LI1L).

$$20. \text{}^{13}\text{C}(\text{t}, \text{n})\text{}^{15}\text{N} \qquad Q_m = 9.903$$

Not reported.

$$21. \text{}^{13}\text{C}(\text{}^3\text{He}, \text{p})\text{}^{15}\text{N} \qquad Q_m = 10.667$$

Observed proton groups and γ -rays corresponding to ^{15}N states are listed in Table 15.9 (1959YO25, 1966GA08, 1966WA08, 1967PH03). Gamma-ray branching ratios obtained by (1965WA16, 1966PE04, 1966WA08) are displayed in Table 15.10 which also shows J^π values obtained from angular correlation measurements. The two states at $E_x = 9.16$ MeV [see reactions 32 and 39] are separated by 2.5 ± 0.5 keV (1968ST10). See also (1959BR79). The τ_m for $^{15}\text{N}^*(7.16, 7.57)$ are 0.115 ± 0.025 psec and 0.15 ± 0.05 psec, respectively: see Table 15.7 (1966LI07). Angular distributions of the ground state protons have been measured for $E(^3\text{He}) = 8.7$ to 10.93 MeV (1962AL01). See also (1959AJ76).

See also (1967YO1C, 1968WE15), (1963CL1A, 1966PR1B) and ^{16}O in (1971AJ02).

$$22. \text{}^{13}\text{C}(\alpha, \text{d})\text{}^{15}\text{N} \qquad Q_m = -7.687$$

This reaction has been studied at $E_\alpha = 40.1$ MeV: see Table 15.9 (1969LU07).

Table 15.9: Energy levels in ^{15}N from $^{13}\text{C}(^3\text{He}, \text{p})^{15}\text{N}$ and $^{13}\text{C}(\alpha, \text{d})^{15}\text{N}$

E_x (MeV \pm keV) ^a				
(1959YO25)	(1966GA08)	(1966WA08) ^b	(1967PH03)	(1969LU07) ^c
5.283 \pm 12				5.266 \pm 20
6.333 \pm 12				6.336 \pm 30
7.169 \pm 12				7.170 \pm 20
7.310 \pm 12				
7.577 \pm 13				7.581 \pm 20
8.318 \pm 12	8.323 \pm 6	8.312		
8.581 \pm 14	8.581 \pm 5	8.570		8.587 \pm 20
9.061 \pm 14	9.056 \pm 5	9.052	9.054 \pm 4	
9.164 \pm 14	9.159 \pm 5			9.169 \pm 30
	9.225 \pm 6		9.225 \pm 3	
	9.760 \pm 5			
	9.827 \pm 6		9.829 \pm 4	9.808 \pm 20
	9.929 \pm 8			
	10.064 \pm 7	10.074	10.072 \pm 4	
	10.454 \pm 6	10.452		10.451 \pm 20
	10.536 \pm 7			
	10.704 \pm 6			10.698 \pm 20
	10.805 \pm 7	10.800		
		(10.94 \pm 30)		d

^a See also (1962SH21, 1966GO1J, 1966PH1B).

^b E_γ , except for $E_x = 10.94$ MeV; errors for E_γ are nominal.

^c $^{13}\text{C}(\alpha, \text{d})^{15}\text{N}$: $E_\alpha = 40.1$ MeV.

^d (1969LU07) also reports levels at $E_x = 11.950 \pm 0.020$ ($J^\pi = (\frac{9}{2}^-)$), 12.318 ± 0.030 and 13.028 ± 0.020 MeV ($J^\pi = (\frac{11}{2}^-)$).

Table 15.10: Radiative decays in ^{15}N

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
5.27	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	100	
5.30	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	100	
6.32	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100	(1965WA16, 1966PE04, 1969SI04)
		5.27	$\frac{5}{2}^+$	< 1	(1965WA16)
				< 3	(1966PE04)
		5.30	$\frac{1}{2}^+$	< 3	(1966PE04)
				< 1	(1965WA16)
7.16	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	< 5	(1965WA16)
				< 3	(1969SI04)
				< 12	(1968GI11)
				< 4	(1966HA30)
		5.27	$\frac{5}{2}^+$	100	(1965WA16, 1966PE04, 1968GI11)
				> 97	(1969SI04)
				< 95	(1967TH05)
		5.30	$\frac{1}{2}^+$	< 4	(1966AL18)
				< 4	(1968GI11)
				< 5	(1966PE04)
7.30	$\frac{3}{2}^+$	6.32	$\frac{3}{2}^-$	< 0.5	(1965WA16)
		0	$\frac{1}{2}^-$	100	(1966PE04, 1969SI04)
				98 ± 1	(1965WA16, 1968GI11)
		5.27	$\frac{5}{2}^+$	2 ± 1	(1965WA16)
				< 1.5	(1968GI11)
		5.30	$\frac{1}{2}^+$	2 ± 1	(1968GI11)
		6.32	$\frac{3}{2}^-$	< 3	(1966PE04)
7.57	$\frac{7}{2}^+$	0	$\frac{1}{2}^-$	< 0.25	(1965WA16)
				< 2	(1965WA16)
				< 4	(1966HA30)
				< 3	(1966PE04)
		5.27	$\frac{5}{2}^+$	100	(1965WA16, 1966PE04, 1968GI11)
		5.30	$\frac{1}{2}^+$	< 5	(1965WA16)
				< 4	(1966AL18)
		< 6	(1968GI11)		

Table 15.10: Radiative decays in ^{15}N (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
8.31	$\frac{1}{2}^+$	6.32	$\frac{3}{2}^-$	< 0.6	(1965WA16)
		0	$\frac{1}{2}^-$	80 ± 3	(1965WA16)
				77 ± 3	(1966WA08)
				70 ± 4	(1966PE04)
				79.1 ± 1.9	(1967PH03)
		5.27	$\frac{5}{2}^+$	< 3	(1965WA16)
		5.30	$\frac{1}{2}^+$	10 ± 2	(1965WA16)
		5.27 + 5.30		12 ± 2	(1966WA08)
				12 ± 3	(1966PE04)
				10.9 ± 1.3	(1967PH03)
		6.32	$\frac{3}{2}^-$	7.8 ± 2	(1965WA16)
				12 ± 3	(1966PE04)
				4.4 ± 1.0	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 1	(1965WA16)
		≤ 6	(1966PE04)		
		1.2 ± 0.6	(1967PH03)		
7.30	$\frac{3}{2}^+$	2.2 ± 0.4	(1965WA16)		
		4.4 ± 0.7	(1967PH03)		
8.58	$\frac{3}{2}^+$	0 ^c	$\frac{1}{2}^-$	34 ± 4	(1965WA16)
				32 ± 3	(1966WA08)
				27 ± 4	(1966PE04)
				33.4 ± 2.0	(1967PH03)
		5.27	$\frac{5}{2}^+$	63 ± 4	(1965WA16)
				65 ± 3	(1966WA08)
		5.30	$\frac{1}{2}^+$	< 12	(1965WA16)
		5.27 + 5.30		66 ± 4	(1966PE04)
				61.6 ± 2.0	(1967PH03)
		6.32	$\frac{3}{2}^-$	3 ± 1	(1965WA16)
				≤ 7	(1966PE04)
				1.4 ± 0.6	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 5	(1965WA16)
				< 4	(1966WA08)
		3.6 ± 0.5	(1967PH03)		

Table 15.10: Radiative decays in ^{15}N (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
9.05	$\frac{1}{2}^+$	7.30	$\frac{3}{2}^+$	< 0.7	(1965WA16)
				< 3	(1966WA08)
		7.57	$\frac{7}{2}^+$	< 3	(1965WA16, 1966WA08)
		0	$\frac{1}{2}^-$	92 ± 3	(1965WA16)
				92 ± 4	(1966WA08)
				91.6 ± 0.9	(1967PH03)
		5.27	$\frac{5}{2}^+$	3.8 ± 1	(1965WA16)
				3.5 ± 1	(1966WA08)
				4.7 ± 0.7	(1967PH03)
		6.32	$\frac{3}{2}^-$	3 ± 2	(1965WA16)
				4.5 ± 1	(1966WA08)
				3.7 ± 0.5	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
		7.30	$\frac{3}{2}^+$	1.2 ± 0.4	(1965WA16)
		7.57	$\frac{7}{2}^+$	< 2	(1965WA16)
8.31	$\frac{1}{2}^+$	< 0.5	(1965WA16)		
9.152 ^a	$\frac{3}{2}^-$	7.16 + 7.30 + 7.57		< 1	(1967PH03)
		0	$\frac{1}{2}^-$	97	(1968ST10)
				100	(1969SI04)
9.155 ^a	$(\frac{5}{2})$	5.27 + 5.30		3	(1968ST06)
		0	$\frac{1}{2}^-$	0 → 17	(1968ST10)
				17	(1967TH05)
		5.27	$\frac{5}{2}^+$	8	(1967TH05)
		5.30	$\frac{1}{2}^+$	10	(1967TH05)
		5.27 + 5.30		14 → 17	(1968ST10)
				23	(1969SI04)
		6.32	$\frac{3}{2}^-$	18 → 22	(1968ST10)
				19	(1969SI04)
				20	(1967TH05)
7.16	$\frac{5}{2}^+$			44 → 52	(1968ST10)
				58	(1969SI04)
				45	(1967TH05)
		7.30	$\frac{3}{2}^+$	7 → 9	(1968ST10)

Table 15.10: Radiative decays in ^{15}N (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
9.23	$\leq \frac{5}{2}^-$	0	$\frac{1}{2}^-$	< 30	(1965WA16)
				41.5 ± 2.2	(1967PH03)
		5.27	$\frac{5}{2}^+$	< 25	(1965WA16)
		5.30	$\frac{1}{2}^+$	100	(1965WA16)
		5.27 + 5.30		31.2 ± 1.7	(1967PH03)
		6.32	$\frac{3}{2}^-$	≤ 25	(1965WA16)
				24.7 ± 1.5	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 30	(1965WA16)
				< 1	(1967PH03)
		7.30	$\frac{3}{2}^+$	< 30	(1965WA16)
				2.6 ± 0.7	(1967PH03)
		7.57	$\frac{7}{2}^+$	< 20	(1965WA16)
		8.31	$\frac{1}{2}^+$	< 5	(1965WA16)
		7.57 + 8.31		< 1	(1967PH03)
9.76	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	100	(1965WA16)
				81.5 ± 2.8	(1967PH03)
		5.27 + 5.30		< 10	(1965WA16)
				7.5 ± 1.5	(1967PH03)
		6.32	$\frac{3}{2}^-$	< 5	(1965WA16)
				3.7 ± 0.8	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
				2.3 ± 0.5	(1967PH03)
		7.30	$\frac{3}{2}^+$	< 3	(1965WA16)
				< 2	(1967PH03)
		7.57	$\frac{7}{2}^+$	< 10	(1965WA16)
				5.0 ± 0.6	(1967PH03)
		8.31	$\frac{1}{2}^+$	< 2	(1965WA16)
				< 1	(1967PH03)
8.58	$\frac{3}{2}^+$	< 2	(1965WA16)		
		< 2	(1967PH03)		
9.83	$\frac{7}{2}$	0	$\frac{1}{2}^-$	< 30	(1965WA16)
				< 4	(1967PH03)
		5.27	$\frac{5}{2}^+$	100	(1965WA16)

Table 15.10: Radiative decays in ^{15}N (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
9.93	$(\frac{1}{2}, \frac{3}{2})^+$	5.30	$\frac{1}{2}^+$	< 15	(1965WA16)
		5.27 + 5.30		84.4 ± 1.8	(1967PH03)
		6.32	$\frac{3}{2}^-$	< 15	(1965WA16)
				2.2 ± 0.9	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
				2.4 ± 1.1	(1967PH03)
		7.30	$\frac{3}{2}^+$	< 10	(1965WA16)
				3.7 ± 0.9	(1967PH03)
		7.57	$\frac{7}{2}^+$	< 10	(1965WA16)
				7.3 ± 1.0	(1967PH03)
		8.31 + 8.58		< 1	(1967PH03)
		0	$\frac{1}{2}^-$	80 ± 10	(1965WA16)
				77.6 ± 1.9	(1967PH03)
		5.27 + 5.30		10 ± 10	(1965WA16)
				15.4 ± 1.5	(1967PH03)
		6.32	$\frac{3}{2}^-$	10 ± 10	(1965WA16)
				4.9 ± 1.2	(1967PH03)
7.16	$\frac{5}{2}^+$	< 10	(1965WA16)		
		< 1	(1967PH03)		
7.30	$\frac{3}{2}^+$	< 3	(1965WA16)		
		2.1 ± 0.8	(1967PH03)		
7.57	$\frac{7}{2}^+$	< 10	(1965WA16)		
8.31	$\frac{1}{2}^+$	< 2	(1965WA16)		
8.58	$\frac{3}{2}^+$	< 2	(1965WA16)		
7.57, 8.31, 8.58		< 1	(1967PH03)		
10.07	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	100	(1965WA16)
				96.0 ± 0.7	(1967PH03)
				94 ± 4	(1966WA08)
		5.27 + 5.30		< 10	(1965WA16)
				4.0 ± 0.7	(1967PH03)
				6 ± 2	(1966WA08)
		6.32	$\frac{3}{2}^-$	< 5	(1965WA16)
				< 2	(1966WA08)

Table 15.10: Radiative decays in ^{15}N (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
10.45	$\frac{3}{2} \rightarrow \frac{7}{2}$	7.16	$\frac{5}{2}^+$	< 7	(1965WA16)
				< 2	(1966WA08)
		7.30	$\frac{3}{2}^+$	< 3	(1965WA16)
				< 2	(1966WA08)
		7.57	$\frac{7}{2}^+$	< 7	(1965WA16)
				< 2	(1966WA08)
		8.31	$\frac{1}{2}^+$	< 2	(1965WA16)
		8.58	$\frac{3}{2}^+$	< 3	(1965WA16)
		0	$\frac{1}{2}^-$	12 ± 12	(1965WA16)
				< 4	(1966WA08, 1967PH03)
		5.27 + 5.30		72 ± 8	(1965WA16)
				66 ± 5	(1966WA08)
				62.4 ± 2.4	(1967PH03)
		6.32	$\frac{3}{2}^-$	12 ± 8	(1965WA16)
				28 ± 5	(1966WA08)
				14.7 ± 1.6	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
				< 6	(1966WA08)
		7.30	$\frac{3}{2}^+$	4 ± 4	(1965WA16)
				< 6	(1966WA08)
7.57	$\frac{7}{2}^+$	< 10	(1965WA16)		
		< 6	(1966WA08)		
8.31	$\frac{1}{2}^+$	1.5 ± 0.5	(1967PH03)		
9.05	$\frac{1}{2}^+$	< 1	(1967PH03)		
9.15 + 9.23		5.0 ± 0.5	(1967PH03)		
9.76	$\frac{5}{2}^-$	1.6 ± 0.7	(1967PH03)		
9.83	$\frac{7}{2}$	2.2 ± 1.5	(1967PH03)		
9.93	$(\frac{1}{2}, \frac{3}{2})^+$	3.7 ± 1.1	(1967PH03)		
10.07	$\frac{3}{2}^+$	< 4	(1967PH03)		
10.54 ^b	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	1 ± 0.3	(1969SI04)
		5.27	$\frac{5}{2}^+$	30 ± 2	(1969SI04)
		6.32	$\frac{3}{2}^-$	7 ± 0.5	(1969SI04)
		7.16	$\frac{5}{2}^+$	23 ± 1.5	(1969SI04)

Table 15.10: Radiative decays in ^{15}N (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Refs.
10.70	$\frac{3}{2}^+$	7.30	$\frac{3}{2}^+$	37 ± 2.5	(1969SI04)
		8.58	$\frac{3}{2}^+$	1.8 ± 0.4	(1969SI04)
		(9.22)	$\leq \frac{5}{2}^-$	< 1	(1969SI04)
		0	$\frac{1}{2}^-$	52 ± 1	(1969SI04)
		5.27	$\frac{5}{2}^+$	38 ± 1	(1969SI04)
		6.32	$\frac{3}{2}^-$	6 ± 0.4	(1969SI04)
		7.16	$\frac{5}{2}^+$	< 1	(1969SI04)
		7.30	$\frac{3}{2}^+$	3 ± 0.3	(1969SI04)
		8.31	$\frac{1}{2}^+$	< 1	(1969SI04)
		9.05	$\frac{1}{2}^+$	< 1	(1969SI04)
10.80	$\frac{3}{2}^+$	(9.22)	$\leq \frac{5}{2}^-$	< 1	(1969SI04)
		0	$\frac{1}{2}^-$	55 ± 5	(1965WA16)
				53 ± 15	(1966WA08)
				47 ± 1	(1969SI04)
		5.27	$\frac{5}{2}^+$	5 ± 0.5	(1969SI04)
		5.30	$\frac{1}{2}^+$	13 ± 0.6	(1969SI04)
		5.27 + 5.30		45 ± 5	(1965WA16)
				47 ± 5	(1966WA08)
		6.32	$\frac{3}{2}^-$	< 5	(1965WA16)
				< 10	(1966WA08)
				7 ± 0.5	(1969SI04)
		7.16	$\frac{5}{2}^+$	< 5	(1965WA16)
				9 ± 0.5	(1969SI04)
		7.30	$\frac{3}{2}^+$	< 5	(1965WA16)
				< 4	(1966WA08)
		8 ± 0.5	(1969SI04)		
7.57	$\frac{7}{2}^+$	< 7	(1965WA16)		
8.31	$\frac{1}{2}^+$	5 ± 0.5	(1969SI04)		
9.05	$\frac{1}{2}^+$	1 ± 0.3	(1969SI04)		
9.152	$\frac{3}{2}^-$	2 ± 0.3	(1969SI04)		
9.155	$(\frac{5}{2})$	4 ± 0.3	(1969SI04)		

^a See also (1965WA16, 1967PH03).

^b See also (1960HE13).

^c See also (1969SI04).

$$23. \text{}^{13}\text{C}(\text{}^6\text{Li}, \alpha)\text{}^{15}\text{N} \quad Q_m = 14.688$$

Angular distributions have been measured for $E(^6\text{Li}) = 3.2$ to 3.8 MeV for the α_0 and α_{1+2} groups (1964BL1B).

$$24. \text{}^{13}\text{C}(\text{}^7\text{Li}, \text{}^5\text{He})\text{}^{15}\text{N} \quad Q_m = 6.478$$

See (1969TH01).

$$25. \text{}^{13}\text{C}(\text{}^{11}\text{B}, \text{}^9\text{Be})\text{}^{15}\text{N} \quad Q_m = 0.341$$

See (1966PO1E, 1967PO1E, 1967VO1A).

$$26. \text{}^{14}\text{C}(\text{p}, \gamma)\text{}^{15}\text{N} \quad Q_m = 10.208$$

Resonances for capture γ -radiation are listed in Table 15.11 (1959FE1C, 1959HE1D, 1968HE12, 1968SI1F, 1969SI04). A combination of $^{15}\text{N}^*(10.80)$ and (9.83) permits a good account of the low energy (n, n) and (n, γ) cross sections (1959HE1D). The thermal (n, p) cross section can be ascribed to the $E_p = 1.5$ MeV resonance ($^{15}\text{N}^*(11.62)$) (1955BA44: see also $^{14}\text{N}(\text{n}, \gamma)^{15}\text{N}$). See also (1959AJ76) and (1969TI05).

Table 15.10 displays branching ratios obtained in this and in other reactions. The angular distributions of γ -rays at the $E_p = 0.35$ MeV resonance ($^{15}\text{N}^*(10.54)$) leads to assignments of $J = \frac{5}{2}$, $(\frac{5}{2})$, $\frac{3}{2}$ and $\frac{5}{2}$, respectively, for $^{15}\text{N}^*(5.27, 7.16, 7.30, 10.54)$ (1960HE13). The angular distribution of the $(10.81 \rightarrow 8.31)$ γ -ray fixes $J = \frac{1}{2}$ for $^{15}\text{N}^*(8.31)$ (1968SI1E). A triple correlation study by (1968SI1E) of the decay of $^{15}\text{N}^*(10.80)$ to the $E_x = 9.15$ MeV states suggests $J = \frac{3}{2}$ for the upper and $J = \frac{5}{2}$ for the lower of these two states. Lifetimes for various ^{15}N states have been measured by (1968COZV): see Table 15.7.

See also (1960FR09, 1961FR1D, 1969ZH1A).

$$27. \text{}^{14}\text{C}(\text{p}, \text{p})\text{}^{14}\text{C} \quad E_b = 10.208$$

Table 15.11: Resonances in $^{14}\text{C} + \text{p}$ ^a

E_p (MeV \pm keV)	Γ (keV)	Γ_n (keV)	Γ_p (keV)	Γ_γ (eV)	J^π ^d	E_x (MeV)	Refs.
0.261 ± 1 ^a						10.451	(1959HE1D, 1968SI1F, 1969SI04)
0.352 ± 1 ^a				$(3.4 \pm 0.4) \times 10^{-2}$ ^c	$\frac{1}{2}^{\pm}$	10.536	(1959HE1D, 1960HE13, 1968SI1F, 1969SI04)
0.527 ± 1 ^a			0.2	1.78 ± 0.3 ^c	$\frac{3}{2}^+$	10.700	(1959HE1D, 1968SI1F, 1969SI04)
0.634 ± 1 ^a				0.23 ± 0.04 ^c	$\frac{3}{2}^+$	10.800	(1959HE1D, 1968SI1F, 1969SI04)
1.162 ± 2	7.9 ± 3	2.3	5.6	0.29	$\frac{1}{2}^-$	11.292	(1951RO16, 1955BA44, 1956SA06, 1968HA27)
1.294 ± 4	34 ± 4	22.6	11.1			^e	(1968HA27)
1.3188 ± 0.5	41.4 ± 1.1	34.6 ± 0.9	6.8 ± 0.5	4.2 ± 0.7	$\frac{1}{2}^+$	11.438	(1951RO16, 1955BA44, 1956SA06, 1968HA27)
1.472 ± 8	418 ± 16	3.7	414.6			^e	(1968HA27)
1.509 ± 4	404.9 ± 6.3	4.0 ± 0.2	400.9 ± 6.3	19.2 ± 0.4	$\frac{1}{2}^+; T = \frac{3}{2}$	11.615	(1959FE1C, 1968HE12)
1.688 ± 3	37	36.5	0.5		$\frac{3}{2}^+$	11.764	(1951RO16, 1955BA44, 1956SA06)
1.788 ± 3	24.5	24.5	0.03		$\frac{3}{2}^-, (\frac{5}{2}^-)$	11.876	(1951RO16, 1955BA44, 1956SA06)
1.884 ± 3	21.5	21.2	0.3		$\frac{1}{2}^-$	11.965	(1951RO16, 1955BA44, 1956SA06)
2.025 ± 4 ^b	14 ± 5	12.0	1.7		$\frac{1}{2}^+$	12.097	(1951RO16, 1955BA44, 1956SA06, 1960BA34, 1968HA27)
2.077 ± 3 ^b	47 ± 7	30.2	16.6		$\frac{3}{2}^-$	12.145	(1951RO16, 1955BA44, 1956SA06, 1960BA34, 1968HA27, 1968HE12)
2.272 ± 4	22	21.7	0.3		$\frac{1}{2}^+$ (+)	12.327	(1951RO16, 1955BA44, 1956SA06, 1960BA34, 1968HA27)
2.450 ± 4 ^b	40	28	0.3		$\frac{1}{2}^{\pm}; T = \frac{1}{2}$	12.493	(1951RO16, 1956SA06, 1960BA34)
2.48 ± 10	80	0	80		$\frac{1}{2}^+; T = \frac{3}{2}$	12.52	(1968HA27, 1968HE12)
2.908 ± 4	75				$\frac{3}{2}^-$	12.921	(1956SA06, 1960BA34, 1961HA12, 1962HA20)
3.19	5.5					13.19	(1960BA34, 1961HA12, 1962HA20)
3.38	22					13.36	(1960BA34, 1961HA12, 1962HA20)

Table 15.11: Resonances in $^{14}\text{C} + \text{p}$ ^a (continued)

E_{p} (MeV \pm keV)	Γ (keV)	Γ_{n} (keV)	Γ_{p} (keV)	Γ_{γ} (eV)	J^{π} ^d	E_{x} (MeV)	Refs.
3.42	61					13.40	(1960BA34, 1961HA12, 1962HA20)
3.55						(13.52)	(1962HA20)
3.63	12					13.60	(1960BA34, 1961HA12, 1962HA20)
3.71						13.67	(1961HA12, 1962HA20)
3.89	33					13.84	(1960BA34, 1961HA12, 1962HA20)
4.19	105					14.12	(1960BA34, 1961HA12, 1962HA20)
4.24	25					14.17	(1960BA34, 1961HA12, 1962HA20)
4.61	130					14.51	(1960BA34, 1961HA12, 1962HA20)
4.93	99					14.81	(1960BA34, 1961HA12)
5.01						14.88	(1961HA12, 1962HA20)
5.14						15.00	(1961HA12, 1962HA20)
5.26						15.11	(1961HA12, 1962HA20)

^a See also (1959HE1B) and Table 15.5 in (1959AJ76).

^b The Γ_{α} are < 0.3 keV for $E_{\text{res}} = 1.16$ to 2.5 MeV, except for the resonances at $E_{\text{p}} = 2.03$, 2.08 and 2.45 MeV for which they are, respectively, 0.6, 2.2 and 5.5 keV (1956SA06).

^c ω_{γ} (in eV) (1968SI1F, 1969SI04).

^d See also (1959AJ76).

^e These states correspond to the levels listed in the line below: a different boundary condition was used to obtain E_{res} .

The elastic scattering has been studied for $E_p = 0.34$ to 2.7 MeV. At the $E_p = 527$ keV resonance (see Table 15.11), the scattering is consistent with d-wave formation of a $J^\pi = \frac{3}{2}^+$ state. No anomalies are observed at $E_p = 0.35$ and 0.63 MeV (1959HE1D). However, anomalies are observed at $E_p = 1.16, 1.29, 1.47, 2.02, 2.07$ and 2.48 MeV: the parameters of these are displayed in Table 15.11 (1968HA27, 1968HE12). The 2.48 MeV anomaly is due to a $J^\pi = \frac{5}{2}^+$; $T = \frac{3}{2}$ state at $E_x = 12.52$ MeV (which is the analog to the first excited state of ^{15}C), and is distinct from the $J^\pi = \frac{5}{2}^+$ resonance in $^{14}\text{C}(p, n)^{14}\text{N}$ observed at $E_p = 2.45$ MeV (1968HE12). See also (1968IW1A, 1969IW1E).

$$28. \ ^{14}\text{C}(p, n)^{14}\text{N} \qquad Q_m = -0.626 \qquad E_b = 10.208$$

Resonances reported by (1951RO16, 1955BA44, 1956SA06, 1959FE1C, 1960BA34, 1961HA12, 1962HA20, 1968HE12) are listed in Table 15.11. In addition to these, (1961HA12) report (in an abstract) a broad resonance corresponding to a $T = \frac{3}{2}$ state of ^{15}N at $E_x = 13.6$ MeV. At $E_p = 1.79$ MeV, the distributions favor $\frac{5}{2}^-$, but $\frac{3}{2}^-$ is not excluded (1955BA44: see also (1953KA20)); a computation of the cross section favors $J = \frac{3}{2}$ (1956SA06). At $E_p = 1.88$ MeV, the angular distribution is consistent with the $J^\pi = \frac{1}{2}^-$ assignment from $^{14}\text{N}(n, n)^{14}\text{N}$ (1955BA44). The $E_p = 2.27$ MeV state has $J = \frac{3}{2}$ or $\frac{5}{2}$; the σ_{nn} clearly indicates the latter (1955BA44, 1956SA06). For $^{15}\text{N}^*(11.62)$ ($E_p = 1.51$ MeV), the proton reduced width indicates a single-particle level, while the neutron reduced width is only 10^{-3} . This is consistent with the assignment $T = \frac{3}{2}$, corresponding to the ground state of ^{15}C (1955BA44, 1956BA16, 1959FE1C). See also (1967VO1B, 1969BA1N, 1969TI05).

Polarization measurements are reported at $E_p = 7.2, 8.8$ and 10.4 MeV by (1969WO1J; n_0, n_1, n_2).

$$29. \ ^{14}\text{C}(d, n)^{15}\text{N} \qquad Q_m = 7.984$$

Angular distributions have been determined at $E_d = 1.3, 1.8$ and 2.8 MeV (1967CO1R; $n_0, n_{1+2}, n_3, n_{4+5+6}$), 1.31 to 3.08 MeV (1963IM01; n_0), 2.10 and 2.33 MeV (1963IM01; n_{1+2}, n_3, n_{4+5+6}) and 3.10 MeV (1967LA11; $n_0, n_{1+2}, n_3, n_4, n_5, n_6$ and $^{15}\text{N}^*(8.31, 9.06, 9.16, 9.22)$). The transitions to $^{15}\text{N}^*(5.30, 9.06)$ involve $l_p = 0$: these states therefore have $J^\pi = \frac{1}{2}^+$ (1967LA11). See also (1959AJ76) and (1961CH14, 1964MO1G).

Studies of the γ -decay of ^{15}N states reached in this and in other reactions are summarized in Table 15.10: see (1965WA16). See also (1967CH19).

$$30. \ ^{14}\text{C}(^3\text{He}, d)^{15}\text{N} \qquad Q_m = 4.715$$

At $E(^3\text{He}) = 14$ MeV, angular distributions of the deuterons corresponding to $^{15}\text{N}^*(0, 5.28, 5.30, 6.33)$ have been measured and analyzed by DWBA: the relative spectroscopic factors for the first four states of ^{15}N are 1, 0.50, 0.34, 0.07 in good agreement with the predictions of (1957HA1E). Angular distributions of the d_0 group are also reported at $E(^3\text{He}) = 1 - 9$ MeV by (1963WE15, 1966DU1B, 1968DA1N).

$$31. \ ^{14}\text{C}(\alpha, t)^{15}\text{N} \quad Q_m = -9.606$$

Not reported.

$$32. \ ^{14}\text{N}(n, \gamma)^{15}\text{N} \quad Q_m = 10.835$$

$$Q_0 = 10.8332 \pm 0.0006 \text{ (1968GR14).}$$

The thermal cross section is 80 ± 20 mb (1957BA18), 75 ± 7.5 mb (1964ST25). This large cross section is not understood in terms of the present level structure in ^{15}N : see (1959AJ76).

Observed capture γ rays are displayed in Table 15.12 (1963MO1C, 1967TH05, 1968GR14). The very accurate γ -ray energy determinations of (1968GR14) show that two states at $E_x \approx 9.15$ MeV are involved in this reaction as previously suggested by (1966WA08). The lower of the two, at $E_x = 9.1518$ MeV, decays predominantly to the ground state. The other state at $E_x = 9.1549$ MeV, which is preferentially fed in this reaction, decays primarily by cascades via $^{15}\text{N}^*(5.27, 5.30, 6.32, 7.16)$ (1968GR14). See also reactions 21 and 39, (1968GRZY, 1968ST10) and Table 15.10.

Recoil Doppler broadening of cascade γ -rays has been measured by (1969WE07): the derived τ_m are listed in Table 15.7. See also (1968CA1J). Observation of non-isotropic correlations in the $\text{C} \rightarrow 6.32$ cascade means that $J \neq \frac{1}{2}$ for $^{15}\text{N}^*(6.32)$: the results are consistent with $J = \frac{3}{2}$ (1964BA02).

The importance of measuring the (n, γ) cross section at $E_n = 0.47$ and 0.66 MeV for astrophysical considerations is suggested by (1968FO1A).

See also (1960CA02, 1965JA09, 1969HO1X) and (1958GR01, 1964LI1C).

$$33. \ ^{14}\text{N}(n, n)^{14}\text{N} \quad E_b = 10.835$$

The thermal (bound) scattering cross section is 5.51 b (1961WI1A). The scattering amplitude (bound) is $a = 9.19 \pm 0.11$ fm (1965DO14) [$a = 9.14 \pm 0.10$ fm is recommended by (1964ST25)]. See also (1969BA1P). The coherent scattering cross section is 10.5 ± 0.3 b (1964ST25).

Recent cross section measurements are listed in Table 15.13. Cross section data are summarized in (1964ST25), while angular distribution data are displayed in (1963GO1M).

Observed resonances are listed in Table 15.14 (1951JO23, 1952HI12, 1955FO27, 1959HA13, 1966FO1D, 1966FO1E, 1966MA2K, 1968BO36, 1968JO1F): for a discussion of the evidence

Table 15.12: Gamma radiation from $^{14}\text{N}(n, \gamma)^{15}\text{N}$

Transition in ^{15}N	E_γ (MeV \pm keV)		E_γ (MeV \pm keV)	I_γ ^b	
	(1967TH05)	(1968GR14)	(1968GR14)	(1967TH05)	(1963MO1C)
C \rightarrow 0	10.830 \pm 2	10.8290	10.8832 \pm 0.6	13.3 \pm 2.0	14
C \rightarrow 5.27	5.562 \pm 1	5.5622 \pm 0.35		10.3 \pm 0.5	11
C \rightarrow 5.30	5.534 \pm 1	5.5332 \pm 0.35		18.8 \pm 0.9	21
C \rightarrow 6.32	4.509 \pm 1	4.5088 \pm 0.3		16.6 \pm 0.8	16
C \rightarrow 7.16	3.678 \pm 1	3.6777 \pm 0.25		15.9 \pm 0.8	< 23
C \rightarrow 7.30	3.532 \pm 1	3.5322 \pm 0.25		9.9 \pm 0.5	9
C \rightarrow 8.31	2.521 \pm 2	2.52055 \pm 0.10		6.1 \pm 0.3	6
C \rightarrow 9.155		1.67819 \pm 0.06			
	1.679 \pm 2			9.2 \pm 0.5	12
C \rightarrow 9.152		1.68141 \pm 0.18		1.4 \pm 0.3 ^d	
5.27 \rightarrow 0	5.270 \pm 1	5.2692 \pm 0.35	5.2701 \pm 0.3	30.6 \pm 1.5	32
5.30 \rightarrow 0	5.298 \pm 1	5.2978 \pm 0.35	5.2989 \pm 0.3	21.4 \pm 1.1	21
6.32 \rightarrow 0	6.323 \pm 1	6.3220 \pm 0.4	6.3235 \pm 0.4	18.8 \pm 0.9	18
7.16 \rightarrow 0			7.1550 \pm 0.4		
7.17 \rightarrow 5.27	1.885 \pm 1	1.88481 \pm 0.06		19.7 \pm 1.0	21
7.16 \rightarrow 5.30	1.857 \pm 2			0.8 \pm 0.2	
7.30 \rightarrow 0	7.299 \pm 1	7.2990 \pm 0.5	7.3009 \pm 0.5	10.0 \pm 0.5	9
8.31 \rightarrow 0	8.311 \pm 2	8.3102 \pm 0.7	8.3124 \pm 0.7	4.4 \pm 0.4	4
8.31 \rightarrow 6.32	1.989 \pm 2			1.5 \pm 0.3	
8.57 \rightarrow 0	8.570 \pm 4		8.573 \pm 4 ^c	0.2 \pm 0.03	
9.05 \rightarrow 0	9.047 \pm 4		9.050 \pm 4 ^c	0.2 \pm 0.03	0.5
9.155 \rightarrow 0			9.1549 \pm 0.5		
9.152 \rightarrow 0	9.151 \pm 2	9.1498 \pm 0.9	9.1518 \pm 0.5	1.7 \pm 0.2	1.4
9.155 \rightarrow 5.27	3.883 \pm 2	3.8837 \pm 0.4		0.8 \pm 0.1	
9.155 \rightarrow 5.30	3.855 \pm 2	3.8552 \pm 0.3		1.0 \pm 0.1	0.5
9.155 \rightarrow 6.32	2.830 \pm 2	2.8311 \pm 0.2		2.0 \pm 0.2	1.5
9.155 \rightarrow 7.16	1.997 \pm 2	1.99965 \pm 0.10		4.6 \pm 0.2	4

^a C = capturing state.

^b In units of photons/100 captures.

^c (1967TH05).

^d (1968GR14).

leading to J^π assignments, see (1959AJ76). See also (1960BA34) and (1966AG1A, 1967BE1F, 1968IW1A, 1969IW1E).

A polarization measurement has been made at $E_n = 3.5$ MeV by (1962OT01). See also (1965TA07).

34. $^{14}\text{N}(n, 2n)^{13}\text{N}$

$Q_m = -10.553$

$E_b = 10.835$

Table 15.13: Recent cross section measurements for $^{14}\text{N} + n$ ^a

E_n (MeV)	Measurements of	Refs.
0.01 – 0.2	σ_t	(1959BI19)
1.7 – 4.0	σ_t	(1960BA34)
2.0 – 4.2	σ_t	(1966MA2K, 1968JO1F)
4.2 – 6.3	σ_{ne}	(1968BO36)
4.2 – 8.7	σ_e	(1968BO36)
4.5, 6, 7	$\sigma_{nn'\gamma}$	(1968CO1W)
5.80 – 8.55	$\sigma_{nn'\gamma}$	(1969DI1B)
6.8 – 14.0	$\sigma_t, \sigma_e, \sigma_{ne}$	(1967BA03)
13.3 – 19.5	σ_e	(1968BO36)
14	σ_e	(1963BA46)
15	σ_{ne}	(1969NY1A)
15.2, 19.8	σ_{ne}	(1965DE1G)
15.3	σ_t, σ_{ne}	(1968HA1V)
17.7 – 29.1	σ_t	(1960PE25)
88 – 152	σ_t	(1966ME14)
10 – 37	$\sigma_{n,2n}$	(1961BR1A)
12.5 – 18	$\sigma_{n,2n}$	(1960FE12)
13.2 – 18.8	$\sigma_{n,2n}$	(1965BO42)
13.3 – 15.2	$\sigma_{n,2n}$	(1960MC05)
14.1	$\sigma_{n,2n}$	(1962CE1B)
14.4	$\sigma_{n,2n}$	(1961RA06)
14.7	$\sigma_{n,2n}$	(1967PA27)
14.8	$\sigma_{n,2n}$	(1965GR41)
1.3 – 4.2	$\sigma_{n,p}$	(1959GA14)

Table 15.13: Recent cross section measurements for $^{14}\text{N} + \text{n}$ ^a (continued)

E_n (MeV)	Measurements of	Refs.
7.4 – 8.6	$\sigma_{n,p}$	(1969DI1B)
14.7	$\sigma_{n,p}, \sigma_{n,np}$	(1966CS1B, 1967CS03)
15	$\sigma_{n,p}, \sigma_{n,d}$	(1969NY1A)
5.6 – 6.4	$\sigma_{n,t}$	(1966SC21)
5.7 – 8.2	$\sigma_{n,t}$	(1959GA14)
14.1	$\sigma_{n,2\alpha}$	(1967MO21)
15	$\sigma_{n,t}$	(1969NY1A)
15.7	$\sigma_{n,t}$	(1967MO21)
1.3 – 8.2	$\sigma_{n,\alpha}$	(1959GA14)
3.9 – 6.4	$\sigma_{n,\alpha}$	(1966SC21)
4.5 – 7.0	$\sigma_{n,\alpha}$	(1968CO1W)
5.7 – 8.2	$\sigma_{n,\alpha}$	(1959HA13)
5.8 – 8.6	$\sigma_{n,\alpha}$	(1969DI1B)
14.7	$\sigma_{n,\alpha}$	(1966CS1B)
15	$\sigma_{n,\alpha}$	(1969NY1A)

^a See also (1959AJ76).

Recent measurements of cross sections for this reaction are listed in Table 15.13 (1960FE12, 1960MC05, 1961BR1A, 1961RA06, 1962CE1B, 1965BO42, 1965GR41, 1967PA27): see the summaries in (1964ST25, 1966JE1B). See also (1964HE18, 1966CS1C, 1967CS02) and (1959AJ76).

35. $^{14}\text{N}(n, p)^{14}\text{C}$

$$Q_m = 0.626$$

$$E_b = 10.835$$

The thermal cross-section is 1.81 ± 0.05 b (1964ST25), 1.83 ± 0.03 b (1961HA43). A number of resonances are reported by (1950JO57), (1959GA14) and (1963EN01): see Table 15.14. The results are summarized in (1964ST25, 1966JE1B). See also (1960BU1C, 1968DA1F).

Recent cross-section measurements are listed in Table 15.13 (1959GA14, 1966CS1B, 1967CS03, 1969DI1B). See also (1964FO1A, 1969BA1N) and (1959AJ76).

36. (a) $^{14}\text{N}(n, d)^{13}\text{C}$

$$Q_m = -5.325$$

$$E_b = 10.835$$

(b) $^{14}\text{N}(n, np)^{13}\text{C}$

$$Q_m = -7.550$$

For reaction (a) see (1967LI06) and (1959AJ76). For reaction (b) see (1966CS1B, 1967CS03).

Table 15.14: Resonances in $^{14}\text{N} + \text{n}$ ^a

E_{res} (MeV \pm keV)	Γ_{lab} (keV)	Γ_{n} (keV)	Γ_{p} (keV)	Γ_{α} (keV)	J^{π}	$^{15}\text{N}^*$ (MeV)	Refs.
0.430 ± 5	3.5	< 3	< 0.01		$\geq \frac{1}{2} \frac{3}{2} \frac{3}{2}$	11.236	(1951JO23, 1952HI12)
0.4926 ± 0.65	7.5	< 3	< 10		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{-}$	11.2943	(1950JO57, 1952HI12, 1963EN01)
0.639 ± 5	43	34	9		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{+}$	11.431	(1950JO57, 1951JO23, 1952HI12)
0.998 ± 5	46	45	0.8		$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	11.766	(1950JO57, 1951JO23, 1952HI12)
1.120 ± 6	19	19	0.20		$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	11.880	(1951JO23, 1952HI12, 1955FO27)
1.188 ± 6	≤ 3.2	< 2	< 0.1		$\geq \frac{1}{2} \frac{1}{2} \frac{1}{2}$	11.943	(1952HI12)
1.211 ± 7	13	12	0.4		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{-}$	11.965	(1952HI12)
1.350 ± 7	21	20	0.9	0.4	$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{+}$	12.094	(1951JO23, 1952HI12, 1955FO27, 1959GA14)
1.401 ± 8	54	41	11	1.8	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	12.142	(1950JO57, 1951JO23, 1952HI12, 1955FO27, 1959GA14)
1.595 ± 8	22	21	0.2	< 0.1	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	12.323	(1952HI12, 1955FO27, 1959GA14, 1966FO1D, 1966FO1E)
1.779 ± 10	47	37	0.5	9.0	$(\frac{3}{2} \frac{3}{2} \frac{3}{2})^{+}$	12.494	(1950JO57, 1952HI12, 1955FO27, 1959GA14, 1966FO1E)
2.23	65	39	7.8	18	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	12.92	(1959GA14, 1966FO1D, 1966FO1E, 1966MA2K, 1968JO1F)
2.47	< 3			r		13.14	(1959GA14)
2.52	≈ 7			r		13.19	(1959GA14)
2.71	40			r	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	13.36	(1959GA14)
2.74	95		r		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{+}$	13.39	(1959GA14)
2.95	20	16	1.1	3.2	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	13.59	(1959GA14, 1966FO1E, 1966MA2K, 1968JO1F)
3.09	60		r	r		13.72	(1959GA14)
3.21	85		r	r	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	13.83	(1959GA14, 1966FO1E, 1966MA2K, 1968JO1F)
3.51	≈ 20		r	r		14.11	(1959GA14)
3.57	30		r	r	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	14.17	(1959GA14, 1966MA2K, 1968JO1F)
≈ 3.8	≈ 2000	≈ 1000	200	≈ 1000		14.4	(1959GA14)
4.09	50		r	r		14.65	(1959GA14, 1966SC21)
≈ 4.1	≈ 300		r	r		14.7	(1959GA14)
4.38	40			r		14.92	(1959GA14, 1966SC21)
4.60				r		15.13	(1959GA14, 1966SC21)
5.03				r		15.52	(1959GA14, 1966SC21, 1968BO36)
5.60	100			r		16.06	(1959GA14, 1966SC21, 1968BO36)

Table 15.14: Resonances in $^{14}\text{N} + \text{n}$ ^a (continued)

E_{res} (MeV \pm keV)	Γ_{lab} (keV)	Γ_{n} (keV)	Γ_{p} (keV)	Γ_{α} (keV)	J^{π}	$^{15}\text{N}^*$ (MeV)	Refs.
5.94				r		16.38	(1959GA14, 1966SC21)
6.16	75			r		16.58	(1959GA14, 1966SC21)
6.26	110	r		r		16.67	(1959GA14, 1959HA13, 1966SC21)
6.55	170	r		r		16.95	(1959GA14, 1959HA13)
6.94				r		17.31	(1959GA14)
	200	r					(1959HA13)
7.16				r		17.51	(1959GA14)
7.34	120			r		17.68	(1959GA14)
7.48	180	r		r		17.81	(1959GA14, 1959HA13)
7.92	170			r		18.22	(1959HA13)
8.00	120			r		18.30	(1959GA14, 1959HA13)

r = resonant channel.

^a See also (1959AJ76).

37. (a) $^{14}\text{N}(\text{n}, \text{t})^{12}\text{C}$	$Q_{\text{m}} = -4.015$	$E_{\text{b}} = 10.835$
(b) $^{14}\text{N}(\text{n}, \text{t})^4\text{He}^4\text{He}^4\text{He}$	$Q_{\text{m}} = -11.289$	
(c) $^{14}\text{N}(\text{n}, 2\alpha)^7\text{Li}$	$Q_{\text{m}} = -8.822$	

For reaction (a) see (1959GA14, 1966SC21, 1967LI06, 1967RE01) and (1968AJ02). For reaction (b) see (1967MO21). For reaction (c) see (1967MO21). See also (1960FA10, 1964SA1E), Table 15.13, the summaries in (1964ST25, 1966JE1B) and (1959AJ76).

38. (a) $^{14}\text{N}(\text{n}, \alpha)^{11}\text{B}$	$Q_{\text{m}} = -0.157$	$E_{\text{b}} = 10.835$
(b) $^{14}\text{N}(\text{n}, \text{n}\alpha)^{10}\text{B}$	$Q_{\text{m}} = -11.613$	

Recent cross-section measurements for reaction (a) are displayed in Table 15.13: these include measurements of cross sections for several different α -groups (1959GA14, 1966SC21) and γ -rays (1959HA13, 1968CO1W, 1969DI1B). For summaries of the experimental evidence, see (1964ST25, 1966JE1B).

Observed resonances are listed in Table 15.14 (1950JO57, 1959GA14, 1966SC21). See also (1960FA10), (1960BU1C, 1963CH1C) and (1964GA1A, 1968GA1M; theor.). For reaction (b), see (1960FA10).

39. $^{14}\text{N}(\text{d}, \text{p})^{15}\text{N}$	$Q_{\text{m}} = 8.610$	
	$Q_0 = 8.614 \pm 0.006$ (1964MA57).	

Proton groups corresponding to levels of ^{15}N are listed in Table 15.15 (1950MA65, 1954SP01, 1956DO41, 1965AL19, 1966GA08, 1966GO1J, 1969PH02). The J^π assignments are based on PWBA and DWBA analyses: see (1959AJ76), (1969PH02) and Table 15.16 for a listing of recent angular distribution studies in the range $E_{\text{d}} = 0.5$ to 27 MeV. See also (1966RO1V). The angular distributions of protons corresponding to $^{15}\text{N}^*(6.32)$ and the p- γ angular correlation fixes $J^\pi = \frac{3}{2}^-$ for that state (1961GO03). (1960HO1B) has looked for additional states of ^{15}N with $E_{\text{x}} \approx 7$ MeV: he finds the upper limits to proton groups corresponding to $7.10 < E_{\text{x}} < 7.31$ to be 1%, and to be 5% of the intensities of groups to known nearby states for $7.31 < E_{\text{x}} < 7.57$ MeV.

Recent very accurate γ -ray energy measurements have been reported by (1967CH19) [$E_{\gamma} = 5299.03 \pm 0.43$ and 5270.60 ± 0.46 keV] by (1966AL18) and by (1965WA16): the derived E_{x} values are displayed in Table 15.15. Branching ratios have been determined by (1965WA16, 1968GI11, 1968ST10) and are shown in Table 15.10 together with the multiplicities determined by (1965WA16). Lifetime measurements are listed in Table 15.7 (1965AL19, 1967BI11, 1968GI11).

The two states of ^{15}N at $E_{\text{x}} = 9.15$ MeV [see Table 15.10 for branching ratios] are separated by 2.5 ± 0.5 keV (1968ST10, 1969YO1C): see also reaction 32.

Table 15.15: ^{15}N levels from $^{14}\text{N}(\text{d}, \text{p})^{15}\text{N}$

E_x (MeV \pm keV)					l_n	J^π
(1950MA65, 1966GO1J)	(1954SP01)	(1956DO41, 1967CH19)	A	(1966GA08)		
0					1 ^b	$\frac{1}{2}^-$, $\frac{3}{2}^-$, $\frac{5}{2}^-$
5.276 \pm 6	5.280 \pm 10	5.27159 \pm 0.46 ^a	5.272 \pm 10		2 ^c	$\leq \frac{7}{2}^+$
5.305 \pm 6		5.30003 \pm 0.43 ^a	5.300 \pm 11		c,d	
6.328 \pm 6	6.330 \pm 10				1 ^{e,i}	$\frac{3}{2}^-$ j
7.164 \pm 6	7.165 \pm 10		7.1555 \pm 1.7		2 ^{f,i}	$\leq \frac{7}{2}^+$
7.309 \pm 6	7.314 \pm 10	7.307 \pm 8			0 ^{f,i}	$\frac{1}{2}^+$, $\frac{3}{2}^+$
	7.575 \pm 10	7.570 \pm 8	7.5671 \pm 1.0		2 ^{g,i}	$\leq \frac{7}{2}^+$
8.315 \pm 6	8.316 \pm 10	8.319 \pm 8	8.309 \pm 4.1		0 ^{e,i}	$\frac{1}{2}^+$, $\frac{3}{2}^+$
	8.571 \pm 10	8.577 \pm 8	8.573 \pm 3.2	8.582 \pm 5	0 + 2 ^{h,i}	$\leq \frac{7}{2}^+$
	9.062 \pm 10			9.056 \pm 5		
	9.165 \pm 10			9.159 \pm 6	i	
9.225 \pm 6				9.226 \pm 6	1 or 2 ⁱ	$(\frac{3}{2}^-)$
9.762 \pm 6				9.764 \pm 6		
	9.834 \pm 10			9.831 \pm 6		
9.929 \pm 7				9.929 \pm 6		
	10.069 \pm 10			10.071 \pm 6	2, 0 ⁱ	$\frac{3}{2}^+$
	10.458 \pm 10			10.456 \pm 7		
	10.544 \pm 10			10.541 \pm 7		
	10.705 \pm 10			10.702 \pm 7	2, 0 ⁱ	$\frac{3}{2}^+$
	10.811 \pm 10			10.809 \pm 9	1 ⁱ	$\frac{1}{2}^-$, $\frac{3}{2}^-$, $\frac{5}{2}^-$
	11.2				1 ^j	$\frac{1}{2}^-$, $\frac{3}{2}^-$, $\frac{5}{2}^-$

A: (1965AL19, 1965WA16, 1966AL18).

^a See also (1965AL19, 1965WA03).

^b (1952GI01, 1957WA01).

^c (1955SH28: see (1958WA1C)).

^d Isotropic: no clear stripping pattern.

^e (1952GI01, 1955SH28, 1956GR37, 1958WA1C).

^f (1955SH28, 1956GR37).

^g (1956GR37): (1957WA01) find a possible $l = 0$ component.

^h (1955SH28, 1957WA01).

ⁱ (1969PH02): absolute spectroscopic factors are also given.

^j (1956GR37).

^k (1961GO03). (This footnote is not labeled in the table content.)

Table 15.16: $^{14}\text{N}(d, p)^{15}\text{N}$ angular distribution studies ^a

E_d (MeV)	Distribution of proton groups	Refs.
0.5 – 0.8	p_0, p_1, p_2	(1957SJ68, 1961SJ1B)
1 – 2.2	$p_0, p_3 \rightarrow p_7$	(1967BE09)
1 – 3.2	$p_4 \rightarrow p_7$	(1969BE08)
1.1 – 3.1	$p_{1+2}, p_3, p_4, p_5, p_6$	(1969GO14)
1.3, 4.5	p_{1+2}, p_3	(1961GO03, 1965FI05, 1966GA09)
1.3 – 5.5	p_0	(1962GO21, 1965FI05, 1966GA09)
1.4 – 3	p_0, p_3	(1969BE08)
1.4 – 3.2	$p_0, p_1 + p_2$	(1961KA05)
2	$p_1 \rightarrow p_5, p_7$	(1962RO13)
7, 8, 9	$p_{3 \rightarrow 8}, p_{10+11}, p_{12}, p_{16}, p_{19}, p_{20}$	(1969PH02)
12	p_0	(1967SC29)
13.8, 15.2	p_0	(1961MO13)
16	p_0	(1960MO03)
27	p_0	(1962ER03)

^a See (1959AJ76) for earlier references.

See also (1961JA23, 1961TE02, 1962ST17, 1963GO1L, 1965HE1B), (1959BO1C, 1960BE1B, 1961KO1E, 1963TA1A, 1964BA1V, 1964ST1J, 1966HO1D; theor.), (1959AJ76) and ^{16}O in (1971AJ02).

$$40. \ ^{14}\text{N}(t, d)^{15}\text{N} \quad Q_m = 4.577$$

The angular distribution of the deuterons corresponding to the ground state of ^{15}N has been measured at $E_t = 1.50, 1.83$ and 1.98 MeV. The cross section at $E_t = 2$ MeV is 48 mb: this large value and the energy and angular behavior of the differential cross section suggest that this reaction may proceed by a cluster exchange process (1964SC09).

$$41. \ ^{14}\text{N}(\alpha, ^3\text{He})^{15}\text{N} \quad Q_m = -9.743$$

At $E_\alpha = 56$ MeV, the angular distribution of the ground state ^3He particles has been measured by (1969GA11) and analyzed by DWBA: the ratio of the $(\alpha, ^3\text{He})$ and (α, t) cross sections at this energy is 1.50 ± 0.15 . See also (1968GA1C).

42. $^{14}\text{N}(^9\text{Be}, ^8\text{Be})^{15}\text{N}$ $Q_m = 9.170$

The lifetime of $^{15}\text{N}^*(5.27)$ is ≈ 1 psec (1969NI09): see Table 15.7 and (1967BI11). See also (1963HO1E).

43. $^{14}\text{N}(^{11}\text{B}, ^{10}\text{B})^{15}\text{N}$ $Q_m = -0.621$

See (1967PO13). See also (1969BR1D).

44. $^{14}\text{N}(^{14}\text{N}, ^{13}\text{N})^{15}\text{N}$ $Q_m = 0.282$

Angular distributions of the transition to the ground state of ^{15}N have been measured for $E(^{14}\text{N})(\text{cm}) = 5.5$ to 16 MeV (1961TO07, 1964JO1A, 1965BE1B, 1965HI1A). See also (1961TO01). Below ≈ 6.5 MeV, the tunneling theory of neutron transfer gives a good account of the data. At higher energies, nuclear absorption of the incident ^{14}N ions occur (1965HI1A, 1966GA04). See also (1968GA03). For reviews of the work on this reaction, see (1964FL1D, 1967DA1E, 1967VO1A, 1969BR1D). For discussions of relevant theories, see (1962BR1G, 1963BR1G, 1964BR1M, 1964GR1G, 1965BR1F, 1966BU1B, 1967BR1M, 1967BR1Q, 1967PE1D, 1968MA2G, 1968NA1F, 1969KA1G).

45. $^{14}\text{N}(^{19}\text{F}, ^{18}\text{F})^{15}\text{N}$ $Q_m = 0.405$

See (1968GA03).

46. $^{15}\text{C}(\beta^-)^{15}\text{N}$ $Q_m = 9.773$

The β^- decay takes place to $^{15}\text{N}^*(0, 5.30, 7.30, 8.31, 9.05)$: see Table 15.2.

Measurements of γ -ray energies give $E_\gamma = 5299.03 \pm 0.43$ keV (1967CH19), 8315 ± 6 and 9048 ± 4 keV (1966AL12). See also ^{15}C .

47. $^{15}\text{N}(\gamma, n)^{14}\text{N}$ $Q_m = -10.835$

See (1967ZH1A, 1968YA1E, 1969ZH1A).

48. $^{15}\text{N}(\gamma, p)^{14}\text{C}$ $Q_m = -10.208$

The integrated cross section for transitions to $^{14}\text{C}(0)$ for E_γ up to 30.5 MeV = 22 ± 3 MeV·mb, assuming an isotropic angular distribution. Pronounced maxima are observed at $E_\gamma = 19.5, 20.4, 22.7$ and 24.5 MeV. In addition a “pigmy” resonance at $E_\gamma = 15.2$ MeV and less pronounced structures at $E_\gamma = 13.6$ and 17.0 MeV are also observed (1964KO10). See also (1963FI04) and (1967ZH1A, 1968YA1E, 1969UB1C, 1969ZH1A).

49. $^{15}\text{N}(e, e)^{15}\text{N}$

Using the harmonic oscillator function, analysis of the scattering of 250 and 400 MeV electrons gives, respectively, $r_{\text{rms}} = 2.68 \pm 0.05$ fm and 2.63 ± 0.05 fm for ^{15}N (1968DA1Q). At $E_e = 50-57$ MeV, excitation of $^{15}\text{N}^*(6.32)$ gives $\Gamma_\gamma^0(\text{M1}) = 3.4 \pm 0.7$ eV and $\Gamma_\gamma^0(\text{E2}) = 0.06 \pm 0.02$ eV: $|\delta(\text{E2/M1})| = 0.13 \pm 0.03$ (1968BE14).

50. $^{15}\text{N}(n, n)^{15}\text{N}$

See ^{16}N .

51. $^{14}\text{N}(p, p)^{15}\text{N}$

The angular distribution of elastically scattered protons has been measured at $E_p = 39.8$ MeV (1969SN04). See also (1963NA1C).

52. $^{15}\text{N}(^3\text{He}, ^3\text{He})^{15}\text{N}$

At $E(^3\text{He}) = 39.8$ MeV, a number of inelastically scattered ^3He groups are observed corresponding to states in ^{15}N : see Table 15.17. Angular distributions were obtained for a number of these, and were analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture (1969BA06). See also (1968BA1E). (1969BO13) measured the angular distribution of elastically scattered ^3He particles at $E(^3\text{He}) = 11$ MeV.

53. $^{15}\text{N}(\alpha, \alpha)^{15}\text{N}$

Table 15.17: ^{15}N levels from $^{15}\text{N}(^3\text{He}, ^3\text{He}')$ and $^{15}\text{N}(\alpha, \alpha')$

$^{15}\text{N}^* \text{ }^{\text{a}}$ (MeV \pm keV)	$L \text{ }^{\text{a}}$	$^{15}\text{N}^* \text{ }^{\text{b}}$ (MeV)	$L \text{ }^{\text{b}}$
0		0	
5.28 ± 30	3	$5.27 + 5.30$	3
6.32	2	6.32	2
7.15	3	7.16	^c
7.30	1	7.30	1
7.56	3	7.57	3
8.31	1	8.31	^c
8.57	1	8.58	^c
9.17 ± 30		9.16	
9.79 ± 40		9.83	
10.03 ± 40		10.07	
10.71 ± 40			
11.34 ± 40			
11.92 ± 40			
12.52 ± 40			
14.12 ± 40			
15.11 ± 40			

^a ($^3\text{He}, ^3\text{He}'$): (1969BA06).

^b (α, α') (1966HA19). The E_x were determined.

^c Weakly excited. See also (1965BU05, 1969BA06).

The surface thickness $a = 0.36$ fm, as determined from analysis of the scattering of 44 MeV α -particles from ^{15}N (1968FA1A). At $E_\alpha = 40.5$ MeV, a number of particle groups have been observed, and angular distributions have been measured: see Table 15.17 (1966HA19). See also (1965BU05), $B(E2)_\downarrow / e^2 = 4.9 \text{ fm}^4$ for $^{15}\text{N}^*(6.32)$: $B(E2)_\downarrow / e^2 = 60 \text{ fm}^6$ for both $^{15}\text{N}^*(5.27, 7.57)$ (1966HA19). See also (1969BA06).

$$54. \ ^{15}\text{O}(\beta^+)^{15}\text{N} \quad Q_m = 2.760$$

See ^{15}O .

$$55. \text{ (a) } ^{16}\text{O}(\gamma, p)^{15}\text{N} \quad Q_m = -12.126$$

$$\text{ (b) } ^{16}\text{O}(e, ep)^{15}\text{N} \quad Q_m = -12.126$$

Over the giant resonance region in ^{16}O , the decay takes place to the odd parity states of ^{15}N at $E_x = 0$ and 6.32 MeV as well as to both of the even parity states at $E_x = 5.27$ and 5.30 MeV. The branching ratios are functions of the excitation energy in ^{16}O and of the authors: see (1965DE24, 1965MA45, 1965MO13, 1966KO1G, 1966OW01, 1967CA1C, 1967CA1P, 1968BA2L, 1969MU07, 1969SH02). The results are in fair agreement with the predictions of the single-particle, single-hole theory of photoexcitation of ^{16}O , although some non-single-particle excitation appears to be necessary in some portions of the ^{16}O giant resonance: see, e.g. (1968BA2L). High-energy γ -rays have also reported from the decay of $^{15}\text{N}^*(7.30, 8.31, 9.05, 9.23)$ (1968BA2L, 1969HO1T, 1969MU07). See also the review in (1968SC1B), (1957JO20, 1959BR69, 1963SC32, 1968TU02, 1969FR20, 1969UL01) and (1968ZH1B; theor.). The τ_m of $^{15}\text{N}^*(5.27) \gg 0.1$ psec; τ_m for $^{15}\text{N}^*(6.33, 9.23) < 0.1$ psec (1969MU07): see also Table 15.7.

For reaction (b), see (1962DO1A).

$$56. \ ^{16}\text{O}(n, d)^{15}\text{N} \quad Q_m = -9.901$$

Angular distribution of the deuterons corresponding to the ground state of ^{15}N have been determined at $E_n = 14.4$ MeV (1964PA11, 1965VA05) and at 14 MeV (1963GA10). See also (1965DI1E; theor.).

$$57. \ ^{16}\text{O}(p, 2p)^{15}\text{N} \quad Q_m = -12.126$$

At $E_p = 460$ MeV, the summed proton spectrum shows two peaks corresponding to the knowckout of $p_{1/2}$ and $p_{3/2}$ protons with binding energies of 12.4 and 19.0 MeV, respectively [$^{15}\text{N}^*(0, 6.32)$] (1966TY01). See also the discussions in (1963CL1B, 1963RI1B), (1962FO03, 1967FU1A), (1963BE1A, 1963BE42; theor.) and (1959AJ76).

58. $^{16}\text{O}(\text{d}, ^3\text{He})^{15}\text{N}$ $Q_{\text{m}} = -6.632$

Angular distributions of the ^3He groups have been measured at $E_{\text{d}} = 20$ MeV (1969PU04: to $^{15}\text{N}^*(0, 5.27, 5.30)$), 28 MeV (1968GA13: to $^{15}\text{N}(0)$), 34.4 MeV (1967HI06: to $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32)$), 52 MeV (1969KA1A, 1969KA1W: to $^{15}\text{N}(0)$), and 82 MeV (1969DO04: to $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32)$) and analyzed by DWBA. The ^3He group to $^{15}\text{N}^*(5.2)$ does not show a stripping pattern (1969PU04). See also (1968BA2J; theor.).

59. $^{16}\text{O}(\text{t}, \alpha)^{15}\text{N}$ $Q_{\text{m}} = 7.688$

Angular distributions have been measured at $E_{\text{t}} = 0.9$ to 1.7 MeV (1967KO1G; α_0), 1.15 to 1.95 MeV (1959JO32; α_0), 1.5 to 3.0 MeV (1966SE1D; α_0), and 13 MeV (1965AJ01; $\alpha_0, \alpha_{1+2}, \alpha_3$). A γ -ray with $E_{\gamma} = 5.272 \pm 0.010$ MeV has been observed in this reaction: see Table 15.7 (1965AL19). See also (1959AJ76).

60. $^{16}\text{O}(^{14}\text{N}, ^{15}\text{O})^{15}\text{N}$ $Q_{\text{m}} = -4.833$

See (1963TO1D, 1969BR1D).

61. $^{17}\text{O}(\text{d}, \alpha)^{15}\text{N}$ $Q_{\text{m}} = 9.803$

See (1954PA39).

62. $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$ $Q_{\text{m}} = 3.981$

Angular distributions of ground state α -particles are reported at $E_{\text{p}} = 0.84$ to 2.00 MeV (1961CA02) and at 7.9, 10.6 and 13.3 MeV (1964EC03). Angular correlation measurements lead to $J = \frac{5}{2}, \frac{3}{2}, (\frac{1}{2}), (\frac{3}{2})$ for $^{15}\text{N}^*(5.28, 6.32, 8.31, 8.58)$ (1965WA06), $J = \frac{5}{2}, \frac{5}{2}, \frac{3}{2}, \frac{7}{2}$ for $^{15}\text{N}^*(5.28, 7.16, 7.30, 7.57)$ (1966HA30), $J = \frac{3}{2}$ for $^{15}\text{N}^*(6.32)$ (1966LO02). The M2/E1 mixing ratio of the 7.30 \rightarrow g.s. transition indicates an unusually large retardation of an E1 transition in a non-self-conjugate nucleus (1966HA30). $J = \frac{5}{2}$ for $^{15}\text{N}^*(5.27)$ and the mixing parameter fix $J = \frac{5}{2}$ for $^{15}\text{N}^*(10.54)$ which fed the first excited state of ^{15}N (1959HE1D) in the $^{14}\text{C}(\text{p}, \gamma)^{15}\text{N}$ reaction (1965WA06).

See also (1960CL02, 1961LO10, 1964AM1A, 1964MA25, 1964MA57, 1964SC01), (1959AJ76) and ^{19}F in (1959AJ76, 1972AJ02).

$$63. {}^{19}\text{F}(\gamma, \alpha){}^{15}\text{N} \quad Q_{\text{m}} = -4.011$$

See (1965HA1G).

$$64. \text{(a) } {}^{19}\text{F}(\text{p}, \text{p}\alpha){}^{15}\text{N} \quad Q_{\text{m}} = -4.011$$

$$\text{(b) } {}^{19}\text{F}(\alpha, 2\alpha){}^{15}\text{N} \quad Q_{\text{m}} = -4.011$$

For reaction (a) see (1962FO03). For reaction (b) see (1963LA02).

$$65. {}^{19}\text{F}(\text{d}, {}^6\text{Li}){}^{15}\text{N} \quad Q_{\text{m}} = -2.539$$

Angular distributions of the ${}^6\text{Li}$ ions corresponding to ${}^{15}\text{N}(0)$ have been measured at $E_{\text{d}} = 9.0$ to 12.5 MeV (1967DE14), 14.5 MeV (1964DA1B), 14.9 MeV (1966DE09) and 21 MeV (1965SL1C). Attempts have been made to fit the data with DWBA: see references above and (1963DR1B). See also (1964BL1C).

$$66. {}^{19}\text{F}({}^{14}\text{N}, {}^{18}\text{F}){}^{15}\text{N} \quad Q_{\text{m}} = 0.405$$

See (1965WI1A).

¹⁵O
(Figs. 11 and 12)

GENERAL:

Model calculations: (1960TA1E, 1960TA1C, 1963CO12, 1963KU1B, 1964AL1L, 1964AM1D, 1964BR1H, 1964RI1A, 1965CO25, 1965GI1B, 1965GR1H, 1965GU1A, 1965HU1D, 1966BO1R, 1966EL08, 1966RI1F, 1966SO05, 1967BO1T, 1967EL03, 1968DE13, 1968EL1A, 1968HO1H, 1968MA2B, 1968SH08, 1968WO1C, 1968ZH1B, 1968ZU02, 1969DE16, 1969EL1B, 1969GU1M, 1969SA1J).

General calculations and reviews: (1964EV1A, 1967FA1A, 1967NE1D, 1968BI1C).

Electromagnetic transitions: (1965RO1N, 1966PO1I, 1966RO1P, 1966WA1E, 1967KU1E, 1967PO1J, 1967WA1C, 1968BI1C, 1968SH08, 1968ZH1B, 1968ZH06, 1969KH1C, 1969ZH1A).

Other: (1966WA1K, 1967AU1B, 1969FO1D, 1969HA1G).

Ground state: $J = \frac{1}{2}$ (1963CO17); $\mu = 0.71898$ nm (1963CO17, 1964LI14, 1967CO1D).

See also (1964ST1B, 1965MA1T, 1966MA1V, 1967NE1D, 1967SH14, 1968PE16, 1968RO1E, 1969FU1I).

1. $^{15}\text{O}(\beta^+)^{15}\text{N}$ $Q_m = 2.760$

Reported half-lives are listed in Table 15.19 (1954KL36, 1955BA83, 1957KI22, 1957PE12, 1959KI99, 1960JA12): the weighted mean is 122.24 ± 0.16 sec. See also (1963CS02, 1963VA31). Using this value for $\tau_{1/2}$ and Q_m , $\log ft = 3.643$. See also (1968BA42) and (1965GA1D, 1966MI1F, 1967AM1H, 1968SH08, 1969LE1D, 1969SU15; theor.).

2. $^7\text{Li}(^{14}\text{N}, ^6\text{He})^{15}\text{O}$ $Q_m = -2.687$

See (1957AL78).

3. (a) $^{10}\text{B}(^6\text{Li}, \text{n})^{15}\text{O}$ $Q_m = 15.209$
(b) $^{10}\text{B}(^7\text{Li}, 2\text{n})^{15}\text{O}$ $Q_m = 7.957$

See (1957NO17).

Table 15.18: Energy levels of ^{15}O ^a

E_x (MeV \pm keV)	J^π	τ_m (psec) or Γ (keV)	Decay	Reactions
0	$\frac{1}{2}^-$	$\tau_{1/2} = 122.24 \pm 0.16$ sec	β^+	1, 2, 3, 4, 6, 7, 8, 9, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29
5.181 \pm 5	$\frac{1}{2}^+$	$\tau_m < 0.1$ psec	γ	8, 9, 16, 17, 22, 24, 25, 26
5.24151 \pm 0.52	$\frac{5}{2}^+$	3.2 \pm 0.5	γ	8, 9, 16, 17, 20, 21, 22, 24, 25, 26
6.176 \pm 3	$\frac{3}{2}^-$	< 0.047	γ	8, 9, 16, 17, 21, 22, 24, 25, 26
6.788 \pm 4	$\frac{3}{2}^+$	< 0.028	γ	8, 9, 16, 17, 26
6.859 \pm 1	$\frac{5}{2}^+$	0.10 \pm 0.06	γ	8, 9, 16, 17, 21, 26
7.2760 \pm 0.6	$\frac{7}{2}^+$		γ	8, 16, 17, 21, 24, 26
7.5522 \pm 0.5	$\frac{1}{2}^+$	$\Gamma = 1.7 \pm 0.5$ keV	γ, p	9, 16, 17, 21
8.2833 \pm 1.5	$\frac{3}{2}^+$	3.6 \pm 0.7	γ, p	9, 17, 21
8.739 \pm 6	$\frac{1}{2}^+$	32	γ, p	9
8.9180 \pm 1.4	$\frac{3}{2}$	3.7 \pm 1	γ, p	8, 9, 21
8.9781 \pm 1.6	$(\frac{1}{2}, \frac{3}{2})^-$	3.9 \pm 0.4	γ, p	8, 9, 21
9.483 \pm 3	$\frac{5}{2}^-$	10.1 \pm 0.5	γ, p	8, 9, 21
9.50 \pm 40	$\frac{3}{2}^+(\frac{1}{2})^+$	280 \pm 24	γ, p	9
9.606 \pm 1.8	$\frac{3}{2}^-$	8.8 \pm 0.5	γ, p	8, 9, 21, 24
9.660 \pm 4	$\frac{1}{2}^-$	2 \pm 1	p	8, 10, 21
9.72 \pm 50	$(\frac{1}{2}, \frac{3}{2})^+$	1190 \pm 50	γ, p	9
10.278 \pm 8	+	16 \pm 4	p	10, 21
10.46 \pm 10		47	γ, p	9, 10, 21, 24
10.91 \pm 15	$\frac{7}{2}^+$	91	p	10, 21
10.939 \pm 7	$\frac{1}{2}^+$	84	γ, p	9, 10, 21
11.023 \pm 7	$\frac{1}{2}^-$	21	p	10
11.15 \pm 15		< 10	p	10
11.20 \pm 15	$(\frac{1}{2}, \frac{3}{2})^+$	36	γ, p	9, 10, 21
11.5 \pm 100	$T = \frac{3}{2}$			8
11.56 \pm 15		< 10	p	10
11.57		140	γ, p	10
11.57 \pm 15	$\frac{5}{2}^-$	25	p	9

Table 15.18: Energy levels of ^{15}O ^a (continued)

E_x (MeV \pm keV)	J^π	τ_m (psec) or Γ (keV)	Decay	Reactions
11.61 \pm 15	$\frac{3}{2}^- (\frac{1}{2}^-)$	25	p	10
11.71 \pm 15		< 10	p	10, 21
11.75 \pm 15	$\frac{5}{2}^+$	80	p	10
11.846 \pm 10	$\frac{5}{2}^-$	50	p	10
11.98 \pm 15	$\frac{5}{2}^-$	30	p	10
12.12 \pm 15	$\frac{5}{2}^+$	160	p	10
12.47 \pm 15	$\frac{5}{2}^- (\frac{3}{2}^-)$	60	p	10
12.8		\approx 230	γ , p	9
12.82	+	9	p	10
13.00	$\frac{5}{2}$	28	p, ^3He , α	5, 10, 15
13.1			p, d, ^3He , α	5
13.4	$\frac{3}{2}^+$	broad	(p, α)	10, 15
13.49	$(\frac{3}{2}^+)$		p	10
13.60	$\frac{5}{2}^+$		p, α	15
13.70	$\frac{3}{2}^-$		p	10
13.79	$\frac{3}{2}^-$		n, p, ^3He , α	5, 10, 15, 21
13.87		\approx 140	γ , p	9
14.03 \pm 40	$(\frac{1}{2}, \frac{3}{2})^-$	160 \pm 20	n, p, ^3He	5
14.17	$\frac{5}{2}^-$		p, α	15
14.27 \pm 40	$\frac{1}{2}^+$	340 \pm 30	n, p, ^3He , α	5, 10, 11, 15
14.34	$\frac{5}{2}^+$		p, α	15
14.460 \pm 10	$\frac{5}{2}^+$	100 \pm 10	n, p, ^3He , α	5, 11
14.69 \pm 40		170 \pm 30	n, p, ^3He	5, 11
14.95 \pm 40		400 \pm 25	n, p, ^3He	5, 11
15.43 \pm 10		65 \pm 15	p, ^3He , α	5
15.56 \pm 40	$\frac{1}{2}^+$	80 \pm 25	p, ^3He , α	5
15.84 \pm 50	$(\frac{1}{2}, \frac{3}{2})^-$	350	n, p, ^3He , α	5, 11
16.04			^3He , α	5
16.09			n, ^3He , α	5
16.19			^3He , α	5
16.43 \pm 50	$\frac{1}{2}^+$	170	^3He , α	5


Table 15.18: Energy levels of ^{15}O ^a (continued)

E_x (MeV \pm keV)	J^π	τ_m (psec) or Γ (keV)	Decay	Reactions
16.48 \pm 50		560 \pm 90	n, p	11
16.77 \pm 50		200	n, ^3He , α	5
17.50 \pm 50	$(\frac{1}{2}, \frac{3}{2})^-$	600	n, p, ^3He , α	5
17.99 \pm 50	$(\frac{1}{2}, \frac{3}{2})^-$	200	^3He , α	5
18.22 \pm 50			n, p, ^3He	5
19.02 \pm 50			n, ^3He	5
19.90 \pm 50			n, p, ^3He	5
\approx 21		broad	γ , p	9

^a See also Tables 15.22 and 15.25.



See (1963HO1E).



Excitation functions for these reactions have been measured over a wide range of energies: see Table 15.20. Observed resonances are displayed in Table 15.21. For discussions of angular distribution measurements which have been measured at many energies over the range displayed in Table 15.20, see the writeups of ^7Be in (1966LA04), ^{11}C and ^{12}C in (1968AJ02), and ^{13}N , ^{14}N , and ^{14}O . See also (1964DE1E).

Consideration of the relative yields at $E(^3\text{He}) = 1.21$ and 2.15 MeV lead to the tentative J^π assignments given in Table 15.21 for $^{15}\text{O}^*(13.00, 13.79)$ (1957BR18). The assignments for $^{15}\text{O}^*(14.03, 14.27, 14.46)$ are derived from analyses of the ^3He elastic scattering, particle and $\gamma_{6.44}$ angular distributions, and the total cross sections of the n_0 and p_1 groups (1964KU05). Above $E(^3\text{He}) \approx 5$ MeV, the excitation functions show broad maxima (see Table 15.21) but it is not

Table 15.19: The half-life of ^{15}O

$\tau_{1/2}$ (sec)	Refs.
123.4 ± 1.3	(1954KL36)
121 ± 3	(1955BA83)
120 ± 2	(1957KI22)
123.95 ± 0.5	(1957PE12)
124.1 ± 0.5	(1959KI99)
122.1 ± 0.1	(1960JA12)
122.24 ± 0.16	weighted average

clear that they correspond to excited states in ^{15}O : for instance the maxima in the yields of n_0 (1964DE1C), p_6 , d_0 , α_0 and α_1 (1960HI07) are not correlated. See, however, (1969WE08). The structures observed in the excitation functions for $p_0 \rightarrow p_9$ at $E(^3\text{He}) = 4.6$ to 11 MeV are attributed to quasi-giant resonances involving single proton orbitals coupled to excited ^{14}N core configurations (1969HA49). Above $E(^3\text{He}) \approx 12$ MeV, the excitation functions do not show clear resonant behavior: see, e.g. (1967GR1L, 1968FO06, 1969FO02).

Broad resonance-like structures in the yield of protons are reported at $E(^3\text{He}) = 7.0, 7.8, 9.6, 10.6, 13.4$ and 14.6 MeV [$\Gamma \approx 1$ MeV] (1970SP1E).

Over the range $E(^3\text{He}) = 6.5 \rightarrow 11$ MeV, the ratio $d\sigma_n/d\sigma_{p_1}$ [to $^{14}\text{O}_{\text{g.s.}}$ and the first $T = 1$ state of ^{14}N], determined at 10° , is approximately constant and ≈ 2 , as would be expected if isospin were conserved and if Coulomb scattering and Q -value differences can be neglected (1965FU16). (1968LA19) find that the compound nucleus level overlap parameter Γ/D_0 and the spin cut-off parameter σ as obtained from elastic scattering and from the statistical model analysis of three proton groups [$E(^3\text{He}) = 5.29 - 5.50$ MeV] are constant. See also (1970CA1G).

For a survey of the energies at which polarization measurements have been made, see Table 15.20. (1968HU1A) find that at $E(^3\text{He}) = 36$ and 42 MeV, the ^3He polarization results are such that optical model calculations require an optical spin-orbit potential depth ≤ 3.5 MeV. See also (1959AJ76).

$$6. \ ^{12}\text{C}(\alpha, n)^{15}\text{O} \quad Q_{\text{m}} = -8.507$$

$$E_{\text{thresh.}} = 11.341 \pm 0.015 \text{ (1963NE05).}$$

Angular distributions of neutrons corresponding to the ground state of ^{15}O have been measured at $E_\alpha = 14$ MeV (1965AL1J) and at 20.0 to 21.8 MeV (1963KO03). See also ^{16}O in (1971AJ02).

Table 15.20: Recent yield and polarization measurements in $^{12}\text{C} + ^3\text{He}$ ^a

 (a) *Excitation functions* ^b

$E(^3\text{He})$	Yield of	Refs.
1.0 – 1.8	$p_0, p_1, p_2, d_0, \alpha_0$	(1965GR1R)
1.4 – 3.5	n	(1964KU06)
1.4 – 6.0	n, α	(1966CI01)
1.4 – 11.45	n_0	(1964OS01)
1.75 – 5.2	α_0	(1963LU05)
2 – 5.5	^{11}C	(1966GO1E)
1.8 – 5.4	$p_0 \rightarrow p_9, ^3\text{He}, \alpha_0$	(1964KU05)
1.8 – 5.5	n	(1964DI02)
2 – 4	$p_0 \rightarrow p_8$	(1967CL1C)
2.0 – 5.7	n_0	(1961TO03)
2 – 6.2	$^3\text{He}, \alpha_0, \alpha_1$	(1966BL01, 1966SC12)
2.4 – 3.3	$^3\text{He}, \alpha_0$	(1970JA1E)
≈ 2.5	p_2	(1963LU01, 1963LU1F)
2.7 – 5.4	$\gamma_{6.44}$	(1964KU05)
3.50 – 3.91	^3He	(1967SC27)
4.0 – 5.4	α_1	(1964KU05)
4.0 – 6.0	$p_2 \rightarrow p_6$	(1966BL01)
4.4 – 8.2	$^3\text{He}, \alpha_0$	(1968WE15, 1969WE03)
4.5 – 6.0	α_0	(1969WE03)
4.6 – 11.0	$p_0 \rightarrow p_9$	(1969HA49)
5 – 30	$^{11}\text{C}, ^{13}\text{N}$	(1965BR42)
5.2 – 7.8	α_1	(1968WE15, 1969WE03)
5.5 – 11	n_0	(1964DE1C)
5.7 – 10.23	$p_6, d_0, \alpha_0, \alpha_1$	(1960HI07)
6 – 17	p	(1970SO1G)
6 – 24	$^7\text{Be}, ^{11}\text{C}, ^{13}\text{N}$	(1962CO31)
8 – 10	$^3\text{He}, \alpha_0$	(1966SC22)
9 – 29	^7Be	(1965EN01)
12.0 – 18.6	$\alpha_0 \rightarrow \alpha_7$	(1967GR1L)
12 – 18.6	$d_0, d_1, d_{2+3}, ^3\text{He}(0), ^3\text{He}(1)$	(1968FO06, 1969FO02)

Table 15.20: Recent yield and polarization measurements in $^{12}\text{C} + ^3\text{He}$ ^a
(continued)

$E(^3\text{He})$	Yield of	Refs.
21 – 31.2	$^{11}\text{C}, ^{13}\text{N}$	(1959MA1D)
21 – 31	^7Be	(1965MA1V)

(b) Polarization measurements ^b

2.24 – 3.70	n_0	(1967SC27)
2.48 – 3.15	p_0	(1967KR1F)
4.1 – 5.9	n	(1967SO1B)
4.5 – 5.5	p_0, p_1	(1967MA1M)
18	^3He	(1969LU1C)
29	^3He	(1961AG1A, 1964BU1D, 1965FR1E)
31.6	^3He	(1969EN03)
36, 42	^3He	(1968HU1A)

^a See also (1959AJ76, 1961FO02).

^b See also (1964BR1G, 1964DI1C, 1965SC1D, 1966HA21, 1966MA1R, 1968PA1F, 1969PA1C).

^c See also (1965PE1H, 1966CA1H).

7. (a) $^{12}\text{C}(^6\text{Li}, t)^{15}\text{O}$ $Q_m = -3.721$
 (b) $^{12}\text{C}(^{14}\text{N}, ^{11}\text{B})^{15}\text{O}$ $Q_m = -8.664$

For reaction (a) see (1969GI1B); for reaction (b) see (1969BR1G).

8. $^{13}\text{C}(^3\text{He}, n)^{15}\text{O}$ $Q_m = 7.125$

Angular distributions of neutrons corresponding to the ground state of ^{15}O have been measured at $E(^3\text{He}) = 1.70$ to 5.35 MeV (1965DI07), 2.6 , 2.8 and 3.1 MeV (1961JO07, 1961JO24), 2.66 MeV (1961DU1B, 1963DU12: also n_3), 5.0 and 6.2 MeV (1969HO1F: also to $^{15}\text{O}^*(6.18, 6.86, 8.92, 8.98, 9.50, 9.60, 9.66)$ and $7.8, 8.6$ and 10.1 MeV (1964DE1C). DWBA analyses have been made: (1969HO1F) find $L = 0$ for $^{15}\text{O}^*(0, 8.92, 8.98, 9.66)$ and $L = 2$ for $^{15}\text{O}^*(6.18, 9.49, 9.60)$. At $E(^3\text{He}) = 11.6$ MeV, a neutron group assigned to a $T = \frac{3}{2}$ state at 11.5 ± 0.10 is reported by (1969BR30).

Table 15.21: Resonances in $^{12}\text{C} + ^3\text{He}$

$E(^3\text{He})$ (MeV \pm keV)	Resonant for	E_x^a (MeV)	J^π	$\Gamma_{\text{c.m.}}$ (keV)	Refs.
1.21	p_0, p_2	13.04	$(\frac{5}{2})^-$		(1957BR18)
1.3	$p_0, p_1, p_2, p_3, d_0, \alpha_0$	12.1			(1957BR18, 1965GR1R)
2.15	n, p_0	13.79	$(> \frac{5}{2})$		(1957BR18)
2.45 ± 40	n_0, p_0, p_1, p_2, p_3	14.03	$(\frac{1}{2}^-, \frac{3}{2}^-)$	160 ± 20	(1957BR18, 1961TO03, 1964KU05, 1964KU06, 1964OS01)
2.75 ± 40	$n_0, p_1, p_2, ^3\text{He}, \alpha$	14.27	$\frac{1}{2}^+$	340 ± 30	(1957BR18, 1964KU05, 1964KU06, 1964OS01, 1966CI01, 1970JA1E)
2.990 ± 10	$n_0, p_0, p_1, p_2, p_4, p_5, p_8, ^3\text{He}, \alpha_0$	14.460	$\frac{3}{2}^+$	100 ± 10	(1958JO20, 1961TO03, 1964KU05, 1964KU06, 1964OS01, 1970JA1E)
3.28 ± 40	p_0	14.69		180 ± 40	(1964KU05)
3.60 ± 40	p_0, p_1, p_2	14.95		400 ± 25	(1958JO20, 1964KU05)
4.20 ± 10	p_5, p_6, α_0	15.43		65 ± 15	(1964KU05)
4.37 ± 40	$p_0, p_1, p_2, p_4, p_7, p_8, \alpha_0$	15.56	$\frac{1}{2}^+$	80 ± 25	(1958JO20, 1961TO03, 1964KU05, 1966CI01)
4.65 ± 50	n_0	15.79			(1961TO03, 1964DI02, 1964KU05, 1964OS01)
4.77	α_0	15.89	$\frac{1}{2}^-, \frac{3}{2}^-$	350	(1969WE03)
4.97 ± 20	α_0	16.04			(1969WE08)
5.03 ± 20	α_0	16.09			{ (1961TO03, 1964DI02, 1964KU05, 1964OS01, 1966SC12, 1969WE08)
5.15 ± 20	α_0	16.19			
5.45	α_0	16.43	$\frac{1}{2}^+$	170	(1969WE03)
5.88 ± 50	$n_0, ^3\text{He}, \alpha_0$	16.77	$\frac{1}{2}^-, \frac{3}{2}^-$	200	(1964OS01, 1966SC12, 1969WE03)
6.80 ± 50	n_0, p	17.50	$\frac{1}{2}^-, \frac{3}{2}^-$	600	(1964OS01, 1969WE03, 1970SO1G)
7.4	α_0	17.99	$\frac{1}{2}^-, \frac{3}{2}^-$	200	(1969WE03)
7.70 ± 50	n_0, p	18.22			(1964OS01, 1970SO1G)
8.70 ± 50	n_0	19.02			(1964OS01)
9.80 ± 50	n_0, p	19.90			(1964OS01, 1970SO1G)

^a See also text.

Table 15.22: Radiative decays in ^{15}O

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Γ_γ ^a (eV)	Refs.
6.18	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100		(1965WA16)
		5.18	$\frac{1}{2}^+$	< 2.5		(1965WA16)
		5.24	$\frac{5}{2}^+$	< 2.5		(1965WA16)
6.79	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	100		(1965WA16, 1968GI11)
		5.18	$\frac{1}{2}^+$	< 6		(1965WA16)
				< 3		(1968GI11)
		5.24	$\frac{5}{2}^+$	< 6		(1965WA16)
		< 3		(1968GI11)		
6.86	$\frac{5}{2}^+$	6.18	$\frac{3}{2}^-$	< 7		(1965WA16)
		0	$\frac{1}{2}^-$	< 10		(1965WA16)
				< 12		(1968GI11)
		5.18	$\frac{1}{2}^+$	< 15		(1965WA16)
				< 4		(1968GI11)
		5.24	$\frac{5}{2}^+$	100		(1965WA16, 1968GI11)
7.28	$\frac{7}{2}^+$	6.18	$\frac{3}{2}^-$	< 0.4		(1965WA16)
		0	$\frac{1}{2}^-$	< 30		(1965WA16)
				< 12		(1968GI11)
				3.8 ± 1.2		(1969KU01)
		5.18	$\frac{1}{2}^+$	< 10		(1965WA16)
				< 4		(1968GI11)
		5.24	$\frac{5}{2}^+$	100		(1965WA16, 1968GI11)
				96.2 ± 1.2		(1969KU01)
7.55	$\frac{1}{2}^+$	6.18	$\frac{3}{2}^-$	< 2		(1965WA16)
		0	$\frac{1}{2}^-$	≈ 3		(1960TA17)
				3.5 ± 0.5		(1963HE11)
		5.18	$\frac{1}{2}^+$	16.2 ± 2		(1960TA17)
				15.8 ± 0.6		(1963HE11)
		6.18	$\frac{3}{2}^-$	57.9 ± 0.6		(1960TA17)
				57.4 ± 0.6		(1963HE11)
		6.79	$\frac{3}{2}^+$	22.9 ± 2		(1960TA17)
				23.3 ± 0.6		(1963HE11)

Table 15.22: Radiative decays in ^{15}O (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Γ_γ ^a (eV)	Refs.
8.28	$\frac{3}{2}^+$	6.86	$\frac{5}{2}^+$	b		
		0	$\frac{1}{2}^-$	53.8 ± 0.25	0.531	(1966EV01)
		5.24	$\frac{5}{2}^+$	42.7 ± 0.5	0.405	(1966EV01)
		6.18	$\frac{3}{2}^-$	2.2 ± 0.6	0.021	(1966EV01)
8.74 ^c	$\frac{1}{2}^+$	6.86	$\frac{5}{2}^+$	1.2 ± 0.3	0.011	(1966EV01)
		5.18	$\frac{1}{2}^+$	67	0.32	(1966EV01)
		6.18	$\frac{3}{2}^-$	33	0.16	(1966EV01)
8.92	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	21 ± 2	0.056	(1966EV01)
		5.18	$\frac{1}{2}^+$	23 ± 6	0.094	(1966EV01)
		6.18	$\frac{3}{2}^-$	30 ± 3	0.094	(1966EV01)
		6.86	$\frac{5}{2}^+$	26 ± 3	0.069	(1966EV01)
8.98	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	93.4	0.74	(1966EV01)
		5.18	$\frac{1}{2}^+$	5.9	0.046	(1966EV01)
9.49	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	86	2.1	(1967EV02)
		5.24	$\frac{5}{2}^+$	6.5	0.15	(1967EV02)
		6.18	$\frac{3}{2}^-$	0.7	0.22	(1967EV02)
		6.86	$\frac{5}{2}^+$	3.4	0.08	(1967EV02)
		7.28	$\frac{7}{2}^+$	5.1	0.11	(1967EV02)
		9.50	$\frac{3}{2}^+(\frac{1}{2}^+)$	0	$\frac{1}{2}^-$	≈ 100
9.60	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	79	4.0	(1967EV02)
		5.24	$\frac{5}{2}^+$	19	1.0	(1967EV02)
		6.18	$\frac{3}{2}^-$	2	0.1	(1967EV02)

^a (1951DU08, 1966EV01). See also (1959HE47).

^b Intensity < 25% of transition to $^{15}\text{O}^*(6.79)$ (1959PO79).

^c See also (1959HE47).

Branching ratios measured by (1965WA16) are listed in Table 15.22. The measured E_γ lead to $E_x = 6.180 \pm 0.004$, 6.857 ± 0.0032 and 7.284 ± 0.007 MeV (1965WA16).

See also (1964BR13, 1968HIIJ, 1968ST19), (1965SH1E, 1966SH1F; theor.) and ^{16}O in (1971AJ02).

$$9. \ ^{14}\text{N}(p, \gamma)^{15}\text{O}$$

$$Q_m = 7.293$$

Table 15.23: Resonances in $^{14}\text{N} + \text{p}$

E_p (keV)	Γ_{lab} (keV)	$\omega\Gamma_\gamma$ (eV)	Particles out	J^π	E_x (MeV)	Refs.
278.1 ± 0.4	1.7 ± 0.5	0.014	γ	$\frac{1}{2}^+$	7.5522	(1951DU08, 1955BA83, 1957PIZZ, 1959PO79, 1960TA17, 1963HE11)
1061.6 ± 1.4	3.9 ± 0.7	0.95	γ	$\frac{3}{2}^+$	8.2833	(1951DU08, 1956TA16, 1957HA03, 1959GA05, 1959HE47, 1959VA04, 1959VA08, 1963HE11, 1966EV01)
1550 ± 6	34	0.16	γ	$\frac{1}{2}^+$	8.739	(1951DU08, 1956TA16, 1957BO58, 1957HA03, 1966EV01)
1742.0 ± 1.2^a	4 ± 1	0.21	γ	$\frac{3}{2}^+$	8.918	(1951DU08, 1956TA16, 1957BO58, 1957HA03, 1959BA16, 1959VA08, 1963CO13, 1966EV01)
1806.4 ± 1.5^a	4.2 ± 0.4	0.52	γ	$(\frac{1}{2}, \frac{3}{2})^-$	8.9781	(1951DU08, 1956TA16, 1957BO58, 1957HA03, 1959VA08, 1963CO13, 1966EV01)
2348 ± 3	10.8 ± 0.5	2.4	γ	$\frac{3}{2}^+$	9.483	(1951DU08, 1957BO58, 1959VA08, 1967EV02, 1967LA05, 1967LA10, 1969OC1B)
2368 ± 32	300 ± 26		γ	$\frac{3}{2}^+$ ($\frac{1}{2}^+$)	9.50	(1957BO58, 1959VA08, 1967EV02, 1967LA05, 1967LA10)
2479 ± 1.7	9.4 ± 0.5	3.3	γ	$\frac{3}{2}^+$	9.606	(1951DU08, 1959VA08, 1967EV02, 1967LA05, 1967LA10, 1969OC1B)
2537 ± 4	2 ± 1		p	$(\frac{7}{2}, \frac{9}{2})^-$	9.660	(1967LA05, 1967LA10)
2600 ± 50	1270 ± 50	46	γ	$(\frac{1}{2}, \frac{3}{2})^+$	9.72	(1951DU08)
3200 ± 8	17 ± 4		p	+	10.278	(1957BO58, 1959VA08, 1967KU1M)
3390 ± 10	50		γ, p		10.46	(1957BO58, 1959VA08, 1969OC1B)
3880 ± 15	97		p_0	$\frac{7}{2}^+$	10.91	(1959BA16, 1967KU1M, 1969WE02)
3908 ± 7	90		$\gamma, \text{p}_0, \text{p}_1$	$\frac{1}{2}^+$	10.939	(1956BA34, 1969OC1B, 1969WE02)
3998 ± 7	22		p_0, p_1	$\frac{1}{2}^-$	11.023	(1956BA34, 1969WE02)
4130 ± 15	< 10		p_0		11.15	(1969WE02)
4190 ± 15	39		γ, p_0	$(\frac{1}{2}, \frac{3}{2})^+$	11.20	(1969OC1B, 1969WE02)
4575 ± 15	< 10		p_0		11.561	(1969WE02)
4580 ± 15	27		p_0	$\frac{5}{2}^-$	11.57	(1969WE02)
4580	150		γ		11.57	(1969OC1B)
4630 ± 15	27		p_0	$\frac{3}{2}^- (\frac{1}{2}^-)$	11.61	(1969WE02)
4740 ± 15	< 10		p_0		11.71	(1969WE02)
4780 ± 15	85		p_0, p_1	$\frac{5}{2}^+$	11.75	(1956BA34, 1969WE02)
4881 ± 10	54		p_0, p_1	$\frac{5}{2}^-$	11.846	(1956BA34, 1969WE02)
5020 ± 15	32		p_0	$\frac{5}{2}^-$	11.98	(1969WE02)
5180 ± 15	172		p_0, p_1	$\frac{5}{2}^+$	12.12	(1969WE02)
5550 ± 15	64		$\text{p}_0, \text{p}_1, \text{p}_2$	$\frac{5}{2}^- (\frac{3}{2}^-)$	12.47	(1969WE02)
5900	≈ 250		γ		12.8	(1969OC1B)
5920	10		p	+	12.82	(1967KU1M)
6100	30		$\text{p}_0 \rightarrow \text{p}_2, \alpha_0$	$\frac{5}{2}^+$	12.9	(1967KU1M, 1968SH11)
6600	broad		(p_2, α_0)	$\frac{5}{2}^+$	13.45	(1968SH11, 1969OC1B)
6640			$(\text{p}_0), \text{p}_2$	$(\frac{5}{2}, \frac{3}{2})^+$	13.49	(1968SH11)
6760			α_0	$\frac{5}{2}^+$	13.60	(1968SH11)
6870			p_2	$\frac{5}{2}^-$	13.70	(1968SH11)

Table 15.23: Resonances in $^{14}\text{N} + \text{p}$ (continued)

E_p (keV)	Γ_{lab} (keV)	$\omega\Gamma_\gamma$ (eV)	Particles out	J^π	E_x (MeV)	Refs.
6960			p1, p2, p4, α_0	$\frac{3}{2}^-$	13.79	(1968SH11)
7050	≈ 150		γ		13.87	(1969OC1B)
7370			α_0	$\frac{5}{2}^-$	14.17	(1968SH11)
7500	≈ 800		n, p2, α_0		14.3	(1964KU06, 1968SH11)
7550			α_0	$\frac{5}{2}^+$	14.34	(1968SH11)
7700			n		14.5	(1964KU06)
7950	170 ± 50		n		14.71	(1964KU06)
8200			n		14.94	(1964KU06)
9050			n		15.73	(1964KU06)
9400			n		16.1	(1964KU06)
9850 ± 50	600 ± 100		n		16.48	(1964KU06)


^a Comparison of analog states in ^{15}N and ^{15}O shows that there should be a (as yet unreported) state in ^{15}O at $E_x \approx 8.95$ MeV. This corresponds to $E_p \approx 1.75$ MeV, and the analysis of the $E_p = 1.74$ and 1.81 MeV resonances is therefore possible in error (E.K. Warburton, private communication).

Observed resonances are listed in Table 15.23 (1951DU08, 1955BA83, 1957PIZZ, 1959GA05, 1959HE47, 1959PO79, 1959VA04, 1959VA08, 1960TA17, 1963HE11, 1966EV01, 1967EV02, 1969OC1B). Branching ratios are displayed in Table 15.22 (1960TA17, 1963HE11, 1966EV01).

The cross section increases from $(8.5 \pm 3.7) \times 10^{-12}$ b at 100 keV to $(140 \pm 30) \times 10^{-12}$ b at 135 keV (1957LA13). Extrapolation from the $E_p = 0.28$ MeV resonance gives $S(0) = 2.75 \pm 0.50$ keV · b, with zero slope to $E_p = 0.05$ MeV (1963HE11). For astrophysical implications, see (1964FO1A, CA65, 1968DU1F). The J^π assignments shown in Table 15.23 arise from considerations of branching ratios, measurements of angular distributions of γ -rays and the studies discussed in reaction 10. Angular correlation measurements also lead to $J^\pi = \frac{3}{2}^+$ and $\frac{3}{2}^-$ for $^{15}\text{O}^*(6.79, 6.18)$. The results are also consistent with $J = \frac{1}{2}$ for $^{15}\text{O}^*(5.18)$ (1959PO79). Energies for states involved in the cascade decays are $E_x = 5.19 \pm 0.01, 6.15 \pm 0.03$ and 6.79 ± 0.01 MeV (1959PO79), $5.168 \pm 0.015, 6.154 \pm 0.010, 6.788 \pm 0.008$ MeV (1960TA17). At $E_p = 0.8$ MeV, the non-resonant radiation goes primarily [(81 ± 3%)] via cascades through $^{15}\text{O}^*(5.2, 6.18, 6.79)$ (1963BA1P). See also (1959HE47).

The excitation function for γ_0 for $E_p = 6.4$ to 19 MeV is characterized by four very pronounced peaks, ≈ 1 MeV, wide, below $E_x = 19$ MeV. The 90° excitation function then shows a giant resonance peak centered at $E_x \approx 21$ MeV (1968KU1F): see, however, (1959CO1C, 1961CO02). See also (1964TA05).

See also (1966ED1A) and (1969ZH1A; theor.).



The yields of elastic and inelastic protons, and of γ -rays have been studied at many energies: see (1959AJ76) and Table 15.24. The scattering anomalies are superposed on a background which decreases less rapidly than the Rutherford cross section; for $E_p < 2.3$ MeV, the background is largely s-wave with some p-wave contribution above $E_p = 1.5$ MeV.

Observed resonances are displayed in Table 15.23: see (1959AJ76) and (1959BA16, 1959VA08, 1963CO13, 1967KU1M, 1967LA05, 1967LA10, 1968SH11, 1969WE02). At $E_p = 9, 10$ and 10.7 MeV, broad structures, possibly intermediate-state resonances, are reported by (1968BO36). See also (1961TA06, 1964DO03, 1966MA02) and (1969AL1H; theor.).

Polarization studies have been made at many energies: see Table 15.24 (1961RO05, 1961RO13, 1965RO22, 1966BE1M, 1966BR09, 1966DR02, 1966ST05, 1968GE04). See also the reviews in (1966RO1R, 1966RO1B, 1969WA11) and (1965TA07; theor.).

Spallation measurements are reported by (1967AU1A, 1967GR1K, 1968JA1M, 1968JU1B). For astrophysical considerations see (1967LI1B).

For reaction (b), see (1966MA02).




Table 15.24: Recent $^{14}\text{N}(p, p)^{14}\text{N}$ yield curves and polarization studies ^a

(a) *Yield curves*

E_p (MeV)	of protons to $^{14}\text{N}^*$	Refs.
0.9 – 4.0	g.s.	(1959BA16)
1.0 – 2.0	g.s.	(1961CO29, 1963CO13)
1.7 – 3.5	g.s.	(1959VA08)
1.9 – 3.0	g.s.	(1967LA05, 1967LA10)
2.6 – 4.8	g.s.	(1966DR02)
3.6 – 14.5	g.s.	(1967KU1M)
3.7 – 5.7	g.s., 2.31	(1967WE1B, 1969WE02)
5.28 – 5.7	3.94	(1967WE1B, 1969WE02)
6 – 9	g.s., 2.31, 3.94	(1968SH11)
6 – 11.5	g.s., 2.31, 3.94	(1968BO36)
6.8 – 9	5.11	(1968SH11)
6.8 – 11.5	5.11	(1968BO36)
8.4 – 11.5	5.83	(1968BO36)
9 – 11.5	4.91, 5.69	(1968BO36)
10 – 12	6.44, 7.03	(1968BO36)

(b) *Polarization measurements*

3.0 – 4.8	g.s.	(1966DR02)
4.9 – 10.1	g.s.	(1966BR09)
7.1 – 8.5	2.31	(1966BR09)
7.6 – 11.8	g.s.	(1966ST05)
7.8	g.s.	(1961RO13)
8.1 – 8.5	3.94	(1966BR09)
10.4	g.s.	(1961RO05)
14.5	g.s.	(1965RO22)
21.0	g.s.	(1966BE1M)
155	g.s.	(1968GE04)

^a See also (1959AJ76).

The excitation function has been measured for $E_p = 6.3$ to 12 MeV. Broad resonances are observed for $E_p = 7.5$ to 9.85 MeV: see Table 15.23 (1964KU06). Broad resonance structure continues in the region up to $E_p \approx 16.5$ MeV (1966KU12). See also (1963VA1C, 1965VA23, 1969VE02), (1969AL1H; theor.) and ^{14}O . See also (1966RE1D, 1969BA1N) for astrophysical considerations.

$$12. \ ^{14}\text{N}(\text{p}, \text{d})^{13}\text{N} \qquad Q_m = -8.328 \qquad E_b = 7.293$$

See ^{13}N and (1963VA1C).

$$13. \ ^{14}\text{N}(\text{p}, \text{t})^{12}\text{N} \qquad Q_m = -22.139 \qquad E_b = 7.293$$

See ^{12}N in (1968AJ02).

$$14. \ ^{14}\text{N}(\text{p}, \ ^3\text{He})^{12}\text{C} \qquad Q_m = -4.779 \qquad E_b = 7.293$$

The integrated cross section for this reaction has been studied for $E_p = 8.2$ to 10.5 MeV (1968SH11). See also (1966MA02).

$$15. \ ^{14}\text{N}(\text{p}, \alpha)^{11}\text{C} \qquad Q_m = -2.920 \qquad E_b = 7.293$$

In the range $E_p = 6$ to 9 MeV, the cross section for ground state α -particles is large and shows many resonances: see Table 15.23 (1968SH11). Integrated cross sections for α_0 and α_1 are also reported for $E_p = 8.2$ to 10.5 MeV by (1968SH11). See also (1952BL64, 1963VA1C, 1966MA02, 1969WE02) and see (1966RE1D) for astrophysical implications.

$$16. \ ^{14}\text{N}(\text{d}, \text{n})^{15}\text{O} \qquad Q_m = 5.068$$

Angular distribution studies have been conducted at many energies: see Table 15.26 (1953EV03, 1960EL04, 1960MO18, 1960RE07, 1962GR18, 1963CH1D, 1963GI16, 1965SI13, 1966LO1N, 1967MU12, 1968RI1T, 1969RI1C). Information derived from DWBA analysis of the angular distributions, and from the very accurate γ -ray measurements of (1965WA16, 1966AL18, 1967CH19) are shown in Table 15.27.

Table 15.25: Lifetimes of some ^{15}O states

E_x (MeV)	τ_m (psec)	Reaction	Refs.
5.18	< 0.3	$^{16}\text{O}(^3\text{He}, \alpha)$	(1965AL19)
	< 0.3	$^{14}\text{N}(\text{d}, \text{n})$	(1965WA03)
	< 0.1	$^{16}\text{O}(\gamma, \text{n})$	(1969MU07)
5.24	> 1	$^{14}\text{N}(\text{d}, \text{n}), ^{16}\text{O}(^3\text{He}, \alpha)$	(1965AL19)
	> 5	$^{14}\text{N}(\text{d}, \text{n})$	(1965WA03)
	3.2 ± 0.5	$^{14}\text{N}(\text{d}, \text{n})$	(1967BI11)
6.18	< 0.047	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)
	< 0.1	$^{16}\text{O}(\gamma, \text{n})$	(1969MU07)
6.79	< 0.028	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)
6.86	0.10 ± 0.06	$^{14}\text{N}(\text{d}, \text{n})$	(1966AL18)
	< 0.018	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)
7.28	(1.25 ± 0.3)	$^{14}\text{N}(\text{d}, \text{n})$	(1966AL18)
	(0.70 ± 0.15)	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)

 Table 15.26: Recent $^{14}\text{N}(\text{d}, \text{n})^{15}\text{O}$ angular distribution studies ^a

E_d (MeV)	Distribution of neutron groups	Refs.
0.91 – 5.27	n_0	(1960RE07)
1.35 – 2.80	n_0	(1960EL04, 1965SI13)
1.53 – 2.90	n_0	(1960MO18)
2.2	n_0	(1966LO1N)
2.55, 2.7	n_{4+5}	(1960EL04)
2.83	$\text{n}_1 + \text{n}_2, \text{n}_3$	(1962GR18, 1963GI16)
2.85	$\text{n}_1, \text{n}_2, \text{n}_4, \text{n}_5$	(1963CH1D)
3.4 – 5.35	$\text{n}_0, \text{n}_2, \text{n}_3, \text{n}_4, \text{n}_5, \text{n}_6, \text{n}_7$	(1968RI1T, 1969RI1C)
5	$\text{n}_0, \text{n}_3, \text{n}_{4+5}, \text{n}_6$	(1967MU12)
7.7	$\text{n}_0, \text{n}_{1+2}, \text{n}_3, \text{n}_{4+5}, \text{n}_7, \text{n}_8$	(1953EV03)

^a See also (1959AJ76).

Table 15.27: Levels of ^{15}O from $^{14}\text{N}(\text{d}, \text{n})^{15}\text{O}$ and $^{14}\text{N}(^3\text{He}, \text{d})^{15}\text{O}$

E_x^a (MeV)	l_p^b	l_p^c	l_p^d	E_x^e (MeV \pm keV)	l_p^f	l_p^g	$J\pi^h$
0	1	1	1		1	1	$\frac{1}{2}^-$
5.18					(0)	(0)	$\frac{1}{2}^+$
5.24	(2)		2	5.24151 ± 0.52	2	2	$\frac{5}{2}^+$
6.18	1	1	1	6.180 ± 4	1	1	$\frac{3}{2}^-$
6.79	0	0 + 2	0		0	0	$\frac{3}{2}^+$
6.86	2	0 + 2	0	6.8598 ± 1.0	2	2	$\frac{5}{2}^+$
7.28	2	2		7.2762 ± 0.6	2	2	$\frac{7}{2}^+$
7.55	0		1			0	$\frac{1}{2}^+$
8.28						0	$\frac{3}{2}^+$

^a Nominal energies.

^b (d, n): (1968RI1T, 1969RI1C).

^c (d, n): (1967MU12).

^d (d, n): (1953EV03).

^e (d, n): (1965WA16, 1966AL18, 1967CH19).

^f (^3He , d): (1968BO14).

^g (^3He , d): (1969AL04).

^h (1968RI1T, 1969AL04, 1969RI1C).

Neutron thresholds have been observed at $E_d = 0.143 \pm 0.004$ and 0.206 ± 0.005 MeV (1963CS02), 1.24 ± 0.02 , 1.967 ± 0.004 and 2.044 ± 0.004 (1955MA85), corresponding to $^{15}\text{O}^*(5.19, 5.25, 6.15, 6.79, 6.86)$. See also (1965MA1K).

Gamma ray branching ratios are shown in Table 15.22 (1965WA16, 1968GI11). Lifetime measurements are listed in Table 15.25 (1965AL19, 1965WA03, 1966AL18, 1967BI11, 1968GI11). See also (1959MO10, 1966AV1A).

17. (a) $^{14}\text{N}(^3\text{He}, \text{d})^{15}\text{O}$ $Q_m = 1.799$
 (b) $^{14}\text{N}(^3\text{He}, \text{np})^{15}\text{O}$ $Q_m = -0.425$
 $Q_0 = 1.803 \pm 0.010$ (1959YO25);
 $Q_0 = 1.802 \pm 0.015$ (1960FO01).

Angular distributions obtained at $E(^3\text{He}) = 11$ MeV (1968BO14) and 14 MeV (1969AL04) have been analyzed by DWBA: see Table 15.27. The angular distribution of deuterons has also

been measured at $E(^3\text{He}) = 5.1$ MeV (1960FO01). For reaction (b) see (1967AD1F) and ^{16}F in (1971AJ02).

$$18. \ ^{14}\text{N}(\alpha, \text{t})^{15}\text{O} \quad Q_{\text{m}} = -12.521$$

Angular distributions of tritons corresponding to the ground state of ^{15}O have been determined at $E_{\alpha} = 43$ MeV (1967DE1K) and at 56 MeV (1968GA1C, 1969GA11). At the higher energy, a detailed comparison is made with the results from the mirror reaction $^{14}\text{N}(\alpha, ^3\text{He})^{15}\text{N}$ (1969GA11).

$$19. \ ^{14}\text{N}(^{11}\text{B}, ^{10}\text{Be})^{15}\text{O} \quad Q_{\text{m}} = -3.936$$

See (1967PO13, 1969BR1D).

$$20. \ ^{15}\text{N}(\text{p}, \text{n})^{15}\text{O} \quad Q_{\text{m}} = -3.542$$

$$E_{\text{thresh.}} = 3.7808 \pm 0.0011 [Q_0 = -3.5425 \pm 0.0011] \text{ (1958JO28, 1958WE1C)}.$$

Angular distributions of ground state neutrons have been measured at $E_{\text{p}} = 3.95$ to 5.99 MeV (1958JO28), 3.95 to 8.99 MeV, 11.4 and 13.6 MeV (1963HA46; also n_2 at $E_{\text{p}} = 5.5$ MeV), 5.5 to 13.6 MeV (1961WO03) and 18.5 MeV (1964AN1B). See also (1961SA01), (1968WO1D), (1964SA1D, 1968HA15; theor.), and ^{16}O in (1971AJ02).

$$21. \ ^{15}\text{N}(^3\text{He}, \text{t})^{15}\text{O} \quad Q_{\text{m}} = -2.778$$

A number of triton groups have been seen in this reaction. Angular distributions of these at $E(^3\text{He}) = 39.8$ and 44.6 MeV, analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture, lead to the L -values shown in Table 15.28 (1967BA13, 1968BA1E, 1969BA06). See also (1966EC1B, 1969BO13).

$$22. \ ^{16}\text{O}(\gamma, \text{n})^{15}\text{O} \quad Q_{\text{m}} = -15.668$$

Table 15.28: Levels of ^{15}O from $^{15}\text{N}(^3\text{He}, t)^{15}\text{O}$ ^a

E_x (MeV \pm keV)	L	E_x (MeV \pm keV)
0	0	9.63 ± 40
5.24 ± 30	1, 3	10.30 ± 40
6.18	0	10.49 ± 40
6.84 ± 40	1, 3	10.97 ± 50
7.28	3	11.21 ± 60
7.55	3	11.69 ± 40
8.28	1	12.34 ± 40
8.94 ± 40		13.78 ± 40
9.47 ± 50		

^a (1967BA13, 1968BA1E, 1969BA06); $E(^3\text{He}) = 39.8, 44.6$ MeV.

The spectrum of photoneutrons has been investigated at many energies. Measurements over the giant dipole resonance region show the predominant strength is to the $J^\pi = \frac{1}{2}^-$ and $\frac{3}{2}^-$ states at $E_x = 0$ and 6.18 MeV, consistent with the basic validity of the single-particle, single-hole theory of photoexcitation in ^{16}O . However, the two positive parity states at $E_x = 5.18$ and 5.24 MeV are also strongly populated, suggesting some non-single-particle excitation in that region in ^{16}O (1965CA14, 1965MA45, 1966OW01, 1967CA1C, 1967CA1P, 1968BA2L, 1969UL01). See also (1963FU05, 1964TA1C, 1965WI03, 1967FI1E, 1968JO1H, 1969CO15, 1969HO1T, 1969JO1N, 1969MU07). See also (1968ZH1B; theor.). For lifetime measurements of $^{15}\text{O}^*$, see Table 15.25 (1969MU07). See also ^{16}O in (1971AJ02).

$$23. \ ^{16}\text{O}(n, 2n)^{15}\text{O} \quad Q_m = -15.668$$

See (1955AJ61).

$$24. \ (a) \ ^{16}\text{O}(p, d)^{15}\text{O} \quad Q_m = -13.443$$

$$(b) \ ^{16}\text{O}(p, pn)^{15}\text{O} \quad Q_m = -15.668$$

Reaction (a) goes primarily to the ground state and 6.18 MeV state ($J^\pi = \frac{1}{2}^-$ and $\frac{3}{2}^-$, respectively). Angular distributions have been reported at many energies: see Table 15.29 (1961LE1A, 1963KA26, 1963LE03, 1966GR1A, 1967CH15, 1968LE01, 1969BA05, 1969SN03). See also

Table 15.29: $^{16}\text{O}(\text{p}, \text{d})^{15}\text{O}$ angular distribution studies

E_p (MeV)	Distribution of deuteron groups to $^{15}\text{O}^*$	Refs.
18.5 – 20	g.s.	(1961LE1A, 1963LE03)
21.3 – 45.3	g.s.	(1969SN03)
25.5 – 45.3	5.18 + 5.24, 6.18	(1969SN03)
30.3	g.s., 6.18	(1967CH15)
35, 40	g.s., 5.18, 6.18	(1966GR1A)
39.8	g.s., 6.18	(1963KA26)
45.3	7.28, 9.60, 10.46 ^a	(1969SN03)
100	g.s., 6.18	(1968LE01, 1968LI1A)
155.6	g.s., 6.18, 7, 10 ^b	(1965DE1A, 1969BA05, 1969TO1A)

^a Partial angular distributions to many excited states.

^b See, however, (1968LE01).

(1968SH08; theor.). The energy of $^{15}\text{O}^*(7.28)$ is 7.285 ± 0.010 MeV (1966MA1A). See also (1964BA04, 1966SH1A).

For reaction (b), see (1962BA1A, 1962FO03, 1963BE1A, 1967FU1A, 1968PU1A).

25. $^{16}\text{O}(\text{d}, \text{t})^{15}\text{O}$ $Q_m = -9.411$

Angular distributions have been measured at $E_d = 15$ MeV (1961KE01; t_0), 20 MeV (1961VL02; t_0), 20 MeV (1969PU04; t_0, t_1, t_2, t_3 ; $I_n = 1, 0, 2, 1$) and 28 MeV (1968GA13; t_0). At $E_d = 28$ MeV, detailed comparison is made with the results from the mirror reaction $^{16}\text{O}(\text{d}, ^3\text{He})^{15}\text{N}$ (1968GA13).

26. $^{16}\text{O}(^3\text{He}, \alpha)^{15}\text{O}$ $Q_m = 4.910$
 $Q_0 = 4.916 \pm 0.010$ (1959HI68);
 $Q_0 = 4.917 \pm 0.015$ (1962SH21).

The $p_{1/2}$ and $p_{3/2}$ hole states at $E_x = 0$ and 6.18 MeV are strongly populated in this reaction, see e.g. (1965WA17). Information on these and other states of ^{15}O observed in this reaction is shown in Table 15.30 (1959HI68, 1959YO25, 1967HE1A, 1968BO14). The J -values are derived from angular correlation measurements (1967HE1A, 1966GA19, 1966GO15, 1967GO07,

Table 15.30: Excited states of ^{15}O from $^{16}\text{O}(^3\text{He}, \alpha)^{15}\text{O}$

E_x (MeV \pm keV)	Refs.	l^b	J	Refs.
0		1		
5.174 ± 10	(1959HI68)	0		
5.167 ± 15	(1959YO25)			
5.193 ± 11	(1965AL19)			
5.233 ± 10	(1959HI68)	2	$\frac{5}{2}$	(1966GO15, 1967GO07)
5.243 ± 10	(1965AL19)			
6.18	^a	1	$\frac{3}{2}$	(1966GO15)
6.79	^a	2	$\frac{3}{2}$	(1966GA19, 1968GI01)
6.86	^a	2	$\frac{5}{2}$	(1966GA19, 1968GI01)
7.2742 ± 1.4	(1967HE1A)		$\frac{7}{2}$	(1967HE1A)

^a Nominal energy.

^b (1968BO14).

(1968GI01): see also (1968BO14). Angular distributions have been measured for $E(^3\text{He}) = 5.2$ to 36.6 MeV: see Table 15.31 (1959HI73, 1960TA12, 1962SE13, 1965AL05, 1965AR07, 1968BO14, 1969BR07). Branching ratios are displayed in Table 15.22 (1965WA16, 1969KU01). See also (1965WA03). The lifetimes of $^{15}\text{O}^*(5.18, 5.24)$ are < 0.3 and > 1 psec, respectively (1965AL19): see also Table 15.25. A comparison of the 5 MeV transitions E3/M2 mixing ratios in ^{15}O and ^{15}N strongly suggest a collective character for the E3 component of these transitions (1968GI01).

The ratio of mixing ratios of the mirror decays $^{15}\text{O}^*(6.18 \rightarrow 0)$ and $^{15}\text{N}^*(6.32 \rightarrow 0)$ are in disagreement with the IPM suggesting a collective contribution to the mirror $\frac{3}{2}^-$ levels (1966RO1U).

The M2/E1 mixing ratio for the $^{15}\text{O}^*(6.79)$ transition indicates an exceptionally high retardation of an E1 transition in a non-self-conjugate nucleus, as is also true of the analog transition in ^{15}N (1968GI01).

The $E_x = 7.28$ MeV state, $J = \frac{7}{2}$, has a negligible effect on the (astrophysical) CNO cycle (1967HE1A).

See also (1961DU02, 1961SI09, 1966AG1B) and ^{19}Ne in (1972AJ02).

$$27. \ ^{16}\text{O}(^{14}\text{N}, ^{15}\text{N})^{15}\text{O} \quad Q_m = -4.833$$

See (1965GA1B, 1969BR1D).

$$28. \ ^{17}\text{O}(p, t)^{15}\text{O} \quad Q_m = -11.329$$

Table 15.31: $^{16}\text{O}(^3\text{He}, \alpha)$ angular distribution studies

$E(^3\text{He})$ (MeV)	Distribution of α groups to $^{15}\text{O}^*$	Refs.
5.2	g.s.	(1960TA12)
5.7, 5.9	g.s.	(1959HI73)
8 – 10	g.s.	(1965AL05)
9.2	g.s., 5.18, 5.24	(1959HI73)
9.8 – 11.7	g.s.	(1966BR13, 1969BR07)
11	g.s., 5.18, 5.24, 6.18, 6.79, 6.86	(1968BO14)
16.6, 25.8, 36.6	g.s., 5.18 + 5.24, 6.18	(1965AR07)
29	5.18 + 5.24, 6.18	(1962SE13)

See (1969ME1M).

29. $^{19}\text{F}(\text{p}, \text{n}\alpha)^{15}\text{O}$ $Q_{\text{m}} = -7.553$

See (1962FO03).

^{15}F

Mass of ^{15}F : A calculation using an isobaric mass formula predicts that ^{15}F is unstable with respect to proton emission by 2.32 MeV: the mass excess of ^{15}F is then 17.62 MeV (1966KE16). See also (1957MU99, 1961BA1C).

References

(Closed 31 December 1969)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1941BE1A Bennett et al., Phys. Rev. 59 (1941) 781
- 1950CU13 C.D. Curling and J.O. Newton, Nature 165 (1950) 609
- 1950JO57 C.H. Johnson and H.H. Barschall, Phys. Rev. 80 (1950) 818
- 1950MA65 R. Malm and W.W. Buechner, Phys. Rev. 80 (1950) 771
- 1950RI57 J.E. Richardson, Phys. Rev. 80 (1950) 850
- 1951DU08 D.B. Duncan and J.E. Perry, Phys. Rev. 82 (1951) 809
- 1951JO23 C.H. Johnson, B. Petree and R.K. Adair, Phys. Rev. 84 (1951) 775
- 1951RO16 W.D. Roseborough, J.J.G. McCue, W.M. Preston and C. Goodman, Phys. Rev. 83 (1951) 1133
- 1951YA1A Yaffe and Stevens, Can. J. Phys. 29 (1951) 186; Phys. Rev. 79 (1950) 893
- 1952BL64 J.P. Blaser, P. Marmier and M. Sempert, Helv. Phys. Acta 25 (1952) 442
- 1952GI01 W.M. Gibson and E.E. Thomas, Proc. Roy. Soc. A210 (1952) 543
- 1952HI12 J.J. Hinchey, P.H. Stelson and W.M. Preston, Phys. Rev. 86 (1952) 483
- 1953EV03 W.H. Evans, T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A66 (1953) 108
- 1953KA20 R. Kay, H. Mark and C. Goodman, Phys. Rev. 91 (1953) 472A; Oral Report
- 1953KO42 B. Koudijs, F.P.G. Valckx and P.M. Endt, Physica 19 (1953) 1133
- 1954BE08 W.E. Bennett, P.A. Roys and B.J. Toppel, Phys. Rev. 93 (1954) 924, K4
- 1954KL36 R.M. Kline and D.J. Zaffarano, Phys. Rev. 96 (1954) 1620
- 1954PA39 R. Pauli, K. Ahnlund and C. Mileikowsky, Ark. Fys. 8 (1954) 213
- 1954SP01 A. Sperduto, W.W. Buechner, C.K. Bockelman and C.P. Browne, Phys. Rev. 96 (1954) 1316
- 1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77
- 1955BA44 G.A. Bartholomew, F. Brown, H.E. Gove, A.E. Litherland and E.B. Paul, Can. J. Phys. 33 (1955) 441
- 1955BA83 S. Bashkin, R.R. Carlson and E.B. Nelson, Phys. Rev. 99 (1955) 107
- 1955FO27 J.L. Fowler and C.H. Johnson, Phys. Rev. 98 (1955) 728

- 1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847.
- 1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46
- 1955SH28 R.D. Sharp, A. Sperduto and W.W. Buechner, Phys. Rev. 99 (1955) 632A
- 1955SH46 E.S. Shire and R.D. Edge, Phil. Mag. 46 (1955) 640
- 1956BA16 G.A. Bartholomew, A.E. Litherland, E.B. Paul and H.E. Gove, Can. J. Phys. 34 (1956) 147
- 1956BA34 J.K. Bair, H.O. Cohn, J.D. Kington and H.B. Willard, Phys. Rev. 104 (1956) 1595
- 1956BO61 T.W. Bonner, A.A. Kraus Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348
- 1956DO37 R.A. Douglas, B.R. Gasten and A. Mukerji, Can. J. Phys. 34 (1956) 1097
- 1956DO41 R.A. Douglas, J.W. Broer, R. Chiba, D.F. Herring and E.A. Silverstein, Phys. Rev. 104 (1956) 1059
- 1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28
- 1956MA35 J.B. Marion and G. Weber, Phys. Rev. 102 (1956) 1355; Erratum Phys. Rev. 103 (1956) 1906
- 1956MA46 J.B. Marion and G. Weber, Phys. Rev. 103 (1956) 167
- 1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434
- 1956TA16 G.W. Tautfest and S. Rubin, Phys. Rev. 103 (1956) 196
- 1956VA17 F.P.G. Valckx, Ph.D. Thesis, Univ. of Utrecht (1956)
- 1957AL78 D.G. Alkhazov, I.P. Gangrskii and I.K. Lemberg, Zh. Eksp. Teor. Fiz. 33 (1957) 1160; Sov. Phys. JETP 6 (1958) 892
- 1957BA18 G.A. Bartholomew and P.J. Campion, Can. J. Phys. 35 (1957) 1347
- 1957BO58 C.R. Bolmgren, G.D. Freier, J.G. Likely and K.F. Famularo, Phys. Rev. 105 (1957) 210
- 1957BR18 D.A. Bromley, E. Almqvist, H.E. Gove, A.E. Litherland, E.B. Paul and A.J. Ferguson, Phys. Rev. 105 (1957) 957
- 1957HA03 F.B. Hagedorn, F.S. Mozer, T.S. Webb, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 105 (1957) 219
- 1957HA1E Halbert and French, Phys. Rev. 105 (1957) 1563
- 1957JO20 S.A.E. Johansson and B. Forkman, Ark. Fys. 12 (1957) 359
- 1957KI22 O.C. Kistner, A. Schwarzschild, B.M. Rustad and D.E. Alburger, Phys. Rev. 105 (1957) 1339
- 1957LA13 W.A.S. Lamb and R.E. Hester, Phys. Rev. 108 (1957) 1304
- 1957MU99 P.G. Murphy, Phys. Rev. 108 (1957) 421

1957NO14 E. Norbeck Jr., Phys. Rev. 105 (1957) 204
 1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
 1957PE12 J.R. Penning and F.H. Schmidt, Phys. Rev. 105 (1957) 647
 1957PIZZ R.E. Pixley, Thesis, CalTech Pasadena (1957)
 1957SJ68 B. Sjogren and K. Ahnlund, Ark. Fys. 12 (1957) 547
 1957WA01 E.K. Warburton and J.N. McGruer, Phys. Rev. 105 (1957) 639
 1958GR01 L.V. Groshev, A.M. Demidov, V.N. Lutsenko and V.I. Pelekhov, Atlas of Thermal Neutron Capture Gamma Rays, Atomizdat, Moscow (1958)
 1958HA1B Haddad, Perry and Smith, Private Communication (1958)
 1958JO20 R.L. Johnston, H.D. Holmgren, E.A. Wolicki and E.G. Illsley, Phys. Rev. 109 (1958) 884
 1958JO28 K.W. Jones, L.J. Lidofsky and J.L. Weil, Phys. Rev. 112 (1958) 1252
 1958WA1C Way, Nucl. Data Cards, Natl. Res. Council, Washington, D.C. (1958)
 1958WE1C Weil, CU-180 (1958)
 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
 1959AL06 D.E. Alburger, A. Gallmann and D.H. Wilkinson, Phys. Rev. 116 (1959) 939
 1959AL97 D.E. Alburger, Phys. Rev. Lett. 3 (1959) 280
 1959BA16 S. Bashkin, R.R. Carlson and R.A. Douglas, Phys. Rev. 114 (1959) 1552
 1959BI19 E.G. Bilpuch, L.W. Weston, C.D. Bowman and H.W. Newson, Bull. Amer. Phys. Soc. 4 (1959) 42, R4
 1959BO1C Bockelman, Nucl. Phys. 13 (1959) 205
 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
 1959BR69 P. Brix and E.K. Maschke, Z. Phys. 155 (1959) 109
 1959BR79 D.A. Bromley, J.A. Kuehner and E. Almqvist, Nucl. Phys. 13 (1959) 1
 1959CO1C Cohen, Fisher and Warburton, Phys. Rev. Lett. 3 (1959) 433; Erratum Phys. Rev. Lett. 4 (1960) 92
 1959FE1B Feingold, Phys. Rev. 114 (1959) 540
 1959FE1C Ferguson and Gove, Can. J. Phys. 37 (1959) 660
 1959GA05 A. Gallmann, Ann. Phys. 4 (1959) 185
 1959GA14 F. Gabbard, H. Bischel and T.W. Bonner, Nucl. Phys. 14 (1959) 277
 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295
 1959HE1B Hess, Ann. Phys. 6 (1959) 115
 1959HE1D Hebbard and Dunbar, Phys. Rev. 115 (1959) 624

- 1959HE47 D.F. Hebbard and B. Povh, Nucl. Phys. 13 (1959) 642
- 1959HI68 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A73 (1959) 727
- 1959HI73 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 775
- 1959JO32 R.L. Johnston, H.D. Holmgren and G.D. Gutsche, Bull. Amer. Phys. Soc. 4 (1959) 403, C4
- 1959KI99 O.C. Kistner and B.M. Rustad, Phys. Rev. 114 (1959) 1329
- 1959LE28 L.L. Lee Jr. and J.P. Schiffer, Phys. Rev. 115 (1959) 160
- 1959MA1D Markowitz and Hall, Bull. Amer. Phys. Soc. 4 (1959) 8
- 1959MO10 E. Moller, Ark. Fys. 15 (1959) 251
- 1959MO1B Morre and McGruer, Bull. Amer. Phys. Soc. 4 (1959) 17, and Private Communication (1959)
- 1959NO38 I. Nonaka, H. Yamaguchi, T. Mikumo, I. Umeda, T. Tabata and S. Hitaka, J. Phys. Soc. Jpn. 14 (1959) 1260; Erratum J. Phys. Soc. Jpn. 15 (1960) 365
- 1959PO79 B. Povh and D.F. Hebbard, Phys. Rev. 115 (1959) 608
- 1959VA04 A.K. Valter, V.Y. Gonchar, A.N. Lvov and S.P. Tsitko, Izv. Akad. Nauk SSSR Ser. Fiz. 23 (1959) 228; Columbia Tech. Transl. 23 (1960) 219
- 1959VA08 A.K. Valter, A.S. Deineko, I.Y. Malakhov, P.V. Sorokin and A.Y. Taranov, Izv. Akad. Nauk SSSR Ser. Fiz. 23 (1959) 839
- 1959YO25 T.E. Young, G.C. Phillips, R.R. Spencer and D.A.A.S.N. Rao, Phys. Rev. 116 (1959) 962
- 1960BA34 J.K. Bair, R.D. Edge and H.B. Willard, Phys. Rev. 119 (1960) 1948
- 1960BE1B Belyaev, Zakhar'ev and Neudachin, Atomnaya Energiya 9 (1960) 298, Sov. J. At. Energy 9 (1961) 833
- 1960BU1C Bullock and Moore, Phys. Rev. 119 (1960) 721
- 1960CA02 R.E. Carter and H.T. Motz, Bull. Amer. Phys. Soc. 5 (1960) 246, HA7
- 1960CL02 R.L. Clarke, E. Almqvist and E.B. Paul, Nucl. Phys. 14 (1960) 472
- 1960EL04 A.J. Elwyn, J.V. Kane, S. Ofer and R. Pixley, Phys. Rev. 120 (1960) 2207
- 1960FA10 R.F. Favreau, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds., D.A. Bromley and E.W. Vogt (1960) 939
- 1960FE12 J.M. Ferguson and W.E. Thompson, Phys. Rev. 118 (1960) 228
- 1960FO01 P.D. Forsyth, F. de S. Barros, A.A. Jaffe, I.J. Taylor and S. Ramavataram, Proc. Phys. Soc. (London) A75 (1960) 291
- 1960FR09 J.B. French, S. Iwao and E.W. Vogt, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds., D.A. Bromley and E.W. Vogt (1960) 480

- 1960HE13 D.F. Hebbard, Nucl. Phys. 19 (1960) 511
- 1960HI07 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) 75 (1960) 745
- 1960HO1B Honsaker, Astrophys. J. 132 (1960) 21
- 1960JA12 J. Janecke, Z. Naturforsch. A15 (1960) 593
- 1960MC05 J.H. McCrary and I.L. Morgan, Bull. Amer. Phys. Soc. 5 (1960) 246, HA5
- 1960MO03 S. Morita, N. Kawai, N. Takano, Y. Goto, R. Hanada, Y. Nakajima, S. Takemoto and Y. Yaegashi, J. Phys. Soc. Jpn. 15 (1960) 361
- 1960MO18 S. Morita, N. Kawai, Y. Goto, T. Maki and M. Mukae, J. Phys. Soc. Jpn. 15 (1960) 2170
- 1960PE25 J.M. Peterson, A. Bratenahl and J.P. Stoering, Phys. Rev. 120 (1960) 521
- 1960PR13 J.R. Priest, D.J. Tendam and E. Bleuler, Phys. Rev. 119 (1960) 1301
- 1960RE07 T. Retz-Schmidt and J.L. Weil, Phys. Rev. 119 (1960) 1079
- 1960SH05 S.M. Shafroth, J. Phys. Rad. 21 (1960) 353
- 1960TA12 I.J. Taylor, F.de S. Barros, P.D. Forsyth, A.A. Jaffe and S. Ramavataram, Proc. Phys. Soc. (London) A75 (1960) 772
- 1960TA17 T. Tabata and K. Okano, J. Phys. Soc. Jpn. 15 (1960) 1552
- 1960TA1C Talmi and Unna, Ann. Rev. Nucl. Sci. 10 (1960) 353
- 1960TA1E Tauber and Wu, Proc. Int. Conf. on Nucl. Struct., Kingston (1960)
- 1960VA11 F.J. Vaughn, L.F. Chase Jr. and R.G. Johnson, Bull. Amer. Phys. Soc. 5 (1960) 404, B2
- 1960ZE03 Ya.B. Zeldovich, Zh. Eksp. Teor. Fiz. 38 (1960) 1123; Sov. Phys. JETP 11 (1960) 812
- 1961AG1A Aguilar and Garcia, An. Real. Soc. Espan. Fis. y Quim. A57 (1961) 29
- 1961BA05 G.A. Bartholomew and J.F. Vervier, Bull. Amer. Phys. Soc. 6 (1961) 237, DA13
- 1961BA1C Baz, Goldanskii and Zeldovich, Sov. Phys. Uspekhi 3 (1961) 729
- 1961BA1E Balashov, Neudachin and Smirnov, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 170; Bull. Acad. Sci. USSR Phys. 25 (1961) 165
- 1961BR13 H.R. Brooker, P.J. Haigh and T.A. Scott, Phys. Rev. 123 (1961) 2143
- 1961BR1A Brill, Vlasov, Kalinin and Sokolov, Dokl. Akad. Nauk SSSR 136 (1961) 55; Sov. Phys. Dokl. 6 (1961) 24
- 1961CA02 R.R. Carlson, C.C. Kim, J.A. Jacobs and A.C.L. Barnard, Phys. Rev. 122 (1961) 607
- 1961CH14 R. Chiba, Phys. Rev. 123 (1961) 1316
- 1961CO02 S.G. Cohen, P.S. Fisher and E.K. Warburton, Phys. Rev. 121 (1961) 858
- 1961CO29 J. Cohen-Ganouna, M. Lambert and J. Schmouker, J. Phys. Rad. 22 (1961) 592

1961DU02 K.L. Dunning and J.W. Butler, Phys. Rev. 123 (1961) 1321
 1961DU1B Duggan, Thesis, Louisiana State Univ. (1961)
 1961FO02 P.D. Forsyth and G.S. Mani, Bull. Amer. Phys. Soc. 6 (1961) 227, AA12
 1961FR1D French, Iwao and Vogt, AECL 1136 (1960)
 1961GO03 S. Gorodetzky, P. Fintz, G. Bassompierre and A. Gallmann, Compt. Rend. 252 (1961) 713
 1961HA12 J. Hanna, T.W. Bonner, R.L. Bramblett and F.A.St. Romain, Bull. Amer. Phys. Soc. 6 (1961) 369, S9
 1961HA43 G.C. Hanna, D.B. Primeau and P.R. Tunncliffe, Can. J. Phys. 39 (1961) 1784
 1961JA09 A.N. James, Nucl. Phys. 24 (1961) 132
 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, Rev. Mex. Fisica 10 (1961) 247
 1961JO07 R.G. Johnson, L.F. Chase Jr. and F.J. Vaughn, Bull. Amer. Phys. Soc. 6 (1961) 236, DA8
 1961JO24 R.G. Johnson, L.F. Chase Jr. and F.J. Vaughn, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed., J.B. Birks (1961) 591
 1961KA05 N. Kawai, J. Phys. Soc. Jpn. 16 (1961) 157
 1961KE01 E.L. Keller, Phys. Rev. 121 (1961) 820
 1961KO04 M. Kondo, T. Yamazaki and S. Yamabe, J. Phys. Soc. Jpn. 16 (1961) 1091
 1961KO1E Kolodziejcki, Acta Phys. Pol. 20 (1961) 275
 1961KR1A Kromminga and McCarthy, Nucl. Phys. 24 (1961) 36
 1961LE01 J.J. Leigh and J.M. Blair, Phys. Rev. 121 (1961) 246
 1961LE1A Legg, Unpublished Thesis, Princeton Univ. (1961)
 1961LO10 G. Lopez and O. Almen, Rev. Mex. Fis. 10 (1961) 239
 1961MO13 S. Morita, T. Ishimatsu, T. Cho, Y. Nakajima, N. Kawai, T. Murata and Y. Hachiya, J. Phys. Soc. Jpn. 16 (1961) 1849; Erratum J. Phys. Soc. Jpn. 17 (1962) 251
 1961NO05 E. Norbeck, Phys. Rev. 121 (1961) 824
 1961PU1B Pullen, Wilkinson and Whitehead, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed., J.B. Birks (1961) 565
 1961RA06 L.A. Rayburn, Phys. Rev. 122 (1961) 168
 1961RO05 L. Rosen, J.E. Brolley Jr. and L. Stewart, Phys. Rev. 121 (1961) 1423
 1961RO13 L. Rosen, J.E. Brolley Jr., M.L. Gursky and L. Stewart, Phys. Rev. 124 (1961) 199
 1961SA01 A. Sayres, D. Lister and D. Lightbody, Bull. Amer. Phys. Soc. 6 (1961) 26, I11
 1961SI09 E.A. Silverstein, S.R. Salisbury, G. Hardie and L.D. Oppliger, Phys. Rev. 124 (1961) 868

- 1961SJ1B Sjogren and Sawa, Ark. Fys. 19 (1961) 417
- 1961TA06 A.E. Taylor and E. Wood, Nucl. Phys. 25 (1961) 642
- 1961TE02 A. Tejera, M. Mazari, A. Jaidar and G. Lopez, Rev. Mex. Fis. 10 (1961) 229
- 1961TO01 K.S. Toth, Phys. Rev. 121 (1961) 1190
- 1961TO03 J.H. Towle and B.E.F. Macefield, Proc. Phys. Soc. (London) 77 (1961) 399
- 1961TO07 K.S. Toth, Phys. Rev. 123 (1961) 582
- 1961VA13 P.I. Vatsset, L.Ya. Kolesnikov and S.G. Tonapetyan, Zh. Eksp. Teor. Fiz. 40 (1961) 1257; Sov. Phys. JETP 13 (1961) 886
- 1961VL02 N.A. Vlasov, S.P. Kalinin, A.A. Ogloblin and V.I. Chuev, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 115; Columbia Tech. Transl. 25 (1961) 111
- 1961WI1A Wilkinson, Wollan and Koehler, Ann. Rev. Nucl. Sci. 11 (1961) 303
- 1961WO03 C. Wong, J.D. Anderson, S.D. Bloom, J.W. McClure and B.D. Walker, Phys. Rev. 123 (1961) 598
- 1962AL01 W.P. Alford, O.M. Bilaniuk and D.B. Marsh, Bull. Amer. Phys. Soc. 7 (1962) 60, R8
- 1962BA1A Balashov and Boyarkina, Nucl. Phys. 38 (1962) 629
- 1962BA63 J.D. Baldeschweiler, J. Chem. Phys. 36 (1962) 152
- 1962BR1G Beit, in Electomag. Lifetimes Props. of Nucl. States, N.A.S.N.R.C. Pub. 974 (1962) 21
- 1962CE1B Celvolani and Petralia, Nuovo Cim. 26 (1962) 1328
- 1962CH14 L.F. Chase Jr., R.G. Johnson, F.J. Vaughn and E.K. Warburton, Phys. Rev. 127 (1962) 859
- 1962CO31 D.R.F. Cochran and J.D. Knight, Phys. Rev. 128 (1962) 1281
- 1962DO1A Dodge and Barber, Phys. Rev. 127 (1962) 1746
- 1962EI03 W.W. Eidson and R.D. Bent, Phys. Rev. 127 (1962) 913
- 1962ER03 H.J. Erramuspe and R.J. Slobodrian, Nucl. Phys. 34 (1962) 532
- 1962FO03 K.J. Foley, G.L. Salmon and A.B. Clegg, Nucl. Phys. 31 (1962) 43
- 1962GO1J Gorshkov, Zyabkin and Tsvetkov, Atomn. Energ. (USSR) 13 (1962) 65
- 1962GO21 S. Gorodetzky, P. Fintz and A. Gallmann, Compt. Rend. 255 (1962) 879
- 1962GR18 W. Gruebler and J. Rossel, Helv. Phys. Acta 35 (1962) 283
- 1962GU01 G.D. Gutsche, H.D. Holmgren, L.M. Cameron and R.L. Johnston, Phys. Rev. 125 (1962) 648
- 1962HA20 J. Hanna, F.R.St. Romain, R.L. Bramblett and T.W. Bonner, Bull. Amer. Phys. Soc. 7 (1962) 453, GA4
- 1962HO06 R.K. Hobbie and F.F. Forbes, Phys. Rev. 126 (1962) 2137

1962HO1D Honda and Ui, Nucl. Phys. 34 (1962) 592
 1962KU09 B. Kuhn, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teopr. Fiz. 43 (1962) 1660; Sov. Phys. JETP 16 (1963) 1171
 1962LI07 A.J. Lieber, F.H. Schmidt and J.B. Gerhart, Phys. Rev. 126 (1962) 1496
 1962LO02 J. Lowe, C.L. McClelland and J.V. Kane, Phys. Rev. 126 (1962) 1811
 1962MC12 R.L. McGrath, Phys. Rev. 127 (1962) 2138
 1962NE1D Nedvedyk, Joint Inst. Nucl. Res. Lab. Neutron Phys. USSR, Rept. No. P-1098 (1962)
 1962OT01 P.S. Otstavnov and V.I. Popov, Zh. Eksp. Teor. Fiz. 43 (1962) 385; Sov. Phys. JETP 16 (1963) 276
 1962RO13 V.M. Rout, W.M. Jones and D.G. Waters, Nucl. Phys. 34 (1962) 628
 1962SE13 H.M. Sen Gupta, J. Rotblat, P.E. Hodgson and J.B.A. England, Nucl. Phys. 38 (1962) 361
 1962SH21 R.D. Sharp, Bull. Amer. Phys. Soc. 7 (1962) 622, W1
 1962ST17 D. Stanojevic and M. Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 13 (1962) 1
 1962TA1E Talmi, Nucl. Spectroscopy; Ed., Racah (1962)
 1962TE1B Teplov, Zh. Eksp. Teor. Fiz. 42 (1962) 211; Sov. Phys. JETP 15 (1962) 150
 1963AL21 D.E. Alburger and E.K. Warburton, Phys. Rev. 132 (1963) 790
 1963BA1P Bailey and Hebbard, Nucl. Phys. 46 (1963) 529
 1963BA1Q R. Ballini and N. Saunier, J. Phys. (France) 24 (1963) 904
 1963BA46 R.W. Bauer, J.D. Anderson and L.J. Christensen, Nucl. Phys. 47 (1963) 241
 1963BE1A Benioff, Phys. Rev. 129 (1963) 1355
 1963BE42 T. Berggren and G. Jacob, Nucl. Phys. 47 (1963) 481
 1963BR1G Breit, in Padua (1963) 480
 1963BU1C Bunakov, Phys. Lett. 7 (1963) 14
 1963CH1C Chatterjee, Nucl. Phys. 49 (1963) 686
 1963CH1D Christiansen and Zeitnitz, in Padua (1963) A557
 1963CL1A Clement, in Padua (1963) 457
 1963CL1B Clegg, CERN 63-28 (1963)
 1963CO12 B.L. Cohen, Phys. Rev. 130 (1963) 227
 1963CO13 J. Cohen-Ganouna, M. Lambert and J. Schmouker, J. Phys. 24 (1963) 43
 1963CO17 E.D. Commins and H.R. Feldman, Phys. Rev. 131 (1963) 700
 1963CS02 J. Csikai and G. Peto, Phys. Lett. 4 (1963) 252
 1963DA1B Dar, Phys. Lett. 7 (1963) 339

- 1963DE19 V.K. Deshpande, Nucl. Phys. 47 (1963) 257
- 1963DR1B Drisko, Satchler and Dassel, 3rd Conf. on Reactions between Complex Nuclei (1963) 85
- 1963DU12 J.L. Duggan, P.D. Miller and R.F. Gabbard, Nucl. Phys. 46 (1963) 336
- 1963ED01 R.D. Edge, Bull. Amer. Phys. Soc. 8 (1963) 11, BA5
- 1963EN01 C.E. Engelke, R.E. Benenson, E. Melkonian and J.M. Lebowitz, Phys. Rev. 129 (1963) 324
- 1963FI04 E. Finckh, R. Kosiek and K. Schlupmann, Naturwiss. 50 (1963) 326
- 1963FU05 H. Fuchs and D. Haag, Z. Physik 171 (1963) 403
- 1963GA10 E. Gadioli and S. Micheletti, Phys. Lett. 6 (1963) 229
- 1963GI16 W.R. Gibbs and W.E. Gruebler, Helv. Phys. Acta 36 (1963) 693
- 1963GO1J Gorshkov and Tsvetkov, Atomnaya Energ.14 (1963) 550
- 1963GO1L Gofman et al., Zh. Eksp. Teor. Fiz. 45 (1963) 1317; Sov. Phys. JETP 18 (1964) 906
- 1963GO1M Goldberg, May and Stehn, BNL-400, 2nd Ed., Vol. 1 (1963)
- 1963HA46 L.F. Hansen and M.L. Stelts, Phys. Rev. 132 (1963) 1123
- 1963HE11 D.F. Hebbard and G.M. Bailey, Nucl. Phys. 49 (1963) 666
- 1963HO1E Hortig, Werner and Gentner, 3rd Conf. on Reactions between Complex Nuclei (1963) 178
- 1963IM01 W.L. Imhof, H.A. Grench and R.G. Johnson, Nucl. Phys. 49 (1963) 503
- 1963KA26 C.D. Kavaloski, G. Bassani and N.M. Hintz, Phys. Rev. 132 (1963) 813
- 1963KI1B Kim, Nucl. Phys. 49 (1963) 383
- 1963KO03 M. Kondo, T. Yamazaki and S. Yamabe, J. Phys. Soc. Jpn. 18 (1963) 22
- 1963KU1B Kunz, Can. J. Phys. 41 (1963) 2187
- 1963LA02 R.A. Lasalle, R.D. Benta and J.G. Cramer Jr., Bull. Amer. Phys. Soc. 8 (1963) 303, D8
- 1963LE03 J.C. Legg, Phys. Rev. 129 (1963) 272
- 1963LU01 C.A. Ludemann, H.D. Holmgren and W.F. Hornyak, Bull. Amer. Phys. Soc. 8 (1963) 12, BA10
- 1963LU05 C.A. Ludemann, H.D. Holmgren and J.E. Etter, Bull. Amer. Phys. Soc. 8 (1963) 303, D6
- 1963LU1F C.A. Ludemann, H.D. Holmgren and W.F. Hornyak, in Padua (1963) 850A
- 1963MA28 G.S. Mani, P.D. Forsyth and R.R. Perry, Nucl. Phys. 44 (1963) 625
- 1963MO1B Morrison, Gale, Hussain and Murray, 3rd Conf. on Reactions between Complex Nuclei (1963) 168

1963MO1C Motz and Journey, Wash-1044 (1963)
 1963NA1C Nagahara et al., in Padua (1963) 850A
 1963NE05 J.W. Nelson, E.B. Carter, G.E. Mitchell and R.H. Davis, Phys. Rev. 129 (1963) 1723
 1963NI04 K. Niedzwiedziuk, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teor. Fiz. 44 (1963) 1450;
 JETP (Sov. Phys.) 17 (1963) 974
 1963RI1B Riou, in Padua (1963) 18
 1963SC32 J.A. Scheer, K. Schlupmann and F. Triantafyllidis, Phys. Lett. 7 (1963) 269
 1963TA1A Tanifuji, Nucl. Phys. 40 (1963) 357
 1963TO1D Toth and Newman, 3rd Conf. on Reactions between Complex Nuclei (1963) 114
 1963VA1C Valentin, Albouy, Cohen and Gusakow, Phys. Lett. 7 (1963) 163
 1963VA31 S.S. Vasilev and L.Y. Shavtvalov, Izv. Akad. Nauk SSSR Ser. Fiz. 27 (1963) 1261;
 Bull. Acad. Sci. USSR Phys. Ser. 27 (1963) 1239
 1963WE15 V.N. Webb, M.M. Duncan, J. Lin, J.L. Duggan, L.E. Akers and H.E. Banta, Bull.
 Amer. Phys. Soc. 9 (1964) 351, S7
 1963YA1C Yamazaki, Knodo and Yamabe, J. Phys. Soc. Jpn. 18 (1963) 620
 1964AL1L Ali and Tauber, Nucl. Phys. 55 (1964) 481
 1964AL21 D.E. Alburger, C. Chasman, K.W. Jones and R.A. Ristinen, Phys. Rev. 136 (1964)
 B913
 1964AM1A Amsel, Ann. Phys. 9 (1964) 297
 1964AM1D Amit and Latz, Nucl. Phys. 58 (1964) 297
 1964AN1B Anderson, Wong, McClure and Walker, Phys. Rev. 136 (1964) B118
 1964BA02 G.A. Bartholomew and J.F. Vervier, Nucl. Phys. 50 (1964) 209
 1964BA04 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz, N.K. Ganguly and P. Radvanyi, Phys.
 Lett. 8 (1964) 56
 1964BA1V Ballot, Dumontet, Picinbono and Saya, Nucl. Phys. 51 (1964) 401
 1964BL1B Blair, Pham-Dihn-Lien and Hobbie, Phys. Rev. 134 (1964) B793
 1964BL1C Blieden, Phys. Lett. 9 (1964) 176
 1964BR13 H.C. Bryant, J.G. Beery, E.R. Flynn and W.T. Leland, Nucl. Phys. 53 (1964) 97
 1964BR1G Brill, Chuev and Ogloblin, 4C/C331, Paris (1964)
 1964BR1H Brown, in Paris (1964) 129
 1964BR1M Breit, Chun and Wahsweiler, Phys. Rev. 133 (1964) B404
 1964BU1D Burcham et al., 4B(1)/C212, Paris (1964)
 1964CA05 R.R. Carlson, E. Norbeck and V. Hart, Bull. Amer. Phys. Soc. 9 (1964) 419, DA9

1964CA18 R.R. Carlson and M. Throop, Phys. Rev. 136 (1964) B630
 1964DA1B W.W. Daehnick and L.J. Denes, Phys. Rev. 136 (1964) B1325.
 1964DA1D Dar, Nucl. Phys. 55 (1964) 305
 1964DE1C Deshpande, Fulbright and Verba, Nucl. Phys. 52 (1964) 457
 1964DE1E Deshpande and Fulbright, UR 875 61 (1964)
 1964DI02 G.U. Din, H.M. Kuan and T.W. Bonner, Nucl. Phys. 50 (1964) 267
 1964DI1C Din, Unpublished Thesis, Rice Univ. (1964)
 1964DO03 P.F. Donovan, J.F. Mollenauer and E.K. Warburton, Phys. Rev. 133 (1964) B113
 1964EC03 S.F. Eccles, H.F. Lutz and J. Stevens, Bull. Amer. Phys. Soc. 9 (1964) 704, B9
 1964EV1A Evans, NOL TR 64-46 (1964)
 1964FL1D Flerov and Karnaukhov, Congres Int. Phys. Nucl., Paris, 1964 (1964) 373
 1964FO1A Fowler and Vogl, Lectures in Theor. Phys., Vol. 6 (1964) 379
 1964GA1A Gardner and Yu, Nucl. Phys. 60 (1964) 49
 1964GR1G Grieder, Phys. Rev. 133 (1964) B1483
 1964GR1H Gromov et al., 4A(11)/C358, Paris (1964)
 1964HA09 V.P. Hart, E. Norbeck and R.R. Carlson, Bull. Amer. Phys. Soc. 9 (1964) 430, EA6
 1964HE18 I. Heertje, I. Delvenne, W. Nagel and A.H.W. Aten Jr., Physica 30 (1964) 1762
 1964JO1A Jobes and McIntyre, Phys. Rev. 133 (1964) B893
 1964KE1C Kelley and Henley, Phys. Lett. 10 (1964) 95
 1964KO10 R. Kosiek, Z. Phys. 179 (1964) 544
 1964KU05 H.-M. Kuan, T.W. Bonner and J.R. Risser, Nucl. Phys. 51 (1964) 481
 1964KU06 H.-M. Kuan and J.R. Risser, Nucl. Phys. 51 (1964) 518
 1964LI14 I. Lindgren, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 379
 1964LI1B Lindskog, Sundstrom and Sparrman, Perturbed Angular Correlations; Eds., E. Karlsson, E. Matthias and K. Siegbahn (1964) 411
 1964LI1C Litherland, Nucl. Instrum. Meth. 28 (1964) 55
 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
 1964MA25 B.S. Madsen and M. Vedelsby, Nucl. Phys. 55 (1964) 477
 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed., W.H. Johnson Jr. (1964) 305
 1964MO1G Moringo, Phys. Rev. 134 (1964) B1243

- 1964NE09 J.B. Nelson, E.L. Hudspeth and E.M. Bernstein, Phys. Rev. 136 (1964) B71
- 1964OS01 D.R. Osgood, J.R. Patterson and E.W. Titterton, Nucl. Phys. 60 (1964) 503
- 1964PA11 G. Paic, I. Slaus and P. Thomas, Phys. Lett. 9 (1964) 147
- 1964RI1A Ripka, 3A(II)/C174, Paris (1964)
- 1964SA1D Satchler, Drisko and Bassel, Phys. Rev. 136 (1964) B637
- 1964SA1E Saint-Pierre, Photographie Corpusculaire III, Ed., Demers (1964) 462
- 1964SC01 J. Schaitman, H.F. Lutz and S.F. Eccles, Bull. Amer. Phys. Soc. 9 (1964) 68, GC10
- 1964SC09 R.B. Schwartz, H.D. Holmgren, L.M. Cameron and A.R. Knudson, Phys. Rev. 134 (1964) B577
- 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
- 1964ST1J Stanojevic, Bull. Boris Kidrich Inst. Nucl. Sci. 15 (1964) 167
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964TA05 N.W. Tanner, G.C. Thomas and E.D. Earle, Nucl. Phys. 52 (1964) 29
- 1964TA1C Tanner and Earle, Phys. Rev. Lett. 13 (1964) 410
- 1964VA1D L. Valentin, G. Albouy, J.P. Cohen and M. Gusakow, J. Phys. (France) 25 (1964) 704
- 1965AJ01 F. Ajzenberg-Selove, J.W. Watson and R. Middleton, Phys. Rev. 139 (1965) B592
- 1965AL05 W.P. Alford, L.M. Blau and D. Cline, Nucl. Phys. 61 (1965) 368
- 1965AL19 T.K. Alexander, A.E. Litherland and C. Broude, Can. J. Phys. 43 (1965) 2310
- 1965AL1J Alevra et al., Stud. Cercetari Fiz. (Rumania) 17 (1965) 761
- 1965AR07 K.P. Artemov, V.Z. Goldberg, B.I. Islamov, V.P. Rudakov and I.N. Serikov, Yad. Fiz. 1 (1965) 1019; Sov. J. Nucl. Phys. 1 (1965) 726
- 1965BE1B Becker and McIntyre, Phys. Rev. 138 (1965) B339
- 1965BO14 R. Bock, M. Grosse-Schulte, W. Von Oertzen and R. Rudel, Phys. Lett. 18 (1965) 45
- 1965BO42 M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, H. Buttner, A. Lindner and H. Meldner, Nucl. Phys. 63 (1965) 438
- 1965BR1F Breit, Ann. Phys. 34 (1965) 377
- 1965BR42 O.D. Brill, Yad. Fiz. 1 (1965) 55; Sov. J. Nucl. Phys. 1 (1965) 37
- 1965BU05 A. Bussiere, N.K. Glendenning, B.G. Harvey, J. Mahoney, J.R. Meriwether and D.J. Horen, Phys. Lett. 16 (1965) 296
- 1965CA14 J.T. Caldwell, R.L. Bramblett, B.L. Berman, R.R. Harvey and S.C. Fultz, Phys. Rev. Lett. 15 (1965) 976
- 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707

1965DE1A Detraz, Cahiers Phys. 19 (1965) 359
 1965DE1G Degtyarev, Atomnaya Energiya 19 (1965) 456
 1965DE24 V.P. Denisov and L.A. Kulchitskii, Yad. Fiz. 2 (1965) 70; Sov. J. Nucl. Phys. 2 (1966) 48
 1965DI07 G.U. Din and J.L. Weil, Nucl. Phys. 73 (1965) 161
 1965DI1E Dickens, Drisko, Perey and Satchler, Phys. Lett. 15 (1965) 337
 1965DO14 R.E. Donaldson, L. Passell, W. Bartolini and D. Groves, Phys. Rev. 138 (1965) B1116
 1965EN01 J.B.A. England and B.L. Reece, Nucl. Phys. 72 (1965) 449
 1965FA1B Fabre de La Ripelle, Prog. Theor. Phys. 33 (1965) 38
 1965FI05 P. Fintz, Ann. Phys. (Paris) 10 (1965) 435
 1965FR1E Frahn and Wiechers, Nucl. Phys. 74 (1965) 65
 1965FU16 H.W. Fulbright, W.P. Alford, O.M. Bilaniuk, V.K. Deshpande and J.W. Verba, Nucl. Phys. 70 (1965) 553
 1965GA1B Gaedke, Toth and Williams, Phys. Rev. 140 (1965) B296
 1965GA1D Gaponov, Yad. Fiz. 2 (1965) 1002
 1965GA1G Garber and Shrader, Bull. Amer. Phys. Soc. 10 (1965) 510
 1965GI1B Giraud, Nucl. Phys. 71 (1965) 373
 1965GL04 R.N. Glover and A.D.W. Jones, Phys. Lett. 16 (1965) 69
 1965GR1H Grechukhin, Nucl. Phys. 62 (1965) 273
 1965GR1R Gromov et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P-2184 (1965)
 1965GR41 B. Grimeland, E. Kjellsby and J. Vines, Phys. Rev. 137 (1965) B878
 1965GU1A Guiasu and Micu, Rev. Phys. (Rumania) 10 (1965) 555
 1965HA1G Havlicek and Modesto, Energ. Nucl. 12 (1965) 77
 1965HE1B Henkel, Bull. Amer. Phys. Soc. 10 (1965) 601
 1965HI1A Hiebert, McIntyre and Couch, Phys. Rev. 138 (1965) B346
 1965HU1D Hull and Shakin, Phys. Lett. 19 (1965) 506
 1965IC1A Ichimura and Yazaki, Nucl. Phys. 63 (1965) 401
 1965JA09 H.E. Jackson, A.I. Namenson and G.E. Thomas, Phys. Lett. 17 (1965) 324
 1965LA09 J.M. Lacambra, D.R. Tilley, N.R. Roberson and R.M. Williamson, Nucl. Phys. 68 (1965) 273
 1965MA1K Marion, Nucl. Phys. 68 (1965) 463
 1965MA1T Mahaux, Nucl. Phys. 67 (1965) 358

1965MA1V Markowitz and Pape, Bull. Amer. Phys. Soc. 10 (1965) 602
 1965MA45 J.M. Maison, M. Langevin and J.M. Loiseaux, Phys. Lett. 19 (1965) 308
 1965MO13 R.C. Morrison, J.R. Stewart and J.S. O'Connell, Phys. Rev. Lett. 15 (1964) 367
 1965NE1D Newton, Phys. Lett. 17 (1965) 132
 1965PE1H Perez Dominguez, Jen 143 DF/1-46 (1965)
 1965RO1N Rose and Lopes, Phys. Lett. 18 (1965) 130
 1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, Ann. Phys. 34 (1965) 96
 1965SC1D Schapira, Ann. Physique 10 (1965) 743
 1965SE05 R. Seltz, C. Gerardin, M. Wery and D. Magnac-Valette, Compt. Rend. 261 (1965) 391
 1965SH1E Shapiro and Timashev, Yad. Fiz. 2 (1965) 459
 1965SI13 R.H. Siemssen, M. Cosack and R. Felst, Nucl. Phys. 69 (1965) 227
 1965SL1C Slee, Bull. Amer. Phys. Soc. 10 (1965) 461
 1965TA07 T. Takemiya, Prog. Theor. Phys. 34 (1965) 433
 1965TS1A Tsenter and Silin, Atomnaya Energ. 19 (1965) 48
 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331
 1965VA23 L. Valentin, Nucl. Phys. 62 (1965) 81
 1965WA03 E.K. Warburton, K.W. Jones, D.E. Alburger, C. Chasman and R.A. Ristinen, Phys. Rev. Lett. 14 (1965) 146
 1965WA06 E.K. Warburton, J.S. Lopes, R.W. Ollerhead, A.R. Poletti and M.F. Thomas, Phys. Rev. 138 (1965) B104
 1965WA16 E.K. Warburton, J.W. Olness and D.E. Alburger, Phys. Rev. 140 (1965) B1202
 1965WA17 E.K. Warburton, P.D. Parker and P.F. Donovan, Phys. Lett. 19 (1965) 397
 1965WI03 E.J. Winhold, R.H. Augustson, N.N. Kaushal, H.A. Medicus, W.R. Moyer and P.F. Yergin, Bull. Amer. Phys. Soc. 10 (1965) 95, HD6
 1965WI1A Williams, Gaedke and Toth, Bull. Amer. Phys. Soc. 10 (1965) 443
 1965ZA1B Zamick, Phys. Lett. 19 (1965) 580
 1966AG1A Agee and Rosen, LA-3538-MS (1966)
 1966AG1B Aguilar, de la Rubia, Sanchez and Martinez, An. Fisica Y Quim. 62 (1966) 279
 1966AL12 D.E. Alburger and K.W. Jones, Phys. Rev. 149 (1966) 743
 1966AL18 D.E. Alburger and E.K. Warburton, Phys. Rev. 152 (1966) 914
 1966AV1A P. Avignon, L.H. Rosier and Y. Deschamps, J. Phys. 27 (1966) C1-40
 1966BA1F Baumgartner et al., Helv. Phys. Acta 39 (1966) 575

1966BE1M Bercaw, Boschitz and Vincent, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (1966) 334

1966BL01 R.S. Blake, D.J. Jacobs, J.O. Newton and J.P. Schapira, Nucl. Phys. 77 (1966) 254

1966BO1R O. Bohigas, J. Phys. 27 (1966) C1-39

1966BO22 R. Bock, M. Grosse-Schulte and W. von Oertzen, Phys. Lett. 22 (1966) 456

1966BR09 R.I. Brown, Nucl. Phys. 78 (1966) 492

1966BR13 K.H. Bray, J. Nurzynski and W.P. Bourke, Phys. Lett. 21 (1966) 536

1966BU1B Buttle and Goldfarb, Nucl. Phys. 78 (1966) 409

1966CA1H Catala, Garcia and Perez, An. Real. Soc. Espan. Fis. y Quim. A61 (1966) 357

1966CI01 S.D. Cirilov, J.O. Newton and J.P. Schapira, Nucl. Phys. 77 (1966) 472

1966CS1B Csikai and Nagy, Acta Phys. Acad. Sci. Hung. 21 (1966) 303

1966CS1C Csikai, Antwerp 1965 Neutron Conf. (1966) 537

1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097

1966DR02 L. Drigo, C. Manduchi, G.C. Nardelli, M.T. Russo-Manduchi, G. Torielli and G. Zannoni, Nuovo Cim. B45 (1966) 206

1966DU1B Duggan et al., Bull. Amer. Phys. Soc. 11 (1966) 831

1966EC1B Eccles, Lutz and Rohn, Bull. Amer. Phys. Soc. 11 (1966) 735

1966ED1A J.A. Edgington and B. Rose, Nucl. Phys. 89 (1966) 523

1966EL08 F. El-Batanoni and A.A. Kresnin, Nucl. Phys. 89 (1966) 577

1966EV01 A.E. Evans, B. Brown and J.B. Marion, Phys. Rev. 149 (1966) 863

1966FO1D Fowler and Johnson, Bull. Amer. Phys. Soc. 11 (1966) 510; ORNL P-2026 (1966)

1966FO1E Fowler, Johnson and Kernell, Bull. Amer. Phys. Soc. 11 (1966) 653

1966GA04 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 141 (1966) 996

1966GA08 A. Gallman, P. Fintz, J.B. Nelson and D.E. Alburger, Phys. Rev. 147 (1966) 753

1966GA09 A. Gallman, P. Fintz and P.E. Hodgson, Nucl. Phys. 82 (1966) 161

1966GA19 A. Gallmann, F. Haas and N. Balaux, Phys. Rev. 151 (1966) 735

1966GA25 G.T. Garvey and I. Kelson, Phys. Rev. Lett. 16 (1966) 197

1966GL01 R.N. Glover and A.D.W. Jones, Nucl. Phys. 84 (1966) 673

1966GO15 S. Gorodetzky, R.M. Freeman, A. Gallman and F. Hass, Phys. Rev. 149 (1966) 801

1966GO1E Gorodetzky et al., Nucl. Instrum. Meth. 42 (1966) 269

1966GO1J S. Gordetzky, A. Gallmann and P. Fintz, J. Phys. 27 (1966) C1-91

1966GR1A Gruhn and Kashy, Bull. Amer. Phys. Soc. 11 (1966) 471

- 1966HA19 B.G. Harvey, J.R. Meriwether, J. Mahoney, A. Bussiere de Nercy and D.J. Horen, Phys. Rev. 146 (1966) 712
- 1966HA21 R.L. Hahn and E. Ricci, Phys. Rev. 146 (1966) 650
- 1966HA30 O. Hausser, R.D. Gill, J.S. Lopes and H.J. Rose, Nucl. Phys. 84 (1966) 683
- 1966HA31 O. Hausser, H.J. Rose, J.S. Lopes and R.D. Gill, Phys. Lett. 22 (1966) 604
- 1966HI1C B. Hird, Nucl. Phys. 86 (1966) 268
- 1966HO1D Hodgson, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 71
- 1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KL06 A.P. Klyucharev and Y.I. Titov, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 435; Bull. Acad. Sci. USSR Phys. 30 (1966) 443
- 1966KL1F Klyucharev, Titov and Vipirailenko, Izv. Vys. Uch. Zaved. Fiz. 3 (1966) 104
- 1966KO1G Komar, Denisov and Kulchitskii, Dokl. Akad. Nauk SSSR 169 (1966) 1307
- 1966KU12 I.V. Kurdyumov, S.H.El Samarai, Y.F. Smirnov and K.V. Shitikova, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 292; Bull. Acad. Sci. USSR Phys. 30 (1966) 297
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966LI07 K.P. Lieb, Nucl. Phys. 85 (1966) 461
- 1966LO02 J.S. Lopes, O. Hausser, H.J. Rose, A.R. Poletti and M.F. Thomas, Nucl. Phys. 76 (1966) 223
- 1966LO1N Louis and Nussbaum, Helv. Phys. Acta 39 (1966) 11
- 1966MA02 A.M. MacLeod and J.M. Reid, Proc. Phys. Soc. (London) 87 (1966) 437
- 1966MA04 G.S. Mani and G.C. Dutt, Nucl. Phys. 78 (1966) 613; Erratum Nucl. Phys. A119 (1968) 691
- 1966MA1A Marion, Ludemann and Roos, Bull. Amer. Phys. Soc. 11 (1966) 332
- 1966MA1R J.H. Manley and W.E. Stein, Phys. Rev. 144 (1966) 956
- 1966MA1V H.A. Mavromatis and L. Zamick, Phys. Lett. 20 (1966) 171
- 1966MA2K Martin, Kernell and Fowler, Bull. Amer. Phys. Soc. 11 (1966) 808
- 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
- 1966ME14 D.F. Measday and J.N. Palmieri, Nucl. Phys. 85 (1966) 129
- 1966MI1F Migdal, Proc. Int. School Enrico Fermi, Course 36; Ed., C. Bloch (1966) 171
- 1966OL1C Olariu, Acad. Repub. Pop. Romine. Inst. de Fiz. Atom. Rept. IS 25 (1966)
- 1966OW01 R.O. Owens and J.E.E. Baglin, Phys. Rev. Lett. 17 (1966) 1268

1966PE04 D. Pelte, B. Povh and W. Scholz, Nucl. Phys. 78 (1966) 241
 1966PH1B Phillips, NASA CR 94632 (1966)
 1966PO09 A.M. Poskanzer, S.W. Cospers, E.K. Hyde and J. Cerny, Phys. Rev. Lett. 17 (1966) 1271
 1966PO11 A.R. Poletti, J.W. Olness and E.K. Warburton, Phys. Rev. 151 (1966) 812
 1966PO1E Poth and Bromley, Bull. Amer. Phys. Soc. 11 (1966) 317
 1966PR1B Prusser, 2nd Symp. on the Struct. of Low-Medium Mass Nucl., April 1966 (1966) 38
 1966RE1D H. Reeves, J. Phys. 27 (1966) C1-28
 1966RI1F Ripka, Lect. in Theor. Phys., Vol. VIII C (1966) 237
 1966RO1B Rosen, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (1966) 253
 1966RO1P H.J. Rose, J.S. Lopes and W. Greiner, Phys. Lett. 19 (1966) 686
 1966RO1R Rosen, Antwerp 1965 Neutron Conf. (1966) 379
 1966RO1U H.J. Rose and J.S. Lopes, Phys. Lett. 22 (1966) 601
 1966RO1V B.A. Robson, Nucl. Phys. 86 (1966) 649
 1966SC12 J.P. Schapira, J.O. Newton, R.S. Blake and D.J. Jacobs, Nucl. Phys. 80 (1966) 565
 1966SC21 W. Scobel, R.W. Fink and M. Bormann, Z. Phys. 197 (1966) 124
 1966SC22 J.J. Schwartz, W.P. Alford, L.M. Blau and D. Cline, Nucl. Phys. 88 (1966) 539
 1966SE1D R. Seltz, C. Gerardin, M. Wery and D. Magnac-Valette, J. Phys. 27 (1966) C1-148
 1966SH1A Shute and Brown, Bull. Amer. Phys. Soc. 11 (1966) 471
 1966SH1F I.S. Shapiro and S.F. Timashev, Nucl. Phys. 79 (1966) 46
 1966SO05 M. Soga, Nucl. Phys. 89 (1966) 697
 1966ST05 J.D. Steben and M.K. Brussel, Phys. Rev. 146 (1966) 780
 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692
 1966WA08 E.K. Warburton and J.W. Olness, Phys. Rev. 147 (1966) 698
 1966WA1E Warburton, F. S. U. Isobaric Spin Conf. (1966) 90
 1966WA1K G.E. Walker, Phys. Rev. 151 (1966) 745
 1966WE1B Weinberg, Antwerp 1965 Neutron Conf. (1966) 37
 1966WI1E D.H. Wilkinson and M.E. Mafethe, Nucl. Phys. 85 (1966) 97
 1967AD1F Adelberger, Cocke and Davids, Bull. Amer. Phys. Soc. 12 (1967) 1194
 1967AL16 A.U. Aldzhauakhiri, A.V. Spasskii, I.B. Teplov and L.N. Fateeva, Yad. Fiz. 6 (1967) 248; Sov. J. Nucl. Phys. 6 (1968) 180

- 1967AM1H Amiet, Ebenhoh and Huguenin, *Helv. Phys. Acta* 40 (1967) 283
- 1967AU1A Audouze, Epherre and Reeves, *High Energy Nucl. Reactions in Astrophys.*, Ed., B.S.P. Shen (1967) 255
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, *Nucl. Phys.* A97 (1967) 144
- 1967BA03 P.W. Bauer, J.D. Anderson, H.F. Lutz, C. Wong, J.W. McClure and B.A. Pohl, *Nucl. Phys.* A93 (1967) 673
- 1967BA13 G.C. Ball and J. Cerny, *Phys. Rev.* 155 (1967) 1170
- 1967BE09 H. Beaumevieille, M. Lambert, M. Yaker and A. Amokrane, *Nuovo Cim.* B47 (1967) 139
- 1967BE1F Beery, Harper, Stovall and Rosen, *Los Alamos Sci. Lab. Rept.* LA-3788 (1967)
- 1967BI11 P.G. Bizzeti, A.M. Bizzeti-Sona, S. Kalbitzer and B. Povh, *Nucl. Phys.* A104 (1967) 577
- 1967BO1T Bohigas, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (1967) 940
- 1967BR1M Breit, Polak and Torchia, *Phys. Rev.* 161 (1967) 993
- 1967BR1Q G. Breit, *Proc. Nat. Acad. Sci.* 67 (1967) 849
- 1967CA1C Caldwell, UCRL 50287 (1967)
- 1967CA1P Caldwell, Fultz and Bramblett, *Bull. Amer. Phys. Soc.* 12 (1967) 197
- 1967CH15 N.S. Chant, P.S. Fisher and D.K. Scott, *Nucl. Phys.* A99 (1967) 669
- 1967CH19 C. Chasman, K.W. Jones, R.A. Ristinen and D.E. Alburger, *Phys. Rev.* 159 (1967) 830
- 1967CH35 R. Chiba, *Nucl. Sci. App.* 3 (1967) 36
- 1967CL1C Cloud, Leonard, Gibson and Wells, *Bull. Amer. Phys. Soc.* 12 (1967) 894
- 1967CO1D Commins, *Ann. Rev. Nucl. Sci.* 17 (1967) 33
- 1967CO1R Coppola and Knitter, *Eur* 3492.E (1967)
- 1967CO32 S. Cohen and D. Kurath, *Nucl. Phys.* 101 (1967) 1
- 1967CS02 J. Csikai and G. Peto, *Acta Phys. Acad. Sci. Hung.* 23 (1967) 87
- 1967CS03 J. Csikai and S. Nagy, *Nucl. Phys.* A91 (1967) 222
- 1967DA1E Dalidtschik and Saiasov, *Proc. Prob. Symp. on Nucl. Phys.*, Tbilishi, April 1967 (1967) 287
- 1967DE14 D. Dehnhard, D.S. Gemmell and Z. Vager, *Nucl. Phys.* A104 (1967) 202
- 1967DE1K Dehnhard and Siemssen, *Bull. Amer. Phys. Soc.* 12 (1967) 17
- 1967EL03 J.P. Elliott, H.A. Mavromatis and E.A. Sanderson, *Phys. Lett.* B24 (1967) 358
- 1967EV02 A.E. Evans, *Phys. Rev.* 155 (1967) 1047

1967FA1A W.M. Fairbairn, Nucl. Phys. A90 (1967) 135
 1967FI1E Firk, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 352
 1967FU1A Funsten and Holt, Bull. Amer. Phys. Soc. 12 (1966) 1176
 1967GO07 S. Gorodetzky, R.M. Freeman, A. Gallmann, F. Haas and B. Heusch, Phys. Rev. 155 (1967) 1119
 1967GR1K Gradsztajn, High Energy Nucl. Reactions in Astrophys., Ed., B.S.P. Shen (1967) 247
 1967GR1L Gray, Fortune, Trost and Fletcher, Bull. Amer. Phys. Soc. 12 (1967) 34
 1967HE1A Hensley, Astrophys. J. 147 (1967) 818
 1967HI06 J.C. Hiebert, E. Newman and R.H. Bassel, Phys. Rev. 154 (1967) 898
 1967IV1B Ivascu, Dumitrescu and Semenescu, Rev. Roum. Phys. 12 (1967) 279
 1967KO1G Kobzev et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P6 3314 (1967)
 1967KR1F Krivopustov, Sizov, Shirmer and Oehler, P15-3504, Dubna (1967)
 1967KU1E D. Kurath and R.D. Lawson, Phys. Rev. 161 (1967) 915
 1967KU1M Kuan, Hanna and Hasinoff, Bull. Amer. Phys. Soc. 12 (1967) 52
 1967LA05 M. Lambert and M. Durand, Phys. Lett. B24 (1967) 287
 1967LA10 M. Lambert and M. Durand, J. Phys. (Paris) 28 (1967) 349
 1967LA11 B. Lawergren and I.V. Mitchell, Nucl. Phys. A98 (1967) 481
 1967LI06 R.H. Lindsay and J.J. Veit, Phys. Rev. 157 (1967) 933
 1967LI1B Lingenfelter and Ramaty, High Energy Nucl. Reactions in Astrophys., Ed., B.S.P. Shen (1967) 99
 1967LO03 P. Loncke and J. Pradal, Nuovo Cim. B48 (1967) 457
 1967MA1M Marr, Kuenhold and Donoghue, Bull. Amer. Phys. Soc. 12 (1967) 501
 1967ME1N Meier, Thomason and Walter, Bull. Amer. Phys. Soc. 12 (1967) 1197
 1967MO21 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. A103 (1967) 238
 1967MU12 S. Mubarakmand and B.E.F. Macefield, Nucl. Phys. A98 (1967) 82
 1967NE06 J.B. Nelson and W.R. Smith, Nucl. Phys. A96 (1967) 671
 1967NE1D Nemirovskii, Sov. J. Nucl. Phys. 4 (1967) 334
 1967PA05 J.C. Parikh and N. Ullah, Nucl. Phys. A99 (1967) 529
 1967PA27 A. Pasquarelli, Nucl. Phys. A93 (1967) 218
 1967PE1D J.-L. Perrenoud and E. Sheldon, Nucl. Phys. A102 (1967) 105
 1967PH03 G.W. Phillips, F.C. Young and J.B. Marion, Phys. Rev. 159 (1967) 891
 1967PO13 J.E. Poth, J.C. Overley and D.A. Bromley, Phys. Rev. 164 (1967) 1295

1967PO1E Poth and Bromley, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (1967) 94
 1967PO1J A.R. Poletti, E.K. Warburton and D. Kurath, Phys. Rev. 155 (1967) 1096
 1967RE01 D. Rendic, Nucl. Phys. A91 (1967) 604
 1967RO1K Robertson, Aust. J. Phys. 20 (1967) 489
 1967SC27 L.A. Schaller, R.S. Thomason, N.R. Roberson, R.L. Walter and R.M. Drisko, Phys. Rev. 163 (1967) 1034
 1967SC29 J.P. Schiffer, G.C. Morrison, R.H. Siemssen and B. Zeidman, Phys. Rev. 164 (1967) 1274
 1967SE08 W.A. Seale, Phys. Rev. 160 (1967) 809
 1967SH14 V.S. Shirley, UCRL-17990 (1967)
 1967SO1B Soltesz et al., Bull. Amer. Phys. Soc. 12 (1967) 1198
 1967ST21 B. Stepanic, R. Popic and M. Aleksic, Phys. Rev. Lett. 19 (1967) 1137
 1967TH05 G.E. Thomas, D.E. Blatchley and L.M. Bollinger, Nucl. Instrum. Meth. 56 (1967) 325
 1967VO1A Volkov, Proc. Problem Symp. on Nucl. Phys., Tbilist, April 1967 (1967) 226
 1967VO1B Vogt, Int. Nucl. Phys. Conf., Gatlinburg 1966 (1967) 748
 1967WA1C Warburton, Nucl. Research with Low Energy Accelerators, Eds., Marion and van Patter (1967) 43
 1967WE1B West, ORNL TM 1926 (1967)
 1967YO1C Young, Nucl. Research with Low Energy Accelerators, Eds., Marion and van Patter (1967) 109
 1967ZH1A Zhusupov, Karapetian and Eramzhian, Joint Inst. Nucl. Res., Lab. Theor. Phys., USSR Rept. No. P4 3177 (1967)
 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
 1968BA1E Ball, UCRL-18263 (1968)
 1968BA2J Bassel and Drisko, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, Sept. 1967 (1968) 13
 1968BA2L Baglin and Thompson, in Tokyo (1968) 388
 1968BA42 A.R. Barnett, Nucl. Phys. A120 (1968) 342
 1968BE14 G.A. Beer, P. Brix, H.-G. Clerc and B. Laube, Phys. Lett. B26 (1968) 506
 1968BI1C Bizzeti, Suppl. Nuovo Cim. 6 (1968) 664
 1968BO14 W. Bohne, H. Homeyer, H. Morgenstern and J. Scheer, Nucl. Phys. A113 (1968) 97
 1968BO36 F. Boreli, P.N. Shrivastava, B.B. Kinsey and V.C. Mistry, Phys. Rev. 174 (1968) 1221
 1968CA1J Campbell, Harvey and Slaughter, Bull. Amer. Phys. Soc. 13 (1968) 1423

- 1968CO04 R.L.A. Cottrell, J.C. Lisle and J.O. Newton, Nucl. Phys. A109 (1968) 288
- 1968CO1W Conde, Bergqvist and Nystrom, Neutron Cross Sect. Tech., NBS Special Pub. 299 (1968) 763
- 1968COZV P.M. Cockburn, R.W. Krone and H.E. Siefken, Bull. Amer. Phys. Soc. 13 (1968) 1423, DC4
- 1968DA1F J.C. Davis and H.H. Barschall, Phys. Lett. B27 (1968) 636
- 1968DA1N Dangle, Duncan, Duggan and Miller, Bull. Amer. Phys. Soc. 13 (1968) 608
- 1968DA1Q Dally, Croissiaux and Schweitz, Bull. Amer. Phys. Soc. 13 (1968) 607
- 1968DE13 N. de Takacsy, Can. J. Phys. 46 (1968) 2091
- 1968DU1F Duorah and Duorah, Ind. J. Pure Appl. Phys. 6 (1968) 389
- 1968EL1A Elliott, 3rd Symp. on the Struct. of Low-Medium Mass Nucl., Ed., Davidson (1968) 48
- 1968FA1A J.C. Faivre, H. Krivine and A.M. Papiou, Nucl. Phys. A108 (1968) 508
- 1968FO06 H.T. Fortune, T.J. Gray, W. Trost and N.R. Fletcher, Phys. Rev. 173 (1968) 1002
- 1968FO1A Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299 (1968) 1
- 1968GA03 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 167 (1968) 957
- 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161
- 1968GA1C Gaillard, Univ. Lyon, Rept. No. Lycen 6828 (1968)
- 1968GA1M E. Gadioli and L. Zetta, Phys. Rev. 167 (1968) 1016
- 1968GE04 B. Geoffrion, N. Marty, M. Morlet, B. Tatischeff and A. Willis, Nucl. Phys. A116 (1968) 209
- 1968GI01 R.D. Gill, J.S. Lopes, B.C. Robertson, R.A.I. Bell and H.J. Rose, Nucl. Phys. A106 (1968) 678
- 1968GI11 R.D. Gill, J.S. Lopes, O. Hausser and H.J. Rose, Nucl. Phys. A121 (1968) 209
- 1968GR14 R.C. Greenwood, Phys. Lett. B27 (1968) 274
- 1968GRZY R.C. Greenwood, Proc. Conf. Slow-Neutron-Capture Gamma-Ray Spectr., Argonne, Ill., 1966; Ed., F.E. Throw, ANL-7282 (1968) 303
- 1968HA15 J.S. Hanna and M.A. Nagarajan, Nucl. Phys. A113 (1968) 412
- 1968HA1V Hansen, Anderson, Stelts and Wong, Neutron Cross Sect. Tech., NBS Special Pub. 299 (1968) 225
- 1968HA27 W.R. Harris and J.C. Armstrong, Phys. Rev. 171 (1968) 1230
- 1968HE12 J.D. Henderson, E.L. Hudspeth and W.R. Smith, Phys. Rev. 172 (1968) 1058
- 1968HI1J Hinderliter and Lochstet, Bull. Amer. Phys. Soc. 13 (1968) 606

1968HO1H Horie and Hsieh, in Tokyo (1968) 51
 1968HU1A R.L. Hutson, S. Hayakawa, M. Chabre, J.J. Kraushaar, B.W. Ridley and E.T. Boschitz,
 Phys. Lett. B27 (1968) 153
 1968IW1A Iwao and Kawakami, Prog. Theor. Phys. 40 (1968) 1046
 1968JA1M Jacquot et al., Compt. Rend. B266 (1968) 1286
 1968JO1F Johnson et al., Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 851
 1968JO1H Jones and Toms, Bull. Amer. Phys. Soc. 13 (1968) 718
 1968JU1B Jung et al., Compt. Rend. 266 (1968) 1154
 1968KU1F Kuan and O'Connell, Bull. Amer. Phys. Soc. 13 (1968) 85
 1968LA19 C.M. Lamba, N. Sarma and N.S. Thampi, Nucl. Phys. A122 (1968) 390
 1968LE01 J.K.P. Lee, S.K. Mark, P.M. Portner and R.B. Moore, Nucl. Phys. A106 (1968) 357
 1968LI1A Li and Mark, Bull. Amer. Phys. Soc. 13 (1966) 679
 1968LI1L Liebenauer, Silverstein, Kibler and Koral, Bull. Amer. Phys. Soc. 13 (1968) 1388
 1968MA2B Manakos, Z. Phys. 214 (1968) 57
 1968MA2G R.M. May, Phys. Lett. B26 (1968) 482
 1968ME08 R.A. Mendelson Jr. and R.T. Carpenter, Phys. Rev. 166 (1968) 988
 1968NA1F Nakamura, in Tokyo (1968) 228
 1968PA1F J.Y. Park, Nucl. Phys. A111 (1968) 433
 1968PE16 A.K. Petrauskas and V.V. Vanagas, Yad. Fiz. 8 (1968) 463; Sov. J. Nucl. Phys. 8
 (1969) 270
 1968PU1A H.G. Pugh, Phys. Rev. Lett. 20 (1968) 601
 1968RI1T Ritter and Parson, Bull. Amer. Phys. Soc. 13 (1968) 607
 1968RO1E Rohl, Z. Phys. 215 (1968) 56
 1968SC1B Schevchenko, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc.
 Jpn. 24 (1968) 397
 1968SH08 A.P. Shukla and G.E. Brown, Nucl. Phys. A112 (1968) 296
 1968SH11 P.N. Shrivastava, F. Boreli and B.B. Kinsey, Phys. Rev. 169 (1968) 842
 1968SI1E Siefken, Cockburn and Krone, Bull. Amer. Phys. Soc. 13 (1968) 1423
 1968SI1F Siefken, COO-1120-75 (1968)
 1968ST06 A.P. Stamp and M.B. Spencer, Nucl. Phys. A111 (1968) 353
 1968ST10 C.E. Steerman and F.C. Young, Phys. Lett. B27 (1968) 8
 1968ST19 T. Stambach, R.S. Thomason, J. Taylor and Jr., R.L. Walter, Phys. Rev. 174 (1968)
 1119

- 1968TU02 P.M. Tutakin, *Yad. Fiz.* 8 (1968) 661; *Sov. J. Nucl. Phys.* 8 (1969) 383
- 1968WA04 G.J. Wagner, *Phys. Lett.* B26 (1968) 429
- 1968WE15 H.R. Weller, N.R. Roberson and D.R. Tilley, *Nucl. Phys.* A122 (1968) 529
- 1968WO1C S.S.M. Wong, *Nucl. Phys.* A120 (1968) 625
- 1968WO1D Wong et al., in *Tokyo* (1968) 249
- 1968YA1E F.B. Yano, *Nucl. Phys.* A118 (1968) 592
- 1968ZH05 M.A. Zhusupov, O. Lkhagva and I. Rotter, *Izv. Akad. Nauk SSSR Ser. Fiz.* 32 (1968) 1714; *Bull. Acad. Sci. USSR Phys. Ser.* 32 (1969) 1579
- 1968ZH06 M.Z. Zhusupov and R.A. Eramzhyan, *Izv. Akad. Nauk SSSR Ser. Fiz.* 32 (1968) 2060; *Bull. Acad. Sci. USSR Phys. Ser.* 32 (1969) 1894
- 1968ZH1B Zhusupov, Karapetyan and Eramzhyan, *Izv. Akad. Nauk SSSR Ser. Fiz.* 32 (1968) 332
- 1968ZU02 A.P. Zuker, B. Buck and J.B. McGrory, *Phys. Rev. Lett.* 21 (1968) 39
- 1969AL04 W.P. Alford and K.H. Purser, *Nucl. Phys.* A132 (1969) 86
- 1969AL1H Alsmiller, Wachter and Moran, *Nucl. Sci. Eng.* 36 (1969) 291
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, *Nucl. Phys.* A137 (1969) 348
- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, *Nucl. Phys.* A126 (1969) 60
- 1969BA06 G.C. Ball and J. Cerny, *Phys. Rev.* 177 (1969) 1466
- 1969BA1N Bahcall and Fowler, *Astrophys. J.* 157 (1969) 659
- 1969BA1P Bacon et al., *Acta Cryst.* A25 (1969) 391
- 1969BA2U Ballini, Delaunay, Fouan and Tellez, *Addendum to Contrib., Montreal* (1969) 6
- 1969BE08 H. Beaumevieille, M. Lambert, M. Yaker, A. Amokrane and Nguyen van Sen, *Nucl. Phys.* A125 (1969) 568
- 1969BO13 W. Bohne, H. Homeyer, H. Lettau, H. Morgenstern, J. Scheer and F. Sichelschmidt, *Nucl. Phys.* A128 (1969) 537
- 1969BR07 K.H. Bray and J. Nurzynski, *Nucl. Phys.* A127 (1969) 622
- 1969BR1D Bromly, *Proc. Enrico Fermi School of Phys., Course XL, Lake Como 1967* (1969) 242
- 1969BR1G Bromley, *Proc. Int. Conf., Montreal* (1969) 147
- 1969BR30 O.D. Brill, A.D. Vongai and A.A. Ogloblin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 33 (1969) 615; *Bull. Acad. Sci. USSR Phys. Ser.* 33 (1970) 567
- 1969CH1R M. Chemtob, *Nucl. Phys.* A123 (1969) 449

- 1969CO15 G.W. Cole Jr., F.W.K. Firk and T.W. Phillips, Phys. Lett. B30 (1969) 91
- 1969DE16 J. Dey, J.P. Elliott, A.D. Jackson, H.A. Mavromatis, E.A. Sanderson and B. Singh, Nucl. Phys. A134 (1969) 385
- 1969DE1H Debenham, Griffith, Irshad and Roman, Contrib., Montreal (1969) 274
- 1969DI1B Dickens and Perey, Nucl. Sci. Eng. 36 (1969) 280
- 1969DO04 H. Doubre, D. Royer, M. Arditi, L. Bimbot, N. Frascaria, J.P. Garron and M. Riou, Phys. Lett. B29 (1969) 355
- 1969EL1B Elliott, Proc. Int. Conf., Montreal (1969) 277
- 1969EN03 J.B.A. England, R.G. Harris, L.H. Watson, D.H. Worledge and J.E. Evans, Phys. Lett. B30 (1969) 476
- 1969ET01 K. Etoh, T. Murata, N. Kawai, R. Chiba and S. Takayanagi, J. Phys. Soc. Jpn. 26 (1969) 1335, and Private Communication (1969)
- 1969FO02 H.T. Fortune, T.J. Gray, W. Trost and N.R. Fletcher, Phys. Rev. 179 (1969) 1033
- 1969FO1D Fowler, Contemp. Phys. 1 (1969) 359
- 1969FR20 D.E. Frederick, R.J.J. Stewart and R.C. Morrison, Phys. Rev. 186 (1969) 992
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA05 A. Gallmann, F. Jundt, E. Aslanides and D.E. Alburger, Phys. Rev. 179 (1969) 921
- 1969GA11 P. Gaillard, R. Bouche, L. Feuvrais, M. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.R. Pizzi, Nucl. Phys. A131 (1969) 353
- 1969GI1B Gillet, Proc. Int. Conf., Montreal (1969) 483
- 1969GL07 Y.A. Glukhov, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta and D.N. Stepanov, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 601; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 554
- 1969GL1D Glenn, Zafiratos and Zaidins, Bull. Amer. Phys. Soc. 14 (1969) 1227
- 1969GO14 V. Gomes Porto, N. Ueta, R.A. Douglas, O. Sala, D. Wilmore, B.A. Robson and P.E. Hodgson, Nucl. Phys. A136 (1969) 385
- 1969GO1R Golestaneh, Schmittroth and Tobocman, Bull. Amer. Phys. Soc. 14 (1969) 572
- 1969GU1M V.N. Guman, L.A. Sliv and Yu.I. Kharitonov, Phys. Lett. B28 (1969) 575
- 1969HA1G Hanna, Proc. Int. Conf., Montreal (1969) 443
- 1969HA49 F. Haas, B. Heusch, A. Gallmann and D.A. Bromley, Phys. Rev. 188 (1969) 1625
- 1969HO1F Honsaker, McDonald, Neilson and Hsu, Contrib., Montreal (1969) 299
- 1969HO1T Horowitz, McConnell and Ssengabi, Bull. Amer. Phys. Soc. 14 (1969) 607
- 1969HO1X Hoot et al., Bull. Amer. Phys. Soc. 14 (1969) 494
- 1969IW1B Iwao, DPKU-027-68 (1969)

1969IW1E Iwao, DPKU-024-68 (1969)
1969JO1N Jones and Toms, Bull. Amer. Phys. Soc. 14 (1969) 607
1969KA1A G.Th. Kaschl, G.J. Wagner, G. Mairle, U. Schmidt-Rohr and P. Turek, Phys. Lett. B29 (1969) 167
1969KA1G T. Kammuri and H. Yoshida, Nucl. Phys. A129 (1969) 625
1969KA1R S.N. Kaplan, R.V. Pyle, L.E. Temple and G.F. Valby, Phys. Rev. Lett. 22 (1969) 795
1969KA1W Kaschl et al., Contrib., Montreal (1969) 262
1969KH1C Khanna and Harvey, Bull. Amer. Phys. Soc. 14 (1969) 604
1969KU01 W. Kutschera, D. Schwalm and B. Povh, Nucl. Phys. A124 (1969) 693
1969LE1D Leonardi and Rosa-Clot, Lett. Nuovo Cim. 1 (1969) 829
1969LU07 C.C. Lu, M.S. Zisman and B.G. Harvey, Phys. Rev. 186 (1969) 1086
1969LU1C Ludwig et al., Bull. Amer. Phys. Soc. 14 (1969) 1229
1969ME1M Mendelson, Hardy and Cerny, Bull. Amer. Phys. Soc. 14 (1969) 529
1969MU07 K.M. Murray and J.C. Ritter, Phys. Rev. 182 (1969) 1097
1969NI09 R.J. Nickles, Nucl. Phys. A134 (1969) 308
1969NY1A Nyberg, Jonsson and Bergqvist, Research Inst. Nat. Defence, Stockholm, NP 6902 (1969)
1969OC1B O'Connell and Close, Bull. Amer. Phys. Soc. 14 (1969) 508
1969PA1C Pape, Chevallier, Sens and Armbruster, Rev. Phys. Appl. 4 (1969) 227
1969PH02 G.W. Phillips and W.W. Jacobs, Phys. Rev. 184 (1969) 1052
1969PU04 K.H. Purser, W.P. Alford, D. Cline, H.W. Fulbright, H.E. Gove and M.S. Krick, Nucl. Phys. A132 (1969) 75
1969RI1C Ritter, Sheldon and Strang, Contrib., Montreal (1969) 256
1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
1969SA1J Saunier and Pearson, Bull. Amer. Phys. Soc. 14 (1969) 36
1969SH02 T. Shintomi and M. Masuda, J. Phys. Soc. Jpn. 26 (1969) 607
1969SI04 H.E. Siefken, P.M. Cockburn and R.W. Krone, Nucl. Phys. A128 (1969) 162, and Private Communication (1969)
1969SN02 F.D. Snyder and M.A. Waggoner, Phys. Rev. 186 (1969) 999
1969SN03 J.L. Snelgrove and E. Kashy, Phys. Rev. 187 (1969) 1246
1969SN04 J.L. Snelgrove and E. Kashy, Phys. Rev. 187 (1969) 1259
1969SU15 K. Sugimoto, Phys. Rev. 182 (1969) 1051
1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011

1969TI05 C.T. Tindle and E. Vogt, *Can. J. Phys.* 47 (1969) 2763
1969TO1A I.S. Towner, *Nucl. Phys.* A126 (1969) 97
1969UB1C Uberall, Albert, Wagner and Werntz, *Bull. Amer. Phys. Soc.* 14 (1969) 606
1969UL01 H. Ullrich and H. Krauth, *Nucl. Phys.* A123 (1969) 641
1969VE02 V.V. Verbinski and W.R. Burrus, *Phys. Rev.* 177 (1969) 1671
1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, *Phys. Rev.* 182 (1969) 977
1969WE02 M.L. West, C.M. Jones, J.K. Bair and H.B. Willard, *Phys. Rev.* 179 (1969) 1047;
Erratum *Phys. Rev. C*1 (1970) 367
1969WE03 H.R. Weller and H.A. Van Rinsvelt, *Nucl. Phys.* A129 (1969) 64
1969WE07 K.J. Wetzel, *Phys. Rev.* 181 (1969) 1465
1969WE08 H.R. Weller, *Phys. Lett.* B30 (1969) 409
1969WO1J Wong et al., *Bull. Amer. Phys. Soc.* 14 (1969) 38
1969YO1C Young and Reisman, *Bull. Amer. Phys. Soc.* 14 (1969) 508
1969ZH1A Zhusupov and Eramzhyan, *Izv. Akad. Nauk SSSR Ser. Fiz.* 33 (1969) 730
1970CA1G Campbell and Ober, *Bull. Amer. Phys. Soc.* 15 (1970) 126
1970JA1E Jackson and Weller, *Bull. Amer. Phys. Soc.* 15 (1970) 163
1970SO1G Sokol, De Luca and Browne, *Bull. Amer. Phys. Soc.* 15 (1970) 36
1970SP1E Spinka and Winkler, *Bull. Amer. Phys. Soc.* 15 (1970) 805
1971AJ02 F. Ajzenberg-Selove, *Nucl. Phys.* A166 (1971) 1
1972AJ02 F. Ajzenberg-Selove, *Nucl. Phys.* A190 (1972) 1
CA65 Unknown Source

