

# Energy Levels of Light Nuclei $A = 15$

F. Ajzenberg-Selove

*University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

**Abstract:** An evaluation of  $A = 13-15$  was published in *Nuclear Physics A152* (1970), p. 1. This version of  $A = 15$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1969)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 15$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides: [<sup>15</sup>B](#), [<sup>15</sup>C](#), [<sup>15</sup>N](#), [<sup>15</sup>O](#), [<sup>15</sup>F](#)

B. Tables of Recommended Level Energies:

[Table 15.1](#): Energy levels of <sup>15</sup>C

[Table 15.4](#): Energy levels of <sup>15</sup>N

[Table 15.18](#): Energy levels of <sup>15</sup>O

C. [References](#)

D. Figures: [<sup>15</sup>C](#), [<sup>15</sup>N](#), [<sup>15</sup>O](#), [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

## <sup>15</sup>B

<sup>15</sup>B has been identified in the 5.3 GeV proton bombardment of uranium. It is particle stable (1966PO09). See also (1960ZE03, 1961BA1C, 1966GA25).

## <sup>15</sup>C

(Figs. 9 and 12)

GENERAL:

See (1960TA1C, 1962TA1E, 1964LI1B, 1964ST1B, 1967LO03). See also (1969AR13).

1. <sup>15</sup>C( $\beta^-$ )<sup>15</sup>N  $Q_m = 9.773$

The half-life is  $2.25 \pm 0.05$  sec (1956DO37),  $2.49 \pm 0.07$  sec (1964NE09). The  $\beta$ -spectrum is complex. Transitions have been observed both to the ground state and to the upper of the 5.3 MeV levels of <sup>15</sup>N: the latter transition is clearly allowed: see Table 15.2 (1959AL06, 1966AL12, 1969GA05). The ground state transition of <sup>15</sup>N\*(5.3) has  $E_\gamma = 5.29903 \pm 0.00043$  MeV (1967CH19);  $J^\pi$  for this state is  $\frac{1}{2}^+$  (see Table 15.4). Thus  $J^\pi(^{15}\text{C}) = \frac{1}{2}^+$  or  $\frac{3}{2}^+$  (1959AL06, 1964AL21: see also (1965WA03)). See also (1963KI1B, 1968ZH1B) and <sup>15</sup>N.

2. <sup>9</sup>Be(<sup>6</sup>He,  $\alpha$ )<sup>11</sup>Be  $Q_m = 6.343$   $E_b = 19.076$

<sup>9</sup>Be was irradiated with 14 MeV neutrons and the decay of <sup>11</sup>Be was observed [<sup>9</sup>Be(n,  $\alpha$ )<sup>6</sup>He, followed by the above reaction]: the cross section (which is determined for a large <sup>6</sup>He energy spread) is  $11 \pm 10$  mb (1967ST21).

3. <sup>9</sup>Be(<sup>7</sup>Li, p)<sup>15</sup>C  $Q_m = 9.095$

Observed proton groups are listed in Table 15.3 (1957MU99, 1964CA05).  $\tau_m$  for <sup>15</sup>C\*(0.75) is  $(3.77 \pm 0.11)$  nsec (1968ME08: delayed coincidence technique). Angular distributions of the protons to <sup>15</sup>C\*(0, 0.75) are reported at  $E(^7\text{Li}) = 5.6, 5.8, 6.0, 6.2$  MeV by (1969SN02).

4. <sup>14</sup>C(d, p)<sup>15</sup>C  $Q_m = -1.006$

Table 15.1: Energy levels of  $^{15}\text{C}$ 

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau$	Decay	Reactions
g.s.	$\frac{1}{2}^+; \frac{3}{2}$	$\tau_{1/2} = 2.33 \pm 0.08$ sec	$\beta^-$	1, 3, 4
$0.747 \pm 7$	$\frac{5}{2}^+; \frac{3}{2}$	$\tau_m = 3.76 \pm 0.10$ nsec	$\gamma$	3, 4
( $2.48 \pm 50$ )				3
$3.08 \pm 40$				3, 4
$4.21 \pm 30$				3, 4
(4.60)				3
$5.93 \pm 30$				3, 4
$6.38 \pm 30$				3, 4
$6.58 \pm 40$				3
(6.84)				3
(7.06)				3
$7.32 \pm 30$				3, 4
(7.69)				3
(8.00)				3
$8.12 \pm 60$				3
(8.47)				3

 Table 15.2: Beta decay of  $^{15}\text{C}$  (1959AL06, 1966AL12, 1969GA05)

Decay to $^{15}\text{N}^*$ (MeV)	$J^\pi$	Branch (%)	$\log ft^a$ (exp)	$\log ft^b$ (theor)
g.s.	$\frac{1}{2}^-$	$32 \pm 2$	$5.96 \pm 0.03$	5.8
5.30	$\frac{1}{2}^+$	$68 \pm 2$	$4.04 \pm 0.02$	4.80
7.30	$\frac{1}{2}^+, \frac{3}{2}^+$	$(0.8 \pm 0.2) \times 10^{-2}$	$6.82 \pm 0.13$	6.49
8.31	$\frac{1}{2}^+, \frac{3}{2}^+$	$(5.0 \pm 0.6) \times 10^{-2}$	$5.06 \pm 0.06$	4.51
8.58	$\frac{1}{2}^+, \frac{3}{2}^+$	$\leq 2.8 \times 10^{-2}$	$\geq 5.0$	
9.05	$\frac{1}{2}^+, \frac{3}{2}^+$	$(3.5 \pm 0.5) \times 10^{-2}$	$3.99 \pm 0.07$	

<sup>a</sup> Using  $\tau_m = 2.25 \pm 0.05$  sec.

<sup>b</sup> See (1959AL97).

Table 15.3: Proton groups from  ${}^9\text{Be}({}^7\text{Li}, \text{p}){}^{15}\text{C}$  and  ${}^{14}\text{C}(\text{d}, \text{p}){}^{15}\text{C}$

$E_x$ (MeV $\pm$ keV)		$E_x$ (MeV)
(1957MU99)	(1959MO1B)	(1964CA05) <sup>c</sup>
0	0	0
$0.62 \pm 60$ <sup>a</sup>	$0.75 \pm 30$	0.74
$2.48 \pm 50$ <sup>b</sup>		
$3.08 \pm 40$	3.09	3.08
$4.26 \pm 40$	4.21	4.16
		4.60
$5.93 \pm 40$	5.94	5.81
	6.38	6.39
$6.58 \pm 40$		6.58
		6.84
		7.06
	7.32	7.31
		7.69
		8.00
$8.16 \pm 60$		8.08
		8.47

<sup>a</sup> (1957NO14) reports  $E_x = 0.70 \pm 0.05$  MeV.

<sup>b</sup> Not observed by (1964CA05).

<sup>c</sup>  $\pm 30$  keV (private communication).

Identification of  ${}^{15}\text{C}_{\text{g.s.}}$  with  ${}^{15}\text{N}^*(11.62)$ ,  $T = \frac{3}{2}$ ,  $J^\pi = \frac{1}{2}^+$ , is suggested by (1956BA16). At  $E_d = 14.9$  MeV, proton groups are observed to the ground state of  ${}^{15}\text{C}$  and to the levels at 0.75, 3.09, 4.21, 5.94, 6.38 and 7.32 MeV ( $\pm 30$  keV) (1959MO1B). The angular distribution of ground state protons implies  $l_n = 0$ ,  $J^\pi = \frac{1}{2}^+$  (1959MO1B: see also (1961PU1B) and ref. <sup>9</sup> there, (1964AL21)); for the first excited state,  $l_n = 2$ .  $\theta^2 = 0.093$  and  $0.032$  (1959MO1B),  $0.16$  and  $0.063$  (1966GL01) for the ground state and the first excited state, respectively. See also (1967NE06).

The 0.75 MeV level has a mean life of  $3.73 \pm 0.23$  nsec (1962LO02);  $E_\gamma = 750 \pm 7$  keV. The angular distribution of the  $\gamma$ -rays requires  $J \geq \frac{5}{2}$ . Since the  $l_n = 2$  stripping requires  $J^\pi = \frac{3}{2}^+$  or  $\frac{5}{2}^+$ ,  $J^\pi = \frac{5}{2}^+$  is established (1962CH14). The observed lifetime excludes  $J > \frac{5}{2}$  (1962LO02). See also (1964NE09, 1965WA03, 1966AL12, 1967CH19).

5.  $^{14}\text{C}(\text{n}, \gamma)^{15}\text{C}$   $Q_{\text{m}} = 1.218$

The capture cross section is  $< 1 \mu\text{b}$  (1951YA1A).

6.  $^{18}\text{O}(\text{n}, \alpha)^{15}\text{C}$   $Q_{\text{m}} = -5.009$

See (1966BA1F).

The following reactions leading to  $^{15}\text{C}$  have not been reported:

$^{13}\text{C}(\text{t}, \text{p})^{15}\text{C}$   $Q_{\text{m}} = 0.912$

$^{14}\text{C}(\text{t}, \text{d})^{15}\text{C}$   $Q_{\text{m}} = -5.039$

$^{14}\text{C}(\alpha, ^3\text{He})^{15}\text{C}$   $Q_{\text{m}} = -19.360$

$^{15}\text{N}(\text{n}, \text{p})^{15}\text{C}$   $Q_{\text{m}} = -8.990$

<sup>15</sup>N  
(Figs. 10 and 12)

GENERAL:

*Model calculations:* (1957HA1E, 1959BR1E, 1959FE1B, 1960TA1C, 1961BA1E, 1963BU1C, 1963KU1B, 1964MA1G, 1965CO25, 1965FA1B, 1965GR1H, 1965GU1A, 1965ZA1B, 1966EL08, 1966SO05, 1967CO32, 1967EL03, 1967PA05, 1968EL1A, 1968HO1H, 1968MA2B, 1968SH08, 1968WA04, 1968ZH05, 1969CH1R, 1969EL1B).

*General calculations and reviews:* (1964EV1A, 1965BE1B, 1966OL1C, 1966WI1E, 1967FA1A, 1967LO03, 1968BI1C, 1968ZH1B, 1969HA1G, 1969IW1B).

*Electromagnetic transitions:* (1965RO1N, 1966HA31, 1966PO11, 1966RO1P, 1966RO1U, 1966WA1E, 1967KU1E, 1967PO1J, 1967WA1C, 1968BI1C, 1968SH08, 1968YA1E, 1968ZH06, 1968ZH1B, 1969KH1C, 1969ZH1A).

*Meson interactions:* (1969KA1R).

*Other:* (1961BA05, 1964VA1D).

*Ground state:*  $\mu = -0.28309$  nm (1962BA63, 1964LI14, 1967CO1D).

See also (1961BR13, 1964ST1B, 1965IC1A, 1965MA1T, 1966MA1V, 1966WI1E, 1967SH14, 1968RO1E, 1969CH1R, 1969FU11, 1969PE1D).

1. (a) ${}^9\text{Be}({}^6\text{Li}, \text{p}){}^{14}\text{B}$	$Q_m = 15.130$	$E_b = 25.339$
(b) ${}^9\text{Be}({}^6\text{Li}, \alpha){}^{11}\text{B}$	$Q_m = 14.347$	
(c) ${}^9\text{Be}({}^6\text{Li}, 2\text{n}){}^{13}\text{N}$	$Q_m = 3.951$	
(d) ${}^9\text{Be}({}^6\text{Li}, {}^8\text{Be}){}^7\text{Li}$	$Q_m = 5.587$	
(e) ${}^9\text{Be}({}^6\text{Li}, {}^5\text{He}){}^{10}\text{B}$	$Q_m = 1.933$	

The yield of  $p_0$  and  $p_1$  (reaction (a)) for  $E({}^6\text{Li}) = 3.84$  to  $6.40$  MeV shows some broad structure: analysis in terms of Ericson fluctuation theory gives a value of  $\approx 0.4$  MeV for the average level width at  $E_x = 28$  MeV in  ${}^{15}\text{N}$  (1967SE08). The excitation functions for  $\alpha_0$ ,  $\alpha_1$  and  $\alpha_{2+3}$  (reaction (b)) [ $E({}^6\text{Li}) = 2$  to  $4$  MeV: (1961LE01)] and the yield of  ${}^{13}\text{N}$  (reaction (c)) [ $E({}^6\text{Li}) = 1.5$  to  $3.5$  MeV: (1961NO05)] show a smooth increase in the cross section with energy. For reactions (d) and (e) see (1962MC12). See also (1963BA1Q).

2. ${}^9\text{Be}({}^7\text{Li}, \text{n}){}^{15}\text{N}$	$Q_m = 18.086$
--	----------------

At  $E(^7\text{Li}) = 2.9$  MeV,  $\gamma$ -rays are observed which are assigned to the ground state decay of  $^{15}\text{N}^*(9.05, 9.83, 10.80)$  (1964CA18). See also (1957NO17).

Table 15.4: Energy levels of  $^{15}\text{N}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions
0	$\frac{1}{2}^-$		stable	2, 3, 4, 8, 9, 10, 12, 13, 14, 21, 22, 23, 24, 25, 26, 29, 30, 32, 39, 40, 41, 42, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66
$5.27055 \pm 0.25$	$\frac{5}{2}^+$	$\tau_m = 2.9 \pm 0.5$	$\gamma$	3, 4, 8, 12, 13, 21, 22, 23, 26, 29, 30, 32, 39, 42, 52, 53, 55, 58, 59, 62
$5.29921 \pm 0.25$	$\frac{1}{2}^+$	$< 0.01$	$\gamma$	3, 4, 8, 12, 13, 21, 22, 23, 26, 29, 30, 32, 39, 46, 52, 53, 55, 58, 59
$6.3235 \pm 0.4$	$\frac{3}{2}^-$	$< 0.040$	$\gamma$	3, 4, 8, 10, 12, 13, 21, 22, 26, 29, 30, 32, 39, 49, 52, 53, 55, 57, 58, 59, 62
$7.1550 \pm 0.4$	$\frac{5}{2}^+$	$< 0.018$	$\gamma$	3, 4, 8, 12, 21, 22, 26, 29, 32, 39, 52, 53, 62
$7.3010 \pm 0.5$	$\frac{3}{2}^+$	$< 0.025$	$\gamma$	3, 4, 8, 12, 13, 21, 26, 29, 32, 39, 46, 52, 53, 55, 62
$7.566 \pm 3$	$\frac{7}{2}^+$	$0.06 \pm 0.02$	$\gamma$	3, 4, 8, 12, 13, 21, 22, 29, 39, 52, 53, 62
$8.3126 \pm 0.7$	$\frac{1}{2}^+$	$< 0.010$	$\gamma$	3, 4, 8, 21, 26, 29, 32, 39, 46, 52, 53, 55, 62
$8.576 \pm 2$	$\frac{3}{2}^+$		$\gamma$	3, 4, 8, 13, 21, 22, 26, 32, 39, 52, 53, 62
$9.053 \pm 2$	$\frac{1}{2}^+$		$\gamma$	2, 3, 4, 8, 21, 26, 29, 32, 39, 46, 55
$9.1518 \pm 0.5$	$\frac{3}{2}^-$		$\gamma$	3, 4, 8, 13, 21, 22, 29, 32, 39, 52, 53
$9.1549 \pm 0.5$	$(\frac{5}{2})$	$< 0.010$	$\gamma$	3, 4, 8, 13, 21, 22, 29, 32, 39, 52, 53
$9.225 \pm 3.5$	$\frac{3}{2}$ or $\frac{1}{2}$ <sup>b</sup>	$< 0.1$	$\gamma$	21, 29, 39, 55
$9.762 \pm 3.5$	$\frac{5}{2}^-$		$\gamma$	21, 39, 52, 53
$9.829 \pm 3$	$\frac{7}{2}$	$< 0.19$	$\gamma$	2, 3, 4, 8, 21, 22, 29, 52, 53
$9.929 \pm 4$	$(\frac{1}{2}, \frac{3}{2})^+$		$\gamma$	21, 39



Table 15.4: Energy levels of  $^{15}\text{N}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions
10.070 $\pm$ 3	$\frac{3}{2}^+$		$\gamma$	13, 21, 39, 52, 53
10.451 $\pm$ 1	$\frac{3}{2} \rightarrow \frac{7}{2}$		$\gamma, p$	21, 22, 26, 39
10.536 $\pm$ 1	$\frac{5}{2}^+$		$\gamma, p$	21, 26, 39
10.700 $\pm$ 1	$\frac{3}{2}^+$		$\gamma, p$	21, 22, 26, 27, 39, 52
10.800 $\pm$ 1	$\frac{3}{2}^-$		$\gamma, p$	2, 3, 4, 8, 13, 21, 26, 39
(10.94 $\pm$ 30)				21
11.236 $\pm$ 5	$\geq \frac{3}{2}$	$\Gamma = 3.3$	n	33
11.2943 $\pm$ 1	$\frac{1}{2}^-$	7.9 $\pm$ 0.3	$\gamma, n, p$	26, 27, 28, 33, 52
11.438 $\pm$ 1	$\frac{1}{2}^+$	41.4 $\pm$ 1.1	$\gamma, n, p, \alpha$	5, 26, 27, 28, 33, 35
11.615 $\pm$ 4	$\frac{1}{2}; \frac{3}{2}$	404.9 $\pm$ 6.3	$\gamma, n, p$	26, 27, 28
11.764 $\pm$ 3	$\frac{3}{2}^+$	40 $\pm$ 3	n, p, $\alpha$	5, 28, 33, 35
11.877 $\pm$ 3	$\frac{3}{2}^-$	21 $\pm$ 4	n, p, $\alpha$	5, 28, 33, 35, 52
11.943 $\pm$ 6	$(\frac{9}{2}^-)$	$\leq 3$	n	22, 33, 52
11.965 $\pm$ 3	$\frac{1}{2}^-$	17 $\pm$ 5	n, p, $\alpha$	5, 28, 33, 35
12.097 $\pm$ 4	$\frac{5}{2}^+$	14 $\pm$ 5	$\gamma, n, p, \alpha$	5, 6, 27, 28, 33, 35, 38
12.145 $\pm$ 3	$\frac{3}{2}^-$	47 $\pm$ 7	$\gamma, n, p, \alpha$	5, 6, 27, 28, 33, 35, 39
12.326 $\pm$ 4	$\frac{5}{2}^+$	22	n, p	22, 28, 33, 35
12.493 $\pm$ 4	$\frac{5}{2}^+; \frac{1}{2}$	42	$\gamma, n, p, \alpha$	5, 6, 28, 33, 35, 38, 52
12.52 $\pm$ 10	$\frac{5}{2}^+; \frac{3}{2}$	80	p	27, 52
12.921 $\pm$ 4	$\frac{3}{2}^-$	67 $\pm$ 8	n, p, $\alpha$	5, 6, 28, 33, 35, 38
12.93	$\frac{7}{2}^-$	30	p, $\alpha$	6
13.028 $\pm$ 20	$(\frac{11}{2}^-)$			22
13.14		$< 3$	n, p, $\alpha$	5, 6, 38
13.19		6	n, p, $\alpha$	5, 6, 28, 38
13.36	$\frac{3}{2}^-$	29 $\pm$ 8	n, p, $\alpha$	5, 6, 28, 38
13.40	$\frac{5}{2}^+$	$\approx 60$	n, p, $\alpha$	6, 28, 35
(13.52)			n, p	28
13.60	$(\frac{5}{2}, \frac{7}{2})^-$	15 $\pm$ 4	n, p, $\alpha$	5, 6, 28, 33, 35, 38
13.67	$\frac{1}{2}^+$	$\approx 80$	n, p, $\alpha$	6, 28
13.71		$\approx 40$	n, p, $\alpha$	5, 35, 38
13.75			n, $\alpha$	5
13.84		$\approx 40$	n, p, $\alpha$	28, 35, 38
13.89			n, $\alpha$	5

Table 15.4: Energy levels of  $^{15}\text{N}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions	
14.03	$\frac{3}{2}^{(+)}$		p, $\alpha$	6	
14.06			n, $\alpha$	5	
14.11			105	n, p, $\alpha$	5, 28, 35, 38, 52
14.17			$30 \pm 5$	n, p, $\alpha$	5, 28, 35, 38
14.18				n, $\alpha$	5
14.23				n, $\alpha$	5
14.4			$\approx 2000$	n, p, $\alpha$	33, 35, 38
14.46			$\approx 180$	p, $\alpha$	6
14.51			130	n, p	28
14.64			$50 \pm 3$	n, p, $\alpha$	5, 35, 38
14.7			$\approx 280$	n, p, $\alpha$	35, 38
14.81			99	n, p	28
14.90			37	n, p, $\alpha$	5, 28, 38
15.00				n, p, $\alpha$	5, 28
15.11				n, p, $\alpha$	5, 28, 38, 52
15.29				n, $\alpha$	5
15.37				n, t, $\alpha$	5, 11
15.52				n, $\alpha$	38
15.61				n, $\alpha$	5
15.74				p, t, $\alpha$	11
15.83			$< 3$	n, p, t, $\alpha$	5, 6, 11
$15.89 \pm 20$			$< 3$	n, t, $\alpha$	5, 11
$15.96 \pm 20$				n, t, $\alpha$	5, 11
15.99				n, $\alpha$	5
16.03				n, p, t, $\alpha$	5, 6, 11, 38
16.08				n, p, t, $\alpha$	5, 6, 11, 38
$16.17 \pm 40$		n, p, t, $\alpha$	5, 6, 11		
16.29		n, p, t, $\alpha$	5, 6, 11		
$16.33 \pm 20$		n, p, t, $\alpha$	5, 6, 11, 38		
$16.43 \pm 20$		n, p, t, $\alpha$	5, 6, 11, 38		
$16.49 \pm 30$		n, p, d, t, $\alpha$	5, 6, 11, 16		
$16.59 \pm 25$	70	n, p, t, $\alpha$	5, 6, 11, 38		
$16.67 \pm 30$	100	n, p, d, t, $\alpha$	5, 6, 11, 15, 16, 33, 38		

Table 15.4: Energy levels of  $^{15}\text{N}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions
16.74			n, p, $\alpha$	5, 6
16.76 $\pm$ 30			n, p, d, t, $\alpha$	5, 6, 11, 15, 16
16.85 $\pm$ 30			t, $\alpha$	11
16.90		$\approx$ 350	n, p, d, t, $\alpha$	5, 6, 11, 15, 33, 38
16.98			n, p, $\alpha$	5, 6, 33, 38
17.05			p, t	11
17.10		broad	d, t, $\alpha$	11, 19
17.16 $\pm$ 50			n, p, t, $\alpha$	5, 6, 11
17.19			n, p, $\alpha$	5, 6, 18
17.30		190	n, p, $\alpha$	5, 6, 18, 38
17.36		350	n, p, d, t, $\alpha$	5, 6, 11, 16, 18, 19
17.50			n, p, $\alpha$	5, 6, 38
17.56			n, p, $\alpha$	5, 6
17.81 $\pm$ 40		$\approx$ 170	n, p, d, t, $\alpha$	5, 6, 18
17.70 $\pm$ 50		$\approx$ 500	n, d, $\alpha$	15, 19
17.72 $\pm$ 10		48 $\pm$ 9	(p), d, t, $\alpha$	16, 18, 19
17.81		170	n, $\alpha$	33, 38
17.95			n, p, $\alpha$	5, 6
18.07 $\pm$ 10		19 $\pm$ 4	(n), d, $\alpha$	15, 19
18.09 $\pm$ 20		$\approx$ 45	(n), p, d, t	15, 16, 18
18.22		160	n, $\alpha$	38
18.28 $\pm$ 30		230 $\pm$ 60	n, p, d, $\alpha$	15, 16, 19, 38
19.16 $\pm$ 30		$\approx$ 130	n, d	15
19.5			$\gamma$ , p	48
20.4			$\gamma$ , p	48
22.7			$\gamma$ , p	48
24.5			$\gamma$ , p	48

<sup>a</sup> See also Tables 15.7 and 15.10.

<sup>b</sup> See (1967PH03).

3.  $^{10}\text{B}(^6\text{Li}, \text{p})^{15}\text{N}$

$$Q_m = 18.751$$

Table 15.5: Resonances in  $^{11}\text{B} + \alpha$ 

$E_\alpha$ (MeV)	$\Gamma_{\text{lab}}$ (keV)	Particle out	$E_x$ (MeV)	$J^\pi$	Refs. <sup>a</sup>
0.60		n	11.43		(1954BE08) <sup>b</sup>
1.03		n	11.75		(1954BE08)
1.18		n	11.86		(1954BE08)
1.30		n	11.95		(1954BE08)
1.51		n, p	12.10		(1955SH46)
1.58		n, p	12.15		(1955SH46)
2.06	66	$n_0, p_0$	12.50	$\frac{5}{2}^+$	(1955SH46, 1956BO61, 1958HA1B, 1959LE28)
2.63	80	$n_0, p_0$	12.92	$\frac{3}{2}^-$	(1956BO61, 1958HA1B, 1959LE28, 1963MA28, 1966MA04)
2.64	40	$p_0$	12.93	$\frac{7}{2}^-$	(1959LE28, 1963MA28)
2.94	< 6	$n_0, p_0$	13.15		(1956BO61, 1958HA1B, 1959LE28)
2.99	8	$n_0, p_0$	13.18		(1956BO61, 1958HA1B, 1959LE28, 1966MA04)
3.23	29	$n_0, p$	13.36	$\frac{3}{2}^-$	(1956BO61, 1958HA1B, 1959LE28, 1963MA28, 1966MA04)
3.31	40	p	13.42	$\frac{5}{2}^+$	(1959LE28, 1963MA28)
3.56	20	$n_0, p$	13.61	$(\frac{5}{2}, \frac{7}{2})^-$	(1956BO61, 1958HA1B, 1959LE28, 1963MA28, 1966MA04)
3.64	$\approx 110$	$p_0$	13.66		(1963MA28)
3.71	40	$n_0$	13.71		(1956BO61, 1958HA1B, 1963MA28, 1966MA04)
3.76		$n_0$	13.75		(1966MA04)
3.95		$n_0$	13.89		(1963MA28, 1966MA04)
4.15		$p_0$	14.03		(1963MA28)
4.19		$n_1$	14.06		(1963MA28, 1966MA04)
4.24		$n_0$	14.10		(1963MA28, 1966MA04)
4.31	35	$n_1$	14.15		(1956BO61, 1958HA1B, 1963MA28, 1966MA04)
4.35		$n_0$	14.18		(1963MA28, 1966MA04)
4.41		$n_0$	14.23		(1963MA28, 1966MA04)
4.56		$p_0$	(14.33)		(1963MA28)

Table 15.5: Energy levels of  $^{15}\text{N}$  <sup>a</sup> (continued)

$E_\alpha$ (MeV)	$\Gamma_{\text{lab}}$ (keV)	Particle out	$E_x$ (MeV)	$J^\pi$	Refs. <sup>a</sup>
4.73	$\approx 250$	p <sub>0</sub>	14.46		(1963MA28)
4.96	72	n <sub>0</sub>	14.63		(1956BO61, 1958HA1B)
5.34		n <sub>0</sub>	14.90		(1958HA1B)
5.49		n <sub>0</sub>	15.01		(1958HA1B)
5.58		n <sub>0</sub>	15.08		(1958HA1B)
5.86		n <sub>0</sub>	15.29		(1958HA1B)
5.98		n <sub>0</sub> , n <sub>2</sub>	15.37		(1958HA1B)
6.30		n <sub>0</sub> , (n <sub>2</sub> )	15.61		(1958HA1B)
6.60		n, p	15.83		(1963ED01) <sup>c</sup>
6.72		n <sub>0</sub> , (n <sub>2</sub> )	15.92		(1958HA1B)
6.74		n <sub>0</sub> , n <sub>2</sub>	15.93		(1958HA1B)
6.82		n <sub>0</sub> , n <sub>2</sub>	15.99		(1958HA1B)
6.89		n, p	16.04		(1958HA1B, 1963ED01)
6.94		n, p	16.08		(1963ED01)
7.08		n, p	16.18		(1963ED01)
7.26		n, p	16.31		(1963ED01)
7.31		n, p	16.35		(1963ED01)
7.41		n, p	16.42		(1963ED01)
7.56		n, p	16.53		(1963ED01)
7.65		n, p	16.60		(1963ED01)
7.77		n, p	16.69		(1963ED01)
7.84		n, p	16.74		(1963ED01)
7.86		n, p	16.75		(1963ED01)
8.07		n, p	16.91		(1963ED01)
8.17		n, p	16.98		(1963ED01)

Table 15.5: Energy levels of  $^{15}\text{N}$  <sup>a</sup> (continued)

$E_\alpha$ (MeV)	$\Gamma_{\text{lab}}$ (keV)	Particle out	$E_x$ (MeV)	$J^\pi$	Refs. <sup>a</sup>
8.44		n, p	17.18		(1963ED01)
8.46		n, p	17.19		(1963ED01)
8.59		n, p	17.29		(1963ED01)
8.68		n, p	17.35		(1963ED01)
8.88		n, p	17.50		(1963ED01)
8.96		n, p	17.56		(1963ED01)
8.99		n, p	17.58		(1963ED01)
9.49		n, p	17.95		(1963ED01)

<sup>a</sup> It should be noted that (1954BE08), (1958HA1B) and (1963ED01) are unpublished. They are quoted here only because of the dearth of published information on these resonances.

<sup>b</sup> And private communication.

<sup>c</sup> Resonant energies listed by (1963ED01) are  $\pm 40$  keV.

At  $E(^6\text{Li}) = 4.9$  MeV, thirty proton groups are observed corresponding to  $^{15}\text{N}$  states with  $E_x < 16.8$  MeV. Angular distributions have been measured for the proton groups corresponding to  $^{15}\text{N}^*(5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.58, 9.05 + 9.15)$  (1966MC05). The ground-state  $\gamma$ -decay of  $^{15}\text{N}^*(9.83, 10.80)$  is reported by (1964CA18). See also (1963MO1B).

4.  $^{10}\text{B}(^7\text{Li}, \text{d})^{15}\text{N}$   $Q_m = 13.723$

At  $E(^7\text{Li}) = 5.2$  MeV, thirty deuteron groups are observed corresponding to  $^{15}\text{N}$  states with  $E_x < 15.1$  MeV. Angular distributions have been measured for the deuteron groups corresponding to  $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.58, 9.05 + 9.15, 9.83)$  (1966MC05). See also (1963MO1B). The ground state  $\gamma$ -decay of  $^{15}\text{N}^*(8.31, 9.05, 9.83, 10.80)$  is reported by (1964CA18).

5.  $^{11}\text{B}(\alpha, \text{n})^{14}\text{N}$   $Q_m = 0.157$   $E_b = 10.992$

Reported resonances are displayed in Table 15.5 (1954BE08, 1955SH46, 1956BO61, 1958HA1B, 1963ED01, 1963MA28, 1966MA04). See also (1962GO1J, 1963GO1J, 1965TS1A) and (1966WE1B). See also  $^{14}\text{N}$ .

6.  $^{11}\text{B}(\alpha, \text{p})^{14}\text{C}$   $Q_m = 0.784$   $E_b = 10.992$

Reported resonances are listed in Table 15.5 (1955SH46, 1959LE28, 1963ED01, 1963MA28). Partial widths for several resonances are listed by (1959LE28, 1963MA28). See also  $^{14}\text{C}$ .

7.  $^{11}\text{B}(\alpha, \text{d})^{13}\text{C}$   $Q_m = -5.168$   $E_b = 10.992$

The yield of ground state deuterons has been measured for  $E_\alpha = 17$  to 22 MeV by (1968AL1C). See also  $^{13}\text{C}$ .

8. (a)  $^{11}\text{B}(^6\text{Li}, \text{d})^{15}\text{N}$   $Q_m = 9.520$

(b)  $^{11}\text{B}(^7\text{Li}, \text{t})^{15}\text{N}$   $Q_m = 8.525$

At  $E(^6\text{Li}) = 4.72$  MeV and at  $E(^7\text{Li}) = 5.00$  MeV, angular distributions are reported for the deuterons and the tritons corresponding to  $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.58, 9.05 + 9.15)$  (1966MC05). See also (1963MO1B). Gamma rays are observed in reaction (a) which are assigned to the ground state transitions of  $^{15}\text{N}^*(9.15, 9.83, 10.80)$  (1964CA18). See also (1963HO1E) and (1964HA09).

9.  $^{11}\text{B}(^9\text{Be}, ^5\text{He})^{15}\text{N}$   $Q_m = 8.464$

See (1963HO1E).

10.  $^{11}\text{B}(^{16}\text{O}, ^{12}\text{C})^{15}\text{N}$   $Q_m = 3.831$

At  $E(^{16}\text{O}) = 27, 30, 32.5$  and  $35$  MeV, angular distributions corresponding to transitions to  $^{15}\text{N}^*(0, 6.32)$  have been measured by (1965BO14, 1966BO22): the ground state angular distributions show strong diffraction structure. See also (1969GO1R, 1969BR1D, 1969KA1G).

11. (a)  $^{12}\text{C}(t, n)^{14}\text{N}$   $Q_m = 4.015$   $E_b = 14.850$   
 (b)  $^{12}\text{C}(t, p)^{14}\text{C}$   $Q_m = 4.641$   
 (c)  $^{12}\text{C}(t, t)^{12}\text{C}$   
 (d)  $^{12}\text{C}(t, \alpha)^{11}\text{B}$   $Q_m = 3.858$

Reported resonances are listed in Table 15.6 (1961VA13, 1962GU01, 1962KU09, 1963NI04, 1965SE05, 1969ET01). The triton yield has been measured for  $E_t = 9$  to  $13$  MeV by (1965GL04). See also (1962NE1D, 1964GR1H, 1967CH35),  $^{11}\text{B}$  and  $^{12}\text{C}$  in (1968AJ02), and  $^{14}\text{C}$  and  $^{14}\text{N}$ .

12.  $^{12}\text{C}(\alpha, p)^{15}\text{N}$   $Q_m = -4.965$

Angular distributions of the protons corresponding to the ground state transition have been measured at  $E_\alpha = 13.4$  to  $16.0$  MeV (1967IV1B),  $16.1$  to  $19.0$  MeV (1960PR13),  $19.7$  to  $22.1$  MeV (1963YA1C),  $20.6$  to  $22.2$  MeV (1961KO04),  $25$  to  $39$  MeV (1959NO38) and at  $42$  MeV (1962LI07). Angular distributions are also reported for the groups to  $^{15}\text{N}^*(5.27+5.30, 6.32, 7.16+7.30+7.57)$  (1959NO38:  $34.6$  MeV) and  $^{15}\text{N}^*(6.32)$  (1962LI07:  $42$  MeV). See also (1962EI03, 1969GL1D), (1961KR1A, 1962HO1D, 1962TE1B, 1963DA1B, 1964DA1D, 1964KE1C, 1965NE1D, 1966HI1C, 1967RO1K; theor.) and (1959AJ76).

13.  $^{12}\text{C}(^7\text{Li}, \alpha)^{15}\text{N}$   $Q_m = 12.382$

Angular distributions have been measured at  $E(^7\text{Li}) = 3.2$  to  $4.0$  MeV (1962HO06;  $\alpha_0, \alpha_{1+2}, \alpha_3$ ) and  $30.3$  MeV (1969GL07;  $\alpha_0, \alpha_{1+2}, \alpha_3$  and the  $\alpha$ 's corresponding to  $^{15}\text{N}^*(7.35, 7.6, 8.6, 9.2, 10.1, 10.9, 12.8, 13.5, 15.2)$ . (1969BA2U) report the excitation of  $^{15}\text{N}^*(12.5, 13.3, 14.9, 15.6, 16.2, 16.8, 18.9, 19.8)$  and (1969GL07) also report  $\alpha$  groups corresponding to  $^{15}\text{N}^*(11.8, 16.5)$



Table 15.6: Resonances in  $^{12}\text{C} + t$ 

$E_t$ (MeV $\pm$ keV)	$E_x$ (MeV)	Particles out	Refs.
0.66	15.38	$\alpha_0$	(1963NI04, 1969ET01)
1.11	15.74	$p_0, t_0, \alpha_1$	(1962GU01, 1962KU09, 1963NI04, 1969ET01)
1.21	15.82	$t_0$	(1969ET01)
$1.30 \pm 20$	15.89	$n, \alpha_0$	(1961VA13, 1969ET01)
$1.39 \pm 20$	15.96	$n, t_0, \alpha_0$	(1961VA13, 1969ET01)
1.46	16.02	$p_0$	(1969ET01)
1.54	16.08	$n, \alpha_0, \alpha_1$	(1961VA13, 1969ET01)
$1.65 \pm 40$	16.17	$n, \alpha_0$	(1961VA13, 1965SE05, 1969ET01)
1.78	16.27	$\alpha_0$	(1969ET01)
$1.85 \pm 20$	16.33	$n, p_0, \alpha_0, \alpha_1$	(1961VA13, 1965SE05, 1969ET01)
$1.98 \pm 20$	16.43	$n, p_0$	(1961VA13, 1969ET01)
$2.05 \pm 30$	16.49	$p_0, t_0, \alpha_0$	(1965SE05, 1969ET01)
$2.18 \pm 25$	16.59	$n, p_0, t_0, \alpha_0, \alpha_1$	(1961VA13, 1969ET01)
$2.28 \pm 30$	16.67	$n, p_0, \alpha_0, \alpha_1$	(1961VA13, 1965SE05, 1969ET01)
$2.39 \pm 30$	16.76	$n, t_0, \alpha_0, \alpha_1$	(1961VA13, 1965SE05, 1969ET01)
$2.50 \pm 30$	16.85	$\alpha_0, \alpha_1$	(1965SE05, 1969ET01)
2.60	16.93	$\alpha_0$	(1969ET01)
2.75	17.05	$p_0$	(1969ET01)
2.82	17.10	$t_0, \alpha_0, \alpha_1$	(1969ET01)
$2.89 \pm 50$	17.16	$\alpha_0$	(1965SE05, 1969ET01)
3.14	17.36	$\alpha_1$	(1969ET01)

and, possibly  $^{15}\text{N}^*(17.1, 18.3)$ . The mean lifetime of  $^{15}\text{N}^*(9.83) < 1.9 \times 10^{-13}$  sec (see Table 15.7):  $E_\gamma = 4562.6 \pm 4.0$  keV ( $^{15}\text{N}^*(9.83) \rightarrow 5.27$ ) (1969TH01). See also (1960SH05) and (1969GI1B, 1969RO1G).

14. (a)  $^{12}\text{C}(^{14}\text{N}, ^{11}\text{C})^{15}\text{N}$   $Q_m = -7.885$   
 (b)  $^{12}\text{C}(^{19}\text{F}, ^{16}\text{O})^{15}\text{N}$   $Q_m = 3.150$

For reaction (a) see (1969BR1D); for reaction (b) see (1969RO1G).

15.  $^{13}\text{C}(d, n)^{14}\text{N}$   $Q_m = 5.325$   $E_b = 16.160$

Table 15.7: Lifetimes of some  $^{15}\text{N}$  states

$E_x$ (MeV)	$\tau_m$ (psec)	Reaction	Refs.
5.27	$> 1$	$^{16}\text{O}(t, \alpha)$	(1965AL19)
	$\approx 1$	$^9\text{Be}(^{14}\text{N}, ^8\text{Be})$	(1969NI09)
	$2.9 \pm 0.5$	$^{14}\text{N}(d, p)$	(1967BI11)
5.30	$< 0.01$	$^{14}\text{C}(p, \gamma)$	(1968COZV)
	$(4.3 \pm 1.8) \times 10^{-2}$	$^{14}\text{N}(d, p)$	(1965AL19)
	$(2.2 \pm 0.7) \times 10^{-2}$	$^{14}\text{N}(n, \gamma)$	(1969WE07)
6.32	$< 0.045$	$^{14}\text{N}(d, p)$	(1968GI11)
	$< 0.040$	$^{14}\text{N}(n, \gamma)$	(1969WE07)
	$< 0.010$	$^{14}\text{C}(p, \gamma)$	(1968COZV)
7.16	$< 0.018$	$^{14}\text{N}(d, p)$	(1968GI11)
	$0.115 \pm 0.025$	$^{13}\text{C}(^3\text{He}, p)$	(1966LI07)
	$0.010 \pm 0.002$	$^{14}\text{C}(p, \gamma)$	(1968COZV)
7.30	$< 0.025$	$^{14}\text{N}(d, p)$	(1968GI11)
	$< 0.030$	$^{14}\text{N}(n, \gamma)$	(1969WE07)
	$< 0.010$	$^{14}\text{C}(p, \gamma)$	(1968COZV)
7.57	$0.15 \pm 0.05$	$^{13}\text{C}(^3\text{He}, p)$	(1966LI07)
	$0.06 \pm 0.02$	$^{14}\text{N}(d, p)$	(1968GI11)
8.31	$< 0.021$	$^{14}\text{N}(d, p)$	(1968GI11)
	$< 0.020$	$^{14}\text{N}(n, \gamma)$	(1969WE07)
	$< 0.010$	$^{14}\text{C}(p, \gamma)$	(1968COZV)
9.155	$< 0.010$	$^{14}\text{C}(p, \gamma)$	(1968COZV)
	$< 0.010$	$^{14}\text{N}(n, \gamma)$	(1969WE07)
9.23	$< 0.1$	$^{16}\text{O}(\gamma, p)$	(1969MU07)
9.83	$< 0.19$	$^{12}\text{C}(^7\text{Li}, \alpha)$	(1969TH01)

Table 15.8: Resonances in  $^{13}\text{C} + \text{d}$

$E_d$ (MeV)	Emitted particles	$\Gamma_{\text{lab}}$ (keV)	$^{15}\text{N}^*$ (MeV)	Refs.
0.37	p		16.47	(1956VA17)
0.64	n, $p_0$	$\approx 100$	16.72	(1950CU13, 1950RI57, 1953KO42, 1956MA46, 1956VA17)
0.85	n	$\approx 400$	16.90	(1950RI57)
1.10	$\alpha_0$	broad	17.11	(1956MA35, 1966KL06)
$1.24 \pm 0.04$	$t_0$	$\approx 200$	17.24	(1956MA35)
$1.40 \pm 0.04$	$p_0, t_0, \alpha_0$	$\approx 400$	17.37	(1956MA35, 1956MA46)
1.55 <sup>a</sup>	n	$\approx 100$		(1941BE1A, 1950RI57, 1966KL06)
$1.64 \pm 0.04$	$t_0$	$\approx 200$	17.58	(1956MA35)
$1.78 \pm 0.05$	n, $\alpha_0$	$\approx 600$	17.70	(1950RI57, 1955MA76, 1956MA35)
$1.80 \pm 0.01$	( $p_0$ ), $t_0, \alpha_1$	$55 \pm 10$	17.72	(1956MA35, 1956MA46)
$2.20 \pm 0.01$	(n), $\alpha_0, \alpha_1$	$22 \pm 4$	18.07	(1956MA35, 1963AL21)
$2.23 \pm 0.02$	(n), $p_0, t$	$\approx 50$	18.09	(1956MA35, 1956MA46, 1963AL21)
$2.45 \pm 0.03$	n, $p_0, \alpha_0$	$270 \pm 70$	18.28	(1955MA76, 1956MA35, 1956MA46)
$3.46 \pm 0.03$	n	$\approx 150$	19.16	(1955MA76, 1963DE19)

<sup>a</sup> Possibly to be identified with 1.40 MeV resonance (1956MA35).

Observed resonances are displayed in Table 15.8 (1950RI57, 1955MA76, 1963AL21, 1963DE19). Excitation functions have recently been measured for  $E_d = 0.8$  to 1.3 MeV (1961JA09;  $n_0, n_1, n_2, n_3, n_5$ ), 2.0 to 3.2 MeV (1963AL21: 7.03 MeV  $\gamma$ -ray) and 3.2 to 4.0 MeV (1963DE19;  $n_0$ ). See also (1960VA11). Polarization measurements are reported for  $E_d = 2.5$  to 4.0 MeV (1967ME1N;  $n_0, n_1, n_2$ ) and 2.8 MeV (1965GA1G;  $n_3, n_4, n_5$ ). See also  $^{14}\text{N}$ .

16.  $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$

$$Q_m = 5.952$$

$$E_b = 16.160$$

Observed resonances are displayed in Table 15.8 (1941BE1A, 1950CU13, 1953KO42, 1956MA46). Excitation functions have recently been measured for  $E_d = 1$  to 3.4 MeV (1968LI1L;  $p_0, p_1$ ), 3.0 to 4.0 MeV (1965LA09:  $^{14}\text{C}^*(6.09, 6.73, 7.34)$ ), 3.1 to 4.1 MeV (1963DE19;  $p_0$ ), and 4.1 to 6.2 MeV (1968CO04;  $p_1$ ). See also (1963AL21). See also  $^{14}\text{C}$ .

17.  $^{13}\text{C}(\text{d}, \text{d})^{13}\text{C}$

$$E_b = 16.160$$

Excitation functions for elastically scattered deuterons have been measured for  $E_d = 1$  to 3.4 MeV (1968LI1L) and 4.5 to 5.7 MeV (1968CO04).

$$18. \text{}^{13}\text{C}(\text{d}, \text{t})\text{}^{12}\text{C} \qquad Q_m = 1.311 \qquad E_b = 16.160$$

Observed resonances are listed in Table 15.8 (1956MA35). (1968LI1L) report measurement of the  $t_0$  excitation function for  $E_d = 1$  to 3.4 MeV. A polarization study has been made at  $E_d = 12.3$  MeV (1969DE1H;  $t_0, t_1$ ). See also  $^{12}\text{C}$ .

$$19. \text{}^{13}\text{C}(\text{d}, \alpha)\text{}^{11}\text{B} \qquad Q_m = 5.168 \qquad E_b = 16.160$$

Observed resonances are listed in Table 15.8 (1956MA35). See also (1966KL06, 1966KL1F, 1968CO04, 1968LI1L).

$$20. \text{}^{13}\text{C}(\text{t}, \text{n})\text{}^{15}\text{N} \qquad Q_m = 9.903$$

Not reported.

$$21. \text{}^{13}\text{C}(\text{}^3\text{He}, \text{p})\text{}^{15}\text{N} \qquad Q_m = 10.667$$

Observed proton groups and  $\gamma$ -rays corresponding to  $^{15}\text{N}$  states are listed in Table 15.9 (1959YO25, 1966GA08, 1966WA08, 1967PH03). Gamma-ray branching ratios obtained by (1965WA16, 1966PE04, 1966WA08) are displayed in Table 15.10 which also shows  $J^\pi$  values obtained from angular correlation measurements. The two states at  $E_x = 9.16$  MeV [see reactions 32 and 39] are separated by  $2.5 \pm 0.5$  keV (1968ST10). See also (1959BR79). The  $\tau_m$  for  $^{15}\text{N}^*(7.16, 7.57)$  are  $0.115 \pm 0.025$  psec and  $0.15 \pm 0.05$  psec, respectively: see Table 15.7 (1966LI07). Angular distributions of the ground state protons have been measured for  $E(^3\text{He}) = 8.7$  to 10.93 MeV (1962AL01). See also (1959AJ76).

See also (1967YO1C, 1968WE15), (1963CL1A, 1966PR1B) and  $^{16}\text{O}$  in (1971AJ02).

$$22. \text{}^{13}\text{C}(\alpha, \text{d})\text{}^{15}\text{N} \qquad Q_m = -7.687$$

This reaction has been studied at  $E_\alpha = 40.1$  MeV: see Table 15.9 (1969LU07).

Table 15.9: Energy levels in  $^{15}\text{N}$  from  $^{13}\text{C}(^3\text{He}, \text{p})^{15}\text{N}$  and  $^{13}\text{C}(\alpha, \text{d})^{15}\text{N}$

$E_x$ (MeV $\pm$ keV) <sup>a</sup>				
(1959YO25)	(1966GA08)	(1966WA08) <sup>b</sup>	(1967PH03)	(1969LU07) <sup>c</sup>
5.283 $\pm$ 12				5.266 $\pm$ 20
6.333 $\pm$ 12				6.336 $\pm$ 30
7.169 $\pm$ 12				7.170 $\pm$ 20
7.310 $\pm$ 12				
7.577 $\pm$ 13				7.581 $\pm$ 20
8.318 $\pm$ 12	8.323 $\pm$ 6	8.312		
8.581 $\pm$ 14	8.581 $\pm$ 5	8.570		8.587 $\pm$ 20
9.061 $\pm$ 14	9.056 $\pm$ 5	9.052	9.054 $\pm$ 4	
9.164 $\pm$ 14	9.159 $\pm$ 5			9.169 $\pm$ 30
	9.225 $\pm$ 6		9.225 $\pm$ 3	
	9.760 $\pm$ 5			
	9.827 $\pm$ 6		9.829 $\pm$ 4	9.808 $\pm$ 20
	9.929 $\pm$ 8			
	10.064 $\pm$ 7	10.074	10.072 $\pm$ 4	
	10.454 $\pm$ 6	10.452		10.451 $\pm$ 20
	10.536 $\pm$ 7			
	10.704 $\pm$ 6			10.698 $\pm$ 20
	10.805 $\pm$ 7	10.800		
		(10.94 $\pm$ 30)		d

<sup>a</sup> See also (1962SH21, 1966GO1J, 1966PH1B).

<sup>b</sup>  $E_\gamma$ , except for  $E_x = 10.94$  MeV; errors for  $E_\gamma$  are nominal.

<sup>c</sup>  $^{13}\text{C}(\alpha, \text{d})^{15}\text{N}$ :  $E_\alpha = 40.1$  MeV.

<sup>d</sup> (1969LU07) also reports levels at  $E_x = 11.950 \pm 0.020$  ( $J^\pi = (\frac{9}{2}^-)$ ),  $12.318 \pm 0.030$  and  $13.028 \pm 0.020$  MeV ( $J^\pi = (\frac{11}{2}^-)$ ).

Table 15.10: Radiative decays in  $^{15}\text{N}$ 

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
5.27	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	100	
5.30	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	100	
6.32	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100	(1965WA16, 1966PE04, 1969SI04)
		5.27	$\frac{5}{2}^+$	< 1	(1965WA16)
				< 3	(1966PE04)
		5.30	$\frac{1}{2}^+$	< 3	(1966PE04)
				< 1	(1965WA16)
7.16	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	< 5	(1965WA16)
				< 3	(1969SI04)
				< 12	(1968GI11)
				< 4	(1966HA30)
		5.27	$\frac{5}{2}^+$	100	(1965WA16, 1966PE04, 1968GI11)
				> 97	(1969SI04)
				< 95	(1967TH05)
		5.30	$\frac{1}{2}^+$	< 4	(1966AL18)
		< 4	(1968GI11)		
7.30	$\frac{3}{2}^+$	6.32	$\frac{3}{2}^-$	< 0.5	(1965WA16)
		0	$\frac{1}{2}^-$	100	(1966PE04, 1969SI04)
				$98 \pm 1$	(1965WA16, 1968GI11)
		5.27	$\frac{5}{2}^+$	$2 \pm 1$	(1965WA16)
				< 1.5	(1968GI11)
		5.30	$\frac{1}{2}^+$	$2 \pm 1$	(1968GI11)
		6.32	$\frac{3}{2}^-$	< 3	(1966PE04)
7.57	$\frac{7}{2}^+$	0	$\frac{1}{2}^-$	< 0.25	(1965WA16)
				< 2	(1965WA16)
				< 4	(1966HA30)
				< 3	(1966PE04)
		5.27	$\frac{5}{2}^+$	100	(1965WA16, 1966PE04, 1968GI11)
		5.30	$\frac{1}{2}^+$	< 5	(1965WA16)
				< 4	(1966AL18)
		< 6	(1968GI11)		

Table 15.10: Radiative decays in  $^{15}\text{N}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
8.31	$\frac{1}{2}^+$	6.32	$\frac{3}{2}^-$	< 0.6	(1965WA16)
		0	$\frac{1}{2}^-$	$80 \pm 3$	(1965WA16)
				$77 \pm 3$	(1966WA08)
				$70 \pm 4$	(1966PE04)
				$79.1 \pm 1.9$	(1967PH03)
		5.27	$\frac{5}{2}^+$	< 3	(1965WA16)
		5.30	$\frac{1}{2}^+$	$10 \pm 2$	(1965WA16)
		5.27 + 5.30		$12 \pm 2$	(1966WA08)
				$12 \pm 3$	(1966PE04)
				$10.9 \pm 1.3$	(1967PH03)
		6.32	$\frac{3}{2}^-$	$7.8 \pm 2$	(1965WA16)
				$12 \pm 3$	(1966PE04)
				$4.4 \pm 1.0$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 1	(1965WA16)
		$\leq 6$	(1966PE04)		
		$1.2 \pm 0.6$	(1967PH03)		
7.30	$\frac{3}{2}^+$	$2.2 \pm 0.4$	(1965WA16)		
		$4.4 \pm 0.7$	(1967PH03)		
8.58	$\frac{3}{2}^+$	0 <sup>c</sup>	$\frac{1}{2}^-$	$34 \pm 4$	(1965WA16)
				$32 \pm 3$	(1966WA08)
				$27 \pm 4$	(1966PE04)
				$33.4 \pm 2.0$	(1967PH03)
		5.27	$\frac{5}{2}^+$	$63 \pm 4$	(1965WA16)
				$65 \pm 3$	(1966WA08)
		5.30	$\frac{1}{2}^+$	< 12	(1965WA16)
		5.27 + 5.30		$66 \pm 4$	(1966PE04)
				$61.6 \pm 2.0$	(1967PH03)
		6.32	$\frac{3}{2}^-$	$3 \pm 1$	(1965WA16)
				$\leq 7$	(1966PE04)
				$1.4 \pm 0.6$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 5	(1965WA16)
				< 4	(1966WA08)
		$3.6 \pm 0.5$	(1967PH03)		

Table 15.10: Radiative decays in  $^{15}\text{N}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
9.05	$\frac{1}{2}^+$	7.30	$\frac{3}{2}^+$	< 0.7	(1965WA16)
				< 3	(1966WA08)
		7.57	$\frac{7}{2}^+$	< 3	(1965WA16, 1966WA08)
		0	$\frac{1}{2}^-$	$92 \pm 3$	(1965WA16)
				$92 \pm 4$	(1966WA08)
				$91.6 \pm 0.9$	(1967PH03)
		5.27	$\frac{5}{2}^+$	$3.8 \pm 1$	(1965WA16)
				$3.5 \pm 1$	(1966WA08)
				$4.7 \pm 0.7$	(1967PH03)
		6.32	$\frac{3}{2}^-$	$3 \pm 2$	(1965WA16)
				$4.5 \pm 1$	(1966WA08)
				$3.7 \pm 0.5$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
		7.30	$\frac{3}{2}^+$	$1.2 \pm 0.4$	(1965WA16)
		7.57	$\frac{7}{2}^+$	< 2	(1965WA16)
8.31	$\frac{1}{2}^+$	< 0.5	(1965WA16)		
9.152 <sup>a</sup>	$\frac{3}{2}^-$	7.16 + 7.30 + 7.57		< 1	(1967PH03)
		0	$\frac{1}{2}^-$	97	(1968ST10)
9.155 <sup>a</sup>	$(\frac{5}{2})$			100	(1969SI04)
		5.27 + 5.30		3	(1968ST06)
		0	$\frac{1}{2}^-$	0 → 17	(1968ST10)
				17	(1967TH05)
		5.27	$\frac{5}{2}^+$	8	(1967TH05)
		5.30	$\frac{1}{2}^+$	10	(1967TH05)
		5.27 + 5.30		14 → 17	(1968ST10)
				23	(1969SI04)
		6.32	$\frac{3}{2}^-$	18 → 22	(1968ST10)
				19	(1969SI04)
		20	(1967TH05)		
7.16	$\frac{5}{2}^+$	44 → 52	(1968ST10)		
		58	(1969SI04)		
		45	(1967TH05)		
7.30	$\frac{3}{2}^+$	7 → 9	(1968ST10)		



Table 15.10: Radiative decays in  $^{15}\text{N}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
9.23	$\leq \frac{5}{2}^-$	0	$\frac{1}{2}^-$	< 30	(1965WA16)
				$41.5 \pm 2.2$	(1967PH03)
		5.27	$\frac{5}{2}^+$	< 25	(1965WA16)
		5.30	$\frac{1}{2}^+$	100	(1965WA16)
		5.27 + 5.30		$31.2 \pm 1.7$	(1967PH03)
		6.32	$\frac{3}{2}^-$	$\leq 25$	(1965WA16)
				$24.7 \pm 1.5$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 30	(1965WA16)
				< 1	(1967PH03)
		7.30	$\frac{3}{2}^+$	< 30	(1965WA16)
				$2.6 \pm 0.7$	(1967PH03)
		7.57	$\frac{7}{2}^+$	< 20	(1965WA16)
		8.31	$\frac{1}{2}^+$	< 5	(1965WA16)
		7.57 + 8.31		< 1	(1967PH03)
9.76	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	100	(1965WA16)
				$81.5 \pm 2.8$	(1967PH03)
		5.27 + 5.30		< 10	(1965WA16)
				$7.5 \pm 1.5$	(1967PH03)
		6.32	$\frac{3}{2}^-$	< 5	(1965WA16)
				$3.7 \pm 0.8$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
				$2.3 \pm 0.5$	(1967PH03)
		7.30	$\frac{3}{2}^+$	< 3	(1965WA16)
				< 2	(1967PH03)
		7.57	$\frac{7}{2}^+$	< 10	(1965WA16)
				$5.0 \pm 0.6$	(1967PH03)
		8.31	$\frac{1}{2}^+$	< 2	(1965WA16)
				< 1	(1967PH03)
8.58	$\frac{3}{2}^+$	< 2	(1965WA16)		
		< 2	(1967PH03)		
9.83	$\frac{7}{2}$	0	$\frac{1}{2}^-$	< 30	(1965WA16)
				< 4	(1967PH03)
		5.27	$\frac{5}{2}^+$	100	(1965WA16)

Table 15.10: Radiative decays in  $^{15}\text{N}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
9.93	$(\frac{1}{2}, \frac{3}{2})^+$	5.30	$\frac{1}{2}^+$	< 15	(1965WA16)
		5.27 + 5.30		$84.4 \pm 1.8$	(1967PH03)
		6.32	$\frac{3}{2}^-$	< 15	(1965WA16)
				$2.2 \pm 0.9$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
				$2.4 \pm 1.1$	(1967PH03)
		7.30	$\frac{3}{2}^+$	< 10	(1965WA16)
				$3.7 \pm 0.9$	(1967PH03)
		7.57	$\frac{7}{2}^+$	< 10	(1965WA16)
				$7.3 \pm 1.0$	(1967PH03)
		8.31 + 8.58		< 1	(1967PH03)
		0	$\frac{1}{2}^-$	$80 \pm 10$	(1965WA16)
				$77.6 \pm 1.9$	(1967PH03)
		5.27 + 5.30		$10 \pm 10$	(1965WA16)
				$15.4 \pm 1.5$	(1967PH03)
		6.32	$\frac{3}{2}^-$	$10 \pm 10$	(1965WA16)
				$4.9 \pm 1.2$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
		< 1	(1967PH03)		
7.30	$\frac{3}{2}^+$	< 3	(1965WA16)		
		$2.1 \pm 0.8$	(1967PH03)		
7.57	$\frac{7}{2}^+$	< 10	(1965WA16)		
8.31	$\frac{1}{2}^+$	< 2	(1965WA16)		
8.58	$\frac{3}{2}^+$	< 2	(1965WA16)		
7.57, 8.31, 8.58		< 1	(1967PH03)		
10.07	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	100	(1965WA16)
				$96.0 \pm 0.7$	(1967PH03)
				$94 \pm 4$	(1966WA08)
		5.27 + 5.30		< 10	(1965WA16)
				$4.0 \pm 0.7$	(1967PH03)
				$6 \pm 2$	(1966WA08)
6.32	$\frac{3}{2}^-$	< 5	(1965WA16)		
		< 2	(1966WA08)		

Table 15.10: Radiative decays in  $^{15}\text{N}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
10.45	$\frac{3}{2} \rightarrow \frac{7}{2}$	7.16	$\frac{5}{2}^+$	< 7	(1965WA16)
				< 2	(1966WA08)
		7.30	$\frac{3}{2}^+$	< 3	(1965WA16)
				< 2	(1966WA08)
		7.57	$\frac{7}{2}^+$	< 7	(1965WA16)
				< 2	(1966WA08)
		8.31	$\frac{1}{2}^+$	< 2	(1965WA16)
		8.58	$\frac{3}{2}^+$	< 3	(1965WA16)
		0	$\frac{1}{2}^-$	$12 \pm 12$	(1965WA16)
				< 4	(1966WA08, 1967PH03)
		5.27 + 5.30		$72 \pm 8$	(1965WA16)
				$66 \pm 5$	(1966WA08)
				$62.4 \pm 2.4$	(1967PH03)
		6.32	$\frac{3}{2}^-$	$12 \pm 8$	(1965WA16)
				$28 \pm 5$	(1966WA08)
				$14.7 \pm 1.6$	(1967PH03)
		7.16	$\frac{5}{2}^+$	< 10	(1965WA16)
				< 6	(1966WA08)
		7.30	$\frac{3}{2}^+$	$4 \pm 4$	(1965WA16)
				< 6	(1966WA08)
		7.57	$\frac{7}{2}^+$	< 10	(1965WA16)
				< 6	(1966WA08)
		8.31	$\frac{1}{2}^+$	$1.5 \pm 0.5$	(1967PH03)
9.05	$\frac{1}{2}^+$	< 1	(1967PH03)		
9.15 + 9.23		$5.0 \pm 0.5$	(1967PH03)		
9.76	$\frac{5}{2}^-$	$1.6 \pm 0.7$	(1967PH03)		
9.83	$\frac{7}{2}$	$2.2 \pm 1.5$	(1967PH03)		
9.93	$(\frac{1}{2}, \frac{3}{2})^+$	$3.7 \pm 1.1$	(1967PH03)		
10.07	$\frac{3}{2}^+$	< 4	(1967PH03)		
10.54 <sup>b</sup>	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	$1 \pm 0.3$	(1969SI04)
		5.27	$\frac{5}{2}^+$	$30 \pm 2$	(1969SI04)
		6.32	$\frac{3}{2}^-$	$7 \pm 0.5$	(1969SI04)
		7.16	$\frac{5}{2}^+$	$23 \pm 1.5$	(1969SI04)

Table 15.10: Radiative decays in  $^{15}\text{N}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	Refs.
10.70	$\frac{3}{2}^+$	7.30	$\frac{3}{2}^+$	$37 \pm 2.5$	(1969SI04)
		8.58	$\frac{3}{2}^+$	$1.8 \pm 0.4$	(1969SI04)
		(9.22)	$\leq \frac{5}{2}^-$	$< 1$	(1969SI04)
		0	$\frac{1}{2}^-$	$52 \pm 1$	(1969SI04)
		5.27	$\frac{5}{2}^+$	$38 \pm 1$	(1969SI04)
		6.32	$\frac{3}{2}^-$	$6 \pm 0.4$	(1969SI04)
		7.16	$\frac{5}{2}^+$	$< 1$	(1969SI04)
		7.30	$\frac{3}{2}^+$	$3 \pm 0.3$	(1969SI04)
		8.31	$\frac{1}{2}^+$	$< 1$	(1969SI04)
		9.05	$\frac{1}{2}^+$	$< 1$	(1969SI04)
10.80	$\frac{3}{2}^+$	(9.22)	$\leq \frac{5}{2}^-$	$< 1$	(1969SI04)
		0	$\frac{1}{2}^-$	$55 \pm 5$	(1965WA16)
				$53 \pm 15$	(1966WA08)
				$47 \pm 1$	(1969SI04)
		5.27	$\frac{5}{2}^+$	$5 \pm 0.5$	(1969SI04)
		5.30	$\frac{1}{2}^+$	$13 \pm 0.6$	(1969SI04)
		5.27 + 5.30		$45 \pm 5$	(1965WA16)
				$47 \pm 5$	(1966WA08)
		6.32	$\frac{3}{2}^-$	$< 5$	(1965WA16)
				$< 10$	(1966WA08)
				$7 \pm 0.5$	(1969SI04)
		7.16	$\frac{5}{2}^+$	$< 5$	(1965WA16)
				$9 \pm 0.5$	(1969SI04)
		7.30	$\frac{3}{2}^+$	$< 5$	(1965WA16)
				$< 4$	(1966WA08)
		$8 \pm 0.5$	(1969SI04)		
7.57	$\frac{7}{2}^+$	$< 7$	(1965WA16)		
8.31	$\frac{1}{2}^+$	$5 \pm 0.5$	(1969SI04)		
9.05	$\frac{1}{2}^+$	$1 \pm 0.3$	(1969SI04)		
9.152	$\frac{3}{2}^-$	$2 \pm 0.3$	(1969SI04)		
9.155	$(\frac{5}{2})$	$4 \pm 0.3$	(1969SI04)		

<sup>a</sup> See also (1965WA16, 1967PH03).

<sup>b</sup> See also (1960HE13).

<sup>c</sup> See also (1969SI04).

$$23. \text{}^{13}\text{C}(\text{}^6\text{Li}, \alpha)\text{}^{15}\text{N} \quad Q_m = 14.688$$

Angular distributions have been measured for  $E(^6\text{Li}) = 3.2$  to  $3.8$  MeV for the  $\alpha_0$  and  $\alpha_{1+2}$  groups (1964BL1B).

$$24. \text{}^{13}\text{C}(\text{}^7\text{Li}, \text{}^5\text{He})\text{}^{15}\text{N} \quad Q_m = 6.478$$

See (1969TH01).

$$25. \text{}^{13}\text{C}(\text{}^{11}\text{B}, \text{}^9\text{Be})\text{}^{15}\text{N} \quad Q_m = 0.341$$

See (1966PO1E, 1967PO1E, 1967VO1A).

$$26. \text{}^{14}\text{C}(\text{p}, \gamma)\text{}^{15}\text{N} \quad Q_m = 10.208$$

Resonances for capture  $\gamma$ -radiation are listed in Table 15.11 (1959FE1C, 1959HE1D, 1968HE12, 1968SI1F, 1969SI04). A combination of  $^{15}\text{N}^*(10.80)$  and  $(9.83)$  permits a good account of the low energy  $(n, n)$  and  $(n, \gamma)$  cross sections (1959HE1D). The thermal  $(n, p)$  cross section can be ascribed to the  $E_p = 1.5$  MeV resonance ( $^{15}\text{N}^*(11.62)$ ) (1955BA44: see also  $^{14}\text{N}(n, \gamma)^{15}\text{N}$ ). See also (1959AJ76) and (1969TI05).

Table 15.10 displays branching ratios obtained in this and in other reactions. The angular distributions of  $\gamma$ -rays at the  $E_p = 0.35$  MeV resonance ( $^{15}\text{N}^*(10.54)$ ) leads to assignments of  $J = \frac{5}{2}$ ,  $(\frac{5}{2})$ ,  $\frac{3}{2}$  and  $\frac{5}{2}$ , respectively, for  $^{15}\text{N}^*(5.27, 7.16, 7.30, 10.54)$  (1960HE13). The angular distribution of the  $(10.81 \rightarrow 8.31)$   $\gamma$ -ray fixes  $J = \frac{1}{2}$  for  $^{15}\text{N}^*(8.31)$  (1968SI1E). A triple correlation study by (1968SI1E) of the decay of  $^{15}\text{N}^*(10.80)$  to the  $E_x = 9.15$  MeV states suggests  $J = \frac{3}{2}$  for the upper and  $J = \frac{5}{2}$  for the lower of these two states. Lifetimes for various  $^{15}\text{N}$  states have been measured by (1968COZV): see Table 15.7.

See also (1960FR09, 1961FR1D, 1969ZH1A).

$$27. \text{}^{14}\text{C}(\text{p}, \text{p})\text{}^{14}\text{C} \quad E_b = 10.208$$

Table 15.11: Resonances in  $^{14}\text{C} + \text{p}$  <sup>a</sup>

$E_p$ (MeV $\pm$ keV)	$\Gamma$ (keV)	$\Gamma_n$ (keV)	$\Gamma_p$ (keV)	$\Gamma_\gamma$ (eV)	$J^\pi$ <sup>d</sup>	$E_x$ (MeV)	Refs.
$0.261 \pm 1$ <sup>a</sup>						10.451	(1959HE1D, 1968SI1F, 1969SI04)
$0.352 \pm 1$ <sup>a</sup>				$(3.4 \pm 0.4) \times 10^{-2}$ <sup>c</sup>	$\frac{1}{2}^0$	10.536	(1959HE1D, 1960HE13, 1968SI1F, 1969SI04)
$0.527 \pm 1$ <sup>a</sup>			0.2	$1.78 \pm 0.3$ <sup>c</sup>	$\frac{3}{2}^+$	10.700	(1959HE1D, 1968SI1F, 1969SI04)
$0.634 \pm 1$ <sup>a</sup>				$0.23 \pm 0.04$ <sup>c</sup>	$\frac{3}{2}^+$	10.800	(1959HE1D, 1968SI1F, 1969SI04)
$1.162 \pm 2$	$7.9 \pm 3$	2.3	5.6	0.29	$\frac{1}{2}^-$	11.292	(1951RO16, 1955BA44, 1956SA06, 1968HA27)
$1.294 \pm 4$	$34 \pm 4$	22.6	11.1			<sup>e</sup>	(1968HA27)
$1.3188 \pm 0.5$	$41.4 \pm 1.1$	$34.6 \pm 0.9$	$6.8 \pm 0.5$	$4.2 \pm 0.7$	$\frac{1}{2}^+$	11.438	(1951RO16, 1955BA44, 1956SA06, 1968HA27)
$1.472 \pm 8$	$418 \pm 16$	3.7	414.6			<sup>e</sup>	(1968HA27)
$1.509 \pm 4$	$404.9 \pm 6.3$	$4.0 \pm 0.2$	$400.9 \pm 6.3$	$19.2 \pm 0.4$	$\frac{1}{2}^+; T = \frac{3}{2}$	11.615	(1959FE1C, 1968HE12)
$1.688 \pm 3$	37	36.5	0.5		$\frac{3}{2}^+$	11.764	(1951RO16, 1955BA44, 1956SA06)
$1.788 \pm 3$	24.5	24.5	0.03		$\frac{3}{2}^-, (\frac{5}{2}^-)$	11.876	(1951RO16, 1955BA44, 1956SA06)
$1.884 \pm 3$	21.5	21.2	0.3		$\frac{1}{2}^-$	11.965	(1951RO16, 1955BA44, 1956SA06)
$2.025 \pm 4$ <sup>b</sup>	$14 \pm 5$	12.0	1.7		$\frac{1}{2}^+$	12.097	(1951RO16, 1955BA44, 1956SA06, 1960BA34, 1968HA27)
$2.077 \pm 3$ <sup>b</sup>	$47 \pm 7$	30.2	16.6		$\frac{3}{2}^-$	12.145	(1951RO16, 1955BA44, 1956SA06, 1960BA34, 1968HA27, 1968HE12)
$2.272 \pm 4$	22	21.7	0.3		$\frac{1}{2}^+(+)$	12.327	(1951RO16, 1955BA44, 1956SA06, 1960BA34, 1968HA27)
$2.450 \pm 4$ <sup>b</sup>	40	28	0.3		$\frac{1}{2}^0; T = \frac{1}{2}$	12.493	(1951RO16, 1956SA06, 1960BA34)
$2.48 \pm 10$	80	0	80		$\frac{1}{2}^+; T = \frac{3}{2}$	12.52	(1968HA27, 1968HE12)
$2.908 \pm 4$	75				$\frac{3}{2}^-$	12.921	(1956SA06, 1960BA34, 1961HA12, 1962HA20)
3.19	5.5					13.19	(1960BA34, 1961HA12, 1962HA20)
3.38	22					13.36	(1960BA34, 1961HA12, 1962HA20)

Table 15.11: Resonances in  $^{14}\text{C} + \text{p}$  <sup>a</sup> (continued)

$E_{\text{p}}$ (MeV $\pm$ keV)	$\Gamma$ (keV)	$\Gamma_{\text{n}}$ (keV)	$\Gamma_{\text{p}}$ (keV)	$\Gamma_{\gamma}$ (eV)	$J^{\pi}$ <sup>d</sup>	$E_{\text{x}}$ (MeV)	Refs.
3.42	61					13.40	(1960BA34, 1961HA12, 1962HA20)
3.55						(13.52)	(1962HA20)
3.63	12					13.60	(1960BA34, 1961HA12, 1962HA20)
3.71						13.67	(1961HA12, 1962HA20)
3.89	33					13.84	(1960BA34, 1961HA12, 1962HA20)
4.19	105					14.12	(1960BA34, 1961HA12, 1962HA20)
4.24	25					14.17	(1960BA34, 1961HA12, 1962HA20)
4.61	130					14.51	(1960BA34, 1961HA12, 1962HA20)
4.93	99					14.81	(1960BA34, 1961HA12)
5.01						14.88	(1961HA12, 1962HA20)
5.14						15.00	(1961HA12, 1962HA20)
5.26						15.11	(1961HA12, 1962HA20)

<sup>a</sup> See also (1959HE1B) and Table 15.5 in (1959AJ76).

<sup>b</sup> The  $\Gamma_{\alpha}$  are  $< 0.3$  keV for  $E_{\text{res}} = 1.16$  to 2.5 MeV, except for the resonances at  $E_{\text{p}} = 2.03, 2.08$  and 2.45 MeV for which they are, respectively, 0.6, 2.2 and 5.5 keV (1956SA06).

<sup>c</sup>  $\omega_{\gamma}$  (in eV) (1968SI1F, 1969SI04).

<sup>d</sup> See also (1959AJ76).

<sup>e</sup> These states correspond to the levels listed in the line below: a different boundary condition was used to obtain  $E_{\text{res}}$ .

The elastic scattering has been studied for  $E_p = 0.34$  to  $2.7$  MeV. At the  $E_p = 527$  keV resonance (see Table 15.11), the scattering is consistent with d-wave formation of a  $J^\pi = \frac{3}{2}^+$  state. No anomalies are observed at  $E_p = 0.35$  and  $0.63$  MeV (1959HE1D). However, anomalies are observed at  $E_p = 1.16, 1.29, 1.47, 2.02, 2.07$  and  $2.48$  MeV: the parameters of these are displayed in Table 15.11 (1968HA27, 1968HE12). The  $2.48$  MeV anomaly is due to a  $J^\pi = \frac{5}{2}^+$ ;  $T = \frac{3}{2}$  state at  $E_x = 12.52$  MeV (which is the analog to the first excited state of  $^{15}\text{C}$ ), and is distinct from the  $J^\pi = \frac{5}{2}^+$  resonance in  $^{14}\text{C}(p, n)^{14}\text{N}$  observed at  $E_p = 2.45$  MeV (1968HE12). See also (1968IW1A, 1969IW1E).

$$28. \ ^{14}\text{C}(p, n)^{14}\text{N} \qquad Q_m = -0.626 \qquad E_b = 10.208$$

Resonances reported by (1951RO16, 1955BA44, 1956SA06, 1959FE1C, 1960BA34, 1961HA12, 1962HA20, 1968HE12) are listed in Table 15.11. In addition to these, (1961HA12) report (in an abstract) a broad resonance corresponding to a  $T = \frac{3}{2}$  state of  $^{15}\text{N}$  at  $E_x = 13.6$  MeV. At  $E_p = 1.79$  MeV, the distributions favor  $\frac{5}{2}^-$ , but  $\frac{3}{2}^-$  is not excluded (1955BA44: see also (1953KA1A)); a computation of the cross section favors  $J = \frac{3}{2}$  (1956SA06). At  $E_p = 1.88$  MeV, the angular distribution is consistent with the  $J^\pi = \frac{1}{2}^-$  assignment from  $^{14}\text{N}(n, n)^{14}\text{N}$  (1955BA44). The  $E_p = 2.27$  MeV state has  $J = \frac{3}{2}$  or  $\frac{5}{2}$ ; the  $\sigma_{nn}$  clearly indicates the latter (1955BA44, 1956SA06). For  $^{15}\text{N}^*(11.62)$  ( $E_p = 1.51$  MeV), the proton reduced width indicates a single-particle level, while the neutron reduced width is only  $10^{-3}$ . This is consistent with the assignment  $T = \frac{3}{2}$ , corresponding to the ground state of  $^{15}\text{C}$  (1955BA44, 1956BA16, 1959FE1C). See also (1967VO1B, 1969BA1N, 1969TI05).

Polarization measurements are reported at  $E_p = 7.2, 8.8$  and  $10.4$  MeV by (1969WO1J;  $n_0, n_1, n_2$ ).

$$29. \ ^{14}\text{C}(d, n)^{15}\text{N} \qquad Q_m = 7.984$$

Angular distributions have been determined at  $E_d = 1.3, 1.8$  and  $2.8$  MeV (1967CO1R;  $n_0, n_{1+2}, n_3, n_{4+5+6}$ ),  $1.31$  to  $3.08$  MeV (1963IM01;  $n_0$ ),  $2.10$  and  $2.33$  MeV (1963IM01;  $n_{1+2}, n_3, n_{4+5+6}$ ) and  $3.10$  MeV (1967LA11;  $n_0, n_{1+2}, n_3, n_4, n_5, n_6$  and  $^{15}\text{N}^*(8.31, 9.06, 9.16, 9.22)$ ). The transitions to  $^{15}\text{N}^*(5.30, 9.06)$  involve  $l_p = 0$ : these states therefore have  $J^\pi = \frac{1}{2}^+$  (1967LA11). See also (1959AJ76) and (1961CH14, 1964MO1G).

Studies of the  $\gamma$ -decay of  $^{15}\text{N}$  states reached in this and in other reactions are summarized in Table 15.10: see (1965WA16). See also (1967CH19).

$$30. \ ^{14}\text{C}(^3\text{He}, d)^{15}\text{N} \qquad Q_m = 4.715$$



At  $E(^3\text{He}) = 14$  MeV, angular distributions of the deuterons corresponding to  $^{15}\text{N}^*(0, 5.28, 5.30, 6.33)$  have been measured and analyzed by DWBA: the relative spectroscopic factors for the first four states of  $^{15}\text{N}$  are 1, 0.50, 0.34, 0.07 in good agreement with the predictions of (1957HA1E). Angular distributions of the  $d_0$  group are also reported at  $E(^3\text{He}) = 1 - 9$  MeV by (1964WE1A, 1966DU1B, 1968DA1N).

$$31. \ ^{14}\text{C}(\alpha, t)^{15}\text{N} \quad Q_m = -9.606$$

Not reported.

$$32. \ ^{14}\text{N}(n, \gamma)^{15}\text{N} \quad Q_m = 10.835$$

$$Q_0 = 10.8332 \pm 0.0006 \text{ (1968GR14).}$$

The thermal cross section is  $80 \pm 20$  mb (1957BA18),  $75 \pm 7.5$  mb (1964ST25). This large cross section is not understood in terms of the present level structure in  $^{15}\text{N}$ : see (1959AJ76).

Observed capture  $\gamma$  rays are displayed in Table 15.12 (1963MO1C, 1967TH05, 1968GR14). The very accurate  $\gamma$ -ray energy determinations of (1968GR14) show that two states at  $E_x \approx 9.15$  MeV are involved in this reaction as previously suggested by (1966WA08). The lower of the two, at  $E_x = 9.1518$  MeV, decays predominantly to the ground state. The other state at  $E_x = 9.1549$  MeV, which is preferentially fed in this reaction, decays primarily by cascades via  $^{15}\text{N}^*(5.27, 5.30, 6.32, 7.16)$  (1968GR14). See also reactions 21 and 39, (1968GRZY, 1968ST10) and Table 15.10.

Recoil Doppler broadening of cascade  $\gamma$ -rays has been measured by (1969WE07): the derived  $\tau_m$  are listed in Table 15.7. See also (1968CA1J). Observation of non-isotropic correlations in the  $\text{C} \rightarrow 6.32$  cascade means that  $J \neq \frac{1}{2}$  for  $^{15}\text{N}^*(6.32)$ : the results are consistent with  $J = \frac{3}{2}$  (1964BA02).

The importance of measuring the  $(n, \gamma)$  cross section at  $E_n = 0.47$  and  $0.66$  MeV for astrophysical considerations is suggested by (1968FO1A).

See also (1960CA02, 1965JA09, 1969HO1X) and (1958GR1B, 1964LI1C).

$$33. \ ^{14}\text{N}(n, n)^{14}\text{N} \quad E_b = 10.835$$

The thermal (bound) scattering cross section is  $5.51$  b (1961WI1A). The scattering amplitude (bound) is  $a = 9.19 \pm 0.11$  fm (1965DO1B) [ $a = 9.14 \pm 0.10$  fm is recommended by (1964ST25)]. See also (1969BA1P). The coherent scattering cross section is  $10.5 \pm 0.3$  b (1964ST25).

Recent cross section measurements are listed in Table 15.13. Cross section data are summarized in (1964ST25), while angular distribution data are displayed in (1963GO1M).

Observed resonances are listed in Table 15.14 (1951JO1A, 1952HI12, 1955FO27, 1959HA13, 1966FO1D, 1966FO1E, 1966MA2K, 1968BO36, 1968JO1F): for a discussion of the evidence

Table 15.12: Gamma radiation from  $^{14}\text{N}(n, \gamma)^{15}\text{N}$ 

Transition in $^{15}\text{N}$	$E_\gamma$ (MeV $\pm$ keV)		$E_\gamma$ (MeV $\pm$ keV)	$I_\gamma$ <sup>b</sup>	
	(1967TH05)	(1968GR14)	(1968GR14)	(1967TH05)	(1963MO1C)
C $\rightarrow$ 0	10.830 $\pm$ 2	10.8290	10.8832 $\pm$ 0.6	13.3 $\pm$ 2.0	14
C $\rightarrow$ 5.27	5.562 $\pm$ 1	5.5622 $\pm$ 0.35		10.3 $\pm$ 0.5	11
C $\rightarrow$ 5.30	5.534 $\pm$ 1	5.5332 $\pm$ 0.35		18.8 $\pm$ 0.9	21
C $\rightarrow$ 6.32	4.509 $\pm$ 1	4.5088 $\pm$ 0.3		16.6 $\pm$ 0.8	16
C $\rightarrow$ 7.16	3.678 $\pm$ 1	3.6777 $\pm$ 0.25		15.9 $\pm$ 0.8	< 23
C $\rightarrow$ 7.30	3.532 $\pm$ 1	3.5322 $\pm$ 0.25		9.9 $\pm$ 0.5	9
C $\rightarrow$ 8.31	2.521 $\pm$ 2	2.52055 $\pm$ 0.10		6.1 $\pm$ 0.3	6
C $\rightarrow$ 9.155		1.67819 $\pm$ 0.06			
	1.679 $\pm$ 2			9.2 $\pm$ 0.5	12
C $\rightarrow$ 9.152		1.68141 $\pm$ 0.18		1.4 $\pm$ 0.3 <sup>d</sup>	
5.27 $\rightarrow$ 0	5.270 $\pm$ 1	5.2692 $\pm$ 0.35	5.2701 $\pm$ 0.3	30.6 $\pm$ 1.5	32
5.30 $\rightarrow$ 0	5.298 $\pm$ 1	5.2978 $\pm$ 0.35	5.2989 $\pm$ 0.3	21.4 $\pm$ 1.1	21
6.32 $\rightarrow$ 0	6.323 $\pm$ 1	6.3220 $\pm$ 0.4	6.3235 $\pm$ 0.4	18.8 $\pm$ 0.9	18
7.16 $\rightarrow$ 0			7.1550 $\pm$ 0.4		
7.17 $\rightarrow$ 5.27	1.885 $\pm$ 1	1.88481 $\pm$ 0.06		19.7 $\pm$ 1.0	21
7.16 $\rightarrow$ 5.30	1.857 $\pm$ 2			0.8 $\pm$ 0.2	
7.30 $\rightarrow$ 0	7.299 $\pm$ 1	7.2990 $\pm$ 0.5	7.3009 $\pm$ 0.5	10.0 $\pm$ 0.5	9
8.31 $\rightarrow$ 0	8.311 $\pm$ 2	8.3102 $\pm$ 0.7	8.3124 $\pm$ 0.7	4.4 $\pm$ 0.4	4
8.31 $\rightarrow$ 6.32	1.989 $\pm$ 2			1.5 $\pm$ 0.3	
8.57 $\rightarrow$ 0	8.570 $\pm$ 4		8.573 $\pm$ 4 <sup>c</sup>	0.2 $\pm$ 0.03	
9.05 $\rightarrow$ 0	9.047 $\pm$ 4		9.050 $\pm$ 4 <sup>c</sup>	0.2 $\pm$ 0.03	0.5
9.155 $\rightarrow$ 0			9.1549 $\pm$ 0.5		
9.152 $\rightarrow$ 0	9.151 $\pm$ 2	9.1498 $\pm$ 0.9	9.1518 $\pm$ 0.5	1.7 $\pm$ 0.2	1.4
9.155 $\rightarrow$ 5.27	3.883 $\pm$ 2	3.8837 $\pm$ 0.4		0.8 $\pm$ 0.1	
9.155 $\rightarrow$ 5.30	3.855 $\pm$ 2	3.8552 $\pm$ 0.3		1.0 $\pm$ 0.1	0.5
9.155 $\rightarrow$ 6.32	2.830 $\pm$ 2	2.8311 $\pm$ 0.2		2.0 $\pm$ 0.2	1.5
9.155 $\rightarrow$ 7.16	1.997 $\pm$ 2	1.99965 $\pm$ 0.10		4.6 $\pm$ 0.2	4

<sup>a</sup> C = capturing state.

<sup>b</sup> In units of photons/100 captures.

<sup>c</sup> (1967TH05).

<sup>d</sup> (1968GR14).

leading to  $J^\pi$  assignments, see (1959AJ76). See also (1960BA34) and (1966AG1A, 1967BE1F, 1968IW1A, 1969IW1E).

A polarization measurement has been made at  $E_n = 3.5$  MeV by (1962OT01). See also (1965TA07).

34.  $^{14}\text{N}(n, 2n)^{13}\text{N}$

$Q_m = -10.553$

$E_b = 10.835$

Table 15.13: Recent cross section measurements for  $^{14}\text{N} + n$  <sup>a</sup>

$E_n$ (MeV)	Measurements of	Refs.
0.01 – 0.2	$\sigma_t$	(1959BI1B)
1.7 – 4.0	$\sigma_t$	(1960BA34)
2.0 – 4.2	$\sigma_t$	(1966MA2K, 1968JO1F)
4.2 – 6.3	$\sigma_{ne}$	(1968BO36)
4.2 – 8.7	$\sigma_e$	(1968BO36)
4.5, 6, 7	$\sigma_{nn'\gamma}$	(1968CO1W)
5.80 – 8.55	$\sigma_{nn'\gamma}$	(1969DI1B)
6.8 – 14.0	$\sigma_t, \sigma_e, \sigma_{ne}$	(1967BA03)
13.3 – 19.5	$\sigma_e$	(1968BO36)
14	$\sigma_e$	(1963BA46)
15	$\sigma_{ne}$	(1969NY1A)
15.2, 19.8	$\sigma_{ne}$	(1965DE1G)
15.3	$\sigma_t, \sigma_{ne}$	(1968HA1V)
17.7 – 29.1	$\sigma_t$	(1960PE1B)
88 – 152	$\sigma_t$	(1966ME14)
10 – 37	$\sigma_{n,2n}$	(1961BR1A)
12.5 – 18	$\sigma_{n,2n}$	(1960FE12)
13.2 – 18.8	$\sigma_{n,2n}$	(1965BO42)
13.3 – 15.2	$\sigma_{n,2n}$	(1960MC05)
14.1	$\sigma_{n,2n}$	(1962CE1B)
14.4	$\sigma_{n,2n}$	(1961RA06)
14.7	$\sigma_{n,2n}$	(1967PA1N)
14.8	$\sigma_{n,2n}$	(1965GR1T)
1.3 – 4.2	$\sigma_{n,p}$	(1959GA14)

Table 15.13: Recent cross section measurements for  $^{14}\text{N} + \text{n}$  <sup>a</sup> (continued)

$E_n$ (MeV)	Measurements of	Refs.
7.4 – 8.6	$\sigma_{n,p}$	(1969DI1B)
14.7	$\sigma_{n,p}, \sigma_{n,np}$	(1966CS1B, 1967CS1A)
15	$\sigma_{n,p}, \sigma_{n,d}$	(1969NY1A)
5.6 – 6.4	$\sigma_{n,t}$	(1966SC21)
5.7 – 8.2	$\sigma_{n,t}$	(1959GA14)
14.1	$\sigma_{n,2\alpha}$	(1967MO21)
15	$\sigma_{n,t}$	(1969NY1A)
15.7	$\sigma_{n,t}$	(1967MO21)
1.3 – 8.2	$\sigma_{n,\alpha}$	(1959GA14)
3.9 – 6.4	$\sigma_{n,\alpha}$	(1966SC21)
4.5 – 7.0	$\sigma_{n,\alpha}$	(1968CO1W)
5.7 – 8.2	$\sigma_{n,\alpha}$	(1959HA13)
5.8 – 8.6	$\sigma_{n,\alpha}$	(1969DI1B)
14.7	$\sigma_{n,\alpha}$	(1966CS1B)
15	$\sigma_{n,\alpha}$	(1969NY1A)

<sup>a</sup> See also (1959AJ76).

Recent measurements of cross sections for this reaction are listed in Table 15.13 (1960FE12, 1960MC05, 1961BR1A, 1961RA06, 1962CE1B, 1965BO42, 1965GR1T, 1967PA1N): see the summaries in (1964ST25, 1966JE1B). See also (1964HE18, 1966CS1C, 1967CS02) and (1959AJ76).

35.  $^{14}\text{N}(n, p)^{14}\text{C}$

$$Q_m = 0.626$$

$$E_b = 10.835$$

The thermal cross-section is  $1.81 \pm 0.05$  b (1964ST25),  $1.83 \pm 0.03$  b (1961HA43). A number of resonances are reported by (1950JO57), (1959GA14) and (1963EN01): see Table 15.14. The results are summarized in (1964ST25, 1966JE1B). See also (1960BU1C, 1968DA1F).

Recent cross-section measurements are listed in Table 15.13 (1959GA14, 1966CS1B, 1967CS1A, 1969DI1B). See also (1964FO1A, 1969BA1N) and (1959AJ76).

36. (a)  $^{14}\text{N}(n, d)^{13}\text{C}$

$$Q_m = -5.325$$

$$E_b = 10.835$$

(b)  $^{14}\text{N}(n, np)^{13}\text{C}$

$$Q_m = -7.550$$

For reaction (a) see (1967LI06) and (1959AJ76). For reaction (b) see (1966CS1B, 1967CS1A).

Table 15.14: Resonances in  $^{14}\text{N} + \text{n}$  <sup>a</sup>

$E_{\text{res}}$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	$\Gamma_{\text{n}}$ (keV)	$\Gamma_{\text{p}}$ (keV)	$\Gamma_{\alpha}$ (keV)	$J^{\pi}$	$^{15}\text{N}^*$ (MeV)	Refs.
$0.430 \pm 5$	3.5	< 3	< 0.01		$\geq \frac{1}{2} \frac{3}{2} \frac{3}{2}$	11.236	(1951JO1A, 1952HI12)
$0.4926 \pm 0.65$	7.5	< 3	< 10		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{-}$	11.2943	(1950JO57, 1952HI12, 1963EN01)
$0.639 \pm 5$	43	34	9		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{+}$	11.431	(1950JO57, 1951JO1A, 1952HI12)
$0.998 \pm 5$	46	45	0.8		$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	11.766	(1950JO57, 1951JO1A, 1952HI12)
$1.120 \pm 6$	19	19	0.20		$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	11.880	(1951JO1A, 1952HI12, 1955FO27)
$1.188 \pm 6$	$\leq 3.2$	< 2	< 0.1		$\geq \frac{1}{2} \frac{1}{2} \frac{1}{2}$	11.943	(1952HI12)
$1.211 \pm 7$	13	12	0.4		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{-}$	11.965	(1952HI12)
$1.350 \pm 7$	21	20	0.9	0.4	$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{+}$	12.094	(1951JO1A, 1952HI12, 1955FO27, 1959GA14)
$1.401 \pm 8$	54	41	11	1.8	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	12.142	(1950JO57, 1951JO1A, 1952HI12, 1955FO27, 1959GA14)
$1.595 \pm 8$	22	21	0.2	< 0.1	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	12.323	(1952HI12, 1955FO27, 1959GA14, 1966FO1D, 1966FO1E)
$1.779 \pm 10$	47	37	0.5	9.0	$(\frac{3}{2} \frac{3}{2} \frac{3}{2})^{+}$	12.494	(1950JO57, 1952HI12, 1955FO27, 1959GA14, 1966FO1E)
2.23	65	39	7.8	18	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	12.92	(1959GA14, 1966FO1D, 1966FO1E, 1966MA2K, 1968JO1F)
2.47	< 3			r		13.14	(1959GA14)
2.52	$\approx 7$			r		13.19	(1959GA14)
2.71	40			r	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{-}$	13.36	(1959GA14)
2.74	95		r		$\frac{1}{2} \frac{1}{2} \frac{1}{2}^{+}$	13.39	(1959GA14)
2.95	20	16	1.1	3.2	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	13.59	(1959GA14, 1966FO1E, 1966MA2K, 1968JO1F)
3.09	60		r	r		13.72	(1959GA14)
3.21	85		r	r	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	13.83	(1959GA14, 1966FO1E, 1966MA2K, 1968JO1F)
3.51	$\approx 20$		r	r		14.11	(1959GA14)
3.57	30		r	r	$\frac{3}{2} \frac{3}{2} \frac{3}{2}^{+}$	14.17	(1959GA14, 1966MA2K, 1968JO1F)
$\approx 3.8$	$\approx 2000$	$\approx 1000$	200	$\approx 1000$		14.4	(1959GA14)
4.09	50		r	r		14.65	(1959GA14, 1966SC21)
$\approx 4.1$	$\approx 300$		r	r		14.7	(1959GA14)
4.38	40			r		14.92	(1959GA14, 1966SC21)
4.60				r		15.13	(1959GA14, 1966SC21)
5.03				r		15.52	(1959GA14, 1966SC21, 1968BO36)
5.60	100			r		16.06	(1959GA14, 1966SC21, 1968BO36)

Table 15.14: Resonances in  $^{14}\text{N} + \text{n}$  <sup>a</sup> (continued)

$E_{\text{res}}$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	$\Gamma_{\text{n}}$ (keV)	$\Gamma_{\text{p}}$ (keV)	$\Gamma_{\alpha}$ (keV)	$J^{\pi}$	$^{15}\text{N}^*$ (MeV)	Refs.
5.94				r		16.38	(1959GA14, 1966SC21)
6.16	75			r		16.58	(1959GA14, 1966SC21)
6.26	110	r		r		16.67	(1959GA14, 1959HA13, 1966SC21)
6.55	170	r		r		16.95	(1959GA14, 1959HA13)
6.94				r		17.31	(1959GA14)
	200	r					(1959HA13)
7.16				r		17.51	(1959GA14)
7.34	120			r		17.68	(1959GA14)
7.48	180	r		r		17.81	(1959GA14, 1959HA13)
7.92	170			r		18.22	(1959HA13)
8.00	120			r		18.30	(1959GA14, 1959HA13)

r = resonant channel.

<sup>a</sup> See also (1959AJ76).

37. (a) $^{14}\text{N}(\text{n}, \text{t})^{12}\text{C}$	$Q_{\text{m}} = -4.015$	$E_{\text{b}} = 10.835$
(b) $^{14}\text{N}(\text{n}, \text{t})^4\text{He}^4\text{He}^4\text{He}$	$Q_{\text{m}} = -11.289$	
(c) $^{14}\text{N}(\text{n}, 2\alpha)^7\text{Li}$	$Q_{\text{m}} = -8.822$	

For reaction (a) see (1959GA14, 1966SC21, 1967LI06, 1967RE01) and (1968AJ02). For reaction (b) see (1967MO21). For reaction (c) see (1967MO21). See also (1960FA10, 1964SA1E), Table 15.13, the summaries in (1964ST25, 1966JE1B) and (1959AJ76).

38. (a) $^{14}\text{N}(\text{n}, \alpha)^{11}\text{B}$	$Q_{\text{m}} = -0.157$	$E_{\text{b}} = 10.835$
(b) $^{14}\text{N}(\text{n}, \text{n}\alpha)^{10}\text{B}$	$Q_{\text{m}} = -11.613$	

Recent cross-section measurements for reaction (a) are displayed in Table 15.13: these include measurements of cross sections for several different  $\alpha$ -groups (1959GA14, 1966SC21) and  $\gamma$ -rays (1959HA13, 1968CO1W, 1969DI1B). For summaries of the experimental evidence, see (1964ST25, 1966JE1B).

Observed resonances are listed in Table 15.14 (1950JO57, 1959GA14, 1966SC21). See also (1960FA10), (1960BU1C, 1963CH1C) and (1964GA1A, 1968GA1M; theor.). For reaction (b), see (1960FA10).

39. $^{14}\text{N}(\text{d}, \text{p})^{15}\text{N}$	$Q_{\text{m}} = 8.610$	
	$Q_0 = 8.614 \pm 0.006$ (1964MA57).	

Proton groups corresponding to levels of  $^{15}\text{N}$  are listed in Table 15.15 (1950MA65, 1954SP01, 1956DO41, 1965AL19, 1966GA08, 1966GO1J, 1969PH02). The  $J^\pi$  assignments are based on PWBA and DWBA analyses: see (1959AJ76), (1969PH02) and Table 15.16 for a listing of recent angular distribution studies in the range  $E_{\text{d}} = 0.5$  to 27 MeV. See also (1966RO1V). The angular distributions of protons corresponding to  $^{15}\text{N}^*(6.32)$  and the p- $\gamma$  angular correlation fixes  $J^\pi = \frac{3}{2}^-$  for that state (1961GO03). (1960HO1B) has looked for additional states of  $^{15}\text{N}$  with  $E_{\text{x}} \approx 7$  MeV: he finds the upper limits to proton groups corresponding to  $7.10 < E_{\text{x}} < 7.31$  to be 1%, and to be 5% of the intensities of groups to known nearby states for  $7.31 < E_{\text{x}} < 7.57$  MeV.

Recent very accurate  $\gamma$ -ray energy measurements have been reported by (1967CH19) [ $E_{\gamma} = 5299.03 \pm 0.43$  and  $5270.60 \pm 0.46$  keV] by (1966AL18) and by (1965WA16): the derived  $E_{\text{x}}$  values are displayed in Table 15.15. Branching ratios have been determined by (1965WA16, 1968GI11, 1968ST10) and are shown in Table 15.10 together with the multiplicities determined by (1965WA16). Lifetime measurements are listed in Table 15.7 (1965AL19, 1967BI11, 1968GI11).

The two states of  $^{15}\text{N}$  at  $E_{\text{x}} = 9.15$  MeV [see Table 15.10 for branching ratios] are separated by  $2.5 \pm 0.5$  keV (1968ST10, 1969YO1C): see also reaction 32.

Table 15.15:  $^{15}\text{N}$  levels from  $^{14}\text{N}(\text{d}, \text{p})^{15}\text{N}$ 

$E_x$ (MeV $\pm$ keV)					$l_n$	$J^\pi$
(1950MA65, 1966GO1J)	(1954SP01)	(1956DO41, 1967CH19)	A	(1966GA08)		
0					1 <sup>b</sup>	$\frac{1}{2}^-$ , $\frac{3}{2}^-$ , $\frac{5}{2}^-$
5.276 $\pm$ 6	5.280 $\pm$ 10	5.27159 $\pm$ 0.46 <sup>a</sup>	5.272 $\pm$ 10		2 <sup>c</sup>	$\leq \frac{7}{2}^+$
5.305 $\pm$ 6		5.30003 $\pm$ 0.43 <sup>a</sup>	5.300 $\pm$ 11		c,d	
6.328 $\pm$ 6	6.330 $\pm$ 10				1 <sup>e,i</sup>	$\frac{3}{2}^-$ j
7.164 $\pm$ 6	7.165 $\pm$ 10		7.1555 $\pm$ 1.7		2 <sup>f,i</sup>	$\leq \frac{7}{2}^+$
7.309 $\pm$ 6	7.314 $\pm$ 10	7.307 $\pm$ 8			0 <sup>f,i</sup>	$\frac{1}{2}^+$ , $\frac{3}{2}^+$
	7.575 $\pm$ 10	7.570 $\pm$ 8	7.5671 $\pm$ 1.0		2 <sup>g,i</sup>	$\leq \frac{7}{2}^+$
8.315 $\pm$ 6	8.316 $\pm$ 10	8.319 $\pm$ 8	8.309 $\pm$ 4.1		0 <sup>e,i</sup>	$\frac{1}{2}^+$ , $\frac{3}{2}^+$
	8.571 $\pm$ 10	8.577 $\pm$ 8	8.573 $\pm$ 3.2	8.582 $\pm$ 5	0 + 2 <sup>h,i</sup>	$\leq \frac{7}{2}^+$
	9.062 $\pm$ 10			9.056 $\pm$ 5		
	9.165 $\pm$ 10			9.159 $\pm$ 6	i	
9.225 $\pm$ 6				9.226 $\pm$ 6	1 or 2 <sup>i</sup>	$(\frac{3}{2}^-)$
9.762 $\pm$ 6				9.764 $\pm$ 6		
	9.834 $\pm$ 10			9.831 $\pm$ 6		
9.929 $\pm$ 7				9.929 $\pm$ 6		
	10.069 $\pm$ 10			10.071 $\pm$ 6	2, 0 <sup>i</sup>	$\frac{3}{2}^+$
	10.458 $\pm$ 10			10.456 $\pm$ 7		
	10.544 $\pm$ 10			10.541 $\pm$ 7		
	10.705 $\pm$ 10			10.702 $\pm$ 7	2, 0 <sup>i</sup>	$\frac{3}{2}^+$
	10.811 $\pm$ 10			10.809 $\pm$ 9	1 <sup>i</sup>	$\frac{1}{2}^-$ , $\frac{3}{2}^-$ , $\frac{5}{2}^-$
	11.2				1 <sup>j</sup>	$\frac{1}{2}^-$ , $\frac{3}{2}^-$ , $\frac{5}{2}^-$



A: (1965AL19, 1965WA16, 1966AL18).

<sup>a</sup> See also (1965AL19, 1965WA03).

<sup>b</sup> (1952GI01, 1957WA01).

<sup>c</sup> (1955SH28: see (1958WA1C)).

<sup>d</sup> Isotropic: no clear stripping pattern.

<sup>e</sup> (1952GI01, 1955SH28, 1956GR37, 1958WA1C).

<sup>f</sup> (1955SH28, 1956GR37).

<sup>g</sup> (1956GR37): (1957WA01) find a possible  $l = 0$  component.

<sup>h</sup> (1955SH28, 1957WA01).

<sup>i</sup> (1969PH02): absolute spectroscopic factors are also given.

<sup>j</sup> (1956GR37).

<sup>k</sup> (1961GO03). (This footnote is not labeled in the table content.)

Table 15.16:  $^{14}\text{N}(d, p)^{15}\text{N}$  angular distribution studies <sup>a</sup>

$E_d$ (MeV)	Distribution of proton groups	Refs.
0.5 – 0.8	$p_0, p_1, p_2$	(1957SJ68, 1961SJ1B)
1 – 2.2	$p_0, p_3 \rightarrow p_7$	(1967BE09)
1 – 3.2	$p_4 \rightarrow p_7$	(1969BE08)
1.1 – 3.1	$p_{1+2}, p_3, p_4, p_5, p_6$	(1969GO14)
1.3, 4.5	$p_{1+2}, p_3$	(1961GO03, 1965FI05, 1966GA09)
1.3 – 5.5	$p_0$	(1962GO21, 1965FI05, 1966GA09)
1.4 – 3	$p_0, p_3$	(1969BE08)
1.4 – 3.2	$p_0, p_1 + p_2$	(1961KA05)
2	$p_1 \rightarrow p_5, p_7$	(1962RO13)
7, 8, 9	$p_{3 \rightarrow 8}, p_{10+11}, p_{12}, p_{16}, p_{19}, p_{20}$	(1969PH02)
12	$p_0$	(1967SC29)
13.8, 15.2	$p_0$	(1961MO13)
16	$p_0$	(1960MO03)
27	$p_0$	(1962ER03)

<sup>a</sup> See (1959AJ76) for earlier references.

See also (1961JA23, 1961TE02, 1962ST17, 1963GO1L, 1965HE1B), (1959BO1C, 1960BE1B, 1961KO1E, 1963TA1A, 1964BA1V, 1964ST1J, 1966HO1D; theor.), (1959AJ76) and  $^{16}\text{O}$  in (1971AJ02).

$$40. \ ^{14}\text{N}(t, d)^{15}\text{N} \quad Q_m = 4.577$$

The angular distribution of the deuterons corresponding to the ground state of  $^{15}\text{N}$  has been measured at  $E_t = 1.50, 1.83$  and  $1.98$  MeV. The cross section at  $E_t = 2$  MeV is 48 mb: this large value and the energy and angular behavior of the differential cross section suggest that this reaction may proceed by a cluster exchange process (1964SC09).

$$41. \ ^{14}\text{N}(\alpha, ^3\text{He})^{15}\text{N} \quad Q_m = -9.743$$

At  $E_\alpha = 56$  MeV, the angular distribution of the ground state  $^3\text{He}$  particles has been measured by (1969GA11) and analyzed by DWBA: the ratio of the  $(\alpha, ^3\text{He})$  and  $(\alpha, t)$  cross sections at this energy is  $1.50 \pm 0.15$ . See also (1968GA1C).

42.  $^{14}\text{N}(^9\text{Be}, ^8\text{Be})^{15}\text{N}$   $Q_m = 9.170$

The lifetime of  $^{15}\text{N}^*(5.27)$  is  $\approx 1$  psec (1969NI09): see Table 15.7 and (1967BI11). See also (1963HO1E).

43.  $^{14}\text{N}(^{11}\text{B}, ^{10}\text{B})^{15}\text{N}$   $Q_m = -0.621$

See (1967PO13). See also (1969BR1D).

44.  $^{14}\text{N}(^{14}\text{N}, ^{13}\text{N})^{15}\text{N}$   $Q_m = 0.282$

Angular distributions of the transition to the ground state of  $^{15}\text{N}$  have been measured for  $E(^{14}\text{N})(\text{cm}) = 5.5$  to 16 MeV (1961TO07, 1964JO1A, 1965BE1B, 1965HI1A). See also (1961TO01). Below  $\approx 6.5$  MeV, the tunneling theory of neutron transfer gives a good account of the data. At higher energies, nuclear absorption of the incident  $^{14}\text{N}$  ions occur (1965HI1A, 1966GA04). See also (1968GA03). For reviews of the work on this reaction, see (1964FL1D, 1967DA1E, 1967VO1A, 1969BR1D). For discussions of relevant theories, see (1962BR1G, 1963BR1G, 1964BR1M, 1964GR1G, 1965BR1F, 1966BU1B, 1967BR1M, 1967BR1Q, 1967PE1D, 1968MA2G, 1968NA1F, 1969KA1G).

45.  $^{14}\text{N}(^{19}\text{F}, ^{18}\text{F})^{15}\text{N}$   $Q_m = 0.405$

See (1968GA03).

46.  $^{15}\text{C}(\beta^-)^{15}\text{N}$   $Q_m = 9.773$

The  $\beta^-$  decay takes place to  $^{15}\text{N}^*(0, 5.30, 7.30, 8.31, 9.05)$ : see Table 15.2.

Measurements of  $\gamma$ -ray energies give  $E_\gamma = 5299.03 \pm 0.43$  keV (1967CH19),  $8315 \pm 6$  and  $9048 \pm 4$  keV (1966AL12). See also  $^{15}\text{C}$ .

47.  $^{15}\text{N}(\gamma, n)^{14}\text{N}$   $Q_m = -10.835$

See (1967ZH1A, 1968YA1E, 1969ZH1A).

48.  $^{15}\text{N}(\gamma, p)^{14}\text{C}$   $Q_m = -10.208$

The integrated cross section for transitions to  $^{14}\text{C}(0)$  for  $E_\gamma$  up to 30.5 MeV =  $22 \pm 3$  MeV·mb, assuming an isotropic angular distribution. Pronounced maxima are observed at  $E_\gamma = 19.5, 20.4, 22.7$  and 24.5 MeV. In addition a “pigmy” resonance at  $E_\gamma = 15.2$  MeV and less pronounced structures at  $E_\gamma = 13.6$  and 17.0 MeV are also observed (1964KO10). See also (1963FI04) and (1967ZH1A, 1968YA1E, 1969UB1C, 1969ZH1A).

49.  $^{15}\text{N}(e, e)^{15}\text{N}$

Using the harmonic oscillator function, analysis of the scattering of 250 and 400 MeV electrons gives, respectively,  $r_{\text{rms}} = 2.68 \pm 0.05$  fm and  $2.63 \pm 0.05$  fm for  $^{15}\text{N}$  (1968DA1Q). At  $E_e = 50-57$  MeV, excitation of  $^{15}\text{N}^*(6.32)$  gives  $\Gamma_\gamma^0(\text{M1}) = 3.4 \pm 0.7$  eV and  $\Gamma_\gamma^0(\text{E2}) = 0.06 \pm 0.02$  eV:  $|\delta(\text{E2/M1})| = 0.13 \pm 0.03$  (1968BE14).

50.  $^{15}\text{N}(n, n)^{15}\text{N}$

See  $^{16}\text{N}$ .

51.  $^{14}\text{N}(p, p)^{15}\text{N}$

The angular distribution of elastically scattered protons has been measured at  $E_p = 39.8$  MeV (1969SN04). See also (1963NA1C).

52.  $^{15}\text{N}(^3\text{He}, ^3\text{He})^{15}\text{N}$

At  $E(^3\text{He}) = 39.8$  MeV, a number of inelastically scattered  $^3\text{He}$  groups are observed corresponding to states in  $^{15}\text{N}$ : see Table 15.17. Angular distributions were obtained for a number of these, and were analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture (1969BA06). See also (1968BA1E). (1969BO13) measured the angular distribution of elastically scattered  $^3\text{He}$  particles at  $E(^3\text{He}) = 11$  MeV.

53.  $^{15}\text{N}(\alpha, \alpha)^{15}\text{N}$

Table 15.17:  $^{15}\text{N}$  levels from  $^{15}\text{N}(^3\text{He}, ^3\text{He}')$  and  $^{15}\text{N}(\alpha, \alpha')$

$^{15}\text{N}^* \text{ }^{\text{a}}$ (MeV $\pm$ keV)	$L \text{ }^{\text{a}}$	$^{15}\text{N}^* \text{ }^{\text{b}}$ (MeV)	$L \text{ }^{\text{b}}$
0		0	
$5.28 \pm 30$	3	$5.27 + 5.30$	3
6.32	2	6.32	2
7.15	3	7.16	<sup>c</sup>
7.30	1	7.30	1
7.56	3	7.57	3
8.31	1	8.31	<sup>c</sup>
8.57	1	8.58	<sup>c</sup>
$9.17 \pm 30$		9.16	
$9.79 \pm 40$		9.83	
$10.03 \pm 40$		10.07	
$10.71 \pm 40$			
$11.34 \pm 40$			
$11.92 \pm 40$			
$12.52 \pm 40$			
$14.12 \pm 40$			
$15.11 \pm 40$			

<sup>a</sup> ( $^3\text{He}, ^3\text{He}'$ ): (1969BA06).

<sup>b</sup> ( $\alpha, \alpha'$ ) (1966HA19). The  $E_x$  were determined.

<sup>c</sup> Weakly excited. See also (1965BU05, 1969BA06).

The surface thickness  $a = 0.36$  fm, as determined from analysis of the scattering of 44 MeV  $\alpha$ -particles from  $^{15}\text{N}$  (1968FA1A). At  $E_\alpha = 40.5$  MeV, a number of particle groups have been observed, and angular distributions have been measured: see Table 15.17 (1966HA19). See also (1965BU05),  $B(E2)_\downarrow / e^2 = 4.9 \text{ fm}^4$  for  $^{15}\text{N}^*(6.32)$ :  $B(E2)_\downarrow / e^2 = 60 \text{ fm}^6$  for both  $^{15}\text{N}^*(5.27, 7.57)$  (1966HA19). See also (1969BA06).

$$54. \ ^{15}\text{O}(\beta^+)^{15}\text{N} \quad Q_m = 2.760$$

See  $^{15}\text{O}$ .

$$55. \text{ (a) } ^{16}\text{O}(\gamma, p)^{15}\text{N} \quad Q_m = -12.126$$

$$\text{ (b) } ^{16}\text{O}(e, ep)^{15}\text{N} \quad Q_m = -12.126$$

Over the giant resonance region in  $^{16}\text{O}$ , the decay takes place to the odd parity states of  $^{15}\text{N}$  at  $E_x = 0$  and 6.32 MeV as well as to both of the even parity states at  $E_x = 5.27$  and 5.30 MeV. The branching ratios are functions of the excitation energy in  $^{16}\text{O}$  and of the authors: see (1965DE24, 1965MA45, 1965MO13, 1966KO1G, 1966OW01, 1967CA1C, 1967CA1P, 1968BA2L, 1969MU07, 1969SH02). The results are in fair agreement with the predictions of the single-particle, single-hole theory of photoexcitation of  $^{16}\text{O}$ , although some non-single-particle excitation appears to be necessary in some portions of the  $^{16}\text{O}$  giant resonance: see, e.g. (1968BA2L). High-energy  $\gamma$ -rays have also reported from the decay of  $^{15}\text{N}^*(7.30, 8.31, 9.05, 9.23)$  (1968BA2L, 1969HO1T, 1969MU07). See also the review in (1968SC1B), (1957JO20, 1959BR69, 1963SC32, 1968TU02, 1969FR20, 1969UL01) and (1968ZH1B; theor.). The  $\tau_m$  of  $^{15}\text{N}^*(5.27) \gg 0.1$  psec;  $\tau_m$  for  $^{15}\text{N}^*(6.33, 9.23) < 0.1$  psec (1969MU07): see also Table 15.7.

For reaction (b), see (1962DO1A).

$$56. \ ^{16}\text{O}(n, d)^{15}\text{N} \quad Q_m = -9.901$$

Angular distribution of the deuterons corresponding to the ground state of  $^{15}\text{N}$  have been determined at  $E_n = 14.4$  MeV (1964PA11, 1965VA05) and at 14 MeV (1963GA10). See also (1965DI1E; theor.).

$$57. \ ^{16}\text{O}(p, 2p)^{15}\text{N} \quad Q_m = -12.126$$

At  $E_p = 460$  MeV, the summed proton spectrum shows two peaks corresponding to the knowckout of  $p_{1/2}$  and  $p_{3/2}$  protons with binding energies of 12.4 and 19.0 MeV, respectively [ $^{15}\text{N}^*(0, 6.32)$ ] (1966TY01). See also the discussions in (1963CL1B, 1963RI1B), (1962FO03, 1967FU1A), (1963BE1A, 1963BE42; theor.) and (1959AJ76).

58.  $^{16}\text{O}(\text{d}, ^3\text{He})^{15}\text{N}$   $Q_{\text{m}} = -6.632$

Angular distributions of the  $^3\text{He}$  groups have been measured at  $E_{\text{d}} = 20$  MeV (1969PU04: to  $^{15}\text{N}^*(0, 5.27, 5.30)$ ), 28 MeV (1968GA13: to  $^{15}\text{N}(0)$ ), 34.4 MeV (1967HI06: to  $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32)$ ), 52 MeV (1969KA1A, 1969KA1W: to  $^{15}\text{N}(0)$ ), and 82 MeV (1969DO04: to  $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32)$ ) and analyzed by DWBA. The  $^3\text{He}$  group to  $^{15}\text{N}^*(5.2)$  does not show a stripping pattern (1969PU04). See also (1968BA2J; theor.).

59.  $^{16}\text{O}(\text{t}, \alpha)^{15}\text{N}$   $Q_{\text{m}} = 7.688$

Angular distributions have been measured at  $E_{\text{t}} = 0.9$  to 1.7 MeV (1967KO1G;  $\alpha_0$ ), 1.15 to 1.95 MeV (1959JO32;  $\alpha_0$ ), 1.5 to 3.0 MeV (1966SE1D;  $\alpha_0$ ), and 13 MeV (1965AJ01;  $\alpha_0, \alpha_{1+2}, \alpha_3$ ). A  $\gamma$ -ray with  $E_{\gamma} = 5.272 \pm 0.010$  MeV has been observed in this reaction: see Table 15.7 (1965AL19). See also (1959AJ76).

60.  $^{16}\text{O}(^{14}\text{N}, ^{15}\text{O})^{15}\text{N}$   $Q_{\text{m}} = -4.833$

See (1963TO1D, 1969BR1D).

61.  $^{17}\text{O}(\text{d}, \alpha)^{15}\text{N}$   $Q_{\text{m}} = 9.803$

See (1954PA39).

62.  $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$   $Q_{\text{m}} = 3.981$

Angular distributions of ground state  $\alpha$ -particles are reported at  $E_{\text{p}} = 0.84$  to 2.00 MeV (1961CA02) and at 7.9, 10.6 and 13.3 MeV (1964EC03). Angular correlation measurements lead to  $J = \frac{5}{2}, \frac{3}{2}, (\frac{1}{2}), (\frac{3}{2})$  for  $^{15}\text{N}^*(5.28, 6.32, 8.31, 8.58)$  (1965WA06),  $J = \frac{5}{2}, \frac{5}{2}, \frac{3}{2}, \frac{7}{2}$  for  $^{15}\text{N}^*(5.28, 7.16, 7.30, 7.57)$  (1966HA30),  $J = \frac{3}{2}$  for  $^{15}\text{N}^*(6.32)$  (1966LO02). The M2/E1 mixing ratio of the  $7.30 \rightarrow$  g.s. transition indicates an unusually large retardation of an E1 transition in a non-self-conjugate nucleus (1966HA30).  $J = \frac{5}{2}$  for  $^{15}\text{N}^*(5.27)$  and the mixing parameter fix  $J = \frac{5}{2}$  for  $^{15}\text{N}^*(10.54)$  which fed the first excited state of  $^{15}\text{N}$  (1959HE1D) in the  $^{14}\text{C}(\text{p}, \gamma)^{15}\text{N}$  reaction (1965WA06).

See also (1960CL02, 1961LO10, 1964AM1A, 1964MA25, 1964MA57, 1964SC01), (1959AJ76) and  $^{19}\text{F}$  in (1959AJ76, 1972AJ02).

$$63. {}^{19}\text{F}(\gamma, \alpha){}^{15}\text{N} \quad Q_m = -4.011$$

See (1965HA1G).

$$64. (a) {}^{19}\text{F}(\text{p}, \text{p}\alpha){}^{15}\text{N} \quad Q_m = -4.011$$

$$(b) {}^{19}\text{F}(\alpha, 2\alpha){}^{15}\text{N} \quad Q_m = -4.011$$

For reaction (a) see (1962FO03). For reaction (b) see (1963LA02).

$$65. {}^{19}\text{F}(\text{d}, {}^6\text{Li}){}^{15}\text{N} \quad Q_m = -2.539$$

Angular distributions of the  ${}^6\text{Li}$  ions corresponding to  ${}^{15}\text{N}(0)$  have been measured at  $E_d = 9.0$  to  $12.5$  MeV (1967DE14),  $14.5$  MeV (1964DA1B),  $14.9$  MeV (1966DE09) and  $21$  MeV (1965SL1C). Attempts have been made to fit the data with DWBA: see references above and (1963DR1B). See also (1964BL1C).

$$66. {}^{19}\text{F}({}^{14}\text{N}, {}^{18}\text{F}){}^{15}\text{N} \quad Q_m = 0.405$$

See (1965WI1A).



**<sup>15</sup>O**  
(Figs. 11 and 12)

GENERAL:

*Model calculations:* (1960TA1E, 1960TA1C, 1963CO12, 1963KU1B, 1964AL1L, 1964AM1D, 1964BR1H, 1964RI1A, 1965CO25, 1965GI1B, 1965GR1H, 1965GU1A, 1965HU1D, 1966BO1R, 1966EL08, 1966RI1F, 1966SO05, 1967BO1T, 1967EL03, 1968DE13, 1968EL1A, 1968HO1H, 1968MA2B, 1968SH08, 1968WO1C, 1968ZH1B, 1968ZU02, 1969DE16, 1969EL1B, 1969GU1M, 1969SA1J).

*General calculations and reviews:* (1964EV1A, 1967FA1A, 1967NE1D, 1968BI1C).

*Electromagnetic transitions:* (1965RO1N, 1966PO11, 1966RO1P, 1966WA1E, 1967KU1E, 1967PO1J, 1967WA1C, 1968BI1C, 1968SH08, 1968ZH1B, 1968ZH06, 1969KH1C, 1969ZH1A).

*Other:* (1966WA1K, 1967AU1B, 1969FO1D, 1969HA1G).

*Ground state:*  $J = \frac{1}{2}$  (1963CO17);  $\mu = 0.71898$  nm (1963CO17, 1964LI14, 1967CO1D).

See also (1964ST1B, 1965MA1T, 1966MA1V, 1967NE1D, 1967SH14, 1968RO1E, 1969FU11, 1969PE1D).

1.  $^{15}\text{O}(\beta^+)^{15}\text{N}$   $Q_m = 2.760$

Reported half-lives are listed in Table 15.19 (1954KL36, 1955BA83, 1957KI22, 1957PE12, 1959KI99, 1960JA12): the weighted mean is  $122.24 \pm 0.16$  sec. See also (1963CS02, 1963VA31). Using this value for  $\tau_{1/2}$  and  $Q_m$ ,  $\log ft = 3.643$ . See also (1968BA42) and (1965GA1D, 1966MI1F, 1967AM1H, 1968SH08, 1969LE1D, 1969SU15; theor.).

2.  $^7\text{Li}(^{14}\text{N}, ^6\text{He})^{15}\text{O}$   $Q_m = -2.687$

See (1958AL1D).

3. (a)  $^{10}\text{B}(^6\text{Li}, \text{n})^{15}\text{O}$   $Q_m = 15.209$   
(b)  $^{10}\text{B}(^7\text{Li}, 2\text{n})^{15}\text{O}$   $Q_m = 7.957$

See (1957NO17).

Table 15.18: Energy levels of  $^{15}\text{O}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$J^\pi$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions
0	$\frac{1}{2}^-$	$\tau_{1/2} = 122.24 \pm 0.16$ sec	$\beta^+$	1, 2, 3, 4, 6, 7, 8, 9, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29
5.181 $\pm$ 5	$\frac{1}{2}^+$	$\tau_m < 0.1$ psec	$\gamma$	8, 9, 16, 17, 22, 24, 25, 26
5.24151 $\pm$ 0.52	$\frac{5}{2}^+$	3.2 $\pm$ 0.5	$\gamma$	8, 9, 16, 17, 20, 21, 22, 24, 25, 26
6.176 $\pm$ 3	$\frac{3}{2}^-$	$< 0.047$	$\gamma$	8, 9, 16, 17, 21, 22, 24, 25, 26
6.788 $\pm$ 4	$\frac{3}{2}^+$	$< 0.028$	$\gamma$	8, 9, 16, 17, 26
6.859 $\pm$ 1	$\frac{5}{2}^+$	0.10 $\pm$ 0.06	$\gamma$	8, 9, 16, 17, 21, 26
7.2760 $\pm$ 0.6	$\frac{7}{2}^+$		$\gamma$	8, 16, 17, 21, 24, 26
7.5522 $\pm$ 0.5	$\frac{1}{2}^+$	$\Gamma = 1.7 \pm 0.5$ keV	$\gamma, p$	9, 16, 17, 21
8.2833 $\pm$ 1.5	$\frac{3}{2}^+$	3.6 $\pm$ 0.7	$\gamma, p$	9, 17, 21
8.739 $\pm$ 6	$\frac{1}{2}^+$	32	$\gamma, p$	9
8.9180 $\pm$ 1.4	$\frac{3}{2}$	3.7 $\pm$ 1	$\gamma, p$	8, 9, 21
8.9781 $\pm$ 1.6	$(\frac{1}{2}, \frac{3}{2})^-$	3.9 $\pm$ 0.4	$\gamma, p$	8, 9, 21
9.483 $\pm$ 3	$\frac{5}{2}^-$	10.1 $\pm$ 0.5	$\gamma, p$	8, 9, 21
9.50 $\pm$ 40	$\frac{3}{2}^+(\frac{1}{2})^+$	280 $\pm$ 24	$\gamma, p$	9
9.606 $\pm$ 1.8	$\frac{3}{2}^-$	8.8 $\pm$ 0.5	$\gamma, p$	8, 9, 21, 24
9.660 $\pm$ 4	$\frac{1}{2}^-$	2 $\pm$ 1	p	8, 10, 21
9.72 $\pm$ 50	$(\frac{1}{2}, \frac{3}{2})^+$	1190 $\pm$ 50	$\gamma, p$	9
10.278 $\pm$ 8	+	16 $\pm$ 4	p	10, 21
10.46 $\pm$ 10		47	$\gamma, p$	9, 10, 21, 24
10.91 $\pm$ 15	$\frac{7}{2}^+$	91	p	10, 21
10.939 $\pm$ 7	$\frac{1}{2}^+$	84	$\gamma, p$	9, 10, 21
11.023 $\pm$ 7	$\frac{1}{2}^-$	21	p	10
11.15 $\pm$ 15		$< 10$	p	10
11.20 $\pm$ 15	$(\frac{1}{2}, \frac{3}{2})^+$	36	$\gamma, p$	9, 10, 21
11.5 $\pm$ 100	$T = \frac{3}{2}$			8
11.56 $\pm$ 15		$< 10$	p	10
11.57		140	$\gamma, p$	10
11.57 $\pm$ 15	$\frac{5}{2}^-$	25	p	9

Table 15.18: Energy levels of  $^{15}\text{O}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions
11.61 $\pm$ 15	$\frac{3}{2}^- (\frac{1}{2}^-)$	25	p	10
11.71 $\pm$ 15		< 10	p	10, 21
11.75 $\pm$ 15	$\frac{5}{2}^+$	80	p	10
11.846 $\pm$ 10	$\frac{5}{2}^-$	50	p	10
11.98 $\pm$ 15	$\frac{5}{2}^-$	30	p	10
12.12 $\pm$ 15	$\frac{5}{2}^+$	160	p	10
12.47 $\pm$ 15	$\frac{5}{2}^- (\frac{3}{2}^-)$	60	p	10
12.8		$\approx$ 230	$\gamma$ , p	9
12.82	+	9	p	10
13.00	$\frac{5}{2}$	28	p, $^3\text{He}$ , $\alpha$	5, 10, 15
13.1			p, d, $^3\text{He}$ , $\alpha$	5
13.4	$\frac{3}{2}^+$	broad	(p, $\alpha$ )	10, 15
13.49	$(\frac{3}{2}^+)$		p	10
13.60	$\frac{5}{2}^+$		p, $\alpha$	15
13.70	$\frac{3}{2}^-$		p	10
13.79	$\frac{3}{2}^-$		n, p, $^3\text{He}$ , $\alpha$	5, 10, 15, 21
13.87		$\approx$ 140	$\gamma$ , p	9
14.03 $\pm$ 40	$(\frac{1}{2}, \frac{3}{2})^-$	160 $\pm$ 20	n, p, $^3\text{He}$	5
14.17	$\frac{5}{2}^-$		p, $\alpha$	15
14.27 $\pm$ 40	$\frac{1}{2}^+$	340 $\pm$ 30	n, p, $^3\text{He}$ , $\alpha$	5, 10, 11, 15
14.34	$\frac{5}{2}^+$		p, $\alpha$	15
14.460 $\pm$ 10	$\frac{5}{2}^+$	100 $\pm$ 10	n, p, $^3\text{He}$ , $\alpha$	5, 11
14.69 $\pm$ 40		170 $\pm$ 30	n, p, $^3\text{He}$	5, 11
14.95 $\pm$ 40		400 $\pm$ 25	n, p, $^3\text{He}$	5, 11
15.43 $\pm$ 10		65 $\pm$ 15	p, $^3\text{He}$ , $\alpha$	5
15.56 $\pm$ 40	$\frac{1}{2}^+$	80 $\pm$ 25	p, $^3\text{He}$ , $\alpha$	5
15.84 $\pm$ 50	$(\frac{1}{2}, \frac{3}{2})^-$	350	n, p, $^3\text{He}$ , $\alpha$	5, 11
16.04			$^3\text{He}$ , $\alpha$	5
16.09			n, $^3\text{He}$ , $\alpha$	5
16.19			$^3\text{He}$ , $\alpha$	5
16.43 $\pm$ 50	$\frac{1}{2}^+$	170	$^3\text{He}$ , $\alpha$	5


Table 15.18: Energy levels of  $^{15}\text{O}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi$	$\tau_m$ (psec) or $\Gamma$ (keV)	Decay	Reactions
16.48 $\pm$ 50		560 $\pm$ 90	n, p	11
16.77 $\pm$ 50		200	n, $^3\text{He}$ , $\alpha$	5
17.50 $\pm$ 50	$(\frac{1}{2}, \frac{3}{2})^-$	600	n, p, $^3\text{He}$ , $\alpha$	5
17.99 $\pm$ 50	$(\frac{1}{2}, \frac{3}{2})^-$	200	$^3\text{He}$ , $\alpha$	5
18.22 $\pm$ 50			n, p, $^3\text{He}$	5
19.02 $\pm$ 50			n, $^3\text{He}$	5
19.90 $\pm$ 50			n, p, $^3\text{He}$	5
$\approx$ 21		broad	$\gamma$ , p	9

<sup>a</sup> See also Tables 15.22 and 15.25.



See (1963HO1E).



Excitation functions for these reactions have been measured over a wide range of energies: see Table 15.20. Observed resonances are displayed in Table 15.21. For discussions of angular distribution measurements which have been measured at many energies over the range displayed in Table 15.20, see the writeups of  $^7\text{Be}$  in (1966LA04),  $^{11}\text{C}$  and  $^{12}\text{C}$  in (1968AJ02), and  $^{13}\text{N}$ ,  $^{14}\text{N}$ , and  $^{14}\text{O}$ . See also (1964DE1E).

Consideration of the relative yields at  $E(^3\text{He}) = 1.21$  and  $2.15$  MeV lead to the tentative  $J^\pi$  assignments given in Table 15.21 for  $^{15}\text{O}^*(13.00, 13.79)$  (1957BR18). The assignments for  $^{15}\text{O}^*(14.03, 14.27, 14.46)$  are derived from analyses of the  $^3\text{He}$  elastic scattering, particle and  $\gamma_{6.44}$  angular distributions, and the total cross sections of the  $n_0$  and  $p_1$  groups (1964KU05). Above  $E(^3\text{He}) \approx 5$  MeV, the excitation functions show broad maxima (see Table 15.21) but it is not

Table 15.19: The half-life of  $^{15}\text{O}$

$\tau_{1/2}$ (sec)	Refs.
$123.4 \pm 1.3$	(1954KL36)
$121 \pm 3$	(1955BA83)
$120 \pm 2$	(1957KI22)
$123.95 \pm 0.5$	(1957PE12)
$124.1 \pm 0.5$	(1959KI99)
$122.1 \pm 0.1$	(1960JA12)
$122.24 \pm 0.16$	weighted average

clear that they correspond to excited states in  $^{15}\text{O}$ : for instance the maxima in the yields of  $n_0$  (1964DE1C),  $p_6$ ,  $d_0$ ,  $\alpha_0$  and  $\alpha_1$  (1960HI07) are not correlated. See, however, (1969WE08). The structures observed in the excitation functions for  $p_0 \rightarrow p_9$  at  $E(^3\text{He}) = 4.6$  to 11 MeV are attributed to quasi-giant resonances involving single proton orbitals coupled to excited  $^{14}\text{N}$  core configurations (1969HA49). Above  $E(^3\text{He}) \approx 12$  MeV, the excitation functions do not show clear resonant behavior: see, e.g. (1967GR1L, 1968FO06, 1969FO02).

Broad resonance-like structures in the yield of protons are reported at  $E(^3\text{He}) = 7.0, 7.8, 9.6, 10.6, 13.4$  and  $14.6$  MeV [ $\Gamma \approx 1$  MeV] (1970SP1E).

Over the range  $E(^3\text{He}) = 6.5 \rightarrow 11$  MeV, the ratio  $d\sigma_n/d\sigma_{p_1}$  [to  $^{14}\text{O}_{\text{g.s.}}$  and the first  $T = 1$  state of  $^{14}\text{N}$ ], determined at  $10^\circ$ , is approximately constant and  $\approx 2$ , as would be expected if isospin were conserved and if Coulomb scattering and  $Q$ -value differences can be neglected (1965FU16). (1969LA1A) find that the compound nucleus level overlap parameter  $\Gamma/D_0$  and the spin cut-off parameter  $\sigma$  as obtained from elastic scattering and from the statistical model analysis of three proton groups [ $E(^3\text{He}) = 5.29 - 5.50$  MeV] are constant. See also (1970CA1G).

For a survey of the energies at which polarization measurements have been made, see Table 15.20. (1968HU1A) find that at  $E(^3\text{He}) = 36$  and  $42$  MeV, the  $^3\text{He}$  polarization results are such that optical model calculations require an optical spin-orbit potential depth  $\leq 3.5$  MeV. See also (1959AJ76).

$$6. \ ^{12}\text{C}(\alpha, n)^{15}\text{O} \quad Q_{\text{m}} = -8.507$$

$$E_{\text{thresh.}} = 11.341 \pm 0.015 \text{ (1963NE05).}$$

Angular distributions of neutrons corresponding to the ground state of  $^{15}\text{O}$  have been measured at  $E_\alpha = 14$  MeV (1965AL1J) and at  $20.0$  to  $21.8$  MeV (1963KO03). See also  $^{16}\text{O}$  in (1971AJ02).

Table 15.20: Recent yield and polarization measurements in  $^{12}\text{C} + ^3\text{He}$  <sup>a</sup>

 (a) *Excitation functions* <sup>b</sup>

$E(^3\text{He})$	Yield of	Refs.
1.0 – 1.8	$p_0, p_1, p_2, d_0, \alpha_0$	(1965GR1R)
1.4 – 3.5	n	(1964KU06)
1.4 – 6.0	$n, \alpha$	(1966CI01)
1.4 – 11.45	$n_0$	(1964OS01)
1.75 – 5.2	$\alpha_0$	(1963LU05)
2 – 5.5	$^{11}\text{C}$	(1966GO1E)
1.8 – 5.4	$p_0 \rightarrow p_9, ^3\text{He}, \alpha_0$	(1964KU05)
1.8 – 5.5	n	(1964DI1D)
2 – 4	$p_0 \rightarrow p_8$	(1967CL1C)
2.0 – 5.7	$n_0$	(1961TO03)
2 – 6.2	$^3\text{He}, \alpha_0, \alpha_1$	(1966BL01, 1966SC12)
2.4 – 3.3	$^3\text{He}, \alpha_0$	(1970JA1E)
$\approx 2.5$	$p_2$	(1963LU01, 1963LU1F)
2.7 – 5.4	$\gamma_{6.44}$	(1964KU05)
3.50 – 3.91	$^3\text{He}$	(1967SC27)
4.0 – 5.4	$\alpha_1$	(1964KU05)
4.0 – 6.0	$p_2 \rightarrow p_6$	(1966BL01)
4.4 – 8.2	$^3\text{He}, \alpha_0$	(1968WE15, 1969WE03)
4.5 – 6.0	$\alpha_0$	(1969WE03)
4.6 – 11.0	$p_0 \rightarrow p_9$	(1969HA49)
5 – 30	$^{11}\text{C}, ^{13}\text{N}$	(1965BR1B)
5.2 – 7.8	$\alpha_1$	(1968WE15, 1969WE03)
5.5 – 11	$n_0$	(1964DE1C)
5.7 – 10.23	$p_6, d_0, \alpha_0, \alpha_1$	(1960HI07)
6 – 17	p	(1970SO1G)
6 – 24	$^7\text{Be}, ^{11}\text{C}, ^{13}\text{N}$	(1962CO1C)
8 – 10	$^3\text{He}, \alpha_0$	(1966SC22)
9 – 29	$^7\text{Be}$	(1965EN01)
12.0 – 18.6	$\alpha_0 \rightarrow \alpha_7$	(1967GR1L)
12 – 18.6	$d_0, d_1, d_{2+3}, ^3\text{He}(0), ^3\text{He}(1)$	(1968FO06, 1969FO02)

Table 15.20: Recent yield and polarization measurements in  $^{12}\text{C} + ^3\text{He}$  <sup>a</sup>  
(continued)

$E(^3\text{He})$	Yield of	Refs.
21 – 31.2	$^{11}\text{C}, ^{13}\text{N}$	(1959MA1D)
21 – 31	$^7\text{Be}$	(1965MA1V)

(b) Polarization measurements <sup>b</sup>

2.24 – 3.70	$n_0$	(1967SC27)
2.48 – 3.15	$p_0$	(1967KR1F)
4.1 – 5.9	$n$	(1967SO1B)
4.5 – 5.5	$p_0, p_1$	(1967MA1M)
18	$^3\text{He}$	(1969LU1C)
29	$^3\text{He}$	(1961AG1A, 1964BU1D, 1965FR1E)
31.6	$^3\text{He}$	(1969EN03)
36, 42	$^3\text{He}$	(1968HU1A)

<sup>a</sup> See also (1959AJ76, 1961FO02).

<sup>b</sup> See also (1964BR1G, 1964DI1C, 1965SC1D, 1966HA21, 1966MA1R, 1968PA1F, 1969PA1C).

<sup>c</sup> See also (1965PE1H, 1966CA1H).

7. (a)  $^{12}\text{C}(^6\text{Li}, t)^{15}\text{O}$   $Q_m = -3.721$   
 (b)  $^{12}\text{C}(^{14}\text{N}, ^{11}\text{B})^{15}\text{O}$   $Q_m = -8.664$

For reaction (a) see (1969GI1B); for reaction (b) see (1969BR1G).

8.  $^{13}\text{C}(^3\text{He}, n)^{15}\text{O}$   $Q_m = 7.125$

Angular distributions of neutrons corresponding to the ground state of  $^{15}\text{O}$  have been measured at  $E(^3\text{He}) = 1.70$  to  $5.35$  MeV (1965DI07),  $2.6$ ,  $2.8$  and  $3.1$  MeV (1961JO07, 1961JO24),  $2.66$  MeV (1961DU1B, 1963DU12: also  $n_3$ ),  $5.0$  and  $6.2$  MeV (1969HO1F: also to  $^{15}\text{O}^*(6.18, 6.86, 8.92, 8.98, 9.50, 9.60, 9.66)$  and  $7.8, 8.6$  and  $10.1$  MeV (1964DE1C). DWBA analyses have been made: (1969HO1F) find  $L = 0$  for  $^{15}\text{O}^*(0, 8.92, 8.98, 9.66)$  and  $L = 2$  for  $^{15}\text{O}^*(6.18, 9.49, 9.60)$ . At  $E(^3\text{He}) = 11.6$  MeV, a neutron group assigned to a  $T = \frac{3}{2}$  state at  $11.5 \pm 0.10$  is reported by (1969BR30).

Table 15.21: Resonances in  $^{12}\text{C} + ^3\text{He}$ 

$E(^3\text{He})$ (MeV $\pm$ keV)	Resonant for	$E_x^a$ (MeV)	$J^\pi$	$\Gamma_{\text{c.m.}}$ (keV)	Refs.
1.21	$p_0, p_2$	13.04	$(\frac{5}{2})^-$		(1957BR18)
1.3	$p_0, p_1, p_2, p_3, d_0, \alpha_0$	12.1			(1957BR18, 1965GR1R)
2.15	$n, p_0$	13.79	$(> \frac{5}{2})$		(1957BR18)
$2.45 \pm 40$	$n_0, p_0, p_1, p_2, p_3$	14.03	$(\frac{1}{2}^-, \frac{3}{2}^-)$	$160 \pm 20$	(1957BR18, 1961TO03, 1964KU05, 1964KU06, 1964OS01)
$2.75 \pm 40$	$n_0, p_1, p_2, ^3\text{He}, \alpha$	14.27	$\frac{1}{2}^+$	$340 \pm 30$	(1957BR18, 1964KU05, 1964KU06, 1964OS01, 1966CI01, 1970JA1E)
$2.990 \pm 10$	$n_0, p_0, p_1, p_2, p_4, p_5, p_8, ^3\text{He}, \alpha_0$	14.460	$\frac{3}{2}^+$	$100 \pm 10$	(1958JO20, 1961TO03, 1964KU05, 1964KU06, 1964OS01, 1970JA1E)
$3.28 \pm 40$	$p_0$	14.69		$180 \pm 40$	(1964KU05)
$3.60 \pm 40$	$p_0, p_1, p_2$	14.95		$400 \pm 25$	(1958JO20, 1964KU05)
$4.20 \pm 10$	$p_5, p_6, \alpha_0$	15.43		$65 \pm 15$	(1964KU05)
$4.37 \pm 40$	$p_0, p_1, p_2, p_4, p_7, p_8, \alpha_0$	15.56	$\frac{1}{2}^+$	$80 \pm 25$	(1958JO20, 1961TO03, 1964KU05, 1966CI01)
$4.65 \pm 50$	$n_0$	15.79			(1961TO03, 1964DI1D, 1964KU05, 1964OS01)
4.77	$\alpha_0$	15.89	$\frac{1}{2}^-, \frac{3}{2}^-$	350	(1969WE03)
$4.97 \pm 20$	$\alpha_0$	16.04			(1969WE08)
$5.03 \pm 20$	$\alpha_0$	16.09			{ (1961TO03, 1964DI1D, 1964KU05, 1964OS01, 1966SC12, 1969WE08)
$5.15 \pm 20$	$\alpha_0$	16.19			
5.45	$\alpha_0$	16.43	$\frac{1}{2}^+$	170	(1969WE03)
$5.88 \pm 50$	$n_0, ^3\text{He}, \alpha_0$	16.77	$\frac{1}{2}^-, \frac{3}{2}^-$	200	(1964OS01, 1966SC12, 1969WE03)
$6.80 \pm 50$	$n_0, p$	17.50	$\frac{1}{2}^-, \frac{3}{2}^-$	600	(1964OS01, 1969WE03, 1970SO1G)
7.4	$\alpha_0$	17.99	$\frac{1}{2}^-, \frac{3}{2}^-$	200	(1969WE03)
$7.70 \pm 50$	$n_0, p$	18.22			(1964OS01, 1970SO1G)
$8.70 \pm 50$	$n_0$	19.02			(1964OS01)
$9.80 \pm 50$	$n_0, p$	19.90			(1964OS01, 1970SO1G)

<sup>a</sup> See also text.



Table 15.22: Radiative decays in  $^{15}\text{O}$ 

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	$\Gamma_\gamma$ <sup>a</sup> (eV)	Refs.
6.18	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100		(1965WA16)
		5.18	$\frac{1}{2}^+$	< 2.5		(1965WA16)
		5.24	$\frac{5}{2}^+$	< 2.5		(1965WA16)
6.79	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	100		(1965WA16, 1968GI11)
		5.18	$\frac{1}{2}^+$	< 6		(1965WA16)
		5.24	$\frac{5}{2}^+$	< 3		(1968GI11)
6.86	$\frac{5}{2}^+$	5.24	$\frac{5}{2}^+$	< 6		(1965WA16)
				< 3		(1968GI11)
		6.18	$\frac{3}{2}^-$	< 7		(1965WA16)
		0	$\frac{1}{2}^-$	< 10		(1965WA16)
		5.18	$\frac{1}{2}^+$	< 12		(1968GI11)
7.28	$\frac{7}{2}^+$	5.18	$\frac{1}{2}^+$	< 15		(1965WA16)
				< 4		(1968GI11)
		5.24	$\frac{5}{2}^+$	100		(1965WA16, 1968GI11)
		6.18	$\frac{3}{2}^-$	< 0.4		(1965WA16)
		0	$\frac{1}{2}^-$	< 30		(1965WA16)
7.55	$\frac{1}{2}^+$			< 12		(1968GI11)
		5.18	$\frac{1}{2}^+$	$3.8 \pm 1.2$		(1969KU01)
		5.24	$\frac{5}{2}^+$	< 10		(1965WA16)
				< 4		(1968GI11)
		5.24	$\frac{5}{2}^+$	100		(1965WA16, 1968GI11)
7.55	$\frac{1}{2}^+$			$96.2 \pm 1.2$		(1969KU01)
		6.18	$\frac{3}{2}^-$	< 2		(1965WA16)
		0	$\frac{1}{2}^-$	$\approx 3$		(1960TA17)
				$3.5 \pm 0.5$		(1963HE11)
		5.18	$\frac{1}{2}^+$	$16.2 \pm 2$		(1960TA17)
				$15.8 \pm 0.6$		(1963HE11)
		6.18	$\frac{3}{2}^-$	$57.9 \pm 0.6$		(1960TA17)
		$57.4 \pm 0.6$		(1963HE11)		
6.79	$\frac{3}{2}^+$	6.79	$\frac{3}{2}^+$	$22.9 \pm 2$		(1960TA17)
				$23.3 \pm 0.6$		(1963HE11)

Table 15.22: Radiative decays in  $^{15}\text{O}$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	$\Gamma_\gamma$ <sup>a</sup> (eV)	Refs.
8.28	$\frac{3}{2}^+$	6.86	$\frac{5}{2}^+$	b		
		0	$\frac{1}{2}^-$	$53.8 \pm 0.25$	0.531	(1966EV01)
		5.24	$\frac{5}{2}^+$	$42.7 \pm 0.5$	0.405	(1966EV01)
		6.18	$\frac{3}{2}^-$	$2.2 \pm 0.6$	0.021	(1966EV01)
8.74 <sup>c</sup>	$\frac{1}{2}^+$	6.86	$\frac{5}{2}^+$	$1.2 \pm 0.3$	0.011	(1966EV01)
		5.18	$\frac{1}{2}^+$	67	0.32	(1966EV01)
		6.18	$\frac{3}{2}^-$	33	0.16	(1966EV01)
8.92	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	$21 \pm 2$	0.056	(1966EV01)
		5.18	$\frac{1}{2}^+$	$23 \pm 6$	0.094	(1966EV01)
		6.18	$\frac{3}{2}^-$	$30 \pm 3$	0.094	(1966EV01)
		6.86	$\frac{5}{2}^+$	$26 \pm 3$	0.069	(1966EV01)
8.98	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	93.4	0.74	(1966EV01)
		5.18	$\frac{1}{2}^+$	5.9	0.046	(1966EV01)
9.49	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	86	2.1	(1967EV02)
		5.24	$\frac{5}{2}^+$	6.5	0.15	(1967EV02)
		6.18	$\frac{3}{2}^-$	0.7	0.22	(1967EV02)
		6.86	$\frac{5}{2}^+$	3.4	0.08	(1967EV02)
		7.28	$\frac{7}{2}^+$	5.1	0.11	(1967EV02)
		9.50	$\frac{3}{2}^+(\frac{1}{2}^+)$	0	$\frac{1}{2}^-$	$\approx 100$
9.60	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	79	4.0	(1967EV02)
		5.24	$\frac{5}{2}^+$	19	1.0	(1967EV02)
		6.18	$\frac{3}{2}^-$	2	0.1	(1967EV02)

<sup>a</sup> (1951DU08, 1966EV01). See also (1959HE47).

<sup>b</sup> Intensity < 25% of transition to  $^{15}\text{O}^*(6.79)$  (1959PO79).

<sup>c</sup> See also (1959HE47).

Branching ratios measured by (1965WA16) are listed in Table 15.22. The measured  $E_\gamma$  lead to  $E_x = 6.180 \pm 0.004$ ,  $6.857 \pm 0.0032$  and  $7.284 \pm 0.007$  MeV (1965WA16).

See also (1964BR13, 1968HIIJ, 1968ST19), (1965SH1E, 1966SH1F; theor.) and  $^{16}\text{O}$  in (1971AJ02).

$$9. \ ^{14}\text{N}(p, \gamma)^{15}\text{O}$$

$$Q_m = 7.293$$

Table 15.23: Resonances in  $^{14}\text{N} + \text{p}$ 

$E_p$ (keV)	$\Gamma_{\text{lab}}$ (keV)	$\omega\Gamma_\gamma$ (eV)	Particles out	$J^\pi$	$E_x$ (MeV)	Refs.
$278.1 \pm 0.4$	$1.7 \pm 0.5$	0.014	$\gamma$	$\frac{1}{2}^+$	7.5522	(1951DU08, 1955BA83, 1957PI1A, 1959PO79, 1960TA17, 1963HE11)
$1061.6 \pm 1.4$	$3.9 \pm 0.7$	0.95	$\gamma$	$\frac{3}{2}^+$	8.2833	(1951DU08, 1956TA16, 1957HA03, 1959GA05, 1959HE47, 1959VA04, 1959VA08, 1963HE11, 1966EV01)
$1550 \pm 6$	34	0.16	$\gamma$	$\frac{1}{2}^+$	8.739	(1951DU08, 1956TA16, 1957BO58, 1957HA03, 1966EV01)
$1742.0 \pm 1.2^a$	$4 \pm 1$	0.21	$\gamma$	$\frac{3}{2}^+$	8.918	(1951DU08, 1956TA16, 1957BO58, 1957HA03, 1959BA16, 1959VA08, 1963CO13, 1966EV01)
$1806.4 \pm 1.5^a$	$4.2 \pm 0.4$	0.52	$\gamma$	$(\frac{1}{2}, \frac{3}{2})^-$	8.9781	(1951DU08, 1956TA16, 1957BO58, 1957HA03, 1959VA08, 1963CO13, 1966EV01)
$2348 \pm 3$	$10.8 \pm 0.5$	2.4	$\gamma$	$\frac{3}{2}^+$	9.483	(1951DU08, 1957BO58, 1959VA08, 1967EV02, 1967LA05, 1967LA10, 1969OC1B)
$2368 \pm 32$	$300 \pm 26$		$\gamma$	$\frac{3}{2}^+$ ( $\frac{1}{2}^+$ )	9.50	(1957BO58, 1959VA08, 1967EV02, 1967LA05, 1967LA10)
$2479 \pm 1.7$	$9.4 \pm 0.5$	3.3	$\gamma$	$\frac{3}{2}^+$	9.606	(1951DU08, 1959VA08, 1967EV02, 1967LA05, 1967LA10, 1969OC1B)
$2537 \pm 4$	$2 \pm 1$		p	$(\frac{7}{2}, \frac{9}{2})^-$	9.660	(1967LA05, 1967LA10)
$2600 \pm 50$	$1270 \pm 50$	46	$\gamma$	$(\frac{1}{2}, \frac{3}{2})^+$	9.72	(1951DU08)
$3200 \pm 8$	$17 \pm 4$		p	+	10.278	(1957BO58, 1959VA08, 1967KU1M)
$3390 \pm 10$	50		$\gamma, \text{p}$		10.46	(1957BO58, 1959VA08, 1969OC1B)
$3880 \pm 15$	97		$\text{p}_0$	$\frac{7}{2}^+$	10.91	(1959BA16, 1967KU1M, 1969WE02)
$3908 \pm 7$	90		$\gamma, \text{p}_0, \text{p}_1$	$\frac{1}{2}^+$	10.939	(1956BA34, 1969OC1B, 1969WE02)
$3998 \pm 7$	22		$\text{p}_0, \text{p}_1$	$\frac{1}{2}^-$	11.023	(1956BA34, 1969WE02)
$4130 \pm 15$	$< 10$		$\text{p}_0$		11.15	(1969WE02)
$4190 \pm 15$	39		$\gamma, \text{p}_0$	$(\frac{1}{2}, \frac{3}{2})^+$	11.20	(1969OC1B, 1969WE02)
$4575 \pm 15$	$< 10$		$\text{p}_0$		11.561	(1969WE02)
$4580 \pm 15$	27		$\text{p}_0$	$\frac{5}{2}^-$	11.57	(1969WE02)
4580	150		$\gamma$		11.57	(1969OC1B)
$4630 \pm 15$	27		$\text{p}_0$	$\frac{3}{2}^- (\frac{1}{2}^-)$	11.61	(1969WE02)
$4740 \pm 15$	$< 10$		$\text{p}_0$		11.71	(1969WE02)
$4780 \pm 15$	85		$\text{p}_0, \text{p}_1$	$\frac{5}{2}^+$	11.75	(1956BA34, 1969WE02)
$4881 \pm 10$	54		$\text{p}_0, \text{p}_1$	$\frac{5}{2}^-$	11.846	(1956BA34, 1969WE02)
$5020 \pm 15$	32		$\text{p}_0$	$\frac{5}{2}^-$	11.98	(1969WE02)
$5180 \pm 15$	172		$\text{p}_0, \text{p}_1$	$\frac{5}{2}^+$	12.12	(1969WE02)
$5550 \pm 15$	64		$\text{p}_0, \text{p}_1, \text{p}_2$	$\frac{5}{2}^- (\frac{3}{2}^-)$	12.47	(1969WE02)
5900	$\approx 250$		$\gamma$		12.8	(1969OC1B)
5920	10		p	+	12.82	(1967KU1M)
6100	30		$\text{p}_0 \rightarrow \text{p}_2, \alpha_0$	$\frac{5}{2}^+$	12.9	(1967KU1M, 1968SH11)
6600	broad		$(\text{p}_2, \alpha_0)$	$\frac{5}{2}^+$	13.45	(1968SH11, 1969OC1B)
6640			$(\text{p}_0), \text{p}_2$	$(\frac{5}{2}, \frac{3}{2})^+$	13.49	(1968SH11)
6760			$\alpha_0$	$\frac{5}{2}^+$	13.60	(1968SH11)
6870			$\text{p}_2$	$\frac{5}{2}^-$	13.70	(1968SH11)

Table 15.23: Resonances in  $^{14}\text{N} + \text{p}$  (continued)

$E_{\text{p}}$ (keV)	$\Gamma_{\text{lab}}$ (keV)	$\omega\Gamma_{\gamma}$ (eV)	Particles out	$J^{\pi}$	$E_{\text{x}}$ (MeV)	Refs.
6960			p1, p2, p4, $\alpha_0$	$\frac{3}{2}^{-}$	13.79	(1968SH11)
7050	$\approx 150$		$\gamma$		13.87	(1969OC1B)
7370			$\alpha_0$	$\frac{5}{2}^{-}$	14.17	(1968SH11)
7500	$\approx 800$		n, p2, $\alpha_0$		14.3	(1964KU06, 1968SH11)
7550			$\alpha_0$	$\frac{5}{2}^{+}$	14.34	(1968SH11)
7700			n		14.5	(1964KU06)
7950	$170 \pm 50$		n		14.71	(1964KU06)
8200			n		14.94	(1964KU06)
9050			n		15.73	(1964KU06)
9400			n		16.1	(1964KU06)
$9850 \pm 50$	$600 \pm 100$		n		16.48	(1964KU06)


<sup>a</sup> Comparison of analog states in  $^{15}\text{N}$  and  $^{15}\text{O}$  shows that there should be a (as yet unreported) state in  $^{15}\text{O}$  at  $E_{\text{x}} \approx 8.95$  MeV. This corresponds to  $E_{\text{p}} \approx 1.75$  MeV, and the analysis of the  $E_{\text{p}} = 1.74$  and 1.81 MeV resonances is therefore possible in error (E.K. Warburton, private communication).

Observed resonances are listed in Table 15.23 (1951DU08, 1955BA83, 1957PI1A, 1959GA05, 1959HE47, 1959PO79, 1959VA04, 1959VA08, 1960TA17, 1963HE11, 1966EV01, 1967EV02, 1969OC1B). Branching ratios are displayed in Table 15.22 (1960TA17, 1963HE11, 1966EV01).

The cross section increases from  $(8.5 \pm 3.7) \times 10^{-12}$  b at 100 keV to  $(140 \pm 30) \times 10^{-12}$  b at 135 keV (1957LA13). Extrapolation from the  $E_p = 0.28$  MeV resonance gives  $S(0) = 2.75 \pm 0.50$  keV · b, with zero slope to  $E_p = 0.05$  MeV (1963HE11). For astrophysical implications, see (1964FO1A, CA65, 1968DU1F). The  $J^\pi$  assignments shown in Table 15.23 arise from considerations of branching ratios, measurements of angular distributions of  $\gamma$ -rays and the studies discussed in reaction 10. Angular correlation measurements also lead to  $J^\pi = \frac{3}{2}^+$  and  $\frac{3}{2}^-$  for  $^{15}\text{O}^*(6.79, 6.18)$ . The results are also consistent with  $J = \frac{1}{2}$  for  $^{15}\text{O}^*(5.18)$  (1959PO79). Energies for states involved in the cascade decays are  $E_x = 5.19 \pm 0.01, 6.15 \pm 0.03$  and  $6.79 \pm 0.01$  MeV (1959PO79),  $5.168 \pm 0.015, 6.154 \pm 0.010, 6.788 \pm 0.008$  MeV (1960TA17). At  $E_p = 0.8$  MeV, the non-resonant radiation goes primarily [(81 ± 3%)] via cascades through  $^{15}\text{O}^*(5.2, 6.18, 6.79)$  (1963BA1P). See also (1959HE47).

The excitation function for  $\gamma_0$  for  $E_p = 6.4$  to 19 MeV is characterized by four very pronounced peaks,  $\approx 1$  MeV, wide, below  $E_x = 19$  MeV. The  $90^\circ$  excitation function then shows a giant resonance peak centered at  $E_x \approx 21$  MeV (1968KU1F): see, however, (1959CO1C, 1961CO02). See also (1964TA05).

See also (1966ED1A) and (1969ZH1A; theor.).



The yields of elastic and inelastic protons, and of  $\gamma$ -rays have been studied at many energies: see (1959AJ76) and Table 15.24. The scattering anomalies are superposed on a background which decreases less rapidly than the Rutherford cross section; for  $E_p < 2.3$  MeV, the background is largely s-wave with some p-wave contribution above  $E_p = 1.5$  MeV.

Observed resonances are displayed in Table 15.23: see (1959AJ76) and (1959BA16, 1959VA08, 1963CO13, 1967KU1M, 1967LA05, 1967LA10, 1968SH11, 1969WE02). At  $E_p = 9, 10$  and 10.7 MeV, broad structures, possibly intermediate-state resonances, are reported by (1968BO36). See also (1961TA06, 1964DO03, 1966MA02) and (1969AL1H; theor.).

Polarization studies have been made at many energies: see Table 15.24 (1961RO05, 1961RO13, 1965RO22, 1966BE1M, 1966BR09, 1966DR02, 1966ST05, 1968GE04). See also the reviews in (1966RO1R, 1966RO1B, 1969WA11) and (1965TA07; theor.).

Spallation measurements are reported by (1967AU1A, 1967GR1K, 1968JA1M, 1968JU1B). For astrophysical considerations see (1967LI1B).

For reaction (b), see (1966MA02).




Table 15.24: Recent  $^{14}\text{N}(p, p)^{14}\text{N}$  yield curves and polarization studies <sup>a</sup>

(a) *Yield curves*

$E_p$ (MeV)	of protons to $^{14}\text{N}^*$	Refs.
0.9 – 4.0	g.s.	(1959BA16)
1.0 – 2.0	g.s.	(1961CO29, 1963CO13)
1.7 – 3.5	g.s.	(1959VA08)
1.9 – 3.0	g.s.	(1967LA05, 1967LA10)
2.6 – 4.8	g.s.	(1966DR02)
3.6 – 14.5	g.s.	(1967KU1M)
3.7 – 5.7	g.s., 2.31	(1967WE1B, 1969WE02)
5.28 – 5.7	3.94	(1967WE1B, 1969WE02)
6 – 9	g.s., 2.31, 3.94	(1968SH11)
6 – 11.5	g.s., 2.31, 3.94	(1968BO36)
6.8 – 9	5.11	(1968SH11)
6.8 – 11.5	5.11	(1968BO36)
8.4 – 11.5	5.83	(1968BO36)
9 – 11.5	4.91, 5.69	(1968BO36)
10 – 12	6.44, 7.03	(1968BO36)

(b) *Polarization measurements*

3.0 – 4.8	g.s.	(1966DR02)
4.9 – 10.1	g.s.	(1966BR09)
7.1 – 8.5	2.31	(1966BR09)
7.6 – 11.8	g.s.	(1966ST05)
7.8	g.s.	(1961RO13)
8.1 – 8.5	3.94	(1966BR09)
10.4	g.s.	(1961RO05)
14.5	g.s.	(1965RO22)
21.0	g.s.	(1966BE1M)
155	g.s.	(1968GE04)

<sup>a</sup> See also (1959AJ76).

The excitation function has been measured for  $E_p = 6.3$  to 12 MeV. Broad resonances are observed for  $E_p = 7.5$  to 9.85 MeV: see Table 15.23 (1964KU06). Broad resonance structure continues in the region up to  $E_p \approx 16.5$  MeV (1966KU12). See also (1963VA1C, 1965VA1E, 1969VE02), (1969AL1H; theor.) and  $^{14}\text{O}$ . See also (1966RE1D, 1969BA1N) for astrophysical considerations.

$$12. \ ^{14}\text{N}(\text{p}, \text{d})^{13}\text{N} \qquad Q_m = -8.328 \qquad E_b = 7.293$$

See  $^{13}\text{N}$  and (1963VA1C).

$$13. \ ^{14}\text{N}(\text{p}, \text{t})^{12}\text{N} \qquad Q_m = -22.139 \qquad E_b = 7.293$$

See  $^{12}\text{N}$  in (1968AJ02).

$$14. \ ^{14}\text{N}(\text{p}, \ ^3\text{He})^{12}\text{C} \qquad Q_m = -4.779 \qquad E_b = 7.293$$

The integrated cross section for this reaction has been studied for  $E_p = 8.2$  to 10.5 MeV (1968SH11). See also (1966MA02).

$$15. \ ^{14}\text{N}(\text{p}, \alpha)^{11}\text{C} \qquad Q_m = -2.920 \qquad E_b = 7.293$$

In the range  $E_p = 6$  to 9 MeV, the cross section for ground state  $\alpha$ -particles is large and shows many resonances: see Table 15.23 (1968SH11). Integrated cross sections for  $\alpha_0$  and  $\alpha_1$  are also reported for  $E_p = 8.2$  to 10.5 MeV by (1968SH11). See also (1952BL64, 1963VA1C, 1966MA02, 1969WE02) and see (1966RE1D) for astrophysical implications.

$$16. \ ^{14}\text{N}(\text{d}, \text{n})^{15}\text{O} \qquad Q_m = 5.068$$

Angular distribution studies have been conducted at many energies: see Table 15.26 (1953EV03, 1960EL04, 1960MO18, 1960RE07, 1962GR18, 1963CH1D, 1963GI16, 1965SI13, 1966LO1N, 1967MU12, 1968RI1T, 1969RI1C). Information derived from DWBA analysis of the angular distributions, and from the very accurate  $\gamma$ -ray measurements of (1965WA16, 1966AL18, 1967CH19) are shown in Table 15.27.

Table 15.25: Lifetimes of some  $^{15}\text{O}$  states

$E_x$ (MeV)	$\tau_m$ (psec)	Reaction	Refs.
5.18	< 0.3	$^{16}\text{O}(^3\text{He}, \alpha)$	(1965AL19)
	< 0.3	$^{14}\text{N}(\text{d}, \text{n})$	(1965WA03)
	< 0.1	$^{16}\text{O}(\gamma, \text{n})$	(1969MU07)
5.24	> 1	$^{14}\text{N}(\text{d}, \text{n}), ^{16}\text{O}(^3\text{He}, \alpha)$	(1965AL19)
	> 5	$^{14}\text{N}(\text{d}, \text{n})$	(1965WA03)
	$3.2 \pm 0.5$	$^{14}\text{N}(\text{d}, \text{n})$	(1967BI11)
6.18	< 0.047	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)
	< 0.1	$^{16}\text{O}(\gamma, \text{n})$	(1969MU07)
6.79	< 0.028	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)
6.86	$0.10 \pm 0.06$	$^{14}\text{N}(\text{d}, \text{n})$	(1966AL18)
	< 0.018	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)
7.28	$(1.25 \pm 0.3)$	$^{14}\text{N}(\text{d}, \text{n})$	(1966AL18)
	$(0.70 \pm 0.15)$	$^{14}\text{N}(\text{d}, \text{n})$	(1968GI11)

 Table 15.26: Recent  $^{14}\text{N}(\text{d}, \text{n})^{15}\text{O}$  angular distribution studies <sup>a</sup>

$E_d$ (MeV)	Distribution of neutron groups	Refs.
0.91 – 5.27	$\text{n}_0$	(1960RE07)
1.35 – 2.80	$\text{n}_0$	(1960EL04, 1965SI13)
1.53 – 2.90	$\text{n}_0$	(1960MO18)
2.2	$\text{n}_0$	(1966LO1N)
2.55, 2.7	$\text{n}_{4+5}$	(1960EL04)
2.83	$\text{n}_1 + \text{n}_2, \text{n}_3$	(1962GR18, 1963GI16)
2.85	$\text{n}_1, \text{n}_2, \text{n}_4, \text{n}_5$	(1963CH1D)
3.4 – 5.35	$\text{n}_0, \text{n}_2, \text{n}_3, \text{n}_4, \text{n}_5, \text{n}_6, \text{n}_7$	(1968RI1T, 1969RI1C)
5	$\text{n}_0, \text{n}_3, \text{n}_{4+5}, \text{n}_6$	(1967MU12)
7.7	$\text{n}_0, \text{n}_{1+2}, \text{n}_3, \text{n}_{4+5}, \text{n}_7, \text{n}_8$	(1953EV03)

<sup>a</sup> See also (1959AJ76).



Table 15.27: Levels of  $^{15}\text{O}$  from  $^{14}\text{N}(\text{d}, \text{n})^{15}\text{O}$  and  $^{14}\text{N}(^3\text{He}, \text{d})^{15}\text{O}$

$E_x^a$ (MeV)	$l_p^b$	$l_p^c$	$l_p^d$	$E_x^e$ (MeV $\pm$ keV)	$l_p^f$	$l_p^g$	$J\pi^h$
0	1	1	1		1	1	$\frac{1}{2}^-$
5.18					(0)	(0)	$\frac{1}{2}^+$
5.24	(2)		2	$5.24151 \pm 0.52$	2	2	$\frac{5}{2}^+$
6.18	1	1	1	$6.180 \pm 4$	1	1	$\frac{3}{2}^-$
6.79	0	0 + 2	0		0	0	$\frac{3}{2}^+$
6.86	2	0 + 2	0	$6.8598 \pm 1.0$	2	2	$\frac{5}{2}^+$
7.28	2	2		$7.2762 \pm 0.6$	2	2	$\frac{7}{2}^+$
7.55	0		1			0	$\frac{1}{2}^+$
8.28						0	$\frac{3}{2}^+$

<sup>a</sup> Nominal energies.

<sup>b</sup> (d, n): (1968RI1T, 1969RI1C).

<sup>c</sup> (d, n): (1967MU12).

<sup>d</sup> (d, n): (1953EV03).

<sup>e</sup> (d, n): (1965WA16, 1966AL18, 1967CH19).

<sup>f</sup> ( $^3\text{He}$ , d): (1968BO14).

<sup>g</sup> ( $^3\text{He}$ , d): (1969AL04).

<sup>h</sup> (1968RI1T, 1969AL04, 1969RI1C).

Neutron thresholds have been observed at  $E_d = 0.143 \pm 0.004$  and  $0.206 \pm 0.005$  MeV (1963CS02),  $1.24 \pm 0.02$ ,  $1.967 \pm 0.004$  and  $2.044 \pm 0.004$  (1955MA85), corresponding to  $^{15}\text{O}^*(5.19, 5.25, 6.15, 6.79, 6.86)$ . See also (1965MA1K).

Gamma ray branching ratios are shown in Table 15.22 (1965WA16, 1968GI11). Lifetime measurements are listed in Table 15.25 (1965AL19, 1965WA03, 1966AL18, 1967BI11, 1968GI11). See also (1959MO10, 1966AV1A).

17. (a)  $^{14}\text{N}(^3\text{He}, \text{d})^{15}\text{O}$   $Q_m = 1.799$   
 (b)  $^{14}\text{N}(^3\text{He}, \text{np})^{15}\text{O}$   $Q_m = -0.425$   
 $Q_0 = 1.803 \pm 0.010$  (1959YO25);  
 $Q_0 = 1.802 \pm 0.015$  (1960FO01).

Angular distributions obtained at  $E(^3\text{He}) = 11$  MeV (1968BO14) and 14 MeV (1969AL04) have been analyzed by DWBA: see Table 15.27. The angular distribution of deuterons has also

been measured at  $E(^3\text{He}) = 5.1$  MeV (1960FO01). For reaction (b) see (1967AD1F) and  $^{16}\text{F}$  in (1971AJ02).

$$18. \ ^{14}\text{N}(\alpha, \text{t})^{15}\text{O} \quad Q_{\text{m}} = -12.521$$

Angular distributions of tritons corresponding to the ground state of  $^{15}\text{O}$  have been determined at  $E_{\alpha} = 43$  MeV (1967DE1K) and at 56 MeV (1968GA1C, 1969GA11). At the higher energy, a detailed comparison is made with the results from the mirror reaction  $^{14}\text{N}(\alpha, ^3\text{He})^{15}\text{N}$  (1969GA11).

$$19. \ ^{14}\text{N}(^{11}\text{B}, ^{10}\text{Be})^{15}\text{O} \quad Q_{\text{m}} = -3.936$$

See (1967PO13, 1969BR1D).

$$20. \ ^{15}\text{N}(\text{p}, \text{n})^{15}\text{O} \quad Q_{\text{m}} = -3.542$$

$$E_{\text{thresh.}} = 3.7808 \pm 0.0011 [Q_0 = -3.5425 \pm 0.0011] \text{ (1958JO28, 1958WE1C)}.$$

Angular distributions of ground state neutrons have been measured at  $E_{\text{p}} = 3.95$  to 5.99 MeV (1958JO28), 3.95 to 8.99 MeV, 11.4 and 13.6 MeV (1963HA46; also  $\text{n}_2$  at  $E_{\text{p}} = 5.5$  MeV), 5.5 to 13.6 MeV (1961WO03) and 18.5 MeV (1964AN1B). See also (1961SA01), (1968WO1D), (1964SA1D, 1968HA15; theor.), and  $^{16}\text{O}$  in (1971AJ02).

$$21. \ ^{15}\text{N}(^3\text{He}, \text{t})^{15}\text{O} \quad Q_{\text{m}} = -2.778$$

A number of triton groups have been seen in this reaction. Angular distributions of these at  $E(^3\text{He}) = 39.8$  and 44.6 MeV, analyzed using a local two-body interaction with an arbitrary spin-isospin exchange mixture, lead to the  $L$ -values shown in Table 15.28 (1967BA13, 1968BA1E, 1969BA06). See also (1966EC1B, 1969BO13).

$$22. \ ^{16}\text{O}(\gamma, \text{n})^{15}\text{O} \quad Q_{\text{m}} = -15.668$$

Table 15.28: Levels of  $^{15}\text{O}$  from  $^{15}\text{N}(^3\text{He}, t)^{15}\text{O}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$L$	$E_x$ (MeV $\pm$ keV)
0	0	$9.63 \pm 40$
$5.24 \pm 30$	1, 3	$10.30 \pm 40$
6.18	0	$10.49 \pm 40$
$6.84 \pm 40$	1, 3	$10.97 \pm 50$
7.28	3	$11.21 \pm 60$
7.55	3	$11.69 \pm 40$
8.28	1	$12.34 \pm 40$
$8.94 \pm 40$		$13.78 \pm 40$
$9.47 \pm 50$		

<sup>a</sup> (1967BA13, 1968BA1E, 1969BA06);  $E(^3\text{He}) = 39.8, 44.6$  MeV.

The spectrum of photoneutrons has been investigated at many energies. Measurements over the giant dipole resonance region show the predominant strength is to the  $J^\pi = \frac{1}{2}^-$  and  $\frac{3}{2}^-$  states at  $E_x = 0$  and 6.18 MeV, consistent with the basic validity of the single-particle, single-hole theory of photoexcitation in  $^{16}\text{O}$ . However, the two positive parity states at  $E_x = 5.18$  and 5.24 MeV are also strongly populated, suggesting some non-single-particle excitation in that region in  $^{16}\text{O}$  (1965CA1B, 1965MA45, 1966OW01, 1967CA1C, 1967CA1P, 1968BA2L, 1969UL01). See also (1963FU05, 1964TA1C, 1965WI03, 1967FI1E, 1968JO1H, 1969CO15, 1969HO1T, 1969JO1N, 1969MU07). See also (1968ZH1B; theor.). For lifetime measurements of  $^{15}\text{O}^*$ , see Table 15.25 (1969MU07). See also  $^{16}\text{O}$  in (1971AJ02).

$$23. \ ^{16}\text{O}(n, 2n)^{15}\text{O} \quad Q_m = -15.668$$

See (1955AJ61).

$$24. \ (a) \ ^{16}\text{O}(p, d)^{15}\text{O} \quad Q_m = -13.443$$

$$(b) \ ^{16}\text{O}(p, pn)^{15}\text{O} \quad Q_m = -15.668$$

Reaction (a) goes primarily to the ground state and 6.18 MeV state ( $J^\pi = \frac{1}{2}^-$  and  $\frac{3}{2}^-$ , respectively). Angular distributions have been reported at many energies: see Table 15.29 (1961LE1A, 1963KA26, 1963LE03, 1966GR1A, 1967CH15, 1968LE01, 1969BA05, 1969SN03). See also

Table 15.29:  $^{16}\text{O}(\text{p}, \text{d})^{15}\text{O}$  angular distribution studies

$E_p$ (MeV)	Distribution of deuteron groups to $^{15}\text{O}^*$	Refs.
18.5 – 20	g.s.	(1961LE1A, 1963LE03)
21.3 – 45.3	g.s.	(1969SN03)
25.5 – 45.3	5.18 + 5.24, 6.18	(1969SN03)
30.3	g.s., 6.18	(1967CH15)
35, 40	g.s., 5.18, 6.18	(1966GR1A)
39.8	g.s., 6.18	(1963KA26)
45.3	7.28, 9.60, 10.46 <sup>a</sup>	(1969SN03)
100	g.s., 6.18	(1968LE01, 1968LI1A)
155.6	g.s., 6.18, 7, 10 <sup>b</sup>	(1965DE1A, 1969BA05, 1969TO1A)

<sup>a</sup> Partial angular distributions to many excited states.

<sup>b</sup> See, however, (1968LE01).

(1968SH08; theor.). The energy of  $^{15}\text{O}^*(7.28)$  is  $7.285 \pm 0.010$  MeV (1966MA1A). See also (1964BA04, 1966SH1A).

For reaction (b), see (1962BA1A, 1962FO03, 1963BE1A, 1967FU1A, 1968PU1A).

25.  $^{16}\text{O}(\text{d}, \text{t})^{15}\text{O}$   $Q_m = -9.411$

Angular distributions have been measured at  $E_d = 15$  MeV (1961KE01;  $t_0$ ), 20 MeV (1961VL02;  $t_0$ ), 20 MeV (1969PU04;  $t_0, t_1, t_2, t_3$ ;  $I_n = 1, 0, 2, 1$ ) and 28 MeV (1968GA13;  $t_0$ ). At  $E_d = 28$  MeV, detailed comparison is made with the results from the mirror reaction  $^{16}\text{O}(\text{d}, ^3\text{He})^{15}\text{N}$  (1968GA13).

26.  $^{16}\text{O}(^3\text{He}, \alpha)^{15}\text{O}$   $Q_m = 4.910$   
 $Q_0 = 4.916 \pm 0.010$  (1959HI68);  
 $Q_0 = 4.917 \pm 0.015$  (1962SH21).

The  $p_{1/2}$  and  $p_{3/2}$  hole states at  $E_x = 0$  and 6.18 MeV are strongly populated in this reaction, see e.g. (1965WA17). Information on these and other states of  $^{15}\text{O}$  observed in this reaction is shown in Table 15.30 (1959HI68, 1959YO25, 1967HE1A, 1968BO14). The  $J$ -values are derived from angular correlation measurements (1967HE1A, 1966GA19, 1966GO15, 1967GO07,

Table 15.30: Excited states of  $^{15}\text{O}$  from  $^{16}\text{O}(^3\text{He}, \alpha)^{15}\text{O}$ 

$E_x$ (MeV $\pm$ keV)	Refs.	$l^b$	$J$	Refs.
0		1		
$5.174 \pm 10$	(1959HI68)	0		
$5.167 \pm 15$	(1959YO25)			
$5.193 \pm 11$	(1965AL19)			
$5.233 \pm 10$	(1959HI68)	2	$\frac{5}{2}$	(1966GO15, 1967GO07)
$5.243 \pm 10$	(1965AL19)			
6.18	<sup>a</sup>	1	$\frac{3}{2}$	(1966GO15)
6.79	<sup>a</sup>	2	$\frac{3}{2}$	(1966GA19, 1968GI01)
6.86	<sup>a</sup>	2	$\frac{5}{2}$	(1966GA19, 1968GI01)
$7.2742 \pm 1.4$	(1967HE1A)		$\frac{7}{2}$	(1967HE1A)

<sup>a</sup> Nominal energy.

<sup>b</sup> (1968BO14).

(1968GI01): see also (1968BO14). Angular distributions have been measured for  $E(^3\text{He}) = 5.2$  to 36.6 MeV: see Table 15.31 (1959HI73, 1960TA12, 1962SE13, 1965AL05, 1965AR07, 1968BO14, 1969BR07). Branching ratios are displayed in Table 15.22 (1965WA16, 1969KU01). See also (1965WA03). The lifetimes of  $^{15}\text{O}^*(5.18, 5.24)$  are  $< 0.3$  and  $> 1$  psec, respectively (1965AL19): see also Table 15.25. A comparison of the 5 MeV transitions E3/M2 mixing ratios in  $^{15}\text{O}$  and  $^{15}\text{N}$  strongly suggest a collective character for the E3 component of these transitions (1968GI01).

The ratio of mixing ratios of the mirror decays  $^{15}\text{O}^*(6.18 \rightarrow 0)$  and  $^{15}\text{N}^*(6.32 \rightarrow 0)$  are in disagreement with the IPM suggesting a collective contribution to the mirror  $\frac{3}{2}^-$  levels (1966RO1U).

The M2/E1 mixing ratio for the  $^{15}\text{O}^*(6.79)$  transition indicates an exceptionally high retardation of an E1 transition in a non-self-conjugate nucleus, as is also true of the analog transition in  $^{15}\text{N}$  (1968GI01).

The  $E_x = 7.28$  MeV state,  $J = \frac{7}{2}$ , has a negligible effect on the (astrophysical) CNO cycle (1967HE1A).

See also (1961DU02, 1961SI09, 1966AG1B) and  $^{19}\text{Ne}$  in (1972AJ02).

$$27. \ ^{16}\text{O}(^{14}\text{N}, ^{15}\text{N})^{15}\text{O} \quad Q_m = -4.833$$

See (1965GA1B, 1969BR1D).

$$28. \ ^{17}\text{O}(p, t)^{15}\text{O} \quad Q_m = -11.329$$

Table 15.31:  $^{16}\text{O}(^3\text{He}, \alpha)$  angular distribution studies

$E(^3\text{He})$ (MeV)	Distribution of $\alpha$ groups to $^{15}\text{O}^*$	Refs.
5.2	g.s.	(1960TA12)
5.7, 5.9	g.s.	(1959HI73)
8 – 10	g.s.	(1965AL05)
9.2	g.s., 5.18, 5.24	(1959HI73)
9.8 – 11.7	g.s.	(1966BR13, 1969BR07)
11	g.s., 5.18, 5.24, 6.18, 6.79, 6.86	(1968BO14)
16.6, 25.8, 36.6	g.s., 5.18 + 5.24, 6.18	(1965AR07)
29	5.18 + 5.24, 6.18	(1962SE13)

See (1969ME1M).

29.  $^{19}\text{F}(\text{p}, \text{n}\alpha)^{15}\text{O}$   $Q_{\text{m}} = -7.553$

See (1962FO03).

### $^{15}\text{F}$

*Mass of  $^{15}\text{F}$ :* A calculation using an isobaric mass formula predicts that  $^{15}\text{F}$  is unstable with respect to proton emission by 2.32 MeV: the mass excess of  $^{15}\text{F}$  is then 17.62 MeV (1966KE16). See also (1957MU99, 1961BA1C).

## References

(Closed 31 December 1969)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1941BE1A Bennett et al., Phys. Rev. 59 (1941) 781
- 1950CU13 C.D. Curling and J.O. Newton, Nature 165 (1950) 609
- 1950JO57 C.H. Johnson and H.H. Barschall, Phys. Rev. 80 (1950) 818
- 1950MA65 R. Malm and W.W. Buechner, Phys. Rev. 80 (1950) 771
- 1950RI57 J.E. Richardson, Phys. Rev. 80 (1950) 850
- 1951DU08 D.B. Duncan and J.E. Perry, Phys. Rev. 82 (1951) 809
- 1951JO1A Johnson, Petree and Adair, Phys. Rev. 84 (1951) 775
- 1951RO16 W.D. Roseborough, J.J.G. McCue, W.M. Preston and C. Goodman, Phys. Rev. 83 (1951) 1133
- 1951YA1A Yaffe and Stevens, Can. J. Phys. 29 (1951) 186; Phys. Rev. 79 (1950) 893
- 1952BL64 J.P. Blaser, P. Marmier and M. Sempert, Helv. Phys. Acta 25 (1952) 442
- 1952GI01 W.M. Gibson and E.E. Thomas, Proc. Roy. Soc. A210 (1952) 543
- 1952HI12 J.J. Hinchey, P.H. Stelson and W.M. Preston, Phys. Rev. 86 (1952) 483
- 1953EV03 W.H. Evans, T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A66 (1953) 108
- 1953KA1A Kay, Mark and Goodman, Phys. Rev. 112 (1958) 503
- 1953KO42 B. Koudijs, F.P.G. Valckx and P.M. Endt, Physica 19 (1953) 1133
- 1954BE08 W.E. Bennett, P.A. Roys and B.J. Toppel, Phys. Rev. 93 (1954) 924, K4
- 1954KL36 R.M. Kline and D.J. Zaffarano, Phys. Rev. 96 (1954) 1620
- 1954PA39 R. Pauli, K. Ahnlund and C. Mileikowsky, Ark. Fys. 8 (1954) 213
- 1954SP01 A. Sperduto, W.W. Buechner, C.K. Bockelman and C.P. Browne, Phys. Rev. 96 (1954) 1316
- 1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77
- 1955BA44 G.A. Bartholomew, F. Brown, H.E. Gove, A.E. Litherland and E.B. Paul, Can. J. Phys. 33 (1955) 441
- 1955BA83 S. Bashkin, R.R. Carlson and E.B. Nelson, Phys. Rev. 99 (1955) 107
- 1955FO27 J.L. Fowler and C.H. Johnson, Phys. Rev. 98 (1955) 728

1955MA76 J.B. Marion, T.W. Bonner and C.F. Cook, Phys. Rev. 100 (1955) 847.  
1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46  
1955SH28 R.D. Sharp, A. Sperduto and W.W. Buechner, Phys. Rev. 99 (1955) 632A  
1955SH46 E.S. Shire and R.D. Edge, Phil. Mag. 46 (1955) 640  
1956BA16 G.A. Bartholomew, A.E. Litherland, E.B. Paul and H.E. Gove, Can. J. Phys. 34 (1956) 147  
1956BA34 J.K. Bair, H.O. Cohn, J.D. Kington and H.B. Willard, Phys. Rev. 104 (1956) 1595  
1956BO61 T.W. Bonner, A.A. Kraus Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348  
1956DO37 R.A. Douglas, B.R. Gasten and A. Mukerji, Can. J. Phys. 34 (1956) 1097  
1956DO41 R.A. Douglas, J.W. Broer, R. Chiba, D.F. Herring and E.A. Silverstein, Phys. Rev. 104 (1956) 1059  
1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28  
1956MA35 J.B. Marion and G. Weber, Phys. Rev. 102 (1956) 1355; Erratum Phys. Rev. 103 (1956) 1906  
1956MA46 J.B. Marion and G. Weber, Phys. Rev. 103 (1956) 167  
1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434  
1956TA16 G.W. Tautfest and S. Rubin, Phys. Rev. 103 (1956) 196  
1956VA17 F.P.G. Valckx, Ph.D. Thesis, Univ. of Utrecht (1956)  
1957BA18 G.A. Bartholomew and P.J. Campion, Can. J. Phys. 35 (1957) 1347  
1957BO58 C.R. Bolmgren, G.D. Freier, J.G. Likely and K.F. Famularo, Phys. Rev. 105 (1957) 210  
1957BR18 D.A. Bromley, E. Almqvist, H.E. Gove, A.E. Litherland, E.B. Paul and A.J. Ferguson, Phys. Rev. 105 (1957) 957  
1957HA03 F.B. Hagedorn, F.S. Mozer, T.S. Webb, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 105 (1957) 219  
1957HA1E Halbert and French, Phys. Rev. 105 (1957) 1563  
1957JO20 S.A.E. Johansson and B. Forkman, Ark. Fys. 12 (1957) 359  
1957KI22 O.C. Kistner, A. Schwarzschild, B.M. Rustad and D.E. Alburger, Phys. Rev. 105 (1957) 1339  
1957LA13 W.A.S. Lamb and R.E. Hester, Phys. Rev. 108 (1957) 1304  
1957MU99 P.G. Murphy, Phys. Rev. 108 (1957) 421  
1957NO14 E. Norbeck Jr., Phys. Rev. 105 (1957) 204  
1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754



1957PE12 J.R. Penning and F.H. Schmidt, Phys. Rev. 105 (1957) 647  
 1957PI1A Pixley, Thesis, CalTech (1957)  
 1957SJ68 B. Sjogren and K. Ahnlund, Ark. Fys. 12 (1957) 547  
 1957WA01 E.K. Warburton and J.N. McGruer, Phys. Rev. 105 (1957) 639  
 1958AL1D Alkhazov, Gangskii and Lemberg, JETP (Sov. Phys.) 6 (1958) 892  
 1958GR1B Groshev, Demidov, Lutsenko and Pelekhov, Atlas of Gamma-Ray Spectra (Moscow, 1958)  
 1958HA1B Haddad, Perry and Smith, Private Communication (1958)  
 1958JO20 R.L. Johnston, H.D. Holmgren, E.A. Wolicki and E.G. Illsley, Phys. Rev. 109 (1958) 884  
 1958JO28 K.W. Jones, L.J. Lidofsky and J.L. Weil, Phys. Rev. 112 (1958) 1252  
 1958WA1C Way, Nucl. Data Cards, Natl. Res. Council, Washington, D.C. (1958)  
 1958WE1C Weil, CU-180 (1958)  
 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1  
 1959AL06 D.E. Alburger, A. Gallmann and D.H. Wilkinson, Phys. Rev. 116 (1959) 939  
 1959AL97 D.E. Alburger, Phys. Rev. Lett. 3 (1959) 280  
 1959BA16 S. Bashkin, R.R. Carlson and R.A. Douglas, Phys. Rev. 114 (1959) 1552  
 1959BI1B Bilpuch, Weston, Bowman and Newson, Bull. Amer. Phys. Soc. 4 (1959) 42  
 1959BO1C Bockelman, Nucl. Phys. 13 (1959) 205  
 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314  
 1959BR69 P. Brix and E.K. Maschke, Z. Phys. 155 (1959) 109  
 1959BR79 D.A. Bromley, J.A. Kuehner and E. Almqvist, Nucl. Phys. 13 (1959) 1  
 1959CO1C Cohen, Fisher and Warburton, Phys. Rev. Lett. 3 (1959) 433; Erratum Phys. Rev. Lett. 4 (1960) 92  
 1959FE1B Feingold, Phys. Rev. 114 (1959) 540  
 1959FE1C Ferguson and Gove, Can. J. Phys. 37 (1959) 660  
 1959GA05 A. Gallmann, Ann. Phys. 4 (1959) 185  
 1959GA14 F. Gabbard, H. Bischel and T.W. Bonner, Nucl. Phys. 14 (1959) 277  
 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295  
 1959HE1B Hess, Ann. Phys. 6 (1959) 115  
 1959HE1D Hebbard and Dunbar, Phys. Rev. 115 (1959) 624  
 1959HE47 D.F. Hebbard and B. Povh, Nucl. Phys. 13 (1959) 642  
 1959HI68 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A73 (1959) 727

- 1959HI73 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 775
- 1959JO32 R.L. Johnston, H.D. Holmgren and G.D. Gutsche, Bull. Amer. Phys. Soc. 4 (1959) 403, C4
- 1959KI99 O.C. Kistner and B.M. Rustad, Phys. Rev. 114 (1959) 1329
- 1959LE28 L.L. Lee Jr. and J.P. Schiffer, Phys. Rev. 115 (1959) 160
- 1959MA1D Markowitz and Hall, Bull. Amer. Phys. Soc. 4 (1959) 8
- 1959MO10 E. Moller, Ark. Fys. 15 (1959) 251
- 1959MO1B Morre and McGruer, Bull. Amer. Phys. Soc. 4 (1959) 17, and Private Communication (1959)
- 1959NO38 I. Nonaka, H. Yamaguchi, T. Mikumo, I. Umeda, T. Tabata and S. Hitaka, J. Phys. Soc. Jpn. 14 (1959) 1260; Erratum J. Phys. Soc. Jpn. 15 (1960) 365
- 1959PO79 B. Povh and D.F. Hebbard, Phys. Rev. 115 (1959) 608
- 1959VA04 A.K. Valter, V.Y. Gonchar, A.N. Lvov and S.P. Tsitko, Izv. Akad. Nauk SSSR Ser. Fiz. 23 (1959) 228; Columbia Tech. Transl. 23 (1960) 219
- 1959VA08 A.K. Valter, A.S. Deineko, I.Y. Malakhov, P.V. Sorokin and A.Y. Taranov, Izv. Akad. Nauk SSSR Ser. Fiz. 23 (1959) 839
- 1959YO25 T.E. Young, G.C. Phillips, R.R. Spencer and D.A.A.S.N. Rao, Phys. Rev. 116 (1959) 962
- 1960BA34 J.K. Bair, R.D. Edge and H.B. Willard, Phys. Rev. 119 (1960) 1948
- 1960BE1B Belyaev, Zakhar'ev and Neudachin, Atomnaya Energiya 9 (1960) 298, Sov. J. At. Energy 9 (1961) 833
- 1960BU1C Bullock and Moore, Phys. Rev. 119 (1960) 721
- 1960CA02 R.E. Carter and H.T. Motz, Bull. Amer. Phys. Soc. 5 (1960) 246, HA7
- 1960CL02 R.L. Clarke, E. Almqvist and E.B. Paul, Nucl. Phys. 14 (1960) 472
- 1960EL04 A.J. Elwyn, J.V. Kane, S. Ofer and R. Pixley, Phys. Rev. 120 (1960) 2207
- 1960FA10 R.F. Favreau, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds. D.A. Bromley and E.W. Vogt, (Univ. of Toronto, Press, 1960) 939
- 1960FE12 J.M. Ferguson and W.E. Thompson, Phys. Rev. 118 (1960) 228
- 1960FO01 P.D. Forsyth, F. de S. Barros, A.A. Jaffe, I.J. Taylor and S. Ramavataram, Proc. Phys. Soc. (London) A75 (1960) 291
- 1960FR09 J.B. French, S. Iwao and E.W. Vogt, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds. D.A. Bromley and E.W. Vogt, (Univ. of Toronto Press, 1960) 480
- 1960HE13 D.F. Hebbard, Nucl. Phys. 19 (1960) 511
- 1960HI07 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) 75 (1960) 745

- 1960HO1B Honsaker, *Astrophys. J.* 132 (1960) 21
- 1960JA12 J. Janecke, *Z. Naturforsch.* A15 (1960) 593
- 1960MC05 J.H. McCrary and I.L. Morgan, *Bull. Amer. Phys. Soc.* 5 (1960) 246, HA5
- 1960MO03 S. Morita, N. Kawai, N. Takano, Y. Goto, R. Hanada, Y. Nakajima, S. Takemoto and Y. Yaegashi, *J. Phys. Soc. Jpn.* 15 (1960) 361
- 1960MO18 S. Morita, N. Kawai, Y. Goto, T. Maki and M. Mukae, *J. Phys. Soc. Jpn.* 15 (1960) 2170
- 1960PE1B Peterson, Bratenel and Stoering, *Phys. Rev.* 120 (1960) 521
- 1960PR13 J.R. Priest, D.J. Tendam and E. Bleuler, *Phys. Rev.* 119 (1960) 1301
- 1960RE07 T. Retz-Schmidt and J.L. Weil, *Phys. Rev.* 119 (1960) 1079
- 1960SH05 S.M. Shafroth, *J. Phys. Rad.* 21 (1960) 353
- 1960TA12 I.J. Taylor, F.de S. Barros, P.D. Forsyth, A.A. Jaffe and S. Ramavataram, *Proc. Phys. Soc. (London)* A75 (1960) 772
- 1960TA17 T. Tabata and K. Okano, *J. Phys. Soc. Jpn.* 15 (1960) 1552
- 1960TA1C Talmi and Unna, *Ann. Rev. Nucl. Sci.* 10 (1960) 353
- 1960TA1E Tauber and Wu, *Proc. Int. Conf. on Nucl. Struct., Kingston* (1960)
- 1960VA11 F.J. Vaughn, L.F. Chase Jr. and R.G. Johnson, *Bull. Amer. Phys. Soc.* 5 (1960) 404, B2
- 1960ZE03 Ya.B. Zeldovich, *Zh. Eksp. Teor. Fiz.* 38 (1960) 1123; *JETP (Sov. Phys.)* 11 (1960) 812
- 1961AG1A Aguilar and Garcia, *An. Real. Soc. Espan. Fis. y Quim.* A57 (1961) 29; *Phys. Abs.* 3611 (1962)
- 1961BA05 G.A. Bartholomew and J.F. Vervier, *Bull. Amer. Phys. Soc.* 6 (1961) 237, DA13
- 1961BA1C Baz, Goldanskii and Zeldovich, *Sov. Phys. Uspekhi* 3 (1961) 729
- 1961BA1E Balashov, Neudachin and Smirnov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 25 (1961) 170; *Bull. Acad. Sci. USSR Phys.* 25 (1961) 165
- 1961BR13 H.R. Brooker, P.J. Haigh and T.A. Scott, *Phys. Rev.* 123 (1961) 2143
- 1961BR1A Brill', Vlasov, Kalinin and Sokolov, *Dokl. Akad. Nauk SSSR* 136 (1961) 55; *Sov. Phys., Dokl.* 6 (1961) 24
- 1961CA02 R.R. Carlson, C.C. Kim, J.A. Jacobs and A.C.L. Barnard, *Phys. Rev.* 122 (1961) 607
- 1961CH14 R. Chiba, *Phys. Rev.* 123 (1961) 1316
- 1961CO02 S.G. Cohen, P.S. Fisher and E.K. Warburton, *Phys. Rev.* 121 (1961) 858
- 1961CO29 J. Cohen-Ganouna, M. Lambert and J. Schmouker, *J. Phys. Rad.* 22 (1961) 592
- 1961DU02 K.L. Dunning and J.W. Butler, *Phys. Rev.* 123 (1961) 1321

- 1961DU1B Duggan, Thesis, Louisiana State Univ. (1961)
- 1961FO02 P.D. Forsyth and G.S. Mani, Bull. Amer. Phys. Soc. 6 (1961) 227, AA12
- 1961FR1D French, Iwao and Vogt, AECL 1136 (1960)
- 1961GO03 S. Gorodetzky, P. Fintz, G. Bassompierre and A. Gallmann, Compt. Rend. 252 (1961) 713
- 1961HA12 J. Hanna, T.W. Bonner, R.L. Bramblett and F.A.St. Romain, Bull. Amer. Phys. Soc. 6 (1961) 369, S9
- 1961HA43 G.C. Hanna, D.B. Primeau and P.R. Tunncliffe, Can. J. Phys. 39 (1961) 1784
- 1961JA09 A.N. James, Nucl. Phys. 24 (1961) 132
- 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, Rev. Mex. Fisica 10 (1961) 247
- 1961JO07 R.G. Johnson, L.F. Chase Jr. and F.J. Vaughn, Bull. Amer. Phys. Soc. 6 (1961) 236, DA8
- 1961JO24 R.G. Johnson, L.F. Chase Jr. and F.J. Vaughn, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 591
- 1961KA05 N. Kawai, J. Phys. Soc. Jpn. 16 (1961) 157
- 1961KE01 E.L. Keller, Phys. Rev. 121 (1961) 820
- 1961KO04 M. Kondo, T. Yamazaki and S. Yamabe, J. Phys. Soc. Jpn. 16 (1961) 1091
- 1961KO1E Kolodziejewski, Acta Phys. Pol. 20 (1961) 275
- 1961KR1A Kromminga and McCarthy, Nucl. Phys. 24 (1961) 36
- 1961LE01 J.J. Leigh and J.M. Blair, Phys. Rev. 121 (1961) 246
- 1961LE1A Legg, Unpublished Thesis, Princeton Univ. (1961)
- 1961LO10 G. Lopez and O. Almen, Rev. Mex. Fis. 10 (1961) 239
- 1961MO13 S. Morita, T. Ishimatsu, T. Cho, Y. Nakajima, N. Kawai, T. Murata and Y. Hachiya, J. Phys. Soc. Jpn. 16 (1961) 1849; Erratum J. Phys. Soc. Jpn. 17 (1962) 251
- 1961NO05 E. Norbeck, Phys. Rev. 121 (1961) 824
- 1961PU1B Pullen, Wilkinson and Whitehead, Proc. Rutherford Jub. Int. Conf., Manchester, England; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 565
- 1961RA06 L.A. Rayburn, Phys. Rev. 122 (1961) 168
- 1961RO05 L. Rosen, J.E. Brolley Jr. and L. Stewart, Phys. Rev. 121 (1961) 1423
- 1961RO13 L. Rosen, J.E. Brolley Jr., M.L. Gursky and L. Stewart, Phys. Rev. 124 (1961) 199
- 1961SA01 A. Sayres, D. Lister and D. Lightbody, Bull. Amer. Phys. Soc. 6 (1961) 26, I11
- 1961SI09 E.A. Silverstein, S.R. Salisbury, G. Hardie and L.D. Oppliger, Phys. Rev. 124 (1961) 868
- 1961SJ1B Sjogren and Sawa, Ark. Fys. 19 (1961) 417

- 1961TA06 A.E. Taylor and E. Wood, Nucl. Phys. 25 (1961) 642
- 1961TE02 A. Tejera, M. Mazari, A. Jaidar and G. Lopez, Rev. Mex. Fis. 10 (1961) 229
- 1961TO01 K.S. Toth, Phys. Rev. 121 (1961) 1190
- 1961TO03 J.H. Towle and B.E.F. Macefield, Proc. Phys. Soc. (London) 77 (1961) 399
- 1961TO07 K.S. Toth, Phys. Rev. 123 (1961) 582
- 1961VA13 P.I. Vatsset, L.Ya. Kolesnikov and S.G. Tonapetyan, Zh. Eksp. Teor. Fiz. 40 (1961) 1257; JETP (Sov. Phys.) 13 (1961) 886
- 1961VL02 N.A. Vlasov, S.P. Kalinin, A.A. Ogloblin and V.I. Chuev, Izv. Akad. Nauk SSSR Ser. Fiz. 25 (1961) 115; Columbia Tech. Transl. 25 (1961) 111
- 1961WI1A Wilkinson, Wollan and Koehler, Ann. Rev. Nucl. Sci. 11 (1961) 303
- 1961WO03 C. Wong, J.D. Anderson, S.D. Bloom, J.W. McClure and B.D. Walker, Phys. Rev. 123 (1961) 598
- 1962AL01 W.P. Alford, O.M. Bilaniuk and D.B. Marsh, Bull. Amer. Phys. Soc. 7 (1962) 60, R8
- 1962BA1A Balashov and Boyarkina, Nucl. Phys. 38 (1962) 629
- 1962BA63 J.D. Baldeschweiler, J. Chem. Phys. 36 (1962) 152
- 1962BR1G Beit, in Electomag. Lifetimes Props. of Nucl. States, N.A.S.N.R.C. Pub. 974 (1962) 21
- 1962CE1B Celvolani and Petralia, Nuovo Cim. 26 (1962) 1328
- 1962CH14 L.F. Chase Jr., R.G. Johnson, F.J. Vaughn and E.K. Warburton, Phys. Rev. 127 (1962) 859
- 1962CO1C Cochran and Knight, Phys. Rev. 128 (1962) 1281
- 1962DO1A Dodge and Barber, Phys. Rev. 127 (1962) 1746
- 1962EI03 W.W. Eidson and R.D. Bent, Phys. Rev. 127 (1962) 913
- 1962ER03 H.J. Erramuspe and R.J. Slobodrian, Nucl. Phys. 34 (1962) 532
- 1962FO03 K.J. Foley, G.L. Salmon and A.B. Clegg, Nucl. Phys. 31 (1962) 43
- 1962GO1J Gorshkov, Zyabkin and Tsvetkov, Atomn. Energ. (USSR) 13 (1962) 65
- 1962GO21 S. Gorodetzky, P. Fintz and A. Gallmann, Compt. Rend. 255 (1962) 879
- 1962GR18 W. Gruebler and J. Rossel, Helv. Phys. Acta 35 (1962) 283
- 1962GU01 G.D. Gutsche, H.D. Holmgren, L.M. Cameron and R.L. Johnston, Phys. Rev. 125 (1962) 648
- 1962HA20 J. Hanna, F.R.St. Romain, R.L. Bramblett and T.W. Bonner, Bull. Amer. Phys. Soc. 7 (1962) 453, GA4
- 1962HO06 R.K. Hobbie and F.F. Forbes, Phys. Rev. 126 (1962) 2137
- 1962HO1D Honda and Ui, Nucl. Phys. 34 (1962) 592

- 1962KU09 B. Kuhn, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teopr. Fiz. 43 (1962) 1660; JETP (Sov. Phys.) 16 (1963) 1171
- 1962LI07 A.J. Lieber, F.H. Schmidt and J.B. Gerhart, Phys. Rev. 126 (1962) 1496
- 1962LO02 J. Lowe, C.L. McClelland and J.V. Kane, Phys. Rev. 126 (1962) 1811
- 1962MC12 R.L. McGrath, Phys. Rev. 127 (1962) 2138
- 1962NE1D Nedvedyk, Joint Inst. Nucl. Res. Lab. Neutron Phys. USSR, Rept. No. P-1098 (1962)
- 1962OT01 P.S. Otstavnov and V.I. Popov, Zh. Eksp. Teor. Fiz. 43 (1962) 385; JETP (Sov. Phys.) 16 (1963) 276
- 1962RO13 V.M. Rout, W.M. Jones and D.G. Waters, Nucl. Phys. 34 (1962) 628
- 1962SE13 H.M. Sen Gupta, J. Rotblat, P.E. Hodgson and J.B.A. England, Nucl. Phys. 38 (1962) 361
- 1962SH21 R.D. Sharp, Bull. Amer. Phys. Soc. 7 (1962) 622, W1
- 1962ST17 D. Stanojevic and M. Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 13 (1962) 1
- 1962TA1E Talmi, Nucl. Spectroscopy; Ed. Racah (Academic Press, 1962)
- 1962TE1B Teplov, Zh. Eksp. Teor. Fiz. 42 (1962) 211; JETP (Sov. Phys.) 15 (1962) 150
- 1963AL21 D.E. Alburger and E.K. Warburton, Phys. Rev. 132 (1963) 790
- 1963BA1P Bailey and Hebbard, Nucl. Phys. 46 (1963) 529
- 1963BA1Q Ballini and Saunier, J. Physique 24 (1963) 904
- 1963BA46 R.W. Bauer, J.D. Anderson and L.J. Christensen, Nucl. Phys. 47 (1963) 241
- 1963BE1A Benioff, Phys. Rev. 129 (1963) 1355
- 1963BE42 T. Berggren and G. Jacob, Nucl. Phys. 47 (1963) 481
- 1963BR1G Breit, Padua (1963) 480
- 1963BU1C Bunakov, Phys. Lett. 7 (1963) 14
- 1963CH1C Chatterjee, Nucl. Phys. 49 (1963) 686
- 1963CH1D Christiansen and Zeitnitz, Padua (1963) A557
- 1963CL1A Clement, Padua (1963) 457
- 1963CL1B Clegg, CERN 63-28 (1963)
- 1963CO12 B.L. Cohen, Phys. Rev. 130 (1963) 227
- 1963CO13 J. Cohen-Ganouna, M. Lambert and J. Schmouker, J. Phys. 24 (1963) 43
- 1963CO17 E.D. Commins and H.R. Feldman, Phys. Rev. 131 (1963) 700
- 1963CS02 J. Csikai and G. Peto, Phys. Lett. 4 (1963) 252
- 1963DA1B Dar, Phys. Lett. 7 (1963) 339
- 1963DE19 V.K. Deshpande, Nucl. Phys. 47 (1963) 257



- 1963DR1B Drisko, Satchler and Dassel, 3rd Conf. on Reactions between Complex Nuclei (1963) 85
- 1963DU12 J.L. Duggan, P.D. Miller and R.F. Gabbard, Nucl. Phys. 46 (1963) 336
- 1963ED01 R.D. Edge, Bull. Amer. Phys. Soc. 8 (1963) 11, BA5
- 1963EN01 C.E. Engelke, R.E. Benenson, E. Melkonian and J.M. Lebowitz, Phys. Rev. 129 (1963) 324
- 1963FI04 E. Finckh, R. Kosiek and K. Schlupmann, Naturwiss. 50 (1963) 326
- 1963FU05 H. Fuchs and D. Haag, Z. Physik 171 (1963) 403
- 1963GA10 E. Gadioli and S. Micheletti, Phys. Lett. 6 (1963) 229
- 1963GI16 W.R. Gibbs and W.E. Gruebler, Helv. Phys. Acta 36 (1963) 693
- 1963GO1J Gorshkov and Tsvetkov, Atomnaya Energ.14 (1963) 550
- 1963GO1L Gofman et al., Zh. Eksp. Teor. Fiz. 45 (1963) 1317; JETP (Sov. Phys) 18 (1964) 906
- 1963GO1M Goldberg, May and Stehn, BNL-400, 2nd Ed., Vol. 1 (1963)
- 1963HA46 L.F. Hansen and M.L. Stelts, Phys. Rev. 132 (1963) 1123
- 1963HE11 D.F. Hebbard and G.M. Bailey, Nucl. Phys. 49 (1963) 666
- 1963HO1E Hortig, Werner and Gentner, 3rd Conf. on Reactions between Complex Nuclei (1963) 178
- 1963IM01 W.L. Imhof, H.A. Grench and R.G. Johnson, Nucl. Phys. 49 (1963) 503
- 1963KA26 C.D. Kavaloski, G. Bassani and N.M. Hintz, Phys. Rev. 132 (1963) 813
- 1963KI1B Kim, Nucl. Phys. 49 (1963) 383
- 1963KO03 M. Kondo, T. Yamazaki and S. Yamabe, J. Phys. Soc. Jpn. 18 (1963) 22
- 1963KU1B Kunz, Can. J. Phys. 41 (1963) 2187
- 1963LA02 R.A. Lasalle, R.D. Benta and J.G. Cramer Jr., Bull. Amer. Phys. Soc. 8 (1963) 303, D8
- 1963LE03 J.C. Legg, Phys. Rev. 129 (1963) 272
- 1963LU01 C.A. Ludemann, H.D. Holmgren and W.F. Hornyak, Bull. Amer. Phys. Soc. 8 (1963) 12, BA10
- 1963LU05 C.A. Ludemann, H.D. Holmgren and J.E. Etter, Bull. Amer. Phys. Soc. 8 (1963) 303, D6
- 1963LU1F C.A. Ludemann, H.D. Holmgren and W.F. Hornyak, Padua (1963) 850A
- 1963MA28 G.S. Mani, P.D. Forsyth and R.R. Perry, Nucl. Phys. 44 (1963) 625
- 1963MO1B Morrison, Gale, Hussain and Murray, 3rd Conf. on Reactions between Complex Nuclei (1963) 168
- 1963MO1C Motz and Journey, Wash-1044 (1963)

1963NA1C Nagahara et al., Padua (1963) 850A  
1963NE05 J.W. Nelson, E.B. Carter, G.E. Mitchell and R.H. Davis, Phys. Rev. 129 (1963) 1723  
1963NI04 K. Niedzwiedziuk, V.I. Salatskii and I.V. Sizov, Zh. Eksp. Teor. Fiz. 44 (1963) 1450;  
JETP (Sov. Phys.) 17 (1963) 974  
1963RI1B Riou, Padua (1963) 18  
1963SC32 J.A. Scheer, K. Schlupmann and F. Triantafyllidis, Phys. Lett. 7 (1963) 269  
1963TA1A Tanifuji, Nucl. Phys. 40 (1963) 357  
1963TO1D Toth and Newman, 3rd Conf. on Reactions between Complex Nuclei (1963) 114  
1963VA1C Valentin, Albouy, Cohen and Gusakow, Phys. Lett. 7 (1963) 163  
1963VA31 S.S. Vasilev and L.Y. Shavtvalov, Izv. Akad. Nauk SSSR Ser. Fiz. 27 (1963) 1261;  
Bull. Acad. Sci. USSR Phys. Ser. 27 (1963) 1239  
1963YA1C Yamazaki, Knodo and Yamabe, J. Phys. Soc. Jpn. 18 (1963) 620  
1964AL1L Ali and Tauber, Nucl. Phys. 55 (1964) 481  
1964AL21 D.E. Alburger, C. Chasman, K.W. Jones and R.A. Ristinen, Phys. Rev. 136 (1964)  
B913  
1964AM1A Amsel, Ann. Phys. 9 (1964) 297  
1964AM1D Amit and Latz, Nucl. Phys. 58 (1964) 297  
1964AN1B Anderson, Wong, McClure and Walker, Phys. Rev. 136 (1964) B118  
1964BA02 G.A. Bartholomew and J.F. Vervier, Nucl. Phys. 50 (1964) 209  
1964BA04 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz, N.K. Ganguly and P. Radvanyi, Phys.  
Lett. 8 (1964) 56  
1964BA1V Ballot, Dumontet, Picinbono and Saya, Nucl. Phys. 51 (1964) 401  
1964BL1B Blair, Pham-Dihn-Lien and Hobbie, Phys. Rev. 134 (1964) B793  
1964BL1C Blieden, Phys. Lett. 9 (1964) 176  
1964BR13 H.C. Bryant, J.G. Beery, E.R. Flynn and W.T. Leland, Nucl. Phys. 53 (1964) 97  
1964BR1G Brill, Chuev and Ogloblin, Paris (1964) 981  
1964BR1H Brown, Paris (1964) 129  
1964BR1M Breit, Chun and Wahsweiler, Phys. Rev. 133 (1964) B404  
1964BU1D Burcham et al., 4B(1)/C212, Paris (1964)  
1964CA05 R.R. Carlson, E. Norbeck and V. Hart, Bull. Amer. Phys. Soc. 9 (1964) 419, DA9  
1964CA18 R.R. Carlson and M. Throop, Phys. Rev. 136 (1964) B630  
1964DA1B W.W. Daehnick and L.J. Denes, Phys. Rev. 136 (1964) B1325.  
1964DA1D Dar, Nucl. Phys. 55 (1964) 305



1964DE1C Deshpande, Fulbright and Verba, Nucl. Phys. 52 (1964) 457  
 1964DE1E Deshpande and Fulbright, UR 875 61 (1964)  
 1964DI1C Din, Unpublished Thesis, Rice Univ. (1964)  
 1964DI1D Din, Kuan and Bonner, Nucl. Phys. 50 (1964) 267  
 1964DO03 P.F. Donovan, J.F. Mollenauer and E.K. Warburton, Phys. Rev. 133 (1964) B113  
 1964EC03 S.F. Eccles, H.F. Lutz and J. Stevens, Bull. Amer. Phys. Soc. 9 (1964) 704, B9  
 1964EV1A Evans, NOL TR 64-46 (1964)  
 1964FL1D Flerov and Karnaukhov, Congres Int. de Phys. Nucl., Paris, 1964 (CRNS, Paris, 1964) 373  
 1964FO1A Fowler and Vogl, Lectures in Theor. Phys., Vol. VI (Univ. of Colorado Press, 1964) 379  
 1964GA1A Gardner and Yu, Nucl. Phys. 60 (1964) 49  
 1964GR1G Grieder, Phys. Rev. 133 (1964) B1483  
 1964GR1H Gromov et al., 4A(11)/C358, Paris (1964)  
 1964HA09 V.P. Hart, E. Norbeck and R.R. Carlson, Bull. Amer. Phys. Soc. 9 (1964) 430, EA6  
 1964HE18 I. Heertje, I. Delvenne, W. Nagel and A.H.W. Aten Jr., Physica 30 (1964) 1762  
 1964JO1A Jobes and McIntyre, Phys. Rev. 133 (1964) B893  
 1964KE1C Kelley and Henley, Phys. Lett. 10 (1964) 95  
 1964KO10 R. Kosiek, Z. Phys. 179 (1964) 544  
 1964KU05 H.-M. Kuan, T.W. Bonner and J.R. Risser, Nucl. Phys. 51 (1964) 481  
 1964KU06 H.-M. Kuan and J.R. Risser, Nucl. Phys. 51 (1964) 518  
 1964LI14 I. Lindgren, Perturbed Angular Correlations; Eds. E. Karlsson, E. Matthias and K. Siegbahn (Amsterdam, North-Holland Publ. Co., 1964) 379  
 1964LI1B Lindskog, Sundstrom and Sparrman, Perturbed Angular Correlations (Amsterdam, North-Holland Publ. Company, 1964) 411  
 1964LI1C Litherland, Nucl. Instrum. Meth. 28 (1964) 55  
 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550  
 1964MA25 B.S. Madsen and M. Vedelsby, Nucl. Phys. 55 (1964) 477  
 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed. W.H. Johnson Jr. (Springer Verlag, Vienna, 1964) 305  
 1964MO1G Moringo, Phys. Rev. 134 (1964) B1243  
 1964NE09 J.B. Nelson, E.L. Hudspeth and E.M. Bernstein, Phys. Rev. 136 (1964) B71

- 1964OS01 D.R. Osgood, J.R. Patterson and E.W. Titterton, Nucl. Phys. 60 (1964) 503
- 1964PA11 G. Paic, I. Slaus and P. Thomas, Phys. Lett. 9 (1964) 147
- 1964RI1A Ripka, 3A ( II )/C174, Paris (1964)
- 1964SA1D Satchler, Drisko and Bassel, Phys. Rev. 136 (1964) B637
- 1964SA1E Saint-Pierre, in Photographie Corpusculaire III, Ed. Demers (Presses Univ. de Montreal, 1964) 462
- 1964SC01 J. Schaitman, H.F. Lutz and S.F. Eccles, Bull. Amer. Phys. Soc. 9 (1964) 68, GC10
- 1964SC09 R.B. Schwartz, H.D. Holmgren, L.M. Cameron and A.R. Knudson, Phys. Rev. 134 (1964) B577
- 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
- 1964ST1J Stanojevic, Bull. Boris Kidrich Inst. Nucl. Sci. 15 (1964) 167
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964TA05 N.W. Tanner, G.C. Thomas and E.D. Earle, Nucl. Phys. 52 (1964) 29
- 1964TA1C Tanner and Earle, Phys. Rev. Lett. 13 (1964) 410
- 1964VA1D Valentin, Albouy, Cohen and Gusakov, J. Physique 25 (1964) 704
- 1964WE1A Webb et al., Bull. Amer. Phys. Soc. 9 (1964) 351
- 1965AJ01 F. Ajzenberg-Selove, J.W. Watson and R. Middleton, Phys. Rev. 139 (1965) B592
- 1965AL05 W.P. Alford, L.M. Blau and D. Cline, Nucl. Phys. 61 (1965) 368
- 1965AL19 T.K. Alexander, A.E. Litherland and C. Broude, Can. J. Phys. 43 (1965) 2310
- 1965AL1J Alevra et al., Stud. Cercetari Fiz. (Rumania) 17 (1965) 761; Phys. Abs. 36453 (1966)
- 1965AR07 K.P. Artemov, V.Z. Goldberg, B.I. Islamov, V.P. Rudakov and I.N. Serikov, Yad. Fiz. 1 (1965) 1019; Sov. J. Nucl. Phys. 1 (1965) 726
- 1965BE1B Becker and McIntyre, Phys. Rev. 138 (1965) B339
- 1965BO14 R. Bock, M. Grosse-Schulte, W. Von Oertzen and R. Rudel, Phys. Lett. 18 (1965) 45
- 1965BO42 M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, H. Buttner, A. Lindner and H. Meldner, Nucl. Phys. 63 (1965) 438
- 1965BR1B Brill, Yad. Fiz. 1 (1965) 55; Sov. J. Nucl. Phys. 1 (1965) 37
- 1965BR1F Breit, Ann. Phys. 34 (1965) 377
- 1965BU05 A. Bussiere, N.K. Glendenning, B.G. Harvey, J. Mahoney, J.R. Meriwether and D.J. Horen, Phys. Lett. 16 (1965) 296
- 1965CA1B Caldwell et al., Phys. Rev. Lett. 15 (1965) 976
- 1965CO25 S. Cohen and D. Kurath, Nucl. Phys. 73 (1965) 1; Erratum Nucl. Phys. 89 (1966) 707

1965DE1A Detraz, Cahiers Phys. 19 (1965) 359  
 1965DE1G Degtyarev, Atomnaya Energiya 19 (1965) 456  
 1965DE24 V.P. Denisov and L.A. Kulchitskii, Yad. Fiz. 2 (1965) 70; Sov. J. Nucl. Phys. 2 (1966) 48  
 1965DI07 G.U. Din and J.L. Weil, Nucl. Phys. 73 (1965) 161  
 1965DI1E Dickens, Drisko, Perey and Satchler, Phys. Lett. 15 (1965) 337  
 1965DO1B Donaldson, Passell, Bartolini and Groves, Phys. Rev. 138 (1965) B1116  
 1965EN01 J.B.A. England and B.L. Reece, Nucl. Phys. 72 (1965) 449  
 1965FA1B Fabre de La Ripelle, Prog. Theor. Phys. 33 (1965) 38  
 1965FI05 P. Fintz, Ann. Phys. (Paris) 10 (1965) 435  
 1965FR1E Frahn and Wiechers, Nucl. Phys. 74 (1965) 65  
 1965FU16 H.W. Fulbright, W.P. Alford, O.M. Bilaniuk, V.K. Deshpande and J.W. Verba, Nucl. Phys. 70 (1965) 553  
 1965GA1B Gaedke, Toth and Williams, Phys. Rev. 140 (1965) B296  
 1965GA1D Gaponov, Yad. Fiz. 2 (1965) 1002  
 1965GA1G Garber and Shrader, Bull. Amer. Phys. Soc. 10 (1965) 510  
 1965GI1B Giraud, Nucl. Phys. 71 (1965) 373  
 1965GL04 R.N. Glover and A.D.W. Jones, Phys. Lett. 16 (1965) 69  
 1965GR1H Grechukhin, Nucl. Phys. 62 (1965) 273  
 1965GR1R Gromov et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P-2184 (1965)  
 1965GR1T Grimeland, Kjellsby and Vines, Phys. Rev. 137 (1965) B878  
 1965GU1A Guiasu and Micu, Rev. Physique (Rumania) 10 (1965) 555  
 1965HA1G Havlicek and Modesto, Energ. Nucl. 12 (1965) 77  
 1965HE1B Henkel, Bull. Amer. Phys. Soc. 10 (1965) 601  
 1965HI1A Hiebert, McIntyre and Couch, Phys. Rev. 138 (1965) B346  
 1965HU1D Hull and Shakin, Phys. Lett. 19 (1965) 506  
 1965IC1A Ichimura and Yazaki, Nucl. Phys. 63 (1965) 401  
 1965JA09 H.E. Jackson, A.I. Namenson and G.E. Thomas, Phys. Lett. 17 (1965) 324  
 1965LA09 J.M. Lacabra, D.R. Tilley, N.R. Roberson and R.M. Williamson, Nucl. Phys. 68 (1965) 273  
 1965MA1K Marion, Nucl. Phys. 68 (1965) 463  
 1965MA1T Mahaux, Nucl. Phys. 67 (1965) 358

1965MA1V Markowitz and Pape, Bull. Amer. Phys. Soc. 10 (1965) 602  
 1965MA45 J.M. Maison, M. Langevin and J.M. Loiseaux, Phys. Lett. 19 (1965) 308  
 1965MO13 R.C. Morrison, J.R. Stewart and J.S. O'Connell, Phys. Rev. Lett. 15 (1964) 367  
 1965NE1D Newton, Phys. Lett. 17 (1965) 132  
 1965PE1H Perez Dominguez, Jen 143 DF/1-46 (1965)  
 1965RO1N Rose and Lopes, Phys. Lett. 18 (1965) 130  
 1965RO22 L. Rosen, J.G. Beery, A.S. Goldhaber and E.H. Auerbach, Ann. Phys. 34 (1965) 96  
 1965SC1D Schapira, Ann. Physique 10 (1965) 743  
 1965SE05 R. Seltz, C. Gerardin, M. Wery and D. Magnac-Valette, Compt. Rend. 261 (1965) 391  
 1965SH1E Shapiro and Timashev, Yad. Fiz. 2 (1965) 459  
 1965SI13 R.H. Siemssen, M. Cosack and R. Felst, Nucl. Phys. 69 (1965) 227  
 1965SL1C Slee, Bull. Amer. Phys. Soc. 10 (1965) 461  
 1965TA07 T. Takemiya, Prog. Theor. Phys. 34 (1965) 433  
 1965TS1A Tsenter and Silin, Atomnaya Energ. 19 (1965) 48  
 1965VA05 V. Valkovic, G. Paic, I. Slaus, P. Tomas, M. Cerineo and G.R. Satchler, Phys. Rev. 139 (1965) B331  
 1965VA1E Valentin, Nucl. Phys. 62 (1965) 81  
 1965WA03 E.K. Warburton, K.W. Jones, D.E. Alburger, C. Chasman and R.A. Ristinen, Phys. Rev. Lett. 14 (1965) 146  
 1965WA06 E.K. Warburton, J.S. Lopes, R.W. Ollerhead, A.R. Poletti and M.F. Thomas, Phys. Rev. 138 (1965) B104  
 1965WA16 E.K. Warburton, J.W. Olness and D.E. Alburger, Phys. Rev. 140 (1965) B1202  
 1965WA17 E.K. Warburton, P.D. Parker and P.F. Donovan, Phys. Lett. 19 (1965) 397  
 1965WI03 E.J. Winhold, R.H. Augustson, N.N. Kaushal, H.A. Medicus, W.R. Moyer and P.F. Yergin, Bull. Amer. Phys. Soc. 10 (1965) 95, HD6  
 1965WI1A Williams, Gaedke and Toth, Bull. Amer. Phys. Soc. 10 (1965) 443  
 1965ZA1B Zamick, Phys. Lett. 19 (1965) 580  
 1966AG1A Agee and Rosen, LA-3538-MS (1966)  
 1966AG1B Aguilar, de la Rubia, Sanchez and Martinez, An. de Fisica Y Quim. 62 (1966) 279  
 1966AL12 D.E. Alburger and K.W. Jones, Phys. Rev. 149 (1966) 743  
 1966AL18 D.E. Alburger and E.K. Warburton, Phys. Rev. 152 (1966) 914  
 1966AV1A Avignon, Rosier and Deschamps, J. Phys. C1-40 (1966)  
 1966BA1F Baumgartner et al., Helv. Phys. Acta 39 (1966) 575

1966BE1M Bercaw, Boschitz and Vincent, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 334

1966BL01 R.S. Blake, D.J. Jacobs, J.O. Newton and J.P. Schapira, Nucl. Phys. 77 (1966) 254

1966BO1R Bohigas, J. Phys. C1-39 (1966)

1966BO22 R. Bock, M. Grosse-Schulte and W. Von Oertzen, Phys. Lett. 22 (1966) 456

1966BR09 R.I. Brown, Nucl. Phys. 78 (1966) 492

1966BR13 K.H. Bray, J. Nurzynski and W.P. Bourke, Phys. Lett. 21 (1966) 536

1966BU1B Buttle and Goldfarb, Nucl. Phys. 78 (1966) 409

1966CA1H Catala, Garcia and Perez, An. Real. Soc. Espan. Fis. y Quim. A61 (1966) 357

1966CI01 S.D. Cirilov, J.O. Newton and J.P. Schapira, Nucl. Phys. 77 (1966) 472

1966CS1B Csikai and Nagy, Acta Phys. Acad. Sci. Hung. 21 (1966) 303

1966CS1C Csikai, Antwerp 1965 Neutron Conf. (North Holland, 1966) 537

1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097

1966DR02 L. Drigo, C. Manduchi, G.C. Nardelli, M.T. Russo-Manduchi, G. Torielli and G. Zannoni, Nuovo Cim. B45 (1966) 206

1966DU1B Duggan et al., Bull. Amer. Phys. Soc. 11 (1966) 831

1966EC1B Eccles, Lutz and Rohn, Bull. Amer. Phys. Soc. 11 (1966) 735

1966ED1A J.A. Edgington and B. Rose, Nucl. Phys. 89 (1966) 523

1966EL08 F. El-Batanoni and A.A. Kresnin, Nucl. Phys. 89 (1966) 577

1966EV01 A.E. Evans, B. Brown and J.B. Marion, Phys. Rev. 149 (1966) 863

1966FO1D Fowler and Johnson, Bull. Amer. Phys. Soc. 11 (1966) 510; ORNL P-2026 (1966)

1966FO1E Fowler, Johnson and Kernell, Bull. Amer. Phys. Soc. 11 (1966) 653

1966GA04 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 141 (1966) 996

1966GA08 A. Gallman, P. Fintz, J.B. Nelson and D.E. Alburger, Phys. Rev. 147 (1966) 753

1966GA09 A. Gallman, P. Fintz and P.E. Hodgson, Nucl. Phys. 82 (1966) 161

1966GA19 A. Gallmann, F. Haas and N. Balaux, Phys. Rev. 151 (1966) 735

1966GA25 G.T. Garvey and I. Kelson, Phys. Rev. Lett. 16 (1966) 197

1966GL01 R.N. Glover and A.D.W. Jones, Nucl. Phys. 84 (1966) 673

1966GO15 S. Gorodetzky, R.M. Freeman, A. Gallman and F. Hass, Phys. Rev. 149 (1966) 801

1966GO1E Gorodetzky et al., Nucl. Instrum. Meth. 42 (1966) 269

1966GO1J Gorodetzky, Gallmann and Fintz, J. Phys. C1-91 (1966)

1966GR1A Gruhn and Kashy, Bull. Amer. Phys. Soc. 11 (1966) 471

- 1966HA19 B.G. Harvey, J.R. Meriwether, J. Mahoney, A. Bussiere de Nercy and D.J. Horen, Phys. Rev. 146 (1966) 712
- 1966HA21 R.L. Hahn and E. Ricci, Phys. Rev. 146 (1966) 650
- 1966HA30 O. Hausser, R.D. Gill, J.S. Lopes and H.J. Rose, Nucl. Phys. 84 (1966) 683
- 1966HA31 O. Hausser, H.J. Rose, J.S. Lopes and R.D. Gill, Phys. Lett. 22 (1966) 604
- 1966HI1C B. Hird, Nucl. Phys. 86 (1966) 268
- 1966HO1D Hodgson, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 71
- 1966JE1B Jessen, Bormann, Dreyer and Neuert, Nucl. Data 1 (1966) 103
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KL06 A.P. Klyucharev and Y.I. Titov, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 435; Bull. Acad. Sci. USSR Phys. 30 (1966) 443
- 1966KL1F Klyucharev, Titov and Vipirailenko, Izv. Vys. Uch. Zaved. Fiz. 3 (1966) 104
- 1966KO1G Komar, Denisov and Kulchitskii, Dokl. Akad. Nauk SSSR 169 (1966) 1307
- 1966KU12 I.V. Kurdyumov, S.H.El Samarai, Y.F. Smirnov and K.V. Shitikova, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 292; Bull. Acad. Sci. USSR Phys. 30 (1966) 297
- 1966LA04 T. Lauritsen and F. Ajzenberg-Selove, Nucl. Phys. 78 (1966) 1
- 1966LI07 K.P. Lieb, Nucl. Phys. 85 (1966) 461
- 1966LO02 J.S. Lopes, O. Hausser, H.J. Rose, A.R. Poletti and M.F. Thomas, Nucl. Phys. 76 (1966) 223
- 1966LO1N Louis and Nussbaum, Helv. Phys. Acta 39 (1966) 11
- 1966MA02 A.M. MacLeod and J.M. Reid, Proc. Phys. Soc. (London) (London) 87 (1966) 437
- 1966MA04 G.S. Mani and G.C. Dutt, Nucl. Phys. 78 (1966) 613; Erratum Nucl. Phys. A119 (1968) 691
- 1966MA1A Marion, Ludemann and Roos, Bull. Amer. Phys. Soc. 11 (1966) 332
- 1966MA1R J.H. Manley and W.E. Stein, Phys. Rev. 144 (1966) 956
- 1966MA1V H.A. Mavromatis and L. Zamick, Phys. Lett. 20 (1966) 171
- 1966MA2K Martin, Kernell and Fowler, Bull. Amer. Phys. Soc. 11 (1966) 808
- 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
- 1966ME14 D.F. Measday and J.N. Palmieri, Nucl. Phys. 85 (1966) 129
- 1966MI1F Migdal, Proc. Int. School Enrico Fermi, Course 36; Ed. C. Bloch (Academic Press, 1966) 171
- 1966OL1C Olariu, Acad. Repub. Pop. Romine. Inst. de Fiz. Atom. Rept. IS 25 (1966)
- 1966OW01 R.O. Owens and J.E.E. Baglin, Phys. Rev. Lett. 17 (1966) 1268



1966PE04 D. Pelte, B. Povh and W. Scholz, Nucl. Phys. 78 (1966) 241  
 1966PH1B Phillips, NASA CR 94632 (1966)  
 1966PO09 A.M. Poskanzer, S.W. Cospers, E.K. Hyde and J. Cerny, Phys. Rev. Lett. 17 (1966) 1271  
 1966PO11 A.R. Poletti, J.W. Olness and E.K. Warburton, Phys. Rev. 151 (1966) 812  
 1966PO1E Poth and Bromley, Bull. Amer. Phys. Soc. 11 (1966) 317  
 1966PR1B Prusser, 2nd Symp. on the Struct. of Low-Medium Mass Nucl., April 1966 (1966) 38  
 1966RE1D Reeves, J. Phys. C1-28 (1966)  
 1966RI1F Ripka, Lect. in Theor. Phys., Vol. VIII C (Univ. of Colorado Press, Boulder, 1966) 237  
 1966RO1B Rosen, Proc. 2nd Int. Symp. on Polariz. Phenom. of Nucleons, Karlsruhe, 1965 (Basel, Birkhauser Verlag, 1966) 253  
 1966RO1P H.J. Rose, J.S. Lopes and W. Greiner, Phys. Lett. 19 (1966) 686  
 1966RO1R Rosen, Antwerp 1965 Neutron Conf. (North Holland, 1966) 379  
 1966RO1U H.J. Rose and J.S. Lopes, Phys. Lett. 22 (1966) 601  
 1966RO1V B.A. Robson, Nucl. Phys. 86 (1966) 649  
 1966SC12 J.P. Schapira, J.O. Newton, R.S. Blake and D.J. Jacobs, Nucl. Phys. 80 (1966) 565  
 1966SC21 W. Scobel, R.W. Fink and M. Bormann, Z. Phys. 197 (1966) 124  
 1966SC22 J.J. Schwartz, W.P. Alford, L.M. Blau and D. Cline, Nucl. Phys. 88 (1966) 539  
 1966SE1D Seltz et al., J. Phys. C1-148 (1966)  
 1966SH1A Shute and Brown, Bull. Amer. Phys. Soc. 11 (1966) 471  
 1966SH1F I.S. Shapiro and S.F. Timashev, Nucl. Phys. 79 (1966) 46  
 1966SO05 M. Soga, Nucl. Phys. 89 (1966) 697  
 1966ST05 J.D. Steben and M.K. Brussel, Phys. Rev. 146 (1966) 780  
 1966TY01 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson and T. Berggren, Nucl. Phys. 79 (1966) 321; Erratum Nucl. Phys. A119 (1968) 692  
 1966WA08 E.K. Warburton and J.W. Olness, Phys. Rev. 147 (1966) 698  
 1966WA1E Warburton, F. S. U. Isobaric Spin Conf. (1966) 90  
 1966WA1K G.E. Walker, Phys. Rev. 151 (1966) 745  
 1966WE1B Weinberg, Antwerp 1965 Neutron Conf. (North Holland, 1966) 37  
 1966WI1E D.H. Wilkinson and M.E. Mafethe, Nucl. Phys. 85 (1966) 97  
 1967AD1F Adelberger, Cocke and Davids, Bull. Amer. Phys. Soc. 12 (1967) 1194  
 1967AM1H Amiet, Ebenhoh and Huguenin, Helv. Phys. Acta 40 (1967) 283

- 1967AU1A Audouze, Epherre and Reeves, High Energy Nucl. Reactions in Astrophys., Ed. B.S.P. Shen (W.A. Benjamin, 1967) 255
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
- 1967BA03 P.W. Bauer, J.D. Anderson, H.F. Lutz, C. Wong, J.W. McClure and B.A. Pohl, Nucl. Phys. A93 (1967) 673
- 1967BA13 G.C. Ball and J. Cerny, Phys. Rev. 155 (1967) 1170
- 1967BE09 H. Beaumevieille, M. Lambert, M. Yaker and A. Amokrane, Nuovo Cim. B47 (1967) 139
- 1967BE1F Beery, Harper, Stovall and Rosen, Los Alamos Sci. Lab. Rept. LA-3788 (1967)
- 1967BI11 P.G. Bizzeti, A.M. Bizzeti-Sona, S. Kalbitzer and B. Povh, Nucl. Phys. A104 (1967) 577
- 1967BO1T Bohigas, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 940
- 1967BR1M Breit, Polak and Torchia, Phys. Rev. 161 (1967) 993
- 1967BR1Q G. Breit, Proc. Nat. Acad. Sci. 67 (1967) 849
- 1967CA1C Caldwell, UCRL 50287 (1967)
- 1967CA1P Caldwell, Fultz and Bramblett, Bull. Amer. Phys. Soc. 12 (1967) 197
- 1967CH15 N.S. Chant, P.S. Fisher and D.K. Scott, Nucl. Phys. A99 (1967) 669
- 1967CH19 C. Chasman, K.W. Jones, R.A. Ristinen and D.E. Alburger, Phys. Rev. 159 (1967) 830
- 1967CH35 R. Chiba, Nucl. Sci. App. 3 (1967) 36
- 1967CL1C Cloud, Leonard, Gibson and Wells, Bull. Amer. Phys. Soc. 12 (1967) 894
- 1967CO1D Commins, Ann. Rev. Nucl. Sci. 17 (1967) 33
- 1967CO1R Coppola and Knitter, Eur 3492.E (1967)
- 1967CO32 S. Cohen and D. Kurath, Nucl. Phys. 101 (1967) 1
- 1967CS02 J. Csikai and G. Peto, Acta Phys. Acad. Sci. Hung. 23 (1967) 87
- 1967CS1A J. Csikai and S. Nagy, Nucl. Phys. A91 (1967) 222
- 1967DA1E Dalidtshik and Saiasov, Proc. Prob. Symp. on Nucl. Phys., Tbilishi, April 1967 (Moscow, 1967) 287
- 1967DE14 D. Dehnhard, D.S. Gemmell and Z. Vager, Nucl. Phys. A104 (1967) 202
- 1967DE1K Dehnhard and Siemssen, Bull. Amer. Phys. Soc. 12 (1967) 17
- 1967EL03 J.P. Elliott, H.A. Mavromatis and E.A. Sanderson, Phys. Lett. B24 (1967) 358
- 1967EV02 A.E. Evans, Phys. Rev. 155 (1967) 1047
- 1967FA1A W.M. Fairbairn, Nucl. Phys. A90 (1967) 135



- 1967FI1E Firk, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 352
- 1967FU1A Funsten and Holt, Bull. Amer. Phys. Soc. 12 (1966) 1176
- 1967GO07 S. Gorodetzky, R.M. Freeman, A. Gallmann, F. Haas and B. Heusch, Phys. Rev. 155 (1967) 1119
- 1967GR1K Gradsztajn, High Energy Nucl. Reactions in Astrophys., Ed. B.S.P. Shen (W.A. Benjamin, 1967) 247
- 1967GR1L Gray, Fortune, Trost and Fletcher, Bull. Amer. Phys. Soc. 12 (1967) 34
- 1967HE1A Hensley, Astrophys. J. 147 (1967) 818
- 1967HI06 J.C. Hiebert, E. Newman and R.H. Bassel, Phys. Rev. 154 (1967) 898
- 1967IV1B Ivascu, Dumitrescu and Semenescu, Rev. Roum. Phys. 12 (1967) 279
- 1967KO1G Kobzev et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P6 3314 (1967)
- 1967KR1F Krivopustov, Sizov, Shirmer and Oehler, P15-3504, Dubna (1967)
- 1967KU1E D. Kurath and R.D. Lawson, Phys. Rev. 161 (1967) 915
- 1967KU1M Kuan, Hanna and Hasinoff, Bull. Amer. Phys. Soc. 12 (1967) 52
- 1967LA05 M. Lambert and M. Durand, Phys. Lett. B24 (1967) 287
- 1967LA10 M. Lambert and M. Durand, J. Phys. (Paris) 28 (1967) 349
- 1967LA11 B. Lawergren and I.V. Mitchell, Nucl. Phys. A98 (1967) 481
- 1967LI06 R.H. Lindsay and J.J. Veit, Phys. Rev. 157 (1967) 933
- 1967LI1B Lingenfelter and Ramaty, High Energy Nucl. Reactions in Astrophys., Ed. B.S.P. Shen (W.A. Benjamin, 1967) 99
- 1967LO03 P. Loncke and J. Pradal, Nuovo Cim. B48 (1967) 457
- 1967MA1M Marr, Kuenhold and Donoghue, Bull. Amer. Phys. Soc. 12 (1967) 501
- 1967ME1N Meier, Thomason and Walter, Bull. Amer. Phys. Soc. 12 (1967) 1197
- 1967MO21 J. Mosner, G. Schmidt and J. Schintlmeister, Nucl. Phys. A103 (1967) 238
- 1967MU12 S. Mubarakmand and B.E.F. Macefield, Nucl. Phys. A98 (1967) 82
- 1967NE06 J.B. Nelson and W.R. Smith, Nucl. Phys. A96 (1967) 671
- 1967NE1D Nemirovskii, Sov. J. Nucl. Phys. 4 (1967) 334
- 1967PA05 J.C. Parikh and N. Ullah, Nucl. Phys. A99 (1967) 529
- 1967PA1N A. Pasquarelli, Nucl. Phys. A93 (1967) 218
- 1967PE1D J.-L. Perrenoud and E. Sheldon, Nucl. Phys. A102 (1967) 105
- 1967PH03 G.W. Phillips, F.C. Young and J.B. Marion, Phys. Rev. 159 (1967) 891
- 1967PO13 J.E. Poth, J.C. Overley and D.A. Bromley, Phys. Rev. 164 (1967) 1295

- 1967PO1E Poth and Bromley, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 94
- 1967PO1J A.R. Poletti, E.K. Warburton and D. Kurath, Phys. Rev. 155 (1967) 1096
- 1967RE01 D. Rendic, Nucl. Phys. A91 (1967) 604
- 1967RO1K Robertson, Aust. J. Phys. 20 (1967) 489
- 1967SC27 L.A. Schaller, R.S. Thomason, N.R. Roberson, R.L. Walter and R.M. Drisko, Phys. Rev. 163 (1967) 1034
- 1967SC29 J.P. Schiffer, G.C. Morrison, R.H. Siemssen and B. Zeidman, Phys. Rev. 164 (1967) 1274
- 1967SE08 W.A. Seale, Phys. Rev. 160 (1967) 809
- 1967SH14 V.S. Shirley, UCRL-17990 (1967)
- 1967SO1B Soltesz et al., Bull. Amer. Phys. Soc. 12 (1967) 1198
- 1967ST21 B. Stepanic, R. Popic and M. Aleksic, Phys. Rev. Lett. 19 (1967) 1137
- 1967TH05 G.E. Thomas, D.E. Blatchley and L.M. Bollinger, Nucl. Instrum. Meth. 56 (1967) 325
- 1967VO1A Volkov, Proc. Problem Symp. on Nucl. Phys., Tbilist, Apr. 1967 (Moscow, 1967) 226
- 1967VO1B Vogt, Int. Nucl. Phys. Conf., Gatlinburg 1966 (Academic Press, 1967) 748
- 1967WA1C Warburton, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 43
- 1967WE1B West, ORNL TM 1926 (1967)
- 1967YO1C Young, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 109
- 1967ZH1A Zhusupov, Karapetian and Eramzhian, Joint Inst. Nucl. Res., Lab. Theor. Phys., USSR Rept. No. P4 3177 (1967)
- 1968AJ02 F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A114 (1968) 1
- 1968AL1C Aldzhauakihiri, Spasskii, Teplov and Fateeva, Sov. J. Nucl. Phys. 6 (1968) 180
- 1968BA1E Ball, UCRL-18263 (1968)
- 1968BA2J Bassel and Drisko, Proc. Symp. on Direct Reactions with  $^3\text{He}$ , IPCR, Japan, Sept. 1967 (1968) 13
- 1968BA2L Baglin and Thompson, Tokyo (1968) 388
- 1968BA42 A.R. Barnett, Nucl. Phys. A120 (1968) 342
- 1968BE14 G.A. Beer, P. Brix, H.-G. Clerc and B. Laube, Phys. Lett. B26 (1968) 506
- 1968BI1C Bizzeti, Suppl. Nuovo Cim. 6 (1968) 664
- 1968BO14 W. Bohne, H. Homeyer, H. Morgenstern and J. Scheer, Nucl. Phys. A113 (1968) 97
- 1968BO36 F. Boreli, P.N. Shrivastava, B.B. Kinsey and V.C. Mistry, Phys. Rev. 174 (1968) 1221

1968CA1J Campbell, Harvey and Slaughter, Bull. Amer. Phys. Soc. 13 (1968) 1423  
 1968CO04 R.L.A. Cottrell, J.C. Lisle and J.O. Newton, Nucl. Phys. A109 (1968) 288  
 1968CO1W Conde, Bergqvist and Nystrom, Neutron Cross Sect. Tech., NBS Special Pub. 299 (1968) 763  
 1968COZV P.M. Cockburn, R.W. Krone and H.E. Siefken, Bull. Amer. Phys. Soc. 13 (1968) 1423, DC4  
 1968DA1F J.C. Davis and H.H. Barschall, Phys. Lett. B27 (1968) 636  
 1968DA1N Dangle, Duncan, Duggan and Miller, Bull. Amer. Phys. Soc. 13 (1968) 608  
 1968DA1Q Dally, Croissiaux and Schweitz, Bull. Amer. Phys. Soc. 13 (1968) 607  
 1968DE13 N. de Takacsy, Can. J. Phys. 46 (1968) 2091  
 1968DU1F Duorah and Duorah, Ind. J. Pure Appl. Phys. 6 (1968) 389  
 1968EL1A Elliott, 3rd Symp. on the Struct. of Low-Medium Mass Nucl., Ed. Davidson (Univ. Press of Kansas, 1968) 48  
 1968FA1A J.C. Faivre, H. Krivine and A.M. Papiou, Nucl. Phys. A108 (1968) 508  
 1968FO06 H.T. Fortune, T.J. Gray, W. Trost and N.R. Fletcher, Phys. Rev. 173 (1968) 1002  
 1968FO1A Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299 (1968) 1  
 1968GA03 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 167 (1968) 957  
 1968GA13 M. Gaillard, R. Bouche, L. Feuvrais, P. Gaillard, A. Guichard, M. Gusakow, J.L. Leonhardt and J.-R. Pizzi, Nucl. Phys. A119 (1968) 161  
 1968GA1C Gaillard, Univ. Lyon, Rept. No. Lycen 6828 (1968)  
 1968GA1M E. Gadioli and L. Zetta, Phys. Rev. 167 (1968) 1016  
 1968GE04 B. Geoffrion, N. Marty, M. Morlet, B. Tatischeff and A. Willis, Nucl. Phys. A116 (1968) 209  
 1968GI01 R.D. Gill, J.S. Lopes, B.C. Robertson, R.A.I. Bell and H.J. Rose, Nucl. Phys. A106 (1968) 678  
 1968GI11 R.D. Gill, J.S. Lopes, O. Hausser and H.J. Rose, Nucl. Phys. A121 (1968) 209  
 1968GR14 R.C. Greenwood, Phys. Lett. B27 (1968) 274  
 1968GRZY Greenwood, Proc. Conf. Slow-Neutron-Capture Gamma-Ray Spectr., Argonne, Ill., 1966; Ed. F.E.Throw, ANL-7282 (1968) 303  
 1968HA15 J.S. Hanna and M.A. Nagarajan, Nucl. Phys. A113 (1968) 412  
 1968HA1V Hansen, Anderson, Stelts and Wong, Neutron Cross Sect. Tech., NBS Special Pub. 299 (1968) 225  
 1968HA27 W.R. Harris and J.C. Armstrong, Phys. Rev. 171 (1968) 1230  
 1968HE12 J.D. Henderson, E.L. Hudspeth and W.R. Smith, Phys. Rev. 172 (1968) 1058

1968HI1J Hinderliter and Lochstet, Bull. Amer. Phys. Soc. 13 (1968) 606  
 1968HO1H Horie and Hsieh, Tokyo (1968) 51  
 1968HU1A R.L. Hutson, S. Hayakawa, M. Chabre, J.J. Kraushaar, B.W. Ridley and E.T. Boschitz,  
 Phys. Lett. B27 (1968) 153  
 1968IW1A Iwao and Kawakami, Prog. Theor. Phys. 40 (1968) 1046  
 1968JA1M Jacquot et al., Compt. Rend. B266 (1968) 1286  
 1968JO1F Johnson et al., Neutron Cross Sections Tech., NBS Special Pub. 299 (1968) 851  
 1968JO1H Jones and Toms, Bull. Amer. Phys. Soc. 13 (1968) 718  
 1968JU1B Jung et al., Compt. Rend. 266 (1968) 1154  
 1968KU1F Kuan and O'Connell, Bull. Amer. Phys. Soc. 13 (1968) 85  
 1968LE01 J.K.P. Lee, S.K. Mark, P.M. Portner and R.B. Moore, Nucl. Phys. A106 (1968) 357  
 1968LI1A Li and Mark, Bull. Amer. Phys. Soc. 13 (1966) 679  
 1968LI1L Liebenauer, Silverstein, Kibler and Koral, Bull. Amer. Phys. Soc. 13 (1968) 1388  
 1968MA2B Manakos, Z. Phys. 214 (1968) 57  
 1968MA2G R.M. May, Phys. Lett. B26 (1968) 482  
 1968ME08 R.A. Mendelson Jr. and R.T. Carpenter, Phys. Rev. 166 (1968) 988  
 1968NA1F Nakamura, Tokyo (1968) 228  
 1968PA1F J.Y. Park, Nucl. Phys. A111 (1968) 433  
 1968PU1A H.G. Pugh, Phys. Rev. Lett. 20 (1968) 601  
 1968RI1T Ritter and Parson, Bull. Amer. Phys. Soc. 13 (1968) 607  
 1968RO1E Rohl, Z. Phys. 215 (1968) 56  
 1968SC1B Schevchenko, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc.  
 Jpn. 24 (1968) 397  
 1968SH08 A.P. Shukla and G.E. Brown, Nucl. Phys. A112 (1968) 296  
 1968SH11 P.N. Shrivastava, F. Boreli and B.B. Kinsey, Phys. Rev. 169 (1968) 842  
 1968SI1E Siefken, Cockburn and Krone, Bull. Amer. Phys. Soc. 13 (1968) 1423  
 1968SI1F Siefken, COO-1120-75 (1968)  
 1968ST06 A.P. Stamp and M.B. Spencer, Nucl. Phys. A111 (1968) 353  
 1968ST10 C.E. Steerman and F.C. Young, Phys. Lett. B27 (1968) 8  
 1968ST19 T. Stammbach, R.S. Thomason, J. Taylor and Jr., R.L. Walter, Phys. Rev. 174 (1968)  
 1119  
 1968TU02 P.M. Tutakin, Yad. Fiz. 8 (1968) 661; Sov. J. Nucl. Phys. 8 (1969) 383  
 1968WA04 G.J. Wagner, Phys. Lett. B26 (1968) 429

- 1968WE15 H.R. Weller, N.R. Roberson and D.R. Tilley, Nucl. Phys. A122 (1968) 529
- 1968WO1C S.S.M. Wong, Nucl. Phys. A120 (1968) 625
- 1968WO1D Wong et al., Tokyo (1968) 249
- 1968YA1E F.B. Yano, Nucl. Phys. A118 (1968) 592
- 1968ZH05 M.A. Zhusupov, O. Lkhagva and I. Rotter, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1714; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 1579
- 1968ZH06 M.Z. Zhusupov and R.A. Eramzhyan, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 2060; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 1894
- 1968ZH1B Zhusupov, Karapetyan and Eramzhyan, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 332
- 1968ZU02 A.P. Zuker, B. Buck and J.B. McGrory, Phys. Rev. Lett. 21 (1968) 39
- 1969AL04 W.P. Alford and K.H. Purser, Nucl. Phys. A132 (1969) 86
- 1969AL1H Alsmiller, Wachter and Moran, Nucl. Sci. Eng. 36 (1969) 291
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A137 (1969) 348
- 1969BA05 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz and P. Radvanyi, Nucl. Phys. A126 (1969) 60
- 1969BA06 G.C. Ball and J. Cerny, Phys. Rev. 177 (1969) 1466
- 1969BA1N Bahcall and Fowler, Astrophys. J. 157 (1969) 659
- 1969BA1P Bacon et al., Acta Cryst. A25 (1969) 391
- 1969BA2U Ballini, Delaunay, Fouan and Tellez, Addendum to Contrib., Montreal (1969) 6
- 1969BE08 H. Beaumevieille, M. Lambert, M. Yaker, A. Amokrane and Nguyen van Sen, Nucl. Phys. A125 (1969) 568
- 1969BO13 W. Bohne, H. Homeyer, H. Lettau, H. Morgenstern, J. Scheer and F. Sichelschmidt, Nucl. Phys. A128 (1969) 537
- 1969BR07 K.H. Bray and J. Nurzynski, Nucl. Phys. A127 (1969) 622
- 1969BR1D Bromly, Proc. Enrico Fermi School of Phys., Course XL, Lake Como 1967 (Academic Press, 1969) 242
- 1969BR1G Bromley, Proc. Int. Conf., Montreal (1969) 147
- 1969BR30 O.D. Brill, A.D. Vongai and A.A. Ogloblin, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 615; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 567
- 1969CH1R M. Chemtob, Nucl. Phys. A123 (1969) 449
- 1969CO15 G.W. Cole Jr., F.W.K. Firk and T.W. Phillips, Phys. Lett. B30 (1969) 91

- 1969DE16 J. Dey, J.P. Elliott, A.D. Jackson, H.A. Mavromatis, E.A. Sanderson and B. Singh, Nucl. Phys. A134 (1969) 385
- 1969DE1H Debenham, Griffith, Irshad and Roman, Contrib., Montreal (1969) 274
- 1969DI1B Dickens and Perey, Nucl. Sci. Eng. 36 (1969) 280
- 1969DO04 H. Doubre, D. Royer, M. Arditì, L. Bimbot, N. Frascaria, J.P. Garron and M. Riou, Phys. Lett. B29 (1969) 355
- 1969EL1B Elliott, Proc. Int. Conf., Montreal (1969) 277
- 1969EN03 J.B.A. England, R.G. Harris, L.H. Watson, D.H. Worledge and J.E. Evans, Phys. Lett. B30 (1969) 476
- 1969ET01 K. Etoh, T. Murata, N. Kawai, R. Chiba and S. Takayanagi, J. Phys. Soc. Jpn. 26 (1969) 1335, and Private Communication (1969)
- 1969FO02 H.T. Fortune, T.J. Gray, W. Trost and N.R. Fletcher, Phys. Rev. 179 (1969) 1033
- 1969FO1D Fowler, Contemp. Phys. 1 (1969) 359
- 1969FR20 D.E. Frederick, R.J.J. Stewart and R.C. Morrison, Phys. Rev. 186 (1969) 992
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA05 A. Gallmann, F. Jundt, E. Aslanides and D.E. Alburger, Phys. Rev. 179 (1969) 921
- 1969GA11 P. Gaillard, R. Bouche, L. Feuvrais, M. Gaillard, A. Guichard, M. Gusakov, J.L. Leonhardt and J.R. Pizzi, Nucl. Phys. A131 (1969) 353
- 1969GI1B Gillet, Proc. Int. Conf., Montreal (1969) 483
- 1969GL07 Y.A. Glukhov, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta and D.N. Stepanov, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 601; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 554
- 1969GL1D Glenn, Zafiratos and Zaidins, Bull. Amer. Phys. Soc. 14 (1969) 1227
- 1969GO14 V. Gomes Porto, N. Ueta, R.A. Douglas, O. Sala, D. Wilmore, B.A. Robson and P.E. Hodgson, Nucl. Phys. A136 (1969) 385
- 1969GO1R Golestaneh, Schmittroth and Tobocman, Bull. Amer. Phys. Soc. 14 (1969) 572
- 1969GU1M V.N. Guman, L.A. Sliv and Yu.I. Kharitonov, Phys. Lett. B28 (1969) 575
- 1969HA1G Hanna, Proc. Int. Conf., Montreal (1969) 443
- 1969HA49 F. Haas, B. Heusch, A. Gallmann and D.A. Bromley, Phys. Rev. 188 (1969) 1625
- 1969HO1F Honsaker, McDonald, Neilson and Hsu, Contrib., Montreal (1969) 299
- 1969HO1T Horowitz, McConnell and Ssengabi, Bull. Amer. Phys. Soc. 14 (1969) 607
- 1969HO1X Hoot et al., Bull. Amer. Phys. Soc. 14 (1969) 494
- 1969IW1B Iwao, DPKU-027-68 (1969)
- 1969IW1E Iwao, DPKU-024-68 (1969)



1969JO1N Jones and Toms, Bull. Amer. Phys. Soc. 14 (1969) 607  
 1969KA1A G.Th. Kaschl, G.J. Wagner, G. Mairle, U. Schmidt-Rohr and P. Turek, Phys. Lett. B29 (1969) 167  
 1969KA1G T. Kammuri and H. Yoshida, Nucl. Phys. A129 (1969) 625  
 1969KA1R S.N. Kaplan, R.V. Pyle, L.E. Temple and G.F. Valby, Phys. Rev. Lett. 22 (1969) 795  
 1969KA1W Kaschl et al., Contrib., Montreal (1969) 262  
 1969KH1C Khanna and Harvey, Bull. Amer. Phys. Soc. 14 (1969) 604  
 1969KU01 W. Kutschera, D. Schwalm and B. Povh, Nucl. Phys. A124 (1969) 693  
 1969LA1A C.M. Lamba, N. Sarma and N.S. Thampi, Nucl. Phys. A122 (1969) 390  
 1969LE1D Leonardi and Rosa-Clot, Lett. Nuovo Cim. 1 (1969) 829  
 1969LU07 C.C. Lu, M.S. Zisman and B.G. Harvey, Phys. Rev. 186 (1969) 1086  
 1969LU1C Ludwig et al., Bull. Amer. Phys. Soc. 14 (1969) 1229  
 1969ME1M Mendelson, Hardy and Cerny, Bull. Amer. Phys. Soc. 14 (1969) 529  
 1969MU07 K.M. Murray and J.C. Ritter, Phys. Rev. 182 (1969) 1097  
 1969NI09 R.J. Nickles, Nucl. Phys. A134 (1969) 308  
 1969NY1A Nyberg, Jonsson and Bergqvist, Research Inst. Nat. Defence, Stockholm, NP 6902 (1969)  
 1969OC1B O'Connell and Close, Bull. Amer. Phys. Soc. 14 (1969) 508  
 1969PA1C Pape, Chevallier, Sens and Armbruster, Rev. Phys. Appl. 4 (1969) 227  
 1969PE1D Petrauskas and Vanagas, Sov. J. Nucl. Phys. 8 (1969) 270  
 1969PH02 G.W. Phillips and W.W. Jacobs, Phys. Rev. 184 (1969) 1052  
 1969PU04 K.H. Purser, W.P. Alford, D. Cline, H.W. Fulbright, H.E. Gove and M.S. Krick, Nucl. Phys. A132 (1969) 75  
 1969RI1C Ritter, Sheldon and Strang, Contrib., Montreal (1969) 256  
 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378  
 1969SA1J Saunier and Pearson, Bull. Amer. Phys. Soc. 14 (1969) 36  
 1969SH02 T. Shintomi and M. Masuda, J. Phys. Soc. Jpn. 26 (1969) 607  
 1969SI04 H.E. Siefken, P.M. Cockburn and R.W. Krone, Nucl. Phys. A128 (1969) 162, and Private Communication (1969)  
 1969SN02 F.D. Snyder and M.A. Waggoner, Phys. Rev. 186 (1969) 999  
 1969SN03 J.L. Snelgrove and E. Kashy, Phys. Rev. 187 (1969) 1246  
 1969SN04 J.L. Snelgrove and E. Kashy, Phys. Rev. 187 (1969) 1259  
 1969SU15 K. Sugimoto, Phys. Rev. 182 (1969) 1051

1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011  
1969TI05 C.T. Tindle and E. Vogt, Can. J. Phys. 47 (1969) 2763  
1969TO1A I.S. Towner, Nucl. Phys. A126 (1969) 97  
1969UB1C Uberall, Albert, Wagner and Werntz, Bull. Amer. Phys. Soc. 14 (1969) 606  
1969UL01 H. Ullrich and H. Krauth, Nucl. Phys. A123 (1969) 641  
1969VE02 V.V. Verbinski and W.R. Burrus, Phys. Rev. 177 (1969) 1671  
1969WA11 B.A. Watson, P.O. Singh and R.E. Segel, Phys. Rev. 182 (1969) 977  
1969WE02 M.L. West, C.M. Jones, J.K. Bair and H.B. Willard, Phys. Rev. 179 (1969) 1047;  
Erratum Phys. Rev. C1 (1970) 367  
1969WE03 H.R. Weller and H.A. Van Rinsvelt, Nucl. Phys. A129 (1969) 64  
1969WE07 K.J. Wetzel, Phys. Rev. 181 (1969) 1465  
1969WE08 H.R. Weller, Phys. Lett. B30 (1969) 409  
1969WI1G D.H. Wilkinson, Nucl. Phys. A133 (1969) 1  
1969WO1J Wong et al., Bull. Amer. Phys. Soc. 14 (1969) 38  
1969YO1C Young and Reisman, Bull. Amer. Phys. Soc. 14 (1969) 508  
1969ZH1A Zhusupov and Eramzhyan, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 730  
1970CA1G Campbell and Ober, Bull. Amer. Phys. Soc. 15 (1970) 126  
1970JA1E Jackson and Weller, Bull. Amer. Phys. Soc. 15 (1970) 163  
1970SO1G Sokol, De Luca and Browne, Bull. Amer. Phys. Soc. 15 (1970) 36  
1970SP1E Spinka and Winkler, Bull. Amer. Phys. Soc. 15 (1970) 805  
1971AJ02 F. Ajzenberg-Selove, Nucl. Phys. A166 (1971) 1  
1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1  
CA65 Unknown Source