

Energy Levels of Light Nuclei $A = 15$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 13-15$ was published in *Nuclear Physics A523* (1991), p. 1. This version of $A = 15$ differs from the published version in that we have corrected some errors discovered after the article went to press. The introduction and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed July 1, 1990)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 15$

Below is a list of links for items found within the PDF document. The introductory [Table 2](#) is available on this website via the link.

A. Nuclides: [\$^{15}\text{Li}\$](#) , [\$^{15}\text{Be}\$](#) , [\$^{15}\text{B}\$](#) , [\$^{15}\text{C}\$](#) , [\$^{15}\text{N}\$](#) , [\$^{15}\text{O}\$](#) , [\$^{15}\text{F}\$](#)

B. Tables of Recommended Level Energies:

[Table 15.1](#): Energy levels of ^{15}C

[Table 15.4](#): Energy levels of ^{15}N

[Table 15.16](#): Energy levels of ^{15}O

[Table 15.22](#): Energy levels of ^{15}F

C. [References](#)

D. Figures: [\$^{15}\text{C}\$](#) , [\$^{15}\text{N}\$](#) , [\$^{15}\text{O}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

¹⁵Li

(Not illustrated)

¹⁵Li has not been observed. Its atomic mass excess is calculated to be 81.60 MeV: see (1981AJ01). It is then unstable with respect to decay into ¹⁴Li+n and ¹³Li+2n by 1.2 and 5.1 MeV, respectively. (1985PO10) calculate [in a (0 + 1) $\hbar\omega$ model space] that the first four states of ¹⁵Li at 0, 0.73, 2.39 and 2.77 MeV have, respectively, $J^\pi = \frac{3}{2}^-, \frac{1}{2}^-, \frac{7}{2}^-$ and $\frac{5}{2}^-$. See also (1988POZS; theor.).

¹⁵Be

(Not illustrated)

¹⁵Be has not been observed. The calculated mass excess is 51.18 MeV: see (1981AJ01). ¹⁵Be is then unstable with respect to ¹⁴Be+n and ¹³Be+2n by 3.4 and 0.04 MeV, respectively. (1985PO10) calculate [in a (0 + 1) $\hbar\omega$ model space] that the first four states of ¹⁵Be at 0, 0.07, 2.32, 3.10 MeV have, respectively, $J^\pi = \frac{5}{2}^+, \frac{3}{2}^+, \frac{9}{2}^+, \frac{7}{2}^+$. See also (1987SA15; theor.).

¹⁵B

(Figs. 1 and 4)

Mass of ¹⁵B: Wapstra adopts 28970 ± 22 keV (1988WA18, and private communication) and so do we: see (1986AJ01). ¹⁵B is then stable with respect to ¹⁴B + n by 2.77 MeV.

Decay of ¹⁵B: ¹⁵B decays by β^- emission to ¹⁵C: Q_{β^-} (max) = 19.10 MeV. The character of the decay is not known but measurements of the half-life are 11 ± 1 ms (1984DU15), 8.8 ± 0.6 ms (1986CU01), 10.4 ± 0.3 ms (1988MU08), 10.8 ± 0.5 ms (1988SA04), $10.3_{-0.5}^{+0.6}$ ms (1989LE16). The weighted mean of these five values is 10.3 ± 0.2 ms. Omitting the low value from (1986CU01) gives 10.5 ± 0.3 ms, which we adopt.

Upper limits have been set on the P_{0n} and P_{2n} : 5% and 1.5%, respectively (1984DU15). See also (1989LE16).

General: (1985PO10) calculate [in a (0 + 1) $\hbar\omega$ model space] that the first four states of ¹⁵B at 0, 1.53, 2.06, 2.71 MeV have, respectively, $J^\pi = \frac{3}{2}^-, \frac{5}{2}^-, \frac{1}{2}^-$ and $\frac{7}{2}^-$.

Interaction cross sections at 790 MeV/A of ¹⁵B ions with Be, C and Al are reported by (1988TA10). The interaction radius and the r.m.s. radius for the nucleon distributions in ¹⁵B have also been derived (1988TA10). See also (1989SA10), (1986DU11, 1989DE52), (1986GU1D, 1988BAYZ, 1988MI1G, 1989DO1I, 1990LO10) and (1986AN07, 1989DO1K, 1989PO1K, 1989SI26, 1990RE04; theor.).

¹⁵C
(Figs. 1 and 4)

GENERAL (See also (1986AJ01)).

Model calculations: (1988MI1J, 1989PO1K, 1989WO1E).

Electromagnetic transitions: (1984VA06).

Astrophysical questions: (1989KA1K).

Complex reactions involving ¹⁵C: (1985PO11, 1986AV1B, 1986BI1A, 1986DU11, 1986HA1P, 1986HA1B, 1986PO06, 1987RI03, 1987SA25, 1987SN01, 1987VI02, 1988CA06, 1988JO1B, 1988MI28, 1988RU01, 1988SA19, 1989AS1B, 1989OG1B, 1989SA10, 1989SI26, 1989YO02).

Hypernuclei: (1988MA1G, 1989TA17).

Other topics: (1985AN28, 1986AN07).

Ground state of ¹⁵C: (1985AN28, 1986AS1B, 1987SA15, 1987VA26, 1988VA03, 1989SA10, 1989WO1E).

$$|g| = 2.63 \pm 0.14 \text{ (1988ASZY; prelim.)}$$

$$\mu_{\text{g.s.}} = 1.315 \pm 0.07 \text{ nm (1989RA17)}$$

$$\mu_{0.74} = -1.758 \pm 0.03 \text{ nm (1989RA17)}$$

1. ¹⁵C(β^-)¹⁵N $Q_m = 9.7717$

The half-life of ¹⁵C is 2.449 ± 0.005 s (1979AL23). Transitions have been observed to ¹⁵N_{g.s.} and to the upper of the 5.3 MeV states in ¹⁵N which has $J^\pi = \frac{1}{2}^+$. The $\log f^t$ to ¹⁵N*(5.30) indicates an allowed transition: therefore J^π (¹⁵C_{g.s.}) = $\frac{1}{2}^+$ or $\frac{3}{2}^+$. Weak transitions are observed to ¹⁵N*(7.30, 8.31, 8.57, 9.05) (1979AL23): see Table 15.14. The shape of the ¹⁵C_{g.s.} \rightarrow ¹⁵N_{g.s.} transition differs appreciably from an allowed shape (1984WA07). See also (1986AS1B, 1988ASZY), (1988WA1E) and (1989BA92, 1989PO1K; theor.).

2. ⁹Be(⁷Li, p)¹⁵C $Q_m = 9.092$

Observed proton groups are displayed in Table 15.2.

3. ¹³C(t, p)¹⁵C $Q_m = 0.9127$

Observed groups are displayed in Table 15.3. See also (1981AJ01).




Fig. 1: Energy levels of ^{15}C . For notation see Fig. 2.

Table 15.1: Energy levels of ^{15}C ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
g.s.	$\frac{1}{2}^+; \frac{3}{2}$	$\tau_{1/2} = 2.449 \pm 0.005$ s $ g = 2.63 \pm 0.14$	β^-	1, 2, 3, 4, 6, 7, 9
0.7400 ± 1.5	$\frac{5}{2}^+$	$\tau_m = 3.76 \pm 0.10$ ns $g = -0.703 \pm 0.012$	γ	2, 3, 4, 7, 8
3.103 ± 4	$\frac{1}{2}^-$	$\Gamma_{\text{c.m.}} \leq 40$		2, 3, 9
4.220 ± 3	$\frac{5}{2}^-$	< 14		2, 3
4.657 ± 9	$\frac{3}{2}^-$			2, 3
4.78 ± 100	$\frac{3}{2}^+$	1740 ± 400		6
5.833 ± 20	$(\frac{3}{2}^+)$	64 ± 8		2, 6
5.866 ± 8	$\frac{1}{2}^-$			2, 3
6.358 ± 6	$(\frac{5}{2}, \frac{7}{2}^+, \frac{9}{2}^+)$	< 20		2, 3
6.417 ± 6	$(\frac{3}{2} \rightarrow \frac{7}{2})$	≈ 50		2, 3
6.449 ± 7	$(\frac{9}{2}^-, \frac{11}{2})$	< 14		2, 3
6.536 ± 4	^a	< 14		2, 3
6.626 ± 8	$(\frac{3}{2})$	20 ± 10		2, 3
6.841 ± 4	^a	< 14		2, 3
6.881 ± 4	$(\frac{9}{2})^{\text{a}}$	< 20		2, 3
7.095 ± 4	$(\frac{3}{2})$	< 15		2, 3
7.352 ± 6	$(\frac{9}{2}, \frac{11}{2})$	20 ± 10		2, 4
7.414 ± 20				2
7.75 ± 30 ^b				2
8.01 ± 30				2
8.11 ± 10 ^b				2
8.47 ± 15	$(\frac{9}{2} \rightarrow \frac{13}{2})$	40 ± 15		2
8.559 ± 15	$(\frac{7}{2} \rightarrow \frac{13}{2})$	40 ± 15		2
9.00 ± 30				2
(9.73 ± 30)				2
9.789 ± 20	$(\frac{9}{2} \rightarrow \frac{15}{2})$	20 ± 15		2
10.248 ± 20	$(\frac{5}{2} \rightarrow \frac{9}{2})$	20 ± 15		2
11.015 ± 25				2
11.123 ± 20	$(\frac{11}{2} \rightarrow \frac{19}{2})$	30 ± 20		2

Table 15.1: Energy levels of ^{15}C ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
(11.68 \pm 30)				2
11.825 \pm 20	$\geq \frac{13}{2}$	70 \pm 30		2

^a See also Tables 15.2 and 15.3 and reaction 8.

^b Broad or unresolved states.

 Table 15.2: Proton groups from $^9\text{Be}(^7\text{Li}, \text{p})^{15}\text{C}$ and $^{14}\text{C}(\text{d}, \text{p})^{15}\text{C}$ ^a

$^9\text{Be}(^7\text{Li}, \text{p})^{15}\text{C}$ ^b			$^{14}\text{C}(\text{d}, \text{p})^{15}\text{C}$ ^c		
E_x (keV)	$\Gamma_{\text{c.m.}}$ (keV)	J^π ^d	E_x (keV)	$\Gamma_{\text{c.m.}}$ (keV)	J^π ^e
g.s. $\equiv 740$ ^f	bound bound		g.s. 744.1 \pm 2 ^j	bound bound	$\frac{1}{2}^+$ ^o $\frac{5}{2}^+$ ^p
3100 \pm 30	< 40	$(\frac{1}{2}^-)$ ^h	3105.3 \pm 5 ^k	≈ 42	$(\frac{1}{2}^-)$
4223 \pm 15	< 15	$(\frac{5}{2}^-)$	4221.1 \pm 3 ^k	< 14	$(\frac{7}{2}^+, \frac{5}{2}^-)$
(4550 \pm 30)			4657 ^k		
			4780 \pm 100 ^l	1740 \pm 400	$\frac{3}{2}^+$
5833 \pm 20		i	5810 \pm 20 ^l	64 \pm 8	$(\frac{3}{2}^+)$ ^q
5858 \pm 20		i			
6370 \pm 15	< 20	$(\frac{5}{2})$	k, m	< 14	$(\frac{7}{2}, \frac{9}{2})^+$
6436 \pm 20			6428.1 \pm 7	≈ 50	$(\frac{3}{2}, \frac{5}{2}, \frac{7}{2})$
6461 \pm 20			m	< 14	$(\frac{9}{2}^-, \frac{11}{2})$
6542 \pm 15	< 20	$(\frac{3}{2})$	6539.8 \pm 5	< 14	$(\frac{9}{2}^-, \frac{11}{2})$
6639 \pm 15	20 \pm 10	$(\frac{3}{2})$			
6847 \pm 15	< 20	$(\frac{11}{2}, \frac{13}{2})$	6844.9 \pm 5	< 14	$(\frac{13}{2}, \frac{11}{2})^+$
6894 \pm 15	< 20	$(\frac{7}{2}, \frac{9}{2})$	6882.4 \pm 5		$((\frac{9}{2}^-, \frac{11}{2}^+, \frac{13}{2}^+))$
7100 \pm 15	< 15	$(\frac{3}{2})$	7097.2 \pm 6		
7354 \pm 15	20 \pm 10	$(\frac{9}{2}, \frac{11}{2})$	7351.3 \pm 6		
7414 \pm 20					
7750 \pm 30 ^g			7.81 \pm 10 ⁿ		
8010 \pm 30					

Table 15.2: Proton groups from ${}^9\text{Be}({}^7\text{Li}, \text{p}){}^{15}\text{C}$ and ${}^{14}\text{C}(\text{d}, \text{p}){}^{15}\text{C}$ ^a (continued)

${}^9\text{Be}({}^7\text{Li}, \text{p}){}^{15}\text{C}$ ^b			${}^{14}\text{C}(\text{d}, \text{p}){}^{15}\text{C}$ ^c		
E_x (keV)	$\Gamma_{\text{c.m.}}$ (keV)	J^π ^d	E_x (keV)	$\Gamma_{\text{c.m.}}$ (keV)	J^π ^e
8130 ± 30 ^g			8.10 ± 10 ⁿ		
8491 ± 15	40 ± 15	$(\frac{9}{2}, \frac{11}{2}, \frac{13}{2})$	8.46 ± 10 ⁿ		
8559 ± 15	40 ± 15	$(\frac{7}{2} \rightarrow \frac{13}{2})$			
9000 ± 30					
(9730 ± 30)					
9789 ± 20	20 ± 15	$(\frac{9}{2} \rightarrow \frac{15}{2})$			
10248 ± 20	20 ± 15	$(\frac{5}{2}, \frac{7}{2}, \frac{9}{2})$			
11015 ± 25					
11123 ± 20	30 ± 20	$(\frac{11}{2} \rightarrow \frac{19}{2})$			
(11680 ± 30)					
11825 ± 20	70 ± 30	$(\frac{13}{2} \rightarrow \frac{31}{2})$			

^a For references see Table 15.2 in (1981AJ01).

^b $E({}^7\text{Li}) = 20$ MeV. E_x based on 740 keV for the first excited state.

^c $E_d = 12 - 14$ MeV.

^d Suggested J^π assignments based on angular distributions (and $2J_f + 1$ dependence) and l_{max} from Γ_n .

^e Analysis of the two bound states is done using DWUCK. For the unbound states DOXY was used.

^f $E_x = 739 \pm 1$ keV [from E_γ]; $\tau_m = 3.77 \pm 0.11$ ns.

^g Broad or unresolved states.

^h $\theta_n^2 = 0.0075 \pm 0.0015$.

ⁱ Sum of the J for these two states is 2 [based on $(2J_f + 1)$ dependence of cross section].

^j $\tau_m = 3.73 \pm 0.23$ ns.

^k See also (1985DA23).

^l See text, reaction 6 (1985DA23).

^m Observed but E_x not determined.

ⁿ Observed at $E_d = 27$ MeV.

^o $S = 0.88$.

^p $S = 0.69$ or 0.55 . $g = -0.77 \pm 0.06$.

^q May be unresolved.

4. ${}^{13}\text{C}(\alpha, 2\text{p}){}^{15}\text{C}$

$Q_m = -18.9013$

See (1981AJ01).

5. $^{14}\text{C}(n, \gamma)^{15}\text{C}$ $Q_m = 1.2181$

$\sigma_\gamma < 1 \mu\text{b}$ (1981MUZQ).

6. $^{14}\text{C}(d, p)^{15}\text{C}$ $Q_m = -1.0065$

At $E_d = 16$ MeV angular distributions and A_y measurements are reported to a state at $E_x = 4.78 \pm 0.10$ MeV ($\Gamma_{\text{c.m.}} = 1.74 \pm 0.40$ MeV); $S = 0.5$. A narrow state at $E_x = 5.81 \pm 0.02$ MeV ($\Gamma_{\text{c.m.}} = 64.3 \pm 8.1$ keV), $S = 0.02$, is also observed. It is suggested that these are 1p2h and 3p4h $\frac{3}{2}^+$ states (1985DA23) [and S.E. Darden, private communication]. For the earlier work see Table 15.2.

Table 15.3: Proton groups from $^{13}\text{C}(t, p)^{15}\text{C}$ ^a

E_x (MeV \pm keV)	J^π	E_x (MeV \pm keV)	J^π
0	$\frac{1}{2}^+$	6.440 ± 6	
0.743 ± 9 ^b	$\frac{5}{2}^+$	6.529 ± 6	
3.100 ± 6 ^b	$\frac{1}{2}^-$	6.622 ± 9	
4.215 ± 9 ^b	$\frac{5}{2}^-$	6.835 ± 6 ^b	$(\frac{7}{2}, \frac{9}{2})^-$
4.657 ± 9 ^b	$\frac{3}{2}^-$	6.876 ± 7	
5.867 ± 8	$\frac{1}{2}^-$	7.093 ± 6	
6.356 ± 6		7.387 ± 7 ^b	$(\frac{9}{2}, \frac{7}{2})^-$
6.404 ± 7			

^a (1983TR12); $E_t = 18$ MeV; DWBA.

^b Strong group.

7. $^{14}\text{C}(^{13}\text{C}, ^{12}\text{C})^{15}\text{C}$ $Q_m = -3.7283$

Angular distributions have been studied at $E(^{13}\text{C}) = 20.0$ to 27.5 MeV to $^{15}\text{C}^*(0, 0.74)$ (1988BI11). See also (1990VO1E).

8. $^{15}\text{N}(\pi^-, \gamma)^{15}\text{C}$ $Q_m = 129.797$

Radiative pion capture shows evidence for $J^\pi = \frac{5}{2}^+$, $T = \frac{3}{2}$ giant magnetic quadrupole states: transitions are reported to $^{15}\text{C}^*(0.74)$ as well as to $^{15}\text{C}^*(6.7, 8.6, 12.0)$ (1983ST04).

9. $^{16}\text{O}(^7\text{Li}, ^8\text{B})^{15}\text{C}$ $Q_m = -22.624$

At $E(^7\text{Li}) = 82 \text{ MeV}$ $^{15}\text{C}^*(0, 3.1)$ are populated (1985AL1G).

^{15}N
(Figs. 2 and 4)

GENERAL (See also (1986AJ01)).

Nuclear models: (1985KW02, 1985PH01, 1987KA09, 1987KI1C, 1987ME1D, 1987ST05, 1988WO04, 1989WO1E, 1990VA01)

Special states: (1985AR1H, 1985GO1A, 1985PH01, 1985SH24, 1987KI1C, 1987ST05, 1988KW02, 1988ZH1B, 1989OR02)

Electromagnetic transitions and giant resonances: (1985BL20, 1985GO1A, 1986ER1A, 1987KI1C, 1987ST05, 1989ASZZ)

Astrophysical questions: (1982BU1A, 1982CA1A, 1982WO1A, 1985PR1D, 1986FR1G, 1987AR1C, 1987AU1A, 1987LE1J, 1987ZI1C, 1988FE1A, 1988KR1G, 1988PI1C, 1988WA1I, 1989CH1X, 1989GU1Q, 1989GU1J, 1989GU1L, 1989JI1A, 1989KA1K, 1989KE1D, 1989ME1C, 1989NO1A, 1989WY1A, 1989YO1H, 1990HA1W, 1990RA1O)

Complex reactions involving ^{15}N : (1985AR1H, 1985BE40, 1985HA1N, 1985PO11, 1985SI19, 1985UT01, 1986AI1A, 1986CH2G, 1986CO1Q, 1986GR1A, 1986HA1B, 1986MA13, 1986MA19, 1986ME06, 1986PO06, 1986PO15, 1986SA30, 1986SC28, 1986SO10, 1986TO10, 1986UT01, 1986VA23, 1987BA38, 1987BE1I, 1987BU07, 1987FE1A, 1987MI27, 1987NA01, 1987OL1A, 1987RI03, 1987ST01, 1987VI02, 1988AR1D, 1988GO11, 1988JO1B, 1988SA19, 1988UT02, 1989BA92, 1989GE11, 1989GRZQ, 1989KI13, 1989PA06, 1989SA10, 1989TE02, 1989YO02, 1990DA03, 1990GL01, 1990WE14)

Applied work: (1986AM1B, 1986CO1Q, 1986EN1A, 1986HE1F, 1986LE1L, 1986NO1C, 1986SA2H, 1986ST1K, 1987SI1D, 1988GR1A, 1988PI12, 1988PR1D, 1988VI1A, 1989KU1P, 1989TA1Y, 1989YO1H, 1990AM1F)

Pion capture and reactions (See also reactions 15, 43, and 46.): (1985LE1E, 1985MA1K, 1986BA1C, 1986SI11, 1987KA09, 1987LE1B, 1988LI23, 1988MI1K, 1988RO1M, 1988TA21, 1989CH31, 1989GE10, 1989JO07, 1989LE1L, 1990ER03, 1990OD1A, 1990TA1K)

Reactions involving other mesons and hyperons: (1985IA01, 1986FE1A, 1989DO1I, 1989DO1K)

Antiproton reactions: (1985BA51)

Hypernuclei: (1984BO1H, 1985IA01, 1986AN1R, 1986DA1G, 1986DA1B, 1986FE1A, 1986GA1H, 1986KO1A, 1986YA1F, 1987MA2A, 1987MI38, 1987PO1H, 1987WU05, 1988MO1L, 1989BA92, 1989BA93, 1989DO1K, 1989KO1H, 1989MI30, 1989TA17)

Other topics: (1985AN28, 1985PH01, 1985SH24, 1986AN07, 1986WI03, 1987CH02, 1988KW02, 1989OR02, 1989PO1K, 1990MU10)

Ground-state properties of ^{15}N : (1985AN28, 1985AR11, 1985BL20, 1985GO1A, 1986BA04, 1986BA49, 1986MC13, 1986WI03, 1986WUZX, 1987DE03, 1987FU06, 1987IC02, 1987KI1C,

1987MI27, 1988AR1B, 1988AR1I, 1988CH1T, 1988DE09, 1988FU04, 1988KE1B, 1988NI05, 1988SH07, 1988VA03, 1988WA08, 1988WO04, 1989CH24, 1989FU05, 1989GOZQ, 1989NE02, 1989SA10, 1989WO1E, 1990VA1G, 1990VA01)

Table 15.4: Energy levels of ^{15}N ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^-; \frac{1}{2}$	-	stable	3, 4, 5, 6, 13, 14, 16, 17, 18, 19, 20, 24, 25, 26, 27, 28, 31, 32, 33, 34, 35, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66
5.270155 ± 0.014 ^b	$\frac{5}{2}^+$	$\tau_m = 2.58 \pm 0.14$ ps $g = +(0.94 \pm 0.07)$	γ	4, 5, 16, 17, 24, 25, 31, 32, 35, 40, 45, 46, 49, 50, 56, 59, 60, 64, 65
5.298822 ± 0.014 ^b	$\frac{1}{2}^+$	25 ± 7 fs	γ	4, 5, 10, 11, 12, 16, 18, 24, 25, 26, 31, 32, 35, 40, 42, 45, 49, 50, 56, 60, 64, 65
6.32378 ± 0.02 ^b	$\frac{3}{2}^-$	0.211 ± 0.012 fs	γ	4, 5, 10, 11, 12, 13, 16, 18, 24, 26, 31, 32, 35, 39, 40, 42, 44, 45, 46, 49, 50, 56, 57, 59, 60, 61, 63, 64, 65
7.15505 ± 0.02 ^b	$\frac{5}{2}^+$	18 ± 8 fs	γ	4, 5, 12, 16, 17, 18, 24, 25, 26, 31, 32, 35, 40, 45, 49, 50, 60
7.30083 ± 0.02 ^b	$\frac{3}{2}^+$	0.61 ± 0.05 fs	γ	4, 5, 12, 16, 18, 24, 25, 26, 31, 32, 35, 40, 42, 45, 49, 50, 60
7.5671 ± 1.0 ^c	$\frac{7}{2}^+$	12_{-6}^{+11} fs	γ	4, 5, 10, 11, 12, 16, 17, 18, 24, 25, 26, 31, 40, 45, 46, 49, 50, 60, 64
8.31262 ± 0.027 ^b	$\frac{1}{2}^+$	1.7 ± 1.1 fs	γ	4, 5, 18, 24, 25, 26, 31, 35, 39, 40, 42, 45, 49, 50, 56
8.5714 ± 0.12	$\frac{3}{2}^+$	0.7 ± 0.7 fs	γ	4, 5, 10, 11, 12, 16, 17, 18, 24, 25, 26, 31, 40, 42, 45, 49, 50
9.04971 ± 0.07	$\frac{1}{2}^+$	0.50 ± 0.08 fs	γ	4, 5, 24, 25, 31, 35, 40, 42, 45, 56
9.15190 ± 0.12 ^b	$\frac{3}{2}^-$	1.40 ± 0.36 fs	γ	4, 5, 10, 11, 24, 25, 31, 35, 40, 45, 49, 50
9.15490 ± 0.03 ^b	$\frac{5}{2}^+$	7_{-3}^{+6} fs	γ	4, 5, 18, 24, 31, 35, 40, 49, 50
9.2221 ± 0.8	$\frac{1}{2}^-$	< 130 fs	γ	24, 26, 31, 35, 40, 56, 60

Table 15.4: Energy levels of ^{15}N ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
9.760 \pm 1	$\frac{1}{2}^{\ominus}$	2.6 \pm 0.9 fs	γ	24, 40, 45
9.829 \pm 3	$\frac{1}{2}^{\ominus}$	17 \pm 7 fs	γ	4, 5, 10, 11, 17, 18, 24, 26, 31, 40, 49, 50
9.9250 \pm 0.2	$\frac{3}{2}^{\ominus}$	0.31 \pm 0.05 fs	γ	18, 24, 31, 35, 40, 45
10.0660 \pm 0.2 ^c	$\frac{3}{2}^{\oplus}$	0.100 \pm 0.006 fs	γ	18, 35, 40, 44, 45, 49, 50
10.4497 \pm 0.3	$\frac{1}{2}^{\ominus}$	$\Gamma < 0.5$ keV	γ, p	5, 10, 11, 24, 28, 40
10.5333 \pm 0.5	$\frac{1}{2}^{\oplus}$		γ, p	5, 10, 11, 18, 24, 25, 28, 31, 40
10.6932 \pm 0.3	$\frac{1}{2}^{\oplus}$	$\tau_m = 18 \pm 9$ fs	γ, p	5, 11, 16, 28, 46
10.7019 \pm 0.3	$\frac{3}{2}^{\ominus}$	$\Gamma = 0.2$ keV	γ, p	10, 11, 17, 18, 24, 26, 28, 60
10.804 \pm 2	$\frac{3}{2}^{\oplus}$	$< 1 \times 10^{-3}$	γ, p	4, 5, 10, 11, 18, 24, 28, 40, 45
11.235 \pm 5 ^b	$\frac{1}{2}^{\oplus}$	3.3	n	16, 31, 36, 40
11.2928 \pm 0.7	$\frac{1}{2}^{\ominus}$	8 \pm 3	γ, n, p	16, 18, 28, 29, 30, 31, 36, 38, 49
11.4376 \pm 0.7	$\frac{1}{2}^{\oplus}$	41.4 \pm 1.1	γ, n, p, α	6, 7, 10, 11, 18, 25, 28, 29, 30, 31, 36, 38, 64
11.615 \pm 4	$\frac{1}{2}^{\oplus}; T = \frac{3}{2}$	405 \pm 6	γ, n, p	28, 29, 30
11.763 \pm 3	$\frac{3}{2}^{\oplus}$	40	n, p, α	7, 29, 30, 36, 38
11.876 \pm 3	$\frac{3}{2}^{\ominus}$	25	γ, n, p, α	7, 29, 30, 36, 38, 48
11.942 \pm 6	$\frac{1}{2}^{\ominus}$	≤ 3.0	n, α	5, 16, 17, 18, 25, 26, 36
11.965 \pm 3	$\frac{1}{2}^{\ominus}$	17	n, p, α	5, 7, 10, 11, 29, 30, 36, 38
12.095 \pm 3	$\frac{1}{2}^{\oplus}$	14 \pm 5	n, p, α	7, 25, 29, 30, 36, 38
12.145 \pm 3	$\frac{3}{2}^{\ominus}$	41 \pm 5	n, p, α	7, 10, 11, 29, 30, 36, 38
12.327 \pm 4	$\frac{1}{2}^{\oplus} (+)$	22	n, p	17, 18, 25, 29, 30, 36, 38
12.493 \pm 4	$\frac{1}{2}^{\oplus}; \frac{1}{2}$	40 \pm 5	n, p, α	7, 18, 25, 29, 30, 36, 38
12.522 \pm 8	$\frac{1}{2}^{\oplus}; \frac{3}{2}$	58 \pm 4	γ, p	28, 45
12.551 \pm 10	$\frac{1}{2}^{\oplus}$			5, 11, 16, 17, 25, 46
12.920 \pm 4	$\frac{3}{2}^{\ominus}$	56 \pm 11	n, p, α	7, 9, 18, 29, 30, 36, 38
12.940 \pm 10	$\frac{1}{2}^{\oplus}$	81	p, α	7, 9, 29, 30
13.004 \pm 10	$\frac{1}{2}^{\ominus}$			5, 10, 11, 16, 18, 25, 26
13.149 \pm 10		7 \pm 3	n, p, α	7, 38
13.174 \pm 7	$(\frac{3}{2})$	7 \pm 3	n, p, α	5, 7, 11, 16, 17, 18, 29, 36, 38
13.362 \pm 8	$\frac{3}{2}^{\ominus}$	16 \pm 8	n, p, α	7, 9, 29, 30, 38
13.390 \pm 10	$\frac{3}{2}^{\oplus}$	56	γ, n, p, α	7, 9, 28, 29, 30, 38
13.537 \pm 10	$\frac{3}{2}^{\ominus}$	85 \pm 30	n, p, α	7, 9, 29, 30

Table 15.4: Energy levels of ^{15}N ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
13.608 \pm 7 (13.612 \pm 10)	$\frac{5}{2}^{(+)}$ $(\frac{1}{2}^+)$	18 \pm 4 90	n, p, α n, p, α	7, 18, 36, 38 9, 29, 30
13.713 \pm 10		26 \pm 8	n, p, α	7, 29, 38
13.84 \pm 30	$\frac{3}{2}^+$	75	n, p, α	5, 7, 9, 11, 25, 36, 38
13.9	$\frac{1}{2}^+$	930	γ , p	28, 29
13.99 \pm 30	$\frac{5}{2}^+$	98 \pm 10	n, p, α	7, 11, 29, 30
14.090 \pm 7	$(\frac{9}{2}^+, \frac{7}{2}^+)$	22 \pm 6	n, p, α	5, 7, 10, 11, 18, 25, 36, 38, 46
14.10 \pm 30	$\frac{3}{2}^+$	\approx 100	n, α	5, 7, 9, 30
14.162 \pm 10	$\frac{3}{2}^{(+)}$	27 \pm 6	n, α	5, 7, 36, 38
14.24 \pm 40	$\frac{5}{2}^+$	150	α	9, 10
14.38 \pm 40	$\frac{7}{2}^+$	100	α	9
14.4		\approx 1900	n, p, α	36, 38
14.55 \pm 20		200 \pm 50	n, (p), α	7
14.647 \pm 10		33 \pm 6	n, p, α	7, 36, 38
14.71		750	γ , p	28
14.720 \pm 10	$\frac{5}{2}^-$	110 \pm 50	γ , n, (p), α	7, 10, 11, 18, 36, 38, 45
14.86 \pm 20		48 \pm 11	n, α	7, 9, 18
14.920 \pm 10		12 \pm 3	n, α	7, 10, 38
15.025 \pm 10		13 \pm 3	n, α	7, 18
15.09 \pm 20		80 \pm 25	n, α	7, 9, 49
15.288 \pm 10		26 \pm 6	n, α	7, 9
15.373 \pm 10	$\frac{13}{2}^+$			5, 10, 11, 16, 17, 18
15.38 \pm 20		75 \pm 25	n, t, α	7, 9, 14
15.43 \pm 20		\approx 100	n, (α)	7, 9
15.45		750	γ , p	28
15.53 \pm 20		\approx 35	n, α	7, 10, 11, 38
15.60 \pm 20		95 \pm 25	n, α	7
15.782 \pm 10			p, t, α	7, 14, 18
15.93 \pm 20		35 \pm 5	n, t, α	7, 14, 17
15.944 \pm 15		21 \pm 6	n, t, α	7, 14
16.026 \pm 10		62 \pm 12	n, p, t, α	7, 9, 14, 18, 38
16.190 \pm 10	$\frac{3}{2}^+$	450 \pm 100	γ , n, p, t, α	10, 14, 18
16.26 \pm 20	$\frac{3}{2}^+$	150 \pm 28	γ , n, t, α	6, 7, 9, 14, 17, 18
16.32 \pm 20		\approx 30	n, p, t, α	7, 14
16.39 \pm 20		44 \pm 11	n, p, t, α	7, 14, 17, 38
16.46		560	γ , p, d	21, 28

Table 15.4: Energy levels of ^{15}N ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
16.576 \pm 15		27 \pm 15	n, α	7, 38
16.59 \pm 25	$\frac{3}{2}^-$	490	γ , n, p, t, α	14
16.677 \pm 15	$\frac{1}{2}^+; \frac{1}{2}$	80 \pm 20	γ , n, p, d, t, α	6, 7, 14, 17, 21, 23, 28, 30, 36, 38, 43
16.85 \pm 30	$\frac{5}{2}$	110 \pm 50	t, α	14
16.91		\approx 350	n, p, d, t, α	14, 21, 36, 38
(17.05)			p, t	14
17.11		broad	d, α	23
17.15 \pm 50	$(\frac{1}{2}^+, \frac{3}{2}^+)$	250 \pm 60	γ , t, α	6, 14
17.23 \pm 40		\approx 175	d, t, (α)	23
17.37 \pm 40		\approx 250	p, d, t, α	14, 21, 23, 36, 38
17.58 \pm 40	$\frac{3}{2}^+$	450 \pm 120	γ , d, t, α	14, 23, 38
17.67 \pm 40	$\frac{3}{2}^+; \frac{1}{2}$	600 \pm 80	γ , n, d, α	6, 20, 21, 23
17.72 \pm 10		48 \pm 10	n, (p), d, t, α	18, 21, 23, 38
17.95 \pm 20		167	n, α	18
18.06 \pm 10		19 \pm 4	(n), d, α	17, 21, 23
18.09 \pm 20		\approx 40	(n), p, d, t	21, 23
18.22		158	n, α	36, 38
18.27 \pm 20		235 \pm 60	n, p, d, α	18, 21, 23, 38
18.70 \pm 20				11, 18
18.91 \pm 150	$\frac{3}{2}^+ + \frac{1}{2}^+$	750 \pm 70	γ , α	6
19.20 \pm 35	$(\frac{1}{2}^+; \frac{1}{2})$	\approx 130	n, d	18, 21
19.5	$\frac{3}{2}^+; (\frac{3}{2})$	\approx 400	γ , p, t	14, 28, 29
19.72 \pm 40		d		11, 17, 18
20.12 \pm 50	$(T = \frac{3}{2})$			16, 46
20.5	$\frac{3}{2}^+$	\approx 400	γ , n, p, d	21, 28
20.96 \pm 65	$\frac{3}{2}^+ + \frac{1}{2}^+$	1740 \pm 150	γ , α	6, 18
21.82		\approx 600	γ , p, d	20, 28, 43
23.19 \pm 60	$(T = \frac{3}{2})$		γ , p	28, 46
23.6		broad	γ , n, d	20, 43
24.75 \pm 150		d		18
25.5	$\frac{3}{2}^-; (T = \frac{3}{2})$		γ , n, p	28, 43
(26.8)			t	14
\approx 37			γ , p	28

^a See also Tables 15.5 and 15.12 here, and Table 15.6 in (1986AJ01) [τ_m].

^b Revisions in the values of the fundamental constants and of the binding energy of the deuteron, as well as a reevaluation of earlier work, lead (1990WA22) to suggest values for E_x which differ from the ones shown by, typically, 40 eV [lower].

^c See also reaction 40.

^d Wide or unresolved.

$$\langle r^2 \rangle^{1/2} = 2.612 \pm 0.009 \text{ fm (1988DE09)}$$

$$\mu = -0.283188842 (45) \text{ nm (see 1989RA17)}$$

Natural abundance: $(0.366 \pm 0.009)\%$ (1984DE53)

$^{15}\text{N}^*(5.27)$: $\mu = +(2.35 \pm 0.18) \text{ nm (see 1989RA17)}$

1. (a) $^9\text{Be}(^6\text{Li}, \text{n})^{14}\text{N}$	$Q_m = 14.4986$	$E_b = 25.3319$
(b) $^9\text{Be}(^6\text{Li}, \text{p})^{14}\text{C}$	$Q_m = 15.1245$	
(c) $^9\text{Be}(^6\text{Li}, \text{t})^{12}\text{C}$	$Q_m = 10.4835$	
(d) $^9\text{Be}(^6\text{Li}, \alpha)^{11}\text{B}$	$Q_m = 14.3403$	

Thick target neutron yields are reported at $E(^6\text{Li}) = 40 \text{ MeV}$ (1987SC11). The yield of p_0 and p_1 (reaction (b)) for $E(^6\text{Li}) = 3.84$ to 6.40 MeV shows some broad structure: analysis in terms of Ericson fluctuation theory gives a value of $\approx 0.4 \text{ MeV}$ for the average level width at $E_x = 28 \text{ MeV}$ in ^{15}N . The excitation functions for t_0 (reaction (c)), α_0 , α_1 and α_2 (reaction (d)) show broad structures for $E(^6\text{Li}) = 4$ to 14 MeV . See (1976AJ04) for the references.

2. $^9\text{Be}(^7\text{Li}, \text{n})^{15}\text{N}$	$Q_m = 18.0818$
--	-----------------

Fig. 2: Energy levels of ^{15}N . In these diagrams, energy values are plotted vertically in MeV, based on the ground state as zero. Uncertain levels or transitions are indicated by dashed lines; levels which are known to be particularly broad are cross-hatched. Values of total angular momentum J , parity, and isobaric spin T which appear to be reasonably well established are indicated on the levels; less certain assignments are enclosed in parentheses. For reactions in which ^{15}N is the compound nucleus, some typical thin-target excitation functions are shown schematically, with the yield plotted horizontally and the bombarding energy vertically. Bombarding energies are indicated in laboratory coordinates and plotted to scale in cm coordinates. Excited states of the residual nuclei involved in these reactions have generally not been shown; where transitions to such excited states are known to occur, a brace is sometimes used to suggest reference to another diagram. For reactions in which the present nucleus occurs as a residual product, excitation functions have not been shown; a vertical arrow with a number indicating some bombarding energy, usually the highest, at which the reaction has been studied, is used instead. Further information on the levels illustrated, including a listing of the reactions in which each has been observed, is contained in the master table, entitled “Energy levels of ^{15}N ”.

Table 15.5: Radiative decays in ^{15}N ^a

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Mult. mixing ratio δ
5.27	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	100	-0.131 ± 0.013
5.30	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	100	
6.32 ^b	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100	$+0.132 \pm 0.004$
7.16 ^c	$\frac{5}{2}^+$	5.27	$\frac{5}{2}^+$	100 ± 0.4	$-0.014^{+0.012}_{-0.015}$
7.30	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	99.3 ± 0.7	$-0.017^{+0.005}_{-0.008}$
		5.27	$\frac{5}{2}^+$	0.6 ± 0.1	$+0.18 \pm 0.15$, or $+2.5 \pm 1.0$
		5.30	$\frac{1}{2}^+$	0.2 ± 0.1	-0.31 ± 0.15 , or $+4.6 \pm 3.4$
		6.32	$\frac{3}{2}^-$	< 0.25	
7.57 ^d	$\frac{7}{2}^+$	0	$\frac{1}{2}^-$	1.3 ± 0.6	
		5.27	$\frac{5}{2}^+$	98.7 ± 1.0	-0.028 ± 0.012
8.31	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	79 ± 2	
		5.27	$\frac{5}{2}^+$	< 3	
		5.30	$\frac{1}{2}^+$	10 ± 2	
		6.32	$\frac{3}{2}^-$	4.4 ± 1.0	
		7.16	$\frac{5}{2}^+$	1.2 ± 0.6	
		7.30	$\frac{3}{2}^+$	4.4 ± 0.7	
8.57 ^e	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	33 ± 2	$-0.085^{+0.005}_{-0.009}$
		5.27	$\frac{5}{2}^+$	65 ± 3	-0.091 ± 0.007
		6.32	$\frac{3}{2}^-$	1.4 ± 0.6	
		7.16	$\frac{5}{2}^+$	3.6 ± 0.5	
		7.30	$\frac{3}{2}^+$	1.2 ± 0.4	
9.05 ^f	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	92 ± 3	
		5.27	$\frac{5}{2}^+$	3.5 ± 1	
		6.32	$\frac{3}{2}^-$	4.5 ± 1	
		7.30	$\frac{3}{2}^+$	1.2 ± 0.4	
9.152	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100 ± 3	$+0.015^{+0.041}_{-0.034}$
9.155	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	< 2	
		5.27	$\frac{5}{2}^+$	11 ± 1	
		5.30	$\frac{1}{2}^+$	10 ± 1	
		6.32	$\frac{3}{2}^-$	22 ± 2	
		7.16	$\frac{5}{2}^+$	57 ± 3	

Table 15.5: Radiative decays in $^{15}\text{N}^a$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Mult. mixing ratio δ
9.22 ^g	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	22 ± 5	
		5.30	$\frac{1}{2}^+$	42 ± 8	
		6.32	$\frac{3}{2}^-$	35 ± 6	
		7.30	$\frac{3}{2}^+$	2.6 ± 0.7	
9.76 ^h	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	81.5 ± 2.8	
		5.27 + 5.30		7.5 ± 1.5	
		6.32	$\frac{3}{2}^-$	3.7 ± 0.8	
		7.16	$\frac{5}{2}^+$	2.3 ± 0.5	
9.83 ⁱ	$\frac{7}{2}^-$	7.57	$\frac{7}{2}^+$	5.0 ± 0.6	
		5.27	$\frac{5}{2}^+$	≈ 85	
		6.32	$\frac{3}{2}^-$	2.2 ± 0.9	
		7.16	$\frac{5}{2}^+$	2.4 ± 1.1	
9.93 ^j	$\frac{3}{2}^-$	7.30	$\frac{3}{2}^+$	3.7 ± 0.9	
		7.57	$\frac{7}{2}^+$	7.3 ± 1.0	
		0	$\frac{1}{2}^-$	77.6 ± 1.9	
		5.27 + 5.30		15.4 ± 1.5	
10.07 ^k	$\frac{3}{2}^+$	6.32	$\frac{3}{2}^-$	4.9 ± 1.2	
		7.30	$\frac{3}{2}^+$	2.1 ± 0.8	
		0	$\frac{1}{2}^-$	96.0 ± 0.7	
10.45 ^l	$\frac{5}{2}^-$	5.27 + 5.30		4.0 ± 0.7	
		5.27	$\frac{5}{2}^+$	55.0 ± 0.8	$+0.021 \pm 0.033$
10.53 ^m	$\frac{5}{2}^+$	6.32	$\frac{3}{2}^-$	31.3 ± 1.7	-0.59 ± 0.13
		7.16	$\frac{5}{2}^+$	5.2 ± 0.1	$+0.13^{+0.03}_{-0.04}$
		8.57	$\frac{3}{2}^+$	3.8 ± 0.6	-0.3 ± 0.4
		9.152	$\frac{3}{2}^-$	4.7 ± 0.1	$-0.32^{+0.09}_{-0.10}$
10.53 ^m	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	< 0.1	
		5.27	$\frac{5}{2}^+$	38.7 ± 0.2	-0.27 ± 0.03
		6.32	$\frac{3}{2}^-$	7.7 ± 0.1	-0.028 ± 0.004
		7.16	$\frac{5}{2}^+$	19.4 ± 0.2	$+0.007^{+0.010}_{-0.008}$
		7.30	$\frac{3}{2}^+$	31.4 ± 0.5	$+0.066 \pm 0.005$

Table 15.5: Radiative decays in $^{15}\text{N}^a$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Mult. mixing ratio δ
10.69 ^m	$\frac{9}{2}^+$	8.57	$\frac{3}{2}^+$	2.4 ± 0.1	$+0.012^{+0.006}_{-0.005}$
		9.152	$\frac{3}{2}^-$	0.3 ± 0.1	$-0.20^{+0.03}_{-0.02}$
		5.27	$\frac{5}{2}^+$	61.6 ± 0.3	
		7.16	$\frac{5}{2}^+$	2.1 ± 0.1	-0.03 ± 0.07
10.70 ^m	$\frac{3}{2}^-$	7.57	$\frac{7}{2}^+$	36.3 ± 0.6	$+0.118 \pm 0.008$
		0	$\frac{1}{2}^-$	52.6 ± 0.8	$+0.180^{+0.006}_{-0.002}$
		5.27	$\frac{5}{2}^+$	37.4 ± 0.6	$-0.24^{+0.004}_{-0.008}$
		5.30	$\frac{1}{2}^+$	0.8 ± 0.1	-0.13 ± 0.07
		6.32	$\frac{3}{2}^-$	3.8 ± 0.1	$+0.135 \pm 0.015$
		7.16	$\frac{5}{2}^+$	0.4 ± 0.1	0.3 ± 0.3
		7.30	$\frac{3}{2}^+$	2.3 ± 0.1	-0.027 ± 0.023
		8.31	$\frac{1}{2}^+$	0.8 ± 0.1	$-0.017^{+0.018}_{-0.016}$
		9.05	$\frac{1}{2}^+$	0.2 ± 0.1	-0.007 ± 0.12
		9.152	$\frac{3}{2}^-$	0.2 ± 0.1	-0.11 ± 0.03
10.80 ⁿ	$\frac{3}{2}^+$	9.23	$\frac{1}{2}^-$	1.5 ± 0.1	$+0.049^{+0.006}_{-0.005}$
		0	$\frac{1}{2}^-$	51.5 ± 0.4	-0.02 ± 0.01
		5.27	$\frac{5}{2}^+$	4.9 ± 0.1	-0.63 ± 0.04
		5.30	$\frac{1}{2}^+$	15.5 ± 0.2	-0.55 ± 0.02
		6.32	$\frac{3}{2}^-$	5.4 ± 0.2	-0.07 ± 0.05
		7.16	$\frac{5}{2}^+$	7.8 ± 0.1	$+0.14 \pm 0.03$
		7.30	$\frac{3}{2}^+$	5.8 ± 0.1	-0.12 ± 0.02
		8.31	$\frac{1}{2}^+$	3.6 ± 0.1	$+0.12 \pm 0.03$
		9.05	$\frac{1}{2}^+$	0.3 ± 0.1	
		9.152	$\frac{3}{2}^-$	0.9 ± 0.1	
11.62 ^o	$\frac{1}{2}^+; T = \frac{3}{2}$	9.155	$\frac{5}{2}$	4.2 ± 0.1	
		0	$\frac{1}{2}^-$	90.7 ± 3.0	
		5.27	$\frac{5}{2}^+$	< 1	
		5.30	$\frac{1}{2}^+$	7.4 ± 1.5	
12.52	$\frac{5}{2}^+; T = \frac{3}{2}$	6.32	$\frac{3}{2}^-$	1.9 ± 1.5	
		0	$\frac{1}{2}^-$	< 1	

Table 15.5: Radiative decays in ^{15}N ^a (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Mult. mixing ratio δ
13.39 ^p	$\frac{3}{2}^+$	5.27	$\frac{5}{2}^+$	94.2 ± 0.6	-0.02 ± 0.04
		5.30	$\frac{1}{2}^+$	< 1	
		6.32	$\frac{3}{2}^-$	5.8 ± 0.6	-0.02 ± 0.04
		0	$\frac{1}{2}^-$	100	

^a See also Tables 15.12 and 15.15, and 15.6 in 1986AJ01). For references see Table 15.4 in (1981AJ01). Please note that (1976BE1B) is an unpublished Ph.D. thesis.

^b Transitions to $^{15}\text{N}^*(5.27, 5.30)$ are < 0.1 and $< 0.05\%$, respectively (1975MO28).

^c Transitions to $^{15}\text{N}^*(0, 5.30, 6.32)$ are < 0.1 , < 4 and $< 0.5\%$.

^d Transitions to $^{15}\text{N}^*(5.30, 6.32)$ are < 4 and $< 0.6\%$.

^e Transitions to $^{15}\text{N}^*(5.30, 7.30, 7.57)$ are < 12 , < 0.7 and $< 3\%$.

^f Transitions to $^{15}\text{N}^*(7.16, 7.57, 8.31)$ are < 10 , < 2 and $< 0.5\%$.

^g Transitions to $^{15}\text{N}^*(7.16, 7.57, 8.31)$ are < 1 , < 20 and $< 5\%$.

^h Transitions to $^{15}\text{N}^*(7.30, 8.31, 8.57)$ are < 2 , < 1 and $< 2\%$.

ⁱ Transitions to $^{15}\text{N}^*(0, 5.30)$ are < 4 and $< 15\%$.

^j Transitions to $^{15}\text{N}^*(7.16, 7.57, 8.31, 8.57)$ are each $< 1\%$.

^k For upper limits for transitions to other states see Table 15.4 in (1981AJ01).

^l Transitions to $^{15}\text{N}^*(0, 5.30, 9.83)$ are < 12 , < 2 and $< 0.1\%$. See also (1990GO25).

^m See also (1990GO25).

ⁿ π is + because if π were $-$ the Γ_γ and δ of the $10.80 \rightarrow 5.30$ MeV transition would lead to an unacceptably high M2 value (33 W.u.) (P.M. Endt, private communication). See also (1990GO25).

^o See footnote ^g in Table 15.4 (1981AJ01).

^p $\Gamma_{\gamma_0} = 3.0 \pm 0.9$ eV, $\Gamma_p \Gamma_{\gamma_0} / \Gamma = 1.70 \pm 0.5$ eV; $\delta = 0.00 \pm 0.04$ (M2/E1); $B(E1) = (1.2 \pm 0.4) 10^{-3} e^2 \cdot \text{fm}^2$. Transitions to $^{15}\text{N}^*(5.27, 5.30)$ are $< 8\%$ and to $^{15}\text{N}^*(6.32, 7.16, 7.30)$ are $< 5\%$.

See (1985MC1C; applied).

$$3. \text{}^9\text{Be}({}^{12}\text{C}, \text{}^6\text{Li}){}^{15}\text{N} \quad Q_m = -2.8395$$

See (1988GO1H; $E({}^{12}\text{C}) = 65$ MeV; prelim.).

$$4. \text{}^{10}\text{B}({}^6\text{Li}, \text{p}){}^{15}\text{N} \quad Q_m = 18.7462$$

Table 15.6: Resonances in $^{11}\text{B}(\alpha, \gamma_0)^{15}\text{N}$ ^a

E_α (MeV)	E_x (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	Γ_γ (eV)	J^π
7.20	16.27 ± 0.04	240 ± 30	≥ 11	$\frac{3}{2}^+$
7.70	16.64 ± 0.04	250 ± 30	≥ 11	$\frac{1}{2}^+$
8.40 ^b	17.15 ± 0.05	250 ± 60	≥ 2	$(\frac{1}{2}^+, \frac{3}{2}^+)$
9.11 ^b	17.67 ± 0.05	600 ± 80	≥ 7	$\frac{3}{2}^+$
10.80 ^c	18.91 ± 0.15	750 ± 70		$\frac{3}{2}^+ + \frac{1}{2}^+$
14.00 ^c	21.25 ± 0.15	1740 ± 150		$\frac{3}{2}^+ + \frac{1}{2}^+$

^aFor references and other information see Table 15.7 in (1986AJ01).

^b These E_α may be 100 keV too high.

^c There is indication of M1/E2 transitions interfering with the predominant E1 transitions.

At $E(^6\text{Li}) = 4.9$ MeV, thirty proton groups are observed corresponding to ^{15}N states with $E_x < 16.8$ MeV. Angular distributions have been measured for the proton groups corresponding to $^{15}\text{N}^*(5.27 + 5.30, 6.32, 7.16 + 7.30, 7.57, 8.31, 8.57, 9.05 + 9.15)$: see (1976AJ04).

5. $^{10}\text{B}(^7\text{Li}, \text{d})^{15}\text{N}$ $Q_m = 13.7207$

At $E(^7\text{Li}) = 24$ MeV angular distributions have been studied to many of the ^{15}N states with $E_x < 15.5$ MeV: see (1981AJ01).

6. $^{11}\text{B}(\alpha, \gamma)^{15}\text{N}$ $Q_m = 10.9916$

The 90° differential cross section for γ_0 production has been measured for $E_\alpha = 5.74$ to 18.0 MeV: see (1981AJ01, 1986AJ01). For the observed resonances see Table 15.6. See also (1988WAZY; prelim.).

7. (a) $^{11}\text{B}(\alpha, \text{n})^{14}\text{N}$ $Q_m = 0.1583$ $E_b = 10.9916$
 (b) $^{11}\text{B}(\alpha, \text{p})^{14}\text{C}$ $Q_m = 0.7842$

Reported resonances are displayed in Table 15.7. Nine resonances have been observed in the total cross section for reaction (a) in the range $E_\alpha = 0.55$ to 2.40 MeV (1988WAZY; prelim.)

[astrophysical reaction rates will be derived]. For thick target neutron yields for $E_\alpha = 1.0$ to 9.8 MeV, see the review in (1989HE04). See also (1987EL1B; applied).

The total cross section for reaction (b) has been measured for $E_\alpha = 0.9$ to 1.7 MeV: resonance information is deduced by (1987TU01). At higher energies (to 25 MeV) the p_0 excitation functions show broad features: see (1981AJ01).

Table 15.7: Resonances in $^{11}\text{B} + \alpha$ ^a

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particle out	J^π	E_x (MeV)
0.606 ^b		γ, n	$\frac{7}{2}$	11.436
01.07 \pm 20 ^c		n, p		11.78
1.20 \pm 10 ^c		n, p		11.87
1.32 \pm 10 ^c		n, p		11.96
1.50 \pm 10 ^c		n, p	$(\frac{5}{2}^+)$	12.09
1.57 \pm 10 ^c	41 \pm 5	n, p	$(\frac{3}{2}^-)$	12.14
2.056 \pm 10	34 \pm 5	n_0, p_0	$\frac{5}{2}^+$	12.499
2.610 \pm 13	56 \pm 11	n_0, p_0, α	$\frac{3}{2}^-$	12.905
2.66 \pm 30	81	p_0, α	$\frac{5}{2}^+$	12.94
2.942 \pm 10	7 \pm 3	n_0, p_0		13.149
2.984 \pm 10	7 \pm 3	n_0, p_0		13.180
3.239 \pm 15	16 \pm 8	n_0, p, α	$\frac{3}{2}^-$	13.366
3.31 \pm 30	61	p, α	$\frac{3}{2}^+$	13.42
3.46 \pm 30	85 \pm 30	n_0, α	$\frac{3}{2}^-$	13.53
3.560 \pm 10	18 \pm 4	n_0, p	$(\frac{5}{2}, \frac{7}{2})^-$	13.602
3.57 \pm 30	94	α	$\frac{1}{2}^+$	13.61
3.712 \pm 10	26 \pm 8	n_0		13.713
(3.78 \pm 30)	70	α	$(\frac{1}{2}^+)$	(13.76)
3.89 \pm 30	\approx 70	n_1, α	$(\frac{3}{2}^+)$	13.84
4.09 \pm 30	\approx 100	n_1		13.99
4.232 \pm 10	22 \pm 6	n_0		14.094
4.24 \pm 30	\approx 100	n_1, α	$\frac{3}{2}^+$	14.10
4.324 \pm 10	27 \pm 6	n_0		14.162
4.43 \pm 40	150	α	$\frac{5}{2}^+$	14.24
4.62 \pm 40	100	α	$\frac{7}{2}^+$	14.38
4.85 \pm 20	200 \pm 50	n_0		14.55

Table 15.7: Resonances in $^{11}\text{B} + \alpha$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particle out	J^π	E_x (MeV)
4.986 \pm 10	33 \pm 6	n ₀		14.647
5.11 \pm 30	110 \pm 50	n ₀		14.74
5.28 \pm 20	48 \pm 11	n ₀ , α		14.86
5.358 \pm 10	12 \pm 3	n ₀		14.920
5.501 \pm 10	13 \pm 3	n ₀		15.025
5.59 \pm 20	80 \pm 25	n ₀ , α		15.09
5.860 \pm 10	22 \pm 6	n ₀ , α		15.288
5.98 \pm 20	75 \pm 25	n ₂ , (α)		15.38
6.06 \pm 20	\approx 100	n ₀ , (α)		15.43
6.19 \pm 20	\approx 35	n ₀		15.53
6.29 \pm 20	95 \pm 25	n ₂		15.60
(6.65 \pm 40)		(α)		(15.87)
6.73 \pm 20	35 \pm 10	n ₀ , n ₂		15.93
6.755 \pm 15	21 \pm 6	n ₁		15.944
6.83 \pm 20	60 \pm 20	n ₂		16.00
6.884 \pm 15	62 \pm 12	n ₀ , α		16.039
(6.98 \pm 40)		(α)		(16.11)
7.18 \pm 20	\approx 100	n ₀ , α		16.26
7.27 \pm 20	\approx 30	n ₀		16.32
7.37 \pm 20	44 \pm 11	n ₂		16.39
7.616 \pm 15	27 \pm 15	n ₀ , (n ₂)		16.576
7.754 \pm 15	60 \pm 10	n ₀ , (n ₂)		16.677

^a For references see Table 15.7 in (1981AJ01).

^b (1988WAZY; prelim.): $\Gamma < 0.2$ keV.

^c (1987TU01); $J^\pi = \frac{3}{2}^{(-)}, \frac{3}{2}^{-}, \frac{1}{2}^{-}, \frac{5}{2}^{+}, \frac{3}{2}^{-}$ [see also for partial widths].

8. (a) $^{11}\text{B}(\alpha, \text{d})^{13}\text{C}$

$$Q_{\text{m}} = -5.1677$$

$$E_{\text{b}} = 10.9916$$

(b) $^{11}\text{B}(\alpha, \text{t})^{12}\text{C}$

$$Q_{\text{m}} = -3.8568$$

The yield of d_0 has been measured for $E_\alpha = 13.5$ to 25 MeV. The excitation functions for t_0 and t_1 (to 25 MeV) show strong uncorrelated structures: see (1976AJ04, 1981AJ01). See also (1989VA07).



Observed resonances are shown in Table 15.7.



At $E(^6\text{Li}) = 34$ MeV angular distributions are reported to the states with $5.3 < E_x < 16.3$ MeV: this reaction appears to be less selective than reaction 11. The most strongly populated states are $^{15}\text{N}^*(9.2, 10.5, 10.7, 13.1, 14.8, 15.5)$. See (1981AJ01). See also (1990AZZZ).



At $E(^7\text{Li}) = 24$ and 34 MeV, angular distributions to states with $5.3 < E_x < 15.6$ MeV have been measured: $^{15}\text{N}^*(9.8, 10.5, 10.7, 15.4, 15.5)$ are particularly strongly populated at 34 MeV. $J^\pi = \frac{9}{2}^+, \frac{9}{2}, \frac{11}{2}, \frac{9}{2}, \frac{11}{2}, \frac{13}{2}, \frac{15}{2}$ are suggested for $^{15}\text{N}^*(10.69, 12.56, 13.03, 13.19, 13.84, 14.11, 15.37)$. Only $^{15}\text{N}^*(15.52)$ appears to have a large cluster component corresponding to $^{11}\text{B} + \alpha$. See (1981AJ01). For a study of the γ -decay, see (1981AJ01). At $E(^7\text{Li}) = 34, 40, 45$ and 55 MeV states at $E_x = 13.88, 17.10, 18.67, 18.81, 19.70, 19.93$ and 22.86 MeV are reported to be strongly populated (1990AZZZ; prelim.). See also (1990DA03).



Gamma-ray cross sections involving $^{15}\text{N}^*(5.3, 6.32, 7.16, 7.30, 7.57, 8.57)$ are reported at $E_{c.m.} = 1.92, 2.30$ and 2.46 MeV (1986CU02). See (1984DA17) for cross sections and S -factors.



Angular distributions have been measured at $E(^{16}\text{O}) = 27$ to 60 MeV involving the two proton-hole states of ^{15}N [$^{15}\text{N}^*(0, 6.32)$; $J^\pi = \frac{1}{2}^-, \frac{3}{2}^-$] and $^{12}\text{C}^*(0, 4.4, 9.6)$: see (1976AJ04). See also (1989KA1N; theor.).

14. (a) $^{12}\text{C}(t, \gamma)^{15}\text{N}$	$Q_m = 14.8484$	
(b) $^{12}\text{C}(t, n)^{14}\text{N}$	$Q_m = 4.0151$	$E_b = 14.8484$
(c) $^{12}\text{C}(t, p)^{14}\text{C}$	$Q_m = 4.6410$	
(d) $^{12}\text{C}(t, t)^{12}\text{C}$		
(e) $^{12}\text{C}(t, \alpha)^{11}\text{B}$	$Q_m = 3.8568$	

The 90° excitation function for γ_0 in the range 1.0 to 6.5 MeV [see (1981AJ01, 1986AJ01)] shows one very strong resonance (at peak, $4.4 \pm 0.5 \mu\text{b/sr}$) corresponding to $^{15}\text{N}^*(16.7)$ as well as two other strong (unresolved and/or broad) resonances at $E_t \approx 3.3$ and 6 MeV: Table 15.8 shows the derived parameters. Table 15.8 also displays the structures observed in reactions (b)→(e). At $E_t = 17$ MeV the polarization and analyzing power for the transition to $^{14}\text{C}_{\text{g.s.}}$ (reaction (c)) are shown to be the same, as required by the conservation of parity. The VAP for the elastic scattering (reaction (d)) has been measured at $E_t = 9$ and 11 MeV: see (1985AJ01). See (1981AJ01) for the earlier work. See also (1985SA31, 1990HA46; theor.).

15. $^{12}\text{C}(^3\text{He}, \pi^+)^{15}\text{N}$ $Q_m = -124.7387$

Individual states have not been resolved in this reaction. The cross section over the bound states of ^{15}N is < 0.03 nb at $E_{\pi^+} = 5$ MeV and 0.8 ± 0.2 nb at $E_{\pi^+} \approx 60$ MeV [$E(^3\text{He}) = 170.2$ and 236.3 MeV, respectively] (1988HO15). For the earlier work see (1984BI08, 1986SC23).

16. $^{12}\text{C}(\alpha, p)^{15}\text{N}$ $Q_m = -4.9656$

Angular distributions have been measured at many energies for $E_\alpha = 13.4$ to 96.8 MeV: see (1976AJ04, 1981AJ01, 1986AJ01). See also (1987MIZY, 1988BRZY; $E_\alpha = 48$ MeV; prelim.), (1987BI1C; $E_\alpha = 218$ MeV; prelim.) and (1989BR1J).

17. $^{12}\text{C}(^6\text{Li}, ^3\text{He})^{15}\text{N}$ $Q_m = -0.9471$

Observed ^3He groups are displayed in Table 15.9 of (1981AJ01). Comparisons of the angular distributions obtained in this reaction at $E(^6\text{Li}) = 60.1$ MeV and in the $(^6\text{Li}, t)$ reaction shows analog correspondence for the following pairs of levels: 5.27 – 5.24, 7.16 – 6.86, 7.57 – 7.28, 8.57 – 8.28, 10.80 – 10.48, 13.15(u) – 12.84, 15.49(u) – 15.05 [first listed is E_x in ^{15}N -second in ^{15}O]. [E_x are nominal; u = unresolved.] For γ -decay measurements see Table 15.5. See also (1990AZZZ).

Table 15.8: Resonances in $^{12}\text{C} + t$ ^a

E_t (MeV \pm keV)	Particles out	J^π	Γ (keV)	E_x (MeV)
0.66	α_0			15.38
1.11	p_0, t_0, α_1			15.74
1.21	t_0			15.82
1.30 ± 20	n, α_0			15.89
1.39 ± 20	n, t_0, α_0			15.96
1.46	p_0			16.02
1.54	n, α_0, α_1			16.08
1.64 ± 40	γ_0, n, α_0	$\frac{3}{2}^+$	450 ± 100	16.16
1.78	α_0			16.27
1.85 ± 20	$n, p_0, \alpha_0, \alpha_1$			16.33
1.98 ± 20	n, p_0			16.43
2.05 ± 30	p_0, t_0, α_0			16.49
2.18 ± 25	$\gamma_0, n, p_0, t_0, \alpha_0, \alpha_1$	$\frac{3}{2}^-$	490	16.59
2.30	$\gamma_0, n, p_0, \alpha_0, \alpha_1$	$\frac{3}{2}^+$	130 ± 15	16.69 ± 0.01
2.39 ± 30	$n, t_0, \alpha_0, \alpha_1$			16.76
2.50 ± 30	α_0, α_1			16.85
2.60	α_0			16.93
2.75	p_0			17.05
2.82	$\gamma_0, t_0, \alpha_0, \alpha_1$	$\frac{3}{2}^-$		17.10
2.89 ± 50	α_0			17.16
3.14	α_1			17.36
3.30	γ_0	$\frac{3}{2}^+$	450 ± 120	17.49 ± 0.09
≈ 6	γ_0			19.6
15.0	t_0			26.8

^a For references see Tables 15.8 in (1976AJ04, 1981AJ01) and 15.9 in (1986AJ01).

Table 15.9: States of ^{15}N from $^{12}\text{C}(^7\text{Li}, \alpha)$

E_x (MeV \pm keV)		E_x (MeV \pm keV)		E_x (MeV \pm keV)	
(1973TS02) ^a	(1980ZE02) ^b	(1973TS02) ^a	(1980ZE02) ^b	(1973TS02) ^a	(1980ZE02) ^b
0		10.808		15.021	15.024
5.295	5.284		11.274	15.373	15.379
6.332	6.323	11.430	11.456	15.782	15.778
7.163	7.157	11.951	11.936	16.026	16.032
7.310	7.299	12.320 ^a	12.328	16.190	16.210
7.566	7.574	12.559 ^{a, c}	12.551		17.735
8.320		12.923			17.949 ^b
8.580 ^a	8.574	13.004 ^a	13.001		18.272
9.163 ^a	9.159	13.173 ^a	13.178		18.698 ^b
9.828 ^a	9.809	13.614			19.27 \pm 40
9.932	9.921	14.087	14.097		19.68 \pm 50 ^{b, d}
10.072	10.075	14.720	14.693		20.93 \pm 50 ^{b, d}
10.524	10.518		14.874		24.75 \pm 150 ^{b, d}
10.700 ^a	10.714				

^a $E(^7\text{Li}) = 35$ MeV; angular distributions have been measured for the states labelled by this footnote; $E_x \pm 10$ keV.

^b $E(^7\text{Li}) = 48$ MeV; angular distributions have been measured for the states labelled by this footnote; $E_x \pm 20$ keV unless otherwise shown.

^c (1973TS02) suggest that this state is not the $T = \frac{3}{2}$ state at 12.52 MeV.

^d Wide or unresolved.

18. $^{12}\text{C}(^7\text{Li}, \alpha)^{15}\text{N}$

$$Q_m = 12.3806$$

Observed α -groups are shown in Table 15.9. Angular distributions have been measured to $E(^7\text{Li}) = 48$ MeV. Comparison of spectra from this reaction ($E(^7\text{Li}) = 34.9$ MeV) with those from $^{13}\text{C}(^6\text{Li}, \alpha)$ (reaction 26) lead to configurations of (d)³ for $^{15}\text{N}^*(10.7, 12.57, 13.20, 15.42)$ and suggest that $^{15}\text{N}^*(12.57, 13.20)$ have lower J than $^{15}\text{N}^*(10.7, 15.5)$, probably $J \leq \frac{7}{2}$. $^{15}\text{N}^*(13.02)$ is shown to be p(d)² in agreement with $J^\pi = \frac{11}{2}^-$: see (1981AJ01).

$^{15}\text{N}^*(9.155)$ [$J = \frac{5}{2}$] decays to $^{15}\text{N}^*(5.30)$ [$J = \frac{1}{2}^+$] by an E2 transition; therefore its parity is positive. It has a large triton cluster parentage. This is not true of $^{15}\text{N}^*(9.152)$: see (1981AJ01). For γ -decay measurements see Table 15.5. See also (1985SA31; theor.).

19. (a) $^{12}\text{C}(^{11}\text{B}, ^8\text{Be})^{15}\text{N}$ $Q_m = 3.6250$
 (b) $^{12}\text{C}(^{13}\text{C}, ^{10}\text{B})^{15}\text{N}$ $Q_m = -9.0275$
 (c) $^{12}\text{C}(^{18}\text{O}, ^{11}\text{B}\alpha)^{15}\text{N}$ $Q_m = -11.9768$

For reaction (a) see (1981AJ01) and (1988MA07). For reaction (b) see (1989VO1D). For reaction (c) see (1984RA07).

20. $^{13}\text{C}(\text{d}, \gamma)^{15}\text{N}$ $Q_m = 16.1594$

The $90^\circ - 95^\circ$ yields of γ_0 have been measured for $E_d = 1$ to 10 MeV: observed resonances are displayed in Table 15.10. The γ -ray angular distributions are consistent with the emission of predominantly E1 radiation except for evidence of M1/E2 transitions in the region $E_x = 20 - 21.5$ MeV: see (1981AJ01). See also (1990HA46).

21. (a) $^{13}\text{C}(\text{d}, \text{n})^{14}\text{N}$ $Q_m = 5.3260$ $E_b = 16.1594$
 (b) $^{13}\text{C}(\text{d}, \text{p})^{14}\text{C}$ $Q_m = 5.9519$
 (c) $^{13}\text{C}(\text{d}, 2\text{p})^{13}\text{B}$ $Q_m = -14.880$

Observed resonances are displayed in Table 15.10. Polarization measurements have been carried out at $E_d = 12.3$ MeV (reaction (a)) and 13 and 56 MeV (reaction (b)): see (1986AJ01). See also (1987AB04). For VAP measurements (reaction (c)) at $E_d = 70$ MeV to $^{13}\text{B}_{\text{g.s.}}$ see (1986MO27).

22. $^{13}\text{C}(\text{d}, \text{d})^{13}\text{C}$ $E_b = 16.1594$

Excitation functions for elastically scattered deuterons have been measured in the range $E_d = 0.4$ to 5.7 MeV: see (1976AJ04). Polarization studies have been reported for $E_d = 12.5$ to 15 MeV and at $E_d = 56$ MeV: see (1981AJ01, 1986AJ01).

23. (a) $^{13}\text{C}(\text{d}, \text{t})^{12}\text{C}$ $Q_m = 1.3109$ $E_b = 16.1594$
 (b) $^{13}\text{C}(\text{d}, ^3\text{He})^{12}\text{B}$ $Q_m = -12.040$
 (c) $^{13}\text{C}(\text{d}, \alpha)^{11}\text{B}$ $Q_m = 5.1677$

Observed resonances are listed in Table 15.10. For polarization measurements to $E_d = 29$ MeV [reactions (a), (b)] see (1981AJ01).

Table 15.10: Resonances in $^{12}\text{C} + \text{d}$ ^a

E_{d} (MeV)	Particles out	Γ_{lab} (keV)	$^{15}\text{N}^*$ (MeV)
0.37	p		16.48
0.64	n, p ₀ , t ₀	≈ 100	16.71
0.85	n, p ₀	≈ 400	16.90
1.10	α ₀	broad	17.11
1.24 ± 0.04	t ₀ , (α ₀)	≈ 200	17.23
1.40 ± 0.04	p ₀ , t ₀ , α ₀	≈ 400	17.37
1.64 ± 0.04	t ₀	≈ 200	17.58
1.74 ± 0.04	γ ₀ , n, α ₀	≈ 600	17.67 ^b
1.80 ± 0.01	(p ₀), t ₀ , α ₁	55 ± 10	17.72
2.20 ± 0.01	(n), α ₀ , α ₁	22 ± 4	18.06
2.23 ± 0.02	(n), p ₀ , t	≈ 50	18.09
2.45 ± 0.03	n, p ₀ , α ₀	270 ± 70	18.28
3.46 ± 0.03	n	≈ 150	19.16
5.1	n ₁ , p ₀	≈ 50	20.6
6.65	γ ₀	≈ 700	21.92
8.8	γ ₀	broad	23.8

^a See references listed in Tables 15.10 (1976AJ04, 1981AJ01).

^b $J^{\pi} = \frac{1}{2}^{-}$ or $\frac{3}{2}^{+}$; $T = \frac{1}{2}$.

$$24. \ ^{13}\text{C}(^3\text{He}, \text{p})^{15}\text{N} \quad Q_{\text{m}} = 10.6658$$

Observed proton groups and γ -rays are listed in Table 15.11 of (1981AJ01). Angular distributions have been reported for $E(^3\text{He}) = 4.37$ to 20 MeV: see (1981AJ01).

$$25. \ ^{13}\text{C}(\alpha, \text{d})^{15}\text{N} \quad Q_{\text{m}} = -7.6874$$

At $E_{\alpha} = 34.9$ MeV a ZRDWBA analysis has been made of the angular distributions to $^{15}\text{N}^*(5.27, 5.30, 7.16, 7.30, 7.56, 8.31, 8.57, 9.05, 9.15, 10.07, 10.53, 10.69, 11.43, 11.94, 12.10, 12.33, 12.49, 12.56, 13.00, 13.83, 14.08)$. $L = 0$ for the group(s) to $^{15}\text{N}^*(9.15, 10.69)$; $L = 2$ for $^{15}\text{N}^*(12.56)$; $L = 3$ for $^{15}\text{N}^*(5.27, 7.16, 7.56)$; $L = 4$ for $^{15}\text{N}^*(11.94, 13.00)$; $L = 1$ for the remaining transitions (1984YA03). See also Table 15.11 of (1976AJ04).

26. $^{13}\text{C}(^6\text{Li}, \alpha)^{15}\text{N}$ $Q_m = 14.6843$

Angular distributions have been measured at $E(^6\text{Li}) = 32$ MeV to $^{15}\text{N}^*(0, 5.30, 6.32, 7.16, 7.30, 7.57, 8.31, 8.57, 9.15, 9.23, 9.83, 10.07, 10.70, 11.94, 13.00)$: the results are consistent with the previously known J^π , with (odd) parity for $^{15}\text{N}^*(9.83)$ and with $J^\pi = \frac{9}{2}^-$ for $^{15}\text{N}^*(11.94)$: see (1981AJ01).

27. (a) $^{13}\text{C}(^{10}\text{B}, ^8\text{Be})^{15}\text{N}$ $Q_m = 10.1328$

(b) $^{13}\text{C}(^{11}\text{B}, ^9\text{Be})^{15}\text{N}$ $Q_m = 0.3440$

For reaction (a) see (1988MA07). For reaction (b) see (1981AJ01).

28. $^{14}\text{C}(\text{p}, \gamma)^{15}\text{N}$ $Q_m = 10.2074$

Observed resonances are displayed in Table 15.11; the branching ratios are shown in Table 15.5. Narrow anomalies (in the γ_0 yield for $E_p = 2.8$ to 30 MeV) are reported at $E_p = 10.0, 11.0, 12.35, 13.6, 16.4$ MeV. A good fit to the total cross section ($E_{\bar{p}} = 7.5$ to 19 MeV) is obtained with the GDR split into peaks at $E_x = 21.0$ and 25.5 MeV with $\Gamma = 6$ and 2 MeV, respectively. The integrated E2 cross section for $E_x = 19.5$ to 27.0 MeV is $(6.8 \pm 1.4)\%$ of the isoscalar sum rule. The reaction thus shows no sign of a collective E2 resonance in that E_x region. [Another study shows no appreciable E2 strength concentration for $E_x = 14.3$ to 23.3 MeV.] Above the GDR region the 90° γ_0 cross section decreases smoothly with energy except for a small peak which would correspond to $^{15}\text{N}^*(37.0)$. See (1981AJ01, 1986AJ01) for the references. See also (1985CA41, 1988CA26, 1990GO25; astrophysics) and (1990HA46; theor.).

29. $^{14}\text{C}(\text{p}, \text{n})^{14}\text{N}$ $Q_m = -0.6259$ $E_b = 10.2074$

Observed resonances are displayed in Table 15.11. Cross sections have recently been measured for $E_p = 0.67$ to 1.20 MeV (1989KEZZ; prelim.). Polarization measurements are reported at $E_{\bar{p}} = 160$ MeV (1984TA07, 1987RA15; A_y ; $D_{\text{NN}}(0^\circ)$; n to $^{14}\text{N}^*(0, 2.31, 3.95, 13.72)$). Forward-angle differential cross sections have been measured at $E_p = 200, 300, (400), 450$ MeV (1986AL18, 1989AL04; $n_1 + n_2$) and at 492 MeV (1989RA09; n_1, n_2). See (1986AJ01) for the earlier work. See also (1985TA23), (1986TA1E, 1987TA22, 1989SU1J), (1985CA41, 1988CA26; astrophysics) and (1987BE1D, 1987LO1D; theor.).

30. (a) $^{14}\text{C}(\text{p}, \text{p})^{14}\text{C}$ $E_b = 10.2074$

(b) $^{14}\text{C}(\text{p}, \alpha)^{11}\text{B}$ $Q_m = -0.7842$

Table 15.11: Resonances in $^{14}\text{C} + \text{p}$ ^a

E_p (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Γ_n (keV)	Γ_p (keV)	Γ_α (keV)	Γ_γ (eV)	J^π	E_x (MeV \pm keV)
0.261 ± 0.6 ^f	< 0.5		$(0.08 \pm 0.01) \times 10^{-6}$		$(0.29 \pm 0.05) \text{ meV}$ ^b	$\frac{5}{2}^-$	10.4497 ± 0.3 ^d
0.352 ± 1 ^f					$37 \pm 6 \text{ meV}$ ^b	$\frac{5}{2}^+$	10.5333 ± 0.5 ^d
0.519 ± 1 ^f			$(0.49 \pm 0.10) \times 10^{-6}$		$3.1 \pm 0.5 \text{ meV}$ ^b	$\frac{9}{2}^+$	10.6932 ± 0.3 ^d
0.527 ± 1 ^f			0.2		0.37 ± 0.07 ^g	$\frac{3}{2}^-$	10.7019 ± 0.3 ^d
0.634 ± 1 ^f			$(0.22 \pm 0.10) \times 10^{-3}$		0.27 ± 0.14 ^h	$\frac{3}{2}^{(+)}$	10.804 ± 2 ^d
1.162 ± 2	7.9 ± 3	2.3	5.6	< 0.3	0.29 ^c	$\frac{1}{2}^-$	11.291
1.3188 ± 0.5	41.4 ± 1.1	34.6 ± 0.9	6.8 ± 0.5	< 0.3	4.2 ± 0.7 ^c	$\frac{1}{2}^+$	11.4376
1.509 ± 4	404.9 ± 6.3	4.0 ± 0.2	400.9 ± 6.3	< 0.3	19.2 ± 0.4 ^c	$\frac{1}{2}^+; T = \frac{3}{2}$	11.615
1.668 ± 3	37	36.5	0.5	< 0.3		$\frac{3}{2}^+$	11.763
1.788 ± 3	24.5	24.5	0.03	< 0.3		$\frac{3}{2}^-, (\frac{5}{2}^-)$	11.875
1.884 ± 3	21.5	21.2	0.3	< 0.3		$\frac{1}{2}^-$	11.965
2.025 ± 4	14 ± 5	12.0	1.7	0.6		$\frac{5}{2}^+$	12.096
2.077 ± 3	47 ± 7	30.2	16.6	2.2		$\frac{3}{2}^-$	12.145
2.272 ± 4	22	21.7	0.3	< 0.3		$\frac{5}{2}^{(+)}$	12.327
2.450 ± 4	44 ± 3	28	0.3	5.5		$\frac{5}{2}^+; T = \frac{1}{2}$	12.493
2.482 ± 8	58 ± 4				4.6 ± 0.7	$\frac{5}{2}^+; T = \frac{3}{2}$	12.523
2.908 ± 4	70	25	9.0	15		$\frac{3}{2}^-$	12.920
2.93 ± 10	81	n.r.	0.5	80		$\frac{5}{2}^+$	12.940
3.19	5.5	r					13.18
3.38 ± 10	24	6	6.0	12		$\frac{3}{2}^-$	13.360
3.421 ± 10	57	20.6	35	5.5	3.0 ± 0.9	$\frac{3}{2}^+$	13.390
3.57 ± 10	124	≈ 75	8.0	≈ 40		$\frac{3}{2}^-$	13.537

Table 15.11: Resonances in $^{14}\text{C} + \text{p}$ ^a (continued)

E_p (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Γ_n (keV)	Γ_p (keV)	Γ_α (keV)	Γ_γ (eV)	J^π	E_x (MeV \pm keV)
3.65 \pm 10	88	≈ 16	12.0	≈ 60		$\frac{1}{2}^+$	13.612
3.71		r					13.67
4.0	930		500		r	$\frac{1}{2}^+$	13.9
4.1 \pm 100	98 \pm 10		25	r		$\frac{5}{2}^+$	14.0
4.2 \pm 100				r		$(\frac{3}{2})$	14.1
4.6 \pm 150	74 \pm 7		20	r	(r)	$\frac{3}{2}^-$	14.5
4.8	149 \pm 18		39	r	(r)	$\frac{3}{2}^+$	14.7
4.83	750				r		14.71
5.08	158 \pm 19		20		r	$\frac{3}{2}^+$	14.95
5.16 \pm 130	28 \pm 3		9.0	r		$\frac{3}{2}^+$	15.0
5.54 \pm 130	39 \pm 5		12	r	(r)	$\frac{3}{2}^-$	15.4
5.62	750				r		15.45
6.4 \pm 150	130 \pm 14		19	r		$\frac{3}{2}^+$	16.2
6.70	560				r		16.46
6.925	90 \pm 10			r	r	$(\frac{3}{2}^+; \frac{1}{2})$	16.67
7.18 \pm 180	110 \pm 50			r		$\frac{5}{2}$	16.9
≈ 9					r	$\frac{1}{2}^+; \frac{1}{2}$	19
10.0	sharp		(1000?)		r	$\frac{3}{2}^+; (T = \frac{3}{2})$	19.5 ^e
11.0	sharp				r	$\frac{3}{2}^+$	20.5
12.35					r		21.72
13.65					r		22.94
16.4					r	$(T = \frac{3}{2})$	25.5 ^e

Table 15.11: Resonances in $^{14}\text{C} + \text{p}$ ^a (continued)

E_{p} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Γ_{n} (keV)	Γ_{p} (keV)	Γ_{α} (keV)	Γ_{γ} (eV)	J^{π}	E_{x} (MeV \pm keV)
≈ 29					r		≈ 37

r = resonant

n.r. = non-resonant

^a See Tables 15.5 in (1959AJ76), 15.11 in (1970AJ04) and 15.12 in (1981AJ01) for references and additional comments.

^b ω_{γ} .

^c Γ_{γ_0} . I am indebted to P.M. Endt for this correction.

^d E_{x} measured directly: see (1981AJ01).

^e Analog not observed in $^{14}\text{N}(\text{p}, \gamma)^{15}\text{O}$.

^f Resonances are observed at $E_{\text{p}} = 262, 351, 520, 528$ and 635 keV [± 1 keV] (1990GO25). See also Table 15.5. I am indebted to Drs. J. Gorres and M. Wiescher for sending me these results prior to publication.

^g $\omega_{\gamma} = 840 \pm 130$ meV (1990GO25).

^h $\omega_{\gamma} = 270 \pm 40$ meV (1990GO25).

Observed resonances and anomalies are displayed in Table 15.11. For polarization measurements see (1981AJ01, 1986AJ01). See also (1989AM01; theor.).

31. $^{14}\text{C}(\text{d}, \text{n})^{15}\text{N}$ $Q_{\text{m}} = 7.9829$

Angular distributions have been measured for $E_{\text{d}} = 1.3$ to 6.5 MeV: see (1976AJ04).

32. $^{14}\text{C}({}^3\text{He}, \text{d})^{15}\text{N}$ $Q_{\text{m}} = 4.7139$

Angular distributions have been studied at $E({}^3\text{He}) = 23$ MeV to $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32, 7.16, 7.30)$: see (1981AJ01). See also (1976AJ04).

33. $^{14}\text{C}({}^7\text{Li}, {}^6\text{He})^{15}\text{N}$ $Q_{\text{m}} = 0.2328$

See (1988AL1G); $E({}^7\text{Li}) = 27$ MeV; prelim.).

34. $^{14}\text{C}({}^{16}\text{O}, {}^{15}\text{N})^{15}\text{N}$ $Q_{\text{m}} = -1.9201$

See (1976AJ04, 1986AJ01).

35. $^{14}\text{N}(\text{n}, \gamma)^{15}\text{N}$ $Q_{\text{m}} = 10.8333$

The thermal cross section is 79.8 ± 1.4 mb (1990ISO5). See also (1988MCZT).

Observed γ -rays from thermal neutron capture are displayed in Table 15.12. See also Table 15.5. The $90^\circ \gamma_0$ yield and angular distributions have been measured for $E_{\text{n}} = 5.6$ to 15.3 MeV. The cross section shows two prominent dips at $E_{\text{x}} = 16.7$ and 18.1 MeV [compare with $^{14}\text{N}(\text{p}, \gamma)$; reaction 9 in ^{15}O] and broad structures at $E_{\text{x}} \approx 17$ and 19 MeV. The angular distribution data are consistent with essentially pure E1 radiation in the region $E_{\text{x}} = 17$ to 24 MeV (1982WE01). See also (1989WO1C) and (1989GU1J; astrophys.).

36. $^{14}\text{N}(\text{n}, \text{n})^{14}\text{N}$ $E_{\text{b}} = 10.8333$

Table 15.12: Gamma radiation from $^{14}\text{N}(n, \gamma)^a$

Transition in ^{15}N	E_γ^b (keV)	E_x^b (keV)	I_γ^c
C → 0	10829.087 (46)	10833.302 (12)	13.65 (21)
C → 5.27	5562.062 (17)		10.65 (12)
C → 5.30	5533.379 (13)		19.75 (21)
C → 6.32	4508.783 (14)		16.54 (17)
C → 7.16	3677.772 (17)		14.89 (15)
C → 7.30	3532.013 (13)		9.24 (9)
C → 8.31	2520.418 (15)		5.79 (7)
C → 9.05			^a
C → 9.152	1681.117 (171)		1.54 (15)
C → 9.155	1678.174 (55)		7.23 (18)
5.27 → 0	5269.169 (12)	5270.155 (10)	30.03 (20)
5.30 → 0	5297.817 (15)	5298.822 (11)	21.31 (18)
6.32 → 0	6322.337 (14)	6323.775 (15)	18.67 (14)
7.16 → 0		7155.051 (16)	
7.16 → 5.27	1884.879 (21)		18.66 (25)
7.16 → 5.30			0.8 (2)
7.30 → 0	7298.914 (33)	7300.832 (16)	9.73 (9)
7.30 → 5.30			^a
8.31 → 0	8310.143 (29)	8312.620 (25)	4.22 (5)
8.31 → 5.30	3013.494 (73)		0.69 (2)
8.31 → 6.32	1988.507 (239)		0.37 (9)
8.57 → 0	8568.920 (230)	8571.412 (120)	0.073 (4)
8.57 → 5.27	3300.728 (113)		0.16 (2)
9.05 → 0	9046.802 (69)	9049.713 (69)	0.186 (5)
9.152 → 0	9149.222 (47)	9151.895 (120)	1.62 (2)
9.155 → 0		9154.895 (23)	
9.155 → 5.27	3884.184 (39)		0.57 (2)
9.155 → 5.30	3855.579 (45)		0.70 (1)
9.155 → 6.32	2830.809 (70)		1.75 (3)
9.155 → 7.16	1999.708 (86)		3.99 (9)

Table 15.12: Gamma radiation from $^{14}\text{N}(n, \gamma)^a$ (continued)

Transition in ^{15}N	E_γ^b (keV)	E_x^b (keV)	I_γ^c
9.222 \rightarrow 0	9219.022 (763)	9222.06 (76)	0.024 (5)
9.925 \rightarrow 0	9921.511 (166)	9925.033 (166)	0.127 (4)
10.066 \rightarrow 0	10062.345 (197)	10065.969 (197)	0.062 (4)

C = capturing state.

^a See also Tables 15.13 in (1981AJ01, 1986AJ01) for earlier references, comments and reports. The previously reported transition to $^{15}\text{N}^*(9.76)$ has not been confirmed: $I_\gamma < 0.01\%$ (T.J. Kennett, private communication). (1990WA22) [see footnote ^b in Table 15.4] recommends different values for E_γ and E_x .

^b Error in Q_m not included. Adjustments due to it require the addition in quadrature of the Q_m error: see (1986KE14).

^c In units of photons/100 captures (1986KE14): errors are statistical only but these are predominant.

 Table 15.13: Resonances in $^{14}\text{N} + n^a$

E_{res} (MeV \pm keV)	Γ_{lab} (keV)	Γ_n (keV)	Γ_p (keV)	Γ_α (keV)	J^π	$^{15}\text{N}^*$ (MeV)
0.430 \pm 5	3.5	< 3	< 0.01		$\geq \frac{3}{2}$	11.235
0.4926 \pm 0.65	7.5	< 3	< 10		$\frac{1}{2}^-$	11.2928
0.639 \pm 5	43	34	9		$\frac{1}{2}^+$	11.429
0.998 \pm 5	46	45	0.8		$\frac{3}{2}^+$	11.764
1.120 \pm 6	19	19	0.20		$\frac{3}{2}^-$	11.878
1.188 \pm 6	≤ 3.2	< 2	< 0.1		$\geq \frac{3}{2}$	11.942
1.211 \pm 7	13	12	0.4		$\frac{1}{2}^-$	11.963
1.350 \pm 7	21	20	0.9	0.4	$\frac{5}{2}^+$	12.093
1.401 \pm 8	54	41	11	1.8	$\frac{5}{2}^+$	12.140
1.595 \pm 8	22	21	0.2	< 0.1	$\frac{5}{2}^-$	12.321
1.779 \pm 10	47	37	0.5	9.0	$(\frac{5}{2}^+)$	12.493
2.23	65	39	7.8	18	$\frac{3}{2}^-$	12.91
2.47	< 3			r		13.14
2.52	≈ 7	r		r		13.18

Table 15.13: Resonances in $^{14}\text{N} + \text{n}$ ^a (continued)

E_{res} (MeV \pm keV)	Γ_{lab} (keV)	Γ_{n} (keV)	Γ_{p} (keV)	Γ_{α} (keV)	J^{π}	$^{15}\text{N}^*$ (MeV)
2.71	40			r	$\frac{3}{2}^-$	13.36
2.74	95		r		$\frac{5}{2}^+$	13.39
2.95	20	16	1.1	3.2	$\frac{5}{2}^+$	13.59
3.09	60		r	r		13.72
3.21	85	r	r	r	$\frac{3}{2}^+$	13.83
3.51	≈ 20	r	r	r		14.11
3.57	30	r	r	r	$\frac{3}{2}^{(+)}$	14.16
≈ 3.8	≈ 2000	≈ 1000	200	≈ 1000		14.4
4.09	50	r	r	r		14.65
≈ 4.2	≈ 300	r	r	r		14.8
4.38	40			r		14.92
4.60		r		r		15.12
5.03				r		15.52
5.60	100			r		16.06
5.94				r		16.37
6.16	75			r		16.58
6.26	100	r		r		16.67
6.55	170	r		r		16.94
6.94	200	r		r		17.31
7.16				r		17.51
7.34	120			r		17.68
7.48	180	r		r		17.81
7.92	170	r		r		18.22
8.00	120			r		18.29

r = resonant.

^a See references in Tables 15.14 in (1970AJ04, 1976AJ04).

The scattering amplitude (bound) $a = 9.37 \pm 0.03$ fm, $\sigma_{\text{free}} = 10.05 \pm 0.12$ b, $\sigma_{\text{inc}}^{\text{spin}}$ (bound nucleus) = 0.49 ± 0.11 b (1979KO26). Observed resonances are displayed in Table 15.13: for a discussion of the evidence leading to J^{π} assignments see (1959AJ76). Cross section curves and a listing of references can be found in (1988MCZT). Recent measurements are reported at

Table 15.14: Beta decay of ^{15}C ^a

Decay to $^{15}\text{N}^*$ (keV)	J^π	Branch (%)	$\log ft$
g.s.	$\frac{1}{2}^-$	36.8 ± 0.8 ^c	5.99 ± 0.03 ^c
5298.87 ± 0.15 ^b	$\frac{1}{2}^+$	63.2 ± 0.8 ^c	4.11 ± 0.01
6323.3 ± 0.6	$\frac{3}{2}^-$	$\leq 0.4 \times 10^{-2}$	≥ 7.8
7301.1 ± 0.5	$\frac{3}{2}^+$	$(0.74 \pm 0.08) \times 10^{-2}$	6.89 ± 0.05
8312.9 ± 0.5	$\frac{1}{2}^+$	$(4.1 \pm 0.5) \times 10^{-2}$	5.18 ± 0.05
8571.4 ± 1.0	$\frac{3}{2}^+$	$(1.3 \pm 0.2) \times 10^{-2}$	5.34 ± 0.07
9050.0 ± 0.7	$\frac{1}{2}^+$	$(3.4 \pm 0.3) \times 10^{-2}$	4.05 ± 0.04

^a (1979AL23).^b (1976AL16). 5297.794 ± 0.035 keV: see (1981WA06).^c (1984WA07).

$E_n = 0.14, 1.3$ and 2.1 MeV (1988KO18; σ_t), 7.67 to 13.50 MeV (1986CH2F; el and inel; prelim.). $10.96, 13.96$ and 16.95 MeV (1985TE01; elastic). The $(n, n'\gamma)$ cross section has been measured for $E_n = 2.5$ to 3.5 MeV (1989STZW; prelim.; applied). Analyzing powers for the n_0 group have been measured for $E_n = 10$ to 17 MeV (1989LI26). See also ^{14}N , (1988BA55, 1986LI1M) and (1988MA1H).

37. $^{14}\text{N}(n, 2n)^{13}\text{N}$

$$Q_m = -10.5535$$

$$E_b = 10.8333$$

Cross sections have been measured for $E_n = 10$ to 37 MeV [see (1988MCZT)] and at $E_n = 13.40$ to 14.87 MeV (1989KA1S). See also (1989KA2B).

38. (a) $^{14}\text{N}(n, p)^{14}\text{C}$

$$Q_m = 0.6259$$

$$E_b = 10.8333$$

(b) $^{14}\text{N}(n, d)^{13}\text{C}$

$$Q_m = -5.3260$$

(c) $^{14}\text{N}(n, t)^{12}\text{C}$

$$Q_m = -4.0151$$

(d) $^{14}\text{N}(n, ^3\text{He})^{12}\text{B}$

$$Q_m = -12.7436$$

(e) $^{14}\text{N}(n, \alpha)^{11}\text{B}$

$$Q_m = -0.1583$$

(1989KO11), using the “white” neutron source LANSCE, have measured the (n, p) cross section from 61 meV to 34.6 keV. Their results support the role of this reaction as a “poison” during s-process nucleosynthesis. [See (1989KO11) for a discussion of other measurements.] See

also (1988BR17, 1989KEZZ). For a display of the measured cross sections for (a) and (c), see (1988MCZT). See also (1988SUZY; $E_n = 5.0$ to 10.6 MeV; $\sigma = 11$ to 30 mb; prelim.) for reaction (c) and (1988MA1H). For resonances in reactions (a) and (e), see Table 15.13. (1986SU15) report double-differential cross sections at $E_n = 27.4, 39.7$ and 60.7 MeV for all five reactions. See also (1989CH2E).

$$39. \text{}^{14}\text{N}(p, \pi^+)\text{}^{15}\text{N} \quad Q_m = -129.518$$

At $E_p = 200$ MeV, angular distributions and A_y have been measured for the transitions to $^{15}\text{N}^*(0, 6.32, 8.31)$ as well as for a number of unresolved transitions. A sharp group at $E_x = 21.5$ MeV is suggested to correspond to a $\frac{15}{2}^-$ state (1988AZ1D, 1987AZZZ; Ph.D. thesis and abstract).

$$40. \text{}^{14}\text{N}(d, p)\text{}^{15}\text{N} \quad Q_m = 8.6087$$

Proton groups (and γ -rays) from this reaction are displayed in Table 15.15 of (1981AJ01). The results include $E_x = 7567.1 \pm 1.0$ keV for $^{15}\text{N}^*(7.57)$. Newer values, derived from measurements of proton groups in a spectrograph, are $5270.2 \pm 1.3, 6324.0 \pm 1.0, 7154.85 \pm 0.17, 7300.80 \pm 0.09, 7563.25 \pm 0.19, 8312.79 \pm 0.12, 8571.53 \pm 0.25, 9050.24 \pm 0.33, 10064.34 \pm 0.31$ and 11235.5 ± 0.5 keV (1990PI05). Angular distributions have been measured for $E_d = 0.32$ to 52 MeV and lead to l_n, J^π and spectroscopic factors: see Table 15.15 in (1981AJ01). Branching ratios and multiplicities are shown in Table 15.15. See also (1985LI1H, 1985ME1E, 1987SI1D, 1988LI1E, 1988VIIA, 1989VI1E; applications).

$$41. \text{}^{14}\text{N}(\text{}^{14}\text{N}, \text{}^{13}\text{N})\text{}^{15}\text{N} \quad Q_m = 0.2799$$

See (1981AJ01) and (1988DA12; theor.).

$$42. \text{}^{15}\text{C}(\beta^-)\text{}^{15}\text{N} \quad Q_m = 9.7717$$

See reaction 1 in ^{15}C and Table 15.14.

$$43. \text{(a) } \text{}^{15}\text{N}(\gamma, n)\text{}^{14}\text{N} \quad Q_m = -10.8333$$

$$\text{(b) } \text{}^{15}\text{N}(\gamma, 2n)\text{}^{13}\text{N} \quad Q_m = -21.3868$$

(c) $^{15}\text{N}(\gamma, \text{p})^{14}\text{C}$	$Q_{\text{m}} = -10.2074$
(d) $^{15}\text{N}(\text{e}, \text{ep}_0)^{14}\text{C}$	$Q_{\text{m}} = -10.2074$
(e) $^{15}\text{N}(\gamma, \text{d})^{13}\text{C}$	$Q_{\text{m}} = -16.1594$
(f) $^{15}\text{N}(\gamma, \text{t})^{12}\text{C}$	$Q_{\text{m}} = -14.8484$
(g) $^{15}\text{N}(\gamma, \pi^-)^{15}\text{O}$	$Q_{\text{m}} = -142.322$

Table 15.15: Radiative widths from $^{15}\text{N}(\gamma, \gamma')$ and $^{15}\text{N}(\text{e}, \text{e}')$ ^a

E_{x} (MeV \pm keV)	J^{π}	Mult.	Γ_{γ_0} (eV)
5.27	$\frac{5}{2}^{+}$	C3	$(4.2 \pm 0.3) \times 10^{-6}$
		M2	$(1.2 \pm 0.7) \times 10^{-4}$
5.30	$\frac{1}{2}^{+}$	C1	2.2 ± 2.3
6.323 ± 1 ^b	$\frac{3}{2}^{-}$	C2	0.050 ± 0.004
		M1	1.9 ± 0.4 ^c
		M1 + E2	3.12 ± 0.18 ^{b, d, e}
7.16	$\frac{5}{2}^{+}$	C3	$(0.86 \pm 0.10) \times 10^{-5}$
7.301 ± 1 ^b	$\frac{3}{2}^{+}$	C1	2.6 ± 1.0
		M2	$(0.3 \pm 0.2) \times 10^{-5}$
		E1 + M2	1.08 ± 0.08 ^b
7.57	$\frac{7}{2}^{+}$	C3	$(1.84 \pm 0.16) \times 10^{-5}$
8.310 ± 4 ^b	$\frac{1}{2}^{+}$	E1	0.3 ± 0.2 ^b
8.575 ± 4 ^b	$\frac{3}{2}^{+}$	E1 + M2	0.3 ± 0.3 ^b
9.048 ± 1 ^b	$\frac{1}{2}^{+}$	E1	1.2 ± 0.2 ^b
9.150 ± 1 ^b	$\frac{3}{2}^{-}$	C2	0.095 ± 0.005 ^f
		M1	0.2 ± 0.8
		M1 + E2	0.47 ± 0.12 ^{b, g}
9.760 ± 1 ^b	$\frac{5}{2}^{-}$	C2	0.20 ± 0.05
		E2	0.21 ± 0.07 ^b
9.924 ± 1 ^b	$\frac{3}{2}^{-}$	M1	1.6 ± 0.2 ^b
10.064 ± 1 ^b	$\frac{3}{2}^{+}$	E1	6.3 ± 0.4 ^b
10.8	$\frac{3}{2}^{+}$	M2	$(1.8 \pm 0.8) \times 10^{-2}$
11.88	$\frac{3}{2}^{-}$	C2	0.44 ± 0.10
		M1	4.4 ± 3.8
12.5	$\frac{5}{2}^{+}$	M2	$(5.2 \pm 2.0) \times 10^{-2}$

Table 15.15: Radiative widths from $^{15}\text{N}(\gamma, \gamma')$ and $^{15}\text{N}(e, e')$ ^a (continued)

E_x (MeV \pm keV)	J^π	Mult.	Γ_{γ_0} (eV)
(13.98)			
14.7	$\frac{5}{2}^-$	C2	1.8 ± 0.2
20.10			
23.25			

^a For references and $B(\lambda)\uparrow$ see Table 15.17 in (1981AJ01). See also Tables 15.5 and 15.6 here. Form factors have also been measured to $^{15}\text{N}^*(9.23, 11.29$ [both $\frac{1}{2}^-$], 10.45 [$\frac{5}{2}^-$], 12.1 [u], 12.9 [u]) (1987DE1Q) [unpublished Ph.D. thesis].

^b (1981MO09): (γ, γ) .

^c See note added in proof in (1975MO28).

^d $\delta(\text{E2/M1}) = 0.137 \pm 0.005$. See, however, Table 15.5.

^e Using $\delta(\text{E2/M1}) = 0.132 \pm 0.004$ [see Table 15.5] $\Gamma_{\gamma_0} = 3.07 \pm 0.18$ eV (M1) and $(5.34 \pm 0.44) \times 10^{-2}$ eV (E2) (D.J. Millener, private communication.)

^f $\delta(\text{E2/M1}) > 0.3$.

^g Mixing ratio is very small [see Table 15.5] and the transition is almost purely M1 (D.J. Millener, private communication).

The total photoneutron cross section from threshold to 38 MeV shows a very broad GDR which extends from ≈ 16 to 30 MeV. Maxima are observed at $E_\gamma \approx 23.5$ and 25.5 MeV ($\sigma \approx 11$ mb): see (1988DI02) [based on (1982JU03; monoenergetic photons)]. However, (1989BA25) report a sharper single peak at $E_\gamma = 25.5$ MeV in the (γ, Sn) reaction with a cross section of ≈ 16 mb. See (1989BA25) for a discussion of the $T_<$ and $T_>$ components of the GDR.

The (γ, n_0) cross section for $E_x = 13$ to 24 MeV shows a broad structure centered at $E_x \approx 14.5$ MeV and a resonance at $E_x = 17.3 \pm 0.1$ MeV. A large fraction of the photoabsorption strength leading to $^{14}\text{N}_{\text{g.s.}}$ is due to the formation of $\frac{3}{2}^+$, $T = \frac{1}{2}$ states in ^{15}N which decay by d-wave emission. The absorption is essentially pure E1 (1983WA03). See also (1981AJ01).

The $(\gamma, 2n)$ reaction has been studied from threshold to 38 MeV [see (1988DI02)] and more recently from 20 to 28 MeV by (1988MC01). The cross section remains near zero for 3 MeV above threshold and then rises sharply at about 24 MeV to a maximum value of 1.7 mb (1988MC01) [see, however, (1982JU03)]. For discussions of the results in terms of the density of $T_>$ states in ^{14}N , see (1988MC01, 1989BA25 (p.511)).

For discussions of the (γ, p) reaction see (1981AJ01) and (1989BA25). The latter show a curve for the total absorption cross section [$(\gamma, n) + (\gamma, p)$] from 10 and 27 MeV dominated by a peak (see above) at $E_\gamma \approx 25$ MeV with $\sigma \approx 23$ mb.

A study at $E_e = 18.8, 20.8, 25.7$ and 29.7 MeV (reaction (d)) shows a ‘‘pigmy’’ resonance at $E_x = 14.8$ MeV, a shoulder at 15.6 MeV, a peak at 16.7 MeV [probably $\frac{1}{2}^+$ but $\frac{3}{2}^+$ is not ruled out], and the giant dipole resonance, which exhibits a great deal of structure, centered at 22 MeV. The data on the pigmy resonance are consistent with an admixture of $\approx 1\%$ $\frac{3}{2}^-$ (E2) or $\frac{1}{2}^-$ (M1)

to a predominantly $\frac{1}{2}^+$ (E1) state. The experiment shows that for $14 < E_x < 28$ MeV the reaction goes predominantly via $\frac{1}{2}^+$ or $\frac{3}{2}^+$ (E1) states in ^{15}N ; the $T = \frac{3}{2}$ strength is concentrated above 18 MeV: see (1981AJ01).

The cross section for d_0 [reaction (e)] is reported at 90° for $E_\gamma \approx 20.5$ to 28.5 MeV: a resonance is observed at $E_x \approx 21.9$ MeV. The (γ, t_0) cross section (reaction (f)) at 90° decreases from a value of $30 \mu\text{b/sr}$ at 20 MeV to $5 \mu\text{b/sr}$ at 22 MeV and remains flat out to 25 MeV. Comparison of this cross section, and those of the other photonuclear reactions, suggest an isospin splitting of ≈ 6 MeV with the $T = \frac{1}{2}$ strength concentrated between 16 and 21 MeV and the $T = \frac{3}{2}$ strength between 21 and 28 MeV. $^{15}\text{N}^*(21.9)$ is not observed. See (1981AJ01) for references. For reaction (g) [to $^{15}\text{O}_{\text{g.s.}}$] see (1988LI23, 1989KO28). See also (1985GO1A, 1987KIIC; theor.).

44. $^{15}\text{N}(\gamma, \gamma)^{15}\text{N}$

See Table 15.15 and (1981AJ01). See also (1987MO03).

45. $^{15}\text{N}(e, e)^{15}\text{N}$

The charge r.m.s. radius of ^{15}N is 2.612 ± 0.009 fm. The C0 elastic scattering form factor of ^{15}N has been measured over $q = 0.4 - 3.2 \text{ fm}^{-1}$ (1988DE09). Inelastic groups are displayed in Table 15.15.

The giant resonance is split into two main peaks at $E_x = 22$ and 25.5 MeV with some structure around 20 MeV. $\Gamma_{\gamma_0}(\text{C1}) = (1.1 \pm 0.3) \times 10^3$ eV (14 - 18.5 MeV), $\Gamma_{\gamma_0}(\text{C2}) = 12.5 \pm 2.0$ eV assuming the states responsible are $\frac{3}{2}^+$ and $\frac{3}{2}^-$, respectively. For $E_x = 18.5$ to 30 MeV, $\Gamma_{\gamma_0}(\text{C1}) = (1.96 \pm 0.04) \times 10^4$ eV while $\Gamma_{\gamma_0} < 0.1$ eV for any C2 strength. See (1981AJ01, 1986AJ01) for references. See also (1988MUZV; (e, e' γ); prelim.), (1986PA1C, 1987DE43, 1988LI23, 1988PA1S, 1989KO28, 1990PA1H) and (1986JE1B, 1987DO12, 1988FU04, 1988GOZM, 1988SH07, 1989FU05, 1989WO1E, 1990BL09, 1990FU04; theor.).

46. $^{15}\text{N}(\pi^\pm, \pi^\pm)^{15}\text{N}$

At $E_{\pi^\pm} = 164$ MeV angular distributions have been studied to states at $E_x = 10.68 \pm 0.03$, 12.52 ± 0.02 , 14.04 ± 0.03 and 17.19 ± 0.03 MeV: $J^\pi = \frac{9}{2}^+, \frac{9}{2}^+, (\frac{9}{2}^+, \frac{7}{2}^+)$ and $(\frac{9}{2}^+, \frac{7}{2}^+)$, respectively, as well as to the $^{15}\text{N}_{\text{g.s.}}$. Additional π^+ cross sections were measured at 120 and 260 MeV: peaks were observed at $E_x = 20.11 \pm 0.06$ and 23.19 ± 0.06 MeV [both are probably $T = \frac{3}{2}$ states]. $^{15}\text{N}^*(5.27, 6.32, 7.57)$ were also populated (1985SE06). At $E_{\pi^+} = 164$ MeV, elastic scattering has been studied from $^{15}\text{N}^{\vec{N}}$: A_y for $\theta = 60^\circ - 100^\circ$ is consistent with zero (1989TA21, 1990TA1L). See also (1990TAZS) and (1988GOZM; theor.).

47. $^{15}\text{N}(n, n)^{15}\text{N}$

See ^{16}N in (1986AJ04). See also (1989FU1J).

48. $^{15}\text{N}(p, p)^{15}\text{N}$

Angular distributions of elastically scattered protons have been measured at E_p to 44.2 MeV: see (1981AJ01, 1986AJ01). For measurements of $K_y^{y'}$ at $E_p = 65$ MeV see (1990NA15). See also ^{16}O in (1986AJ04) and (1990DU01; theor.).

49. (a) $^{15}\text{N}(d, d)^{15}\text{N}$

(b) $^{15}\text{N}(^3\text{He}, ^3\text{He})^{15}\text{N}$

Angular distributions of elastically scattered deuterons have been measured at $E_d = 5 - 6$ MeV. Elastic and inelastic ^3He distributions have been studied for $E(^3\text{He}) = 11$ to 39.8 MeV: see (1976AJ04). Elastic distributions and A_y have also been measured at $E(^3\text{He}) = 33$ MeV (1986DR03).

50. $^{15}\text{N}(\alpha, \alpha)^{15}\text{N}$

At $E_\alpha = 40.5$ MeV, a number of particle groups have been observed and angular distributions have been measured: see Table 15.17 of (1976AJ04). At $E_\alpha = 48.7$ and 54.1 MeV elastic angular distributions have been reported by (1987AB03). See also (1981AJ01) for additional information and (1985SH1D; theor.).

51. $^{15}\text{N}(^7\text{Li}, ^7\text{Li})^{15}\text{N}$

The elastic scattering has been studied at $E(^7\text{Li}) = 28.8$ MeV: see (1986AJ01).

52. (a) $^{15}\text{N}(^{12}\text{C}, ^{12}\text{C})^{15}\text{N}$

(b) $^{15}\text{N}(^{13}\text{C}, ^{13}\text{C})^{15}\text{N}$

(c) $^{15}\text{N}(^{14}\text{C}, ^{14}\text{C})^{15}\text{N}$

Angular distributions of elastic scattering have been measured at $E(^{15}\text{N}) = 31.5$ to 47 MeV [reaction (a)] and $E(^{13}\text{C}) = 105$ MeV [reaction (b)]: see (1981AJ01, 1986AJ01) [also for yield measurements]. See also (1986HA1F, 1989BEZC) and (1985HU04, 1986BA69, 1986HA13; theor.).

53. $^{15}\text{N}(^{16}\text{O}, ^{16}\text{O})^{15}\text{N}$

Elastic angular distributions have been measured at $E(^{16}\text{O}) = 35.1$ and 42.6 MeV: see (1986AJ01). For fusion cross sections and yields see (1976AJ04, 1986AJ01) and (1985NO1C, 1986HA1F). See also (1985HU04; theor.).

54. (a) $^{15}\text{N}(^{27}\text{Al}, ^{27}\text{Al})^{15}\text{N}$
 (b) $^{15}\text{N}(^{28}\text{Si}, ^{28}\text{Si})^{15}\text{N}$

Elastic distributions have been measured in the range $E(^{15}\text{N}) = 32.8$ to 69.8 MeV [reaction (a)] and at 44 MeV [reaction (b)]: see (1981AJ01, 1986AJ01). See also (1988SN1A).

55. $^{15}\text{O}(\beta^+)^{15}\text{N}$ $Q_m = 2.7539$

See ^{15}O .

56. (a) $^{16}\text{O}(\gamma, p)^{15}\text{N}$ $Q_m = -12.1276$
 (b) $^{16}\text{O}(e, ep)^{15}\text{N}$ $Q_m = -12.1276$

Over the giant resonance region in ^{16}O , the decay takes place to the odd-parity states $^{15}\text{N}^*(0, 6.32)$ and less strongly to the even-parity states $^{15}\text{N}^*(5.27, 5.30, 8.31, 9.05)$ and to $^{15}\text{N}^*(9.22)$: see (1970AJ04, 1976AJ04). At $E_e = 500$ MeV most of the $1p$ hole strength is concentrated in the groups to $^{15}\text{N}^*(0, 6.32)$. The $1s$ state shows up as a very wide asymmetric structure centered at $E_x \approx 41$ MeV: see (1981AJ01). See also (1990LE1P). In the range $E_\gamma = 101.5$ to 382 MeV differential cross sections are reported for the p_0 , (p_{1+2}) and p_3 groups at $\theta = 45^\circ, 90^\circ$ and 135° (1985LE07). Differential cross sections have also been measured at $E_\gamma = 196, 257, 312, 316$ MeV for the p_0 and $p_{1\rightarrow 3}$ groups [the latter not at 316 MeV] (1988AD07, 1985TU02). $^{15}\text{N}^*(0, 6.3, 10.8)$ have been populated at $E_e = 500$ MeV (1982BE02). See also (1986AJ01), (1987VOZR, 1988LEZW, 1989LEZY; prelim.), (1987MA1K) and (1985CA17, 1985GO1B, 1986CH05, 1986LU1A, 1986PO14, 1987RY03, 1988CA10, 1988DUZX, 1988HO10, 1988YA08, 1988LO07, 1988MC03, 1988RY03, 1989RY01, 1990BRZY, 1990FL1A, 1990OWZZ; theor.).

$$57. {}^{16}\text{O}(\mu^-, \nu\text{n})^{15}\text{N} \quad Q_{\text{m}} = 92.7492$$

Gamma rays from the decay of one of the states at 5.3 MeV and from ${}^{15}\text{N}^*(6.3)$ are reported by (1983VA1E).

$$58. {}^{16}\text{O}(\text{n}, \text{d})^{15}\text{N} \quad Q_{\text{m}} = -9.9030$$

Angular distributions of the d_0 group have been reported at $E_{\text{n}} = 14$ and 14.4 MeV: see (1976AJ04). See also (1988YOZX; $E_{\text{n}} = 60$ MeV; prelim.) and (1990MC04; applied).

$$59. \text{(a) } {}^{16}\text{O}(\pi^\pm, \pi^\pm\text{p})^{15}\text{N} \quad Q_{\text{m}} = -12.1276$$

$$\text{(b) } {}^{16}\text{O}(\text{p}, 2\text{p})^{15}\text{N} \quad Q_{\text{m}} = -12.1276$$

At $E_{\pi^\pm} = 240$ MeV, the spectra are dominated by ${}^{15}\text{N}^*(0, \approx 6.5)$. The π^+/π^- ratio has been measured for the ground-state transitions (1984KY01). At $E_{\pi^+} = 2.0$ GeV/ c differential cross sections have been determined for the transition to ${}^{15}\text{N}^*(6.3)$ (1983KI01).

At $E_{\text{p}} = 505$ MeV the summed proton spectrum shows two peaks corresponding to the knockout of $\text{p}_{1/2}$ and $\text{p}_{3/2}$ protons with binding energies of 12.12 and 18.44 MeV [${}^{15}\text{N}^*(0, 6.32)$]. Differential cross sections and (p, 2p)/(p, pn) ratios are also reported (1986MC10). For work at 1 GeV involving the knockout of $\text{s}_{1/2}$ protons see (1985BE30). For γ -ray production (${}^{15}\text{N}^*(5.27)$) at $E_{\text{p}} = 30 - 40$ MeV see (1988LE08). See also (1985KI1A, 1986CHI1J, 1987VD1A), (1987LA11, 1988LE08; astrophys.) and (1986BO1A; theor.). For earlier work see (1976AJ04).

$$60. {}^{16}\text{O}(\text{d}, {}^3\text{He})^{15}\text{N} \quad Q_{\text{m}} = -6.6340$$

Angular distributions of ${}^3\text{He}$ groups have been measured for $E_{\text{d}} = 20$ to 82 MeV: see (1976AJ04, 1981AJ01). The spectra are dominated by the transitions to ${}^{15}\text{N}^*(0, 6.32)$. A ZRDWBA analysis leads to $C^2S = 2.25$ and 3.25 for these two states [and to 2.37 and 3.31 for the analog states in ${}^{15}\text{O}$ studied with the (d, t) reaction]. $J^\pi = \frac{3}{2}^-$ for both ${}^{15}\text{N}^*(9.93, 10.70)$: see (1981AJ01). See also (1987MO03) for a re-analysis of C^2S .

$$61. {}^{16}\text{O}(\alpha, \text{ap})^{15}\text{N} \quad Q_{\text{m}} = -12.1276$$

At $E_{\alpha} = 139.2$ MeV the absolute spectroscopic factors $S = 5.4$ and 6.9 for ${}^{15}\text{N}^*(0, 6.32)$ (1987SA01).

62. (a) $^{16}\text{O}(^6\text{Li}, ^7\text{Be})^{15}\text{N}$ $Q_m = -6.522$
 (b) $^{16}\text{O}(^7\text{Li}, ^8\text{Be})^{15}\text{N}$ $Q_m = 5.1268$
 (c) $^{16}\text{O}(^{16}\text{O}, ^{17}\text{F})^{15}\text{N}$ $Q_m = -11.5274$

For reaction (a) see (1986GL1E). For reaction (b) see (1988MA07) and for (c) see (1988AU03). For other heavy-ion reactions see (1981AJ01, 1986AJ01).

63. $^{17}\text{O}(\text{p}, ^3\text{He})^{15}\text{N}$ $Q_m = -8.5529$

At $E_p = 39.8$ MeV angular distributions of the groups to $^{15}\text{N}^*(0, 6.32)$ have been compared with those to the analog states in ^{15}O reached in the (p, t) reaction: see (1976AJ04).

64. $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$ $Q_m = 3.9804$

Angular distributions of α_0 have been measured for $E_p = 0.125$ to 42.2 MeV: see (1976AJ04, 1981AJ01) and ^{19}F in (1987AJ02). At $E_p = 40.9$ MeV angular distributions have also been studied to $^{15}\text{N}^*(0, 5.27 + 5.30, 6.32, 7.57, 9.15[\text{u}], 9.83[\text{u}], 10.7[\text{u}], 11.24[\text{u}], 11.44, 12.52[\text{u}])$ (1987CA15; see for C^2S).

For $^{15}\text{N}^*(5.27)$, $\tau_m = 2.49 \pm 0.24$ ps, $|g| = 0.94 \pm 0.07$ (1983BI10). See also (1986CO1F), (1988FI1C, 1989NW1A, 1990MI15; applied) and (1986BA89, 1988CA26; astrophys.).

65. $^{19}\text{F}(\text{d}, ^6\text{Li})^{15}\text{N}$ $Q_m = -2.5388$

Angular distributions involving $^{15}\text{N}^*(0, 5.3, 6.3)$ have been measured in the range $E_d = 9.0$ to 28 MeV [see (1976AJ04, 1981AJ01)].

66. $^{19}\text{F}(^3\text{He}, ^7\text{Be})^{15}\text{N}$ $Q_m = -2.4265$

See (1976AJ04) and (1986HA1E; theor.).

¹⁵O
(Figs. 3 and 4)

GENERAL (See also (1986AJ01)).

Nuclear models: (1987ST05)

Special states: (1985SH24, 1986LI1B, 1987ST05, 1988AN1E, 1989WU1C)

Electromagnetic transitions: (1984VA06, 1987HO1L, 1987ST05)

Astrophysical questions: (1985TA1A, 1987RA1D, 1989JI1A, 1989ST14, 1990RA1O)

Complex reactions involving ¹⁵O: (1985FI08, 1985HU1C, 1985PO11, 1985SI19, 1986GR1A, 1986ME06, 1986PO06, 1986TO10, 1986UT01, 1987BA38, 1987BE1I, 1987BU07, 1987NA01, 1987RI03, 1987ST01, 1988BE02, 1988BE56, 1988MI28, 1989BA92, 1989CA25, 1989DR03, 1989KI13, 1989PO07, 1989SA10, 1989TA1O, 1989YO02)

Applied work: (1985BO1P, 1985HA40, 1987HI1B, 1988HI1F, 1988VO1D, 1989AR1J, 1989WO1B)

Pion and other mesons capture and reactions (See also reactions 8, 16, 17 and 21.): (1986LI1B, 1987LE1B, 1988CH49, 1988MI1K, 1989LE1L, 1990OD1A)

Hypernuclei: (1984ZH1B, 1986DA1G, 1989BA93, 1989KO1H, 1989TA17)

Other topics: (1985AN28, 1985SH24, 1986AN07, 1986WI03, 1987CH02, 1989WU1C, 1990MU1O)

Ground-state properties of ¹⁵O: (1985AN28, 1985AR11, 1986MC13, 1986WI03, 1986WUZX, 1987FU06, 1987SA15, 1988AR1B, 1988CH1T, 1988FU04, 1988NI05, 1988SH07, 1988VA03, 1988WA08, 1989CH24, 1989FU05, 1989NE02, 1989SA10)

$\mu = 0.7189 (8) \text{ nm}$ (1978LEZA). See also (1989RA17).

¹⁵O*(5.24): $\mu = +(0.65 \pm 0.07) \text{ nm}$: see (1989RA17).

1. ¹⁵O(β^+)¹⁵N $Q_m = 2.7539$

The half-life of ¹⁵O is $122.24 \pm 0.16 \text{ s}$: see (1981AJ01); $\log f_0 t = 3.637$. The K/β^+ ratio is $(10.7 \pm 0.6) \times 10^{-4}$: see (1976AJ04). The β -anisotropy has been measured by (1988SE11, 1989SE07). See also (1986AJ01), (1990ST08), (1985BA1N, 1985BA1M, 1986GR04, 1987BA89, 1987FR1C, 1987RI1E, 1987WE1C, 1988BA86, 1988BA1Y, 1989BA2P, 1989DA1H; astrophysics) and (1988TA09, 1988TO1C, 1989WO1E; theor.).

2. (a) ¹¹B(⁷Li, 3n)¹⁵O $Q_m = -3.494$
 (b) ¹¹B(⁹Be, t2n)¹⁵O $Q_m = -13.932$




Fig. 3: Energy levels of ^{15}O . For notation see Fig. 2.

Table 15.16: Energy levels of ^{15}O ^a

E_x in ^{15}O (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^-; \frac{1}{2}$	$\tau_{1/2} = 122.24 \pm 0.16$ s	β^+	1, 3, 4, 5, 6, 7, 8, 9, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 27, 28
5.183 ± 1	$\frac{1}{2}^+$	$\tau_m = 8.2 \pm 1.0$ fs	γ	5, 7, 9, 14, 15, 19, 20, 23, 24, 25
5.2409 ± 0.3	$\frac{5}{2}^+$	3.25 ± 0.30 ps	γ	4, 5, 6, 7, 9, 14, 15, 18, 19, 20, 23, 24, 25, 27
6.1763 ± 1.7	$\frac{3}{2}^-$	$g = +0.248 \pm 0.026$ < 2.5 fs	γ	5, 7, 9, 14, 15, 18, 19, 20, 21, 22, 23, 24, 25, 27
6.7931 ± 1.7	$\frac{3}{2}^+$	< 28 fs	γ	5, 7, 9, 14, 15, 19, 25
6.8594 ± 0.9	$\frac{5}{2}^+$	16.0 ± 2.5 fs	γ	4, 5, 7, 9, 14, 15, 19, 20, 25, 27
7.2759 ± 0.6	$\frac{7}{2}^+$	0.70 ± 0.15 ps	γ	4, 5, 6, 7, 8, 14, 15, 18, 19, 23, 25, 27
7.5565 ± 0.4	$\frac{1}{2}^+$	$\Gamma = 0.99 \pm 0.10$ keV	γ, p	7, 9, 14, 15, 18, 19, 23, 25
8.2840 ± 0.5	$\frac{3}{2}^+$	3.6 ± 0.7	γ, p	5, 7, 9, 14, 15, 25
8.743 ± 6	$\frac{1}{2}^+$	32	γ, p	7, 9, 25
8.922 ± 2	$\frac{5}{2}^+$	3.3 ± 0.3	γ, p	4, 5, 7, 9, 23, 25
8.922 ± 2	$\frac{1}{2}^+$	7.5	γ, p	4, 7, 9, 23, 25
8.9821 ± 1.7	$(\frac{1}{2})^-$	3.9 ± 0.4	γ, p	5, 7, 9, 25
9.484 ± 8	$(\frac{3}{2})^+$	≈ 200	γ, p	9, 25
9.488 ± 3	$\frac{5}{2}^-$	10.1 ± 0.5	γ, p	5, 7, 9, 25
9.609 ± 2	$\frac{3}{2}^-$	8.8 ± 0.5	γ, p	4, 5, 7, 9, 25
9.662 ± 3	$(\frac{7}{2}, \frac{9}{2})^-$	2 ± 1	p	4, 5, 7, 11, 25
10.29^b	$(\frac{5}{2})^-$	3 ± 1	p	5, 7, 11, 25
10.30^b	$\frac{5}{2}^+$	11 ± 2	p	5, 7, 11, 25

Table 15.16: Energy levels of ^{15}O ^a (continued)

E_x in ^{15}O (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
10.461 \pm 5	$(\frac{9}{2}^+)$	< 2	γ, p	4, 5, 6, 7, 9, 25
10.48	$(\frac{3}{2}^-)$	25 ± 5	γ, p	4, 7, 9, 11, 24
(10.506)	$(\frac{3}{2}^+)$	140 ± 40	γ, p	9, 11
10.917 \pm 12	$\frac{7}{2}^+$	90	p	11, 25
10.938 \pm 3	$\frac{1}{2}^+$	99 ± 5	γ, p	9, 11, 25
11.025 \pm 3	$\frac{1}{2}^-$	25 ± 2	γ, p	9, 11, 25
11.151 \pm 7		< 10	p	5, 11, 25
11.218 \pm 3	$\frac{3}{2}^+$	40 ± 4	γ, p	9, 11, 25
11.565 \pm 15		< 10	p	5, 11, 25
11.569 \pm 15	$\frac{5}{2}^-$	20 ± 15	γ, p	5, 9, 11
11.616 \pm 15	$(\frac{3}{2}, \frac{1}{2})^-$	80 ± 50	γ, p	9, 11
11.719 \pm 8		< 10	p	4, 5, 11, 25
11.748 \pm 3	$\frac{5}{2}^+$	99 ± 5	γ, p	9, 11
11.846 \pm 3	$\frac{5}{2}^-$	65 ± 3	γ, p	9, 11
11.980 \pm 10	$\frac{5}{2}^-$	20 ± 5	p	5, 11, 25
12.129 \pm 15	$\frac{5}{2}^+$	200 ± 50	p	11
12.222 \pm 20		100 ± 50	p	11
12.255 \pm 13	$\frac{5}{2}^+; \frac{3}{2}$	135 ± 15	p	27
12.295 \pm 10				5
12.471 \pm 3	$\frac{5}{2}^-, (\frac{3}{2}^-)$	77 ± 4	p	11
12.60 \pm 10				5
12.80		≈ 250	γ, p	9
12.835 \pm 3		16 ± 1	p	4, 5, 6, 11
13.008 \pm 3		215 ± 3	p	11
13.025 \pm 3		40 ± 30	p, (^3He)	3, 11
13.45	$(\frac{1}{2}, \frac{3}{2})^+$	≈ 1000	$\gamma, \text{p}, (\alpha)$	9, 11, 13
(13.49)	$(\frac{3}{2}^+)$		(p)	11
13.60	$\frac{5}{2}^+$		p, α	13
13.70	$\frac{3}{2}^-$		p	11
13.79	$\frac{3}{2}^-$		n, p, $^3\text{He}, \alpha$	3, 11, 13

Table 15.16: Energy levels of ^{15}O ^a (continued)

E_x in ^{15}O (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
13.87		≈ 150	γ, p	9
14.03 \pm 40	$(\frac{1}{2}^-, \frac{3}{2}^-)$	160 \pm 20	n, p, ^3He	3
14.17	$\frac{5}{2}^-$		p, α	13
14.27 \pm 10	$\frac{1}{2}^+$	340 \pm 30	n, p, $^3\text{He}, \alpha$	3, 4, 5, 10, 11, 12, 13
14.34	$\frac{5}{2}^+$	(240)	p, (^3He), α	3, 13
14.465 \pm 10	$\frac{3}{2}^+, \frac{5}{2}^+$	100 \pm 10	n, p, $^3\text{He}, \alpha$	3, 10, 11, 13
14.70 \pm 40		170 \pm 35	n, p, ^3He	3, 10
14.95 \pm 40		400 \pm 25	n, p, $^3\text{He}, \alpha$	3, 10, 11, 12, 13
15.05 \pm 10	$((\frac{13}{2}^+))$			4, 5, 6
15.1	$(\frac{1}{2}, \frac{3}{2})^+$	≈ 1000	γ, p	9
15.45 \pm 30		70 \pm 20	p, $^3\text{He}, \alpha$	3
15.54 \pm 10			(p, $^3\text{He}, \alpha$)	3, 5
15.60 \pm 10			(p, $^3\text{He}, \alpha$)	3, 5
15.65 \pm 10				4, 5
15.80 \pm 10			n, ^3He	3, 5
15.90 \pm 15	$\frac{1}{2}^-, \frac{3}{2}^-$	350	$^3\text{He}, \alpha$	3
16.05 \pm 20		≈ 185	n, p, $^3\text{He}, \alpha$	3, 10, 11, 13
16.10 \pm 20			(n), $^3\text{He}, \alpha$	3
16.21 \pm 20		≈ 140	(n), p, $^3\text{He}, \alpha$	3, 11, 12, 13
16.43 \pm 75	$\frac{1}{2}^+$	560 \pm 100	n, $^3\text{He}, \alpha$	3, 10, 12
16.75 \pm 50			n, ^3He	3, 25
17.05 \pm 60	$(\frac{1}{2}, \frac{3}{2})^+; \frac{1}{2}$	700 \pm 70	$\gamma, \text{p}, ^3\text{He}$	3, 9, 11, 13
17.46 \pm 20				5
17.51 \pm 20	$\frac{1}{2}^-, \frac{3}{2}^-$	640 \pm 120	$\gamma, \text{n}, ^3\text{He}, \alpha$	3, 5
17.99 \pm 50	$\frac{1}{2}^-, \frac{3}{2}^-$	200	^3He	3
18.23 \pm 50			n, p, ^3He	3
18.67 \pm 60	$(\frac{1}{2}, \frac{3}{2})^+; \frac{1}{2}$	520 \pm 110	$\gamma, ^3\text{He}$	3, 9
19.03 \pm 50		1120 \pm 300	$\gamma, \text{n}, ^3\text{He}$	3, 23
19.57 \pm 80	$(\frac{1}{2}, \frac{3}{2})^+; \frac{1}{2}$	780 \pm 270	$\gamma, ^3\text{He}$	3

Table 15.16: Energy levels of ^{15}O ^a (continued)

E_x in ^{15}O (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
19.91 \pm 50			n, ^3He	3
20.42 \pm 70	$(\frac{3}{2}, \frac{1}{2})^+; \frac{1}{2}$	970 \pm 240	γ , p, ^3He	3, 9
21.56 \pm 70	$(\frac{3}{2}, \frac{1}{2})^+; \frac{1}{2}$	730 \pm 120	γ , p, ^3He	3, 9, 23
23.8 \pm 0.1		\lesssim 500	γ , ^3He	3
(26.0)	$(\frac{13}{2}^-)$	\approx 600	^3He	3
(28.0)	$(\frac{9}{2}^-, \frac{11}{2}^-)$	\approx 2500	^3He	3
(29.0)		\approx 2500	^3He	3

^aSee also Table 15.17.

^b It is possible that these two are in fact a single state: see (1976AJ04).

For reaction (a) see (1986BE54); for reaction (b) see (1986BE35). For other heavy-ion reactions see (1986AJ01).

3. (a) $^{12}\text{C}(^3\text{He}, \gamma)^{15}\text{O}$	$Q_m = 12.0759$	
(b) $^{12}\text{C}(^3\text{He}, \text{n})^{14}\text{O}$	$Q_m = -1.1466$	$E_b = 12.0759$
(c) $^{12}\text{C}(^3\text{He}, \text{p})^{14}\text{N}$	$Q_m = 4.7789$	
(d) $^{12}\text{C}(^3\text{He}, \text{d})^{13}\text{N}$	$Q_m = -3.5500$	
(e) $^{12}\text{C}(^3\text{He}, \text{t})^{12}\text{N}$	$Q_m = -17.357$	
(f) $^{12}\text{C}(^3\text{He}, ^3\text{He})^{12}\text{C}$		
(g) $^{12}\text{C}(^3\text{He}, \alpha)^{11}\text{C}$	$Q_m = 1.8560$	

Excitation functions and polarization measurements for these reactions have been measured over a wide range of energies: see Tables 15.20 in (1970AJ04, 1976AJ04, 1981AJ01) and the text below. Observed resonances are displayed in Table 15.18 here.

The 90° yield and the angular distributions of γ_0 , measured from $E(^3\text{He}) = 5.24$ to 13.95 MeV show five resonances attributed to E1 transitions from $J^\pi = \frac{1}{2}^+$ or $\frac{3}{2}^+$, $T = \frac{1}{2}$ states in the GDR characterized by a considerable 3p4h admixture (1978DE33 [also for ω_γ], 1984DE09). Yields of γ_{1+2} at 90° have also been reported at $E(^3\text{He}) = 5.3$ to 16.7 MeV: the cross section is some eight times greater than that for $(^3\text{He}, \gamma_0)$ and is similar to that for the $^{14}\text{N}(\text{p}, \gamma_0)^{15}\text{O}$ reaction over the same excitation range. Three resonances are reported [see Table 15.18 (1989KI09)]: it is suggested that they are due to cluster states with a large 3p4h component. See also (1988BLZY; prelim.; $E(^3\text{He}) = 12.0$ to 24.6 MeV; γ to many states of ^{15}O).

Table 15.17: Radiative decays in ^{15}O ^a

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	δ^b
5.24	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	100	$+0.10 \pm 0.04$ (E3/M2)
6.18 ^c	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100	-0.125 ± 0.007 (E2/M1) ^k
6.79 ^d	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	100	-0.02 ± 0.02 (M2/E1)
6.86 ^e	$\frac{5}{2}^+$	5.24	$\frac{5}{2}^+$	100	$+0.04 \pm 0.03$ (E2/M1)
7.28 ^f	$\frac{7}{2}^+$	0	$\frac{1}{2}^-$	3.8 ± 1.2	
		5.24	$\frac{5}{2}^+$	96.2 ± 1.2	
7.56 ^g	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	3.5 ± 0.5	
		5.18	$\frac{1}{2}^+$	15.8 ± 0.6	
		6.18	$\frac{3}{2}^-$	57.5 ± 0.4	
		6.79	$\frac{3}{2}^+$	23.2 ± 0.6	
		6.86	$\frac{5}{2}^+$	1	Γ_γ (eV)
8.28 ^h	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	53.2 ± 0.25 ^m	0.24
		5.18	$\frac{1}{2}^+$	1.2 ± 0.1	0.006
		5.24	$\frac{5}{2}^+$	42.2 ± 0.5 ^m	0.20
		6.18	$\frac{3}{2}^-$	2.2 ± 0.6 ^m	0.01
		6.86	$\frac{5}{2}^+$	1.2 ± 0.3 ^m	0.006
8.74 ^h	$\frac{1}{2}^+$	5.18	$\frac{1}{2}^+$	64 ± 3	0.18
		6.18	$\frac{3}{2}^-$	36 ± 3	0.10
8.922 ⁱ	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	9 ± 4	
		5.18	$\frac{1}{2}^+$	39 ± 3	
		6.18	$\frac{3}{2}^-$	24 ± 3	
		6.86	$\frac{5}{2}^+$	28 ± 3	
8.922 ⁱ	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	50 ± 25	
		5.18	$\frac{1}{2}^+$	20 ± 10	
		6.18	$\frac{3}{2}^-$	20 ± 10	
		6.86	$\frac{5}{2}^+$	(10 ± 10)	
8.982 ^j	$(\frac{3}{2})^-$	0	$\frac{1}{2}^-$	94 ± 1	
		5.18	$\frac{1}{2}^+$	6 ± 1	
9.48 ^h	$(\frac{3}{2})^+$	0	$\frac{1}{2}^-$	100	9.1 ± 2.0 ⁿ
9.49	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	86	2.1

Table 15.17: Radiative decays in ^{15}O ^a (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	δ^b
9.61	$\frac{3}{2}^-$	5.24	$\frac{5}{2}^+$	6.5	0.15
		6.18	$\frac{3}{2}^-$	0.7	0.22
		6.86	$\frac{5}{2}^+$	3.4	0.08
		7.28	$\frac{7}{2}^+$	5.1	0.11
		0	$\frac{1}{2}^-$	79	4.0
		5.24	$\frac{5}{2}^+$	19	1.0
10.46	$(\frac{9}{2}^+)$	6.18	$\frac{3}{2}^-$	2	0.1
		5.24	$\frac{5}{2}^+$	62 ± 6	18 ± 6^n
		6.86	$\frac{5}{2}^+$	< 4	< 1.5
10.48	$(\frac{3}{2})^-$	7.28	$\frac{7}{2}^+$	38 ± 6	11 ± 4^n
		0	$\frac{1}{2}^-$	60 ± 8	0.21 ± 0.07^n
		5.24	$\frac{5}{2}^+$	40 ± 6	0.14 ± 0.01^n
10.94	$\frac{1}{2}^+$	6.18	$\frac{3}{2}^-$	< 4	< 0.02
		9.79	$\frac{3}{2}^+$	< 4	< 0.02
		0	$\frac{1}{2}^-$	44 ± 8	14 ± 4
		5.18	$\frac{1}{2}^+$	34 ± 3	11 ± 2
		6.18	$\frac{3}{2}^-$	22 ± 8	7 ± 2
		6.79	$\frac{3}{2}^+$	< 8	< 3
11.03 ^a	$\frac{1}{2}^-$	0	$\frac{1}{2}^-$	100	1.4 ± 0.4
11.22	$\frac{3}{2}^+$	0	$\frac{1}{2}^-$	74 ± 5	5.5 ± 0.5
		5.18	$\frac{1}{2}^+$	14 ± 5	1.0 ± 0.2
		5.24	$\frac{5}{2}^+$	12 ± 5	0.9 ± 0.2
		6.79	$\frac{3}{2}^+$	< 4	< 0.4
11.57	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	18 ± 9	0.3 ± 0.2
		5.24	$\frac{5}{2}^+$	63 ± 9	1.2 ± 0.1
		6.18	$\frac{3}{2}^-$	20 ± 9	0.4 ± 0.2
		6.79	$\frac{3}{2}^+$	< 3	< 0.1
		5.24	$\frac{5}{2}^+$	47 ± 7	5 ± 1
11.75 ^a	$\frac{5}{2}^+$	6.18	$\frac{3}{2}^-$	53 ± 7	5 ± 1
		5.24	$\frac{5}{2}^+$	100	1.4 ± 0.6

- ^a For references and other comments see Table 15.19 in (1981AJ01).
- ^b δ =multipole mixing ratio.
- ^c Branches to $^{15}\text{O}^*(5.18, 5.24)$ are $< 2.5\%$ each.
- ^d Branches to $^{15}\text{O}^*(5.18, 5.24, 6.18)$ are < 3 , < 3 and $< 7\%$, respectively.
- ^e Branches to $^{15}\text{O}^*(0, 5.18, 6.18)$ are < 10 , < 4 and $< 0.4\%$, respectively.
- ^f Branches to $^{15}\text{O}^*(5.18, 6.18)$ are < 4 and $< 2\%$, respectively.
- ^g Branchings shown to $^{15}\text{O}^*(5.18, 6.18, 6.79)$ are weighted means of values shown in Table 15.19 of (1981AJ01), recalculated to sum to 100% for all the transitions.
- ^h (1987SC1H).
- ⁱ See, however, the comments in reaction 14 of (1981AJ01).
- ^j Branchings to $^{15}\text{O}^*(6.18, 6.86)$ are $< 1\%$ each.
- ^k Weighted mean of values shown in Table 15.19 of (1981AJ01).
- ^l Intensity $< 25\%$ of transition to $^{15}\text{O}^*(6.79)$.
- ^m Recalculated because of new transition to $^{15}\text{O}^*(\frac{1}{2}^+)$ (1987SC1H).
- ⁿ Γ_γ values assume J -values in column 2.

The yield of n_0 (reaction (b)) shows resonances for $E(^3\text{He}) < 10$ MeV and little structure above, to 30.6 MeV: see (1981AJ01) [n_1 and n_{2+3+4} yields are also reported].

The yield of protons (reaction (c)) shows some clear resonances below $E(^3\text{He}) = 4.5$ MeV and some uncorrelated structures at higher energies (to $E(^3\text{He}) = 12$ MeV) with the possible exception of states at $E_{\text{res}} = 7.8, 9.2 - 9.6$ and (10.5) MeV. For $E(^3\text{He}) = 16$ to 30.6 MeV no appreciable structure is observed in the p_0, p_1 and p_2 yields: see (1976AJ04). At $E(^3\vec{\text{He}}) = 33$ MeV A_y has been measured for $^{14}\text{N}^*(0, 2.31, 3.95)$: see (1986AJ01). For polarization effects in the ($^3\text{He}, 2p$) reaction at $E(^3\vec{\text{He}}) = 33$ MeV see (1986KA44). For reactions (d) and (e) see (1976AJ04, 1981AJ01, 1986AJ01).

The elastic scattering (reaction (f)) shows some resonant structure near 3, 5 and 6 MeV and some largely uncorrelated structures in the range $E(^3\text{He}) = 16.5$ to 24 MeV. There is some suggestion, however, of two resonances at $E(^3\text{He}) = 17$ and 20 MeV: see (1976AJ04). Resonance-like behavior is also reported at $E(^3\vec{\text{He}}) = 29$ MeV. Polarization measurements are reported for $E(^3\vec{\text{He}}) = 20.5$ to 32.6 MeV: see (1981AJ01). See also (1986AJ01). The yield of α -particles displays resonance structure below 8 MeV, and broad fluctuations for $E(^3\text{He}) = 12$ to 18.6 MeV: see (1976AJ04). Polarization measurements are reported for $E(^3\vec{\text{He}}) = 33.3$ MeV for the α_0 and α_1 groups: see (1981AJ01). For A_y measurements of the ($^3\text{He}, ^7\text{Be}$) and ($^3\text{He}, ^6\text{Li}$) reactions see (1989SI02, 1986CL1B). For π^\pm production see (1986MI25). For a search for subthreshold K^+ production see (1985IA01). For work at very high energies see (1985AB1B, 1985AD1C). See also (1986AJ01), (1984NA1F), (1990TO10; applied) and (1986EV01, 1986SI02; theor.).

$$4. \ ^{12}\text{C}(\alpha, n)^{15}\text{O} \quad Q_m = -8.5019$$

Angular distributions of the n_0 group have been measured for $E_\alpha = 18.4$ to 23.1 MeV: see (1976AJ04). At $E_\alpha = 41$ MeV angular distributions are reported to $^{15}\text{O}^*(5.24, 6.86+7.28, 9.63[\text{u}],$

10.48[u], 11.72[u], 12.85[u], 15.05[u]). $^{15}\text{O}^*(8.92, 11.1, 12.3, 13.45, 13.72, 14.27, 15.65)$ are also populated (1981OV01 [uncertainties in E_x are not shown; unresolved states are a problem]). At $E_\alpha = 47.4$ MeV groups are populated at $\theta = 0^\circ$ corresponding to $^{15}\text{O}_{\text{g.s.}}$ and to unresolved states at 5.2, 7.3, 10.0, 12.5 and 15.3 MeV (1988LU02). See also (1988CA26; astrophys.).

Table 15.18: Resonances in $^{12}\text{C} + ^3\text{He}$ ^a

$E(^3\text{He})$ (MeV \pm keV)	Resonant for	$\Gamma_{\text{c.m.}}$ (keV)	J^π	E_x (MeV)
1.21	p_0, p_2		$(\frac{5}{2})^-$	13.04
1.3	$p_0 \rightarrow p_3$			13.1
2.15	n, p_0		$(> \frac{5}{2})$	13.79
2.45 ± 40	$n_0, p_0 \rightarrow p_3$	160 ± 20	$(\frac{1}{2}^-, \frac{3}{2}^-)$	14.03
2.75 ± 40	$n_0, p_1, p_2, ^3\text{He}, \alpha_0$	340 ± 30	$\frac{1}{2}^+$	14.27
(2.87)	p_0, p_2	240		(14.37)
2.990 ± 10	$n_0, p_0, p_1, p_2, p_4,$ $p_5, p_8, ^3\text{He}, \alpha_0$	100 ± 10	$\frac{3}{2}^+, \frac{5}{2}^+$	14.465
3.28 ± 40	$p_0, (p_1, p_2)$	180 ± 40		14.70
3.60 ± 40	p_0, p_1, p_2	400 ± 25		14.95
4.20 ± 10	p_5, p_6, α_0	65 ± 15		15.43
4.37 ± 40	$p_0, p_1, p_2, p_4, p_7,$ p_8, α_0	80 ± 25		15.57
4.65 ± 50	n_0			15.79
4.78 ± 50	$^3\text{He}, \alpha_0$	350	$\frac{1}{2}^-, \frac{3}{2}^-$	15.90
4.97 ± 20	α_0			16.05
5.03 ± 20	$n_0, ^3\text{He}, \alpha_0$			16.10
5.15 ± 20	$n_0, ^3\text{He}, \alpha_0$			16.19
5.45 ± 50	$^3\text{He}, \alpha_0$	170	$\frac{1}{2}^+$	16.43
5.85 ± 50	$n_0, ^3\text{He}$			16.75
6.23 ± 70	γ_0	700 ± 70	$(\frac{1}{2}, \frac{3}{2})^+$	17.05 ± 0.06 ^b
6.83 ± 40	$\gamma_{1+2}, n_0, ^3\text{He}, \alpha_0$	640 ± 120	$\frac{1}{2}^-, \frac{3}{2}^-$	17.53 ^c
7.40 ± 50	^3He	200	$\frac{1}{2}^-, \frac{3}{2}^-$	17.99
7.70 ± 50	n_0, p_0			18.23
8.25 ± 70	γ_0	520 ± 110	$(\frac{1}{2}, \frac{3}{2})^+$	18.67 ± 0.06 ^b
8.70 ± 50	γ_{1+2}, n_0	1120 ± 300		19.03 ^c

Table 15.18: Resonances in $^{12}\text{C} + ^3\text{He}$ ^a (continued)

$E(^3\text{He})$ (MeV \pm keV)	Resonant for	$\Gamma_{\text{c.m.}}$ (keV)	J^π	E_x (MeV)
9.38 ± 100	γ_0	780 ± 270	$(\frac{1}{2}, \frac{3}{2})^+$	19.57 ± 0.08
9.80 ± 50	\mathbf{n}_0			19.91
10.45 ± 90	$\gamma_0, (\text{p}_0)$	970 ± 240	$(\frac{3}{2}, \frac{1}{2})^+$	20.42 ± 0.07 ^b
11.87 ± 80	γ_0	730 ± 120	$(\frac{3}{2}, \frac{1}{2})^+$	21.56 ± 0.07 ^b
14.7	γ_{1+2}	$\lesssim 0.5 \text{ MeV}$ ^e		23.8 ± 0.1 ^c
(17.0) ^d	^3He	≈ 600	$(\frac{13}{2}^-)$	(26.0)
(20.0) ^d	^3He	≈ 2500	$(\frac{9}{2}^-, \frac{11}{2}^-)$	(28.0)
(21.5)	^3He to $^{12}\text{C}^*(15.1)$	≈ 2500		(29.0)

^a For references see Table 15.21 in (1976AJ04).

^b (1978DE33, 1984DE09 [see p.290]); $T = \frac{1}{2}$; $\Gamma_{^3\text{He}}/\Gamma_{\text{p}} = 0.17 \pm 0.07$ and 0.09 ± 0.04 for $^{15}\text{O}^*(17.05, 18.67)$.

^c (1989KI09). See also for ω_γ . See also Table 15.19 in (1986AJ01); $T = \frac{1}{2}$ if they are 3p4h cluster states.

^d $\Gamma_{\text{p}} = 0.06$ and $\geq 0.1 \text{ MeV}$ for $^{15}\text{O}^*(26, 28)$.

^e Estimated by reviewer.

5. $^{12}\text{C}(^6\text{Li}, \text{t})^{15}\text{O}$ $Q_{\text{m}} = -3.7196$

States observed in this reaction are displayed in Table 15.19 (1975BI06: $E(^6\text{Li}) = 59.8 \text{ MeV}$). Comparisons of angular distributions of the triton groups in this reaction and of the ^3He groups to analog states in ^{15}N have been made: analog correspondence is established for (10.48 – 10.70), (12.84 – 13.15 (u)) and (15.05 – 15.49 (u)) [E_x in ^{15}O , E_x in ^{15}N ; u = unresolved] (1975BI06). See also (1976AJ04) for the earlier work.

6. $^{12}\text{C}(^{12}\text{C}, ^9\text{Be})^{15}\text{O}$ $Q_{\text{m}} = -14.2031$

At $E(^{12}\text{C}) = 187 \text{ MeV}$, $\theta_{\text{lab}} = 8^\circ$, the spectrum is dominated by $^{15}\text{O}^*(12.84, 15.05)$ [assumed $J^\pi = \frac{1}{2}^-, \frac{13}{2}^+$, respectively]. $^{15}\text{O}^*(7.28)$ [$J^\pi = \frac{7}{2}^+$] is populated but $^{15}\text{O}^*(0, 6.79)$ are not observed. The situation is similar at $E(^{12}\text{C}) = 114 \text{ MeV}$ but at $E(^{12}\text{C}) = 72 \text{ MeV}$ ($\theta_{\text{lab}} = 11^\circ$) $^{15}\text{O}^*(0, 5.2, 7.28)$ are populated with comparable intensities: see (1976AJ04).

At $E(^{12}\text{C}) = 480 \text{ MeV}$ the three most strongly excited states in the forward direction are $^{15}\text{O}^*(10.46, 12.83[\text{u}], 15.05[\text{u}])$ [$J^\pi = \frac{9}{2}^+, \frac{11}{2}^-, \frac{13}{2}^+$] and forward angle $\sigma(\theta)$ have been measured. $^{15}\text{O}^*(0, 5.24, 7.3, 8.9[\text{u}], 16.7[\text{u}], 18.2[\text{u}], 21.1[\text{u}], 22.1[\text{u}])$ are also populated (1988KR11).

Table 15.19: Levels of ^{15}O from $^{12}\text{C}(^6\text{Li}, t)^{15}\text{O}$ ^a

E_x (MeV \pm keV)	L	E_x (MeV \pm keV)	L
5.180 \pm 5		11.72 \pm 10	c
5.242 \pm 5	b	11.98 \pm 10	
6.179 \pm 5		12.295 \pm 10	c
6.790 \pm 5		12.60 \pm 10	
6.865 \pm 5	b	12.835 \pm 10 ^e	3
7.275 \pm 5	b	13.55 \pm 10	c, d
8.285 \pm 5	b	13.75 \pm 10	c, d
8.918 \pm 5	c	14.27 \pm 10	c
8.978 \pm 5		15.05 \pm 10 ^e	3
9.485 \pm 5		15.48 \pm 10	
9.610 \pm 5	c, d	15.54 \pm 10	
9.658 \pm 5	c, d	15.60 \pm 10	c, d
9.76 \pm 5		15.65 \pm 10	
10.27 \pm 5		15.80 \pm 10	
10.45 \pm 5 ^e	3	17.46 \pm 20	
11.145 \pm 10		17.51 \pm 20	
11.56 \pm 10			

^a (1975BI06): $E(^6\text{Li}) = 59.8$ MeV.

^b Angular distributions measured and compared with those of the $(^6\text{Li}, ^3\text{He})$ reaction to analog states in ^{15}N .

^c Angular distribution measured: analog states in ^{15}N not known.

^d Unresolved in angular distribution.

^e $\Gamma_\gamma/\Gamma < 0.13$.

7. $^{13}\text{C}(^3\text{He}, n)^{15}\text{O}$ $Q_m = 7.1296$

Observed groups are displayed in Table 15.22 of (1981AJ01).

8. $^{14}\text{C}(p, \pi^-)^{15}\text{O}$ $Q_m = -132.115$

At $E_p = 183$ MeV differential cross sections and A_γ are reported for the transitions to $^{15}\text{O}^*(0, 7.3)$, the two states strongly populated in the reaction ([1982JA05](#), [1982VI05](#)). See also ([1986JA1H](#)) and ([1986KU1J](#), [1990KU1H](#); theor.).

9. $^{14}\text{N}(p, \gamma)^{15}\text{O}$ $Q_m = 7.2971$

Observed resonances in the yield of γ -rays are listed in Table [15.20](#). Branching ratios are displayed in Table [15.17](#). Measurements of E_γ lead to $E_x = 5183 \pm 1$, 5240.9 ± 0.4 , 6175 ± 2 , 6794 ± 2 , 6858 ± 2 , 8284.1 ± 0.8 , 8922 ± 2 and 8978 ± 2 keV: see [1981AJ01](#)). For τ_m see ([1981AJ01](#)).

([1987SC1H](#)) have studied absolute cross sections, γ -ray angular distributions and excitation functions for $E_p = 0.2$ to 3.6 MeV: $S(0)$ is determined to be 3.20 ± 0.54 keV \cdot b. C^2S are derived for the first eight states of ^{15}O ([1987SC1H](#)).

The 90° γ_0 yield has been measured for $E_p = 2.2$ to 19 MeV: resonances are observed over most of the range. The $(\gamma_1 + \gamma_2)$ yield is relatively weak. For $E_p = 18$ to 28 MeV the excitation function for γ_0 decreases smoothly with energy: there is no evidence for structures [see ([1981AJ01](#))]. See also ([1985CA41](#), [1987KR04](#), [1987WE1C](#), [1988CA1J](#), [1988CA26](#), [1989BA2P](#), [1989TH1C](#), [1990MA1P](#); astrophysics) and ([1990HA46](#); theor.).

10. $^{14}\text{N}(p, n)^{14}\text{O}$ $Q_m = -5.9255$ $E_b = 7.2971$

The excitation function has been measured for $E_p = 6.3$ to 12 MeV: see ([1970AJ04](#)). Observed resonances are displayed in Table [15.20](#). The cross section [obtained by measuring the 2.31 MeV γ -rays from the ^{14}O (β^+) decay] is reported at 12 energies in the range $E_p = 7$ to 22 MeV ([1981DY03](#)). Production cross sections for the 2.31 MeV γ -rays have been measured at $E_p = 8.9$, 20, 30, 33 and 40 MeV by ([1988LE08](#)). The ratio of the cross section to $^{14}\text{O}_{\text{g.s.}}$ to that for the analog state $^{14}\text{N}^*(2.31)$ [from the (p, p') reaction] has been determined at $E_p = 35$ MeV ([1984TA02](#)). Forward-angle differential cross sections (n_0) are reported by ([1979MO16](#)) at $E_p = 144$ MeV. See also ([1985CA41](#); astrophys.) and ^{14}O .

11. $^{14}\text{N}(p, p)^{14}\text{N}$ $E_b = 7.2971$

The yields of elastic and inelastic protons, and of 2.31 MeV γ -rays, have been studied at many energies: see ([1959AJ76](#), [1970AJ04](#), [1976AJ04](#)). Observed resonances are displayed in Table [15.20](#). At higher energies excitation functions have been measured for the p_0 , p_1 and p_2 groups for $E_p = 17$ to 26.5 MeV: there is no evidence for resonant behavior but the p_1 yield shows a large increase between $E_p = 20$ and 23 MeV. Total cross sections for the $p_0 \rightarrow p_9$ groups have been measured at $E_p = 8.6$, 10.6, 12.6 and 14.6 MeV [see ([1981AJ01](#))]. Total reaction cross sections

have also been measured in the range $E_p = 22.9$ to 49.0 MeV by (1985CA36). (1988LE08) report 2.31–MeV γ -ray production cross sections at $E_p = 8.9, 20, 30, 33$ and 40 MeV. For measurements at $E(^{14}\text{N}) = 516$ MeV/A see (1990WE14).

Polarization measurements have been carried out at $E_p = 3.0$ to 159.4 MeV [see (1970AJ04, 1976AJ04, 1986AJ01)] and at $E_{\bar{p}} = 35$ MeV (1990IE01; p_1) and 0.8 GeV (1985BL22; A_y ; elastic). See also ^{14}N , (1986BA2U), (1986BA88, 1987HU01) and (1989AM01; theor.).

Table 15.20: Resonances in $^{14}\text{N} + p$ ^a

E_p (keV)	Γ_{lab} (keV)	$\omega\Gamma_\gamma$ (eV)	Particles out	J^π	E_x (MeV)
278.1 ± 0.4	1.06 ± 0.11 ^b	$(14 \pm 1) \times 10^{-3}$ ^{a, b}	γ	$\frac{1}{2}^+$	7.5565
1058.0 ± 0.5	3.9 ± 0.7	0.31 ± 0.04 ^{a, b}	γ	$\frac{3}{2}^+$	8.2840
1550 ± 6	34	$(93 \pm 20) \times 10^{-3}$ ^{a, b}	γ	$\frac{1}{2}^+$	8.743
1742 ± 2 ^c	3.5 ± 0.3	0.16	γ, p_0	$\frac{5}{2}^+$	8.922
1742 ± 2 ^c	8	0.06	γ, p_0	$\frac{1}{2}^+$	8.922
1806.4 ± 1.5	4.2 ± 0.4	0.52	γ	$(\frac{3}{2})^-$	8.9821
2344 ± 8 ^b	205 ^b	6.1 ± 1.3 ^b	γ, p_0	$(\frac{3}{2})^+$	9.484
2348 ± 3	10.8 ± 0.5	2.4	γ	$\frac{5}{2}^-$	9.488
2.479 ± 1.7	9.4 ± 0.5	3.3	γ	$\frac{3}{2}^-$	9.609
2537 ± 4	2 ± 1		p_0	$(\frac{7}{2}, \frac{9}{2})^-$	9.664
3209	3 ± 1		p_0	$(\frac{5}{2})^-$	10.291
3215	12 ± 2		p_0	$\frac{5}{2}^+$	10.296
3392 ± 5	< 2	0.029 ± 0.010	γ_2, γ_6	$(\frac{9}{2})^+$	10.461
3410	27 ± 5		γ_0, γ_2, p_0	$(\frac{3}{2})^-$	10.478
3440	150 ± 45		γ, p_0	$(\frac{3}{2})^+$	10.506
3880 ± 15	97		p_0	$\frac{7}{2}^+$	10.916
		Γ_{γ_0} (eV)			
3903 ± 3	106 ± 5	14 ± 3	γ, p_0, p_1	$\frac{1}{2}^+$	10.938
3996 ± 3	27 ± 2	1.4 ± 0.4	γ, p_0, p_1	$\frac{1}{2}^-$	11.025
4130 ± 15	< 10		p_0		11.150
4203 ± 3	43 ± 4	5.2 ± 0.4	γ, p_0	$\frac{3}{2}^+$	11.218
4575 ± 15	< 10		p_0		11.565
4580 ± 15	21 ± 15	0.7 ± 0.2	γ, p_0	$\frac{5}{2}^-$	11.569
4580	150		γ		11.57
4630 ± 15	86 ± 50		γ, p_0	$(\frac{3}{2}, \frac{1}{2})^-$	11.616

Table 15.20: Resonances in $^{14}\text{N} + \text{p}^a$ (continued)

E_p (keV)	Γ_{lab} (keV)	$\omega\Gamma_\gamma$ (eV)	Particles out	J^π	E_x (MeV)
4740 \pm 15	< 10		p ₀		11.718
4772 \pm 3	106 \pm 5		γ , p ₀ , p ₁	$\frac{5}{2}^+$	11.748
4877 \pm 3	70 \pm 3		γ , p ₀ , p ₁	$\frac{5}{2}^-$	11.846
5025 \pm 15	21 \pm 5		p ₀ , p ₁	$\frac{5}{2}^-$	11.984
5180 \pm 15	214 \pm 50		p ₀ , p ₁	$\frac{5}{2}^+$	12.129
5280 \pm 20	106 \pm 50		p ₁ ^d		12.222
5547 \pm 3	82 \pm 4		p ₁ , p ₂	$\frac{5}{2}^- \left(\frac{3}{2}^- \right)$	12.471
5900	\approx 250		γ		12.80
5937 \pm 3	17 \pm 1		p ₂ ^e		12.835
(6100)	30		p ₀ \rightarrow p ₂ , α_0	$\frac{5}{2}^+$	(12.99)
6123 \pm 3	230 \pm 30		p ₂ ^e		13.008
6141 \pm 3	43 \pm 30		p ₂ ^e		13.025
6600	\approx 1000		γ , (p ₂ , α_0)	$\left(\frac{1}{2}, \frac{3}{2} \right)^+$	13.45
6640			(p ₀), (p ₂)	$\left(\frac{3}{2}^+ \right)$	13.49
6760			α_0	$\frac{5}{2}^+$	13.60
6870			p ₂	$\frac{3}{2}^-$	13.70
6960			p ₁ , p ₂ , p ₄ , α_0	$\frac{3}{2}^-$	13.79
7050	\approx 150		γ		13.87
7370			α_0	$\frac{5}{2}^-$	14.17
7500	\approx 500		n, p ₀ \rightarrow p ₂ , ³ He, α		14.29
7550			α_0	$\frac{5}{2}^+$	14.34
7700			n, p ₀ , α_0		14.48
7950	170 \pm 50		n		14.71
8200			n, p ₂ \rightarrow p ₆ , ³ He, α_0 , α_1		14.94
8400	\approx 1000		γ	$\left(\frac{1}{2}, \frac{3}{2} \right)^+$	15.1
9050			n		15.74
f					
9370 \pm 20	\approx 200		n, p ₂ , p ₈ , α_1		16.04

Table 15.20: Resonances in $^{14}\text{N} + \text{p}$ ^a (continued)

E_p (keV)	Γ_{lab} (keV)	$\omega\Gamma_\gamma$ (eV)	Particles out	J^π	E_x (MeV)
9580 ± 20	≈ 150		p_0, p_1, p_3 $\rightarrow p_7, p_9, {}^3\text{He},$ α_1		16.23
9850 ± 50	600 ± 100		$n, {}^3\text{He}$		16.48
10300	≈ 1000		γ	$(\frac{1}{2}, \frac{3}{2})^+$	16.9
10600			$p_4 \rightarrow p_9, \alpha_0, \alpha_1$		17.2
11900	≈ 1000		γ	$(\frac{1}{2}, \frac{3}{2})^+$	18.4
14200	≈ 2000		γ	$(\frac{1}{2}, \frac{3}{2})^+$	20.5
15800	≈ 2000		γ	$(\frac{1}{2}, \frac{3}{2})^+$	22.0

^a For references see (1970AJ04, 1976AJ04, 1981AJ01). See also Table 15.17 here.


^b (1987SC1H). See also (1987KR04; theor.).

^c Separated by 0.5 ± 0.5 keV: see, however, reaction 14 in (1981AJ01).

^d Weak.

^e Strong.

^f See footnote ^e in Table 15.23 of (1981AJ01).



$$Q_m = -4.7789$$

$$E_b = 7.2971$$


Excitation functions for the ground-state group have been measured at $E_p = 7$ to 11 MeV: some resonant structure is indicated [see Table 15.20]. See also (1976AJ04).



$$Q_m = -2.9228$$

$$E_b = 7.2971$$

Excitation functions and total cross-section measurements have been measured for the α_0 group for $E_p = 3.8$ to 45 MeV: see (1976AJ04). Fairly sharp structures persist until $E_p = 15$ MeV: see Table 15.20 here and footnote ^e in Table 15.23 of (1981AJ01). See also (1986AI04; applied) and (1985CA41; astrophys.).



$$Q_m = 5.0725$$

Table 15.21: Levels of ^{15}O from $^{14}\text{N}(\text{d}, \text{n})$ and $^{14}\text{N}({}^3\text{He}, \text{d})$ ^a

E_x in ^{15}O ^b (MeV \pm keV)	l_p	S	J^π
0	1 ^d	0.87	$\frac{1}{2}^-$
5.18	(0) ^e	0	$\frac{1}{2}^+$
5.2410 ± 0.5 ^c	2 ^d	(0.03)	$\frac{5}{2}^+$
6.180 ± 4 ^c	1 ^d	0.04	$\frac{3}{2}^-$
6.79	0 ^d	≤ 0.3	$\frac{3}{2}^+$
6.8598 ± 1.0 ^c	2 ^d	0.4	$\frac{5}{2}^+$
7.2762 ± 0.6 ^c	2 ^d	0.42	$\frac{7}{2}^+$
7.56	0 ^d	≤ 0.4	$\frac{1}{2}^+$
8.28	0 ^e		$\frac{3}{2}^+$

^a See Tables 15.27 in (1970AJ04) and 15.26 in (1976AJ04) for references and additional information.

^b Nominal energies if uncertainty is not indicated.

^c From γ -ray measurements.

^d From both (d, n) and (${}^3\text{He}$, d) work: see (1976AJ04).

^e From (${}^3\text{He}$, d).

Angular distributions have been studied at many energies in the range $E_d = 0.9$ to 11.8 MeV: see Tables 15.27 and 15.28 in (1970AJ04) and Table 15.26 in (1976AJ04). For τ_m measurements see (1970AJ04). See also Table 15.21 here, ^{16}O in (1986AJ04), (1987HI1B; applied) and (1984BL21; theor.).

$$15. \quad {}^{14}\text{N}({}^3\text{He}, \text{d})^{15}\text{O} \quad Q_m = 1.8035$$

See Table 15.28 in (1970AJ04). See also Table 15.21 here.

$$16. \quad {}^{15}\text{N}(\gamma, \pi^-)^{15}\text{O} \quad Q_m = -142.322$$

At $E_\gamma = 170$ MeV four-point angular distributions of the π^- to $^{15}\text{O}_{\text{g.s.}}$ have been measured by (1988LI23) and (1989KO28): the two studies are not in good agreement. See also (1990ER03; theor.).

17. $^{15}\text{N}(\pi^+, \pi^0)^{15}\text{O}$ $Q_m = 1.850$

Angular distributions of the π^0 to $^{15}\text{O}_{\text{g.s.}}$ have been studied at $E_{\pi^+} = 32.4$ and 55.5 MeV (1985IR02), at 48 MeV (1984CO04, 1986LE01) and at 165 MeV (1982DO10). Forward-angle differential cross sections of the π^0 to $^{15}\text{O}_{\text{g.s.}}$ have also been measured at $E_{\pi^+} = 20$ MeV (1987IR01) and at 40.7 and 63.6 MeV (1985IR02). See also (1989LE1L).

18. $^{15}\text{N}(\text{p}, \text{n})^{15}\text{O}$ $E_m = -3.5363$

Angular distributions have been measured for $E_p = 3.95$ to 18.5 MeV [see (1981AJ01, 1986AJ01)], at 35 MeV (1987OR01; $^{15}\text{O}^*(7.56)$ [$J^\pi = \frac{1}{2}^+$]), at 135 MeV (1985WA24; $^{15}\text{O}^*(0, 6.2)$) and at 200 and 494 MeV (1988CIZZ; prelim.; also A_y). Forward-angle differential cross sections at $E_p = 200, 300$ and 400 MeV to $^{15}\text{O}^*(0, 6.2)$ are reported by (1987ALZW; prelim.). The ratio of the population of $^{15}\text{O}_{\text{g.s.}}$ to that of $^{15}\text{O}^*(6.2)$ has been determined at $E_p = 800$ MeV (1986KI12). $^{15}\text{O}^*(6.2)$ contains only about $\frac{1}{3}$ of the expected GT strength (1985GO02). [I am indebted to Prof. C.D. Goodman for his comments.] For a discussion of GT strengths, see (1987TA13). $^{15}\text{O}^*(5.2[\text{u}], 6.8[\text{u}], 7.28)$ have also been populated (1987OR01). For the earlier work see (1986AJ01). See also ^{16}O in (1986AJ04), (1985GO1Q, 1986VO1G, 1987BE25, 1988RO17, 1988WA1Q), (1986MA1P, 1987HI1B, 1988HI1F; applied) and (1989RA15; theor.).

19. $^{15}\text{N}(^3\text{He}, \text{t})^{15}\text{O}$ $Q_m = -2.7725$

Angular distributions for the $t_0, t_{1+2}, t_3, t_{4+5}, t_6$ and t_7 groups have been studied for $E(^3\text{He}) = 16.5$ to 44.6 MeV: see (1976AJ04).

20. $^{16}\text{O}(\gamma, \text{n})^{15}\text{O}$ $Q_m = -15.6638$

The spectrum of photoneutrons has been investigated at many energies. Measurements over the giant dipole resonance region show the predominant strength is to the $J^\pi = \frac{1}{2}^-$ and $\frac{3}{2}^-$ states $E_x = 0$ and 6.18 MeV, consistent with the basic validity of the single-particle, single-hole theory of photoexcitation in ^{16}O . However, the positive-parity states at $E_x = 5.18, 5.24, 6.86$ MeV are also populated suggesting some more complicated excitations in ^{16}O : see (1970AJ04, 1976AJ04). Differential cross sections for the n_0 group have been measured from threshold to $E_\gamma = 28$ MeV [see (1976AJ04)], at $E_\gamma = 60$ to 160 MeV (also $^{15}\text{O}^*(6.18)$; no appreciable strength in the 5.2 MeV doublet) [see (1986AJ01)] and at $150, 200$ and 250 MeV (1989BE14). See also ^{16}O in (1986AJ04) and (1987RY03, 1988CA10, 1988DUZX, 1988RY03, 1990BRZY, 1990FL1A; theor.).

21. (a) $^{16}\text{O}(\pi^+, p)^{15}\text{O}$ $Q_m = 124.687$
 (b) $^{16}\text{O}(\pi^+, \pi^+n)^{15}\text{O}$ $Q_m = -15.6638$
 (c) $^{16}\text{O}(\pi^+, \pi^0p)^{15}\text{O}$ $Q_m = -10.277$

For reaction (a) see (1982DO01). At $E_{\pi^+} = 2.0$ GeV/c differential cross sections have been determined for the transition to $^{15}\text{N}^*(6.2)$ (1983KI01) in reaction (b). For reaction (c), see (1986GI1A).

22. $^{16}\text{O}(p, pn)^{15}\text{O}$ $Q_m = -15.6638$

At $E_p = 505$ MeV the summed spectra show two peaks corresponding to the $p_{1/2}$ and $p_{3/2}$ knockouts [$^{15}\text{O}^*(0, 6.18)$] (binding energies of 15.64 and 21.82 MeV). Differential cross sections are also reported (1986MC10) [see also reaction 59 in ^{15}N]. For work at 1 GeV, see (1985BE30). See also (1983WA1C) and (1987HI1B; applied).

23. $^{16}\text{O}(p, d)^{15}\text{O}$ $Q_m = -13.4392$

Angular distributions have been reported at many energies for $E_p = 18.5$ to 155.6 MeV [see Table 15.30 in (1970AJ04), (1976AJ04), (1986AJ01)]: at those energies $^{15}\text{O}^*(0, 6.18)$ are preferentially populated. At $E_p = 200$ MeV angular distributions have been studied for $^{15}\text{O}^*(0, 6.18)$ (1989AB01) [also A_y and C^2S]. At $E_p = 800$ MeV $^{15}\text{O}^*(0, 5.2[\text{u}], 6.18, 7.4[\text{u}], 9.0[\text{u}], 10.42 \pm 0.15, 10.87 \pm 0.15, 12.21 \pm 0.15, 13.59 \pm 0.15, 19.0 \pm 0.2, 21.1 \pm 0.2)$ [the last two states have $\Gamma \geq 0.8$ MeV] have been populated (1984SM04).

For γ -ray production [$^{15}\text{O}^*(5.24)$] see (1988LE08). See also ^{17}F in (1986AJ04), (1989WA16) and (1987LA11, 1988LE08, 1989GU28; astrophysics).

24. $^{16}\text{O}(d, t)^{15}\text{O}$ $Q_m = -9.4065$

Angular distributions have been reported at a number of energies in the range $E_d = 20$ to 52 MeV [see (1981AJ01) and reaction 60 in ^{15}N here] and at $E_d = 89$ MeV (1990SA27; $^{15}\text{O}^*(0, 6.18)$). See also ^{18}F in (1983AJ01).

25. $^{16}\text{O}(^3\text{He}, \alpha)^{15}\text{O}$ $Q_m = 4.9140$

The $p_{1/2}$ and $p_{3/2}$ hole states $^{15}\text{O}^*(0, 6.18)$ are strongly populated. Information on these and other states are displayed in Table 15.25 of (1981AJ01). Angular distributions have been measured at energies up to $E(^3\text{He}) = 217$ MeV: see (1981AJ01). Branching ratios and multipole mixing ratios are displayed in Table 15.17. (1978BE73) report τ_m of $^{15}\text{O}^*(5.24) = 3.25 \pm 0.30$ ps, $|g| = 0.260 \pm 0.028$. (1983BI10) determine $g = +0.17 \pm 0.07$. See also (1986AJ01), ^{19}Ne in (1983AJ01) and (1990AB1G; applied).

$$26. \ ^{16}\text{O}(^6\text{Li}, ^7\text{Li})^{15}\text{O} \quad Q_m = -8.414$$

See (1986GL1E; prelim.).

$$27. \ ^{17}\text{O}(p, t)^{15}\text{O} \quad Q_m = -11.3254$$

At $E_p = 39.8$ MeV angular distributions of t_0 and t_3 groups have been compared to those of the ^3He groups to the analog states in ^{15}N . At $E_p = 45$ MeV a state, assumed to be the $J^\pi = \frac{5}{2}^+$, $T = \frac{3}{2}$ analog of $^{15}\text{C}^*(0.74)$, is observed at $E_x = 12.255 \pm 0.013$ MeV, $\Gamma_{\text{c.m.}} = 135 \pm 15$ keV. The state decays by proton emission to the $T = 1, 0^+$ state $^{14}\text{N}^*(2.31)$ [the population of some $T = \frac{1}{2}$ states is also reported]: see (1981AJ01).

$E_{\bar{p}} = 89.7$ MeV angular distributions and A_y measurements have been reported to $^{15}\text{O}^*(0, 5.24[\text{u}], 6.18, 6.86, 7.28)$ (1985VO12).

$$28. \ ^{19}\text{F}(^3\text{He}, ^7\text{Li})^{15}\text{O} \quad Q_m = -4.3185$$

See (1976AJ04). See also (1986HA1E; theor.).

^{15}F
(Fig. 4)

GENERAL (See also (1986AJ01)).

See (1989AYZU, 1989OG1B) and (1985AN28, 1986AN07, 1988CO15; theor.). See (1986AN07) for comments on ^{15}Ne .

Mass of ^{15}F : The atomic mass excess of ^{15}F is 16.77 ± 0.13 MeV. ^{15}F is unstable with respect to breakup into $^{14}\text{O} + p$ by 1.47 MeV: see (1981AJ01).

$$1. \ ^{12}\text{C}(^3\text{He}, \pi^-)^{15}\text{F} \quad Q_m = -141.41$$

Table 15.22: Energy levels of ^{15}F

E_x in ^{15}F (MeV)	$J^\pi; T$	$\Gamma_{\text{c.m.}}$ (MeV)	Decay	Reaction
g.s.	$(\frac{1}{2}^+); \frac{3}{2}$	1.0 ± 0.2	p	2
1.3 ± 0.1	$(\frac{5}{2}^+); \frac{3}{2}$	0.24 ± 0.03	p	2

This reaction is not observed at $E(^3\text{He}) = 235$ MeV, $\theta_{\text{lab}} = 20^\circ$: the differential cross section (c.m.) is $\leq 4 \times 10^{-11}$ b (1984BI08).

2. $^{20}\text{Ne}(^3\text{He}, ^8\text{Li})^{15}\text{F}$ $Q_{\text{m}} = -29.83$

This reaction has been studied at $E(^3\text{He}) = 74.5$ MeV (1978BE26) and 75.4 and 87.8 MeV (1978KE06). Two groups are observed: the ground state [$\Gamma_{\text{c.m.}} = 0.8 \pm 0.3$ MeV (1978KE06), 1.2 ± 0.3 MeV (1978BE26)] and a relatively strongly populated state, presumed to be the mirror of $^{15}\text{C}^*(0.74)$ [$J^\pi = \frac{5}{2}^+$], with $E_x = 1.3 \pm 0.1$ MeV (1978KE06), 1.2 ± 0.2 MeV (1978BE26) and $\Gamma_{\text{c.m.}} = 0.5 \pm 0.2$ MeV (1978KE06), 0.24 ± 0.03 MeV (1978BE26). The differential cross section for populating $^{15}\text{F}^*(1.3)$ is 250 ± 20 nb/sr at 10° and $E(^3\text{He}) = 74.5$ MeV (1978BE26) and 80 ± 25 nb/sr at 9° , 87.8 MeV (1978KE06). At $E(^3\text{He}) = 75.4$ MeV, $\theta = 9^\circ$, the ground state is populated with a differential cross section of 8 ± 4 nb/sr (1978KE06).

References

(Closed 01 July 1990)

- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- 1973TS02 I. Tserruya, B. Rosner and K. Bethge, Nucl. Phys. A213 (1973) 22
- 1975BI06 H.G. Bingham, M.L. Halbert, D.C. Hensley, E. Newman, K.W. Kemper and L.A. Charlton, Phys. Rev. C11 (1975) 1913
- 1975MO28 R. Moreh and O. Shahal, Nucl. Phys. A252 (1975) 429
- 1976AJ04 F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1
- 1976AL16 D.E. Alburger, Nucl. Instrum. Meth.136 (1976) 323
- 1976BE1B Beukens, Unpublished Ph.D. Thesis, Univ. Toronto (1976)
- 1978BE26 W. Benenson, E. Kashy, A.G. Ledebuhr, R.C. Pardo, R.G.H. Robertson and L.W. Robinson, Phys. Rev. C17 (1978) 1939
- 1978BE73 F.A. Beck, T. Byrski, G.J. Costa, W.L. Randolph and J.P. Vivien, Hyperfine Interactions 4 (1978) 181
- 1978DE33 W. Del Bianco, J.C. Kim and G. Kajrys, Can. J. Phys. 56 (1978) 1054
- 1978KE06 G.J. KeKelis, M.S. Zisman, D.K. Scott, R. Jahn, D.J. Vieira, J. Cerny and F. Ajzenberg-Selove, Phys. Rev. C17 (1978) 1929
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buym, Table of Isotopes 7th ed. (New York: John Wiley & Sons, 1978)
- 1979AL23 D.E. Alburger and D.J. Millener, Phys. Rev. C20 (1979) 1891
- 1979KO26 L. Koester, K. Knopf and W. Waschkowski, Z. Phys. A292 (1979) 95
- 1979MO16 G.L. Moake, L.J. Gutay, R.P. Scharenberg, P.T. Debevec and P.A. Quin, Phys. Rev. Lett. 43 (1979) 910
- 1980ZE02 A.F. Zeller, K.W. Kemper, T.R. Ophel and A. Johnston, Nucl. Phys. A344 (1980) 307
- 1981AJ01 F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1
- 1981DY03 P. Dyer, D. Bodansky, A.G. Seamster, E.B. Norman and D.R. Maxson, Phys. Rev. C23 (1981) 1865
- 1981MO09 R. Moreh, W.C. Sellyey and R. Vodhanel, Phys. Rev. C23 (1981) 988; Erratum Phys. Rev. C24 (1981) 2394
- 1981MUZQ S.F. Mughabghab, M. Divadeenam and N.E. Holden, Neutron Cross Sections 1A (1981) Z=1-60

- 1981OV01 D.J. Overway and W.C. Parkinson, Nucl. Phys. A363 (1981) 93
- 1981WA06 E.K. Warburton and D.E. Alburger, Phys. Rev. C23 (1981) 1234
- 1982BE02 M. Bernheim, A. Bussiere, J. Mougey, D. Royer, D. Tarnowski, S. Turck-Chieze, S. Frullani, S. Boffi, C. Giusti, F.D. Pacati et al, Nucl. Phys. A375 (1982) 381
- 1982BU1A Burbidge and Burbidge, Essays in Nucl. Astrophys. (1982) 11
- 1982CA1A Cameron, Essays in Nucl. Astrophys. (1982) 23
- 1982DO01 K.G.R. Doss, P.D. Barnes, N. Colella, S.A. Dytman, R.A. Eisenstein, C. Ellegaard, F. Takeuchi, W.R. Wharton, J.F. Amann, R.H. Pehl et al, Phys. Rev. C25 (1982) 962
- 1982DO10 A. Doron, J. Alster, A. Erell, M.A. Moinester, R.A. Anderson, H.W. Baer, J.D. Bowman, M.D. Cooper, F.H. Cverna, C.M. Hoffman et al, Phys. Rev. C26 (1982) 189
- 1982JA05 W.W. Jacobs, T.G. Throwe, S.E. Vigdor, M.C. Green, J.R. Hall, H.O. Meyer, W.K. Pitts and M. Dillig, Phys. Rev. Lett. 49 (1982) 855
- 1982JU03 J.W. Jury, B.L. Berman, J.G. Woodworth, M.N. Thompson, R.E. Pywell and K.G. McNeill, Phys. Rev. C26 (1982) 777
- 1982VI05 S.E. Vigdor, T.G. Throwe, M.C. Green, W.W. Jacobs, R.D. Bent, J.J. Kehayias, W.K. Pitts and T.E. Ward, Phys. Rev. Lett. 49 (1982) 1314
- 1982WE01 S.A. Wender, H.R. Weller, N.R. Roberson, D.R. Tilley and R.G. Seyler, Phys. Rev. C25 (1982) 89
- 1982WO1A Woosley and Weaver, Essays in Nucl. Astrophys. (1982) p. 377
- 1983AJ01 F. Ajzenberg-Selove, Nucl. Phys. A392 (1983) 1; Errata Nucl. Phys. A413 (1984) 168
- 1983BI10 J. Billowes, J. Burde, J.A.G. De Raedt, M.A. Grace, W.R. Kolbl and A. Pakou, J. Phys. G9 (1983) 1407
- 1983KI01 I.V. Kirpichnikov, V.A. Kuznetsov, A.S. Starostin, E.F. Kislyakov, V.L. Korotkikh and D.E. Lansky, Nucl. Phys. A392 (1983) 352
- 1983ST04 U. Straumann, M. Lebrun, C.J. Martoff, P. Truol, H.R. Kissener, J.P. Perroud, C. Joseph, W. Dahme and K.M. Crowe, Phys. Rev. C27 (1983) 2771
- 1983TR12 S. Truong and H.T. Fortune, Phys. Rev. C28 (1983) 977
- 1983VA1E A. Van Der Schaaf, E.A. Hermes, R.J. Powers, F.W. Schlegel, R.G. Winter, A. Zglin-ski, T. Kozlowski, W. Bertl, L. Felawka, W. H.A. Hesselink et al, Nucl. Phys. A408 (1983) 573
- 1983WA03 J.D. Watson, J.W. Jury, P.C.-K. Kuo, W.F. Davidson, N.K. Sherman and K.G. McNeill, Phys. Rev. C27 (1983) 506
- 1983WA1C Watson et al, in Florence (1983) 473
- 1984BI08 L. Bimbot, T. Hennino, J.C. Jourdain, Y. Le Bornec, F. Reide, N. Willis and J.F. Germond, Phys. Rev. C30 (1984) 739

- 1984BL21 L.D. Blokhintsev, A.M. Mukjamedzhanov and A.N. Safronov, *Fiz. Elem. Chastits At. Yad.* 15 (1984) 1296; *Sov. J. Part. Nucl* 15 (1984) 580
- 1984BO1H Bogdanova and Markushin, *Sov. J. Part. Nucl.* 15 (1984) 361
- 1984CO04 M.D. Cooper, H.W. Baer, R. Bolton, J.D. Bowman, F. Cverna, N.S.P. King, M. Leitch, J. Alster, A. Doron, A. Erell et al, *Phys. Rev. Lett.* 52 (1984) 1100
- 1984DA17 B. Dasmahapatra, B. Cujec and F. Lahlou, *Nucl. Phys.* A427 (1984) 186
- 1984DE09 W. Del Bianco, G. Kajrys, J. Kim, S. Kundu, S. Landsberger, R. Lecomte and S. Monaro, *Can. J. Phys.* 62 (1984) 288
- 1984DE53 P. De Bievre, M. Gallet, N.E. Holden and I.L. Barnes, *J. Phys. Chem. Ref. Data* 13 (1984) 809
- 1984DU15 J.P. Dufour, S. Beraud-Sudreau, R. Del Moral, H. Emmermann, A. Fleury, F. Hubert, C. Poinot, M. Pravikoff, J. Frehaut, M. Beau et al, *Z. Phys.* A319 (1984) 237
- 1984KY01 G.S. Kyle, P.-A. Amaudruz, Th.S. Bauer, J.J. Domingo, C.H.Q. Ingram, J. Jansen, D. Renker, J. Zichy, R. Stamminger and F. Vogler, *Phys. Rev. Lett.* 52 (1984) 974
- 1984NA1F Nakamura, *Indian J. Phys.* 58A (1984) 12
- 1984RA07 W.D.M. Rae and R.K. Bhowmik, *Nucl. Phys.* A420 (1984) 320
- 1984SM04 G.R. Smith, J.R. Shepard, R.L. Boudrie, R.J. Peterson, G.S. Adams, T.S. Bauer, G.J. Igo, G. Pauletta, C.A. Whitten, Jr., A. Wriekat et al, *Phys. Rev.* C30 (1984) 593
- 1984TA02 T.N. Taddeucci, R.R. Doering, A. Galonsky and S.M. Austin, *Phys. Rev.* C29 (1984) 764
- 1984TA07 T.N. Taddeucci, T.A. Carey, C. Gaarde, J. Larsen, C.D. Goodman, D.J. Horen, T. Masterson, J. Rapaport, T.P. Welch and E. Sugarbaker, *Phys. Rev. Lett.* 52 (1984) 1960
- 1984VA06 A.G.M. van Hees and P.W.M. Glaudemans, *Z. Phys.* A315 (1984) 223
- 1984WA07 E.K. Warburton, D.E. Alburger and D.J. Millener, *Phys. Rev.* C29 (1984) 2281
- 1984YA03 M. Yasue, J.J. Hamill, H.C. Bhang, M.A. Rumore and R.J. Peterson, *Phys. Rev.* C30 (1984) 770
- 1984ZH1B Zhuang Fei, Chen Hua-Zhong and Jin Xing-nan, *Phys. Energ. Fortis Phys. Nucl.* 8 (1984) 215
- 1985AB1B Ableev et al, *Sov. J. Nucl. Phys.* 42 (1985) 129
- 1985AD1C B.P. Adyasevich, V.G. Antonenko, Yu.I. Grigoryan, M.S. Ippolitov, K.V. Karadzhev, A.L. Lebedev, V.I. Manko, S.A. Nikolaev, Yu.P. Polunin, A.A. Vinogradov and A.A. Tsvetkov, *Phys. Lett.* B161 (1985) 55
- 1985AJ01 F. Ajzenberg-Selove, *Nucl. Phys.* A433 (1985) 1; Erratum *Nucl. Phys.* A449 (1986) 155
- 1985AL1G Aleksandrov et al, in *Questions in At. Phys. and in Tech.*, USSR (1985) 3

- 1985AN28 M.S. Antony, J. Britz, J.B. Bueb and A. Pape, *At. Data Nucl. Data Tables* 33 (1985) 447
- 1985AR11 A. Arima, T. Cheon and K. Shimizu, *Hyperfine Interactions* 21 (1985) 79
- 1985AR1H A. Arima, *Nucl. Phys.* A446 (1985) 45
- 1985BA1M Bahcall, *Solar Phys.* 100 (1985) 53
- 1985BA1N Bahcall, *AIP Conf. Proc.* 126 (1985) 60
- 1985BA51 A.J. Baltz, C.B. Dover, M.E. Sainio, A. Gal and G. Toker, *Phys. Rev.* C32 (1985) 1272
- 1985BE30 S.L. Belostotsky, S.S. Volkov, A.A. Vorobyev, Yu.V. Dotsenko, L.G. Kudin, N.P. Kuropatkin, O.V. Miklukho, V.N. Nikulin and O.E. Prokofyev, *Yad. Fiz.* 41 (1985) 1425; *Sov. J. Nucl. Phys.* 41 (1985) 903
- 1985BE40 C. Beck, F. Haas, R.M. Freeman, B. Heusch, J.P. Coffin, G. Guillaume, F. Rami and P. Wagner, *Nucl. Phys.* A442 (1985) 320
- 1985BL20 P.G. Blunden and B. Castel, *Nucl. Phys.* A445 (1985) 742
- 1985BL22 G.S. Blanpied, B.G. Ritchie, M.L. Barlett, G.W. Hoffmann, J.A. McGill, M.A. Franey and M. Gazzaly, *Phys. Rev.* C32 (1985) 2152
- 1985BO1P Boyd et al, *Nucl. Instrum. Meth. Phys. Res.* B10-11 (1985) 378
- 1985CA17 G.P. Capitani, E. De Sanctis, P. Levi-Sandri, M. Bernheim, S. Turck-Chieze, S. Frulani and J. Mougey, *Nuovo Cim.* 85A (1985) 37
- 1985CA36 R.F. Carlson, A.J. Cox, T.N. Nasr, M.S. De Jong, D.L. Ginther, D.K. Hasell, A.M. Sourkes, W.T.H. Van Oers and D.J. Margaziotis, *Nucl. Phys.* A445 (1985) 57
- 1985CA41 G.R. Caughlan, W. A. Fowler, M.J. Harris and B.A. Zimmerman, *At. Data Nucl. Data Tables* 32 (1985) 197
- 1985DA23 S.E. Darden, G. Murillo and S. Sen, *Phys. Rev.* C32 (1985) 1764
- 1985FI08 L.K. Fifield, P.V. Drumm, M.A.C. Hotchkis, T.R. Ophel and C.L. Woods, *Nucl. Phys.* A437 (1985) 141
- 1985GO02 C.D. Goodman, R.C. Byrd, I.J. Van Heerden, T.A. Carey, D.J. Horen, J.S. Larsen, C. Gaarde, J. Rapaport, T.P. Welch, E. Sugarbaker et al, *Phys. Rev. Lett.* 54 (1985) 877; *Erratum Phys. Rev. Lett.* 54 (1985) 2060
- 1985GO1A Goncharova, Kissener and Eramzhyan, *Sov. J. Part. Nucl.* 16 (1985) 337
- 1985GO1B Goncharova, Kissener and Eramzhyan, *Izv. Akad. Nauk SSSR Ser. Fiz.* 49 (1985) 1032
- 1985GO1Q Goodman, in *AIP Conf. proc.* 126 (1985) 109
- 1985HA1N B.G. Harvey, *Nucl. Phys.* A444 (1985) 498
- 1985HA40 R.C. Haight, G.J. Mathews and R.W. Bauer, *Nucl. Instr. Meth. Phys. Res.* B10-11 (1985) 361

- 1985HU04 M.S. Hussein, B.V. Carlson, O. Civitarese and A. Szanto De Toledo, Phys. Rev. Lett. 54 (1985) 2659
- 1985HU1C Hufner, Phys. Rep. 125 (1985) 129
- 1985IA01 F. Iazzi, B. Minetti, T. Bressani, E. Chiavassa, S. Costa, G. Dellacasa, M. Gallio, A. Musso, G. Puddu and S. Serici, Lett. Nuovo Cim. 43 (1985) 305
- 1985IR02 F. Irom, M.J. Leitch, H.W. Baer, J.D. Bowman, M.D. Cooper, B.J. Dropesky, E. Pisetzky and J.N. Knudson, Phys. Rev. Lett. 55 (1985) 1862
- 1985KI1A Kitching, McDonald, Maris and Vasconcellos, Adv. Nucl. Phys. 15 (1985) 43
- 1985KW02 E. Kwasniewicz and L. Jarczyk, Nucl. Phys. A441 (1985) 77
- 1985LE07 M.J. Leitch, J.L. Matthews, W.W. Sapp, C.P. Sargent, S.A. Wood, D.J.S. Findlay, R.O. Owens and B.L. Roberts, Phys. Rev. C31 (1985) 1633
- 1985LE1E Lenz, Lecture Notes in Phys. 234 (1985) 336
- 1985LI1H Liubinskii, Melenevskii, Nemikin and Tikhii, in Leningrad (1985) 500
- 1985MA1K Masutani, in AIP Conf. Proc. 133 (1985) 312
- 1985MC1C McNally, Fusion Tech. 7 (1985) 331
- 1985ME1E Melenevskii, Muzika, Romanova and Tikhii, in Leningrad (1985) 499
- 1985NO1C Nomura, J. Phys. Soc. Jpn. Suppl. 54 (1985) 295
- 1985PH01 R.J. Philpott, Nucl. Phys. A439 (1985) 397
- 1985PO10 N.A.F.M. Poppelier, L.D. Wood and P.W.M. Glaudemans, Phys. Lett. B157 (1985) 120
- 1985PO11 D.N. Poenaru, M. Ivascu, A. Sandulescu and W. Greiner, Phys. Rev. C32 (1985) 572
- 1985PR1D Prombo and Clayton, Science 230 (1985) 935
- 1985SA31 D.P. Sanderson, J.A. Carr and K.W. Kemper, Phys. Rev. C32 (1985) 1169
- 1985SE06 S.J. Seestrom-Morris, D. Dehnhard, C.L. Morris, L.C. Bland, R. Gilman, H.T. Fortune, D.J. Millener, D.P. Saunders, P.A. Seidl, R.R. Kiziah et al, Phys. Rev. C31 (1985) 923
- 1985SH1D Shvedov and Nemets, in Leningrad (1985) 317
- 1985SH24 R. Sherr and G. Bertsch, Phys. Rev. C32 (1985) 1809
- 1985SI19 K. Siwek-Wilczynska, R.A. Blue, L.H. Harwood, R.M. Ronningen, H. Utsunomiya, J. Wilczynski and D.J. Morrissey, Phys. Rev. C32 (1985) 1450
- 1985TA1A Taam, Ann. Rev. Nucl. Part. Sci. 35 (1985) 1
- 1985TA23 T.N. Taddeucci, C.D. Goodman, R.C. Byrd, T.A. Carey, D.J. Horen, J. Rapaport and E. Sugarbaker, Nucl. Instrum. Meth. A241 (1985) 448
- 1985TE01 J.A. Templon, J.H. Dave, C.R. Gould and S. Singkarat, Nucl. Sci. Eng. 91 (1985) 451

- 1985TU02 R.S. Turley, E.R. Kinney, J.L. Matthews, W.W. Sapp, E.J. Scheidker, R.A. Schumacher, S.A. Wood, G.S. Adams and R.O. Owens, Phys. Lett. B157 (1985) 19
- 1985UT01 H. Utsunomiya, Phys. Rev. C32 (1985) 849
- 1985VO12 K.F. von Reden, W.W. Daehnick, S.A. Dytman, R.D. Rosa, J.D. Brown, C.C. Foster, W.W. Jacobs and J.R. Comfort, Phys. Rev. C32 (1985) 1465
- 1985WA24 J.W. Watson, W. Pairsuwan, B.D. Anderson, A.R. Baldwin, B.S. Flanders, R. Madey, R.J. McCarthy, B.A. Brown, B.H. Wildenthal and C.C. Foster, Phys. Rev. Lett. 55 (1985) 1369
- 1986AI04 A. Ait Haddou, M. Berrada and G. Paic, J. Radioanal. Nucl. Chem. 102 (1986) 159
- 1986AI1A Aichelin and Stocker, Phys. Lett. B176 (1986) 14
- 1986AJ01 F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1
- 1986AJ04 F. Ajzenberg-Selove, Nucl. Phys. A460 (1986) 1
- 1986AL18 W.P. Alford, R.L. Helmer, R. Abegg, A. Celler, O. Hausser, K. Hicks, K.P. Jackson, C.A. Miller, S. Yen, R.E. Azuma et al, Phys. Lett. B179 (1986) 20
- 1986AM1B Amsel, Cohen and Maurel, Nucl. Instrum. Meth. Phys. Res. B14 (1986) 226
- 1986AN07 M.S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables 34 (1986) 279
- 1986AN1R Ansari, Shoeb and Rahman Khan, J. Phys. G12 (1986) 1369
- 1986AS1B Asahi et al, J. Phys. Soc. Jpn. Suppl. 55 (1986) 1032
- 1986AV1B Avdeichikov, in Dubna (1986) 122
- 1986BA04 M. Bawin and G.L. Strobel, Phys. Rev. C33 (1986) 732
- 1986BA1C Baer and Miller, Comments Nucl. and Part. Phys. 15 (1986) 269
- 1986BA2U Baker, Bull. Amer. Phys. Soc. 31 (1986) 1282
- 1986BA49 M. Bawin, Phys. Rev. C34 (1986) 1487
- 1986BA69 D. Baye, Nucl. Phys. A460 (1986) 581
- 1986BA88 W. Bauhoff, At. Data Nucl. Data Tables 35 (1986) 429
- 1986BA89 G. Baur, Phys. Lett. B178 (1986) 135
- 1986BE35 A.V. Belozyorov, C. Borcea, Z. Dlouhy, A.M. Kalinin, R. Kalpakchieva, Nguyen Hoai Chau, Yu.Ts. Oganessian and Yu.E. Penionzhkevich, Nucl. Phys. A460 (1986) 352
- 1986BE54 A.V. Belozorov, K. Borcha, Z. Dlouhy, A.M. Kalinin, Nguyen Hoai Tyau and Yu.E. Penionzhkevich, Pisma Zh. Eksp. Teor. Fiz. 44 (1986) 498; JETP Lett. (USSR) 44 (1986) 641
- 1986BI1A Bimbot et al, J. Phys. (France) 47 (1986) C4-241
- 1986BO1A Boikova et al, Sov. J. Nucl. Phys. 43 (1986) 173
- 1986CH05 C.Y. Cheung, B.D. Keister, Phys. Rev. C33 (1986) 776

1986CH1J Chant, AIP Conf. Proc. 142 (1986) 246
 1986CH2F Chardine, Haouat, Seguin and Humeau, CEA-N-2506 (1986)
 1986CH2G Chbihi et al, J. Physique 47 (1986) C4-87
 1986CL1B Clarke et al, J. Phys. Soc. Jpn. Suppl. 55 (1986) 756
 1986CO1F D.D. Cohen, A. Katsaros and S. Frisken, 11th Ains Nucl. Phys. Conf., Melbourne, Australia, 1986 (Australian Inst. Nucl. Sci. Eng., 1986) 16; Phys. Abs. 79276 (1986)
 1986CO1Q Court and Heyes, Nucl. Instr. Meth. Phys. Res. A243 (1986) 37
 1986CU01 M.S. Curtin, L.H. Harwood, J.A. Nolen, B. Sherrill, Z.Q. Xie and B.A. Brown, Phys. Rev. Lett. 56 (1986) 34
 1986CU02 B. Cujec, B. Dasmahapatra, Q. Haider, F. Lahlou and R.A. Dayras, Nucl. Phys. A453 (1986) 505
 1986DA1B Davis and Pniewski, Contemp. Phys. 27 (1986) 91
 1986DA1G Dalitz, Davis and Tovee, Nucl. Phys. A450 (1986) 311C
 1986DR03 P.V. Drumm, O. Karban, A.K. Basak, P.M. Lewis, S. Roman and G.C. Morrison, Nucl. Phys. A448 (1986) 93
 1986DU11 J.P. Dufour, R. Del Moral, H. Emmermann, F. Hubert, D. Jean, C. Poinot, M.S. Pravikoff, A. Fleury, H. Delagrang and K.-H. Schmidt, Nucl. Instrum. Meth. Phys. Res. A248 (1986) 267
 1986EN1A Engelmann and Bardy, Rep. CEA-R-5340 (1986)
 1986ER1A Eramzhyan, Ishkhanov, Kapitonov and Neudatchin, Phys. Rep. 136 (1986) 229
 1986EV01 M.V. Evlanov and A.M. Sokolov, Nucl. Phys. A452 (1986) 477
 1986FE1A Fetisov et al, Czech. J. Phys. 36 (1986) 451
 1986FR1G Franchi, Wright and Pillinger, Nature 323 (1986) 138
 1986GA1H Gal, AIP Conf. Proc. 150 (1986) 127
 1986GI1A Gilad, Bull. Amer. Phys. Soc. 31 (1986) 1203
 1986GL1E Glukhov et al, in Kharkov 86 (1986) 377, 378
 1986GR04 K. Grotz, H.V. Klapdor and J. Metzinger, Phys. Rev. C33 (1986) 1263
 1986GR1A Gregoire and Tamain, Ann. Physique 11 (1986) 323
 1986GU1D Guerreau, J. Phys. (France) 47 (1986) C4-207
 1986HA13 Q. Haider and F.B. Malik, J. Phys. G12 (1986) 537
 1986HA1B Harvey, J. Phys. 47 (1986) C4-29
 1986HA1E Harney, Richter and Weidenmuller, Rev. Mod. Phys. 58 (1986) 607
 1986HA1F Haas et al, in Harrogate (1986) C184

- 1986HA1P Hanna, J. Phys. Soc. Jpn. Suppl. 55 (1986) 528
- 1986HE1F Heaton et al, Nature 322 (1986) 822
- 1986JA1H Jacobs, AIP Conf. Proc. 142 (1986) 181
- 1986JE1B Jeong and Cheon, J. Korean Phys. Soc. 19 (1986) 157; Phys. Abs. 89541 (1987)
- 1986KA44 O. Karban, L. Potvin and I.M. Turkiewicz, Nucl. Phys. A460 (1986) 529
- 1986KE14 T.J. Kennett, W.V. Prestwich and J.S. Tsai, Nucl. Instrum. Meth. Phys. Res. A249 (1986) 366
- 1986KI12 N.S.P. King, P.W. Lisowski, G.L. Morgan, P.N. Craig, R.G. Jeppesen, D.A. Lind, J.R. Shepard, J.L. Ullmann, C.D. Zafiratos, C.D. Goodman et al, Phys. Lett. B175 (1986) 279
- 1986KO1A Kolesnikov et al, in Kharkov (1986) 225
- 1986KU1J Kume, J. Phys. Soc. Jpn. Suppl. 55 (1986) 920
- 1986LE01 M.J. Leitch, H.W. Baer, J.D. Bowman, M.D. Cooper, E. Piassetzky, U. Sennhauser, H.J. Ziock, F. Irom, P.B. Siegel and M.A. Moinester, Phys. Rev. C33 (1986) 278
- 1986LE1L Leavitt, Bull. Amer. Phys. Soc. 31 (1986)1319
- 1986LI1B Liu and Haider, AIP Conf. Proc. 150 (1986) 930
- 1986LI1M Li et al, J. Phys. Soc. Jpn. Suppl. 55 (1986) 564
- 1986LU1A Ludeking and Cotanch, AIP Conf. Proc. 150 (1986) 542
- 1986MA13 J.F. Mateja, A.D. Frawley, R.A. Parker and K. Sartor, Phys. Rev. C33 (1986) 1307
- 1986MA19 J.F. Mateja, A.D. Frawley, L.C. Dennis and K. Sartor, Phys. Rev. C33 (1986) 1649
- 1986MA1P Martin et al, Bull. Amer. Phys. Soc. 31 (1986) 1304
- 1986MC10 W.J. McDonald, R.N. MacDonald, W.C. Olsen, R. Dymarz, F. Khanna, L. Antonuk, J.M. Cameron, P. Kitching, G.C. Neilson, D.M. Sheppard et al, Nucl. Phys. A456 (1986) 577
- 1986MC13 J.A. McNeil, R.D. Amado, C.J. Horowitz, M. Oka, J.R. Shepard and D.A. Sparrow, Phys. Rev. C34 (1986) 746
- 1986ME06 M.C. Mermaz, T. Suomijarvi, R. Lucas, B. Berthier, J. Matuszek, J.P. Coffin, G. Guillaume, B. Heusch, F. Jundt and F. Rami, Nucl. Phys. A456 (1986) 186
- 1986MI25 B. Million, V. Bellini, L. Bimbot, M. Bolore, X. Charlot, J. Girard, J.M. Hisleur, J.C. Jourdain, J. Julien, A. Palmeri et al, Nucl. Phys. A459 (1986) 594
- 1986MO27 T. Motobayashi, H. Sakai, N. Matsuoka, T. Saito, K. Hosono, A. Okihana, M. Ishihara, S. Shimoura and A. Sakaguchi, Phys. Rev. C34 (1986) 2365
- 1986NO1C Nojiri et al, J. Phys. Soc. Jpn. Suppl. 55 (1986) 391
- 1986PA1C Papanicolas, AIP Conf. Proc. 142 (1986) 110

- 1986PO06 D.N. Poenaru, W. Greiner, K. Depta, M. Ivascu, D. Mazilu and A. Sandulescu, *At. Data Nucl. Data Tables* 34 (1986) 423
- 1986PO14 M. Potokar and A. Ramsak, *Phys. Rev. C* 34 (1986) 2338
- 1986PO15 D.N. Poenaru, W. Greiner, M. Ivascu, D. Mazilu and I.H. Plonski, *Z. Phys. A* 325 (1986) 435
- 1986SA2H Sawicki, Davies and Jackman, *Nucl. Instrum. Meth. Phys. Res. B* 15 (1986) 530
- 1986SA30 H. Sato and Y. Okuhara, *Phys. Rev. C* 34 (1986) 2171
- 1986SC23 W. Schott, W. Wagner, P. Kienle, R. Pollock, R. Bent, M. Fatyga, J. Kehayias, M. Green and K. Rehm, *Phys. Rev. C* 34 (1986) 1406
- 1986SC28 C.J.S. Scholz, L. Ricken and E. Kuhlmann, *Z. Phys. A* 325 (1986) 203
- 1986SI02 A.G. Sitenko, M.V. Evlanov, A.D. Polozov and A.M. Sokolov, *Yad. Fiz.* 43 (1986) 78; *Sov. J. Nucl. Phys.* 53 (1986) 50
- 1986SI11 E.R. Siciliano, M.D. Cooper, M.B. Johnson and M.J. Leitch, *Phys. Rev. C* 34 (1986) 267
- 1986SO10 L.G. Sobotka, D.G. Sarantites, H. Puchta, F.A. Dilmanian, M. Jaaskelainen, M.L. Halbert, J.H. Barker, J.R. Beene, R.L. Ferguson, D.C. Hensley et al, *Phys. Rev. C* 34 (1986) 917
- 1986ST1K Stevenson, Espe, Reiter and Lovett, *Nature* 323 (1986) 522
- 1986SU15 T.S. Subramanian, J.L. Romero, F.P. Brady, D.H. Fitzgerald, R. Garrett, G.A. Needham, J.L. Ullmann, J.W. Watson, C.I. Zanelli, D.J. Brenner et al, *Phys. Rev. C* 34 (1986) 1580
- 1986TA1E Taddeucci, *J. Phys. Soc. Jpn. Suppl.* 55 (1986) 156
- 1986TO10 L.W. Townsend, J.W. Wilson, F.A. Cucinotta and J.W. Norbury, *Phys. Rev. C* 34 (1986) 1491
- 1986UT01 H. Utsunomiya, E.C. Deci, R.A. Blue, L.H. Harwood, R.M. Ronningen, K. Siwek-Wilczynska, J. Wilczynski and D.J. Morrissey, *Phys. Rev. C* 33 (1986) 185
- 1986VA23 C.P.M. van Engelen, E.A. Bakkum, R.J. Meijer and R. Kamermans, *Nucl. Phys. A* 457 (1986) 375
- 1986VO1G Vogt, *Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., (1986) No. 68, Vol. 2, p. 23; Publ. by Inst. of Phys., Bristol, U.K.*
- 1986WI03 A.G. Williams and A.W. Thomas, *Phys. Rev. C* 33 (1986) 1070
- 1986WUZX Shi-Shu Wu and Xin-Yi Chen, *Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., (1986) 293*
- 1986YA1F Yamamoto, *Prog. Theor. Phys.* 75 (1986) 639
- 1987AB03 H. Abele, H.J. Hauser, A. Korber, W. Leitner, R. Neu, H. Plappert, T. Rohwer, G. Staudt, M. Strasser, S. Welte et al, *Z. Phys. A* 326 (1987) 373

- 1987AB04 K. Abe, K. Maeda, T. Ishimatsu, T. Kawamura, T. Furukawa, H. Orihara and H. Ohnuma, Nucl. Phys. A466 (1987) 109
- 1987AJ02 F. Ajzenberg-Selove, Nucl. Phys. A475 (1987) 1
- 1987ALZW W.P. Alford, R. Helmer, J.W. Watson, C. Zafiratos, R. Abegg, A. Celler, S. El-Kateb, D. Frekers, O. Hausser, R. Henderson et al, Bull. Amer. Phys. Soc. 32 (1987) 1578
- 1987AR1C Arnould, Phil. Trans. Roy. Soc. (London) 323 (1987) 251
- 1987AU1A Audouze, J. Astrophys. Astron. 8 (1987) 147
- 1987AZZZ S.M. Aziz, A.D. Bacher, L.C. Bland, G.T. Emery, W.W. Jacobs, E. Korkmaz, H. Nann, P.W. Park, J. Templon, P.L. Walden et al, Bull. Amer. Phys. Soc. 32 (1987) 1602
- 1987BA38 G.J. Balster, P.C.N. Crouzen, P.B. Goldhoorn, R.H. Siemssen and H.W. Wilschut, Nucl. Phys. A468 (1987) 93
- 1987BA89 Bahcall, Rev. Mod. Phys. 59 (1987) 505
- 1987BE1D Bertsch and Esbensen, Rep. Prog. Phys. 50 (1987) 607
- 1987BE1I M. Bedjidian, D. Contardo, E. Descroix, S. Gardien, J.Y. Grossiord, A. Guichard, M. Gusakow, R. Haroutunian, M. Jacquin, J.R. Pizzi et al, Z. Phys. A327 (1987) 337
- 1987BE25 I. Bergqvist, A. Brockstedt, L. Carlen, L.P. Ekstrom, B. Jakobsson, C. Ellegaard, C. Gaarde, J.S. Larsen, C. Goodman, M. Bedjidian et al, Nucl. Phys. A469 (1987) 648
- 1987BI1C Bimbot et al, in Panic (1987) 370
- 1987BU07 M. Burgel, H. Fuchs, H. Homeyer, G. Ingold, U. Jahnke and G. Thoma, Phys. Rev. C36 (1987) 90
- 1987CA15 J.R. Campbell, O.A. Abou-Zeid, W.R. Falk, R. Abegg, S.K. Datta and S.P. Kwan, Nucl. Phys. A467 (1987) 205
- 1987CH02 G. Chanfray and H.J. Pirner, Phys. Rev. C35 (1987) 760
- 1987DE03 B. Desplanques, Z. Phys. A326 (1987) 147
- 1987DE1Q de Vries, Ph. D. Thesis (1987)
- 1987DE43 H. De Vries, C.W. De Jager and C. De Vries, At. Data Nucl. Data Tables 36 (1987) 495
- 1987DO12 T.W. Donnelly, A.S. Raskin and J. Dubach, Nucl. Phys. A474 (1987) 307
- 1987EL1B Elevant and Andersson, Phys. Scr. T16 (1987) 148
- 1987FE1A Feng et al, Chin. Phys. 7 (1987) 121
- 1987FR1C Friedlander and Weneser, Science 235 (1987) 760
- 1987FU06 R.J. Furnstahl and B.D. Serot, Nucl. Phys. A468 (1987) 539
- 1987HI1B R.D. Hichwa, E.A. Hugel, J.J. Moskwa and R.R. Raylman, Nucl. Instrum. Meth. Phys. Res. B24-25 (1987) 932

- 1987HO1L Ho, Chin. Phys. Lett. 4 (1987) 69
- 1987HU01 R.L. Huffman, J. Dubach, R.S. Hicks and M.A. Plum, Phys. Rev. C35 (1987) 1
- 1987IC02 S. Ichii, W. Bentz and A. Arima, Nucl. Phys. A464 (1987) 575
- 1987IR01 F. Irom, H.W. Baer, A.G. Bergmann, J.D. Bowman, P. Heusi, D.H. Fitzgerald, C.J. Seftor, M.E. Sadler, J.N. Knudson, W.J. Briscoe et al, Phys. Rev. C36 (1987) 1453
- 1987KA09 T. Kajino, H. Toki and K.-I. Kubo, Phys. Rev. C35 (1987) 1370
- 1987KI1C Kissener, Rotter and Goncharova, Fortschr. Phys. 35 (1987) 277
- 1987KR04 D. Krolle and K. Langanke, Z. Phys. A327 (1987) 71
- 1987LA11 F.L. Lang, C.W. Werntz, C.J. Crannell, J.I. Trombka and C.C. Chang, Phys. Rev. C35 (1987) 1214
- 1987LE1B F. Lenz, Prog. Theor. Phys. Suppl. 91 (1987) 27
- 1987LE1J Letaw, Astrophys. J. 317 (1987) L69
- 1987LO1D Love et al, Can. J. Phys. 65 (1987) 536
- 1987MA1K Matthews, Bull. Amer. Phys. Soc. 32 (1987) 1575
- 1987MA2A Majling and Zofka, in Panic (1987) 596
- 1987ME1D Merchant, Rev. Bras. Fis. (1987) 182
- 1987MI27 W. Mittig, J.M. Chouvel, W.L. Zhan, L. Bianchi, A. Cunsolo, B. Fernandez, A. Foti, J. Gastebois, A. Gillibert, C. Gregoire et al, Phys. Rev. Lett. 59 (1987) 1889
- 1987MI38 Mian, Phys. Rev. C35 (1987) 1463
- 1987MIZY A. Middleton, J.D. Brown, L. Herold, K.E. Luther, M.L. Pitt, D. Barker, H.S. Camarda and S. Aziz, Bull. Amer. Phys. Soc. 32 (1987) 1578
- 1987MO03 R. Moreh and W.C. Sellyey, Phys. Lett. B185 (1987) 11
- 1987NA01 M.N. Namboodiri, R.K. Choudhury, L. Adler, J.D. Bronson, D. Fabris, U. Garg, P.L. Gonthier, K. Hagel, D.R. Haenni, Y.W. Lui et al, Phys. Rev. C35 (1987) 149
- 1987OL1A Olson et al, Bull. Amer. Phys. Soc. 32 (1987) 1015
- 1987OR01 H. Orihara, M. Kabasawa, K. Furukawa, T. Kawamura, Y. Takahashi, A. Satoh, T. Niizeki, T. Nakagawa, K. Maeda, K. Ishii et al, Phys. Lett. B187 (1987) 240
- 1987PO1H Povh, Prog. Part. Nucl. Phys. 18 (1987) 183
- 1987RA15 J. Rapaport, D. Wang, J.A. Carr, F. Petrovich, C.C. Foster, C.D. Goodman, C. Gaarde, J. Larsen, C.A. Goulding, T.N. Taddeucci et al, Phys. Rev. C36 (1987) 500
- 1987RA1D R. Ramaty and R.J. Murphy, Space Sci. Rev. 45 (1987) 213
- 1987RI03 J. Richert and P. Wagner, Nucl. Phys. A466 (1987) 132
- 1987RI1E Rich, Owen and Spiro, Phys. Rep. 151 (1987) 239

- 1987RY03 J. Ryckebusch, M. Waroquier, K. Heyde and D. Ryckbosch, Phys. Lett. B194 (1987) 453
- 1987SA01 C. Samanta, N.S. Chant, P.G. Roos, A. Nadasen and A.A. Cowley, Phys. Rev. C35 (1987) 333
- 1987SA15 H. Sagawa and H. Toki, J. Phys. G13 (1987) 453
- 1987SA25 M.G. Saint-Laurent, Nucl. Instrum. Meth. Phys. Res. B26 (1987) 273
- 1987SC11 L. Schmieder, D. Hilscher, H. Rossner, U. Jahnke, M. Lehmann, K. Ziegler and H.-H. Knitter, Nucl. Instrum. Meth. Phys. Res. A256 (1987) 457
- 1987SC1H Scfroder et al, Nucl. Phys. A467 (1987) 240
- 1987SI1D Simpson, Earwaker and Khan, Nucl. Instrum. Meth. Phys. Res. B24-25 (1987) 701
- 1987SN01 K. Sneppen, Nucl. Phys. A470 (1987) 213
- 1987ST01 G.S.F. Stephans, R.V.F. Janssens, D.G. Kovar and B.D. Wilkins, Phys. Rev. C35 (1987) 614
- 1987ST05 R.A. Stratton and R.A. Baldock, J. Phys. G13 (1987) 361
- 1987TA13 T.N. Taddeucci, C.A. Goulding, T.A. Carey, R.C. Byrd, C.D. Goodman, C. Gaarde, J. Larsen, D. Horen, J. Rapaport and E. Sugarbaker, Nucl. Phys. A469 (1993) 125
- 1987TA22 T.N. Taddeucci, Can. J. Phys. 65 (1987) 557
- 1987TU01 A. Turowiecki, A. Saganek, M. Sieminski, E. Wesolowski and Z. Wilhelmi, Nucl. Phys. A468 (1987) 29
- 1987VA26 A.G.M. van Hees, A.A. Wolters and P.W.M. Glaudemans, Phys. Lett. B196 (1987) 19
- 1987VD1A A.I. Vdovin, A.V. Golovin and I.I. Loschakov, Sov. J. Part. Nucl. 18 (1987) 573
- 1987VI02 F. Videbaek, S.G. Steadman, G.G. Batrouni and J. Karp, Phys. Rev. C35 (1987) 2333
- 1987VOZR B.B. Voitsekhovskiy, D.M. Nikolenko, S.G. Popov, I.A. Rachek, D.K. Toporkov, E.P. Tsentalovich, R. Vyunsh, V.F. Dmitriev and V.B. Telitsyn, Program and Theses, Proc. 37th Ann. Conf. Nucl. Spectrosc. Struct. At. Nuclei, Yurmala (1987) 362
- 1987WE1C Weneser and Friedlander, Science 235 (1987) 755
- 1987WU05 R. Wunsch and J. Zofka, Phys. Lett. B193 (1987) 7
- 1987ZI1C Zinner, Tang and Anders, Nature 330 (1987) 730
- 1988AD07 G.S. Adams, E.R. Kinney, J.L. Matthews, W.W. Sapp, T. Soos, R.O. Owens, R.S. Turley and G. Pignault, Phys. Rev. C38 (1988) 2771
- 1988AL1G Aleksandrov et al, in Baku (1988) 377
- 1988AN1E Antony and Britz, Indian J. Phys. A62 (1988) 411
- 1988AR1B Arima, Interactions and Structures in Nuclei, Proc. in Honor of D.H. Wilkinson, Sussex, 9/87, A. Hilger Publ. (1988) 15

- 1988AR1D Ardito et al, *Europhys. Lett.* 6 (1988) 131
- 1988AR1I A. Arima, *Hyperfine Interactions* 43 (1988) 47
- 1988ASZY K. Asahi, M. Ishihara, T. Shimoda, T. Fukuda, N. Takahashi, K. Katori, K. Hanakawa, T. Itahashi, Y. Nojiri, T. Minamisono et al, *AIP Conf. Proc.* 164 (1988) 165; *Proc. 5th Int. Conf. Nucl. Far from Stability, Rosseau Lake, Canada 1987*, Ed. I.S. Towner (1988) 165
- 1988AU03 F. Auger and B. Fernandez, *Nucl. Phys.* A481 (1988) 577
- 1988AZ1D Aziz, Ph. D. Thesis, Indiana Univ. (1988)
- 1988BA1Y Bahcall, Davis and Wolfenstein, *Nature* 334 (1988) 487
- 1988BA55 P.H. Barker and S.M. Ferguson, *Phys. Rev.* C38 (1988) 1936
- 1988BA86 Bahcall and Ulrich, *Rev. Mod. Phys.* 60 (1988) 297
- 1988BAYZ D. Bazin, R. Anne, D. Guerreau, D. Guillemaud-Mueller, A.C. Mueller, M.G. Saint-Laurent and W.D. Schmidt-Ott, *AIP Conf. Proc.* 164 (1988) 722; *Proc. 5th Int. Conf. Nuclei Far from Stability, Rosseau Lake, Canada 1987*
- 1988BE02 A.V. Belozyorov, C. Borcea, Z. Dlouhy, A.M. Kalinin, Nguyen Hoai Chau and Yu.E. Penionzhkevich, *Nucl. Phys.* A477 (1988) 131
- 1988BE56 A.V. Belozorov, K.C. Borcea, J. Wincour, M. Lewitowicz, N.H. Chau, Yu.E. Penionzhkevich, N.K. Skobelev and A. Chasha, *Izv. Akad. Nauk SSSR* 52 (1988) 2171; *Bull. Acad. Sci. USSR, Phys.Ser.* 52 (1988) 90
- 1988BI11 N. Bischof, W. Tiereth, I. Weitzenfelder, H. Voit, W. von Oertzen and H.H. Wolter, *Nucl. Phys.* A490 (1988) 485
- 1988BLZY S.L. Blatt, A. Abduljalil, D.G. Marchlenski, H.J. Hausman, J.D. Kalen, W. Kim, H.R. Weller and G. Feldman, *Bull. Amer. Phys. Soc.* 33 (1988) 1575
- 1988BR17 K. Brehm, H.W. Becker, C. Rolfs, H.P. Trautvetter, F. Kappeler and W. Ratynski, *Z. Phys.* A330 (1988) 167
- 1988BRZY J.D. Brown, A. Middleton and S.M. Aziz, *Bull. Amer. Phys. Soc.* 33 (1988) 1022
- 1988CA06 G. Caskey, L. Heilbronn, B. Remington, A. Galonsky, F. Deak, A. Kiss and Z. Seres, *Phys. Rev.* C37 (1988) 696
- 1988CA10 M. Cavinato, M. Marangoni and A.M. Saruis, *Phys. Rev.* C37 (1988) 1823
- 1988CA1J C. Carraro, A. Schafer and S.E. Koonin, *Astrophys. J.* 331 (1988) 565
- 1988CA26 G.R. Caughlan and W.A. Fowler, *At. Data Nucl. Data Tables* 40 (1988) 283
- 1988CH1T Chen, Yang and Wu, *High Energy Phys. Nucl. Phys.* 12 (1988) 822
- 1988CH49 R.E. Chrien, S. Bart, P. Pile, R. Sutter, N. Tsoupas, H.O. Funsten, J.M. Finn, C. Lyndon, V. Punjabi, C.F. Perdriat et al, *Phys. Rev. Lett.* 60 (1988) 2595

- 1988CIZZ D.E. Ciskowski, M.R. Barlett, T.A. Carey, J.B. McClelland, L.J. Rybarcyk, T.N. Taddeucci, E. Sugarbaker, D. Marchlenski, C.D. Goodman, W. Huang et al, Bull. Amer. Phys. Soc. 33 (1988) 1583
- 1988CO15 E. Comay, I. Kelson and A. Zidon, Phys. Lett. B210 (1988) 31
- 1988DA12 R. da Silveira and Ch. Leclercq-Willain, Phys. Rev. C38 (1988) 543
- 1988DE09 J.W. de Vries, D. Doornhof, C.W. de Jager, R.P. Singhal, S. Salem, G.A. Peterson and R.S. Hicks, Phys. Lett. B205 (1988) 22
- 1988DI02 S.S. Dietrich and B.L. Berman, At. Data Nucl. Data Tables 38 (1988) 199
- 1988DUZX E.I. Dubovoi, V.I. Pryazhinsky and G.I. Chitanava, Prog. and Theses, Proc. 38th Ann. Conf. Nucl. Spectrosc. Struct. At. Nucl., Baku (1988) 317
- 1988FE1A Ferrando et al, Phys. Rev. C37 (1988) 1490
- 1988FI1C H.J. Fischbeck, Bull. Amer. Phys. Soc. 33 (1988) 1691
- 1988FU04 R.J. Furnstahl, Phys. Rev. C38 (1988) 370
- 1988GO11 J. Gomez del Campo, J.L. Charvet, A. D'Onofrio, R.L. Auble, J.R. Beene, M.L. Halbert and H.J. Kim, Phys. Rev. Lett. 61 (1988) 290
- 1988GO1H Goryonov et al, in Baku (1988) 367
- 1988GOZM N.G. Goncharova and V.Ya. Spevak, Prog. and Theses, Proc. 38th Ann. Conf. Nucl. Spectrosc. Struct. At. Nucl., Baku (1988) 140
- 1988GR1A Grady and Pillinger, Nature 331 (1988) 321
- 1988HI1F R.D. Hichwa, Bull. Amer. Phys. Soc. 33 (1988) 1747
- 1988HO10 T. Hoshino, H. Sagawa and A. Arima, Nucl. Phys. A481 (1988) 458
- 1988HO15 J. Homolka, W. Schott, W. Wagner, W. Wilhelm, R.D. Bent, M. Fatyga, R.E. Pollock, M. Saber, R.E. Segel and P. Kienle, Phys. Rev. C38 (1988) 2686
- 1988JO1B G. A. Jones, Interactions and Structures in Nuclei, Proc. in Honor of D.H. Wilkinson, Sussex, 9/87, A. Hilger Publ. (1988) 9
- 1988KE1B Kessler et al, J. Phys. G14 (1988) S167
- 1988KO18 L. Koester, W. Waschkowski, J. Meier, G. Rau and M. Salehi, Z. Phys. A330 (1988) 387
- 1988KR11 L. Kraus, A. Boucenna, I. Linck, B. Lott, R. Rebmeister, N. Schulz, J.C. Sens, M.C. Mermaz, B. Berthier, R. Lucas et al, Phys. Rev. C37 (1988) 2529
- 1988KR1G Krombel and Wiedenbeck, Astrophys. J. 328 (1988) 940
- 1988KW02 E. Kwasniewicz and J. Kisiel, Acta Phys. Pol. B19 (1988) 141
- 1988LE08 K.T. Lesko, E.B. Norman, R.-M. Larimer, S. Kuhn, D.M. Meekhof, S.G. Crane and H.G. Bussell, Phys. Rev. C37 (1988) 1808

- 1988LEZW M.B. Leuschner, F.W. Hersman, J.R. Calarco, J.E. Wise, L. Lapikas, P.K.A. de Witt-Huberts, E. Jans, G. Kramer and H.P. Blok, *Bull. Amer. Phys. Soc.* 33 (1988) 1097
- 1988LI1E Liobarsky, Melenevsky, Nemykin and Shimanov, *Baku* (1988) 566
- 1988LI23 A. Liesenfeld, B. Alberti, D. Eyl, G. Kobschall, A.W. Richter, K. Rohrich, Ch. Schmitt, L. Tiator and V.H. Walther, *Nucl. Phys.* A485 (1988) 580
- 1988LO07 G.M. Lotz and H.S. Sherif, *Phys. Lett.* B210 (1988) 45
- 1988LU02 K.E. Luther, J.D. Brown and R.T. Kouzes, *Phys. Rev.* C38 (1988) 529
- 1988MA07 J.F. Mateja, G.L. Gentry, N.R. Fletcher, L.C. Dennis and A.D. Frawley, *Phys. Rev.* C37 (1988) 1004
- 1988MA1G Majling et al, *Phys. Lett.* B202 (1988) 489
- 1988MA1H Manokhin, INDC(CCP)-283 (1988)
- 1988MC01 K.G. McNeill, A.D. Bates, R.P. Rassool, E.A. Milne and M.N. Thompson, *Phys. Rev.* C37 (1988) 1403
- 1988MC03 J.P. McDermott, E. Rost, J.R. Shepard and C.Y. Cheung, *Phys. Rev. Lett.* 61 (1988) 814
- 1988MCZT V. McLane, C.L. Dunford and P.F. Rose, *Neutron Cross Sections*, Vol. 2 (Academic Press, New York, 1988)
- 1988MI1G D. Mikolas, B.A. Brown, W. Benenson, Y. Chen, M.S. Curtin, L.H. Harwood, E. Kashy, J.A. Nolen, Jr., M. Samuel, B. Sherrill et al, *AIP Conf. Proc.* 164 (1988) 708
- 1988MI1J D.J. Millener, *AIP Conf. Proc.* 163 (1988) 402
- 1988MI1K Miller, *AIP Conf. Proc.* 163 (198) 438
- 1988MI28 M. Mishra, M. Satpathy and L. Satpathy, *J. Phys.* G14 (1988) 1115
- 1988MO1L Motoba, Itonaga and Bando, *Nucl. Phys.* A489 (1988) 683
- 1988MU08 A.C. Mueller, D. Bazin, W.D. Schmidt-Ott, R. Anne, D. Guerreau, D. Guillemaud-Mueller, M.G. Saint-Laurent, V. Borrel, J.D. Jacmart, F. Pougheon et al, *Z. Phys.* A330 (1988) 63
- 1988MUZV P.E. Mueller, C.N. Papanicolas, S.E. Williamson, H. Rothhaas and R. Nick, *Bull. Amer. Phys. Soc.* 33 (1988) 1594
- 1988NI05 S. Nishizaki, H. Kurasawa and T. Suzuki, *Phys. Lett.* B209 (1988) 6
- 1988PA1S Papanicolas et al, *AIP Conf. Proc.* 176 (1988) 464
- 1988PI12 D. Pietersen and W.J. Strydom, *Nucl. Instrum. Meth. Phys. Res.* B35 (1988) 467
- 1988PI1C C.T. Pillinger, *Phil. Trans. Roy. Soc. London* A325 (1988) 525
- 1988POZS N.A.F.M. Poppelier, J.H. de Vries, A.A. Wolters and P.W.M. Glaudemans, in *AIP Conf. Proc.* 164 (1988) 334
- 1988PR1D Price et al, *Bull. Amer. Phys. Soc.* 33 (1988) 1781

1988RO17 M. Roy-Stephan, Nucl. Phys. A488 (1988) 187C
 1988RO1M Roos, in AIP Conf. Proc. 163 (1988) 210
 1988RU01 V.A. Rubchenya and S.G. Yavshits, Z. Phys. A329 (1988) 217
 1988RY03 J. Ryckebusch, M. Waroquier, K. Heyde, J. Moreau and D. Ryckbosch, Nucl. Phys. A476 (1988) 237
 1988SA04 M. Samuel, B.A. Brown, D. Mikolas, J. Nolen, B. Sherrill, J. Stevenson, J.S. Winfield and Z.Q. Xie, Phys. Rev. C37 (1988) 1314
 1988SA19 H. Sato, Phys. Rev. C37 (1988) 2902
 1988SE11 N. Severijns, J. Wouters, j. Vanhaverbeke, W. Vanderpoorten and L. Vanneste, Hyperfine Interact. 34 (1988) 415
 1988SH07 J.R. Shepard, E. Rost, C.-Y. Cheung and J.A. McNeil, Phys. Rev. C37 (1988) 1130
 1988SN1A K.A. Snover, Nucl. Phys. A482 (1988) 13C
 1988SUZY A. Suhaimi, JUL-2196 (1988)
 1988TA09 S. Takeuchi, K. Shimizu and K. Yazaki, Nucl. Phys. A481 (1988) 693
 1988TA10 I. Tanihata, T. Kobayashi, O. Yamakawa, S. Shimoura, K. Ekuni, K. Sugimoto, N. Takahashi, T. Shimoda and H. Sato, Phys. Lett. B206 (1988) 592
 1988TA21 T. Takaki and M. Thies, Phys. Rev. C38 (1988) 2230
 1988TO1C Towner, in AIP Conf. Proc. 164 (1988) 593
 1988UT02 H. Utsunomiya and R.P. Schmitt, Nucl. Phys. A487 (1988) 162
 1988VA03 A.G.M. van Hees, A.A. Wolters and P.W.M. Glaudemans, Nucl. Phys. A476 (1988) 61
 1988VI1A Vinogradova et al, Baku (1988) 567
 1988VO1D J.R. Votaw, Bull. Amer. Phys. Soc. 33 (1988) 1748
 1988WA08 F. Wang, C.W. Wong and S.-Q. Lu, Nucl. Phys. A480 (1988) 490
 1988WA18 A.H. Wapstra, G. Audi and R. Hoekstra, At. Data Nucl. Data Tables 39 (1988) 281
 1988WA1E E.K. Warburton, in Brighton 87 (1988) 81
 1988WA1I Wanke and Dreibus, Phil. Trans. Roy. Soc. London A325 (1988) 545
 1988WA1Q Watson, Nucl. Instrum. Meth. Phys. Res. B40-41 (1988) 481
 1988WAZY T.R. Wang, R.B. Vogelaar and R.W. Kavanagh, Bull. Amer. Phys. Soc. 33 (1988) 1563
 1988WO04 A.A. Wolters, A.G.M. van Hees and P.W.M. Glaudemans, Europhys. Lett. 5 (1988) 7
 1988YA08 Yanhe Jin and D.S. Onley, Phys. Rev. C38 (1988) 813
 1988YOZX J.C. Young, F.P. Brady, J.L. Romero, G.A. Needham and J.L. Ullmann, Bull. Amer. Phys. Soc. 33 (1988) 1568

- 1988ZH1B Zhusupov and Usmanov, in Baku (1988) 167
- 1989AB01 R. Abegg, D.A. Hutcheon, C.A. Miller, L. Antonuk, J.M. Cameron, G. Gaillard, J.M. Greben, P. Kitching, R.P. Liljestr and, W.J. McDonald et al, Phys. Rev. C39 (1989) 65
- 1989AL04 W.P. Alford, R. Helmer, R. Abegg, A. Celler, D. Frekers, O. Hausser, R. Henderson, K. Hicks, K.P. Jackson, R. Jeppesen et al, Phys. Rev. C39 (1989) 1189
- 1989AM01 K. Amos, D. Koetsier and D. Kurath, Phys. Rev. C40 (1989) 374
- 1989AR1J M. Arnould, F. Baeten, D. Darquennes, Th. Delbar, C. Dom, M. Huysse, Y. Jongen, P. Leleux, M. Lacroix, P. Lipnik et al, Nucl. Instrum. Meth. Phys. Res. B40-41 (1989) 498
- 1989AS1B Asahi et al, Tokyo (1988) 173
- 1989ASZZ R.M. Asherova, Yu.F. Smirnov and D.V. Fursa, in Tashkent (1989) 158
- 1989AYZU J. Aysto and J. Cerny, Treatise on Heavy-Ion Sci.8 (1989) 207
- 1989BA25 A.D. Bates, R.P. Rassool, E.A. Milne, M.N. Thompson and K.G. McNeill, Phys. Rev. C40 (1989) 506
- 1989BA2P J.N. Bahcall, Neutrino Astrophys. (Publ. Cambridge Univ. Press 1989)
- 1989BA92 H. Bando, Nuovo Cim. A102 (1989) 627
- 1989BA93 H. Bando, M. Sano, J. Zoofka and M. Wakai, Nucl. Phys. A501 (1989) 900
- 1989BE14 E.J. Beise, G. Dodson, M. Garcon, S. Hoibraten, C. Maher, D.L. Pham, R.P. Redwine, W. Sapp, K.E. Wilson, S.A. Wood and M. Deady, Phys. Rev. Lett. 62 (1989) 2593
- 1989BEZC J.A. Behr, K.A. Snover, C.A. Gossett, J.H. Gundlach, W. Hering, Bull. Amer. Phys. Soc. 34 (1989) 1832
- 1989BR1J Brown, in Sao Paulo (1989) 187
- 1989CA25 W.N. Catford, L.K. Fifield, N.A. Orr and C.L. Woods, Nucl. Phys. A503 (1989) 263
- 1989CH1X Chen and Li, Astrophys. Space Sci. 158 (1989) 153
- 1989CH24 M. Chiapparini and A.O. Gattone, Phys. Lett. 224B (1989) 243
- 1989CH2E Cheng, Proc. Specialists Mtg. Neutron Activation Cross Sections for Fission and Fusion Energy Applications, Eds. Wagner and Vonach (Argonne 1989) p. 29
- 1989CH31 A.A. Chumbalov, R.A. Eramzhyan and S.S. Kamalov, Czech. J. Phys. B39 (1989) 853
- 1989DA1H Davis, Mann and Wolfenstein, Ann. Rev. Nucl. Part. Sci. 39 (1989) 467
- 1989DE52 C. Detraz and D.J. Vieira, Ann. Rev. Nucl. Part. Sci. 39 (1989) 407
- 1989DO1I C.B. Dover, D.J. Millener and A. Gal, Phys. Rept. 184 (1989) 1
- 1989DO1K Dover, Millener and Gal, Phys. Rept. 184 (1989) 1

- 1989DR03 P.V. Drumm, L.K. Fifield, R.A. Bark, M.A.C. Hotchkis and C.L. Woods, Nucl. Phys. A496 (1989) 530
- 1989FU05 R.J. Furnstahl and C.E. Price, Phys. Rev. C40 (1989) 1398
- 1989FU1J Fukahori, JAERI-M 89-047 (1989)
- 1989GE10 P.M. Gensini, Nuovo Cim. A102 (1989) 1563
- 1989GE11 C.K. Gelbke, Nucl. Phys. A495 (1989) C27
- 1989GOZQ N.G. Goncharova, Prog. and Thesis, Proc. 39th Ann. Conf. Nucl. Spectrosc. Struct. At. Nucl., Tashkent (1989) 154
- 1989GRZQ W. Greiner, M. Ivascu, D.N. Poenaru and A. Sandulescu, Treatise on Heavy-Ion Sci. 8 (1989) 641; Ed. Bromlev, published by Plenum Publ. Corp. 1989
- 1989GU1J N. Guessoum, Astrophys. J. 345 (1989) 363
- 1989GU1L Gustafsson, Ann. Astron. Astrophys. 27 (1989) 701
- 1989GU1Q Gupta and Webber, Astrophys. J. 340 (1989) 1124
- 1989GU28 N. Guessoum and R.J. Gould, Astrophys. J. 345 (1989) 356
- 1989HE04 R. Heaton, H. Lee, P. Skensved and B.C. Robertson, Nucl. Instrum. Meth. Phys. Res. A276 (1989) 529
- 1989JI1A L. Jin, W.D. Arnett and S.K. Chakrabarti, Astrophys. J. 336 (1989) 572
- 1989JO07 M.B. Johnson, Czech. J. Phys. B39 (1989) 822
- 1989KA1K Kajino, Mathews and Fuller, in Tokyo 1988 (1989) 51
- 1989KA1N Kayomov, Mukhamedzhanov and Yarmukhamedov, Tashkent (1989) 408
- 1989KA1S Katoh, Kawade and Yamamoto, JAERI-M89-083 (1989)
- 1989KA2B Kawade et al, Proc. Specialists Mtg. Neutron Activation Cross Sections for Fission and Fusion Energy Applications, Eds. Wagner and Vonach (Argonne, 1989) p.99
- 1989KE1D Kerridge, Science 45 (1989) 480
- 1989KEZZ S.E. Kellogg, R.B. Vogelaar and R.W. Kavanagh, Bull. Amer. Phys. Soc. 34 (1989) 1192
- 1989KI09 J.C. Kim, G. Kajrys and W. Del Bianco, Can. J. Phys. 67 (1989) 135
- 1989KI13 A. Kiss, F. Deák, Z. Seres, G. Caskey, A. Galonsky, B. Remington and L. Heilbronn, Nucl. Phys. A499 (1989) 131
- 1989KO11 P.E. Koehler and H.A. O'Brien, Phys. Rev. C39 (1989) 1655
- 1989KO1H Kouthoulos and Grypeos, Phys. Rev. C40 (1989) 275
- 1989KO28 T. Kobayashi, K. Takeshita, A. Kagaya, H. Tsubota and K. Shoda, J. Phys. Soc. Jpn. 58 (1989) 1570
- 1989KU1P Kudziev, Nucl. Instrum. Meth. Phys. Res. A282 (1989) 267

- 1989LE16 M. Lewitowicz, Yu.E. Penionzhkevich, A.G. Artukh, A.M. Kalinin, V.V. Kamanin, S.M. Lukyanov, Nguyen Hoai Chau, A.C. Mueller, D. Guillemaud-Mueller, R. Anne et al, Nucl. Phys. A496 (1989) 477
- 1989LE1L Leitch, "Fund. Symm. and Nucl. Struct.", Eds. Ginocchio and Rosen, in Santa Fe, NM 1988 (World Scientific: 1989) 163
- 1989LEZY M. Leuschner, F.W. Hersman, J.R. Calarco, J.E. Wise, L. Lapikas, P.K.A. de Witt-Huberts, E. Jans, G. Kramer and H.P. Blok, Bull. Amer. Phys. Soc. 34 (1989) 1153
- 1989LI26 A. Li, H.G. Pfutzner, C.R. Howell and R.L. Walter, Chin. J. Nucl. Phys. 11 (1989) 1
- 1989ME1C Mewaldt and Stone, Astrophys. J. 337 (1989) 959
- 1989MI30 M. Mian, Phys. Rev. C39 (1989) 279
- 1989NE02 Y. Nedjadi and J.R. Rook, J. Phys. G15 (1989) 589
- 1989NO1A Nomoto, Hashimoto, Arai and Kaminisi, Proc. int. Symp. on Heavy Ion Phys. and Nucl. Astrophys. Problems, Tokyo, 21-23 July 1988, ed. S. Kubono, M. Ishihara, T. Nomura (World Scientific, 1989) 9
- 1989NW1A Nwosu and Fischbeck, Nucl. Instrum. Meth. Phys. Res. B40-41 (1989) 833
- 1989OG1B A.A. Oglobin and Y.E. Penionzhkevich, Treatise On Heavy-Ion Science, Vol. 8, Ed. D.A. Bromley (Plenum Publ. Corp., 1989) p.261
- 1989OR02 W.E. Ormand and B.A. Brown, Nucl. Phys. A491 (1989) 1
- 1989PA06 D.J. Parker, J.J. Hogan and J. Asher, Phys. Rev. C39 (1989) 2256
- 1989PO07 J. Pouliot, Y. Chan, A. Dacal, D.E. DiGregorio, B.A. Harmon, R. Knop, M.E. Ortiz, E. Plagnol, R.G. Stokstad, C. Moisan et al, Phys. Lett. B223 (1989) 16
- 1989PO1K Poppelier, Ph.D. Thesis, Univ. of Utrecht (1989)
- 1989RA09 J. Rapaport, P.W. Lisowski, J.L. Ullmann, R.C. Byrd, T.A. Carey, J.B. McClelland, L.J. Rybarcyk, T.N. Taddeucci, R.C. Haight, N.S.P. King et al, Phys. Rev. C39 (1989) 1929
- 1989RA15 L. Ray and J.R. Shepard, Phys. Rev. C40 (1989) 237
- 1989RA17 P. Raghavan, At. Data Nucl. Data Tables 42 (1989) 189
- 1989RY01 J. Ryckebusch, K. Heyde, D. Van Neck and M. Waroquier, Phys. Lett. B216 (1989) 252
- 1989SA10 M.G. Saint-Laurent, R. Anne, D. Bazin, D. Guillemaud-Mueller, U. Jahnke, Jin Gen-Ming, A.C. Mueller, J.F. Bruandet, F. Glasser, S. Kox et al, Z. Phys. A332 (1989) 457
- 1989SE07 N. Severijns, J. Wouters, J. Vanhaverbeke and L. Vanneste, Phys. Rev. Lett. 63 (1989) 1050
- 1989SI02 P.J. Simmonds, N.M. Clarke, K.I. Pearce, R.J. Griffiths, B. Stanley, S. Roman, A. Farooq, G. Rai, M.C. Mannion and C.A. Ogilvie, J. Phys. G15 (1989) 353

- 1989SI26 A.K. Sinha, D.M. Nadkarni and G.K. Mehta, *Pramana* 33 (1989) 85
- 1989ST14 A. Staudt, E. Bender, K. Muto and H.V. Klapdor, *Z. Phys.* A334 (1989) 47
- 1989STZW P.A. Staples, J.J. Egan, G.H.R. Kegel, A. Mittler and D.J. Desimone, *Bull. Amer. Phys. Soc.* 34 (1989) 1831
- 1989SU1J Sugarbaker and Marchlenski, *Wein* 89 (1989) Paper PB09
- 1989TA17 H. Tamura, T. Yamazaki, M. Sano, Y. Yamamoto, M. Wakai and H. Bando, *Phys. Rev.* C40 (1989) R483
- 1989TA10 I. Tanihata, *Treatise On Heavy-Ion Science*, Vol. 8, Ed. D.A. Bromley, published by Plenum Publ. Corp. (1989) 443
- 1989TA1Y Tanaka et al, *Nature* 341 (1989) 727
- 1989TA21 R. Tacik, E.T. Boschitz, R. Meier, S. Ritt, M. Wessler, K. Junker, J.A. Konter, S. Mango, D. Renker, B. van den Brandt et al, *Phys. Rev. Lett.* 63 (1989) 1784
- 1989TE02 F. Terrasi, A. Brondi, G. La Rana, G. De Angelis, A. D'Onofrio, R. Moro, E. Perillo and M. Romano, *Phys. Rev.* C40 (1989) 742
- 1989TH1C Thielemann and Wiescher, *Proc. int. Symp. on Heavy Ion Phys. and Nucl. Astrophys. Problems*, Tokyo, 21-23 July 1988, ed. S. Kubono, M. Ishihara, T. Nomura (World Scientific, 1989) 27
- 1989VA07 O.I. Vasileva, G.S. Gurevich, A.V. Ignatenko, V.M. Lebedev, N.V. Orlova, A.V. Spassky, I.B. Teplov, G.V. Shakhvorostova and I.K. Shestakova, *Yad. Fiz.* 49 (1989) 625; *Sov. J. Nucl. Phys.* 49 (1989) 387
- 1989VIIE Vinogradova et al, *Tashkent* (1989) 556
- 1989VO1D von Oertzen, *Tokyo* (1988) 373
- 1989WA16 S. Wa-Kitwanga, P. Leleux, P. Lipnik and J. Vanhorenbeeck, *Phys. Rev.* C40 (1989) 35
- 1989WO1B Wood, *Bull. Amer. Phys. Soc.* 34 (1989) 1133
- 1989WO1C Wolfs et al, *Wein* 89 (1989) Paper PD02
- 1989WO1E Wolters, *Ph.D. Thesis*, Univ. of Utrecht (1989)
- 1989WU1C Wu, Yang and Li, *High Energy Phys. Nucl. Phys.* 13 (1989) 75
- 1989WY1A Wyckoff et al, *Astrophys. J.* 339 (1989) 488
- 1989YO02 A. Yokoyama, T. Saito, H. Baba, K. Hata, Y. Nagame, S. Ichikawa, S. Baba, A. Shinohara and N. Imanishi, *Z. Phys.* A332 (1989) 71
- 1989YO1H Yoshida et al, *Nature* 342 (1989) 895
- 1990AB1G F. Abel, G. Amsel, E. d'Artemare, C. Ortega, J. Siejka and G. Vizkelethy, *Nucl. Instrum. Meth. Phys. Res.* B45 (1990) 100

- 1990AM1F G. Amsel, M. Menu, J. Moulin and J. Salomon, Nucl. Instrum. Meth. Phys. Res. B45 (1990) 296
- 1990AZZZ S.M. Aziz, K.W. Kemper and E. Reber, Bull. Amer. Phys. Soc. 35 (1990) 1059
- 1990BL09 P.G. Blunden and C.J. Horowitz, Phys. Lett. B240 (1990) 6
- 1990BRZY T.B. Bright and S.R. Cotanch, Bull. Amer. Phys. Soc. 35 (1990) 927
- 1990DA03 B. Dasmahapatra, B. Cujec, F. Lahlou, I.M. Szoghy, S.C. Gujrathi, G. Kajrys and J.A. Cameron, Nucl. Phys. A509 (1990) 393
- 1990DU01 O. Dumitrescu, M. Horoi, F. Carstoiu and G. Stratan, Phys. Rev. C41 (1990) 1462
- 1990ER03 R.A. Eramzhyan, M. Gmitro and S.S. Kamalov, Phys. Rev. C41 (1990) 2865
- 1990FL1A Fleischhauer and Scheid, Nucl. Phys. A510 (1990) 817
- 1990FU04 R.J. Furnstahl and C.E. Price, Phys. Rev. C41 (1990) 1792
- 1990GL01 A. Glaesner, W. Dunnweber, M. Bantel, W. Hering, D. Konnerth, R. Ritzka, W. Trautmann, W. Trombik and W. Zipper, Nucl. Phys. A509 (1990) 331
- 1990GO25 J. Gorres, S. Graff, M. Wiescher, R.E. Azuma, C.A. Barnes, H.W. Becker and T.R. Wang, Nucl. Phys. A517 (1990) 329
- 1990HA1W Haxton, Panic XII (1990) Paper XI-2
- 1990HA46 D. Harley, B. Muller and J. Rafelski, J. Phys. G16 (1990) 281
- 1990IE01 K. Ieki, J. Iimura, M. Iwase, H. Ohnuma, H. Shimizu, H. Toyokawa, K. Furukawa, H. Kabasawa, T. Nakagawa, T. Tohei et al., Phys. Rev. C42 (1990) 457
- 1990ISO5 M.A. Islam, T.J. Kennett and W.V. Prestwich, Nucl. Instrum. Meth. Phys. Res. A287 (1990) 460
- 1990KU1H Kume and Nose, Int. Conf. on Particles and Nucl., Cambridge, Mass., 25-29 June 1990 (Organizing Committee, 1990) Paper III-81, 82
- 1990LE1P Leuschner et al., Panic XII (1990) Paper I-80
- 1990LO10 R.J. Lombard, Europhys. Lett. 12 (1990) 119
- 1990MA1P Malaney, Meyer and Butler, Astrophys. J. 352 (1990) 767
- 1990MC04 W.R. McMurray, K. Bharuth-Ram and D.G. Aschman, Nucl. Instrum. Meth. Phys. Res. A288 (1990) 421
- 1990MI15 I.V. Mitchell, G.R. Massoumi, W.N. Lennard, S.Y. Tong, P.F.A. Alkemade, K. Griffiths, S.J. Bushby and P.R. Norton, Nucl. Instrum. Meth. Phys. Res. B45 (1990) 107
- 1990MU10 A.M. Mukhamedzhanov and N.K. Timofeyuk, Yad. Fiz. 51 (1990) 679; Sov. J. Nucl. Phys. 51 (1990) 431
- 1990NA15 T. Nakano, M. Nakamura, H. Sakaguchi, M. Yosoi, M. Ieiri, H. Togawa, S. Hirata, O. Kamigaito, H.M. Shimizu, M. Iwaki et al., Phys. Lett. B240 (1990) 301
- 1990OD1A Odagawa, Sato and Ohtsubo, Panic XII (1990) Paper I-62

- 1990OWZZ R.O. Owens, J.L. Matthews and G.S. Adams, Bull. Amer. Phys. Soc. 35 (1990) 928
- 1990PA1H Papanicolas, Bull. Amer. Phys. Soc. 35 (1990) 994
- 1990PI05 S. Piskor and W. Schaferlingova, Nucl. Phys. A510 (1990) 301
- 1990RA1O Ramaty, in Sao Paulo (1990) 763
- 1990RE04 Z. Ren and G.Xu, Phys. Lett. B237 (1990) 1
- 1990SA27 S.K. Saha, W.W. Daehnick, S.A. Dytman, P.C. Li, J.G. Hardie, G.P.A. Berg, C.C. Foster, W.P. Jones, D.W. Miller and E.J. Stephenson, Phys. Rev. C42 (1990) 922
- 1990ST08 A. Staudt, E. Bender, K. Muto and H.V. Klapdor-Kleingrothaus, At. Data Nucl. Data Tables 44 (1990) 79
- 1990TA1K Tacik, Panic XII (1990) Paper III-8
- 1990TA1L Tacik et al., Panic XII (1990) Paper III-67
- 1990TAZS Tacik, Bull. Amer. Phys. Soc. 35 (1990) 994
- 1990TO10 S.Y. Tong, W.N. Lennard, P.F.A. Alkemade and I.V. Mitchell, Nucl. Instrum. Meth. Phys. Res. B45 (1990) 91
- 1990VA01 A.G.M. Van Hees, J.G.L. Booten and P.W.M. Glaudemans, Nucl. Phys. A507 (1990) 55c
- 1990VA1G van Dyck, Bull. Amer. Phys. Soc. 35 (1990) 944
- 1990VO1E von Oertzen, Nucl. Instrum. Meth. Phys. Res. A287 (1990) 188
- 1990WA22 A.H. Wapstra, Nucl. Instrum. Meth. Phys. Res. A292 (1990) 671
- 1990WE14 W.R. Webber, J.C. Kish and D.A. Schrier, Phys. Rev C41 (1990) 520