

Energy Levels of Light Nuclei $A = 16$

F. Ajzenberg-Selove^a and T. Lauritsen^b

^a *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

^b *California Institute of Technology, Pasadena, California*

Abstract: An evaluation of $A = 5-24$ was published in *Nuclear Physics* 11 (1959), p. 1. This version of $A = 16$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 1, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 16$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{16}\text{N}\$](#) , [\$^{16}\text{O}\$](#) , [\$^{16}\text{F}\$](#)

B. Tables of Recommended Level Energies:

[Table 16.1](#): Energy levels of ^{16}N

[Table 16.3](#): Energy levels of ^{16}O

C. [References](#)

D. Figures: [\$^{16}\text{N}\$](#) , [\$^{16}\text{O}\$](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

^{16}N
(Fig. 31)

GENERAL:

Theory: See (1954FL1A, 1956KA1C, 1957EL1B).

1. $^{16}\text{N}(\beta^-)^{16}\text{O}$ $Q_m = 10.402$

From the character of the beta decay, it is concluded that $^{16}\text{N}(0)$ has $J^\pi = 2^-$. See ^{16}O .

2. $^7\text{Li}(^{14}\text{N}, ^5\text{Li})^{16}\text{N}$ $Q_m = 0.589$

See (1958AL1D).

3. $^9\text{Be}(^7\text{Li}, ^8\text{Li})^8\text{Be}$ $Q_m = 0.368$ $E_b = 20.585$

The cross section for this reaction has been measured from $E(^7\text{Li}) = 1.1$ to 2.0 MeV. At the higher energy it is approximately 1 mb (1957NO17).

4. $^{13}\text{C}(\alpha, p)^{16}\text{N}$ $Q_m = -7.416$

Not reported.

5. $^{14}\text{C}(\text{d}, \gamma)^{16}\text{N}$ $Q_m = 10.481$

At $E_d = 1.16$ MeV, the cross section is $\lesssim 0.1$ mb (1957BO04). At $E_d = 2.0$ MeV, $\sigma \leq 0.6$ mb (1956DO37).

6. $^{14}\text{C}(\text{d}, \text{n})^{15}\text{N}$ $Q_m = 7.987$ $E_b = 10.481$

See (1950HU72) and ^{15}N .

Table 16.1: Energy levels of ^{16}N

E_x (MeV \pm keV)	J^π	$\tau_{1/2}$ or Γ (keV)	Decay	Reactions
0	2^-	$\tau_{1/2} = 7.37 \pm 0.04$ sec	β^-	1, 2, 5, 12, 13, 16, 19, 24
0.120 ± 1	0^-	$\tau_{1/2} = 5.7$ μsec	γ	16, 24
0.295 ± 4	3^-		γ	16, 24, 25
0.392 ± 3	1^-		γ	16, 24, 25
(3.53 ± 30)		sharp		16
3.980 ± 20		sharp		16
4.80 ± 50		230 ± 40		16
(5.01 ± 50)				16
5.25 ± 50	$2^-, 3^\pm$	200	n	14, 16
12.2		270	d, p	7
12.62		190	d, p	7
12.8		165	d, p	7

7. $^{14}\text{C}(\text{d}, \text{p})^{15}\text{C}$

$$Q_m = -1.007$$

$$E_b = 10.481$$

The excitation function has been studied from $E_d = 0.6$ to 3.0 MeV (1954RI1B, 1956DO37). Resonances are observed at 2.0 , 2.45 and 2.7 MeV, with c.m. widths of 270 , 190 and 165 keV, respectively, corresponding to $^{16}\text{N}^*(12.2, 12.62, 12.8$ MeV) (1956DO37). See also (1950HU72, 1954RI1B) and ^{15}C .

8. $^{14}\text{C}(\text{d}, \alpha)^{12}\text{B}$

$$Q_m = 0.358$$

$$E_b = 10.481$$

See (1950HU72).

9. $^{14}\text{C}(\text{t}, \text{n})^{16}\text{N}$

$$Q_m = 4.223$$

Not reported.

10. $^{14}\text{C}(\text{}^3\text{He}, \text{p})^{16}\text{N}$

$$Q_m = 4.988$$

Not reported.

11. $^{14}\text{C}(\alpha, \text{d})^{16}\text{N}$ $Q_{\text{m}} = -13.363$

Not reported.

12. $^{14}\text{N}(\text{t}, \text{p})^{16}\text{N}$ $Q_{\text{m}} = 4.851$

See (1953CU1D).

13. $^{15}\text{N}(\text{n}, \gamma)^{16}\text{N}$ $Q_{\text{m}} = 2.494$

The thermal cross section is $24 \pm 8 \mu\text{b}$ (1958HU18), $80 \pm 30 \mu\text{b}$ (1952FE1A).

14. $^{15}\text{N}(\text{n}, \text{n})^{15}\text{N}$ $E_{\text{b}} = 2.494$

The total cross section has been measured for $E_{\text{n}} = 2.8$ to 3.3 MeV: a resonance is observed at $E_{\text{n}} \approx 2.95$ MeV. $\sigma_{\text{max}} = 4$ b, $\Gamma = 140 \pm 40$ keV, corresponding to $E_{\text{x}} \approx 5.26$ MeV, $J = 2^-$ or 3^{\pm} (1956BA1A).

15. $^{15}\text{N}(\text{n}, \alpha)^{12}\text{B}$ $Q_{\text{m}} = -7.629$ $E_{\text{b}} = 2.494$

See (1948JE03).

16. $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$ $Q_{\text{m}} = 0.267$
 $Q_0 = 0.286$ (1957VA11);
 $Q_0 = 0.269 \pm 0.010$ (1957WA01).

Levels derived from observed proton groups and γ -rays are listed in Table 16.2. Gamma transitions are shown in the inset of Fig. 31 (Energy Level Diagram of ^{16}N). Even at $E_{\text{d}} = 2.7$ MeV, the stripping angular distribution patterns to the low-lying states are well-developed, and the theory yields quite good fits (1956ZI1A, 1957WA01). Shell-model theory in intermediate coupling predicts a close group of 4 levels within about 700 keV, with $J = 0, 1, 2, 3$, odd parity (order

Table 16.2: Levels of ^{16}N from $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$

E_x^a (keV)	$\Gamma_{c.m.}$ (keV)	l_n^b	J^π^b	θ^2^c
0	sharp	2	2^-	0.05
120 ± 1	sharp	0	0^-	0.19
294 ± 5	sharp	2	3^-	0.05
392 ± 3	sharp	0	1^-	0.18
(3530 ± 30)	sharp			
3980 ± 20	sharp			
4800 ± 50	230 ± 40			
(5010 ± 50)				
5250 ± 50	290 ± 50			

^a (1957WA01: $E_d = 14.8$ MeV). No other proton groups are observed corresponding to $^{16}\text{N}^* < 9$ MeV. Energies of first three excited states are from γ -ray measurements (1957FR56, 1957WI1B).

^b (1957WA01) and (1956ZI1A: $E_d = 2.75$ MeV); J assignments from stripping and gamma decay data: see text.

^c (1957WA01).

uncertain), arising from the configurations $p_{\frac{1}{2}}^{-1}s_{\frac{1}{2}}$ and $p_{\frac{1}{2}}^{-1}d_{\frac{3}{2}}$. Levels from $p_{\frac{1}{2}}^{-1}d_{\frac{3}{2}}$ should lie several MeV higher (1953IN1A, 1957EL1B). These results are strikingly confirmed by the experimental evidence. The ground state is assigned $J = 2^-$ on the basis of the β -decay (see $^{16}\text{N}(\beta^-)^{16}\text{O}$). The first excited state may have $J = 0^-$ or 1^- from the stripping pattern; however, the half-life $\tau_{1/2} = 6.7 \pm 0.5 \mu\text{sec}$ (1957FR56), $5.43 \pm 0.22 \mu\text{sec}$ (1959ZI18) is much too long for dipole radiation, and $J = 0^-$ is indicated (1957WI1B). The third excited state ($E_x = 392$ keV), again limited to $J = 0^-$ or 1^- by the stripping pattern, decays to both $^{16}\text{N}(0)$ and $^{16}\text{N}^*(0.12)$; $J = 0^-$ is therefore excluded and $J = 1^-$ indicated. Of the possibilities $J = 1^-, 2^-, 3^-$ for the second excited state, $^{16}\text{N}^*(0.285)$, $J = 1^-$ is rendered unlikely by the low intensity of γ -decay to the second excited state and by the absence of an $l_n = 0$ component in the stripping pattern (1956ZI1A, 1957WA01, 1957WI1B). The assignment $J = 3^-$ is strongly favored by the (p- γ) angular correlation (1957FR56).

The observed γ -branching of $^{16}\text{N}^*(0.395)$ is in accord with the theory, which predicts lifetimes in the range 1 to 7×10^{-11} sec. A considerable enhancement through collective excitation is required to account for the lifetime of the first excited state. The reduced neutron widths are expected to be of the order of the single-particle limit. Calculation of level shifts and comparison of observed reduced widths suggests that the $^{16}\text{O}^*$ analogues to the first four states of ^{16}N are

$^{16}\text{O}^*(12.95, 12.78, 13.24, 13.09 \text{ MeV})$ (1957EL1B): see also (1957WI1B) and $^{15}\text{N}(p, p)^{15}\text{N}$.

17. $^{15}\text{N}(t, d)^{16}\text{N}$ $Q_m = -3.765$

Not reported.

18. $^{15}\text{N}(\alpha, ^3\text{He})^{16}\text{N}$ $Q_m = 18.084$

Not reported.

19. $^{16}\text{O}(n, p)^{16}\text{N}$ $Q_m = -9.619$

The possibility that ^{16}N might have a long-lived ($\approx \text{sec}$) isomeric state has been examined, with negative result (1956TO1A). See also (1952LI24) and ^{17}O .

20. $^{16}\text{O}(t, ^3\text{He})^{16}\text{N}$ $Q_m = -10.384$

Not reported.

21. $^{17}\text{O}(d, ^3\text{He})^{16}\text{N}$ $Q_m = -8.271$

Not reported.

22. $^{18}\text{O}(n, t)^{16}\text{N}$ $Q_m = -13.349$

Not reported.

23. $^{18}\text{O}(p, ^3\text{He})^{16}\text{N}$ $Q_m = -14.114$

Not reported.

24. $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$

$$Q_{\text{m}} = 4.237$$

$$Q_0 = 4.237 \pm 0.009 \text{ (1955PA50)}.$$

Alpha groups are observed at $E_{\text{d}} = 0.85 \text{ MeV}$ corresponding to $^{16}\text{N}^*(0, 116 \pm 6, 300 \pm 12, 391 \pm 12 \text{ keV})$. No other states are observed below $E_{\text{x}} = 1.24 \text{ MeV}$ (1955PA50). See also (1957BO04) and ^{20}F .

25. $^{19}\text{F}(\text{n}, \alpha)^{16}\text{N}$

$$Q_{\text{m}} = -1.500$$

At $E_{\text{n}} = 4.87 \text{ MeV}$, an α -group has been observed with $Q = -1.77 \pm 0.15 \text{ MeV}$, probably corresponding to a transition to one or more excited states of ^{16}N (1955JA18). See also (1956TO1A) and ^{20}F .

¹⁶O
(Fig. 32)

GENERAL:

Theory: See (1954DE1E, 1954FL1A, 1955HE1E, 1955JA1D, 1955MA1K, 1955MA1L, 1955SC1B, 1955WI1F, 1956EL1C, 1956FE1E, 1956JA1C, 1956KA1D, 1956MO1D, 1956PE1A, 1956RE1C, 1956WI1D, 1957EL1B, 1957FE1D, 1957GR1G, 1957HE1B, 1957RE1A, 1957TA1C, 1957TO1A, 1958CA1G, 1958DA1E, 1958DA1F, 1958FE1C, 1958FE1D, 1958HA1F, 1958MO17, 1958RA1F, 1958UM1A, 1958WI1E).

1. $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ $Q_m = 7.148$

Resonant capture radiation to $^{16}\text{O}_{\text{g.s.}}$ is observed at $E_\alpha \approx 3.24$ MeV, corresponding to the known $J = 1^-$ state at 9.58 MeV: see $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$. The radiative width of 6×10^{-3} eV implies a $T = 1$ admixture of the order of 3×10^{-4} , an amount slightly lower than usual for ^{16}O states. It is suggested that the $T = 1$ admixture may derive from $^{16}\text{O}^*(13.09)$, $J = 1^-$; $T = 1$ (1957BL01): see also (1953JO1A, 1956WI1D). This state does not arise in any natural way from $p^{-1}s$, or $p^{-1}d$ (1957EL1B). The integrated capture cross section to $E_\alpha = 1.60$ MeV is $< 30 \mu\text{b-MeV}$ (1955AL16). See also (1958PH37). The relevance of the capture of alpha particles by ^{12}C to the buildup of the elements in stars is discussed by (1956CA1F, 1956HA1C, 1956HA1D, 1957BU66, 1957SA1B).

2. $^{12}\text{C}(\alpha, n)^{15}\text{O}$ $Q_m = -8.452$ $E_b = 7.148$

See (1939KI1A).

3. $^{12}\text{C}(\alpha, \alpha n)^{11}\text{C}$ $Q_m = -18.722$ $E_b = 7.148$

See (1953LI28).

4. $^{12}\text{C}(\alpha, p)^{15}\text{N}$ $Q_m = -4.965$ $E_b = 7.148$

See ^{15}N .

5. $^{12}\text{C}(\alpha, d)^{14}\text{N}$ $Q_m = -13.579$ $E_b = 7.148$

Table 16.3: Energy Levels of ^{16}O

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or Γ (keV)	Decay	Reactions
0	$0^+; 0$	—	stable	1, 18, 24, 27, 34, 36, 37, 38, 43
6.056 ± 10	$0^+; 0$	$\tau_m = 72 \pm 7$ psec	π	9, 18, 35, 36, 38, 43
6.135 ± 10	$3^-; 0$	$\tau_m = 12 \pm 6$ psec	γ	18, 24, 27, 35, 36, 38, 43
6.923 ± 10	$2^+; 0$	$\tau_m = 12 \pm 3$ fsec	γ	18, 33, 36, 38, 43
7.121 ± 10	$1^-; 0$	$\tau_m = 10 \pm 3$ fsec	γ	18, 24, 27, 33, 36, 38, 43
8.875 ± 10	$2^-; 0$		γ	18, 24, 27, 36, 38, 43
9.58	$1^-; 0$	650	α, γ	1, 6
9.843 ± 12	$2^+; 0$	0.8	α	6, 36, 43
10.363 ± 14	$4^+; 0$	27	α	6, 36, 43
(10.804)				(43)
10.937 ± 10	$0^-; 0$		γ	(18), 24, (43)
11.070 ± 10	$3^+; (0)$		γ	(6), 18, 24, 36, 43
11.25	$0^+; 0$	2500	α	6
11.51 ± 30	$2^+; 0$	80	α	6, 36
11.62	$3^-; 0$	1200	α	6
12.02 ± 30			(γ)	36
(12.29)		40	γ	20
12.43 ± 10	$1^-; 0$	89	α, γ, p	6, 20, 22
12.52 ± 10	2^-	0.8	p, γ, α	20, 22, 36
12.78 ± 10	$0^-; 1$	38	p, γ	20, 21
12.96 ± 10	$2^-; 1$	2 ± 0.2	p, α	21, 22
13.09 ± 10	$1^-; 1$	130 ± 10	p, γ, α	20, 21, 22, 36
13.25 ± 10	$3^-; 1$	21 ± 1	p, α	21, 22
13.65 ± 10	$1^+; 0$	64 ± 3	p, α	21, 22
13.97 ± 10	2^-	22 ± 2	p, α	21, 22
14.93 ± 40	4^+	43 ± 10	p, α	22
15.21 ± 40	$2^-, 3^+$	72 ± 15	p, α	22
15.25 ± 60	2^+	720 ± 100	p, α	22
15.41 ± 50		96 ± 25	p, α	22
15.79		30	p, α	22

Table 16.3: Energy Levels of ^{16}O (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m or Γ (keV)	Decay	Reactions
16.21	1^+	23	p, n	23
16.3	0^-	250	p, n	23
16.44		24	p, α	22
(16.82)			p, α	22
(16.93)			p, α	22
17.0		≈ 200	p, n	23
17.12		41	p, n	23
17.29		84	p, n	23
17.5		≈ 250	p, n	23
17.63		66	p, n	23
17.85		100	p, n	23
17.96		49	p, n	23
18.05		38	p, n	23
22.05		broad	d, n	11
23.05		broad	d, n	11
23.54		300	α	6
24.38		broad	d, n	11
(25.7)			^3He , p, α	7, 8
(26.4)			^3He , p	7

The angular distribution at $E_\alpha = 42$ MeV shows three peaks and three valleys in the range $\theta_{\text{c.m.}} = 40^\circ$ to 150° . The location of peaks and the absolute cross sections agree closely with those obtained in the inverse reaction at $E_d = 20$ MeV. The experiment provides a test of the hypothesis of invariance under time reversal and places an upper limit of a few per cent on possible T-R odd forces (1958BO71).

 6. $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$

$$E_b = 7.148$$

Resonances derived from a phase-shift analysis of the elastic scattering are exhibited in Table 16.4 (1953HI05, 1954BI96). At the upper limit of these experiments ($^{16}\text{O}^*(12.5)$), the existence of higher $J = 0^+$ and 2^+ levels is indicated by a pronounced increase in the $l = 0$ and $l = 2$ phase shifts (1953HI05, 1954BI96). The inelastic scattering for $E_\alpha = 20.4$ to 22.6 MeV indicates

Table 16.4: Resonances in $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$

E_α (MeV)	Γ_{lab} (keV)	θ_α^2	$^{16}\text{O}^*$ (MeV)	J^π
3.24 ^a	860	0.85	9.58	1 ⁻
3.582 ^a	1	0.0015	9.835	2 ⁺
4.28 ^b	36	0.26	10.36	4 ⁺
5.27 ^b	10		(11.10) ^d	
5.47 ^b	3300	0.76	11.25	0 ⁺
5.82 ^b	106	0.03	11.51	2 ⁺
5.96 ^b	1600	0.73	11.62	3 ⁻
7.04 ^b	230	0.04	12.43	1 ⁻
21.85 ^c	400		23.54	

^a (1953HI05).

^b (1954BI96).

^c (1955RA1B).

^d Assignment to present reaction not certain.

a resonance at $E_\alpha = 21.85 \pm 0.1$ MeV, $\Gamma \approx 0.4$ MeV ($^{16}\text{O}^*(23.54)$). The asymmetry of the angular distribution is said to indicate at least one other level, of opposite parity (1955RA1B). See also (1954DE1E; theor.) and ^{12}C .

7. $^{13}\text{C}(^3\text{He}, \text{p})^{15}\text{N}$

$$Q_m = 10.668$$

$$E_b = 22.780$$

The yields of ground state protons have been studied at several angles in the region $E(^3\text{He}) = 1.4$ to 4.8 MeV. There are indications of resonances at $E(^3\text{He}) \approx 3.6$ and 4.5 MeV ($^{16}\text{O}^*(25.7$ and $26.4)$) (1956SC01, 1957IL01). See also ^{15}N .

8. $^{13}\text{C}(^3\text{He}, \alpha)^{12}\text{C}$

$$Q_m = 15.632$$

$$E_b = 22.780$$

The ground state α -group shows a broad weak resonance at $E(^3\text{He}) \approx 3.6$ MeV (1956WO1C). See also ^{12}C .

9. $^{13}\text{C}(\alpha, \text{n})^{16}\text{O}$

$$Q_m = 2.203$$

A threshold for $^{16}\text{O}^*(6.05)$ is observed at $E_\alpha = 5.05$ MeV (1956BO61). See also (1951JO1C) and ^{17}O .

10. $^{14}\text{C}(^3\text{He}, \text{n})^{16}\text{O}$ $Q_m = 14.607$

Not reported.

11. $^{14}\text{N}(\text{d}, \text{n})^{15}\text{O}$ $Q_m = 5.073$ $E_b = 20.728$

Excitation functions have been measured for $E_d = 0.6$ to 4.5 MeV (1955MA85, 1956JO1D, 1957JO1C, 1957NO1C, 1958MO14, 1958WE31). The yield of ground-state neutrons shows broad but well-defined peaks at $E_d = 1.52, 2.62,$ and 4.19 MeV; $E_x = 22.05, 23.05,$ and 24.38 MeV. It is not clear whether the structure is to be attributed to resonances or to surface interaction (1958WE31). See also ^{15}O .

12. $^{14}\text{N}(\text{d}, \text{p})^{15}\text{N}$ $Q_m = 8.615$ $E_b = 20.728$

Excitation functions and angular distributions are reported by (1954JO1F: $E_d = 0.4$ to 0.6 MeV) (1956KO26, 1956VA17: $E_d = 0.2$ to 0.7 MeV), and (1958BO18: $E_d = 0.6$ to 1.0 MeV): see also (1957JA37). At the lower energies, both stripping and compound nucleus effects are reported, although the distributions for $E_d = 0.6$ to 1.0 MeV (1958BO18) appear to be explicable entirely on the basis of overlapping resonance levels. See also ^{15}N and (1954ST1C).

13. $^{14}\text{N}(\text{d}, \text{d})^{14}\text{N}$ $E_b = 20.728$

See ^{14}N .

14. $^{14}\text{N}(\text{d}, \text{t})^{13}\text{N}$ $Q_m = -4.292$ $E_b = 20.728$

See (1942BO1A).

15. $^{14}\text{N}(\text{d}, \alpha)^{12}\text{C}$ $Q_m = 13.579$ $E_b = 20.728$

The cross section rises gradually from $E_d = 0.45$ to 0.90 MeV (1954CA1D). Angular distributions for $E_d = 0.6$ to 1.0 MeV indicate that the stripping contribution is small in this range (1958BO18). The angular distribution at $E_d = 20$ MeV agrees closely with that of the inverse reaction at $E_\alpha = 42$ MeV (1958BO71): see $^{12}\text{C}(\alpha, d)^{14}\text{N}$, (1957FI1C) and ^{12}C .

$$16. \ ^{14}\text{N}(d, 4\alpha) \qquad Q_m = 6.298 \qquad E_b = 20.728$$

See (1947FO1A).

$$17. \ ^{14}\text{N}(t, n)^{16}\text{O} \qquad Q_m = 14.470$$

Not reported.

$$18. \ ^{14}\text{N}(^3\text{He}, p)^{16}\text{O} \qquad Q_m = 15.235$$

At $E(^3\text{He}) = 2.1$ MeV, proton groups corresponding to ^{16}O levels up to $E_x = 13.6$ MeV have been identified. In the region $E_x \approx 11$ MeV, four groups are resolved, corresponding to $^{16}\text{O}^*(10.94, 11.087 \pm 0.020, 11.25, 11.51)$. The first two are presumably those observed in $^{15}\text{N}(d, n)^{16}\text{O}$ at 10.937 and 11.063 MeV. The gamma decay of $^{16}\text{O}^*(8.88, 10.94$ and $11.07)$ has been studied with coincidence techniques: branching ratios are given in Fig. 33 (Gamma-ray transitions in ^{16}O). From the observed transition intensities and from (p- γ) and (γ - γ) correlations, the assignment $J = 2^-$ for $^{16}\text{O}^*(8.88)$ is confirmed, and assignments of $J = 0^-$ and 3^+ are fixed for $^{16}\text{O}^*(10.94$ and $11.07)$, respectively. For the 10.94 MeV state, $\Gamma_\alpha/\Gamma_\gamma < 0.2$, indicating an upper limit for the intensity of possible opposite-parity admixture of $\approx 2 \times 10^{-9}$ (1959BR68, 1959KU78). See also (1957BR17, 1957LI1D, 1958BR1D).

$$19. \ ^{14}\text{N}(\alpha, d)^{16}\text{O} \qquad Q_m = -3.116$$

Not reported.

$$20. \ ^{15}\text{N}(p, \gamma)^{16}\text{O} \qquad Q_m = 12.113$$

Ground-state capture radiation resonances occur at $E_p = 190, 338$ keV (1958HE52) and 1050 keV (1952SC28, 1957HA98, 1958HE52): see Table 16.5. The large radiative width, $\Gamma_\gamma = 150$ eV, of the latter indicates E1 radiation and $J = 1^-$; $T = 1$ for $^{16}\text{O}^*(13.09)$; on the other hand, the large

α -width speaks for a strong admixture of $T = 0$ (1953WI1B, 1956WI1D: see also (1957BA03)). The 338 keV resonance is relatively weak, $\Gamma_\gamma = 8$ eV. The Breit-Wigner formula with destructive interference between these two $J = 1^-$ states gives a good account of the γ_0 yield from $E_p = 200$ to 1200 keV (1958HE52). Cascade radiation, via $^{16}\text{O}^*(6$ and 7), is weakly resonant at $E_p = 340$ keV (1958HE52): at the higher resonance, the relative amount of cascade radiation is $< 1.3 \times 10^{-3}$ (1953DE1A, 1954GO1F). No further resonances for ground-state radiation appear for $E_p < 3.3$ MeV with intensity $> 2\%$ of that at $E_p = 1050$ keV (1957BA03). Within a few microbarns, the cross section for $E_p < 3$ MeV is completely accounted for by the 13.09 MeV state; a small rise from $E_p = 3$ to 4 MeV may reflect higher resonances. This result is in disagreement with that of (1955SP1B) on the inverse reaction in which a level at $^{16}\text{O}^*(14.7)$ ($E_p = 2.8$ MeV) is reported. The isotropy of the radiation over the entire range from $E_p = 1$ to 4 MeV places an upper limit of 3% on the relative intensity of $p^{-1}d$ in the 13.09 MeV state (1957WI1H); see also (1957EL1B).

At $E_p = 429$ keV (1958HE52) and 710 keV (1957HA98), resonances for cascade radiation are found; neither produces any detectable ground state radiation. See also (1958CA13).

21. $^{15}\text{N}(p, p)^{15}\text{N}$

$$E_b = 12.113$$

Elastic scattering studies are reported for $E_p = 600$ to 1800 keV by (1957HA98) and for $E_p = 950$ to 3960 keV by (1956BA1H); see Table 16.5. The $E_p = 710$ keV state, ($^{16}\text{O}^*(12.78)$) having $J = 0^-$, does not appear in the (p, α) reaction or in the ground-state capture: see $^{15}\text{N}(p, \gamma)^{16}\text{O}$. The assignment $J = 3^-$ to the state at $E_p = 1210$ keV ($^{16}\text{O}^*(13.25)$) is in disagreement with the observed angular distribution in $^{15}\text{N}(p, \alpha_1)^{12}\text{C}^*$ but is confirmed by the α_1 - γ correlation. The reason for this discrepancy is not known (1957HA98).

By comparison of J^π assignments and reduced widths, it is concluded that the ^{16}O levels formed at $E_p = 898, 710, 1210,$ and 1028 keV are the analogues of the first four states of ^{16}N (1956WI1G, 1956WI1H, 1957EL1B, 1957HA98). It is pointed out that the $E_p = 340$ and 429 keV states are also possible candidates, and that in any case appreciable isobaric spin mixing is to be expected (1957HA98). See also (1956WI1D).

Elastic scattering at $\theta = 132^\circ$ shows structure at $E_p = 3000$ keV and a large peak of undetermined origin at $E_p = 3920 \pm 40$ keV (1956BA1H). See also (1958CA13).

22. $^{15}\text{N}(p, \alpha)^{12}\text{C}$

$$Q_m = 4.964$$

$$E_b = 12.113$$

Two groups of α -particles occur, to $^{12}\text{C}(0)$ (α_0) and to $^{12}\text{C}^*(4.43, J = 2^+)$ (α_1, γ). Observed resonances are exhibited in Table 16.5. The cross section for (p, α_0) is 5×10^{-7} b at $E_p = 100$ keV (1950SC1A: see also (1952SC28, 1957BA03)). See (1957JA37).

Angular distributions $(p-\alpha_0)$ have been studied in the range $E_p = 230$ to 960 keV by (1953CO1D), $E_p = 500$ to 1000 keV by (1953NE1B), $E_p = 920$ to 1260 keV by (1957HG01). At and below the 338 keV resonance, the distribution is isotropic. Strong $\cos \theta$ terms develop for $E_p > 400$

Table 16.5: Levels of ^{16}O from $^{15}\text{N}(p, p)^{15}\text{N}$, $^{15}\text{N}(p, \gamma)^{16}\text{O}$ and $^{15}\text{N}(p, \alpha)^{12}\text{C}$

E_p (keV)	Γ_{lab} (keV)	Γ_p (keV)	$\sigma(\alpha_0)$ (mb)	$\sigma(\alpha_1)$ (mb)	$\sigma(\gamma_0)$ (μb)	$\sigma(\gamma_1)$ (μb)	θ_p^2 (%)	$J^\pi; T$	E_x (MeV)
(190)	40				res ^f				12.29
338 ^a	94	0.8 ^{a, f}	75 ^{a, f}	0.03 ^f	7 ^f	0.4 ^f	4 ^{f, c}	1 ⁻ ; 0 ^c	12.430
429 ± 1 ^a	0.9	0.020 ^f	n.r. ^g	200 ^{a, f}		1300 ^f	4 ^b	2 ⁻	12.515
710 ± 7 ^b	40 ± 4	40 ^b	n.r.	n.r.	n.r.	res	11	0 ⁻ ; 1 ^b	12.779
898 ± 1 ^a	2.2 ± 0.2	1.2	n.r.	800			8.8	2 ⁻ ; 1	12.955
1028 ± 10 ^b	140 ± 10	110	500 ⁱ	15	1000 ^a	< 1.3 ^h	10	1 ⁻ ; 1	13.088
1210 ± 3 ^a	22.5 ± 1	4.1	600 ⁱ	425 ^d			7.4	3 ⁻ ; 1	13.247
1640 ± 3 ^c	68 ± 3	10	n.r. ^d	340			0.8	1 ⁺ ; 0	13.651
1979 ± 3	23 ± 2	0.5 ^c	n.r.	35				2 ⁻ ^d	13.968
3000 ± 30 ^d	45 ± 10		50	600				4 ⁺	14.93
3300 ± 35	75 ± 15		n.r.	270				2 ⁻ , 3 ⁺	15.21
3350 ± 50	750 ± 100		50	205				2 ⁺	15.25
3520 ± 40	100 ± 25		100	80					15.41
3920 ^e	30			res ^e					15.79
4610	25			res					16.44
(5020)				(res)					(16.82)
(5140)				(res)					(16.93)

^a (1952SC28); values for 338-keV resonance corrected for s-wave penetration: observed $E_{\text{max}} = 360$ keV, $\sigma_{\alpha_0}(\text{max}) = 90$ mb.

^b (1957HA98).

^c (1957HG01).

^d (1957BA03).

^e (1956LI1C); see also Table 16.6.

^f (1958HE52).

^g n.r. = non-resonant.

^h (1954GO1F).

ⁱ (1952SC28): (1957HG01) find 340 and 300 mb respectively.

keV, with $\cos^2 \theta$ terms gradually increasing above 700 keV, and higher-order terms above 1 MeV. The terms in $\cos \theta$ indicate interference between states of opposite parity. Analysis of this effect led (1953CO1D) to the conclusion that the resonances in question were $E_p = 338$ keV, assigned $J = 0^+$, and $E_p = 1028$ keV, $J = 1^-$. Subsequent work has shown, however, that the lower state is $J = 1^-$, formed by s-wave protons: evidently the interfering even parity state or states remain to be identified; see also $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$. The 1028 keV resonance has a large width both for α -particles and E1 radiation and appears to have a mixed character as regards isobaric spin (1953WI1A, 1953WI1B, 1956WI1D, 1957HA98).

Analysis of the $(p-\alpha_0)$ distribution at the $E_p = 1210$ keV resonance indicates $J = 3^-$, although $J = 4^+$ is not clearly excluded (1957HG01). Angular distributions of (p, α_1) and the 4.4 MeV γ -ray have been studied at the $E_p = 429, 898,$ and 1210 keV resonances by (1952BA1C, 1952SE1B, 1953KR1B). Channel-spin ratios have been interpreted in terms of L - S and j - j coupling models by (1953CH1A). The α_1 -distribution at $E_p = 1210$ keV appears to require $J = 4^+$ (1953KR1B, 1957HA98), but the $(\alpha_1-\gamma)$ correlation requires $J = 3^-$, in agreement with elastic scattering results; the reason for this discrepancy is not clear (1957HA98). Angular distributions of the 4.4 MeV γ -rays are reported for $E_p = 1050$ and 1640 keV by (1954KR1C) and for various energies from $E_p = 1210$ to 3900 keV by (1957BA03). See also (1958CA13).

A pronounced anomaly appears at $E_p = 1.87$ MeV in the (p, α_0) excitation curve which is faintly reproduced in (p, α_1) but not in (p, p) (1957HG01).

23. $^{15}\text{N}(p, n)^{15}\text{O}$

$$Q_m = -3.542$$

$$E_b = 12.113$$

The excitation function has been measured from threshold to 6.4 MeV, and angular distributions have been studied at several energies: see Table 16.6. Only four levels are found with $\Gamma \lesssim 40$ keV, as contrasted with the ten or more reported in $^{16}\text{O}(\gamma, n)^{15}\text{O}$ (1958JO28).

24. $^{15}\text{N}(d, n)^{16}\text{O}$

$$Q_m = 9.886$$

Slow neutron thresholds have been observed at $E_d = 1.192$ and 1.335 MeV, corresponding to $^{16}\text{O}^*(10.937 \pm 0.010, 11.063 \pm 0.015)$ (1957WE1A). At $E_d = 5.1$ MeV, in this reaction, and at $E_p = 5.77$ MeV in $^{19}\text{F}(p, \alpha)^{16}\text{O}$, four γ -ray lines are observed which are assigned to ^{16}O : $E_\gamma = 2.73$ ($8.87 \rightarrow 6.14$), 3.86 ± 0.04 ($10.94 \rightarrow 7.12$), 6.14 ($6.14 \rightarrow 0$), and 7.1 MeV ($6.9 + 7.1 \rightarrow 0$). The 10.94 MeV state is not observed to decay in any way other than through the $J = 1^-$ state at 7.12 MeV; upper limits to transitions to $^{16}\text{O}^*(0, 6.06, 6.14, 6.92, 8.87)$ are, respectively, 5, 1, 6, 20, and 40% of the observed cascade. The strong transition to $^{16}\text{O}^*(7.1)$, $J = 1^-$, suggests $J = 0^-$ for $^{16}\text{O}^*(10.94)$, although $J = 1^+$ is not ruled out. The γ - γ correlation strongly favors $J = 0^-$. It is suggested that this γ -emitting state is to be identified with the first neutron threshold: $^{16}\text{O}^*(10.94)$, above, and is distinct from either the (doubtful) $^{16}\text{O}^*(11.1)$ reported in $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$ or the $^{16}\text{O}^*(11.08)$ reported in $^{16}\text{O}(p, p')^{16}\text{O}^*$. Possibly both $^{16}\text{O}^*(10.94$ and 11.08) are involved

Table 16.6: Resonances in $^{15}\text{N}(\text{p}, \text{n})^{15}\text{O}$ (1958JO28)

E_p (MeV)	Γ_{lab} (keV)	J^π	E_x in ^{16}O (MeV)
4.372	24	1^+	16.21
≈ 4.5	≈ 250	0^-	16.3
≈ 5.3	≈ 200		17.0
5.35	44		17.12
5.52	90		17.29
≈ 5.8	≈ 250		17.5
5.88	70		17.63
6.12	105		17.85
6.24	52		17.96
6.33	40		18.05

in $^{14}\text{N}({}^3\text{He}, \text{p})^{16}\text{O}$ (1957BE61). It is noted that the $^{19}\text{F}(\text{p}, \alpha)^{16}\text{O}$ spectrum of (1956SQ1A) shows three groups in this range, of which one is definitely attributed to $^{16}\text{O}^*(11.085)$ (1957WE1A). The 8.87 MeV state has a $7 \pm 2\%$ direct transition to $^{16}\text{O}_{\text{g.s.}}$ (1957BE61). Angular distributions of ground-state neutrons for $E_d = 1.1$ to 5.2 MeV are well accounted for by the exchange stripping theory of (1957OW03) (1958WE31: see ^{17}O). See also (1955AJ61) and (1957EL1B; theor.).

25. $^{15}\text{N}({}^3\text{He}, \text{d})^{16}\text{O}$ $Q_m = 6.619$

Not reported.

26. $^{15}\text{N}(\alpha, \text{t})^{16}\text{O}$ $Q_m = -7.700$

Not reported.

27. $^{16}\text{N}(\beta^-)^{16}\text{O}$ $Q_m = 10.402$

^{16}N decays to several states of ^{16}O : reported branching fractions are listed in Table 16.7. The ground-state transition, with $E_\beta(\text{max}) = 10.33 \pm 0.08$ MeV (1957MO1A), 10.40 ± 0.05 MeV (1958BR95), has the unique first-forbidden shape corresponding to $\Delta J = 2$, yes, fixing J^π of ^{16}N as 2^- (1957MO1A, 1958BR95, 1959AL1M). This assignment is also indicated by the fact

Table 16.7: Beta-decay of ^{16}N

Final State		A		B		C		D
$^{16}\text{O}^*$ (MeV)	J^π	Branch (%)	$\log f_0t$	Branch (%)	$\log f_0t$	Branch (%)	$\log f_0t$	
0	0^+	26 ± 2	6.7	28	6.67^a	24 ± 2	8.2^b	
6.06	0^+	≤ 0.015	≥ 8.2					
6.14	3^-	68	4.5	54	4.6	55 ± 4	4.5	14 ± 1
6.92	2^+							
7.12	1^-	4.9	5.1	18	4.6	21 ± 4	4.4	1
8.88	2^-	1.1	4.4					

A: (1956W11A, 1958AL13, 1959AL1M): includes data of B, C, D.

B: (1958BR95).

C: (1957MO1A).

D: γ -rays (1951MI1B, 1956TO1A, 1957BO04).

^a $\log(f_1t) = 8.0$.

^b $\log(f_1t)$.

that the transitions to $^{16}\text{O}^*(6.1$ and $7.1)$ are both allowed (1951MI1B). There appears to be some discrepancy between the branching ratios of $^{16}\text{O}^*(6.1$ and $7.1)$ as determined by decomposition of the β -spectrum (1958BR95) and by comparison of the γ -ray intensities (1951MI1B, 1957BO04: see, however, (1959AL1M)). The low ft -value for $^{16}\text{O}^*(7.1)$ presents some difficulty for the theory (1957EL1B). Transitions to the nuclear pair emitting state are $< 1.5 \times 10^{-4}$: $\log ft > 8.2$ (1958AL13: see also (1956EL1C, 1957EL1B)). A 1.1% branch leads to $^{16}\text{O}^*(8.87)$ which decays via the 7.1, 6.9, and 6.1 MeV levels in the ratio 3 : 1 : 30. Since the β -transition is allowed, J^π of $^{16}\text{O}^*(8.8)$ is 1^- , 2^- or 3^- ; the first and last would permit α -decay, so $J^\pi = 2^-$. The γ -branching and the γ - γ correlation ($8.88 \rightarrow 6.1 \rightarrow \text{g.s.}$) are consistent with this assignment (1956W11A).

The half-life is 7.35 ± 0.05 sec (1947BL1A), 7.38 ± 0.05 sec (1954MA97). (1956TO1A) finds no evidence for β -decay of excited states of ^{16}N .

28. $^{16}\text{O}(\gamma, n)^{15}\text{O}$

$$Q_m = -15.655$$

$$E_{\text{thresh.}} = 15.60 \pm 0.05 \text{ (1957BA27: see also (1955BA1P))}.$$

The cross section exhibits a slow rise for ≈ 3 MeV above threshold followed by the usual giant resonance. Characteristics of the giant resonance are: $E_\gamma = 24$ MeV, $\Gamma = 3.5$ MeV, $\sigma_{\text{max}} = 10$ mb; $\int \sigma dE = 40$ MeV-mb (1951JO1B, 1953MO1B, 1954FE16, 1957CA1D).

Discontinuities in the activation curve are said to indicate absorption into discrete levels of ^{16}O : see Table 16.8 (1955BA1P, 1955CO1C, 1955PE1D, 1957BA27, 1958BE74, 1958KA1D). From the cross section it would appear that a substantial fraction of the absorption takes place through discrete levels (1954KA1A, 1955PE1D). However, in a measurement of the absorption in water for $E_\gamma = 15$ to 25 MeV, using various detectors including ^{16}O , (1958SI1A) conclude that the absorption is generally continuous, with only a small contribution from discrete, narrow levels. See also (1954BI04, 1955CA1E, 1955SA1F, 1957SP1A, 1957SV1A, 1958LI1D, 1958WO51), (1955MO1C, 1956WI1D, 1957BA1H, 1957EL1B, 1958FE1C, 1958FE1D, 1958WI1E; theor.) and (1955AJ61).

$$29. \ ^{16}\text{O}(\gamma, p)^{15}\text{N} \quad Q_m = -12.113$$

Resonances are observed at $E_\gamma = 19.6, 20.6, 22.4$ MeV; $\int \sigma dE = 2, 2,$ and 20 MeV-mb for ground-state protons (1956CO59), $E_\gamma = 20.7, 21.9, 24.0$ MeV (1956LI1D). See also (1955JO1D). The 22 MeV resonance appears to be the giant resonance. Proton angular distributions have been measured by (1956CO59, 1956LI1D, 1957MI1C). A resonance reported at $E_\gamma = 14.7$ MeV (1955SP1B: see also (1956CO59)) does not appear in the inverse reaction $^{15}\text{N}(p, \gamma)^{16}\text{O}$ (1955WI1F, 1957WI1H). (1958SI1A) find that the absorption in the range $E_\gamma = 15$ to 25 MeV is generally continuous, with only small contributions from narrow resonances: see also $^{16}\text{O}(\gamma, n)^{15}\text{O}$. See also (1955ST1D, 1956JO1C, 1957BR55, 1957MI1B, 1957SV1A, 1958LI1D, 1958MI89, 1958PE1A) and (1955MO1B, 1956GO1G, 1957BA1K, 1957WI1J; theor.).

$$30. \ ^{16}\text{O}(\gamma, \alpha)^{12}\text{C} \quad Q_m = -7.148$$

The cross section for production of ^{12}C exhibits a maximum near 17.5 MeV ($\Gamma \approx 5$ MeV), $\sigma(\text{max}) \approx 50 \mu\text{b}$ (1953MI31). Evidence is also reported for excited states of $^{16}\text{O}^*((14.2), 16.75, 17.3, 22.6, (23.15), \text{ and } 24.6)$ with $J = 2^+; T = 0$ (1954ST89). See also (1955HA1D, 1955TI1A, 1956DA1C) and (1955AJ61).

$$31. \ ^{16}\text{O}(\gamma, 4\alpha) \quad Q_m = -14.429$$

The cross section for production of 4-pronged stars shows a number of maxima: see Table 16.9 (1952GO1A, CO54I, 1956DA1C). An appreciable fraction of the stars appear to involve excited states of ^{12}C and ^8Be (1955AJ61, 1956DA1C: see, however, (1953MI31)). See also (1955RA1E, 1955HA1D) and (1955AJ61).

$$32. \text{ (a) } \ ^{16}\text{O}(\gamma, n\alpha)^{11}\text{C} \quad Q_m = -25.870$$

$$\text{ (b) } \ ^{16}\text{O}(\gamma, t)^{13}\text{N} \quad Q_m = -25.020$$

Table 16.8: Levels in ^{16}O from $^{16}\text{O}(\gamma, n)^{15}\text{O}$

E_γ (MeV)			
(1955PE1D)	(1954KA1A)	(1955BA1P, 1957BA27)	(1955CO1C, 1958BE74)
15.85	15.9 ^a	(15.8)	15.93 ^b
16.03		16.03	16.17 ^b
16.47	16.4	16.50	16.37 ^c
16.75	16.7	(16.7)	
16.95	16.9	16.82	16.84 ^c
17.02		17.04	
17.13	17.1		
17.18			
17.55			
17.68			
17.84			
18.04			
18.70			
19.01	18.9		
19.18	19.3		
20.33			
20.58			
20.79	20.7		
20.93			
21.21			
21.52	21.54 ± 0.05		
	21.96 ± 0.05		
22.37			
22.54			
22.76			
23.02			

^a See also (1958KA1D).

^b (1958BE74).

^c (1955CO1C).

Table 16.9: Maxima in $^{16}\text{O}(\gamma, 4\alpha)$ ^a

E_γ (MeV)		
(1952GO1A)	(1956DA1C)	(CO54I)
22.6	22.5	23.2
25.8	(26.0)	24.7
	(27.5)	27.2
29.5	(29.5)	29.2
	32.5	
	(35.2)	

^a Maxima in number of 4-pronged stars.

See (1955SC36, 1957ER24).

33. $^{16}\text{O}(\gamma, \gamma')^{16}\text{O}^*$

Measurement of the resonance scattering cross section for 6.9 and 7.1 MeV γ -rays from $^{19}\text{F}(\text{p}, \alpha)^{16}\text{O}$ yields mean lifetimes of $(1.2 \pm 0.3) \times 10^{-14}$ and $(1.0 \pm 0.3) \times 10^{-14}$ sec for $^{16}\text{O}^*(6.92$ and $7.12)$, respectively. Values obtained from self absorption measurements are consistent with these. The observed life of the $7.1 \rightarrow \text{g.s.}$, E1 transition is consistent with an isobaric spin inhibition of the order of 300 (see (1955MA1K)). The lifetime of the 6.9 MeV state is longer than that expected from a collective model but can be accounted for by one version of the α -particle model (1957SW17). See also (1956KA1D, 1957GR1G; theor.) and $^{19}\text{F}(\text{p}, \alpha)^{16}\text{O}$.

34. $^{16}\text{O}(\text{e}, \text{e}')^{16}\text{O}^*$

Elastic scattering angular distributions at $E_e = 240, 360,$ and 420 MeV strongly favor a shell-model charge distribution based on a harmonic well with a length parameter $a = (1.76 \pm 0.02) \times 10^{-13}$ cm. The rms radius of the charge distribution is 2.70×10^{-13} cm (1958EH1B); see also (1956FE1B, 1957HO1E, 1958FE1D, 1958RA43).

35. $^{16}\text{O}(\text{n}, \text{n}')^{16}\text{O}^*$

At $E_n = 7.06$ MeV, a 6.094 ± 0.06 MeV γ -ray is observed with a cross section of 104 ± 25 mb (1956DA23). See also (1954TH42, 1955BE1H).

36. $^{16}\text{O}(\text{p}, \text{p}')^{16}\text{O}^*$

At $E_p = 19$ MeV, proton groups are observed corresponding to $^{16}\text{O}^*(6.14, 7.02, 8.87, 9.85, 10.34, 11.08, 11.51, 12.02, 12.53, 13.06, \text{ and } (13.39) \text{ MeV}) (\pm 30 \text{ keV})$. Of these, three have not been reported elsewhere: $^{16}\text{O}^*(11.08, 12.02, 13.39)$. The level at 11.08 MeV decays via cascade through the 6 to 7 MeV states; it is presumably not to be identified with the (doubtful) 11.10 MeV state reported by (1954BI96) in $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$ since the 10 keV width of the latter would preclude observation of the γ -decay. If it is a pure γ -emitting state, $J = 2^-$ is favored. The 12.02 MeV state appears to give weak γ -radiation; the state at 12.53 MeV yields 4.4 MeV γ -rays via $^{16}\text{O}(\text{p}, \text{p}'\alpha)^{12}\text{C}^*$ as expected from its behavior in $^{15}\text{N} + \text{p}$. The 8.87 MeV state cascades 80% via $^{16}\text{O}^*(6.1)$ and 20% via $^{16}\text{O}^*(6.9 - 7.1)$ (1955HO68).

At higher bombarding energies, evidence is reported for the excitation of states at 6 to 7, 12.5, and ≈ 20 MeV (1956ST30, 1957TY36: $E_p = 96$ and 177 MeV). The elastic scattering at high energies is generally characterized by direct interaction and may be described in terms of the optical model: see (1953BU72, 1955FU1A, 1955KI43, 1956BU95, 1956KI54, 1957CH32, 1957GI14, 1957VA1B). Polarization of scattered protons has been studied at $E_p = 173$ MeV by (1957AL39, 1957HI98, 1957MA58). See also (1955KO1A, 1957JA1B, 1958MA1B, 1958TY49).

37. $^{16}\text{O}(\text{d}, \text{d})^{16}\text{O}$

See ^{18}F .

38. $^{16}\text{O}(\alpha, \alpha')^{16}\text{O}^*$

Both elastic and inelastic scattering distributions ($^{16}\text{O}^*(6 - 7, 8.8)$) have been studied at $E_\alpha = 19$ MeV by (1957CO1H, 1958CO59). The elastic scattering shows strong diffraction effects.

39. $^{17}\text{O}(\text{p}, \text{d})^{16}\text{O}$

$$Q_m = -1.919$$

Not reported.

40. $^{17}\text{O}(\text{d}, \text{t})^{16}\text{O}$

$$Q_m = 2.113$$

Not reported.

41. $^{17}\text{O}({}^3\text{He}, \alpha)^{16}\text{O}$ $Q_m = 16.432$

Not reported.

42. $^{18}\text{O}(\text{p}, \text{t})^{16}\text{O}$ $Q_m = -3.730$

Not reported.

43. $^{19}\text{F}(\text{p}, \alpha)^{16}\text{O}$ $Q_m = 8.119$
 $Q_0 = 8.110 \pm 0.010$ (1956SQ1A).

Levels derived from observed α -particle groups are listed in Table 16.10 (1956SQ1A, 1957YO04: see also (1952AJ38)). There is some evidence for a broad level near 9.58 MeV (compare $^{12}\text{C}(\alpha, \alpha)^{12}\text{C}$) and for two additional unidentified groups which may represent levels near 11 MeV (see $^{15}\text{N}(\text{d}, \text{n})^{16}\text{O}$); no other groups appear with intensity $> 5\%$ of that corresponding to $^{16}\text{O}^*(8.87)$ (1956SQ1A). (1957YO04) find no other groups corresponding to $^{16}\text{O}^*(6 - 9)$ with $d\sigma/d\Omega > 0.4$ mb/sr. The indicated assignments for the first five states derive from a variety of evidence; see (1955AJ61).

6.06-MeV state

$W_\pi = 6.065 \pm 0.009$ MeV (1956AL1G). The angular correlation of the monopole pairs is given by $W(\theta) = 1 + (0.955 \pm 0.007) \cos \theta$, consistent with $J = 0^+ \rightarrow 0^+$ (1955GO1E, 1958AR1B: see also (1954DE36)). The mean life is $(7.2 \pm 0.7) \times 10^{-11}$ sec, considerably longer than is given by the α -model (1954DE36). Calculations on various models are summarized by (1957GR1G); see also (1956EL1C, 1957EL1B, 1957FE1D).

6.14-MeV state

The mean life is $(1.2 \pm 0.6) \times 10^{-11}$ sec (1958KO63), 0.5 to 1×10^{-11} sec (1955DE51). This transition rate is somewhat faster than that predicted on the α -particle model (1956KA1D) and is an order of magnitude faster than indicated by shell-model calculations (1957EL1B). It accounts for about one third of the total E3 width to the ground state from $T = 0$ states (1958KO63). See also (1958KN1B).

6.92-MeV state

$E_\gamma = 6.97 \pm 0.05$ MeV (1955BE62: Doppler corrected). The mean life is $\leq 2.5 \times 10^{-14}$ sec (1956DE22: see also (1955DE51)). Compare $^{16}\text{O}(\gamma, \gamma')^{16}\text{O}^*$: (1957SW17).

7.12-MeV state

Table 16.10: ^{16}O levels from $^{19}\text{F}(p, \alpha)^{16}\text{O}$

A		B		
E_x (MeV)	Γ (keV)	E_x (MeV)	$\frac{d\sigma}{d\Omega}$ ^a	J^π
0	< 20	0		0^+
6.051 ± 0.010	< 20	6.058 ± 0.017	0.1	0^+
6.131 ± 0.010	< 20	6.138 ± 0.011	0.8	3^-
6.920 ± 0.010	< 20	6.926 ± 0.011	1	2^+
7.120 ± 0.010	< 20	7.122 ± 0.011	2	1^-
8.874 ± 0.012	< 20	8.882 ± 0.011	0.6	2^-
9.852 ± 0.012	< 20			
10.363 ± 0.014	$\approx 25 - 30$			
11.085 ± 0.014	$\approx 25 - 30$			

A: (1956SQ1A): $E_p = 7$ MeV.

B: (1957YO04): $E_p = 5.2$ MeV; E_x calculated from reported Q -values and Q_m .

^a Cross sections in mb/sr at $\theta = 180^\circ$.

The mean life is $\leq 1.2 \times 10^{-14}$ sec (1955DE51); compare $^{16}\text{O}(\gamma, \gamma')^{16}\text{O}^*$: (1957SW17). A search for circular polarization of the γ -ray yields an upper limit of 2.0×10^{-3} indicating a maximum of $F^2 = 3 \times 10^{-8}$ for the intensity of any parity non-conserving part of the wave function (1958WI41).

8.87-MeV state

The direct ground state transition is 7 ± 2 % (1957BE61), 9 ± 4 % (1957MC35), 0.6% (1957WA1B) of the cascade decays. The cascade decays take place via the 6.14, 6.92, and 7.12 MeV states with relative intensities 27 : 1 : 3. The three γ -rays have energies 2.75 ± 0.02 , 1.90 ± 0.03 , and 1.72 ± 0.03 MeV, fixing the energy of the state as 8.87 ± 0.02 MeV. The observed branching ratios, together with the absence of α -decay fix the assignment as $J^\pi = 2^-$ (1956WI1A, 1957MC35). The separation of this state from the nearest 2^+ state ($^{16}\text{O}^*(9.85)$) is too large to be accounted for in the α -model of (1954DE1E) (1956WI1A: see also (1956KA1D)).

10.94-MeV state

At $E_p = 5.77$ MeV, a γ -ray of energy 3.86 ± 0.03 MeV is observed in coincidence with the 7.1 MeV radiation, indicating a level energy of 10.98 ± 0.04 MeV. It is presumed that this level is to be identified with one of the unassigned groups observed by (1956SQ1A, 1957BE61). The direct ground state decay is $< 5\%$ and cascades via $^{16}\text{O}^*(6.0, 6.1, 6.9, \text{ and } 8.9)$ are less than 1, 6,

20, and 40%, respectively. The strong branch to $^{16}\text{O}^*(7.1)$, $J = 1^-$, and the weakness of other branches suggests $J(10.94) = 0^-$, although 1^+ is not excluded. The γ - γ correlation favors $J = 0^-$ (1957BE61: see also $^{15}\text{N}(d, n)^{16}\text{O}$). ((1957WA1B) find the γ -ray energy to be 3.86 ± 0.02 MeV and report a weak 4.9 MeV γ -ray). A search for possible nuclear pairs from the $0^- \rightarrow 0^+$ ground-state transition yielded an upper limit of $< 2 \times 10^{-5}$ for the pair branch (1958EK36). The importance of a definitive assignment of J^π for this state is stressed by (1957EL1B).

The influence of isobaric spin selection rules on cascade decays in ^{16}O has been discussed by (1953WI1E, 1956WI1D: see also (1955MA1K, 1955MA1L)). At $E_p = 18.5$ MeV, the α_0 angular distribution has been studied by (1956LI37): see ^{19}F .

The reaction exhibits a large number of resonances: see ^{20}Ne . See also (1956IS1A, 1957BI75, MO57C, 1958MO1J).

^{16}F
(Not illustrated)

Mass of ^{16}F : From the threshold of the $^{14}\text{N}(^3\text{He}, \text{n})^{16}\text{F}$ reaction and the Wapstra (1955WA1A) masses for ^{14}N , ^3He and n , the mass excess of ^{16}F is 15.63 ± 0.02 MeV. A semi-empirical computation of the level shifts of the low $T = 1$ levels of ^{16}N and ^{16}O suggests that the ground state of ^{16}F should have $J = 0^-$, and be unstable to proton emission by about 1 MeV (1957EL1B). Using the $(M - A)$ stated above, ^{16}F is unstable with respect to proton emission by 0.81 MeV. The binding energies of a deuteron, a ^3He -particle, and an α -particle in ^{16}F are, respectively, 10.24, 9.37 and 8.98 MeV.

1. $^{14}\text{N}(^3\text{He}, \text{n})^{16}\text{F}$ $Q_m = -1.18$

A slow neutron threshold has been observed at $E(^3\text{He}) = 1.434 \pm 0.015$ MeV corresponding to the ground state of ^{16}F ($Q_0 = -1.18 \pm 0.01$ MeV). There is some indication of another threshold corresponding to an excited state at ≈ 60 keV. The lifetime of ^{16}F for decay into $^{15}\text{O} + \text{p}$ is computed to be $\approx 10^{-19}$ sec (H. Bichsel, private communication). See also (1956BU22).

2. $^{16}\text{O}(\text{p}, \text{n})^{16}\text{F}$ $Q_m = -16.41$

See ^{17}F .

3. $^{16}\text{O}(^3\text{He}, \text{t})^{16}\text{F}$ $Q_m = -15.61$

Not reported.

References

(Closed 01 December 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1939KI1A L.D.P. King, W.J. Henderson and J.R. Risser, *Phys. Rev.* 55 (1939) 1118; 41
- 1942BO1A L.B. Borst, *Phys. Rev.* 61 (1942) 106; 20
- 1947BL1A Bleuler, Scherrer, Walter and Zunti, *Helv. Phys. Acta* 20 (1947) 96
- 1947FO1A Fowler, Burrows and Curry, *Nature* 159 (1947) 569
- 1948JE03 J.V. Jelley and E.B. Paul, *Proc. Cambridge Phil. Soc.* 44 (1948) 133
- 1950HU72 E.L. Hudspeth, C.P. Swann and N.P. Heydenburg, *Phys. Rev.* 80 (1950) 643
- 1950SC1A A.W. Schardt, W.A. Fowler and C.C. Lauritsen, *Phys. Rev.* 80 (1950) 136; N5
- 1951JO1B H.E. Johns, R.J. Horsley, R.N. Haslam and A. Quinton, *Phys. Rev.* 84 (1951) 856
- 1951JO1C Jones and Wilkinson, *Proc. Phys. Soc. (London)* A64 (1951)756
- 1951MI1B C.H. Millar, G.A. Bartholomew and B.B. Kinsey, *Phys. Rev.* 81 (1951) 150
- 1952AJ38 F. Ajzenberg and T. Lauritsen, *Rev. Mod. Phys.* 24 (1952) 321
- 1952BA1C Barnes, James and Neilson, *Can. J. Phys.* 30 (1952) 717
- 1952FE1A A.J. Ferguson and J.H. Montague, *Phys. Rev.* 87 (1952) 215; Q3
- 1952GO1A Goward and Wilkins, *Proc. Phys. Soc. (London)* A65 (1952) 671
- 1952LI24 A.B. Lillie, *Phys. Rev.* 87 (1952) 716
- 1952SC28 A. Schardt, W.A. Fowler and C.C. Lauritsen, *Phys. Rev.* 86 (1952) 527
- 1952SE1B Seed and French, *Phil. Mag.* 43 (1952) 1214
- 1953BU72 W.E. Burcham, W.M. Gibson, A. Hossain and J. Rotblat, *Phys. Rev.* 92 (1953) 1266
- 1953CH1A Christy, *Phys. Rev.* 89 (1953) 839
- 1953CO1D Cohen and French, *Phil. Mag.* 44 (1953) 1259
- 1953CU1D Cuer and Magnac-Valette, *J. Phys. Rad.* 14 (1953) 15S
- 1953DE1A Devons, *Rept. of Birmingham Conf. on Nucl. Phys.* (1953)
- 1953HI05 R.W. Hill, *Phys. Rev.* 90 (1953) 845
- 1953IN1A Inglis, *Rev. Mod. Phys.* 25 (1953) 390
- 1953JO1A Jones and Wilkinson, *Proc. Phys. Soc. (London)* A66 (1953) 1176

1953KR1B A.A. Kraus, A.P. French, W.A. Fowler and C.C. Lauritsen, Phys. Rev. 89 (1953) 299
1953LI28 M. Lindner and R.N. Osborne, Phys. Rev. 91 (1953) 1501
1953MI31 C.H. Millar and A.G.W. Cameron, Can. J. Phys. 31 (1953) 723
1953MO1B Montalbetti, Katz and Goldemberg, Phys. Rev. 91 (1953) 659
1953NE1B Neilson, James and Barnes, Phys. Rev. 92 (1953) 1084
1953WI1A Wilkinson, Phil. Mag. 44 (1953) 450
1953WI1B Wilkinson, Phys. Rev. 90 (1953) 721
1953WI1E Wilkinson and Jones, Phil. Mag. 44 (1953) 542
1954BI04 M. Birnbaum, Phys. Rev. 93 (1954) 146
1954BI96 J.W. Bittner and R.D. Moffat, Phys. Rev. 96 (1954) 374
1954CA1D Cartwright, Green and Willmott, Phil. Mag. 45 (1954) 742
1954DE1E D.M. Dennison, Phys. Rev. 96 (1954) 378
1954DE36 S. Devons, G. Goldring and G.R. Lindsey, Proc. Phys. Soc. (London) A67 (1954) 134
1954FE16 G.A. Ferguson, J. Halpern, R. Nathans and P.F. Yergin, Phys. Rev. 95 (1954) 776
1954FL1A Florian, Urban and Wildermuth, Z. Naturforsch. A9 (1954) 748
1954GO1F Goldring, Proc. Phys. Soc. (London) A67 (1954) 930
1954JO1F Jongerius, Valckx and Endt, Physica 20 (1954) 29
1954KA1A L. Katz, R.N. Haslam, R.J. Horsley, A.G. Cameron and R. Montalbetti, Phys. Rev. 95 (1954) 464
1954KR1C A.A. Kraus, Phys. Rev. 94 (1954) 975
1954MA97 H.C. Martin, Phys. Rev. 93 (1954) 498
1954RI1B J.A. Rickard, E.L. Hudspeth and W.W. Clendenin, Phys. Rev. 96 (1954) 1272
1954ST1C A.G. Stanley, Phil. Mag. 45 (1954) 807
1954ST89 P. Stoll, Helv. Phys. Acta 27 (1954) 395
1954TH42 L.C. Thompson and J.R. Risser, Phys. Rev. 94 (1954) 941
1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77
1955AL16 H.R. Allan and N. Sarma, Proc. Phys. Soc. (London) A68 (1955) 535
1955BA1P Basile and Schuhl, Compt. Rend. 240 (1955) 2399
1955BE1H Beghian, Hicks and Milman, Phil. Mag. 464 (1955) 924
1955BE62 R.D. Bent, T.W. Bonner and R.F. Sippel, Phys. Rev. 98 (1955) 1237
1955CA1E Campbell, Aust. J. Phys. 8 (1955) 449
1955CO1C Collie, Proc. Glasgow Conf. (Pergamon Press, 1955)

1955DE51 S. Devons, G. Manning and D.St.P. Bunbury, Proc. Phys. Soc. (London) A68 (1955) 18

1955FU1A Fujimoto and Hossain, Phil. Mag. 46 (1955) 542

1955GO1E Gorodetzky, Armbruster, Chevallier, Gallman and Manquenouille, Compt. Rend. 241 (1955) 482, 698; J. Phys. Rad. 16 (1955) 594

1955HA1D Havlicek and Dobovisek, Phys. Rev. 100 (1955) 1355

1955HE1E Herzenberg, Nuovo Cim. 1 (1955) 986, 1008

1955HO68 W.F. Hornyak and R. Sherr, Phys. Rev. 100 (1955) 1409

1955JA18 D.B. James, W. Kubelka, S.A. Heiberg and J.B. Warren, Can. J. Phys. 33 (1955) 219

1955JA1D Jancovici, Compt. Rend. 240 (1955) 1608

1955JO1D S.A. Johansson and B. Forkman, Phys. Rev. 99 (1955) 1031

1955KI43 B.B. Kinsey, Phys. Rev. 99 (1955) 332

1955KO1A Kolar, UCRL 3012 (1955)

1955MA1K W.M. MacDonald, Phys. Rev. 98 (1955) 60

1955MA1L W.M. MacDonald, Phys. Rev. 100 (1955) 51

1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46

1955MO1B H. Morinaga, Phys. Rev. 97 (1955) 1185

1955MO1C H. Morinaga, Phys. Rev. 97 (1955) 444

1955PA50 R.T. Pauli, Ark. Fys. 9 (1955) 571

1955PE1D A.S. Penfold and B.M. Spicer, Phys. Rev. 100 (1955) 1377

1955RA1B Rasmussen, Miller and Sampson, Phys. Rev. 100 (1955) 181

1955RA1E D. Raymond, D. Cooper and Dan J. Zaffarano, Phys. Rev. 98 (1955) 1199, X13

1955SA1F Santos, Goldemberg, Pieroni, Silva, Borelli, Villaca and Leite Lopes, Anais Acad. Brasil. Ciencias 27 (1955) 437

1955SC1B L.I. Schiff, Phys. Rev. 98 (1955) 1281

1955SC36 J. Schmouker, P. Erdos, P. Jordan and P. Stoll, J. Phys. Rad. 16 (1955) 169

1955SP1B B.M. Spicer, Phys. Rev. 99 (1955) 33

1955ST1D W.E. Stephens, A.K. Mann, B.J. Patton and E.J. Winhold, Phys. Rev. 98 (1955) 839

1955TI1A Titterton, Prog. Nucl. Phys. 4 (1955) 1

1955WA1A Wapstra, Physica 21 (1955) 367

1955WI1F D.H. Wilkinson, Phys. Rev. 99 (1955) 1347

1956AL1G Alburger, Rev. Sci. Instrum. 27 (1956) 991

1956BA1A Baumgartner et al., Helv. Phys. Acta 29 (1956) 255

- 1956BA1H Bashkin, Carlson and Jacobs, Bull. Amer. Phys. Soc. 1 (1956) 212; Physica 22 (1956) 1124A
- 1956BO61 T.W. Bonner, A.A. Kraus Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348
- 1956BU22 J.W. Butler, Bull. Amer. Phys. Soc. 1 (1956) 94
- 1956BU95 E.J. Burge, Y. Fujimoto and A. Hossain, Phil. Mag. 1 (1956) 19
- 1956CA1F Cameron, Bull. Amer. Phys. Soc. 1 (1956) 191
- 1956CO59 L. Cohen, A.K. Mann, B.J. Patton, K. Reibel, W.E. Stephens and E.J. Winhold, Phys. Rev. 104 (1956) 108
- 1956DA1C Dawson and Livesey, Can. J. Phys. 34 (1956) 241
- 1956DA23 R.B. Day, Phys. Rev. 102 (1956) 767
- 1956DE22 S. Devons, G. Manning and J.H. Towle, Proc. Phys. Soc. (London) A69 (1956) 173
- 1956DO37 R.A. Douglas, B.R. Gasten and A. Mukerji, Can. J. Phys. 34 (1956) 1097
- 1956EL1C Elliott, Phys. Rev. 101 (1956) 1212
- 1956FE1B R.A. Ferrell and W.M. Visscher, Phys. Rev. 104 (1956) 475
- 1956FE1E R.A. Ferrell and W.M. Visscher, Phys. Rev. 102 (1956) 450
- 1956GO1G Goldanskii, Zh. Eksp. Teor. Fiz. 30 (1956) 969; JETP (Sov. Phys.) 3 (1956) 791
- 1956HA1C Hayakawa, Hayashi, Imoto and Kikuchi, Prog. Theor. Phys. 16 (1956) 507
- 1956HA1D Hayashi and Nishida, Prog. Theor. Phys. 16 (1956) 613
- 1956IS1A Isoya, Goto and Momota, J. Phys. Soc. Jpn. 11 (1956) 899
- 1956JA1C Jancovici, Compt. Rend. 242 (1956) 883
- 1956JO1C Johansson, Physica 22 (1956) 1144A
- 1956JO1D Jones, Weil, Kruse, Baicker and Lidofsky, Bull. Amer. Phys. Soc. 1 (1956) 326
- 1956KA1C Kassecker and Urban, Acta Phys. Aust. 10 (1956) 95
- 1956KA1D S.L. Kameny, Phys. Rev. 103 (1956) 358
- 1956KI54 B.B. Kinsey and T. Stone, Phys. Rev. 103 (1956) 975
- 1956KO26 B. Koudijs, Ph.D. Thesis, Univ. of Utrecht (1956)
- 1956LI1C Lidofsky, Jones, Bent, Weil, Kruse, Bardon and Havens, Bull. Amer. Phys. Soc. 1 (1956) 212
- 1956LI1D Livesey, Can. J. Phys. 34 (1956) 1022
- 1956LI37 J.G. Likely and F.P. Brady, Phys. Rev. 104 (1956) 118
- 1956MO1D H. Morinaga, Phys. Rev. 101 (1956) 254
- 1956PE1A Perring and Skyrme, Proc. Phys. Soc. (London) A69 (1956) 600

1956RE1C P.J. Redmond, Phys. Rev. 101 (1956) 751
 1956SC01 J.P. Schiffer, T.W. Bonner, R.H. Davis and F.W. Prosser Jr., Phys. Rev. 104 (1956) 1064
 1956SQ1A Squires, Bockelman and Buechner, Phys. Rev. 104 (1956) 413
 1956ST30 K. Strauch and F. Titus, Phys. Rev. 104 (1956) 191
 1956TO1A B.J. Toppel, Phys. Rev. 103 (1956) 141
 1956VA17 F.P.G. Valckx, Ph.D. Thesis, Univ. of Utrecht (1956)
 1956WI1A Wilkinson, Toppel and Alburger, Phys. Rev. 101 (1956) 673
 1956WI1D Wilkinson, Phil. Mag. 1 (1956) 379
 1956WI1G Wilkinson, Phil. Mag. 1 (1956) 127
 1956WI1H Wilkinson, Phil. Mag. 1 (1956) 1031
 1956WO1C Wolicki, Holmgren, Johnston and Geer, Bull. Amer. Phys. Soc. 1 (1956) 325
 1956ZI1A Zimmerman, Phys. Rev. 104 (1956) 387
 1957AL39 R. Alphonse, A. Johansson and G. Tibell, Nucl. Phys. 4 (1957) 672
 1957BA03 S. Bashkin and R.R. Carlson, Phys. Rev. 106 (1957) 261
 1957BA1H Barker, Phil. Mag. 2 (1957) 780
 1957BA1K Barker and Mann, Phil. Mag. 2 (1957) 5
 1957BA27 R. Basile, Ann. Phys. 2 (1957) 267
 1957BE61 R.D. Bent and T.H. Kruse, Phys. Rev. 108 (1957) 802
 1957BI75 G.R. Bishop and J.C. Bizot, J. Phys. Rad. 18 (1957) 434
 1957BL01 S.D. Bloom, B.J. Toppel and D.H. Wilkinson, Phil. Mag. 2 (1957) 57
 1957BO04 N.A. Bostrom, E.L. Hudspeth and I.L. Morgan, Phys. Rev. 105 (1957) 1545
 1957BR17 D.A. Bromley, A.J. Ferguson, H.E. Gove, A.E. Litherland and E. Almqvist, Bull. Amer. Phys. Soc. 2 (1957) 51, RA4
 1957BR55 P. Brix and E.K. Maschke, Z. Naturforsch. A12 (1957) 1013
 1957BU66 E.M. Burbidge, G.R. Burbidge, W.A. Fowler and F. Hoyle, Rev. Mod. Phys. 29 (1957) 547
 1957CA1D Carver and Lokan, Aust. J. Phys. 10 (1957) 312
 1957CH32 R.H. Chow and B.T. Wright, Can. J. Phys. 35 (1957) 184
 1957CO1H Corelli, Bleuler and Tendam, Bull. Amer. Phys. Soc. 2 (1957) 34
 1957EL1B Elliott and Flowers, Proc. Roy. Soc. A242 (1957) 57
 1957ER24 P. Erdos, P. Scherrer and P. Stoll, Helv. Phys. Acta 30 (1957) 639
 1957FE1D R.A. Ferrell, Phys. Rev. 107 (1957) 1631

1957FI1C Fischer and Fischer, Bull. Amer. Phys. Soc. 2 (1957) 182
1957FR56 J.M. Freeman and R.C. Hanna, Nucl. Phys. 4 (1957) 599
1957GI14 W.M. Gibson, D.J. Prowse and J. Rotblat, Proc. Roy. Soc. A243 (1957) 237
1957GR1G J.J. Griffin, Phys. Rev. 108 (1957) 328
1957HA98 F.B. Hagedorn, Phys. Rev. 108 (1957) 735
1957HE1B A. Herzenberg, Nucl. Phys. 3 (1957) 1
1957HG01 F.B. Hagedorn and J.B. Marion, Phys. Rev. 108 (1957) 1015
1957HI98 P. Hillman, A. Johansson and H. Tyren, Nucl. Phys. 4 (1957) 648
1957HO1E Hofstadter, Ann. Rev. Nucl. Sci., Vol. 7 (1957)
1957IL01 E.G. Illsley, H.D. Holmgren, R.L. Johnston and E.A. Wolicki, Phys. Rev. 107 (1957) 538
1957JA1B Jannelli and Mezzanares, Nuovo Cim. 5 (1957) 1047
1957JA37 N. Jarmie, J.D. Seagrave et al., LA-2014 (1957)
1957JO1C Johnston and Bostrom, Bull. Amer. Phys. Soc. 2 (1957) 104
1957LI1D Litherland, Almqvist, Bromley, Gove and Ferguson, Bull. Amer. Phys. Soc. 2 (1957) 51
1957MA58 Th.A.J. Maris and H. Tyren, Nucl. Phys. 4 (1957) 662
1957MC35 J.H. McCrary, T.W. Bonner and W.A. Ranken, Phys. Rev. 108 (1957) 392
1957MI1B Milone, Ricamo and Rinzivillo, Nuovo Cim. 5 (1957) 532
1957MI1C Milone and Ricamo, Nuovo Cim. 5 (1957) 1338
1957MO1A Morton and Lewis, Bull. Amer. Phys. Soc. 2 (1957) 286
1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
1957NO1C Nonaka, Morita, Kawai, Ishimatsu, Takeshita, Nakajima and Takano, J. Phys. Soc. Jpn. 12 (1957) 841
1957OW03 G.E. Owen and L. Madansky, Phys. Rev. 105 (1957) 1766; Erratum Phys. Rev. 108 (1957) 1647
1957RE1A Reiner, Physica 23 (1957) 338
1957SA1B E.E. Salpeter, Phys. Rev. 107 (1957) 516
1957SP1A Spicer, Aust. J. Phys. 10 (1957) 326
1957SV1A N.L. Svantesson, Nucl. Phys. 3 (1957) 273
1957SW17 C.P. Swann and F.R. Metzger, Phys. Rev. 108 (1957) 982
1957TA1C G.E. Tauber and T.-Y. Wu, Phys. Rev. 105 (1957) 1772
1957TO1A Touchard, Compt. Rend. 244 (1957) 2499

1957TY36 H. Tyren and Th.A.J. Maris, Nucl. Phys. 4 (1957) 637
 1957VA11 D.M. Van Patter and W. Whaling, Rev. Mod. Phys. 29 (1957) 757
 1957VA1B Vanetsian and Fedchenko, Sov. J. At. Energy 2 (1957) 141
 1957WA01 E.K. Warburton and J.N. McGruer, Phys. Rev. 105 (1957) 639
 1957WA1B Wakatsuki, Hirao, Okada and Miura, J. Phys. Soc. Jpn. 12 (1957) 1178
 1957WE1A Weil, Jones and Lidofsky, Phys. Rev. 108 (1957) 800
 1957WI1B Wilkinson, Phys. Rev. 105 (1957) 686
 1957WI1H D.H. Wilkinson and S.D. Bloom, Phys. Rev. 105 (1957) 683
 1957WI1J H. Wilhelmsson and M. Nilsson, Nucl. Phys. 4 (1957) 234
 1957YO04 T.E. Young, G.C. Phillips and R.R. Spencer, Phys. Rev. 108 (1957) 72
 1958AL13 D.E. Alburger, Phys. Rev. 111 (1958) 1586
 1958AL1D Alkhazov, Gangpskii and Lemberg, JETP (Sov. Phys.) 6 (1958) 892
 1958AR1B Armbuster, Ann. Phys. (France) 3 (1958) 88
 1958BE74 W.L. Bendel, J. McElhinney and R.A. Tobin, Phys. Rev. 111 (1958) 1297
 1958BO18 D.L. Booth, F.V. Price, D. Roaf and G.L. Salmon, Proc. Phys. Soc. (London) A71 (1958) 325
 1958BO71 D. Bodansky, S.F. Eccles, G.W. Farwell, M. Rickey and P.C. Robinson, Bull. Amer. Phys. Soc. 3 (1958) 327, K9
 1958BR1D Bromley, Proc. Rehovoth Conf. (North-Holland Pub. Co., Amsterdam, 1958)
 1958BR95 G. Brunhart, V.P. Kenney and B.C. Kern, Phys. Rev. 110 (1958) 924
 1958CA13 R.R. Carlson, R.A. Douglas, S. Bashkin and C. Broude, Bull. Amer. Phys. Soc. 3 (1958) 199, P3
 1958CA1G Cameron, Bull. Amer. Phys. Soc. 3 (1958) 269
 1958CO59 J.C. Corelli, E. Bleuler and D.J. Tendam, Bull. Amer. Phys. Soc. 3 (1958) 200, P9
 1958DA1E Dabrowski, Proc. Phys. Soc. (London) A71 (1958) 658
 1958DA1F Dabrowski, Proc. Phys. Soc. (London) A72 (1958) 499
 1958EH1B Ehrenberg, Hofstadter, Meyer-Berkhout, Ravenhall and Sobottka, HEPL 149 (1958)
 1958EK36 K.E. Eklund and R.D. Bent, Phys. Rev. 112 (1958) 488
 1958FE1C Ferrell, Bull. Amer. Phys. Soc. 3 (1958) 49
 1958FE1D Ferrell, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)
 1958HA1F R.B. Hall and R.J. Eden, Nucl. Phys. 6 (1958) 157
 1958HE52 D.F. Hebbard, Bull. Amer. Phys. Soc. 3 (1958) 406, F6

1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)

1958JO28 K.W. Jones, L.J. Lidofsky and J.L. Weil, Phys. Rev. 112 (1958) 1252

1958KA1D Katz, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)

1958KN1B Knipper, Ph.D. Thesis, Univ. of Strasbourg (1958)

1958KO63 D. Kohler and H.H. Hilton, Phys. Rev. 110 (1958) 1094

1958LI1D Lindenberger, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)

1958MA1B Th.A.J. Maris, P. Hillman and H. Tyren, Nucl. Phys. 7 (1958) 1

1958MI89 C. Milone, S. Milone-Tamburino, R. Rinzivillo, A. Rubbino and C. Tribuno, Nuovo Cim. 7 (1958) 729

1958MO14 S. Morita, J. Phys. Soc. Jpn. 13 (1958) 126

1958MO17 G. Morpurgo, Phys. Rev. 110 (1958) 721

1958MO1J Morinaga and Johansson, Proc. Rehovoth Conf. (North-Holland Pub. Co., Amsterdam, 1958)

1958PE1A Penner and Leiss, Bull. Amer. Phys. Soc. 3 (1958) 56

1958PH37 W.R. Phillips, Phys. Rev. 110 (1958) 1408

1958RA1F Rapoport and Butusov, JETP (Sov. Phys.) 6 (1958) 1163

1958RA43 D.G. Ravenhall, Rev. Mod. Phys. 30 (1958) 430

1958SI1A Siddig and Haslam, Can. J. Phys. 36 (1958) 963

1958TY49 H. Tyren, P. Hillman and T.A.J. Marris, Nucl. Phys. 7 (1958) 10

1958UM1A Umezawa, Proc. Rehovoth Conf. (North-Holland Pub. Co., Amsterdam, 1958)

1958WE31 J.L. Weil and K.W. Jones, Phys. Rev. 112 (1958) 1975

1958WI1E Wildermuth and Kanellopoulos, Nucl. Phys. 7 (1958) 150

1958WI41 D.H. Wilkinson, Phys. Rev. 109 (1958) 1610

1958WO51 M. Wolff, E.J. Winhold, W.E. Stephens and E.E. Carroll, Bull. Amer. Phys. Soc. 3 (1958) 173, E8; see Halpern in Conf. on Photonucle. Reactions, National Bureau of Standards (1958)

1959AL1M Alburger, Wilkinson and Gallmann, Bull. Amer. Phys. Soc. 4 (1959) 56

1959BR68 D.A. Bromley, H.E. Gove, J.A. Kuehner, A.E. Litherland and E. Almqvist, Phys. Rev. 114 (1959) 758

1959KU78 J.A. Kuehner, A.E. Litherland, E. Almqvist, D.A. Bromley and H.E. Gove, Phys. Rev. 114 (1959) 775

1959ZI18 W. Zimmermann Jr., Phys. Rev. 114 (1959) 867

CO54I Unknown Source

MO57C Unknown Source