

Energy Levels of Light Nuclei $A = 17$

F. Ajzenberg-Selove^a and T. Lauritsen^b

^a *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

^b *California Institute of Technology, Pasadena, California*

Abstract: An evaluation of $A = 5-24$ was published in *Nuclear Physics* 11 (1959), p. 1. This version of $A = 17$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 1, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 17$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{17}\text{N}\$](#) , [\$^{17}\text{O}\$](#) , [\$^{17}\text{F}\$](#)

B. Tables of Recommended Level Energies:

[Table 17.1](#): Energy levels of ^{17}N

[Table 17.2](#): Energy levels of ^{17}O

[Table 17.6](#): Energy levels of ^{17}F

C. [References](#)

D. Figures: [\$^{17}\text{N}\$](#) , [\$^{17}\text{O}\$](#) , [\$^{17}\text{F}\$](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

Table 17.1: Energy levels of ^{17}N

E_x (MeV)	$J^\pi; T$	$\tau_{1/2}$ (sec)	Decay	Reactions
0	$;$ $\frac{3}{2}$	4.14 ± 0.04	β^-	1, 2, 3, 4, 5, 6, 9, 11
1.32 ± 0.08				2
1.89 ± 0.08				2
2.50 ± 0.08				2
2.82 ± 0.08				2
(3.27 ± 0.09)				2
(3.57 ± 0.09)				2
(3.86 ± 0.09)				2
(4.18 ± 0.09)				2

^{17}N
(Fig. 34)

GENERAL:

Mass of ^{17}N : From the Q_0 of the $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ reaction and the Wapstra masses (1955WA1A) for ^{11}B , ^7Li and ^1H , the mass excess ($M - A$) of ^{17}N is 12.93 ± 0.06 MeV.

1. $^{17}\text{N}(\beta^-)^{17}\text{O}^* \rightarrow ^{16}\text{O} + \text{n}$ $Q_m = 4.57$

The decay is complex. See ^{17}O .

2. $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ $Q_m = 8.38$
 $Q_0 = 8.38 \pm 0.06$ (C. Littlejohn, private communication).

At $E(^7\text{Li}) = 2$ MeV, proton groups are observed corresponding to the ground state of ^{17}N and to excited states at 1.32 ± 0.08 , 1.89 ± 0.08 , 2.50 ± 0.08 , 2.82 ± 0.08 , 3.27 ± 0.09 , 3.57 ± 0.09 , 3.86 ± 0.09 , and 4.18 ± 0.09 MeV. The data on the four highest levels are preliminary (C. Littlejohn, private communication).

3. $^{14}\text{C}(\alpha, \text{p})^{17}\text{N}$ $Q_m = -9.75$

See ^{18}O .

4. $^{15}\text{N}(\text{t}, \text{p})^{17}\text{N}$ $Q_{\text{m}} = -0.15$

See (1956SH1A).

5. $^{17}\text{O}(\text{n}, \text{p})^{17}\text{N}$ $Q_{\text{m}} = -7.93$

See (1956FA1C).

6. $^{18}\text{O}(\gamma, \text{p})^{17}\text{N}$ $Q_{\text{m}} = -15.99$

See (1955RE1C).

7. $^{18}\text{O}(\text{n}, \text{d})^{17}\text{N}$ $Q_{\text{m}} = -13.77$

Not reported.

8. $^{18}\text{O}(\text{d}, ^3\text{He})^{17}\text{N}$ $Q_{\text{m}} = -10.50$

Not reported.

9. $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$ $Q_{\text{m}} = 3.82$

See (1956SH1A).

10. $^{19}\text{F}(\text{n}, ^3\text{He})^{17}\text{N}$ $Q_{\text{m}} = -16.24$

Not reported.

11. Photospallation reactions.

A number of spallation reactions yielding ^{17}N have been observed by (1955RE1C) and (1955CH1C). See also (1958BA60).

^{17}O
(Fig. 35)

GENERAL:

Theory: (1956BA1J, 1956KA1C, 1956KI1C, 1956LE1C, 1956SC1E, 1956VI1A, 1957AM1B, 1957AM1C, 1957FA1B, 1957FE1A, 1957PE1D, 1957RA1B, 1957TA1C, 1958BA32, 1958NI1D).

1. $^{13}\text{C}(\alpha, n)^{16}\text{O}$ $Q_m = 2.203$ $E_b = 6.348$

Observed resonances for $E_\alpha = 1$ to 5 MeV are listed in Table 17.3 (1953JO1A, 1954TR09, 1956BE1D, 1956BO61, 1956RU1A, 1957WA1C), together with those observed in $^{16}\text{O}(n, \alpha)^{13}\text{C}$. The observed cross section of $^{13}\text{C}(\alpha, n)^{16}\text{O}$ is closely matched by that calculated from the inverse reaction, for $E_\alpha = 4.0$ to 5.2 MeV (1957WA1C). Assignments for J^π in the range $^{16}\text{O}^*(7.1$ to 9.0 MeV) are derived from angular distributions (1956SC1C, 1957WA1C). The existence of additional broad states underlying this region is indicated by the analysis (1956RU1A, 1957WA1C). Reduced widths have been tabulated for a number of levels by (1957WA1C); it is of interest that θ_n^2 and θ_α^2 generally differ by less than one order of magnitude (1957WA1C). The implications of the present reaction for element synthesis in stars are discussed by (1955FO1C, 1957BU1B, 1957SA1B). See also (1954NA1B, 1956BA1K).

2. $^{14}\text{C}(\alpha, n)^{17}\text{O}$ $Q_m = -1.825$
 $Q_0 = -1.820 \pm 0.002$ (1956SA06).

3. $^{14}\text{N}(\alpha, p)^{17}\text{O}$ $Q_m = -1.197$

Proton groups have been observed to the ground and first excited states of ^{17}O (1951RO1D, 1953HJ1A, 1953HJ1B).

4. $^{15}\text{N}(d, n)^{16}\text{O}$ $Q_m = 9.886$ $E_b = 14.032$

The excitation function has been measured from 0.5 to 5.3 MeV. Above $E_d = 1.0$ MeV pronounced peaks are observed, presumably to be ascribed to numerous overlapping resonances. Angular distributions are generally well accounted for by the exchange stripping theory of (1957OW03), with an admixture of heavy-particle stripping which is almost constant for $E_d > 1$ MeV (1958WE31).

Table 17.2: Energy levels of ^{17}O

E_x in ^{17}O (MeV \pm keV)	J^π	τ_m or Γ (keV)	Decay	Reactions
0	$\frac{5}{2}^+$		stable	2, 3, 17, 21, 22, 23, 28, 29
0.871 ± 4	$\frac{1}{2}^+$	$\tau_m = (2.5 \pm 1) \times 10^{-10}$ sec	γ	3, 17, 28, 29
3.058 ± 6	$(\frac{1}{2}^-)$		γ	17, 28
3.846 ± 5	$(\frac{7}{2}^-)$		γ	17, 28
4.555 ± 5	$\frac{3}{2}^-$	$\Gamma = 40 \pm 3$	n	11, 17, 28
5.083 ± 5	$\frac{3}{2}^+$	90 ± 5	n	11, 17, 28
5.217 ± 5		< 8		17, 28
5.378 ± 7	$\frac{3}{2}^-$	31 ± 4	n	11, 17, 28
5.697 ± 4	$\frac{7}{2}^-$	≤ 7	n	11, 17
5.729 ± 5		< 8		17, 28
5.866 ± 5	$\geq \frac{3}{2}$	≤ 8	n	11, 17, 28
5.940 ± 10	$\frac{1}{2}^-$	26 ± 5	n	11, 17, 28
6.24 ± 20				17, 28
6.38 ± 30	$\frac{1}{2}^+$	130	n	11
6.87 ± 20				17, 28
(6.99 ± 20)				28
7.161 ± 6	$\frac{5}{2}$	2.7 ± 1	n, α	1
7.28 ± 30	$\frac{3}{2}^+$	210	n	11
7.373 ± 6	$\frac{5}{2}$	≤ 2	n, α	1, 28
7.560 ± 10	$\geq \frac{7}{2}$	≤ 4	n, α	1, 17, 28
7.676 ± 7	$\geq \frac{5}{2}$	22	n, α	1
(7.72 ± 80)	$(\frac{3}{2}^-)$	750	n	11
7.94 ± 20	$\frac{1}{2}$	90	n, α	1, 16
8.07 ± 20	$\frac{3}{2}$	75	n, α	1, 11, 16
8.20 ± 20	$\frac{3}{2}$	60	n, α	1, 11, 16
8.27 ± 40				28
8.340 ± 7	$\frac{1}{2}$	≤ 5	n, α	1, 11, 16, 28
8.390 ± 7	$\frac{5}{2}$	8	n, α	1, 11, 16
8.460 ± 7	$\frac{7}{2}$	9	n, α	1, 11, 16
8.493 ± 7	$(\frac{3}{2})$	≤ 6	n, α	1, 11, 16

Table 17.2: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	J^π	τ_m or Γ (keV)	Decay	Reactions
(8.59 \pm 40)				28
8.70 \pm 20	$\frac{3}{2}$	70	n, α	1, 11, 16
8.89 \pm 20	$\frac{3}{2}$	110	n, α	1, 11, 16
8.96 \pm 20	$\frac{7}{2}$	30	n, α	1, 11, 16
9.06 \pm 40				28
9.15 \pm 20		≤ 6	n, α	1, 11
9.20	$\frac{5}{2}$	≤ 4	n, α	1
9.50	$\frac{7}{2}$	11	n, α	1, 11
9.73	$\frac{7}{2}$	19	n, α	1, 11
9.78		55	n, α	1
9.89	$\frac{9}{2}$	12	n, α	1
9.98	$(\frac{5}{2}, \frac{7}{2})$	150	n, α	1
10.21		50	n, α	1, 11
10.5			n	11, 12
10.9			n	11, 12
11.5			n	11, 12
12.1			n	11, 12
15.2			d, p	5
15.8			d, p	5

5. $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$

$$Q_m = 0.267$$

$$E_b = 14.032$$

The excitation curve has been obtained for $E_d = 0.3$ to 2.7 MeV. There is some resonance structure at $E_d = 1.3$ and 1.9 MeV corresponding to $^{17}\text{O}^*(15.2$ and 15.8 MeV) ([1957BO04](#)). See also ^{16}N .

6. $^{15}\text{N}(\text{d}, \alpha)^{13}\text{C}$

$$Q_m = 7.683$$

$$E_b = 14.032$$

See ([1940HO1A](#)).

Table 17.3: Resonances in $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{16}\text{O}(n, \alpha)^{13}\text{C}$

E_α (MeV \pm keV)	E_n	$\Gamma_{\text{c.m.}}$ (keV)	l_α	l_n	J^π	E_x in ^{17}O (MeV)	Refs.
1.063 \pm 4		2.7 \pm 1	3, 2	2, 3	$\frac{5}{2}$	7.161	a
1.340 \pm 4		≤ 2	3, 2	2, 3	$\frac{5}{2}$	7.373	a
1.585 \pm 7		≤ 4			$\geq \frac{7}{2}$	7.560	b
1.736 \pm 5		22			$\geq \frac{5}{2}$	7.676	b
2.080 \pm 20	4.0	90	1, 0		$\frac{1}{2}^\pm$	7.94	c
2.250 \pm 20	4.15	75	2, 1		$\frac{3}{2}^\mp$	8.07	d
2.420 \pm 20	4.25	60	1, 2	2, 1	$\frac{3}{2}^\pm$	8.20	d
2.603 \pm 5	4.46	≤ 5	1, 0		$\frac{1}{2}$	8.34	e
2.675 \pm 5	4.53	8	3, 2		$\frac{5}{2}^\pm$	8.39	f
2.760 \pm 5	4.59	9	3, 4		$\frac{7}{2}^\pm$	8.46	f
2.805 \pm 5	4.62	≤ 6	1, 2		$(\frac{3}{2}^\mp)$	8.493	e
3.08 \pm 15	4.85	70	1, 2		$\frac{3}{2}^\pm$	8.70	e
3.33 \pm 15	5.05	110	2, 1		$\frac{3}{2}^\mp$	8.89	f
3.42 \pm 15	5.13	30	3, 4		$\frac{7}{2}^\pm$	8.96	f
3.67 \pm 15		≤ 6				9.15	g
3.73		≤ 4			$\frac{5}{2}$	9.20	h
4.125		11			$\frac{7}{2}$	9.502	h
4.42		19			$\frac{7}{2}$	9.73	h
(4.50)		55				9.78	h
4.63		12			$\frac{9}{2}$	9.89	h
4.75		150			$\frac{5}{2}, \frac{7}{2}$	9.98	h
5.05		50				10.21	h

a (1953JO1A, 1956RU1A, 1957WA1C).

b (1956RU1A, 1957WA1C).

c (1955SE1A, 1956BO61, 1956RU1A, 1956SC1C, 1957WA1C).

d (1954TR09, 1955SE1A, 1956BE1D, 1956BO61, 1956RU1A, 1956SC1C, 1957WA1C).

e (1956BE1D, 1956BO61, 1956SC1C, 1957WA1C).

f (1954TR09, 1956BE1D, 1956BO61, 1956SC1C, 1957WA1C).

g (1956BO61).

h (1956BO61, 1956SC1C).

7. $^{15}\text{N}(t, n)^{17}\text{O}$ $Q_m = 7.774$

Not reported.

8. $^{15}\text{N}(^3\text{He}, p)^{17}\text{O}$ $Q_m = 8.539$

Not reported.

9. $^{15}\text{N}(\alpha, d)^{17}\text{O}$ $Q_m = -9.812$

Not reported.

10. $^{16}\text{O}(n, \gamma)^{17}\text{O}$ $Q_m = 4.146$
 $\sigma_{\text{capt.}} < 0.2 \text{ mb}$ (1955HU1B).

11. $^{16}\text{O}(n, n)^{16}\text{O}$ $E_b = 4.146$

The cross section is almost constant at 3.76 b from thermal energies to $E_n = 0.3 \text{ MeV}$ (1958HU18). Observed resonances are indicated in Table 17.4 (1952BA1D, 1955OK1A, 1956BE1D, 1957WA1C, 1958HU18, 1958LA09, 1958ST28). (1958FO67) have measured angular distributions at six energies in the range $E_n = 0.7$ to 2.2 MeV and have derived the phase shifts up to 2.4 MeV. From measurements in the range $E_n = 0.1$ to 1.7 MeV, (1958LA09) find that the $E_n = 1.65 \text{ MeV}$ resonance is due to a $\frac{7}{2}^-$ state. Phase shifts for $E_n = 0.2$ to 1.4 MeV derived by (1958ST28) from total cross section measurements are in good agreement with those of (1958FO67) except for a 5° discrepancy in $\delta_{\frac{3}{2}^+}$ at $E_n = 0.73 \text{ MeV}$. According to (1958ST28), the s-wave shifts from 0.2 to 2.3 MeV fit hard-sphere scattering with $R = 5.6 \times 10^{-13} \text{ cm}$. (1958FO67) find that the s- and d-wave phase shifts and the locations of $^{17}\text{O}(\text{g.s.}; d_{\frac{5}{2}})$, $^{17}\text{O}^*(0.87; s_{\frac{1}{2}})$ and $^{17}\text{O}^*(5.08; d_{\frac{3}{2}})$ can be adequately described by a potential well 40 – 45 MeV deep, with a diffuse boundary. See also (1956VIIA).

The angular distribution has also been measured at $E_n = 5 \text{ MeV}$ (1958HI68). Polarization studies have been carried out for $E_n = 0.25$ to 0.6 MeV by (1954WI42, 1955OK1A) and for $E_n = 0.7$ to 1.4 MeV by (1958ST28). Cross sections at $E_n = 14 \text{ MeV}$ are tabulated by (1958HU18). See also (1956BA1K, 1956FL1B, 1957MC1B) and (1955AJ61).

12. $^{16}\text{O}(n, n')^{16}\text{O}^*$ $E_b = 4.146$

Table 17.4: Resonances in $^{16}\text{O}(n, n)^{16}\text{O}$

E_n (MeV \pm keV)	Γ_{lab} (keV)	l_n	J^π	E_x in ^{17}O (MeV)	Refs.
0.433 \pm 4	41 \pm 3	1	$\frac{3}{2}^{\frac{3}{2}-}$	4.554	(1955OK1A, 1958HU18)
0.998 \pm 8	96 \pm 5	2	$\frac{3}{2}^{\frac{3}{2}+}$	5.085	(1958HU18, 1958LA09, 1958ST28)
1.310 \pm 8	33 \pm 4	1	$\frac{3}{2}^{\frac{3}{2}-}$	5.379	(1958HU18, 1958ST28)
1.651 \pm 6	\leq 7	\geq 1	$\frac{7}{2}^{\frac{7}{2}-}$	5.700	(1958HU18, 1958LA09)
1.830 \pm 6	\leq 8	\geq 1	$\geq \frac{3}{2}^{\frac{3}{2}}$	5.868	(1958HU18)
1.903 \pm 11	28 \pm 5	1	$\frac{1}{2}^{\frac{1}{2}-}$	5.937	(1952BA1D, 1958HU18)
2.370 \pm 20	140	0	$\frac{1}{2}^{\frac{1}{2}+}$	6.377	(1952BA1D, 1958HU18)
3.330 \pm 30	220	2	$\frac{3}{2}^{\frac{3}{2}+}$	7.28	(1952BA1D, 1958HU18)
3.750 \pm 20	$<$ 100			7.68	(1957WA1C, 1958HU18)
(3.800)	800	1	$(\frac{3}{2}^{\frac{3}{2}-})$	(7.72)	(1952BA1D, 1958HU18)
4.190 \pm 20	$<$ 100			8.09	(1957WA1C, 1958HU18)
4.320 \pm 20	$<$ 100			8.21	(1952BA1D, 1957WA1C, 1958HU18)
4.440 \pm 20	$<$ 100			8.32	(1956BE1D, 1957WA1C, 1958HU18)
4.53			$\geq \frac{3}{2}^{\frac{3}{2}}$	8.41	(1956BE1D)
4.59				8.47	(1956BE1D)
4.62			$\geq \frac{3}{2}^{\frac{3}{2}}$	8.49	(1956BE1D)
4.85				8.71	(1956BE1D)
5.08				8.93	(1956BE1D)
5.13				8.97	(1956BE1D)
5.37				9.20	(1956BE1D)
5.7				9.5	(1958HU18)
5.9				9.7	(1958HU18)
6.4				10.2	(1958HU18)
6.8				10.5	(1958HU18)
7.2				10.9	(1958HU18)
7.8				11.5	(1956BE1D, 1958HU18)
8.4				12.1	(1956BE1D, 1958HU18)

The yield of γ -rays shows marked structure in the range $E_n = 7 - 10$ MeV (1958HU18). At $E_n = 14.1$ MeV, the inelastic scattering cross section is 0.5 b (1953CO1E).

$$13. \text{}^{16}\text{O}(n, 2n)\text{}^{15}\text{O} \quad Q_m = -15.655 \quad E_b = 4.146$$

See (1955AJ61).

$$14. \text{}^{16}\text{O}(n, p)\text{}^{16}\text{N} \quad Q_m = -9.619 \quad E_b = 4.146$$

The cross section rises from threshold to a peak of 89 ± 30 mb at $E_n = 14$ MeV, falling to ≈ 60 mb at 18 MeV (1954MA1B). See also (1955AJ61) and (1953KI1A, 1956FA1C, 1958HU18).

$$15. \text{}^{16}\text{O}(n, d)\text{}^{15}\text{N} \quad Q_m = -9.884 \quad E_b = 4.146$$

See (1953KI1A, 1955HU1B).

$$16. \text{}^{16}\text{O}(n, \alpha)\text{}^{13}\text{C} \quad Q_m = -2.203 \quad E_b = 4.146$$

The cross section has been measured from threshold to 4.2 MeV by (1955SE1A) and from 4.0 to 5.2 MeV by (1957WA1C). All of the ^{16}O levels observed in $^{13}\text{C}(\alpha, n)^{16}\text{O}$ from $^{17}\text{O}^*(7.92$ to 8.96 MeV) are also observed in this reaction, with good agreement in level position: see Table 17.3. Observed cross sections correspond closely to those calculated from the inverse reaction (1957WA1C). At higher energies, using continuous fast neutrons, (1956KI1B) report 18 levels in ^{17}O in the range $E_x = 6.8$ to 9.7 MeV, and (1953GI1B) reports 32 levels in the range $E_x = 6.8$ to 12.8 MeV. See also (1953KI1A, 1955HU1B, 1955HU1C, 1956BA1K).

$$17. \text{}^{16}\text{O}(d, p)\text{}^{17}\text{O} \quad Q_m = 1.919$$

$$Q_0 = 1.918 \pm 0.004 \text{ (1957BR82);}$$

$$Q_0 = 1.915 \pm 0.010 \text{ (1954SP01);}$$

$$Q_0 = 1.885 \text{ (1955KH35).}$$

^{17}O levels derived from observed proton groups are listed in Table 17.5 (1951BU1A, 1957BR82: see also (1953KH1A, 1954SP01, 1955KH35)), together with those from the α -groups from $^{19}\text{F}(d, \alpha)^{17}\text{O}$. The J^π assignments are derived from stripping analysis: see (1955AJ61, 1956GR37,

Table 17.5: States of ^{17}O from $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ and $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$

E_x^a (MeV \pm keV)	$\Gamma_{\text{c.m.}}^a$ (keV)	l_n^b	J^π^b	$\frac{\Lambda^c}{2J+1}$	E_x^d (MeV \pm keV)	E_x^e (MeV \pm keV)
0	< 8	2	$\frac{5}{2}^+$	72	0	0
0.871 ± 4	< 8	0	$\frac{1}{2}^+$	68	0.870 ± 20	0.883 ± 11
3.055 ± 4	< 8	iso.	$(\frac{1}{2}^-)$		3.060 ± 30	3.069 ± 10
3.846 ± 5	< 8	3	$(\frac{7}{2}^-)$	15	3.850 ± 30	3.856 ± 11
4.553 ± 6	40 ± 5	1	$\frac{3}{2}^-$	2.2	4.580 ± 20	4.567 ± 14
5.083 ± 10	95 ± 5	2	$\frac{3}{2}^+$	25	5.070 ± 20	
5.215 ± 5	< 8					5.229 ± 13
5.378 ± 7	28 ± 7				5.310 ± 20	5.397 ± 14
5.695 ± 5	< 8					
5.731 ± 5	< 8				5.760 ± 20	5.723 ± 14
5.866 ± 5	< 8					5.875 ± 15
5.940 ± 15	23 ± 10					5.947 ± 15
					6.240 ± 20	
					6.890 ± 30	6.869 ± 14
						(6.986 ± 15)
						(7.371 ± 15)
					7.510 ± 30	
					8.270 ± 40	
					(8.590 ± 40)	
					9.060 ± 40	

^a $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$: (1957BR82); $E_d = 6.5$ to 7.5 MeV. No other proton groups appear with intensities greater than 0.1 of 4.55 MeV group.

^b $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$: (1956GR37); $E_d = 9$ MeV.

^c (1956GR37); relative capture probabilities.

^d $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ and $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$: (1951BU1A); $E_d = 7.9$ MeV.

^e $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$: (1952WA1A); $E_d = 1.8$ MeV.

1958RI1A). It is noted that the relatively large capture probabilities corresponding to $^{17}\text{O}^*(0, 0.87, 3.8, 5.08)$ are consistent with the assumption that these are relatively pure single particle states formed from an ^{16}O core with 1d, 2s, 1f and 1d neutrons, respectively. The states at $E_x = 3.06$ and 4.55 MeV must be presumed to arise from more complicated configurations (1956GR37).

An earlier report that the 871 keV level might be a close doublet (1956DO41) is not confirmed by (1957BR82, 1957MA1C, 1957MO1B). Q for this level is 1049.0 ± 2.2 keV (1957MO1B), 1048 ± 2 keV (1956DO41), 1047 ± 4 keV (1957BR82). The γ -ray energy is 870.5 ± 2.0 keV (1952TH24), 869 ± 3 keV (1955MA36); the internal conversion coefficient is consistent with E2 (1952TH24). The lifetime is $\tau_m = (2.5 \pm 1) \times 10^{-10}$ sec (1953TH14): this value indicates a considerable enhancement of the single particle transition rate. Possible models are discussed by (1956BA1J, 1956KA1D, 1956SC1E, 1957AM1B, 1957AM1C, 1957FA1B, 1957RA1B, 1958BA32).

Observed widths and shapes of $^{17}\text{O}^*(4.55$ and 5.08 MeV) are in good accord with $^{16}\text{O}(n, n)^{16}\text{O}$ results (1957BR82). The ratio of ground state reduced widths is 1.30 (1956CA1D). See also (1955AJ61, 1955AL1D, 1955KH31, 1956EL1A), (1956IV1A; theor.) and ^{18}F .

18. $^{16}\text{O}(t, d)^{17}\text{O}$ $Q_m = -2.113$

Not reported.

19. $^{16}\text{O}(\alpha, ^3\text{He})^{17}\text{O}$ $Q_m = -16.436$

Not reported.

20. $^{17}\text{N}(\beta^-)^{17}\text{O}^* \rightarrow ^{16}\text{O} + n$ $Q_m = 4.57$

The half-life is 4.14 ± 0.04 sec (1948KN1A), 4.15 ± 0.1 sec (1951ST1E), 4.16 ± 0.15 sec (1956FA1C). $E_\beta(\text{max}) = 3.7 \pm 0.2$ MeV (1949AL04): $\log ft = 3.8$. The β -decay proceeds to an excited state or states of ^{17}O which in turn decay to ^{16}O by neutron emission: the neutron spectrum has a maximum at 0.92 ± 0.07 MeV (1949AL04, 1949HA1B) and a half-width < 0.5 MeV. See also (1957JO1D; theor.).

21. $^{17}\text{O}(p, p')^{17}\text{O}^*$

See (1954WA1A).

22. $^{17}\text{F}(\beta^+)^{17}\text{O}$ $Q_m = 2.767$

See ^{17}F .

23. $^{18}\text{O}(\text{p}, \text{d})^{17}\text{O}$ $Q_m = -5.842$

See (1956TS1A).

24. $^{18}\text{O}(\text{d}, \text{t})^{17}\text{O}$ $Q_m = -1.810$

Not reported.

25. $^{18}\text{O}(\text{}^3\text{He}, \alpha)^{17}\text{O}$ $Q_m = 12.509$

Not reported.

26. $^{19}\text{F}(\text{n}, \text{t})^{17}\text{O}$ $Q_m = -7.548$

Not reported.

27. $^{19}\text{F}(\text{p}, \text{}^3\text{He})^{17}\text{O}$ $Q_m = -8.313$

Not reported.

28. $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$ $Q_m = 10.038$

Observed alpha groups are displayed in Table 17.5 (1951BU1A, 1952WA1A). At $E_d = 1.2$ MeV, two γ -rays are observed with $E_\gamma = 2.2$ MeV ($3.06 \rightarrow 0.87$) and 3.81 MeV. The absence of the direct ground state decay of the 3.06 MeV state is consistent with $J = \frac{1}{2}$ (1956GO1Q, 1957MU1A).

29. $^{20}\text{Ne}(\text{n}, \alpha)^{17}\text{O}$ $Q_m = -0.608$

See ^{21}Ne .

^{17}F
(Fig. 36)

GENERAL:

Theory: See (1956KA1C, 1958BA32).

1. $^{17}\text{F}(\beta^+)^{17}\text{O}$ $Q_m = 2.767$

The decay proceeds to the ground state of ^{17}O : $E_{\beta^+}(\text{max}) = 1.748 \pm 0.006$ MeV. The spectrum has the allowed shape down to 570 keV, and an upper limit of 1% is placed on possible transitions to the 875 keV state of ^{17}O (1954WO20). The weighted mean value of the half-life from the determinations of (1949BR27, 1951LA1A, 1954KO54, 1954WA1A, 1954WO20, 1958AR15) is 66.0 ± 0.3 sec. $\text{Log } ft = 3.38$. See also (1958BE1G; theor.).

2. $^{14}\text{N}(\alpha, n)^{17}\text{F}$ $Q_m = -4.747$
 $Q_0 = -4.76 \pm 0.07$ (1956DO1C).

Slow neutron threshold and proton recoil measurements indicate an excited state at 0.53 ± 0.04 MeV (1956DO1C).

3. $^{15}\text{N}({}^3\text{He}, n)^{17}\text{F}$ $Q_m = 4.989$

Not observed.

4. $^{16}\text{O}(\text{p}, \gamma)^{17}\text{F}$ $Q_m = 0.596$

Non-resonant capture has been studied for $E_p = 0.8$ to 2.1 MeV. This work indicates, in addition to the direct ground-state transition (γ_1), a transition (γ_2) to a state of ^{17}F at 487 ± 10 keV which then radiates (γ_3) to the ground state. The ratio of γ_2 to γ_1 is ≈ 10 over the energy region studied. The γ_1 and γ_3 radiations are approximately isotropic, while γ_2 has an almost pure ($\sin^2 \theta$) distribution ($E_p = 1.0$ to 1.9 MeV). At $E_p = 1.35$ MeV, the cross section for production of γ_2 radiation is 6 ± 3 μb . The fact that most transitions involve the 0.50 MeV state (γ_2) indicates

Table 17.6: Energy levels of ^{17}F

E_x (MeV)	J^π	τ (sec) or Γ (keV)	Decay	Reactions
0	$(\frac{5}{2}^+)$	$\tau_{1/2} = 66.0 \pm 0.3$	β^+	1, 2, 4, 8, 13
0.500 ± 0.003	$\frac{1}{2}^+$	$\tau_m = (2.9 \pm 0.7) \times 10^{-10}$	γ	2, 4, 8
3.10	$\frac{1}{2}^-$	18.7	p	6, 8
3.86	$\frac{7}{2}^-$	< 3.3	p, γ	4, 6
4.4	$(\frac{3}{2}^-)$	≈ 400	p	6
4.7	$(\frac{3}{2}^+)$	(300)	p	6, 8
(5.1)	$(\frac{1}{2}^+)$	190	p	6
5.45		≈ 40	p	6
5.66		≈ 40	p	6
5.83		≈ 40	p	6
(6.15)		< 25	p	6
(6.6)	$(\frac{1}{2}^+)$	140	p	6
(6.75)		< 25	p	6
(6.90)		< 25	p	6

that the non-resonant yield is not to be attributed to this state; it is suggested that direct, one-stage capture is involved (1954WA1A).

The relative ^{17}F activity has been measured from $E_p = 1.1$ to 4.1 MeV by (1951LA1A). The cross section increases almost linearly with proton energy from 1.1 to 3.75 MeV, except for a sharp resonance at 3.47 MeV. There is some indication of a broad resonance at 3.99 MeV (1951LA1A, 1951LA1B).

For $E_p = 140$ to 170 keV, the cross section varies from 0.46 to 2.34×10^{-10} b (1958HE57). At $E_p = 616$ keV, $\sigma = 0.29 \pm 0.03 \mu\text{b}$; both the magnitude of the cross section and the character of the excitation function from $E_p = 275$ to 616 keV are consistent with a direct capture process (1959TA03). See also (1953WI1A, 1957BU1B, 1958LI1A, 1959CH1F).

The excitation energy of the first level of ^{17}F is 495 ± 15 keV (1958HO97), 505 ± 10 keV (1957LE28). The lifetime of $^{17}\text{F}^*(0.5)$ is $\tau_m = (2.5 \pm 0.7) \times 10^{-10}$ sec (1957LE28), $(3.5 \pm 1.5) \times 10^{-10}$ sec (1958HO97). Using the weighted mean, 2.9×10^{-10} sec, the ratio $\tau_m(^{17}\text{F}^*(0.5))/\tau_m(^{17}\text{O}^*(0.87)) = 1.2$ (1958HO97); see $^{16}\text{O}(d, p)^{17}\text{O}$ and (1958BA32).

5. $^{16}\text{O}(p, n)^{16}\text{F}$

$$Q_m = -16.41$$

$$E_b = 0.596$$

See (1958TA03).

6. $^{16}\text{O}(p, p)^{16}\text{O}$

$$E_b = 0.596$$

Observed anomalies in the scattering are exhibited in Table 17.7. The low energy scattering data require a large s-wave phase shift, probably reflecting the bound level at $^{17}\text{F}^*(0.55)$, and a relatively large d-wave shift (1951LA1B, 1953EP1A). A study of proton polarization for $E_p = 2.5$ to 3.0 MeV confirms the assignment $J = \frac{1}{2}^-$ to $^{17}\text{F}^*(3.10)$ and excludes $\frac{1}{2}^+$, $\frac{3}{2}^-$ (1957SO1B). A phase shift analysis for $E_p = 3$ to 7 MeV requires four broad resonances corresponding to $^{17}\text{F}^*(4.5)$ ($P_{\frac{3}{2}}$), $^*(4.6)$ ($D_{\frac{3}{2}}$), $^*(5.15)$ ($S_{\frac{1}{2}}$) and $^*(6.65)$ MeV ($S_{\frac{1}{2}}$). Eight additional sharp resonances are found in this region (1954SE1B: see Table 17.7): however, the curves of (1956HE1B) show no evidence of fine structure from $E_p = 2.5$ to 5.5 MeV beyond that indicated in column A of Table 17.7. See also $^{16}\text{O}(d, n)^{17}\text{F}$.

Table 17.7: Levels of ^{17}F from $^{16}\text{O}(p, p)^{16}\text{O}$

A			B				$^{17}\text{F}^*$ (MeV)
E_p (MeV)	Γ (keV)	J^π	E_p (MeV)	Γ (keV)	J^π	θ_p^2 (%)	
2.66	19.9	$\frac{1}{2}^-$					3.10
3.47	< 3.5	$\frac{7}{2}^-$	3.50	< 25			3.86
(3.99)	≈ 500	$(\frac{3}{2}^+)$	4.1	400	$\frac{3}{2}^-$	20	4.4
(4.39)	240	$(\frac{3}{2}^-)$	4.2	350	$\frac{3}{2}^+$	32	4.7
			4.70	< 25			(5.05)
			4.8	200	$\frac{1}{2}^+$	5.5	5.1
			4.95	< 25			(5.30)
5.16	60		5.20	< 25			5.45
5.38	50		5.40	< 25			5.66
5.56	60		5.60	< 25			5.83
			5.85	< 25			(6.15)
			6.50	< 25			(6.75)
			6.4	150	$\frac{1}{2}^+$	3.7	(6.6)
			6.65	< 25			(6.90)

A: (1951LA1A, 1951LA1B, 1953EP1A, 1956HE1B).

B: (1954SE1A, 1954SE1B).

Scattering at $E_p > 10$ MeV appears to be dominated by optical model effects: see ^{16}O .

7. $^{16}\text{O}(p, \alpha)^{13}\text{N}$

$$Q_m = -5.208$$

$$E_b = 0.596$$

See (1958WH34).

8. $^{16}\text{O}(\text{d}, \text{n})^{17}\text{F}$ $Q_{\text{m}} = -1.631$
 $E_{\text{thresh.}} = 1.830 \pm 0.004$; $Q_0 = -1.626 \pm 0.004$ (1955MA85).

Neutron groups have been observed corresponding to $^{17}\text{F}^*(0, 0.536 \pm 0.010)$ (1951AJ1B), 0.53 ± 0.06 (1951EL1A, 1953MI10), 3.03 ± 0.06 , and 4.74 ± 0.07 MeV (1957GR1A)). Angular distributions of the n_0 and n_1 groups, analyzed by stripping theory, indicate $J = \frac{3}{2}^+$ or $\frac{5}{2}^+$ for the ground state and $J = \frac{1}{2}^+$ for the first excited state (1951AJ1B, 1951EL1A, 1953MI10). The ratio of the reduced widths of the ^{17}F and ^{17}O ground states is 1.30 (1956CA1D). The width of the neutron group corresponding to $^{17}\text{F}^*(4.74)$ is < 90 keV, considerably smaller than is reported in $^{16}\text{O}(\text{p}, \text{p})^{16}\text{O}$ (1957GR1A).

Slow neutron thresholds have been observed corresponding to the ground and first excited states of ^{17}F : $Q_1 = -2.125 \pm 0.004$ MeV corresponding to $^{17}\text{F}^*(499 \pm 3 \text{ keV})$ (1955MA85). See also (1958BE03, 1958GO77).

9. $^{16}\text{O}(^3\text{He}, \text{d})^{17}\text{F}$ $Q_{\text{m}} = -4.898$

Not reported.

10. $^{16}\text{O}(\alpha, \text{t})^{17}\text{F}$ $Q_{\text{m}} = -19.217$

Not reported.

11. $^{17}\text{O}(\text{p}, \text{n})^{17}\text{F}$ $Q_{\text{m}} = -3.550$

Not reported.

12. $^{17}\text{O}(^3\text{He}, \text{t})^{17}\text{F}$ $Q_{\text{m}} = -2.785$

Not reported.

13. $^{19}\text{F}(\gamma, 2\text{n})^{17}\text{F}$ $Q_{\text{m}} = -19.582$

See ^{19}F .

14. $^{19}\text{F}(\text{p}, \text{t})^{17}\text{F}$ $Q_{\text{m}} = -11.098$

Not reported.

15. $^{20}\text{Ne}(\text{p}, \alpha)^{17}\text{F}$ $Q_{\text{m}} = -4.158$

Not reported.

References

(Closed 01 December 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1940HO1A M.G. Holloway and B.L. Moore, *Phys. Rev.* 58 (1940) 847
- 1948KN1A Knable, Lawrence, Leith, Moyer and Thornton, *Phys. Rev.* 74 (1948) 1217A
- 1949AL04 L.W. Alvarez, *Phys. Rev.* 75 (1949) 1127
- 1949BR27 H. Brown and V. Perez-Mendez, *Phys. Rev.* 75 (1949) 1286A
- 1949HA1B E. Hayward, *Phys. Rev.* 75 (1949) 917
- 1951AJ1B F. Ajzenberg, *Phys. Rev.* 83 (1951) 693
- 1951BU1A Burrows, Powell and Rotblat, *Proc. Roy. Soc.* A209 (1951) 478
- 1951EL1A El-Bedewi, Middleton and Tai, *Proc. Phys. Soc. (London)* A64 (1951) 757
- 1951LA1A Laubenstein, Laubenstein, Koester and Mobley, *Phys. Rev.* 84 (1951) 2
- 1951LA1B Laubenstein and Laubenstein, *Phys. Rev.* 84 (1951) 18
- 1951RO1D R.R. Roy, *Phys. Rev.* 82 (1951) 227
- 1951ST1E W.E. Stephens, J. Halpern and R. Sher, *Phys. Rev.* 82 (1951) 511
- 1952BA1D Baldinger, Huber and Proctor, *Helv. Phys. Acta* 25 (1952) 142
- 1952TH24 R.G. Thomas and T. Lauritsen, *Phys. Rev.* 88 (1952) 969
- 1952WA1A Watson and Buechner, *Phys. Rev.* 88 (1952) 1324
- 1953CO1E J.P. Conner, *Phys. Rev.* 89 (1953) 712
- 1953EP1A Eppling, Cameron, Davis, Divatia, Galonsky, Goldberg and Hill, *Phys. Rev.* 91 (1953) 438A
- 1953GI1B V. Gierke, *Z. Naturforsch.* A8 (1953) 567
- 1953HJ1A E. Hjalmar and H. Slatis, *Phys. Rev.* 89 (1953) 1151
- 1953HJ1B Hjalmar and Slatis, *Ark. Fysik* 6 (1953) 451
- 1953JO1A Jones and Wilkinson, *Proc. Phys. Soc. (London)* A66 (1953) 1176
- 1953KH1A Khromchenko, *Dokl. Akad. Nauk. SSSR* 93 (1953) 451
- 1953KI1A Kimura, Ishiwara, Sakisaka, Kumabe, Yamashita and Miyake, *Bull. Inst. Chem. Res., Kyoto Univ.* 31 (1953) 204

1953MI10 R. Middleton, F.A. El Bedewi and C.T. Tai, Proc. Phys. Soc. (London) A66 (1953) 95
1953TH14 J. Thirion and V.L. Telegdi, Phy. Rev. 92 (1953) 1253
1953WI1A Wilkinson, Phil. Mag. 44 (1953) 450
1954KO54 M.I. Korsunskii, Usp. Fiz. Nauk 52 (1954) 3
1954MA1B Martin, Phys. Rev. 93 (1954) 498
1954NA1B Nahmias and Savel, Compt. Rend. 239 (1954) 761
1954SE1A Sempert, Schneider and Martin, Helv. Phys. Acta 27 (1954) 313
1954SE1B Sempert, Ph.D. Thesis, Zurich ETH (1954)
1954SP01 A. Sperduto, W.W. Buechner, C.K. Bockelman and C.P. Browne, Phys. Rev. 96 (1954) 1316
1954TR09 R.E. Trumble, Jr., Phys. Rev. 94 (1954) 748A
1954WA1A Warren, Laurie, James and Erdman, Can. J. Phys. 32 (1954) 563
1954WI42 H.B. Willard, J.K. Bair and J.D. Kington, Phys. Rev. 95 (1954) 1359
1954WO20 C. Wong, Phys. Rev. 95 (1954) 765
1955AJ61 F. Ajzenberg and T. Lauritsen, Revs. Mod. Phys. 27 (1955) 77
1955AL1D Alba, Fernandez, Mazari, Serment and Vazquez, Rev. Mex. Fis. 4 (1955) 207
1955CH1C Chupp, UCRL 2955 (1955)
1955FO1C Fowler, Burbidge and Burbidge, Astro. J. 122 (1955) 271
1955HU1B Hughes and Harvey, BNL-325 (1955)
1955HU1C Huber and Seitz, U. N. A/Conf. 8/P/1033 (1955)
1955KH31 L.M. Khromchenko and V.A. Blinov, Zh. Eksp. Teor. Fiz. 28 (1955) 741; JETP (Sov. Phys.) 1 (1955) 596
1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252
1955MA36 R.J. Mackin, Jr., W.B. Mims and W.R. Mills, Jr., Phys. Rev. 98 (1955) 43
1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46
1955OK1A Okazaki, Phys. Rev. 99 (1955) 55
1955RE1C D. Reagan, Phys. Rev. 100 (1955) 113
1955SE1A Seitz and Huber, Helv. Phys. Acta 28 (1955) 227
1955WA1A Wapstra, Physica 21 (1955) 367
1956BA1J Barker, Phil. Mag. 1 (1956) 329
1956BA1K Barschall, Helv. Phys. Acta 29 (1956) 145
1956BE1D Becker and Barschall, Phys. Rev. 102 (1956) 1384

1956BO61 T.W. Bonner, A.A. Kraus, Jr., J.B. Marion and J.P. Schiffer, Phys. Rev. 102 (1956) 1348

1956CA1D Calvert, Jaffe and Maslin, Phys. Rev. 101 (1956) 501

1956DO1C Doyle and Robbins, Phys. Rev. 101 (1956) 1056

1956DO41 R.A. Douglas, J.W. Broer, R. Chiba, D.F. Herring and E.A. Silverstein, Phys. Rev. 104 (1956) 1059

1956EL1A El-Bedewi, Proc. Phys. Soc. (London) A69 (1956) 221

1956FA1C Faull, AERE R/R 1919 (1956); Phys. Abs. 7393 (1956)

1956FL1B Flerov and Talyzin, Sov. J. Nucl. Energy 4 (1956) 617

1956GO1Q Gorodetzky, Muller and Port, Physica 22 (1956) 1159A

1956GR37 T.S. Green and R. Middleton, Proc. Phys. Soc. (London) A69 (1956) 28

1956HE1B Henry, Phillips, Reich and Russell, Bull. Amer. Phys. Soc. 1 (1956) 96

1956IV1A Ivash, Bull. Amer. Phys. Soc. 1 (1956) 84

1956KA1C Kassecker and Urban, Acta Phys. Aust. 10 (1956) 95

1956KA1D S.L. Kameny, Phys. Rev. 103 (1956) 358

1956KI1B Kimura, Kumabe, Miyake, Ogata and Miyasita, J. Phys. Soc. Jpn. 11 (1956) 1211

1956KI1C L.S. Kisslinger, Phys. Rev. 104 (1956) 1077

1956LE1C Levintov, Zh. Eksp. Teor. Fiz. 30 (1956) 987; JETP (Sov. Phys.) 3 (1956) 796

1956RU1A Rusbridge, Proc. Phys. Soc. (London) A69 (1956) 830

1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434

1956SC1C Schiffer, Kraus and Risser, Phys. Rev. 105 (1956) 1811

1956SC1E Schwartz and Gartenhaus, Bull. Amer. Phys. Soc. 1 (1956) 383

1956SH1A Sher and Floyd, Phys. Rev. 102 (1956) 242

1956TS1A Tsai, Bull. Amer. Phys. Soc. 1 (1956) 327

1956VI1A Visscher and Ferrell, Bull. Amer. Phys. Soc. 1 (1956) 16

1957AM1B Amado and Blin-Stoyle, Proc. Phys. Soc. (London) A70 (1957) 532

1957AM1C R.D. Amado, Phys. Rev. 108 (1957) 1462

1957BO04 N.A. Bostrom, E.L. Hudspeth and I.L. Morgan, Phys. Rev. 105 (1957) 1545

1957BR82 C.P. Browne, Phys. Rev. 108 (1957) 1007

1957BU1B Burbidge, Burbidge, Fowler and Hoyle, Revs. Mod. Phys. 29 (1957) 547

1957FA1B Fallieros and Ferrell, Bull. Amer. Phys. Soc. 2 (1957) 26

1957FE1A Feingold, Bull. Amer. Phys. Soc. 2 (1957) 392

1957GR1A Grismore and Parkinson, *Rev. Sci. Instrum.* 28 (1957) 245
 1957JO1D G.A. Jones and F. Mandl, *Nucl. Phys.* 4 (1957) 690
 1957LE28 P. Lehmann, A. Leveque, T. Grjebine, J.-L. Picou and R. Barloutaud, *Compt. Rend.* 245 (1957) 2259
 1957MA1C Mazari, Castro, Alba and Vazquez, *Rev. Mex. Fis.* 6 (1957) 9
 1957MC1B McCormac, Steuer, Bond and Hereford, *Phys. Rev.* 108 (1957) 116
 1957MO1B F.S. Mozer and F.B. Hagedorn, *Phys. Rev.* 105 (1957) 1270
 1957MU1A Muller, Ph.D. Thesis, Univ. of Strasbourg (1957)
 1957OW03 G.E. Owen and L. Madansky, *Phys. Rev.* 105 (1957) 1766; Erratum *Phys. Rev.* 108 (1957) 1647
 1957PE1D C.A. Pearse, *Phys. Rev.* 106 (1957) 545
 1957RA1B B.J. Raz, *Phys. Rev.* 107 (1957) 1201
 1957SA1B E.E. Salpeter, *Phys. Rev.* 107 (1957) 516
 1957SO1B Sorokin, Valter, Gavrilovskii, Karadshev, Manko and Taranov, *Zh. Eksp. Teor. Fiz.* 33 (1957) 106; *JETP (Sov. Phys.)* 6 (1958) 466
 1957TA1C G.E. Tauber and T.-Y. Wu, *Phys. Rev.* 105 (1957) 1772
 1957WA1C Walton, Clement and Boreli, *Phys. Rev.* 107 (1957) 1065
 1958AR15 S.E. Arnell, J. Dubois and O. Almen, *Nucl. Phys.* 6 (1958) 196
 1958BA32 G. Barton, *Nucl. Phys.* 6 (1958) 100
 1958BA60 W.C. Barber, *Phys. Rev.* 111 (1958) 1642
 1958BE03 R.E. Benenson and M.B. Shurman, *Rev. Sci. Instrum.* 29 (1958) 1
 1958BE1G J.S. Bell and R.J. Blin-Stoyle, *Nucl. Phys.* 6 (1958) 87
 1958FO67 J.L. Fowler and H.O. Cohn, *Phys. Rev.* 109 (1958) 89
 1958GO77 H.E. Gove, J.A. Kuehner, A.E. Litherland, E. Almqvist, D.A. Bromley, A.J. Ferguson, P.H. Rose, R.P. Bastide, N. Brooks and R.J. Connor, *Phys. Rev. Lett.* 1 (1958) 251
 1958HE57 R.E. Hester, R.E. Pixley and W.A.S. Lamb, *Phys. Rev.* 111 (1958) 1604
 1958HI68 R.W. Hill, *Phys. Rev.* 109 (1958) 2105
 1958HO97 R.E. Holland, F.J. Lynch and S.S. Hanna, *Phys. Rev.* 112 (1958) 903
 1958HU18 D.J. Hughes and R.B. Schwartz, *BNL-325*, 2nd Ed. (1958); *BNL-325*, 2nd Ed., Suppl. I (1960)
 1958LA09 R.O. Lane and J.E. Monahan, *Bull. Amer. Phys. Soc.* 3 (1958) 364, G6
 1958LI1A Livesey, *Conf. on Photonucle. Reactions*, National Bureau of Standards (1958)
 1958NI1D Nigam and Sundaresan, *Can. J. Phys.* 36 (1958) 571

1958RI1A Richter and Ivash, Phys. Rev. 111 (1958) 265
1958ST28 H.R. Striebel, S.E. Darden and W. Haeberli, Nucl. Phys. 6 (1958) 188
1958TA03 Y.-K. Tai, G.P. Millburn, S.N. Kaplan and B.J. Moyer, Phys. Rev. 109 (1958) 2086
1958WE31 J.L. Weil and K.W. Jones, Phys. Rev. 112 (1958) 1975
1958WH34 A.B. Whitehead and J.S. Foster, Can. J. Phys. 36 (1958) 1276
1959CH1F R.F. Christy and I. Duck, Nucl. Phys. 24 (1961) 89
1959TA03 N.Tanner, Phys. Rev. 114 (1959) 1060