

Energy Levels of Light Nuclei $A = 17$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 16-17$ was published in *Nuclear Physics A281* (1977), p. 1. This version of $A = 17$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed November 1, 1976)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 17$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{17}\text{Be}\$](#) , [\$^{17}\text{B}\$](#) , [\$^{17}\text{C}\$](#) , [\$^{17}\text{N}\$](#) , [\$^{17}\text{O}\$](#) , [\$^{17}\text{F}\$](#) , [\$^{17}\text{Ne}\$](#) , [\$^{17}\text{Na}\$](#)

B. Tables of Recommended Level Energies:

[Table 17.1](#): Energy levels of ^{17}N

[Table 17.6](#): Energy levels of ^{17}O

[Table 17.17](#): Energy levels of ^{17}F

[Table 17.20](#): Energy levels of ^{17}Ne

C. [References](#)

D. Figures: [\$^{17}\text{N}\$](#) , [\$^{17}\text{O}\$](#) , [\$^{17}\text{F}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

^{17}Be

(Not illustrated)

This nucleus has not been observed. Its atomic mass excess is calculated to be 70.67 MeV. It is then unstable with respect to breakup into $^{16}\text{Be} + \text{n}$ and $^{15}\text{Be} + 2\text{n}$ by 3.37 and 3.34 MeV, respectively (1974TH01). See also (1976CA1R; theor.).

^{17}B

(Not illustrated)

^{17}B has been observed in the 4.8 GeV proton bombardment of uranium: it is particle stable and its ground state J^π is probably $\frac{3}{2}^-$ (1973BO30, 1974BO05). Its atomic mass excess is calculated to be 44.37 MeV (transverse form of the mass equation): it is then stable with respect to decay into $^{15}\text{B} + 2\text{n}$ by 0.53 MeV (1974TH01, 1975JE02). The $E_{\beta^-}(\text{max})$ for the decay to ^{17}C would then be 23.1 MeV. See also (1971AJ02) and (1972GA1F, 1972TH13, 1972WI1C, 1975BE31).

^{17}C

(Not illustrated)

^{17}C has been observed in the 5.5 GeV proton bombardment of uranium: it is particle stable (1968PO04). Its atomic mass excess is calculated to be 21.27 MeV (transverse form of the mass equation): it is then stable with respect to decay into $^{16}\text{C} + \text{n}$ by 0.50 MeV (1974TH01, 1975JE02). The $E_{\beta^-}(\text{max})$ for the decay to ^{17}N would then be 13.4 MeV. See also (1971AJ02), (1971AR02, 1971BU1E), (1973TO16) and (1972TH13, 1973WI15, 1975BE31, 1975WI1E; theor.).

^{17}N

(Figs. 6 and 9)

GENERAL: (See also (1971AJ02).)

Theory and reviews: (1973PA1F, 1973RE17, 1973TO16, 1973WI15, 1974HA61, 1975BE31).

Experimental papers: (1971AR02, 1973KO1D, 1976DE1P).

1. (a) $^{17}\text{N}(\beta^-)^{17}\text{O}^* \rightarrow ^{16}\text{O} + \text{n}$ $Q_m = 4.536$

(b) $^{17}\text{N}(\beta^-)^{17}\text{O}$ $Q_m = 8.682$

Table 17.1: Energy levels of ^{17}N ^a

E_x in ^{17}N (MeV \pm keV)	$J^\pi; T$	τ or Γ	Decay	Reactions
0	$\frac{1}{2}^-; \frac{3}{2}$	$\tau_{1/2} = 4.169 \pm 0.008$ sec	β^- ^b	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
1.3739 \pm 0.3	$\frac{3}{2}^-$	$\tau_m = 93 \pm 35$ fsec	γ	3, 11, 12
1.8496 \pm 0.3	$\frac{1}{2}^+$	41_{-9}^{+20} psec	γ	3, 11, 12
1.9068 \pm 0.3	$\frac{5}{2}^-$	11 ± 2 psec	γ	3, 11, 12
2.5260 \pm 0.5	$\frac{5}{2}^+$	33 ± 3 psec	γ	3, 11, 12
3.1289 \pm 0.5	$\frac{7}{2}^{(-)}$	275 ± 80 fsec	γ	3, 12
3.2042 \pm 0.9	$\frac{3}{2}^-$	< 30 fsec	γ	3, 11, 12
3.6287 \pm 0.7	$(\frac{7}{2}^-, \frac{9}{2}^-)$	12 ± 2 psec	γ	3, 11, 12
3.663 \pm 4	$(\frac{1}{2}, \frac{3}{2})^-$	< 350 fsec	γ	3, 11, 12
3.9060 \pm 2.0	$\leq \frac{7}{2}$	52 ± 22 fsec	γ	3
4.0064 \pm 2.0	$(\frac{3}{2})$	< 15 fsec	γ	3, 11, 12
4.208 \pm 3	$\leq \frac{5}{2}$	< 70 fsec	γ	3, 12
4.415 \pm 3	$\leq \frac{7}{2}$	$(< 60$ fsec)	γ	3
5.170 \pm 2	$(\frac{7}{2}^+, \frac{9}{2}^+)$	< 60 fsec	γ	3
5.195 \pm 3	$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$	< 95 fsec	γ	3
5.514 \pm 3	$(\frac{3}{2})^-$	< 100 fsec	γ	3, 11
5.770 \pm 3	$\leq \frac{7}{2}$	< 120 fsec	γ	3, 11
6.08 \pm 30				3
6.24 \pm 25				3
6.43 \pm 30				3
6.61 \pm 25				3
6.99 \pm 20				3, 11
7.17 \pm 40				3
7.37 \pm 40				3
7.63 \pm 40				3
7.73 \pm 40				3
8.00 \pm 25				3
8.14 \pm 40				3
8.55 \pm 40		broad		3
8.93 \pm 40		broad		3
9.26 \pm 40		broad		3
9.74 \pm 40		broad		3

^a See also Tables 17.4 and 17.5.

^b See also Tables 17.2 and 17.3.

The half-life of ^{17}N is 4.169 ± 0.008 sec (1972AL42): the mean of earlier values (see (1971AJ02)) was 4.16 ± 0.01 sec. See also (1976FI03). The decay is principally [see Table 17.2] to the neutron unbound states $^{17}\text{O}^*(4.55, 5.38, 5.94)$ [$J^\pi = \frac{3}{2}^-, \frac{3}{2}^-$ and $\frac{1}{2}^-$, respectively]. The nature of the decay is in agreement with $J^\pi = \frac{1}{2}^-$ for $^{17}\text{N}_{\text{g.s.}}$ (1973PO11, 1976AL02). For measurements of the neutron energies see (1973PO11). Excitation energies of ^{17}O states have been determined by (1973PO11) from E_n [e.g., 4.552 ± 0.004 MeV] and by (1976AL02). From E_γ to the first excited state whose E_x was taken to be 870.8 ± 0.2 keV, E_x for the second excited state of ^{17}O is 3055.2 ± 0.3 keV; the direct ground state decay of that state is $< 1.5\%$ (1976AL02). See also (1972SH1F) and (1972TO03; theor.). For a comparison of the ^{17}N and ^{17}Ne decays see Table 17.3 (1976AL02): the large values of δ show the importance of final state nuclear asymmetry in odd- A nuclei. See also (1972AL42).

2. $^{10}\text{Be}(^{11}\text{B}, \alpha)^{17}\text{N}$ $Q_m = 10.980$

See (1976FI03).

3. $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ $Q_m = 8.416$

Proton groups and γ -rays have been studied by several groups: see Tables 17.4 and 17.5. Angular distributions of γ -rays, branching ratio and lifetime measurements lead to most of the J^π shown in these two tables; transition energies have also been very accurately determined (1974RO27, 1974RO28). See also (1971AJ02).

4. $^{12}\text{C}(^7\text{Li}, 2\text{p})^{17}\text{N}$ $Q_m = -7.541$

See (1971HO26, 1976CE1E).

5. $^{14}\text{C}(\alpha, \text{p})^{17}\text{N}$ $Q_m = -9.715$

See (1961PE1A).

6. $^{14}\text{C}(^{18}\text{O}, ^{15}\text{N})^{17}\text{N}$ $Q_m = -5.735$

See (1972EY01).

7. $^{15}\text{N}(\text{t}, \text{p})^{17}\text{N}$ $Q_m = -0.108$

See (1971AJ02).

Table 17.2: Beta decay of ^{17}N

Decay to $^{17}\text{O}^*$ (MeV)	J^π	Branch (%) ^a		$\log ft$ ^b
		(1973PO11)	(1976AL02) ^A	
0	$\frac{5}{2}^+$	1.7 ± 0.5	1.6 ± 0.5 ^c	7.29 ± 0.11 ^d
0.87	$\frac{1}{2}^+$	2.9 ± 0.5	3.0 ± 0.5	6.80 ± 0.07
3.06	$\frac{1}{2}^-$	0.54 ± 0.08	0.34 ± 0.06	7.08 ± 0.08
3.84	$\frac{5}{2}^-$		$< 7 \times 10^{-3}$	> 8.5
4.55	$\frac{3}{2}^-$	37.9 ± 1.8	39.2 ± 2.0	4.40 ± 0.02
5.38	$\frac{3}{2}^-$	51.1 ± 1.5	48.0 ± 1.5	3.88 ± 0.02
5.94	$\frac{1}{2}^-$	5.8 ± 0.6	7.9 ± 0.7	4.31 ± 0.04

A = adopted.

^a See also (1964SI06, 1973DE32).

^b (1976AL02).

^c (1964SI06).

^d $\log f_1 t = 9.56 \pm 0.13$ (1971TO08).

Table 17.3: Comparison of ^{17}N and ^{17}Ne β -decay^a

Final state in		J^π	Γ_n ^b (keV)	Γ_p ^b (keV)	$(ft)^-$ ^c	$(ft)^+$ ^c	δ ^d
^{17}O	^{17}F						
3.06	3.10	$\frac{1}{2}^-$	0	19	$(8.0 \pm 1.1) \times 10^6$ ^e	$(2.78 \pm 0.40) \times 10^6$	-0.65 ± 0.07
4.55	4.70	$\frac{3}{2}^-$	40	230	$(2.53 \pm 0.14) \times 10^4$ ^f	$(3.92 \pm 0.18) \times 10^4$	0.55 ± 0.11
5.38	5.52	$\frac{3}{2}^-$	28	69	$(7.59 \pm 0.28) \times 10^3$ ^f	$(7.22 \pm 0.15) \times 10^3$	-0.55 ± 0.04
5.94	6.04	$\frac{1}{2}^-$	23	28	$(2.04 \pm 0.19) \times 10^4$ ^f	$(2.61 \pm 0.07) \times 10^4$	0.28 ± 0.12

^a (1976AL02).

^b Γ_n and Γ_p are the neutron and proton widths of the ^{17}O and ^{17}F states, respectively.

^c $(ft)^-$ and $(ft)^+$ are for the ^{17}N and ^{17}Ne decays, respectively.

^d $\delta = [(ft)^+/(ft)^-] - 1$.

^e From mean of branching values (see Table 17.2).

^f From branching values of (1976AL02).

Table 17.4: Excited states of ^{17}N from $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$, $^{18}\text{O}(\text{d}, ^3\text{He})^{17}\text{N}$ and $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$ ^a

E_x (keV)				l^c	J^π ^d
(1974RO27) ^b	(1965HA05) ^b	(1966MC05) ^b	(1971HA48) ^b		
			0	1	$\frac{1}{2}^-$
1373.7 ± 0.5	1374.1 ± 0.4 ⁱ		1370 ± 20	1	$\frac{3}{2}^-$
1850.0 ± 0.5	1849.5 ± 0.3 ⁱ				$\frac{1}{2}^+$
			1870 ± 20	0	
1906.8 ± 0.4	1906.9 ± 0.5 ⁱ				$\frac{5}{2}^-$ ⁱ
2526.3 ± 1.0	2525.9 ± 0.6 ⁱ		2540 ± 20	2	$\frac{5}{2}^+$
3128.7 ± 0.6	3129.2 ± 0.6 ⁱ				$\frac{7}{2}^{(-)}$ ⁱ
			3180 ± 30	1	
3203 ± 2	3204.4 ± 0.9 ⁱ				$\frac{3}{2}^-$ ⁱ
3628.7 ± 0.7					$\geq \frac{3}{2}^g$
			3660 ± 30	1	
3663 ± 4					$(\frac{1}{2}, \frac{3}{2})^-$
3906.0 ± 2.0					$\leq \frac{7}{2}$
4006.4 ± 2.0			4020 ± 40		$\frac{3}{2}, \frac{5}{2}, \frac{7}{2}$
4208 ± 3					$\leq \frac{5}{2}$
4415 ± 3	4470 ± 10	4470 ± 40			$\leq \frac{7}{2}$
5170 ± 2					$\frac{3}{2} \leq J \leq \frac{9}{2}$ ^h
5195 ± 3	5210 ± 20	5230 ± 40			$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$
5514 ± 3	5530 ± 20	5510 ± 40	$\equiv 5523$ ^e	1	$(\frac{1}{2}, \frac{3}{2})^-$
5770 ± 3	5830 ± 20	5830 ± 40	5820 ± 40		$\leq \frac{7}{2}$
	6070 ± 50	6090 ± 40			
	6250 ± 30	6230 ± 40			
	6450 ± 40	6410 ± 40			
	6600 ± 30	6620 ± 40			
	6990 ± 30	6990 ± 40	6990 ± 30	1	$(\frac{3}{2}, \frac{1}{2})^-$
	(7260 ± 50)	7170 ± 40			
	(7510 ± 70)	7370 ± 40			

Table 17.4: Excited states of ^{17}N from $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$, $^{18}\text{O}(\text{d}, ^3\text{He})^{17}\text{N}$ and $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$ ^a (continued)

E_x (keV)				l ^c	J^π ^d
(1974RO27) ^b	(1965HA05) ^b	(1966MC05) ^b	(1971HA48) ^b		
		7630 ± 40			
	7790 ± 20	7730 ± 40			
	8000 ± 30	8000 ± 40			
	(8250 ± 30)	8140 ± 40			
		8550 ± 40 ^f			
		8930 ± 40			
		9260 ± 40			
		9740 ± 40			

^a See also Table 17.3 in (1971AJ02) for the earlier work. The work reported in col. A in that table has not been published.

^b $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$.

^c $^{18}\text{O}(\text{d}, ^3\text{He})^{17}\text{N}$.

^d (1971HA48, 1974RO27), except for values labeled ⁱ.

^e Used as calibration point.

^f This state and the ones below are broad.

^g Probably $(\frac{7}{2}, \frac{9}{2})^-$ (1974RO27).

^h Probably $(\frac{7}{2}, \frac{9}{2})^+$ (1974RO27).

ⁱ $^{18}\text{O}(\text{t}, \alpha)^{17}\text{N}$ (1976GU14).

8. $^{17}\text{O}(\text{n}, \text{p})^{17}\text{N}$ $Q_m = -7.896$

See ^{18}O in (1978AJ03).

9. $^{18}\text{O}(\gamma, \text{p})^{17}\text{N}$ $Q_m = -15.943$

See ^{18}O in (1978AJ03).

10. $^{18}\text{O}(\text{n}, \text{d})^{17}\text{N}$ $Q_m = -13.718$

See ^{19}O in (1972AJ02).

Table 17.5: Radiative transitions and lifetimes of ^{17}N states

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Γ_γ/Γ_w^e (W.u.)	δ	τ_m	Refs.
1.37	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	100	$0.13^{+0.08}_{-0.04}$ (M1)		93 ± 35 fsec	(1974RO27)
				100	0.13 ± 0.05 (M1)	0.00 ± 0.03		(1976GU14)
1.85	$\frac{1}{2}^+$	0	$\frac{1}{2}^-$	90 ± 3	$(5.0 \pm 1.6) \times 10^{-6}$ (E1)		41^{+20}_{-9} psec	(1974RO27, 1974RO28)
				83 ± 3	$(5.0 \pm 2.0) \times 10^{-6}$ (E1)			(1976GU14)
		1.37	$\frac{3}{2}^-$	10 ± 3	$(3.2 \pm 1.5) \times 10^{-5}$ (E1)			(1974RO27, 1974RO28)
				17 ± 3	$(6 \pm 3) \times 10^{-5}$ (E1)	0.00 ± 0.02		(1976GU14)
1.91	$\frac{5}{2}^-$	0	$\frac{1}{2}^-$	78 ± 3	1.0 ± 0.2 (E2)		11 ± 2 psec	(1974RO27, 1974RO28)
				74 ± 4	0.8 ± 0.2 (E2)	0.00 ± 0.05		(1976GU14)
		1.37	$\frac{3}{2}^-$	22 ± 3	$(4.2 \pm 1.5) \times 10^{-3}$ (M1)			(1974RO27, 1974RO28)
				26 ± 4	$(5 \pm 1) \times 10^{-3}$ (M1) ^g	$-0.05^{+0.03}_{-0.14}$		(1976GU14)
2.53	$\frac{5}{2}^+$	0	$\frac{1}{2}^-$	14 ± 4	0.28 ± 0.11 (M2)		33 ± 3 psec	(1974RO27, 1974RO28)
				11 ± 1	0.22 ± 0.04 (M2)	-0.07 ± 0.18		(1976GU14)
		1.37	$\frac{3}{2}^-$	34 ± 4	$(1.0 \pm 0.2) \times 10^{-5}$ (E1)			(1974RO27, 1974RO28)
				34 ± 4	$(1.0 \pm 0.2) \times 10^{-5}$ (E1)	0.0 ± 0.1		(1976GU14)
		1.85	$\frac{1}{2}^+$	12 ± 2	9 ± 2 (E2)			(1974RO27, 1974RO28)
				12 ± 2	8.1 ± 1.6 (E2)	0.00 ± 0.06		(1976GU14)
		1.91	$\frac{5}{2}^-$	40 ± 3	$(8 \pm 1) \times 10^{-5}$ (E1)			(1974RO27, 1974RO28)
				43 ± 4	$(2.3 \pm 0.4) \times 10^{-5}$ (E1)	0.07 ± 0.07		(1976GU14)
3.13 ^a	$\frac{7}{2}^{(-)}$	1.91	$\frac{5}{2}^-$	100	$0.063^{+0.036}_{-0.016}$ (M1)		275 ± 80 fsec	(1974RO27)
				100	0.06 ± 0.02 (M1)	0.00 ± 0.04		(1976GU14)
3.20 ^b	$\frac{3}{2}^-$	0	$\frac{1}{2}^-$	88 ± 6	> 0.03 (M1)		< 30 fsec	(1974RO27)
				88 ± 4	> 0.025 (M1)	-0.06 ± 0.08 ^f		(1976GU14)
		1.91	$\frac{5}{2}^-$	12 ± 6	> 0.05 (M1)			(1974RO27)
				12 ± 4	> 0.05 (M1)			(1976GU14)
3.63 ^c	$(\frac{7}{2}^-, \frac{9}{2}^-)$	1.91	$\frac{5}{2}^-$	47 ± 10	0.8 ± 0.2 (E2)		12 ± 2 psec	(1974RO27, 1974RO28)
		3.13	$\frac{7}{2}^{(-)}$	53 ± 10	0.010 ± 0.03 (M1)			(1974RO27, 1974RO28)
3.66	$(\frac{1}{2}, \frac{3}{2})^-$	1.85	$\frac{1}{2}^+$	100	$> 7 \times 10^{-4}$ (E1)		< 350 fsec	(1974RO27)

Table 17.5: Radiative transitions and lifetimes of ^{17}N states (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	Γ_γ/Γ_w ^e (W.u.)	δ	τ_m	Refs.
3.91	$\leq \frac{7}{2}$	1.91	$\frac{5}{2}^-$	100	$(8_{-3}^{+5}) \times 10^{-2}$ (M1) ^h		52 ± 22 fsec	(1974RO27)
4.01	$(\frac{3}{2})$	1.85	$\frac{1}{2}^+$	$\leq 15 \pm 5$ ^d				< 15 fsec
		2.53	$\frac{5}{2}^+$	85 ± 5	0.55 (M1)			(1974RO27)
4.21	$\leq \frac{5}{2}$	1.37	$\frac{3}{2}^-$	100			< 70 fsec	(1974RO27)
4.42	$\leq \frac{7}{2}$	1.91	$\frac{5}{2}^-$	100			(< 60) fsec	(1974RO27)
5.17	$(\frac{7}{2}^+, \frac{9}{2}^+)$	2.53	$\frac{5}{2}^+$	37 ± 7	> 15 (E2)		< 60 fsec	(1974RO27)
		3.13	$\frac{7}{2}^{(-)}$	63 ± 7				(1974RO27)
5.20	$(\frac{1}{2}, \frac{3}{2}, \frac{5}{2})^+$	1.85	$\frac{1}{2}^+$	≈ 42			< 95 fsec	(1974RO27)
		1.91	$\frac{5}{2}^-$	≈ 58				(1974RO27)
5.51	$(\frac{3}{2})^-$	0	$\frac{1}{2}^-$	≈ 50			< 100 fsec	(1974RO27)
		1.37	$\frac{3}{2}^-$	≈ 50				(1974RO27)
5.77	$\leq \frac{7}{2}$	1.37	$\frac{3}{2}^-$	≈ 25			< 120 fsec	(1974RO27)
		1.91	$\frac{5}{2}^-$	≈ 25				(1974RO27)
		4.01	$(\frac{3}{2})$	≈ 50 ^d				(1974RO27)

^a Branches to $^{17}\text{N}^*(0, 1.37, 1.85, 2.53)$ are, respectively, < 2, < 5, < 2 and < 3% (1976GU14).

^b Branches to $^{17}\text{N}^*(1.37, 1.85, 2.53)$ are, respectively, < 5, < 6 and < 3% (1976GU14).

^c Branches to $^{17}\text{N}^*(0, 1.37, 1.85, 2.53, 3.20)$ are, respectively, < 10, < 10, < 7, < 3, < 2% (1974RO28).

^d This branch is uncertain.

^e Assuming pure multipole transitions and J^π shown: see also Table 2 in the Introduction.

^f Or $\delta = 2.1 \pm 0.4$ (1976GU14).

^g $\Gamma_\gamma/\Gamma_w = 0.4_{-1.3}^{+0.4}$ (E2) (1976GU14).

^h This number appears to be in error: see Table 2 in the Introduction.

$$11. \text{}^{18}\text{O}(\text{d}, \text{}^3\text{He})\text{}^{17}\text{N} \quad Q_{\text{m}} = -10.449$$

Angular distributions of eight ${}^3\text{He}$ groups have been measured at $E_{\text{d}} = 52 \text{ MeV}$; see Table [17.4](#) ([1971HA48](#)).

$$12. \text{}^{18}\text{O}(\text{t}, \alpha)\text{}^{17}\text{N} \quad Q_{\text{m}} = 3.872$$

Alpha particle groups corresponding to ${}^{17}\text{N}$ states with $E_{\text{x}} < 4.3 \text{ MeV}$ have been studied by ([1960JA13](#)): see Table 17.3 in ([1971AJ02](#)). ([1976GU14](#): $E_{\text{t}} = 3.5 \text{ MeV}$) have studied α - γ angular correlations and γ -branching ratios for the first six excited states of ${}^{17}\text{N}$: see Tables [17.4](#) and [17.5](#).

17O
(Figs. 7 and 9)

GENERAL: (See also (1971AJ02).)

Shell model: (1968KA1C, 1969FE1A, 1970HA49, 1970IR01, 1971AR1R, 1971HS02, 1971JE02, 1971KA40, 1971LE30, 1971MU23, 1971WI01, 1971WI1F, 1972BE22, 1972EL1C, 1972EN03, 1972HA1Q, 1972KA38, 1972LE1L, 1973BA2J, 1973DE13, 1973JU1A, 1973KU04, 1973LA1D, 1973RE17, 1973SM1C, 1974LO04, 1974RI09, 1976PO01).

Collective and cluster models: (1969FE1A, 1971AR1R, 1972LE1L, 1972NE1B).

Special levels: (1968KA1C, 1969FE1A, 1969WI1C, 1971AR1R, 1971BE59, 1971BE2D, 1971HS02, 1971KO12, 1971MU23, 1971SE1C, 1972BE22, 1972BE1E, 1972EN03, 1972HI17, 1972NI15, 1973JU1A, 1974RI09).

Electromagnetic transitions: (1969FE1A, 1970AL1D, 1970HA49, 1970SI1J, 1972EN03, 1972SE1G, 1973HA53, 1973RE17, 1973ZA1D, 1974KO1R, 1974LO04, 1974MC1F, 1976SH04).

Special reactions: (1971AR02, 1972PU1B, 1973WI15, 1975KU1K, 1975TS01, 1975UD01, 1975VO09, 1976DA1T, 1976DE1P, 1976HI05, 1977PE08).

Astrophysical questions: (1972CL1A, 1973AR1E, 1973AU1B, 1973AU1D, 1973AU1C, 1973EN1A, 1973SM1A, 1973TA1D, 1973TR1B, 1974DE1M, 1975AR1E, 1975AU1D, 1975CO1J, 1975EN1A, 1975LA1E, 1975NO1D, 1975SC1H, 1975TR1A, 1976FI1E, 1976KO1K, 1976ME1H, 1976WA1M).

Pion capture and reactions (See also reactions 40 and 69.): (1973EI01, 1974DA23, 1974LI1D, 1975PA06, 1976EN02).

Other topics: (1968KA1C, 1969EL1A, 1970AL1E, 1970SI1J, 1971AU08, 1971BA2Y, 1971BE59, 1971ER1C, 1971JE02, 1971KA40, 1971KO12, 1971LA1D, 1971PL1D, 1971RY1A, 1971SE1C, 1971NG01, 1972CA37, 1972CH16, 1972DA21, 1972HA57, 1972KA38, 1972LE1L, 1972MA57, 1972NI15, 1972SH32, 1973AR1K, 1973BA2J, 1973BE1N, 1973DE13, 1973GO1H, 1973HY1A, 1973KO1J, 1973KU04, 1973MA48, 1973OS1A, 1973PA1F, 1973RA1E, 1973RE17, 1973RO1R, 1973RO1P, 1973SP1A, 1973YO1A, 1974AU03, 1974BR1E, 1974RE03, 1974SA05, 1974SL1C, 1975DR1D, 1975HE10, 1975MI02, 1975SH1H, 1976FE1B, 1976MA04, 1976MA05, 1976VA1C).

Ground state:

$$\mu = -1.89379 \pm 0.00009 \text{ nm (see (1974SHYR))};$$

$$Q = -25.6 \text{ mb (1968SC18)};$$

$$Q = -25.78 \text{ mb (1969SC34)}.$$

See also (1973CO1P, 1974SHYR).

See also (1970AL1D, 1970SI1J, 1971RY1A, 1971SH26, 1971TA1A, 1971WI01, 1972GL06, 1972LE1L, 1972VA36, 1972YO1B, 1973AR1J, 1973HI1A, 1973HO32, 1973LE07, 1973MI1C,

1973RE17, 1973RO1P, 1973SU1B, 1973SU1C, 1974DE1E, 1974HA27, 1974MC1F, 1974NE1B, 1974RE03, 1975BE31, 1975MI02, 1976CH1T, 1976PO01).

Table 17.6: Energy levels of ^{17}O

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{1}{2}$		stable	1, 2, 5, 6, 7, 8, 13, 14, 15, 16, 18, 19, 20, 21, 22, 30, 31, 32, 40, 41, 42, 43, 44, 45, 46, 48, 49, 50, 52, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 70, 71, 72, 73, 74
0.87081 ± 0.22	$\frac{1}{2}^+$	$\tau_m = 258.6 \pm 2.6$ psec ^b	γ	1, 2, 5, 6, 7, 13, 14, 15, 16, 18, 19, 20, 21, 22, 30, 31, 32, 40, 41, 45, 46, 47, 48, 49, 50, 58, 61, 62, 63, 69, 70, 71, 72, 73, 74
3.0552 ± 0.3	$\frac{1}{2}^-$	$\tau_m = 120^{+80}_{-60}$ fsec ^c	γ	5, 6, 7, 13, 14, 18, 21, 22, 30, 32, 41, 45, 50, 52, 61, 71, 72, 74
3.841 ± 3	$\frac{5}{2}^-$	$\tau_m \leq 25$ fsec ^c	γ	5, 6, 7, 13, 14, 18, 21, 22, 30, 31, 40, 41, 45, 52, 61, 62, 71, 72
4.553 ± 2	$\frac{3}{2}^-$	$\Gamma = 40 \pm 5$	n	5, 6, 7, 13, 14, 21, 22, 30, 31, 34, 41, 45, 50, 52, 61, 62, 72
5.086 ± 2	$\frac{3}{2}^+$	95 ± 5	n	6, 7, 13, 14, 21, 22, 30, 34, 41, 61, 62
5.215 ± 5	$(\frac{9}{2}^-)$	< 0.1		6, 13, 14, 21, 22, 30, 31, 41, 52, 61, 72
5.380 ± 2	$\frac{3}{2}^-$	28 ± 7	n	21, 22, 30, 32, 34, 41, 50, 52, 61, 62, 72
5.698 ± 2	$\frac{7}{2}^-$	3.4 ± 0.3	n	6, 13, 14, 21, 22, 30, 31, 34, 41, 52

Table 17.6: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
5.734 ± 2		< 1	n	5, 6, 13, 14, 21, 22, 34, 41, 72
5.870 ± 2	$\frac{3}{2}^+$	6.6 ± 0.7	n	6, 14, 21, 22, 30, 34, 41, 72
5.940 ± 4	$\frac{1}{2}^-$	32 ± 3	n	5, 6, 14, 21, 22, 30, 34, 41, 50, 52, 72
6.357 ± 8	$\frac{1}{2}^+$	124 ± 12	n	5, 21, 30, 34
6.863 ± 2	$(\frac{1}{2}^-)$	< 1	n	5, 6, 13, 14, 21, 22, 30, 34, 41, 52, 72
6.973 ± 2		< 1	n	6, 13, 14, 21, 22, 30, 34, 72
7.1687 ± 1.5	$\frac{5}{2}^-$	1.5 ± 0.2	n, α	5, 6, 9, 13, 14, 21, 30, 34, 39, 52
7.202 ± 10	$\frac{3}{2}^+$	280 ± 30	n, α	34, 39
7.3831 ± 1.5	$\frac{5}{2}^+$	0.6 ± 0.2	n, α	5, 6, 9, 13, 14, 21, 30, 31, 34, 39
7.3860 ± 1.5	$\frac{5}{2}^-$	0.9 ± 0.3	n, α	5, 9, 21, 30, 31, 34, 39, 52
7.560 ± 20	$\frac{3}{2}^-$	500 ± 50	n, α	34, 39, 41
7.577 ± 2	$\frac{7}{2}^-$	≤ 1	n, α	5, 6, 9, 13, 14, 21, 30, 34, 52
7.690 ± 4	$\frac{7}{2}^-$	18 ± 2	n, α	5, 6, 9, 14, 30, 34, 39
7.75 ± 20	$\frac{11}{2}^-$			13, 14, 23, 30, 31, 32, 52
7.956 ± 6	$\frac{1}{2}^+$	90 ± 9	n, α	9, 30, 34, 39
7.99 ± 50	$\frac{1}{2}^-$	270 ± 30	n, α	34, 39
8.070 ± 10	$\frac{3}{2}^+$	85 ± 9	n, α	9, 30, 34, 39
(8.18 ± 20)	$\frac{1}{2}^-$	69 ± 7	n, α	34, 39
8.200 ± 7	$\frac{3}{2}^-$	60	n, α	9, 30, 31, 34, 39
8.352 ± 4	$\frac{1}{2}^+$	9 ± 3	n, α	9, 30, 34, 39
8.410 ± 3	$\frac{5}{2}^+$	4 ± 3	n, α	6, 9, 13, 14, 30, 34, 39
8.474 ± 3	$\frac{7}{2}^+$	7 ± 3	n, α	5, 6, 9, 13, 14, 31, 39

Table 17.6: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
8.508 ± 3	$\frac{5}{2}^-$	5 ± 3	n, α	6, 9, 13, 14, 30, 34, 39
8.700 ± 5	$\frac{3}{2}^-$	50 ± 3	n, α	9, 30, 34, 39
8.898 ± 8	$\frac{3}{2}^+$	101 ± 3	n, α	6, 9, 13, 14, 30, 31, 34, 39
8.972 ± 4	$\frac{7}{2}^-$	21 ± 3	n, α	6, 9, 14, 30, 34, 39
9.148 ± 4	$\frac{1}{2}^-$	4 ± 3	n, α	6, 9, 14, 34
9.15 ± 20	$\frac{9}{2}^-$			23, 30, 31, 32
9.187	$\frac{7}{2}^-$	3	n, α	9, 34
9.201 ± 4	$\frac{5}{2}^+$	5.5 ± 1	n, α	9, 34
9.422	$\frac{3}{2}^-$	120	n	34
9.493 ± 4	$\frac{5}{2}^-$	15 ± 1	n, α	5, 9, 14, 30
9.720 ± 5	$\frac{7}{2}^+$	16 ± 1	n, α	9, 14, 30, 34
9.775 ± 15	$\frac{3}{2}^+$	≈ 25	n, α	9, 34
9.865 ± 5		14	n, α	9, 14, 30, 34
9.878 ± 15		≈ 10	n, α	9, 34
9.977 ± 20	$\frac{5}{2}^+$	≈ 80	n, α	9
10.046 ± 20		≈ 100	n, α	9
10.178 ± 5	$\frac{7}{2}^-$	40	n, α	9, 34
10.337 ± 15	$\frac{5}{2}^+, \frac{7}{2}^-$	150	n, α	9, 30
10.429 ± 7		14 ± 3	n, α	9
10.49	$\frac{5}{2}^+, \frac{7}{2}^-$	75 ± 30	n, α	9
10.563 ± 10	$(\frac{7}{2}^-)$	47 ± 15	n, α	9, 30, 34, 35
10.773 ± 10	$\frac{1}{2}^+, \frac{7}{2}^-$	80 ± 20	n, α	9, 14, 30, 35
10.910 ± 7	$\frac{5}{2}$	57 ± 15	n, α	9, 30, 34, 35
11.030 ± 4	$T = \frac{1}{2}$	45 ± 10	n, α	9, 30
11.076 ± 4^a	$\frac{1}{2}^-; \frac{3}{2}$	5 ± 1	n, α	9, 30, 35, 63
11.229 ± 10		100 ± 30	n, α	5, 9
11.52	$(\frac{3}{2}, \frac{5}{2})$	190	n	34, 35
11.619 ± 10		120 ± 30	n, α	9
11.752 ± 10		40 ± 25	n, α	9

Table 17.6: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
11.817 \pm 15		12 \pm 3	n, α	9, 14
12.006 \pm 15	$\geq \frac{3}{2}$	270	n, α	9, 34, 35
12.11 \pm 20		150 \pm 50	n, α	9, 35
12.275 \pm 15		100 \pm 30	n, α	9
12.39 \pm 20		130	n, α	9, 34
12.421 \pm 15			n, α	9, 35
12.464 \pm 5	$(\frac{3}{2})^-; \frac{3}{2}$	8 \pm 2	n, α	9, 35, 63
12.596 \pm 15		75 \pm 30	n, α	9
(12.656)		95	n	34, 35
12.670 \pm 15		\approx 5	n, α	9, 35
12.81 \pm 25			n, α	9
12.93 \pm 20		\gtrsim 150	n, α	9, 35
12.946 \pm 5	$\frac{1}{2}^+; \frac{3}{2}$	6 \pm 2	n, α	9, 63
12.993 \pm 5	$T = \frac{3}{2}$	\leq 3	n, α	9, 63
13.077 \pm 15		16 \pm 4	n, α	9
13.485 \pm 15		\approx 120	n, α	9
13.610 \pm 15		250 \pm 100	n, α	9
13.640 \pm 5	$(\frac{5}{2})^+; \frac{3}{2}$			63
(13.672)		400	n	34
14.219 \pm 8	$T = \frac{3}{2}$			63
14.282 \pm 12	$T = \frac{3}{2}$			63
14.621		340	n	34
(14.98)	$\frac{5}{2}^+$	\approx 150	n, d, α	28, 34
15.101 \pm 8	$T = \frac{3}{2}$			63
(15.15)	$(\frac{5}{2}, \frac{7}{2})^-$	\approx 200	p, d	26
(15.5)		broad	p, d α	26, 28
20.45			γ , t	20
21.7 \pm 100	$\frac{5}{2}^+$	750	γ , d, ^3He , α	16, 17, 24
22.1 \pm 100	$\frac{7}{2}^-$	750	γ , n, d, ^3He , α	16, 17, 24
22.5 \pm 200	$\frac{3}{2}^-$	\approx 1000	γ , d, ^3He	16, 24, 51

Table 17.6: Energy levels of ^{17}O (continued)

E_x in ^{17}O (MeV \pm keV)	$J^\pi; T$	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
23.0 ± 100	$\frac{1}{2}^+$	≈ 400	$\gamma, d, ^3\text{He}$	16, 17, 24
23.5 ± 100			$\gamma, ^3\text{He}$	16
24.4 ± 100			$\gamma, ^3\text{He}$	16

^a See also Table 17.11.

^b See Table 17.7 in (1971AJ02).

^c See (1964AL11).

Mass of ^{17}O : the atomic mass excess of ^{17}O is -810.5 ± 0.8 keV (A.H. Wapstra, private communication).

$$1. \ ^7\text{Li}(^{14}\text{N}, \alpha)^{17}\text{O} \quad Q_m = 16.158$$

The angular distribution of the α_{0+1} group has been measured at $E(^{14}\text{N}) = 27.6$ MeV (1964WA1B).

$$2. \ ^9\text{Be}(^{16}\text{O}, ^8\text{Be})^{17}\text{O} \quad Q_m = 2.480$$

Angular distributions have been studied at $E(^{16}\text{O}) = 11$ MeV ($^{17}\text{O}_{g.s.}$) and 15 and 18 MeV ($^{17}\text{O}^*(0, 0.87)$) (1970BA49, 1971BA68). The excitation curve for the one-neutron transfer to $^{17}\text{O}^*(0.87)$ has been measured for $E(^{16}\text{O}) = 6$ to 22 MeV (1970BA55). See also (1971NI04) and (1971AL1D; theor.).

$$\begin{aligned}
 3. \text{ (a) } & \ ^{10}\text{B}(^7\text{Li}, p)^{16}\text{N} & Q_m = 13.990 & E_b = 27.771 \\
 & \text{ (b) } & \ ^{10}\text{B}(^7\text{Li}, d)^{15}\text{N} & Q_m = 13.723 \\
 & \text{ (c) } & \ ^{10}\text{B}(^7\text{Li}, t)^{14}\text{N} & Q_m = 9.147 \\
 & \text{ (d) } & \ ^{10}\text{B}(^7\text{Li}, \alpha)^{13}\text{C} & Q_m = 21.411
 \end{aligned}$$

Cross sections to various of the final states have been measured at $E(^7\text{Li}) = 5.20$ MeV (1966MC05).

4. (a) $^{11}\text{B}(^6\text{Li}, \text{p})^{16}\text{N}$ $Q_m = 9.785$ $E_b = 23.566$
 (b) $^{11}\text{B}(^6\text{Li}, \text{d})^{15}\text{N}$ $Q_m = 9.517$
 (c) $^{11}\text{B}(^6\text{Li}, \text{t})^{14}\text{N}$ $Q_m = 4.9413$
 (d) $^{11}\text{B}(^6\text{Li}, \alpha)^{13}\text{C}$ $Q_m = 17.205$

Cross sections to various of the final states have been measured at $E(^6\text{Li}) = 4.72$ MeV (1966MC05).

5. $^{12}\text{C}(^6\text{Li}, \text{p})^{17}\text{O}$ $Q_m = 7.609$

Proton groups have been identified to various states of ^{17}O with $E_x \leq 15.8$ MeV and angular distributions of proton groups have been studied for $E(^6\text{Li}) = 3$ to 20 MeV: see (1971AJ02) for a complete listing. See also (1970JO1D) and ^{18}F in (1978AJ03).

6. $^{12}\text{C}(^7\text{Li}, \text{d})^{17}\text{O}$ $Q_m = 2.583$

Angular distributions have been measured at $E(^7\text{Li}) = 3.24 - 3.64$ MeV (1967MO23: d_0, d_1) and at 21.1 MeV (1971SC21: see Table 17.7). The latter have been analyzed by a Hauser-Feshbach method. See also (1971AJ02) and ^{19}F in (1978AJ03).

7. $^{12}\text{C}(^9\text{Be}, \alpha)^{17}\text{O}$ $Q_m = 9.734$

Cross sections have been measured for the population of $^{17}\text{O}^*(0, 0.87, 3.06, 3.84)$ for $E(^{12}\text{C})_{\text{c.m.}} = 2.40$ to 6.34 MeV (1974HA25). At $E(^9\text{Be}) = 26.3$ MeV, α -groups are observed to the first six states of ^{17}O and to $^{17}\text{O}^*(7.5, 8.4, 9.8, 11.0, 11.8, 13.6)$. Some of the higher states (unresolved) may be formed by some sort of direct mechanism (1975VE10).

8. $^{12}\text{C}(^{13}\text{C}, 2\alpha)^{17}\text{O}$ $Q_m = -0.914$

See (1976DA13).

9. (a) $^{13}\text{C}(\alpha, \text{n})^{16}\text{O}$ $Q_m = 2.2152$ $E_b = 6.361$
 (b) $^{13}\text{C}(\alpha, \alpha)^{13}\text{C}$

The yield of neutrons increases monotonically for $E_\alpha = 0.475$ to 0.700 MeV: $S(E) = [(5.48 \pm$

Table 17.7: States of ^{17}O from $^{12}\text{C}(^7\text{Li}, \text{d})^{17}\text{O}$, $^{13}\text{C}(^6\text{Li}, \text{d})^{17}\text{O}$ and $^{13}\text{C}(^7\text{Li}, \text{t})^{17}\text{O}$ ^a

$^{17}\text{O}^*$ (MeV)	σ^b (mb)	σ^c (mb)	$d\sigma/d\Omega$ (in $\mu\text{b}/\text{sr}$) ^d		J^e
			($^6\text{Li}, \text{d}$)	($^7\text{Li}, \text{t}$)	
0	0.67	0.83	105	75	$\frac{5}{2}$
0.87	0.33	0.32	180	92	$\frac{1}{2}$
3.06	1.05	0.50	560	750	$\frac{1}{2}$
3.84	1.83	1.24	340	1400	$\frac{5}{2}$
4.55	2.02	0.76	285	1350	$\frac{3}{2}$
5.08	0.72		180	250	
5.22	1.87	2.40	245	230	$\frac{7}{2}$
5.70 } 5.73 } 5.87 }	2.69	1.93	230	530	
5.94	1.10	1.22			$\frac{5}{2} + \frac{1}{2}$
6.86	1.30	1.40	92	125	$\frac{7}{2}$
6.97	1.79	1.20	200	320	$\frac{5}{2}$
7.17 + 7.20	2.62	1.22	350	1050	$\frac{5}{2}$
7.38 + 7.39	4.52	1.96	720	2000	$\frac{9}{2}$
7.58	1.67	2.06	98	310	$\frac{9}{2}$
7.69	3.47	2.89	620	1100	$\frac{3}{2} + \frac{7}{2} + \frac{3}{2}$
7.76 \pm 0.02			f	f	$(\frac{11}{2})^f$
8.41 } 8.47 } 8.51 }	7.52	4.39	940	2400	$\frac{5}{2} + \frac{9}{2} + \frac{5}{2}$
8.90 } 8.97 }	8.85	3.92			$\frac{3}{2} + \frac{7}{2} + \frac{7}{2}$
9.15 + 9.19	4.26	2.63			
9.49		1.65			
9.72 + 9.78		2.51			

Table 17.7: States of ^{17}O from $^{12}\text{C}(^7\text{Li}, \text{d})^{17}\text{O}$, $^{13}\text{C}(^6\text{Li}, \text{d})^{17}\text{O}$ and $^{13}\text{C}(^7\text{Li}, \text{t})^{17}\text{O}$ ^a (continued)

$^{17}\text{O}^*$ (MeV)	σ^b (mb)	σ^c (mb)	$d\sigma/d\Omega$ (in $\mu\text{b}/\text{sr}$) ^d		J^e
			($^6\text{Li}, \text{d}$)	($^7\text{Li}, \text{t}$)	
9.87 } 9.88 }		3.03			
10.77		3.44			
11.91		4.72			

^a See also Table 17.8 in (1971AJ02).

^b From integration over angular distributions of deuteron groups from ($^7\text{Li}, \text{d}$) (1971SC21).

^c From integration over angular distributions of triton groups (1971SC21).

^d (1970BE31): $d\sigma/d\Omega$ taken at maximum of angular distribution. See also Table 17.8 in (1971AJ02) for $d\sigma/d\Omega$ at 30 for other states.

^e (1971SC21).

^f Angular distribution obtained by (1970BE31).

1.77) + (12.05 ± 3.91)E] × 10⁵ MeV · b (1968DA05). Astrophysical considerations are discussed by (1968DA1D, 1973BA10, 1975FO19, 1976DE1G). Yield curves for reaction (a) have been measured for $E_\alpha = 1.0$ to 22.5 MeV: see (1971AJ02) for a listing of the earlier work and (1973BA10: total neutron yield; $E_\alpha = 1.0$ to 5.4 MeV), (1976MC11: n_0 ; $E_\alpha = 4.2$ to 8.7 MeV; $T = \frac{3}{2}$ states). Elastic scattering studies (reaction (b)) have been studied at $E_\alpha = 2.0$ to 26.6 MeV: see (1971AJ02) for the earlier results and (1973KU18, 1973LE28: 18 to 26.6 MeV). See also (1974WE1P). Observed resonances in the neutron yields and anomalies in the elastic scattering are displayed in Table 17.8. Some of the J^π values derived from the polarization studies of (1971BA06: n_0 ; $E_\alpha = 3.36$ to 4.80 MeV) and (1973BU14: n_0 ; $E_\alpha = 2.08, 2.25, 2.43$ MeV). See also (1970RO08, 1971SE1E, 1974LO1B) and (1972HA2A; theor.).

$$10. \ ^{13}\text{C}(\alpha, \text{p})^{16}\text{N} \qquad Q_m = -7.421 \qquad E_b = 6.361$$

See (1974SC1L; theor.).

$$11. \ ^{13}\text{C}(\alpha, \text{d})^{15}\text{N} \qquad Q_m = -7.6879 \qquad E_b = 6.361$$

Excitation functions for elastically scattered deuterons have been measured for $E_\alpha = 13$ to 25 MeV: strong fluctuations are observed (1973LE28). See also (1976LE1K).

Table 17.8: Resonances in $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{13}\text{C}(\alpha, \alpha)^{13}\text{C}$ ^a

E_{res} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Γ_{α}/Γ	J^{π}	E_x (MeV)	Refs.
1.0563 \pm 1.5	1.5 \pm 0.2		$\frac{5}{2}$	7.1687	(1973BA10), A
1.3367 \pm 1.5	0.6 ^{+0.2} _{-0.1}			7.3831	(1973BA10), A
1.3406 \pm 1.5	0.8 ^{+0.3} _{-0.2}			7.3860	(1973BA10), A
1.590 \pm 2	≤ 1		$\frac{7}{2}^{-}$	7.577	(1973BA10), A
1.745 \pm 6	≤ 15		$\frac{5}{2}^{+}$	7.695	(1973BA10), A
2.083 \pm 8	75	0.03	$\frac{1}{2}^{-}$	7.954	(1973BA10, 1973BU14), A
2.250 \pm 8	110	0.05	$\frac{3}{2}^{+}$	8.081	(1973BA10, 1973BU14), A
2.407 \pm 8	70	0.11	$\frac{3}{2}^{-}$	8.201	(1973BA10, 1973BU14), A
2.604 \pm 4	9 \pm 3	0.44	$\frac{1}{2}^{+}$	8.352	(1973BA10), A
2.680 \pm 3	4 \pm 3	0.08	$\frac{5}{2}^{+}$	8.410	(1973BA10), A
2.763 \pm 3	7 \pm 3	0.97	$\frac{7}{2}^{+}$	8.474	(1973BA10), A
2.808 \pm 3	5 \pm 3	0.26	$\frac{5}{2}^{-}$	8.508	(1973BA10), A
3.059 \pm 5	50 \pm 3	0.06	$\frac{3}{2}^{-}$	8.700	(1971BA06, 1973BA10), A
(3.1)	broad		$\frac{1}{2}^{-}$	(8.7)	(1971BA06)
3.318 \pm 8	101 \pm 3	0.50	$\frac{3}{2}^{+}$	8.898	(1971BA06, 1973BA10), A
3.415 \pm 4	21 \pm 3	0.04	$\frac{7}{2}^{-}$	8.972	(1971BA06, 1973BA10), A
3.645 \pm 4	4 \pm 3	0.45	$\frac{1}{2}^{-}$	9.148	(1971BA06, 1973BA10), A
(3.69)	3	1.00	$\frac{7}{2}^{-}$	(9.18)	(1968KE02)
3.714 \pm 4	5.5 \pm 1	0.20	$\frac{5}{2}^{+}$	9.201	(1971BA06, 1973BA10), A
4.096 \pm 4	15 \pm 1	0.85	$\frac{5}{2}^{-}$	9.493	(1973BA10), A
(4.3)			$\frac{3}{2}^{-}$	(9.6)	(1971BA06)
4.394 \pm 5	16 \pm 1	0.70	$\frac{7}{2}^{+}$	9.720	(1971BA06, 1973BA10), A
4.465 \pm 15	≈ 25	0.90	$\frac{3}{2}^{+}$	9.775	(1971BA06, 1973BA10), A
4.583 \pm 5	14			9.865	(1971BA06, 1973BA10), A
4.600 \pm 15	≈ 10			9.878	(1971BA06, 1973BA10), A
4.730 \pm 20	≈ 80	0.78	$\frac{5}{2}^{+}$	9.977	(1971BA06, 1973BA10), A
4.820 \pm 20	≈ 100			10.046	(1973BA10)
(4.94)	138	0.85	$\frac{5}{2}^{+}$	(10.14)	(1968KE02)
4.993 \pm 5	45	0.15	$\frac{7}{2}^{-}$	10.178	(1971BA06, 1973BA10), A
(5.08)	122	0.60	$\frac{7}{2}^{+}$	(10.2)	A

Table 17.8: Resonances in $^{13}\text{C}(\alpha, n)^{16}\text{O}$ and $^{13}\text{C}(\alpha, \alpha)^{13}\text{C}$ ^a (continued)

E_{res} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Γ_{α}/Γ	J^{π}	E_x (MeV)	Refs.
5.200 \pm 15	150		$\frac{5}{2}^+, \frac{7}{2}^-$	10.337	(1973BA10), A
5.321 \pm 7	14 \pm 3			10.429	(1973BA10), A
5.40	75 \pm 30		$\frac{5}{2}^+, \frac{7}{2}^-$	10.49	A
5.496 \pm 10	47 \pm 15		$\frac{7}{2}^-, \frac{9}{2}^+$	10.563	A
(5.68)	\leq 25	1.00	$(\frac{7}{2}^+)$	(10.70)	(1968KE02)
5.771 \pm 10	80 \pm 20		$\frac{1}{2}^+, \frac{7}{2}^-$	10.773	A
5.945 \pm 10	57 \pm 15		$\frac{5}{2}$	10.906	A
6.107 \pm 10	45 \pm 10			11.030	A
6.167	5.0 \pm 1.1		$\frac{1}{2}^-; T = \frac{3}{2}$	11.076 \pm 0.005	(1976MC11)
6.367 \pm 10	100 \pm 30			11.229	A
6.878 \pm 10	120 \pm 30			11.619	A
7.051 \pm 10	40 \pm 25			11.752	A
7.136 \pm 15	12 \pm 3			11.817	A
7.384 \pm 15				12.006	A
7.52 \pm 20	150 \pm 50			12.11	(1963SP02)
7.736 \pm 15	100 \pm 30			12.275	A
7.88 \pm 20				12.39	(1963SP02)
7.927 \pm 15				12.421	(1963SP02)
7.975	8 \pm 2		$\frac{3}{2}^-; T = \frac{3}{2}$	12.458 \pm 0.005	(1976MC11)
8.156 \pm 15	75 \pm 30			12.596	(1963SP02)
8.253 \pm 15	\approx 5			12.670	(1963SP02)
8.44 \pm 25				12.81	(1963SP02)
8.59 \pm 20	\gtrsim 150			12.93	(1963SP02)
8.611	6 \pm 2		$\frac{1}{2}^+; T = \frac{3}{2}$	12.944 \pm 0.006	(1976MC11)
8.675	\leq 3		$\frac{5}{2}^-; T = \frac{3}{2}$	12.993 \pm 0.006	(1976MC11)
8.72 \pm 20				13.03	(1963SP02)
8.785 \pm 15	16 \pm 4			13.077	(1963SP02)
9.319 \pm 15	\approx 120			13.485	(1963SP02)
9.483 \pm 15	250 \pm 100			13.610	(1963SP02)

A: see references listed for this state in Table 17.6 of (1971AJ02).

^a See also Table 17.6 in (1971AJ02) and Table 17.12 here.

$$12. \text{}^{13}\text{C}(\alpha, t)\text{}^{14}\text{N} \quad Q_m = -12.2640 \quad E_b = 6.361$$

See (1974DM01). See also (1976LE1K).

$$13. \text{}^{13}\text{C}({}^6\text{Li}, d)\text{}^{17}\text{O} \quad Q_m = 4.887$$

At $E({}^6\text{Li}) = 18$ MeV angular distributions have been measured to many ${}^{17}\text{O}$ states: see Table 17.7 here and Table 17.8 in (1971AJ02) (1970BE31, 1971BE2D). Angular distributions are also reported at $E({}^6\text{Li}) = 25.6$ MeV (1971GO1C: $d_0 \rightarrow d_3$ and d to ${}^{17}\text{O}^*(7.5, 8.9)$). See also (1973OG1A).

$$14. \text{}^{13}\text{C}({}^7\text{Li}, t)\text{}^{17}\text{O} \quad Q_m = 3.894$$

Angular distributions have been measured at $E({}^7\text{Li}) = 17$ MeV (1970BE31), 20.5 MeV (1971BE2D, 1971SC21; DWBA analysis) and 30.1 MeV (1971GO1C): see Table 17.7 here and Table 17.8 in (1971AJ02).

$$15. \text{}^{13}\text{C}({}^{16}\text{O}, \text{}^{12}\text{C})\text{}^{17}\text{O} \quad Q_m = -0.801$$

Angular distributions have been studied at $E({}^{16}\text{O}) = 14$ MeV (${}^{17}\text{O}_{\text{g.s.}}$) and 17 and 20 MeV (${}^{17}\text{O}^*(0, 0.87)$) (1970BA49). The excitation curve for the one-neutron transfer to ${}^{17}\text{O}^*(0.87)$ has been measured for $E({}^{16}\text{O}) = 12$ to 25 MeV (1970BA55). See also (1971BA68) and (1973DE21).

$$16. \text{}^{14}\text{C}({}^3\text{He}, \gamma)\text{}^{17}\text{O} \quad Q_m = 18.762$$

The capture cross sections at 90° for γ_0 and for γ_1 have been studied for $E({}^3\text{He}) = 3.2$ to 7.5 MeV and angular distributions of the γ -rays have been studied at the six observed resonances: see Table 17.9 (1976CH04).

Table 17.9: States of ^{17}O from $^{14}\text{C} + ^3\text{He}$

E_{res} (MeV)	Resonant for	$\Gamma_{\text{c.m.}}$ (MeV)	E_x	J^π	Refs.
3.6 ± 0.1	$\gamma_0, (\gamma_1), \alpha_0, \alpha_1$	0.75	21.7	$\frac{5}{2}^+$	(1972KE08, 1976CH04)
4.1 ± 0.1	$\gamma_0, n_0, n_{3+4}, \alpha_0, \alpha_1$	0.75	22.1	$\frac{7}{2}^-$	(1970HO08, 1972KE08, 1976CH04)
4.6 ± 0.2	γ_1	≈ 1	22.5	$\frac{3}{2}^{(-)}$	(1976CH04)
5.1 ± 0.1	$\gamma_0, ^3\text{He}$	≈ 0.4	23.0	$\frac{1}{2}^+$	(1972KE08, 1976CH04)
5.7 ± 0.1	γ_1		23.5		(1976CH04)
6.9 ± 0.1	γ_1		24.4		(1976CH04)

17. (a) $^{14}\text{C}(^3\text{He}, n)^{16}\text{O}$ $Q_m = 14.6167$ $E_b = 18.762$
 (b) $^{14}\text{C}(^3\text{He}, p)^{16}\text{N}$ $Q_m = 4.981$
 (c) $^{14}\text{C}(^3\text{He}, d)^{15}\text{N}$ $Q_m = 4.7136$
 (d) $^{14}\text{C}(^3\text{He}, ^3\text{He})^{14}\text{C}$
 (e) $^{14}\text{C}(^3\text{He}, \alpha)^{13}\text{C}$ $Q_m = 12.4015$

The excitation function for reaction (a) shows a resonance in the n_0 and n_{3+4} yields at $E(^3\text{He}) = 4.1$ MeV but not in the n_{1+2} yield: $J^\pi = \frac{1}{2}^-$ or $\frac{3}{2}^-$ is suggested [see, however, Table 17.9] (1970HO08).

Resonances are observed in the ^3He yield (reaction (d)) at $E(^3\text{He}) = 5.1$ MeV and in the α_0 and α_1 yield (reaction (e)) at 3.6 and 4.1 MeV. On the basis of a two-level analysis of the α -channel and an optical model plus resonance analysis of the elastic data, the corresponding ^{17}O states [21.7, 22.1, 23.0 MeV] are assigned $J^\pi = \frac{5}{2}^+, \frac{7}{2}^-$ and $\frac{1}{2}^+$, respectively (1971KE08, 1972KE08): see Table 17.9. (1972KE08) also report excitation functions in the range $E(^3\text{He}) = 2.2 - 7.0$ MeV ($\alpha_0 \rightarrow \alpha_3$), 3.2 - 4.4 MeV ($p_0 \rightarrow p_3$), 3.2 - 5.5 MeV (d) and 4.0 to 6.1 MeV (^3He): angular distributions for the α -groups have been measured at a number of energies. See also (1971AJ02).

18. $^{14}\text{C}(\alpha, n)^{17}\text{O}$ $Q_m = -1.816$

The upper limits to the decays $3.06 \rightarrow 0$ and $3.84 \rightarrow 0.87$ are, respectively, 2 and 5%. A study of n - γ correlations leads to $J^\pi = \frac{1}{2}^-$ and $(\frac{5}{2}^-)$ for $^{17}\text{O}^*(3.06, 3.84)$ (1964AL11). See also (1973CL1E; astrophys. considerations).

19. $^{14}\text{C}(^{16}\text{O}, ^{13}\text{C})^{17}\text{O}$ $Q_m = -4.032$

Angular distributions have been measured at $E(^{16}\text{O}) = 20, 25$ and 30 MeV to $^{17}\text{O}_{\text{g.s.}}$ and at 30 MeV to $^{17}\text{O}^*(0.87)$ (1975SC35, 1975SC42). See also (1973BR1C).

20. (a) $^{14}\text{N}(t, \gamma)^{17}\text{O}$	$Q_{\text{m}} = 18.625$	
(b) $^{14}\text{N}(t, p)^{16}\text{N}$	$Q_{\text{m}} = 4.843$	$E_{\text{b}} = 18.625$
(c) $^{14}\text{N}(t, d)^{15}\text{N}$	$Q_{\text{m}} = 4.5761$	
(d) $^{14}\text{N}(t, t)^{14}\text{N}$		
(e) $^{14}\text{N}(t, \alpha)^{13}\text{C}$	$Q_{\text{m}} = 12.2640$	

The excitation functions for γ_0 and γ_1 have been measured at 90° for $E_{\text{t}} = 1.5$ to 3.6 MeV: a broad resonance is observed in the γ_0 yield corresponding to $^{17}\text{O}^*(20.45)$. Some evidence is also reported for structures in the γ_1 yield (1973LI1G: prelim. results). Excitation functions have also been measured for the $p_0 \rightarrow p_3$, d_0 , t_0 and α_0 and α_1 groups for $E_{\text{t}} = 1.0$ to 2.0 MeV: the reactions appear to proceed primarily via a direct interaction mechanism (1964SC09). See also (1974FA1A; theor.).

21. (a) $^{14}\text{N}(\alpha, p)^{17}\text{O}$	$Q_{\text{m}} = -1.190$
(b) $^{14}\text{N}(\alpha, p\alpha)^{13}\text{C}$	$Q_{\text{m}} = -7.551$

Angular distributions have been measured for ^{17}O states with $E_{\text{x}} < 7.6$ MeV in the range $E_{\text{p}} = 8.1 - 33.3$ MeV: see a listing of the references in (1971AJ02). The sequential decay (reaction (b)) appears to take place via ^{17}O states with $8.46 \leq E_{\text{x}} \leq 13.57$ MeV. Those involved are believed to have $J \geq \frac{5}{2}$, $\Gamma_{\alpha}/\Gamma \geq 0.6$ (1969BA17). See also ^{18}F in (1978AJ03) and (1971BU1K; theor.).

22. $^{14}\text{N}(^6\text{Li}, ^3\text{He})^{17}\text{O}$	$Q_{\text{m}} = 2.830$
--	------------------------

At $E(^6\text{Li}) = 18$ MeV, the ^3He groups in this reaction and the triton groups in the mirror reaction (see ^{17}F , reaction 7) have been compared: $^{17}\text{O}^*(3.84, 4.55, 5.22, 5.70 + 5.73)$ are strongly excited. It is suggested that $^{17}\text{O}^*(5.22)$ and $^{17}\text{F}^*(5.21)$ are analogs with $J^\pi = \frac{9}{2}^-$. $^{17}\text{O}^*(0, 0.87, 3.06, 5.08, 5.38, 5.87 + 5.94, 6.86, 6.97)$ are also populated (1973BI01).

23. $^{14}\text{N}(^{10}\text{B}, ^7\text{Be})^{17}\text{O}$	$Q_{\text{m}} = -0.044$
--	-------------------------

At $E(^{10}\text{B}) = 100$ MeV $^{17}\text{O}^*(7.75, 9.15)$ [$J^\pi = \frac{11}{2}^-$ and $\frac{9}{2}^-$, respectively] are populated (1976HA1W).

24. $^{15}\text{N}(\text{d}, \gamma)^{17}\text{O}$ $Q_{\text{m}} = 14.049$

Peaks in the excitation functions of both γ_0 and γ_1 have been observed corresponding to $^{17}\text{O}^*(21.3, 21.9, 22.8)$ (1972CA1M; abstract).

25. $^{15}\text{N}(\text{d}, \text{n})^{16}\text{O}$ $Q_{\text{m}} = 9.9031$ $E_{\text{b}} = 14.049$

The excitation function has been measured for $E_{\text{d}} = 0.5$ to 5.3 MeV. Above $E_{\text{d}} = 1$ MeV, pronounced peaks are observed, presumably to be ascribed to numerous overlapping resonances (1958WE31). The differential cross sections at 10° have been measured for $E_{\text{d}} = 2.2$ to 5.9 MeV for the neutrons to $^{16}\text{O}^*(0, 6.13, 7.12, 8.87, 9.85, 10.95, 11.08)$ (1971MU09). Polarization measurements have been reported for $E_{\text{n}} = 1.6$ to 5.5 MeV (n_0 : see (1971AJ02)) and at 4.83 MeV (1972FO17: n to $^{16}\text{O}^*(0, 6.13, 7.12, 8.87, 9.85, 10.95+11.08, 11.52+11.60, 12.05, 12.44+12.53, 12.80)$) and 10.0 and 11.8 MeV (1971HI09; n_0). See also (1973BO1G) and ^{16}O , and (1971HA1R).

26. $^{15}\text{N}(\text{d}, \text{p})^{16}\text{N}$ $Q_{\text{m}} = 0.267$ $E_{\text{b}} = 14.049$

Excitation functions have been obtained for $E_{\text{d}} = 0.3$ to 2.7 MeV (1957BO04) and 4.25 to 6.25 MeV (1973BO1G: $p_0 \rightarrow p_3$). Resonant structure at $E_{\text{d}} = 1.3$ and 1.9 MeV [$^{17}\text{O}^*(15.2, 15.7)$] is reported by (1957BO04).

27. $^{15}\text{N}(\text{d}, \text{d})^{15}\text{N}$ $E_{\text{b}} = 14.049$

The excitation function for the d_0 group has been measured for $E_{\text{d}} = 4.25$ to 6.25 MeV (1973BO1G).

28. $^{15}\text{N}(\text{d}, \alpha)^{13}\text{C}$ $Q_{\text{m}} = 7.6879$ $E_{\text{b}} = 14.049$

The α_0 yield curve for $E_{\text{d}} = 0.8$ to 1.8 MeV indicates two resonances at $E_{\text{d}} = 1.06$ and 1.25 MeV [$\Gamma \approx 100$ and 200 keV, respectively], attributed to an ^{17}O state at $E_{\text{x}} = 14.98$ MeV [$J^\pi = \frac{5}{2}^+$] and to one or more ^{17}O states at $E_{\text{x}} = 15.15$ MeV [$J^\pi = \frac{5}{2}^+$ or $\frac{7}{2}^-$] (1966TI03). In the range $E_{\text{d}} = 1.2$ to 2.5 MeV a broad maximum is observed in both the α_0 and α_1 yields at $E_{\text{d}} \approx 1.7$ MeV (1965MA1A). The excitation function has also been studied for $E_{\text{d}} = 5$ to 10 MeV (1974WE1P).

29. $^{15}\text{N}(t, n)^{17}\text{O}$ $Q_m = 7.791$

Not reported.

30. $^{15}\text{N}(^3\text{He}, p)^{17}\text{O}$ $Q_m = 8.555$

At $E(^3\text{He}) = 18$ MeV angular distributions analyzed by DWBA, have been obtained for a number of ^{17}O states as displayed in Table 17.10: the wave functions of Zuker, Buck and McGroary appear to provide a good description of most of the states below $E_x = 9$ MeV (1972LE01). Angular distributions have also been reported at $E(^3\text{He}) = 2.4$ to 4.1 MeV (1973AB1D: $p_0 \rightarrow p_2$) and at 15 MeV (1975HA33: p_0, p_2). For the decay of the first $T = \frac{3}{2}$ state at $E_x = 11.08$ MeV, see Table 17.11 (1973AD02).

31. $^{15}\text{N}(\alpha, d)^{17}\text{O}$ $Q_m = -9.799$

At $E_\alpha = 45.4$ MeV, the deuteron spectrum is dominated by the groups corresponding to states with $E_x = 7.742 \pm 0.020$ and 9.137 ± 0.030 MeV. These states are assigned $J^\pi = (\frac{11}{2}^-)$ and $(\frac{9}{2}^-)$ and arise from a dominant $(d_{5/2})^2_5 p_{1/2}^{-1}$ configuration. Angular distributions were measured as well for the deuterons corresponding to $^{17}\text{O}(0)$ and to states with $E_x = 0.87 \pm 0.05, 5.208 \pm 0.030, 5.690 \pm 0.030, 7.367 \pm 0.030, 8.459 \pm 0.030, 8.890 \pm 0.030$ and 9.814 ± 0.030 MeV. In addition the excitation of states with $E_x = 3.85 \pm 0.05, 4.57 \pm 0.05$ and 8.147 ± 0.030 MeV is also reported (1969LU07).

32. $^{15}\text{N}(^{11}\text{B}, ^9\text{Be})^{17}\text{O}$ $Q_m = -1.768$

At $E(^{11}\text{B}) = 113.5$ MeV, ^{17}O states at 7.6 and 9.0 MeV are strongly populated. The excitation of $^{17}\text{O}^*(0, 0.9, 3.0, 5.4)$ is also reported (1975PO10).

33. $^{16}\text{O}(n, \gamma)^{17}\text{O}$ $Q_m = 4.146$

$$\sigma_{\text{capt.}} = 178 \pm 25 \mu\text{b} [\text{see } (1973\text{MU}14)], 202 \pm 27 \mu\text{b} (1976\text{EA}1\text{B}).$$

At the $E_n = 426 \pm 10$ keV resonance [see Table 17.12], $\Gamma_\gamma < 4.0$ eV, $\Gamma_n = 60 \pm 15$ keV (1971AL09). At thermal energies the branching via $^{17}\text{O}^*(3.05)$ is $82 \pm 3\%$ (1976EA1B). For astrophysical considerations see (1968FO1A, 1970CL1C, 1971AL09, 1973CL1E).

Table 17.10: Levels of ^{17}O from $^{15}\text{N}(^3\text{He}, \text{p})^{17}\text{O}$ ^a

E_x ^b (MeV)	L	E_x ^b (MeV)	L
0	(1 + 3)	8.192	0
0.874	1	8.322	
3.053	0	8.390	
3.845	2	8.492	(2)
4.549	0	8.682	
5.081	(1)	8.900	
5.215	(4)	8.955	
5.381	0	9.16	(4)
5.698	2	9.495	
5.873	(1)	9.712	
5.938	0	9.856	
6.37		(10.24)	
6.861	(0)	10.33	
6.973	(1 + 3)	10.57	
7.162	2	10.782	
7.382	2	10.913	
7.561		11.032 ± 0.004 ^c	
7.687		11.075 ± 0.004 ^d	
7.761	4		
7.938			
8.054	(1)		

^a (1972LE01).

^b ± 10 keV, except where shown otherwise.

^c See also (1970MC02): $T = \frac{1}{2}$.

^d $J^\pi = \frac{1}{2}^-$; $T = \frac{3}{2}$: see Table 17.11 (1972LE01, 1973AD02).

Table 17.11: Decay properties of the lowest $T = \frac{3}{2}$ states in $A = 17$ ^a

		$^{17}\text{O}^*(11.076 \pm 0.004)$	$^{17}\text{F}^*(11.1931 \pm 0.0023)$ ^c
J^π		$\frac{1}{2}^-$	$\frac{1}{2}^-$
$\Gamma_{\text{c.m.}}$ (keV)		5.0 ± 1.1 ^b	0.20 ± 0.04
Branching ratio (%) to $^{16}\text{O}^*$ (MeV)	J^π		
0	0^+	91 ± 15	8.8 ± 1.6
6.05	0^+	} 5 ± 2	< 3
6.13	3^-		22 ± 2
6.92	2^+		24 ± 6
7.12	1^-		44 ± 4
$^{13}\text{C} + \alpha_0$ or $^{13}\text{N} + \alpha_0$		6	< 7
Partial widths [Γ_p or Γ_n] to			
$^{16}\text{O}(0)$		4.5 ± 1.2 keV	19 ± 3 eV
$^{16}\text{O}^*(6.05)$		} 0.25 ± 0.11 keV	< 17 eV
$^{16}\text{O}^*(6.13)$			95_{-50}^{+30} eV ^d
$^{16}\text{O}^*(6.92)$			100_{-60}^{+40} eV ^d
$^{16}\text{O}^*(7.12)$			190_{-100}^{+60} eV ^d
Γ_{α_0}		0.3 keV	< 40 eV
Γ_{γ_1}			6.0 ± 2.5 eV
$\theta^2(\text{g.s.})/\theta^2(6.13)$		0.31 ± 0.14	0.065 ± 0.019

^a See also Table 2 in (1973AD02) and reaction 63.

^b (1976MC11).

^c (1971HA05, 1973AD02, 1974SK02, 1975HA06, 1976HI09).

^d Note that the total width is 200 ± 40 eV.

34. $^{16}\text{O}(n, n)^{16}\text{O}$ $E_b = 4.146$

The scattering amplitude bound is $a = 5.80 \pm 0.05$ fm (1965DO1B). The coherent scattering cross section is 4.23 ± 0.07 b (1964ST25). Earlier cross section, angular distribution and polarization measurements are listed in Table 17.10 of (1971AJ02). Recent measurements of σ_t are reported by (1972PE1E: 0.2 – 20 MeV), (1974SC1C: 0.5 – 20.5 MeV), (1973FO11: 0.6 – 0.9, 1.12 – 1.16, 1.39 – 4.33 MeV), (1971FO1A: 2.5 – 15.0 MeV), (1972KI1D: elastic; 4.34 – 8.56 MeV), (1971AN16: 14.7 MeV), (1972AU01: 24.63 – 58.90 MeV) and (1974BA52, 1975BA2L: 28 – 54 GeV). See also (1976GAYV). (1973BU1D, 1974BU19) report $\sigma(\theta)$ at small angles for $E_n = 7.40$ to 9.50 MeV. Polarization studies are reported by (1973HI09, 1974HI1B: $E_n = 1 - 4$ MeV) and (1976DR08: 2.25 to 3.90 MeV).

Recent high resolution cross section measurements and analyses of the elastic scattering and of the (n, α) and $^{13}\text{C}(\alpha, n)$ data have led to a much better understanding of the ^{17}O structure below $E_x = 9.5$ MeV: see Table 17.12 (1973FO11, 1973JO01). (1973JO01) has performed a multilevel two-channel R -matrix analysis. Five states contain nearly 100% of the $1d_{3/2}$ strength and have their eigenenergy at $E_x \approx 5.7$ MeV [the dominant state is $^{17}\text{O}^*(5.08)$]. Spectroscopic factors are deduced for 26 states in ^{17}O for $4.5 < E_x < 9.5$ MeV [see Table 17.12]: the sum of these factors is 1% for $J^\pi = \frac{1}{2}^+$, 5% for $\frac{1}{2}^-$, 12% for $\frac{3}{2}^-$, 99% for $\frac{3}{2}^+$, 0.1% for $\frac{5}{2}^+$, 1% for $\frac{5}{2}^-$ and 14% for $\frac{7}{2}^-$ (1973JO01).

See also ^{16}O , (1971SE1D), (1971AJ02, 1972LA1F, 1973MU14, 1974SA1N) and (1970GN1A, 1970TI02, 1970VA1M, 1971DO15, 1971LE1B, 1971MA05, 1971MA62, 1971SC22, 1972HA2A, 1972LE1M, 1972PH06, 1972SC45, 1973LE1K, 1973SC1R, 1973ST05, 1973WE06, 1974GE03, 1974HI04, 1974MU1F, 1974PH03, 1975BE06, 1975CA05, 1975GE08, 1975PH01, 1975TH12; theor.).

35. $^{16}\text{O}(n, n')^{16}\text{O}^*$ $E_b = 4.146$

Earlier cross section measurements are listed in Table 17.10 of (1971AJ02). The cross sections for production of 6.13 and $(6.92 + 7.12)$ γ -rays in the range $E_n = 6.5$ to 10 MeV show a number of resonances: see Table 17.13 (1959HA13). See also the more recent studies of (1970LU16: $E_n = 6.74$ to 8.20 MeV; $\gamma_{6.13}$) and (1970OR1B: $E_n = 6.35$ to 16.52 MeV; $\gamma_{6.13}$, $\gamma_{6.92}$, $\gamma_{7.12}$). See also (1969RO1F, 1972KI1D, 1976NO1F).

36. $^{16}\text{O}(n, 2n)^{15}\text{O}$ $Q_m = -15.664$ $E_b = 4.146$

The cross section has been measured for $E_n = 17$ to 37 MeV (1961BR1A). See also (1974CA1J) and (1976GAYV).

37. $^{16}\text{O}(n, p)^{16}\text{N}$ $Q_m = -9.636$ $E_b = 4.146$

Table 17.12: Resonances ^a in ¹⁶O(n, n)¹⁶O and ¹⁶O(n, α)¹³C

E_n ^b (keV)	$\Gamma_{c.m.}$ ^b (keV)	$\Gamma_{\lambda n}$ ^c (keV)	$\Gamma_{\lambda\alpha}$ ^c (keV)	$\theta_{\lambda n}^2$ ^c (%)	$J\pi$ ^b	E_x (MeV)
433 ± 2 ^j	45	45		4.4	$\frac{3}{2}^-$	4.553
1000 ± 2 ^j	96	96		68.9 ⁱ	$\frac{3}{2}^+$	5.086
1140 ^d	< 0.1					5.218
1312 ± 2 ^j	42	41.5		0.91	$\frac{3}{2}^-$	5.380
1651 ± 2	3.4 ± 0.3	3.4		9.4	$\frac{7}{2}^-$	5.698
1689 ± 2	< 1				e	5.734
1833 ± 2	6.6 ± 0.7	6.6		0.95	$\frac{3}{2}^+$	5.870
1908 ± 4	32 ± 3	31.5		0.51	$\frac{1}{2}^-$	5.940
2351 ± 8 ^h	124 ± 12	124		0.81	$\frac{1}{2}^+$	6.357
2889 ± 2	< 1				e	6.863
3006 ± 2	< 1				e	6.973
3211 ± 3	1.3 ± 0.4	1.3	0.0033	0.44	$(\frac{5}{2}^-)$	7.166
3250 ± 10	280 ± 30	280	0.07	16.9	$\frac{3}{2}^+$	7.202
3438 ± 3	0.5 ± 0.2	0.5	0.01	0.01	$(\frac{5}{2}^+)$	7.379
3441 ± 3	1.1 ± 0.4	1.1	0.003	0.31	$\frac{5}{2}^-$	7.382
3630 ± 20	500 ± 50	500	0.08	5.1	$\frac{3}{2}^-$	7.560
3647 ^d	< 0.1					7.576
3766 ± 4	18 ± 2	18	0.01	2.9	$\frac{7}{2}^-$	7.688
4053 ± 8 ^f	90 ± 9	84	6.7	0.47	$\frac{1}{2}^+$	7.958
4090 ± 50 ^f	270 ± 30	250	16	3.4	$\frac{1}{2}^-$	7.99
4162 ± 8 ^f	85 ± 9	71	15	8.5	$\frac{3}{2}^+$	8.060
4290 ± 20 ^f	69 ± 7	68	0.8	0.68	$\frac{1}{2}^-$	(8.18)
4310 ± 10 ^f	52	48	4.0	0.43	$(\frac{3}{2}^-)$	8.199
4470	12	10	2.2	0.06	$\frac{1}{2}^+$	8.350
4532	5	4.8	0.54	0.8	$\frac{5}{2}^+$	8.408
4600	8		7.6	0.53	$\frac{7}{2}^+$	8.472
4610	≤ 11				≥ $\frac{3}{2}^+$	8.481
4637	5	3.4	1.9	0.40	$\frac{5}{2}^-$	8.507
4830 ^f	44	42	1.8	0.36	$\frac{3}{2}^-$	8.688
5050 ^f	78	68	9.5	4.2	$\frac{3}{2}^+$	8.895

Table 17.12: Resonances ^a in ¹⁶O(n, n)¹⁶O and ¹⁶O(n, α)¹³C (continued)

E_n ^b (keV)	$\Gamma_{c.m.}$ ^b (keV)	$\Gamma_{\lambda n}$ ^c (keV)	$\Gamma_{\lambda\alpha}$ ^c (keV)	$\theta_{\lambda n}^2$ ^c (%)	J^π ^b	E_x (MeV)
5131 ^f	25	23	2.3	1.5	$\frac{7}{2}^-$	8.971
5320	7				$\frac{1}{2}^-$	9.149
5360	3				$\frac{7}{2}^-$	9.187
5370	7				$\frac{5}{2}^+$	9.196
5610 ^f	120	120		0.91	$\frac{3}{2}^-$	9.422
5640 ^f	15				$\frac{5}{2}^-$	9.450
5640 ^g	140				$\geq \frac{3}{2}^-$	9.450
5914 ± 5	28				$\geq \frac{5}{2}^-$	9.708
6010	28				$\geq \frac{3}{2}^-$	9.798
6100	25				$\geq \frac{1}{2}^-$	9.88
6395 ± 7	38				$\geq \frac{3}{2}^-$	10.160
6807 ± 7	40				$\geq \frac{3}{2}^-$	10.547
7200 ± 8	70				$\geq \frac{3}{2}^-$	10.917
7830	190				$\geq \frac{3}{2}^-$	11.509
8320	270				$\geq \frac{3}{2}^-$	11.970
8740	130					12.365
9050	95					12.656
10130	400					13.672
11140	340				$(\geq \frac{3}{2}^-)$	14.621
11540	180					14.997

^a See also Table 17.8 here and Table 17.11 in (1971AJ02).

^b See (1973FO11, 1973JO01). See (1971AJ02) for the earlier values.

^c See (1973JO01).

^d Not observed in σ_1 : see (1973FO11).

^e Not $\frac{1}{2}^+$ (1973FO11).

^f See also (1974SC1C).

^g For this resonance and all the following ones see (1961FO07, 1969DA13, 1974SC1C).

^h See also (1971WE08, 1973WE06).

ⁱ $S = 0.8 \pm 0.1$ (1974CO10).

^j C.H. Johnson, private communication.

Table 17.13: Resonances in $^{16}\text{O}(n, n'\gamma)^{16}\text{O}$
(1959HA13, 1970LU16)

E_n (MeV)	E_x (MeV)
6.85 ^a	10.59
7.07	10.79
7.24	10.95
7.42	11.12
7.85 ^b	11.53
8.35	12.00
8.50	12.14
8.84	12.46
9.10	12.70
9.34	12.93

^a Proposed $J^\pi = \frac{7}{2}^-$ (1970LU16).

^b Proposed $J = \frac{3}{2}$ or $\frac{5}{2}$ (1970LU16).

Cross section measurements are listed in Table 17.10 of (1971AJ02) and in (1975SA1D; $E_n = 14.1$ MeV). See also (1971AJ02, 1971CU1B, 1971PR09, 1972ED01, 1974BO1E, 1974CA1J, 1976GAYV).

$$38. \ ^{16}\text{O}(n, d)^{15}\text{N} \qquad Q_m = -9.9031 \qquad E_b = 4.146$$

See ^{15}N in (1976AJ04).

$$39. \ ^{16}\text{O}(n, \alpha)^{13}\text{C} \qquad Q_m = -2.2152 \qquad E_b = 4.146$$

The cross section has been measured from threshold to 20 MeV: see Table 17.10 in (1971AJ02) and (1970OR1B: 8–15.5 MeV; $\gamma_{3.09}, \gamma_{3.68}, \gamma_{3.85}$) and (1973BO26: 14.1 MeV). See also (1968DA1E, 1971BR33, 1971NY03). Table 17.12 reflects the results from a multilevel two-channel R -matrix analysis of the data from this reaction and from the elastic scattering of neutrons (1973JO01). See also (1969RO1F, 1976NO1F), (1971AJ02, 1976GAYV) and (1972HA2A; theor.).

$$40. \ ^{16}\text{O}(p, \pi^+)^{17}\text{O} \qquad Q_m = -136.205$$

At $E_p = 185$ MeV angular distributions of pions to $^{17}\text{O}^*(0, 0.88 \pm 0.08, 3.85 \pm 0.08)$ are reported by (1974DA23). See also (1976NO1D; theor.).

$$41. \ ^{16}\text{O}(d, p)^{17}\text{O} \quad Q_m = 1.921$$

Observed proton groups are displayed in Table 17.14. Angular distributions have been measured at many energies in the range of $E_d = 0.3$ to 63.2 MeV: see Table 17.13 in (1971AJ02) for the earlier references and (1973CA30: $E_d = 1.0$ to 2.0 MeV; p_0, p_1), (DeForest, quoted in (1971KO21): 8 MeV; p_0, p_1), (1972CO15, 1973DA17: 9.3 and 13.3 MeV; $p_0, p_1, p_3, p_4, p_5, p_8$), (1972BR12: 12.3 MeV; p_0, p_1) and (1974CO04: 25.4, 36.0, 63.2 MeV: p_0, p_1, p_5).

The lifetime of $^{17}\text{O}^*(0.87)$ is 258.6 ± 2.6 psec: see Table 17.7 in (1971AJ02). See also (1971DO13, 1973DO1D). $E_\gamma = 870.81 \pm 0.22$ keV (1966WI01). The width of $^{17}\text{O}^*(5.08)$ observed in this reaction ($\Gamma = 70$ keV) is ≈ 0.7 that observed in $^{16}\text{O}(n, n)$: see (1974CO04, 1974FO17).

See also ^{18}F in (1978AJ03), (1970CA1C, 1971GR2B, 1972PR1D), (1976SC1G; applied work) and (1970DO10, 1970KI15, 1970KU1B, 1970OH1C, 1971BO50, 1971CO1B, 1971DO1A, 1972BU23, 1972DZ06, 1972FR1E, 1972GO04, 1972PH06, 1972SC45, 1972SC20, 1973BA74, 1973DO02, 1974BA19, 1974CO10, 1974GO02, 1974IM01, 1974OR1A, 1975CO12, 1976BO15, 1976SH13; theor.).

$$42. \ ^{16}\text{O}(t, d)^{17}\text{O} \quad Q_m = -2.112$$

The angular distribution of the d_0 group has been studied at $E_t = 5.5$ MeV (1961BA10).

$$43. \ ^{16}\text{O}(\alpha, ^3\text{He})^{17}\text{O} \quad Q_m = -16.433$$

See (1971AJ02, 1972PR1D, 1976HA27).

$$44. \ ^{16}\text{O}(^6\text{Li}, p\alpha)^{17}\text{O} \quad Q_m = 0.447$$

See (1974MI1F).

$$45. \ ^{16}\text{O}(^7\text{Li}, ^6\text{Li})^{17}\text{O} \quad Q_m = -3.105$$

Table 17.14: States of ^{17}O from $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ and $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$

E_x^a (MeV \pm keV)	$\Gamma_{\text{c.m.}}^a$ (keV)	E_x^b (MeV \pm keV)	E_x^c (MeV \pm keV)	S^d	J^π
0	< 8	0	0	≈ 0.9	$\frac{5}{2}^+$
0.871 ± 4^e	< 8	0.870 ± 20	0.883 ± 11	≈ 0.9	$\frac{1}{2}^+$
3.055 ± 4^e	< 8	3.060 ± 30	3.069 ± 10		$\frac{1}{2}^-$
3.846 ± 5^e	< 8	3.850 ± 30	3.856 ± 11		$\frac{5}{2}^-^g$
4.553 ± 6	40 ± 5	4.580 ± 20	4.567 ± 14	0.23	$\frac{3}{2}^-$
5.083 ± 10	95 ± 5	5.070 ± 20		1.25	$\frac{3}{2}^+$
5.215 ± 5	< 8		5.229 ± 13		$\frac{3}{2}^h$
5.378 ± 7	28 ± 7	5.310 ± 20	5.397 ± 14		$\frac{3}{2}^-$
5.695 ± 5^f	< 8			≈ 0.15	$\frac{7}{2}^-$
5.731 ± 5^f	< 8	5.760 ± 20	5.723 ± 14		
5.866 ± 5	< 8		5.875 ± 15		
5.940 ± 15	23 ± 10		5.957 ± 15		
		6.240 ± 20			
		6.890 ± 30	6.869 ± 14		
			(6.986 ± 15)		
			(7.371 ± 15)		
		7.510 ± 30			
		8.270 ± 40			
		(8.590 ± 40)			
		9.060 ± 40			

^a $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ (1957BR82).

^b $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ and $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$ (1951BU1A).

^c $^{19}\text{F}(\text{d}, \alpha)^{17}\text{O}$ (1952WA1A).

^d (1973DA17, 1974CO04) [$^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$].

^e (1965GA1A) report $E_x = 873 \pm 5, 3056 \pm 4$ and 3838 ± 4 keV.

^f ΔE_x between $^{17}\text{O}^*(5.73, 5.70) = 34 \pm 2$ keV (1968BI1A).

^g (1965CO07, 1965CO09).

^h (1968BI1A).

The angular distribution involving $^{17}\text{O}_{\text{g.s.}}$ has been studied at $E(^7\text{Li}) = 36$ MeV. The population of $^{17}\text{O}^*(0.87, 3.06, 3.84, 4.55, 5.38)$ is also reported (1973SC26).

$$46. \ ^{16}\text{O}(^{11}\text{B}, ^{10}\text{B})^{17}\text{O} \quad Q_{\text{m}} = -7.311$$

The excitation of $^{17}\text{O}^*(0, 0.87)$ is reported at $E(^{11}\text{B}) = 113.1$ MeV (1967PO13).

$$47. \ ^{16}\text{O}(^{13}\text{C}, ^{12}\text{C})^{17}\text{O} \quad Q_{\text{m}} = -0.801$$

The product of the spectroscopic factors in the initial and final states [$^{17}\text{O}^*(0.87)$] is 0.72 (1975SE03: from measurements of σ_{t} below Coulomb barrier). The angular distribution to $^{17}\text{O}^*(0.87)$ has been measured at $E(^{13}\text{C}) = 36$ MeV (1976WE21). See also (1973BR1C) and (1974BE1J; theor.).

$$48. \ ^{16}\text{O}(^{14}\text{N}, ^{13}\text{N})^{17}\text{O} \quad Q_{\text{m}} = -6.408$$

At $E(^{14}\text{N}) = 79$ MeV angular distributions involving $^{17}\text{O}^*(0, 0.87)$ have been studied: an anomaly is observed in the phase behavior of the distribution to the excited state. From this and other studies it is concluded that a multistep process via inelastic scattering is unlikely to occur in the excitation of $2s_{1/2}$ states (1976MO03). An angular distribution involving $^{16}\text{O}^*(0 + 0.87)$ has also been reported at $E(^{14}\text{N}) = 155$ MeV (1976NA09). See also (1975NA15).

$$49. \ ^{16}\text{O}(^{18}\text{O}, ^{17}\text{O})^{17}\text{O} \quad Q_{\text{m}} = -3.898$$

At $E(^{18}\text{O}) = 42$ and 52 MeV, the angular distributions involving $^{17}\text{O}^*(0, 0.87)$ have been studied (1975RE15).

$$50. \ ^{17}\text{N}(\beta^-)^{17}\text{O} \quad Q_{\text{m}} = 8.682$$

^{17}N decays principally to $^{17}\text{O}^*(4.55, 5.38)$: see reaction 1 and Table 17.2 in ^{17}N .

$$\begin{aligned} 51. \text{ (a) } & \ ^{17}\text{O}(\gamma, \text{n})^{16}\text{O} & Q_{\text{m}} &= -4.146 \\ & \text{ (b) } & \ ^{17}\text{O}(\gamma, 2\text{n})^{15}\text{O} & Q_{\text{m}} &= -19.809 \\ & \text{ (c) } & \ ^{17}\text{O}(\gamma, \alpha)^{13}\text{C} & Q_{\text{m}} &= -6.361 \end{aligned}$$

Table 17.15: $B(E3)$ values from $^{17}\text{O}(e, e')$ ^a

$^{17}\text{O}^*$ (MeV)	J^π	$B(E3)\uparrow$ ($e^2 \cdot \text{fm}^6$)
3.06	$\frac{1}{2}^-$	31 ± 6
3.84	$\frac{5}{2}^-$	153 ± 6
4.55	$\frac{3}{2}^-$	98 ± 8
5.22	$(\frac{9}{2}^-)$	360 ± 11
5.38	$\frac{3}{2}^-$	45 ± 12
5.70	$\frac{7}{2}^-$	270 ± 32
5.94	$\frac{1}{2}^-$	17 ± 10
6.86	$(\frac{1}{2}^-)$	(147 ± 34)
7.17	$\frac{5}{2}^-$	22 ± 25
7.39	$\frac{5}{2}^-$	47 ± 38
7.58	$\frac{7}{2}^-$	109 ± 26
7.75	$(\frac{11}{2}^-)$	369 ± 15

^a (1975KI15).

The (γ, n_0) differential cross section at 98° has been measured for $E_{\text{bs}} = 8$ to 33 MeV by (1976WO1D): both narrow and broad structures are observed throughout this energy range. The (γ, n) cross section has also been measured for $E_\gamma = 7.6$ to 30 MeV: it is dominated by a broad giant resonance centered at $E_x \approx 23$ MeV (1976ME1K). The $^{17}\text{O}(\gamma, 2n)$ cross section is very small for these energies (1976ME1K). For reaction (c) see (1964GR08).

52. $^{17}\text{O}(e, e)^{17}\text{O}$

The ^{17}O charge radius, $r_{\text{rms}} = 2.662 \pm 0.026$ (using a distorted wave approximation), and 2.700 ± 0.026 fm (using a Born approximation) (1970SI02, 1970SI1K). Inelastic scattering in a range of momentum transfer $q = 0.6 - 1.1 \text{ fm}^{-1}$ has led to calculation of $B(E3)$ values from the measured Coulomb form factors: see Table 17.15 (1975KI15). See also (1976AU1G).

53. $^{17}\text{O}(n, n)^{17}\text{O}$

See (1973IS07; theor.).

54. $^{17}\text{O}(p, p)^{17}\text{O}$

Angular distributions of elastically scattered protons have been studied at $E_p = 8.62, 9.45$ and 10.5 MeV (1975CR04; also p_1 and p_2), 11 MeV (1967AL06) and 65.8 MeV (1972LE27, 1972LE28, 1972LE1G). The matter rms radius of ^{17}O is 0.04 ± 0.03 fm greater than that for ^{16}O (1973LE07). See also ^{18}F in (1978AJ03) and (1970OH1C, 1971DO1A, 1975CO12, 1976CO01; theor.).

55. $^{17}\text{O}(d, d)^{17}\text{O}$

The angular distribution of elastically scattered deuterons has been studied at $E_d = 18$ MeV (1976LI01).

56. (a) $^{17}\text{O}(^3\text{He}, ^3\text{He})^{17}\text{O}$
 (b) $^{17}\text{O}(\alpha, \alpha)^{17}\text{O}$

Elastic angular distributions have been measured at $E(^3\text{He}) = 11.0$ MeV (1970BO25) and 17.3 MeV (1968HA30). For reaction (b) see (1976CO27; theor.).

57. (a) $^{17}\text{O}(^{12}\text{C}, ^{12}\text{C})^{17}\text{O}$
 (b) $^{17}\text{O}(^{13}\text{C}, ^{13}\text{C})^{17}\text{O}$

For reaction (a) see (1976EY01). For reaction (b) see (1974CH1Q).

58. $^{17}\text{O}(^{16}\text{O}, ^{16}\text{O})^{17}\text{O}$

Angular distributions involving $^{17}\text{O}^*(0, 0.87)$ have been studied at $E(^{16}\text{O}) = 22, 24, 28$ and 32 MeV (1973GE04, 1974GE01) and at $E(^{17}\text{O}) = 25.7, 27.7, 29.8$ and 32.0 MeV (1975KA24): second-order transfer contributions are found to be important (1975KA24). See also (1973FI1C, 1974GO1L, 1975VO1B) and (1973BA2F, 1974BA46, 1974BE1J, 1974BO13, 1974YU01, 1975IM04, 1975WO1E; theor.).

59. $^{17}\text{F}(\beta^+)^{17}\text{O}$ $Q_m = 2.762$

See ¹⁷F.

60. ¹⁸O(γ , n)¹⁷O $Q_m = -8.044$

See (1975AL03) and ¹⁸O in (1978AJ03).

61. ¹⁸O(p, d)¹⁷O $Q_m = -5.819$

Angular distributions have been measured at $E_p = 17.6$ MeV (1963LE03: $d_0 \rightarrow d_2$), 18.2 MeV (1967LU05: $d_0 \rightarrow d_3$), 20.0, 24.4, 29.8, 37.5 and 43.6 MeV (1974PI05: $d_0 \rightarrow d_4$) and 24.4 MeV (1973PI09: $d_0 \rightarrow d_7$; polarized protons). See also ¹⁹F in (1978AJ03), (1976DA1K) and (1973IG02, 1973OR09, 1973YA1B; theor.).

62. ¹⁸O(d, t)¹⁷O $Q_m = -1.786$

Angular distributions of the tritons corresponding to ¹⁷O*(0, 0.87, 3.84, 4.55, 5.08, 5.38) have been studied at $E_d = 15$ MeV (1961AR06). See also (1975HS01, 1976LA13; theor.).

63. ¹⁸O(³He, α)¹⁷O $Q_m = 12.535$

Angular distributions of alpha particles are reported by (1965WA1D: α_0, α_1) at $E(^3\text{He}) = 2.68$ to 6.47 MeV and by (1969DE06: see Table 17.16) at $E(^3\text{He}) = 16$ MeV. The $T = \frac{3}{2}$ states reported by (1969DE06) are displayed in Table 17.16 [the isospin identification is based on the enhanced excitation and the narrow widths of these states]. The branching ratios for the various decays of ¹⁷O*(11.08) [the lowest $T = \frac{3}{2}$ state in ¹⁷O] and for the analog state in ¹⁷F are displayed in Table 17.11: the decay width of the ¹⁷O state is approximately 200 times greater than that of the ¹⁷F state (1973AD02).

64. ¹⁸O(⁹Be, ¹⁰Be)¹⁷O $Q_m = -1.232$

Angular distributions have been studied at $E(^{18}\text{O}) = 16$ and 20 MeV (1971BA68, 1971KN05).

65. (a) ¹⁸O(¹⁰B, ¹¹B)¹⁷O $Q_m = 3.412$

(b) ¹⁸O(¹¹B, ¹²B)¹⁷O $Q_m = -4.674$

Table 17.16: $T = \frac{3}{2}$ states of ^{17}O from $^{18}\text{O}(^3\text{He}, \alpha)^{17}\text{O}$ ^{a,b}

E_x (MeV \pm keV)	l_n	J^π	C^2S^c
11.082 ± 6	1	$(\frac{1}{2})^-$	0.49
12.471 ± 5	1	$(\frac{3}{2})^-$	0.27
12.950 ± 8	0	$\frac{1}{2}^+$	0.096
12.994 ± 8			
13.640 ± 5	2	$(\frac{5}{2})^+$	0.39
14.219 ± 8			
14.282 ± 12			
15.101 ± 8			

^a See also Table 17.11.

^b (1969DE06).

^c Calculated assuming $C^2S = 4$ for $^{15}\text{O}^*(6.18)$.

Angular distributions (reaction (a)) have been measured at $E(^{18}\text{O}) = 20$ and 24 MeV (1971BA68, 1971KN05). For S -factor measurements see (1974SW04). Cross sections for reaction (b) are several orders of magnitude less than those for reaction (a) for $E(^{18}\text{O})_{\text{c.m.}} = 3-7.7$ MeV (1974SW04).

$$66. \text{ (a) } ^{18}\text{O}(^{12}\text{C}, ^{13}\text{C})^{17}\text{O} \quad Q_m = -3.097$$

$$\text{ (b) } ^{18}\text{O}(^{13}\text{C}, ^{14}\text{C})^{17}\text{O} \quad Q_m = 0.133$$

See (1974CH1Q).

$$67. ^{18}\text{O}(^{14}\text{N}, ^{15}\text{N})^{17}\text{O} \quad Q_m = 2.790$$

See (1974SW04).

$$68. ^{18}\text{O}(^{18}\text{O}, ^{19}\text{O})^{17}\text{O} \quad Q_m = -4.087$$

See (1972EY01).

$$69. ^{19}\text{F}(\pi^-, 2n)^{17}\text{O} \quad Q_m = 122.822$$

The 0.87 MeV γ -ray is observed when ^{19}F captures π^- mesons (1976EN02).

$$70. \ ^{19}\text{F}(\text{n}, \text{t})^{17}\text{O} \quad Q_{\text{m}} = -7.554$$

Angular distributions of the t_0 and t_1 groups are reported at $E_{\text{n}} = 14.4$ MeV (1968RE07).

$$71. \text{ (a) } ^{19}\text{F}(\text{p}, ^3\text{He})^{17}\text{O} \quad Q_{\text{m}} = -8.318$$

$$\text{ (b) } ^{19}\text{F}(\text{p}, \text{pd})^{17}\text{O} \quad Q_{\text{m}} = -13.812$$

Angular distributions have been measured at $E_{\text{p}} = 30.5$ MeV (1967CO05: to $^{17}\text{O}^*(0, 0.87)$) and at 42.4 MeV (1974NE03: to $^{17}\text{O}^*(0, 0.87, 3.06, 3.84)$): comparisons have been made with the analog transitions in the mirror reaction $^{19}\text{F}(\text{p}, \text{t})^{17}\text{F}$. See also (1972HU1B, 1972PR1D), ^{20}Ne in (1978AJ03) and (1971AJ02). For reaction (b) see (1971DE1F).

$$72. \ ^{19}\text{F}(\text{d}, \alpha)^{17}\text{O} \quad Q_{\text{m}} = 10.036$$

Observed α -groups are displayed in Table 17.14. Angular distributions have been measured at many energies in the range $E_{\text{d}} = 0.3$ to 27.5 MeV: see Table 17.16 in (1971AJ02) for the earlier work and (1969ZA1A: 1.10 – 3.65 (α_0, α_1) and 1.40 to 3.65 MeV (α_2, α_3)), (1971BE2F: 1.35 – 2.0 MeV; $\alpha_0 \rightarrow \alpha_3$) and (1972LA18: 3 MeV; $\alpha_0 \rightarrow \alpha_3$). See also (1976BI03, 1976NE1D) and (1970SO12; applied).

$$73. \ ^{19}\text{F}(\alpha, ^6\text{Li})^{17}\text{O} \quad Q_{\text{m}} = -12.338$$

Angular distributions are reported at $E_{\alpha} = 28$ MeV (1971KL1E: $^{17}\text{O}^*(0, 0.87) + ^6\text{Li}_{\text{g.s.}}$; $^{17}\text{O}_{\text{g.s.}} + ^6\text{Li}^*(3.56)$) and 42 MeV (1968MI05: $^{17}\text{O}^*(0, 0.87)$).

$$74. \ ^{20}\text{Ne}(\text{n}, \alpha)^{17}\text{O} \quad Q_{\text{m}} = -0.584$$

At $E_{\text{n}} = 14.1$ MeV angular distributions are reported for α_0 and α_1 by (1966MC14) and (1971BA82: also α_2). (1971KA18) report the excitation of a number of states of ^{17}O with $E_{\text{x}} < 8$ MeV. See also (1972LI30) and (1973CL1E; astrophys. questions).

¹⁷F
(Figs. 8 and 9)

GENERAL: (See also (1971AJ02).)

Shell and cluster models: (1970HA49, 1972EN03, 1972LE1L, 1973DE13, 1973KU04, 1973LA1D, 1973RE17, 1973SM1C, 1974LO04).

Special levels: (1969WI1C, 1971SE1C, 1972BE1E, 1972EN03, 1973LE06, 1974VA24).

Electromagnetic transitions: (1970AL1D, 1970HA49, 1970SI1J, 1972EN03, 1972SE1G, 1973HA53, 1973LE06, 1974HA1C, 1974LO04, 1974MC1F, 1976SH04).

Special reactions: (1971AR02, 1974LI1D, 1975WI07).

Other topics: (1970RY04, 1970SI1J, 1971AU02, 1971SE1C, 1971NG01, 1972BA25, 1972CA37, 1972CH16, 1972LE1L, 1972SH32, 1973DE13, 1973GO1H, 1973OS1A, 1973RE17, 1973RO1R, 1974BR1E, 1974RE03, 1974SL1C, 1974VA24, 1975SH20, 1975SH1H).

Ground state:

$$\mu = 4.7223 \pm 0.0012 \text{ nm (1974SHYR);}$$

$$Q = 0.10 \pm 0.02 \text{ b (1974MI21).}$$

See also (1970AL1D, 1970SI1J, 1971SH26, 1971TA1A, 1972LE1L, 1972VA36, 1972YO1B, 1973NO06, 1973RE17, 1974AN1F, 1974HA27, 1974MC1F, 1974RE03, 1974WI1N, 1975BE31, 1976CH1T).

1. $^{17}\text{F}(\beta^+)^{17}\text{O}$ $Q_m = 2.762$

The half-life of ^{17}F is 65.2 ± 0.2 sec (1969WO09), 64.50 ± 0.25 sec (1972AL42). The mean of previous values [see Table 17.18 in (1971AJ02)] was 66.0 ± 0.2 sec which was the value reported by (1960JA12). We adopt $\tau_{1/2} = 64.50 \pm 0.25$ sec, $\log ft = 3.488 \pm 0.001$ (1972AL42) but suggest that another measurement of this half-life is in order. The upper limit for the β^+ decay to $^{17}\text{O}^*(0.87)$ is $< 3.4 \times 10^{-4}$ per decay (1969GA05) [$\log ft > 5.6$]. See also (1970KO41, 1970MC23, 1971BH04, 1971LI1H, 1971WI18, 1972WI28, 1972WI1C, 1973LA03, 1973MU1D, 1973WI04, 1973WI11, 1974WI1M, 1975BA59, 1975BL1G, 1975KR14, 1975WI1E; theor.).

2. $^{12}\text{C}(^{12}\text{C}, ^7\text{Li})^{17}\text{F}$ $Q_m = -16.861$

See (1971AJ02).

Table 17.17: Energy levels of ^{17}F

E_x in ^{17}F (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{1}{2}$	$\tau_{1/2} = 64.50 \pm 0.25$ sec	β^+	1, 2, 3, 4, 6, 7, 8, 9, 10, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 28, 29, 30
0.49533 ± 0.10	$\frac{1}{2}^+; \frac{1}{2}$	$\tau_m = 412 \pm 9$ psec	γ	4, 6, 7, 9, 10, 16, 17, 20, 21, 23, 25, 26, 29
3.104 ± 3	$\frac{1}{2}^-; \frac{1}{2}$	$\Gamma = 19 \pm 1$	γ, p	7, 9, 10, 11, 16, 17, 27, 29
3.857 ± 4	$\frac{5}{2}^-; \frac{1}{2}$	$\tau_m = 6 \pm 1$ fsec	γ, p	7, 9, 10, 11, 16, 17, 29
4.696 ± 10	$\frac{3}{2}^-; \frac{1}{2}$	$\Gamma = 225$	p	7, 9, 11, 16, 27
5.103 ± 10	$\frac{3}{2}^+; \frac{1}{2}$	1530	p	11, 20
5.212 ± 11	$(\frac{9}{2}); \frac{1}{2}$			7, 9
5.521 ± 10	$\frac{3}{2}^-; \frac{1}{2}$	68	p	7, 9, 11, 27
5.672 ± 10	$\frac{7}{2}^-; \frac{1}{2}$	40	p	7, 9, 11
5.682 ± 10	$\frac{1}{2}^+; \frac{1}{2}$	< 0.6	p	7, 9, 11
5.817 ± 10	$\frac{3}{2}^+; \frac{1}{2}$	180	p	7, 11
6.036 ± 10	$\frac{1}{2}^-; \frac{1}{2}$	30	p	7, 9, 11, 27
6.556 ± 10	$\frac{1}{2}^+; \frac{1}{2}$	200	p	11
6.699 ± 10	$\frac{3}{2}^-; \frac{1}{2}$	< 3	p	7, 9, 11
6.774 ± 10	$\frac{3}{2}^+; \frac{1}{2}$	4.5	p	11
7.027 ± 10	$\frac{5}{2}^-; \frac{1}{2}$	3.8	p	9, 11
7.356 ± 10	$\frac{3}{2}^+; \frac{1}{2}$	10 ± 2	p, α	9, 11, 15
7.448 ± 7		≤ 5	p	11
7.454 ± 7		7 ± 2	p, α	11, 15
7.471 ± 7		5 ± 2	p	11
7.479 ± 10	$\frac{3}{2}^+; \frac{1}{2}$	795	p	11
7.546 ± 10	$\frac{7}{2}^-; \frac{1}{2}$	30	p	11
7.75 ± 20	$\frac{1}{2}^+; \frac{1}{2}$	179 ± 3	p, α	11, 15, 27
7.95 ± 15		10 ± 3	p	11
8.01 ± 20		50 ± 20	p, α	11, 15
8.075 ± 10	$\frac{5}{2}^+; \frac{1}{2}$	100 ± 20	p, α	9, 11, 15, 27
8.2	$\frac{3}{2}^-; \frac{1}{2}$	700 ± 250	p, α	11, 15
8.383 ± 5	$\frac{5}{2}^-; \frac{1}{2}$	11 ± 5	p, α	11, 15

Table 17.17: Energy levels of ^{17}F (continued)

E_x in ^{17}F (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
8.416 \pm 10	$\frac{7}{2}^+; \frac{1}{2}$	45 \pm 10	p, α	11, 15, 27
8.75 \pm 30	$\frac{5}{2}^+; \frac{1}{2}$	170 \pm 30	p, α	11, 15
8.76	$\frac{3}{2}^+; \frac{1}{2}$	90 \pm 20	p	11
8.97	$\frac{7}{2}^-; \frac{1}{2}$	165 \pm 30	p, α	11, 15
9.27	$\frac{3}{2}^-; \frac{1}{2}$	140 \pm 30	p, α	11, 15
9.91	$\frac{9}{2}^+; \frac{1}{2}$	90 \pm 30	p, α	11, 15
10.04 \pm 20	$\frac{7}{2}; \frac{1}{2}$	280 \pm 100	p	11
10.22 \pm 20		250 \pm 80	p, α	15
10.40 \pm 20	$(\frac{5}{2}^+); \frac{1}{2}$	160 \pm 40	p	11
10.499 \pm 15	$\frac{7}{2}^-; \frac{1}{2}$	165 \pm 25	p, α	11, 15
10.79 \pm 20		120 \pm 40	p	11
10.91 \pm 100	$\frac{1}{2}^-$	560 \pm 100	p	11
10.95 \pm 20		190 \pm 50	p, α	11, 15
11.1931 \pm 2.3	$\frac{1}{2}^-; \frac{3}{2}$	0.20 \pm 0.04	γ , p, α	9, 10, 11, 15, 27
11.43 \pm 20		240 \pm 50	p, α	11, 15
11.58 \pm 40		160 \pm 30	p	11
12.00 \pm 20		120 \pm 40	p, α	11, 15
12.25 \pm 20	$\frac{3}{2}^-$	300 \pm 30	p	11
12.355 \pm 10	$\frac{1}{2}^-$	190 \pm 20	p	11
\approx 12.50	$\frac{7}{2}^-$	\approx 660	p	11
12.550 \pm 1.4	$\frac{3}{2}^-; \frac{3}{2}$	2.83 \pm 0.12	γ , p, α	9, 10, 11, 15
13.061 \pm 4	$\frac{5}{2}^-; \frac{3}{2}$	2 \pm 1	γ , p, α	9, 10, 11, 15
13.080 \pm 4	$(\frac{1}{2}^+); \frac{3}{2}$	2 \pm 1	p, α	11, 15
13.13	$\frac{5}{2}^-$	520 \pm 50	p	11
13.781 \pm 4	$\frac{5}{2}^+; \frac{3}{2}$	12 \pm 5	p, α	11, 15
14.00 \pm 50	$\frac{7}{2}^-$	260 \pm 30	p	11
14.176 \pm 6	$\frac{3}{2}^-; \frac{3}{2}$	30 \pm 5	γ , p	10, 11
14.3040 \pm 3.3	$\frac{7}{2}^-; \frac{3}{2}$	19.3 \pm 1.6	γ , p, α	10, 11, 15
14.38 \pm 50	$\frac{5}{2}^-$	610 \pm 50	p	11
14.71 \pm 100	$\frac{1}{2}^-$	470 \pm 100	p	11

Table 17.17: Energy levels of ^{17}F (continued)

E_x in ^{17}F (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
14.809 \pm 20	$\frac{1}{2}^+$	190 \pm 25	p	11
15.6		\approx 550	p	11
(16.9)	$\frac{7}{2}^-$		γ, p	10
17.1	$\frac{5}{2}^-$	1500	p	11
(18.0)	$\frac{7}{2}^-$		γ, p	10
19.42 \pm 50	$\frac{5}{2}^+$	\approx 300	$\gamma, ^3\text{He}, \alpha$	4, 5
20.25 \pm 50	$\frac{7}{2}^-$	\approx 350	$\gamma, ^3\text{He}, \alpha$	4, 5
20.9	$\frac{9}{2}^+$	600	p	11
21.01 \pm 50	$\frac{1}{2}^+$	\approx 280	$\gamma, ^3\text{He}$	4
21.8	$(\frac{9}{2}^+)$	400	p	11
22		\approx 5000	γ, p, α	10, 15
22.7	$\frac{7}{2}^+$	600	p	11
23.8	$\frac{7}{2}^+$	600	p	11
25.4	$\frac{7}{2}^-$	1500	p, α	11, 15
27.2	$\frac{5}{2}^-$	1500	p	11
28.9	$\frac{5}{2}^+$	2000	p	11

3. $^{12}\text{C}(^{14}\text{N}, ^9\text{Be})^{17}\text{F}$ $Q_m = -10.436$

See (1975ZE1C).

4. $^{14}\text{N}(^3\text{He}, \gamma)^{17}\text{F}$ $Q_m = 15.8437$

The yield of γ_{0+1} at 90° shows resonant structures at $E(^3\text{He}) = 4.35 \pm 0.05, 5.36 \pm 0.05, 5.7$ and 6.28 ± 0.05 MeV, corresponding to $^{17}\text{F}^*(19.42, 20.25, 20.5, 21.01)$, respectively. Angular distribution measurements are consistent with $J^\pi = \frac{5}{2}^+, \frac{7}{2}^-, (\frac{3}{2}^-)$ and $\frac{1}{2}^+$ for these states; the widths of the three highest ones are $\approx 350, 210$ and 280 keV (1973MO1C).

5. (a) $^{14}\text{N}(^3\text{He}, \text{n})^{16}\text{F}$	$Q_{\text{m}} = -0.969$	$E_{\text{b}} = 15.8437$
(b) $^{14}\text{N}(^3\text{He}, \text{p})^{16}\text{O}$	$Q_{\text{m}} = 15.2430$	
(c) $^{14}\text{N}(^3\text{He}, \text{d})^{15}\text{O}$	$Q_{\text{m}} = 1.8037$	
(d) $^{14}\text{N}(^3\text{He}, \text{np})^{15}\text{O}$	$Q_{\text{m}} = -0.4210$	
(e) $^{14}\text{N}(^3\text{He}, ^3\text{He})^{14}\text{N}$		
(f) $^{14}\text{N}(^3\text{He}, \alpha)^{13}\text{N}$	$Q_{\text{m}} = 10.025$	
(g) $^{14}\text{N}(^3\text{He}, 2\alpha)^{11}\text{C}$	$Q_{\text{m}} = 2.2955$	

Excitation functions for p_0 have been measured for $E(^3\text{He}) = 2.5$ to 5.5 MeV (1963GO09), 3.5 to 18 MeV (1972BI01) and 5.5 to 10.5 MeV (1971GU22; also p_{1+2} , p_3 , p_4): some large structures are observed. The elastic scattering of ^3He (reaction (e)) has been studied for $E(^3\text{He}) = 4$ to 7 MeV: there is some evidence in the excitation functions of the resonances reported in ($^3\text{He}, \gamma$) (reaction 4) (1973MO1C). The yields of α_0 , α_1 and α_{2+3} (reaction (f)) in the range $E(^3\text{He}) = 2.5$ to 8.5 MeV show broad un-correlated fluctuations, except for a structure at $E(^3\text{He}) = 4.5$ MeV (1970KN01). However, a study of α_0 [$E(^3\text{He}) = 3.5$ to 7 MeV] shows, at $\theta = 150^\circ$, strong structures corresponding to the resonances at $E(^3\text{He}) = 4.35$ and 5.36 MeV reported in ($^3\text{He}, \gamma$): their analysis suggests $\Gamma = 300$ and 350 keV (1973MO1C). For reaction (d) see (1973AD02). See also (1971ADZZ), (1971AJ02, 1974LO1B), (1974FA1A; theor.), ^{16}O , ^{16}F , and ^{13}N , ^{14}N and ^{15}O in (1976AJ04).

6. $^{14}\text{N}(\alpha, \text{n})^{17}\text{F}$ $Q_{\text{m}} = -4.7347$

See ^{18}F in (1972AJ02). See also (1973DO1K).

7. $^{14}\text{N}(^6\text{Li}, \text{t})^{17}\text{F}$ $Q_{\text{m}} = 0.049$

At $E(^6\text{Li}) = 18$ MeV the triton groups in this reaction and the ^3He groups in the mirror reaction have been compared. $^{17}\text{F}^*(3.86, 5.22, 5.67 + 5.68)$ are strongly excited. A state at $E_{\text{x}} = 5.220 \pm 0.010$ MeV is identified as the mirror state to $^{17}\text{O}^*(5.22)$: its probable J^π is $\frac{9}{2}^-$. $^{17}\text{F}^*(0, 0.50, 3.10, 4.70, 5.52, 5.82, 6.04, 6.70)$ are also populated (1973BI01). See also (1972BA1P).

8. $^{14}\text{N}(^{10}\text{B}, ^7\text{Li})^{17}\text{F}$ $Q_{\text{m}} = -1.944$

See (1976HA1W).

9. $^{15}\text{N}(^3\text{He}, \text{n})^{17}\text{F}$

$$Q_m = 5.0101$$

Angular distributions have been measured for the neutron groups to $^{17}\text{F}^*(0, 0.50, 3.10, 3.86)$ at $E(^3\text{He}) = 3.8$ and 4.8 MeV and to $^{17}\text{F}^*(4.70, 5.52, 5.68)$ at the higher energy. The population of a state at $E_x = 5.179 \pm 0.020$ MeV and of $^{17}\text{F}^*(6.04, 6.70, 7.03, 7.36, 8.08)$ is also reported. $^{17}\text{F}^*(5.18)$ probably has $J^\pi = \frac{3}{2}^+$ or $\frac{9}{2}^+$ ([1972TH07](#), [1973ET01](#)). Neutron groups have also been reported to ^{17}F states at $E_x = 11.195 \pm 0.007$, 12.540 ± 0.010 and 13.059 MeV, with $\Gamma < 20$, < 25 and < 25 keV, respectively. Angular distributions at $E(^3\text{He}) = 10.36$ and 11.88 MeV lead to $J^\pi = \frac{1}{2}^-$ for $^{17}\text{F}^*(11.20)$ [$L = 0$], $\frac{3}{2}^-$ or $\frac{5}{2}^-$ for $^{17}\text{F}^*(12.54)$ and $(\frac{3}{2}^-, \frac{5}{2}^-)$ for $^{17}\text{F}^*(13.06)$. These three states are probably the first three $T = \frac{3}{2}$ states in ^{17}F ([1969AD02](#)). The branching ratios for transitions to $^{16}\text{O}^*(0, 6.05, 6.13)$ for $^{17}\text{F}^*(11.20)$ and for the analog $T = \frac{3}{2}$ state in ^{17}O are displayed in Table [17.11](#): the ratios of the reduced widths are quite different in the two mirror nuclei ([1970MC02](#), [1973AD02](#)).

10. $^{16}\text{O}(\text{p}, \gamma)^{17}\text{F}$

$$Q_m = 0.6007$$

$$Q_0 = 600.35 \pm 0.28 \text{ keV} \text{ ([1975RO05](#))}.$$

At low energies the direct capture to $^{17}\text{F}^*(0, 0.50)$ has been observed: see ([1971AJ02](#)) for a summary of the earlier measurements and ([1973RO34](#): $0.3 - 3.1$ MeV), ([1975CH34](#): $0.85 - 2.55$ MeV). Extrapolation of cross section data leads to $S(0) = 9 \text{ keV} \cdot \text{b}$ ([1973RO34](#)), $7.45 \text{ keV} \cdot \text{b}$ ([1971BA1A](#)). See also ([1973CL1E](#), [1975CH34](#)). An anomaly is observed at $E_p = 2.66$ MeV [$^{17}\text{F}^*(3.10)$] in the yield of the γ -ray transition to $^{17}\text{F}^*(0.50)$: $\Gamma_\gamma = 12 \pm 2 \text{ meV}$; $C^2S = 0.90$ and 1.00 , respectively, for $^{17}\text{F}^*(0, 0.50)$ ([1973RO34](#)). Another resonance is observed at $E_p = 3.47$ MeV [$^{17}\text{F}^*(3.86)$]: angular distributions of the γ -rays are characteristic of an almost pure dipole transition. The data lead to $J^\pi = \frac{5}{2}^-$, $\Gamma < 1.5 \text{ keV}$, $\Gamma_\gamma = 0.11 \pm 0.02 \text{ eV}$, $\tau_m = 6 \pm 1 \text{ fsec}$ ([1963SE14](#)). For $^{17}\text{F}^*(0.50)$, $\tau_m = 445 \pm 22 \text{ psec}$ ([1960KA10](#)).

Five resonances corresponding to $T = \frac{3}{2}$ states are observed in the γ_1 and $\gamma_0 + \gamma_1$ yields: see Table [17.18](#) for the reported parameters ([1975HA06](#)). The lowest $T = \frac{3}{2}$ states of even parity at $E_x = 13.27$ and 14.02 MeV [$J^\pi = (\frac{1}{2}^+)$ and $\frac{5}{2}^+$] (see Table [17.19](#)) are not observed here: $\Gamma_\gamma \leq 7$ and $\leq 11.8 \text{ eV}$, respectively ([1975HA06](#)). The (E1) values for the $T = \frac{3}{2}$ states are in good agreement with shell model 2p-1h calculations using realistic Kuo-Brown interaction matrix elements ([1975HA06](#)).

The $(\gamma_0 + \gamma_1)$ yield at 90° has been studied for $E_p = 15.75$ to 31.66 MeV: it shows the giant dipole resonance centered at $E_x = 22$ MeV with a width of ≈ 5 MeV and a pigmy resonance centered at 17.5 MeV. The integrated strength of the, mainly $T = \frac{1}{2}$, giant resonance is $10 \text{ MeV} \cdot \text{mb}$: the observed strength distribution is in good agreement with odd parity 2p-1h, 1p excitation calculations. The pigmy resonance is due to $f_{7/2} \rightarrow d_{5/2}$: the main $f_{7/2}$ strength lies in two states at $E_x = 16.9$ and 18.0 MeV ([1975HA07](#)). See also ([1971JO1D](#)).

Table 17.18: Resonances in $^{16}\text{O}(p, \gamma)^{17}\text{F}$ ^a

E_p (MeV \pm keV)	Resonant ^b in	Γ_γ (eV)	Γ (keV)	E_x (MeV)	$J^\pi; T$	Refs.
2.66	γ_1	$(12 \pm 2) \times 10^{-3}$		3.10	$\frac{1}{2}^-; \frac{1}{2}$	A, (1973RO34)
3.47	γ_0	0.11 ± 0.02	< 1.5	3.86	$\frac{5}{2}^-; \frac{1}{2}$	(1963SE14)
11.275 ± 6	γ_1	6.0 ± 2.5 ^c	≤ 1.6	11.204	$\frac{1}{2}^-; \frac{3}{2}$	(1975HA06)
12.707 ± 1	$\gamma_0 + \gamma_1$	11.3 ± 3.4 ^c	1.8 ± 0.5	12.550	$\frac{3}{2}^-; \frac{3}{2}$	(1975HA06) ^e
13.255 ± 6	$\gamma_0 + \gamma_1$	2.8 ± 1.8 ^c	5.0 ± 1.5	13.065	$\frac{5}{2}^-; \frac{3}{2}$	(1975HA06)
14.435 ± 10	$\gamma_0 + \gamma_1$	81 ± 54 ^{c,f}	41 ± 10	14.174	$\frac{3}{2}^-; \frac{3}{2}$	(1975HA06)
14.583 ± 6 d	$\gamma_0 + \gamma_1$	13.4 ± 7.0 ^c	28 ± 5	14.313	$\frac{7}{2}^-; \frac{3}{2}$	(1975HA06)

A: See (1971AJ02).

^a See also Table 17.19.

^b γ_0 and γ_1 correspond to transitions to $^{17}\text{F}^*(0, 0.50)$, respectively.

^c These Γ_γ are based on J^π and Γ_{p_0}/Γ determinations by (1974SK02) and R.G. Van Bree (unpublished) [quoted by (1975HA06)]. The $B(E1)$ values for these five states are 4.7 ± 2.0 , 5.4 ± 1.6 , 1.2 ± 0.8 , 27 ± 18 and 4.4 ± 2.3 [$\times 10^{-3}$] $e^2 \cdot \text{fm}^2$.

^d See the text of reaction 10 for discussion of the observed pigmy and giant resonances (1975HA07).

^e J. Lowe, private communication.

^f $\Gamma(\gamma_1)/\Gamma(\gamma_0) \leq 0.14$ (J. Lowe, private communication).

11. (a) $^{16}\text{O}(p, p)^{16}\text{O}$

$$E_b = 0.6007$$

(b) $^{16}\text{O}(p, p\alpha)^{12}\text{C}$

$$Q_m = -7.1616$$

(c) $^{16}\text{O}(p, pn)^{15}\text{O}$

$$Q_m = -15.6640$$

Yield curves for elastic protons, protons scattered to $^{16}\text{O}^*(6.05, 6.13, 6.92, 7.12, 8.87)$ and for γ -rays from $^{16}\text{O}^*(6.13, 6.92)$ have been studied at many energies up to $E_p = 46$ MeV: see Table 17.19 in (1971AJ02) for a listing of the earlier measurements and (1975CH34: p_0 , 171.5_{c.m.}, $E_p = 0.39$ to 1.99 MeV), (1971AU04: p_{1+2} , p_5 ; σ_t ; $E_p = 17.0$ to 46.1 MeV) and (1969BU1B, 1971BU05: p_0 , p_{1+2} , p_5 ; $E_p = 21.3$ to 38.5 MeV). The observed resonances are displayed in Table 17.19. Phase-shift analyses of elastic scattering polarization and cross section data have led to an increased understanding of the $T = \frac{1}{2}$ states of ^{17}F (1971PR05, 1975HI02) and the work of (1974SK02, 1976HI09) has fixed the characteristics of the first seven $T = \frac{3}{2}$ states of ^{17}F . (1974SK02) has studied the isospin multiplet mass equation by comparing analog $T = \frac{3}{2}$ states in $A = 17$. See also (1976IK01).

Table 17.19: Resonances in $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$

E_p (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	Γ_{p_0}/Γ	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$	Refs.
2.663 ± 7	19 ± 1	p_0		3.106	$\frac{1}{2}^-$	A
3.47	1.53 ± 0.2	p_0		3.86	$\frac{5}{2}^-$	A, (1974DA04)
4.354 ± 10	225	p_0		4.696	$\frac{3}{2}^-$	A
4.787 ± 10	1530	p_0		5.103	$\frac{3}{2}^+$	A
5.231 ± 10	68	p_0		5.521	$\frac{3}{2}^-$	A
5.392 ± 10	40	p_0		5.672	$\frac{7}{2}^-$	A
5.402 ± 10	< 0.6	p_0		5.682	$\frac{1}{2}^+$	A
5.546 ± 10	180	p_0		5.817	$\frac{3}{2}^+$	A
5.779 ± 10	30	p_0		6.036	$\frac{1}{2}^-$	A
6.332 ± 10	200	p_0		6.556	$\frac{1}{2}^+$	A
6.484 ± 10	< 3	p_0		6.699	$\frac{3}{2}^-$	A
6.564 ± 10	4.5	p_0		6.774	$\frac{3}{2}^+$	A
6.833 ± 10	3.8	$p_0, \gamma_{6.13}$		7.027	$\frac{5}{2}^-$	A, (1974DA04)
7.183 ± 10	10 ± 2	p_0, p_2, α_0		7.356	$\frac{3}{2}^+$	A
7.280 ± 7	≤ 5	p_0		7.448		A
7.287 ± 7	7 ± 2	p_0, p_1, p_2, α		7.454		A
7.305 ± 7	5 ± 2	p_0, p_2		7.471		A
7.313 ± 10	795	p_0		7.479	$\frac{3}{2}^+$	A
7.385 ± 10	30	$p_0, p_2, \gamma_{6.13}$		7.546	$\frac{7}{2}^-$	A
7.60 ± 20	179 ± 3	p_0, p_1, α_0		7.75	$\frac{1}{2}^+$	A
7.81 ± 15	10 ± 3	p_2		7.95		A
7.88 ± 20	50 ± 20	$p_0, \gamma_{6.13}, \gamma_{6.92}, \alpha_0$		8.01		A

Table 17.19: Resonances in $^{16}\text{O}(\text{p}, \text{p})^{16}\text{O}$ and $^{16}\text{O}(\text{p}, \alpha)^{13}\text{N}$ (continued)

E_p (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	Γ_{p_0}/Γ	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$	Refs.
7.94 ± 15	100 ± 20	p_0, p_1, α_0		8.07	$\frac{5}{2}^+$	A
8.1	700 ± 250	$(p_0), p_1, \alpha_0$		8.2	$\frac{3}{2}^-$	A
8.275 ± 5	11 ± 5	$p_0 \rightarrow p_3, \alpha_0$		8.383	$\frac{5}{2}^-$	A
8.310 ± 10	45 ± 10	$p_0 \rightarrow p_3, \gamma_{6.13}, \gamma_{6.92}, \alpha_0$		8.416	$\frac{7}{2}^+$	A
8.66 ± 30	170 ± 30	p_2, p_3, p_4, α_0		8.75	$\frac{5}{2}^+$	A
8.68	90 ± 20	p_0	0.2	8.76	$\frac{3}{2}^+$	(1971PR05)
8.90	165 ± 30	$p_0 \rightarrow p_4, \gamma_{6.13}, \gamma_{6.92}, \alpha_0$	0.3	8.97	$\frac{7}{2}^-$	A, (1971PR05)
9.22	140 ± 30	$p_0 \rightarrow p_4, \gamma_{6.13}, \gamma_{6.92}, \alpha_0$	$0.5 - 0.6$	9.27	$\frac{3}{2}^-$	A, (1971PR05)
(9.59 ± 20)	310 ± 70	p_0, p_1, p_4		(9.62)		(1964DA02)
9.90	90 ± 30	p_0, p_2, α_0	0.05	9.91	$\frac{9}{2}^+$	A, (1971PR05)
10.04 ± 20	280 ± 100	p_0, p_1		10.04	$\frac{7}{2}$	A
10.23 ± 20	250 ± 80	α_0		10.22		(1964DA02)
10.42 ± 20	160 ± 40	p_0, p_1, p_3		10.40	$(\frac{5}{2}^+)$	(1964DA02, 1975HI02)
10.525 ± 15	165 ± 25	p_0, p_2, α_0	0.28 ± 0.03	10.499	$\frac{7}{2}^-$	A, (1971PR05, 1975HI02)
(10.75 ± 50)		p_0, p_1, α_0		(10.71)	$(\frac{7}{2}^-)$	(1964DA02, 1975HI02)
10.83 ± 20	120 ± 40	$p_0, p_2, (p_3), (\alpha_0)$		10.79		A
10.96 ± 100	560 ± 100	p_0	0.25 ± 0.07	10.91	$\frac{1}{2}^-$	(1975HI02)
11.00 ± 20	190 ± 50	$(p_2), p_3, (\alpha_0)$		10.95		A
11.2636 ± 2.0^a	0.20 ± 0.04	p_0, p_2, p_4, α_0		11.1931 ± 2.3	$\frac{1}{2}^-; \frac{3}{2}$	A, (1974SK02, 1976HI09)
11.52 ± 20	240 ± 50	p_2, α_0		11.43		A
11.67 ± 40	160 ± 30	p_0, p_3		11.58		A
12.12 ± 20	120 ± 40	p_2, α_0		12.00		A

Table 17.19: Resonances in $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$ (continued)

E_p (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	Γ_{p_0}/Γ	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$	Refs.
12.39 ± 20	300 ± 30	p_0, p_2	0.26 ± 0.03	12.25	$\frac{3}{2}^-$	A, (1971PR05, 1975HI02)
12.500 ± 10	190 ± 20	p_0, p_1, p_4	0.31 ± 0.03	12.355	$\frac{1}{2}^-$	A, (1975HI02)
≈ 12.65	≈ 600	p_0	≈ 0.09	≈ 12.50	$\frac{7}{2}^-$	(1975HI02)
12.7077 ± 2.0^b	2.83 ± 0.12	$p_0, p_2, p_4, p_5, \alpha_0, \alpha_1$	0.26 ± 0.04	12.5507 ± 2.3	$\frac{3}{2}^-; \frac{3}{2}$	A, (1974SK02, 1976HI09)
(13.06 \pm 100)		p_0		(12.88)	$(\frac{7}{2}^-)$	(1975HI02)
(13.06 \pm 50)		p_0		(12.88)	$(\frac{1}{2}^+)$	(1975HI02)
13.250 ± 4	2 ± 1	$p_0, p_{1+2}, p_{3+4}, p_5, \alpha_0$	0.15 ± 0.04	13.060	$\frac{5}{2}^-; \frac{3}{2}$	A, (1974SK02)
13.271 ± 4	2 ± 1	$p_0 \rightarrow p_4, \alpha_0$	0.04 ± 0.02	13.080	$(\frac{1}{2}^+); \frac{3}{2}$	A, (1974SK02)
13.32 ± 100	520 ± 50	p_0	0.163 ± 0.016	13.13	$\frac{5}{2}^-$	A, (1975HI02)
14.017 ± 4	12 ± 5	$p_0, p_{1+2}, p_{3+4}, \alpha_0$	0.02 ± 0.01	13.781	$\frac{5}{2}^+; \frac{3}{2}$	A, (1974SK02)
(14.20 \pm 50)		p_0		(13.95)	$(\frac{1}{2}^+)$	(1975HI02)
14.25 ± 50	260 ± 30	p_0	0.08 ± 0.01	14.00	$\frac{7}{2}^-$	(1975HI02)
14.438 ± 6	27 ± 5	p_0, p_{3+4}	0.04 ± 0.02	14.177	$\frac{3}{2}^-; \frac{3}{2}$	(1974SK02)
14.5730 ± 3.0^c	19.3 ± 1.6	$p_0, p_{1+2}, p_{3+4}, p_5, \alpha_0$	0.11 ± 0.03	14.3040 ± 3.3	$\frac{7}{2}^-; \frac{3}{2}$	A, (1974SK02, 1976HI09)
14.65 ± 50	610 ± 50	p_0	0.10 ± 0.01	14.38	$\frac{5}{2}^-$	(1975HI02)
(14.94 \pm 100)		p_0			$(\frac{3}{2}^-)$	(1975HI02)
15.00 ± 100	470 ± 100	p_0	0.25 ± 0.03	14.71	$\frac{1}{2}^-$	(1975HI02)
15.110 ± 20	190 ± 25	p_0	0.150 ± 0.015	14.809	$\frac{1}{2}^+$	(1975HI02)
(15.245 \pm 100)		p_0		(14.94)	$(\frac{5}{2}^+)$	(1975HI02)
(15.30 \pm 50)		p_0		(14.98)	$(\frac{3}{2}^+)$	(1975HI02)
(15.37 \pm 100)		p_0		(15.05)	$(\frac{3}{2}^-)$	(1975HI02)
(15.545 \pm 100)		p_0		(15.22)	$(\frac{7}{2}^-)$	(1975HI02)

Table 17.19: Resonances in $^{16}\text{O}(p, p)^{16}\text{O}$ and $^{16}\text{O}(p, \alpha)^{13}\text{N}$ (continued)

E_p (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Particles out	Γ_{p_0}/Γ	$^{17}\text{F}^*$ (MeV)	$J^\pi; T$	Refs.
15.9 ^d	≈ 550	p_0, p_{1+2}		15.6		A
17.6	1500	p_0, p_{3+4}		17.1	$\frac{5}{2}^-$	A, (1971BU05)
20.4	600	p_0		19.8	$\frac{3}{2}^+$	A, (1971BU05)
21.6	600	$p_0, (\alpha)$		20.9	$\frac{9}{2}^+$	A, (1971BU05)
22.6	400	$p_0, (\alpha)$		21.8	$(\frac{9}{2}^+)$	(1971BU05)
23.5	600	p_0, p_5		22.7	$\frac{7}{2}^+$	A, (1971BU05)
24.7	600	$p_0, (\alpha)$		23.8	$\frac{7}{2}^+$	(1971BU05)
26.4	1500	$p_0, (\alpha)$		25.4	$\frac{7}{2}^-$	A, (1971BU05)
28.3	1500	p_0		27.2	$\frac{5}{2}^-$	A, (1971BU05)
30.1	2000	p_0		28.9	$\frac{5}{2}^+$	(1971BU05)

A: See references listed for this state in Table 17.20 (1971AJ02).

^a $\Gamma_{p_0} = 19 \pm 3$ eV (1976HI09).

^b $\Gamma_{p_0} = 0.94 \pm 0.06$ keV, $\Gamma_{\alpha_0} = 62 \pm 16$ eV, $\Gamma_{\alpha_1} = 53 \pm 22$ eV (1976HI09); J. Lowe, private communication.

^c $\Gamma_{p_0} = 1.65 \pm 0.12$ keV, $\Gamma_{\alpha_0} = 2.6 \pm 0.7$ keV (1976HI09).

^d See also Table 17.20 of (1971AJ02), for possible other resonances.

Other polarization measurements are reported at $E_p = 3.47$ and 6.83 MeV (1974DA04; p_0), 20 MeV (1974PL05, 1976PL1C; p_3, p_4, p_6, p_7), 30.3 MeV (1976DE12; p_0, p_2, p_3, p_5), 30.4 MeV (1972GR02; p_0, p_5), 30.4 to 39.9 MeV (1976LE16; p_5) and 49.4 MeV (1970CL10; p_0). See also (1971KL03). Total reaction cross section measurements are reported at $E_p = 18.8$ to 47.7 MeV (1975CA26), 21 to 43 MeV (1975SL02), and $231, 345, 464$ and 552 MeV (1972RE06). See also (1971BE1D, 1971CR1A, 1973GO27, 1976WU1A).

For reaction (b) see (1971EP03: 46.8 MeV). For reaction (c) see (1965VA1E). For spallation measurements see (1973BE36, 1974LA18). See also (1970SH1J, 1971EP1D, 1971ST30, 1972AM04, 1973BA81).

See also ^{16}O , (1971AJ02, 1971PA1H), (1975ME1E; astrophys. considerations) and (1972GE07, 1972PH06, 1973CL01, 1973MO1E, 1973RU07, 1973TH02, 1974SI13, 1975BA05, 1975CA05, 1975GE08, 1975SM01, 1975TA1A, 1975TH03, 1975TH12, 1976CO2A; theor.).

$$12. \text{}^{16}\text{O}(p, n)\text{}^{16}\text{F} \qquad Q_m = -16.212 \qquad E_b = 0.6007$$

See ^{16}F and (1974MO06; theor.).

$$13. \text{}^{16}\text{O}(p, d)\text{}^{15}\text{O} \qquad Q_m = -13.4393 \qquad E_b = 0.6007$$

The excitation function for d_0 at $\theta = 70^\circ$ has been measured for $E_p = 21$ to 38.5 MeV: a strong resonance is observed at $E_p \approx 24$ MeV (1969BU1B, 1971BU05): see Table 17.19. For polarization measurements see (1967CH15; 30.3 MeV; d_0, d_3). See also ^{15}O in (1976AJ04) and (1970DE38, 1971SA1D; theor.).

$$14. \text{(a) } \text{}^{16}\text{O}(p, t)\text{}^{14}\text{O} \qquad Q_m = -20.4064 \qquad E_b = 0.6007$$

$$\text{(b) } \text{}^{16}\text{O}(p, \text{}^3\text{He})\text{}^{14}\text{N} \qquad Q_m = -15.2430$$

The excitation function at $\theta = 70^\circ$ for tritons (reaction (a)) has been measured for $E_p = 32$ to 39.5 MeV: no structure is observed (1969BU1B, 1971BU05). Differential cross sections and analyzing powers have been studied at $E_p = 43.8$ MeV for the transitions to $^{14}\text{O}^*(0, 5.17, 6.29, 6.59, 7.78, 9.72)$ and $^{14}\text{N}^*(0, 2.31, 3.95, 5.11, 7.03, 9.17)$: attempts to fit the analyzing powers with zero-range DWBA were only successful for the first pair of analog states [$^{14}\text{O}_{\text{g.s.}}, ^{14}\text{N}^*(2.31)$] (1974MA12). For other polarization measurements see (1970NE17: $E_p = 49.5$ MeV; $t_0, ^3\text{He}$ to $^{14}\text{N}^*(0, 2.31)$). See also (1975MI01, 1976PL1C), (1971AJ02, 1972HA1X, 1976DA1K) and ^{14}N and ^{14}O in (1976AJ04).

$$15. \text{}^{16}\text{O}(p, \alpha)\text{}^{13}\text{N} \qquad Q_m = -5.218 \qquad E_b = 0.6007$$

Excitation functions of various α -groups and activation functions have been measured from threshold to $E_p = 44$ MeV: see Table 17.19 in (1971AJ02) for the earlier work and (1973MC12, 1973MC1H: 5.6 to 7.7 MeV; σ_t), (1973NE12: 5.7 to 10.5 MeV; σ_t for α_0). (1971GU23: 19 to 44 MeV; σ_t for $\alpha_0, \alpha_1, \alpha_{2+3}, \alpha_4$) and (1969BU1B, 1971BU05: 21.5 to 38.5 MeV). In addition, the work of (1974SK02, 1976HI09) has established the parameters of the first seven $T = \frac{3}{2}$ states of ^{17}F : see Table 17.19. No other resonances corresponding to $T = \frac{3}{2}$ states have been observed below $E_x = 16.21$ MeV (1974SK02). Resonances corresponding to $T = \frac{1}{2}$ states are also reported in Table 17.19. Some broad structures have been reported above $E_p \approx 15$ MeV: see (1971AJ02) and (1971BU05, 1971GU23): particularly strong peaks appear at $E_p \approx 22$ and 25.5 MeV (1971BU05).

This reaction is involved in explosive oxygen burning in stars: cross sections, and reaction rates determined as a function of the temperature of the star are discussed in (1973NE12, 1973MC12, 1973WO1C). See also (1973AR1E).

$$16. \quad ^{16}\text{O}(d, n)^{17}\text{F} \quad Q_m = -1.6239$$

$$E_{\text{thresh.}} \text{ for } ^2\text{H}(^{16}\text{O}, n)^{17}\text{F} = 14530 \pm 3.2 \text{ keV (1973MA1V).}$$

Slow neutron thresholds have been observed corresponding to the ground and first excited states of ^{17}F : see (1971AJ02). The E_x of $^{17}\text{F}^*(0.50)$ is 495.33 ± 0.10 keV (1966AL10; from γ -measurement); $\tau_m = 407 \pm 9$ psec (1964BE15). Neutron groups have been observed corresponding to $^{17}\text{F}^*(0, 0.50, 3.10, 3.86, 4.70)$. Angular measurements have been obtained for the n_0 and n_1 groups ($l_p = 2$ and 0 ; $J^\pi = \frac{5}{2}^+$ and $\frac{1}{2}^+$, respectively) for $E_d \leq 12$ MeV: see Table 17.21 in (1971AJ02). For polarization measurements see (1972AN1G; n_0, n_1 ; $E_d = 3 - 4$ MeV). See also (1971TH1E, 1976FR13), (1976LO1B; applied) and (1972DZ1A, 1976SH13; theor.).

$$17. \quad (a) \quad ^{16}\text{O}(^3\text{He}, d)^{17}\text{F} \quad Q_m = -4.8930$$

$$(b) \quad ^{16}\text{O}(^3\text{He}, dp)^{16}\text{O} \quad Q_m = -5.4938$$

At $E(^3\text{He}) = 18$ MeV, angular distributions of the deuterons to $^{17}\text{F}^*(0, 0.50, 3.104 \pm 0.003, 3.857 \pm 0.004)$ have been measured. The spectroscopic factors for $^{17}\text{F}^*(0, 0.50)$ are 0.94 and 0.83. Two step processes appear to be involved in the excitation of $^{17}\text{F}^*(3.10, 3.86)$ (1975FO16). See also (1972PR1D). For reaction (b) see (1967HO14). See also (1971AJ02).

$$18. \quad ^{16}\text{O}(\alpha, t)^{17}\text{F} \quad Q_m = -19.2139$$

See (1972PR1D). See also (1971AJ02).

$$19. \quad ^{16}\text{O}(^7\text{Li}, ^6\text{He})^{17}\text{F} \quad Q_m = -9.377$$

The angular distribution involving $^{17}\text{F}_{\text{g.s.}}$ has been measured at $E(^7\text{Li}) = 36 \text{ MeV}$ (1973SC26).

20. $^{16}\text{O}(^{10}\text{B}, ^9\text{Be})^{17}\text{F}$ $Q_{\text{m}} = -5.985$

Angular distributions have been measured at $E(^{10}\text{B}) = 100 \text{ MeV}$ for the transitions to $^{17}\text{F}^*(0+0.50, 5.1, 8.1)$ (1975NA15).

21. $^{16}\text{O}(^{11}\text{B}, ^{10}\text{Be})^{17}\text{F}$ $Q_{\text{m}} = -10.629$

$^{17}\text{F}^*(0+0.50)$ are populated at $E(^{11}\text{B}) = 113 \text{ MeV}$ (1967PO13). See also (1971AL1D; theor.).

22. $^{16}\text{O}(^{12}\text{C}, ^{11}\text{B})^{17}\text{F}$ $Q_{\text{m}} = -15.357$

See (1972SC21). See also (1971SC1F).

23. $^{16}\text{O}(^{14}\text{N}, ^{13}\text{C})^{17}\text{F}$ $Q_{\text{m}} = -6.950$

Angular distributions involving $^{17}\text{F}^*(0, 0.50)$ have been measured at $E(^{14}\text{N}) = 79 \text{ MeV}$ (1976MO03) and 155 MeV (1975NA15, 1976NA09).

24. $^{16}\text{O}(^{16}\text{O}, ^{15}\text{N})^{17}\text{F}$ $Q_{\text{m}} = -11.527$

See (1974RO04).

25. $^{17}\text{O}(\text{p}, \text{n})^{17}\text{F}$ $Q_{\text{m}} = -3.542$

$$E_{\text{thresh.}} = 3743 \pm 6 \text{ keV (1973BA31).}$$

Angular distributions of the n_0 and n_1 groups have been obtained for $E_{\text{p}} = 6.95$ to 13.50 MeV (n_0) and 6.95 to 12.45 MeV (n_1). There appears to be collective enhancement in the $L = 2$ transition to $^{17}\text{F}^*(0.5)$. A large spin-flip term in the effective two-body force is necessary to account for the strength of the ground state transition (1969AN06). See also ^{18}F in (1978AJ03).

26. $^{17}\text{O}(^3\text{He}, \text{t})^{17}\text{F}$ $Q_{\text{m}} = -2.778$

At $E(^3\text{He}) = 17.3$ MeV, angular distributions have been obtained for the tritons corresponding to $^{17}\text{F}^*(0, 0.50)$. The data have been analyzed using DWBA and a two-body interaction between the incident and target nucleons. An exact coupled-channel-equation calculation was also made for the ground state transition (1968HA30). See also (1971AJ02).

27. $^{17}\text{Ne}(\beta^+)^{17}\text{F}$ $Q_{\text{m}} = 14.53$

See ^{17}Ne .

28. $^{19}\text{F}(\gamma, 2\text{n})^{17}\text{F}$ $Q_{\text{m}} = -19.581$

See (1959OC07, 1976AN06) and ^{19}F in (1978AJ03).

29. $^{19}\text{F}(\text{p}, \text{t})^{17}\text{F}$ $Q_{\text{m}} = -11.099$

Angular distributions have been measured for the t_0 , t_1 , t_2 and t_3 groups at $E_{\text{p}} = 22.8$ MeV (1963HO24), 42.4 MeV (1974NE03) and 45 MeV (1972HU1B). See also (1968AN1A, 1972PR1D), (1971AJ02) and ^{20}Ne in (1978AJ03).

30. $^{20}\text{Ne}(\text{p}, \alpha)^{17}\text{F}$ $Q_{\text{m}} = -4.1293$

See (1971AJ02). See also (1973CL1E; astrophys. considerations).

¹⁷Ne
(Figs. 8 and 9)

GENERAL: (See also (1971AJ02).)

Theory and reviews: (1971HA1Y, 1973HA77, 1973RE17, 1975BE31).

Mass of ¹⁷Ne: The mass excess of ¹⁷Ne, determined from a measurement of the Q -value of ²⁰Ne(³He, ⁶He)¹⁷Ne is 16.48 ± 0.05 MeV (1970ME11, 1972CE1A). Then $^{17}\text{Ne} - ^{17}\text{F} = 14.53$ MeV and E_b for p, ³He and α are, respectively, 1.50, 6.46 and 9.05 MeV. See also (1971AJ02).

1. (a) $^{17}\text{Ne}(\beta^+)^{17}\text{F}^* \rightarrow ^{16}\text{O} + \text{p}$ $Q_m = 13.93$
 (b) $^{17}\text{Ne}(\beta^+)^{17}\text{F}$ $Q_m = 14.53$

The half-life of ¹⁷Ne is 109.0 ± 1.0 msec (1971HA05). Earlier values (see (1971AJ02)) gave a mean value of 108.0 ± 2.7 msec. The decay is primarily to the proton unstable states of ¹⁷F at 4.70, 5.52 and 6.04 MeV with $J^\pi = \frac{3}{2}^-$, $\frac{3}{2}^-$ and $\frac{1}{2}^-$, respectively: see Table 17.21. The super-allowed decay to the analog state [¹⁷F*(11.20)] has $\log ft = 3.29^{+0.04}_{-0.07}$. The character of the decay leads to $J^\pi = \frac{1}{2}^-$ for ¹⁷Ne_{g.s.} (1971HA05). See Table 17.3 for a comparison of the mirror ¹⁷N and ¹⁷Ne decays (1976AL02) and Table 17.11 for the decay of ¹⁷F*(11.20). See also (1971HO1D, 1972TO03; theor.).

Table 17.20: Energy levels of ¹⁷Ne

E_x in ¹⁷ Ne (MeV)	$J^\pi; T$	$\tau_{1/2}$ (msec)	Decay	Reactions
0	$\frac{1}{2}^-; \frac{3}{2}$	109.0 ± 1.0	β^+ ^a	1, 2
1.35 ^b				2
(1.84)				2
(2.77)				2
(3.70)				2
(5.28)				2

^a See Tables 17.3 and 17.21.

^b The evidence for the excited states of ¹⁷Ne is preliminary.

2. ²⁰Ne(³He, ⁶He)¹⁷Ne $Q_m = -26.19$

Table 17.21: β^+ decay of ^{17}Ne ^a

Decay to $^{17}\text{F}^*$ (MeV \pm keV)	J^π	Branching (%)	$\log ft$ ^b	Decay to $^{16}\text{O}^*$ (MeV)	Decay (%)
0	$\frac{5}{2}^+$	0.5 ± 0.2 ^c	6.95 ± 0.13		
0.50	$\frac{1}{2}^+$	1.1 ± 0.5 ^c	6.55 ± 0.21		
3.084 ± 30	$\frac{1}{2}^-$	0.48 ± 0.07	6.44 ± 0.06	0	100
4.609 ± 15	$\frac{3}{2}^-$	16.2 ± 0.7	4.59 ± 0.02	0	100
5.480 ± 10	$\frac{3}{2}^-$	54.0 ± 0.7	3.86 ± 0.01	0	100
6.037 ± 10	$\frac{1}{2}^-$	10.6 ± 0.2	4.42 ± 0.01	0	100
6.406 ± 30		0.35 ± 0.10	5.80 ± 0.13	0	100
7.708 ± 30	$\frac{1}{2}^+$	0.18 ± 0.05	5.67 ± 0.12	0	> 95
				6.05	< 5
8.075 ± 10	$\frac{5}{2}^+$	6.83 ± 0.11	3.96 ± 0.01	0	99.5
				6.05	0.49 ± 0.02
8.436 ± 10		6.51 ± 0.26	3.85 ± 0.02	0	94.3
				6.05	5.7 ± 0.5
8.825 ± 25		1.90 ± 0.06	4.23 ± 0.02	0	92.4
				6.05	7.6 ± 1.1
11.20 ^e	$\frac{1}{2}^-; T = \frac{3}{2}$	$0.71^{+0.10}_{-0.05}$	$3.29^{+0.04}_{-0.07}$	0	10 ± 2
				6.13	22 ± 2
				6.92	24 ± 6
				7.12	44 ± 4
d			0.54 ± 0.05		

^a (1971HA05). See also Table 17.23 in (1971AJ02).

^b $\log ft$ values calculated by (1971HA05) using an atomic mass excess of 16.517 ± 0.026 MeV [and $\tau_{1/2} = 109.0 \pm 1.0$ msec] rather than the presently adopted 16.48 ± 0.05 MeV. Since this energy difference leads to quite small changes, the original calculations are quoted here. However, Table 17.3 (which compares the analog decays) shows corrected ft values.

^c Calculated branchings, based on the mirror ^{17}N decay.

^d A proton group with $E_{c.m.} = 2.83$ MeV has been observed: the level in ^{17}F to which it corresponds is not known.

^e See also Table 17.11.

At $E(^6\text{He}) = 62.6$ MeV, ^6He groups are observed to $^{17}\text{Ne}^*(0, 1.35, 1.84, 2.77, 3.70, 5.28)$ [from Fig. 3 in (1972CE1A)]. See also (1970WO1D).

^{17}Na

(Not illustrated)

^{17}Na has not been observed: its mass excess is predicted to be 35.61 MeV by (1966KE16). It is then unbound with respect to breakup into $^{16}\text{Ne} + \text{p}$ by 3.2 MeV and with respect to breakup into $^{14}\text{O} + 3\text{p}$ by 5.8 MeV. See also (1976CA1R; theor.).

References

(Closed 01 November 1976)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1951BU1A Burrows, Powell and Rotblat, Proc. Roy. Soc. A209 (1951) 478
- 1952WA1A Watson and Buechner, Phys. Rev. 88 (1952) 1324
- 1957BO04 N.A. Bostrom, E.L. Hudspeth and I.L. Morgan, Phys. Rev. 105 (1957) 1545
- 1957BR82 C.P. Browne, Phys. Rev. 108 (1957) 1007
- 1958WE31 J.L. Weil and K.W. Jones, Phys. Rev. 112 (1958) 1975
- 1959HA13 H.E. Hall and T.W. Bonner, Nucl. Phys. 14 (1959) 295
- 1959OC07 J. O'Connell, P. Dyal and J. Goldemberg, Phys. Rev. 116 (1959) 173
- 1960JA12 J. Janecke, Z. Naturforsch. A15 (1960) 593
- 1960JA13 N. Jarmie and M.G. Silbert, Phys. Rev. 120 (1960) 914
- 1960KA10 J.V. Kane, R.E. Pixley, R.B. Schwartz and A. Schwarzschild, Phys. Rev. 120 (1960) 162
- 1961AR06 J.C. Armstrong and K.S. Quisenberry, Phys. Rev. 122 (1961) 150
- 1961BA10 F.De S. Barros, P.D. Forsyth, A.A. Jaffe and I.J. Taylor, Proc. Phys. Soc. (London) 77 (1961) 853
- 1961BR1A Brill', Vlasov, Kalinin and Sokolov, Dokl. Akad. Nauk SSSR 136 (1961) 55; Sov. Phys., Dokl. 6 (1961) 24
- 1961FO07 D.B. Fossan, R.L. Walter, W.E. Wilson and H.H. Barschall, Phys. Rev. 123 (1961) 209
- 1961PE1A Perlow, Ramler, Stehney and Yntema, Phys. Rev. 122 (1961) 899
- 1963GO09 S. Gorodetzky, G. Bassompierre, C.St. Pierre, A. Gallmann and P. Wagner, Nucl. Phys. 43 (1963) 92
- 1963HO24 H.D. Holmgren and C.B. Fulmer, Phys. Rev. 132 (1963) 2644
- 1963LE03 J.C. Legg, Phys. Rev. 129 (1963) 272
- 1963SE14 B.E. Segel, P.P. Singh, R.G. Allas and S.S. Hanna, Phys. Rev. Lett. 10 (1963) 345
- 1963SP02 R.H. Spear, J.D. Larson and J.D. Pearson, Nucl. Phys. 41 (1963) 353
- 1964AL11 T.K. Alexander, C. Broude and A.E. Litherland, Nucl. Phys. 53 (1964) 593

- 1964BE15 J.A. Becker and D.H. Wilkinson, Phys. Rev. 134 (1964) B1200
- 1964DA02 R.L. Dangle, L.D. Oppliger and G. Hardie, Phys. Rev. 133 (1964) B647
- 1964GR08 L.H. Greenberg, J.P. Roalsvig and R.N.H. Haslam, Can. J. Phys. 42 (1964) 731
- 1964SC09 R.B. Schwartz, H.D. Holmgren, L.M. Cameron and A.R. Knudson, Phys. Rev. 134 (1964) B577
- 1964SI06 M.G. Silbert and J.C. Hopkins, Phys. Rev. 134 (1964) B16
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964WA1B Wachter, Phys. Rev. 135 (1964) B1180
- 1965CO07 S.W. Cospers, B.T. Lucas and O.E. Johnson, Phys. Rev. 138 (1965) B51
- 1965CO09 S.W. Cospers and O.E. Johnson, Phys. Rev. 138 (1965) B610
- 1965DO1B Donaldson, Passell, Bartolini and Groves, Phys. Rev. 138 (1965) B1116
- 1965GA1A Garcia, Bolta, Lopez and Senent, An. Real Soc. Espan. Fiz. Quim 61 (1965) 341
- 1965HA05 V.P. Hart, E. Norbeck and R.R. Carlson, Phys. Rev. 137 (1965) B17
- 1965MA1A Mansour et al., Nucl. Phys. 65 (1965) 433
- 1965VA1E Valentin, Nucl. Phys. 62 (1965) 81
- 1965WA1D Waggoner and Jaffe, Nucl. Phys. 69 (1965) 305
- 1966AL10 D.E. Alburger, A. Gallmann, J.B. Nelson, J.T. Sample and E.K. Warburton, Phys. Rev. 148 (1966) 1050
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
- 1966MC14 W.N. McDicken and W. Jack, Nucl. Phys. 88 (1966) 457
- 1966TI03 Y.I. Titov, A.P. Klyucharev and V.D. Vypirailenko, Yad. Fiz. 4 (1966) 308; Sov. J. Nucl. Phys. 4 (1967) 221
- 1966WI01 H.H. Williams, E.K. Warburton, K.W. Jones and J.W. Olness, Phys. Rev. 144 (1966) 801
- 1967AL06 J.L. Alty, L.L. Green, R. Huby, G.D. Jones, J.R. Mines and J.F. Sharpey-Schafer, Nucl. Phys. A97 (1967) 541
- 1967CH15 N.S. Chant, P.S. Fisher and D.K. Scott, Nucl. Phys. A99 (1967) 669
- 1967CO05 R.K. Cole, R. Dittman, H.S. Sandhu, C.N. Waddell and J.K. Dickens, Nucl. Phys. A91 (1967) 665
- 1967HO14 R.A. Hoffswell, D. Jamnik, T.M. Noweir and A.I. Yavin, Phys. Rev. Lett. 19 (1967) 754

- 1967LU05 H.F. Lutz, J.J. Wesolowski, S.F. Eccles and L.F. Hansen, Nucl. Phys. A101 (1967) 241
- 1967MO23 J.F. Morgan and R.K. Hobbie, Phys. Rev. 163 (1967) 992
- 1967PO13 J.E. Poth, J.C. Overley and D.A. Bromley, Phys. Rev. 164 (1967) 1295
- 1968AN1A Anderson, Thesis, Stanford Univ. (1968)
- 1968BI1A Bister, Fontell and Vikberg, Ann. Acad. Sci. Fennicae, Ser. A6 (1968) 3
- 1968DA05 C.N. Davids, Nucl. Phys. A110 (1968) 619
- 1968DA1D Davids, Astrophys. J. 151 (1968) 775
- 1968DA1E Dandy, Wankling and Parnell, AWRE 060/68 (1968)
- 1968FO1A Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299 (1968) 1
- 1968HA30 L.F. Hansen, M.L. Stelts, J.G. Vidal, J.J. Wesolowski and V.A. Madsen, Phys. Rev. 174 (1968) 1155
- 1968KA1C Kahana, Nucl. Part. Phys., Montreal, 1967 (New York, Benjamin. 1968) 75
- 1968KE02 G.W. Kerr, J.M. Morris and J.R. Risser, Nucl. Phys. A110 (1968) 637
- 1968MI05 P.F. Mizera and J.B. Gerhart, Phys. Rev. 170 (1968) 839
- 1968PO04 A.M. Poskanzer, G.W. Butler, E.K. Hyde, J. Cerny, D.A. Landis and F.S. Goulding, Phys. Lett. B27 (1968) 414
- 1968RE07 D. Rendic, B. Antolkovic, G. Paic, M. Turk and P. Tomas, Nucl. Phys. A117 (1968) 113
- 1968SC18 H.F. Schaefer III, R.A. Klemm and F.E. Harris, Phys. Rev. 176 (1968) 49
- 1969AD02 E.G. Adelberger, A.B. McDonald and C.A. Barnes, Nucl. Phys. A124 (1969) 49
- 1969AN06 J.D. Anderson, S.D. Bloom, C. Wong, W.F. Hornyak and V.A. Madsen, Phys. Rev. 177 (1969) 1416
- 1969BA17 K. Bahr, T. Becker, R. Jahr and W.R. Kuhlmann, Nucl. Phys. A129 (1969) 388
- 1969BU1B Bunker, UCLA 10 P18-16 (1969)
- 1969DA13 J.C. Davis and F.T. Noda, Nucl. Phys. A134 (1969) 361
- 1969DE06 C. Detraz and H.H. Duhm, Phys. Lett. B29 (1969) 29
- 1969EL1A Elliott, Cargese Lecture in Phys., Ed. M. Jean, Vol. 3 (Gordon & Breach, 1969) 337
- 1969FE1A Federman, Cargese Lecture in Physics, Vol. 3, Ed. M. Jean (Gordon and Breach, 1969) 21
- 1969GA05 A. Gallmann, F. Jundt, E. Aslanides and D.E. Alburger, Phys. Rev. 179 (1969) 921
- 1969LU07 C.C. Lu, M.S. Zisman and B.G. Harvey, Phys. Rev. 186 (1969) 1086
- 1969RO1F Roturier, Ann. Phys. 4 (1969) 289

- 1969SC34 H.F. Schaefer, III, R.A. Klemm and F.E. Harris, Phys. Rev. 181 (1969) 137.
- 1969WI1C Wildermuth, Bochum Conf. STI/PUB/232 IAEA (1969) 3
- 1969WO09 V.K. Wohlleben and E. Schuster, Radiochim. Acta 12 (1969) 75
- 1969ZA1A Zalyubovski, Lobkovski and Visotzki, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 2056
- 1970AL1D Alaga, Theor. of Nucl. Struct., Trieste 1969, IAEA STI/PUB/249 (1970) 195
- 1970AL1E Albert, Wagner, Uberall and Werntz, Proc. 3rd Int. Conf. on High Energy Phys. and Nucl. Struct., New York, 1969 (Plenum, 1970) 89; Phys. Abs. 43447 (1971)
- 1970BA49 P.H. Barker, A. Huber, H. Knoth, U. Matter, A. Gobbi and P. Marmier, Nucl. Phys. A155 (1970) 401
- 1970BA55 P.H. Barker, P.M. Cockburn, H.P. Seiler and P. Marmier, Phys. Rev. Lett. 25 (1970) 1350
- 1970BE31 K. Bethge, D.J. Pullen and R. Middleton, Phys. Rev. C2 (1970) 395
- 1970BO25 W. Bohne, H. Homeyer, H. Lettau, H. Morgenstern and J. Scheer, Nucl. Phys. A156 (1970) 93
- 1970CA1C Cavallaro, Cunsolo, Potenza and Rubbino, Programme of 56th National Congress of Italian Phys. Soc., Venice, 1970 (Italian Phys. Soc. 1970) 36; Phys. Abs. 51269 (1971)
- 1970CL10 N.M. Clarke, E.J. Burge, D.A. Smith and J.C. Dore, Nucl. Phys. A157 (1970) 145
- 1970CL1C Clayton, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki (1970) 35
- 1970DE38 G. Delic, Nucl. Phys. A158 (1970) 117
- 1970DO10 J. Dobes, Nucl. Phys. A157 (1970) 661
- 1970GN1A Gnedin, Dolginov, Silantyev and Shibanov, Yad. Fiz. 12 (1970) 815
- 1970HA49 M. Harvey and F.C. Khanna, Nucl. Phys. A155 (1970) 337
- 1970HO08 J.L. Honsaker, T.H. Hsu, W.J. McDonald and G.C. Neilson, Nucl. Phys. A144 (1970) 473
- 1970IR01 J.M. Irvine and V.F.E. Pucknell, Nucl. Phys. A159 (1970) 513
- 1970JO1D Johnson, Thesis, Univ. Iowa (1970); Phys. Abs. 60802 (1971)
- 1970KI15 K. King and B.H.J. McKellar, Aust. J. Phys. 23 (1970) 453
- 1970KN01 A.R. Knudson and F.C. Young, Nucl. Phys. A149 (1970) 323
- 1970KO41 S.-I. Koyama, K. Takahashi and M. Yamada, Prog. Theor. Phys. 44 (1970) 663
- 1970KU1B Kubo, Prog. Theor. Phys. 44 (1970) 929
- 1970LU16 B. Lundberg, L.G. Stromberg and H. Conde, Phys. Scr. 2 (1970) 273
- 1970MC02 A.B. McDonald, E.G. Adelberger, H.B. Mak, D. Ashery, A.P. Shukla, C.L. Cocke and C.N. Davids, Phys. Lett. B31 (1970) 119
- 1970MC23 J.B. McGrory, Phys. Lett. B33 (1970) 327

- 1970ME11 R. Mendelson, G.J. Wozniak, A.D. Bacher, J.M. Loiseaux and J. Cerny, Phys. Rev. Lett. 25 (1970) 533
- 1970NE17 J.M. Nelson, N.S. Chant and P.S. Fisher, Nucl. Phys. A156 (1970) 406
- 1970OH1C Ohmura, Imanishi, Ichimura and Kawai, Prog. Theor. Phys. 44 (1970) 1242
- 1970OR1B Orphan, Hoot and John, Nucl. Sci. Eng. 42 (1970) 352
- 1970RO08 A.D. Robb, W.A. Schier and E. Sheldon, Nucl. Phys. A147 (1970) 423
- 1970RY04 V.F. Rybachenko and A.A. Sadovoi, Izv. Akad. Nauk SSSR Ser. Fiz. 34 (1970) 1784; Bull. Acad. Sci. USSR Phys. Ser. 34 (1971) 1586
- 1970SH1J Shima and Alsmiller, Nucl. Sci. Eng. 41 (1970) 47
- 1970SI02 R.P. Singhal, J.R. Moreira and H.S. Caplan, Phys. Rev. Lett. 24 (1970) 73
- 1970SI1J Siegel, Thesis, Rutgers State Univ. (1970); Phys. Abs. 60723 (1971)
- 1970SI1K Singhal, Thesis, Univ. Saskatchewan (1970); Phys. Abs. 71860 (1971)
- 1970SO12 G. Somogyi and B. Schlenk, Radiat. Eff. 5 (1970) 61
- 1970TI02 C.T. Tindle, Can. J. Phys. 48 (1970) 1747
- 1970VA1M Vautherin, Theory of Nucl. Struct. - Trieste 1969, IAEA STI/PUB/249 (1970) 767
- 1970WO1D Wozniak, Mendelson, Loiseaux and Cerny, Bull. Amer. Phys. Soc. 15 (1970) 1599
- 1971ADZZ J.M. Adams, A. Adams and J.M. Calvert, AERE-R-6473 (1971)
- 1971AJ02 F. Ajzenberg-Selove, Nucl. Phys. A166 (1971) 1
- 1971AL09 B.J. Allen and R.L. Macklin, Phys. Rev. C3 (1971) 1737
- 1971AL1D Alder and Trautmann, Nucl. Phys. A178 (1971) 60
- 1971AN16 I. Angeli, J. Csikai, J.L. Nagy, T. Scharbert, T. Sztaricskai and D. Novak, Acta Phys. (Hungary) 30 (1971) 115
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, Nucl. Phys. A160 (1971) 511
- 1971AR1R A. Arima and I. Hamamoto, Ann. Rev. Nucl. Sci. 21 (1971) 55
- 1971AU02 N. Auerbach and A. Lev, Phys. Lett. B34 (1971) 13
- 1971AU04 S.M. Austin, P.J. Locard, S.N. Bunker, J.M. Cameron, J.R. Richardson, J.W. Verba and W.T.H. van Oers, Phys. Rev. C3 (1971) 1514
- 1971AU08 N. Auerbach, Phys. Lett. B36 (1971) 293
- 1971BA06 W.L. Baker, C.E. Busch, J.A. Keane and T.R. Donoghue, Phys. Rev. C3 (1971) 494
- 1971BA1A Barnes, Advances in Nucl. Phys., Eds. Baranger and Vogt, Vol. 4 (Plenum Press, 1971) p. 133
- 1971BA2Y Baus-Baghdikian, U. Libre Bruxelles, Bull. No. 45 (1971)

- 1971BA68 P.H. Barker, P.M. Cockburn, A. Huber, H. Knoth, U. Matter, H.-P. Seiler and P. Marmier, *Ann. Phys. (New York)* 66 (1971) 705
- 1971BA82 R. Bachinger and M. Uhl, *Acta Phys. Aust.* 33 (1971) 317
- 1971BE1D Benenson, Locard, Escudie and Moss, *Bull. Amer. Phys. Soc.* 16 (1971) 509
- 1971BE2D Bethge, *Suppl. J. Phys.* 32 (1971) C6-87
- 1971BE2F Beckert, *Zentral. Kernf. Rossendorf Bei Dresden, Rept. No. Zfk 222* (1971)
- 1971BE59 R.L. Becker and M.R. Patterson, *Nucl. Phys. A178* (1971) 88
- 1971BH04 S.K. Bhattacharjee, *Ann. Phys. (New York)* 63 (1971) 613
- 1971BO50 I. Borbely, *Phys. Lett. B37* (1971) 243
- 1971BR33 H.J. Brede, M. Morike, B. Schurenberg, G. Staudt and F. Weng, *Z. Phys.* 245 (1971) 1
- 1971BU05 S.N. Bunker, H. Appel, J.M. Cameron, M.B. Epstein, J.R. Quinn, J.R. Richardson and J.W. Verba, *Nucl. Phys. A163* (1971) 378
- 1971BU1E Butler, Poskanzer and Landis, *Proc. Int. Conf. on Heavy Ion Phys., Dubna* (1971) 210
- 1971BU1K Burymov et al., *Izv. Akad. Nauk SSSR Ser. Fiz.* 35 (1971) 159
- 1971CO1B Cooper, Lerner and Redish, *Bull. Amer. Phys. Soc.* 16 (1971) 99
- 1971CR1A Cramer et al., *Bull. Amer. Phys. Soc.* 16 (1971) 829
- 1971CU1B Cuzzocrea, Perillo and Notarrigo, *Nuovo Cim. A4* (1971) 251
- 1971DE1F Devins, High, Bedi and Shapiro, *Bull. Amer. Phys. Soc.* 16 (1971) 511
- 1971DO13 D.J. Donahue and R.L. Hershberger, *Phys. Rev. C4* (1971) 1693
- 1971DO15 C.B. Dover and N. Van Giai, *Nucl. Phys. A177* (1971) 559
- 1971DO1A Dohnert, *Ann. Phys.* 62 (1971) 422
- 1971EP03 M.B. Epstein, J.R. Quinn, S.N. Bunker, J.W. Verba and J.R. Richardson, *Nucl. Phys. A169* (1971) 337
- 1971EP1D Epherre, *Symp. Isotopic Composition of the Primary Cosmic Radiation, Lyngby, Denmark, 1971* (Lyngby, Denmark: Danish Space Res. Inst., 1971) p. 135; *Phys. Abs.* 6843 (1973)
- 1971ER1C Ericson, *Ann. Phys.* 63 (1971) 562
- 1971FO1A D.G. Foster and D.W. Glasgow, *Phys. Rev. C3* (1971) 576, 604
- 1971GO1C Goldberg et al., *Sov. J. Nucl. Phys.* 12 (1971) 16
- 1971GR2B Grambole et al., *Zentralinst. Kernfor. Rossendorf Dresden, Rept. No. Zfk 212* (1971)
- 1971GU22 P. Guazzoni, S. Micheletti, M. Pignanelli, F. Gentilin and F. Pellegrini, *Phys. Rev. C4* (1971) 1086

- 1971GU23 P. Guazzoni, I. Iori, S. Micheletti, N. Molho, M. Pignanelli and G. Semenescu, Phys. Rev. C4 (1971) 1092
- 1971HA05 J.C. Hardy, J.E. Esterl, R.G. Sextro and J. Cerny, Phys. Rev. C3 (1971) 700
- 1971HA1R Haeberli, Polarization, Madison, 1970 (Univ. Wisconsin Press, 1971) p. 235
- 1971HA1Y Hardy, Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) p. 261
- 1971HA48 D. Hartwig, G.T. Kaschl, G. Mairle and G.J. Wagner, Z. Phys. 246 (1971) 418
- 1971HI09 D. Hilscher, J.C. Davis and P.A. Quin, Nucl. Phys. A174 (1971) 417
- 1971HO1D B.R. Holstein, Phys. Rev. C4 (1971) 764
- 1971HO26 H.H. Howard, R.H. Stokes and B.H. Erkkila, Phys. Rev. Lett. 27 (1971) 1086
- 1971HS02 S.T. Hsieh, T.Y. Lee and C.T. Chen-Tsai, Phys. Rev. C4 (1971) 105
- 1971JE02 A.S. Jensen and M. Harvey, Can. J. Phys. 49 (1971) 1837
- 1971JO1D T. Joy and D.G. Barnes, Nucl. Instrum. Meth. 95 (1971) 199
- 1971KA18 S. Kardonsky, H.L. Finston and E.T. Williams, Phys. Rev. C4 (1971) 846
- 1971KA40 N. Kassis, H.A. Mavromatis and B. Singh, Phys. Lett. B37 (1971) 15
- 1971KE08 R.M. Keyser, R.A. Blue and H.R. Weller, Phys. Lett. B34 (1971) 602
- 1971KL03 A.P. Klyucharev and R.P. Slabospitskii, Izv. Akad. Nauk SSSR Ser. Fiz. 35 (1971) 71; Bull. Acad. Sci. USSR Phys. Ser. 35 (1972) 65
- 1971KL1E Klages et al., Suppl. J. Phys. 32 (1971) C6-209
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971KO12 H.S. Kohler and Y.C. Lin, Nucl. Phys. A167 (1971) 305
- 1971KO21 D.C. Kocher, P.J. Bjorkholm and W. Haeberli, Nucl. Phys. A172 (1971) 663
- 1971LA1D Lang, Z. Phys. 242 (1971) 179
- 1971LE1B Lerner and Redish, Bull. Amer. Phys. Soc. 16 (1971) 99
- 1971LE30 T.Y. Lee, C.T. Chen-Tsai, W.W. Yeh and S.T. Hsieh, Phys. Rev. C4 (1971) 1991
- 1971LI1H H.J. Lipkin, Phys. Rev. Lett. 27 (1971) 432
- 1971MA05 A.D. MacKellar, J.F. Reading and A.K. Kerman, Phys. Rev. C3 (1971) 460
- 1971MA62 A.D. Mackellar and R.E. Schenter, Nucl. Phys. A178 (1971) 249
- 1971MU09 B.D. Murphy, R.M. Strang and R.C. Ritter, Nucl. Phys. A168 (1971) 328
- 1971MU23 A. Muller-Arnke, Z. Phys. 247 (1971) 408
- 1971NG01 Nguyen Van Giai, D. Vautherin, M. Veneroni and D.M. Brink, Phys. Lett. B35 (1971) 135

- 1971NI04 R.J. Nickles, P.L. Jolivette and G.M. Klody, *Izv. Akad. Nauk SSSR Ser. Fiz.* 35 (1971) 65; *Bull. Acad. Sci. USSR Phys. Ser.* 35 (1972) 60
- 1971NY03 K. Nyberg-Ponnert, B. Jonsson and I. Bergqvist, *Phys. Scr.* 4 (1971) 165
- 1971PA1H Palevsky, *Acta Phys. Pol.* B2 (1971) 79
- 1971PL1D Pluhar, *Nucl. Phys.* A167 (1971) 33
- 1971PR05 R.M. Prior, K.W. Corrigan, E.D. Berners and S.E. Darden, *Nucl. Phys.* A167 (1971) 143
- 1971PR09 R. Prasad and D.C. Sarkar, *Nuovo Cim.* A3 (1971) 467
- 1971RY1A Rybachenko and Sadovoi, *Sov. J. Nucl. Phys.* 12 (1971) 384
- 1971SA1D Santos, *Polarization Phenomena in Nucl. Reactions*, Madison, 1970, Eds. H.H. Barschall and W. Haeberli (Univ. Wisconsin Press, 1971) p. 758
- 1971SC1F Scott et al., *Suppl. J. Phys.* 32 (1971) C6-275
- 1971SC21 H. Schmidt-Bocking, G. Brommundt and K. Bethge, *Z. Phys.* 246 (1971) 431
- 1971SC22 R.E. Schenter and A.D. MacKellar, *Phys. Rev.* C4 (1971) 2020
- 1971SE1C D. Sen and C.S. Shastry, *Phys. Rev.* C4 (1971) 349
- 1971SE1D Sene, Delpierre, Kahane and de Billy de Crespin, *Polarization Phenomena in Nucl. Reactions*, Madison, 1970, Eds. H.H. Barschall and W. Haeberli (Univ. Wisconsin Press, 1971) p. 611
- 1971SE1E Seyer, *Polarization Phenomena in Nucl. Reactions*, Madison, 1970, Eds. H.H. Barschall and W. Haeberli (Univ. Wisconsin Press, 1971) p. 624
- 1971SH26 V.S. Shirley, *Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation*, Israel, 1970 (London, Gordon & Breach, 1971) 1255
- 1971ST30 G.B. Stapleton and R.H. Thomas, *Nucl. Phys.* A175 (1971) 124; Erratum *Nucl. Phys.* A196 (1972) 635
- 1971TA1A Talmi, *Proc. Conf. Hyperfins Interactions Detected by Nucl. Radiation*, Israel, 1970 (London, Gordon & Breach, 1971) 1133
- 1971TH1E Thornton et al., *Polarization Phenomena in Nucl. Reactions*, Madison, 1970, Eds. H.H. Barschall and W. Haeberli (Univ. Wisconsin Press, 1971) p. 871
- 1971TO08 I.S. Towner, E.K. Warburton and G.T. Garvey, *Ann. Phys. (N.Y.)* 66 (1971) 674
- 1971WE08 G.D. Westin and J.L. Adams, *Phys. Rev.* C4 (1971) 363; Erratum *Phys. Rev.* C4 (1972) 1971
- 1971WI01 WildeB.H. Wildenthal, J.B. McGrory and P.W.M. Glaudemans, *Phys. Rev. Lett.* 26 (1971) 96
- 1971WI18 D.H. Wilkinson, *Phys. Rev. Lett.* 27 (1971) 1018
- 1971WI1F Wildenthal, Halbert, McGrory and Kuo, Private Communication (1971)

- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1
- 1972AL42 D.E. Alburger and D.H. Wilkinson, Phys. Rev. C6 (1972) 2019
- 1972AM04 B.S. Amin, S. Biswas, D. Lal and B.L.K. Somayajulu, Nucl. Phys. A195 (1972) 311
- 1972AN1G Anderson, Thesis, Case Western Reserve Univ. (1972); Phys. Abs. 83014 (1972)
- 1972AU01 M. Auman, F.P. Brady, J.A. Jungerman, W.J. Knox, M.R. McGie and T.C. Montgomery, Phys. Rev. C5 (1972) 1
- 1972BA1P Bassani et al., Communications, Proc. Aix-En-Provence Conf. 2 (1972) 68
- 1972BA25 W.H. Bassichis and I. Kelson, Phys. Rev. C5 (1972) 1169
- 1972BE1E Bertsch and Mekjian, Ann. Rev. Nucl. Sci., 1972 (Ann. Rev. Inc., Palo Alto, California) 25
- 1972BE22 A.M. Bernstein, Ann. Phys. (New York) 69 (1972) 19
- 1972BI01 O.M. Bilaniuk, H.T. Fortune, J.D. Garrett, R. Middleton and W.P. Alford, Nucl. Phys. A180 (1972) 69
- 1972BR12 R.C. Brown, I. Govil, J.A.R. Griffith, G. Hudson, O. Karban and S. Roman, Nucl. Phys. A185 (1972) 49
- 1972BU23 V.E. Bunakov, K.A. Gridnev and L.V. Krasnov, Yad. Fiz. 15 (1972) 906; Sov. J. Nucl. Phys. 15 (1972) 508
- 1972CA1M Calarco et al., Bull. Amer. Phys. Soc. 17 (1972) 915
- 1972CA37 P. Camiz, E. Olivieri, M. Scalia and A. D'Andrea, Nuovo Cim. A12 (1972) 71
- 1972CE1A Cerny, At. Masses & Fund. Constants, Teddington, 1971 (Plenum Press 1972) 26
- 1972CH16 E.F. Chaffin and N.V.V.J. Swamy, Nucl. Phys. A187 (1972) 593
- 1972CL1A Clayton, Encyclopedia of the Twentieth Century (1972)
- 1972CO15 K.W. Corrigan, R.M. Prior and S.E. Darden, Nucl. Phys. A188 (1972) 164
- 1972DA21 A.V. Davydov and I.M. Narodetskii, Pisma Zh. Eksp. Teor. Fiz. 15 (1972) 741; JETP Lett. (USSR) 15 (1972) 525
- 1972DZ06 P.O. Dzhamalov, E.I. Dolinskii and A.M. Mukhamedzhanov, Yad. Fiz. 15 (1972) 258; Sov. J. Nucl. Phys. 15 (1972) 147
- 1972DZ1A Dzhamalov and Dolinskii, Sov. J. Nucl. Phys. 14 (1972) 423
- 1972ED01 G. Eder, G. Winkler and P. Hille, Z. Phys. 253 (1972) 335
- 1972EL1C Elliott, Proc. Roy. Soc. 326 (1972) 199
- 1972EN03 T. Engeland and P.J. Ellis, Nucl. Phys. A181 (1972) 368
- 1972EY01 Y. Eyal, I. Dostrovsky and Z. Fraenkel, Nucl. Phys. A180 (1972) 545
- 1972FO17 R.P. Fogel and S.T. Thornton, Nucl. Phys. A192 (1972) 391
- 1972FR1E W.A. Friedman, Phys. Rev. C6 (1972) 87

1972GA1F Garvey, Comments Nucl. Part. Phys. 5 (1972) 85
 1972GE07 H.V. Geramb and G.L. Strobel, Phys. Lett. B39 (1972) 611
 1972GL06 D. Glas, Z. Phys. 255 (1972) 175
 1972GO04 L.J.B. Goldfarb and K. Takeuchi, Nucl. Phys. A181 (1972) 609
 1972GR02 P.D. Greaves, V. Hnizdo, J. Lowe and O. Karban, Nucl. Phys. A179 (1972) 1
 1972HA1Q Harvey, 1970 4th Symp. Struct. of Low-Medium Mass Nuclei (Univ. Press of Kansas, 1972) 1
 1972HA1X Hardy, Symp. on Two-Nucleon Transfer and Pairing Excitations, Argonne, Conf-720309 (1972) 203
 1972HA2A Hale, Young and Foster, Bull. Amer. Phys. Soc. 17 (1972) 915
 1972HA57 H.H. Hackenbroich and T.H. Seligman, Phys. Lett. B41 (1972) 102
 1972HI17 J. Hiura, F. Nemoto and H. Bando, Suppl. Prog. Theor. Phys. 52 (1972) 173
 1972HU1B Huber, Thesis, Univ. Southern California (1972); Phys. Abs. 82972 (1973)
 1972KA38 N.I. Kassis, Nucl. Phys. A194 (1972) 205
 1972KE08 R.M. Keyser, R.A. Blue and H.R. Weller, Nucl. Phys. A186 (1972) 528
 1972KI1D Kinney and Perey, ORNL-4780 (1972)
 1972LA18 S. Laribi, H. Beaumevieille, N. Bendjaballah, D. Lalanne, J.F. Allard and B. Faid, Nucl. Phys. A191 (1972) 368
 1972LA1F Lane, COO-1717-3 (1972)
 1972LE01 M.-C. Lemaire, M.C. Mermaz and K.K. Seth, Phys. Rev. C5 (1972) 328
 1972LE1G Lerner, Thesis, Univ. Maryland (1972); Phys. Abs. 70442 (1972)
 1972LE1L Lee and Cusson, Ann. Phys. 72 (1972) 353
 1972LE1M Leung, Bull. Amer. Phys. Soc. 17 (1972) 58
 1972LE27 G.M. Lerner and E.F. Redish, Nucl. Phys. A193 (1972) 565
 1972LE28 G.M. Lerner and J.B. Marion, Nucl. Phys. A193 (1972) 593
 1972LI30 S.-Y. Lin, Y.-C. Hsu, M.-C. Chou, C.-Y. Huang, J.-L. Hwang and Y.-C. Hsu, Chin. J. Phys. (Taiwan) 10 (1972) 69
 1972MA57 H.A. Mavromatis and B. Singh, Phys. Lett. B41 (1972) 251
 1972NE1B Nemoto and Bando, Prog. Theor. Phys. 47 (1972) 1210
 1972NI15 H. Nissimov and J.P. Elliott, Nucl. Phys. A198 (1972) 1
 1972PE1E Perey, Love and Kinney, ORNL-4823 (1972)
 1972PH06 R.J. Philpott, Phys. Rev. C5 (1972) 1457
 1972PR1D Proctor, Bennson and Bayer, Bull. Amer. Phys. Soc. 17 (1972) 442

- 1972PU1B Puhlhofer et al., Communications, Proc. of Aix-En-Provence Conf., 2 (1972) 55
- 1972RE06 P.U. Renberg, D.F. Measday, M. Pepin, P. Schwaller, B. Favier and C. Richard-Serre, Nucl. Phys. A183 (1972) 81
- 1972SC20 B. Schwesinger and W. Stocker, Phys. Lett. B39 (1972) 475
- 1972SC21 D.K. Scott, P.N. Hudson, P.S. Fisher, C.U. Cardinal, N. Anyas-Weiss, A.D. Panagiotou and P.J. Ellis, Phys. Rev. Lett. 28 (1972) 1659
- 1972SC45 L. Schlessinger and G.L. Payne, Phys. Rev. C6 (1972) 2047
- 1972SE1G Segel, 1970 4th Symp. on the Struct. of Low-Medium Mass Nuclei (Univ. Press of Kansas, 1972) 112
- 1972SH1F Shalev, Cuttler and Ben Yaalov, Conf. Nucl. Struct. Study with Neutrons, Budapest, 1972 (Budapest, Central Res. Inst. Phys. 1972) 108; Phys. Abs. 67194 (1972)
- 1972SH32 S. Shlomo, Phys. Lett. B42 (1972) 146
- 1972TH07 S.T. Thornton, Phys. Lett. B39 (1972) 623
- 1972TH13 C. Thibault and R. Klapisch, Phys. Rev. C6 (1972) 1509
- 1972TO03 I.S. Towner and J.C. Hardy, Nucl. Phys. A179 (1972) 489
- 1972VA36 J.F.A. Van Hienen and P.W.M. Glaudemans, Phys. Lett. B42 (1972) 301
- 1972WI1C Wilkinson, Few Particle Problems, UCLA, 1972 (North-Holland, 1972) 191
- 1972WI28 D.H. Wilkinson, Proc. Roy. Soc. Edinburgh A70 (1972) 307
- 1972YO1B Yoshida and Zamick, Ann. Rev. Nucl. Sci. (Ann. Rev. Inc., Palo Alto, Calif., 1972) 121
- 1973AB1D Abuzeid, Osetinsky, Tyke and Fryshchin, Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR, Rept. P15 (1973) 7216
- 1973AD02 E.G. Adelberger, A.B. McDonald, C.L. Cocke, C.N. Davids, A.P. Shukla, H.B. Mak and D. Ashery, Phys. Rev. C7 (1973) 889
- 1973AR1E Arnett, Ann. Rev. Astron. Astrophys. 11 (1973) 73
- 1973AR1J Arima, J. Phys. Soc. Jpn. Suppl. 34 (1973) 205
- 1973AR1K Arima, Ichimura and Shimizu, J. Phys. Soc. Jpn. Suppl. 34 (1973) 526
- 1973AU1B Audouze, Truran and Zimmerman, OAP-315 (1973)
- 1973AU1C Audouze, in Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 47
- 1973AU1D Audouze and Fricke, Astrophys. J. 186 (1973) 239
- 1973BA10 J.K. Bair and F.X. Haas, Phys. Rev. C7 (1973) 1356
- 1973BA2F Baur and Gelbke, in Munich 1 (1973) 396
- 1973BA2J Baz et al., Sov. J. Nucl. Phys. 16 (1973) 179
- 1973BA31 J.K. Bair, Phys. Rev. C8 (1973) 120

- 1973BA74 H.W. Barz, V.E. Bunakov and A.M. El-Naiem, Nucl. Phys. A217 (1973) 141
- 1973BA81 L.K. Batist, E.E. Berlovich, Y.S. Blinnikov, Y.V. Elkin, Y.N. Novikov, B.M. Ovchinnikov and V.K. Tarasov, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1944; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 124
- 1973BE1N Belyaev, Dmitriev and Romyantsev, in Munich 1 (1973) 556
- 1973BE36 F.E. Bertrand and R.W. Peelle, Phys. Rev. C8 (1973) 1045
- 1973BI01 H.G. Bingham, H.T. Fortune, J.D. Garrett and R. Middleton, Phys. Rev. C7 (1973) 57
- 1973BO1G Bohne et al., Nucl. Phys. A196 (1973) 41
- 1973BO26 M. Bormann, D. Kaack, V. Schroder, W. Scobel and L. Wilde, Z. Phys. 258 (1973) 285
- 1973BO30 J.D. Bowman, A.M. Poskanzer, R.G. Korteling and G.W. Butler, Phys. Rev. Lett. 31 (1973) 614
- 1973BR1C Bromley, in Munich 2 (1973) 22
- 1973BU14 C.E. Busch, T.R. Donoghue, J.A. Keane, H. Paetz gen. Schieck and R.G. Seyler, Phys. Rev. C8 (1973) 848
- 1973BU1D Bucher, Hollandsworth and Lamoreaux, Nucl. Instrum. Meth. 111 (1973) 237
- 1973CA30 S. Cavallaro, A. Cunsolo, R. Potenza and A. Rubbino, Nuovo Cim. A14 (1973) 692
- 1973CL01 B.C. Clark, R.L. Mercer, D.G. Ravenhall and A.M. Saperstein, Phys. Rev. C7 (1973) 466
- 1973CL1E Clayton and Woosley, in Munich 2 (1973) 718
- 1973CO1P Cohen, J. Phys. Soc. Jpn. Suppl. 34 (1973) 63
- 1973DA17 S.E. Darden, S. Sen, H.R. Hiddleston, J.A. Aymar and W.A. Yoh, Nucl. Phys. A208 (1973) 77
- 1973DE13 R.J. de Meijer, Nucl. Phys. A204 (1973) 427
- 1973DE21 P.T. Debevec, H.J. Korner and J.P. Schiffer, Phys. Rev. Lett. 31 (1973) 171
- 1973DE32 R.J. de Meijer, C. Delaune, D. McShan, J.W. Nelson and H.A. van Rinsvelt, Nucl. Phys. A209 (1973) 424
- 1973DO02 E.I. Dolinsky, P.O. Dzhamalov and A.M. Mukhamedzhanov, Nucl. Phys. A202 (1973) 97
- 1973DO1D D.J. Donahue, Phys. Rev. C8 (1973) 846
- 1973DO1K Donahue, in 5th Symp. Struct. Low-Medium Mass Nuclei, Univ. Press of Kentucky (1973) 215
- 1973EI01 J.M. Eisenberg, R. Guy, J.V. Noble and H.J. Weber, Phys. Lett. B43 (1973) 20; Erratum Phys. Lett. B45 (1973) 93
- 1973EN1A Encrenaz et al., Astrophys. J. 186 (1973) L77

- 1973ET01 M.P. Etten and S.T. Thornton, *Z. Phys.* 264 (1973) 187
- 1973FI1C Fick, MPI H-1973-V27 (1973)
- 1973FO11 J.L. Fowler, C.H. Johnson and R.M. Feezel, *Phys. Rev. C* 8 (1973) 545
- 1973GE04 C.K. Gelbke, R. Bock, P. Braun-Munzinger, D. Fick, K.D. Hildenbrand, W. Weiss, S. Wenneis and G. Baur, *Phys. Lett.* B43 (1973) 284
- 1973GO1H J.M.G. Gomez and J. Sesma, *Phys. Lett.* B44 (1973) 231
- 1973GO27 Y.M. Goryachev, V.P. Kanavets, I.V. Kirpichnikov, I.I. Levintov, B.V. Morozov, N.A. Nikiforov and A.S. Starostin, *Yad. Fiz.* 17 (1973) 910; *Sov. J. Nucl. Phys.* 17 (1974) 476
- 1973HA53 M. Harvey and F.C. Khanna, *Phys. Lett.* B47 (1973) 8
- 1973HA77 J.C. Hardy, *Nucl. Data Tables* A11 (1973) 327
- 1973HI09 G.T. Hickey, F.W.K. Firk, R.J. Holt, R. Nath and H.L. Schultz, *Phys. Lett.* B47 (1973) 348
- 1973HI1A Hirata, Yamaji and Fujita, *J. Phys. Soc. Jpn. Suppl.* 34 (1973) 541
- 1973HO32 V. Horsfjord, *Phys. Lett.* B45 (1973) 455
- 1973HY1A Hyuga and Arima, *J. Phys. Soc. Jpn. Suppl.* 34 (1973) 538
- 1973IG02 M. Igarashi, M. Kawai and K. Yazaki, *Prog. Theor. Phys.* 49 (1973) 825
- 1973IS07 M. Ismail, *J. Phys. (Paris)* 34 (1973) 369
- 1973JO01 C.H. Johnson, *Phys. Rev. C* 7 (1973) 561; *Erratum Phys. Rev. C* 8 (1973) 851
- 1973JU1A Junkin and Suen, in *Munich 1* (1973) 77
- 1973KO1D Kovar, in *Symp. on Heavy Ion Transfer Reactions*, ANL Phys. B1 (1973) 59
- 1973KO1J Kochetov and Khrylin, *JETP Lett.* 18 (1973) 79
- 1973KU04 T.T.S. Kuo and E. Osnes, *Nucl. Phys.* A205 (1973) 1
- 1973KU18 B.I. Kuznetsov, I.P. Chernov and R.E. Ovsyannikova, *Yad. Fiz.* 18 (1973) 950; *Sov. J. Nucl. Phys.* 18 (1974) 490
- 1973LA03 W.A. Lanford and B.H. Wildenthal, *Phys. Rev. C* 7 (1973) 668
- 1973LA1D Lanford and Wildenthal, *Bull. Amer. Phys. Soc.* 18 (1973) 578
- 1973LE06 A. Lev and N. Auerbach, *Nucl. Phys.* A206 (1973) 563
- 1973LE07 C.M. Lerner, *Nucl. Phys.* A205 (1973) 385
- 1973LE1K Leung, ARL 73-0011 (1973)
- 1973LE28 V.M. Lebedev, A.V. Spasskii and I.B. Teplov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 37 (1973) 2663; *Bull. Acad. Sci. USSR Phys. Ser.* 37 (1974) 177
- 1973LI1G Linnck, Kraus and Blatt, *Asilomar* (1973) Paper 8B5

1973MA1V Mak, Jensen and Barnes, LAP-112 (1973)
1973MA48 F. Malaguti and P.E. Hodgson, Nucl. Phys. A215 (1973) 243
1973MC12 R.H. McCamis, G.A. Moss and J.M. Cameron, Can. J. Phys. 51 (1973) 1689
1973MC1H McCamis, Moss and Cameron, J. Roy. Astronom. Soc. Canada 67 (1973) 197
1973MI1C Millener and Strottman, in Munich 1 (1973) 66
1973MO1C Mo, Blue and Weller, Nucl. Phys. A197 (1973) 290
1973MO1E Moffa and Walker, Bull. Amer. Phys. Soc. 18 (1973) 624
1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
1973MU1D N.C. Mukhopadhyay and L.D. Miller, Phys. Lett. B47 (1973) 415
1973NE12 A.V. Nero and A.J. Howard, Nucl. Phys. A210 (1973) 60
1973NO06 J.V. Noble, Phys. Lett. B47 (1973) 99
1973OG1A Ogloblin, Sov. J. Part. Nucl. 3 (1973) 467
1973OR09 Y.V. Orlov, Yad. Fiz. 18 (1973) 1028; Sov. J. Nucl. Phys. 18 (1974) 529
1973OS1A Osnes, Phys. Norvegica 7 (1973) 103
1973PA1F Parker, Baglin and Skwiersky, Bull. Amer. Phys. Soc. 18 (1973) 550
1973PI09 M. Pignanelli, J. Gosset, F. Resmini, B. Mayer and J.L. Escudie, Phys. Rev. C8 (1973) 2120
1973PO11 A.R. Poletti and J.G. Pronko, Phys. Rev. C8 (1973) 1285
1973RA1E Rayet, Nucl. Phys. B57 (1973) 269
1973RE17 B.S. Reehal and B.H. Wildenthal, Part. Nucl. 6 (1973) 137
1973RO1P Rosina, in Nucl. Many-Body Problem, Vol. 1, 1972 (Ed. Compositori, Bologna, 1973) 490
1973RO1R Robson, Nucl. Phys. A204 (1973) 523
1973RO34 C. Rolfs, Nucl. Phys. A217 (1973) 29
1973RU07 F. Ruzzene and K. Amos, Aust. J. Phys. 26 (1973) 315
1973SC1R Scheerbaum, Shakin and Thaler, Bull. Amer. Phys. Soc. 18 (1973) 547
1973SC26 P. Schumacher, N. Ueta, H.H. Duhm, K.-I. Kubo and W.J. Klages, Nucl. Phys. A212 (1973) 573
1973SM1A Smith, Sackmann and Despain, Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 168
1973SM1C Smirnov, Shitikova and Orlova, Moscow Univ. Phys. Bull. 28:5 (1973) 32
1973SP1A Sprung, in Nucl. Many-Body Problem, Vol. 2, 1972 (Ed. Compositori, Bologna, 1973) 123

- 1973ST05 F. Stancu, Nucl. Phys. A205 (1973) 561
- 1973SU1B Sugimoto, J. Phys. Soc. Jpn. Suppl. 34 (1973) 197
- 1973SU1C Sugimoto and Tanihata, J. Phys. Soc. Jpn. Suppl. 34 (1973) 245
- 1973TA1D Talbot, in Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 47
- 1973TH02 G.L. Thomas, B.C. Sinha and F. Duggan, Nucl. Phys. A203 (1973) 305
- 1973TO16 L. Tomlinson, At. Data Nucl. Data Tables 12 (1973) 179
- 1973TR1B Truran, in Cosmochem., Ed. Cameron (Reidel Pub. Co., 1973) 23
- 1973WE06 G.D. Westin and J.L. Adams, Phys. Rev. C8 (1973) 1
- 1973WI04 D.H. Wilkinson, Phys. Rev. C7 (1973) 930
- 1973WI11 D.H. Wilkinson, Nucl. Phys. A209 (1973) 470
- 1973WI15 J. Wilczynski, Phys. Lett. B47 (1973) 124
- 1973WO1C Woosley, Astrophys. J. 186 (1973) 601
- 1973YA1B Yazaki, in Munich 1 (1973) 431
- 1973YO1A Yord and Une, J. Phys. Soc. Jpn. Suppl. 34 (1973) 535
- 1973ZA1D Zamick, J. Phys. Soc. Jpn. Suppl. 34 (1973) 470
- 1974AN1F Anagnostatos and Touliatos, Bull. Amer. Phys. Soc. 19 (1974) 991
- 1974AU03 N. Auerbach, Nucl. Phys. A229 (1974) 447
- 1974BA19 G. Baur and D. Trautmann, Z. Phys. 267 (1974) 103
- 1974BA46 G. Baur and H.H. Wolter, Phys. Lett. B51 (1974) 205
- 1974BA52 A. Babaev, E. Brackman, G. Eliseev, A. Ermilov, Y. Galaktionov, Y. Kamyshkov, V. Lubimov, N. Lugetsy, V. Nagovitzin, V. Nozik et al., Phys. Lett. B51 (1974) 501
- 1974BE1J Becker, Joffily, Beccaria and Baron, Nucl. Phys. A221 (1974) 475
- 1974BO05 J.D. Bowman, A.M. Poskanzer, R.G. Korteling and G.W. Butler, Phys. Rev. C9 (1974) 836
- 1974BO13 H.G. Bohlen and W. Norenberg, Phys. Lett. B49 (1974) 227
- 1974BO1E Bormann, Neuert and Scobel, IAEA, STI/DOC/10/156 (1974) 87
- 1974BR1E Brussard, Int. Conf. Nucl. Struct. Spectroscopic, Pt. II, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 489
- 1974BU19 W.P. Bucher, C.E. Hollandsworth, D. McNatt, R. Lamoreaux, A. Niller and J.E. Youngblood, Nucl. Sci. Eng. 54 (1974) 416
- 1974CA1J Calamand, IAEA, STI/DOC/10/156 (1974) 273
- 1974CH1Q Chechik, Stocker, Eyal and Fraenkel, in Proc. Conf. Reactions Between Complex Nuclei, Vol. 1 (North-Holland, 1974) p. 6

- 1974CO04 M.D. Cooper, W.F. Hornyak and P.G. Roos, Nucl. Phys. A218 (1974) 249
- 1974CO10 M.D. Cooper, W. Galati and W.F. Hornyak, Nucl. Phys. A221 (1974) 528
- 1974DA04 S.E. Darden and H.R. Hiddleston, Phys. Rev. C9 (1974) 800
- 1974DA23 S. Dahlgren, P. Grafstrom, B. Hoistad and A. Asberg, Nucl. Phys. A227 (1974) 245
- 1974DE1E De Jager, De Vries and De Vries, At. Data Nucl. Data Tables 14 (1974) 479
- 1974DE1M Dearborn and Schramm, Astrophys. J. 194 (1974) L67
- 1974DM01 T.A. Dmitrieva, V.M. Lebedev, V.G. Markman, A.V. Spasskii and I.B. Teplov, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2567; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 91
- 1974FA1A Faldt, Pilkhun and Schlails, Ann. Phys. 82 (1974) 326
- 1974FO17 H.T. Fortune and C.M. Vincent, Phys. Rev. C10 (1974) 1233
- 1974GE01 C.K. Gelbke, G. Baur, R. Bock, P. Braun-Munzinger, W. Grochulski, H.L. Harney and R. Stock, Nucl. Phys. A219 (1974) 253
- 1974GE03 J. George and R.J. Philpott, Phys. Rev. Lett. 32 (1974) 901
- 1974GO02 L.J.B. Goldfarb and K. Takeuchi, Nucl. Phys. A218 (1974) 396
- 1974GO1L Gobbi, in Nashville, Vol. 2 (1974) 211
- 1974HA1C Hanna, Int. Conf. Nucl. Struct. Spectrosc., Pt. II, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 249
- 1974HA25 D.L. Hanson, R.G. Stokstad, K.A. Erb, C. Olmer, M.W. Sachs and D.A. Bromley, Phys. Rev. C9 (1974) 1760
- 1974HA27 S.R. Habbal and H.A. Mavromatis, Nucl. Phys. A223 (1974) 174
- 1974HA61 G.D. Harp, Phys. Rev. C10 (1974) 2387
- 1974HI04 G.T. Hickey, F.W.K. Firk, R.J. Holt and R. Nath, Nucl. Phys. A225 (1974) 470
- 1974HI1B Hickey, Thesis, Yale Univ. (1974); Phys. Abs. 40781 (1975)
- 1974IM01 B. Imanishi, M. Ichimura and M. Kawai, Phys. Lett. B52 (1974) 267
- 1974KO1R Kopytin and Gvozdenko, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2657
- 1974LA18 H. Laumer, S.M. Austin and L.M. Panggabean, Phys. Rev. C10 (1974) 1045
- 1974LI1D Lieb et al., Bull. Amer. Phys. Soc. 19 (1974) 594
- 1974LO04 N. Lo Iudice, D.J. Rowe and S.S.M. Wong, Nucl. Phys. A219 (1974) 171
- 1974LO1B Lorenzen and Brune, IAEA, STI/DOC/10/156 (1974) 325
- 1974MA12 J.A. Macdonald, J. Cerny, J.C. Hardy, H.L. Harney, A.D. Bacher and G.R. Plattner, Phys. Rev. C9 (1974) 1694
- 1974MC1F McGrory, Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 73

- 1974MI1F Miljanic, Zabel and Phillips, Bull. Amer. Phys. Soc. 19 (1974) 547
- 1974MI21 T. Minamisono, Y. Nojiri, A. Mizobuchi and K. Sugimoto, Nucl. Phys. A236 (1974) 416
- 1974MO06 P.J. Moffa and G.E. Walker, Nucl. Phys. A222 (1974) 140
- 1974MU1F Munro and Coz, Bull. Amer. Phys. Soc. 19 (1974) 1079
- 1974NE03 J.M. Nelson and W.R. Falk, Nucl. Phys. A218 (1974) 441
- 1974NE1B Negele, Int. Conf. Nucl. Struct. Spectrosc., Pt. II, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 618
- 1974OR1A Orlov, Vestn. Mos. Univ. Fiz. Astron. (USSR) 15 (1974) 259; Phys. Abs. 81502 (1974)
- 1974PH03 R.J. Philpott and J. George, Nucl. Phys. A233 (1974) 164
- 1974PI05 M. Pignanelli, S. Micheletti, I. Iori, P. Guazzoni, F.G. Resmini and J.L. Escudie, Phys. Rev. C10 (1974) 445
- 1974PL05 A.V. Plavko, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2618; Bull. Acad. Sci. USSR Phys. Ser. 38 (1974) 137
- 1974RE03 N.E. Reid, N.E. Davison and J.P. Svenne, Phys. Rev. C9 (1974) 1882
- 1974RI09 P. Ring and P. Schuck, Z. Phys. 269 (1974) 323
- 1974RO04 H.H. Rossner, G. Hinderer, A. Weidinger and K.A. Eberhard, Nucl. Phys. A218 (1974) 606
- 1974RO27 D.W.O. Rogers, J.A. Becker, T.K. Alexander, N. Anyas-Weiss, T.A. Belote, S.P. Dolan, N.A. Jelley and W.L. Randolph, Nucl. Phys. A226 (1974) 424
- 1974RO28 D.W.O. Rogers, N. Anyas-Weiss, J.A. Becker, T.A. Belote, S.P. Dolan and W.L. Randolph, Nucl. Phys. A226 (1974) 445
- 1974SA05 E.A. Sanderson, J.P. Elliott, H.A. Mavromatis and B. Singh, Nucl. Phys. A219 (1974) 190
- 1974SA1N Sanna, Trans. Amer. Nucl. Soc. (USA) 18 (1974) 362
- 1974SC1C Schwartz, Schrack and Heaton, NBS Monograph 138 (1974)
- 1974SC1L Schier et al., Bull. Amer. Phys. Soc. 19 (1974) 553
- 1974SHYR V.S. Shirley and C.M. Lederer, LBL-3450 (1974)
- 1974SI13 B. Sinha and F. Duggan, Nucl. Phys. A226 (1974) 31
- 1974SK02 B.M. Skwiersky, C.M. Baglin and P.D. Parker, Phys. Rev. C9 (1974) 910
- 1974SL1C Sliv, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 2
- 1974SW04 Z.E. Switkowski, R.M. Wieland and A. Winther, Phys. Rev. Lett. 33 (1974) 840
- 1974TH01 C. Thibault and R. Klapisch, Phys. Rev. C9 (1974) 793

- 1974VA24 J.H. van der Merwe and C.M. Villet, Phys. Rev. C10 (1974) 1498
- 1974WE1P Weller et al., Bull. Amer. Phys. Soc. 19 (1974) 432
- 1974WI1M D.H. Wilkinson, Nucl. Phys. A225 (1974) 365
- 1974WI1N D.H. Wilkinson, Phys. Lett. B48 (1974) 407
- 1974YU01 T. Yukawa, Phys. Rev. C9 (1974) 864
- 1975AL03 J.D. Allan, J.W. Jury, R.G. Johnson, K.G. McNeill, J.G. Woodworth and Y.S. Horowitz, Can. J. Phys. 53 (1975) 786
- 1975AR1E Arnould and Norgaard, Astron. Astrophys. 42 (1976) 55
- 1975AU1D Audouze, Lequeux and Vigroux, Astron. Astrophys. 43 (1975) 71
- 1975BA05 S. Barshay, C.B. Dover and J.P. Vary, Phys. Rev. C11 (1975) 360
- 1975BA2L Babaev et al., Sov. J. Nucl. Phys. 20 (1975) 37
- 1975BA59 A. Barroso and R.J. Blin-Stoyle, Nucl. Phys. A251 (1975) 446
- 1975BE06 M. Betz, J.P. Jeukenne and A. Lejeune, Can. J. Phys. 53 (1975) 203
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, Nucl. Phys. A249 (1975) 1
- 1975BL1G R.J. Blin-Stoyle, Nucl. Phys. A254 (1975) 353
- 1975CA05 G. Cattapan, G. Pisent and V. Vanzani, Nucl. Phys. A241 (1975) 204
- 1975CA26 R.F. Carlson, A.J. Cox, J.R. Nimmo, N.E. Davison, S.A. Elbakt, J.L. Horton, A. Houdayer, A.M. Sourkes, W.T.H. van Oers and D.J. Margaziotis, Phys. Rev. C12 (1975) 1167
- 1975CH34 H.C. Chow, G.M. Griffiths and T.H. Hall, Can. J. Phys. 53 (1975) 1672
- 1975CO12 S.R. Cotanch, Phys. Lett. B57 (1975) 123
- 1975CO1J Cowan and Rose, Astrophys. J. 201 (1975) L45
- 1975CR04 G.B. Crinean, L.W.J. Wild and B.M. Spicer, Nucl. Phys. A244 (1975) 67
- 1975DR1D J.P. Draayer, J.B. French, M. Prasad and V. Potbhare and S.S.M. Wong, Phys. Lett. B57 (1975) 130
- 1975EN1A Endal, Astrophys. J. 195 (1975) 187
- 1975FO16 H.T. Fortune, L.R. Medsker, J.D. Garrett and H.G. Bingham, Phys. Rev. C12 (1975) 1723
- 1975FO19 W.A. Fowler, G.R. Caughlan and B.A. Zimmerman, Ann. Rev. Astron. Astrophys. 13 (1975) 69
- 1975GE08 J. George and R.J. Philpott, Phys. Rev. C12 (1975) 658
- 1975HA06 M.N. Harakeh, P. Paul and K.A. Snover, Phys. Rev. C11 (1975) 998
- 1975HA07 M.N. Harakeh, P. Paul and P. Gorodetzky, Phys. Rev. C11 (1975) 1008

- 1975HA33 J. Havloujian, G. Ravalli and B.M. Spicer, *Aust. J. Phys.* 28 (1975) 247
- 1975HE10 K.T. Hecht and D. Braunschweig, *Nucl. Phys.* A244 (1975) 365
- 1975HI02 H.R. Hiddleston, J.A. Aymar and S.E. Darden, *Nucl. Phys.* A242 (1975) 323
- 1975HS01 S.T. Hsieh, K.T. Knopfle, G. Mairle and G.J. Wagner, *Nucl. Phys.* A243 (1975) 380
- 1975IM04 B. Imanishi, H. Onishi and O. Tanimura, *Phys. Lett.* B57 (1975) 309
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, *Phys. Rev.* C11 (1975) 2049
- 1975KA24 D. Kalinsky, D. Melnik, U. Smilansky, N. Trautner, B.A. Watson, Y. Horowitz, S. Mordechai, G. Baur and D. Pelte, *Nucl. Phys.* A250 (1975) 364
- 1975KI15 J.C. Kim, R. Yen, I.P. Auer and H.S. Caplan, *Phys. Lett.* B57 (1975) 341
- 1975KR14 S. Krol and S. Szpikowski, *Acta Phys. Pol.* B6 (1975) 409
- 1975KU1K Kuznetsov and Chernov, *Sov. J. Nucl. Phys.* 20 (1975) 340
- 1975LA1E Lavrukhina, *Izv. Akad. Nauk SSSR Ser. Fiz.* 39 (1975) 395
- 1975ME1E Meneguizzi and Reeves, *Astron. Astrophys.* 40 (1975) 91, 99
- 1975MI01 S. Micheletti, M. Pignanelli and P. Guazzoni, *Phys. Rev.* C11 (1975) 64
- 1975MI02 M.V. Mihailovic and M. Rosina, *Nucl. Phys.* A237 (1975) 221
- 1975NA15 K.G. Nair, H. Voit, C.W. Towsley, M. Hamm, J.D. Bronson and K. Nagatani, *Phys. Rev.* C12 (1975) 1575
- 1975NO1D Norgaard and Arnould, *Astron. Astrophys.* 40 (1975) 331
- 1975PA06 A. Partensky and C. Thevenet, *Phys. Lett.* B56 (1975) 258
- 1975PH01 R.J. Philpott, *Nucl. Phys.* A243 (1975) 260
- 1975PO10 J.E. Poth, M.W. Sachs and D.A. Bromley, *Nuovo Cim.* A28 (1975) 215
- 1975RE15 W.N. Reisdorf, P.H. Lau and R. Vandenbosch, *Nucl. Phys.* A253 (1975) 490
- 1975RO05 C. Rolfs, W.S. Rodney, S. Durrance and H. Winkler, *Nucl. Phys.* A240 (1975) 221
- 1975SA1D Sanchez, Casanova and Casanova, *An. Fis. (Spain)* 71 (1975) 119
- 1975SC1H J.M. Scalo, K.H. Despain and R.K. Ulrich, *Astrophys. J.* 196 (1975) 805
- 1975SC35 W.F.W. Schneider, B. Kohlmeyer, F. Puhlhofer and R. Bock, *Nucl. Phys.* A251 (1975) 331
- 1975SC42 W.F.W. Schneider, B. Kohlmeyer and F. Puhlhofer, *Z. Phys.* A275 (1975) 249
- 1975SE03 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, *Nucl. Phys.* A241 (1975) 151
- 1975SH1H S. Shlomo and D.O. Riska, *Nucl. Phys.* A254 (1975) 281
- 1975SH20 R. Sherr and G. Bertsch, *Phys. Rev.* C12 (1975) 1671

- 1975SL02 I. Slaus, D.J. Margaziotis, R.F. Carlson, W.T.H. van Oers and J.R. Richardson, Phys. Rev. C12 (1975) 1093
- 1975SM01 R. Smith and K. Amos, Phys. Lett. B55 (1975) 162
- 1975TA1A Tang and Thompson, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 119
- 1975TH03 D.R. Thompson and Y.C. Tang, Phys. Rev. C11 (1975) 1473; Erratum Phys. Rev. C13 (1976) 2597
- 1975TH12 D.R. Thompson and Y.C. Tang, Phys. Rev. C12 (1975) 1432; Erratum Phys. Rev. C13 (1976) 2597
- 1975TR1A Trimble, Rev. Mod. Phys. 47 (1975) 877
- 1975TS01 I. Tserruya, W. Bohne, P. Braun-Munzinger, C.K. Gelbke, W. Grochulski, H.L. Harney and J. Kuzminski, Nucl. Phys. A242 (1975) 345
- 1975UD01 T. Udagawa, T. Tamura and K.S. Low, Phys. Rev. Lett. 34 (1975) 30
- 1975VE10 N.I. Venikov, Y.A. Glukhov, V.I. Manko, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta, D.N. Stepanov, V.N. Unezhev, V.I. Chuev and N.I. Chumakov, Yad. Fiz. 22 (1975) 924; Sov. J. Nucl. Phys. 22 (1976) 481
- 1975VO09 V.V. Volkov, Fiz. Elem. Chastits At. Yad. (USSR) 6 (1975) 1040; Sov. J. Part. Nucl. 6 (1976) 420
- 1975VO1B Von Oertzen and Bohlen, Phys. Rept. 19 (1975) 1
- 1975WI07 J. Wilczynski, K. Siwek-Wilczynska, J.S. Larsen, J.C. Acquadro and P.R. Christensen, Nucl. Phys. A244 (1975) 147
- 1975WI1E Wilkinson, Proc. Int. Symp. on Inter. Studies in Nucl., Germany, 1975 (Amsterdam, North-Holland, 1975) 147
- 1975WO1E Wolter, in Lect. Notes in Phys. 33 (Springer-Verlag, 1975) 140
- 1975ZE1C Zeller et al., Bull. Amer. Phys. Soc. 20 (1975) 718
- 1976AJ04 F. Ajzenberg-Selove, Nucl. Phys. A268 (1976) 1
- 1976AL02 D.E. Alburger and D.H. Wilkinson, Phys. Rev. C13 (1976) 835
- 1976AN06 D.W. Anderson, R.F. Petry and H.J. Fishbeck, Nucl. Phys. A262 (1976) 91
- 1976AU1G Auer et al., Bull. Amer. Phys. Soc. 21 (1976) 517
- 1976BI03 R.E. Bigler, S.A.A. Zaidi, J.L. Horton and H. Seitz, Phys. Rev. C13 (1976) 528
- 1976BO15 I. Borbely, Nucl. Phys. A262 (1976) 244
- 1976CA1R Calboreanu, in Cargese Conf., CERN 76-13 (1976) 166
- 1976CE1E Cerny, in Cargese Conf., CERN 76-13 (1976) 225
- 1976CH04 C.C. Chang, E.M. Diener and E. Ventura, Nucl. Phys. A258 (1976) 91
- 1976CH1T Chung and Wildenthal, MSUCL-214 (1976)

1976CO01 S.R. Cotanch and C.M. Vincent, Phys. Rev. Lett. 36 (1976) 21
 1976CO27 H.T. Coelho, F.A.B. Coutinho and A.F.R. de Toledo Piza, Phys. Rev. C14 (1976) 1280
 1976CO2A Compani-Tabrizi and Malik, Bull. Amer. Phys. Soc. 21 (1976) 29
 1976DA13 R.A. Dayras, R.G. Stokstad, Z.E. Switkowski and R.M. Wieland, Nucl. Phys. A263 (1976) 153
 1976DA1K Darden and Haerberli, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 229
 1976DA1T Dayras, Stokstad, Switkowski and Wieland, LAP-146 (1976)
 1976DE12 R. de Swiniarski, F.G. Resmini, D.L. Hendrie and A.D. Bacher, Nucl. Phys. A261 (1976) 111
 1976DE1G Despain, OAP-459 (1976)
 1976DE1P Detraz et al., in Cargese Conf., CERN 76-13 (1976) 248
 1976DR08 L. Drigo, G. Tornielli and G. Zannoni, Nuovo Cim. A31 (1976) 1
 1976EA1B Earle et al., in Contrib., Icin, Lowell (1976)
 1976EN02 H.D. Engelhardt, C.W. Lewis and H. Ullrich, Nucl. Phys. A258 (1976) 480
 1976EY01 Y. Eyal, M. Beckerman, R. Chechik, Z. Fraenkel and H. Stocker, Phys. Rev. C13 (1976) 1527
 1976FE1B Feshbach, Meson-Nucl. Phys (AIP, 1976) 521
 1976FI03 P. Fintz, G. Guillaume, F. Jundt, I. Ordonez, A. Gallmann and D.E. Alburger, Nucl. Phys. A259 (1976) 493
 1976FI1E Fisher et al., Atstrophys. J. 205 (1976) 938
 1976FR13 J.M. Freeman, Nucl. Instrum. Meth. Phys. Res. 134 (1976) 153
 1976GAYV D.I. Garber and R.R. Kinsey, BNL 325, Vol. 2 (1976)
 1976GU14 G. Guillaume, B. Rastegar, P. Fintz and A. Gallmann, Nucl. Phys. A272 (1976) 338
 1976HA1W Hamm et al., Bull. Amer. Phys. Soc. 21 (1976) 554
 1976HA27 M.N. Harakeh, A.R. Arends, M.J.A. de Voigt, A.G. Drentje, S.Y. van der Werf and A. van der Woude, Nucl. Phys. A265 (1976) 189; Erratum Nucl. Phys. A280 (1977) 500
 1976HI05 M.D. High and B. Cujec, Nucl. Phys. A259 (1976) 513
 1976HI09 F. Hinterberger, P. Von Rossen, B. Schuller, J. Bisping and R. Jahn, Nucl. Phys. A263 (1976) 460
 1976IK01 P.G. Ikossi, W.J. Thompson, T.B. Clegg, W.W. Jacobs and E.J. Ludwig, Phys. Rev. Lett. 36 (1976) 1357
 1976KO1K Kostik, Astronom. Zh. 53 (1976) 125
 1976LA13 R.D. Lawson, F.J.D. Serduke and H.T. Fortune, Phys. Rev. C14 (1976) 1245

- 1976LE16 D. Lebrun, G. Perrin, P. Martin, P. de Saintignon, J. Arvieux, M. Buenerd, H.V. Geramb and K.A. Amos, Nucl. Phys. A265 (1976) 291
- 1976LE1K Lebedev et al., in Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) 98
- 1976LI01 T.K. Li, D. Dehnhard, R.E. Brown and P.J. Ellis, Phys. Rev. C13 (1976) 55
- 1976LO1B Lorenzen, Nucl. Instrum. Meth. Phys. Res. 136 (1976) 289
- 1976MA04 F. Malaguti and P.E. Hodgson, Nucl. Phys. A257 (1976) 37
- 1976MA05 H.A. Mavromatis, Nucl. Phys. A257 (1976) 109
- 1976MC11 A.B. McDonald, T.K. Alexander and O. Hausser, Nucl. Phys. A273 (1976) 464
- 1976ME1H Mewaldt, Stone, Vidor and Vogt, Astrophys. J. 205 (1976) 931
- 1976ME1K Meyer et al., Bull. Amer. Phys. Soc. 21 (1976) 68
- 1976MO03 T. Motobayashi, I. Kohno, K. Katori, M. Yoshie, T. Ohi and H. Kamitsubo, Phys. Rev. Lett. 36 (1976) 390
- 1976NA09 K.G. Nair, K. Nagatani and H. Voit, Phys. Rev. Lett. 36 (1976) 1293
- 1976NE1D Nelson, Karban, Ludwig and Roman, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 691
- 1976NO1D Noble, Meson-Nucl. Phys.-1976 (AIP, 1976) 221
- 1976NO1F Nordberg, Nilsson, Conde and Stromberg, in Lowell Conf., CONF-760715-P2 (1976) 1318
- 1976PL1C Plavko et al., Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 828
- 1976PO01 V. Potbhare and S.P. Pandya, Nucl. Phys. A256 (1976) 253
- 1976SC1G Schulte, Kamykowski, Pacawer and Stauber, Bull. Amer. Phys. Soc. 21 (1976) 109
- 1976SH04 S. Shlomo, Phys. Lett. B60 (1976) 244
- 1976SH13 R. Shyam and S. Mukherjee, Phys. Rev. C13 (1976) 2099
- 1976VA1C Van Giai, Meson-Nucl., 1976 (AIP, 1976) 507
- 1976WA1M Wannier et al., Astrophys. J. 205 (1976) L169
- 1976WE21 G.D. Westfall and S.A.A. Zaidi, Phys. Rev. C14 (1976) 610
- 1976WO1D Woodworth, Johnson, McNeill and Jury, Bull. Amer. Phys. Soc. 21 (1976) 68
- 1976WU1A Wu et al., Bull. Amer. Phys. Soc. 21 (1976) 514
- 1977PE08 J.F. Petersen, D. Dehnhard and B.F. Bayman, Phys. Rev. C15 (1977) 1719
- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1