

# Energy Levels of Light Nuclei $A = 18$

F. Ajzenberg-Selove<sup>a</sup> and T. Lauritsen<sup>b</sup>

<sup>a</sup> *University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

<sup>b</sup> *California Institute of Technology, Pasadena, California*

**Abstract:** An evaluation of  $A = 5-24$  was published in *Nuclear Physics* 11 (1959), p. 1. This version of  $A = 18$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 1, 1958)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 18$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides:  [\$^{18}\text{O}\$](#) ,  [\$^{18}\text{F}\$](#) ,  [\$^{18}\text{Ne}\$](#)

B. Tables of Recommended Level Energies:

[Table 18.1](#): Energy levels of  $^{18}\text{O}$

[Table 18.4](#): Energy levels of  $^{18}\text{F}$

C. [References](#)

D. Figures:  [\$^{18}\text{O}\$](#) ,  [\$^{18}\text{F}\$](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

<sup>18</sup>O  
(Fig. 37)

GENERAL:

*Theory:* See (1955EL1B, 1955RE1D, 1956TH1B, 1957RA1C, 1958RE1B).

1. <sup>14</sup>C(α, γ)<sup>18</sup>O  $Q_m = 6.243$

Three resonances are reported, at  $E_\alpha = 1.13, 1.79,$  and  $2.33$  MeV (1958GO69, 1958PH37): see Table 18.2. Angular distribution measurements of the capture radiation at the resonances permit the assignments  $J = 0^+, 2^+, 4^+$  for the first three states of <sup>18</sup>O and  $J = 4^+, 1^-$  and  $1^-$  to the resonance levels, <sup>18</sup>O\*(7.12, 7.64, 8.06) (1958GO69, 1958PH37). The branching ratios for the upper two levels do not agree with values estimated from a simple collective model (1958PH37). The ground state transition from <sup>18</sup>O\*(3.55) is less than 3% of the transition  $3.55 \rightarrow 1.98$  (1958GO69).

2. <sup>14</sup>C(α, n)<sup>17</sup>O  $Q_m = -1.825$   $E_b = 6.243$   
 $E_{\text{thresh.}} = 2.340 \pm 0.003;$   
 $Q_0 = -1.820 \pm 0.002$  (1956SA06).

Observed resonances in the 0° yield curve for  $E_\alpha = 2.3$  to  $3.6$  MeV are shown in Table 18.3. The angular distribution of the neutrons at the 2.64 MeV resonance indicates  $J = 1^-$  or  $3^-$  for the 8.29 MeV state of <sup>18</sup>O (1956SA06).

3. <sup>14</sup>C(α, α)<sup>14</sup>C  $E_b = 6.243$

Observed anomalies in the scattering for  $E_\alpha = 2$  to  $3.9$  MeV are shown in Table 18.3. The resonances at  $E_\alpha = (2.8), 3.3,$  and  $3.5$  MeV appear to have  $\Gamma_n \gg \Gamma_\alpha$  (1958WE29).

4. <sup>14</sup>C(α, p)<sup>17</sup>N  $Q_m = -9.75$   $E_b = 6.238$

See (1951SU1A).

5. <sup>15</sup>N(t, p)<sup>17</sup>N  $Q_m = -0.15$   $E_b = 15.842$

Table 18.1: Energy levels of  $^{18}\text{O}$ 

$E_x$ (MeV $\pm$ keV)	$J^\pi$	$\Gamma$ (keV)	Decay	Reactions
0	$0^+$		stable	1, 7, 10, 12, 13, 14, 15, 16, 18
$1.982 \pm 4$	$2^+$		$\gamma$	1, 10, 12, 13, 16, 18
$3.55 \pm 20$	$4^+$		$\gamma$	1, 10, 13, 18
$3.929 \pm 40$				18
$5.007 \pm 40$				18
$5.170 \pm 40$				18
$5.311 \pm 40$				18
$5.456 \pm 40$				18
$6.190 \pm 40$				18
$6.328 \pm 40$				18
7.12	$4^+$		$\gamma, \alpha$	1
$7.638 \pm 15$	$1^-$	$< 3$	$\gamma, \alpha$	1
$8.057 \pm 15$	$1^-$	$< 3$	$\gamma, \alpha$	1, 3
$8.229 \pm 15$	$2^+$	$1.3 \pm 0.8$	$\alpha, n$	2, 3
$8.302 \pm 15$	$3^-$	$8 \pm 1$	$\alpha, n$	2, 3
$(8.42 \pm 20)$		$17 \pm 8$	$\alpha, n$	2
$8.84 \pm 20$		$80 \pm 16$	$\alpha, n$	2
$8.971 \pm 15$	$\geq 2^+$	$42 \pm 2$	$\alpha, n$	2, 3
$9.0 - 9.2$ (two levels)	$\left\{ \begin{array}{l} 2^+, 3^- \\ 4^+, 3^- \end{array} \right.$ or	$> 150$	$\alpha$	3

 Table 18.2: Resonances in  $^{14}\text{C}(\alpha, \gamma)^{18}\text{O}$  (1958GO69, 1958PH37)

$E_{\text{res}}$ (MeV $\pm$ keV)	$\Gamma$ (keV)	$E_x$ in $^{18}\text{O}$ (MeV)	$J^\pi$	$\Gamma_s^a$ (eV)			
				(g.s.)	(1.98)	(3.55)	(3.93)
1.13		7.12	$4^+$		0.01		
$1.794 \pm 6$	$< 3$	7.638	$1^-$	0.12	0.24	$< 0.02$	$< 0.02$
$2.334 \pm 6$	$< 3$	8.058	$1^-$	0.09	0.23	$< 0.05$	$< 0.05$

<sup>a</sup>  $\Gamma_\alpha \Gamma_\gamma / \Gamma$  for  $\gamma$ -transitions leading to the states indicated.

Table 18.3: Resonances in  $^{14}\text{C}(\alpha, n)^{17}\text{O}$  and  $^{14}\text{C}(\alpha, \alpha)^{14}\text{C}$

$E_{\text{res}}$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (total) (keV)	$\theta_{\alpha}^2$ (%)	$\theta_n^2$ (%)	$E_x$ (MeV)	$J^{\pi}$ <sup>c</sup>
$2.331 \pm 5$ <sup>a</sup>	$< 6 \pm 3$ <sup>a, d</sup>	$< 2.8$		8.056	$0^+, 1^-$ <sup>d</sup>
$2.553 \pm 4$ <sup>a, b</sup>	$1.6 \pm 1$ <sup>b</sup>	0.90	$6.2 \times 10^{-4}$	8.229	$2^+$
$2.642 \pm 3$ <sup>a, b</sup>	$10 \pm 1$ <sup>b</sup>	20	0.16	8.302	$3^-$
$(2.798 \pm 11)$ <sup>b</sup>	$22 \pm 10$ <sup>b</sup>			(8.419)	
$3.335 \pm 15$ <sup>a, b</sup>	$100 \pm 20$ <sup>b</sup>			8.837	
$3.508 \pm 4$ <sup>a, b</sup>	$54 \pm 3$ <sup>b</sup>			8.971	$\geq 2^+$
$3.6 - 3.9$ <sup>a</sup>	$> 200$ <sup>a</sup>			9.0 - 9.2 (two levels)	$\left\{ \begin{array}{l} 2^+, 3^- \\ 4^+, 3^- \end{array} \right.$ or

<sup>a</sup> (1958WE29):  $^{14}\text{C}(\alpha, \alpha)^{14}\text{C}$ ; measurements at several back angles.

<sup>b</sup> (1956SA06):  $^{14}\text{C}(\alpha, n)^{17}\text{O}$ ;  $0^\circ$ .

<sup>c</sup> From analysis of  $(\alpha, \alpha)$  and  $(\alpha, n)$  data (1958WE29).

<sup>d</sup> Compare  $^{14}\text{C}(\alpha, \gamma)^{18}\text{O}$ .

See (1956SH1A).

6.  $^{15}\text{N}(\alpha, p)^{18}\text{O}$   $Q_m = -3.971$

Not observed.

7.  $^{16}\text{O}(t, p)^{18}\text{O}$   $Q_m = 3.730$

See (1953CU1D).

8.  $^{17}\text{O}(n, p)^{17}\text{N}$   $Q_m = -7.93$   $E_b = 8.069$

See (1949CH1A).

9.  $^{17}\text{O}(n, \alpha)^{14}\text{C}$   $Q_m = 1.825$   $E_b = 8.069$

The thermal neutron cross section is  $0.4 \pm 0.1$  b (1958HU18).

10.  $^{17}\text{O}(\text{d}, \text{p})^{18}\text{O}$   $Q_{\text{m}} = 5.842$   
 $Q_0 = 5.821 \pm 0.010$  (1954AH37, 1954AH47).

Proton groups have been observed corresponding to excited states at  $1.986 \pm 0.013$  (1954AH37, 1954AH47: based on  $Q_0$ ),  $1.981 \pm 0.02$  MeV, at  $2.449 \pm 0.02$  MeV (1955HO28: based on  $Q_{\text{m}}$ ) and at  $3.56 \pm 0.02$  MeV (1956BA1L: based on  $Q_0$ ). The reported level at 2.45 MeV is not found by (1956BA1L): an upper limit of 0.2 – 0.3 is given for the relative intensity of groups leading to states in the range  $E_{\text{x}} = 2$  to 3.5 MeV compared to the ground-state group; see also  $^{18}\text{O}(\text{p}, \text{p}')^{18}\text{O}^*$  and  $^{19}\text{F}(\text{t}, \alpha)^{18}\text{O}$ . At  $E_{\text{d}} = 7.77$  MeV, angular distributions of protons to the first three states of  $^{18}\text{O}$  are fitted by  $l_{\text{n}} = 2, 0$  and  $2$ , respectively. Assignments of  $J^{\pi} = 0^+, 2^+, 4^+$  are indicated for  $^{18}\text{O}^*(0, 1.98, \text{ and } 3.56 \text{ MeV})$ . Derived coefficients for the first excited state wave function are in good agreement with the theories of (1954RE1B and 1955EL1B) (1957BI80).

11.  $^{18}\text{O}(\gamma, \text{p})^{17}\text{N}$   $Q_{\text{m}} = -15.99$

See (1955AJ61) and (1955RE1C).

12.  $^{18}\text{O}(\text{p}, \text{p}')^{18}\text{O}^*$

At  $E_{\text{p}} = 4.6$  to  $5.3$  MeV, a group is reported corresponding to  $^{18}\text{O}^*(1.981 \pm 0.004)$ . No evidence is found for a level at 2.45 MeV reported in  $^{17}\text{O}(\text{d}, \text{p})^{18}\text{O}$ : intensity less than 0.1 of 1.98 MeV group (1957YO04). A 1.9 MeV  $\gamma$ -ray is reported by (1956NA1C), presumably arising from the decay of the first excited state.

13.  $^{18}\text{O}(\text{d}, \text{d}')^{18}\text{O}^*$

At  $E_{\text{d}} = 7.8$  MeV, deuteron groups are reported to states at 0, 1.98 and  $3.551 \pm 0.019$  MeV. Upper limits of 3 to 8% are given for other possible groups in the range  $E_{\text{x}} = 2$  to 3.5 MeV (1956BA1L). At  $E_{\text{d}} = 4.98$  MeV, the  $Q = -1.98$  MeV group is  $< 3\%$  of the ground-state group (1957YO04).

14.  $^{18}\text{F}(\beta^+)^{18}\text{O}$   $Q_{\text{m}} = 1.667$

See  $^{18}\text{F}$ .

$$15. \text{}^{19}\text{F}(\gamma, \text{p})^{18}\text{O} \quad Q_{\text{m}} = -7.964$$

See  $^{19}\text{F}$ .

$$16. \text{}^{19}\text{F}(\text{n}, \text{d})^{18}\text{O} \quad Q_{\text{m}} = -5.737$$
$$Q_0 = -5.79 \pm 0.08 \text{ (1957RI44)}.$$

At  $E_{\text{n}} = 14.1$  MeV, deuteron groups are observed to the ground state, to an excited state at  $1.9 \pm 0.1$  MeV and , possibly, to others near 3 to 4 MeV. Angular distributions indicate  $l_{\text{p}} = 0$  for the ground state transition, consistent with  $J = 0^+$  or  $1^+$ , and most probably  $l_{\text{p}} = 2$  for the 1.9 MeV state ( $J = 1^+, 2^+, 3^+$ ). The ratio of reduced widths  $\theta_{\text{p}}^2(1.9)/\theta_{\text{p}}^2(0) = 0.87$ , in good agreement with shell model calculations of (1955EL1B and 1957RI44).

$$17. \text{}^{19}\text{F}(\text{d}, \text{}^3\text{He})^{18}\text{O} \quad Q_{\text{m}} = -2.470$$

Not observed.

$$18. \text{}^{19}\text{F}(\text{t}, \alpha)^{18}\text{O} \quad Q_{\text{m}} = 11.849$$

At  $E_{\text{t}} = 1.8$  MeV,  $\alpha$ -groups are reported to levels at  $1.989 \pm 0.024$ ,  $3.504 \pm 0.034$ ,  $3.929 \pm 0.040$ ,  $5.007 \pm 0.040$ ,  $5.170 \pm 0.040$ ,  $5.311 \pm 0.040$ ,  $5.456 \pm 0.040$ ,  $6.190 \pm 0.040$  and  $6.328 \pm 0.040$  MeV. No other states below  $E_{\text{x}} = 6.8$  MeV are observed (1956JA31).

$$19. \text{}^{21}\text{Ne}(\text{n}, \alpha)^{18}\text{O} \quad Q_{\text{m}} = 0.704$$

Not observed.

**<sup>18</sup>F**  
(Fig. 38)

GENERAL:

*Theory:* See (1954RE1B, 1955EL1A, 1955EL1B, 1956NE1B, 1957GR1D, 1959WA16).

1.  $^{18}\text{F}(\beta^+)^{18}\text{O}$   $Q_m = 1.677$

The positron end point is  $635 \pm 15$  (1949BL26),  $649 \pm 9$  keV (1951RU24). The spectrum is simple (see (1956DR38)). The half-life is  $112 \pm 1$  min (1949BL26),  $111 \pm 1$  min (1955JA1A),  $110 \pm 1$  min (1958BE74):  $\log ft = 3.62$ . The fact that the  $\beta$  transition to the ground state of  $^{18}\text{O}$  is allowed indicates  $J = 1^+$  for  $^{18}\text{F}$  (assumed  $T = 0$ ) (MO54). The ratio  $\epsilon_K/\beta^+ = 0.030 \pm 0.002$  (1956DR38). See also (1956DZ1A) and (1958RE1B; theor.).

2.  $^{12}\text{C}(^7\text{Li}, n)^{18}\text{F}$   $Q_m = 5.963$

See (1957NO17).

3.  $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$   $Q_m = 4.421$

Resonances have been observed at  $E_\alpha = 1.53, 1.62, 2.35,$  and  $2.88$  MeV, corresponding to  $^{18}\text{F}^* = 5.61, 5.68, 6.25,$  and  $6.651$  MeV (1955PR1A, 1958AL03, 1958BR29, 1958PH37): see Table 18.5.

The 5.61 MeV state decays to the  $^{18}\text{F}$  ground state (16%), and to states at 1.08 MeV (41%) and 3.06 MeV (45%). Gamma rays with  $E_\gamma = 2.12$  and 0.94 MeV are also observed (1958AL03): see Fig. 39. The radiative widths  $\omega\Gamma_s$  to the ground state and  $^{18}\text{F}^*(1.08)$  are 0.7 and 2.2 eV, respectively; all others amount to 3 eV (1955PR1A). The branching ratios favor  $J = 1^-$ ;  $T = 0$  for  $^{18}\text{F}^*(5.61)$  and  $T = 1$  for  $^{18}\text{F}^*(1.08$  and  $3.06)$ . Angular distributions of the cascade  $\gamma$ -rays are consistent with  $J = 0^+$  and  $2^+$  for the states at 1.08 and 3.06 MeV. Angular correlations rule out  $J = 0$  for the 0.94 MeV state. The 3.06 MeV state shows a 25% ground-state branch and a 75% cascade through the 0.94 MeV level (1958AL03: see also (1958KU81)). Delayed coincidence measurements on the  $\gamma$ -decay of the 0.94 MeV state show  $\tau \leq 5 \times 10^{-9}$  sec. This implies  $1 \leq J \leq 3$  when taken together with the results of the  $^{16}\text{O}(^3\text{He}, p\gamma)^{18}\text{F}$  reaction (1958BR29).

The 5.68 MeV state decays to the 1.08 MeV state ( $1.075 \pm 0.010$  MeV) with  $\omega\Gamma_\gamma = 2.2 \pm 0.3$  eV. The ground-state transition is  $< 0.2$  eV; others amount to about 2 eV (1955PR1A). The strength of the transition  $5.68 \rightarrow 1.08$  suggests dipole radiation,  $J(5.68) = 1$ ;  $T = 0$  (1959WA16).

Table 18.4: Energy levels of  $^{18}\text{F}$

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma$ (keV)	Decay	Reactions
0	$1^+; 0$	$\tau_{1/2} = 111 \pm 1 \text{ min}$	$\beta^+$	1, 2, 3, 8, 13, 14, 15, 16, 17, 19, 22, 23, 24, 25, 26, 27, 28, 29, 32, 34
$0.940 \pm 10$	$(2^+, 3^+; 0)$	$\tau_m \leq 5 \times 10^{-9} \text{ sec}$	$\gamma$	3, 14, 22, 27, 28, 29
$1.043 \pm 10$	$(0; 0)$			3, 14, 22, 29, 32
$1.085 \pm 10$	$0^+; 1$	$\tau_m < 3 \times 10^{-13} \text{ sec}$	$\gamma$	3, 14, 27, 28, 29
$1.127 \pm 10$	$(4^+, 5^+)$			14, (22), 29
$1.700 \pm 10$	$1^+; 0$		$\gamma$	3, 14, 27, 29, 32
$2.104 \pm 10$	$(1, 2); 0$		$\gamma$	14, 29, 32
$2.525 \pm 10$	$(1, 2, 3); 0$		$\gamma$	14, 29, 33
$3.063 \pm 10$	$(2^+; 1)$		$\gamma$	3, 14, 29
$3.133 \pm 10$				14, 29, 33
$3.354 \pm 10$				14, 27, 29, 33
$3.725 \pm 10$				14, 29
$3.790 \pm 10$				14, 29
$3.839 \pm 10$				14, 27, 29, 32
$4.115 \pm 10$	$3^+; 0$			14, 27, 29, 32
$4.226 \pm 10$				14, 29
$4.360 \pm 10$				14, 29
$4.400 \pm 10$				29, 32
$4.649 \pm 10$				29
$4.741 \pm 10$				29
$4.840 \pm 10$				29
$4.965 \pm 13$				29, 32
$5.292 \pm 10$				29
$5.500 \pm 10$				29
$5.608 \pm 15$	$(1^-); 0$	$< 1.5$	$\alpha, \gamma$	3, 29
$5.670 \pm 10$	$(1^-); 0$	$< 1.0$	$\alpha, \gamma$	3, 29
$5.785 \pm 10$				29
$6.093 \pm 10$				29

Table 18.4: Energy levels of  $^{18}\text{F}$  (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma$ (keV)	Decay	Reactions
6.137 $\pm$ 10				29
6.240 $\pm$ 10	(2 $^\pm$ ; 1)	< 1.0	$\alpha, \gamma$	3, 6, 29
6.264 $\pm$ 10	(1 $^+$ )	< 3	$\alpha$	6, 29
6.374 $\pm$ 10				29
6.470 $\pm$ 10				29
6.560 $\pm$ 10	(3 $^+$ , 4 $^+$ , 5 $^+$ )	< 0.8	$\alpha$	6, 29
6.640 $\pm$ 10		< 2.0	p, $\alpha, \gamma$	3, 6, 18, 29
6.653 $\pm$ 10	1 $^-$	93 $\pm$ 5	$\alpha$	5, 6
6.765 $\pm$ 10				29
6.790 $\pm$ 10				29
6.817 $\pm$ 10	2 $^-$	101 $\pm$ 5	$\alpha, \text{p}$	5, 6, 18
6.857 $\pm$ 10				29
7.190 $\pm$ 10	(4 $^+$ )	< 4	$\alpha, \text{p}$	6, 29
7.29 $\pm$ 20	(1 $^+$ )	45	$\alpha, \text{p}$	5, 6, 18
7.313 $\pm$ 10	(3 $^-$ )	53	$\alpha, \text{p}$	5, 6, 18, 29
7.50 $\pm$ 20	(3 $^-$ )	25	$\alpha, \text{p}$	5, 6, 18, 29
7.56 $\pm$ 20		75	$\alpha, \text{p}$	5, 6, 18
7.62 $\pm$ 20		40	$\alpha, \text{p}$	5, 6
7.71 $\pm$ 20		11	$\alpha, \text{p}$	18
7.74 $\pm$ 20		110	$\alpha, \text{p}$	5, 6, 18
7.91 $\pm$ 20	(2 $^-$ )	$\approx$ 25	$\alpha, \text{p}$	5, 6, 18
7.96 $\pm$ 20	(1 $^+$ )	70	$\alpha, \text{p}$	5, 6
8.10 $\pm$ 20		$\approx$ 40	d, p, $\alpha$	10, 18
8.22 $\pm$ 20		$\approx$ 15	$\alpha, \text{p}$	18
8.24 $\pm$ 20		$\approx$ 10	$\alpha, \text{p}$	18
8.38 $\pm$ 20		$\approx$ 50	$\alpha, \text{p}$	6, 18
9.19 – 13.8	29 levels reported in $^{16}\text{O} + \text{d}$ : see Table 18.6			

The 6.25 MeV state decays mainly to  $^{18}\text{F}^*(1.7)$ :  $E_\gamma = 4.45 \pm 0.05$  MeV ( $\omega\Gamma_\gamma = 7.9 \pm 1.0$  eV), and  $^{18}\text{F}^*(0.94)$ :  $E_\gamma = 5.30 \pm 0.10$  MeV ( $\omega\Gamma_\gamma = 0.8 \pm 0.15$  eV). Transitions to the ground state and to levels at  $\approx 3$  MeV have upper limits of 5% and 20%. The 1.7 MeV state decays through

Table 18.5: Resonances in  $^{14}\text{N} + \alpha$ 

$E_\alpha$ (MeV $\pm$ keV)	Particle Out	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	$\theta_\alpha^2$	$E_x$ (MeV)	References
$1.530 \pm 3$	$\gamma$	$< 1.5$	$(1^-); 0$		5.611	(1955PR1A, 1958AL03, 1958PH37)
$1.617 \pm 3$	$\gamma$	$< 1.0$	$(1^-; 0)$		5.679	(1955PR1A, 1958PH37, 1959WA16)
$2.351 \pm 3$	$\gamma, \alpha$	$< 1.0$	$2^\pm; 1$		6.250	(1958HE54, 1958HE56, 1958PH37, 1959WA16)
$2.370 \pm 4$	$\alpha$	$< 3$	$(1^+)$	$< 0.033$	6.264	(1958HE54, 1958HE56)
$2.767 \pm 4$	$\alpha$	$(< 0.8)$	$(3^+, 4^+, 5^+)$		6.573	(1958HE54, 1958HE56)
$2.868 \pm 4$	$\gamma, p_0$	$< 2.0$			6.651	(1958HE54, 1958HE56, 1958PH37)
$2.870 \pm 6$	$\alpha$	$93 \pm 5$	$1^-$	0.715	6.653	(1958HE54, 1958HE56, 1958KA32)
$3.080 \pm 6$	$\alpha, p_0$	$101 \pm 5$	$2^-$	0.50	6.817	(1953HE58, 1958HE54, 1958HE56, 1958KA32)
$3.576 \pm 4$	$\alpha$	$< 4$	$(4^+)$		7.202	(1958HE54, 1958HE56, 1958KA32)
3.67	$\alpha$	$45 \pm 10$	$(1^+)$		7.28	(1958HE54, 1958HE56, 1958KA32)
3.72	$\alpha, p_0$	$53 \pm 6$	$(3^-)$		7.32	(1958HE54, 1958HE56, 1958KA32)
4.00	$\alpha, p_0$	35	$(3^-)$		7.53	(1958KA32)
4.05	$\alpha, p_0$	60			7.57	(1958KA32)
4.11	$\alpha, p_0$	40			7.62	(1958KA32)
4.28	$\alpha, p_0$	120			7.75	(1958KA32)
4.50	$\alpha, p_0$	30	$(2^-)$		7.92	(1958KA32)
4.55	$\alpha, p_0$	70	$(1^+)$		7.96	(1958KA32)
5.2	$\alpha$				8.5	(1939BR1A, 1939DE1A)

$^{18}\text{F}^*(1.05)$ . The  $(6.25 \rightarrow 1.7)$  and  $(1.05 \rightarrow 0)$   $\gamma$ - $\gamma$  correlation is isotropic within 10%, consistent with  $J = 0^+$  for the 1.05 MeV state. The 1.7 MeV state is most likely to be  $1^\pm$  or  $(2^+)$ , and is assumed to be  $T = 0$  from the  $^{18}\text{O}$  level structure (compare  $^{20}\text{Ne}(d, \alpha)^{18}\text{F}$ ). The 6.25 MeV state is then  $2^\pm$  or  $3^-$  (1958PH37). If the 1.7 MeV state is  $1^+$ ,  $J(6.25)$  is  $2^+$  or  $2^-$ ; the strength of the transition  $(6.25 \rightarrow 1.7)$  indicates  $T = 1$  for  $^{18}\text{F}^*(6.25)$  (1959WA16).

The 6.65 MeV state decays with  $E_\gamma = 5.80 \pm 0.10$  (7%),  $\omega\Gamma_\gamma = 0.25$  eV and  $4.90 \pm 0.05$  MeV (70%),  $\omega\Gamma_\gamma = 2.1$  eV to the 0.94 and 1.70 MeV states, respectively. Transitions with relative intensity  $\approx 23\%$  are also reported to states with  $E_x \approx 3$  MeV (1958PH37). See also (1958BA59, 1958BR1D, 1958HE1F) and  $^{16}\text{O}(^3\text{He}, p)^{18}\text{F}$ .

$$4. \ ^{14}\text{N}(\alpha, n)^{17}\text{F} \qquad Q_m = -4.747 \qquad E_b = 4.421$$

See (1938FU01).

$$5. \ ^{14}\text{N}(\alpha, p)^{17}\text{O} \qquad Q_m = -1.197 \qquad E_b = 4.421$$

Observed resonances are displayed in Table 18.5 (1953HE58, 1958HE54, 1958KA32). Absolute cross sections are given by (1958HE54) for  $E_\alpha = 2.7$  to 3.6 MeV. The yield of protons to  $^{17}\text{O}^*(0.7)$  is at least a factor of 10 smaller than the ground-state yield. See also (1954MA1F, 1955GR1F) and (1952AJ38).

$$6. \ ^{14}\text{N}(\alpha, \alpha)^{14}\text{N} \qquad E_b = 4.421$$

Observed anomalies in the elastic scattering are exhibited in Table 18.5 (1939BR1A, 1939DE1A, 1953HE58, 1958HE54, 1958HE56, 1958KA32). The indicated assignments to the narrow resonances at  $E_\alpha = 2.35, 2.37, 2.77, 2.88,$  and  $3.58$  MeV are obtained from qualitative analysis of the excitation functions. The two broad resonances at  $E_\alpha = 2.870$  and  $3.080$  MeV have been analyzed in detail: both are formed with p-waves,  $J = 1^-$  and  $2^-$ , respectively. For the former,  $\Gamma_\alpha/\Gamma = 0.85$ . See also (1958KA32). A broad anomaly at  $E_\alpha = 3.7$  MeV requires two levels, and possible admixture of higher  $l$ -values (1958HE56, 1958KA32).

$$7. \ ^{15}\text{N}(\alpha, n)^{18}\text{F} \qquad Q_m = -6.420$$

Not reported.

$$8. \ ^{16}\text{O}(d, \gamma)^{18}\text{F} \qquad Q_m = 7.538$$

At  $E_d = 1.7$  MeV, the capture cross section is  $\approx 10 \mu\text{b}$  (1955BU1C). See also (1954SI07).

$$9. \text{}^{16}\text{O}(\text{d}, \text{n})\text{}^{17}\text{F} \qquad Q_m = -1.631 \qquad E_b = 7.538$$

Observed resonances in the forward yield of neutrons are displayed in Table 18.6 (1955MA85). See also (1952AJ38) and (1955BU1C).

$$10. \text{}^{16}\text{O}(\text{d}, \text{p})\text{}^{17}\text{O} \qquad Q_m = 1.919 \qquad E_b = 7.538$$

The yield shows strong variations with energy, suggesting compound nucleus formation (1948HE1C, 1955BE1J, 1955ST1A, 1956RO1A, 1956VA17): see Table 18.6. On the other hand, angular distribution measurements show strong stripping peaks for both  $p_0$  and  $p_1$ , for  $E_d = 0.5$  to 4.1 MeV (1955AL1E, 1955BE1J, 1955JU1C, 1955ST1A, 1956GR1F, 1956JU1G, 1956KO26, 1957BA14, 1957JU1A). See also  $^{17}\text{O}$ .

$$11. \text{}^{16}\text{O}(\text{d}, \text{d})\text{}^{16}\text{O} \qquad E_b = 7.538$$

Observed maxima in the  $\theta = 135^\circ$  and  $97^\circ$  cross section are listed in Table 18.6 (1956BE1B). Structure in the yield of elastically scattered deuterons is also reported by (1957BA14). See also (1955KH31, 1955KH35).

$$12. \text{}^{16}\text{O}(\text{d}, \alpha)\text{}^{14}\text{N} \qquad Q_m = 3.116 \qquad E_b = 7.538$$

Two resonances are reported for ground-state  $\alpha$ -particles at  $E_d = 3.85$  MeV,  $\Gamma = 35$  keV, and 4.0 MeV,  $\Gamma = 100$  keV (1957BA14): see Table 18.6. Yield curves if the  $\alpha_0$ ,  $\alpha_1$  and  $\alpha_2$  groups are reported for  $E_d = 5.5$  to 7.5 MeV by (1956BR36). The  $\alpha_1$ -group, leading to the  $T = 1$ , 2.31 MeV state of  $^{14}\text{N}$ , is greatly inhibited: the average intensity is only a few per cent of the average yield of the  $\alpha_0$  group. Broad resonances are observed in the yields of all three  $\alpha$  groups. The resonances in the  $\alpha_1$ -yield occur at  $\approx 6.2$ , 6.6, and 7.0 MeV; they are presumably due to predominantly  $T = 1$  states at  $E_x \approx 13.0$ , 13.4, and 13.8 MeV (1956BR36). At  $E_d = 7.2$  MeV, the intensity ratio  $\alpha_1/\alpha_0$  is 16% ( $\theta = 35^\circ$ ), at 6.8 and 7.1 MeV,  $\alpha_1/\alpha_0 < 6\%$ , and at 8.9 MeV,  $< 2\%$  (1958DA16). The observations are consistent with expected isobaric spin impurities in the states involved (1956BR36). Angular distributions for  $\alpha_0$ ,  $\alpha_2$  may give evidence of compound nucleus effects (1958DA16): see also  $^{14}\text{N}$ .

$$13. \text{}^{16}\text{O}(\text{t}, \text{n})\text{}^{18}\text{F} \qquad Q_m = 1.280$$

Table 18.6: Maxima in the yield of ( $^{16}\text{O} + \text{d}$ ) reactions

$E_d$ (MeV)	Particle Out	$E_x$ in $^{18}\text{F}$ (MeV)	References
(0.68)	$p_0$	(8.14)	(1956VA17)
1.86	$p_0, p_1$	9.19	(1948HE1C, 1955BE1J, 1956RO1A)
2.06	$p_1$	9.37	(1955BE1J, 1956RO1A)
2.14	$p_1$	9.44	(1956RO1A)
2.22	$n$	9.51	(1955MA85)
2.34	$n, p_1$	9.62	(1955MA85, 1956RO1A)
2.42 <sup>a</sup>	$n$	9.69	(1955MA85)
2.55	$p_1$	9.81	(1955ST1A, 1956RO1A)
2.75	$p_1$	9.98	(1956RO1A)
2.92	$n, p_0, p_1$	10.13	(1948HE1C, 1955MA85, 1955ST1A, 1956RO1A)
3.10	$n, p_1$	10.29	(1955MA85, 1956RO1A)
3.22	$n$	10.40	(1955MA85)
3.36	$n, p_0, p_1$	10.53	(1955MA85, 1955ST1A, 1956RO1A)
(3.43)	$p_1$	(10.59)	(1956RO1A)
3.65	$n, p_0, p_1$	10.78	(1955MA85, 1955ST1A, 1956RO1A)
(3.75)	$p_1$	(10.87)	(1956RO1A)
3.84 <sup>b</sup>	$p_1, \alpha_0$	10.95	(1956RO1A, 1957BA14)
3.94	$n, p_1$	11.04	(1955MA85, 1956RO1A)
4.00 <sup>c</sup>	$p_1, \alpha_0$	11.09	(1956RO1A, 1957BA14)
4.10	$n, p_1$	11.18	(1955MA85, 1956RO1A)
4.35	$p_1$	11.41	(1956RO1A)
4.79	$d_0$	11.80	(1956BE1B)
5.08	$d_0$	12.05	(1956BE1B)
5.617	$d_0$	12.53	(1956BE1B)
5.78	$d_0$	12.68	(1956BE1B)
6.10	$d_0$	12.96	(1956BE1B)
6.2 <sup>d</sup>	$\alpha_1$	13.0	(1956BR36)
6.40	$d_0$	13.23	(1956BE1B)
6.57 <sup>d</sup>	$d_0, \alpha_1$	13.38	(1956BE1B, 1956BR36)
6.74	$d_0$	13.53	(1956BE1B)
7.0 <sup>d</sup>	$\alpha_1$	13.8	(1956BR36)

<sup>a</sup>  $J^\pi = 2^+$  or  $3^+$ .

<sup>b</sup>  $\Gamma \approx 100$  keV (1957BA14).

<sup>c</sup>  $\Gamma \approx 35$  keV (1957BA14).

<sup>d</sup> ( $T = 1$ ).

See (1951PO1A, 1955BA1Q, 1956SH1A).

$$14. \text{}^{16}\text{O}(\text{}^3\text{He}, \text{p})\text{}^{18}\text{F} \quad Q_{\text{m}} = 2.045$$
$$Q_0 = 2.052 \pm 0.015 \text{ (1959HI67)}.$$

Seventeen proton groups corresponding to excited states of  $^{18}\text{F}$  from 0 to  $E_{\text{x}} = 4.36$  MeV are reported by (1959HI67:  $E(^3\text{He}) = 5.7$  to  $5.9$  MeV): see Table 18.7. See also (1958KU81). Angular distributions for the first five states, analyzed by direct interaction theory, indicate  $J = 1^+$ ,  $2^+$  or  $3^+$ ,  $0^+$ , (?), and  $4^+$  or  $5^+$  for  $^{18}\text{F}^*(0, 0.94, 1.04, 1.08$  and  $1.13)$ , respectively (1959HI67: tentative assignments). Gamma transitions to  $^{18}\text{F}(0)$  have been observed from the levels at 0.94, 1.04 and 1.08 MeV; the decay of the 1.13 MeV level has not been detected (1958KU81). Proton-gamma coincidence techniques have been used to determine branching ratios and angular correlations involving  $^{18}\text{F}$  states at 0.94, 1.08, 1.7, 2.1, 2.5, and 3.07 MeV (1958KU1D: see also (1956BU1F)): see Fig. 39. All of these states show at least one anisotropic radiation except for  $^{18}\text{F}^*(1.04$  and  $1.08)$ , excluding  $J = 0$  for all but these two ((1958KU1D) and E. Almqvist, private communication).

The mean life of the 0.94 MeV state is  $\leq 5 \times 10^{-9}$  sec, indicating  $J \leq 3$  (1958BR29). The polarization of the  $\gamma$ -radiation indicates positive parity for the state (1958LI41). The strong transition from  $^{18}\text{F}^*(3.07: T = 1)$  suggests  $T = 0$ . For the 1.04 MeV state,  $J = 0^+$  is indicated by the proton angular distributions (1959HI67);  $J = 0^-; T = 0$  is suggested by the  $\gamma$ -branching (E. Almqvist, private communication). The 1.08 MeV state appears to have  $J = 0^+; T = 1$  (see  $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$ ). The absence of the ground-state transition from  $^{18}\text{F}^*(1.12)$  is consistent with its expected  $J = 5^+$  character. The 1.7 MeV state is fixed as  $J = 1^+$  from the present reaction and  $^{19}\text{F}(\text{p}, \text{d})^{18}\text{F}$ . For  $^{18}\text{F}^*(2.1$  and  $2.54)$  assignments of  $J = 1, 2$  and  $1, 2, 3$  are indicated by the branching ratios (E. Almqvist, private communication). See also (1958BR1D, 1958BR86).

$$15. \text{}^{16}\text{O}(\alpha, \text{pn})\text{}^{18}\text{F} \quad Q_{\text{m}} = -18.533$$

See (1947TE01).

$$16. \text{}^{16}\text{O}(\text{}^6\text{Li}, \alpha)\text{}^{18}\text{F} \quad Q_{\text{m}} = 6.067$$

See (1957NO17).

$$17. \text{}^{17}\text{O}(\text{p}, \gamma)\text{}^{18}\text{F} \quad Q_{\text{m}} = 5.619$$

A resonance in the capture cross section may be indicated near 1.24 MeV ( $E_{\text{x}} = 6.79$  MeV),  $\sigma \approx 1$  mb (1956NI1C).

Table 18.7: Energy levels of  $^{18}\text{F}$  from  $^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$ ,  $^{19}\text{F}(^3\text{He}, \alpha)^{18}\text{F}$  and  $^{20}\text{Ne}(\text{d}, \alpha)^{18}\text{F}$

(1959HI67) <sup>a</sup>		(1958KU81) <sup>b</sup>	(1951MI1A) <sup>c</sup>	Assignment <sup>d</sup>	
( $^3\text{He}, \text{p}$ ) (MeV)	( $^3\text{He}, \alpha$ ) (MeV)	( $^3\text{He}, \text{p}$ ) (MeV)	( $\text{d}, \alpha$ ) (MeV $\pm$ keV)	$l_{\text{p}}$	$J^{\pi}; T$
0	0	0	0	0 + 2	1 <sup>+</sup> ; 0
0.941	0.940	0.940		2	2 <sup>+</sup> , 3 <sup>+</sup> ; (0)
1.044	1.042	1.045	1.05 $\pm$ 30	0	0 <sup>(+)</sup> e; (0)
1.084	1.087	1.080 <sup>g</sup>			0 <sup>+</sup> ; 1 <sup>h</sup>
1.126	1.129	1.125		(4)	(4 <sup>+</sup> , 5 <sup>+</sup> )
1.704	1.699		1.83 $\pm$ 30		1 <sup>+</sup> ; 0
2.104	2.105		2.20 $\pm$ 60		1, 2; 0
2.525	2.525		2.61 $\pm$ 50		1, 2, 3; 0
3.064	3.063				2 <sup>+</sup> ; 1
3.135	3.131		3.23 $\pm$ 80		
3.357	3.352				
3.723	3.727				
3.791	3.790				
3.837	3.841				
4.115	4.116		3.92 $\pm$ 40		
4.226	4.227				
4.361	4.358				
	4.400		4.42 $\pm$ 100		
	4.649				
	4.741				
	4.840				
	4.965 <sup>f</sup>		5.01 $\pm$ 90		
	5.292				
	5.500				
	5.603 <sup>f</sup>		5.61 $\pm$ 100		
	5.666				
	5.785				
	6.093				

Table 18.7: Energy levels of  $^{18}\text{F}$  from  $^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$ ,  $^{19}\text{F}(^3\text{He}, \alpha)^{18}\text{F}$  and  $^{20}\text{Ne}(\text{d}, \alpha)^{18}\text{F}$  (continued)

(1959HI67) <sup>a</sup>		(1958KU81) <sup>b</sup>	(1951MI1A) <sup>c</sup>	Assignment <sup>d</sup>	
( $^3\text{He}, \text{p}$ ) (MeV)	( $^3\text{He}, \alpha$ ) (MeV)	( $^3\text{He}, \text{p}$ ) (MeV)	( $\text{d}, \alpha$ ) (MeV $\pm$ keV)	$l_{\text{p}}$	$J^{\pi}; T$
	6.137				
	6.232				
	6.264 <sup>f</sup>				
	6.374				
	6.470				
	6.551				
	6.633				
	6.765				
	6.790				
	6.857				
	7.183				
	7.313				
	7.495				

<sup>a</sup>  $E(^3\text{He}) = 5.7$  and  $5.9$  MeV: energies  $\pm 10$  keV.

<sup>b</sup>  $E(^3\text{He}) = 2.4$  to  $2.9$  MeV.

<sup>c</sup>  $E_{\text{d}} = 7.8$  MeV.

<sup>d</sup>  $l_{\text{p}}$  from proton distributions (1959HI67).

<sup>e</sup>  $J = 0^{-}$ ,  $T = 0$  suggested by  $\gamma$ -branching.

<sup>f</sup>  $\pm 13$  keV.

<sup>g</sup>  $\pm 10$  keV.

<sup>h</sup> Assignment from  $\gamma$ -branching, proton group weak but suggests odd parity (1959HI67).

18.  $^{17}\text{O}(\text{p}, \alpha)^{14}\text{N}$

$$Q_{\text{m}} = 1.197$$

$$E_{\text{b}} = 5.619$$

Resonances in the yield of ground state  $\alpha$ -particles are displayed in Table 18.8 (1957AH20).

19.  $^{17}\text{O}(\text{d}, \text{n})^{18}\text{F}$

$$Q_{\text{m}} = 3.392$$

Table 18.8: Resonances in  $^{17}\text{O}(p, \alpha_0)^{14}\text{N}$  (1957AH20)

$E_p$ (MeV)	$\Gamma_{\text{c.m.}}$ (keV)	$E_x$ (MeV)
1.110	9	6.667
1.273	90	6.821
1.786	$\approx 65$	7.306
2.021	11	7.528
2.048	90	7.553
2.218	11	7.714
2.235	100	7.730
2.406	$\approx 25$	7.891
2.435	$\approx 25$	7.919
2.623	$\approx 40$	8.096
2.753	$\approx 15$	8.219
2.775	$\approx 10$	8.240
2.928	$\approx 50$	8.384

See  $^{19}\text{F}$ .

20.  $^{17}\text{O}(^3\text{He}, d)^{18}\text{F}$

$$Q_m = 0.125$$

Not reported.

21.  $^{17}\text{O}(\alpha, t)^{18}\text{F}$

$$Q_m = -14.194$$

Not reported.

22.  $^{18}\text{O}(p, n)^{18}\text{F}$

$$Q_m = -2.450$$

$$E_{\text{thresh.}} = 2.590 \pm 0.004 \text{ (1950RI59);}$$

$$E_{\text{thresh.}} = 2.584 \pm 0.010 \text{ (1956MA18);}$$

$$E_{\text{thresh.}} = 2.577 \pm 0.008 \text{ (1956HI35 ).}$$

Slow neutron thresholds are reported corresponding to  $E_x = 960 \pm 10$ ,  $1065 \pm 15$ , and, possibly,  $1245 \pm 10$  keV; no others appear with  $E_x < 1.65$  MeV (1956NA1B). Gamma rays are observed with  $E_\gamma = 940 \pm 20$  and  $1040 \pm 20$  keV at  $E_p = 4.3$  and  $4.4$  MeV: the 1040 keV  $\gamma$ -ray exhibits a Doppler shift ( $\tau < 3 \times 10^{-13}$  sec, consistent with  $J = 0^+$ ), the 940 keV  $\gamma$ -ray does not, which is consistent with  $J = 3^+$ ,  $5^+$ . No  $\gamma$ -ray is observed from the 1245 keV state (1956NA1C, 1956RO1C). See also  $^{16}\text{O}(^3\text{He}, p)^{18}\text{F}$  and  $^{19}\text{F}$ .

$$23. \ ^{18}\text{O}(^3\text{He}, t)^{18}\text{F} \quad Q_m = -1.685$$

See (1953KU08).

$$24. \ ^{18}\text{Ne}(\beta^+)^{18}\text{F} \quad Q_m = 4.227$$

See  $^{18}\text{Ne}$ .

$$25. \ ^{19}\text{F}(\gamma, n)^{18}\text{F} \quad Q_m = -10.414$$

See  $^{19}\text{F}$ .

$$26. \ ^{19}\text{F}(n, 2n)^{18}\text{F} \quad Q_m = -10.414$$

See  $^{20}\text{F}$ .

$$27. \ ^{19}\text{F}(p, d)^{18}\text{F} \quad Q_m = -8.187$$

$$Q_0 = -8.12 \pm 0.2 \text{ (1956RE04)}.$$

At  $E_p = 18$  MeV, angular distributions have been measured to the ground state of  $^{18}\text{F}$  (1956RE04, 1958BE28) and to levels at 0.94, 1.05, 1.7, 3.4 and 4.1 MeV (1958BE28). The results indicate  $J = 0^+$  or  $1^+$  for the ground state (1956RE04), even parity for the 0.94, 1.04, 1.7 and 4.1 MeV states; probably assignments are stated to be  $3^+$ ,  $0^+$ ,  $0^+$  and  $2^+$ , respectively, and odd parity,  $J \leq 2$  for the 3.4 MeV state (1958BE28). The reduced width for the ground state reaction is  $\theta^2 = 0.009$  (1956RE04). According to (1959WA16), the distributions permit  $J = 0^+$  or  $1^+$  for  $^{18}\text{F}^*(1.7)$ ;  $J = 0$  is excluded by the observed  $\gamma$ -decay to  $^{18}\text{F}^*(1.04)$ . See also  $^{19}\text{F}$  and (1958EL1A; theor.).

28.  $^{19}\text{F}(\text{d}, \text{t})^{18}\text{F}$   $Q_{\text{m}} = -4.155$   
 $Q_0 = -4.17 \pm 0.02$  (1957EL12).

At  $E_{\text{d}} = 8.9$  MeV proton groups are observed to the ground state and to states at  $0.94 \pm 0.02$  and  $1.07 \pm 0.02$  MeV. The angular distributions are consistent with  $l_{\text{n}} = 0, 1$  and  $0$ , respectively (1957EL12). The  $l_{\text{n}} = 1$ , odd parity assignment for the  $0.94$  MeV state is not in accord with other experiments (see, e.g.,  $^{16}\text{O}(\text{}^3\text{He}, \text{p})^{18}\text{F}$  and  $^{19}\text{F}(\text{p}, \text{d})^{18}\text{F}$ ). See also (1950BO1A, 1951SH1B).

29.  $^{19}\text{F}(\text{}^3\text{He}, \alpha)^{18}\text{F}$   $Q_{\text{m}} = 10.164$   
 $Q_0 = 10.162 \pm 0.015$  (1959HI67).

At  $E(\text{}^3\text{He}) = 5.9$  MeV, 41  $\alpha$ -particle groups have been observed, corresponding to the ground state of  $^{18}\text{F}$  and to excited states with  $E_{\text{x}} < 7.5$  MeV (1959HI67): see Table 18.7.

30.  $^{20}\text{Ne}(\text{n}, \text{t})^{18}\text{F}$   $Q_{\text{m}} = -14.802$

Not reported.

31.  $^{20}\text{Ne}(\text{p}, \text{}^3\text{He})^{18}\text{F}$   $Q_{\text{m}} = -15.567$

Not reported.

32.  $^{20}\text{Ne}(\text{d}, \alpha)^{18}\text{F}$   $Q_{\text{m}} = 2.784$   
 $Q_0 = 2.791 \pm 0.009$  (1954MI61).

Observed alpha-particle groups are listed in Table 18.7 (1951MI1A).

33.  $^{21}\text{Ne}(\text{p}, \alpha)^{18}\text{F}$   $Q_{\text{m}} = -1.746$

Not reported.

34.  $^{25}\text{Mg}(\text{p}, 2\alpha)^{18}\text{F}$   $Q_{\text{m}} = -11.636$

See (1954CO72).

<sup>18</sup>Ne  
(Not illustrated)

GENERAL:

*Theory:* see (1957RA1C).

1. <sup>18</sup>Ne( $\beta^+$ )<sup>18</sup>F  $Q_m = 4.227$

The maximum energy of the positrons is  $3.2 \pm 0.2$  MeV, the half-life is  $1.6 \pm 0.2$  sec:  $\log ft = 2.9 \pm 0.2$  (1954GO17). See also (1956DZ1A).

2. <sup>16</sup>O(<sup>3</sup>He, n)<sup>18</sup>Ne  $Q_m = -2.966$

See (1953KU08).

3. <sup>19</sup>F(p, 2n)<sup>18</sup>Ne  $Q_m = -15.424$

See (1954GO17).

4. <sup>20</sup>Ne(p, t)<sup>18</sup>Ne  $Q_m = -19.812$

Not reported.

## References

(Closed 01 December 1958)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1938FU01 E. Funfer, *Ann. Phys. (Paris)* 32 (1938) 313
- 1939BR1A Brubaker, *Phys. Rev.* 56 (1939) 1181
- 1939DE1A Devons, *Proc. Roy. Soc. A*172 (1939) 127
- 1947TE01 D.H. Templeton, J.J. Howland and I. Perlman, *Phys. Rev.* 72 (1947) 758
- 1948HE1C N.P. Heydenburg and D.R. Inglis, *Phys. Rev.* 73 (1948) 230
- 1949BL26 J.P. Blaser, F. Boehm and P. Marmier, *Phys. Rev.* 75 (1949) 1953
- 1949CH1A R.A. Charpie, K.-H. Sun, B. Jennings and J.F. Nechaj, *Phys. Rev.* 76 (1949) 1255
- 1950BO1A Boyer, M.I.T. Prog. Rept. LNSE (July, 1950)
- 1950RI59 H.T. Richards, R.V. Smith and C.P. Browne, *Phys. Rev.* 80 (1950) 524
- 1951MI1A Middleton and Tai, *Proc. Phys. Soc. (London)* A64 (1951) 801
- 1951PO1A M.L. Pool, D.N. Kundu, P.E. Weiler and T.W. Donaven, *Phys. Rev.* 82 (1951) 305, DA12
- 1951RU24 L. Ruby and J.R. Richardson, *Phys. Rev.* 83 (1951) 698
- 1951SH1B Shull, *Phys. Rev.* 83 (1951) 875, F9
- 1951SU1A K.-H. Sun, B. Jennings, W.E. Shoupp and A.J. Allen, *Phys. Rev.* 82 (1951) 267
- 1952AJ38 F. Ajzenberg and T. Lauritsen, *Rev. Mod. Phys.* 24 (1952) 321
- 1953CU1D Cuer and Magnac-Valette, *J. Phys. Rad.* 14 (1953) 15S
- 1953HE58 N.P. Heydenburg and G.M. Temmer, *Phys. Rev.* 92 (1953) 89
- 1953KU08 D.N. Kundu, T.W. Donaven, M.L. Pool and J.K. Long, *Phys. Rev.* 89 (1953) 1200
- 1954AH37 K. Ahnlund, *Phys. Rev.* 96 (1954) 999
- 1954AH47 K. Ahnlund, S. Thulin and R. Pauli, *Ark. Fys.* 8 (1954) 489
- 1954CO72 B.L. Cohen, H.L. Reynolds and A. Zucker, *Phys. Rev.* 96 (1954) 1617
- 1954GO17 J.D. Gow and L.W. Alvarez, *Phys. Rev.* 94 (1954) 365
- 1954MA1F Mani and Pandhi, *Proc. Indian Acad. Sci.* 40 (1954) 61
- 1954MI61 C. Mileikowsky, *Ark. Fys.* 7 (1954) 117

1954RE1B M.G. Redlich, Phys. Rev. 95 (1954) 448  
 1954SI07 R.M. Sinclair, Phys. Rev. 93 (1954) 1082  
 1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77  
 1955AL1E Alba, Brody, Fernandez, Mazari, Serment and Vazquez, An. del Inst. Fis., Univ. Nacional. Autonoma de Mexico 1 (1955) 1  
 1955BA1Q Banks, Nucleonics 13 (1955) 62  
 1955BE1J Berthelot, Cohen, Cotton, Faraggi, Grjebine, Leveque, Naggiar, Roclawski-Conjeaud and Steinszneider, J. Phys. Rad. 16 (1955) 241  
 1955BU1C Butler, Phys. Rev. 99 (1955) 643A  
 1955EL1A Elliot and Flowers, Proc. Glasgow Conf. (Pergamon Press, 1955)  
 1955EL1B Elliott and Flowers, Proc. Roy. Soc. A229 (1955) 536  
 1955GR1F Graetzer and Pollard, Phys. Rev. 98 (1955) 1184, RA10  
 1955HO28 H.D. Holmgren, T.D. Hanscome and D.K. Willett, Phys. Rev. 98 (1955) 241A  
 1955JA1A Jarmie, Phys. Rev. 98 (1955) 41  
 1955JU1C Juric and Petrovic, Bull. Inst. Nucl. Sci. Boris Kidrich 5 (1955) 1  
 1955KH31 L.M. Khromchenko and V.A. Blinov, Zh. Eksp. Teor. Fiz. 28 (1955) 741; JETP (Sov. Phys.) 1 (1955) 596  
 1955KH35 L.M. Khromchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 19 (1955) 277; Columbia Tech. Transl. 19 (1956) 252  
 1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46  
 1955PR1A Price, Proc. Phys. Soc. (London) A68 (1955) 553  
 1955RE1C D. Reagan, Phys. Rev. 100 (1955) 113  
 1955RE1D M.G. Redlich, Phys. Rev. 99 (1955) 1427  
 1955ST1A Stratton, Blair, Famularo and Stuart, Phys. Rev. 98 (1955) 629  
 1956BA1L D.R. Bach and P.V. Hough, Phys. Rev. 102 (1956) 1341  
 1956BE1B Berger and Loper, Phys. Rev. 104 (1956) 1603  
 1956BR36 C.P. Browne, Phys. Rev. 104 (1956) 1598  
 1956BU1F Butler, Holmgren and Kunz, Bull. Amer. Phys. Soc. 1 (1956) 29  
 1956DR38 R.W.B. Drever, A. Moljk and J. Scobie, Phil. Mag. 1 (1956) 942  
 1956DZ1A Dzhelepov and Peker, Sov. Phys. Dokl. 1 (1956) 71  
 1956GR1F J.C. Grosskreutz, Phys. Rev. 101 (1956) 706  
 1956HI35 H.A. Hill and J.M. BlPhys. Rev. 104 (1956) 198  
 1956JA31 N. Jarmie, Phys. Rev. 104 (1956) 1683

1956JU1G Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 6 (1956) 41  
 1956KO26 B. Koudijs, Ph.D. Thesis, Univ. of Utrecht (1956)  
 1956MA18 H. Mark and C. Goodman, Phys. Rev. 101 (1956) 768  
 1956NA1B Naggiar, Roclawski-Conjeaud, Szteinszneider and Thirion, Compt. Rend. 242 (1956) 1443  
 1956NA1C Naggiar, Roclawski-Conjeaud, Szteinszneider and Thirion, J. Phys. Rad. 17 (1956) 561  
 1956NE1B Neudachin, Zh. Eksp. Teor. Fiz. 31 (1956) 892; JETP (Sov. Phys.) 4 (1957) 756  
 1956NI1C Nikolic, Povh and Zupancic, Bull. Inst. Nucl. Sci. Boris Kidrich 6 (1955) 51  
 1956RE04 J.B. Reynolds and K.G. Standing, Phys. Rev. 101 (1956) 158  
 1956RO1A Roclawski-Conjeaud and Cotton, Nucl. Phys. 1 (1956) 603; J. Phys. Rad. 17 (1956) 552; Physica 22 (1956) 1155A  
 1956RO1C Roclawski-Conjeaud, Naggiar, Szteinszneider and Thirion, Physica 22 (1956) 1158A  
 1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434  
 1956SH1A Sher and Floyd, Phys. Rev. 102 (1956) 242  
 1956TH1B R. Thieberger and I. Talmi, Phys. Rev. 102 (1956) 923  
 1956VA17 F.P.G. Valckx, Ph.D. Thesis, Univ. of Utrecht (1956)  
 1957AH20 K. Ahnlund, Phys. Rev. 106 (1957) 124  
 1957BA14 E. Baumgartner and H.W. Fulbright, Phys. Rev. 107 (1957) 219  
 1957BI80 O.M. Bilaniuk and P.V.C. Hough, Phys. Rev. 108 (1957) 305  
 1957EL12 F.A. El Bedewi and I. Hussein, Proc. Phys. Soc. (London) A70 (1957) 233  
 1957GR1D Groshev and Demidov, Sov. J. At. Energy 3 (1957) 853  
 1957JU1A Juric, Bull. Inst. Nucl. Sci. Boris Kidrich 7 (1957) 1  
 1957NO17 E. Norbeck Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754  
 1957RA1C G. Rakavy, Nucl. Phys. 4 (1957) 375  
 1957RI44 F.L. Ribe, Phys. Rev. 106 (1957) 767  
 1957YO04 T.E. Young, G.C. Phillips and R.R. Spencer, Phys. Rev. 108 (1957) 72  
 1958AL03 E. Almqvist, D.A. Bromley and J.A. Kuehner, Bull. Amer. Phys. Soc. 3 (1958) 27, J12  
 1958BA59 J.B. Ball, A.W. Fairhall and I. Halpern, Bull. Amer. Phys. Soc. 3 (1958) 322, H6  
 1958BE28 E.F. Bennett, Bull. Amer. Phys. Soc. 3 (1958) 26, J8  
 1958BE74 W.L. Bendel, J. McElhinney and R.A. Tobin, Phys. Rev. 111 (1958) 1297  
 1958BR1D Bromley, Proc. Rehovoth Conf. (North-Holland Pub. Co., Amsterdam, 1958)

- 1958BR29 D.A. Bromley, J.A. Kuehner and E. Almqvist, Bull. Amer. Phys. Soc. 3 (1958) 27, J13
- 1958BR86 D.A. Bromley, J.A. Kuehner and E. Almqvist, Bull. Amer. Phys. Soc. 3 (1958) 199, P5
- 1958DA16 A.W. Dalton, S. Hinds and G. Parry, Proc. Phys. Soc. (London) A71 (1958) 252
- 1958EL1A El Nadi and Hadid, Nucl. Phys. 8 (1958) 51
- 1958GO69 H.E. Gove and A.E. Litherland, Bull. Amer. Phys. Soc. 3 (1958) 199, P7
- 1958HE1F Henley, Conf. on Photonucle. Reactions, National Bureau of Standards (1958)
- 1958HE54 D.F. Herring, R. Chiba, B.R. Gasten and H.T. Richards, Phys. Rev. 112 (1958) 1210
- 1958HE56 D.F. Herring, Phys. Rev. 112 (1958) 1217
- 1958HU18 D.J. Hughes and R.B. Schwartz, BNL-325, 2nd Ed. (1958); BNL-325, 2nd Ed., Suppl. I (1960)
- 1958KA32 E. Kashy, P.D. Miller and J.R. Risser, Phys. Rev. 112 (1958) 547
- 1958KU1D Kuehner, Almqvist and Bromley, Bull. Amer. Phys. Soc. 3 (1958) 27
- 1958KU81 J.A. Kuehner, E. Almqvist and D.A. Bromley, Phys. Rev. Lett. 1 (1958) 260
- 1958LI41 A.E. Litherland and H.E. Gove, Bull. Amer. Phys. Soc. 3 (1958) 200, P12
- 1958PH37 W.R. Phillips, Phys. Rev. 110 (1958) 1408
- 1958RE1B M.G. Redlich, Phys. Rev. 110 (1958) 468
- 1958WE29 J.A. Weinman and E.A. Silverstein, Phys. Rev. 11 (1958) 277
- 1959HI67 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A73 (1959) 721
- 1959WA16 E.K. Warburton, Phys. Rev. 113 (1959) 595
- MO54 Unknown Source