

Energy Levels of Light Nuclei $A = 18$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 18$ – 20 was published in *Nuclear Physics A190* (1972), p. 1. This version of $A = 18$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1971)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 18$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{18}\text{C}\$](#) , [\$^{18}\text{N}\$](#) , [\$^{18}\text{O}\$](#) , [\$^{18}\text{F}\$](#) , [\$^{18}\text{Ne}\$](#) , [\$^{18}\text{Na}\$](#)

B. Tables of Recommended Level Energies:

[Table 18.1](#): Energy levels of ^{18}O

[Table 18.10](#): Energy levels of ^{18}F

[Table 18.23](#): Energy levels of ^{18}Ne

C. [References](#)

D. Figures: [\$^{18}\text{O}\$](#) , [\$^{18}\text{F}\$](#) , [\$^{18}\text{Ne}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

^{18}C
(Not illustrated)

^{18}C is particle stable. Therefore its atomic mass excess, $M - A$, must be < 29.84 MeV [$^{16}\text{C} + 2n$] (1970WA1G). ^{18}C has been observed in the bombardment of ^{232}Th by 122 MeV ^{18}O ions (1969AR13, 1970AR1D) and in the 3 GeV proton bombardment of Au (1970RA1A). See also (1960ZE03, 1968PO04, 1971BU1E).

^{18}N
(Fig. 4)

GENERAL: See (1960ZE03, 1961BA1C, 1962GO1B, 1969AR13, 1971AR02).

Mass of ^{18}N : From the Q -value of the $^{18}\text{O}(t, ^3\text{He})^{18}\text{N}$ reaction, $M - A$ for ^{18}N is 13.274 ± 0.030 MeV (1969ST07).

1. $^{18}\text{N}(\beta^-)^{18}\text{O}$ $Q_m = 14.06$

The half-life of ^{18}N is 0.63 ± 0.03 sec (1964CH19). $\log ft = 4.88^\dagger$. $E_\beta(\text{max}) = 9.4 \pm 0.4$ MeV. The decay is to $^{18}\text{O}^*(4.45)$, whose subsequent γ -decay has been studied using $\beta - \gamma$ coincidences: see ^{18}O . The allowed nature of the decay to the 1^- state at 4.45 MeV leads to $J^\pi = 0^-, 1^-$ or 2^- for the ^{18}N ground state (1964CH19).

2. $^{18}\text{O}(n, p)^{18}\text{N}$ $Q_m = -13.27$

This reaction has been studied at $E_n = 19$ MeV: see above (1964CH19).

3. $^{18}\text{O}(t, ^3\text{He})^{18}\text{N}$ $Q_m = -14.04$
 $Q_0 = -14.038 \pm 0.030$ (1969ST07).

At $E_t = 22$ MeV, ^3He groups corresponding to the ground state of ^{18}N have been observed by (1969ST07).

[†] B.A. Zimmerman, private communication.

GENERAL: (See also (1959AJ76).)

Shell model: (1957WIIE, 1960TA1C, 1962HO1C, 1962TA1B, 1962TA1D, 1963HA05, 1963PA03, 1963SA07, 1964CO24, 1964IN03, 1964MC1A, 1964PA1D, 1964WA1F, 1965BA1J, 1965BE1T, 1965DE1H, 1965EL06, 1965EN02, 1965FE1B, 1965FE02, 1965NA1A, 1965ZA1B, 1966AR10, 1966BA2E, 1966BA2C, 1966BO25, 1966BR1R, 1966HU09, 1966IN01, 1966KU05, 1966LA1E, 1966LE11, 1966RI1F, 1966RO01, 1967BA04, 1967BR1G, 1967EN01, 1967FE01, 1967FL01, 1967GR09, 1967GR1D, 1967HO11, 1967IN03, 1967KU09, 1967KU13, 1967LA1H, 1967LY02, 1967MO1J, 1967PA05, 1967PA1K, 1967PI1B, 1967RO1F, 1967ST02, 1967VI1B, 1967WO1C, 1968AR02, 1968BE1T, 1968BH1B, 1968CL03, 1968CO1N, 1968DE1M, 1968DE13, 1968DW1A, 1968EL1C, 1968FR03, 1968GA06, 1968GA16, 1968GU1E, 1968GU1C, 1968GU1G, 1968HA17, 1968HA1P, 1968HE1H, 1968KA1E, 1968KA1C, 1968KO1L, 1968KU1E, 1968NA09, 1968NA19, 1969BA2F, 1969BE1T, 1969BU15, 1969EL1B, 1969FE1A, 1969GU1E, 1969HO32, 1969IG1A, 1969KA07, 1969KA09, 1969KU1G, 1969MA1T, 1969ME06, 1969SA1F, 1969ST1G, 1969UL03, 1969VO1E, 1969ZU03, 1970BA2E, 1970EL1G, 1970EL08, 1970GA1J, 1970HA49, 1970KA32, 1970TA1J, 1970TR08, 1970TR07, 1971AR1R, 1971EL07, 1971HO12, 1971JA06, 1971LO23, 1971MO05, 1971PR16, 1971QU01, 1971UL04, 1971VI08, 1971WI01, 1972GA02).

Cluster, collective and deformed models: (1959YU1A, 1963MA1D, 1964BR1H, 1965FE1B, 1965FE02, 1966BE29, 1966BR1Q, 1966HA1K, 1966RI1F, 1966RO01, 1967BH07, 1967BR1G, 1967FE01, 1967GR1D, 1967PA1K, 1967RO1F, 1968EL1C, 1969BA2E, 1969DR1B, 1969FE1A, 1969ZA1D, 1970BA2B, 1970TR07, 1971AR1R, 1971UL04).

Astrophysical questions: (1970BA1M).

Electromagnetic transitions: (1963BU11, 1964BR1H, 1965EL06, 1965EN02, 1965FE02, 1965ST22, 1966BO25, 1967GR1D, 1967IN03, 1967LA1H, 1968EL1C, 1968HA17, 1968LA1G, 1968LE02, 1968SH06, 1969BE1T, 1969BE22, 1970EL08, 1970HA49, 1970TR07).

Special levels: (1960EV1A, 1962DA04, 1962TA1B, 1964BR1H, 1964EN1A, 1964MC1A, 1965BE1T, 1965LE1C, 1966BO25, 1966BR1R, 1966HU09, 1966MI1G, 1967BH07, 1967EN01, 1967FL01, 1967GR09, 1967PA1K, 1967VI1B, 1968AR02, 1968BH1B, 1968CO1N, 1968FR03, 1968GA06, 1968GA16, 1968HA1P, 1968KA1C, 1968LE02, 1968NA09, 1969BA1Z, 1969BE1T, 1969FE1A, 1969KA09, 1969KA29, 1969MA1T, 1969ME06, 1969OS01, 1970EL1G, 1970EL08, 1970GA1J, 1970HO17, 1971AR1R, 1971EL07, 1971JA06, 1971PR16, 1971QU01, 1971UL01, 1971WI01, 1972GA02).

Other theoretical topics: (1961BA1D, 1962DA04, 1963BU11, 1963VL1A, 1964EN1A, 1964HE1C, 1964IN03, 1964TR1A, 1964WA1E, 1965DE1H, 1965ER04, 1965GO1F, 1965KA1B, 1965KA1C, 1965NA1A, 1965NI1A, 1965ZA1B, 1966BR1R, 1966BR1P, 1966DA1E, 1966DO1C, 1966GI1A, 1966KU05, 1966LE11, 1966OL1C, 1966SU1D, 1966WA1H, 1966YO1B, 1967FE01, 1967FL01, 1967GR09, 1967KU1G, 1967KU13, 1967MO1J, 1967ST02, 1967WO1C, 1968BA2H, 1968BE1T, 1968BL1G, 1968CO1N, 1968DE1M, 1968DW1A, 1968EL1C, 1968GU1E, 1968GU1C, 1968GU1F,

1968GU1G, 1968JO1E, 1968KA1C, 1968KO1K, 1968LE02, 1968MU1B, 1968NE1C, 1968SU1C, 1968SU1D, 1968VA1J, 1968VA24, 1969DE06, 1969JA1P, 1969JO1L, 1969KA29, 1969MU09, 1969OS01, 1969RA28, 1969SO08, 1970AG1C, 1970BA1Z, 1970DI1G, 1970EL08, 1970GA1J, 1970PR1D, 1970SU1B, 1971BA14, 1971BO21, 1971DZ06, 1971EL07, 1971LA1D, 1971LE1H, 1971LO23, 1971PR16, 1971SC01, 1971UL01, 1971VI08, 1972GA02).

Table 18.1: Energy levels of ^{18}O ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$0^+; 1$	stable		2, 3, 4, 7, 10, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 36
1.9821 ± 0.8	2^+	$\tau_m = 3.8 \pm 0.6$	γ	2, 3, 4, 7, 10, 15, 17, 19, 20, 21, 22, 25, 32, 34, 35
3.5529 ± 2.1	4^+	> 3	γ	3, 4, 7, 10, 15, 22, 25, 34, 35
3.6317 ± 2.0	0^+	3 ± 1	γ	2, 3, 4, 7, 10, 15, 17, 21, 22, 25, 34, 35
3.9191 ± 2.0	2^+	0.18 ± 0.06	γ	2, 3, 4, 7, 10, 15, 21, 22, 25, 34, 35
4.4488 ± 3.5	1^-	$0.11^{+0.08}_{-0.06}$	γ	2, 7, 10, 15, 17, 19, 21, 22, 25, 34, 35
5.090 ± 5	3^-		γ	7, 10, 15, 21, 22, 25, 35
5.250 ± 6	2^+		γ	4, 7, 10, 15, 21, 25, 34, 35
5.329 ± 7	0^+		γ	10, 15, 21, 22, 25, 34, 35
5.372 ± 7	3^+		γ	10, 15, 21, 22, 25, 35
5.517 ± 9	2^-		γ	10, 15, 19, 21, 25, 35
6.191 ± 9	1^-		γ	10, 15, 25, 34, 35
6.341 ± 10			(γ)	10, 15, 25, 34, 35
6.391 ± 10	3^-		γ	10, 25, 35
6.86 ± 20	(0^-)		γ	25, 35
7.114 ± 2	4^+		γ, α	4, 7, 15, 25, 34, 35

Table 18.1: Energy levels of ^{18}O ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
7.620 \pm 2	1^-	$\Gamma < 2.5$ keV	γ, α	4, 25, 34, 35
7.75 \pm 20			γ	35
7.848 \pm 14				7, 15, 25, 34, 35
7.961 \pm 14	$(3^+, 4^-)$		γ	15, 25, 35
8.039 \pm 2	1^-	< 2.5	γ, α	4, 25, 35
8.122 \pm 10	5^-		γ, α	4, 7, 25, 35
8.213 \pm 4	2^+	1.0 ± 0.8	γ, n, α	4, 5, 6, 25, 35
8.283 \pm 3	3^-	8 ± 1	n, α	5, 6, 7, 25, 35
8.403 \pm 7		8 ± 6	n, α	5, 35
8.480 \pm 20				25, 35
8.640 \pm 20				25, 35
8.817 \pm 12		70 ± 12	n, α	5, 6, 25
8.955 \pm 4		43 ± 3	n, α	5, 6, 25
9.03			n, α	6, 25
9.10			n, α	6, 25
9.361 \pm 15		27 ± 15	n, α	5, 6, 7, 25
9.39 \pm 40		≈ 120	n, α	5, 6, 7, 25
9.47 \pm 40		≈ 65	n, α	5, 6
9.675 \pm 10		60 ± 30	n, α	5, 6, 25, 34
9.72 \pm 30			(n, α)	5, 25
9.88 \pm 40		≈ 150	n, α	5, 6, 25, 34
10.119 \pm 10	3^-	16 ± 4	n, α	5, 6, 25
10.29 \pm 20	4^+		n, α	5, 6, 7, 25
10.38 \pm 20	3^-		n, α	5, 6, 25
10.58 \pm 20			n, α	5, 6
10.82 \pm 20			n, α	5, 6
10.91 \pm 20			n, α	5, 6, 7
10.99 \pm 20			n, α	5, 6
11.14 \pm 20	$(0, 1, 2)^-$		n, α	5, 6, 7, 34
11.39 \pm 20	(2^+)		n, α	5, 6

Table 18.1: Energy levels of ^{18}O ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ_m (psec) or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
11.41 \pm 20	(4 ⁺)		n, α	5, 6
11.62 \pm 20	5 ⁻		n, α	5, 6, 7, 25
11.69 \pm 20	6 ⁺		n, α	5, 6, 7, 25, 34
11.82 \pm 20	(3 ⁻)		n, α	5, 6, 34
12.04 \pm 20	(2 ⁺)		n, α	5, 6
12.25 \pm 20	1 ⁻		n, α	5, 6, 34
12.33 \pm 20	5 ⁻		n, α	5, 6, 7
12.49 \pm 20	4 ⁺		n, α	5, 6
12.53 \pm 20	6 ⁺		n, α	5, 6, 7
14.05 \pm 140				19, 34
14.56 \pm 100				34
15.1 \pm 200			γ , n	19
15.95 \pm 200				19
18.4 \pm 200				19
20.2 \pm 200				19
21.9 \pm 200				19
23.7 \pm 200				19
25.2 \pm 200				19
26.9 \pm 200				19

^a See also Tables 18.2 and 18.3.

Reactions involving pions and muons: (1965PA1F, 1967BA78, 1968BA2G, 1968TA1C, 1968WI1B, 1969CH1C, 1969KO1F, 1969WU1A, 1970BA1E, 1970CH25, 1970CH1C, 1970HA46).

Complex reactions involving ^{18}O : (1968GA03, 1969AR13, 1971AR02).

Ground state: (1971MO05).

1. (a) $^{11}\text{B}(^7\text{Li}, \text{p})^{17}\text{N}$ $Q_m = 8.417$ $E_b = 24.359$
- (b) $^{11}\text{B}(^7\text{Li}, \text{d})^{16}\text{N}$ $Q_m = 4.758$
- (c) $^{11}\text{B}(^7\text{Li}, \text{t})^{15}\text{N}$ $Q_m = 8.525$
- (d) $^{11}\text{B}(^7\text{Li}, \alpha)^{14}\text{C}$ $Q_m = 18.131$

Cross sections to various of the final states have been measured at $E(^7\text{Li}) = 5.00$ MeV (1966MC05).

$$2. \text{}^{12}\text{C}(^7\text{Li}, \text{p})^{18}\text{O} \quad Q_{\text{m}} = 8.402$$

Radiative decay and lifetime measurements are reported by (1964ES02): see Tables 18.2 and 18.3. Angular distributions have been measured at $E(^7\text{Li}) = 3.24$ to 3.64 MeV (1967MO23; p_0 , p_1). See also (1960SH05, 1970CA1N).

$$3. \text{}^{13}\text{C}(^7\text{Li}, \text{d})^{18}\text{O} \quad Q_{\text{m}} = 5.681$$

Gamma-ray energies of 1982.3 ± 0.8 [$1.98 \rightarrow 0$], 1571.9 ± 2.5 [$3.55 \rightarrow 1.98$], 1649.8 ± 2.2 [$3.63 \rightarrow 1.98$] and 1938.6 ± 2.1 keV [$3.92 \rightarrow 1.98$], due to the transitions shown in brackets, have been measured by (1969TH01).

$$4. \text{}^{14}\text{C}(\alpha, \gamma)^{18}\text{O} \quad Q_{\text{m}} = 6.228$$

Four resonances in the yield of capture γ -rays are observed for $E_{\alpha} = 0.50$ to 3.00 MeV at $E_{\alpha} = 1.14$, 1.79 , 2.33 and 2.44 MeV: see Table 18.4. Gamma-ray angular distribution and correlation measurements lead to $J^{\pi} = 4^+$, 1^- , 1^- and 5^- for $^{18}\text{O}^*(7.11, 7.62, 8.04, 8.12)$, as well as to J^{π} assignments for lower states involved in the cascade decay: see Table 18.2 (1958GO1A, 1958PH37, 1966LE1B, 1967LE02). (1970DU1D) report resonance at $E_{\text{p}} = 2.56$ MeV ($E_{\text{x}} = 8.22$) with $J^{\pi} = 2^+$.

$$5. \text{}^{14}\text{C}(\alpha, \text{n})^{17}\text{O} \quad Q_{\text{m}} = -1.819 \quad E_{\text{b}} = 6.228$$

The relative neutron yield has been measured for $E_{\alpha} = 2.3$ to 8.5 MeV: the parameters of observed resonances are displayed in Table 18.4 (1956SA06, 1966BA03, 1970MO13).

$$6. \text{}^{14}\text{C}(\alpha, \alpha)^{14}\text{C} \quad E_{\text{b}} = 6.228$$

Observed anomalies in the scattering for $E_{\alpha} = 2$ to 8.2 MeV shown in Table 18.4 (1958WE29, 1970MO13). The yield of elastic scattering has been measured up to $E_{\alpha} = 16.5$ MeV (1970MO13).

Table 18.2: Radiative decays in ^{18}O

E_i (MeV)	J_i^π	E_f (MeV)	Branch ^a (%)	$\Gamma_\alpha \Gamma_\gamma / \Gamma$ (eV)	Refs.
1.98	2^+	0	100	$B(E2) = (4.9 \pm 1.1)^d$ $= (4.6 \pm 1.4)^d$	(1967DEZW) (1968AN20)
3.55	4^+	0	< 4		(1959GO74)
		1.98	> 96		(1959GO74, 1961LI03)
			100		(1967LE02, 1967CH1D, 1971BE45)
3.63	0^+	0		$\Gamma_\pi = 1.9 \times 10^{-3} \Gamma_\gamma$	(1963GO32)
		1.98	> 85		(1961LI03)
			100		(1967LE02, 1967CH1D, 1971BE45)
3.92 ^b	2^+	0	6.5		(1964ES02)
			15		(1967MO09)
			15 ± 5		(1964GO11)
			15 ± 2		(1965OL02)
			12 ± 5		(1971BE45)
		1.98	93.5		(1964ES02)
			85		(1967MO09)
			> 85		(1961LI03)
			85 ± 5		(1964GO11)
			85 ± 2		(1965OL02)
			88 ± 5		(1971BE45)
4.45 ^c	1^-	0	< 2		(1964ES02, 1964GO11)
			< 4		(1965OL02)
			< 5		(1971BE45)
			< 1		(1967LE02, 1967CH1D)
		1.98	26		(1964ES02)
			28		(1967CH1D)
			37 ± 5		(1964GO11)
			36 ± 5		(1965OL02)
			32 ± 5		(1971BE45)
		3.63	74		(1964ES02)
			72		(1967CH1D)
			63 ± 5		(1964GO11)
			64 ± 5		(1965OL02)

Table 18.2: Radiative decays in ^{18}O (continued)

E_i (MeV)	J_i^π	E_f (MeV)	Branch ^a (%)	$\Gamma_\alpha \Gamma_\gamma / \Gamma$ (eV)	Refs.			
5.09	3^-	0	68 ± 5		(1971BE45)			
			< 10		(1965OL02)			
			< 5		(1971BE45)			
		1.98	> 90		(1965OL02)			
			78		(1967CH1D)			
			81		(1966LO12)			
			75 ± 3		(1971BE45)			
			3.55		6	(1967CH1D)		
					5	(1966LO12)		
			3.92		9 ± 3	(1971BE45)		
					14	(1966LO12)		
			5.25 ^b		2^+	0	16 ± 3	(1971BE45)
							32	(1967CH1D)
40	(1966LO12)							
1.98	40 ± 8	(1965OL02)						
	41 ± 6	(1967MO09)						
	32 ± 3	(1971BE45)						
	68	(1967CH1D)						
	60	(1966LO12)						
	60 ± 8	(1965OL02)						
	59 ± 6	(1967MO09)						
	68 ± 3	(1971BE45)						
	3.55	< 4		(1967MO09)				
	3.63	< 3		(1967MO09)				
3.92	< 5	(1967MO09)						
5.33 ^b	0^+	4.45	< 4	(1967MO09)				
			1.98	63	(1966LO12)			
		3.92	61 ± 2	(1971BE45)				
			10	(1966LO12)				
			27	(1966LO12)				
5.37	3^+	0	39 ± 2	(1971BE45)				
			< 2	(1967MO09)				

Table 18.2: Radiative decays in ^{18}O (continued)

E_i (MeV)	J_i^π	E_f (MeV)	Branch ^a (%)	$\Gamma_\alpha \Gamma_\gamma / \Gamma$ (eV)	Refs.
5.52	(2^-)	1.98	85		(1966LO12)
			88 ± 3		(1967MO09)
		3.55	< 3		(1967MO09)
		3.92	15		(1966LO12)
			12 ± 3		(1967MO09)
		4.45	< 2		(1967MO09)
		1.98	48		(1967CH1D)
			65		(1966LO12)
		3.92	23		(1967CH1D)
			10		(1966LO12)
6.19	1^-	4.45	29	(1967CH1D)	
			25	(1966LO12)	
		0	> 90	(1967LE02)	
			88	(1967CH1D)	
			100	(1971BE45)	
		1.98	< 5	(1967LE02)	
			≤ 10	(1967CH1D)	
		4.45	< 5	(1967LE02)	
			12	(1967CH1D)	
		6.34 + 6.39			
6.39	3^-	1.98	90 ± 5	(1971BE45)	
		3.92	10 ± 5	(1971BE45)	
6.86	(0^-)	4.45	100	(1967CH1D)	
7.11	4^+	1.98	44	1.2×10^{-2}	(1959GO74, 1961LI03)
			29	(1967CH1D)	
			26 ± 2	(1967LE02)	
		3.55	56	1.5×10^{-2}	(1959GO74, 1961LI03)
			71	(1967CH1D)	
			70 ± 3	4.2×10^{-2} ^e	(1967LE02)
		3.92	4 ± 1	(1967LE02)	
		4.45	≤ 15	(1967CH1D)	
		5.09	≤ 15	(1967CH1D)	

Table 18.2: Radiative decays in ^{18}O (continued)

E_i (MeV)	J_i^π	E_f (MeV)	Branch ^a (%)	$\Gamma_\alpha\Gamma_\gamma/\Gamma$ (eV)	Refs.	
7.62	1^-	0	35	8×10^{-2}	(1958PH37)	
			33		(1959GO74, 1961LI03)	
			24 ± 2		(1967LE02)	
		1.98	65	0.16	(1958PH37)	
			67		(1959GO74, 1961LI03)	
			62 ± 3		(1967LE02)	
			3.55		0.34 ^e	(1958PH37)
			3.92			(1958PH37)
			4.45			(1967LE02)
			5.33			(1967LE02)
		6.19	(1967LE02)			
		7.75	1^-	1.98	50	(1967CH1D)
				4.45	11	(1967CH1D)
5.09	39			(1967CH1D)		
5.25	≤ 10			(1967CH1D)		
5.37	≤ 10			(1967CH1D)		
6.34	≤ 10			(1967CH1D)		
7.96	$(3^+, 4^-)$	3.55	67	(1967CH1D)		
		5.09	12	(1967CH1D)		
		5.37	21	(1967CH1D)		
		6.39	< 15	(1967CH1D)		
8.04	1^-	0	30	0.89 ^e	(1958PH37)	
			16 ± 1		(1967LE02)	
		1.98	70		(1958PH37)	
		68 ± 3	(1967LE02)			
		3.55	< 15		(1958PH37)	
		3.63	11 ± 1		(1967LE02)	
		3.92	< 15		(1958PH37)	
8.12	5^-	5.25	5 ± 1	(1967LE02)		
		3.55	> 95	(1967LE02)		
		8.21	2^+	0	100	(1970DU1D)

^a The last value listed for each transition is believed to be the most reliable.

^b See also (1967LE02).

^c See also (1964CH19, 1967LE02).

^d $e^2 \times 10^{-51} \text{ cm}^4$.

^e $\Gamma_\gamma \Gamma_\alpha / \Gamma$ for *all* transitions from this state.

7. $^{14}\text{C}(^7\text{Li}, \text{t})^{18}\text{O}$ $Q_m = 3.761$

At $E(^7\text{Li}) = 20.4 \text{ MeV}$, triton groups are observed corresponding to a number of states of ^{18}O with $E_x < 12.6 \text{ MeV}$. Angular distributions were obtained for some of these, including $^{18}\text{O}^*(0, 1.98, 7.11, 11.69)$ with $J^\pi = 0^+, 2^+, 4^+, 6^+$. The latter two are the most strongly populated in this reaction: they appear to be part of the ground state rotational band (1970MO17).

8. $^{15}\text{N}(\text{t}, \text{p})^{17}\text{N}$ $Q_m = -0.108$ $E_b = 15.834$

See (1956SH1A).

9. $^{15}\text{N}(\alpha, \text{p})^{18}\text{O}$ $Q_m = -3.980$

Not reported.

10. $^{16}\text{O}(\text{t}, \text{p})^{18}\text{O}$ $Q_m = 3.707$

Proton groups corresponding to many states of ^{18}O have been reported by (1960JA13, 1960JA17, 1962HI06): see Table 18.5. The L assignments displayed there have been derived from PWBA analysis of angular distributions, principally those obtained by (1964MI05). For a complete summary of angular distribution measurements, see Table 18.6.

Measurements of the angular correlation of cascade γ -rays require $J = 0$ and 2 for $^{18}\text{O}^*(3.63, 3.92)$. Lifetime measurements are reported by (1961LI03, 1963LI07): see Table 18.3.

See also (1964GA1B), (1967CH1L, 1967OG1A) and (1963RO1C, 1964TR1A, 1965GL07, 1965MA1L, 1965SH1E, 1966GL1C, 1966GL1D, 1966SH1F, 1966ST1F, 1967DO1B, 1967MA1E, 1967VE1B, 1968KO04, 1968KO1K, 1969LI1G, 1969SO08; theor.).

11. $^{17}\text{O}(\text{n}, \gamma)^{18}\text{O}$ $Q_m = 8.047$

Table 18.3: Lifetime measurements of some ^{18}O states

$^{18}\text{O}^*$ (MeV)	τ_m (psec) ^a	Reaction	Refs.
1.98 ^b	> 1	$^{17}\text{O}(\text{d}, \text{p})$	(1969NI09)
	$6.1^{+5.0}_{-2.0}$	$^{12}\text{C}(^7\text{Li}, \text{p})$	(1964ES02)
	3.3 ± 1.5	$^{18}\text{O}(\text{e}, \text{e}')$	(1961LA09)
	3.7 ± 0.7	$^{16}\text{O}(\text{t}, \text{p})$	(1963LI07)
	$4.2^{+1.5}_{-1.0}$	$^{16}\text{O}(\text{t}, \text{p})$	(1971HI06)
3.55	> 3	$^{16}\text{O}(\text{t}, \text{p})$	(1963LI07)
3.63	> 1	$^{16}\text{O}(\text{t}, \text{p})$	(1961LI03)
	≥ 2.6	$^{12}\text{C}(^7\text{Li}, \text{p})$	(1964ES02)
	3 ± 1	$^{16}\text{O}(\text{t}, \text{p})$	(1963LI07)
3.92	$0.35^{+0.3}_{-0.2}$	$^{12}\text{C}(^7\text{Li}, \text{p})$	(1964ES02)
	0.18 ± 0.06	$^{16}\text{O}(\text{t}, \text{p})$	(1963LI07)
4.45	< 0.1	$^{16}\text{O}(\text{t}, \text{p})$	(1963LI07)
	$0.11^{+0.08}_{-0.06}$	$^{12}\text{C}(^7\text{Li}, \text{p})$	(1964ES02)

^a The last value quoted for each state is believed to be the most reliable.

^b See also (1968AN20).

See (1968FO1A, 1970CL1C).

12. $^{17}\text{O}(\text{n}, \text{n})^{17}\text{O}$

$$E_b = 8.047$$

The coherent scattering length (thermal, bound) is 5.79 fm (1969BA1P). See also (1970TA1E; theor.).

13. $^{17}\text{O}(\text{n}, \text{p})^{17}\text{N}$

$$Q_m = -7.896$$

$$E_b = 8.047$$

At $E_n = 14.1$ MeV, the cross section is 21.5 ± 1.7 mb (1970ME31). See also (1958RO1A, 1964AM02).

14. $^{17}\text{O}(\text{n}, \alpha)^{14}\text{C}$

$$Q_m = 1.819$$

$$E_b = 8.047$$

The thermal cross section is 235 ± 5 mb (1961HA43). See also (1964ST25), (1964GA1A; theor.) and (1968FO1A) for a discussion of astrophysical implications.

Table 18.4: Resonances in $^{14}\text{C}(\alpha, \gamma)^{18}\text{O}$, $^{14}\text{C}(\alpha, n)^{17}\text{O}$ and $^{14}\text{C}(\alpha, \alpha)^{14}\text{C}$

E_α (MeV \pm keV)	Γ_{lab} (keV)	Particles out ^a	$^{18}\text{O}^*$ (MeV)	J^π	Refs.
1.140 \pm 2		γ	7.114	4 ⁺	(1959GO74, 1966LE1B, 1967LE02)
1.790 \pm 2	< 3	γ	7.620	1 ⁻	(1958PH37, 1959GO74, 1966LE1B, 1967LE02)
2.330 \pm 2	< 3	γ, α_0	8.039	1 ⁻	(1958PH37, 1958WE29, 1966LE1B, 1967LE02)
2.440 \pm 12		γ	8.125	5 ⁻	(1966LE1B, 1967LE02)
2.554 \pm 4	1.3 \pm 1	γ, n, α_0	8.214	2 ⁺	(1956SA06, 1958WE29, 1966BA03)
2.643 \pm 3	10 \pm 1	n, α_0	8.283	3 ⁻	(1956SA06, 1958WE29, 1966BA03)
2.800 \pm 7	10 \pm 7	n	8.405		(1956SA06, 1966BA03)
3.330 \pm 12	90 \pm 15	n, α_0	8.817		(1956SA06, 1958WE29, 1966BA03)
3.508 \pm 4	55 \pm 3	n, α_0	8.955 ^g		(1956SA06, 1958WE29, 1966BA03)
4.030 \pm 15	35 \pm 20	n, (α_0)	9.361		(1966BA03, 1970MO13)
4.07 \pm 40	\approx 150	n, (α_0)	9.39		(1966BA03, 1970MO13)
4.17 \pm 40	\approx 70	n, (α_0)	9.47		(1966BA03, 1970MO13)
4.434 \pm 10	80 \pm 40	n, (α_0)	9.675		(1966BA03, 1970MO13)
4.70 \pm 40	\approx 200	n, (α_0)	9.88		(1966BA03, 1970MO13)
5.004 \pm 10	21 \pm 5	n, α_0	10.119	3 ⁻	(1966BA03, 1970MO13)
5.23 ^f	b	n, α_0	10.29	4 ⁺	(1970MO13)
5.34	b	n, α_0	10.38	3 ⁻	(1970MO13)
5.60	c	n, α_0	10.58		(1970MO13)
5.90	d	n, α_0	10.82		(1970MO13)
6.02	d	n, α_0	10.91		(1970MO13)
6.13	d	n, α_0	10.99		(1970MO13)
6.30	c	n, α_0	11.13		(1970MO13)
6.64	b	n, α_0	11.39	(2 ⁺)	(1970MO13)
6.67	b	n, α_0	11.41	(4 ⁺)	(1970MO13)
6.93	b	n, α_0	11.62	5 ⁻	(1970MO13)
7.03	b	n, α_0	11.69	6 ⁺	(1970MO13)
7.19	d	n, α_0	11.82	(3 ⁻)	(1970MO13)
7.47	d	n, α_0	12.04	(2 ⁺)	(1970MO13)
7.75	e	n, α_0	12.25	(0 ⁺ , 1 ⁻)	(1970MO13)
7.85	b	n, α_0	12.33	5 ⁻	(1970MO13)
8.06	b	n, α_0	12.49	4 ⁺	(1970MO13)
8.10	b	n, α_0	12.53	6 ⁺	(1970MO13)

^a For the first four states, see also Table 18.2.

^b Γ_α , large; Γ_n , large.

^c Γ_α , small; Γ_n , small.

^d Γ_α , small; Γ_n , large.

^e Γ_α , large; Γ_n , small.

^f $\pm 10 - 20$ keV for this and all higher resonances (G.E. Mitchell, private communication).

^g Two states with $E_x = 9.0$ to 9.2 MeV and $J^\pi = (2^+, 3^-)$ or $(4^+, 3^-)$ are reported by (1958WE29).

$$15. \text{}^{17}\text{O}(\text{d}, \text{p})\text{}^{18}\text{O} \quad Q_m = 5.822$$
$$Q_0 = 5.820 \pm 0.010 \text{ (1965MO16)}.$$

Proton groups have been observed to most of the ^{18}O states with $E_x < 8$ MeV (1963YA03, 1965MO16, 1966WI07): see Table 18.7. See also (1959AJ76) and (1964HE11, 1965WI1B). Angular distributions of the protons to the relatively strongly populated states have been measured at $E_d = 5.55$ to 14.95 MeV and analyzed by PWBA (1957BI80, 1964MO25, 1965MO16, 1966WI07). Proton- γ coincidence measurements by (1967MO09) are displayed in Table 18.2 and a lifetime measurement for $^{18}\text{O}^*(1.98)$ by (1969NI09) is shown in Table 18.3.

See also (1963EL04, 1964BA1G, 1966AR10, 1968SU1C; theor.).

$$16. \text{}^{17}\text{O}(\text{}^{17}\text{O}, \text{}^{16}\text{O})\text{}^{18}\text{O} \quad Q_m = 3.904$$

See (1967DA1E).

$$17. \text{}^{18}\text{N}(\beta^-)\text{}^{18}\text{O} \quad Q_m = 14.06$$

The decay is to $^{18}\text{O}^*(4.45)$: γ -rays with $E_\gamma = 0.82 \pm 0.02, 1.65 \pm 0.02, 1.98 \pm 0.02$ and 2.47 ± 0.03 MeV with intensities $0.60 \pm 0.03, 0.63 \pm 0.04, 1.00 \pm 0.04$ and 0.43 ± 0.03 have been observed. These are due to the decay of $^{18}\text{O}^*(4.45)$ to $^{18}\text{O}^*(3.63, 1.98)$ and the subsequent decay of these two states (1964CH19).

$$18. \text{ (a) } \text{}^{18}\text{O}(\gamma, \text{n})\text{}^{17}\text{O} \quad Q_m = -8.047$$
$$\text{ (b) } \text{}^{18}\text{O}(\gamma, \text{p})\text{}^{17}\text{N} \quad Q_m = -15.943$$
$$\text{ (c) } \text{}^{18}\text{O}(\gamma, \alpha)\text{}^{14}\text{C} \quad Q_m = -6.227$$

For reaction (a) see (1963FU06, 1964MU1A). For reaction (b) see (1955AJ61, 1964KO09, 1969HO16). See also (1965DO1E). For reaction (c), see (1964GR08).

Table 18.5: States in ^{18}O from $^{16}\text{O}(t, p)^{18}\text{O}$

E_x (MeV \pm keV)			L^b	J^π
(1960JA17)	(1960JA13)	(1962HI06)		
0	0	0	0	0^+
1.979 ± 5	^a	1.980 ± 5	2	2^+
3.552 ± 5	3.560 ± 15	3.549 ± 5	3 or 4	3^- or 4^+
3.634 ± 5	3.639 ± 15	3.627 ± 5	0	0^+
3.915 ± 5	3.925 ± 15	3.915 ± 5	2	2^+
4.448 ± 5	4.457 ± 15	4.449 ± 5	(3)	^c
	5.084 ± 18	5.090 ± 5	3	3^-
		5.247 ± 7	2	2^+
		5.329 ± 7	0	0^+
		5.368 ± 10	(2)	$(2^+)^d$
		5.521 ± 10		
		6.189 ± 10		
		6.341 ± 10		
		6.391 ± 10		

^a Observed but not measured.

^b From PWBA analysis of angular distributions: see (1964MI05). See also (1960JA17).

^c $J^\pi = 1^-$. See, e.g. discussion in (1964MI05).

^d See, however, reaction 15.

19. $^{18}\text{O}(e, e)^{18}\text{O}$

The ^{18}O charge radius, $r_{\text{rms}} = 2.727 \pm 0.020$ fm (using a distorted wave approximation), 2.766 ± 0.020 fm (using a Born approximation) (1970SI02). See also (1967DA1D, 1970SI1M).

At $E_e = 69$ MeV, the 180° scattering spectrum shows, in addition to the elastic peak, some weak structure corresponding to $E_x = 2.0, 4.45$ and 5.6 MeV. Somewhat stronger inelastic peaks are found corresponding to $E_x = 10.9, 12.6, 14.0$ and 15.1 MeV. In addition six prominent peaks, each of about the same strength and assumed to arise from giant dipole excitations, are observed at $E_x = 16.6, 18.4, 20.2, 21.9, 23.7, 25.2$ and 26.9 MeV [all E_x values, ± 0.2 MeV] (1965VA1H). See also (1961LA09, 1970CA1P).

See also (1962BA1D) and (1963BI05, 1963WI1B, 1966GR1H, 1967GR1B; theor.).

20. $^{18}\text{O}(n, n)^{18}\text{O}$

Table 18.6: $^{16}\text{O}(t, p)^{18}\text{O}$ angular distribution studies

E_t (MeV)	Distribution of proton groups	Refs.
0.67 – 1.40	p	(1963KU07)
0.9 – 1.7	p_0, p_1	(1967KO1G)
1.15 – 1.95	p_0, p_1	(1959JO32)
5	p_1	(1963RE06)
5.5	$p_0 \rightarrow p_5$	(1960JA17)
5.5	p_0	(1960JA11)
5.55	$p_0 \rightarrow p_5$	(1965MO19)
10.0	$p_0 \rightarrow p_9$	(1964MI05)
10.15	p_0, p_1	(1962PU01)

Angular distribution measurements are listed in Table 18.8 (1964EX1A, 1969ME15).

21. $^{18}\text{O}(p, p)^{18}\text{O}$

A proton group is observed to the first excited state: $E_x = 1.981 \pm 0.004$ MeV (1957YO04). Angular distribution measurements reported by (1961CA02, 1965PR1D, 1971LE1R, 1971RE20) are displayed in Table 18.8. The transition from $^{18}\text{O}^*(3.63)$ to the ground state has been determined to be EO: $J^\pi = 0^+$ for the 3.63 MeV state (1963GO32, 1965GO11, 1967CO30). Proton- γ angular correlation measurements establish $J = 2$ and 1 for $^{18}\text{O}^*(3.92, 4.45)$ (1965OL02), and $J = 3, 2, 0, 3$ and (2) for $^{18}\text{O}^*(5.09, 5.25, 5.33, 5.37, 5.52)$, respectively (1966LO12): see Table 18.2 for branching ratios obtained by (1964GO11, 1965OL02, 1966LO12). Angular distributions to $^{18}\text{O}^*(0, 1.98, 7.11)$ [$J^\pi = 0^+, 2^+, 4^+$, respectively] using polarized protons with $E_p = 22.5$ and 24.5 MeV have been analyzed with the rotational model: $\beta_2 = 0.37 \pm 0.03$ and $\beta_4 = 0.18 \pm 0.04$. The results confirm the assignment of $^{18}\text{O}^*(7.11)$ to the ground state rotational band (1971RE20).

See also (1964SC01, 1968BE34) and (1966RE1C, 1967LO1F, 1967SC16, 1970AG1C; theor.).

22. $^{18}\text{O}(d, d)^{18}\text{O}$

At $E_d = 15$ MeV, inelastically scattered deuteron groups are reported to ^{18}O states with $E_x = 1.982, 3.564 \pm 0.025, 3.655 \pm 0.030, 3.949 \pm 0.030, 4.501 \pm 0.040, 5.139 \pm 0.030, 5.301 \pm 0.030, 5.413 \pm 0.035$ MeV (1961AR06). Angular distribution measurements by (1961AR06, 1964WI05, 1965DI1C, 1965LU1A, 1966DE09) are listed in Table 18.8. See also (1959AJ76).

Table 18.7: States of ^{18}O from $^{17}\text{O}(\text{d}, \text{p})^{18}\text{O}$

E_x (MeV \pm keV)			l_n ^f	J^π ^g
(1963YA03) ^a	(1965MO16) ^d	(1966WI07) ^e		
	0	0	2	0 ⁺
	1.982 \pm 10	1.98	0 + 2	2 ⁺
	3.552 \pm 10	3.55	2	4 ⁺
		3.63	(2)	0 ⁺
		3.92	0	2 ⁺
4.440 \pm 25 ^b		4.45	1	1 ⁻
5.092 \pm 25		5.09		
^c	5.255 \pm 10	5.25	0	2 ⁺ ^h
^c		5.33		
(5.375 \pm 25)	5.375 \pm 10	5.37	0	3 ⁺ ^h
5.492 \pm 25				
6.200 \pm 25		6.19		
^c		6.34		
	7.110 \pm 15	7.10	2	4 ⁺
	7.855 \pm 20			
	7.962 \pm 20			

^a $E_d = 14.95$ MeV. Angular distribution measured for protons to $^{18}\text{O}^*(4.45)$.

^b Lower energy states were not investigated.

^c This part of the spectrum could not be observed.

^d $E_d = 5.56$ MeV. Angular distributions measured for protons to the first six states shown.

^e $E_d = 10.0$ MeV. The energies shown are nominal. Proton groups to $^{18}\text{O}^*(5.52, 6.39, 6.86)$ were not observed. Angular distributions were measured for all the listed states.

^f (1965MO16, 1966WI07), except for the assignment for $^{18}\text{O}^*(4.45)$ (1963YA03). The levels for which l -values are not given by (1965MO16, 1966WI07) are weakly populated.

^g Known J^π from this and other experiments.

^h See also (1964MO25, 1967MO09).

Table 18.8: $^{18}\text{O}(n, n)$, (p, p) , (d, d) , (t, t) , $(^3\text{He}, ^3\text{He})$, (α, α) angular distribution studies

E (MeV)	Angular distribution of groups	Refs.
2.9 – 4.2	n_{0+1}	(1964EX1A)
14.1	$n_0, n_1, n_{2 \rightarrow 4}, n_{6 \rightarrow 10}$	(1969ME15)
0.84 – 2.00	p_0	(1961CA02)
2.64 – 2.93	$\gamma_{1.98}$	(1965PR1D)
22.5, 24.5	p_0, p_1, p to $^{18}\text{O}^*(7.11)$	(1971RE20)
65	p_0	(1971LE1R)
7.0	d_0	(1964WI05)
12.3	d_0	(1965LU1A)
15	d_0	(1965DI1C, 1966DE09)
15	d_1	(1961AR06)
6.4, 7.2	t_0	(1964PU01)
11.0	$^3\text{He}(0)$	(1970BO25, 1970GR04)
15	$^3\text{He}(0)$	(1969ZU02)
17.3	$^3\text{He}(0)$	(1968HA30, 1969HA1U)
21.4	α to $^{18}\text{O}^*(0^a, 1.98, 3.55 + 3.63, 3.92, 4.45, 5.1, 6.19, 7.1)$	(1966LU05)
40.5	α to $^{18}\text{O}^*(0, 1.98, 3.92, 4.45, 5.09, 6.34 + 6.39, 7.96 + 8.04 + 8.08)$	(1966HA19)

^a See also (1966WE04).

23. $^{18}\text{O}(t, t)^{18}\text{O}$

The angular distributions of elastically scattered tritons have been measured at $E_t = 6.4$ and 7.2 MeV (1964PU01). See also (1968HO1C).

24. $^{18}\text{O}(^3\text{He}, ^3\text{He})^{18}\text{O}$

Angular distributions of elastically scattered ^3He ions have been studied at $E(^3\text{He}) = 11.0$ to 17.3 MeV: see Table 18.8 (1968HA30, 1969HA1U, 1969ZU02, 1970BO25, 1970GR04). See also (1968HO1C) and (1969MA1G; theor.).

25. $^{18}\text{O}(\alpha, \alpha)^{18}\text{O}$

Angular distributions of elastically scattered α -particles have been measured at $E_\alpha = 21.4$ (1966LU05) and 40.5 MeV (1966HA19): see Table 18.8. The transitions to $^{18}\text{O}^*(4.45, 5.09)$ are $L = 1$ and 3, respectively, fixing $J^\pi = 1^-$ and 3^- for these states. $B(E2)$ values for $^{18}\text{O}^*(1.98, 4.45)$ and $B(E3)$ for $^{18}\text{O}^*(5.09)$ are also reported (1966HA19).

Measurements of α groups near 180° for $E_\alpha = 20.0$ to 23.4 MeV confirm assignments of natural parity for $^{18}\text{O}^*(1.98, 3.55, 3.63, 3.92, 4.45, 5.09, 5.25, 5.33, 6.19, 6.39, 7.11, 7.62, 7.85, 8.21, 8.28, 8.82, 8.96, 9.03, 9.10, 9.36, 9.39, 9.68, 9.72 \pm 0.03, 9.88, 10.12, 10.29, 10.38, 11.62, 11.69)$. The levels at $E_x = 5.37, 8.48$ and 8.64 MeV were not observed, and those at 5.52, 6.34 and 6.86 MeV were populated weakly indicating unnatural parity: $J^\pi = 3^+$ and 2^- , respectively for $^{18}\text{O}^*(5.37, 5.52)$ (1971OL06).

Alpha- γ correlation measurements involving ^{18}O states below $E_x = 6.4$ MeV [see Table 18.2] lead to $J^\pi = 1^-$ and 3^- for $^{18}\text{O}^*(6.19, 6.39)$. Other J^π values agree with previous assignments. The transitions $3.92 \rightarrow 1.98$ and $5.25 \rightarrow 1.98$ are almost pure M1 (1971BE45). See also (1968FA1A).

26. (a) $^{18}\text{O}(^7\text{Li}, ^7\text{Li})^{18}\text{O}$
 (b) $^{18}\text{O}(^9\text{Be}, ^9\text{Be})^{18}\text{O}$
 (c) $^{18}\text{O}(^{10}\text{B}, ^{10}\text{B})^{18}\text{O}$

For reaction (a) see (1969NE1E). For reactions (b) and (c) see (1971KN05).

27. (a) $^{18}\text{O}(^{12}\text{C}, ^{12}\text{C})^{18}\text{O}$
 (b) $^{18}\text{O}(^{13}\text{C}, ^{13}\text{C})^{18}\text{O}$

For reaction (a) see (1967GO1A, 1968GO1H). For reaction (b) see (1971KN05).

28. $^{18}\text{O}(^{16}\text{O}, ^{16}\text{O})^{18}\text{O}$

See (1968GO1H, 1970FO1F, 1970SI09). See also (1965GO1G).

29. $^{18}\text{O}(^{18}\text{O}, ^{18}\text{O})^{18}\text{O}$

See (1970MO35, 1970SH07, 1971GO1T, 1971RA1C, 1971VA1H).

30. $^{18}\text{F}(\beta^+)^{18}\text{O}$ $Q_m = 1.655$

Table 18.9: ^{18}O states from $^{19}\text{F}(d, ^3\text{He})^{18}\text{O}$ ^a

E_x (MeV \pm keV)	l	C^2S ^d
0	0 ^c	1.00
1.97 \pm 20	2 ^c	1.39
3.63 \pm 100 ^b	0 + 2 + 4	\approx 0.2
4.45 \pm 10	1 ^c	1.31
5.28 \pm 30 ^b	0 + 2	\approx 1.2
6.27 \pm 10 ^b	1	0.70
6.90 \pm 20	1	1.03
7.67 \pm 30 ^b	1	0.42
9.76 \pm 150 ^b		
11.14 \pm 70	1	0.65
11.75 \pm 70 ^b	1	0.72
12.25 \pm 70	1	0.89
14.10 \pm 200		
14.56 \pm 100		

^a (1969KA1A, 1970KA31): $E_d = 51.7$ MeV; DWBA analysis.

^b Corresponds to unresolved states.

^c See also (1965ZE04).

^d Normalized to 1.00 for the ground-state transition.

See ^{18}F .

31. (a) $^{19}\text{F}(\gamma, p)^{18}\text{O}$ $Q_m = -7.993$

(b) $^{19}\text{F}(e, ep)^{18}\text{O}$ $Q_m = -7.993$

See (1962DO1A). See also (1968AB09).

32. $^{19}\text{F}(n, d)^{18}\text{O}$ $Q_m = -5.768$

Angular distributions are reported at $E_n = 14$ MeV (1965SA14; d_0), 14.1 MeV [(1957RI1A; d_0, d_1), (1959VE19, 1960VE06; d_0) and (1968FA01; d_0, d_1)] and at 14.4 MeV (1968AN1F, 1968RE07; d_0, d_1). See also (1961BO1A, 1961KO06, 1961LE1D, 1969BE1R, 1971MI1H).

$$33. \text{}^{19}\text{F}(\text{p}, 2\text{p})^{18}\text{O} \quad Q_{\text{m}} = -7.993$$

See (1971DE1F) and ^{19}F .

$$34. \text{}^{19}\text{F}(\text{d}, \text{}^3\text{He})^{18}\text{O} \quad Q_{\text{m}} = -2.499$$

Many states of ^{18}O have been populated in this reaction: see Table 18.9 (1965ZE04, 1969KA1A, 1970KA31). See also (1967WI1F) and (1968BA2J; theor.).

$$35. \text{}^{19}\text{F}(\text{t}, \alpha)^{18}\text{O} \quad Q_{\text{m}} = 11.821$$

Alpha groups are reported at $E_{\text{t}} = 5.75$ to the thirteen excited states reported in the $^{16}\text{O}(\text{t}, \text{p})^{18}\text{O}$ reaction [see Table 18.5] and to states with $E_{\text{x}} = 6.86, 7.10, 7.60, 7.75, 7.84, 7.96, 8.02, 8.11, 8.19, 8.26, 8.39, 8.48, 8.64$ MeV (± 20 keV) (1962HI06). Branching ratios are reported by (1967CH1D): see Table 18.2. See also (1966BE1G, 1967CH1L, 1968BE1U, 1968CH1K). See also (1967CA1L).

$$36. \text{}^{21}\text{Ne}(\text{n}, \alpha)^{18}\text{O} \quad Q_{\text{m}} = 0.698$$

See (1961AB05).

^{18}F
(Figs. 2 and 4)

GENERAL: (See also (1959AJ76).)

Shell model: (1957WI1E, 1959BR1E, 1960TA1C, 1961TR1B, 1962TA1D, 1964FE02, 1964IN03, 1964PA1D, 1964YO1B, 1965BA1J, 1965DE1H, 1965GI1B, 1966BA2E, 1966BA2C, 1966HU09, 1966IN01, 1966KU05, 1966RI1F, 1967EN01, 1967EV1C, 1967FE01, 1967FL01, 1967HO11, 1967IN03, 1967KU09, 1967KU13, 1967LY02, 1967MO1J, 1967PA1K, 1967PI1B, 1967VI1B, 1967WO1C, 1968AR02, 1968BE1T, 1968BH1B, 1968CL03, 1968CO11, 1968DE1M, 1968DE13, 1968EL1C, 1968FR03, 1968GA16, 1968GU1E, 1968HA17, 1968HA1P, 1968HE1H, 1968KA1C, 1968KU1E, 1968NA09, 1968ZU02, 1969BA2F, 1969BE1T, 1969BU15, 1969KA09, 1969KU1G, 1969MA1T, 1969ME06, 1969ST1G, 1969ZU1B, 1969ZU03, 1970BA2E, 1970EL1G, 1970EL08, 1970HA49, 1970KA32, 1970TR08, 1970WA1T, 1971AR1R, 1971EL07, 1971GU21, 1971HO12, 1971JA06, 1971LO23, 1971PE1A, 1971PR16, 1971QU01, 1971WI01, 1972KA01, 1972LE1L).

Cluster, collective and deformed models: (1964KE1A, 1964MA1G, 1966PI1B, 1966RI1F, 1967FE01, 1967PA1K, 1968EL1C, 1969BA2E, 1969ZA1D, 1970BA2B, 1970WA1T, 1971AR1R, 1972LE1L).

Astrophysical questions: (1970BA1M).

Electromagnetic transitions: (1962MO1A, 1963BU11, 1966HA31, 1967IN03, 1968EL1C, 1968HA17, 1968HE1G, 1968LE02, 1969BE1T, 1969HA1F, 1969WA1C, 1970EL08, 1970HA49, 1970WA31, 1970WA1T).

Special levels: (1964KE1A, 1965GI1B, 1966HU09, 1966MI1G, 1966NA1B, 1967EN01, 1967EV1C, 1967FL01, 1967PA1K, 1967VI1B, 1968AR02, 1968BH1B, 1968CO11, 1968FR03, 1968GA16, 1968HA1P, 1968HE1G, 1968KA1C, 1968LE02, 1968NA09, 1968NO1C, 1969BA1Z, 1969BE1T, 1969CH1K, 1969EL1A, 1969FE1A, 1969HA1F, 1969KA09, 1969KA29, 1969MA1T, 1969ME06, 1969OS01, 1970EL1G, 1970EL08, 1971AR1R, 1971EL07, 1971GU21, 1971JA06, 1971PE1A, 1971PR16, 1971QU01, 1972KA01).

Other theoretical topics: (1961MA1B, 1962IN1A, 1963BU11, 1963VL1A, 1964IN03, 1964ST1B, 1965DE1H, 1965FO1E, 1965KA1B, 1966GI1A, 1966HE1E, 1966KU05, 1967FE01, 1967FL01, 1967KU13, 1967MA1B, 1967MO1J, 1967WO1C, 1968BA2H, 1968BE1T, 1968DE1M, 1968EL1C, 1968GU1E, 1968JO1C, 1968KA1C, 1968KU1E, 1968LE02, 1968MU1B, 1968VA1J, 1968VA24, 1969DE16, 1969GA1Q, 1969KA29, 1969MU09, 1969OS01, 1969PE1D, 1969RA28, 1969WA1F, 1970BA1Z, 1970EL08, 1970PR1D, 1971BA14, 1971BO21, 1971EL07, 1971LO23, 1971PR16, 1972DE05, 1972KA01, 1972LE1L).

Reactions involving pions: (1968BE1F).

Complex reactions involving ^{18}F : (1965LO1D, 1966LA1G, 1967AU1B, 1967WI16, 1967WI20, 1968MO1C, 1968SH1H, 1968ST1L, 1969FE04, 1969KR21, 1970BR13, 1970FA01, 1970FE05, 1970KO1P, 1970KR12, 1970KR1C, 1971AR02, 1971MO1H).

Table 18.10: Energy levels of ^{18}F ^a

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$1^+; 0$	0^+	$\tau_{1/2} = 109.77 \pm 0.05$ min	β^+	1, 2, 3, 4, 5, 9, 10, 11, 12, 17, 18, 19, 20, 21, 22, 23, 27, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42
0.9370 ± 0.2	$3^+; 0$	0^+	$\tau_m = 68 \pm 7$ psec	γ	5, 9, 10, 17, 18, 19, 23, 27, 28, 30, 31, 35, 36, 37, 42
1.0419 ± 0.8	$0^+; 1$		4_{-2}^{+3} fsec	γ	5, 9, 18, 28, 30, 31, 32, 35, 37, 41
1.0809 ± 0.8	$0^-; 0$		30 ± 3 psec	γ	5, 9, 10, 17, 18, 30, 31, 35, 36, 37
1.1218 ± 0.7	$5^+; 0$	0^+	218 ± 9 nsec	γ	5, 9, 10, 17, 18, 19, 23, 28, 30, 31, 35, 37, 42
1.7007 ± 0.8	$1^+; 0$	1^+	0.86 ± 0.20 psec	γ	5, 9, 10, 17, 18, 19, 23, 30, 31, 32, 35, 37, 41, 42
2.1013 ± 0.5	$2^-; 0$		4.3 ± 1.4 psec	γ	9, 10, 17, 18, 19, 23, 31, 35, 37, 42
2.5240 ± 0.8	$2^+; 0$	1^+	0.65 ± 0.13 psec	γ	5, 10, 17, 18, 19, 23, 27, 28, 35, 37, 42
3.0598 ± 2.6	$2^+; 1$		≤ 1.2 fsec	γ	5, 18, 20, 23, 27, 28, 31, 35, 37, 41, 42
3.1349 ± 1.6	$1^-; 0$		$0.37_{-0.10}^{+0.15}$ psec	γ	5, 10, 18, 19, 23, 31, 35, 37, 42
3.3569 ± 2.6	$3^+; 0$	1^+	0.48 ± 0.09 psec	γ	5, 10, 18, 19, 23, 27, 31, 35, 37, 42
3.7242 ± 2.7	$1^+; 0$		< 0.08 psec	γ	10, 18, 23, 31, 37, 42
3.787 ± 7	$(3^-); 0$			γ	10, 18, 23, 31, 35, 37
3.836 ± 3	$2^+; 0$		< 0.073 psec	γ	10, 18, 19, 23, 27, 28, 31, 35, 37, 42
4.119 ± 4	$3^+; 0$			γ	10, 18, 19, 23, 28, 31, 35, 37, 42
4.229 ± 4	(2)			γ	10, 18, 23, 31, 35, 37, 42
4.361 ± 3	2, 3		< 0.61 psec	γ	10, 18, 20, 31, 35, 37, 42
4.402 ± 5	$\geq 2; 0$			γ	10, 18, 19, 31, 35, 37, 42
4.6503 ± 1.0	$4^+; 1$			γ	5, 10, 18, 23, 28, 31, 35, 37
4.739 ± 4	$0^+; 1$			γ	18, 31, 37
4.849 ± 4	$1; 0$			γ	5, 10, 18, 19, 37, 42
4.957 ± 5	$2^+; 1$			γ	18, 23, 27, 28, 37
5.301 ± 4	$4^+; 0$	1^+	31 ± 6 fsec	γ, α	5, 10, 18, 19, 23, 37, 42
5.501 ± 5				γ, α	5, 10, 37, 42
5.599 ± 11	$(4^+; 0)$				18, 19, 35, 37
5.606 ± 2	$1^-; 0, 1$		$\Gamma < 1.2$ keV	γ, α	5, 8, 10, 27, 35, 37, 42
5.674 ± 2	$1^-; 0, 1$		< 0.8	γ, α	5, 8, 10, 27, 37

Table 18.10: Energy levels of ^{18}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
5.785 \pm 10	($T = 0$)				19, 37, 42
6.095 \pm 2	$4^-; 0$		< 2.0	γ, p, α	8, 10, 18, 19, 23, 26, 37, 42
6.135 \pm 2	$0^+; 1$			γ	23, 37, 42
6.161 \pm 2	$3^+; 1$		15	γ	23, 25
6.240 \pm 2	(3^-) ^b		< 0.8	γ, p, α	5, 8, 23, 25, 26, 37, 41, 42
6.261 \pm 3	(1)		< 3	α	8, 10, 18, 37, 41, 42
6.280 \pm 2	$2^+; 1$		7.5	γ, p	23, 25
6.309 \pm 2	$2^+, 3^+$		3.1 ± 1.4	γ, p, α	23, 25, 26
6.385 \pm 3	(1^+), $2, 3^-$		< 4.5	γ, p, α	23, 26, 37
6.483 \pm 3	(1^+), $2, 3^-$		< 1.2	γ, p, α	23, 26, 37
6.565 \pm 4	$5^+; 0$	1^+	(< 0.8)	γ, α	5, 8, 10, 19, 23, 37, 42
6.646 \pm 4	1^-		89 ± 4	p, α	8, 10, 26, 37, 42
6.6474 \pm 2.5	(2^-) ^b		< 1.6	γ, p, α	5, 7, 10, 23, 25, 26, 37, 42
6.780 \pm 2	$4^+, 5^+; 0$		10 ± 3	γ, p, α	23, 25, 26, 37, 42
6.808 \pm 3	$2^+, 3^+$		5 ± 3	γ, p, α	10, 19, 23, 25, 26, 37, 42
6.808 \pm 5	$2^-; 0$		90 ± 10	p, α	7, 8, 18, 26
6.871 \pm 2				γ, p, α	23, 26, 37, 42
7.194 \pm 3	(4^+); 0		< 4	α	8, 10, 19, 37
7.206 \pm 9	(1^+)		45 ± 10	(p), α	8, 10, 18, 26
7.313 \pm 10	(3^-)		53 ± 6	p, α	7, 8, 10, 37
7.395 \pm 10	($3^-; 0$)		35	p, α	7, 8, 19, 26, 37
7.57			60	p, α	7, 8, 26
7.653 \pm 9	$T = 0$		40	p, α	7, 8, 18, 19, 26
7.74			120	p, α	7, 8, 26
7.872 \pm 10	(2^-); 0		30	p, α	7, 8, 18, 19, 26
7.95	(1^+)		70	p, α	7, 8, 26
(8.21)			≈ 15	p, α	26
(8.23)			≈ 10	p, α	26
(8.37)			≈ 50	p, α	26
8.46				p, d, α	8, 14, 15, 16
8.596 \pm 19	$T = 0$				19
8.861 \pm 190	$T = 0$				19
9.145 \pm 32	$3, 4^-; 0$		108 ± 12	p, d, α	14, 15, 16, 18
9.26	3^- ^b		≈ 30	d, α	16
9.32	2^+ ^b		≈ 40	(p), d, α	14, 16, 18
9.494 \pm 15	(6^-); 0				19
9.50	$2, 3^+; 0$			n, d, α	13, 16, 18
(9.55)	$2, 3^+; 0$			d, α	16

Table 18.10: Energy levels of ^{18}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
(9.70)	$3^- b$		370	n, d, α	13, 16
9.82 \pm 40	$2^+ b$		200	d, α	16, 18, 19
10.06 \pm 45				n, p, d	13, 14, 18, 19
(10.21)	$1^- b$		90	d, α	16
(10.24)	$3, 4^-$			d, α	16
10.268 \pm 12	$3^- b$		180	n, p, d, α	13, 14, 16, 19
(10.31)	$2^+ b$		570	d, α	16
10.352 \pm 25	$1^- b$		270	(n), d, α	13, 16, 18
10.42	$3^- b$		50	(n), d, α	13, 16
10.541 \pm 10	$2^+ b$		50	n, p, d, α	13, 14, 16, 19
10.59	$1^- b$		160	d, α	16
(10.61)	$4, 5^+; 0$			d, α	16
10.61	$2^+ b$		150	d, α	16
(10.63)	$1^- b$		60	d, α	16
10.64	$3^- b$		240	d, α	16
(10.68)	$2^+ b$		60	d, α	16
10.69	$3^- b$		75	d, α	16
(10.71)	$2^+ b$		375	d, α	16
10.73 \pm 30	$(2^-); 0$			(n, p), α	6, 7
10.80	$2^+ b$		85	n, p, d, α	13, 14, 16
(10.90)	$(2, 3)^+; 0$			p, d, α	14, 16
10.91	$1^- b$		240	d, α	16
(10.95)	$2^+ b$		400	d, α	16, 18
10.98	$3^- b$		120	d, α	16, 18
(10.99)	$2^+ b$		220	d, α	16
(11.03)	$2^+ b$		80	d, α	16
(11.03)	$3, 4^-; 0$		≈ 35	p, d, α	14, 16
11.073 \pm 37	2^+		50	d, α	16, 19
(11.11)	$1^- b$		1000	d, α	16
11.13 \pm 50	$3^- b$		35	n, p, d, α	13, 14, 16, 18
11.26	$3^- b$		240	d, α	16
(11.30)	$2^+ b$		475	d, α	16
11.31	$4^+ b$		40	d, α	16
(11.32)	$2^+ b$		95	d, α	16
11.32	$3^- b$		65	d, α	16
11.384 \pm 18	$4, 5^+; 0$			p, d, α	14, 16, 19
(11.51)	$2^+ b$		125	d, α	16
11.56	$3^- b$		70	d, α	16

Table 18.10: Energy levels of ^{18}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
(11.59)	$5, 6^-; 0$			d, α	16
(11.66)	2^+ ^b		240	d, α	16
(11.79)	$\geq 3; 0$			$(n, p), d, \alpha$	6, 7, 15, 16
11.83	1^- ^b		180	$(n, p), d, \alpha$	6, 7, 16
(11.91)	$5, 6^-; 0$			d, α	16
11.96	3^- ^b		30	d, α	16
(11.97)	2^+ ^b		125	d, α	16
12.01			40	d, α	16
12.055 ± 16	$T = 0$		60	d, α	16, 19
12.12	$T = 0$		55	d, α	16
12.12	3^- ^b		35	d, α	16
(12.14)	2^+ ^b		220	d, α	16
12.25			70	d, α	16
12.25 ± 30	$(2^-, 3); 0$		170	$(n, p), d, \alpha$	6, 7, 16
12.32			130	d, α	16
(12.31)	1^- ^b		100	d, α	16
12.34	3^- ^b		190	d, α	16
12.36	2^+ ^b		80	d, α	16
12.38	2^+ ^b		120	d, α	16
(12.47)	1^- ^b		35	d, α	16
(12.48)	(3^-)		120	d, α	16
12.484 ± 15	4^+ ^b		60	d, α	8, 16
12.526 ± 15	3^- ^b		80	d, α	8, 16
12.565 ± 15	2^+ ^b		≈ 300	$(n, p), d, \alpha$	6, 7, 8, 16
(12.63)	2^+		210	d, α	16
(12.67)	2^+ ^b		290	d, α	16
12.68			160	d, α	16
12.683 ± 15	5^- ^b		45	n, p, d, α	6, 7, 8, 16, 19
12.702 ± 15	3^- ^b		70	d, α	8, 16
(12.78)			160	d, α	16
(12.81)	2^+ ^b		440	d, α	16
(12.86)			120	d, α	16
(12.90)	3^- ^b		120	d, α	16
(12.96)			120	d, α	16
(12.99)	2^+ ^b		650	d, α	16
(13.03)	3^- ^b		250	d, α	16
13.06 ± 15	4^+		40	d, α	8, 16
(13.08)	$T = 0$		150	d, α	16

Table 18.10: Energy levels of ^{18}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
(13.12)			160	d, α	16
13.129 \pm 15	(4 ⁺)		35	d, α	8, 16
(13.16)			160	d, α	16
(13.17)	1 ⁻ ^b		185	d, α	16
13.198 \pm 15	5 ⁻ ^b		140	d, α	8, 16
(13.20)	$T = 0$		145	d, α	16
(13.20)	4 ⁺ ^b		255	d, α	16
(13.24)			120	d, α	15, 16
(13.26)			100	d, α	16
(13.26)	4 ⁺ ^b		360	d, α	16
(13.29)	2 ⁺ ^b		100	d, α	16
13.332 \pm 15	5 ⁻ ^b		50	d, α	8, 16
(13.34)			135	d, α	16
(13.40)			120	d, α	16
13.398 \pm 15	6 ⁺		85	α	8
(13.43)			200	d, α	16
(13.43)	3 ⁻ ^b		235	d, α	16
(13.44)	1 ⁻ ^b		100	d, α	16
13.464 \pm 15	4 ⁺		100	d, α	8, 16
13.50	5 ⁻ ^b		65	d, α	16
(13.50)	2 ⁺ ^b		145	d, α	16
(13.57)			140	d, α	16
13.597 \pm 15	6 ⁺		100	α	8
(13.60)			150	d, α	16
(13.60)	4 ⁺ ^b		40	d, α	16
(13.61)	1 ⁻ ^b		90	d, α	16
13.65	5 ⁻ ^b		300	d, α	16
13.652 \pm 15	3 ⁻		60	α	8
13.671 \pm 15	(2 ⁺)		60	α	8
13.679 \pm 15	4 ⁺ ^b		70	d, α	8, 16
(13.70)	2 ⁺ ^b		80	d, α	16
(13.77)	3 ⁻ ^b		205	d, α	16
13.780 \pm 15	4 ⁺ ^b		130	d, α	8, 16
(13.81)	2 ⁺ ^b		265	d, α	16
(13.85)			60	d, α	16
(13.92)			210	d, α	16
13.917 \pm 15	(4 ⁺ , 2 ⁺)			α	8
(13.96)	1 ⁻ ^b		70	d, α	16

Table 18.10: Energy levels of ^{18}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
(13.99)			150	d, α	16
13.980 \pm 15	5 ⁻ ^b		170	d, α	8, 16
(14.01)			110	d, α	16
(14.05)			200	d, α	16
(14.08)			210	d, α	16
(14.10)	2 ⁺ ^b		170	d, α	16
(14.10)	4 ⁺ ^b		75	d, α	16
14.162 \pm 5	5 ⁻ ^b		200	d, α	8, 16
(14.18)			220	d, α	16
(14.26)			200	d, α	16
(14.30)			85	d, α	16
(14.33)	$T = 0$		130	d, α	16
14.323 \pm 15	4 ⁺		95	α	8
(14.34)	$T = 0$		250	d, α	16
d					
16.513 \pm 15 ^c	6 ⁺ (7 ⁻)		\approx 50	α	8
16.606 \pm 15 ^c	5 ⁻		\approx 50	α	8
16.723 \pm 15 ^c	6 ⁺		\approx 150	α	8
16.870 \pm 15 ^c	7 ⁻		\approx 200	α	8
16.948 \pm 15 ^c	6 ⁺		\approx 150	α	8

^a See also Tables 18.14, 18.15 and 18.17.

^b Isospin mixed state: see (1970JO1C) and Table 18.20.

^c State not reported in $^{16}\text{O}(\text{d}, \alpha)^{14}\text{N}$: see Table 18.17.

^d [92 additional states with $14.36 < E_x < 20.80$ MeV are reported in $^{16}\text{O}(\text{d}, \alpha)^{14}\text{N}$: see Table 18.17].

Ground state: (1970WA1T, 1971GU21, 1971WI01, 1972LE1L).

$$1. \ ^{18}\text{F}(\beta^+)^{18}\text{O} \quad Q_m = 1.655$$

The positron decay is entirely to the ground state if ^{18}O [$J^\pi = 0^+; T = 1$]: $E_\beta(\text{max}) = 635 \pm 2$ keV (1964HO28). See also (1959AJ76). The half-life is 109.77 ± 0.05 min: see Table 18.11. $\log ft = 3.554$: see also (1969KA1B). The fact that the β^+ transition to $^{18}\text{O}(0)$ is allowed indicates $J^\pi = 1^+$ for $^{18}\text{F}(0)$. The ratio $\epsilon_K/\beta^+ = 0.030 \pm 0.002$ (1956DR38). See also (1959PE1B, 1966KU1F, 1968DA1J, 1968LE02, 1970MC23, 1971BL12, 1971DE1E, 1971VA1C, 1971WI18; theor.).

Table 18.11: The half-life of ^{18}F

$\tau_{1/2}$ (min)	Refs.
112 ± 1	(1949BL26)
111 ± 1	(1955JA1A)
110 ± 1	(1958BE74)
110.5 ± 0.6	(1964HO28)
109.72 ± 0.06	(1964MA12)
111.0 ± 2.0	(1965BO42)
109.87 ± 0.12	(1965EB01)
109.77 ± 0.05	mean

2. (a) $^{12}\text{C}(^6\text{Li}, \text{n})^{17}\text{F}$	$Q_{\text{m}} = 4.064$	$E_{\text{b}} = 13.215$
(b) $^{12}\text{C}(^6\text{Li}, \text{p})^{17}\text{O}$	$Q_{\text{m}} = 7.606$	
(c) $^{12}\text{C}(^6\text{Li}, \text{d})^{16}\text{O}$	$Q_{\text{m}} = 5.688$	
(d) $^{12}\text{C}(^6\text{Li}, \text{t})^{15}\text{O}$	$Q_{\text{m}} = -3.723$	
(e) $^{12}\text{C}(^6\text{Li}, \alpha)^{14}\text{N}$	$Q_{\text{m}} = 8.799$	

Cross section measurements have been carried out in the range $E(^6\text{Li}) = 1.9$ to 14.0 MeV: see Table 18.12. The cross sections for reactions (b), (c) and (e) rise monotonically and rapidly with energy up to $E(^6\text{Li}) = 4$ MeV due to Coulomb barrier effects. At higher energies, Ericson-type fluctuations are observed (1967DZ01, 1970JO09). Neither the direct reaction nor the statistical compound nucleus model alone is adequate to describe the data (1970JO09). The cross section for the isospin-forbidden α_1 group [to $^{14}\text{N}^*(2.31)$, $J^\pi = 0^+$; $T = 1$] shows an energy dependence very unlike those for the α_0 and α_2 groups, characterized by a maximum at ≈ 4 MeV, a minimum at ≈ 5 MeV, and a further rise to 6 MeV. Typically the cross section for the α_1 group is two orders of magnitude lower than those for the α_0 or α_2 groups (1965CA06, 1967DZ01). See also (1968NO1C; theor.). For angular distribution studies see ^{16}O and ^{17}O in (1971AJ02) and ^{14}N in (1970AJ04).

3. $^{12}\text{C}(^7\text{Li}, \text{n})^{18}\text{F}$	$Q_{\text{m}} = 5.964$
--	------------------------

See (1957NO17, 1961NO05) and ^{19}F .

4. $^{12}\text{C}(^{12}\text{C}, ^6\text{Li})^{18}\text{F}$	$Q_{\text{m}} = -14.960$
---	--------------------------

See (1962CH01).

Table 18.12: $^{12}\text{C} + ^6\text{Li}$ studies ^a

$E(^6\text{Li})$ (MeV)	Measurement of	Refs.
1.9 – 3.5	α_0, α_2	(1966BE07)
2.2 – 3.4	^{17}F yield	(1961NO05)
2.4 – 8.5	$p_0 \rightarrow p_3, d_0, \alpha_0$	(1967DZ01)
3.0 – 5.5	$p_0, d_0, \alpha_0, \alpha_2$ (σ_t)	(1965CA06)
3.2 – 4.0	α_0, α_2	(1962HO06)
3.4 – 4.0	$p_0 \rightarrow p_3, d_0$ (σ_t)	(1962BL13)
3.5 – 6.5	α_1	(1967DZ01)
3.5 – 7.5	α_2	(1967DZ01)
3.8 – 4.0	α_1	(1962HO06)
3.8 – 5.5	α_1 (σ_t)	(1965CA06)
4.5 – 5.5	$p_0 \rightarrow p_4, d_0, d_{1+2}, d_3, d_4, \alpha_0, \alpha_2$ (σ_t)	(1966HE05)
5.6 – 14.0	$p_0 \rightarrow p_3, d_0, d_{1+2}, d_{3+4}, \alpha_0, \alpha_2, \alpha_{3+4}$ (σ_t)	(1970JO09)
9.0 – 14.0	p_4, d_5 (σ_t)	(1970JO09)
12.0 – 14.0	t_0 (σ_t)	(1970JO09)

^a At $E_{c.m.} = 7$ and 8 MeV, σ_t for various deuteron groups are reported by (1967LO01).

Table 18.13: Resonances in $^{14}\text{N} + \alpha$

E_α (MeV)	Particle out	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	$(2J+1)\Gamma_\gamma\Gamma_\alpha/\Gamma$ (eV)	E_x (MeV)	Refs.
0.559	γ		1; 0	$(2.79 \pm 0.5) \times 10^{-4}$	4.851	(1971CO27)
1.140 ± 5	γ		$4^+; 0$	0.084 ± 0.004^b	5.303	(1968PA10, 1971CO27, 1971RO25)
1.395 ± 5	γ			0.027 ± 0.003	5.501	(1968PA10)
1.531 ± 2	γ, α_0	< 1.2	$1^-; 0, 1$	4.80 ± 0.40	5.606	(1955PR1A, 1958AL03, 1958PH37, 1961SI09, 1968PA10, 1971CH1F)
1.618 ± 2	γ, α_0	< 0.8	$1^-; 0, 1$	1.35 ± 0.15	5.674	(1955PR1A, 1958PH37, 1959WA16, 1961SI09, 1968PA10, 1971CH1F)
2.165 ± 3	α_0		2, 3, 4		6.100	(1961SI09)
2.351 ± 3	γ, α_0	< 0.8	$2^{(+)}; 1$	≈ 9	6.244	(1958HE54, 1958HE56, 1958PH37, 1959WA16, 1961SI09)
2.372 ± 3	α_0	< 3	1		6.261	(1961SI09)
2.767 ± 4	γ, α_0	(< 0.8)	$5^+; 0$	c	6.568	(1958HE54, 1958HE56, 1971RO25)
2.870 ± 4	γ, p_0	< 1.6		≈ 3	6.648	(1958HE54, 1958HE56, 1958PH37)
2.870 ± 6	α_0	93 ± 5	1^-	$\Gamma_\alpha/\Gamma = 0.85$	6.648	(1958HE54, 1958HE56, 1958KA32)
3.080 ± 6	p_0, α_0	101 ± 5	2^-		6.811	(1958HE54, 1958HE56)
3.576 ± 4	α_0	< 4	(4^+)		7.197	(1958HE54, 1958HE56, 1958KA32)
3.67	α_0	45 ± 10	(1^+)		7.27	(1958HE54, 1958HE56, 1958KA32)
3.72	p_0, α_0	53 ± 6	(3^-)		7.31	(1958HE54, 1958HE56, 1958KA32)
4.00	p_0, α_0	35	(3^-)		7.53	(1958KA32)
4.05	p_0, α_0	60			7.57	(1958KA32)
4.11	p_0, α_0	40			7.61	(1958KA32)
4.28	p_0, α_0	120			7.74	(1958KA32)
4.50	p_0, α_0	30	(2^-)		7.92	(1958KA32)
4.55	p_0, α_0	70	(1^+)		7.95	(1958KA32)
5.2	α				8.5	(1939BR1A, 1939DE1A)
8.12 ± 30	(n_1, p_1)		$(2^-); 0$		10.73	(1969SC21)
9.5	(n_1, p_1)				11.8	(1969SC21)
10.07 ± 30	(n_1, p_1)		$(2^-, 3); 0$		12.25	(1969SC21)
10.376 ± 15	α_1^d	75	4^+		12.484	(1966CH1E, 1970TO03)
10.431 ± 15	α_1	75	3^-		12.526	(1970TO03)
10.481 ± 15^a	α_1	350	2^+		12.565	(1970TO03)

Table 18.13: Resonances in $^{14}\text{N} + \alpha$ (continued)

E_α (MeV)	Particle out	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	$(2J + 1)\Gamma_\gamma\Gamma_\alpha/\Gamma$ (eV)	E_x (MeV)	Refs.
10.632 ± 15	α_1	54	5^-		12.683	(1970TO03)
10.657 ± 15	α_1	74	3^-		12.702	(1970TO03)
11.110 ± 15	α_1	37	(4^+)		13.054	(1970TO03)
11.206 ± 15	α_1	43	(4^+)		13.129	(1970TO03)
11.295 ± 15	α_1		5^-		13.198	(1970TO03)
11.467 ± 15	α_1	60	5^-		13.332	(1966CH1E, 1970TO03)
11.553 ± 15	α_1	85	6^+		13.398	(1970TO03)
11.638 ± 15	α_1	93	4^+		13.464	(1970TO03)
11.809 ± 15	α_1	100	6^+		13.597	(1970TO03)
11.879 ± 15	α_1	62	3^-		13.652	(1970TO03)
11.904 ± 15	α_1	62	(2^+)		13.671	(1970TO03)
11.914 ± 15	α_1	78	4^+		13.679	(1970TO03)
12.044 ± 15	α_1		4^+		13.780	(1970TO03)
12.220 ± 15	α_1		$(4^+, 2^+)$		13.917	(1970TO03)
12.301 ± 15	α_1		5^-		13.980	(1970TO03)
12.535 ± 15	α_1		5^-		14.162	(1966CH1E, 1970TO03)
12.743 ± 15	α_1	93	4^+		14.323	(1966CH1E, 1970TO03)
15.56 ± 15	α_1	≈ 50	$6^+(7^-)$		16.513	(1970TO03)
15.68 ± 15	α_1	≈ 50	5^-		16.606	(1970TO03)
15.83 ± 15	α_1	≈ 150	6^+		16.723	(1970TO03)
16.02 ± 15	α_1	≈ 200	7^-		16.870	(1970TO03)
16.12 ± 15	α_1	≈ 150	6^+		16.948	(1970TO03)
19.8	α	broad			19.8	(1962JO14)

^a May be a cluster of levels (1970TO03).

^b $\Gamma_\gamma = \Gamma_\alpha = 12 \pm 4$ meV (1971RO25).

^c $\Gamma_\gamma = 26 \pm 6$ meV, $\Gamma_\alpha \gg \Gamma_\gamma$ (1971RO25).

^d This and the following resonances are observed in the isospin-forbidden channel (H.T. Richards, private communication).

Table 18.14: Radiative decays in ^{18}F

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.			
0.94	$3^+; 0$	0	100	$(2.2 \pm 0.2) \times 10^{-5}$	(1966AL04, 1966CH12, 1967GO07)			
1.04	$0^+; 1$	0	100					
1.08	$0^-; 0$	0	100					
1.13	$5^+; 0$	0.94	100			$ M(E2) ^2 = 4.62 \pm 0.16$ W.u.	(1960RA18, 1966AL04, 1967OL03, 1967PO09, 1967WA19)	
1.70 ^h	$1^+; 0$	0 ^a	35 ± 3	$(4.6 \pm 2.2) \times 10^{-5}$	(1965PO01)			
			28 ± 3			(1966OL03)		
		0.94	< 1			(1967WA06)		
		1.04 ^a	65 ± 3			(1965PO01)		
			72 ± 3			(1966OL03)		
		1.08	< 3.5			(1967WA06)		
		1.13	< 3			(1967WA06)		
		2.10 ^h	$2^-; 0$			0 ^b	38 ± 3	(1965PO01)
						36 ± 2	(1966OL03)	
						33 ± 3	(1967GO07)	
	36	(1967WA06)						
	35.8 ± 1.5	mean						
	0.94 ^b	30 ± 4		(1965CH10, 1965PO01)				
		32 ± 2		(1966OL03)				
		33 ± 3		(1967GO07)				
		31		(1967WA06)				
		32.0 ± 1.5		mean				
	1.04	< 3	(1966OL03, 1967GO07)					
		< 0.8	(1967WA06)					
	1.08 ^b	32 ± 4	(1965CH10, 1965PO01)					
		32 ± 2	(1966OL03)					
		34 ± 3	(1967GO07)					
		32.5 ± 1.5	mean					
	1.13	< 3	(1966OL03, 1967GO07)					
		< 2	(1967WA06)					
	1.70	< 4	(1967WA06)					
2.53 ^h	$2^+; 0$	0 ^c	73 ± 4	$ M(E2) ^2 = 20 \pm 7$ W.u.	(1965PO01)			
			75 ± 4			(1966OL03)		
		0.94 ^c	23 ± 2			(1965PO01)		
			21 ± 4			(1966OL03)		
		1.04	< 0.5			(1967WA06)		

Table 18.14: Radiative decays in ^{18}F (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.	
3.06	$2^+; 1$	1.08	< 0.5		(1967WA06)	
		1.13	< 4		(1967WA06)	
		1.70 ^b	4 ± 2		(1965PO01)	
			4 ± 1.5		(1966OL03)	
			5 ± 2		(1967WA06)	
			4.3 ± 1.0		mean	
		2.10	< 4		(1967WA06)	
		0	24 ± 4		(1965PO01)	
			25 ± 2		(1966OL03)	
		0.94	76 ± 4		(1965PO01)	
	75 ± 2	(1966OL03)				
1.04	< 8	(1967WA06)				
1.08	< 8	(1967WA06)				
1.13	< 14	(1967WA06)				
1.70	< 8	(1967WA06)				
2.10	< 5	(1967WA06)				
2.53	< 5	(1967WA06)				
3.13	$1^-; 0$	0	31 ± 3	$(5.7 \pm 2) \times 10^{-4}$	(1965PO01)	
			32 ± 2		(1966OL03)	
		0.94	< 2		(1967WA06)	
		1.04 ^d	41 ± 4		$(7.3 \pm 2.7) \times 10^{-4}$	(1967WA06)
		1.08 ^d	27 ± 4		$(4.8 \pm 1.8) \times 10^{-4}$	(1967WA06)
		1.13	< 8		(1967WA06)	
		1.70	< 6		(1967WA06)	
		2.10	< 6		(1967WA06)	
		2.53	< 1		(1967WA06)	
		3.36	$3^+; 0$		0	53 ± 5
	56 ± 6			(1966OL03)		
0.94	4 ± 4			(1965PO01)		
	7 ± 4			(1966OL03)		
	7 ± 3			(1967WA06)		
	6 ± 2			mean		
1.04	< 5			(1967WA06)		
1.08	< 5			(1967WA06)		
1.13	< 25			(1967WA06)		
1.70	38 ± 4			(1965PO01)		

Table 18.14: Radiative decays in ^{18}F (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.			
3.72	$1^+; 0^f$	2.10	30 ± 4	$> 8.2 \times 10^{-3}$	(1966OL03)			
			5 ± 4		(1965PO01)			
			7 ± 4		(1966OL03)			
		2.53	< 25		(1967WA06)			
			0		6 ± 2	(1967GO07)		
		3.79	$(3^-); 0$		0	6 ± 3	(1967OL03)	
						16 ± 8	(1967WA06)	
						6.4 ± 1.6	mean	
						0.94	< 6	(1967WA06)
						1.04	94 ± 2	(1967GO07)
	94 ± 3					(1967OL03)		
1.08	< 15					(1967WA06)		
1.13	< 10					(1967OL03)		
1.70	< 6					(1967OL03)		
2.10	< 10					(1967OL03)		
3.84	2^+	0	2.53		< 3	(1967OL03)		
			3.06		< 5	(1967OL03)		
			3.13		< 3	(1967OL03)		
			3.36		< 4	(1967OL03)		
			0.94		< 15	(1967OL03)		
			1.04	< 10	(1967OL03)			
			1.04	< 15	(1967OL03)			
			1.08	< 15	(1967OL03)			
			1.13	< 10	(1967OL03)			
			1.70	< 10	(1967OL03)			
3.84	2^+	0	2.10	76 ± 10	(1967OL03)			
			2.53	4 ± 4	(1967OL03)			
			3.06	20 ± 8	(1967OL03)			
			3.13	< 10	(1967WA06)			
			3.36	< 20	(1967WA06)			
			0.94	39 ± 6	(1967GO07)			
				41 ± 4	(1967OL03)			
			0.94	5 ± 4	(1967GO07)			
				5 ± 3	(1967OL03)			
			1.04	< 4	(1967OL03)			
1.08	< 3	(1967OL03)						
1.13	< 3	(1967OL03)						
		$> 2.2 \times 10^{-3}$						

Table 18.14: Radiative decays in ^{18}F (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.
4.12	3^+ g	1.70	4 ± 4		(1967GO07)
			4 ± 3		(1967OL03)
		2.10	< 5		(1967OL03)
		2.53	< 2		(1967OL03)
		3.06	61 ± 6		(1967GO07)
			50 ± 4		(1967OL03)
		3.13	< 6		(1967OL03)
		3.36	< 9		(1967OL03)
		0	5 ± 3		(1967OL03)
		0.94	< 8		(1967OL03)
		1.04	< 8		(1967OL03)
		1.08	< 8		(1967OL03)
		1.13	< 8		(1967OL03)
		1.70	< 8		(1967OL03)
		2.10	< 15		(1967OL03)
2.53	< 15		(1967OL03)		
3.06	95 ± 3		(1967OL03)		
3.13	< 13		(1967OL03)		
3.36	< 10		(1967OL03)		
4.23	(2)	0^e	32 ± 5		(1967GO07)
		0.94^e	55 ± 5		(1967GO07)
		1.04	< 5		(1967OL03)
		1.08	(≤ 5)		(1967OL03)
		1.13	< 10		(1967OL03)
		1.70	5 ± 3		(1967OL03)
		2.10^e	13 ± 5		(1967GO07)
		2.53	< 4		(1967OL03)
		3.06	< 3		(1967OL03)
		3.13	(≤ 5)		(1967OL03)
		3.36	< 5		(1967OL03)
		4.36	2, 3	0.94	< 20
3.06	(100)				(1967GO07)
4.40	$\geq 2; 0$		> 60		(1967OL03)
		0.94	< 40		(1967GO07)
			≤ 30		(1967OL03)
		1.13	(100)		(1967GO07)
			> 70		(1967OL03)

Table 18.14: Radiative decays in ^{18}F (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.		
4.66	$4^+; 1^g$	0	< 5		(1967OL03)		
		0.94	15 ± 5		(1967OL03)		
		1.04	< 5		(1967OL03)		
		1.08	< 5		(1967OL03)		
		1.13	85 ± 5		(1967OL03)		
		1.70	< 5		(1967OL03)		
		2.10	< 10		(1967OL03)		
		2.53	< 10		(1967OL03)		
		3.06	< 4		(1967OL03)		
		3.13	< 4		(1967OL03)		
		3.36	< 3		(1967OL03)		
		4.74	$0^+; 1$		0	(100)	(1967GO07)
		4.85	$1; 0$		0	< 6	
0.94	< 4			(1967OL03)			
1.04	65 ± 4			(1967GO07)			
	60 ± 10			(1967OL03)			
1.08	< 15			(1967OL03)			
1.13	< 15			(1967OL03)			
1.70	< 10			(1967OL03)			
2.10	< 15			(1967OL03)			
2.53	< 10			(1967OL03)			
3.06	35 ± 4			(1967GO07)			
	40 ± 10			(1967OL03)			
3.13	< 15			(1967OL03)			
3.36	< 15			(1967OL03)			
4.96	$2^+; 1$	0	(100)	(1967GO07)			
5.30	$4^+; 1$	0.94	13 ± 3		(1968PA10)		
			9		(1971RO25)		
		1.13	7		(1971RO25)		
		2.53	87 ± 3		(1968PA10)		
			78		(1971RO25)		
		3.36	5		(1971RO25)		
	4.66	1	(1971RO25)				
5.50				$\Gamma_\gamma/\Gamma = 0.38 \pm 0.12$	(1967GO07)		
5.61 ⁱ	$1^-; 0, 1$	0	10 ± 1		(1968PA10)		
		1.04	4 ± 1		(1971CH1F)		

Table 18.14: Radiative decays in ^{18}F (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.
5.67	$1^-; 0, 1$	1.08	35 ± 2	$\left\{ \begin{array}{l} 0.17 \pm 0.09 \\ \Gamma_\gamma = 66 \pm 21 \text{ meV} \\ 0.063 \end{array} \right.$	(1971CH1F)
		3.06	30 ± 2		(1971CH1F)
		3.13	21 ± 2		(1971CH1F)
		0	6 ± 1		(1971CH1F)
		1.04	8 ± 1		(1971CH1F)
		1.08	52 ± 3		(1971CH1F)
		3.06	4 ± 1		(1971CH1F)
6.10	$4^-; 0$	3.13	29 ± 2		(1971CH1F)
		0.94	6		(1971BE1E)
			6		(1971SE1H)
		1.13	57		(1971BE1E)
			66		(1971SE1H)
		2.10	25		(1971BE1E)
			28		(1971SE1H)
6.14	0^+	3.79	2	(1971BE1E)	
		4.12	2	(1971BE1E)	
		4.66	8	(1971BE1E)	
		0	54	0.57 (1971SE1H)	
		1.70	10	(1971SE1H)	
6.16	3^+	3.72	36	(1971SE1H)	
		0.94	51	1.72 (1971SE1H)	
		2.53	8	(1971SE1H)	
		3.79	13	(1971SE1H)	
6.24 ^j	3^-	3.84	28	(1971SE1H)	
		0.94	5	1.00 (1971SE1H)	
		2.10	76	(1971SE1H)	
		3.79	12	(1971SE1H)	
6.27	2^+	4.23	7	(1971SE1H)	
		0.94	70	1.91 (1971SE1H)	
		1.70	4	(1971SE1H)	
		2.10	5	(1971SE1H)	
		3.36	3	(1971SE1H)	
		3.84	15	(1971SE1H)	
6.31	$2^+, 3^+$	4.12	3	(1971SE1H)	
		0	4	(1971SE1H)	
		0.94	33	(1971SE1H)	

Table 18.14: Radiative decays in ^{18}F (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (eV)	Refs.
6.39		3.06	55	0.18	(1971SE1H)
		4.96	8		(1971SE1H)
6.48		0.94	80	0.16	(1971SE1H)
		1.70	7		(1971SE1H)
		3.84	13		(1971SE1H)
		0	15		(1971SE1H)
		0.94	31	0.094	(1971SE1H)
6.57	$5^+; 0$	1.13	14		(1971SE1H)
		1.70	4		(1971SE1H)
		3.06	28		(1971SE1H)
		3.84	8		(1971SE1H)
		0.94	(100)	0.043	(1971SE1H)
6.65 ^j	(2^-)		15		(1971RO25)
		3.36	83		(1971RO25)
		5.30	2		(1971RO25)
		0.94	12		(1971SE1H)
6.78	$4^+, 5^+$	2.10	72	1.23	(1971SE1H)
		3.13	16		(1971SE1H)
		0.94	15		(1971SE1H)
6.81	2^-	1.13	38		(1971SE1H)
		4.66	47	0.48	(1971SE1H)
		0	24		(1971SE1H)
6.86		0.94	35		(1971SE1H)
		3.06	41	0.17	(1971SE1H)
		0.94	12		(1971SE1H)
		4.66	88	0.15	(1971SE1H)

^a See also (1960RA18, 1961KU02, 1963LI07, 1967WA06).

^b See also (1961KU02, 1966CH12).

^c See also (1961KU02, 1967WA06).

^d See also (1965PO01, 1966OL03).

^e See also (1967OL03).

^f Parity is positive: see (1970RO1F).

^g Parity is positive: see (1970DU08).

^h See also (1966HA31).

ⁱ See also (1955PR1A, 1958AL03).

^j See also (1958PH37).

5. $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$

$$Q_m = 4.416$$

Resonances have been observed at $E_\alpha = 0.56, 1.14, 1.40, 1.53, 1.62, 2.35, 2.77$ and 2.88 MeV, corresponding to $^{18}\text{F}^* = 4.85, 5.30, 5.50, 5.61, 5.68, 6.24, 6.57$ and 6.65 MeV: see Table 18.13 (1955PR1A, 1958PH37, 1968PA10, 1971CH1F, 1971CO27, 1971RO25). See also (1970AL1F). Capture resonances are not observed corresponding to $^{18}\text{F}^*(4.66, 4.96), (2J + 1)\Gamma_\alpha\Gamma_\gamma/\Gamma < 2 \times 10^{-5}$ and $< 0.5 \times 10^{-4}$ eV, respectively (1971CO27).

The branching ratios for $^{18}\text{F}^*(5.30, 5.61, 5.68, 6.57)$ are displayed in Table 18.14.

The sequence of the ^{18}F states with $E_x = 1.70 [J^\pi = 1^+], 2.53 [2^+], 3.36 [3^+], 5.30 [4^+]$ and 6.57 MeV $[5^+]$ as well as the enhanced E2 strengths of the γ -transitions show that these states can be considered to be members of a predominantly 4p-2h $K^\pi = 1^+$ rotational band. $^{18}\text{F}^*(3.36)$ is found to have $J = 3$ (1971RO25). The M1 transitions between the $J^\pi = 1^-$ states at $E_x = 5.61$ and 5.68 MeV and $^{18}\text{F}^*(1.08, 3.13)$ have $|M|^2$ between 0.1 and 1 W.u. implying $T = 1$ for these resonant states but both also decay by E1 transitions to the $T = 1$ states at $E_x = 1.04$ and 3.06 MeV, implying $T = 0$. This result is in agreement with $^{18}\text{F}^*(5.61, 5.68)$ both being 1^- members of a $T = 0, 1$ isospin doublet. The $T = 1$ strength is split nearly equally between the two levels (1971CH1F). See also (1967CH1J, 1967CH1K).

The non-resonant S factor for this reaction, $S \leq 1.5 \times 10^6$ keV \cdot b (1968PA10), ≈ 0.7 MeV \cdot b (1971CO27). See also (1966IB1A, 1971CO1G).

6. $^{14}\text{N}(\alpha, n)^{17}\text{F}$

$$Q_m = -4.735$$

$$E_b = 4.416$$

The total cross section ratios of $^{14}\text{N}(\alpha, n)^{17}\text{F}(0.50)/^{14}\text{N}(\alpha, p)^{17}\text{O}(0.87)$ were measured for $E_\alpha = 7$ to 12 MeV: major maxima are observed at $E_\alpha = 8.12, 9.5, 10.07$ and 11.52 MeV. Symmetric angular distributions at three of these maxima lead to $J^\pi = (2^-), (2^-, 3)$ and $(2^-, 3^+)$ at $E_x = 10.73, 12.25$ and 13.36 MeV. These states are primarily $T = 0$ (1969SC21): see Table 18.13.

7. $^{14}\text{N}(\alpha, p)^{17}\text{O}$

$$Q_m = -1.193$$

$$E_b = 4.416$$

Observed resonances are displayed in Table 18.13 (1953HE1A, 1958HE54, 1958KA32). Excitation functions showing a number of structures have been measured for $E_\alpha = 10$ to 25 MeV (p_0, p_1), 13 to 25 MeV (p_2) and 15 to 25 MeV (p_3) (1970ZE01). See also (1961YA02).

8. (a) $^{14}\text{N}(\alpha, \alpha)^{14}\text{N}$

$$E_b = 4.416$$

(b) $^{14}\text{N}(\alpha, \alpha p)^{13}\text{C}$

$$Q_m = -7.551$$

(c) $^{14}\text{N}(\alpha, d)^{16}\text{O}$

$$Q_m = -3.111$$

Table 18.15: Lifetime measurements of some ^{18}F states

$^{18}\text{F}^*$ (MeV)	τ_m	Reaction	Refs.
0.94	68 ± 7 psec	$^3\text{H}(^{16}\text{O}, ^{18}\text{F})\text{n}$	(1966AL04) ^a
1.04	4_{-2}^{+3} fsec	$^{18}\text{O}(\text{p}, \text{n})^{18}\text{F}$	(1967BL18) ^{b,c}
1.08	30 ± 3 psec	$^3\text{H}(^{16}\text{O}, ^{18}\text{F})\text{n}$	(1966AL04) ^b
1.13	220 nsec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1970BL1F)
	190 ± 45 nsec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1959AL99)
	221 ± 21 nsec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967BE14)
	219 ± 10 nsec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967PO09)
	218 ± 9 nsec		mean value
1.70	2.0 ± 1.0 psec	$^3\text{H}(^{16}\text{O}, ^{18}\text{F})\text{n}$	(1963LI07)
	0.86 ± 0.20 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1966OL03)
	0.86 ± 0.20 psec		best value
2.10	0.70 ± 0.20 psec	$^3\text{H}(^{16}\text{O}, ^{18}\text{F})\text{n}$	(1963LI07)
	5 ± 3 psec	$^3\text{H}(^{16}\text{O}, ^{18}\text{F})\text{n}$	(1966AL04)
	4.1 ± 1.6 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1966OL03)
	4.3 ± 1.4 psec		best value
2.53	1.1 ± 0.2 psec	$^3\text{H}(^{16}\text{O}, ^{18}\text{F})\text{n}$	(1963LI07)
	$0.63_{-0.14}^{+0.18}$ psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967WA06)
	0.67 ± 0.18 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1966OL03)
	0.65 ± 0.13 psec		best value
3.06	< 0.20 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967WA06)
	< 0.17 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1966OL03)
	≤ 1.2 fsec	$^{17}\text{O}(\text{d}, \text{n})^{18}\text{F}$	(1971LE29)
	$0.37_{-0.10}^{+0.15}$ psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967WA06)
3.36	$0.55_{-0.15}^{+0.20}$ psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967WA06)
	0.46 ± 0.10 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1966OL03)
	0.48 ± 0.09 psec		mean value
3.72	< 0.08 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967WA06)
3.84	< 0.1 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1967WA06)
	< 0.073 psec	$^{16}\text{O}(^3\text{He}, \text{p})^{18}\text{F}$	(1966OL03)
4.36	< 0.61 psec	$^{16}\text{O}(^6\text{Li}, \alpha)^{18}\text{F}$	(1969TH01)
5.30	31 ± 6 fsec	$^{14}\text{N}(\alpha, \gamma)^{18}\text{F}$	(1971RO25)

^a See also (1963LI07, 1963LO03).

^b See also (1963LI07).

^c See also (1966OL03).

Observed anomalies in the elastic scattering are exhibited in Table 18.13 (1939BR1A, 1939DE1A, 1953HE1A, 1958HE54, 1958HE56, 1958KA32, 1961SI09, 1962JO14, 1970TO03). Resonances in the α_1 (isospin “forbidden”) and α_2 yield are also displayed (1970TO03). See also (1966CH1E). Excitation functions have recently been reported for $E_\alpha = 1.0$ to 2.4 MeV (1961SI09; α_0), 10.2 to 18.3 MeV (1970TO03; $\alpha_0, \alpha_1, \alpha_2$), 15 to 23 MeV (1962JO14; α_0) and 18.2 to 23.5 MeV (1970CH1D; α_1). See also (1963NO1B, 1967BO1M, 1969FE10).

In the range $E_\alpha = 10.2$ to 17.3 MeV compound nucleus formation dominates the α_0, α_1 and α_2 channels. The α_1 channel [to the $J^\pi = 0^+; T = 1$ state at 2.31 MeV] displays relatively isolated resonances which imply ^{18}F states with large isospin impurities [$\approx 10 - 25\%$]. Comparison of this work with the results from $^{16}\text{O}(d, \alpha_1)^{18}\text{F}$ [reaction 16] suggests that the same ^{18}F states are usually involved in both reactions (1970TO03). These data do not support the suggestions by (1968NO1C, 1969NO1B) concerning the importance of direct reactions in isospin violating reactions (1970TO03). See also (1970JO1G).

For reaction (b), see (1967BE30). For spallation cross sections see (1968JA1J, 1968JU1A, 1970BA48, 1970JA1Q, 1970JU1B). For reaction (c), see (1971TH03) and reaction 16. See also (1970VI02) and ^{14}N in (1970AJ04).

$$9. \ ^{14}\text{N}(^6\text{Li}, d)^{18}\text{F} \quad Q_m = 2.943$$

Angular distributions have been measured for $E(^6\text{Li}) = 5.3$ to 6.0 MeV (1968RI13; $d_0, d_{1 \rightarrow 4}$).

$$10. \ ^{14}\text{N}(^7\text{Li}, t)^{18}\text{F} \quad Q_m = 1.950$$

At $E(^7\text{Li}) = 15$ MeV, triton groups are observed to the known $T = 0$ states with $E_x < 7.4$ MeV: the $T = 1$ states are not excited although such transitions are not forbidden in principle, suggesting a direct α -transfer mechanism. The transitions to $^{18}\text{F}^*(1.70, 2.53, 3.36, 4.23, 5.30, 6.57)$ account for more than one half of the summed cross section at 15° . It is proposed that these states (which are only weakly excited in $^{16}\text{O}(^3\text{He}, p)^{18}\text{F}$ and $^{17}\text{O}(^3\text{He}, d)^{18}\text{F}$) are predominantly of a 4p-2h nature and are excited by transfer of four nucleons into the (2s, 1d) shell (1968MI09). [See also reaction 5]. See also (1968OG1A).

$$11. \ ^{15}\text{N}(\alpha, n)^{18}\text{F} \quad Q_m = -6.418$$

See (1966AD07).

$$12. \ ^{16}\text{O}(d, \gamma)^{18}\text{F} \quad Q_m = 7.527$$

Table 18.16: Recent $^{16}\text{O} + \text{d}$ yield curves and polarization studies

E_d (MeV)	Yield of	Refs.
0.4 – 3.5	γ	(1965OW01)
13.0 – 18.5	γ	(1970AL1F)
thresh. – 2.4	n_0	(1961DI06)
thresh. – 12	n	(1968MA1C)
2.0 – 4.2	n_0, n_1	(1970BA31)
2.1 – 3.4	n_0	(1968DI06)
2.5 – 5.5	n_0, n_1	(1970LO01)
2.8 – 3.4	n_1	(1968DI06)
4.5 – 6.0	n_0, n_1	(1970DA14)
0.3 – 1.0	p_0	(1965LO03)
0.32 – 1.07	p_1	(1968NG1B)
0.35 – 1.05	p_1	(1969DU11)
0.45 – 1.0	p_1	(1965LO03)
0.7 – 2.1	p_0, p_1	(1963SE12)
0.8 – 1.7	p_0, p_1	(1964KI1B)
0.8 – 2.0	p_0, p_1	(1962CA20, 1963AM1A, 1964AM1A)
1.55 – 1.85	p_0	(1959LO59, 1961LO1C)
1.9 – 3.6	p_0, p_1	(1968DI06)
4.0 – 6.0	p_0, p_1	(1970DA14)
4.4 – 8.4	p_0, p_1	(1969CO12)
4.5 – 6.0	p_3	(1970DA14)
6.0 – 11.0	p_0, p_1	(1968NA06)
10.5 – 13.0	p_0, p_1, p_3, p_5	(1967AL06)
0.65 – 2.0	d_0	(1963SE12)
0.8 – 2.0	d_0	(1963AM1A, 1964AM1A)
1.0 – 2.5	d_0	(1968MA53)
1.8 – 3.6	d_0	(1968DI06)
4.0 – 6.0	d_0	(1970DA14)
4.4 – 8.4	d_0	(1969CO12)
10.0 – 13.0	d_0	(1967AL06)
0.3 – 1.0	α_0	(1965LO03)

Table 18.16: Recent $^{16}\text{O} + \text{d}$ yield curves and polarization studies (continued)

E_d (MeV)	Yield of	Refs.
0.7 – 2.1	α_0	(1963SE12)
0.8 – 1.7	α_0	(1964KI1B)
0.8 – 2.0	α_0	(1962CA20, 1963AM1A, 1964AM1A)
0.85 – 2.0	α_0	(1960AM03)
1.1 – 2.5	α_0	(1965MA1A)
1.8 – 14	α_0, α_1	(1970JO1C)
1.9 – 3.6	α_0	(1968DI06)
3 – 5	α_0, α_1	(1969JO1M, 1970JO1G)
3 – 15	α_1	(1969JO09)
3.9 – 5.3	α_0	(1967TH1E, 1968TH1J, 1971TH03)
4.4 – 8.4	α_0	(1969CO12)
5.0 – 9.0	$\alpha_0, \alpha_2 \rightarrow \alpha_5, \alpha_7 \rightarrow \alpha_{10}$	(1968JO07)
5.5 – 7.0	α_0, α_2	(1965SA18)
7.0 – 12.5	α_0	(1970AL1F)
7 – 14	α_2	(1970JO1C)
9.0 – 12.5	α_2	(1970AL1F)
9.0 – 15.0	$\alpha_0, \alpha_2, \alpha_3, \alpha_4$	(1968JO07)
14.0 – 18.1	α_1	(1971JA04)
15 – 20	$\alpha_0, \alpha_1, \alpha_2$	(1963YA1B)
E_d (MeV)	Polarization measurements of	Refs.
3 – 4	n_0, n_1	(1971AN1A)
3.96 – 5.35	n_0, n_1	(1971TH10)
6.5 – 9.5	p_1	(1963AL1D)
6.5 – 9.55	p_1	(1963EV05)
8	p_0, p_1	(1971KO21)
9.0 – 10.3	p_0, p_1	(1969CU10)
9.3, 13.3	p_0, p_1, p_3, p_4, p_5	(1970CO1P)
12.3	p_0, p_1	(1971BR44, 1971HU1C)
6.34	d_0	(1969CO12)
8	d_0	(1971KO21)

Table 18.16: Recent $^{16}\text{O} + \text{d}$ yield curves and polarization studies (continued)

E_d (MeV)	Yield of	Refs.
9.3, 13.3	d_0	(1970CO1P)

The capture cross section rises from $0.1 \mu\text{b}$ at $E_d = 0.4 \text{ MeV}$ to $25 \mu\text{b}$ at 3.5 MeV : Γ_γ over this range is $\approx 2 \text{ eV}$. The results can be interpreted satisfactorily in terms of compound nucleus formation (1965OW01). See also (1970AL1F) and Table 18.16.

13. (a) $^{16}\text{O}(\text{d}, \text{n})^{17}\text{F}$ $Q_m = -1.624$ $E_b = 7.527$
 (b) $^{16}\text{O}(\text{d}, \text{np})^{16}\text{O}$ $Q_m = -2.225$

Excitation functions have been measured for the n_0 and n_1 groups from threshold to 12 MeV : see Table 18.16 (1961DI06, 1968DI06, 1968MA1C, 1970BA31, 1970DA14, 1970LO01). Some structure is observed: that which is attributed to states in ^{18}F is displayed in Table 18.17 (1955MA85, 1961DI06, 1968MA1C). The coherence energy determined from the yield for $E_d = 4.5$ to 6.0 MeV is 70 keV for n_0 and 67 keV for n_1 (1970DA14). See also ^{17}F in (1971AJ02). For polarization measurements see (1971AN1A, 1971TH10) and Table 18.16.

For reaction (b) see (1968CU04).

14. $^{16}\text{O}(\text{d}, \text{p})^{17}\text{O}$ $Q_m = 1.918$ $E_b = 7.527$

Excitation functions have been reported recently for several proton groups up to $E_d = 13.0 \text{ MeV}$: see Table 18.16 (1959LO59, 1961LO1C, 1962CA20, 1963AM1A, 1963SE12, 1964AM1A, 1964KI1B, 1965LO03, 1968DI06, 1968NG1B, 1969CO12, 1969DU11, 1970DA14). See also (1963DO1B, 1970CA1C).

Some of the maxima in the yield are interpreted in terms of resonances: these are shown in Table 18.17 (1955ST1A, 1956RO1A, 1963AM1A, 1964AM1A, 1968MA53). See also (1959AJ76). The coherence energy determined from the yields is 75 keV for p_0 , 63 keV for p_1 and 62 keV for p_3 in the range $E_d = 4.0$ to 6.0 MeV (1970DA14).

Polarization measurements are reported by (1963AL1D, 1963EV05, 1969CU10, 1970CO1P, 1971BR44, 1971HU1C, 1971KO21): see Table 18.16. Large values of the polarization are observed for $E_d = 6.5$ to 12.3 MeV : the results are interpreted in terms of a direct interaction mechanism (1963EV05). See also (1967BA1R, 1967MA1F, 1970PE1B; theor.) See also ^{17}O in (1971AJ02).

15. $^{16}\text{O}(\text{d}, \text{d})^{16}\text{O}$ $E_b = 7.527$

Table 18.17: Maxima in the yield of $^{16}\text{O} + \text{d}$ reactions

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Refs.
0.895 ^a	p_1, α_0	210 ± 25		(8.322)	(1963AM1A, 1964AM1A)
1.048 ^b	p_1, d_0, α_0	88 ± 10	1^+	8.458	(1960AM03, 1963AM1A, 1964AM1A, 1968MA53)
1.199	α_0	230 ± 30		(8.592)	(1963AM1A, 1964AM1A, 1965MA1A)
1.298	p_1, d_0, α_0	13 ± 3		(8.680)	(1960AM03, 1963AM1A, 1964AM1A)
1.325	d_0, α_0			(8.704)	(1963AM1A, 1964AM1A)
1.482	α_0	40 ± 5		(8.843)	(1963AM1A, 1964AM1A)
1.563	d_0, α_0	121 ± 15		(8.915)	(1960AM03, 1963AM1A, 1964AM1A)
1.616 ^c	α_0	19 ± 15		(8.962)	(1960AM03, 1963AM1A, 1964AM1A)
1.765 ^d	d_0, α_0	141 ± 10		(9.095)	(1960AM03, 1963AM1A, 1964AM1A)
1.885 ^{d,e}	p_0, p_1, d_0, α_0	108 ± 12	$3, 4^-; 0$	9.201	(1956RO1A, 1963AM1A, 1964AM1A, 1965MA1A, 1970JO1C)
1.95 ^f	α_1	≈ 30	3^-	9.26	(1970JO1C)
2.02 ^f	$(p_1), \alpha_1$	≈ 40	2^+	9.32	(1956RO1A, 1970JO1C)
2.22	n_0, α_0		$2, 3^+; 0$	9.50	(1955MA85, 1961DI06, 1970JO1C)
2.28	α_0		$2, 3^+; 0$	9.55	(1970JO1C)
2.34	n_0, p_1			(9.61)	(1955MA85, 1956RO1A, 1961DI06)
2.448 ^f	n, α_1	372	3^-	9.701	(1955MA85, 1970JO1C)
2.55	p_1			(9.79)	(1955ST1A, 1956RO1A)
2.620 ^f	α_1	195	2^+	9.854	(1970JO1C)
2.92	n, p_0, p_1			10.12	(1955MA85, 1955ST1A, 1956RO1A)
3.021 ^f	α_1	93	1^-	10.210	(1970JO1C)
3.05	α_0		$3, 4^-$	10.24	(1970JO1C)
3.126 ^f	$n, p_1, \alpha_0, \alpha_1$	179	3^-	10.303	(1955MA85, 1956RO1A, 1970JO1C)
3.138 ^f	α_1	572	2^+	10.314	(1970JO1C)
3.179 ^f	$(n), \alpha_1$	269	1^-	10.350	(1955MA85, 1970JO1C)
3.254 ^f	$(n), \alpha_1$	48	3^-	10.417	(1955MA85, 1969JO09, 1970JO1C)
3.366 ^f	n, p_0, p_1, α_1	46	2^+	10.516	(1955MA85, 1955ST1A, 1956RO1A, 1969JO09, 1970JO1C)
3.454 ^f	α_1	157	1^-	10.594	(1970JO1C)
3.47	α_0		$4, 5^+; 0$	10.61	(1970JO1C)
3.471 ^{f,g}	α_1	151	2^+	10.610	(1969JO09, 1970JO1C)
3.492 ^f	α_1	58	1^-	10.628	(1970JO1C)
3.508 ^f	α_1	236	3^-	10.642	(1970JO1C)
3.547 ^{f,g}	α_1	62	2^+	10.677	(1970JO1C)
3.562 ^f	α_1	75	3^-	10.690	(1970JO1C)
3.578 ^{f,g}	α_1	375	2^+	10.705	(1970JO1C)
3.684 ^{f,g}	n, p_0, p_1, α_1	85	2^+	10.799	(1955MA85, 1955ST1A, 1956RO1A,

Table 18.17: Maxima in the yield of $^{16}\text{O}+d$ reactions (continued)

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Res.
					1968MA1C, 1969JO09, 1969JO1C, 1970JO1C)
3.80	p_0, α_0		$(2, 3)^+; 0$	10.90	(1956RO1A, 1957BA14, 1970JO1C)
3.809 ^f	α_1	241	1^-	10.910	(1970JO1C)
3.849 ^{f,g}	α_1	402	2^+	10.945	(1969JO09, 1969JO1C, 1970JO1C)
3.892 ^f	α_1	119	3^-	10.983	(1970JO1C)
3.904 ^{f,g}	α_1	222	2^+	10.994	(1970JO1C)
3.944 ^{f,g}	n, p_1, α_1	78	2^+	11.030	(1955MA85, 1956RO1A, 1970JO1C)
3.95	p_1, α_0	≈ 35	$3, 4^-; 0$	11.03	(1956RO1A, 1957BA14, 1970JO1C)
3.989 ^{f,g}	α_1	51	2^+	11.070	(1969JO09, 1969JO1C, 1970JO1C)
4.031 ^{f,h}	α_1	1009	1^-	11.107	(1970JO1C)
4.069 ^f	n, p_1, α_1	35	3^-	11.141	(1955MA85, 1956RO1A, 1969JO09, 1969JO1C, 1970JO1C)
4.208 ^f	α_1	238	3^-	11.264	(1969JO09, 1969JO1C, 1970JO1C)
4.253 ^f	α_1	475	2^+	11.304	(1970JO1C)
4.264 ^f	α_1	38	4^+	11.314	(1970JO1C)
4.267 ^f	α_1	94	2^+	11.316	(1970JO1C)
4.276 ^f	α_1	65	3^-	11.324	(1970JO1C)
4.38	p_1, α_0		$4, 5^+; 0$	11.42	(1956RO1A, 1970JO1C)
4.480 ^f	α_1	126	2^+	11.505	(1970JO1C)
4.543 ^f	α_1	67	3^-	11.561	(1970JO1C)
4.57	α_0		$5, 6^-; 0$	11.59	(1970JO1C)
4.655 ^f	α_1	240	2^+	11.661	(1970JO1C)
4.80	d_0, α_0		$\geq 3; 0$	11.79	(1956BE1B, 1970JO1C)
4.847 ^f	α_1	179	1^-	11.831	(1970JO1C)
4.93	α_0		$5, 6^-; 0$	11.91	(1970JO1C)
4.993 ^f	α_1	32	3^-	11.961	(1969JO09, 1969JO1C, 1970JO1C)
5.000 ^f	α_1	124	2^+	11.967	(1970JO1C)
5.05 ± 15	α_4	40		12.01	(1968JO07)
5.10	$\alpha_0, \alpha_2, \alpha_4$	60	$T = 0$	12.06	(1968JO07, 1970JO1C)
5.17	α_0	55	$T = 0$	12.12	(1968JO07, 1970JO1C)
5.175 ^f	α_1	36	3^-	12.123	(1970JO1C)
5.190 ^f	α_1	219	2^+	12.136	(1970JO1C)
5.32	α_0	70		12.25	(1968JO07)
5.34	α_0, α_2	170		12.27	(1968JO07)
5.40	α_0, α_4	130		12.32	(1968JO07)
5.390 ^f	α_1	100	1^-	12.314	(1970JO1C)
5.414 ^f	α_1	187	3^-	12.335	(1970JO1C)

Table 18.17: Maxima in the yield of $^{16}\text{O}+d$ reactions (continued)

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Res.
5.441 ^f	α_1, α_4	80	2^+	12.359	(1968JO07, 1970JO1C)
5.462 ^f	$\alpha_1, \alpha_2, \alpha_3, \alpha_4$	120	2^+	12.377	(1968JO07, 1970JO1C)
5.58	$\alpha_0, \alpha_1, \alpha_2$	120	(3^-)	12.48	(1968JO07, 1969JO09, 1969JO1C, 1970JO1C)
5.571 ^f	α_1	35	1^-	12.474	(1970JO1C)
5.606 ^f	$(\alpha_0), \alpha_1, (\alpha_2)$	88	3^-	12.505	(1968JO07, 1969JO09, 1969JO1C, 1970JO1C)
5.617 ^f	$(\alpha_0), \alpha_1, (\alpha_2)$	47	4^+	12.515	(1968JO07, 1969JO09, 1969JO1C, 1970JO1C)
5.742 ^f	α_0, α_1	210	2^+	12.626	(1968JO07, 1969JO1C, 1970JO1C)
5.786 ^f	α_1	288	2^+	12.665	(1970JO1C)
5.80	$\alpha_0, \alpha_2, \alpha_4$	160		12.68	(1968JO07)
5.804 ^f	$\alpha_1, \alpha_3, \alpha_4$	36	5^-	12.681	(1968JO07, 1969JO09, 1969JO1C, 1970JO1C)
5.867 ^f	α_1	63	3^-	12.737	(1969JO1C, 1970JO1C)
5.91	α_2	160		12.78	(1968JO07)
5.954 ^f	α_1	441	2^+	12.814	(1970JO1C)
6.00	α_0	120		12.86	(1968JO07)
6.048 ^f	α_1	120	3^-	12.898	(1970JO1C)
6.11	$\alpha_0, \alpha_1, \alpha_4$	120		12.96	(1968JO07)
6.14 ⁱ	α_1	100		(12.98)	(1968JO07)
6.151 ^f	α_1	648	2^+	12.989	(1970JO1C)
6.198 ^f	$\alpha_1, \alpha_2, \alpha_3$	250	3^-	13.031	(1968JO07, 1970JO1C)
6.25	α_0, α_4	150	$T = 0$	13.08	(1968JO07, 1970JO1C)
6.25	α_1	≈ 10	$4^+{}^i$	13.08	(1956BR36, 1968JO07, 1969JO09)
6.30	α_0, α_2	160		13.12	(1968JO07)
6.31	α_1	35		13.13	(1968JO07)
6.34	α_0, α_3	160		13.16	(1968JO07)
6.353 ^f	α_1	184	1^-	13.169	(1970JO1C)
6.369 ^f	α_1	137	5^-	13.183	(1968JO07, 1970JO1C)
6.38	α_0, α_3	145	$T = 0$	13.20	(1968JO07, 1970JO1C)
6.393 ^f	α_1	254	4^+	13.204	(1970JO1C)
6.43	d_0, α_2	120		13.24	(1956BE1B, 1968JO07)
6.46	$\alpha_0, \alpha_1, \alpha_4$	100		13.26	(1968JO07)
6.461 ^f	α_1	359	4^+	13.264	(1970JO1C)
6.487 ^f	α_1	99	2^+	13.288	(1968JO07, 1970JO1C)
6.54	α_0, α_2	135		13.34	(1968JO07)
6.563 ^f	α_1	42	5^-	13.355	(1970JO1C)

Table 18.17: Maxima in the yield of $^{16}\text{O}+d$ reactions (continued)

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Res.
6.61	$(\alpha_1), \alpha_2, \alpha_3, \alpha_4$	120		13.40	(1956BR36, 1968JO07)
6.64	α_0, α_2	200		13.43	(1968JO07)
6.650 ^f	α_1	233	3^-	13.432	(1970JO1C)
6.662 ^f	α_0, α_1	100	1^-	13.443	(1968JO07, 1970JO1C)
6.72	α_2	100		13.49	(1968JO07)
6.724 ^f	α_1	65	5^-	13.498	(1968JO07, 1970JO1C)
6.726 ^f	α_1, α_2	146	2^+	13.500	(1968JO07, 1970JO1C)
6.80	$\alpha_1, \alpha_2, \alpha_3$	140		13.57	(1968JO07)
6.84	$\alpha_0, \alpha_2, \alpha_4$	150		13.60	(1968JO07)
6.839 ^f	α_1	41	4^+	13.600	(1970JO1C)
6.847 ^f	α_1	88	1^-	13.607	(1970JO1C)
6.894 ^f	α_1	297	5^-	13.649	(1970JO1C)
6.936 ^f	$\alpha_0, \alpha_1, \alpha_3$	60	4^+	13.686	(1968JO07, 1969JO09, 1970JO1C)
6.954 ^f	α_1	78	2^+	13.702	(1970JO1C)
7.025 ^f	α_1	206	3^-	13.765	(1969JO09, 1970JO1C)
7.067 ^f	α_1	132	4^+	13.802	(1970JO1C)
7.074 ^f	α_1	263	2^+	13.809	(1968JO07, 1970JO1C)
7.12	α_3, α_4	60		13.85	(1968JO07)
7.19	α_1	210		13.92	(1968JO07)
7.244 ^f	α_1	69	1^-	13.960	(1970JO1C)
7.27	α_1, α_3	150		13.99	(1968JO07)
7.286 ^f	α_1	168	5^-	13.997	(1970JO1C)
7.30	α_2	110		14.01	(1968JO07)
7.34	$\alpha_0, \alpha_1, \alpha_3, \alpha_4$	200		14.05	(1968JO07)
7.38	$\alpha_0, \alpha_1, \alpha_3$	210		14.08	(1968JO07)
7.400 ^f	α_1, α_3	170	2^+	14.098	(1968JO07, 1970JO1C)
7.406 ^f	α_1	73	4^+	14.103	(1970JO1C)
7.410 ^f	α_1	200	5^-	14.107	(1969JO09, 1970JO1C)
7.49	α_0	220		14.18	(1968JO07)
7.58	α_0	200		14.26	(1968JO07)
7.62	α_4	85		14.30	(1968JO07)
7.66	$\alpha_0, \alpha_2, \alpha_4$	130	$T = 0$	14.33	(1968JO07, 1970JO1C)
7.67	$\alpha_0, \alpha_2, \alpha_3, \alpha_4$	250	$T = 0$	14.34	(1968JO07, 1970JO1C)
7.73	α_1	145		14.39	(1968JO07)
7.74	α_3	235		14.40	(1968JO07)
7.80	α_0, α_4	70		14.46	(1968JO07)
7.82	α_0, α_2	225		14.48	(1968JO07)
7.832 ^f	α_1	203	1^-	14.482	(1968JO07, 1970JO1C)

Table 18.17: Maxima in the yield of $^{16}\text{O}+d$ reactions (continued)

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Res.
7.839 ^f	α_1	76	4^+	14.488	(1969JO09, 1970JO1C)
7.99	α_4	200		14.63	(1968JO07)
8.02	α_0	150		14.65	(1968JO07)
8.03	α_3	310		14.66	(1968JO07)
8.07	α_0, α_1	120		14.70	(1968JO07)
8.08	α_3, α_4	310		14.70	(1968JO07)
8.082 ^f	α_1	169		14.704	(1968JO07, 1970JO1C)
8.14	α_1	90		14.77	(1968JO07)
8.21	α_2	250		14.82	(1968JO07)
8.216 ^f	α_1	91	6^+	14.823	(1970JO1C)
8.228 ^f	α_1	124	2^+	14.833	(1970JO1C)
8.25	α_4	380		14.86	(1968JO07)
8.30	$\alpha_0, \alpha_2, \alpha_3$	210		14.90	(1968JO07)
8.308 ^f	α_1	304	4^+	14.904	(1970JO1C)
8.310 ^f	α_1	1201	2^+	14.906	(1970JO1C)
8.34	α_4	115		14.93	(1968JO07)
8.340 ^f	α_1	119	5^-	14.933	(1970JO1C)
8.37	α_0, α_1	130		14.96	(1968JO07)
8.37	α_0, α_3	250		14.96	(1968JO07)
8.385 ^f	α_1	491	5^-	14.973	(1970JO1C)
8.40	α_0	310		14.99	(1968JO07)
8.43	α_1, α_4	120		15.02	(1968JO07)
8.50	α_3, α_4	190		15.08	(1968JO07)
8.52	α_2	150		15.10	(1968JO07)
8.530 ^f	α_1	115	3^-	15.101	(1968JO07, 1970JO1C)
8.56	α_2	220		15.13	(1968JO07)
8.570 ^f	α_1	156	4^+	15.137	(1970JO1C)
8.58	α_4	180		15.15	(1968JO07)
8.61	α_0, α_3	200		15.17	(1968JO07)
8.616 ^f	α_1	299	3^-	15.178	(1970JO1C)
8.65	α_0, α_2	135		15.21	(1968JO07)
8.72	α_2, α_4	120		15.28	(1968JO07)
8.76	α_2	160		15.31	(1968JO07)
8.766 ^f	α_1	100	6^+	15.311	(1970JO1C)
8.79	α_0	200		15.34	(1968JO07)
8.82	$\alpha_0, \alpha_3, \alpha_4$	230		15.36	(1968JO07)
8.865 ^f	α_1	157	1^-	15.399	(1970JO1C)
8.89	α_3	110		15.43	(1968JO07)

Table 18.17: Maxima in the yield of $^{16}\text{O}+d$ reactions (continued)

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Res.
8.906 ^f	α_1	123	3^-	15.435	(1969JO09, 1970JO1C)
8.93	α_3, α_4	190		15.46	(1968JO07)
8.97	α_2, α_4	210		15.50	(1968JO07)
9.00	$\alpha_0, \alpha_1, \alpha_2$	190		15.52	(1968JO07)
9.032 ^f	α_1	319	4^+	15.547	(1970JO1C)
9.073 ^f	α_1	174	5^-	15.583	(1970JO1C)
9.200 ^f	α_1	174	6^+	15.696	(1970JO1C)
9.223 ^f	α_1	237	4^+	15.717	(1970JO1C)
9.255 ^f	α_1	249	1^-	15.745	(1970JO1C)
9.329 ^f	α_1	1044	5^-	15.811	(1970JO1C)
9.349 ^f	α_1	20	2^+	15.828	(1970JO1C)
9.403 ^f	α_1	272	3^-	15.876	(1970JO1C)
9.476 ^f	α_1	153	4^+	15.941	(1970JO1C)
9.643 ^f	α_1	103	2^+	16.089	(1970JO1C)
9.692 ^f	α_1	355	1^-	16.133	(1970JO1C)
9.748 ^f	α_1	149	4^+	16.183	(1970JO1C)
9.771 ^f	α_1	77	3^-	16.203	(1970JO1C)
9.781 ^f	α_1	181	1^-	16.212	(1970JO1C)
9.909 ^f	α_1	292	2^+	16.325	(1970JO1C)
10.049 ^f	α_1	52	4^+	16.450	(1969JO09, 1970JO1C)
10.333 ^f	α_1	296	3^-	16.703	(1970JO1C)
10.357 ^f	α_1	212	5^-	16.723	(1970JO1C)
10.406 ^f	α_1	135	4^+	16.767	(1970JO1C)
10.431 ^f	α_1	209	5^-	16.789	(1970JO1C)
10.457 ^f	α_1	86	6^+	16.812	(1970JO1C)
10.494 ^f	α_1	47	2^+	16.845	(1970JO1C)
10.499 ^f	α_1	78	1^-	16.849	(1970JO1C)
10.533 ^f	α_1	251	3^-	16.879	(1970JO1C)
10.728 ^f	α_1	180	5^-	17.052	(1970JO1C)
10.777 ^f	α_1	86	4^+	17.096	(1970JO1C)
10.888 ^f	α_1	88	6^+	17.195	(1970JO1C)
10.911 ^f	α_1	112	3^-	17.215	(1970JO1C)
11.332 ^f	α_1	208	4^+	17.580	(1970JO1C)
11.367 ^f	α_1	313	1^-	17.620	(1970JO1C)
11.567 ^f	α_1	78	2^+	17.797	(1970JO1C)
11.592 ^f	α_1	273	3^-	17.819	(1970JO1C)
11.704 ^f	α_1	155	6^+	17.919	(1970JO1C)
11.799 ^f	α_1	203	1^-	18.003	(1970JO1C)

Table 18.17: Maxima in the yield of $^{16}\text{O}+d$ reactions (continued)

E_d	Particles out	$\Gamma_{\text{c.m.}}$	$J^\pi; T$	E_x	Res.
11.869 ^f	α_1	223	7^-	18.065	(1970JO1C)
12.135 ^f	α_1	554	5^-	18.301	(1970JO1C)
12.495 ^f	α_1	395	7^-	18.621	(1970JO1C)
12.556 ^f	α_1	208	5^-	18.675	(1970JO1C)
12.682 ^{f,h}	α_1	1072	1^-	18.787	(1970JO1C)
12.951 ^{f,h}	α_1	1894	2^+	19.026	(1970JO1C)
12.990 ^f	α_1	416	4^+	19.060	(1970JO1C)
13.080 ^f	α_1	477	7^-	19.140	(1970JO1C)
13.307 ^f	α_1	911	6^+	19.342	(1970JO1C)
13.366 ^f	α_1	483	2^+	19.394	(1970JO1C)
14.35 ± 100	α_1	≈ 300		20.27	(1971JA04)
14.95 ± 100	α_1	≈ 550		20.80	(1971JA04)

^a Maxima at lower energies are reported by (1965LO03, 1968NG1B).

^b See also (1962CA20, 1965LO03, 1968NG1B, 1969DU11).

^c See also (1959LO59).

^d See also (1968MA53).

^e For this and the following levels, see also (1959AJ76).

^f Isospin mixed state: See (1970JO1C). Resonances in italics are definitely real: their influence on the S -matrix elements is certain and the ambiguities in the S -matrix elements do not allow the effect to be redistributed into other partial waves; and the interference effects are relatively small. All widths for states identified by the footnote ^f are uncertain to $\geq 10\%$ and the resonance energies are uncertain to $\geq 10\%$ of the widths (P. L. Jolivet, private communication).

^g These are the main components of the 2^+ strength from $3.4 < E_d < 4$ MeV. The resonances are not well defined and the shape can probably be built up from a different set of levels (1970JO1C).

^h Possibly should be broken up into several more levels but data are insufficient (1970JO1C).

ⁱ P.L. Jolivet, private communication.

The yield of elastically scattered deuterons has been measured for $E_d \leq 13.0$ MeV: see Table 18.16 (1963AM1A, 1963SE12, 1964AM1A, 1967AL06, 1968DI06, 1968MA53, 1969CO12, 1970DA14). Fluctuations are observed. Some of the maxima are interpreted in terms of ^{18}F states: for these see Table 18.17 (1956BE1B, 1963AM1A, 1964AM1A, 1968MA53). See also (1963DO1B, 1966AL09, 1970VE06, 1971GA1D, 1971SC1Q) and ^{16}O in (1971AJ02). For polarization measurements, see (1969CO12, 1970CO1P, 1971KO21) and Table 18.16.

$$16. \text{}^{16}\text{O}(d, \alpha)^{14}\text{N}$$

$$Q_m = 3.110$$

$$E_b = 7.527$$

The yields of various groups of α -particles have been measured for $E_d \leq 20$ MeV: see Table 18.16 (1960AM03, 1962CA20, 1963AM1A, 1963SE12, 1963YA1B, 1964AM1A, 1964KI1B, 1965LO03, 1965MA1A, 1965SA18, 1967TH1E, 1968DI06, 1968JO07, 1968TH1J, 1969CO12,

1969JO09, 1969JO1M, 1970AL1F, 1970JO1G, 1970JO1C, 1971JA04, 1971TH03). See also (1963DO1B, 1970CA1C) and (1969NO1C, 1971SC1Q; theor.).

The yield curves have been fitted in terms of a large number of states in ^{18}F : see Table 18.17 (1956BR36, 1957BA14, 1960AM03, 1963AM1A, 1964AM1A, 1965MA1A, 1968JO07, 1969JO09, 1969JO1C, 1970JO1C, 1971JO11). See also (1970JO1F). A detailed study by (1970JO1C) of the isospin-forbidden α_1 yield shows that there are no very strong states seen for $J > 5$, and those seen in $J = 7$, the highest partial wave needed to fit the yield, are wide, $\Gamma \approx 300$ keV. This contradicts the prediction that isospin violation should be negligible for low spin states while large for those with high J . See also (1971JO11). The average coherence width is ≈ 150 keV for $12 < E_x < 15.5$ MeV (1968JO07).

For the isospin-forbidden α_1 yield, virtually all the observed levels overlap several others of the same spin and parity in the tail of the resonance. The α_1 yield has a smaller cross section than the yield of $^{14}\text{N}(\alpha, \alpha_1)^{14}\text{N}$ at the same E_x in ^{18}F [see (1970TO03)]. This is most pronounced at high excitation energy and is thought to be related to the very small binding energy of the deuteron. Thus reaction 8 appears to be a better tool to study ^{18}F at high E_x (1970JO1C).

For polarization studies, see (1970PR1C, 1971KE1E). See also ^{14}N . Very accurate cross section measurements of the $^{16}\text{O}(d, \alpha)^{14}\text{N}$ reaction and of its inverse, $^{14}\text{N}(\alpha, d)^{16}\text{O}$, are consistent with the principle of detailed balance. The lowest uncertainty was $\pm 0.5\%$. An upper limit of 0.2% is assigned to the time-reversal non-invariant part of the reaction amplitudes (1971TH03).

$$17. \ ^{16}\text{O}(t, n)^{18}\text{F} \qquad Q_m = 1.269$$

Measurements of the strengths and of lifetimes of the radiative decays are displayed in Tables 18.14 and 18.15 (1963LI07, 1966AL04). (1966AL04) report that ΔE_x ($1.13 \rightarrow 0.94$) = 194 ± 1 keV: E_x of $^{18}\text{F}^*(1.13)$ is then 1131.0 ± 1 keV [based on $E_x = 937.0 \pm 0.2$ keV]. See also (1967WA1C) and (1959AJ76).

$$18. \ ^{16}\text{O}(^3\text{He}, p)^{18}\text{F} \qquad Q_m = 2.033$$

Excitation energies derived from measurements of proton spectra (1959HI67, 1959YO25, 1967MA1G, 1968GR1G, 1968GR1H, 1968MA33) and of γ -rays (1960RA18, 1961DU02, 1965CH10, 1967WA06) are displayed in Table 18.18.

Angular distributions of proton groups have been obtained at $E(^3\text{He}) = 4.00$ MeV (1964MA50), 5.9 and 9.2 MeV (1959HI74), 15 MeV (1968PO1B, 1969PO11), 18 MeV (1967PU03, 1971BE19) and 19.8 MeV (1967MA1G, 1968MA33). Distorted wave analyses lead to the l -assignments shown in Table 18.18. (1960JA11) have compared the angular distribution they obtained for the $^{16}\text{O}(t, p)^{18}\text{O}$ reaction to $^{18}\text{O}(0)$ with the angular distributions obtained by (1959HI74) for $^{18}\text{F}^*(1.04, 1.08)$ in this reaction. It is clearly $^{18}\text{F}^*(1.04)$ which is the analog to the ground state of ^{18}O : J^π is then 0^+ and $T = 1$.

Table 18.18: States of ^{18}F from $^{16}\text{O}(^3\text{He}, p)^{18}\text{F}$

E_x (MeV \pm keV)							l^f	$J^\pi; T^g$
(1959HI67) ^a	(1959YO25) ^a	(1960RA18, 1961DU02) ^c	(1965CH10) ^c	(1967WA06) ^c	(1968GR1G, 1968GR1H) ^a	(1967MA1G, 1968MA33) ^a		
0	0					0	0	1 ⁺ ; 0
0.934 \pm 10	0.943 \pm 7	0.939 \pm 5	0.9374 \pm 1.5	0.9370 \pm 1	e	0.937 \pm 8	2	3 ⁺ ; 0
1.038 \pm 10	1.047 \pm 7	1.041 \pm 5	1.0446 \pm 1.5	1.0413 \pm 1.5			0	0 ⁺ ; 1
1.076 \pm 10 ^b	1.089 \pm 7	1.070 \pm 20	1.0817 \pm 1.5	1.0803 \pm 1				0 ⁻ ; 0
1.119 \pm 10	1.128 \pm 7	1.170 \pm 10 ^d	(1.1197 \pm 2)		1.120 \pm 5	1.111 \pm 7	4	5 ⁺ ; 0
1.698 \pm 10	1.708 \pm 7	1.680 \pm 20	1.7035 \pm 2	1.7003 \pm 1.5	e	1.680 \pm 24	0	1 ⁺ ; 0
2.096 \pm 10	2.102 \pm 7	2.090 \pm 10	2.1026 \pm 2	2.1005 \pm 1.5	e	2.096 \pm 13		2 ⁻ ; 0
2.517 \pm 10	2.521 \pm 10	2.510 \pm 10	2.5297 \pm 2	2.5235 \pm 1.5	e	2.509 \pm 18	2	2 ⁺ ; 0
3.055 \pm 10	3.058 \pm 10	3.060 \pm 50		3.0603 \pm 3		3.062 \pm 15	2	2 ⁺ ; 1
3.128 \pm 10	3.130 \pm 10	3.110 \pm 50		3.1339 \pm 3				1 ⁻ ; 0
3.352 \pm 10	3.355 \pm 10	3.350 \pm 100	(3.3505 \pm 3)	3.3581 \pm 3		3.352 \pm 16		(3) ⁺
3.715 \pm 10	3.724 \pm 10			3.7248 \pm 3				1 ⁺ ; 0
3.783 \pm 10								(3 ⁻); 0
3.830 \pm 10	3.843 \pm 10	3.840 \pm 100		3.8385 \pm 3.5	e	3.830 \pm 12	2	2 ⁺ ; 0
4.108 \pm 10					4.120 \pm 5	4.134 \pm 11		\leq 3; 0
4.218 \pm 10					4.232 \pm 5			(2)
4.350 \pm 15					4.362 \pm 5	4.378 \pm 9		2, 3
					4.403 \pm 5			\geq 2; 0
					4.659 \pm 5	4.651 \pm 12	4	4 ⁺ ; 1
					4.739 \pm 5			0 ⁺ ; 1
					4.852 \pm 5	4.843 \pm 12		1; 0
					4.955 \pm 5	4.967 \pm 21		2 ⁺ ; 1
						5.297 \pm 25		T = 0
						5.601 \pm 12		(4 ⁺); 0
						6.105 \pm 8		
						6.265 \pm 13		(1 ⁺)
						6.779 \pm 7		2 ⁻ ; 0
						7.206 \pm 9		(1 ⁺)
						7.646 \pm 14		T = 0
						7.874 \pm 22		(2 ⁻); 0
						9.145 \pm 32		
						9.404 \pm 31		
						9.82 \pm 40		
						10.06 \pm 45		
						10.352 \pm 25		
						10.96 \pm 60		
						11.13 \pm 50		

^a From measurements of proton groups.^b (1958KU81) report $E_x = 1.080 \pm 0.010$ MeV.^c From measurements of γ -rays.^d The transition 1.13 \rightarrow 0.94 is observed: $E_\gamma = 189 \pm 4$ keV (1960RA18).^e Observed but E_x not determined.^f (1959HI74, 1967PU03, 1968PO1B, 1969PO11).^g See discussion in (1968MA33) and Table 18.14.

The magnetic moment of $^{18}\text{F}^*(1.13)$, $\mu = +(0.568 \pm 0.013)J$ (1967PO09), $+(0.572 \pm 0.006)J$ (1967SC09). If $J = 5$, and all data are consistent with this assignment, $\mu = 2.855 \pm 0.030$ nm. This value is in agreement with shell-model predictions for a 5^+ state remaining from $1d_{5/2}^2$ (1967PO09).

The γ -decay of many states has been studied: Table 18.14 displays observed branching ratios and radiative widths. Studies of these parameters, coupled with angular correlation and γ -ray angular distribution studies (1961KU02, 1965PO01, 1966CH06, 1966CH12, 1966OL03, 1967GO07, 1967OL03, 1967PO02, 1967WA06), lifetime measurements [see Table 18.15 (1959AL99, 1963LO03, 1966OL03, 1967BE14, 1967PO09, 1967WA06, 1970BL1F)] and the l -values shown in Table 18.18 lead to the J^π assignments displayed in Table 18.14. See also (1963HI06).

The linear polarization of the 2.10 MeV γ -ray [$2.10 \rightarrow 0$], together with the mixing ratio obtained by (1967GO07), leads to an assignment of negative parity to $^{18}\text{F}^*(2.10)$, and also to $^{18}\text{F}^*(1.08)$. The latter results from the value of τ_m for $^{18}\text{F}^*(2.10)$ and the observation of a $P_4(\cos \theta)$ term in the $2.10 \rightarrow 1.08$ transition (1967PO02).

See also (1964MA57, 1966AG1B, 1970CA28), (1959FA1A) and (1960NE1A; theor.).

$$19. \ ^{16}\text{O}(\alpha, d)^{18}\text{F} \quad Q_m = -16.321$$

At $E_\alpha = 40$ to 52 MeV, deuteron spectra are dominated by the groups to $^{18}\text{F}^*(1.13)$, $J^\pi = 5^+$ [$1d_{5/2}^2$ configuration] (1962HA40, 1966RI04, 1967MA1G, 1968MA33). Many other states of ^{18}F have also been observed: see Table 18.19 (1966RI04, 1967MA1G, 1968MA33). For configuration assignments, see (1968MA33). See also (1960AG01), (1969BR1D) and (1963GL1C, 1965GR1F, 1967IM1A; theor.).

$$20. \ ^{16}\text{O}(^6\text{Li}, \alpha)^{18}\text{F} \quad Q_m = 6.053$$

Angular distributions of the α -particles to the ground state of ^{18}F have been measured at $E(^6\text{Li}) = 5.5$ to 13.3 MeV (1968GR1H, 1968GR22) and at 26 MeV (1969DA19). The lifetime of $^{18}\text{F}^*(4.36)$ is < 0.61 psec (1969TH01): the γ -ray energy for the transition $4.36 \rightarrow 3.06$, $E_\gamma = 1297.4 \pm 2.5$ keV. Assuming $E_x = 3.0598$, the energy of $^{18}\text{F}^*(4.36)$, $E_x = 4.357 \pm 0.004$ MeV (1969TH01). See also (1967CA1D).

$$21. \ ^{16}\text{O}(^{11}\text{B}, ^9\text{Be})^{18}\text{F} \quad Q_m = -8.290$$

See (1966PO1E, 1967PO1E).

$$22. \ ^{16}\text{O}(^{14}\text{N}, ^{12}\text{C})^{18}\text{F} \quad Q_m = -2.746$$

See (1966GA10, 1969BR1D, 1969RO1G, 1970AN1D).

Table 18.19: States of ^{18}F from $^{16}\text{O}(\alpha, d)^{18}\text{F}$ ^a

E_x (MeV \pm keV)	$J^\pi; T$
0.0	$1^+; 0$
0.934 ± 14	$3^+; 0$
1.119 ± 16	$5^+; 0$
1.716 ± 18	$1^+; 0$
2.100 ± 11	$2^-; 0$
2.541 ± 19	$2^+; 0$
3.122 ± 14	$1^-; 0$
3.363 ± 20	$(3)^+; 0$
3.808 ± 12	$2^+; 0$
4.140 ± 12	$\leq 3; 0$
4.393 ± 9	$\geq 2; 0$
4.852 ± 10	$1; 0$
5.266 ± 34	$T = 0$
5.590 ± 27	$(4^+); 0$
5.785 ± 31	$T = 0$
6.139 ± 12	$T = 0$
6.548 ± 18	$(\leq 5^+); 0$
6.807 ± 10	$2^-; 0$
7.191 ± 8	$(4^+); 0$
7.434 ± 13	$(3^-); (0)$
7.658 ± 12	$(T = 0)$
7.871 ± 11	$(2^-); 0$
8.596 ± 19	$T = 0$
8.861 ± 190	$(T = 0)$
9.494 ± 15	$T = 0$ ^b
9.96 ± 120	$T = 0$
10.268 ± 12	
10.541 ± 10	$(T = 0)$
11.073 ± 37	$T = 0$
11.384 ± 18	$T = 0$
12.055 ± 16	$T = 0$

Table 18.19: States of ^{18}F from $^{16}\text{O}(\alpha, d)^{18}\text{F}$ ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$
12.67 ± 60	$T = 0$

^a (1967MA1G, 1968MA33). See also (1962HA40, 1966RI04).

^b $J^\pi = (6^-)$ (1966RI04).

23. $^{17}\text{O}(\text{p}, \gamma)^{18}\text{F}$ $Q_m = 5.609$

The yield of 0.94 MeV γ -rays (γ_1) has been measured for $E_p = 0.2$ to 1.5 MeV. A number of resonances are observed: see Table 18.20 (1969ZA1C, 1971BE1E, 1971SE1H). A strong resonance for 1.04 MeV γ -rays (γ_2) is observed, corresponding to $^{18}\text{F}^*(6.14)$ (1969ZA1C, 1971SE1H). A study of this resonance shows that $^{18}\text{F}^*(6.14)$ has $J^\pi = 0^+$. The parity of $^{18}\text{F}^*(3.72)$ is found to be even (1970RO1F).

Branching ratios for the decay of $^{18}\text{F}^*$ states with $6.09 < E_x < 6.87$ MeV are displayed in Table 18.15. DSAM measurements show $\tau_m < 15$ fsec for these states. From the γ -decay of the resonant states, accurate excitation energies have been determined for a number of low-lying states: $E_x = 0.9369 \pm 0.2, 1.041 \pm 1, 1.121 \pm 0.4, 1.700 \pm 1, 2.1013 \pm 0.5, 2.523 \pm 1, 3.136 \pm 2, 4.650 \pm 1$ and 4.965 ± 1 MeV (\pm keV) (1971SE1H).

24. $^{17}\text{O}(\text{p}, \text{n})^{17}\text{F}$ $Q_m = -3.542$ $E_b = 5.609$

The yields of n_0 , measured for $E_p = 7$ to 13.5 MeV, and of n_1 , measured for $E_p = 7$ to 12.5 MeV, show some gross structures (1969AN06). See also (1965BL07).

25. $^{17}\text{O}(\text{p}, \text{p})^{17}\text{O}$ $E_b = 5.609$

The elastic scattering has been studied for $E_p = 0.5$ to 1.33 MeV (1971SE1J) and for $E_p = 11.0$ to 13.0 MeV (1967AL06): observed anomalies are displayed in Table 18.20 (1971SE1J).

26. $^{17}\text{O}(\text{p}, \alpha)^{14}\text{N}$ $Q_m = 1.193$ $E_b = 5.609$

The yield of ground state α -particles shows a number of resonances for $E_p = 0.49$ to 3.0 MeV: see Table 18.20 (1957AH20, 1962BR08, 1971SE1J). Astrophysical considerations are discussed by (1962BR08, 1967FO1B).

Table 18.20: Resonances in $^{17}\text{O} + \text{p}$

E_p (MeV \pm keV)	Yield of	$\Gamma_{\text{c.m.}}$ (keV)	$J^\pi; T$	E_x (MeV)	Refs.
0.515 \pm 2	γ_1, α_0	< 2.0	4 ⁻ ; 0	6.095	(1962BR08, 1969ZA1C, 1971BE1E, 1971SE1H)
0.557 \pm 2	γ_2		0 ⁺ ; (1)	6.135	(1969ZA1C, 1970RO1F, 1971SE1H, 1971SE1J)
0.585 \pm 1	γ_1, p_0	15	3 ⁺ ; 1	6.161	(1969ZA1C, 1971SE1H, 1971SE1J)
0.668 \pm 2	$\gamma_1, \alpha_0, \text{p}_0$	< 2.0	3 ⁻ ^a	6.240	(1962BR08, 1969ZA1C, 1971SE1H, 1971SE1J)
0.690	α_0			6.260	(1971SE1J)
0.711 \pm 1	γ_1, p_0	7.5	2 ⁺ ; 1	6.280	(1969ZA1C, 1971SE1H, 1971SE1J)
0.742 \pm 2	$\gamma_1, \alpha_0, \text{p}_0$	3.1 \pm 1.4	2 ⁺ , 3 ⁺	6.309	(1962BR08, 1969ZA1C, 1971SE1H, 1971SE1J)
0.822 \pm 3	γ_1, α_0	< 4.5	(1 ⁺), 2, 3 ⁻	6.385	(1962BR08, 1969ZA1C, 1971SE1H, 1971SE1J)
0.926 \pm 3	γ_1, α_0	< 1.2	(1 ⁺), 2, 3 ⁻	6.483	(1962BR08, 1969ZA1C, 1971SE1H, 1971SE1J)
1.096 \pm 6	α_0	85 \pm 5		6.644	(1962BR08)
1.100 \pm 1	$\gamma_1, \alpha_0, \text{p}_0$	< 3.0	(2 ⁻) ^a	6.647	(1957AH20, 1962BR08, 1969ZA1C, 1971SE1H, 1971SE1J)
1.245 \pm 1	$\gamma_1, \alpha_0, \text{p}_0, \text{p}_1$	10 \pm 3	4 ⁺ , 5 ⁺ ; 0	6.780	(1962BR08, 1969ZA1C, 1971SE1H, 1971SE1J)
1.270 \pm 3	$\gamma_1, \alpha_0, \text{p}_0, \text{p}_1$	5 \pm 3	2 ⁺ , 3 ⁺	6.808	(1969ZA1C, 1971SE1H, 1971SE1J)
1.270 \pm 5	α_0	79 \pm 5		6.808	(1957AH20, 1962BR08, 1971SE1J)
1.337 \pm 2	$\gamma_1, \alpha_0, \text{p}_1$			6.871	(1962BR08, 1971SE1H)
1.786	α_0	\approx 65		7.295	(1957AH20)
2.021	α_0	11		7.517	(1957AH20)
2.048	α_0	90		7.542	(1957AH20)
2.218	α_0	11		7.703	(1957AH20)
2.235	α_0	100		7.719	(1957AH20)
2.406	α_0	\approx 25		7.880	(1957AH20)
2.435	α_0	\approx 25		7.908	(1957AH20)
2.623	α_0	\approx 40		8.085	(1957AH20)
2.753	α_0	\approx 15		8.208	(1957AH20)
2.775	α_0	\approx 10		8.228	(1957AH20)
2.928	α_0	\approx 50		8.373	(1957AH20)

^a Mixed $T = 0$ and 1.

27. $^{17}\text{O}(\text{d}, \text{n})^{18}\text{F}$ $Q_{\text{m}} = 3.384$

At $E_{\text{d}} = 4.4$ MeV angular distributions of the neutrons corresponding to $^{18}\text{F}^*(0.94, 2.52, 3.06, 3.36, 3.84, 4.96, 5.61, 5.67)$ have been measured. The cross section for formation of $^{18}\text{F}^*(5.61)$ is an order of magnitude greater than for $^{18}\text{F}^*(5.67)$, consistent with the interpretation that these (1^-) states contain a substantial isospin admixture (1971GA1B, 1971LI1D). See also (1971AR33). For a lifetime measurement, see Table 18.15 (1971LE29). See also ^{19}F .

28. $^{17}\text{O}(^3\text{He}, \text{d})^{18}\text{F}$ $Q_{\text{m}} = 0.115$

At $E(^3\text{He}) = 15$ MeV, DWBA analysis of angular distributions of deuteron groups corresponding to the ground state of ^{18}F [$l = 2$] and to the excited states of 0.94 [$l = 0 + 2$], 1.04 [2], 1.12 [2], 2.53 [$0 + 2$], 3.06 [$0 + 2$], 3.84 [$0 + 2$], 4.12 [$0 + 2$], 4.66 [2] and 4.96 MeV [$0 + 2$] have been obtained by (1968PO1B, 1969PO11) who also report spectroscopic information. Thus all these states have even parity and $^{18}\text{F}^*(4.12)$ may be assigned $J^\pi = (2^+)$ or 3^+ . Since $l = 2$ for $^{18}\text{F}^*(4.66)$, $J^\pi \leq 5^+$, with 4^+ most likely (1969PO11).

29. $^{17}\text{O}(\alpha, \text{t})^{18}\text{F}$ $Q_{\text{m}} = -14.205$

Not reported.

30. $^{18}\text{O}(\text{p}, \text{n})^{18}\text{F}$ $Q_{\text{m}} = -2.438$

$$E_{\text{thresh.}} = 2.57372 \pm 0.00077 \text{ (1964BO13);}$$

$$E_{\text{thresh.}} = 2.577 \pm 0.003 \text{ (1967PR04);}$$

$$E_{\text{thresh.}} = 2.576 \pm 0.001, Q_0 = -2.439 \pm 0.001 \text{ (1969BE57). See also (1959AJ76).}$$

Gamma rays are observed with $E_\gamma = 938 \pm 6, 1043 \pm 8$ and 1082 ± 10 keV (1960RA18). See also (1968BE34). The lifetime of $^{18}\text{F}^*(1.04)$ is 4_{-2}^{+3} fsec (1967BL18). Angular distributions have been measured at $E_{\text{p}} = 6.9$ to 13.5 MeV ($n_0, n_{1 \rightarrow 4}$), 6.9 to 9.25 MeV (n_5), 7.55 to 9.25 MeV (n_6), 7.95 to 9.25 MeV (n_7) and 8.4 to 13.5 MeV (n_8) (1969AN06). See also (1965BL07, 1967WI1H, 1971TH1D) and (1969SC1H; theor.). See also ^{19}F .

31. $^{18}\text{O}(^3\text{He}, \text{t})^{18}\text{F}$ $Q_{\text{m}} = -1.674$

Table 18.21: Branching in $^{18}\text{Ne}(\beta^+)^{18}\text{F}$

Decay to $^{18}\text{F}^*$	E_γ (keV)	Branch (%)	$\log ft^a$	Refs.
g.s.		92.5 ± 0.2		(1970AS06)
1.04	1041 ± 5	7 ± 2		(1961BU05)
	1035 ± 5			(1961EC02)
	1030 ± 20	9 ± 3		(1963FR10, 1964FR1B, 1965FR09)
	1043 ± 1			(1968GO05)
		7.3 ± 0.2		(1970AS06)
1.08	1042.6 ± 1.0	7.3 ± 0.2		mean value (1968GO05)
1.70		< 0.7		(1970AS06)
2.10		0.17 ± 0.05		(1970AS06)
		< 1.5		(1968GO05)

^a Based on Q_m and $\tau_{1/2} = 1.67 \pm 0.02$ sec: see Table 18.24.

At $E(^3\text{He}) = 16$ MeV, the triton spectrum is dominated by strong groups to the ground and 0.94 MeV excited states and to the 0^+ and 2^+ ; $T = 1$ states at $E_x = 1.04$ and 3.06 MeV. Angular distributions have been measured and analyzed by DWBA for the tritons corresponding to these states and to $^{18}\text{F}^*(1.08, 1.13, 1.70, 2.10, 3.13, 3.36, 3.72, 3.79, 3.84, 4.12, 4.23, 4.36, 4.40, 4.66, 4.74)$. The angular distributions are consistent with the J^π assignments shown in Table 18.14, except for the distribution to $^{18}\text{F}^*(1.04)$ (1970DU08). At $E(^3\text{He}) = 17.3$ MeV, angular distributions to ^{18}F states with $E_x < 4$ MeV have been analyzed using DWBA and a two-body interaction between the incident and target nucleons. An exact coupled-channel calculation was also made for the transition to $^{18}\text{F}^*(1.04)$ (1968HA30, 1969HA1U, 1969MA1G). See also (1971HE1F).

$$32. \ ^{18}\text{Ne}(\beta^+)^{18}\text{F} \quad Q_m = 4.447$$

The decay is to $^{18}\text{F}^*(0, 1.04, 1.70)$: see Table 18.21 for the branching ratios and $\log ft$ values. See also ^{18}Ne .

$$33. \ ^{19}\text{F}(\gamma, n)^{18}\text{F} \quad Q_m = -10.431$$

See ^{19}F .

34. $^{19}\text{F}(\text{n}, 2\text{n})^{18}\text{F}$ $Q_{\text{m}} = -10.431$

See ^{20}F .

35. (a) $^{19}\text{F}(\text{p}, \text{d})^{18}\text{F}$ $Q_{\text{m}} = -8.206$

(b) $^{19}\text{F}(\text{p}, \text{pn})^{18}\text{F}$ $Q_{\text{m}} = -10.431$

Angular distributions have been measured of the ground state deuterons at $E_{\text{p}} = 16$ and 18 (1967AN1B, 1968AN1A, 1971AN1B), 17.5 (1969HA1T), 18 (1956RE04), 18.6 (1961BE12), 30.3 (1967DI1C) and 155.6 MeV (1966BA44): $l_{\text{n}} = 0$. (1969HA1T) also reports angular distributions at $E_{\text{p}} = 17.5$ MeV to the ^{18}F states at $E_{\text{x}} = 0.94$ [$l_{\text{n}} = 2$], 1.04 [0], 1.08 [1], 1.13, 1.70 [0], 2.10 [(1)], 2.53 [2], 3.06 [2], 3.13 [1], 3.36 [2], 3.8 (unresolved), 4.12, 4.23, 4.36 + 4.40, 4.66 and 5.6 MeV. Spectroscopic factors are also listed. Similar results are reported by (1967AN1B, 1968AN1A, 1971AN1B). See also (1961BE12, 1966BA44). See also (1961LE1A, 1962CO17) and (1969BE1T, 1969DO08; theor.). For reaction (b), see (1968DE21).

36. $^{19}\text{F}(\text{d}, \text{t})^{18}\text{F}$ $Q_{\text{m}} = -4.173$

Angular distributions of triton groups are reported at $E_{\text{d}} = 8.9$ MeV (1957EL12; t_0, t_1, t_3) and 14.8 MeV (1959HA1E, 1960HA22; t_0). See also (1959VL23) and (1963DA1B, 1963OG1A, 1964DA1D, 1964EL1B; theor.).

37. $^{19}\text{F}(^3\text{He}, \alpha)^{18}\text{F}$ $Q_{\text{m}} = 10.147$

At $E(^3\text{He}) = 5.9$ MeV, 41 α -particle groups have been observed, corresponding to the ground state of ^{18}F and to excited states with $E_{\text{x}} < 7.5$ MeV (1959HI67): see Table 18.22. Angular distributions of the α -particles corresponding to $^{18}\text{F}^*(3.06, 3.13)$ are reported by (1966MA43; $E(^3\text{He}) = 4.0, 6.0$ and 8.0 MeV): $l = 2$ and 1, respectively. See also (1964BR1G, 1966HA21, 1966HO1E, 1971BA2A).

Alpha- α_0 angular correlations measured in the range $E(^3\text{He}) = 5.0$ to 6.0 MeV establish $J^{\pi} = 1^-$ for both $^{18}\text{F}^*(5.61, 5.67)$: $\Gamma_{\alpha}/\Gamma \approx 1$ (1971LI27).

38. $^{19}\text{F}(^{10}\text{B}, ^{11}\text{B})^{18}\text{F}$ $Q_{\text{m}} = 1.025$

See (1968GA03, 1970GO1B, 1971KN05).

Table 18.22: Energy levels of ^{18}F from $^{19}\text{F}(^3\text{He}, \alpha)^{18}\text{F}$

E_x^a (MeV \pm keV)	E_x^a (MeV \pm keV)
0	4.965 ± 13
0.940 ± 10	5.292 ± 10
1.042 ± 10	5.500 ± 10
1.087 ± 10	5.603 ± 13
1.129 ± 10	5.666 ± 10
1.699 ± 10	5.785 ± 10
2.105 ± 10	6.093 ± 10
2.525 ± 10	6.137 ± 10
3.063 ± 10	6.232 ± 10
3.131 ± 10	6.264 ± 13
3.352 ± 10	6.374 ± 10
3.727 ± 10	6.470 ± 10
3.790 ± 10	6.551 ± 10
3.841 ± 10	6.633 ± 10
4.116 ± 10	6.765 ± 10
4.227 ± 10	6.790 ± 10
4.358 ± 10	6.857 ± 10
4.400 ± 10	7.183 ± 10
4.649 ± 10	7.313 ± 10
4.741 ± 10	7.495 ± 10
4.840 ± 10	

^a (1959HI67): $E(^3\text{He}) = 5.9$ MeV.

39. (a) $^{19}\text{F}(^{14}\text{N}, ^{15}\text{N})^{18}\text{F}$ $Q_m = 0.403$
 (b) $^{19}\text{F}(^{19}\text{F}, ^{20}\text{F})^{18}\text{F}$ $Q_m = -3.829$

For reaction (a) see ([1965GA1B](#), [1971GA13](#)); for reaction (b) see ([1971GA13](#)).

40. $^{20}\text{Ne}(n, t)^{18}\text{F}$ $Q_m = -14.793$

Not reported.

41. $^{20}\text{Ne}(p, ^3\text{He})^{18}\text{F}$ $Q_m = -15.557$

At $E_p = 45$ MeV, ^3He groups are observed to $^{18}\text{F}^*(0, 1.04, 1.70, 3.06, 6.27 \pm 0.03)$ ([1969HA38](#)). See also ([1968FA1G](#), [1970OL1B](#)).

42. $^{20}\text{Ne}(d, \alpha)^{18}\text{F}$ $Q_m = 2.796$

At $E_d = 11$ MeV α -groups are observed to many states of ^{18}F with $E_x < 7$ MeV. Weak or absent (each $\leq 0.3\%$ of the total yield at 30°) are the groups corresponding to $^{18}\text{F}^*(1.04, 3.06, 4.66, 4.74, 4.96)$: $T = 1$ for these states ([1968PO1B](#), [1969PO11](#)). However, ([1971HR1A](#)) report a significant yield (as high as 30% of the yield to the $J^\pi = 2^+$; $T = 0$ state at 2.52 MeV) for the 2^+ ; 1 state at 3.06 MeV in the range $E_d = 6$ to 10 MeV. The yield is particularly large in the range $E_d = 6$ to 8 MeV. Angular distributions are reported at 2 MeV ([1966LA15](#); $\alpha_0, \alpha_1, \alpha_2, \alpha_5, \alpha_6$), 4 MeV ([1964MA50](#); $\alpha_0, \alpha_5, \alpha_6, \alpha_7, \alpha_8, \alpha_9, \alpha_{10}$), 6.75, 7.37 and 8 MeV ([1971HR1A](#); $\alpha_7, \alpha_8, \alpha_9$), and 14.7 MeV ([1962TA07](#); $\alpha_0, \alpha_{1 \rightarrow 4}, \alpha_5, \alpha_6, \alpha_7$). See also ([1951MI1A](#), [1961LO10](#), [1964MA57](#)) and ([1966BR1G](#), [1966KU1D](#), [1969DE29](#)).

43. $^{21}\text{Ne}(p, \alpha)^{18}\text{F}$ $Q_m = -1.740$

Not reported.

¹⁸Ne
(Figs. 3 and 4)

GENERAL:

Shell and cluster model calculations: (1957WI1E, 1969BE1T, 1970BA2E, 1970EL08, 1970HA49, 1972KA01).

Electromagnetic transitions: (1970EL08, 1970HA49).

Special levels: (1966MI1G, 1969KA29, 1972KA01).

Pion reactions: (1965PA1F).

Other theoretical calculations: (1965GO1F, 1966KE16, 1968BA2H, 1968BE1V, 1968MU1B, 1968NE1C, 1968VA1J, 1968VA24, 1969BA1Z, 1969GA1G, 1969KA29, 1969MU09, 1969RA28, 1969SO08, 1970BA1Z, 1970DI1G, 1970EL08, 1972KA01).

1. ¹⁸Ne(β^+)¹⁸F $Q_m = 4.447$

The half-life of ¹⁸Ne is 1.67 ± 0.02 sec: see Table 18.24 (1970AL11, 1970AS06). The decay is to ¹⁸F*(0, 1.04, 1.70) with $J^\pi = (1^+)$; $T = (0), (0^+; 1), (1^+; 0)$, respectively: the branching ratios and log ft values are displayed in Table 18.21. The ratio of the ft values for the ¹⁸Ne \rightarrow ¹⁸F(0) and ¹⁸F \rightarrow ¹⁸O transitions is 0.992 ± 0.015 (1970AL11): see also (1969KA1B, 1970AS06). See also (1966ZA03, 1968BO1U, 1968FR03, 1970MC23, 1971BL12, 1971DE1E).

2. ¹⁶O(³He, n)¹⁸Ne $Q_m = -3.196$
 $Q_0 = -3.206 \pm 0.013$ (1961DU02);
 $Q_0 = -3.199 \pm 0.006$ (1961TO03).

Excitation energies of ¹⁸Ne states derived from neutron spectra and γ -ray measurements are displayed in Table 18.25 (1961TO03, 1968GI09, 1969RO08, 1970AD02, 1970NE1J, 1970NE1N, 1970SH04). Branching ratios and lifetimes of the first four excited states of ¹⁸Ne are shown in Table 18.26 (1968GI09, 1969RO08, 1969RO22, 1970SH04, 1971RO18, 1972GI01). Table 18.27 summarizes the neutron angular distribution studies. Angular correlation studies, together with PWBA and DWBA double stripping analysis of neutron angular distributions, and the branching ratio and lifetime measurements, lead to assignments of $J^\pi = 0^+, 2^+, 4^+, 0^+$ and 2^+ for ¹⁸Ne*(0, 1.89, 3.38, 3.58, 3.62), respectively (1968GI09, 1968TO09, 1969RO08, 1969RO22, 1970AD02, 1970SH04, 1971RO18, 1972GI01). It is also suggested that ¹⁸Ne*(4.58) has $J^\pi = 0^+$ and a predominantly two-particle structure (1970NE1J). See also (1961BU05, 1961EC02, 1963FR10, 1964BR13, 1964FR1B, 1965BR1H, 1965FR09, 1968GO05, 1970AL11, 1970AS06), (1969BA1Z) and (1964HE06; theor.).

Table 18.23: Energy levels of ^{18}Ne ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$0^+; 1$	$\tau_{1/2} = 1.67 \pm 0.02$ sec	β^+	1, 2, 3
1.8873 ± 0.2	2^+	$\tau_m = 0.49^{+0.17}_{-0.09}$ psec	γ	2, 3
3.3762 ± 0.4	4^+	4.4 ± 0.6 psec	γ	2, 3
3.576 ± 2	0^+	$2 < \tau_m < 6$ psec	γ	2, 3
3.6164 ± 0.6	2^+	$0.063^{+0.030}_{-0.020}$ psec	γ	2, 3
4.510 ± 10	(1^-)	$\Gamma \leq 40$		2, 3
4.580 ± 10	(0^+)	≤ 40		2, 3
5.075 ± 13		≤ 60		2, 3
5.135 ± 12		≤ 60		2, 3
6.30 ± 20		180 ± 60		2, 3
7.062 ± 12	$(2^+, 1^-)$	180 ± 50		2
7.712 ± 20		≤ 50		2
7.915 ± 12	$(2^+, 1^-)$	≤ 50		2, 3
8.100 ± 14		≤ 50		2, 3
8.50 ± 30		≤ 120		2
9.20 ± 20				3

^a See also Table 18.26.

Table 18.24: The half-life of ^{18}Ne

$\tau_{1/2}$ (sec)	Refs.
1.6 ± 0.2	(1954GO17)
1.7 ± 0.4	(1961EC02)
1.46 ± 0.07	(1961BU05)
1.47 ± 0.10	(1963FR10, 1964FR1B, 1965FR09)
1.67 ± 0.02	(1970AL11)
1.69 ± 0.04	(1970AS06)
1.67 ± 0.02	Weighted average of last two values

Table 18.25: Excitation energies from $^{16}\text{O}(^3\text{He}, n)^{18}\text{Ne}$

E_x (MeV \pm keV)					Γ (keV) ^d	J^π ^d
(1961TO03)	(1968GI09, 1969RO08)	(1970AD02)	(1970NE1J, 1970NE1N) ^d	(1970SH04)	c.m.	
0	0	0		0		
1.880 ± 10	1.8873 ± 0.2	^a		1.890 ± 2		
3.362 ± 11	3.3762 ± 0.4	3.375 ± 15		3.383 ± 4		
	3.5763 ± 2.0	3.564 ± 20				
3.608 ± 12	3.6164 ± 0.6	3.610 ± 15		3.623 ± 3		
		4.505 ± 15 ^b	4.513 ± 13			
		4.571 ± 15 ^b	4.587 ± 13 ^e			$0^+(1^-)$
			5.075 ± 13 ^f		≤ 60	
		5.14 ± 20 ^c	5.135 ± 12 ^f		≤ 60	
			6.291 ± 30		180 ± 60	
			7.062 ± 12		180 ± 50	$(2^+, 1^-)$
			7.712 ± 20		≤ 50	
			7.915 ± 12		≤ 50	$(2^+, 1^-)$
			8.100 ± 14		≤ 50	
			8.50 ± 30		≤ 120	

^a Observed but energy not determined.

^b $\Gamma \leq 40$ keV.

^c No other narrow states observed with $E_x < 7.5$ MeV (1970AD02).

^d A. Nero, private communication.

^e (1968TO09) report $E_x = 4.59 \pm 0.03$ MeV, $\Gamma \leq 130$ keV; the 4.51 MeV state was not resolved.

^f One of these two states has $J^\pi = 3^-$.

Table 18.26: Branching ratios and lifetimes of ^{18}Ne states

E_i (MeV)	J_i^π	E_f (MeV)	Branch (%)			τ_m (psec)
			(1968GI09, 1969RO08, 1972GI01)	(1969RO22)	(1970SH04)	
1.887	2^+	0	100			$0.49^{+0.17}_{-0.09}$ ^a
3.376	4^+	0	< 4	< 1	< 1	$\left\{ \begin{array}{l} 1.9^{+1.0}_{-0.4}$ ^a \\ 4.4 ± 0.6 ^b \end{array} \right.
		1.887	100	100	100	
3.576	0^+	0	< 17	< 5		$\left\{ \begin{array}{l} > 2$ ^a \\ < 6 ^b \end{array} \right.
		1.887	100	100		
3.616	2^+	0	12.5 ± 2.5	7 ± 2	< 3	$0.063^{+0.030}_{-0.020}$ ^a
		1.887	87.5 ± 2.5	93 ± 2	100	

^a (1968GI09, 1969RO08).

^b (1972GI01).

 Table 18.27: $^{16}\text{O}(^3\text{He}, n)^{18}\text{Ne}$ angular distribution studies

$E(^3\text{He})$ (MeV)	Distribution of neutron groups to $^{18}\text{Ne}^*$	Refs.
4.9, 5.2, 5.6	g.s.	(1967MC03)
5.51	g.s.	(1960AJ03)
5.6, 6.1	g.s.	(1961GA01)
9.0 – 11.5	3.38, 3.58 + 3.62	(1970AD02)
9.0 – 12.5	g.s., 1.89	(1970AD02)
9.15	g.s., 3.38, 3.58 + 3.62	(1968TO09)
10.15	g.s., 3.38, 3.58 + 3.62, 4.58	(1968TO09)
11	g.s., 1.89	(1966KR05, 1967KR1B)
11.5, 12.5	4.51 + 4.58, 5.08 + 5.13	(1970AD02)
13.8 – 17.8	g.s., 1.89, 3.5, 4.5, 5.1, 6.29, 7.06, 7.71, 7.92, 8.10, 8.50	(1970NE1N)

Table 18.28: States in ^{18}Ne from $^{20}\text{Ne}(\text{p}, \text{t})^{18}\text{Ne}$ ^a

E_x (MeV \pm keV)				L ^b	J^π
(1969HA38)	(1970FA17)	(1970LE08)	(1971PA1L)		
0	0	0	0	0	0^+
1.890 ± 20	1.89	1.83 ± 50	1.894 ± 10	2	2^+
3.375 ± 30	3.38	3.36 ± 50	3.390 ± 14	4	4^+
3.588 ± 25	3.61	3.58 ± 50	3.614 ± 13	0	0^+
4.580 ± 30	4.53 ± 20	4.46 ± 50	4.576 ± 17	1	1^-
5.115 ± 25	5.10 ± 20	5.12 ± 50	5.150 ± 14	2 + 3	$(2^+ + 3^-)$
	6.28 ± 20		6.326 ± 18	4	(4^+)
			7.957 ± 25		
	9.17 ± 30		9.215 ± 20		

^a See, however, Table 18.25 for states not resolved in this reaction.

^b (1970FA17, 1970LE08, 1971PA1L). See also (1969HA38).

3. $^{20}\text{Ne}(\text{p}, \text{t})^{18}\text{Ne}$ $Q_m = -20.022$

Transitions have been reported to nine states of ^{18}Ne at $E_p = 42.6$ to 50 MeV. L values derived from DWBA analysis of angular distributions are displayed in Table 18.28 (1969HA38, 1970FA17, 1970LE08, 1971PA1L). See also (1968PA1R, 1970OL1B) and (1969SO08; theor.).

^{18}Na
(Not illustrated)

A calculation using an isobaric mass formula predicts that the mass excess of ^{18}Na is 25.4 ± 0.4 MeV (1966KE16): ^{18}Na is then unbound with respect to proton emission by 1.6 MeV. See also (1965JA1C).

References

(Closed 31 December 1971)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1939BR1A Brubaker, Phys. Rev. 56 (1939) 1181
1939DE1A Devons, Proc. Roy. Soc. A172 (1939) 127
1949BL26 J.P. Blaser, F. Boehm and P. Marmier, Phys. Rev. 75 (1949) 1953
1951MI1A Middleton and Tai, Proc. Phys. Soc. (London) A64 (1951) 801
1953HE1A Heydenberg and Temmer, Phys. Rev. 92 (1953) 89
1954GO17 J.D. Gow and L.W. Alvarez, Phys. Rev. 94 (1954) 365
1955AJ61 F. Ajzenberg and T. Lauritsen, Rev. Mod. Phys. 27 (1955) 77
1955JA1A Jarmie, Phys. Rev. 98 (1955) 41
1955MA85 J.B. Marion, R.M. Brugger and T.W. Bonner, Phys. Rev. 100 (1955) 46
1955PR1A Price, Proc. Phys. Soc. (London) A68 (1955) 553
1955ST1A Stratton, Blair, Famularo and Stuart, Phys. Rev. 98 (1955) 629
1956BE1B Berger and Loper, Phys. Rev. 104 (1956) 1603
1956BR36 C.P. Browne, Phys. Rev. 104 (1956) 1598
1956DR38 R.W.B. Drever, A. Moljk and J. Scobie, Phil. Mag. 1 (1956) 942
1956RE04 J.B. Reynolds and K.G. Standing, Phys. Rev. 101 (1956) 158
1956RO1A Roclawski-Conjeaud and Cotton, Nucl. Phys. 1 (1956) 603; J. Phys. Rad. 17 (1956) 552; Physica 22 (1956) 1155A
1956SA06 R.M. Sanders, Phys. Rev. 104 (1956) 1434
1956SH1A Sher and Floyd, Phys. Rev. 102 (1956) 242
1957AH20 K. Ahnlund, Phys. Rev. 106 (1957) 124
1957BA14 E. Baumgartner and H.W. Fulbright, Phys. Rev. 107 (1957) 219
1957BI80 O.M. Bilaniuk and P.V.C. Hough, Phys. Rev. 108 (1957) 305
1957EL12 F.A. El Bedewi and I. Hussein, Proc. Phys. Soc. (London) A70 (1957) 233
1957NO17 E. Norbeck, Jr. and C.S. Littlejohn, Phys. Rev. 108 (1957) 754
1957RI1A Ribe, Phys. Rev. 106 (1957) 767

1957WI1E Winterberg, Z. Naturforsch. A12 (1957) 271
 1957YO04 T.E. Young, G.C. Phillips and R.R. Spencer, Phys. Rev. 108 (1957) 72
 1958AL03 E. Almqvist, D.A. Bromley and J.A. Kuehner, Bull. Amer. Phys. Soc. 3 (1958) 27, J12
 1958BE74 W.L. Bendel, J. McElhinney and R.A. Tobin, Phys. Rev. 111 (1958) 1297
 1958GO1A Goodman and Lee, Bull. Amer. Phys. Soc. 3 (1958) 37
 1958HE54 D.F. Herring, R. Chiba, B.R. Gasten and H.T. Richards, Phys. Rev. 112 (1958) 1210
 1958HE56 D.F. Herring, Phys. Rev. 112 (1958) 1217
 1958KA32 E. Kashy, P.D. Miller and J.R. Risser, Phys. Rev. 112 (1958) 547
 1958KU81 J.A. Kuehner, E. Almqvist and D.A. Bromley, Phys.. Rev. Lett. 1 (1958) 260
 1958PH37 W.R. Phillips, Phys. Rev. 110 (1958) 1408
 1958RO1A Roys and Shure, Nucl. Sci. Eng. 4 (1958) 536
 1958WE29 J.A. Weinman and E.A. Silverstein, Phys. Rev. 11 (1958) 277
 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
 1959AL99 K.W. Allen, D. Eccleshall and M.J.L. Yates, Proc. Phys. Soc. (London) A74 (1959) 660
 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
 1959FA1A Fagg and Hanna, Rev. Mod. Phys. 31 (1959) 711
 1959GO74 H.E. Gove and A.E. Litherland, Phys. Rev. 113 (1959) 1078
 1959HA1E Hamburger, M.S. Thesis, Univ. of Pittsburgh (1959)
 1959HI67 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A73 (1959) 721
 1959HI74 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 762
 1959JO32 R.L. Johnston, H.D. Holmgren and G.D. Gutsche, Bull. Amer. Phys. Soc. 4 (1959) 403, C4
 1959LO59 G. Lopez, F. Alba, M. Mazari and M.E. Ortiz, Rev. Mex. Fisica 8 (1959) 17
 1959PE1B Perlman, Welker and Wolfsberg, Congr. Int. De Phys. Nucleaire, 1958 (Dunod, 1959) 856
 1959VE19 G.E. Velyukhov, A.N. Prokofev and S.V. Starodubtsev, Dokl. Akad. Nauk SSSR 127 (1959) 781; Sov. Phys. Dokl. 4 (1960) 837
 1959VL23 N.A. Vlasov, S.P. Kalinin, A.A. Ogloblin and V.J. Chuev, Zh. Eksp. Teor. Fiz. 37 (1959) 1187; JETP (Sov. Phys.) 10 (1960) 844
 1959WA16 E.K. Warburton, Phys. Rev. 113 (1959) 595
 1959YO25 T.E. Young, G.C. Phillips, R.R. Spencer and D.A.A.S.N. Rao, Phys. Rev. 116 (1959) 962

1959YU1A Yu, Teng, Chou and Lee, *Scientia Sinica* 8 (1959) 935; *Phys. Abs.* 1350 (1960)

1960AG01 J. Aguilar, W.E. Burcham, J. Catala, J.B.A. England, J.S.C. McKee and J. Rotblat, *Proc. Roy. Soc.* A254 (1960) 395

1960AJ03 F. Ajzenberg-Selove and K.L. Dunning, *Phys. Rev.* 119 (1960) 1681

1960AM03 G. Amsel and O. Smulkowski, *Compt. Rend.* 251 (1960) 950

1960EV1A Everling, *Z. Naturforsch.* A15 (1960) 84

1960HA22 A.I. Hamburger, *Phys. Rev.* 118 (1960) 1271

1960JA11 A.A. Jaffe, I.J. Taylor and P.D. Forsyth, *Proc. Phys. Soc. (London)* 75 (1960) 940

1960JA13 N. Jarmie and M.G. Silbert, *Phys. Rev.* 120 (1960) 914

1960JA17 A.A. Jaffe, F. de S. Barros, P.D. Forsyth, J. Muto, I.J. Taylor and S. Ramavataram, *Prog. Phys. Soc.* 76 (1960) 914

1960NE1A Newns, *Proc. Phys. Soc. (London)* 76 (1960) 489

1960RA18 T.A. Rabson, T.W. Bonner, R. Castillo-Bahena, M.V. Harlow, Jr., H.P. Haenni and W.A. Ranken, *Nucl. Phys.* 19 (1960) 314

1960SH05 S.M. Shafroth, *J. Phys. Rad.* 21 (1960) 353

1960TA1C Talmi and Unna, *Ann. Rev. Nucl. Sci.* 10 (1960) 353

1960VE06 G.E. Velyukhov, A.N. Prokofev and S.V. Starodubtsev, *Zh. Skep. Teor. Fiz.* 39 (1960) 563; *JETP (Sov. Phys.)* 12 (1961) 395

1960ZE03 Ya.B. Zeldovich, *Zh. Eksp. Teor. Fiz.* 38 (1960) 1123; *JETP (Sov. Phys.)* 11 (1960) 812

1961AB05 A.K. Abramov and M.G. Yutkin, *Zh. Eksp. Teor. Fiz.* 41 (1961) 1023; *JETP (Sov. Phys.)* 14 (1962) 728

1961AR06 J.C. Armstrong and K.S. Quisenberry, *Phys. Rev.* 122 (1961) 150

1961BA1C Baz, Goldanskii and Zeldovich, *Sov. Phys. Uspekhi* 3 (1961) 729

1961BA1D Badalyan and Baz, *Zh. Eksp. Teor. Fiz.* 40 (1961) 549; *JETP (Sov. Phys.)* 13 (1961) 383

1961BE12 E.F. Bennett, *Phys. Rev.* 122 (1961) 595

1961BO1A Bonnel and Levy, *Compt. Rend.* 252 (1961) 2214

1961BU05 J.W. Butler and K.L. Dunning, *Phys. Rev.* 121 (1961) 1782

1961CA02 R.R. Carlson, C.C. Kim, J.A. Jacobs and A.C.L. Barnard, *Phys. Rev.* 122 (1961) 607

1961DI06 O. Dietzsch, Y. Hama, E.W. Hamburger and F.C. Zawislak, *Nucl. Phys.* 27 (1961) 103

1961DU02 K.L. Dunning and J.W. Butler, *Phys. Rev.* 123 (1961) 1321

1961EC02 D. Eccleshall and M.J.L. Yates, *Proc. Phys. Soc. (London)* 77 (1961) 93

1961GA01 N.H. Gale, J.B. Garg and K. Ramavataram, *Nucl. Phys.* 22 (1961) 500

1961HA43 G.C. Hanna, D.B. Primeau and P.R. Tunncliffe, *Can. J. Phys.* 39 (1961) 1784
 1961KO06 E. Kondaiyah, M.L. Jhingan and C. Badrinathan, *Nucl. Phys.* 27 (1961) 166
 1961KU02 J.A. Kuehner, E. Almqvist and D.A. Bromley, *Phys. Rev.* 122 (1961) 908
 1961LA09 F. Lacoste and G.R. Bishop, *Nucl. Phys.* 26 (1961) 511
 1961LE1A Legg, Unpublished Thesis, Princeton Univ. (1961)
 1961LE1D Levy and Bonnel, *J. Phys. Rad.* 22 (1961) 489
 1961LI03 A.E. Litherland, R. Batchelor, A.J. Ferguson and H.E. Gove, *Can. J. Phys.* 39 (1961) 276
 1961LO10 G. Lopez and O. Almen, *Rev. Mex. Fis.* 10 (1961) 239
 1961LO1C Lopez, *Rev. Mex. Fis.* 10 (1961) 283
 1961MA1B MacDonald and Douglas, *Nucl. Phys.* 24 (1961) 614
 1961NO05 E. Norbeck, *Phys. Rev.* 121 (1961) 824
 1961SI09 E.A. Silverstein, S.R. Salisbury, G. Hardie and L.D. Oppliger, *Phys. Rev.* 124 (1961) 868
 1961TO03 J.H. Towle and B.E.F. Macefield, *Proc. Phys. Soc. (London)* 77 (1961) 399
 1961TR1B True and Warburton, *Nucl. Phys.* 22 (1961) 426
 1961YA02 H. Yamaguchi, *J. Phys. Soc. Jpn.* 16 (1961) 583
 1962BA1D Barber, *Ann. Rev. Nucl. Sci.* 12 (1962) 1
 1962BL13 J.M. Blair and R.K. Hobbie, *Phys. Rev.* 128 (1962) 2282
 1962BR08 R.E. Brown, *Phys. Rev.* 125 (1962) 347
 1962CA20 G. Calvi, A. Rubbino and D. Zubke, *Nucl. Phys.* 38 (1962) 436
 1962CH01 C. Chasman and D.A. Bromley, *Bull. Amer. Phys. Soc.* 7 (1962) 36, JA4
 1962CO17 C.F. Coleman, P.E. Cavanagh, B.W. Ridley and J.F. Turner, *Conf. Low Energy Nucl. Phys. Harwell, AERE-R-4131*, 8, 3.3; Sept. 1962 (1962)
 1962DA04 J.F. Dawson, I. Talmi and J.D. Walecka, *Ann. Phys.* 18 (1962) 339, 350
 1962DO1A Dodge and Barber, *Phys. Rev.* 127 (1962) 1746
 1962GO1B Golsanskii, *Dokl. Akad. Nauk SSSR* 146 (1962) 1309
 1962HA40 B.G. Harvey, J. Cerny, R.H. Pehl and E. Rivet, *Nucl. Phys.* 39 (1962) 160
 1962HI06 S. Hinds, H. Marchant and R. Middleton, *Nucl. Phys.* 38 (1962) 81
 1962HO06 R.K. Hobbie and F.F. Forbes, *Phys. Rev.* 126 (1962) 2137
 1962HO1C Honda and Ui, *Nucl. Phys.* 34 (1962) 609
 1962IN1A Inglis, *Rev. Mod. Phys.* 34 (1962) 165
 1962JO14 J.C. Jodogne, P.C. Macq and J. Steyaert, *Phys. Lett.* 2 (1962) 325

1962MO1A Morpurgo, Nucl. Spectroscopy, Ed. Racah (Academic Press, 1962)
 1962PU01 D.J. Pullen, A.E. Litherland, S. Hinds and R. Middleton, Nucl. Phys. 36 (1962) 1
 1962TA07 K. Takamatsu, J. Phys. Soc. Jpn. 17 (1962) 896; Erratum. J. Phys. Soc. Jpn. 18 (1963) 1234
 1962TA1B Talmi and Unna, Nucl. Phys. 30 (1962) 280
 1962TA1D Talmi, Rev. Mod. Phys. 34 (1962) 704
 1963AL1D Almqvist, Evans and Kuehner, Padua (1963) 629
 1963AM1A Amsel, Thesis, Univ. of Paris (1963)
 1963BI05 G.R. Bishop, Nucl. Phys. 41 (1963) 118
 1963BU11 G.M. Bukat, L.A. Sliv and G.A. Sogomonova, Zh. Eksp. Teor. Fiz. 44 (1963) 316; JETP (Sov. Phys.) 17 (1963) 216
 1963DA1B Dar, Phys. Lett. 7 (1963) 339
 1963DO1B Douglas, Sala, Gomes and Polga, Padua (1963) 558A
 1963EL04 E. El-Baz, Nucl. Phys. 45 (1963) 383
 1963EV05 J.E. Evans, Phys. Rev. 131 (1963) 1642
 1963FR10 G. Frick, A. Gallmann, D.E. Alburger, D.H. Wilkinson and J.P. Coffin, Phys. Rev. 132 (1963) 2169
 1963FU06 H. Fuchs, Z. Phys. 171 (1963) 416
 1963GL1C Glendenning, Ann. Rev. Nucl. Sci. 13 (1963) 191
 1963GO32 S. Gorodetzky, R.E. Pixley, A. Gallman, G. Frick, J.P. Coffin and G. Sibille, J. Phys. 24 (1963) 892
 1963HA05 M. Harvey, Phys. Lett. 3 (1963) 209
 1963HI06 S. Hinds and B.M. Hinds, Nucl. Phys. 48 (1963) 690
 1963KU07 B. Kuhn and B. Schlenk, Phys. Lett. 5 (1963) 91
 1963LI07 A.E. Litherland, M.J.L. Yates, B.M. Hinds and D. Eccleshall, Nucl. Phys. 44 (1963) 220
 1963LO03 J. Lowe and C.L. McClelland, Phys. Rev. 132 (1963) 367
 1963MA1D Matthies, Neudachin and Smirnov, Nucl. Phys. 49 (1963) 97
 1963NO1B Nonaka et al., INSJ-57 (1963)
 1963OG1A Ogloblin, Nucl. Phys. 47 (1963) 408
 1963PA03 S.P. Pandya, Nucl. Phys. 43 (1963) 636
 1963RE06 F.H. Read and J.M. Calvert, Nucl. Phys. 41 (1963) 210
 1963RO1C Rook and Mitra, Padua (1963) 201

1963SA07 K. Sasaki, Prog. Theor. Phys. 29 (1963) 383
1963SE12 R.F. Seiler, C.H. Jones, W.J. Ansick, D.F. Herring and K.W. Jones, Nucl. Phys. 45 (1963) 647
1963VL1A Vlasov, Zh. Eksp. Teor. Fiz. 45 (1963) 160; JETP (Sov. Phys.) 18 (1964) 160
1963WI1B Willey, Nucl. Phys. 40 (1963) 529
1963YA03 K. Yagi, Y. Nakajima, K. Katori, Y. Awaya and M. Fujioka, Nucl. Phys. 41 (1963) 584
1963YA1B Yanabu et al., J. Phys. Soc. Jpn. 18 (1963) 747
1964AM02 S. Amiel and J. Gilat, Nucl. Sci. Eng. 18 (1964) 105
1964AM1A Amsel, Ann. Phys. 9 (1964) 297
1964BA1G Barz, Ann. Phys. 13 (1964) 164
1964BO13 R.O. Bondelid and J.W. Butler, Nucl. Phys. 53 (1964) 618
1964BR13 H.C. Bryant, J.G. Beery, E.R. Flynn and W.T. Leland, Nucl. Phys. 53 (1964) 97
1964BR1G Brill, Chuev and Ogloblin, Paris (1964) 981
1964BR1H Brown, Paris (1964) 129
1964CH19 L.F. Chase, Jr., H.A. Grench, R.E. McDonald and F.J. Vaughn, Phys. Rev. Lett. 13 (1964) 665
1964CO24 S. Cohen, R.D. Lawson, M.H. Macfarlane and M. Soga, Phys. Lett. 9 (1964) 180
1964DA1D Dar, Nucl. Phys. 55 (1964) 305
1964EL1B El-Nadi and Riad, Nucl. Phys. 50 (1964) 33
1964EN1A Engeland and Kallio, Nucl. Phys. 59 (1964) 211
1964ES02 M.A. Eswaran and C. Broude, Can. J. Phys. 42 (1964) 1311
1964EX1A Extermann, Baumgartner and Huber, Helv. Phys. Acta 37 (1964) 505
1964FE02 J.M. Ferguson, Nucl. Phys. 59 (1964) 97
1964FR1B Frick, Thesis, Univ. of Strasbourg (1964)
1964GA1A Gardner and Yu, Nucl. Phys. 60 (1964) 49
1964GA1B Gallman, Nucl. Instrum. Meth. 28 (1964) 33
1964GO11 A. Gobbi, A. Ruh, B. Gobbi and R.E. Pixley, Helv. Phys. Acta 37 (1964) 104
1964GR08 L.H. Greenberg, J.P. Roalsvig and R.N.H. Haslam, Can. J. Phys. 42 (1964) 731
1964HE06 E.M. Henley and D.U.L. Yu, Phys. Rev. 133 (1964) B1445
1964HE11 P. Hewka, R. Middleton and J. Wiza, Phys. Lett. 10 (1964) 93
1964HE1C Herbut, 3A(II)/C135, Paris (1964)
1964HO28 Hofmann, Acta Phys. Aust. 18 (1964) 309

- 1964IN03 T. Inoue, T. Sebe, H. Hagiwara and A. Arima, Nucl. Phys. 59 (1964) 1
- 1964KE1A Kelson, Phys. Rev. 134 (1964) B267
- 1964KI1B Kim, Seiler, Herring and Jones, Nucl. Phys. 57 (1964) 526
- 1964KO09 R. Kosiek, K. Schlupmann, H.W. Siebert and R. Wendling, Z. Phys. 179 (1964) 9
- 1964MA12 J.D. Mahoney and S.S. Markowitz, J. Inorg. Nucl. Chem. 26 (1964) 907
- 1964MA1G Mamasakhlisov, Izv. Akad. Nauk SSSR Ser. Fiz. 28 (1964) 1550
- 1964MA50 G.M. Matous and C.P. Browne, Phys. Rev. 136 (1964) B399
- 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, Proc. 2nd Int. Conf. on Nucl. Masses, Vienna, Austria, 1963; Ed. W.H. Johnson, Jr. (Springer Verlag, Vienna, 1964) 305
- 1964MC1A McKellar, Phys. Rev. 134 (1964) B1190
- 1964MI05 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 63
- 1964MO25 R. Moreh, T. Daniels and A.A. Jaffe, Proc. Phys. Soc. (London) 84 (1964) 332; Erratum Proc. Phys. Soc. (London) 84 (1964) 835
- 1964MU1A Mughabghab and Stephens, Phys. Rev. 133 (1964) 8660
- 1964PA1D Pandya and Green, Nucl. Phys. 57 (1964) 658
- 1964PU01 D.J. Pullen, J.R. Rook and R. Middleton, Nucl. Phys. 51 (1964) 88
- 1964SC01 J. Schaitman, H.F. Lutz and S.F. Eccles, Bull. Amer. Phys. Soc. 9 (1964) 68, GC10
- 1964ST1B Stovall, Phys. Rev. 133 (1964) B268
- 1964ST25 J.R. Stehn, M.D. Goldberg, B.N. Magurno and R. Wiener-Chasman, BNL-325, 2nd Ed., Suppl. 2, Vol. 1 (1964)
- 1964TR1A Trlifaj, Nucl. Phys. 52 (1964) 114
- 1964WA1E Walecka, Congres Int. de Phys. Nucl., Paris (1964)
- 1964WA1F Waghmare, Phys. Rev. 134 (1964) B1185
- 1964WI05 G. Wickenberg, S. Hjorth, N.G.E. Johansson and B. Sjogren, Ark. Fys. 25 (1964) 191
- 1964YO1B Young, Nucl. Phys. 55 (1964) 84
- 1965BA1J Bassichis, Giraud and Ripka, Phys. Rev. Lett. 15 (1965) 980
- 1965BE1T Berstch, Nucl. Phys. 74 (1965) 234
- 1965BL07 S.D. Bloom, J.D. Anderson, W.F. Hornyak and C. Wong, Phys. Rev. Lett. 15 (1965) 264
- 1965BO42 M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, H. Buttner, A. Lindner and H. Meldner, Nucl. Phys. 63 (1965) 438
- 1965BR1H Bryant and Flynn, Bull. Amer. Phys. Soc. 10 (1965) 515

1965CA06 R.R. Carlson and D.W. Heikkinen, Phys. Lett. 17 (1965) 305
1965CH10 C. Chasman, K.W. Jones, R.A. Ristinen and E.K. Warburton, Phys. Rev. 137 (1965) B1445
1965DE1H De Llano et al., Nucl. Phys. 72 (1965) 379
1965DI1C Dittmer et al., Bull. Amer. Phys. Soc. 10 (1965) 122
1965DO1E Dorosh, Kostyu and Shkodul-Yanov, Ukrayin. Fiz. Nauk USSR 9 (1965) 1040; Phys. Abs. 31902 (1965)
1965EB01 T.G. Ebrey and P.R. Gray, Nucl. Phys. 61 (1965) 479
1965EL06 P.J. Ellis and D. Wilmore, Proc. Phys. Soc. (London) 86 (1965) 422
1965EN02 T. Engeland, Nucl. Phys. 72 (1965) 68
1965ER04 J.R. Erskine, R.E. Holland, R.D. Lawson, M.H. Macfarlane and J.P. Schiffer, Phys. Rev. Lett. 14 (1965) 915
1965FE02 P. Federman and I. Talmi, Phys. Lett. 19 (1965) 490
1965FE1B Federman and Talmi, Phys. Lett. 15 (1965) 165
1965FO1E Fosado and Wolf, Rev. Mex. Fisica 14 (1965) 57
1965FR09 G. Frick, Ann. Phys. (Paris) 10 (1965) 155
1965GA1B Gaedke, Toth and Williams, Phys. Rev. 140 (1965) B296
1965GI1B Giraud, Nucl. Phys. 71 (1965) 373
1965GL07 R.N. Glover and A.D.W. Jones, Phys. Lett. 18 (1965) 165
1965GO11 S. Gorodetzky, A. Gallmann, J.P. Coffin, G. Frick and F. Jundt, Nucl. Phys. 73 (1965) 74
1965GO1F Goswami and Kisslinger, Phys. Rev. 140 (1965) B26
1965GO1G Gobbi, Matter and Thievent, Helv. Phys. Acta 38 (1965) 371
1965GR1F Greider, Ann. Rev. Nucl. Sci. 15 (1965) 291
1965JA1C Janecke, Nucl. Phys. 61 (1965) 383
1965KA1B Kallio, Phys. Lett. 18 (1965) 51
1965KA1C Kahana and Tomusiak, Nucl. Phys. 71 (1965) 402
1965LE1C Lee, Acta Phys. Sinica 21 (1965) 720
1965LO03 N. Longequeue, H. Beaumevielle, E. Ligeon, J.P. Longequeue and M. Sandon, J. Phys. (Paris) 26 (1965) 367
1965LO1D Lozynski, Nucl. Phys. 64 (1965) 321
1965LU1A Lutz, Eccles and Mason, Bull. Amer. Phys. Soc. 10 (1965) 122
1965MA1A Mansour et al., Nucl. Phys. 65 (1965) 433

1965MA1L Maguin, Compt. Rend. 260 (1965) 1149
 1965MO16 R. Moreh and T. Daniels, Nucl. Phys. 74 (1965) 403
 1965MO19 R. Moreh, Nucl. Phys. 70 (1965) 293
 1965NA1A Narasimhan and Shah, Nucl. Phys. 69 (1965) 204
 1965NI1A Nissim-Sabat, Thesis, Columbia Univ. (1965)
 1965OL02 R.W. Ollerhead, J.S. Lopes, A.R. Poletti, M.F. Thomas and E.K. Warburton, Nucl. Phys. 66 (1965) 161
 1965OW01 R.O. Owens and R.G. Winter, Nucl. Phys. 71 (1965) 625
 1965PA1F Parsons, Trefil and Drell, Phys. Rev. 138 (1965) 8847
 1965PO01 A.R. Poletti and E.K. Warburton, Phys. Rev. 137 (1965) B595
 1965PR1D Prosser, Din and Tolbert, Bull. Amer. Phys. Soc. 10 (1965) 462
 1965SA14 B. Saeki, Nucl. Phys. 73 (1965) 631
 1965SA18 V.S. Sadkovskii, E.D. Teterin, K.A. Gridnev, A.E. Denisov, R.P. Kolalis and Y.A. Nemilov, Yad. Fiz. 2 (1965) 843; Sov. J. Nucl. Phys. 2 (1966) 601
 1965SH1E Shapiro and Timashev, Yad. Fiz. 2 (1965) 459
 1965ST22 P.H. Stelson and L. Grodzins, Nucl. Data A1 (1965) 21
 1965VA1H Van Praet, Phys. Lett. 17 (1965) 120
 1965WI1B Wiza, Thesis, Univ. of Pennsylvania (1965) Unpublished
 1965ZA1B Zamick, Phys. Lett. 19 (1965) 580
 1965ZE04 B. Zeidman and T.H. Braid, Phys. Lett. 16 (1965) 139
 1966AD07 E.G. Adelberger and C.A. Barnes, Phys. Lett. 23 (1966) 474
 1966AG1B Aguilar, de la Rubia, Sanchez and Martinez, An. de Fisica Y Quim. 62 (1966) 279
 1966AL04 T.K. Alexander, K.W. Allen and D.C. Healey, Phys. Lett. 20 (1966) 402
 1966AL09 J.L. Alty, L.L. Green, R. Huby, G.D. Jones, J.R. Mines and J.F. Sharpey-Schafer, Phys. Lett. 20 (1966) 664
 1966AR10 A. Armigliato, F. Brandolini, F. Pellegrini and E. Crescenti, Nuovo Cim. B45 (1966) 92
 1966BA03 J.K. Bair, J.L.C. Ford, Jr. and C.M. Jones, Phys. Rev. 144 (1966) 799
 1966BA2C Bassichis, Giraud and Ripka, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintmeister, ZFK-122 (1966) 354
 1966BA2E Bassichis, Giraud and Ripka, J. Phys. C1-46 (1966)
 1966BA44 D. Bachelier, M. Bernas, I. Brissaud, P. Radvanyi and M. Roy, Nucl. Phys. 88 (1966) 307

1966BE07 C. Bergman and R.K. Hobbie, Phys. Rev. 142 (1966) 575
1966BE1G Becker, Chase, Kohler and McDonald, Bull. Amer. Phys. Soc. 11 (1966) 405
1966BE29 H.G. Benson and J.M. Irvine, Proc. Phys. Soc. (London) 89 (1966) 249
1966BO25 A. Bottino and B. Carezza, Nuovo Cim. B46 (1966) 137
1966BR1G Browne, F. S. U. Isobaric Spin Conf. (1966) 136
1966BR1P Brown, Proc. Int. School Enrico Fermi, Course 36, Ed. C. Bloch (Academic Press, 1969) 524
1966BR1Q Brown, Proc. Int. School Enrico Fermi, Course 36, Ed. C. Bloch (Academic Press, 1969) 513
1966BR1R Brown, F.S.U. Isobaric Spin Conf. (1966) 136
1966CH06 P.R. Chagnon, Nucl. Phys. 78 (1966) 193
1966CH12 P.R. Chagnon, Nucl. Phys. 81 (1966) 433
1966CH1E Chesterfield and Spicer, Bull. Amer. Phys. Soc. 11 (1966) 628; F.S.U. Isobaric Spin Conf. (1966) 433
1966DA1E Dalidchik and Sayasov, Yad. Fiz. 3 (1966) 820
1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097
1966DO1C Donau, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 20
1966GA10 R.M. Gaedke and K.S. Toth, Nucl. Phys. 83 (1966) 353
1966GI1A Gillet, Proc. Int. School Enrico Fermi, Course 36; Ed. C. Bloch (Academic Press, 1966) 43
1966GL1C R.N. Glover and A.D.W. Jones, Nucl. Phys. 81 (1966) 277
1966GL1D R.N. Glover, A.D.W. Jones and J.R. Rock, Nucl. Phys. 81 (1966) 289
1966GR1H L. Grunbaum and N.K. Ganguly, Phys. Lett. 23 (1966) 129
1966HA19 B.G. Harvey, J.R. Meriwether, J. Mahoney, A. Bussiere de Nercy and D.J. Horen, Phys. Rev. 146 (1966) 712
1966HA1K J. Hayward, Nucl. Phys. 81 (1966) 193
1966HA21 R.L. Hahn and E. Ricci, Phys. Rev. 146 (1966) 650
1966HA31 O. Hausser, H.J. Rose, J.S. Lopes and R.D. Gill, Phys. Lett. 22 (1966) 604
1966HE05 D.W. Heikkinen, Phys. Rev. 141 (1966) 1007; Erratum Phys. Rev. 149 (1966) 990
1966HE1E Hehl and Reidel, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 322
1966HO1E Holbrow, Polsky and Middleton, Bull. Amer. Phys. Soc. 11 (1966) 472
1966HU09 T.A. Hughes, R. Snow and W.T. Pinkston, Nucl. Phys. 82 (1966) 129

1966IB1A Iben, Bull. Amer. Phys. Soc. 11 (1966) 708
 1966IN01 T. Inoue, T. Sebe, H. Hagiwara and A. Arima, Nucl. Phys. 85 (1966) 184
 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
 1966KR05 M. Krick and G.J.F. Legge, Nucl. Phys. 89 (1966) 63
 1966KU05 T.T.S. Kuo and G.E. Brown, Nucl. Phys. 85 (1966) 40
 1966KU1D Kuehner, Nucl. Spin-Parity Assignments; Ed. Gove (Academic Press, 1966) 146
 1966KU1F M. Kushner and J. Quintanilla, Phys. Lett. 23 (1966) 572
 1966LA15 M. Lambert, G. Dumazet and H. Beaumevieille, Compt. Rend. B263 (1966) 580
 1966LA1E Lawson, 2nd Symp. on the Struct. of Low-Medium Mass Nuclei, Apr. 1966 (1966) 6
 1966LA1G I.M. Ladenbauer-Bellis, R.I. Morse and I.L. Preiss, Nucl. Phys. 88 (1966) 21
 1966LE11 C.W. Lee and E. Baranger, Nucl. Phys. 79 (1966) 385
 1966LE1B Lee, Thesis, Univ. of Kansas (1966)
 1966LO12 J.S. Lopes, O. Hausser, R.D. Gill and H.J. Rose, Nucl. Phys. 89 (1966) 127
 1966LU05 H.F. Lutz and S.F. Eccles, Nucl. Phys. 81 (1966) 423
 1966MA43 G.M. Matous, G.H. Herling and E.A. Wolicki, Phys. Rev. 152 (1966) 908
 1966MC05 R.L. McGrath, Phys. Rev. 145 (1966) 802
 1966MI1G Mikulinsky, Yad. Fiz. 3 (1966) 245
 1966NA1B Narayana, Current Sci. (India) 35 (1966) 38
 1966OL03 J.W. Olness and E.K. Warburton, Phys. Rev. 151 (1966) 792
 1966OL1C Olariu, Acad. Repub. Pop. Romine. Inst. de Fiz. Atom. Rept. IS 25 (1966)
 1966PI1B Picard and De Pinho, J. Phys. C1-42 (1966)
 1966PO1E Poth and Bromley, Bull. Amer. Phys. Soc. 11 (1966) 317
 1966RE1C R. Reif, Phys. Lett. 23 (1966) 125
 1966RI04 E. Rivet, R.H. Pehl, J. Cerny and B.G. Harvey, Phys. Rev. 141 (1966) 1021
 1966RI1F Ripka, Lect. in Theor. Phys., Vol. VIII C (Univ. of Colorado Press, Boulder, 1966) 237
 1966RO01 E. Rost, Phys. Lett. 21 (1966) 87
 1966SH1F I.S. Shapiro and S.F. Timashev, Nucl. Phys. 79 (1966) 46
 1966ST1F A.P. Stamp, Nucl. Phys. 83 (1966) 232
 1966SU1D Suzuki and Nagata, Prog. Theor. Phys. 36 (1966) 37
 1966WA1H Waghmare, Shakin and Svenne, Bull. Amer. Phys. Soc. 11 (1966) 321
 1966WE04 J.J. Wesolowski, L.F. Hansen, J.G. Vidal and M.L. Stelts, Phys. Rev. 148 (1966) 1063
 1966WI07 J.L. Wiza, R. Middleton and P.V. Hewka, Phys. Rev. 141 (1966) 975

- 1966YO1B Yoccoz, J. Phys. C1-3 (1966)
- 1966ZA03 L. Zamick, Phys. Lett. 21 (1966) 194
- 1967AL06 J.L. Alty, L.L. Green, R. Huby, G.D. Jones, J.R. Mines and J.F. Sharpey-Schafer, Nucl. Phys. A97 (1967) 541
- 1967AN1B Anderson and Bevington, Bull. Amer. Phys. Soc. 12 (1967) 1144
- 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
- 1967BA04 B. Banerjee, J.C. Parikh and Y.S.T. Rao, Nucl. Phys. A94 (1967) 481
- 1967BA1R J.M. Bang, C.A. Pearson and L. Pocs, Nucl. Phys. A100 (1967) 24
- 1967BA78 G. Backenstoss, S. Charalambus, H. Daniel, H. Koch, G. Poelz, H. Schmitt and L. Tauscher, Phys. Lett. B25 (1967) 547
- 1967BE14 J.A. Becker, J.W. Olness and D.H. Wilkinson, Phys. Rev. 155 (1967) 1089
- 1967BE30 T. Becker, K. Bahr, R. Jahr and W. Kuhlmann, Phys. Lett. B24 (1967) 458
- 1967BH07 K.H. Bhatt and J.C. Parikh, Nucl. Phys. A98 (1967) 113
- 1967BL18 A.E. Blaugrund, D.H. Youngblood, G.C. Morrison and R.E. Segel, Phys. Rev. 158 (1967) 893
- 1967BO1M Bolta, Garcia and Senent, An. Real. Soc. Espan. Fis. y Quim. 63 (1967) 35
- 1967BR1G B. Bremond, Nucl. Phys. A90 (1967) 241
- 1967CA1D Carlson, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 475
- 1967CA1L Cameron, Knudson, Cohen and Herling, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 171
- 1967CH1D Chase, Kohler, Becker and McDonald, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 930
- 1967CH1J Charlesworth and Azuma, Bull. Amer. Phys. Soc. 12 (1967) 53
- 1967CH1K Charlesworth, Azuma, Cassel and Kuehner, Bull. Amer. Phys. Soc. 12 (1967) 663
- 1967CH1L Chase, Nucl. Research With Low Energy Accelerators, Eds. Marion and Van Patter (Academic Press, 1967) 445
- 1967CO30 J.-P. Coffin, Ann. Phys. (Paris) 2 (1967) 273
- 1967DA1D Daniel et al., Z. Phys. 205 (1967) 472
- 1967DA1E Dalidtschik and Saiasov, Proc. Prob. Symp. on Nucl. Phys., Tbilishi, April 1967 (Moscow, 1967) 287
- 1967DEZW J. de Boer, A.M. Kleinfeld, R. Covello-Moro and H.P. Lie, Bull. Amer. Phys. Soc. 12 (1967) 535, GD8
- 1967DI1C Dittman, Cole, Waddell and Sandhu, Bull. Amer. Phys. Soc. 12 (1967) 663

- 1967DO1B F. Donau, K. Hehl, C. Riedel, R.A. Broglia and P. Federman, Nucl. Phys. A101 (1967) 495
- 1967DZ01 T.G. Dzubay, Phys. Rev. 158 (1967) 977
- 1967EN01 T. Engeland and P.J. Ellis, Phys. Lett. B25 (1967) 57
- 1967EV1C J.A. Evans and R.P.J. Perazzo, Nucl. Phys. A103 (1967) 225
- 1967FE01 P. Federman, Nucl. Phys. A95 (1967) 443
- 1967FL01 J. Flores and M. Moshinsky, Nucl. Phys. A93 (1967) 81
- 1967FO1B Fowler, Caughlan and Zimmerman, Ann. Rev. Astron. Astrophys. 5 (1967) 525
- 1967GO07 S. Gorodetzky, R.M. Freeman, A. Gallmann, F. Haas and B. Heusch, Phys. Rev. 155 (1967) 1119
- 1967GO1A G. Goldring, H.M. Loebenstein, I. Plesser and M.W. Sachs, Phys. Lett. B25 (1967) 538
- 1967GR09 H. Grote, Z. Naturforsch. A22 (1967) 415
- 1967GR1B Grunbaum and Ganguly, Bull. Amer. Phys. Soc. 12 (1967) 52
- 1967GR1D L. Grunbaum and N.K. Ganguly, Nucl. Phys. A100 (1967) 645
- 1967HO11 R.J.W. Hodgson, Phys. Rev. 156 (1967) 1173
- 1967IM1A Imanishi, Prog. Theor. Phys. 38 (1967) 1095
- 1967IN03 T. Inoue, T. Sebe, K.K. Huang and A. Arima, Nucl. Phys. A99 (1967) 305
- 1967KO1G Kobzev et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P6 3314 (1967)
- 1967KR1B Krick, Dissertation Abstr. B28 (1967); Phys. Abs. 43086 (1968)
- 1967KU09 T.T.S. Kuo, Nucl. Phys. A103 (1967) 71
- 1967KU13 T.T.S. Kuo, Phys. Lett. B26 (1967) 63
- 1967KU1G Kuo and Brown, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 526
- 1967LA1H Lawson and Soper, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 511
- 1967LE02 F.D. Lee, R.W. Krone and F.W. Prosser, Jr., Nucl. Phys. A96 (1967) 209
- 1967LO01 H.M. Loebenstein, D.W. Mingay, H. Winkler and C.S. Zaidins, Nucl. Phys. A91 (1967) 481
- 1967LO1F Love and Satchler, Nucl. Phys. A101 (1967) 424
- 1967LY02 R.P. Lynch and T.T.S. Kuo, Nucl. Phys. A95 (1967) 561
- 1967MA1B Marion, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 497
- 1967MA1E Mathur and Rook, Nucl. Phys. A91 (1967) 305

- 1967MA1F May and Truelove, *Ann. Phys.* 43 (1967) 322
- 1967MA1G Magelson, Thesis, Univ. of California, Berkeley (1967)
- 1967MC03 W.R. McMurray, P. Van Der Merwe and I.J. Van Heerden, *Nucl. Phys.* A92 (1967) 401
- 1967MO09 R. Moreh, *Nucl. Phys.* A97 (1967) 106
- 1967MO1J Moszkowski, *Rev. Mod. Phys.* 39 (1967) 657
- 1967MO23 J.F. Morgan and R.K. Hobbie, *Phys. Rev.* 163 (1967) 992
- 1967OG1A Ogloblin, Proc. Problem Symp. on Nucl. Phys., Tbilisi, Apr. 1967 (Moscow, 1967) 169
- 1967OL03 J.W. Olness and E.K. Warburton, *Phys. Rev.* 156 (1967) 1145
- 1967PA05 J.C. Parikh and N. Ullah, *Nucl. Phys.* A99 (1967) 529
- 1967PA1K J.C. Parikh and K.H. Bhatt, *Nucl. Phys.* A103 (1967) 496
- 1967PI1B Pinkston, *Nucl. Research with Low Energy Accelerators*, Eds. Marion and van Patter (Academic Press, 1967) 419
- 1967PO02 A.R. Poletti, *Phys. Rev.* 153 (1967) 1108
- 1967PO09 A.R. Poletti and D.B. Fossan, *Phys. Rev.* 160 (1967) 883
- 1967PO1E Poth and Bromley, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (Academic Press, 1967) 94
- 1967PR04 F.W. Prosser, Jr., G.U. Din and D.D. Tolbert, *Phys. Rev.* 157 (1967) 779
- 1967PU03 F. Puhlhofer and R. Bock, *Phys. Lett.* B25 (1967) 117
- 1967RO1F E. Rost, *Phys. Rev.* 154 (1967) 994
- 1967SC09 H. Schmidt, J. Morgenstern, H.J. Korner, J. Braunsfurth and S.J. Skorcka, *Phys. Lett.* B24 (1967) 457
- 1967SC16 H. Schulz, H.J. Wiebicke and R. Reif, *Nucl. Phys.* A101 (1967) 577
- 1967ST02 J.E. Stover, *Nucl. Phys.* A92 (1967) 209
- 1967TH1E Thornton, ORNL TM 1917 (1967)
- 1967VE1B Vervier, *Int. Nucl. Phys. Conf., Gatlinburg, 1966* (Academic Press, 1967) 558
- 1967VI1B C.M. Vincent, *Phys. Rev.* 159 (1967) 869
- 1967WA06 E.K. Warburton, J.W. Olness and A.R. Poletti, *Phys. Rev.* 155 (1967) 1164
- 1967WA19 E.K. Warburton, *Phys. Rev.* 163 (1967) 1032
- 1967WA1C Warburton, *Nucl. Research with Low Energy Accelerators*, Eds. Marion and van Patter (Academic Press, 1967) 43
- 1967WI16 J. Wilczynski, V.V. Volkov and P. Decowski, *Yad. Fiz.* 5 (1967) 942; *Sov. J. Nucl. Phys.* 5 (1967) 672

- 1967WI1F Wildenthal, Caies and Newman, Bull. Amer. Phys. Soc. 12 (1967) 536
- 1967WI1H Wilkniss, Int. J. Appl. Rad. Isotopes 18 (1967) 809
- 1967WI20 I.R. Williams and C.B. Fulmer, Phys. Rev. 162 (1967) 1055
- 1967WO1C C.W. Wong, Nucl. Phys. A91 (1967) 399
- 1968AB09 K. Abe, E. Tanaka, N. Kawamura, M. Kanazawa and N. Mutsuro, J. Phys. Soc. Jpn. 25 (1968) 1507
- 1968AN1A Anderson, Thesis, Stanford Univ. (1968)
- 1968AN1F Antolkovic et al., Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1658
- 1968AN20 D.S. Andreev, O.F. Afonin, V.K. Bondarev, A.P. Grinberg, K.I. Erokhina and I.K. Lemberg, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1671; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 1543
- 1968AR02 A. Arima, S. Cohen, R.D. Lawson and M.H. Macfarlane, Nucl. Phys. A108 (1968) 94
- 1968BA2G Backenstoss et al., Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 500
- 1968BA2H B.R. Barrett and M.W. Kirson, Phys. Lett. B27 (1968) 544
- 1968BA2J Bassel and Drisko, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, Sept. 1967 (1968) 13
- 1968BE1F H.W. Bertini, Phys. Rev. 171 (1968) 1261
- 1968BE1T Bertsch, Phys. Rev. Lett. 21 (1968) 1694
- 1968BE1U Becker, 3rd Symp. on the Struct. of Low-Medium Mass Nucl, Ed. Davidson (Univ. Press of Kansas, 1968) 1
- 1968BE1V G.F. Bertsch, Phys. Rev. 174 (1968) 1313
- 1968BE34 A.-M. Bergdolt and G. Bergdolt, Compt. Rend. B267 (1968) 455
- 1968BH1B Bhatt and Parikh, Tokyo (1968)
- 1968BL1G Blann et al., Bull. Amer. Phys. Soc. 13 (1968) 608
- 1968BO1U Bohr, Damgard and Mottelson, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 178
- 1968CH1K Chase, Becker, Kohler and McDonald, Tokyo (1968)
- 1968CL03 D.M. Clement and E.U. Baranger, Nucl. Phys. A108 (1968) 27
- 1968CO11 S. Cohen, E.C. Halbert and S.P. Pandya, Nucl. Phys. A114 (1968) 353
- 1968CO1N S. Cohen, R.D. Lawson and S.P. Pandya, Nucl. Phys. A114 (1968) 541
- 1968CU04 N. Cue and P. Richard, Phys. Rev. 173 (1968) 1108
- 1968DA1J Daniel, Rev. Mod. Phys. 40 (1968) 659
- 1968DE13 N. de Takacsy, Can. J. Phys. 46 (1968) 2091

- 1968DE1M Demos, Bull. Amer. Phys. Soc. 13 (1968) 628
- 1968DE21 F.P. Denisov, V.P. Milovanov, V.N. Pokrovskii, P.A. Cerenkov and I.A. Yutlandov, Yad. Fiz. 7 (1968) 743; Sov. J. Nucl. Phys. 7 (1968) 452
- 1968DI06 O. Dietzsch, R.A. Douglas, E.F. Pessoa, V.G. Porto, E.W. Hamburger, T. Polga, O. Sala, S.M. Perez and P.E. Hodgson, Nucl. Phys. A114 (1968) 330
- 1968DW1A M. Dworzecka and C.S. Warke, Phys. Rev. 175 (1968) 1346
- 1968EL1C Elliott and Wilsdon, Proc. Roy. Soc. 302 (1968) 509
- 1968FA01 M. Fazio, P. Guazzoni, S. Micheletti, M. Pignanelli and L. Zetta, Nucl. Phys. A111 (1968) 255
- 1968FA1A J.C. Faivre, H. Krivine and A.M. Papiou, Nucl. Phys. A108 (1968) 508
- 1968FA1G Falk, Kidney, Tandon and Kulisic, Bull. Amer. Phys. Soc. 13 (1968) 1464
- 1968FO1A Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299 (1968) 1
- 1968FR03 N. Freed and P. Ostrander, Nucl. Phys. A111 (1968) 63
- 1968GA03 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 167 (1968) 957
- 1968GA06 P.S. Ganas, Phys. Rev. 170 (1968) 879
- 1968GA16 P.S. Ganas and B.H.J. McKellar, Nucl. Phys. A120 (1968) 545
- 1968GI09 R.D. Gill, B.C. Robertson, J. L'Ecuyer, R.A.I. Bell and H.J. Rose, Phys. Lett. B28 (1968) 116
- 1968GO05 S. Gorodetzky, E. Aslanides, A. Gallmann and G. Frick, Nucl. Phys. A109 (1968) 417
- 1968GO1H A. Gobbi, U. Matter, J.-L. Perrenoud and P. Marmier, Nucl. Phys. A112 (1968) 537
- 1968GR1G Greene, Bull. Amer. Phys. Soc. 13 (1968) 606
- 1968GR1H Greene, Thesis, Univ. of Iowa (1968); Phys. Abs. 36614 (1969)
- 1968GR22 M.W. Greene, Phys. Rev. 176 (1968) 1204
- 1968GU1C M.R. Gunye, Nucl. Phys. A118 (1968) 174
- 1968GU1E M.R. Gunye, Phys. Lett. B27 (1968) 136
- 1968GU1F R.K. Gupta and L.E.H. Trainor, Nucl. Phys. A120 (1968) 273
- 1968GU1G M.R. Gunye, Nucl. Phys. A120 (1968) 691
- 1968HA17 E.C. Halbert, J.B. McGrory and B.H. Wildenthal, Phys. Rev. Lett. 20 (1968) 1112
- 1968HA1P Halbert, 3rd Symp. on the Struct. of Low-Medium Mass Nuclei; Ed. Davidson (U. Press of Kansas, 1968) 128
- 1968HA30 L.F. Hansen, M.L. Stelts, J.G. Vidal, J.J. Wesolowski and V.A. Madsen, Phys. Rev. 174 (1968) 1155
- 1968HE1G E.M. Henley, Phys. Lett. B28 (1968) 1

1968HE1H K.T. Hecht, P.J. Ellis and T. Engeland, Phys. Lett. B27 (1968) 479
 1968HO1C Hodgson, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, Sept. 1967 (1968) 41
 1968JA1J Jacquot et al., Compt. Rend. B266 (1968) 963
 1968JO07 J.E. Jobst, Phys. Rev. 168 (1968) 1156
 1968JO1C Joshi, Nucl. Phys. and Solid State Phys. Symp. (Powai, India, 1968); Phys. Abs. 38853 (1970)
 1968JO1E K.P. Joshi, G.K. Mehta and Y.R. Waghmare, Phys. Rev. 176 (1968) 1154
 1968JU1A Jung et al., Compt. Rend. B266 (1968) 815
 1968KA1C Kahana, Nucl. Part. Phys., Montreal, 1967 (New York, Benjamin. 1968) 75
 1968KA1E Kawai and Yazaki, Tokyo (1968) 238
 1968KO04 K. Kolltveit, Phys. Rev. 166 (1968) 1057
 1968KO1K K. Kolltveit, R. Muthukrishnan and R. Trilling, Phys. Lett. B26 (1969) 423
 1968KO1L Koren and Kanestrom, Phys. Norvegica 3 (1968) 107
 1968KU1E Kuriyama, Nagata and Bando, Tokyo (1968) 58
 1968LA1G R.D. Lawson, Nucl. Phys. A116 (1968) 363
 1968LE02 E. Ley-Koo, M. De Llano and J. Quintanilla, Nucl. Phys. A107 (1968) 35
 1968MA1C Marmier, Gobbi, Huber and Matter, Helv. Phys. Acta 41 (1968) 1028
 1968MA33 N.F. Mangelson, B.G. Harvey and N.K. Glendenning, Nucl. Phys. A119 (1968) 79; Erratum Nucl. Phys. A127 (1969) 693
 1968MA53 F. Machali, Z.A. Saleh, A.T. Baranik, F. Asfour, I. Bondouk and V.E. Storizhko, Ann. Phys. (Leipzig) 21 (1968) 1
 1968MI09 R. Middleton, L.M. Polsky, C.H. Holbrow and K. Bethge, Phys. Rev. Lett. 21 (1968) 1398
 1968MO1C Moskaleva, Fedoseev and Shcherbovskii, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 703
 1968MU1B Murthy, Nucl. Phys. Solid State Phys. Symp., Powai, India (1968); Phys. Abs. 38861 (1970)
 1968NA06 I.M. Naqib and L.L. Green, Nucl. Phys. A112 (1968) 76
 1968NA09 G.F. Nash, Nucl. Phys. A117 (1968) 615
 1968NA19 K.L. Narayana, Current Sci. (India) 37 (1968) 132
 1968NE1C Nemirovskii, Sov. J. Nucl. Phys. 6 (1968) 29
 1968NG1B Nguyen Van Sen et al., J. Phys. 29 (Paris) (1968) 15
 1968NO1C J.V. Noble, Phys. Rev. 173 (1968) 1034

- 1968OG1A Ogloblin, Nucl. Struct., Dubna Symp., 1968 (IAEA, 1968) 204
- 1968PA10 P.D. Parker, Phys. Rev. 173 (1968) 1021
- 1968PA1R Paddock, Benesen, Locard and Proctor, Bull. Amer. Phys. Soc. 13 (1968) 1464
- 1968PO04 A.M. Poskanzer, G.W. Butler, E.K. Hyde, J. Cerny, D.A. Landis and F.S. Goulding, Phys. Lett. B27 (1968) 414
- 1968PO1B Polsky, Thesis, Univ. of Penn. (1968)
- 1968RE07 D. Rendic, B. Antolkovic, G. Paic, M. Turk and P. Tomas, Nucl. Phys. A117 (1968) 113
- 1968RI13 L.R. Rice and R.T. Carpenter, Nucl. Phys. A120 (1968) 220
- 1968SH06 S. Shlomo and R. Moreh, Nucl. Phys. A110 (1968) 204
- 1968SH1H D.W. Sheffey, I.R. Williams and C.B. Fulmer, Phys. Rev. 172 (1968) 1094
- 1968ST1L Stehney and Steinberg, Nucl. Instrum. Meth. 59 (1968) 102
- 1968SU1C K. Sugawara, Nucl. Phys. A110 (1968) 305
- 1968SU1D Suekane, Prog. Theor. Phys. 39 (1968) 662
- 1968TA1C Tanner, Proc. Symp. on Use of Nimrod, 1968, RHEL/R166 (1968) 91
- 1968TH1J S.T. Thornton, C.M. Jones, J.K. Bair, M.D. Mancusi and H.B. Willard, Phys. Rev. Lett. 21 (1968) 447
- 1968TO09 J.H. Towle and G.J. Wall, Nucl. Phys. A118 (1968) 500
- 1968VA1J Vashakidze, Sov. J. Nucl. Phys. 6 (1968) 337
- 1968VA24 I.S. Vashakidze, Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 551; Bull. Acad. Sci. USSR Phys. Ser. 32 (1969) 507
- 1968WI1B Wilkinson, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 469
- 1968ZU02 A.P. Zuker, B. Buck and J.B. McGrory, Phys. Rev. Lett. 21 (1968) 39
- 1969AN06 J.D. Anderson, S.D. Bloom, C. Wong, W.F. Hornyak and V.A. Madsen, Phys. Rev. 177 (1969) 1416
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A137 (1969) 348
- 1969BA1P Bacon et al., Acta Cryst. A25 (1969) 391
- 1969BA1Z Barnes, Nucl. Isospin, Proc. 1969 Asilomar Conf. (Academic Press, 1969) 179
- 1969BA2E A.I. Baz and V.I. Manko, Phys. Lett. B28 (1969) 541
- 1969BA2F Baranger, Proc. Enrico Fermi School of Phys., Course XL, lake Como, 1967 (Academic Press, 1969) 643

- 1969BE1R Betigeri et al., Nucl. Phys. Solid State Phys. Symp., Roorke, India (1969); Phys. Abs. 38756 (1970)
- 1969BE1T H.G. Benson and B.H. Flowers, Nucl. Phys. A126 (1969) 332
- 1969BE22 A.M. Bernstein, Phys. Lett. B29 (1969) 335
- 1969BE57 P.M. Beard, P.B. Parks, E.G. Bilpuch and H.W. Newson, Ann. Phys. 54 (1969) 566
- 1969BR1D Bromly, Proc. Enrico Fermi School of Phys., Course XL, Lake Como 1967 (Academic Press, 1969) 242
- 1969BU15 S.T. Butler, R.G.L. Hewitt, B.H.J. McKellar, I.R. Nicholls and J.S. Truelove, Phys. Rev. 186 (1969) 963
- 1969CH1C D.T. Chivers, E.M. Rimmer, B.W. Allardyce, R.C. Witcomb, J.J. Domingo and N.W. Tanner, Nucl. Phys. A126 (1969) 129
- 1969CH1K Il-T. Cheon and H.M. Sen Gupta, Phys. Lett. B29 (1969) 268
- 1969CO12 H. Cords, G.U. Din and B.A. Robson, Nucl. Phys. A134 (1969) 561
- 1969CU10 H.-H. Cuno, G. Clausnitzer and R. Fleischmann, Nucl. Phys. A139 (1969) 657
- 1969DA19 V.V. Davydov, A.A. Ogloblin, S.B. Sakuta and V.I. Chuev, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 597; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 551
- 1969DE06 C. Detraz and H.H. Duhm, Phys. Lett. B29 (1969) 29
- 1969DE16 J. Dey, J.P. Elliott, A.D. Jackson, H.A. Mavromatis, E.A. Sanderson and B. Singh, Nucl. Phys. A134 (1969) 385
- 1969DE29 A.E. Denisov, R.P. Kolalis, Y.A. Nemilov, V.S. Sadkovskii and E.U. Teterin, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 1376; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 1271
- 1969DO08 E.I. Dolinskii and V.V. Turovtsev, Yad. Fiz. 9 (1969) 515; Sov. J. Nucl. Phys. 9 (1969) 295
- 1969DR1B Draayer, Thesis, Iowa State Univ. (1969); Phys. Abs. 38791 (1970)
- 1969DU11 P.D. Dumont and M. Huez, Bull. Soc. Roy. Sci. Liege 38 (1969) 516
- 1969EL1A Elliott, Cargese Lecture in Phys., Ed. M. Jean, Vol. 3 (Gordon & Breach, 1969) 337
- 1969EL1B Elliott, Proc. Int. Conf., Montreal (1969) 277
- 1969FE04 G.A. Fedoseev, L.P. Moskaleva and B.Y. Shcherbovskii, Yad. Fiz. 9 (1969) 1155; Sov. J. Nucl. Phys. 9 (1969) 675
- 1969FE10 J.L. Ferrero, A. Garcia, J. Milio and F. Senent, An. Fis. 65 (1969) 307
- 1969FE1A Federman, Cargese Lecture in Physics, Vol. 3, Ed. M. Jean (Gordon and Breach, 1969) 21
- 1969GA1G Garvey, Ann. Rev. Nucl. Sci. 19 (1969) 433
- 1969GA1Q E. Gadioli, E. Recami and L. Zetta, Nucl. Phys. A128 (1969) 339

1969GU1E M.R. Gunye, Nucl. Phys. A128 (1969) 457
 1969HA1F Hanna, Isospin in Nucl. Phys., Ed. D.H. Wilkinson (North-Holland, 1969) 591
 1969HA1T Haight, Thesis, Princeton Univ. (1969)
 1969HA1U Hansen and Stelts, Nucl. Isospin, Proc. 1969 Asilomar Conf. (Academic Press, 1969) 269
 1969HA38 J.C. Hardy, H. Brunnader, J. Cerny and J. Janecke, Phys. Rev. 183 (1969) 854
 1969HO16 W. Hofmann, R. Kosiek, G. Kraft and R. Mundhenke, Z. Phys. 225 (1969) 303
 1969HO32 R.J.W. Hodgson, Can. J. Phys. 47 (1969) 2269
 1969IG1A Igarashi, Kawai and Yazaki, Prog. Theor. Phys. 42 (1969) 245
 1969JA1P R.L. Jaffe and W.J. Gerace, Nucl. Phys. A125 (1969) 1
 1969JO09 J. Jobst, S. Messelt and H.T. Richards, Phys. Rev. 178 (1969) 1663
 1969JO1C P.L. Jolivette and H.T. Richards, Phys. Rev. 188 (1969) 1660
 1969JO1L Jones, Chase and Nightingale, Bull. Amer. Phys. Soc. 14 (1969) 1221
 1969JO1M Jolivette, Bull. Amer. Phys. Soc. 14 (1969) 1220
 1969KA07 I. Kanestrom and H. Koren, Nucl. Phys. A130 (1969) 527
 1969KA09 S. Kahana, H.C. Lee and C.K. Scott, Phys. Rev. 180 (1969) 956
 1969KA1A G.Th. Kaschl, G.J. Wagner, G. Mairle, U. Schmidt-Rohr and P. Turek, Phys. Lett. B29 (1969) 167
 1969KA1B R.W. Kavanagh, Nucl. Phys. A129 (1969) 172
 1969KA29 S. Kahana, H.C. Lee and C.K. Scott, Phys. Rev. 185 (1969) 1378
 1969KO1F V.M. Kolybasov and N.Ya. Smorodinskaya, Phys. Lett. B30 (1969) 11
 1969KR21 N.N. Krasnov, P.P. Dmitriev, Z.P. Dmitrieva, I.O. Konstantinov and G.A. Molin, Atomn. Energ. (USSR) 27 (1969) 125
 1969KU1G Kulkarni and Pandya, Nuovo Cim. B60 (1969) 199
 1969LI1G T.K. Lim, Nucl. Phys. A124 (1969) 475
 1969MA1G Madsen, Nucl. Isospin, Proc. 1969 Asilomar Conf. (Academic Press, 1969) 149
 1969MA1T MacFarlane, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (Academic Press, 1969) 457
 1969ME06 R. Mercier, E.U. Baranger and R.J. McCarthy, Nucl. Phys. A130 (1969) 322
 1969ME15 D. Meier, M. Brullmann, H. Jung and P. Marmier, Helv. Phys. Acta 42 (1969) 813
 1969MU09 M. Murthy and Y.R. Waghmare, Nucl. Phys. A137 (1969) 579
 1969NE1E Nettles, Hensley and Tombrello, Nucl. Isospin, Proc. 1969 Asilomar Conf. (Academic Press, 1969) 819

1969NI09 R.J. Nickles, Nucl. Phys. A134 (1969) 308
 1969NO1B J.V. Noble, Phys. Rev. Lett. 22 (1969) 473, Erratum Phys. Rev. Lett. 22 (1969) 1028
 1969NO1C Noble, Nucl. Isospin, Proc. 1969 Asilomar Conf. (Academic Press, 1969) 233
 1969OS01 E. Osnes and C.S. Warke, Phys. Lett. B30 (1969) 306
 1969PE1D Petrauskas and Vanagas, Sov. J. Nucl. Phys. 8 (1969) 270
 1969PO11 L.M. Polsky, C.H. Holbrow and R. Middleton, Phys. Rev. 186 (1969) 966
 1969RA28 C.A. Rappleyea and P.D. Kunz, Nucl. Phys. A139 (1969) 24
 1969RO08 B.C. Robertson, R.A.I. Bell, J. L'Ecuyer, R.D. Gill and H.J. Rose, Nucl. Phys. A126 (1969) 431
 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
 1969RO22 C. Rolfs, W. Trost, F. Riess, R. Kramer and E. Kuhlmann, Nucl. Phys. A137 (1969) 481
 1969SA1F P.U. Sauer, A. Faessler, H.H. Wolter and M.M. Stingl, Nucl. Phys. A125 (1969) 257
 1969SC1H Schmittroth, Thesis, Oregon State Univ. (1969); Phys. Abs. 42732 (1970)
 1969SC21 W.A. Schier and J.D. Reber, Phys. Rev. 181 (1969) 1371
 1969SO08 B. Sorensen, Nucl. Phys. A134 (1969) 1
 1969ST07 R.H. Stokes and P.G. Young, Phys. Rev. 178 (1969) 1789
 1969ST1G Stocker, Lett. Nuovo Cim. 1 (1969) 605
 1969TH01 M.J. Throop, Phys. Rev. 179 (1969) 1011
 1969UL03 N. Ullah and S.S.M. Wong, Phys. Rev. 188 (1969) 1645
 1969VO1E Vogt, Proc. Int. Conf., Montreal (1969) 5
 1969WA1C Warburton and Weneser, Isospin in Nucl. Phys., Ed. D.H. Wilkinson (North-Holland, 1969) 173
 1969WA1F Waghmare, Joshi and Mehta, Contrib., Montreal (1969) 186
 1969WU1A Wu and Wilets, Ann. Rev. Nucl. Sci. 19 (1969) 527
 1969ZA1C Zaremba and Roush, Bull. Amer. Phys. Soc. 14 (1969) 529
 1969ZA1D L. Zamick, Phys. Rev. Lett. 23 (1969) 1406
 1969ZU02 R.W. Zurmuhle and C.M. Fou, Nucl. Phys. A129 (1969) 502
 1969ZU03 A.P. Zucker, Phys. Rev. Lett. 23 (1969) 983
 1969ZU1B Zuker, Buck and McGrory, Bull. Amer. Phys. Soc. 14 (1969) 35
 1970AD02 E.G. Adelberger and A.B. McDonald, Nucl. Phys. A145 (1970) 497
 1970AG1C Agassi and Schaeffer, Nucl. Phys. A145 (1970) 401
 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1

- 1970AL11 D.E. Alburger and D.H. Wilkinson, Phys. Lett. B32 (1970) 190
- 1970AL1F Alvesh et al., Sov. J. Nucl. Phys. 10 (1970) 276
- 1970AN1D Anni and Taffara, Rivista Nuovo Cim. 2 (1970) 1
- 1970AR1D Artukh et al., Nucl. Reactions Included by Heavy Ions, Heidelberg, 1969 (North-Holland, 1970) 140
- 1970AS06 E. Aslanides, F. Jundt and A. Gallmann, Nucl. Phys. A152 (1970) 251
- 1970BA1E Backenstoss, Ann. Rev. Nucl. Sci. 20 (1970) 467
- 1970BA1M Bahcall and Fowler, Astrophys. J. 161 (1970) 119
- 1970BA1Z Barrett and Kirson, Nucl. Phys. A148 (1970) 145
- 1970BA2B Baz and Manko, Sov. J. Nucl. Phys. 10 (1970) 46
- 1970BA2E Barrett, Hewitt and McCarthy, Bull. Amer. Phys. Soc. 15 (1970) 1658
- 1970BA31 R.M. Bahnsen, W.R. Wylie and H.W. Lefevre, Phys. Rev. C2 (1970) 859
- 1970BA48 C. Baixeras-Aiguabella, M. Jung, C. Jacquot, L. Girardin and R. Schmitt, Phys. Rev. C2 (1970) 1194
- 1970BL1F Bleck, Haag and Ribbe, Z. Phys. 233 (1970) 65
- 1970BO25 W. Bohne, H. Homeyer, H. Lettau, H. Morgenstern and J. Scheer, Nucl. Phys. A156 (1970) 93
- 1970BR13 R.L. Brodzinski and N.A. Wogman, Phys. Rev. C1 (1970) 1955
- 1970CA1C Cavallaro, Cunsolo, Potenza and Rubbino, Programme of 56th National Congress of Italian Phys. Soc., Venice, 1970 (Italian Phys. Soc. 1970) 36; Phys. Abs. 51269 (1971)
- 1970CA1N Carlson and Johnson, Bull. Amer. Phys. Soc. 15 (1970) 629
- 1970CA1P Caplan, Singhal and Groh, Bull. Amer. Phys. Soc. 15 (1970) 803
- 1970CA28 J. Catala, A. Garcia, V. Martinez and F. Senent, An. Fis. 66 (1970) 297
- 1970CH1C Chung, Huber and Danos, Z. Phys. 240 (1970) 195
- 1970CH1D Chesterfield and Parker, Bull. Amer. Phys. Soc. 15 (1970) 1654
- 1970CH25 K. Chung, M.G. Huber, B. Blum and M. Danos, Phys. Lett. B32 (1970) 536
- 1970CL1C Clayton, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki (1970) 35
- 1970CO1P Corrigan, Prior and Darden, Bull. Amer. Phys. Soc. 15 (1970) 483
- 1970DA14 N.E. Davison, W.K. Dawson, G. Roy and W.J. McDonald, Can. J. Phys. 48 (1970) 2235
- 1970DI1G Dietrich, Nero and Walker, Bull. Amer. Phys. Soc. 15 (1970) 1658
- 1970DU08 H.H. Duhm, K. Peterseim, R. Seehars, R. Finlay and C. Detraz, Nucl. Phys. A151 (1970) 579

- 1970DU1D Dunnam et al., Bull. Amer. Phys. Soc. 15 (1970) 1686
- 1970EL08 P.J. Ellis and S. Siegel, Nucl. Phys. A152 (1970) 547
- 1970EL1G Ellis and Engeland, Nucl. Phys. A144 (1970) 161
- 1970FA01 W.R. Falk, A. Huber, U. Matter, R.W. Benjamin and P. Marmier, Nucl. Phys. A140 (1970) 548
- 1970FA17 W.R. Falk, R.J. Kidney, P. Kulisic and G.K. Tandon, Nucl. Phys. A157 (1970) 241
- 1970FE05 G.A. Fedoseev and L.P. Moskaleva, Yad. Fiz. 11 (1970) 1146; Sov. J. Nucl. Phys. 11 (1970) 636
- 1970FO1F Fortune, Richter, Siemssen and Tippie, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (North-Holland, 1970) 69
- 1970GA1J P.S. Ganas, Phys. Rev. C2 (1970) 93
- 1970GO1B Goldfarb, Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (North-Holland, 1970) 115
- 1970GR04 L.L. Green, C.O. Lennon and I.M. Naqib, Nucl. Phys. A142 (1970) 137
- 1970HA46 P.K. Haff and J.M. Eisenberg, Phys. Lett. B33 (1970) 133
- 1970HA49 M. Harvey and F.C. Khanna, Nucl. Phys. A155 (1970) 337
- 1970HO17 A.J. Howard, R.G. Hirko, D.A. Bromley, K. Bethge and J.W. Olness, Phys. Rev. C1 (1970) 1446
- 1970JA1Q Jacquot et al., Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (North-Holland, 1970) 701
- 1970JO09 D.J. Johnson and M.A. Waggoner, Phys. Rev. C2 (1970) 41
- 1970JO1C Jolivette, Thesis, Univ. Wisconsin (1970)
- 1970JO1F Jolivette, Bull. Amer. Phys. Soc. 15 (1970) 529
- 1970JO1G Jolivette, Bull. Amer. Phys. Soc. 15 (1970) 1686
- 1970JU1B M. Jung, C. Jacquot, C. Baixeras-Aiguabella, R. Schmitt and H. Braun, Phys. Rev. C1 (1970) 435
- 1970KA31 G.T. Kaschl, G.J. Wagner, G. Mairle and U. Schmidt-Rohr, Nucl. Phys. A155 (1970) 417
- 1970KA32 I. Kakkar, G.K. Mehta and Y.R. Waghmare, Nucl. Phys. A156 (1970) 199
- 1970KO1P R.R. Korteling and A.A. Caretto, Phys. Rev. C1 (1970) 193
- 1970KR12 N.N. Krasnov, P.P. Dmitriev, Z.P. Dmitrieva, I.O. Konstantinov and G.A. Molin, At. Energ. 28 (1970) 257; Sov. At. Energy 28 (1970) 333
- 1970KR1C Krasnov et al., At. Energ. 28 (1970) 503

- 1970LE08 J. L'Ecuyer, R.D. Gill, K. Ramavataram, N.S. Chant and D.G. Montague, Phys. Rev. C2 (1970) 116
- 1970LO01 G. Lodin and L. Nilsson, Z. Phys. 233 (1970) 181
- 1970MC23 J.B. McGrory, Phys. Lett. B33 (1970) 327
- 1970ME31 H.O. Menlove, R.H. Augustson, C.N. Henry, Nucl. Sci. Eng. 40 (1970) 136.
- 1970MO13 G.L. Morgan, D.R. Tilley, G.E. Mitchell, R.A. Hilko and N.R. Roberson, Nucl. Phys. A148 (1970) 480
- 1970MO17 G.L. Morgan, D.R. Tilley, G.E. Mitchell, R.A. Hilko and N.R. Roberson, Phys. Lett. B32 (1970) 353
- 1970MO35 U. Mosel, T.D. Thomas and P. Riesenfeldt, Phys. Lett. B33 (1970) 565
- 1970NE1J Nero and Adelberger, Bull. Amer. Phys. Soc. 15 (1970) 1686
- 1970NE1N Nero, Thesis, Stanford Univ. (1970)
- 1970OL1B Olsen, Thesis, Univ. Minnesota (1970); Phys. Abs. 47335 (1971)
- 1970PE1B Pearson, Rickel and Zissermann, Nucl. Phys. A148 (1970) 273
- 1970PR1C Prior, Corrigan and Darden, Bull. Amer. Phys. Soc. 15 (1970) 35
- 1970PR1D Preston and Yip, Bull. Amer. Phys. Soc. 15 (1970) 817
- 1970RA1A Raisbeck et al., PUC-937-369 (1970)
- 1970RO1F Rolfs, Kieser, Azuma and Litherland, Bull. Amer. Phys. Soc. 15 (1970) 782
- 1970SH04 M.H. Shapiro, A. Adams, C. Moss and W.M. Denny, Nucl. Phys. A144 (1970) 17
- 1970SH07 R.W. Shaw, Jr., R. Vandenbosch and M.K. Mehta, Phys. Rev. Lett. 25 (1970) 457
- 1970SI02 R.P. Singhal, J.R. Moreira and H.S. Caplan, Phys. Rev. Lett. 24 (1970) 73
- 1970SI09 R.H. Siemssen, H.T. Fortune, R. Malmin, A. Richter, J.W. Tippie and P.P. Singh, Phys. Rev. Lett. 25 (1970) 536
- 1970SI1M Singhal, SAL 17 (1970)
- 1970SU1B Sukhoruchkin, Sov. J. Nucl. Phys. 10 (1970) 144
- 1970TA1E K. Takeuchi and P.A. Moldauer, Phys. Rev. C2 (1970) 925
- 1970TA1J Talmi, Theory of Nucl. Struct. - Trieste 1969, IAEA STI/PUB/249 (1970) 455
- 1970TO03 P.B. Tollefsrud and P.L. Jolivet, Phys. Rev. C1 (1970) 398
- 1970TR07 L. Trlifaj and M. Sotona, Czech. J. Phys. B20 (1970) 865
- 1970TR08 J.S. Truelove and I.R. Nicholls, Aust. J. Phys. 23 (1970) 231
- 1970VE06 A.N. Vereshchagin, I.N. Korostova and I.P. Chernov, Izv. Vys. Ucheb. Zav. Fiz. 8 (1970) 105; Sov. Phys. J. 13 (1973) 1071

- 1970VI02 A.I. Vikhrov, V.E. Dudkin, I.I. Pyanov and O.V. Sysoeva, *Yad. Fiz.* 11 (1970) 36; *Sov. J. Nucl. Phys.* 11 (1970) 19
- 1970WA1G Walker and Stokes, LA-DC 11224 (1970)
- 1970WA1T Wasielewski, Thesis, Yale Univ. (1970); *Phys. Abs.* 51131 (1971)
- 1970WA31 U.M. Wambach, M. Gari and H. Kummel, *Phys. Lett.* B33 (1970) 253
- 1970ZE01 N.S. Zelenskaya, N.V. Karabanov, A.V. Spassky, K.G. Tarov, I.B. Teplov and L.N. Fateeva, *Yad. Fiz.* 11 (1970) 722; *Sov. J. Nucl. Phys.* 11 (1970) 405
- 1971AJ02 F. Ajzenberg-Selove, *Nucl. Phys.* A166 (1971) 1
- 1971AN1A Anderson, Velkley and Willard, *Bull. Amer. Phys. Soc.* 16 (1971) 36
- 1971AN1B Anderson and Bevington, Private Communication (1971)
- 1971AR02 A.G. Artukh, V.V. Avdeichikov, J. Ero, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski, *Nucl. Phys.* A160 (1971) 511
- 1971AR1R A. Arima and I. Hamamoto, *Ann. Rev. Nucl. Sci.* 21 (1971) 55
- 1971AR33 D. Ardouin, J. Uzureau, R. Tamisier, L.-H. Rosier and P. Avignon, *Compt. Rend.* B273 (1971) 561
- 1971BA14 B.R. Barrett, R.G.L. Hewitt and R.J. McCarthy, *Phys. Rev.* C3 (1971) 1137
- 1971BA2A Bainum, Lindgren and Park, *Bull. Amer. Phys. Soc.* 16 (1971) 1155
- 1971BE19 R.R. Betts, H.T. Fortune, J.D. Garrett, R. Middleton, D.J. Pullen and O. Hansen, *Phys. Rev. Lett.* 26 (1971) 1121
- 1971BE1E Berka et al., *Bull. Amer. Phys. Soc.* 16 (1971) 510
- 1971BE45 Z. Berant, M.B. Goldberg, M. Popp, J.S. Sokolowski, P.N. Tandon and Y. Wolfson, *Nucl. Phys.* A173 (1971) 401
- 1971BL12 R.J. Blin-Stoyle, J.A. Evans and A.M. Khan, *Phys. Lett.* B36 (1971) 202
- 1971BO21 A. Bouyssy and N. Vinh Mau, *Phys. Lett.* B35 (1971) 269
- 1971BR44 R.C. Brown, A.A. Debenham, G.W. Greenlees, J.A.R. Griffith, O. Karban, D.C. Kocher and S. Roman, *Phys. Rev. Lett.* 27 (1971) 1446
- 1971BU1E Butler, Poskanzer and Landis, *Proc. Int. Conf. on Heavy Ion Phys.*, Dubna (1971) 210
- 1971CH1F Charlesworth, Azuma, Cassell and Kuehner, Private Communication (1971); Unknown Source
- 1971CO1G Couch, Spinka, Tombrello and Weaver, OAP 263 (1971)
- 1971CO27 R.G. Couch, H. Spinka, T.A. Tombrello and T.A. Weaver, *Nucl. Phys.* A175 (1971) 300
- 1971DE1E J. Delorme and M. Rho, *Phys. Lett.* B34 (1971) 238
- 1971DE1F Devins, High, Bedi and Shapiro, *Bull. Amer. Phys. Soc.* 16 (1971) 511

- 1971DZ06 R.I. Dzhibuti, *Yad. Fiz.* 13 (1971) 1008; *Sov. J. Nucl. Phys.* 13 (1971) 579
- 1971EL07 P.J. Ellis and H.A. Mavromatis, *Nucl. Phys.* A175 (1971) 309
- 1971GA13 R.M. Gaedke, K.S. Toth and I.R. Williams, *Phys. Rev.* C4 (1971) 98
- 1971GA1B Galati et al., *Bull. Amer. Phys. Soc.* 16 (1971) 36
- 1971GA1D Gambhir and Griffin, *Bull. Amer. Phys. Soc.* 16 (1971) 580
- 1971GO1T Gobbi et al., *Proc. Int. Conf. on Heavy Ion Phys.*, Dubna (1971) 443
- 1971GU21 M.R. Gunye and S.B. Khadkikar, *Phys. Rev.* C4 (1971) 1073
- 1971HE1F Hensley, Goodman and Saltmarsh, *Bull. Amer. Phys. Soc.* 16 (1971) 510
- 1971HI06 F. Hibou, P. Fintz, B. Rastegar and A. Gallmann, *Nucl. Phys.* A171 (1971) 603
- 1971HO12 R.J.W. Hodgson, *Can. J. Phys.* 49 (1971) 1401
- 1971HR1A Hrejsa and Browne, *Bull. Amer. Phys. Soc.* 16 (1971) 1164
- 1971HU1C Hudson et al., *Polarization*, Madison, 1970 (Univ. Wisconsin Press, 1971) 777
- 1971JA04 J. Janecke, T.F. Yang, W.S. Gray and R.M. Polichar, *Phys. Rev.* C3 (1971) 79
- 1971JA06 M.C. Jain, G.K. Mehta and Y.R. Waghmare, *Phys. Rev.* C3 (1971) 1466
- 1971JO11 P.L. Jolivet, *Phys. Rev. Lett.* 26 (1971) 1383
- 1971KE1E Keaton et al., *Polarization*, Madison, 1970 (Univ. Wisconsin Press, 1971) 849
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, *Nucl. Phys.* A172 (1971) 25
- 1971KO21 D.C. Kocher, P.J. Bjorkholm and W. Haeberli, *Nucl. Phys.* A172 (1971) 663
- 1971LA1D Lang, *Z. Phys.* 242 (1971) 179
- 1971LE1H T.Y. Lee, S.T. Hsieh, W.W. Yeh and C.T. Chen-Tsai, *Phys. Rev.* C4 (1971) 64
- 1971LE1R Lerner and Marion, *Bull. Amer. Phys. Soc.* 16 (1971) 1163
- 1971LE29 C. Lebrun, F. Guilbault and Y. Deschamps, *Compt. Rend.* B273 (1971) 41
- 1971LI1D Lindgren et al., *Bull. Amer. Phys. Soc.* 16 (1971) 36
- 1971LI27 R.A. Lindgren, F.C. Young and B. Cotton, *Phys. Lett.* B37 (1971) 358
- 1971LO23 N. Lo Iudice, D.J. Rowe and S.S.M. Wong, *Phys. Lett.* B37 (1971) 44
- 1971MI1H Milyanich, Tomash, Furich and Valkovich, *Izv. Akad. Nauk SSSR Ser. Fiz.* 35 (1971) 739
- 1971MO05 R. Mohan, M. Danos and L.C. Biedenharn, *Phys. Rev.* C3 (1971) 468
- 1971MO1H Moskaleva, Fedoseev and Khalemskii, *Sov. J. Nucl. Phys.* 12 (1971) 472
- 1971OL06 R.W. Ollerhead, G.F.R. Allen, A.M. Baxter and J.A. Kuehner, *Can. J. Phys.* 49 (1971) 2589
- 1971PA1L Paddock, MSUCL-36 (1971)

1971PE1A Pearson and Quang-Hoc, Bull. Amer. Phys. Soc. 16 (1971) 44
 1971PR16 H.C. Pradhan and C.M. Shakin, Phys. Lett. B37 (1971) 151
 1971QU01 N. Quang-Hoc and J.M. Pearson, Nucl. Phys. A164 (1971) 631
 1971RA1C Rawitscher, Bull. Amer. Phys. Soc. 16 (1971) 100
 1971RE20 F. Resmini, R.M. Lombard, M. Pignanelli, J.L. Escudie and A. Tarrats, Phys. Lett. B37 (1971) 275
 1971RO18 H.J. Rose, R.D. Gill, J.M.G. Caraca and A.J. Cox, Part. Nucl. 1 (1971) 453
 1971RO25 C. Rolfs, H.P. Trautvetter, R.E. Azuma and A.E. Litherland, Phys. Rev. Lett. 27 (1971) 1007
 1971SC01 D. Schutte, Nucl. Phys. A161 (1971) 73
 1971SC1Q Schuck, Z. Phys. 241 (1971) 395
 1971SE1H Sens, Pape and Armbruster, Private Commication (1971)
 1971SE1J Sens, Rietsch, Pape and Armbruster, Private Communication (1971)
 1971TH03 S.T. Thornton, C.M. Jones, J.K. Bair, M.D. Mancusi and H.B. Willard, Phys. Rev. C3 (1971) 1065
 1971TH10 S.T. Thornton, R.P. Fogel and C.L. Morris, Nucl. Phys. A170 (1971) 485
 1971TH1D Thirion, Conf. on Certain Microscopic Aspects of Nucl. Reactions, La Toursinei, 1971 (Vieleurbanne, France, Inst. Nucl. Phys. 1971); Phys. Abs. 43602 (1971)
 1971UL01 N. Ullah, Nucl. Phys. A164 (1971) 658
 1971UL04 N. Ullah and K.K. Gupta, Phys. Lett. B36 (1971) 196
 1971VA1C E. Vatai, Phys. Lett. B34 (1971) 395
 1971VA1H Vandenbosch, Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 337
 1971VI08 B.N. Vinogradov and P.E. Nemirovskii, Yad. Fiz. 13 (1971) 1016; Sov. J. Nucl. Phys. 13 (1971) 583
 1971WI01 WildeB.H. Wildenthal, J.B. McGrory and P.W.M. Glaudemans, Phys. Rev. Lett. 26 (1971) 96
 1971WI18 D.H. Wilkinson, Phys. Rev. Lett. 27 (1971) 1018
 1972DE05 J.R. Demos and M.K. Banerjee, Phys. Rev. C5 (1972) 75
 1972GA02 P.S. Ganas, Phys. Rev. C5 (1972) 36
 1972GI01 R.D. Gill, J.M.G. Caraca, A.J. Cox and H.J. Rose, Nucl. Phys. A180 (1972) 79
 1972KA01 S. Kahana, Phys. Rev. C5 (1972) 63
 1972LE1L Lee and Cusson, Ann. Phys. 72 (1972) 353