

Energy Levels of Light Nuclei $A = 19$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 18-20$ was published in *Nuclear Physics A190* (1972), p. 1. This version of $A = 19$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed December 31, 1971)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 19$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{19}\text{C}\$](#) , [\$^{19}\text{N}\$](#) , [\$^{19}\text{O}\$](#) , [\$^{19}\text{F}\$](#) , [\$^{19}\text{Ne}\$](#) , [\$^{19}\text{Na}\$](#) , [\$^{19}\text{Mg}\$](#)

B. Tables of Recommended Level Energies:

[Table 19.1](#): Energy levels of ^{19}O

[Table 19.6](#): Energy levels of ^{19}F

[Table 19.21](#): Energy levels of ^{19}Ne

C. [References](#)

D. Figures: [\$^{19}\text{O}\$](#) , [\$^{19}\text{F}\$](#) , [\$^{19}\text{Ne}\$](#) , [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

¹⁹C
(Not illustrated)

¹⁹C may have been observed in the 3 GeV proton bombardment of a ¹⁹⁷Au target: if so it is particle stable (1970RA1A). Its mass excess must then be < 37.9 MeV (¹⁸C + n). See also (1960ZE03, 1971BU1E).

¹⁹N
(Not illustrated)

¹⁹N has been observed in the 3 GeV proton bombardment of a ¹⁹⁷Au target: it is particle stable (1968TH04). See also (1969AR13, 1970AR1D). The mass excess of ¹⁹N is 15.01 < (M - A) < 18.68 MeV (1970WA1G). See also (1960ZE03, 1961BA1C, 1962GO1B, 1966GA25, 1969ST07).

¹⁹O
(Figs. 5 and 8)

GENERAL: (See also (1959AJ76).)

Shell model: (1957WI1E, 1959BR1E, 1960TA1C, 1962SH1B, 1962TA1B, 1963PA03, 1964CO24, 1964IN03, 1965FE02, 1966AR10, 1966IN01, 1966LA1E, 1967BO09, 1967EN01, 1967FE01, 1967FL13, 1967IN03, 1967PI1B, 1968AR02, 1968CO1N, 1968EL1C, 1968GU1E, 1968HA17, 1968HA1P, 1968MO1G, 1969FE1A, 1969HO32, 1969KU1G, 1969MA1T, 1970TA1J, 1971AR25, 1971WI01).

Cluster, collective and deformed models: (1963CH02, 1965FE02, 1969FE1A).

Astrophysical questions: (1970BA1M).

Electromagnetic transitions: (1963CH02, 1965FE02, 1967FE01, 1967IN03, 1968EL1C, 1968HA17, 1971AR25).

Special levels: (1959LA1A, 1960GO1C, 1967EN01, 1967PR04, 1968AR02, 1968CO1N, 1968HA1P, 1968YA03, 1969FE1A).

Other theoretical topics: (1964IN03, 1965FO1D, 1967FL13, 1968CO1N, 1968GU1E, 1968MO1G, 1968MU1B, 1968PA1Q, 1968SU1C, 1969MA1T, 1971LA1D, 1971LE1H).

Ground state properties: (1964ST1B, 1968EL1C, 1971AR25, 1971WI01).

1. ¹⁹O(β^-)-¹⁹F $Q_m = 4.819$

The half-life of ¹⁹O is 29.12 ± 0.26 sec (1962MA38), 27.2 ± 0.3 sec (1965BO42), 26.76 ± 0.08 sec (1967YU01), 27.1 ± 0.1 sec (1970AL21): the weighted mean of the last three values is 26.91 ± 0.08 sec. The decay is complex: see ¹⁹F.

Table 19.1: Energy levels of ^{19}O ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{3}{2}$	$\tau_{1/2} = 26.91 \pm 0.08$ sec	β^-	1, 3, 8, 11, 12
0.0960 ± 0.5	$\frac{3}{2}^+$	$\tau_m = 1.98 \pm 0.06$ nsec	γ	2, 3, 8, 12
1.4717 ± 0.4	$\frac{1}{2}^+$	$\tau_m = 1.13 \pm 0.14$ psec	γ	2, 3, 8, 12
2.3711 ± 1.0	$(\frac{9}{2})$	$\tau_m \geq 1$ psec	γ	3, 8
2.7787 ± 0.8	$(\frac{7}{2}^+)$	$\tau_m = 93 \pm 19$ fsec	γ	3, 8
3.0671 ± 2.6	$(\frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+)$	$\tau_m \geq 1$ psec	γ	3, 8
3.1545 ± 2.0	$\frac{5}{2}^+$	$\tau_m \geq 1$ psec	γ	3, 8
3.235 ± 4	$\frac{1}{2}^+$		γ	8
3.9453 ± 2.5	$(\frac{1}{2}^-)$		γ	3, 8
4.116 ± 4	$(\frac{3}{2}, \frac{5}{2})^+$	$\Gamma < 15$ keV		3, 8
4.333 ± 12		< 15		8
4.400 ± 9		< 15		3, 8
4.583 ± 8	$\frac{3}{2}^-$	53 ± 3	γ, n	4, 5, 8
4.707 ± 12	$\frac{5}{2}^+$	< 15		3, 8
4.998 ± 12		< 15		3, 8
5.086 ± 10	$\frac{1}{2}^-$	49 ± 5	n	5
5.149 ± 7	$\frac{3}{2}$	3.4 ± 1.0	n	3, 5, 8
5.455 ± 9	$\frac{5}{2}^+$	330 ± 40	n	5, 8
5.502 ± 12		< 15		3, 8
5.53	$(\frac{1}{2}^-)$	380 ± 100	n	5
5.706 ± 8	$\frac{3}{2}$	7.8 ± 1.4	n	5, 8
6.13	$\frac{3}{2}^+$	150 ± 40	n	5
6.20	$\frac{1}{2}^-$	140 ± 40	n	5
6.276 ± 7	$\frac{7}{2}^-$	19.2 ± 2.4	n	5, 8
6.480 ± 15			(n)	5, 8
6.560 ± 15				8
6.899 ± 15				8
6.997 ± 15				8
7.117 ± 15				8
7.248 ± 15				8
11.25 ± 50		240	n, α	7
11.58 ± 50		330	n, α	7

^a See also Table 19.5.

Table 19.2: Lifetime measurements of some ^{19}O states

$^{19}\text{O}^*$ (MeV)	τ_m	Reaction	Refs.
0.096 ^a	1.75 ± 0.3 nsec	$^{18}\text{O}(\text{d}, \text{p})$	(1959ZI16)
	1.89 ± 0.2 nsec	$^9\text{Be}(^{18}\text{O}, ^8\text{Be})$	(1969NI09)
	2.00 ± 0.07 nsec	$^{18}\text{O}(\text{d}, \text{p})$	(1965MC10)
1.472 ^b	1.98 ± 0.06 nsec		mean
	1.80 ± 0.29 psec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971BR02)
	1.22 ± 0.36 psec	$^{18}\text{O}(\text{d}, \text{p})$	(1971MC11)
	0.90 ± 0.23 psec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971HI06, 1971HIZF)
	1.27 ± 0.2 psec		A
2.37	≥ 1 psec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971BR02)
	> 3.5 psec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971HI06, 1971HIZF)
2.78	70 ± 26 fsec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971BR02)
	117 ± 26 fsec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971HI06, 1971HIZF)
3.07 ^c	≥ 1 psec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971BR02)
3.16 ^c	≥ 1 psec	$^{17}\text{O}(\text{t}, \text{p}), ^{18}\text{O}(\text{d}, \text{p})$	(1971BR02)

A: E.K. Warburton, private communication.

^a See also (1965AL13, 1967WA1C).

^b See also (1965MC10, 1969NI09).

^c See also (1971HIZF).

2. $^9\text{Be}(^{18}\text{O}, ^8\text{Be})^{19}\text{O}$ $Q_m = 2.292$

The lifetimes of the excited states of ^{19}O at 95.1 ± 0.1 and 1468 keV are, respectively, $\tau_m = 1.89 \pm 0.2$ nsec and 18 ± 10 psec (1969NI09): see, however, Table 19.2. $|M|^2$ values are also listed. See also (1968FA04, 1970BA1Y, 1970FA01, 1971KN05).

3. $^{17}\text{O}(\text{t}, \text{p})^{19}\text{O}$ $Q_m = 3.521$

Proton groups corresponding to ^{19}O states with $E_x < 5.6$ MeV have been observed by (1964MO24, 1965MO19: $E_t = 5.55$ MeV) and (1965WI1B, 1966WI05: $E_t = 12.0$ MeV): level parameters and L values derived from angular distribution analyses are displayed in Table 19.3. Excitation energies from E_γ measurements, and lifetime measurements are reported in Tables 19.3 and 19.2, respectively (1971BR02, 1971HI06).

Table 19.3: Levels of ^{19}O from $^{17}\text{O}(t, p)^{19}\text{O}$ and $^{18}\text{O}(d, p)^{19}\text{O}$

E_x (MeV \pm keV)											Γ^e (keV)	l_n a,e,f	L b,e,f	J^π
(1961AR06) a	(1961SJ01) a	(1963YA03) a	(1964EL1A) a	(1964MO24, 1965MO19) b	(1965MO16) a	(1971BR02) c,d	(1965WI1B, 1966WI05) a,b	(1969FI07) a	(1969FI07, 1970FI08) c	(1971HI06) c,d				
0	0 0.098 \pm 6	0 h	0	0	0	0.096 \pm 2	0	0	0.097 \pm 2 ⁱ		2	0	$\frac{1}{2}^+$	
1.469 \pm 11	(1.257 \pm 30) 1.469 \pm 28 2.353 \pm 29	1.468 \pm 15	1.467 \pm 10 2.371 \pm 10 2.617 \pm 10	1.468 \pm 10 2.367 \pm 10	1.468 \pm 10	1.470 \pm 2 2.367 \pm 4	1.467 \pm 12 2.373 \pm 12	1.468 \pm 4 2.370 \pm 5	1.470 \pm 3 ^j 2.369 \pm 4	1.4719 \pm 0.5 2.3715 \pm 1.0	0 2	(2) (2)	$\frac{1}{2}^+$ ($\frac{3}{2}^+$) ^m	
3.164 \pm 30	2.765 \pm 30 (3.047 \pm 30) 3.144 \pm 29	2.612 \pm 15 3.171 \pm 15	2.794 \pm 10 3.161 \pm 10 3.243 \pm 10	2.775 \pm 10 3.061 \pm 10 3.153 \pm 10 3.223 \pm 15	3.064 \pm 10 3.155 \pm 10 3.233 \pm 15	2.774 \pm 4 (2.794 \pm 3) 3.066 \pm 3 3.150 \pm 3	2.779 \pm 12 3.070 \pm 12 3.156 \pm 12 3.229 \pm 12	2.780 \pm 5 3.070 \pm 5 3.160 \pm 5 3.237 \pm 5	2.777 \pm 3 3.157 \pm 3	2.7790 \pm 0.9	2	0 (2)	$\frac{3}{2}^+$ ($\frac{1}{2}^+$) ^l ($\frac{7}{2}^+$) ^m	
3.948 \pm 30 (4.123 \pm 40)	(3.791 \pm 31) 3.942 \pm 30 4.109 \pm 20	3.942 \pm 15 4.111 \pm 15	3.953 \pm 10 4.116 \pm 10	3.946 \pm 15 4.107 \pm 15 (4.328 \pm 15) 4.396 \pm 15	4.055 \pm 15		3.945 \pm 12 4.111 \pm 12 4.333 \pm 12 4.402 \pm 12	3.951 \pm 6 4.118 \pm 5	3.944 \pm 3		< 15 2	k 2	($\frac{1}{2}^-$) ($\frac{3}{2}^+$, $\frac{5}{2}^+$) ⁺	
(4.586 \pm 40) (4.706 \pm 40)			4.421 \pm 10 4.599 \pm 10 4.725 \pm 10 4.954 \pm 10 5.107 \pm 10	4.396 \pm 15			4.584 \pm 12 4.707 \pm 12 4.998 \pm 12				75 \pm 15 0 k	k 0 k	$\frac{5}{2}^+$	
(5.165 \pm 40)		5.153 \pm 15 5.447 \pm 15	5.172 \pm 10				5.148 \pm 12 5.460 \pm 12 5.502 \pm 12				< 15 310 \pm 30 k	0 (2) k	$\frac{3}{2}^+$	
5.45 ^g			5.582 \pm 10				5.714 \pm 12 6.280 \pm 12				< 15 k (3)	k (3)		
5.707 \pm 35 6.279 \pm 30		(5.708 \pm 15) 6.282 \pm 15 6.480 \pm 15 6.560 \pm 15 6.899 \pm 15 6.997 \pm 15 7.117 \pm 15 7.248 \pm 15									< 15 k (3)	k (3)		

9

a $^{18}\text{O}(d, p)^{19}\text{O}$: proton spectra measurements. b $^{17}\text{O}(t, p)^{19}\text{O}$: proton spectra measurements. c $^{18}\text{O}(d, p)^{19}\text{O}$: γ -ray measurements. d $^{17}\text{O}(t, p)^{19}\text{O}$: γ -ray measurements. e (1965WI1B, 1966WI05).

f See also (1961AR06, 1963YA03, 1964MO24, 1965MO16, 1965MO19, 1969FI07). g Unresolved. h Observed but group was weak.

i Other values for this level are $E_x = 96 \pm 11$ (1959AJ76), 96 ± 6 (1961JA23), 95 ± 2 (1963GI12) and 96.0 ± 0.5 keV (1971MC11). j (1971MC11) find $E_x = 1472 \pm 1$ keV. k Angular distribution for l_n or L not obtained.

^l See also (1969FI07). ^m See however (1971HI06).

Table 19.4: Resonances in $^{18}\text{O}(n, n)^{18}\text{O}$ ^a

E_{res} (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	$^{19}\text{O}^*$ (MeV)	$J\pi$	Refs.
0.661 ± 10	52 ± 3	4.583	$\frac{3}{2}^-$	(1964DO08, 1965VA03)
1.192 ± 10	49 ± 5	5.086	$\frac{1}{2}^-$	(1964DO08, 1965VA03)
1.256 ± 10	3.4 ± 1.0	5.146	$\frac{3}{2}$	(1965VA03)
1.56	470 ± 80	5.43	$\frac{5}{2}^+$	(1964DO08, 1965VA03)
1.66	380 ± 100	5.53	$(\frac{1}{2}^-)$	(1964DO08)
1.840 ± 10	7.8 ± 1.4	5.699	$\frac{3}{2}$	(1965VA03)
2.30	150 ± 40	6.13	$\frac{3}{2}^+$	(1964DO08, 1965VA03)
2.37	140 ± 40	6.20	$\frac{1}{2}^-$	(1964DO08, 1965VA03)
2.445 ± 10	19.2 ± 2.4	6.272	$\frac{7}{2}^-$	(1964DO08, 1965VA03)
≈ 2.58		(6.40)		(1965SA1A)
(2.63)		(6.45)		(1965SA1A)

^a See also (1959SC30, 1965SA1A) for evidence of additional structures.

4. $^{18}\text{O}(n, \gamma)^{19}\text{O}$

$$Q_m = 3.957$$

In the energy interval $E_n = 0.2$ to 1.0 MeV, the capture cross section displays a resonance at $E_n \approx 0.6$ MeV (1964VA02). Astrophysical considerations are discussed by (1968FO1A, 1970CL1C).

5. $^{18}\text{O}(n, n)^{18}\text{O}$

$$E_b = 3.957$$

The coherent scattering amplitude (bound) is $a = 6.00$ fm (1969BA1P). The total cross section measured for $E_n = 0.14$ to 2.47 MeV shows four resonances at $E_n = 0.66, 1.19, 1.26, 1.84$ and 2.45 MeV [see Table 19.4], in addition to a broad maximum at $E_n = 1.6$ MeV and resonance structure near 2.3 MeV (1965VA03). A phase-shift analysis by (1964DO08) is consistent with the (1965VA03) results and suggests that the two broad structures can each be accounted for in terms of two levels whose parameters are displayed in Table 19.4. At higher energies [$E_n = 2.82$ to 4.17 MeV (1959SC30), 2.45 to 8.50 MeV and 10.6 to 19.0 MeV (1965SA1A)] the total cross section shows additional structures. At $E_n = 14.1$ MeV, $\sigma_t = 1610 \pm 70$ mb (1969ME15). See also (1964EX1A) and ^{18}O .

6. $^{18}\text{O}(n, d)^{17}\text{N}$

$$Q_m = -13.718$$

$$E_b = 3.957$$

See (1964AM02).

$$7. \text{}^{18}\text{O}(\text{n}, \alpha)\text{}^{15}\text{C} \quad Q_{\text{m}} = -5.009 \quad E_{\text{b}} = 3.957$$

The total cross sections for the α_0 and α_1 groups have been measured for $E_{\text{n}} = 7.5$ to 8.6 MeV: resonance structure is reported at $E_{\text{n}} = 7.70 \pm 0.05$ and 8.05 ± 0.05 MeV with $\Gamma_{\text{lab}} = 0.25$ and 0.35 MeV, respectively [$^{19}\text{O}^*(11.25, 11.58)$] (1967ST28). See also (1964GA1A).

$$8. \text{}^{18}\text{O}(\text{d}, \text{p})\text{}^{19}\text{O} \quad Q_{\text{m}} = 1.732 \\ Q_0 = 1.734 \pm 0.010 \text{ (1965MO16);} \\ Q_0 = 1.727 \pm 0.008 \text{ (1967SP09).}$$

Angular distributions of proton groups have been measured at $E_{\text{d}} = 0.8$ MeV (1957SJ66), 5 MeV (1969FI07), 5.56 MeV (1964MO25, 1965MO16), 7.0 MeV (1964WI05), 10 MeV (1964EL1A), 12 MeV (1965WI1B, 1966WI05, 1971FO18), 12.3 MeV (1965LU1A), 14.95 MeV (1963YA03) and 15 MeV (1961AR06). l_{n} values derived from these measurements, as well as other level parameters for ^{19}O states below $E_{\text{x}} = 7.3$ MeV, are displayed in Table 19.3. See also (1959AJ76) and reaction 23 in ^{19}F .

Branching ratios for the decays of $^{19}\text{O}^*(0.096, 1.47, 2.37, 2.78, 3.16, 3.95)$ are shown in Table 19.5 (1965AL13, 1969FI07, 1970FI08, 1971BR02, 1971HI06, 1971MC11). Angular correlation measurements are consistent with $J^{\pi} = \frac{5}{2}^{+}$ for the ground state and unambiguously fix $J^{\pi} = \frac{3}{2}^{+}$ and $\frac{1}{2}^{+}$, respectively, for $^{19}\text{O}^*(0.096, 1.47)$ (1965AL13). They also suggest $J = \frac{9}{2}$ for $^{19}\text{O}^*(2.37)$ and $\frac{7}{2}^{+}$ for $^{19}\text{O}^*(2.78)$ (1971HI06). These assignments are also consistent with the lifetime measurements for these two states shown in Table 19.2 (1959ZI16, 1965MC10, 1971BR02, 1971HI06, 1971MC11). Measurement of the internal conversion coefficient for the 0.096 MeV transition shows that it is M1 (1963GI12).

See also (1959WI39, 1961LO10, 1964MA57) and (1963RO12, 1966AR10, 1967BU23, 1967ST1H, 1968ST1K, 1968SU1C, 1968YA03; theor.).

$$9. \text{}^{18}\text{O}(\text{t}, \text{d})\text{}^{19}\text{O} \quad Q_{\text{m}} = -2.301$$

Not reported.

$$10. \text{}^{18}\text{O}(\alpha, \text{}^3\text{He})\text{}^{19}\text{O} \quad Q_{\text{m}} = -16.621$$

Table 19.5: Radiative decays in ^{19}O

E_i (MeV)	J_i^π	E_f (MeV)	Branch ^a (%)	Refs.
0.096	$\frac{3}{2}^+$	0	100	
1.47	$\frac{1}{2}^+$	0	3.5 ± 1.6	(1965AL13)
			1.5 ± 0.6	(1970FI08, 1971HIZF)
			1.4 ± 0.2	(1971MC11)
			2.0 ± 0.2	(1971BR02)
		0.096	97 ± 2	(1969FI07)
			96.5 ± 1.6	(1965AL13)
			98.5 ± 0.6	(1970FI08, 1971HIZF)
			98.6 ± 0.2	(1971MC11)
2.37 ^b	$(\frac{3}{2}^+)$	0	100	(1970FI08, 1971HIZF)
		0.096	< 11	(1970FI08, 1971HIZF)
		1.47	< 12	(1970FI08, 1971HIZF)
2.78	$\frac{9}{2}^+$	0	100	(1970FI08, 1971HIZF)
		0.096	< 10	(1970FI08, 1971HIZF)
		1.47	< 5	(1970FI08, 1971HIZF)
		2.37	< 7	(1970FI08, 1971HIZF)
3.07		0.096		(1971HI06)
		1.47		(1971BR02)
3.16	$\frac{5}{2}^+$	0	8 ± 4	(1970FI08, 1971HIZF)
		0.096	92 ± 4	(1970FI08, 1971HIZF)
		1.47	< 6 ^c	(1970FI08, 1971HIZF)
		2.37	< 9	(1970FI08, 1971HIZF)
		2.78	< 10	(1970FI08, 1971HIZF)
3.95	$(\frac{1}{2}^-)$	0	24 ± 8	(1970FI08)
			33 ± 8	(1971HI06, 1971HIZF)
		0.096	48 ± 8	(1970FI08)
			39 ± 8	(1971HI06, 1971HIZF)
		1.47	28 ± 4	(1970FI08)
			28 ± 4	(1971HI06, 1971HIZF)
		2.37	< 15	(1970FI08)
		2.78	< 15	(1970FI08)
		3.16	< 15	(1970FI08)

^a The last value listed is believed to be the most reliable.

^b See also (1969FI07).

^c See, however, (1971BR02).

Not reported.

$$11. {}^{18}\text{O}({}^{13}\text{C}, {}^{12}\text{C}){}^{19}\text{O} \quad Q_{\text{m}} = -0.990$$

See (1971KN05).

$$12. {}^{19}\text{F}(\text{n}, \text{p}){}^{19}\text{O} \quad Q_{\text{m}} = -4.036$$

Angular distributions are reported at $E_{\text{n}} = 14.1$ MeV (1968FA01; $p_0, p_1, p_2, p_{\text{unres.}}$) and 14.4 MeV (1968AN1F, 1968RE07; $p_{0+1}, p_2, p_{\text{unres.}}$). See also (1961KO06, 1970FO1G), (1959AJ76) and ${}^{20}\text{F}$.

$$13. {}^{19}\text{F}(\text{t}, {}^3\text{He}){}^{19}\text{O} \quad Q_{\text{m}} = -4.800$$

Not reported.

$$14. {}^{21}\text{Ne}(\text{n}, {}^3\text{He}){}^{19}\text{O} \quad Q_{\text{m}} = -15.924$$

Not reported.

$$15. {}^{22}\text{Ne}(\text{n}, \alpha){}^{19}\text{O} \quad Q_{\text{m}} = -5.711$$

Not reported.

¹⁹F
(Figs. 6 and 8)

GENERAL: (See also (1959AJ76) and (1969BE1V).)

Shell model: (1957WI1E, 1959BR1E, 1960GO1C, 1960TA1C, 1961BA1E, 1962BH09, 1962HO1C, 1963HA05, 1964BR1H, 1964DR1A, 1964HA21, 1964IN03, 1965BA1J, 1965YU1A, 1966BA2E, 1966IN01, 1966LA1E, 1966RI1F, 1967AR02, 1967BO09, 1967EN01, 1967FL13, 1967GU05, 1967GU1D, 1967HA1M, 1967IN03, 1968AR02, 1968CO11, 1968EI1B, 1968EL1C, 1968GU1E, 1968GU1F, 1968GU1G, 1968HA17, 1968HA16, 1968HA26, 1968HA1P, 1968HI08, 1968MO1G, 1968WO1A, 1969BE1V, 1969GU1E, 1969KU1G, 1969MA1T, 1969SE02, 1969SE03, 1969TA1C, 1970EL1G, 1970GI11, 1970MC1J, 1970OL1A, 1971AR25, 1971AR1R, 1971GU1N, 1971LE30, 1971WI01, 1972LE1L).

Collective model: (1960RA1A, 1962DA1B, 1963CH02, 1964HA21, 1966RI1F, 1966ZA1B, 1966ZA1C, 1967BR1H, 1967RI1B, 1967SH1G, 1967ZA1C, 1968EI1B, 1968HA16, 1968RI1M, 1968UN1A, 1969BE1V, 1969BH03, 1969CU07, 1969KR02, 1970WA02, 1972LE1L).

Cluster model: (1959WI1B, 1960SH1A, 1963MA1D, 1964MA1G, 1968ME13, 1969BA2E, 1969HI1B, 1969ME1C, 1969TA1C, 1970BA2B).

Electromagnetic transitions: (1960RA1A, 1960RO1C, 1963BE03, 1963CH02, 1964BR1H, 1964DR1A, 1964HA21, 1966MA44, 1967GU1D, 1967IN03, 1967ZA1C, 1968BA2K, 1968EL1C, 1968GU1G, 1968HA17, 1969BE1V, 1969CU07, 1969HI1B, 1971AR25, 1971GU1N).

Pion and muon reactions [†]: (1967BA78, 1967KA1C, 1967MI1B, 1968BA2G, 1968WI1B, 1969CH1C, 1969MO1E, 1969PL1B, 1969PO07, 1970BA1E, 1970BL07, 1971FA09, 1971TH05).

Special levels: (1959LA1A, 1960SH1A, 1961TR1B, 1964DR1A, 1964HA21, 1966BR1H, 1966MA44, 1967EN01, 1967SH1G, 1968AR02, 1968BA2K, 1968CE1A, 1968HA1P, 1968SO1A, 1968YA03, 1969BE1V, 1969CU07, 1969GA1G, 1969HA1G, 1969HA1F, 1971AR1R).

Special reactions: (1967AU1B, 1968MI05).

Astrophysical questions: (1970BA1M).

Other theoretical topics: (1964IN03, 1964KA1A, 1965AM1A, 1965FA1B, 1965GO1F, 1965OK1A, 1965RO1H, 1965WO1C, 1965YO1A, 1966DA1E, 1966DO1C, 1966GI1A, 1966SU1D, 1967FL13, 1967GU05, 1967NE1D, 1967OS02, 1968DW1A, 1968GA03, 1968GU1E, 1968GU1C, 1968GU1G, 1968MO1G, 1968PA1Q, 1969LO06, 1969MA1T, 1971LE1H, 1972LE1L).

Ground state: $\mu = +2.628383 \pm 0.000005$ nm (1964BA11: see also (1969FU11)). See (1963BE36, 1963CA1B, 1964BO37, 1964BR1H, 1964DR1A, 1964LI14, 1964ST1B, 1965HU13, 1966RI12, 1967CO1D, 1967GU1D, 1967PA1G, 1967SH14, 1968EL1C, 1968HI08, 1968LE1L, 1969HI1B, 1969PE1D, 1969WU1A, 1970LE1A, 1971AR25, 1971GU1N, 1971WI01, 1972LE1L).

[†] Gamma rays with $E_\gamma = 109.8 \pm 0.1$ and 197.0 ± 0.1 keV (1969PO07), 109.8 ± 0.2 and 197.98 ± 0.19 keV (1970BL07) are reported in pion-induced reactions.

Table 19.6: Energy levels of ^{19}F ^a

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^+; \frac{1}{2}$	$\frac{1}{2}^+$			1, 3, 4, 9, 11, 12, 13, 23, 27, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 49, 50, 52
0.109893 ± 0.004	$\frac{1}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 0.849 \pm 0.010$ nsec	γ	9, 11, 12, 13, 23, 25, 26, 27, 32, 33, 34, 36, 37, 38, 40, 41, 50
0.197147 ± 0.012	$\frac{5}{2}^+$	$\frac{1}{2}^+$	$\tau_m = 128.8 \pm 1.5$ nsec $\mu = 3.69 \pm 0.04$ nm $Q = -(0.10 \pm 0.02)$ bn	γ	7, 11, 12, 13, 23, 26, 31, 32, 33, 34, 38, 40, 41, 50
1.34572 ± 0.11	$\frac{5}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 4.8 \pm 0.4$ psec	γ	9, 11, 12, 13, 23, 31, 32, 33, 34, 37
1.4585 ± 0.4	$\frac{3}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 78 \pm 12$ fsec	γ	9, 11, 12, 13, 23, 27, 32, 33, 34
1.5541 ± 0.2	$\frac{3}{2}^+$	$\frac{1}{2}^+$	$\tau_m = 4.4_{-2.0}^{+2.4}$ fsec	γ	11, 12, 13, 23, 26, 31, 32, 33, 34, 40
2.7798 ± 0.6	$\frac{9}{2}^+$	$\frac{1}{2}^+$	$\tau_m = 261 \pm 24$ fsec	γ	5, 7, 11, 12, 13, 18, 22, 23, 32, 33, 46, 47, 49
3.9071 ± 1.0	$\frac{3}{2}^+(+)$	$\frac{3}{2}^+$	$\tau_m \leq 20$ fsec	γ	12, 13, 18, 23, 33, 47
3.9985 ± 0.8	$\frac{7}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 23 \pm 10$ fsec	γ	9, 12, 13, 23, 33
4.0325 ± 1.0	$\frac{9}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 73 \pm 11$ fsec	γ	9, 12, 13, 23, 33, 47
4.3777 ± 1.0	$\frac{7}{2}^+$	$\frac{1}{2}^+$		γ	7, 12, 13, 23, 26, 33, 47, 49
4.555 ± 5	$\frac{5}{2}^+$	$\frac{3}{2}^+$	$\tau_m \leq 20$ fsec	γ	12, 23, 47
4.5575 ± 1.0	$\frac{3}{2}^-$		$\tau_m \leq 20$ fsec	γ	12, 33, 46, 47
4.648	$\frac{13}{2}^+$	$\frac{1}{2}^+$	$\tau_m = 2.2 \pm 0.3$ psec	γ	12, 37
4.683 ± 1	$\frac{5}{2}^-$		$\tau_m = 15.4 \pm 3.0$ fsec	γ, α	7, 23, 33, 47
5.106 ± 3	$\frac{5}{2}^(-)$			γ, α	7, 12, 22, 23, 33, 47, 49
5.340 ± 5	$\frac{1}{2}^-$		$\tau_m \leq 15$ fsec $\Gamma < 2$ keV	γ, α	7, 12, 23, 33, 47
5.428 ± 8	$\frac{7}{2}^-$			γ, α	7, 9, 12, 23, 33
5.464 ± 2	$\frac{7}{2}^+$	$\frac{3}{2}^+$	$\tau_m \leq 19$ fsec $\Gamma < 1$ keV	γ, α	7, 12, 22, 23
5.499 ± 3	$\frac{3}{2}^+$			γ, α	7, 8, 22, 23, 33
5.540 ± 5	$\frac{5}{2}^+, \frac{7}{2}^-$			γ, α	7, 23, 33, 47
5.630 ± 10	$(\frac{1}{2}, \frac{3}{2})^-$			γ, α	7, 22, 33, 46, 47
5.943 ± 5	$\frac{1}{2}^(+)$			γ, α	7, 22, 23, 33, 47
6.076 ± 6	$(\frac{7}{2}^+)$		1.2	γ, α	7, 8, 22, 33
6.093 ± 4	$(\frac{3}{2}^-)$		4	γ, α	7, 8, 22, 23, 33, 47
6.167 ± 5	$\frac{7}{2}^(-)$			γ, α	7, 23, 33, 47
6.250 ± 5	$\frac{1}{2}^+$		8	α	8, 22, 23, 33, 47
6.290 ± 7	$\frac{5}{2}^+$		2.4	γ, α	7, 8, 22, 33

Table 19.6: Energy levels of ^{19}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
6.332 \pm 7	$\frac{7}{2}^+$		2.4	γ, α	7, 8, 33
6.43	$\frac{1}{2}^-$		280	α	8
6.497 \pm 2	$\frac{3}{2}^+(+)$			γ, α	7, 23
6.499 \pm 2	$\frac{11}{2}^+$	$\frac{1}{2}^+$		γ, α	7
6.525 \pm 2	$\frac{3}{2}^+$		4	γ, α	7, 8
6.553 \pm 2	$\frac{7}{2}^+$		1.6	γ, α	7, 8
6.592 \pm 2	$\frac{9}{2}^+$	$\frac{3}{2}^+$		γ, α	7, 23
6.788 \pm 3	$\frac{3}{2}^-$		2.4	γ, α	7, 8, 22, 23, 46
6.838 \pm 5	$(\frac{5}{2}, \frac{3}{2})^+$		1.2	γ, α	7, 8
6.890 \pm 10	$\frac{5}{2}^-$		28	α	8, 22
6.926 \pm 5	$\frac{7}{2}^-(+)$		2.4	γ, α	7, 8, 22, 23
6.99	$\frac{1}{2}^-$		51	α	8
7.11	$\frac{7}{2}^+$		≈ 32	α	8, 23
7.167 \pm 3	$\frac{11}{2}^- (\frac{7}{2}^-)$			γ, α	7
7.12	$\frac{3}{2}^+$		≈ 8	α	8, 23
7.257 \pm 5			≈ 6	γ, α	7, 8, 22, 23
7.364 \pm 5	$\frac{1}{2}^+$		(≈ 63)	(α)	8, 22, 23
7.539 \pm 2	$\frac{5}{2}^+; \frac{3}{2}^+$			γ, α	7, 23
7.660 \pm 2	$\frac{3}{2}^+; \frac{3}{2}^+$			γ, α	7, 23, 31, 49
7.702 \pm 5	$(\frac{3}{2}^-)$		≈ 24	α	8, 22, 23
7.762 \pm 5			≈ 6	γ, α	7, 8
7.90			≈ 210	α	8
7.928 \pm 3	$(\frac{7}{2}^+, \frac{9}{2}^+)$			γ, α	7
7.935 \pm 3	$\frac{11}{2}^+$	$\frac{3}{2}^+$		γ, α	7
(7.961)			(≈ 6)	α	8
(7.972)			(≈ 4)	α	8
8.014 \pm 5	$\frac{5}{2}^+$		≈ 6	γ, α	7, 8, 23
8.086 \pm 5	$(\frac{5}{2}^+)$		≈ 6	α	8, 23
8.136 \pm 5	$(\frac{1}{2}^+)$		≈ 5	α	8, 22, 23
8.155			≈ 51	α	8
8.195 \pm 5	$(\frac{5}{2}^+)$		≈ 8	α	8, 23
8.255 \pm 5	$(\frac{5}{2}^+)$				23
8.287 \pm 3	$(\frac{13}{2}^-)$				7
8.320 \pm 10			≈ 8	α	8, 23
8.590 \pm 1	$\frac{3}{2}^+$		2.0 \pm 0.1	γ, p	18, 20, 23
8.637	$\frac{1}{2}^+$		95	p	20
8.795 \pm 1.3	$\frac{1}{2}^+; \frac{3}{2}^+$		45 \pm 1	γ, p, α	18, 20, 21, 23
8.928 \pm 0.8	$\frac{3}{2}^-$		3.6 \pm 0.2	p, α	20, 21

Table 19.6: Energy levels of ^{19}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
8.942 \pm 6	$(\frac{11}{2}^-)$				7, 9
9.0984 \pm 0.6	$\frac{7}{2}^+; \frac{3}{2}$		$(20 \pm 19) \times 10^{-3}$	γ, p, α	18, 20, 21, 23
9.166 \pm 1	$\frac{1}{2}^+$		5.8 \pm 0.3	p, (α)	20, 21, 23
9.321 \pm 1	$\frac{1}{2}^+$		4.9 \pm 0.2	γ, p, α	18, 20, 21
9.527 \pm 6	$(\frac{5}{2})$		29	p, α	20, 21
9.578 \pm 7	$\frac{3}{2}$		26	p, α	20, 21, 23
9.668 \pm 2	$\frac{3}{2}^+$		3.8 \pm 1.0	γ, p, α	18, 20, 21, 23
9.819 \pm 0.8	$\frac{5}{2}^-$		0.29 \pm 0.05	γ, p, α	18, 20, 21
9.888 \pm 4	$\frac{1}{2}^+$		31	p, α	20, 21
10.136 \pm 0.8	$\frac{3}{2}^-$		4.7 \pm 1.0	γ, p, α	18, 21
10.161 \pm 3	$\frac{1}{2}^+$		31	p, α	20, 21
10.231 \pm 3	$\frac{1}{2}^+$		4.3	(γ), p, α	18, 20, 21
10.253 \pm 3	$\frac{3}{2}^+$		23	(γ), p, α	18, 20, 21, 23
10.306 \pm 3	$\frac{3}{2}^+$		9.2	(γ), p, α	18, 20, 21, 23
10.496 \pm 1	$\frac{3}{2}^+$		6.2 \pm 0.5	(γ), n, p, α	18, 19, 20, 21
10.541 \pm 1			2.5 \pm 0.2	n, p	19, 23
10.554 \pm 3	$(\frac{3}{2}^+; \frac{3}{2})$		7.6 \pm 2	(γ), p, α	18, 20, 21
10.566 \pm 1.0			5.2 \pm 0.5	n, p	19
(10.580 \pm 4)	$(\frac{5}{2}^+)$		22 \pm 3	(γ), p, α	18, 20, 21
10.613 \pm 1.5	$\frac{5}{2}^+; \frac{3}{2}$		4.8 \pm 0.5	n, p, α	19, 20, 21
10.763 \pm 3	$\frac{1}{2}^-$		5.4	p, α	20, 21
10.858 \pm 2	$\frac{5}{2}^+$		25 \pm 1.5	n, p, α	19, 20, 21
10.972 \pm 3	$(\frac{3}{2}, \frac{5}{2})^+$		\approx 11	n, p, α	19, 20, 21
10.988 \pm 3			\approx 10	n, p	19
11.070 \pm 3	$\frac{1}{2}^+$		31 \pm 5	n, p, α	19, 20, 21
11.199 \pm 2	$(\frac{1}{2}^-)$		43 \pm 2	n, p, α	19, 20, 21
11.288 \pm 9			24	n, p, α	19, 20, 21
11.310 \pm 9				n, p, α	19, 21
11.43 \pm 10			81 \pm 19	n, p, α	19, 21
11.51				n, p, α	19, 20, 21
11.56 \pm 10			< 48	n, p, α	19, 20, 21
11.65 \pm 10			38	n, p, α	19, 20, 21
11.68				p, α	20, 21
11.78				n, p, α	19, 20, 21
11.84 \pm 10			< 48	n, p	19
11.93 \pm 10			85	n, p, α	19, 20, 21
12.05				p, α	21
12.09				n, p, α	19, 20, 21

Table 19.6: Energy levels of ^{19}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	$\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
12.13 \pm 10	$T = \frac{3}{2}$		95	n, p, α	19, 20, 21
12.23			48	n, p, α	19, 20, 21
12.33			n, p, α	19, 20, 21	
12.44			n, p, t, α	10, 19, 20, 21	
12.52			14	p	20
12.57			n, p, α	19, 20, 21	
12.63			n, p, α	19, 20, 21	
12.69			p, t, α	10, 20, 21	
12.75			p, t, α	10, 21	
12.77 \pm 10			190	n, p, (t), α	10, 19, 20, 21
12.92			n, p, t, α	10, 19, 20, 21	
13.06			n, p, t, α	10, 19, 20, 21	
13.17 \pm 15			67	n, p, t, α	10, 19, 21
13.31 \pm 15			20	n, p, α	19, 20, 21
13.45			n, p, α	19, 21	
13.58			p, α	20, 21	
13.73 \pm 15			95	n, p, α	19, 20, 21
14.24 \pm 15			330	n, p	19
14.29			p, α	20, 21	
14.78 \pm 20			280	n, p	19
15.00 \pm 20	n, p	19			
15.75 \pm 25	150	n, p	19		
16.32 \pm 25	190	n, p, α	19, 21		
16.80 \pm 30	n, p	19			
17.9	p, α	20, 21			

^a See also Tables 19.9 and 19.10 and (1969BE1V; theor.).

1. ${}^9\text{Be}({}^{14}\text{N}, \alpha){}^{19}\text{F}$ $Q_m = 13.273$

See (1958GO71).

2. (a) ${}^{12}\text{C}({}^7\text{Li}, \text{n}){}^{18}\text{F}$ $Q_m = 5.964$ $E_b = 16.395$

(b) ${}^{12}\text{C}({}^7\text{Li}, \text{p}){}^{18}\text{O}$ $Q_m = 8.402$

(c) ${}^{12}\text{C}({}^7\text{Li}, \text{d}){}^{17}\text{O}$ $Q_m = 2.580$

- (d) $^{12}\text{C}(^7\text{Li}, \text{t})^{16}\text{O}$ $Q_m = 4.695$
 (e) $^{12}\text{C}(^7\text{Li}, \alpha)^{15}\text{N}$ $Q_m = 12.382$

The yield of ^{18}F (reaction (a)) has been determined for $E(^7\text{Li}) = 2.5$ to 3.5 MeV (1961NO05). The yields of the α_0 , α_{1+2} and α_3 groups (reaction (e)) have been measured for $E(^7\text{Li}) = 3.2$ to 4.0 MeV by (1962HO06), while (1970CA1N) report that the cross sections for reactions (b), (c), (d) and (e) vary strongly over the range $E(^7\text{Li}) = 4$ to 14 MeV but with little, if any, cross-correlation. A coherence energy of about 0.4 MeV is reported (1970CA1N). See also (1967MO23), ^{18}O , ^{16}O and ^{17}O in (1971AJ02) and ^{15}N in (1970AJ04).

3. $^{12}\text{C}(^{11}\text{B}, \alpha)^{19}\text{F}$ $Q_m = 7.730$

See (1966KU1D).

4. $^{12}\text{C}(^{19}\text{F}, ^{12}\text{C})^{19}\text{F}$

The angular distributions corresponding to the reaction $^{19}\text{F}(0)$ and $^{12}\text{C}(0)$ have been measured at $E(^{19}\text{F}) = 40$ and 60 MeV (1968VO1A, 1969VO10). See also (1971BO1V).

5. $^{14}\text{N}(^6\text{Li}, \text{p})^{19}\text{F}$ $Q_m = 11.148$

At $E(^6\text{Li}) = 5.3$, 5.6 and 6.0 MeV angular distributions have been obtained for the protons corresponding to $^{19}\text{F}^*(2.78)$ (1968RI13).

6. $^{14}\text{N}(^7\text{Li}, \text{d})^{19}\text{F}$ $Q_m = 6.123$

See (1968MI09).

7. $^{15}\text{N}(\alpha, \gamma)^{19}\text{F}$ $Q_m = 4.013$

Thirty resonances in the yield of capture γ -rays are observed below $E_\alpha = 5.1$ MeV: the parameters for these are displayed in Table 19.7 (1957PR1A, 1969AI01, 1969AI02, 1970AI01, 1971DI09, 1971DI18, 1972RO01). Branching ratios are shown in Table 19.9 (1969AI01, 1969AI02, 1970AI01, 1971DI18, 1972RO01), and lifetime measurements are listed in Table 19.10 (1968TO11, 1968TOZU, 1972RO01). See also (1967TO1C).

Table 19.7: Resonances in $^{15}\text{N}(\alpha, \gamma)^{19}\text{F}$ ^a

E_α (MeV \pm keV)	Γ (keV)	$\frac{1}{2}(2J+1)\Gamma_\gamma\Gamma_\alpha/\Gamma$ (eV)	J^π ^c	E_x (MeV)	Refs.
0.850 \pm 1	$(43.8 \pm 8.5) \times 10^{-6}$	$(6 \pm 1) \times 10^{-3}$ ^b	$\frac{5}{2}^-$	4.684	(1970AI01, 1972RO01)
1.385 \pm 3		$(13 \pm 8) \times 10^{-3}$ ^b	$\frac{5}{2}^+$	5.106	(1970AI01)
1.681 \pm 5	< 2	1.64 ± 0.16	$\frac{1}{2}^-$	5.340	(1957PR1A, 1970AI01, 1971DI09)
1.790		0.42 ± 0.09 ^b	$\frac{7}{2}^-$	5.426	(1970AI01)
1.839 \pm 2	< 1	2.5 ± 0.4 ^b	$\frac{7}{2}^+$	5.464	(1957PR1A, 1971DI18)
1.883 \pm 3	4 ± 1	4.2 ± 1.1 ^b	$\frac{3}{2}^+$	5.499	(1957PR1A)
1.930		0.48 ± 0.11 ^b	$\frac{5}{2}^+$	5.54	(1970AI01)
2.05				5.63	(1970AI01)
2.44			$\frac{1}{2}^-$	5.94	A, (1970AI01)
2.61			$(\frac{7}{2}^+)$	6.07	A, (1970AI01)
2.64			$(\frac{3}{2}^-)$	6.10	A, (1970AI01)
2.74			$\frac{7}{2}^-$	6.18	A, (1970AI01)
2.88			$\frac{5}{2}^+$	6.29	(1970AI01)
2.95			$\frac{7}{2}^+$	6.34	(1970AI01)
3.147 \pm 1.5		1.7 ± 0.3	$\frac{3}{2}^+$	6.497	B, (1970AI01)
3.150 \pm 1.5		2.3 ± 0.4	$\frac{11}{2}^+$	6.499	B, (1969AI01, 1970AI01)
3.183 \pm 1.5		2.35 ± 0.35	$\frac{3}{2}^+$	6.525	B
3.218 \pm 1.5		0.57 ± 0.12	$\frac{7}{2}^-$	6.553	B
3.268 \pm 1.5		1.64 ± 0.25	$\frac{9}{2}^+$	6.592	B, (1971DI18)
3.515 \pm 3		10.9 ± 1.5	$\frac{3}{2}^-$	6.787	B
3.58 \pm 5				6.838	B
3.691 \pm 3		9.7 ± 1.4	$\frac{7}{2}^-$	6.926	B
3.996 \pm 3		1.15 ± 0.23	$\frac{11}{2}^-$, $(\frac{7}{2}^-)$	7.167	B
4.11 \pm 5				7.257	B
4.468 \pm 3		17.0 ± 2.7	$\frac{5}{2}^+$; $T = \frac{3}{2}$	7.539	A, B, (1969AI02)
4.623 \pm 3		3.7 ± 0.9	$\frac{3}{2}^+$; $T = \frac{3}{2}$	7.662	A, B, (1969AI02)
4.75 \pm 5				7.762	B
4.96 \pm 3		2.3 ± 0.4	$\frac{7}{2}^+$; $\frac{9}{2}$	7.928	B, (1971DI18)
4.97 \pm 3		3.1 ± 0.5	$\frac{11}{2}^+$	7.935	A, B, (1971DI18)
5.07 \pm 5				8.014	B
d				8.287	C
d				8.942	C

A: D. Rogers, private communication.

B: W.R. Dixon and R.S. Storey, private communication.

C: K. Bharuth-Ram, K.P. Jackson, N.A. Jelley, P.G. Lawson and K.W. Allen, private communication.

^a See also Table 19.9.

^b Recalculated by D. Rogers (private communication) on basis of results of (1971DI09) for $^{19}\text{F}^*(5.34)$.

^c See (1970AI01) for tentative assignments to rotational bands.

^d Resonant energies determined to ± 3 and ± 6 keV, respectively. $^{19}\text{F}^*(8.29)$ decays only to the $J^\pi = \frac{9}{2}^-$ state at 4.03 MeV and γ -ray angular distribution studies establish $J = \frac{13}{2}$ or $\frac{9}{2}$. The former is favored since no other branches are observed. Of $J^\pi = \frac{13}{2}^-$, and with $\omega\gamma = 1.02 \pm 0.21$ eV, the strength of the transition to $^{19}\text{F}^*(4.03)$ is 21.1 ± 4.3 W.u. $^{19}\text{F}^*(8.94)$ is weaker than expected for the $\frac{11}{2}^-$ member of the $K^\pi = \frac{1}{2}^-$ band (K.W. Allen, private communication).

Correlation experiments on the transitions $5.47 \rightarrow 2.78 \rightarrow 0.197$ fix $J = \frac{9}{2}$ for $^{19}\text{F}^*(2.78)$ and $J = \frac{7}{2}$ and $\frac{1}{2}$ for $^{19}\text{F}^*(5.46, 5.34)$ (1968TO11, 1968TOZU). See, however, reaction 23. Measurements of γ -ray angular distributions and radiative widths lead to the J^π values listed in Table 19.7 (1969AI01, 1969AI02, 1970AI01, 1971DI18) as well as to $J = \frac{7}{2}$ for $^{19}\text{F}^*(4.39)$ (1969AI02). The capture yield for $E_\alpha = 4.2$ to 4.7 MeV is dominated by two resonances corresponding to the first two $T = \frac{3}{2}$ states [$^{19}\text{F}^*(7.54, 7.66)$] with $J^\pi = \frac{5}{2}^+$ and $\frac{3}{2}^+$ analogous to $^{19}\text{O}^*(0, 0.097)$ (1969AI02). The resonance at $E_\alpha = 4.97$ MeV corresponds to a $J = \frac{11}{2}$ state at 7.935 MeV: it appears to be the $\frac{11}{2}^+$ member of a $K = \frac{3}{2}^+$ rotational band based on the $\frac{3}{2}^+$ state at $E_x = 3.91$ MeV (1971DI18). $^{19}\text{F}^*(6.09)$ has a very definite non-isotropic decay and hence $J = \frac{1}{2}$ is ruled out: J^π is probably $\frac{3}{2}^-$. $^{19}\text{F}^*(5.95, 6.08, 6.17)$ are tentatively assigned $J^\pi = \frac{1}{2}^{(-)}, (\frac{7}{2}^+), \frac{7}{2}^{(-)}$ (D. Rogers, private communication) [see, however, Table 19.8].

8. $^{15}\text{N}(\alpha, \alpha)^{15}\text{N}$

$$E_b = 4.013$$

The elastic scattering has been studied for $E_\alpha = 1.75$ to 5.50 MeV: observed anomalies are shown in Table 19.8 (1959SM02, 1961SM02).

9. $^{15}\text{N}(^7\text{Li}, t)^{19}\text{F}$

$$Q_m = 1.546$$

This reaction has been studied at $E(^7\text{Li}) = 15$ and 20 MeV. The spectra have been compared with those from the $^{16}\text{O}(^7\text{Li}, t)^{20}\text{Ne}$ reaction. The similarities suggest that $^{19}\text{F}^*(0.11)$ [$J^\pi = \frac{1}{2}^-$] arise from the coupling of a $p_{1/2}$ hole with $^{20}\text{Ne}(0)$; that $^{19}\text{F}(1.46, 1.35)$ [$J^\pi = \frac{3}{2}^-$ and $\frac{5}{2}^-$] and $^{19}\text{F}^*(4.00, 4.04)$ [$J^\pi = \frac{7}{2}^-$ and $\frac{9}{2}^-$] correspond respectively to the 2^+ and 4^+ members of the ^{20}Ne ground state band. The $J^\pi = \frac{11}{2}^-$ and $\frac{13}{2}^-$ states in ^{19}F based on the 6^+ state of ^{20}Ne at 8.79 MeV are reported at 8.25 and 8.91 MeV (± 50 keV). [See, however, reaction 7.] Angular distributions for $^{19}\text{F}^*(0.11, 1.35, 1.46, 4.00 + 4.04, 5.43, 8.25, 8.91)$ have been measured (1970MI1E). See also (1971AR1R; theor.).

Table 19.8: Levels of ^{19}F from $^{15}\text{N}(\alpha, \alpha_0)^{15}\text{N}$ ^a

E_α (MeV \pm keV)	Γ_{lab} (keV)	J^π	E_x (MeV)
1.878 ± 10	4	$\frac{3}{2}^+$	5.495
2.614 ± 10	1.5	$\frac{5}{2}^+$	6.076
2.635 ± 10	5	$\frac{5}{2}^-$	6.093
2.833 ± 10	10	$\frac{1}{2}^+$	6.249
2.883 ± 10	3	$\frac{5}{2}^+$	6.288
2.944 ± 10	3	$\frac{7}{2}^+$	6.336
3.07	360	$\frac{1}{2}^-$	6.44
3.194 ± 10	5	$\frac{1}{2}^+$	6.534
3.229 ± 10	2	$\frac{5}{2}^+$	6.561
3.525 ± 10	3	$\frac{3}{2}^-$	6.795
3.587 ± 10	1.5	$(\frac{5}{2}, \frac{3}{2})^+$	6.844
3.648 ± 10	35	$\frac{5}{2}^-$	6.892
3.705 ± 10	3	$(\frac{9}{2}, \frac{7}{2})^-$	6.937
3.78	64	$\frac{1}{2}^-$	7.00
3.92	≈ 40	$\frac{7}{2}^+$	7.11
3.94	≈ 10	$\frac{3}{2}^+$	7.12
4.127 ± 10	≈ 8		7.270
4.22	≈ 80		7.34
4.49	≈ 90		7.56
(4.53)			(7.59)
4.700 ± 10	≈ 30		7.722
4.780 ± 10	≈ 8		7.785
4.93	≈ 260		7.90
(5.005)	(≈ 8)		(7.963)
(5.018)	(≈ 5)		(7.973)
5.116 ± 10	≈ 8		8.050
5.203 ± 10	≈ 8		8.119
5.232 ± 10	≈ 6		8.142
5.25	≈ 65		8.156
5.284 ± 10	≈ 10		8.183
5.481 ± 10	≈ 10		8.339

^a (1959SM02, 1961SM02).

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.
0.110	$\frac{1}{2}^-$	0	$\frac{1}{2}^+$	100		
0.197	$\frac{5}{2}^+$	0	$\frac{1}{2}^+$	100		
		0.110	$\frac{1}{2}^-$	< 0.06		(1970CO22)
1.35 ^a	$\frac{5}{2}^-$	0	$\frac{1}{2}^+$	< 3	$ M ^2 = 7.6 \pm 1.3$ W.u. (E3)	(1963LI01, 1969AL15)
		0.110	$\frac{1}{2}^-$	96.8 ± 1^b		C
				96.4 ± 3.0		(1969PO03)
		0.197	$\frac{5}{2}^+$	3.2 ± 1		C
				3.6 ± 1.0		(1969PO03)
1.46 ^{a,b}	$\frac{3}{2}^-$	0	$\frac{1}{2}^+$	22.5 ± 2		C
				21 ± 1		(1970LA02)
				20.0 ± 1		(1971HA30)
		0.110	$\frac{1}{2}^-$	66.6 ± 3		(1969PO03)
				68 ± 3		(1970LA02)
				69.7 ± 1		(1971HA30)
		0.197	$\frac{5}{2}^+$	10.9 ± 2		(1969PO03)
				10.2 ± 1		(1971HA30)
				11 ± 0.5		(1970LA02)
1.55 ^{a,b}	$\frac{3}{2}^+$	0	$\frac{1}{2}^+$	2.0 ± 0.7		(1969PO03)
				2.4 ± 0.5		(1966OL01)
					$ M ^2 = 6.8 \pm 0.7$ W.u. (E2)	(1969AL15)
		0.110	$\frac{1}{2}^-$	4.6 ± 0.5		(1959JO26)

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.						
2.78	$\frac{9}{2}^+$	0.197	$\frac{5}{2}^+$	5.3 ± 1	$ M ^2 = 7.7 \pm 1.5$ W.u.	(1969PO03)						
				5.2 ± 0.7		(1966OL01)						
				(95.4 ± 0.5)		(1954JO21)						
				92.7 ± 1		(1969PO03)						
		1.35	$\frac{5}{2}^-$	0.197		$\frac{5}{2}^+$	92.4 ± 0.9	(1966OL01)				
							≤ 4	(1954JO21)				
		3.91 ^c	$\frac{3}{2}^{(+)}$	0		$\frac{1}{2}^+$	100	(1966TH02, 1968SP01, 1969JA09, 1970CO22)				
							62	(1969WO1F)				
							79 ± 3	(1965AL20)				
							47 ± 5	C				
							0.110	$\frac{1}{2}^-$	0.110	$\frac{1}{2}^-$	12	(1969WO1F)
											< 10	(1965AL20)
							0.197	$\frac{5}{2}^+$	0.197	$\frac{5}{2}^+$	18 ± 4	C
											10	(1969WO1F)
1.35	$\frac{5}{2}^-$				1.35		$\frac{5}{2}^-$	< 2	(1965AL20)			
								16 ± 4	C			
		4	(1969WO1F)									
		< 4	(1965AL20), C									
		1.46	$\frac{3}{2}^-$	1.46		$\frac{3}{2}^-$		3	(1969WO1F)			
								< 4	(1965AL20), C			
1.55	$\frac{3}{2}^+$	1.55	$\frac{3}{2}^+$	16	(1969WO1F)							
				21 ± 3	(1965AL20)							

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	Refs.
4.00 ^c	$\frac{7}{2}^-$	2.78	$\frac{9}{2}^+$	18 ± 4	C
		0.110	$\frac{1}{2}^+$	< 2	(1965AL20)
		0.197	$\frac{5}{2}^+$	(22)	(1965AL20)
		1.35	$\frac{5}{2}^-$	18 ± 4	C
				78	(1965AL20)
				85	(1969WO1F)
				90	(1970AI01)
				70 ± 4	C
		1.46	$\frac{3}{2}^-$	15	(1969WO1F)
				10	(1970AI01)
4.03	$\frac{9}{2}^-$ $\frac{7}{2}^+$	1.35	$\frac{5}{2}^-$	12 ± 2	C
4.38		0	$\frac{1}{2}^+$	100	(1966TH02, 1970AI01)
		0.110	$\frac{1}{2}^-$	< 5	(1966OL01)
		0.197	$\frac{5}{2}^+$	< 2	(1966OL01)
				85 ± 5	(1966TH02)
				82 ± 7	(1966OL01)
		1.35 + 1.46		≤ 2	(1966OL01)
		1.55	$\frac{3}{2}^+$	< 4	(1966OL01)
		2.78	$\frac{9}{2}^+$	15 ± 5	(1966TH02)
				18 ± 7	(1966OL01)
4.555 ^c	$\frac{5}{2}^+$	0	$\frac{1}{2}^+$	< 5	C
		0.110	$\frac{1}{2}^-$	< 5	C
		0.197	$\frac{5}{2}^+$	69 ± 7	C

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.
4.558 ^c	$\frac{3}{2}^-$	1.35	$\frac{5}{2}^-$	5 ± 3		C
		1.46	$\frac{3}{2}^-$	8 ± 3		C
		1.55	$\frac{3}{2}^+$	18 ± 4		C
		0	$\frac{1}{2}^+$	36 ± 4		C
		0.110	$\frac{1}{2}^-$	45 ± 5		C
		0.197	$\frac{5}{2}^+$	69 ± 7		C
		1.35	$\frac{5}{2}^-$	5 ± 3		C
		1.46	$\frac{3}{2}^-$	< 4		C
4.65	$\frac{13}{2}^+$	1.55	$\frac{3}{2}^+$	6 ± 3	$ M ^2 = 5.5 \pm 1.8$ W.u.	C
		2.78	$\frac{9}{2}^+$	100		(1969BH01, 1969JA09, 1970AI01)
4.68 ^{c,d,m}	$\frac{5}{2}^-$	0.197	$\frac{5}{2}^+$	6	$\Gamma_\gamma = \Gamma = 0.95 \pm 0.05$	(1969WO1F, 1970AI01, 1972RO01)
			6 ± 1	(1972RO01)		
			4 ± 2	C		
		1.35	$\frac{5}{2}^-$	60		(1969WO1F)
			63	(1970AI01)		
			63 ± 6	(1972RO01)		
		1.46	$\frac{3}{2}^-$	64 ± 5		C
			34	(1969WO1F)		
			30	(1970AI01)		
			31 ± 3	(1972RO01)		
2.78	$\frac{9}{2}^+$	32 ± 3	C			
		< 1	(1970AI01)			

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	Refs.
5.11 ^d	$\frac{5}{2}^{(-)}$	0.197	$\frac{5}{2}^{+}$	< 2	(1972RO01)
		1.46	$\frac{3}{2}^{-}$	80	(1970AI01)
5.34 ^{d,e}	$\frac{1}{2}^f$	0	$\frac{1}{2}^{+}$	20	(1970AI01)
		0.110	$\frac{1}{2}^{-}$	37	(1970AI01)
		0.197	$\frac{5}{2}^{+}$	42	(1970AI01)
		1.46	$\frac{3}{2}^{-}$	< 2	(1970AI01)
5.43 ^d	$\frac{7}{2}^{-}$	1.35	$\frac{5}{2}^{-}$	20	(1970AI01)
		1.46	$\frac{3}{2}^{-}$	70	(1970AI01)
		4.00	$\frac{7}{2}^{-}$	13	(1970AI01)
		4.03	$\frac{9}{2}^{-}$	10	(1970AI01)
		0.197	$\frac{5}{2}^{+}$	6	(1970AI01)
5.46 ^{d,e}	$\frac{7}{2}^{-}$	1.35	$\frac{5}{2}^{-}$	4	(1971DII8)
		1.55	$\frac{3}{2}^{+}$	32	(1971DII8)
		2.78	$\frac{9}{2}^{+}$	5	(1971DII8)
		0.110	$\frac{1}{2}^{-}$	59	(1971DII8)
5.50 ^d	$\frac{3}{2}^{+}$	0.197	$\frac{5}{2}^{+}$	25 ⁿ	(1970AI01)
		1.35	$\frac{5}{2}^{-}$	49	(1970AI01)
		1.55	$\frac{3}{2}^{+}$	16	(1970AI01)
		0	$\frac{1}{2}^{+}$	11	(1970AI01)
5.54	$\frac{5}{2}^{+}$	0.197	$\frac{5}{2}^{+}$	7	(1970AI01)
		1.46	$\frac{3}{2}^{-}$	47	(1970AI01)
5.63		0.197	$\frac{5}{2}^{+}$	45	(1970AI01)
				33 ⁿ	(1970AI01)

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	Refs.
5.94	$\frac{1}{2}^{(+)}$	1.35	$\frac{5}{2}^-$	66	(1970AI01)
		0	$\frac{1}{2}^+$	15	(1970AI01)
		0.110	$\frac{1}{2}^-$	38	(1970AI01)
		1.46	$\frac{3}{2}^-$	38	(1970AI01)
		3.91	$\frac{3}{2}^{(+)}$	9	(1970AI01)
6.08	$(\frac{7}{2}^+)$	0.197	$\frac{5}{2}^+$	40 ⁿ	(1970AI01)
		1.35	$\frac{5}{2}^-$	20	(1970AI01)
		2.78	$\frac{9}{2}^+$	40	(1970AI01)
6.09	$(\frac{3}{2}^-)$	0	$\frac{1}{2}^+$	25 ⁿ	(1970AI01)
		0.110	$\frac{1}{2}^-$	62	(1970AI01)
		0.197	$\frac{5}{2}^+$	14	(1970AI01)
6.17	$\frac{7}{2}^{(-)}$	0.197	$\frac{5}{2}^+$	32 ⁿ	(1970AI01)
		1.35	$\frac{5}{2}^-$	68	(1970AI01)
6.29	$\frac{5}{2}^+$	1.35	$\frac{5}{2}^-$	31 ⁿ	(1970AI01)
		1.46	$\frac{3}{2}^-$	47	(1970AI01)
		1.55	$\frac{3}{2}^+$	22	(1970AI01)
6.33	$\frac{7}{2}^+$	0.197	$\frac{5}{2}^+$	84 ⁿ	(1970AI01)
		1.35	$\frac{5}{2}^-$	16	(1970AI01)
6.497 ^d	$\frac{3}{2}^{(+)}$	0	$\frac{1}{2}^+$	38	(1970AI01)
		0.110	$\frac{1}{2}^-$	14	(1970AI01)
		0.197	$\frac{5}{2}^+$	9	(1970AI01)
		1.35	$\frac{5}{2}^-$	14	(1970AI01)
		1.46	$\frac{3}{2}^-$	25	(1969AI01, 1970AI01)

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	Refs.
6.499 ^d	$\frac{11}{2}^+$	2.78	$\frac{9}{2}^+$	55	(1969AI01)
		4.00	$\frac{7}{2}^-$	< 3	(1969AI01)
		4.03	$\frac{9}{2}^-$	< 3	(1969AI01)
		4.39	$\frac{7}{2}^+$	< 3	(1969AI01)
		4.65	$\frac{13}{2}^+$	45	(1969AI01, 1970AI01)
		5.47	$\frac{7}{2}^+$	< 2	(1969AI01)
6.53	$\frac{3}{2}^+$	0	$\frac{1}{2}^+$	29	B
		0.110	$\frac{1}{2}^-$	59	B
		4.56	$\frac{5}{2}^+$	12	B
6.55	$\frac{7}{2}$	0.197	$\frac{5}{2}^+$	20	B
		1.35	$\frac{5}{2}^-$	55	B
		2.78	$\frac{9}{2}^+$	25	B
6.59	$\frac{9}{2}^+$	0.197	$\frac{5}{2}^+$	13	(1971DI18)
		2.78	$\frac{9}{2}^+$	63	(1971DI18)
		4.00	$\frac{7}{2}^-$	< 4	(1971DI18)
		4.03	$\frac{9}{2}^-$	< 2	(1971DI18)
		4.39	$\frac{7}{2}^+$	24	(1971DI18)
		4.56	$\frac{5}{2}^+$	< 2	(1971DI18)
		4.65	$\frac{13}{2}^+$	< 2	(1971DI18)
		5.43	$\frac{7}{2}^-$	< 3	(1971DI18)
		5.46	$\frac{7}{2}^+$	< 8	(1971DI18)
6.79	$\frac{3}{2}^-$	0	$\frac{1}{2}^+$	16	B
		0.110	$\frac{1}{2}^-$	40	B

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.		
6.93	$\frac{7}{2}^{(-)}$	0.197	$\frac{5}{2}^+$	13	$\Gamma_\gamma/\Gamma \lesssim 0.05$	B		
		1.35	$\frac{5}{2}^-$	5		B		
		1.46	$\frac{3}{2}^-$	26		B		
		0.197	$\frac{5}{2}^+$	73		B		
		1.35	$\frac{5}{2}^-$	5		B		
		2.78	$\frac{9}{2}^+$	2.4		B		
		4.00	$\frac{7}{2}^-$	1.3		B		
		4.03	$\frac{9}{2}^-$	1.3		B		
		7.17	$\frac{11}{2}^-, (\frac{7}{2}^-)$	4.00		$\frac{7}{2}^-$	6	B
				4.03		$\frac{9}{2}^-$	94	B
7.54 ^{d,g}	$\frac{5}{2}^+; T = \frac{3}{2}$	0	$\frac{1}{2}^+$	< 1		(1969AI02)		
		0.197	$\frac{5}{2}^+$	30		(1969AI02)		
7.66 ^{d,g}	$\frac{3}{2}^+; T = \frac{3}{2}$	1.55	$\frac{3}{2}^+$	42		A (1969AI02)		
		3.91	$\frac{3}{2}^{(+)}$	< 1		(1969AI02)		
		4.39	$\frac{7}{2}^+$	28		(1969AI02)		
		4.56	$\frac{3}{2}^{(+)}$	< 1		(1969AI02)		
		5.34	$\frac{1}{2}^-$	< 1		(1969AI02)		
		5.47	$\frac{7}{2}^+$	< 1		(1969AI02)		
		0	$\frac{1}{2}^+$	43		$\Gamma_\gamma = 4.7 \text{ eV}$ (1963BA19, 1969AI02)		
		0.197	$\frac{5}{2}^+$	17		$\Gamma_\gamma/\Gamma = 0.65 \pm 0.10$ A		
		1.55	$\frac{3}{2}^+$	40		(1969AI02)		

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	Refs.		
7.93	$(\frac{7^+}{2}, \frac{9}{2})$	3.91	$\frac{3}{2}^{(+)}$	< 3	(1969AI02)		
		4.39	$\frac{7}{2}^+$	< 7	(1969AI02)		
		4.56	$\frac{3}{2}^{(+)}$	< 5	(1969AI02)		
		5.34	$\frac{1}{2}$	< 10	(1969AI02)		
		5.47	$\frac{7}{2}^+$	< 10	(1969AI02)		
		0.197	$\frac{5}{2}^+$	4	(1971DI18)		
		2.78	$\frac{9}{2}^+$	96	(1971DI18)		
		7.94 ^d	$\frac{11}{2}^+$	2.78	$\frac{9}{2}^+$	11	(1970RO1C)
						10	(1971DI18)
				4.00	$\frac{7}{2}^-$	< 7	(1971DI18)
4.03	$\frac{9}{2}^-$			< 7	(1971DI18)		
4.39	$\frac{7}{2}^+$			< 7	(1971DI18)		
4.65	$\frac{13}{2}^+$			89	(1970RO1C)		
				90	(1971DI18)		
5.43	$\frac{7}{2}^-$			< 9	(1971DI18)		
5.46	$\frac{7}{2}^+$			< 10	(1971DI18)		
6.50	$\frac{11}{2}^+$			< 7	(1971DI18)		
6.59	$\frac{9}{2}^+$			< 7	(1971DI18)		
8.29	$(\frac{13}{2}^-)$			4.03	$\frac{9}{2}^-$	100	C
8.59	$\frac{3}{2}$	0	$\frac{1}{2}^+$	51 ± 4	(1962NE03)		
		0.110	$\frac{1}{2}^-$	9 ± 3	(1962NE03)		
		0.197	$\frac{5}{2}^+$	40 ± 3	(1965AL20)		
8.80 ^h	$\frac{1}{2}^+$	0	$\frac{1}{2}^+$	< 10	(1965AL20)		

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.
		0.110	$\frac{1}{2}^-$	42 ± 4		(1965AL20)
		0.197	$\frac{5}{2}^+$	< 5		(1965AL20)
		1.35	$\frac{5}{2}^-$	< 5		(1965AL20)
		1.46	$\frac{3}{2}^-$	21 ± 5		(1965AL20)
		1.55	$\frac{3}{2}^+$	19 ± 5		(1965AL20)
		2.78	$\frac{9}{2}^+$	< 1		(1965AL20)
		3.91	$\frac{3}{2}^{(+)}$	18 ± 2		(1965AL20)
E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	Γ_γ (meV)	Refs.
8.94	$(\frac{11}{2}^-)$	2.78	$\frac{9}{2}^+$	36 ± 10	54 ± 17	C
		4.00	$\frac{7}{2}^-$	19 ± 7	29 ± 12	C
		4.03	$\frac{9}{2}^-$	45 ± 13	68 ± 17	C
9.09 ⁱ	$\frac{7}{2}^+; T = \frac{3}{2}$	0.110	$\frac{1}{2}^-$	(< 0.5)		(1965AL20)
		0.197	$\frac{5}{2}^+$	11 ± 2		(1965AL20)
		1.35	$\frac{5}{2}^-$	4 ± 1		(1965AL20)
		2.78	$\frac{9}{2}^+$	64 ± 4^j		(1965AL20)
		4.00	$\frac{7}{2}^-$	8 ± 2		(1965AL20)
		5.43	$\frac{7}{2}^-$	(8 ± 2)		(1965AL20)
		6.08	$(\frac{7}{2}^+)$	(5 ± 2)		(1962NE03)
9.32	$\frac{1}{2}^+$	0	$\frac{1}{2}^+$	86 ± 4		(1962NE03)
		0.110	$\frac{1}{2}^-$	4 ± 2		(1962NE03)
		0.197	$\frac{5}{2}^+$	10 ± 2		(1962NE03)
9.67 ^{,h,k}	$\frac{3}{2}^+$	0	$\frac{1}{2}^+$	25		(1969WO1F)
		0.110	$\frac{1}{2}^-$	21		(1969WO1F)

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.
9.82 ^{h,l}	$\frac{5}{2}^-$	0.197	$\frac{5}{2}^+$	12		(1969WO1F)
		1.35	$\frac{5}{2}^-$	8		(1969WO1F)
		1.46	$\frac{3}{2}^-$	6		(1969WO1F)
		1.55	$\frac{3}{2}^+$	11		(1969WO1F)
		3.91	$\frac{3}{2}^{(+)}$	6		(1969WO1F)
		4.56	$\frac{5}{2}^+$	5		(1969WO1F)
		7.54	$\frac{5}{2}^+$	3		(1969WO1F)
		7.66	$\frac{3}{2}^+$	3		(1969WO1F)
		0.197	$\frac{5}{2}^+$	41		(1969WO1F)
		1.35	$\frac{5}{2}^-$	4		(1969WO1F)
		1.46	$\frac{3}{2}^-$	8		(1969WO1F)
		1.55	$\frac{3}{2}^+$	30		(1969WO1F)
		4.00	$\frac{7}{2}^-$	2		(1969WO1F)
		4.68	$\frac{5}{2}^-$	5		(1969WO1F)
		5.43	$\frac{7}{2}^-$	10		(1969WO1F)
E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)	$ M ^2 \times 10^{-3}$ W.u.	Refs.
10.136	$\frac{3}{2}^-$	0	$\frac{1}{2}^+$	84 ± 3		(1962NE03)
				64	4.9	A
		0.110	$\frac{1}{2}^-$	4 ± 2		(1962NE03)
					< 3	A
		0.197	$\frac{5}{2}^+$	12 ± 2		(1962NE03)
				29	2.4	A
		1.35	$\frac{5}{2}^-$		< 2	A

Table 19.9: Radiative transitions in $^{19}\text{F}^\dagger$ (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching ratio (%)		Refs.
		1.46	$\frac{3}{2}^-$	2	4.6	A
		1.55	$\frac{3}{2}^+$	2	0.20	A
		3.91	$\frac{3}{2}^{(+)}$	4	0.57	A

$^\dagger E_x = 1345.4 \pm 0.6, 1458.6 \pm 0.6, 1553.5 \pm 0.6, 2779.8 \pm 0.6, 3907.1 \pm 1.0, 3998.5 \pm 0.8, 4032.5 \pm 1.0, 4377.7 \pm 1.0, 4557.5 \pm 1.0$ and 4682.5 ± 1.0 keV (K. Bharuth-Ram, K.P. Jackson, N.A. Jelley, P.G. Lawson and K.W. Allen, private communication).

A: I.F. Wright and M.R. Wormald, private communication.

B: W.R. Dixon and R.S. Storey, private communication.

C: K. Bharuth-Ram, K.P. Jackson, N.A. Jelley, P.G. Lawson and K.W. Allen, private communication.

^a For partial widths, see (1969PO03).

^b See also (1968SP01, 1969WO1F).

^c See also (1966TH02, 1969WO1F).

^d See also Table 19.7.

^e See also (1967TO1C) and (1970AI01).

^f See, however, (1970SC25).

^g See also (1969WO1F).

^h See also (1962NE03).

ⁱ See also (1959BU05, 1963HU07).

^j $\Gamma_\gamma = 0.84 \pm 0.19$ eV. The total $\Gamma_\gamma(9.07) = 1.31 \pm 0.31$ eV.

^k $\Gamma_p \Gamma_\gamma / \Gamma = 1.3 \pm 0.3$ eV (1969WO1F).

^l $\Gamma_p \Gamma_\gamma / \Gamma = 1.7 \pm 0.3$ eV (1969WO1F).

^m The branching ratios to $^{19}\text{F}^*(0, 0.11, 1.55)$ are $< 5, < 1.5$ and $< 5\%$ respectively (1972RO01).

ⁿ Values shown are no true branching ratios but are relative intensities observed at 90° . I am indebted to W.R. Dixon and R.S. Storey for calling this to my attention.

Table 19.10: Lifetimes of some ^{19}F states

$^{19}\text{F}^*$ (MeV)	τ_m	Refs.
0.110 ^a	0.52 ± 0.13 nsec	(1960ST17)
	1.3 ± 0.2 nsec	(1962BO1A, 1962SE12)
	0.92 ± 0.04 nsec	(1963GA14)
	0.87 ± 0.14 nsec	(1963VA01, 1963VA12)
	0.84 ± 0.04 nsec	(1965BO28)
	0.91 ± 0.05 nsec	(1969ME13)
	0.847 ± 0.010 nsec	(1969NI09)
0.197 ^a	0.853 ± 0.010 nsec	mean of last five values
	100 ± 20 nsec	(1954JO21)
	125 ± 25 nsec	(1956JO35)
	114 ± 25 nsec	(1960ST17)
	129.9 ± 2.3 nsec	(1967BE14)
	128 ± 2 nsec	(1968KL05)
1.35 ^b	128.8 ± 1.5 nsec	mean of last two values
	5.2 ± 0.9 psec	(1968PA11)
	4.7 ± 0.6 psec	(1969JO10)
	$5.3^{+1.5}_{-0.9}$ psec	(1969PO03)
	$3.5^{+1.5}_{-1.1}$ psec	A
1.46 ^c	4.8 ± 0.4 psec	mean
	62 ± 20 fsec	(1964BO22)
	102 ± 35 fsec	(1968TOZU)
	57 ± 24 fsec	(1969PO03)
	100 ± 20 fsec	A
1.55 ^d	78 ± 12 fsec	mean value
	≤ 30 fsec	(1968TOZU)
	< 17 fsec	(1969PO03)
2.78	$4.4^{+2.4}_{-2.0}$ fsec	“best” value: see (1969PO03)
	< 190 fsec	(1968RO07)
	246 ± 44 fsec	(1968TO11, 1968TOZU)
	300 ± 60 fsec	(1969JA09)
	270 ± 40 fsec	A

Table 19.10: Lifetimes of some ^{19}F states (continued)

$^{19}\text{F}^*$ (MeV)	τ_m	Refs.
	230 ± 60 fsec	see (1969JA09)
	261 ± 24 fsec	mean of the last four values
3.91	≤ 20 fsec	A
4.00	23 ± 10 fsec	A
4.03	60 ± 15 fsec	A
	87 ± 16 fsec	B
4.555, 4.558	≤ 20 fsec	A
4.65	2.3 ± 0.5 psec	(1969BH01)
	2.1 ± 0.4 psec	(1969JA09)
	2.2 ± 0.3 psec	mean
4.68	15.4 ± 3.0 fsec	(1972RO01)
5.34	≤ 15 fsec	(1969PO03)
5.46	≤ 19 fsec	(1969PO03)
5.50	≤ 43 fsec	(1969PO03)

A: Bharuth-Ram, Jackson, Jelley, Lawson and Allen, private communication.

B: W.R. Dixon and R.S. Storey, private communication.

^a See also (1964LI1B).

^b See also (1968TOZU, 1971NI05).

^c See also (1971NI05).

^d See also (1967WA13, 1969NI09).

10. (a) $^{16}\text{O}(t, n)^{18}\text{F}$ $Q_m = 1.269$ $E_b = 11.700$
 (b) $^{16}\text{O}(t, p)^{18}\text{O}$ $Q_m = 3.707$
 (c) $^{16}\text{O}(t, t)^{16}\text{O}$
 (d) $^{16}\text{O}(t, \alpha)^{15}\text{N}$ $Q_m = 7.686$

The excitation function for neutrons (reaction (a)) has been measured for $E_t = 0.3$ to 2.2 MeV (1961LO14) and 0.7 to 2.1 MeV by (1955JA1A). See also (1967CH35). Resonances in the yields of p_0 , p_1 , α_0 and α_{1+2} (reactions (b) and (d)) are reported by (1967KO11), corresponding to states with $E_x = 12.42, 12.67, (12.75), 12.83, 12.91, 12.97, (13.06)$ and (13.14) MeV ($\pm \approx 25$ keV)

[not all resonances observed in every channel]. Additional states from the α_0 yield are reported by (1965GE05). See also (1959JO32, 1967KO1G). Two pronounced peaks are observed in the elastic scattering (reaction (c)) at $E_t = 2.6$ and 3.2 MeV [$^{19}\text{F}^*(13.9, 14.4)$] (1968ET1A: $E_t = 1.4$ to 3.5 MeV). The elastic scattering has also been studied for $E_t = 9$ to 13 MeV (1965GL04).

$$11. \text{}^{16}\text{O}(\alpha, \text{p})^{19}\text{F} \quad Q_m = -8.115$$

At $E_\alpha = 26.7$ and 33.1 MeV, angular distributions have been measured for the $\text{p}_{0 \rightarrow 2}$, $\text{p}_{3 \rightarrow 5}$ and p_6 groups (1961YA02).

$$12. \text{}^{16}\text{O}({}^6\text{Li}, {}^3\text{He})^{19}\text{F} \quad Q_m = -4.094$$

This reaction (and its mirror reaction $^{16}\text{O}({}^6\text{Li}, \text{t})^{19}\text{Ne}$: see ^{19}Ne) have been studied at $E({}^6\text{Li}) = 24$ MeV. The states most strongly excited by both reactions correspond to members of the $K^\pi = \frac{1}{2}^+$ (g.s.) rotational band, with the exception of the $\frac{7}{2}^+$ state. It is suggested that it is the second $\frac{7}{2}^+$ state (at $E_x = 5.46$ MeV) which is a member of the g.s. rotational band although it is more weakly populated in both reactions than predicted [see, however, Table 19.11, footnote ^d]. Up to $J^\pi = \frac{9}{2}^+$, cross sections are observed to increase with increasing spin. Members of a $K^\pi = \frac{1}{2}^-$ rotational band based primarily on a configuration of four s-d shell particles and a p-shell hole are also populated in these reactions: again cross sections increase with increasing spin. Table 19.11 presents the data on the $^{16}\text{O}({}^6\text{Li}, {}^3\text{He})^{19}\text{F}$ and $^{16}\text{O}({}^6\text{Li}, \text{t})^{19}\text{Ne}$ reactions (1971BI06). See also (1971GA1H).

$$13. \text{}^{16}\text{O}({}^7\text{Li}, \alpha)^{19}\text{F} \quad Q_m = 9.233$$

At $E({}^7\text{Li}) = 20$ MeV, α -particles to the ^{19}F states with $E_x < 4.5$ MeV have been observed (1969BI1A). At $E({}^7\text{Li}) = 30.3$ MeV, the excitation of $^{19}\text{F}^*(7.25, 10.2, 12.1, 13.4)$, in addition to lower states, is reported by (1969GL06). See also (1969RO1G; theor.).

$$\begin{array}{lll} 14. \text{(a) } {}^{17}\text{O}(\text{d}, \text{n})^{18}\text{F} & Q_m = 3.384 & E_b = 13.815 \\ \text{(b) } {}^{17}\text{O}(\text{d}, \text{p})^{18}\text{O} & Q_m = 5.822 & \\ \text{(c) } {}^{17}\text{O}(\text{d}, \alpha)^{15}\text{N} & Q_m = 9.802 & \end{array}$$

For reaction (a) see ^{18}F and (1950HO1A). For reaction (b) see ^{18}O ; for reaction (c) see ^{15}N in (1970AJ04).

15. $^{17}\text{O}(t, n)^{19}\text{F}$

$$Q_m = 7.557$$

Table 19.11: Levels of ^{19}F and ^{19}Ne from $^{16}\text{O}(^6\text{Li}, ^3\text{He})$ and $^{16}\text{O}(^6\text{Li}, t)$ (1971BI06)

J^π	E_x^a in ^{19}F (MeV)			$\sigma_{\max}(\theta)$ (mb/sr)	E_x^a in ^{19}Ne (MeV)			$\sigma_{\max}(\theta)$ (mb/sr)
	$K^\pi = \frac{1}{2}^+$	$K^\pi = \frac{1}{2}^-$	other		$K^\pi = \frac{1}{2}^+$	$K^\pi = \frac{1}{2}^-$	other	
$\frac{1}{2}^+$	0.0			0.106	0.0			0.095
$\frac{5}{2}^+$	0.197			0.682	0.238			0.612
$\frac{3}{2}^+$	1.55			0.279	1.54 ^b			0.294
$\frac{9}{2}^+$	2.78			1.16	2.79 ^b			0.763
$\frac{13}{2}^+$	4.65			0.197	4.62 ^b			0.125
$\frac{7}{2}^+$	5.46 ^d			0.347	5.43 ^b			0.169
$\frac{1}{2}^-$		0.110		0.026		0.275		0.026
$\frac{5}{2}^-$		1.35		0.173		1.51 ^b		0.082
$\frac{3}{2}^-$		1.46		0.065		1.62 ^b		0.052
$\frac{7}{2}^-$		4.00		0.357		4.14 ^b		0.072
$\frac{9}{2}^-$		4.04					4.20 ^b	
$\frac{3}{2}^+$			3.91 ^b	0.012			4.03 ^b	0.008
$\frac{7}{2}^+$			4.39	0.037			4.37 ^b	0.016
$\frac{5}{2}^+$ (+)			4.55	0.059				
$\frac{3}{2}^-$, ($\frac{1}{2}^-$)			4.56				4.55	0.006
$\frac{5}{2}^-$			4.68	c				
$\frac{5}{2}^-$ (-)			5.11	0.012				
$\frac{5}{2}^+$			5.34	0.005				
$\frac{7}{2}^-$			5.43	0.215				

^a Energies are nominal.

^b J^π assignment based on similarity in σ_{\max} in both reactions and on known spin of analog state.

^c Transition masked by strong group to $E_x = 4.65$ MeV.

^d See, however, (1971DI18).

Not reported.

16. $^{17}\text{O}(^3\text{He}, p)^{19}\text{F}$

$$Q_m = 8.321$$

Not reported.

Table 19.12: Resonances in $^{18}\text{O}(p, \gamma)^{19}\text{F}$ ^a

E_p (keV)	Γ_{lab} (keV)	J^π	E_x (keV)	Refs.
629.6 ± 0.3	2.0 ± 0.3	$\frac{3}{2}^+$	8.5892	(1959BU05, 1962NE03, 1963HU07, 1969DU1A)
848 ± 2	40 ± 5	$\frac{3}{2}$	8.796	(1959BU05, 1962NE03, 1963HU07, 1965AL20)
1166.5 ± 0.4	$(25 \pm 24) \times 10^{-3}$	$\frac{7}{2}^+$ ^b	9.0976	(1959BU05, 1963HU07, 1969DU1A)
1398 ± 3	4		9.317	(1959BU05, 1962NE03, 1963HU07)
1685 ± 5 ^d	< 15		(9.589)	(1959BU05)
1769 ± 2	4.0 ± 1.0	$\frac{3}{2}^+$	9.668	(1959BU05, 1962NE03, 1969WO1F)
1778			(9.677)	(1962NE03)
1790			(9.688)	(1962NE03)
1928.4 ± 0.6 ^c	0.3 ± 0.05	$\frac{5}{2}$ ^f	9.819	(1959BU05, 1962NE03, 1969DU1A, 1969WO1F)
2263.0 ± 0.7	5.0 ± 1.0	$\frac{3}{2}^-$	10.136	A, (1962NE03, 1969DU1A)
2.36 ^e			(10.23)	(1962NE03)
2.39	47 ± 10		(10.26)	(1962NE03)
2.41	10 ± 5		(10.28)	(1962NE03)
2.44			(10.30)	(1962NE03)
(2.60)			(10.45)	(1962NE03)
(2.66)			(10.51)	(1962NE03)
(2.68)			(10.53)	(1962NE03)
(2.73)			(10.58)	(1962NE03)
2.77			(10.62)	(1962NE03)
(2.80)			(10.64)	(1962NE03)
(2.84)			(10.68)	(1962NE03)

A: M.R. Wormald and I.F. Wright, private communication.

^a See also Table 19.9.

^b Most probable value, although $J^\pi = \frac{9}{2}^+$ is also possible: see text. $T = \frac{3}{2}$ (1965AL20).

^c Γ_γ and Γ_p are \lesssim few eV (1969DU1A).

^d See, however, (1962NE03).

^e See (1962NE03) for additional resonant structure between $E_p = 2.33$ and 2.78 MeV.

^f From γ -ray angular distributions (I.F. Wright, private communication).

17. $^{17}\text{O}(\alpha, d)^{19}\text{F}$ $Q_m = -10.032$

See (1963HA1C).

18. $^{18}\text{O}(p, \gamma)^{19}\text{F}$ $Q_m = 7.993$

Resonances for capture radiation observed for $E_p = 0.3$ to 3.0 MeV are displayed in Table 19.12 (1959BU05, 1962NE03, 1963HU07, 1965AL20, 1969DU1A, 1969WO1F). At the $E_p = 0.85$ MeV resonance [$^{19}\text{F}^*(8.80)$], the intensity of the transition $8.80 \rightarrow 3.91$ and the anisotropy of the $8.80 \rightarrow 3.91 \rightarrow 0$ γ -rays limit J of $^{19}\text{F}^*(3.91)$ to $\frac{3}{2}$ or $\frac{5}{2}$. $J = \frac{5}{2}$ is ruled out by the angular distribution of the γ -rays. At the $E_p = 1.17$ MeV resonance [$^{19}\text{F}^*(9.10)$], the angular distribution of the γ -rays to the $\frac{5}{2}^+$ state at 0.197 MeV indicate $J^\pi = \frac{7}{2}^+$ [$\frac{9}{2}^+$ not completely excluded] for $^{19}\text{F}^*(9.10)$. $J = \frac{5}{2}$ or $\frac{9}{2}$ is suggested for $^{19}\text{F}^*(2.78)$. The γ -decay of the resonances at $E_p = 1.77$ and 1.93 MeV is very complex (1965AL20). For branching ratios and Γ_γ , see Table 19.9 (1962NE03, 1965AL20, 1969WO1F). See also (1969SK1A) and (1959LA1A).

19. $^{18}\text{O}(p, n)^{18}\text{F}$ $Q_m = -2.438$ $E_b = 7.993$

Recent yield measurements are reported for 2.5 to 3.0 MeV (1967PR04), 2.6 to 3.0 MeV (1960BL05; σ_t), 2.6 to 3.3 MeV (1969BE57: rel. total yield, high resolution), 3.0 to 7.0 MeV (1969DI07; $n_1\gamma$, $n_2\gamma$, $n_3\gamma$ and n), 3.5 to 10 MeV (1964BA16; σ_t), 3.6 to 4.0 MeV (1968BE34; $n_1\gamma$, $n_2\gamma$, $n_3\gamma$) and 6.9 to 13.5 MeV (1965BL07, 1969AN06; σ_t for n_0 , n_2 (unresolved), n_5 , n_6 , n_7 , n_8 (unresolved); not all groups over the entire range). See also (1959BR06). Observed resonances are displayed in Table 19.13 (1956MA18, 1964BA16, 1967PR04, 1968BE34, 1969BE57, 1969DI07).

(1969AN06) find that the rather large cross section for the reaction to $^{18}\text{F}(0)$ [$\Delta T = 1$, $\Delta J = 1$] over the range $E_p = 6.9$ to 13.5 MeV and the cross section for the reaction to $^{18}\text{F}^*(3.06)$ [$\Delta T = 0$, $\Delta J = 2$] indicate the operation of a spin-flip mechanism in the effective two-body force, and a sizable value for the quadrupole term in the multipole expansion of that force.

For a summary of threshold measurements and angular distribution studies, see ^{18}F .

20. $^{18}\text{O}(p, p)^{18}\text{O}$ $E_b = 7.993$

Scattering studies have been carried out at $E_p = 0.60$ to 1.45 MeV (1962YA03, 1962YA1B; p_0), 0.79 to 3.55 MeV (1961CA02; p_0 (and $p_1\gamma$ at higher energies)), 1.39 to 3.20 MeV (1969SE02, 1969SE03; p_0 , high resolution), 1.7 to 3.5 MeV (1962SO01, 1964ST1E; p_0), 2.5 to 3.0 MeV (1967PR04; $p_1\gamma$), 3.0 to 7.0 MeV (1969DI07; $p_1\gamma$), 3.2 to 5.4 MeV (1968BE34, 1969BE13; $p_1\gamma$), and 7.9 to 16.3 MeV (1964ST1D, 1966ST04; p_0 , p_1). For a spallation study, see (1969EP1B).

Table 19.13: Resonances in $^{18}\text{O}(p, n)^{18}\text{F}$ ^a

E_p (MeV \pm keV)	Γ_{lab} (keV)	Res. ^b in yield of	J^π	E_x in ^{19}F (MeV)	Refs.
2.643 \pm 1.0	6.5 \pm 0.5	n	$(\frac{3}{2})$	10.496	(1967PR04, 1969BE57)
2.691 \pm 1.0	2.6 \pm 0.2	n		10.541	(1967PR04, 1969BE57)
2.717 \pm 1.0	5.5 \pm 0.5	n		10.566	(1967PR04, 1969BE57)
2.767 \pm 1.5	5.0 \pm 0.5	n	$\frac{5}{2}^{(+)}$	10.613	(1967PR04, 1969BE57)
3.025 \pm 2.0	26.0 \pm 1.5	n	$\frac{3}{2}$	10.857	(1956MA18, 1969BE57)
3.145 \pm 3.0	\approx 12	n		10.971	(1969BE57)
3.163 \pm 3.0	\approx 11	n		10.988	(1969BE57)
3.248 \pm 3.0 ^c	33.0 \pm 5.0	n	$\frac{3}{2}$	11.069	(1956MA18, 1969BE57)
3.386 \pm 2	45 \pm 2	n	$(\frac{1}{2})$	11.199	(1956MA18)
3.495 \pm 4		n		11.302	(1956MA18)
3.63 \pm 10 ^e	85 \pm 20	n, n ₁		11.43	(1956MA18, 1964BA16, 1969DI07)
3.71		n, n ₁		(11.51)	(1964BA16, 1969DI07)
3.77 \pm 10	< 50	n, n ₂		11.56 ^d	(1964MA12, 1968BE34, 1969DI07)
3.83		n ₁		(11.62)	(1964BA16, 1969DI07)
3.86 \pm 10	40	n, n ₁		11.65	(1964BA16, 1969DI07)
4.00		n ₁ , n ₃		(11.78)	(1969DI07)
4.06 \pm 10	< 50	n, n ₁		11.84	(1964BA16, 1969DI07)
4.11		n ₁		(11.88)	(1969DI07)
4.16 \pm 10	90	n, n ₁		11.93	(1964BA16, 1969DI07)
4.33		n ₁ , n ₃		(12.09)	(1969DI07)
4.37 \pm 10	100	n, n ₁ , n ₂		12.13	(1964BA16, 1969DI07)
4.47	50	n, n ₁ , n ₂ , n ₃		12.23	(1964BA16, 1969DI07)
4.58 \pm 10		n ₁		(12.33)	(1969DI07)
4.70		n ₃		(12.44)	(1969DI07)
4.83		n ₁ , n ₂ , n ₃		(12.57)	(1969DI07)
4.90		n ₂		(12.63)	(1969DI07)
5.05 \pm 10	200	n, n ₁		12.77	(1964BA16, 1969DI07)
5.10		n ₁ , n ₂		(12.82)	(1969DI07)
5.20		n ₂ , n ₃		(12.92)	(1969DI07)
5.35		n, n ₁ , n ₂ , n ₃		13.06	(1964BA16, 1969DI07)

Table 19.13: Resonances in $^{18}\text{O}(\text{p}, \text{n})^{18}\text{F}$ ^a (continued)

E_p (MeV \pm keV)	Γ_{lab} (keV)	Res. ^b in yield of	J^π	E_x in ^{19}F (MeV)	Refs.
5.47 ± 15	70	n, n ₁		13.17	(1964BA16, 1969DI07)
5.62 ± 15	20	n, n ₁		13.31	(1964BA16, 1969DI07)
5.63		n ₁		(13.32)	(1969DI07)
5.76		n ₁ , n ₃		(13.45)	(1969DI07)
6.06 ± 15	100	n, n ₁ , n ₂		13.73	(1964BA16, 1969DI07)
6.60 ± 15	350	n		14.24	(1964BA16)
(6.70 ± 15)		n		(14.34)	(1964BA16)
7.17 ± 20	300	n		14.78	(1964BA16)
7.40 ± 20		n		15.00	(1964BA16)
(7.8)		n		(15.4)	(1964BA16)
(7.98)		n		(15.55)	(1964BA16)
8.19 ± 25	160	n		15.75	(1964BA16)
8.74 ± 25	200	n		16.27	(1964BA16)
9.30 ± 30		n		16.80	(1964BA16)

^a See also (1959AJ76).

^b n means in the total yield.

^c See also (1956MA18).

^d $T = \frac{3}{2}$: see (1968BE34).

^e Errors here and below are estimated from published data of (1964BA16) by H.B. Willard, private communication.

Observed resonances are shown in Table 19.14 (1961CA02, 1962SO01, 1962YA03, 1962YA1B, 1964ST1D, 1964ST1E, 1966ST04, 1967PR04, 1969DI07, 1969SE02, 1969SE03). All the positive-parity states with large proton width observed experimentally up to $E_p = 3.2$ MeV can be associated with a doorway state. Both the experimental data and shell-model calculations indicate that the $\frac{5}{2}^+$ resonance at $E_p = 2.768$ MeV [$^{19}\text{F}^*(10.613)$] may be identified as the analog of $^{19}\text{O}^*(3.16)$ although isospin impurities are present in $^{19}\text{F}^*(10.613)$ (1969SE02, 1969SE03).

21. $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$

$$Q_m = 3.980$$

$$E_b = 7.993$$

Yield measurements are reported for $E_p = 0.50$ to 0.75 MeV (1971SK1C; α_0), 0.73 to 1.05 MeV (1963KA25; α_0), 0.79 to 3.55 MeV (1961CA02; α_0 (and $\alpha_{1+2}\gamma$ at the higher energies)), 0.9 to 2.1 MeV (1963AM1A, 1964AM1A; α_0), 1 to 3 MeV (1960CL02; α_0), 1.1 to 2.6 MeV

Table 19.14: Energy levels of ^{19}F from $^{18}\text{O}(p, p)^{18}\text{O}$ and $^{18}\text{O}(p, \alpha)^{15}\text{N}$

E_p^a (MeV \pm keV)	Γ_{lab} (keV)	Particles out	Γ_p (keV)	Γ_α (keV)	J^π	E_x (MeV)	Refs.
0.6326 ± 0.4	2.1 ± 0.1	p_0	0.065 ± 0.006	2.0 ± 0.2	$\frac{3}{2}^-$	8.5921	(1962YA03, 1962YA1B)
0.680	100	p_0	5	95	$\frac{1}{2}^+$	8.637	(1962YA03, 1962YA1B)
0.846 ± 1.5	47 ± 1	p_0, α_0	26 ± 1.5	21 ± 1	$\frac{1}{2}^+; T = \frac{3}{2}$	8.794	(1960CL02, 1961CA02, 1962YA03, 1962YA1B, 1963KA25, 1968YA03)
0.9870 ± 0.7	3.8 ± 0.2	p_0, α_0	0.080 ± 0.007	3.7 ± 0.3	$\frac{3}{2}^-$	8.928	(1960CL02, 1961CA02, 1962YA03, 1962YA1B, 1963AM1A, 1963KA25, 1964AM1A)
(1.135)	140					(9.068)	(1963AM1A, 1964AM1A)
1.1685 ± 0.5	0.60 ± 0.03	p_0, α_0	0.005 ± 0.0006	0.595 ± 0.08	$\frac{7}{2}^+$	9.0995	(1960CL02, 1961AM01, 1961CA02, 1962YA03, 1962YA1B, 1963AM1A, 1964AM1A)
1.2390 ± 1	6.1 ± 0.3	$p_0, (\alpha_0)$	0.40 ± 0.03	5.7 ± 0.4	$\frac{1}{2}^+$	9.166	(1962YA03, 1962YA1B, 1963AM1A, 1964AM1A)
1.4025 ± 1	5.2 ± 0.2	p_0, α_0	0.23 ± 0.02	5.0 ± 0.4	$\frac{1}{2}^+$	9.321	(1960CL02, 1961CA02, 1962YA03, 1962YA1B, 1963AM1A, 1964AM1A, 1964KA1B)
1.620 ± 6	30	p_0, α_0			$(\frac{5}{2})$	9.527	(1960CL02, 1961CA02, 1963AM1A, 1964AM1A, 1964KA1B)
1.668 ± 6	27	p_0, α_0			$\frac{3}{2}$	9.572	(1960CL02, 1961CA02, 1963AM1A, 1964AM1A, 1964KA1B)
1.766 ± 3	3.6	p_0, α_0	2.1	1.5	$\frac{3}{2}^+$	9.665	(1960CL02, 1961CA02, 1962SO01, 1963AM1A, 1964AM1A, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
1.928 ± 3	0.16	p_0, α_0	0.09	0.07	$(\frac{5}{2}, \frac{7}{2})^-$	9.819	(1960CL02, 1961CA02, 1962SO01, 1963AM1A, 1964AM1A, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
2.001 ± 4	31	p_0, α_0	12	19	$\frac{1}{2}^+$	9.888	(1960CL02, 1961CA02, 1962SO01,

Table 19.14: Energy levels of ^{19}F from $^{18}\text{O}(p, p)^{18}\text{O}$ and $^{18}\text{O}(p, \alpha)^{15}\text{N}$ (continued)

E_p^a (MeV \pm keV)	Γ_{lab} (keV)	Particles out	Γ_p (keV)	Γ_α (keV)	J^π	E_x (MeV)	Refs.
2.2630 ± 0.7	5.0 ± 1.0	$\alpha_0, \alpha_1, \alpha_2$	≈ 5	0.004^d	$\frac{3}{2}^-$	10.136	1963AM1A, 1964AM1A, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
2.289 ± 3	33	p_0, α_0	2.3	(1.0)	$\frac{1}{2}^+$	10.161	A (1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
2.363 ± 3	4.5	p_0, α_0	2.8	1.7	$\frac{1}{2}^+$	10.231	(1960CL02, 1962SO01, 1963GO08, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
2.387 ± 3	24	p_0, α_0	11	13	$\frac{3}{2}^+$	10.253	(1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
2.443 ± 4	9.7	p_0, α_0	5.2	4.5	$\frac{3}{2}^+$	10.306	(1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964KA1B, 1964ST1E, 1969SE02, 1969SE03)
2.644 ± 3	4.6	$p_0, p_1, \alpha_0, \alpha_{1+2}$	2.4	(1.0)	$\frac{3}{2}^+$	10.497	(1960BL05, 1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1967PR04, 1969SE02, 1969SE03)
2.705 ± 3	8 ± 2	p_1, α_0			$\frac{3}{2}(^+); (T = \frac{3}{2})$	10.554	(1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1967PR04)
2.732 ± 4	23 ± 3	p_1, α_0			$(\frac{5}{2}^+)$	10.580	(1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1967PR04)
2.768 ± 3	4.0	$p_0, p_1, \alpha_0, \alpha_{1+2}$	0.7	(1.0)	$\frac{5}{2}^+; T = \frac{3}{2}^b$	10.614	(1960BL05, 1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1967PR04, 1969SE02, 1969SE03)
2.925 ± 3	5.7	$p_0, p_1, \alpha_0, \alpha_{1+2}$	4.5	1.2	$\frac{1}{2}^-$	10.763	(1960BL05, 1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1967PR04, 1969SE02, 1969SE03)

Table 19.14: Energy levels of ^{19}F from $^{18}\text{O}(p, p)^{18}\text{O}$ and $^{18}\text{O}(p, \alpha)^{15}\text{N}$ (continued)

E_p^a (MeV \pm keV)	Γ_{lab} (keV)	Particles out	Γ_p (keV)	Γ_α (keV)	J^π	E_x (MeV)	Refs.
3.029 ± 4	19.5	$p_0, p_1, \alpha_0, \alpha_{1+2}$	13.0		$\frac{5}{2}^+$	10.861	(1960CL02, 1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1969DI07, 1969SE02, 1969SE03)
(3.06)		α_0				(10.89)	(1963GO08)
3.148 ± 4	(14)	$p_0, p_1, \alpha_0, \alpha_{1+2}$	(4.5)	(4.5)	$(\frac{3}{2}, \frac{5}{2})^+$	10.974	(1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1969DI07, 1969SE02, 1969SE03)
3.266 ± 9	35	$p_0, p_1, \alpha_0, \alpha_{1+2}$			$\frac{1}{2}^+$	11.086	(1961CA02, 1962SO01, 1963GO08, 1964ST1E, 1969DI07)
3.386 ± 9	20	$p_0, p_1, \alpha_0, \alpha_{1+2}$			$(\frac{1}{2}^-)$	11.199	(1961CA02, 1962SO01, 1964ST1E, 1969DI07)
3.480 ± 9	25	$p_0, p_1, \alpha_0, \alpha_{1+2}$				11.288	(1961CA02, 1963GO08, 1969DI07)
3.503 ± 9		α_0, α_{1+2}				11.310	(1961CA02, 1963GO08, 1969DI07)
3.67		α_{1+2}				11.47	(1963GO08, 1969DI07)
3.71		p_1, α_0				11.51	(1963GO08, 1969DI07)
3.75		p_1, α_0				11.54	(1963GO08, 1969DI07)
(3.83)		α_{1+2}				(11.62)	(1969DI07)
3.86	≈ 40	p_1, α_{1+2}				11.65	(1963GO08, 1969DI07)
3.89		$p_1, \alpha_0, \alpha_{1+2}$				11.68	(1963GO08, 1969DI07)
4.00		p_0, α_0				11.78	(1963GO08, 1969DI07)
4.06		$p_1, \alpha_0, \alpha_{1+2}$				11.84	(1963GO08, 1969DI07)
4.11		p_1, α_0				11.88	(1963GO08, 1969DI07)
4.14		p_1, α_0				11.91	(1963GO08, 1969DI07)
4.19		p_1, α_{1+2}				11.96	(1969DI07)
4.28		α_0, α_{1+2}				12.05	(1963GO08, 1969DI07)
4.33		$p_1, \alpha_0, \alpha_{1+2}$				12.09	(1963GO08, 1969DI07)
4.38		$p_1, \alpha_0, \alpha_{1+2}$				12.14	(1963GO08, 1969DI07)
4.48	≈ 80	$p_1, \alpha_0, \alpha_{1+2}$				12.23	(1963GO08, 1969DI07)

Table 19.14: Energy levels of ^{19}F from $^{18}\text{O}(p, p)^{18}\text{O}$ and $^{18}\text{O}(p, \alpha)^{15}\text{N}$ (continued)

E_p^a (MeV \pm keV)	Γ_{lab} (keV)	Particles out	Γ_p (keV)	Γ_α (keV)	J^π	E_x (MeV)	Refs.	
4.58	15	p_1, α_0			c	12.33	(1963GO08, 1969DI07)	
4.70		p_1, α_0				12.44	(1963GO08, 1969DI07)	
4.78		p_1				12.52	(1969DI07)	
4.83		p_1, α_{1+2}				12.57	(1969DI07)	
4.90		p_1, α_0				12.63	(1963GO08, 1969DI07)	
4.96		$p_1, \alpha_0, \alpha_{1+2}$				12.69	(1963GO08, 1969DI07)	
5.02		α_{1+2}				12.75	(1969DI07)	
5.06		p_1, α_0				12.78	(1963GO08, 1969DI07)	
5.20		p_1, α_{1+2}				12.92	(1969DI07)	
5.35		p_1, α_{1+2}				13.06	(1969DI07)	
5.47		α_{1+2}						
5.64		30	$p_1, \alpha_0, \alpha_{1+2}$				13.17	(1969DI07)
5.76			α_{1+2}				13.33	(1963GO08, 1969DI07)
5.90			p_1, α_{1+2}				13.45	(1969DI07)
6.08			p_1, α_{1+2}				13.58	(1969DI07)
6.65	p_1, α_{1+2}				13.75	(1969DI07)		
8.85	α_0				14.29	(1969DI07)		
10.5	p_0, p_1, α_0				16.37	(1964ST1D)		
					17.9	(1964ST1D, 1966ST04)		

43

A: M.R. Wormald and I.F. Wright, private communication.

^a Additional resonances are reported by (1956HI1A, 1960CL02, 1961CA02, 1962SO01, 1963AM1A, 1963GO08, 1964AM1A, 1964KA1B, 1964ST1E). See also (1959AJ76).

^b $T = \frac{3}{2}$ (1969SE02, 1969SE03).

^c $T = \frac{3}{2}$: see (1969DI07). See, however, (1963GO08).

^d $\alpha_1 + \alpha_2$ only.

Table 19.15: Thresholds in the $^{18}\text{O}(\text{d}, \text{n})^{19}\text{F}$ reaction

E_d (keV)		E_x
(1956HA1A) ^a	(1958BU12, 1959BU81) ^b	(MeV)
	346 ± 8	6.079
	(370)	(6.101)
497 ± 15	525 ± 8	6.240
	584 ± 10	6.293
	1850 ± 50	7.43
	2150 ± 50	7.70
	2640 ± 50	8.14
3050 ± 20		8.51
	3160 ± 30	8.61

^a Slow neutron thresholds. See also (1965MA1K).

^b Thresholds for γ -rays.

(1964KA1B; α_0), 1.39 to 3.20 MeV (1969SE02, 1969SE03; α_0 , high resolution), 2.2 to 6 MeV (1963GO08; α_0), 2.5 to 3.0 MeV (1967PR04; $\alpha_{1+2}\gamma$), 3.0 to 7.0 MeV (1969DI07; $\alpha_{1+2}\gamma$), 3.2 to 5.5 MeV (1969BE13; $\alpha_{1+2}\gamma$, $\alpha_3\gamma$), and 7.9 to 14 MeV (1964ST1D; α_0). See also (1960BL05, 1961AM01, 1964EC03, 1964SC01). Observed resonances are displayed in Table 19.14 (1960CL02, 1961AM01, 1961CA02, 1963AM1A, 1963GO08, 1964AM1A, 1964KA1B, 1964ST1D, 1967PR04, 1969DI07, 1969SE02, 1969SE03). See also ^{15}N in (1970AJ04).

$$22. \text{ (a) } ^{18}\text{O}(\text{d}, \text{n})^{19}\text{F} \quad Q_m = 5.768$$

$$\text{ (b) } ^{18}\text{O}(\text{d}, \text{n}\alpha)^{15}\text{N} \quad Q_m = 1.755$$

Thresholds for slow neutron production and for γ -rays are shown in Table 19.15 (1956HA1A, 1958BU12, 1959BU81). Angular distributions have been measured at $E_d = 3$ MeV (and analyzed by DWBA) to many of the states with $E_x \leq 7.4$ MeV: the closely spaced states were not resolved; however, $l_p = 4$ for the neutrons corresponding to $^{19}\text{F}^*(2.78)$, and therefore $J^\pi = \frac{9}{2}^+, \frac{7}{2}^+$. In addition, neutron groups corresponding to ^{19}F states at 5.11 [$\frac{7}{2}^-, \frac{5}{2}^-$], 5.49 [u], 5.62, 5.94, 6.09 [u], 6.26 [u] [$\frac{1}{2}^+$], 6.53 [u] [$\frac{5}{2}^+, \frac{3}{2}^+, \frac{1}{2}^+$], 6.80 [u] [$\frac{3}{2}^-, \frac{1}{2}^-$], 6.93 [u], 7.26 and 7.40 MeV (± 0.03 MeV) are reported [(u = unresolved states)] [J^π of dominant state(s)] (1968GU07). See also (1963BE05, 1964MO23, 1969HE1P). The lifetime of $^{19}\text{F}^*(1.55)$ is < 1 psec (1969NI09); see, however, Table 19.10.

Table 19.16: Energy levels of ^{19}F from $^{18}\text{O}(^3\text{He}, \text{d})^{19}\text{F}$

E_x (MeV \pm keV)		l^a	$C^2S(2J_f + 1)^{a,b}$	J^π^a
(1970GR04)	(1970SC25)			
0	0	0	0.42	$\frac{1}{2}^+$
0.110 ± 10	0.112 ± 3	1	0.224	$\frac{1}{2}^-$
0.198 ± 10	0.199 ± 3	2	2.45	$\frac{5}{2}^+$
	1.347 ± 5			
1.46 ± 15	1.460 ± 5	1	0.098	$\frac{3}{2}^-$
1.56 ± 10	1.556 ± 5	2	1.01	$\frac{3}{2}^+$
2.790 ± 15	2.784 ± 5	4^b	$< 0.027^b$	$\frac{9}{2}^+$
	3.912 ± 5			
4.00 ± 20^c	4.002 ± 5	(3)	(0.019)	$(\frac{7}{2}^-)$
	4.036 ± 10			
4.39 ± 20	4.385 ± 5	$(4)^b$	$(0.048)^b$	$(\frac{7}{2}^+)$
4.56 ± 20	4.555 ± 5	2	0.31	$\frac{3}{2}^+^d$
	4.675 ± 10^c			
5.10 ± 20	5.113 ± 5	$(2, 3)^{a,b}$		$\frac{5}{2}^-$
	5.34 ± 5	(2, 3)	0.0065	$\frac{5}{2}^+$
5.43 ± 30^c	5.428 ± 8	(2, 3)	(0.042)	$(\frac{3}{2}^+)$
	5.495 ± 5^c			
5.55 ± 30^c	5.54 ± 5	3	0.14	$\frac{7}{2}^-$
	5.943 ± 5	0	0.014	$\frac{1}{2}^+$
6.08 ± 30^c	6.095 ± 5	1	0.12	$\frac{1}{2}^-$
	6.167 ± 5			
6.25 ± 30^c	6.255 ± 8	(0)	0.19	$(\frac{1}{2}^+)$
6.47 ± 30^c	6.503 ± 5^c	2^b	0.133^b	$\frac{3}{2}^+$
	6.595 ± 10			
6.76 ± 30	6.792 ± 5	1^b	0.29^b	$\frac{3}{2}^-$
6.90 ± 30^c	6.93 ± 5	(2, 3)		$(\frac{5}{2}^+, \frac{7}{2}^-)$
7.10 ± 30^c	7.112 ± 8^c	2	0.087	$\frac{5}{2}^+$
	7.26 ± 5			
	7.364 ± 5	0	0.091	$\frac{1}{2}^+$
7.56 ± 30	7.540 ± 3	2	0.665	$\frac{5}{2}^+; T = \frac{3}{2}$

Table 19.16: Energy levels of ^{19}F from $^{18}\text{O}(^3\text{He}, \text{d})^{19}\text{F}$ (continued)

E_x (MeV \pm keV)		l^a	$C^2S(2J_f + 1)^{a,b}$	J^π^a
(1970GR04)	(1970SC25)			
7.74 \pm 30	7.665 \pm 5	(2)	0.035	$(\frac{3}{2}^+)$
	7.702 \pm 5	(0, 1)	(0.052)	$(\frac{3}{2}^-)$
	8.015 \pm 5	2	0.26	$(\frac{5}{2}^+)$
	8.086 \pm 5	(2, 3)	0.097	$(\frac{5}{2}^+)$
	8.135 \pm 5	(0, 1)	0.156	$(\frac{1}{2}^+)$
	8.198 \pm 5	(2, 3)	0.035	$(\frac{5}{2}^+)$
	8.255 \pm 5	(2)	0.035	$(\frac{5}{2}^+)$
	8.31 \pm 5			
	8.592 \pm 10	(2, 3)		
	8.795 \pm 15	0	(0.13)	$\frac{1}{2}^+; T = \frac{3}{2}$
	9.113 \pm 10			
	9.18 \pm 15			
	9.596 \pm 10			
	9.682 \pm 15			
	10.275 \pm 15			
10.33 \pm 15				
10.525 \pm 15				

^a (1970SC25): $E(^3\text{He}) = 16$ MeV.

^b See also (1970GR04).

^c Unresolved.

^d J probably $\frac{5}{2}$. This appears to be a different state from the one involved in the $^{20}\text{Ne}(\text{d}, ^3\text{He})^{19}\text{F}$ reaction, with $J^\pi = \frac{3}{2}^-$; see (1970KA31).

At $E_d = 5.0$ and 5.2 MeV, reaction (b) appears to involve ^{19}F states at ≈ 8 to 10 MeV (1970BO08).

23. $^{18}\text{O}(^3\text{He}, \text{d})^{19}\text{F}$

$$Q_m = 2.499$$

Angular distributions of the deuterons corresponding to many states of ^{19}F have been analyzed by DWBA: the results are shown in Table 19.16 (1970SC25: $E(^3\text{He}) = 16$ MeV; (1968LE18, 1970GR04): $E(^3\text{He}) = 11$ MeV). See also (1965ER04).

The spectroscopic factors obtained by DWBA for $^{19}\text{F}^*(7.54, 8.80)$, the $T = \frac{3}{2}$, $J^\pi = \frac{5}{2}^+$ and $\frac{1}{2}^+$ analogs of $^{19}\text{O}^*(0, 1.47)$ are in good agreement with those obtained for the ^{19}O states in the $^{18}\text{O}(\text{d}, \text{p})^{19}\text{O}$ reaction (1971FO18). See also (1970SC25). See also (1966LA1E, 1966RO01, 1967RO1F, 1970WA02; theor.).

24. $^{18}\text{O}(\alpha, \text{t})^{19}\text{F}$ $Q_{\text{m}} = -11.822$

Not reported.

25. $^{18}\text{O}({}^6\text{Li}, \alpha\text{n})^{19}\text{F}$ $Q_{\text{m}} = 4.295$

τ_{m} of $^{19}\text{F}^*(0.110)$ is 847 ± 10 psec: see Table 19.10 (1969NI09).

26. $^{19}\text{O}(\beta^-)^{19}\text{F}$ $Q_{\text{m}} = 4.819$

The decay is primarily by allowed transitions to $^{19}\text{F}^*(0.197, 1.55)$, $J^\pi = \frac{5}{2}^+, \frac{3}{2}^+$, with very weak branches also observed to $^{19}\text{F}^*(0.11)$, $J^\pi = \frac{1}{2}^-$ ($\log f_1 t = 10.16$) and $^{19}\text{F}^*(4.39)$, $J^\pi = \frac{7}{2}^+$ ($\log ft = 3.60$): see Table 19.17 (1959AL06, 1959JO26, 1966OL01, 1970CO22). The half-life is 26.91 ± 0.08 sec: see reaction 1 in ^{19}O . The character of the allowed decay to the $\frac{5}{2}^+$ and $\frac{3}{2}^+$ states, and the forbiddenness of the decay to the ground state of ^{19}F are consistent with $J^\pi = \frac{5}{2}^+$ for the ground state of ^{19}O and then with $(\frac{7}{2}^+)$ for $^{19}\text{F}^*(4.39)$: see (1966OL01). Gamma-ray branching ratios are displayed in Table 19.9 (1966OL01, 1970CO22). See also (1961OK1A) and (1970MC23; theor.).

27. $^{19}\text{F}(\gamma, \gamma)^{19}\text{F}$

The energy of the first excited state is 109.894 ± 0.005 keV; its width is $(5.1 \pm 0.7) \times 10^{-7}$ eV (1962BO1A, 1962SE12). Lifetimes of $^{19}\text{F}^*(0.11, 1.46)$ are shown in Table 19.10 (1962SE12, 1964BO22). The scattering cross section is relatively small and structureless in the range $E_\gamma = 14$ to 30 MeV (1967LO1B). See also (1960BO23, 1960RE05, 1962BO17).

28. $^{19}\text{F}(\gamma, \text{n})^{18}\text{F}$ $Q_{\text{m}} = -10.431$

Table 19.17: Branching in $^{19}\text{O}(\beta^-)^{19}\text{F}$ ^a

Decay to $^{19}\text{F}^*$	J^π	Branch (%)	$\log ft$	Refs.
0	$\frac{1}{2}^+$	≤ 4	≥ 6.5	(1959AL06, 1959JO26)
0.110 ^b	$\frac{1}{2}^-$	$0.048^{+0.012}_{-0.033}$	$10.16^{+0.51}_{-0.10}$ ^e	(1970CO22)
0.197 ^c	$\frac{5}{2}^+$	41.5^{+2}_{-5}	$5.42^{+0.06}_{-0.03}$ ^f	(1959AL06)
1.35	$\frac{5}{2}^-$		≥ 7.1	(1959JO26)
1.46	$\frac{3}{2}^-$		≥ 6.7	(1959JO26)
1.55 ^d	$\frac{3}{2}^+$	58.5 ± 2	4.59 ± 0.03 ^f	(1959AL06)
2.78	$\frac{9}{2}^+$	≤ 0.15	> 7.4	(1959AL06, 1966OL01)
4.39	$\frac{7}{2}^+$	0.16 ± 0.012	3.60 ^f	(1966OL01)

^a See also (1959AJ76).

^b $E_\gamma = 111.5 \pm 1.5$ keV (1954JO21), 112 ± 2 keV (1959JO26).

^c $E_\gamma = 199.6 \pm 1.5$ keV (1954JO21).

^d $E_x = 1.5539 \pm 0.0013$ MeV (1966AL12).

^e $\log f_1 t$ (1970CO22).

^f Based on Q_m and $\tau_{1/2} = 26.91 \pm 0.08$ sec: see ^{19}O , reaction 1 (B.A. Zimmerman, private communication).

Maxima are reported at (10.6), 12.4, 14.0, 16.1, 17.2 and 19.3 MeV in the (γ, n) cross section (1960KI06). Three ground state photoneutron groups are reported, corresponding to $^{19}\text{F}^*(12.10, 12.38, 16.24)$ ((1971SH1G), and N.K. Sherman, private communication). These states are thought to have $J^\pi = \frac{1}{2}^-$ or $\frac{3}{2}^-$; $T = \frac{1}{2}$. See also (1960GE06, 1967BA28) and (1959AJ76). At $E_\gamma = 20.5$ MeV, the (γ, n) cross section is 3.30 ± 0.41 b (1961DE1A, 1962DE03).

The nuclear absorption cross section shows a peak at $E_\gamma = 20.09 \pm 0.05$ MeV with an integrated cross section of 3.5 MeV \cdot mb (1964TE04) and a very broad giant resonance extending beyond 30 MeV and with considerable strength also below 20 MeV (1969BE92). These results disagree with those of (1966DO1D, 1967NI1B) which show considerable structure. See also (1960SA09, 1965DE1J, 1970JO1H, 1971DI1F, 1971FR11) and (1958SP1A, 1959FU1A, 1971AN08; theor.).

$$29. \text{ (a) } ^{19}\text{F}(\gamma, 2n)^{17}\text{F} \quad Q_m = -19.581$$

$$\text{ (b) } ^{19}\text{F}(\gamma, 2pn)^{16}\text{N} \quad Q_m = -29.820$$

For reaction (a) see (1970JO1H); for reaction (b) see (1968ME23).

$$30. ^{19}\text{F}(\gamma, p)^{18}\text{O} \quad Q_m = -7.993$$

Peaks are reported corresponding to ^{19}F states at $E_x = 10.4, 11.4, 11.9, (12.8), 13.6, 15.4, 16.5$ and (18.1) MeV (1960FO10). See also (1963MU08, 1964SE09). See also (1961DO08, 1962BR1E) and (1958SP1A, 1965SH1D; theor.).

31. (a) $^{19}\text{F}(e, e)^{19}\text{F}$

(b) $^{19}\text{F}(e, ep)^{18}\text{O}$ $Q_m = -7.993$

At $E_e = 41.5$ MeV (180°) the elastic scattering cross section has been measured by (1963GO04) and the excitation of a $\frac{3}{2}^+$ state of ^{19}F at $E_x = 7.7$ MeV [$T = \frac{3}{2}$; see Table 19.9] is reported by (1963BA19). The excitation of $^{19}\text{F}^*(0.197, 1.35, 1.55)$ has been observed by (1970HA1R): $\Gamma_\gamma(1.35 \rightarrow 0) = (8.1 \pm 1.7) \times 10^{-10}$ eV (E3) and $\Gamma_\gamma(1.55 \rightarrow 0) = (1.76 \pm 0.15) \times 10^{-4}$ eV (E2) (1970WA08). See also (1970WA1N) and (1965PR1C, 1966GO1C; theor.).

Reaction (b), studied at $E_e = 18, 24.5$ and 30 MeV, shows resonances (assuming ground state transitions) at $E_x = 11.42, 11.90, 12.74, 15.70$ and 18.7 MeV in ^{19}F (1962DO1A).

32. $^{19}\text{F}(n, n)^{19}\text{F}$

Angular distributions have been reported at $E_n = 2.56$ MeV (1960DA08; $\gamma_{0.110}, \gamma_{0.197}$), 14.1 MeV (1970CL03; n_{0+1+2}, n_{3+4+5}) and 14.2 MeV (1966BO21; $n_{0+1+2}, n_{3+4+5}, n_6$ and n to $^{19}\text{F}^*(4.7, 5.4)$). Gamma-ray energy measurements are reported by (1968SP01) and (1969WH1A); see Table 19.18. Branching ratios are listed in Table 19.9 (1968SP01). See also (1959BI1A, 1959KO60, 1961AN1A, 1961RO01, 1966VE1B, 1971NI05) and (1963DO1C; theor.).

33. (a) $^{19}\text{F}(p, p)^{19}\text{F}$

(b) $^{19}\text{F}(p, 2p)^{18}\text{O}$ $Q_m = -7.993$

Table 19.19 displays energy levels of ^{19}F derived from measurements of inelastically scattered protons and γ -rays (1956SQ1A, 1961VA28, 1962VA28, 1968GU07, 1969PO03). See also (1959AJ76). Angular distributions of various proton groups have been measured at $E_p = 4.26, 5.96$ and 6.87 MeV (1967TH06; p_0, p_1, p_2), 6.6 MeV (1961VA28, 1962VA28; $p_2 \rightarrow p_6$), 6.78 to 7.41 MeV (1961KO08; p_{3+4+5}, p_6), 13.9 MeV (1966LU03; p_0, p_2, p_5) and 17.5 MeV (1968CR03; $p_0, p_2 \rightarrow p_7, p_{8+9}, p_{10}, p_{11}$ and p to $^{19}\text{F}^*(5.43)$). (1967TH06) report that the scattering below $E_p = 7$ MeV is adequately described as scattering from an axially symmetric quadrupole-deformed potential, and that only positive parity states are strongly excited. Branching ratios for observed transitions are shown in Table 19.9 and lifetimes in Table 19.10 (1969PO03, 1970LA02). (1969PO03) find a pronounced inhibition of the four E1 transitions involved in the decay of

Table 19.18: Energy levels of ^{19}F from $^{19}\text{F}(n, n')^{19}\text{F}$ ^a

E_x (keV) ^b	
(1968SP01)	(1969WH1A) ^c
109.8 ± 0.2	109.844 ± 0.030
197.2 ± 0.2	197.147 ± 0.012
1345.4 ± 0.3	1345.81 ± 0.20
1456.9 ± 1.1	1458.16 ± 0.50
1554.0 ± 0.3	1554.01 ± 0.40
2775.1 ± 3.5	

^a See also (1960MI1A) and (1959AJ76).

^b From γ -ray measurements.

^c And D.H. White, private communication.

$^{19}\text{F}^*(1.35, 1.46, 1.55)$ in accord with the predictions of (1964HA21). Partial widths have been determined by (1969PO03). (1970LA02) have studied the decay of $^{19}\text{F}^*(1.46)$: from the angular distribution of the γ -rays, measurement of their linear polarization and from τ_m , $|M|^2(\text{E}2) = 14_{-8}^{+26}$, $|M|^2(\text{M}1) = 0.10 \pm 0.03$ W.u. The mixing ratios in the γ -decay of $^{19}\text{F}^*(1.46)$ are $-0.1 < \delta_{\text{M}2/\text{E}1} < 0.0$ for $1.46 \rightarrow 0.20$, $0.30 < \delta_{\text{E}2/\text{M}1} < 0.38$ for $1.46 \rightarrow 0.11$ and $|\delta_{\text{M}2/\text{E}1}| < 0.06$ for $1.46 \rightarrow 0$. These results indicate that $^{19}\text{F}^*(1.46)$ cannot be explained as a $p_{1/2}$ proton hole coupled to the pure ground state rotational band (1971HA30). See also (PR63A) for other linear polarization data and (1965KL01, 1966RO03, 1966TH02, 1967HI1C, 1968HI1F) for other angular correlation work. See also (1964NE08).

Additional parameters measured for the $\frac{5}{2}^+$ state at 0.197 MeV are : $Q = 0.11 \pm 0.02$ b (1964SU01: see also (1958SU58)), $Q = -(0.10 \pm 0.02)$ b (1968RI1N); $\mu = 3.69 \pm 0.04$ nm (1961FR07): [see also (1964BO40)]; $\tau_m = 128 \pm 2$ nsec (1968KL05), 129.9 ± 2.3 nsec (1967BE14): see also Table 19.10 for additional measurements of τ_m for this and for other ^{19}F states (1963GA14, 1963VA01, 1963VA12, 1965BO28, 1969ME13, 1969PO03). See also (PR63A).

See also (1959TR1A, 1968JA1H) and (1967AM1D; theor.). For reaction (b) see the review of (1965RI1A). See also (1961PU1A, 1964GO1C) and (1966JA1A; theor.).

34. $^{19}\text{F}(d, d)^{19}\text{F}$

Angular distributions of elastically scattered deuterons have been measured at $E_d = 2.0$ and 2.2 MeV (1968BI1A), 2.6 to 4.0 MeV (1969ZA01) and at 15 MeV (1965DI1C, 1966DE09, 1970DE06). In addition, angular distributions have been measured at $E_d = 15$ MeV for five groups of inelastically scattered deuterons (1970DE06; $d_1 \rightarrow d_5$). Analysis by DWBA leads to

Table 19.19: States in ^{19}F from $^{19}\text{F}(p, p')^{19}\text{F}$ ^a

E_x (MeV \pm keV)			
(1956SQ1A)	(1961VA28, 1962VA28)	(1968GU07)	(1969PO03)
0.111 \pm 2			
0.197 \pm 2	0.198 \pm 12		0.1976 \pm 0.6
1.350 \pm 5	1.344 \pm 12		1.3458 \pm 0.2
1.462 \pm 5	1.458 \pm 12		1.4591 \pm 0.5
1.558 \pm 5	1.553 \pm 12		1.5542 \pm 0.4
2.784 \pm 8	2.789 \pm 6		
3.912 \pm 10		3.92 \pm 10	
4.002 \pm 10		4.01 \pm 10	
4.036 \pm 10		4.04 \pm 10	
		4.39 \pm 10	
		4.56 \pm 10	
		4.69 \pm 10	
		5.11 \pm 10	
		5.34 \pm 10	
		5.42 \pm 10	
		5.47 \pm 10	
		5.50 \pm 10	
		5.54 \pm 10	
		5.63 \pm 10	
		5.94 \pm 10	
		(6.08)	
		6.09 \pm 10	
		6.17 \pm 10	
		6.25 \pm 10	
		6.29 \pm 10	
		6.33 \pm 10	

^a See also Table 19.8 in (1959AJ76).

$B(E2)_{\downarrow} = 9 \pm 3$ W.u. for the $0.197 \rightarrow 0$ transition and 10 ± 3 W.u. for the $1.56 \rightarrow 0$ transition, and to $B(E3)_{\downarrow} = 1.4 \pm 0.6$ W.u. for the $1.35 \rightarrow 0$ transition (1970DE06). See also (1969VE09, 1970VE06; theor.).

35. $^{19}\text{F}(t, t)^{19}\text{F}$

Angular distributions of elastically scattered tritons have been measured at $E_t = 2$ MeV (1969HE08) and at 7.2 MeV (1964PU01). See also (1968HO1C; theor.).

36. $^{19}\text{F}(^3\text{He}, ^3\text{He})^{19}\text{F}$

Angular distributions of elastically scattered ^3He particles are reported at $E(^3\text{He}) = 4.0, 6.0$ and 8.0 MeV (1966MA43), at 9 MeV (1965SI18), at 11.0 MeV (1970SC23) and at 29 MeV (1961CA19, 1962GA17). See also (1959BR72) and (1968HO1C; theor.). (1967CO1J) also report the excitation of $^{19}\text{F}^*(0.11)$.

37. (a) $^{19}\text{F}(\alpha, \alpha)^{19}\text{F}$

(b) $^{19}\text{F}(\alpha, 2\alpha)^{15}\text{N} \quad Q_m = -4.013$

Angular distributions of elastically scattered α -particles have been measured at $E_\alpha = 19.9$ to 23.3 MeV (1968AG1D, 1970AG08, 1970FE1E), and at 38 MeV (1960AG01). The energy of the γ -ray from the $1.35 \rightarrow 0.11$ transition is 1235.8 ± 0.2 keV. Using $E_x = 109.893 \pm 0.004$ keV for the energy of the first excited state, E_x for $^{19}\text{F}^*$ is then 1345.7 ± 0.2 keV (1967WA13).

At $E_\alpha = 12.7$ MeV, a state at 4.648 MeV is populated which is then observed to γ -decay to the $\frac{9}{2}^+$ state at 2.78 MeV. The angular distribution of the cascade γ -rays and the lifetime of $^{19}\text{F}^*(4.65)$, set $J^\pi = \frac{13}{2}^+$ for $^{19}\text{F}^*(4.65)$ (1969JA09). For a listing of the τ_m for this state and $^{19}\text{F}^*(1.35, 2.78)$ see Table 19.10 (1968PA11, 1968RO07, 1969JA09, 1969JO10). See also (1958CA97, 1959BR72, 1963FR12, 1964FR03, 1970BL1F). For reaction (b) see (1963LA02).

38. (a) $^{19}\text{F}(^6\text{Li}, ^6\text{Li})^{19}\text{F}$

(b) $^{19}\text{F}(^7\text{Li}, ^7\text{Li})^{19}\text{F}$

Elastic scattering angular distributions have been reported for both reactions at $E(\text{Li}) = 20$ MeV (1969BE90). See also (1959LI55).

39. (a) $^{19}\text{F}(^{10}\text{B}, ^{10}\text{B})^{19}\text{F}$
 (b) $^{19}\text{F}(^{12}\text{C}, ^{12}\text{C})^{19}\text{F}$

For reaction (a) see (1971KN05). For reaction (b), see reaction 4.

40. $^{19}\text{F}(^{14}\text{N}, ^{14}\text{N})^{19}\text{F}$

$B(\text{E}2)\uparrow$, from the Coulomb excitation of $^{19}\text{F}^*(1.55) = 2.44 \times 10^{-51} e^2 \cdot \text{cm}^4$ (1966AF01). See also (1956AL1D, 1956AL36, 1956AL55).

41. $^{19}\text{F}(^{20}\text{Ne}, ^{20}\text{Ne})^{19}\text{F}$

$B(\text{E}1)\uparrow$, from the Coulomb excitation of $^{19}\text{F}^*(0.110) = 6.3 \times 10^{-30} e^2 \cdot \text{cm}^2$; $B(\text{E}2)\uparrow$ for $^{19}\text{F}^*(0.197) = 5 \times 10^{-51} e^2 \cdot \text{cm}^4$. Reported mean lifetimes for these states are displayed in Table 19.10 (1960ST17, 1962RI09).

42. $^{19}\text{Ne}(\beta^+)^{19}\text{F}$ $Q_{\text{m}} = 3.238$

See ^{19}Ne .

43. $^{20}\text{Ne}(\gamma, \text{p})^{19}\text{F}$ $Q_{\text{m}} = -12.845$

See (1962DO1A, 1963FI07, 1969HO16). See also (1962NU1A) and ^{20}Ne .

44. $^{20}\text{Ne}(\text{p}, \text{p}\alpha)^{19}\text{F}$ $Q_{\text{m}} = -12.845$ †

See (1967CH04).

† This reaction should be completely deleted. Please refer to the Errata for this evaluation found directly after the list of references in (1974AJ01).

45. $^{20}\text{Ne}(\text{n}, \text{d})^{19}\text{F}$ $Q_{\text{m}} = -10.621$

Table 19.20: States of ^{19}F from $^{20}\text{Ne}(t, \alpha)^{19}\text{F}$ (1961SI03)

E_x (MeV \pm keV)	E_x (MeV \pm keV)
2.794 ± 15	5.481 ± 15
3.917 ± 15	5.539 ± 15
4.032 ± 15	5.628 ± 15
4.385 ± 15	5.937 ± 20
4.563 ± 15	6.092 ± 15
(4.690 ± 40)	6.169 ± 30
5.102 ± 15	6.247 ± 25
5.343 ± 15	6.501 ± 25

See (1960VE06).

46. $^{20}\text{Ne}(d, ^3\text{He})^{19}\text{F}$ $Q_m = -7.351$

At $E_d = 52$ MeV, ^3He groups are observed, and angular distributions are reported, corresponding to states at $E_x = 0.15 \pm 0.04$, 1.51 ± 0.03 , 2.83 ± 0.04 ($l = 4$), 3.99 ± 0.07 , 4.56 ± 0.02 ($l = 1$), 5.44 ± 0.05 , 5.69 ± 0.07 ($l = 1$), 6.10 ± 0.03 , 6.78 ± 0.02 ($l = 1$) and 10.42 ± 0.15 MeV (1970KA31). See also (1971DU12; theor.).

47. $^{20}\text{Ne}(t, \alpha)^{19}\text{F}$ $Q_m = 6.969$

At $E_t = 2.6$ MeV, alpha groups are observed to states with $2.7 < E_x < 6.6$ MeV: see Table 19.20 (1961SI03). The lifetime, τ_m , of $^{19}\text{F}^*(4.65)$, $J^\pi = \frac{13}{2}^+$ [$K^\pi = \frac{1}{2}^+$ rotational band] is 2.3 ± 0.5 psec (see also Table 19.10): the E2 strength of the transition to $^{19}\text{F}^*(2.78)$ is 5.0 ± 1.1 W.u. (1969BH01).

48. $^{21}\text{Ne}(n, t)^{19}\text{F}$ $Q_m = -11.124$

Not reported.

49. $^{21}\text{Ne}(p, ^3\text{He})^{19}\text{F}$ $Q_m = -11.888$

At $E_p = 45$ MeV, ^3He groups are observed to some $T = \frac{1}{2}$ states in ^{19}F and to the $J^\pi = \frac{3}{2}^+$; $T = \frac{3}{2}$ analog of $^{19}\text{O}^*(0.095)$: $E_x = 7.660 \pm 0.035$ MeV (1969HA38): see also reaction 12 in ^{19}Ne .

$$50. \ ^{21}\text{Ne}(d, \alpha)^{19}\text{F} \quad Q_m = 6.465$$

Alpha-particle groups have been observed to $^{19}\text{F}^* = 0, 113 \pm 8$ and 192 ± 12 keV (1952MI54).

$$51. \ ^{22}\text{Ne}(p, \alpha)^{19}\text{F} \quad Q_m = -1.675$$

Not reported.

$$52. \ ^{23}\text{Na}(n, n'\alpha)^{19}\text{F} \quad Q_m = -10.468$$

See (1966WO03).

¹⁹Ne
(Figs. 7 and 8)

GENERAL: (See also (1959AJ76).)

Shell model: (1957WI1E, 1960TA1C, 1962BH09, 1967BO09, 1967GU1D, 1968EL1C, 1968WA04, 1971AR25, 1972LE1L).

Cluster, collective and deformed models: (1960RA1A, 1969BA2E, 1970BA2B, 1972LE1L).

Astrophysical questions: (1970BA1M).

Electromagnetic transitions: (1960RA1A, 1967GU1D, 1968EL1C, 1971AR25).

Special levels: (1960GO1C, 1968CE1A, 1969GA1G).

Other theoretical topics: (1968MU1B, 1972LE1L).

Ground state: $\mu = -1.887$ nm (1963CO22, 1967CO1D). See also (1964LI14, 1964ST1B, 1967GU1D, 1967SH14, 1968EL1C, 1968LE1L, 1969PE1D, 1970LE1A, 1971AR25, 1972LE1L).

1. ¹⁹Ne(β^+)¹⁹F $Q_m = 3.238$

The half-life is 17.40 ± 0.04 sec: see Table 19.22 (1962EA02, 1968GO10). The decay is to ¹⁹F(0) (1954JO21). $\log ft = 3.23$ ³. See also (1965HA31, 1970ST04). The allowed nature of the decay sets $J^\pi = \frac{1}{2}^+$ for ¹⁹Ne(0).

A test of time reversal invariance and a measurement of positron and neutrino asymmetry has been made by (1969CA14). See also (1959AL1E, 1959AL10, 1961JO03, 1963CO22, 1967CA1M, 1967CO1D, 1969CA1H) and (1965GA1D, 1966MI1F, 1967CA1N, 1969LE1D, 1969SU15, 1970KO41, 1970MC23, 1970PR1G, 1971WE1F; theor.).

2. ¹⁰B(¹⁴N, α n)¹⁹Ne $Q_m = 2.667$

The lifetime, τ_m , of ¹⁹Ne*(0.28) is 61_{-20}^{+4} psec (1969NI09) [see Table 19.23].

3. (a) ¹⁶O(³He, γ)¹⁹Ne $Q_m = 8.443$
 (b) ¹⁶O(³He, n)¹⁸Ne $Q_m = -3.196$ $E_b = 8.443$
 (c) ¹⁶O(³He, p)¹⁸F $Q_m = 2.033$
 (d) ¹⁶O(³He, ³He)¹⁶O

³ B.A. Zimmerman, private communication.

Table 19.21: Energy levels of ^{19}Ne

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^+; \frac{1}{2}$	$\tau_{1/2} = 17.40 \pm 0.04$ sec	β^+	1, 5, 7, 8, 9, 10, 12, 13
0.23833 ± 0.11	$\frac{5}{2}^+$	$\tau_m = 26.0 \pm 0.8$ nsec	γ	5, 6, 7, 8, 12, 13
0.27520 ± 0.14	$\frac{1}{2}^-$	$\tau_m = 61.4 \pm 3.0$ psec	γ	2, 5, 7, 12
1.5078 ± 0.4	$\frac{5}{2}^-$	$\tau_m = 4.1_{-1.4}^{+3.5}$ psec	γ	4, 5, 6, 7, 12
1.5362 ± 0.5	$\frac{3}{2}^+$	$\tau_m = 28 \pm 15$ fsec	γ	4, 5, 6, 7, 8, 12
1.6152 ± 0.7	$\frac{3}{2}^-$	$\tau_m = 180 \pm 60$ fsec	γ	4, 5, 6, 7, 12
2.7940 ± 1.3	$\frac{9}{2}^+$	$\tau_m = 330 \pm 130$ fsec	γ	4, 5, 6, 7, 8, 12, 13
(3.841 ± 15)				12
4.026 ± 8	$(\frac{3}{2}, \frac{5}{2})^+$			5, 6, 12, 13
4.146 ± 8	$(\frac{7}{2}^-)$		γ	5, 6, 12
4.200 ± 10	$(\frac{9}{2}^-)$			5, 12
4.368 ± 8	$(\frac{7}{2}^+)$			5, 6, 12
4.549 ± 8	$(\frac{1}{2}, \frac{3}{2})^-$			5, 6, 12
4.624 ± 10				5, 6, 12
4.710 ± 9				5, 6, 12
4.783 ± 20				5, 12
5.087 ± 6				5, 6, 12
5.351 ± 10	$\frac{1}{2}^+$			5, 12
5.425 ± 7				5, 12
5.463 ± 20				12
5.545 ± 10				12
5.832 ± 9				12
6.013 ± 7	$(\frac{3}{2}, \frac{1}{2})^-$			12
6.096 ± 7				12
6.149 ± 20				12
6.290 ± 7				12
6.437 ± 9				12
6.743 ± 7	$(\frac{3}{2}, \frac{1}{2})^-$			12
6.862 ± 7				12
7.067 ± 9				12
(7.178 ± 15)				12
7.253 ± 10				12
(7.326 ± 15)				12

Table 19.21: Energy levels of ^{19}Ne (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions			
(7.531 \pm 15)	$\frac{3}{2}^+; \frac{3}{2}$			12			
7.616 \pm 16				12, 13			
7.700 \pm 10				12			
(7.788 \pm 10)				12			
7.994 \pm 15				12			
8.063 \pm 15				12			
8.236 \pm 10 ^a				12			
8.440 \pm 10				12			
8.523 \pm 10				12			
(8.810 \pm 25)				12			
8.915 \pm 10				12			
9.013 \pm 10				12			
9.100 \pm 20				12			
9.240 \pm 20				12			
9.489 \pm 25				12			
9.886 \pm 50 ^a				12			
10.46				$\frac{1}{2}^+$	$\Gamma = 355$	p, ^3He , α	3, 12
10.48				$\frac{3}{2}^+$	45	p, ^3He , α	3, 12
10.613 \pm 20	$\frac{3}{2}^- (\frac{1}{2}^-)$			12			
11.51				^3He , α	3		
12.23				^3He , α	3		
12.50				^3He , α	3		
12.69 \pm 50				$\frac{1}{2}^+$	180 ± 40	p, ^3He , α	3

^a Broad or unresolved states.

$$(e) \text{}^{16}\text{O}({}^3\text{He}, \alpha){}^{15}\text{O} \quad Q_m = 4.909$$

$$(f) \text{}^{16}\text{O}({}^3\text{He}, 2n){}^{17}\text{Ne} \quad Q_m = -22.430$$

$$(g) \text{}^{16}\text{O}({}^3\text{He}, 2\alpha){}^{11}\text{C} \quad Q_m = 5.305$$

Studies of the excitation functions for reactions (c) and (e) [α_0 , $p_{1+2+3+4}$, p_5 , p_6 , p_7] for $E({}^3\text{He}) = 2.0$ to 3.0 MeV are interpreted in terms of two resonances at $E({}^3\text{He}) = 2.400$ and

Table 19.22: The half-life of ^{19}Ne ^a

$\tau_{1/2}$ (sec)	Refs.
18.5 ± 0.5	(1952SC15)
18.3 ± 0.5	(1957AL29)
17.7 ± 0.1	(1957PE12)
16.72 ± 0.05	(1960JA12)
19.5 ± 1.0	(1960WA04)
$17.43 \pm 0.06^*$	(1962EA02)
16.5 ± 1	(1964VA23)
$17.36 \pm 0.06^*$	(1968GO10)
17.40 ± 0.04	mean of values marked by asterisk

^a See also (1959AJ76, 1964BAZZ).

2.425 MeV ($^{19}\text{Ne}^*(10.46, 10.48)$) with $\Gamma = 355$ and 45 keV, $J^\pi = \frac{1}{2}^+$ and $\frac{3}{2}^+$, respectively (1959BR79, 1961SI09). The capture cross section at the 2.40 MeV resonance (reaction (a)) is $< 0.8 \mu\text{b}$ (1959BR79). Studies of the (p_0, p_1, p_5) yields, of the elastic yield (reaction (d)), and the analysis of angular distributions in the range $E(^3\text{He}) = 4.3$ to 5.6 MeV show the presence of a single resonance at $E(^3\text{He}) = 5.05 \pm 0.05$ MeV [$^{19}\text{Ne}^*(12.69)$], $\Gamma = 180 \pm 40$ keV, $J^\pi = \frac{1}{2}^+$ (1967RO10). A resonance-like structure is also reported at $E(^3\text{He}) \approx 9.5$ MeV in the yields of neutron groups (reaction (b)) (1970AD02).

The elastic scattering (reaction (d)) and the α_0 yield (reaction (e)) have been studied for $E(^3\text{He}) = 3.2$ to 7.0 MeV by (1971OT1C). They report a state at $E_x = 11.51$ MeV [$E(^3\text{He}) = 3.65$ MeV] with $J^\pi = \frac{3}{2}^-$ or $(\frac{1}{2}^-)$, $\Gamma_{\text{c.m.}} = 24$ keV. In addition, two states at $E_x = 12.23$ and 12.50 MeV [$E(^3\text{He}) = 4.50$ and 4.82 MeV], $J^\pi = \frac{5}{2}^+$ and $\frac{7}{2}^+$, $\Gamma_{\text{c.m.}} = 200$ and 150 keV respectively, are indicated by the work of (1971OT1C). See also (1970OT1B, 1970WE1K).

The α_0 yield for $E(^3\text{He}) = 4.0$ to 9.0 MeV shows fluctuations which are analyzed with a coherence width of 130 ± 20 keV (1969DA08). These fluctuations continue at least to 11.8 MeV (1969BR07).

For reaction (b), see also (1961DU02: 3.4 to 5.6 MeV), (1961TO03: 3.9 to 5.7 MeV), (1968TO09: 9.1 to 10.4 MeV) and (1964BR1G, 1965BR1B: 11 to 29 MeV). For reaction (c), see also (1966GO1E: 2.0 to 5.5 MeV), (1966HA21: 3.0 to 9.5 MeV) and (1964BR1G, 1965BR1B: 11 to 29 MeV). For reaction (d), see also (1965AL05: 8.1 to 10 MeV) and (1969BR07: 9.8 to 11.7 MeV). For polarization measurements, see (1970MC1F, 1971MC1M). For reaction (e), see also (1966HA21: 3 to 9.5 MeV), (1959HI73: 5.6 to 5.9 MeV), (1965AL05: 8.0 to 10.0 MeV). For reaction (f) see (1967ES02) and for reaction (g) see (1964BR1G, 1965BR1B). See also (1965JI1A, 1966MA1R, 1969PA1C).

Table 19.23: Radiative decays of ^{19}Ne levels

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branch (%)	τ_m^a and Γ_γ	Refs.
0.24	$\frac{5}{2}^+$	0	$\frac{1}{2}^+$	100	26.6 ± 1.2 nsec	(1967BE14)
0.28	$\frac{1}{2}^-$	0	$\frac{1}{2}^+$	(100) ^c	25.5 ± 1.0 nsec 61_{-20}^{+4} psec 61.4 ± 3.0 psec	(1969BL02) (1969NI09) (1970BH02)
1.51 ^b	$\frac{5}{2}^-$	0	$\frac{1}{2}^+$	< 3	$\left\{ \begin{array}{l} 4.1_{-1.4}^{+3.5} \text{ psec} \\ \Gamma_\gamma = 0.17 \pm 0.08 \text{ meV} \end{array} \right.$	(1970GI09)
		0.24	$\frac{5}{2}^+$	12 ± 3		
		0.28	$\frac{1}{2}^-$	88 ± 3		
1.54 ^b	$\frac{3}{2}^+$	0	$\frac{1}{2}^+$	< 6	$\left\{ \begin{array}{l} 28 \pm 15 \text{ fsec} \\ \Gamma_\gamma = 24_{-8}^{+27} \text{ meV} \end{array} \right.$	(1970GI09)
		0.24	$\frac{5}{2}^+$	95 ± 3		
		0.28	$\frac{1}{2}^-$	5 ± 3		
1.62 ^b	$\frac{3}{2}^-$	0	$\frac{1}{2}^+$	20 ± 3	$\left\{ \begin{array}{l} 180 \pm 60 \text{ fsec} \\ \Gamma_\gamma = 3.7_{-0.9}^{+1.8} \text{ meV} \end{array} \right.$	(1970GI09)
		0.24	$\frac{5}{2}^+$	10 ± 3		
		0.28	$\frac{1}{2}^-$	70 ± 4		
2.79 ^b	$\frac{9}{2}^+$	0	$\frac{1}{2}^+$	< 10	$\left\{ \begin{array}{l} 330 \pm 130 \text{ fsec} \\ \Gamma_\gamma = 2.0_{-0.6}^{+1.3} \text{ meV} \end{array} \right.$	(1970GI09)
		0.24	$\frac{5}{2}^+$	100		
		0.28	$\frac{1}{2}^-$	< 10		
		1.51	$\frac{5}{2}^-$	< 12		
		1.54	$\frac{3}{2}^+$	< 10		
		1.62	$\frac{3}{2}^-$	< 10		
4.15	$(\frac{7}{2}^-)$	1.51	$\frac{5}{2}^-$	principal		(1967OL05)

^a τ_m for this state; Γ_γ (total).

^b See also (1967OL05).

^c $B(E1) = (1.06 \pm 0.05) \times 10^{-3}$ W.u. (1970BH02). See also (1969NI09).

4. $^{16}\text{O}(\alpha, n)^{19}\text{Ne}$ $Q_m = -12.135$

At $E_\alpha = 15$ to 19 MeV, lifetime measurements have been obtained for $^{19}\text{Ne}^*(1.51, 1.54, 1.62, 2.79)$: see Table 19.23 (1970GI09). See also (1964VA23).

5. $^{16}\text{O}(^6\text{Li}, t)^{19}\text{Ne}$ $Q_m = -7.350$

This reaction (and its mirror reaction $^{16}\text{O}(^6\text{Li}, ^3\text{He})^{19}\text{F}$) have been studied at $E(^6\text{Li}) = 24$ MeV: see reaction 12 in ^{19}F and Table 19.11 for a summary of the results (1971BI06). See also (1971GA1H, 1971GO1B).

6. $^{17}\text{O}(^3\text{He}, n)^{19}\text{Ne}$ $Q_m = 4.300$

At $E(^3\text{He}) = 3.1$ MeV, neutron groups have been observed corresponding to states at $E_x = 0.24 \pm 0.07, 1.55 \pm 0.06, 2.78 \pm 0.05, (3.70 \pm 0.04), 4.01 \pm 0.02, 4.13 \pm 0.03, 4.36 \pm 0.03, 4.54 \pm 0.03, 4.61 \pm 0.04, 4.69 \pm 0.03$ and 5.09 ± 0.03 MeV (1968GU07). Angular distributions have been obtained for n_{0+1+2}, n_{3+4+5} and n_6 . The latter suggest $L = 2$ and $J^\pi = (\frac{1}{2}, \frac{3}{2}, \frac{7}{2}, \frac{9}{2})^+$ for $^{19}\text{Ne}^*(2.79)$ (1968GU07).

7. $^{19}\text{F}(p, n)^{19}\text{Ne}$ $Q_m = -4.021$

$E_{\text{thresh.}} = 4233.2 \pm 2.0$ keV (1961BE13: see also (1966MA60)), 4234.7 ± 1.0 keV (1961RY04), 4234.3 ± 0.8 keV (1966MA60), 4233.7 ± 0.7 keV (1969OV01). See also (1959BR06) and (1959AJ76).

Excited states of ^{19}Ne determined from neutron threshold measurements and the measurement of neutron and γ spectra are displayed in Table 19.24 (1955MA84, 1962FR09, 1963GI09, 1965WE05, 1970GI09). The radiative decay of the first six states has been studied by (1970GI09) who also studied the lifetimes and J^π of $^{19}\text{Ne}^*(1.51, 1.54, 1.62, 2.79)$: see Table 19.23. See also (1957BA09). The g-factor of $^{19}\text{Ne}^*(0.24) = -0.296 \pm 0.003$ (1969BL02). An angular distribution measurement is reported at $E_p = 6.8$ MeV (1967DR08; $n_{0+1+2+3?}$). See also (1964ST1C). Astrophysical considerations are discussed in (1969BA1N). See also (1962EA02).

8. $^{19}\text{F}(^3\text{He}, t)^{19}\text{Ne}$ $Q_m = -3.257$

At $E(^3\text{He}) = 26$ MeV, angular distributions of the triton groups to $^{19}\text{Ne}^*(0, 0.24, 1.54, 2.79)$ have been obtained by (1970SC05); those to $^{19}\text{Ne}^*(0, 0.24)$ appear to proceed primarily via $L = 0$ and $L = 2$, respectively.

Table 19.24: Excited states of ^{19}Ne from $^{19}\text{F}(p, n)^{19}\text{Ne}$

E_x (MeV \pm keV)					J^π ^a
(1955MA84)	(1962FR09)	(1963GI09)	(1965WE05)	(1970GI09)	
0.241 \pm 4		0.241 \pm 4		0.2382 \pm 0.2	
			0.25 \pm 20		
0.280 \pm 4		0.271 \pm 4		0.2751 \pm 0.2	
	1.506 \pm 5			1.5079 \pm 0.4	$\frac{5}{2}^-$
			1.51 \pm 20		
	1.538 \pm 4			1.5363 \pm 0.5	$\frac{3}{2}^+$
	1.612 \pm 5			1.6155 \pm 0.7	$\frac{3}{2}^-$
			2.78 \pm 30	2.7946 \pm 1.5	$\frac{9}{2}^+$

^a (1970GI09).

9. $^{20}\text{Ne}(\gamma, n)^{19}\text{Ne}$ $Q_m = -16.866$

See ^{20}Ne .

10. $^{20}\text{Ne}(p, d)^{19}\text{Ne}$ $Q_m = -14.641$

See (1970PR1B).

11. $^{20}\text{Ne}(d, t)^{19}\text{Ne}$ $Q_m = -10.608$

Not reported.

12. $^{20}\text{Ne}(^3\text{He}, \alpha)^{19}\text{Ne}$ $Q_m = 3.712$
 $Q_0 = 3.750 \pm 0.013$ (1967SP09).

Alpha groups have been observed to many states of ^{19}Ne with $0 < E_x < 9.7$ MeV: see Table 19.25 (1967GR04, 1970GA18, 1972HA03). Angular distributions have been measured at $E(^3\text{He}) = 10$ MeV (1969BA62; α_0, α_6), 15 MeV (1969BA62; α_6), 15 MeV (1970GA18; α to $0 < E_x < 6.87$), 18 MeV (1972HA03) and 35 MeV (1971AR1E). DWBA analysis of the strongest

transitions leads to the l and J^π values shown in Table 19.25. Relative spectroscopic factors were also extracted. $^{19}\text{Ne}^*(0, 0.24, 1.54, 2.79)$ are identified as members of the $K = \frac{1}{2}^+$ rotational band [with $^{19}\text{Ne}^*(4.38)$ as the $\frac{7}{2}^+$ member] and $^{19}\text{Ne}^*(0.28, 1.51, 1.62)$ with the $K = \frac{1}{2}^-$ band. Candidates for the $\frac{7}{2}^-$ and $\frac{9}{2}^-$ members of the $K = \frac{1}{2}^-$ band are thought to be $^{19}\text{Ne}^*(4.15, 4.20)$. Possible matching of other ^{19}Ne states with those in ^{19}F is also discussed (1970GA18).

Table 19.25: ^{19}Ne levels from $^{20}\text{Ne}(^3\text{He}, \alpha)^{19}\text{Ne}$

E_x (MeV \pm keV)				l_n^e	J^π^f
(1967GR04) ^a	(1970GA18) ^a	(1967OL05) ^b	(1972HA03) ^{a,c}		
	0			0	$\frac{1}{2}^+$
	0.238 \pm 10	0.2384 \pm 0.3 ^d	0.2397 \pm 2 ^c	2	$\frac{5}{2}^+$
	0.273 \pm 10	0.2748 \pm 0.3 ^d	0.2766 \pm 2 ^c	1	$\frac{1}{2}^-$
		h	1.5040 \pm 3 ^c		$(\frac{5}{2}^-)$
1.521 \pm 20	1.524 \pm 20				
		h	1.5324 \pm 3 ^c	2	$(\frac{3}{2})^+$
1.620 \pm 15	1.615 \pm 10	h	1.6115 \pm 3 ^c	1	$(\frac{3}{2})^-$
2.778 \pm 15	2.793 \pm 10	h	2.7917 \pm 3	4, 5 ^g	$(\frac{9}{2})^+^g$
(3.841 \pm 15)					
4.013 \pm 15	4.036 \pm 10			2	$(\frac{3}{2}, \frac{5}{2})^+$
4.152 \pm 15	4.142 \pm 10	4.160 \pm 20.0			
	4.200 \pm 10				
4.344 \pm 15	4.379 \pm 10				
4.547 \pm 15	4.551 \pm 10			1	$(\frac{1}{2}, \frac{3}{2})^-$
	4.625 \pm 10				
4.689 \pm 15					
	4.712 \pm 10				
	4.783 \pm 20				
5.077 \pm 15	5.093 \pm 10		5.086 \pm 10		
	5.351 \pm 10			0	$\frac{1}{2}^+$
	5.426 \pm 10		5.423 \pm 10		
	5.463 \pm 20				
	5.545 \pm 10		5.517 \pm 20		
	5.831 \pm 10		5.837 \pm 20		
	6.012 \pm 10		6.014 \pm 10	1	$(\frac{3}{2}, \frac{1}{2})^-$

Table 19.25: ^{19}Ne levels from $^{20}\text{Ne}(^3\text{He}, \alpha)^{19}\text{Ne}$ (continued)

E_x (MeV \pm keV)				l_n^e	J^π^f
(1967GR04) ^a	(1970GA18) ^a	(1967OL05) ^b	(1972HA03) ^{a,c}		
	6.089 \pm 10		6.104 \pm 15	1	$(\frac{3}{2}, \frac{1}{2})^-$
	6.149 \pm 20				
	6.290 \pm 10		6.289 \pm 10		
	6.433 \pm 20		6.438 \pm 10		
	6.744 \pm 10		6.741 \pm 10		
	6.866 \pm 10		6.858 \pm 10		
	7.064 \pm 20		7.068 \pm 10		
			(7.178 \pm 15)		
			7.253 \pm 10		
			(7.326 \pm 15)		
			(7.531 \pm 15)		
			7.614 \pm 20		
			7.700 \pm 10		
			(7.788 \pm 10)		
			7.994 \pm 15		
			8.063 \pm 15		
			8.236 \pm 10 ⁱ		
			8.440 \pm 10		
			8.523 \pm 10		
			(8.810 \pm 25)		
			8.915 \pm 10		
			9.013 \pm 10		
			9.100 \pm 20		
			9.240 \pm 20		
			9.489 \pm 25		
			9.886 \pm 50 ⁱ		
			10.407 \pm 30 ⁱ		
			10.613 \pm 20		

- ^a From measurements of α -groups.
^b From measurements of de-excitation γ -rays.
^c The energy separations within each multiplet are fixed at the values determined by (1968GU07) (1972HA03).
^d (1970BH02) report $E_x = 238.34 \pm 0.15$ and 275.30 ± 0.2 keV for these states.
^e (1970GA18). Spectroscopic factors have also been calculated.
^f (1967OL05, 1970GA18). See also (1971AR1E).
^g (1969BA62).
^h The excitation energies obtained by (1967OL05) for $^{19}\text{Ne}^*(1.51, 1.54, 1.62, 2.79)$ are superseded by those of (1970GI09): see Table 19.24.
ⁱ Unresolved states.

The mean lifetimes, τ_m , of $^{19}\text{Ne}^*(0.24, 0.28)$ are, respectively, 26.6 ± 1.2 nsec (1967BE14) and 61.4 ± 3.0 psec (1970BH02). $^{19}\text{Ne}^*(0.28)$ decays to the ground state by E1 emission: $B(\text{E1}) = (1.06 \pm 0.05) \times 10^{-3}$ W.u. The ratio of this value to that for the analog state in ^{19}F is 0.95 ± 0.06 (1970BH02). The γ -decay of seven states of ^{19}Ne has been studied by (1967OL05): see Table 19.23.



At $E_p = 45$ MeV triton groups are observed corresponding to $^{19}\text{Ne}^*(0, 0.24, 2.79, 4.03, 7.620 \pm 0.025)$. The latter has an angular distribution [$L = 0$] similar to that for $^{19}\text{F}^*(7.66)$: both are thought to be the analogs of the $J^\pi = \frac{3}{2}^+$; $T = \frac{3}{2}$, 0.096 MeV first excited state of ^{19}O . The ground state of ^{19}O has $J^\pi = \frac{5}{2}^+$: L for the tritons to that state should be 2 (1969HA38).

^{19}Na
(Fig. 8)

A study of the reaction $^{24}\text{Mg}(\text{p}, ^6\text{He})^{19}\text{Na}$ at $E_p = 54.7$ MeV reveals a group of ^6He particles corresponding to a state in ^{19}Na with $M - A = 12.974 \pm 0.070$ MeV. It is presumed to be the ground state of ^{19}Na , although the close proximity of the second $T = \frac{3}{2}$ state in ^{19}O from the first (96 keV), does not permit a definite assignment. If it is assumed that $^{19}\text{Na}(0)$ has $M - A = 12.974 \pm 0.070$ MeV, then ^{19}Na is unbound with respect to decay into $^{18}\text{Ne} + \text{p}$ by 0.366 ± 0.070 MeV (1969CE01). The mass excess of ^{19}Na is calculated to be 12.965 ± 0.025 MeV, using a seniority scheme, and 12.968 ± 0.025 MeV, using a supermultiplet scheme [both these values were calculated assuming that the lowest $J^\pi = \frac{3}{2}^+$; $T = \frac{3}{2}$ state in ^{19}Na is 0.096 MeV above the $J^\pi = \frac{5}{2}^+$; $T = \frac{3}{2}$ state, as in ^{19}O] (1969HA38). See also (1960BA1F, 1966KE16, 1968MU1B, 1971CE1A).

^{19}Mg
(Not illustrated)

See (1965GO1D).

References

(Closed 31 December 1971)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1950HO1A W.F. Hornyak, T. Lauritsen, P. Morrison and W.A. Fowler, *Rev. Mod. Phys.* 22 (1950) 291
- 1952MI54 C. Mileikowsky and W. Whaling, *Phys. Rev.* 88 (1952) 1254
- 1952SC15 G. Schrank and J.R. Richardson, *Phys. Rev.* 86 (1952) 248
- 1954JO21 G.A. Jones, W.R. Phillips, C.M.P. Johnson and D.H. Wilkinson, *Phys. Rev.* 96 (1954) 547
- 1955JA1A Jarmie, *Phys. Rev.* 98 (1955) 41
- 1955MA84 J.B. Marion, T.W. Bonner and C.F. Cook, *Phys. Rev.* 100 (1955) 91
- 1956AL1D D.G. Alkhazov, D.S. Andreyev, A.P. Greenberg and I.K. Lemberg, *Physica* 22 (1956) 1129A
- 1956AL36 D.G. Alkhazov, D.S. Andreyev, A.P. Greenberg and I.K. Lemberg, *Nucl. Phys.* 2 (1956) 65
- 1956AL55 D.G. Alkhazov, D.S. Andreyev, A.P. Greenberg and I.K. Lemberg, *Zh. Eksp. Teor. Fiz.* 30 (1956) 809; *JETP (Sov. Phys.)* 3 (1957) 964
- 1956HA1A Harlow, Marion, Chapman and Bonner, *Phys. Rev.* 101 (1956) 214
- 1956HI1A Hill and Blair, *Phys. Rev.* 104 (1956) 198
- 1956JO35 C.M.P. Johnson, *Phil. Mag.* 1 (1956) 573
- 1956MA18 H. Mark and C. Goodman, *Phys. Rev.* 101 (1956) 768
- 1956SQ1A Squires, Bockelman and Buechner, *Phys. Rev.* 104 (1956) 413
- 1957AL29 W.P. Alford and D.R. Hamilton, *Phys. Rev.* 105 (1957) 673
- 1957BA09 R. Barloutaud, P. Lehmann, A. Leveque, G.C. Phillips and J. Quidort, *Compt. Rend.* 245 (1957) 422
- 1957PE12 J.R. Penning and F.H. Schmidt, *Phys. Rev.* 105 (1957) 647
- 1957PR1A Price, *Proc. Phys. Soc. (London)* A70 (1957) 661
- 1957SJ66 B. Sjogren, *Ark. Fys.* 12 (1957) 537
- 1957WI1E Winterberg, *Z. Naturforsch.* A12 (1957) 271

- 1958BU12 J.W. Butler and H.D. Holmgren, Phys. Rev. 112 (1958) 461
- 1958CA97 J. Catala, L. Miralles, J. Aguilar and M. de la Cuadra, An. Real. Soc. Espan. Fis. y Quim. A54 (1958) 255
- 1958GO71 C.D. Goodman and J.L. Need, Phys. Rev. 110 (1958) 676
- 1958SP1A Spicer, Aust. J. Phys. 11 (1958) 490
- 1958SU58 K. Sugimoto, A. Mizobuchi and H. Yamamoto, J. Phys. Soc. Jpn. 13 (1958) 1548
- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1959AL06 D.E. Alburger, A. Gallmann and D.H. Wilkinson, Phys. Rev. 116 (1959) 939
- 1959AL10 J.S. Allen, R.L. Burman, W.B. Hermannsfeldt, P. Stahelm and T.H. Braid, Phys. Rev. 116 (1959) 134
- 1959AL1E Allen, Rev. Mod. Phys. 31 (1959) 791
- 1959BI1A Bilman, J. Phys. Rad. 20 (1959) 51
- 1959BR06 D.A. Bromley, A.J. Ferguson, H.E. Gove, J.A. Kuehner, A.E. Litherland, E. Almqvist and R. Batchelor, Can. J. Phys. 37 (1959) 1514
- 1959BR1E Brink and Kerman, Nucl. Phys. 12 (1959) 314
- 1959BR72 D.A. Bromley, J.A. Kuehner and E. Almqvist, Phys. Rev. 115 (1959) 586
- 1959BR79 D.A. Bromley, J.A. Kuehner and E. Almqvist, Nucl. Phys. 13 (1959) 1
- 1959BU05 J.W. Butler and J.D. Holmgren, Phys. Rev. 116 (1959) 1485
- 1959BU81 J.W. Butler, L.W. Fagg and H.D. Holmgren, Phys. Rev. 113 (1959) 268
- 1959FU1A Fujii, Prog. Theor. Phys. 21 (1959) 511
- 1959HI73 S. Hinds and R. Middleton, Proc. Phys. Soc. (London) A74 (1959) 775
- 1959JO26 C.M.P. Johnson, G.A. Jones, W.R. Phillips and D.H. Wilkinson, Proc. Roy. Soc. A252 (1959) 1
- 1959JO32 R.L. Johnston, H.D. Holmgren and G.D. Gutsche, Bull. Amer. Phys. Soc. 4 (1959) 403, C4
- 1959KO60 E. Kondaiah, C. Badrinathan and K.V.K. Iyengar, Nucl. Phys. 9 (1959) 561
- 1959LA1A Lane, Cong. Int. de Phys. Nucl., 1958, Dundo, Paris (1959) 177
- 1959LI55 C.S. Littlejohn and G.C. Morrison, Phys. Rev. 116 (1959) 1526
- 1959SC30 L. Schellenberg, E. Baumgartner, P. Huber and F. Seiler, Helv. Phys. Acta 32 (1959) 357
- 1959SM02 H.H. Smotrich, Thesis, Columbia Univ. (1959); CU-188 (1959)
- 1959TR1A Trail and Raboy, Bull. Amer. Phys. Soc. 4 (1959) 220
- 1959WI1B Wildermuth and Kanellopoulous, Congres. Int. de Phys. Nucl., 1958(Dunod, Paris, 1959) 793

- 1959WI39 W. Williams, Jr. and P.V.C. Hough, Bull. Amer. Phys. Soc. 4 (1959) 219, AB11
- 1959ZI16 W. Zimmermann, Jr., Phys. Rev. 114 (1959) 837
- 1960AG01 J. Aguilar, W.E. Burcham, J. Catala, J.B.A. England, J.S.C. McKee and J. Rotblat, Proc. Roy. Soc. A254 (1960) 395
- 1960BA1F Baz, Sov. J. At. Energy 6 (1960) 422
- 1960BL05 J.M. Blair and J.J. Leigh, Phys. Rev. 118 (1960) 495
- 1960BO23 E.C. Booth, Nucl. Phys. 19 (1960) 426
- 1960CL02 R.L. Clarke, E. Almqvist and E.B. Paul, Nucl. Phys. 14 (1960) 472
- 1960DA08 R.B. Day and M. Walt, Phys. Rev. 117 (1960) 1330
- 1960FO10 B. Forkman and I. Wahlstrom, Ark. Fys. 18 (1960) 339
- 1960GE06 K.N. Geller, J. Halpern and E.G. Muirhead, Phys. Rev. 119 (1960) 716
- 1960GO1C Gove, Proc. Int. Conf. on Nucl. Struct., Kingston, Canada; Eds. D.A. Bromley and E.W. Vogt (Univ. of Toronto Press, 1960) 438
- 1960JA12 J. Janecke, Z. Naturforsch. A15 (1960) 593
- 1960KI06 J.D. King, R.N.H. Haslam and W.J. McDonald, Can. J. Phys. 38 (1960) 1069
- 1960MI1A Milman, Amsel and Loyau, J. Phys. Rad. 20 (1959) 51
- 1960RA1A Raz, Phys. Rev. 120 (1960) 169
- 1960RE05 K. Reibel and A.K. Mann, Phys. Rev. 118 (1960) 701
- 1960RO1C Roth and Wildermuth, Nucl. Phys. 20 (1960) 10
- 1960SA09 D. Sadeh, Compt. Rend. 250 (1960) 1632
- 1960SH1A Sheline and Wildermuth, Nucl. Phys. 21 (1960) 196
- 1960ST17 P.H. Stelson and F.K. McGowan, Nucl. Phys. 16 (1960) 92
- 1960TA1C Talmi and Unna, Ann. Rev. Nucl. Sci. 10 (1960) 353
- 1960VE06 G.E. Velyukhov, A.N. Prokofev and S.V. Starodubtsev, Zh. Skep. Teor. Fiz. 39 (1960) 563; JETP (Sov. Phys.) 12 (1961) 395
- 1960WA04 R. Wallace and J.A. Welch, Jr., Phys. Rev. 117 (1960) 1297
- 1960ZE03 Ya.B. Zeldovich, Zh. Eksp. Teor. Fiz. 38 (1960) 1123; JETP (Sov. Phys.) 11 (1960) 812
- 1961AM01 G. Amsel and G.R. Bishop, Phys. Rev. 123 (1961) 957
- 1961AN1A Androsenko, Broder and Lashuk, Sov. J. At. Energy 9 (1961) 945
- 1961AR06 J.C. Armstrong and K.S. Quisenberry, Phys. Rev. 122 (1961) 150
- 1961BA1C Baz, Goldanskii and Zeldovich, Sov. Phys. Uspekhi 3 (1961) 729

- 1961BA1E Balashov, Neudachin and Smirnov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 25 (1961) 170; *Bull. Acad. Sci. USSR Phys.* 25 (1961) 165
- 1961BE13 E.H. Beckner, R.L. Bramblett, G.C. Phillips and T.A. Eastwood, *Phys. Rev.* 123 (1961) 2100
- 1961CA02 R.R. Carlson, C.C. Kim, J.A. Jacobs and A.C.L. Barnard, *Phys. Rev.* 122 (1961) 607
- 1961CA19 J. Catala, J. Aguilar, A. Garcia, G. Gil and J. De la Rubia, *An. Real. Soc. Espan. Fis. y Quim. (Madrid)* A57 (1961) 201; *Phys. Abs.* 5965 (1962)
- 1961DE1A Del Bianco, Thesis, Univ. of Pennsylvania (1961)
- 1961DO08 W.R. Dodge, Thesis, Stanford Univ. (1961); HEPL-246 (1961); *Nucl. Sci. Abstr.* 16, 750, Abstr. 5921 (1962)
- 1961DU02 K.L. Dunning and J.W. Butler, *Phys. Rev.* 123 (1961) 1321
- 1961FR07 R.M. Freeman, *Nucl. Phys.* 26 (1961) 446
- 1961JA23 A. Jaidar, G. Lopez, M. Mazari and R. Dominguez, *Rev. Mex. Fisica* 10 (1961) 247
- 1961JO03 J.V. Jovanovich, E.D. Lambe, T.A. Pond and J.B. Reynolds, *Bull. Amer. Phys. Soc.* 6 (1961) 50, QA5
- 1961KO06 E. Kondaiah, M.L. Jhingan and C. Badrinathan, *Nucl. Phys.* 27 (1961) 166
- 1961KO08 J. Kokame, *J. Phys. Soc. Jpn.* 16 (1961) 2101
- 1961LO10 G. Lopez and O. Almen, *Rev. Mex. Fis.* 10 (1961) 239
- 1961LO14 K. Lorenzen and D. Konig, *Z. Naturforsch.* A16 (1961) 933
- 1961NO05 E. Norbeck, *Phys. Rev.* 121 (1961) 824
- 1961OK1A Okada, *Nucleonics* 19 (1961) 79
- 1961PU1A Pugh and Riley, *Proc. Rutherford Jub. Int. Conf., Manchester, England*; Ed. J.B. Birks (Academic Press Inc., New York, 1961) 195
- 1961RO01 W.L. Rogers, D.I. Garber and E.F. Shrader, *Bull. Amer. Phys. Soc.* 6 (1961) 61, TA7
- 1961RY04 A. Rytz, H. Winkler, F. Zamboni and W. Zych, *Helv. Phys. Acta* 34 (1961) 819
- 1961SI03 M.G. Silbert and N. Jarmie, *Phys. Rev.* 123 (1961) 221
- 1961SI09 E.A. Silverstein, S.R. Salisbury, G. Hardie and L.D. Oppliger, *Phys. Rev.* 124 (1961) 868
- 1961SJ01 B. Sjogren, G. Wickenberg and N. Johansson, *Ark. Fys.* 20 (1961) 117
- 1961SM02 H. Smotrich, K.W. Jones, L.C. McDermott and R.E. Benenson, *Phys. Rev.* 122 (1961) 232
- 1961TO03 J.H. Towle and B.E.F. Macefield, *Proc. Phys. Soc. (London)* 77 (1961) 399
- 1961TR1B True and Warburton, *Nucl. Phys.* 22 (1961) 426

- 1961VA28 S.S. Vasilev, E.A. Romanovskii and G.F. Timushev, Zh. Skep. Teor. Fiz. 41 (1961) 1040; JETP (Sov. Phys.) 14 (1962) 741
- 1961YA02 H. Yamaguchi, J. Phys. Soc. Jpn. 16 (1961) 583
- 1962BH09 K.H. Bhatt, Nucl. Phys. 39 (1962) 375
- 1962BO17 E.C. Booth and K.A. Wright, Nucl. Phys. 35 (1962) 472
- 1962BO1A Boehm, Dumond, Hendrikson and Seppi, Electromag. Lifetimes Prop. of Nucl. States, N.A.S.-N.R.C. Publ. 974 (1962) 106
- 1962BR1E Braun, Z. Phys. 166 (1962) 62
- 1962DA1B Davidson and Chi, in Electromag. Lifetimes and Properties of Nucl. States; N.A.S.-N.R.C. Publ. 974 (1962) 237
- 1962DE03 W.E. Del Bianco and W.E. Stephens, Phys. Rev. 126 (1962) 709
- 1962DO1A Dodge and Barber, Phys. Rev. 127 (1962) 1746
- 1962EA02 L.G. Earwaker, J.G. Jenkin and E.W. Titterton, Nature 195 (1962) 271
- 1962FR09 J.M. Freeman and D. West, Nucl. Phys. 38 (1962) 89
- 1962GA17 A. Garcia, J. Kirk, J.B.A. England and P.E. Hodgson, Nucl. Phys. 38 (1962) 372
- 1962GO1B Golsanskii, Dokl. Akad. Nauk SSSR 146 (1962) 1309
- 1962HO06 R.K. Hobbie and F.F. Forbes, Phys. Rev. 126 (1962) 2137
- 1962HO1C Honda and Ui, Nucl. Phys. 34 (1962) 609
- 1962MA38 S. Malmskog and J. Konijn, Nucl. Phys. 38 (1962) 196
- 1962NE03 J.W. Nelson and E.L. Hudspeth, Phys. Rev. 125 (1962) 301
- 1962NU1A Nung Sung, J. Phys. Soc. Jpn. 17 (1962) 1811
- 1962RI09 R.C. Ritter, P.H. Stelson, F.K. McGowan and R.L. Robinson, Phys. Rev. 128 (1962) 2320
- 1962SE12 E.J. Seppi and F. Boehm, Phys. Rev. 128 (1962) 2334
- 1962SH1B Shah, Phys. Lett. 1 (1962) 261
- 1962SO01 P.V. Sorokin, A.I. Popov, V.E. Storizhko and A.Y. Taranov, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1962) 1084; Columbia Tech. Transl. 26 (1963) 1091
- 1962TA1B Talmi and Unna, Nucl. Phys. 30 (1962) 280
- 1962VA28 S.S. Vasilev, E.A. Romanovskii and G.F. Timushev, Izv. Akad. Nauk SSSR Ser. Fiz. 26 (1962) 1508; Columbia Tech. Transl. 26 (1963) 1533
- 1962YA03 K. Yagi, K. Katori, H. Ohnuma, Y. Hashimoto and Y. Nogami, J. Phys. Soc. Jpn. 17 (1962) 595
- 1962YA1B Yagi, J. Phys. Soc. Jpn. 17 (1962) 604
- 1963AM1A Amsel, Thesis, Univ. of Paris (1963)

- 1963BA19 W.C. Barber, J. Goldemberg, G.A. Peterson and Y. Torizuka, Nucl. Phys. 41 (1963) 461; Erratum Nucl. Phys. 47 (1963) 527
- 1963BE03 D. Beder, Phys. Lett. 3 (1963) 206
- 1963BE05 R.E. Benenson and B. Yaramis, Phys. Rev. 129 (1963) 720
- 1963BE36 N. Bessis, H. Lefebvre-Brion and C.M. Moser, Rev. Mod. Phys. 35 (1963) 548
- 1963CA1B Casabella and Gicking, Bull. Amer. Phys. Soc. 8 (1963) 345
- 1963CH02 B.E. Chi and J.P. Davidson, Phys. Rev. 131 (1963) 366
- 1963CO22 E.D. Commins and D.A. Dobson, Phys. Rev. Lett. 10 (1963) 347
- 1963DO1C Dodd and McCarthy, Padua (1963) 309
- 1963FI07 P.S. Fisher, D.F. Measday, F.A. Nikolaev, A. Kalmykov and A.B. Clegg, Nucl. Phys. 45 (1963) 113
- 1963FR12 R. Freeman, J.M.F. Jeronimo, G.S. Mani and A. Sadeghi, J. Phys. 24 (1963) 894
- 1963GA14 N.H. Gale and J.M. Calvert, Phys. Lett. 7 (1963) 348
- 1963GI09 W.W. Givens, R.C. Bearnse, G.C. Phillips and A.A. Rollefson, Nucl. Phys. 43 (1963) 553
- 1963GI12 W.W. Givens, R.C. Bearnse, G.C. Phillips and A.A. Rollefson, Nucl. Phys. 46 (1963) 519
- 1963GO04 J. Goldemberg and Y. Torizuka, Phys. Rev. 129 (1963) 312
- 1963GO08 S. Gorodetzky, M. Port, J. Graff and J.M. Thirion, Nucl. Phys. 42 (1963) 462
- 1963HA05 M. Harvey, Phys. Lett. 3 (1963) 209
- 1963HA1C Harvey et al., Padua (1963) 974
- 1963HU07 K.K. Huang, K. Yagi, T. Awaya, H. Ohnuma, M. Fujioka and Y. Nogami, J. Phys. Soc. Jpn. 18 (1963) 646
- 1963KA25 K.V. Karadzhev, V.I. Manko and F.E. Chukreev, Zh. Eksp. Teor. Fiz. 44 (1963) 870; JETP (Sov. Phys.) 17 (1963) 593
- 1963LA02 R.A. Lasalle, R.D. Benta and J.G. Cramer, Jr., Bull. Amer. Phys. Soc. 8 (1963) 303, D8
- 1963LI01 A.E. Litherland, M.A. Clark and C. Broude, Phys. Lett. 3 (1963) 204
- 1963MA1D Matthies, Neudachin and Smirnov, Nucl. Phys. 49 (1963) 97
- 1963MU08 K.M. Murray and W.L. Bendel, Phys. Rev. 132 (1963) 1134
- 1963PA03 S.P. Pandya, Nucl. Phys. 43 (1963) 636
- 1963RO12 J.R. Rook, Nucl. Phys. 41 (1963) 343
- 1963VA01 F.M. Vazquez, C. Taguena and V.C. de la Pena, Bull. Amer. Phys. Soc. 8 (1963) 126, U14

- 1963VA12 M. Vazquez, C. Taguena and V.C. de la Pena, *Rev. Mex. Fis.* 12 (1963) 27
- 1963YA03 K. Yagi, Y. Nakajima, K. Katori, Y. Awaya and M. Fujioka, *Nucl. Phys.* 41 (1963) 584
- 1964AM02 S. Amiel and J. Gilat, *Nucl. Sci. Eng.* 18 (1964) 105
- 1964AM1A Amsel, *Ann. Phys.* 9 (1964) 297
- 1964BA11 M.R. Baker, C.H. Anderson and N.F. Ramsey, *Phys. Rev.* 133 (1964) A1533
- 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, *Nucl. Phys.* 53 (1964) 209
- 1964BAZZ K.A. Baskova, S.S. Vasilyev and L.Y. Shavtvalov, *Compt. Rend. Congr. Int. Phys. Nucl., Paris, Vol. II* (1964) 523
- 1964BO22 E.C. Booth, B. Chasan and K.A. Wright, *Nucl. Phys.* 57 (1964) 403
- 1964BO37 R.V. Boeckh, G. Graff and R. Ley, *Z. Phys.* 179 (1964) 285
- 1964BO40 E. Bodenstedt and J.D. Rogers, *Perturbed Angular Correlations*; Eds. E. Karlsson, E. Matthias and K. Siegbahn (Amsterdam, North-Holland Publ. Company, 1964) p. 91
- 1964BR1G Brill, Chuev and Ogloblin, Paris (1964) 981
- 1964BR1H Brown, Paris (1964) 129
- 1964CO24 S. Cohen, R.D. Lawson, M.H. Macfarlane and M. Soga, *Phys. Lett.* 9 (1964) 180
- 1964DO08 T.R. Donoghue, A.F. Behof and S.E. Darden, *Nucl. Phys.* 54 (1964) 33
- 1964DR1A Dreizler, *Phys. Rev.* 136 (1964) B321
- 1964EC03 S.F. Eccles, H.F. Lutz and J. Stevens, *Bull. Amer. Phys. Soc.* 9 (1964) 704, B9
- 1964EL1A El Bedewi, Fawzi and Rizk, Paris (1964) 432
- 1964EX1A Extermann, Baumgartner and Huber, *Helv. Phys. Acta* 37 (1964) 505
- 1964FR03 R.M. Freeman and G.S. Mani, *Nucl. Phys.* 51 (1964) 593
- 1964GA1A Gardner and Yu, *Nucl. Phys.* 60 (1964) 49
- 1964GO1C Gottschalk, Strauch and Wang, Paris (1964) 324
- 1964HA21 M. Harvey, *Nucl. Phys.* 52 (1964) 542
- 1964IN03 T. Inoue, T. Sebe, H. Hagiwara and A. Arima, *Nucl. Phys.* 59 (1964) 1
- 1964KA1A Kawai, Terasawa and Izumo, *Nucl. Phys.* 59 (1964) 289
- 1964KA1B Karadzhev, Manko and Chukreev, *Zh. Eksp. Teor. Fiz.* 47 (1964) 1185; *JETP (Sov. Phys.)* 20 (1965) 800
- 1964LI14 I. Lindgren, *Perturbed Angular Correlations*; Eds. E. Karlsson, E. Matthias and K. Siegbahn (Amsterdam, North-Holland Publ. Co., 1964) 379
- 1964LI1B Lindskog, Sundstrom and Sparrman, *Perturbed Angular Correlations* (Amsterdam, North-Holland Publ. Company, 1964) 411

- 1964MA12 J.D. Mahoney and S.S. Markowitz, *J. Inorg. Nucl. Chem.* 26 (1964) 907
- 1964MA1G Mamasakhlisov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 28 (1964) 1550
- 1964MA57 M. Mazari, A. Jaidar, G. Lopez, A. Tejera, J. Careaga, R. Dominguez and F. Alba, *Proc. 2nd Int. Conf. on Nucl. Masses*, Vienna, Austria, 1963; Ed. W.H. Johnson, Jr. (Springer Verlag, Vienna, 1964) 305
- 1964MO23 J.H. Montague, R.E. Lee and E.B. Paul, *Bull. Amer. Phys. Soc.* 9 (1964) 705, B13
- 1964MO24 R. Moreh and A.A. Jaffe, *Proc. Phys. Soc. (London)* 84 (1964) 330
- 1964MO25 R. Moreh, T. Daniels and A.A. Jaffe, *Proc. Phys. Soc. (London)* 84 (1964) 332; Erratum *Proc. Phys. Soc. (London)* 84 (1964) 835
- 1964NE08 D. Newton, A.B. Clegg and G.L. Salmon, *Nucl. Phys.* 55 (1964) 353
- 1964PU01 D.J. Pullen, J.R. Rook and R. Middleton, *Nucl. Phys.* 51 (1964) 88
- 1964SC01 J. Schaitman, H.F. Lutz and S.F. Eccles, *Bull. Amer. Phys. Soc.* 9 (1964) 68, GC10
- 1964SE09 S. Seki, M. Yamanouchi and M. Miwa, *J. Phys. Soc. Jpn.* 19 (1964) 1999
- 1964ST1B Stovall, *Phys. Rev.* 133 (1964) B268
- 1964ST1C Stromberg, Wiedling and Holmqvist, *AE* 139 (1964)
- 1964ST1D Stevens, UCRL 7648 (1964)
- 1964ST1E Storizhko, NP 14092 (1964)
- 1964SU01 K. Sugimoto, A. Mizobuchi and K. Nakai, *Phys. Rev.* 134 (1964) B539
- 1964TE04 G. Tessler and W.E. Stephens, *Phys. Rev.* 135 (1964) B129
- 1964VA02 F.J. Vaughn, L.F. Chase, Jr., H.A. Grench and J.W. Hilson, *Bull. Amer. Phys. Soc.* 9 (1964) 20, BC10; Oral Report
- 1964VA23 S.S. Vasilev, E.T. George and L.Y. Shavtvalov, *Zh. Eksp. Teor. Fiz.* 47 (1964) 1164; *JETP (Sov. Phys.)* 20 (1965) 783
- 1964WI05 G. Wickenberg, S. Hjorth, N.G.E. Johansson and B. Sjogren, *Ark. Fys.* 25 (1964) 191
- 1965AL05 W.P. Alford, L.M. Blau and D. Cline, *Nucl. Phys.* 61 (1965) 368
- 1965AL13 J.P. Allen, A.J. Howard, D.A. Bromley and J.W. Olness, *Nucl. Phys.* 68 (1965) 426
- 1965AL20 J.P. Allen, A.J. Howard, D.A. Bromley and J.W. Olness, *Phys. Rev.* 140 (1965) B1245
- 1965AM1A Amos and McCarthy, *Nucl. Phys.* 65 (1965) 81
- 1965BA1J Bassichis, Giraud and Ripka, *Phys. Rev. Lett.* 15 (1965) 980
- 1965BL07 S.D. Bloom, J.D. Anderson, W.F. Hornyak and C. Wong, *Phys. Rev. Lett.* 15 (1965) 264
- 1965BO28 G. Bougnot, G. Espi, J.C. Poizat, J.J. Samueli and A. Sarazin, *J. Phys. (Paris)* 26 (1965) 546

- 1965BO42 M. Bormann, E. Fretwurst, P. Schehka, G. Wrege, H. Buttner, A. Lindner and H. Meldner, Nucl. Phys. 63 (1965) 438
- 1965BR1B Brill, Yad. Fiz. 1 (1965) 55; Sov. J. Nucl. Phys. 1 (1965) 37
- 1965DE1J Denisov, Duisebaev, Kosareva and Cherenkov, Yad. Fiz. 2 (1965) 82
- 1965DI1C Dittmer et al., Bull. Amer. Phys. Soc. 10 (1965) 122
- 1965ER04 J.R. Erskine, R.E. Holland, R.D. Lawson, M.H. Macfarlane and J.P. Schiffer, Phys. Rev. Lett. 14 (1965) 915
- 1965FA1B Fabre de La Ripelle, Prog. Theor. Phys. 33 (1965) 38
- 1965FE02 P. Federman and I. Talmi, Phys. Lett. 19 (1965) 490
- 1965FO1D Fosado and Wolf, Rev. Mex. Fisica 14 (1965) 181A
- 1965GA1D Gaponov, Yad. Fiz. 2 (1965) 1002
- 1965GE05 C. Gerardin, M. Wery, R. Seltz and D. Magnac-Valette, Compt. Rend. 261 (1965) 1512
- 1965GL04 R.N. Glover and A.D.W. Jones, Phys. Lett. 16 (1965) 69
- 1965GO1D Goldanskii, Usp. Fiz. Nauk 87 (1965) 255
- 1965GO1F Goswami and Kisslinger, Phys. Rev. 140 (1965) B26
- 1965HA31 J.C. Hardy and B. Margolis, Phys. Lett. 15 (1965) 276
- 1965HU13 R.W. Huggins and J.H. Sanders, Proc. Phys. Soc. (London) 86 (1965) 53
- 1965JI1A Jin and Harris, Bull. Amer. Phys. Soc. 10 (1965) 638, 1195
- 1965KL01 O. Klepper and H. Spehl, Nucl. Phys. 64 (1965) 393
- 1965LU1A Lutz, Eccles and Mason, Bull. Amer. Phys. Soc. 10 (1965) 122
- 1965MA1K Marion, Nucl. Phys. 68 (1965) 463
- 1965MC10 R.E. McDonald, D.B. Fossan, L.F. Chase, Jr. and J.A. Becker, Phys. Rev. 140 (1965) B1198
- 1965MO16 R. Moreh and T. Daniels, Nucl. Phys. 74 (1965) 403
- 1965MO19 R. Moreh, Nucl. Phys. 70 (1965) 293
- 1965OK1A Okai, Park and Wildermuth, Z. F. Phys. 184 (1965) 451
- 1965PR1C Pratt, Walecka and Griffy, Nucl. Phys. 64 (1965) 677
- 1965RI1A Riou, Rev. Mod. Phys. 37 (1965) 375
- 1965RO1H Rook, Nucl. Phys. 61 (1965) 219
- 1965SA1A Salisbury, Fossan and Vaughn, Nucl. Phys. 64 (1965) 343
- 1965SH1D Shoda, Nucl. Phys. 72 (1965) 305
- 1965SI18 R.H. Siemssen, L.L. Lee, Jr. and D. Cline, Phys. Rev. 140 (1965) B1258

- 1965VA03 F.J. Vaughn, H.A. Grench, W.L. Imhof, J.H. Rowland and M. Walt, Nucl. Phys. 64 (1965) 336
- 1965WE05 J.J. Wesolowski, J.D. Anderson, L.F. Hansen, C. Wong and J.W. McClure, Nucl. Phys. 71 (1965) 586
- 1965WI1B Wiza, Thesis, Univ. of Pennsylvania (1965) Unpublished
- 1965WO1C Wolf and Fosadc, Rev. Mex. Fis. 14 (1965) 180A
- 1965YO1A You-Wen, Tsung-Hua and Min, Acta Phys. Sin. 21 (1965) 542
- 1965YU1A Yu, Chang and Yu, Acta Phys. Sin. 21 (1965) 545; Phys. Abs. 990 (1966)
- 1966AF01 O.F. Afonin, A.P. Grinberg and I.K. Lemberg, Izv. Akad. Nauk SSSR Ser. Fiz. 30 (1966) 1142; Bull. Acad. Sci. USSR Phys. Ser. 30 (1967) 1193
- 1966AL12 D.E. Alburger and K.W. Jones, Phys. Rev. 149 (1966) 743
- 1966AR10 A. Armigliato, F. Brandolini, F. Pellegrini and E. Crescenti, Nuovo Cim. B45 (1966) 92
- 1966BA2E Bassichis, Giraud and Ripka, J. Phys. C1-46 (1966)
- 1966BO21 G.C. Bonazzola, E. Chiavassa and T. Bressani, Nucl. Phys. 86 (1966) 378
- 1966BR1H Brink, Struct. of Complex Nuclei, Ed. N.N. Bogolyubov, Atomizdat, 1966 (Consultants Bureau, New York, 1969) 69
- 1966DA1E Dalidchik and Sayasov, Yad. Fiz. 3 (1966) 820
- 1966DE09 L.J. Denes, W.W. Daehnick and R.M. Drisko, Phys. Rev. 148 (1966) 1097
- 1966DO1C Donau, Proc. Conf. Nucl. Reactions, Jan. 1966, Rossendorf, Ed. J. Schintlmeister, ZFK-122 (1966) 20
- 1966DO1D Dolbilkin et al., Bull. Acad. Sci. Usssr Phys. 30 (1966) 354
- 1966GA25 G.T. Garvey and I. Kelson, Phy. Rev. Lett. 16 (1966) 197
- 1966GI1A Gillet, Proc. Int. School Enrico Fermi, Course 36; Ed. C. Bloch (Academic Press, 1966) 43
- 1966GO1C Goldemberg and Pratt, Rev. Mod. Phys. 38 (1966) 311
- 1966GO1E Gorodetzky et al., Nucl. Instrum. Meth. 42 (1966) 269
- 1966HA21 R.L. Hahn and E. Ricci, Phys. Rev. 146 (1966) 650
- 1966IN01 T. Inoue, T. Sebe, H. Hagiwara and A. Arima, Nucl. Phys. 85 (1966) 184
- 1966JA1A Jacob and Maris, Rev. Mod. Phys. 38 (1966) 121
- 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
- 1966KU1D Kuehner, Nucl. Spin-Parity Assignments; Ed. Gove (Academic Press, 1966) 146
- 1966LA1E Lawson, 2nd Symp. on the Struct. of Low-Medium Mass Nuclei, Apr. 1966 (1966) 6
- 1966LU03 H.F. Lutz, J.J. Wesolowski, L.F. Hansen and S.F. Eccles, Phys. Lett. 20 (1966) 410

1966MA1R J.H. Manley and W.E. Stein, Phys. Rev. 144 (1966) 956
 1966MA43 G.M. Matous, G.H. Herling and E.A. Wolicki, Phys. Rev. 152 (1966) 908
 1966MA44 E. Maqueda, Phys. Lett. 23 (1966) 571
 1966MA60 J.B. Marion, Rev. Mod. Phys. 38 (1966) 660
 1966MI1F Migdal, Proc. Int. School Enrico Fermi, Course 36; Ed. C. Bloch (Academic Press, 1966) 171
 1966OL01 J.W. Olness and D.H. Wilkinson, Phys. Rev. 141 (1966) 966
 1966RI12 G. Ripka and L. Zamick, Phys. Lett. 23 (1966) 347
 1966RI1F Ripka, Lect. in Theor. Phys., Vol. VIII C (Univ. of Colorado Press, Boulder, 1966) 237
 1966RO01 E. Rost, Phys. Lett. 21 (1966) 87
 1966RO03 H. Ropke, Nucl. Phys. 80 (1966) 476
 1966ST04 J. Stevens, H.F. Lutz and S.F. Eccles, Nucl. Phys. 76 (1966) 129
 1966SU1D Suzuki and Nagata, Prog. Theor. Phys. 36 (1966) 37
 1966TH02 M.F. Thomas, J.S. Lopes, R.W. Ollerhead, A.R. Poletti and E.K. Warburton, Nucl. Phys. 78 (1966) 298
 1966VE1B Velkley, Brandenberger and McEllistrem, Bull. Amer. Phys. Soc. 11 (1966) 832
 1966WI05 J.L. Wiza and R. Middleton, Phys. Rev. 143 (1966) 676
 1966WO03 G. Wolfer and M. Bormann, Z. Phys. 194 (1966) 75
 1966ZA1B D.A. Zaikin, Phys. Lett. 22 (1966) 158
 1966ZA1C D.A. Zaikin, Nucl. Phys. 86 (1966) 638
 1967AM1D K.A. Amos, V.A. Madsen and I.E. McCarthy, Nucl. Phys. A94 (1967) 103
 1967AR02 A. Arima, H. Horiuchi and T. Sebe, Phys. Lett. B24 (1967) 129
 1967AU1B J. Audouze, M. Epherre and H. Reeves, Nucl. Phys. A97 (1967) 144
 1967BA28 G. Baciu, D. Catana, C. Deberth and I. Raileanu, Rev. Roum. Phys. 12 (1967) 385
 1967BA78 G. Backenstoss, S. Charalambus, H. Daniel, H. Koch, G. Poelz, H. Schmitt and L. Tauscher, Phys. Lett. B25 (1967) 547
 1967BE14 J.A. Becker, J.W. Olness and D.H. Wilkinson, Phys. Rev. 155 (1967) 1089
 1967BO09 M.C. Bouten, J.P. Elliott and J.A. Pullen, Nucl. Phys. A97 (1967) 113
 1967BR1H C. Brihaye and G. Reidemeister, Nucl. Phys. A100 (1967) 65
 1967BU23 S.T. Butler, R.G.L. Hewitt, B.H.J. McKellar and R.M. May, Ann. Phys. 43 (1967) 282
 1967CA1M F.P. Calaprice, E.D. Commins, H.M. Gibbs, G.L. Wick and D.A. Dobson, Phys. Rev. Lett. 18 (1967) 918
 1967CA1N C.G. Callan and S.B. Treiman, Phys. Rev. 162 (1967) 1494

- 1967CH04 R. Chapman and A.M. Macleod, Nucl. Phys. A94 (1967) 324
- 1967CH35 R. Chiba, Nucl. Sci. App. 3 (1967) 36
- 1967CO1D Commins, Ann. Rev. Nucl. Sci. 17 (1967) 33
- 1967CO1J Coop, Poate and Titterton, Aust. J. Phys. 20 (1967) 609
- 1967DR08 I.P. Dryapachenko, V.A. Kornilov, O.F. Nemetz and V.A. Pilipchenko, Yad. Fiz. 6 (1967) 440; Sov. J. Nucl. Phys. 6 (1968) 321
- 1967EN01 T. Engeland and P.J. Ellis, Phys. Lett. B25 (1967) 57
- 1967ES02 R.A. Esterlund, R. McPherson, A.M. Poskanzer and P.L. Reeder, Phys. Rev. 156 (1967) 1094
- 1967FE01 P. Federman, Nucl. Phys. A95 (1967) 443
- 1967FL13 J. Flores and R. Perez, Phys. Lett. B26 (1967) 55
- 1967GR04 M.W. Greene and E.B. Nelson, Phys. Rev. 153 (1967) 1068
- 1967GU05 M.R. Gunye and C.S. Warke, Phys. Rev. 156 (1967) 1087
- 1967GU1D M.R. Gunye and C.S. Warke, Phys. Rev. 159 (1967) 885
- 1967HA1M Halbert, Kim, McGrory and Kuo, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 531
- 1967HI1C Hirko et al., Bull. Amer. Phys. Soc. 12 (1967) 536
- 1967IN03 T. Inoue, T. Sebe, K.K. Huang and A. Arima, Nucl. Phys. A99 (1967) 305
- 1967KA1C S. Kaufman and C.O. Hower, Phys. Rev. 154 (1967) 924
- 1967KO11 A.P. Kobzev, A.V. Gromov, K. Kashlik, K. Nedvedyuk, V.I. Salatskii and S.A. Telezhnikov, Yad. Fiz. 5 (1967) 510; Sov. J. Nucl. Phys. 5 (1967) 359
- 1967KO1G Kobzev et al., Joint Inst. Nucl. Res., Lab. Neutron Phys., USSR Rept. No. P6 3314 (1967)
- 1967LO1B Loiseaux, Maison and Langevin, J. Phys. 28 (1967) 11
- 1967MI1B Miller, Thesis, Princeton Univ. (1967); Phys. Abs. 9047 (1969)
- 1967MO23 J.F. Morgan and R.K. Hobbie, Phys. Rev. 163 (1967) 992
- 1967NE1D Nemirovskii, Sov. J. Nucl. Phys. 4 (1967) 334
- 1967NI1B Nikolaev, Proc. P. N. Lebedev Inst. 36 (1967) 77
- 1967OL05 J.W. Olness, A.R. Poletti and E.K. Warburton, Phys. Rev. 161 (1967) 1131
- 1967OS02 V.M. Osadchiev and M.A. Troitskii, Yad. Fiz. 5 (1967) 85; Sov. J. Nucl. Phys. 5 (1967) 58
- 1967PA1G Pauling, Proc. Natl. Acad. Sci. U. S. 58 (1967) 2175
- 1967PI1B Pinkston, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 419

- 1967PR04 F.W. Prosser, Jr., G.U. Din and D.D. Tolbert, Phys. Rev. 157 (1967) 779
- 1967RI1B Ripka, Int. Nucl. Phys. Conf., Gatlinburg, 1966 (Academic Press, 1967) 833
- 1967RO10 H. Ropke, K.P. Lieb and R. Konig, Nucl. Phys. A97 (1967) 609
- 1967RO1F E. Rost, Phys. Rev. 154 (1967) 994
- 1967SH14 V.S. Shirley, UCRL-17990 (1967)
- 1967SH1G Shima and Sakisaka, Mem. Fac. Engng. Kyoto Univ. (Japan) 29 (1967) 483; Phys. Abs. 29297 (1968)
- 1967SP09 A. Sperduto, Proc. 3rd Int. Conf. on Atomic Masses, Winnipeg, Canada, 1967; Ed. R.C. Barber (Univ. Manitoba Press, 1967) 657
- 1967ST1H G.L. Strobel, Phys. Rev. 154 (1967) 941
- 1967ST28 T. Stambach, S.E. Darden, P. Huber and I. Sick, Helv. Phys. Acta 40 (1967) 915
- 1967TH06 W.J. Thompson, S. Edwards and T. Tamura, Nucl. Phys. A105 (1967) 678
- 1967TO1C Tolbert and Prosser, Bull. Amer. Phys. Soc. 12 (1967) 1033
- 1967WA13 E.K. Warburton, J.W. Olness and A.R. Poletti, Phys. Rev. 160 (1967) 938
- 1967WA1C Warburton, Nucl. Research with Low Energy Accelerators, Eds. Marion and van Patter (Academic Press, 1967) 43
- 1967YU01 H.P. Yule, Nucl. Phys. A94 (1967) 442
- 1967ZA1C Zaikin, Proc. P. N. Lebedev Inst. 36 (1967) 9
- 1968AG1D Aguilar, Lineda and de La Rubia, An. Fis. 64 (1968) 93
- 1968AN1F Antolkovic et al., Izv. Akad. Nauk SSSR Ser. Fiz. 32 (1968) 1658
- 1968AR02 A. Arima, S. Cohen, R.D. Lawson and M.H. Macfarlane, Nucl. Phys. A108 (1968) 94
- 1968BA2G Backenstoss et al., Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 500
- 1968BA2K Barnes, Moline, Morris and Adams, National Acad. Sci. Autumn Meeting, CalTech (1968)
- 1968BE34 A.-M. Bergdolt and G. Bergdolt, Compt. Rend. B267 (1968) 455
- 1968BI1A Bister, Fontell and Vikberg, Ann. Acad. Sci. Fennicae, Ser. A6 (1968) 3
- 1968CE1A Cerny, Ann. Rev. Nucl. Sci. 18 (1968) 27
- 1968CO11 S. Cohen, E.C. Halbert and S.P. Pandya, Nucl. Phys. A114 (1968) 353
- 1968CO1N S. Cohen, R.D. Lawson and S.P. Pandya, Nucl. Phys. A114 (1968) 541
- 1968CR03 G.M. Crawley and G.T. Garvey, Phys. Rev. 167 (1968) 1070
- 1968DW1A M. Dworzecka and C.S. Warke, Phys. Rev. 175 (1968) 1346
- 1968EI1B Eichler, Marumori and Takada, Prog. Theor. Phys. 40 (1968) 60

1968EL1C Elliott and Wilsdon, Proc. Roy. Soc. 302 (1968) 509
1968ET1A Eto, Kawai, Matsuda and Mirata, J. Phys. Soc. Jpn. 24 (1968) 422
1968FA01 M. Fazio, P. Guazzoni, S. Micheletti, M. Pignanelli and L. Zetta, Nucl. Phys. A111 (1968) 255
1968FA04 W.R. Falk, U. Matter, A. Huber, R.W. Benjamin and P. Marmier, Nucl. Phys. A117 (1968) 353
1968FO1A Fowler, Neutron Cross Sections Tech., NBS Special Publ. 299 (1968) 1
1968GA03 R.M. Gaedke, K.S. Toth and I.R. Williams, Phys. Rev. 167 (1968) 957
1968GO10 J.D. Goss, F.L. Riffle, D.R. Parsignault and J.C. Harris, Nucl. Phys. A115 (1968) 113
1968GU07 K. Gul, B.H. Armitage and B.W. Hooton, Nucl. Phys. A122 (1968) 81
1968GU1C M.R. Gunye, Nucl. Phys. A118 (1968) 174
1968GU1E M.R. Gunye, Phys. Lett. B27 (1968) 136
1968GU1F R.K. Gupta and L.E.H. Trainor, Nucl. Phys. A120 (1968) 273
1968GU1G M.R. Gunye, Nucl. Phys. A120 (1968) 691
1968HA16 I. Hamamoto and A. Arima, Nucl. Phys. A112 (1968) 481
1968HA17 E.C. Halbert, J.B. McGroory and B.H. Wildenthal, Phys. Rev. Lett. 20 (1968) 1112
1968HA1P Halbert, 3rd Symp. on the Struct. of Low-Medium Mass Nuclei; Ed. Davidson (U. Press of Kansas, 1968) 128
1968HA26 M. Harvey and T. Sebe, Nucl. Phys. A117 (1968) 289
1968HI08 B. Hird and T.Y. Li, Can. J. Phys. 46 (1968) 1273
1968HI1F Hirko et al., Tokyo (1968)
1968HO1C Hodgson, Proc. Symp. on Direct Reactions with ^3He , IPCR, Japan, Sept. 1967 (1968) 41
1968JA1H Jackson et al., Bull. Amer. Phys. Soc. 13 (1968) 1370
1968KL05 O. Klepper and H. Spehl, Z. Phys. 215 (1968) 17
1968LE18 C.O. Lennon, P.R. Alderson, J.L. Durell, L.L. Green and I.M. Naqib, Phys. Lett. B28 (1968) 253
1968LE1L R. Leonardi and M. Rosa-Clot, Phys. Rev. Lett. 21 (1968) 377
1968ME13 R.O. Mead and F.C. Young, Nucl. Phys. A115 (1968) 161
1968ME23 R.A. Meyer, W.B. Walters and J.P. Hummel, Nucl. Phys. A122 (1968) 606
1968MI05 P.F. Mizera and J.B. Gerhart, Phys. Rev. 170 (1968) 839
1968MI09 R. Middleton, L.M. Polsky, C.H. Holbrow and K. Bethge, Phys. Rev. Lett. 21 (1968) 1398

- 1968MO1G Moshinsky, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 105
- 1968MU1B Murthy, Nucl. Phys. Solid State Phys. Symp., Powai, India (1968); Phys. Abs. 38861 (1970)
- 1968PA11 P. Paul, J.W. Olness and E.K. Warburton, Phys. Rev. 173 (1968) 1063
- 1968PA1Q Pascual and Flores, Rev. Mex. Fisica 17 (1968) 86
- 1968RE07 D. Rendic, B. Antolkovic, G. Paic, M. Turk and P. Tomas, Nucl. Phys. A117 (1968) 113
- 1968RI13 L.R. Rice and R.T. Carpenter, Nucl. Phys. A120 (1968) 220
- 1968RI1M Ripka, 3rd Symp. on the Struct. of Low-Medium Mass Nuclei; Ed. Davidson (U. Press of Kansas, 1968) 167
- 1968RI1N Richter and Wiegandt, Z. Phys. 217 (1968) 225
- 1968RO07 S.W. Robinson and R.D. Bent, Phys. Rev. 169 (1968) 780
- 1968SO1A Sotona, Czech. J. Phys. 18 (1968) 714
- 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, Nucl. Phys. A113 (1968) 395
- 1968ST1K G.L. Strobel, Nucl. Phys. A115 (1968) 364
- 1968SU1C K. Sugawara, Nucl. Phys. A110 (1968) 305
- 1968TH04 T.D. Thomas, G.M. Raisbeck, P. Boerstling, G.T. Garvey and R.P. Lynch, Phys. Lett. B27 (1968) 504
- 1968TO09 J.H. Towle and G.J. Wall, Nucl. Phys. A118 (1968) 500
- 1968TO11 D.D. Tolbert, P.M. Cockburn and F.W. Prosser, Jr., Phys. Rev. Lett. 21 (1968) 1535
- 1968TOZU D.D. Tolbert, COO-1120-73 (1968)
- 1968UN1A Une and Yoshida, Tokyo (1968) p. 63
- 1968VO1A W. Von Oertzen, H.H. Gutbrod, M. Muller, U. Voos and R. Bock, Phys. Lett. B26 (1968) 291
- 1968WA04 G.J. Wagner, Phys. Lett. B26 (1968) 429
- 1968WI1B Wilkinson, Proc. Int. Conf. Nucl. Struct., Tokyo, Japan, 1967; Suppl. J. Phys. Soc. Jpn. 24 (1968) 469
- 1968WO1A S.S.M. Wong, Nucl. Phys. A116 (1968) 609
- 1968YA03 K. Yagi, J. Phys. Soc. Jpn. 24 (1968) 947
- 1969AI01 J.H. Aitken, K.W. Allen, R.E. Azuma, A.E. Litherland and D.W.O. Rogers, Phys. Lett. B28 (1969) 653
- 1969AI02 J.H. Aitken, A.E. Litherland, W.R. Dixon and R.S. Storey, Phys. Lett. B30 (1969) 473

- 1969AL15 T.K. Alexander, O. Hausser, K.W. Allen and A.E. Litherland, *Can. J. Phys.* 47 (1969) 2335
- 1969AN06 J.D. Anderson, S.D. Bloom, C. Wong, W.F. Hornyak and V.A. Madsen, *Phys. Rev.* 177 (1969) 1416
- 1969AR13 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, *Nucl. Phys.* A137 (1969) 348
- 1969BA1N Bahcall and Fowler, *Astrophys. J.* 157 (1969) 659
- 1969BA1P Bacon et al., *Acta Cryst.* A25 (1969) 391
- 1969BA2E A.I. Baz and V.I. Manko, *Phys. Lett.* B28 (1969) 541
- 1969BA62 D.J. Baugh, J. Nurzynski, D.M. Rosalky and C.H. Osman, *Aust. J. Phys.* 22 (1969) 555
- 1969BE13 G. Bergdolt and G. Guillaume, *J. Phys. (Paris)* 30 (1969) 145
- 1969BE1V H.G. Benson and B.H. Flowers, *Nucl. Phys.* A126 (1969) 305
- 1969BE57 P.M. Beard, P.B. Parks, E.G. Bilpuch and H.W. Newson, *Ann. Phys.* 54 (1969) 566
- 1969BE90 K. Bethge, C.M. Fou and R.W. Zurmu, *Nucl. Phys.* A123 (1969) 521
- 1969BE92 N. Bezic, D. Brajnik, D. Jamnik and G. Kernel, *Nucl. Phys.* A128 (1969) 426
- 1969BH01 K. Bharuth-Ram, K.P. Jackson, K.W. Jones and E.K. Warburton, *Nucl. Phys.* A137 (1969) 262
- 1969BH03 K.L. Bhatia and N. De Takacsy, *Phys. Rev.* 187 (1969) 1347
- 1969BI1A Bishop and Middleton, *Bull. Amer. Phys. Soc.* 14 (1969) 530
- 1969BL02 J. Bleck, D.W. Haag, W. Leitz, R. Michaelsen, W. Ribbe and F. Sichelschmidt, *Nucl. Phys.* A123 (1969) 65
- 1969BR07 K.H. Bray and J. Nurzynski, *Nucl. Phys.* A127 (1969) 622
- 1969CA14 F.P. Calaprice, E.D. Commins, H.M. Gibbs, G.L. Wick and D.A. Dobson, *Phys. Rev.* 184 (1969) 1117
- 1969CA1H Calaprice, *Bull. Amer. Phys. Soc.* 14 (1969) 1217
- 1969CE01 J. Cerny, R.A. Mendelson, Jr., G.J. Wozniak, J.E. Esterl and J.C. Hardy, *Phys. Rev. Lett.* 22 (1969) 612
- 1969CH1C D.T. Chivers, E.M. Rimmer, B.W. Allardyce, R.C. Witcomb, J.J. Domingo and N.W. Tanner, *Nucl. Phys.* A126 (1969) 129
- 1969CU07 R.Y. Cusson, *Can. J. Phys.* 47 (1969) 2835
- 1969DA08 C.M. da Silva, J.O. Newton, J.C. Lisle and M.F. da Silva, *Nucl. Phys.* A132 (1969) 9
- 1969DI07 G.U. Din, *Nucl. Phys.* A134 (1969) 655
- 1969DU1A Dunning, Thesis, Catholic Univ. (1969); *Phys. Abs.* 42836 (1970)

- 1969EP1B M. Epherre, E. Gradsztajn, R. Klapisch and H. Reeves, Nucl. Phys. A139 (1969) 545
- 1969FE1A Federman, Cargese Lecture in Physics, Vol. 3, Ed. M. Jean (Gordon and Breach, 1969) 21
- 1969FI07 P. Fintz, F. Hibou, B. Rastegar and A. Gallmann, Nucl. Phys. A132 (1969) 265
- 1969FU11 G.H. Fuller and V.W. Cohen, Nucl. Data Tables A5 (1969) 433
- 1969GA1G Garvey, Ann. Rev. Nucl. Sci. 19 (1969) 433
- 1969GL06 Y.A. Glukhov, B.G. Novatskii, A.A. Ogloblin, S.B. Sakuta and D.N. Stepanov, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 609; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 561
- 1969GU1E M.R. Gunye, Nucl. Phys. A128 (1969) 457
- 1969HA1F Hanna, Isospin in Nucl. Phys., Ed. D.H. Wilkinson (North-Holland, 1969) 591
- 1969HA1G Hanna, Proc. Int. Conf., Montreal (1969) 443
- 1969HA38 J.C. Hardy, H. Brunnader, J. Cerny and J. Janecke, Phys. Rev. 183 (1969) 854
- 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, Phys. Rev. 178 (1969) 1551
- 1969HE1P Hevezi, Klopčic, Darden and Lietz, Bull. Amer. Phys. Soc. 14 (1969) 186
- 1969HI1B Hiura, Abe, Saito and Endo, Prog. Theor. Phys. 42 (1969) 555
- 1969HO16 W. Hofmann, R. Kosiek, G. Kraft and R. Mundhenke, Z. Phys. 225 (1969) 303
- 1969HO32 R.J.W. Hodgson, Can. J. Phys. 47 (1969) 2269
- 1969JA09 K.P. Jackson, K. Bharuth Ram, P.G. Lawson, N.G. Chapman and K.W. Allen, Phys. Lett. B30 (1969) 162
- 1969JO10 K.W. Jones, A.Z. Schwarzschild, E.K. Warburton and D.B. Fossan, Phys. Rev. 178 (1969) 1773
- 1969KR02 H.J. Krappe and U. Wille, Nucl. Phys. A124 (1969) 641
- 1969KU1G Kulkarni and Pandya, Nuovo Cim. B60 (1969) 199
- 1969LE1D Leonardi and Rosa-Clot, Lett. Nuovo Cim. 1 (1969) 829
- 1969LO06 M.A.K. Lodhi, Phys. Rev. 182 (1969) 1061
- 1969MA1T MacFarlane, Proc. Enrico Fermi School of Phys., Course XL, Lake Como, 1967 (Academic Press, 1969) 457
- 1969ME13 R.A. Mendelson, Jr. and R.T. Carpenter, Phys. Rev. 183 (1969) 924
- 1969ME15 D. Meier, M. Brullmann, H. Jung and P. Marmier, Helv. Phys. Acta 42 (1969) 813
- 1969ME1C Meyer, Amer. J. Phys. 37 (1969) 296
- 1969MO1E L. Moyer and D.S. Koltun, Phys. Rev. 182 (1969) 999
- 1969NI09 R.J. Nickles, Nucl. Phys. A134 (1969) 308

- 1969OV01 J.C. Overley, P.D. Parker and D.A. Bromley, Nucl. Instrum. Meth. 68 (1969) 61
- 1969PA1C Pape, Chevallier, Sens and Armbruster, Rev. Phys. Appl. 4 (1969) 227
- 1969PE1D Petrauskas and Vanagas, Sov. J. Nucl. Phys. 8 (1969) 270
- 1969PL1B Plendl et al., Bull. Amer. Phys. Soc. 14 (1969) 1241
- 1969PO03 A.R. Poletti, J.A. Becker and R.E. McDonald, Phys. Rev. 182 (1969) 1054
- 1969PO07 G. Poelz, H. Schmitt, L. Tauscher, G. Backenstoss, S. Charalambus, H. Daniel and H. Koch, Z. Phys. 227 (1969) 311
- 1969RO1G I. Rotter, Nucl. Phys. A135 (1969) 378
- 1969SE02 D.L. Sellin, H.W. Newson and E.G. Bilpuch, Ann. Phys. (N.Y.) 51 (1969) 461
- 1969SE03 D.L. Sellin, W.P. Beres and E.G. Bilpuch, Ann. Phys. (N.Y.) 51 (1969) 476
- 1969SK1A Skakun and Kharkov, Atomn. Energ. (USSR) 27 (1969) 351
- 1969ST07 R.H. Stokes and P.G. Young, Phys. Rev. 178 (1969) 1789
- 1969SU15 K. Sugimoto, Phys. Rev. 182 (1969) 1051
- 1969TA1C Tanaka, Bochum Conf. STI/PUB/232 IAEA (1969) 137
- 1969VE09 A.N. Vereshchagin, I.N. Simonov, K.O. Terenetskii, V.V. Tokarevskii and I.P. Chernov, Izv. Akad. Nauk SSSR Ser. Fiz. 33 (1969) 2064; Bull. Acad. Sci. USSR Phys. Ser. 33 (1970) 1880
- 1969VO10 U.C. Voos, W. von Oertzen and R. Bock, Nucl. Phys. A135 (1969) 207
- 1969WH1A White, Bull. Amer. Phys. Soc. 14 (1969) 1214
- 1969WO1F Wormald and Wright, Contrib., Montreal (1969) 111
- 1969WU1A Wu and Wilets, Ann. Rev. Nucl. Sci. 19 (1969) 527
- 1969ZA01 V.M. Zabegai, A.E. Melenevskii, O.F. Nemetz, V.N. Nemykin, Y.S. Stryuk and K.O. Terenetskiim, Yad. Fiz. 10 (1969) 227; Sov. J. Nucl. Phys. 10 (1970) 129
- 1970AD02 E.G. Adelberger and A.B. McDonald, Nucl. Phys. A145 (1970) 497
- 1970AG08 J. Aguilar, C.F. Pineda, C.R. Bauza, J.L. Ferrero, A. Garcia and F. Senent, An. Fis. 66 (1970) 279
- 1970AI01 J.H. Aitken, R.E. Azuma, A.E. Litherland, A.M. Charlesworth, D.W.O. Rogers and J.J. Simpson, Can. J. Phys. 48 (1970) 1617
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- 1970AL21 D.E. Alburger and G.A.P. Engelbertink, Phys. Rev. C2 (1970) 1594
- 1970AR1D Artukh et al., Nucl. Reactions Included by Heavy Ions, Heidelberg, 1969 (North-Holland, 1970) 140
- 1970BA1E Backenstoss, Ann. Rev. Nucl. Sci. 20 (1970) 467
- 1970BA1M Bahcall and Fowler, Astrophys. J. 161 (1970) 119

- 1970BA1Y Barker, Cockburn, Seiler and Marmier, *Helv. Phys. Acta* 43 (1970) 744
- 1970BA2B Baz and Manko, *Sov. J. Nucl. Phys.* 10 (1970) 46
- 1970BH02 K. Bharuth-Ram, R.D. Gill, K.P. Jackson, B. Povh and E.K. Warburton, *Phys. Rev. C2* (1970) 1210
- 1970BL07 R. Blomquist, M. Eckhause, F.R. Kane, J.R. Kane, P. Martin, G.H. Miller, W.W. Sapp, C.B. Spence and R.E. Welsh, *Nucl. Phys. B19* (1970) 632
- 1970BL1F Bleck, Haag and Ribbe, *Z. Phys.* 233 (1970) 65
- 1970BO08 T.I. Bonner, *Phys. Rev. C1* (1970) 1699
- 1970CA1N Carlson and Johnson, *Bull. Amer. Phys. Soc.* 15 (1970) 629
- 1970CL03 R.L. Clarke and P. Perrin, *Nucl. Phys. A147* (1970) 174
- 1970CL1C Clayton, 2nd Int. Conf. on Nucl. Data for Reactors, Helsinki (1970) 35
- 1970CO22 J.C. Cooper and B. Crasemann, *Phys. Rev. C2* (1970) 451
- 1970DE06 D. Dehnhard and N.M. Hintz, *Phys. Rev. C1* (1970) 460
- 1970EL1G Ellis and Engeland, *Nucl. Phys. A144* (1970) 161
- 1970FA01 W.R. Falk, A. Huber, U. Matter, R.W. Benjamin and P. Marmier, *Nucl. Phys. A140* (1970) 548
- 1970FE1E Fernandez et al., *An. Fis.* 66 (1970) 81
- 1970FI08 P. Fintz, F. Hibou, B. Rastegar and A. Gallman, *Nucl. Phys. A150* (1970) 49
- 1970FO1G Foroughi, *Helv. Phys. Acta* 43 (1970) 748
- 1970GA18 J.D. Garrett, R. Middleton and H.T. Fortune, *Phys. Rev. C2* (1970) 1243
- 1970GI09 R.D. Gill, K. Bharuth-Ram, K.P. Jackson, R.A.I. Bell, B.C. Robertson, J. L'Ecuyer, N.G. Chapman and H.J. Rose, *Nucl. Phys. A152* (1970) 369
- 1970GI11 B. Giraud and P.U. Sauer, *Nucl. Phys. A154* (1970) 587
- 1970GR04 L.L. Green, C.O. Lennon and I.M. Naqib, *Nucl. Phys. A142* (1970) 137
- 1970HA1R Hallowell et al., *Bull. Amer. Phys. Soc.* 15 (1970) 501
- 1970JO1H Jones, Thesis, Iowa State Univ.; *Phys. Abs.* 6448 (1970)
- 1970KA31 G.T. Kaschl, G.J. Wagner, G. Mairle and U. Schmidt-Rohr, *Nucl. Phys. A155* (1970) 417
- 1970KO41 S.-I. Koyama, K. Takahashi and M. Yamada, *Prog. Theor. Phys.* 44 (1970) 663
- 1970LA02 S.T. Lam, A.E. Litherland and J.J. Simpson, *Can. J. Phys.* 48 (1970) 827
- 1970LE1A R. Leonardi and M. Rosa-Clot, *Phys. Rev. Lett.* 24 (1970) 407
- 1970MC1F McEver et al., *Bull. Amer. Phys. Soc.* 15 (1970) 653
- 1970MC1J J.B. McGrory, *Phys. Lett. B31* (1970) 339

- 1970MC23 J.B. McGrory, Phys. Lett. B33 (1970) 327
- 1970MI1E Middleton, Proc. Int. Conf. Nucl. Reactions Induced by Heavy Ions, Heidelberg, 1969 (North-Holland, 1970) p. 263
- 1970OL1A Olivieri and Salusti, Nuovo Cim. 69 (1970) 551
- 1970OT1B Ott and Weller, Bull. Amer. Phys. Soc. 15 (1970) 163
- 1970PR1B Preedom, Wildenthal, Snelgrove and Kashy, Bull. Amer. Phys. Soc. 15 (1970) 36
- 1970PR1G Primakoff, Springer Tracts in Modern Phys. 53 (1970)
- 1970RA1A Raisbeck et al., PUC-937-369 (1970)
- 1970RO1C Rogers et al., Bull. Amer. Phys. Soc. 15 (1970) 542
- 1970SC05 J.J. Schwartz and B.A. Watson, Phys. Rev. Lett. 24 (1970) 322
- 1970SC23 G. Schekliniski, U. Strohbush and B. Goel, Nucl. Phys. A153 (1970) 97
- 1970SC25 C. Schmidt and H.H. Duhm, Nucl. Phys. A155 (1970) 644
- 1970ST04 H.J. Strubbe and D.K. Callebaut, Nucl. Phys. A143 (1970) 537
- 1970TA1J Talmi, Theory of Nucl. Struct. - Trieste 1969, IAEA STI/PUB/249 (1970) 455
- 1970VE06 A.N. Vereshchagin, I.N. Korostova and I.P. Chernov, Izv. Vys. Ucheb. Zav. Fiz. 8 (1970) 105; Sov. Phys. J. 13 (1973) 1071
- 1970WA02 B.C. Walsh and I.M. Naqib, Nucl. Phys. A140 (1970) 571
- 1970WA08 T. Walcher and P. Strehl, Z. Phys. 232 (1970) 342
- 1970WA1G Walker and Stokes, LA-DC 11224 (1970)
- 1970WA1N Walcher, Private Communication (1970)
- 1970WE1K Weller and Ott, Bull. Amer. Phys. Soc. 15 (1970) 1686
- 1971AJ02 F. Ajzenberg-Selove, Nucl. Phys. A166 (1971) 1
- 1971AN08 G. Andersson, B. Forkman and B. Friberg, Nucl. Phys. A171 (1971) 529
- 1971AR1E Artemov, Gol'dberg, Islamov and Rudakov, Sov. J. Nucl. Phys. 12 (1971) 130
- 1971AR1R A. Arima and I. Hamamoto, Ann. Rev. Nucl. Sci. 21 (1971) 55
- 1971AR25 A. Arima, M. Sakakura and T. Sebe, Nucl. Phys. A170 (1971) 273
- 1971BI06 H.G. Bingham, H.T. Fortune, J.D. Garrett and R. Middleton, Phys. Rev. Lett. 26 (1971) 1448
- 1971BO1V Bock et al., Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 494
- 1971BR02 C. Broude, U. Karfunkel and Y. Wolfson, Nucl. Phys. A161 (1971) 241
- 1971BU1E Butler, Poskanzer and Landis, Proc. Int. Conf. on Heavy Ion Phys., Dubna (1971) 210
- 1971CE1A Cerny, LBL-234 (1971)
- 1971DI09 W.R. Dixon and R.S. Storey, Can. J. Phys. 49 (1971) 1714

- 1971DI18 W.R. Dixon, R.S. Storey, J.H. Aitken, A.E. Litherland and D.W.O. Rogers, Phys. Rev. Lett. 27 (1971) 1460
- 1971DI1F di Napoli et al., Nucl. Instrum. Meth. 93 (1971) 77
- 1971DU12 A. Dudek and D.J. Edens, Phys. Lett. B36 (1971) 309
- 1971FA09 J. Favier, T. Bressani, G. Charpak, L. Massonnet, W.E. Meyerhof and C. Zupancic, Nucl. Phys. A169 (1971) 540
- 1971FO18 H.T. Fortune and C.M. Vincent, Phys. Rev. C4 (1971) 1994
- 1971FR11 B. Friberg, G. Andersson and B. Forkman, Nucl. Phys. A171 (1971) 551
- 1971GA1H Garrett, Bingham, Fortune and Middleton, Bull. Amer. Phys. Soc. 16 (1971) 1148
- 1971GO1B Gove and Panagiotou, Bull. Amer. Phys. Soc. 16 (1971) 490
- 1971GU1N M.R. Gunye, Phys. Lett. B37 (1971) 125
- 1971HA30 K.A. Hardy, A.H. Lumpkin, Y.K. Lee and G.E. Owen, Phys. Rev. C4 (1971) 317
- 1971HI06 F. Hibou, P. Fintz, B. Rastegar and A. Gallmann, Nucl. Phys. A171 (1971) 603
- 1971HIZF F. Hibou, Thesis, Univ. Louis Pasteur, Strasbourg (1971)
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1971LA1D Lang, Z. Phys. 242 (1971) 179
- 1971LE1H T.Y. Lee, S.T. Hsieh, W.W. Yeh and C.T. Chen-Tsai, Phys. Rev. C4 (1971) 64
- 1971LE30 T.Y. Lee, C.T. Chen-Tsai, W.W. Yeh and S.T. Hsieh, Phys. Rev. C4 (1971) 1991
- 1971MC11 R.E. McDonald, J.A. Becker, A.D.W. Jones and A.R. Poletti, Phys. Rev. C4 (1971) 377
- 1971MC1M McEver et al., Polarization Phenomena in Nucl. Reactions, Madison, 1970; Eds. H.H. Barschall and W. Haerberli (Univ. Wisconsin Press, 1971) p. 603
- 1971NI05 L. Nichol and T.J. Kennett, Can. J. Phys. 49 (1971) 1461
- 1971OT1C Ott and Weller, Private Communication (1971)
- 1971SH1G Sherman, Lokan and Gellie, Bull. Amer. Phys. Soc. 16 (1971) 16; Phys. Canada 27 (1971) 59
- 1971SK1C Skakun et al., At. Energ. 30 (1971) 456
- 1971TH05 S.O. Thompson, L. Husain and S. Katcoff, Phys. Rev. C3 (1971) 1538
- 1971WE1F Weidenmuller, Polarization Phenomena in Nucl. Reactions, Madison, 1970, Eds. H.H. Barshall and W. Haerberli (Univ. Wisconsin Press, 1971) p. 15
- 1971WI01 WildeB.H. Wildenthal, J.B. McGrory and P.W.M. Glaudemans, Phys. Rev. Lett. 26 (1971) 96
- 1972HA03 D.S. Haynes, K.W. Kemper and N.R. Fletcher, Phys. Rev. C5 (1972) 5

1972LE1L Lee and Cusson, *Ann. Phys.* 72 (1972) 353
1972RO01 D.W.O. Rogers, J.G. Aitken and A.E. Litherland, *Can. J. Phys.* 50 (1972) 268
1974AJ01 F. Ajzenberg-Selove and T. Lauritsen, *Nucl. Phys.* A227 (1974) 1
PR63A Unknown Source