

# Energy Levels of Light Nuclei $A = 19$

F. Ajzenberg-Selove

*University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396*

**Abstract:** An evaluation of  $A = 18-20$  was published in *Nuclear Physics A392* (1983), p. 1. This version of  $A = 19$  differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed May 1, 1982)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

## Table of Contents for $A = 19$

*Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.*

A. Nuclides: [<sup>19</sup>B](#), [<sup>19</sup>C](#), [<sup>19</sup>N](#), [<sup>19</sup>O](#), [<sup>19</sup>F](#), [<sup>19</sup>Ne](#), [<sup>19</sup>Na](#), [<sup>19</sup>Mg](#)

B. Tables of Recommended Level Energies:

[Table 19.1](#): Energy levels of <sup>19</sup>O

[Table 19.6](#): Energy levels of <sup>19</sup>F

[Table 19.23](#): Energy levels of <sup>19</sup>Ne

C. [References](#)

D. Figures: [<sup>19</sup>O](#), [<sup>19</sup>F](#), [<sup>19</sup>Ne](#), [Isobar diagram](#)

E. Erratum to this Publication: [PS](#) or [PDF](#)

**<sup>19</sup>B**  
(Not illustrated)

Assuming the atomic mass excess to be 60.1 MeV [see (1978AJ03)], <sup>19</sup>B is stable with respect to breakup into <sup>18</sup>B + n by 1.8 MeV and into <sup>17</sup>B + 2n by 0.4 MeV.

**<sup>19</sup>C**  
(Not illustrated)

<sup>19</sup>C has been observed in the 4.8 GeV proton bombardment of uranium: it is particle stable (1974BO05). The calculated mass excess of <sup>19</sup>C is 32.45 MeV using the modified form of the IMME (1975JE02): <sup>19</sup>C would then be stable with respect to decay into <sup>18</sup>C + n by 0.53 MeV and into <sup>17</sup>C + 2n by 4.72 MeV [see (1982AJ01) for the mass of <sup>17</sup>C. See also (1978AJ03).

**<sup>19</sup>N**  
(Fig. 8)

Studies of the <sup>18</sup>O(<sup>18</sup>O, <sup>17</sup>F)<sup>19</sup>N and <sup>208</sup>Pb(<sup>18</sup>O, <sup>207</sup>Bi)<sup>19</sup>N reactions at  $E(^{18}\text{O}) = 91$  and 93 MeV, respectively, lead to values of the atomic mass excess of <sup>19</sup>N of  $15.856 \pm 0.050$  (1982NA08) and  $15.96 \pm 0.15$  MeV (1979BA31). The adopted value is  $15.866 \pm 0.048$  MeV. <sup>19</sup>N is then stable with respect to decay into <sup>18</sup>N + n by 5.45 MeV. Differential cross sections for the two reactions in which <sup>19</sup>N has been observed are  $\approx 500$  nb/sr ( $6^\circ$ ) (1982NA08) and  $\approx 120$  nb/sr ( $85^\circ$ ) (1979BA31). In addition to the ground-state transition, (1982NA08) report the population of states at  $E_x = 1.12 \pm 0.04$  and  $1.59 \pm 0.04$  MeV. See also (1980AL1F, 1980NA12) and (1978AJ03).

**<sup>19</sup>O**  
(Figs. 5 and 8)

GENERAL: (See also (1978AJ03).)

*Shell model:* (1977GR16, 1979DA15, 1980KU05, 1982KI02).

*Electromagnetic transitions:* (1976MC1G, 1978KR19, 1980KU05).

*Special states:* (1977GR16, 1977SH18, 1979DA15, 1982KI02).

*Astrophysical questions:* (1978WO1E).

*Complex reactions involving <sup>19</sup>O:* (1978KO01, 1979AL22, 1981GR08).

*Other topics:* (1977GR16, 1977SH18, 1979BE1H, 1979CO09, 1980SH1H, 1982KI02).

*Ground-state properties of <sup>19</sup>O:* (1976MC1G).

Table 19.1: Energy levels of  $^{19}\text{O}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau^b$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{5}{2}^+; \frac{3}{2}$	$\tau_{1/2} = 26.91 \pm 0.08$ sec	$\beta^-$	1, 2, 3, 4, 5, 9, 10, 11, 12
$0.0960 \pm 0.5$	$\frac{3}{2}^+$	$\tau_m = 2.00 \pm 0.07$ nsec $g = -0.48 \pm 0.06$	$\gamma$	3, 4, 9, 10, 12
$1.4717 \pm 0.4$	$\frac{1}{2}^+$	$\tau_m = 1.27 \pm 0.17$ psec	$\gamma$	3, 4, 9
$2.3715 \pm 1.0$	$\frac{9}{2}^+$	$> 3.5$ psec	$\gamma$	3, 4, 9
$2.7790 \pm 0.9$	$\frac{7}{2}^+$	$93 \pm 19$ fsec	$\gamma$	3, 4, 9
$3.0674 \pm 1.6$	$\frac{3}{2}^+$	$\geq 1$ psec	$\gamma$	3, 4, 9
$3.1535 \pm 1.7$	$\frac{5}{2}^+$	$(\geq 1$ psec)	$\gamma$	3, 4, 9
$3.2316 \pm 2.3$	$\frac{3}{2}^+$			3, 4, 9
$3.9449 \pm 1.4$ <sup>c</sup>	$\frac{3}{2}^-$		$\gamma$	3, 4, 9
$4.1093 \pm 1.9$	$\frac{3}{2}^+$	$\Gamma < 15$ keV		3, 4, 9
$4.3281 \pm 2.4$	$\frac{3}{2}, \frac{5}{2}$	$< 15$		3, 4, 9
$4.4025 \pm 2.7$	$\frac{3}{2} \rightarrow \frac{7}{2}$	$< 15$		3, 4, 9
$4.5820 \pm 4.6$	$\frac{3}{2}^-$	$52 \pm 3$	n	3, 4, 6, 9
$4.7026 \pm 2.7$	$\frac{5}{2}^+$	$< 15$		3, 4, 9
$4.9683 \pm 5.5$	$\frac{5}{2}, \frac{7}{2}$			3
$5.0070 \pm 4.5$	$\frac{3}{2}, \frac{5}{2}$	$< 15$		3, 4, 9
$5.0820 \pm 5.4$	$\frac{1}{2}^-$	$49 \pm 5$	n	3, 6
$5.1484 \pm 3.2$	$\geq \frac{5}{2}^+$	$3.4 \pm 1.0$	n	3, 4, 6, 9
5.33	$\frac{3}{2}^+$	330	n	6
$5.3840 \pm 2.8$	$(\frac{9}{2} \rightarrow \frac{13}{2})$			3
$5.455 \pm 9$	$\frac{5}{2}^+$	280	n	6
$5.5035 \pm 3.1$ <sup>c</sup>		$< 15$		3, 4, 9
$5.7046 \pm 4.3$ <sup>c</sup>		$7.8 \pm 1.4$	n	3, 4, 6, 9
$6.1196 \pm 3.2$ <sup>c</sup>				3
6.13	$\frac{3}{2}^+$	190	n	6
$6.1916 \pm 5.5$	$\frac{1}{2}^-$	120	n	3, 6
$6.2693 \pm 2.6$	$\frac{7}{2}^-$	$19.2 \pm 2.4$	n	3, 4, 6, 9
$6.4058 \pm 3.1$ <sup>c</sup>				3
$6.4662 \pm 4.8$	$(\frac{7}{2} \rightarrow \frac{11}{2})$		(n)	3, 6, 9

Table 19.1: Energy levels of  $^{19}\text{O}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$\tau$ <sup>b</sup> or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
6.583 $\pm$ 6 <sup>c</sup>				3, 9
6.903 $\pm$ 8				3, 9
6.988 $\pm$ 9				3, 9
7.118 $\pm$ 10				3, 9
7.242 $\pm$ 8				3, 9
7.508 $\pm$ 10				3
8.048 $\pm$ 20				3
8.132 $\pm$ 20				3
8.247 $\pm$ 20				3
8.450 $\pm$ 20				3
8.561 $\pm$ 20				3
8.591 $\pm$ 20				3
8.916 $\pm$ 20				3
8.923 $\pm$ 20				3
9.022 $\pm$ 20				3
9.064 $\pm$ 20				3
9.253 $\pm$ 20				3
9.324 $\pm$ 20				3
9.43				3
9.56				3
9.77				3
9.88				3
9.93				3
9.98				3
11.25 $\pm$ 50		240	n, $\alpha$	6
11.58 $\pm$ 50		330	n, $\alpha$	6

<sup>a</sup> See also Table 19.2.

<sup>b</sup> See also reaction 1, and Table 19.2 in (1978AJ03).

<sup>c</sup> See footnotes to Table 19.3.

Table 19.2: Radiative decays in  $^{19}\text{O}$  <sup>a</sup>

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branch (%) <sup>a</sup>	$\delta$
0.096	$\frac{3}{2}^+$	0	100	
1.47	$\frac{1}{2}^+$	0	$2.0 \pm 0.2$	
		0.096	$98.0 \pm 0.2$	
2.37	$\frac{9}{2}^+$	0	100	$0.002 \pm 0.05$
2.78	$\frac{7}{2}^+$	0	100	$0.8 \pm 0.5$
3.07	$\frac{3}{2}^+$	1.47	100	
3.16	$\frac{5}{2}^+$	0	$8 \pm 4$	
		0.096	$92 \pm 4$	$0.03 < \delta < 2.3$
3.94	$\frac{3}{2}^-$	0	$33 \pm 8$	
		0.096	$39 \pm 8$	
		1.47	$28 \pm 4$	

<sup>a</sup> For other values and for references see Table 19.5 in (1978AJ03).

1.  $^{19}\text{O}(\beta^-)^{19}\text{F}$   $Q_m = 4.819$

The weighted mean of several reported half-lives is  $26.91 \pm 0.08$  sec: see (1972AJ02). The decay is complex: see  $^{19}\text{F}$ , reaction 36 and Table 19.20. See also (1977OK1A; theor.).

2.  $^9\text{Be}(^{18}\text{O}, ^8\text{Be})^{19}\text{O}$   $Q_m = 2.292$

Cross sections and  $S$ -factors are reported for  $E_{\text{cm}} = 1.77$  to  $6.71$  MeV (1980RO13). See also (1978AJ03).

3.  $^{13}\text{C}(^7\text{Li}, \text{p})^{19}\text{O}$   $Q_m = 7.413$

States of  $^{19}\text{O}$  reported in this reaction are displayed in Table 19.3 (1977FO10).

4.  $^{17}\text{O}(\text{t}, \text{p})^{19}\text{O}$   $Q_m = 3.520$

Proton groups corresponding to  $^{19}\text{O}$  states with  $E_x < 5.6$  MeV and  $E_\gamma$  measurements are displayed in Table 19.4.

Table 19.3: States in  $^{19}\text{O}$  from  $^{13}\text{C}(^7\text{Li}, \text{p})^a$

$E_x$ (MeV $\pm$ keV)	$J^b$	$E_x$ (MeV $\pm$ keV)
0	$\frac{5}{2}$	$6.4662 \pm 4.8^i$
$0.0944 \pm 1.1$	$\frac{3}{2}$	$6.5827 \pm 6.0^j$
$1.4716 \pm 1.8$	$\frac{1}{2}$	$6.903 \pm 8$
$2.3711 \pm 1.9$	$\frac{9}{2}$	$6.988 \pm 9$
$2.7776 \pm 1.9$	$\frac{7}{2}$	$7.118 \pm 10$
$3.0674 \pm 1.6$	$\frac{3}{2}$	$7.242 \pm 8$
$3.1536 \pm 2.8$	$\frac{5}{2}$	$7.508 \pm 10$
$3.2316 \pm 2.3$	$\frac{3}{2}$	$8.048 \pm 20$
$3.9449 \pm 1.4^c$	$\frac{3}{2}$	$8.132 \pm 20$
$4.1093 \pm 1.9$	$\frac{3}{2}$	$8.247 \pm 20$
$4.3281 \pm 2.4$	$\frac{3}{2}, \frac{5}{2}$	$8.450 \pm 20$
$4.4025 \pm 2.7$	$\frac{3}{2}, \frac{5}{2}, \frac{7}{2}$	$8.561 \pm 20$
$4.5820 \pm 4.6$	$\frac{3}{2}$	$8.591 \pm 20$
$4.7026 \pm 2.7^d$		$8.916 \pm 20$
$4.9683 \pm 5.5$	$\frac{5}{2}, \frac{7}{2}$	$8.923 \pm 20$
$5.0070 \pm 4.5$	$\frac{3}{2}, \frac{5}{2}$	$9.022 \pm 20$
$5.0820 \pm 5.4$	$\frac{1}{2}$	$9.064 \pm 20$
$5.1484 \pm 3.2$	$\frac{5}{2}$	$9.253 \pm 20$
$5.3840 \pm 2.8$	$\frac{9}{2}, \frac{11}{2}, \frac{13}{2}^e$	$9.324 \pm 20$
$5.5035 \pm 3.1^f$		9.43
$5.7046 \pm 4.3^g$		9.56
$6.1196 \pm 3.2^h$		9.77
$6.1916 \pm 5.5$	$\frac{1}{2}$	9.88
$6.2693 \pm 2.6$	$\frac{7}{2}$	9.93
$6.4058 \pm 3.1^h$		9.98

<sup>a</sup> (1977FO10);  $E(^7\text{Li}) = 16.0$  MeV. Angular distributions have been reported to all states with  $E_x < 6.8$  MeV. See also (1978AJ03).

<sup>b</sup> Derived from total cross section and  $2J + 1$  analysis.

<sup>c</sup> Corresponds to unresolved states. Assuming one of these to be a  $\frac{3}{2}^-$  state (see Table 19.5), the other should have  $J = \frac{7}{2} \rightarrow \frac{13}{2}$ .

<sup>d</sup> May correspond to unresolved states.

<sup>e</sup> If this group corresponds to a single state.

<sup>f</sup> Narrow unresolved states: see discussion in (1977FO10).

<sup>g</sup> Cross section is too large for the known state at this energy with  $J^\pi = \frac{3}{2}^+$ . If this group corresponds to a doublet, the other member should have  $J = \frac{1}{2} \rightarrow \frac{5}{2}$ .

<sup>h</sup> Sharp group; if due to a single state  $J = \frac{11}{2} \rightarrow \frac{17}{2}$ .

<sup>i</sup>  $J = (\frac{7}{2}, \frac{9}{2}, \frac{11}{2})$ .

<sup>j</sup> The total cross section to this state is very high implying unresolved states: if there are two states one must have  $J \geq \frac{13}{2}$ .

5.  $^{18}\text{O}(n, \gamma)^{19}\text{O}$   $Q_m = 3.957$

The thermal capture cross section is  $0.16 \pm 0.01$  mb (1981MUZQ). See also (1978AJ03).

6.  $^{18}\text{O}(n, n)^{18}\text{O}$   $E_b = 3.957$

The scattering amplitude (bound)  $a = 5.84 \pm 0.07$  fm,  $\sigma_{\text{free}} = 3.86 \pm 0.10$  b (1979KO26). The total cross section measured for  $E_n = 0.14$  to 2.47 MeV shows five resonances at  $E_n = 0.66, 1.19, 1.26, 1.84$  and 2.45 MeV [see Table 19.5] in addition to a broad maximum at  $E_n = 1.6$  MeV and resonance structure near 2.3 MeV (1965VA03). A phase-shift analysis by (1964DO08, 1973DO05) is consistent with the (1965VA03) results and suggests that the two broad structures can each be accounted for in terms of two levels whose parameters are displayed in Table 19.5. At higher energies [ $E_n = 2.45$  to 8.50 MeV and 10.6 to 19.0 MeV] the total cross section shows additional structures. See also (1978AJ03, 1981MUZQ).

7. (a)  $^{18}\text{O}(n, p)^{18}\text{N}$   $Q_m = -13.25$   $E_b = 3.957$   
 (b)  $^{18}\text{O}(n, d)^{17}\text{N}$   $Q_m = -13.717$

See (1978AJ03).



Table 19.4: Levels of  $^{19}\text{O}$  from  $^{17}\text{O}(t, p)$  and  $^{18}\text{O}(d, p)$ 

$E_x$ (MeV $\pm$ keV) <sup>a</sup>	$\Gamma_{\text{c.m.}}$ (keV) <sup>a</sup>	$l_n$ <sup>b</sup>	$L$ <sup>c</sup>	$S$ <sup>d</sup>	$J^\pi$
0		2	0	0.57	$\frac{5}{2}^+$
$0.0960 \pm 0.5$		2	2		$\frac{3}{2}^+$
$1.4719 \pm 0.5$		0	2	1.00	$\frac{1}{2}^+$
$2.3715 \pm 1.0$		2	(2 + 4)		$\frac{9}{2}^+$
$2.7790 \pm 0.9$		(2)	2		$\frac{7}{2}^+$
$3.0671 \pm 2.6$		2	(0 + 2)	(0.06)	$\frac{5}{2}^+$
$3.1535 \pm 2.4$		a			$\frac{3}{2}^+$
$3.237 \pm 5$		1		0.11	$\frac{3}{2}^-$
$3.944 \pm 3$	< 15	2	(2)	0.03	$\frac{3}{2}^+$
$4.118 \pm 5$	< 15				
$4.333 \pm 12$					
$4.402 \pm 12$	$75 \pm 5$	1		0.15	$\frac{3}{2}^-$
$4.584 \pm 12$	< 15	2	a	0.02	$\frac{5}{2}^+$
$4.707 \pm 12$	< 15				
$4.998 \pm 12$	< 15	2	a	0.08	$\frac{5}{2}^+$
$5.150 \pm 10$	$320 \pm 25$	2	(2 + 4)	0.85	$\frac{3}{2}^+$
$5.455 \pm 10$	45				
$5.502 \pm 12$	< 15	2		0.17	$(\frac{3}{2})^+$
$5.714 \pm 12$	< 15	3		0.13	$(\frac{7}{2}^-)$
$6.280 \pm 12$					
$6.480 \pm 15$					
$6.560 \pm 15$					
$6.899 \pm 15$					
$6.997 \pm 15$					
$7.117 \pm 15$					
$7.248 \pm 15$					

<sup>a</sup> For references and other values see Table 19.3 in (1978AJ03).

<sup>b</sup>  $^{18}\text{O}(d, p)^{19}\text{O}$ .

<sup>c</sup>  $^{17}\text{O}(t, p)^{19}\text{O}$ .

<sup>d</sup>  $E_{\bar{d}} = 14.8$  MeV: polarization and differential cross section measurements. The spectroscopic factors for the states with  $E_x > 4.1$  MeV have been calculated in the weakly bound approximation.

Table 19.5: Resonances in  $^{18}\text{O}(n, n)^{18}\text{O}$  <sup>a</sup>

$E_{\text{res}}$ (MeV $\pm$ keV)	$\Gamma_{\text{cm}}$ (keV)	$^{19}\text{O}^*$ (MeV)	$J^\pi$
0.661 $\pm$ 10	52 $\pm$ 3	4.583	$\frac{3}{2}^-$
1.192 $\pm$ 10	49 $\pm$ 5	5.086	$\frac{1}{2}^-$
1.256 $\pm$ 10	3.4 $\pm$ 1.0	5.146	$\frac{3}{2}^-$
1.45	330	5.33	$\frac{3}{2}^+$
1.60	280	5.47	$\frac{5}{2}^+$
1.840 $\pm$ 10	7.8 $\pm$ 1.4	5.699	$\frac{3}{2}^-$
2.30	190	6.13	$\frac{3}{2}^+$
2.37	120	6.20	$\frac{1}{2}^-$
2.445 $\pm$ 10	19.2 $\pm$ 2.4	6.272	$\frac{7}{2}^-$
$\approx$ 2.58		(6.40)	
(2.63)		(6.45)	

<sup>a</sup> See Table 19.4 in (1978AJ03) for references.

8.  $^{18}\text{O}(n, \alpha)^{15}\text{C}$

$$Q_m = -5.0097$$

$$E_b = 3.957$$

The total cross sections for the  $\alpha_0$  and  $\alpha_1$  groups have been measured for  $E_n = 7.5$  to 8.6 MeV: resonance structure is reported at  $E_n = 7.70 \pm 0.05$  and  $8.05 \pm 0.05$  MeV with  $\Gamma_{\text{lab}} = 0.25$  and 0.35 MeV, respectively [ $^{19}\text{O}^*(11.25, 11.58)$ ] (1967ST28).

9.  $^{18}\text{O}(d, p)^{19}\text{O}$

$$Q_m = 1.732$$

Angular distributions have been measured at  $E_d = 0.8$  to 15 MeV: see (1978AJ03) for the earlier work and (1979ST21;  $E_d = 8$  to 12 MeV). The  $l_n$  values and spectroscopic factors derived from these measurements are displayed in Table 19.4.

Branching ratios are shown in Table 19.3.  $^{19}\text{O}^*(0.096)$  has  $g = -0.48 \pm 0.06$ ; its configuration appears to be mainly  $d_{5/2}^3$  and  $B(\text{M1}) = (0.040 \pm 0.015) \mu_N^2$  (1976GO09). The  $\Delta E$  value for the  $1.47 \rightarrow 0.096$  transition is  $1375.3 \pm 0.5$  keV. Assuming  $E_x = 96.0 \pm 0.5$  keV (Table 19.1)  $E_x = 1471.4 \pm 0.7$  keV (1973WA10). Angular correlations are consistent with  $J^\pi = \frac{5}{2}^+$  for the ground state and unambiguously fix  $J^\pi = \frac{3}{2}^+$  and  $\frac{1}{2}^+$ , respectively, for  $^{19}\text{O}^*(0.096, 1.47)$  (1965AL13).

10. (a)  $^{18}\text{O}(^{13}\text{C}, ^{12}\text{C})^{19}\text{O}$   $Q_{\text{m}} = -0.989$   
 (b)  $^{18}\text{O}(^{14}\text{C}, ^{13}\text{C})^{19}\text{O}$   $Q_{\text{m}} = -4.219$   
 (c)  $^{18}\text{O}(^{17}\text{O}, ^{16}\text{O})^{19}\text{O}$   $Q_{\text{m}} = -0.186$   
 (d)  $^{18}\text{O}(^{18}\text{O}, ^{17}\text{O})^{19}\text{O}$   $Q_{\text{m}} = -4.087$

An angular distribution is reported for reaction (a) at  $E(^{18}\text{O}) = 31$  MeV to  $^{19}\text{O}_{\text{g.s.}+0.096}$  ([1978CH16](#)). At  $E(^{18}\text{O}) = 36$  MeV angular distributions for reaction (c) to  $^{19}\text{O}^*(0 + 0.096, 1.47)$  have been studied by ([1977KA1Y](#)). Differential cross sections are reported for  $E(^{18}\text{O}) = 30$  to 36 MeV for reaction (d) ([1977KA21](#)). See also ([1978AJ03](#)).

11.  $^{19}\text{N}(\beta^-)^{19}\text{O}$   $Q_{\text{m}} = 12.54$

The  $\beta$ -decay of  $^{19}\text{N}$  has not been observed.

12.  $^{19}\text{F}(\text{n}, \text{p})^{19}\text{O}$   $Q_{\text{m}} = -4.036$

Angular distributions have been reported at  $E_{\text{n}} = 14.1$  and 14.4 MeV for the  $p_0 \rightarrow p_2$  groups: see ([1972AJ02](#)) and  $^{20}\text{F}$ .

**<sup>19</sup>F**  
(Figs. 6 and 8)

GENERAL: (See also (1978AJ03).)

*Shell model:* (1978CH26, 1978DA1N, 1978MA2H, 1979DA15, 1980KU05, 1980MC1L, 1981ER03, 1981GR06, 1982KI02).

*Cluster, collective and rotational models:* (1977BU22, 1977FO1E, 1978BR21, 1978CH26, 1978PE09, 1978PI1E, 1978TA1A, 1978TH1A, 1978ZE07, 1979FO03, 1979MA27, 1979PE16, 1979SA41, 1979SA43, 1980FU1G).

*Electromagnetic transitions:* (1976MC1G, 1977BI1D, 1977CL03, 1977HE1L, 1977MA2P, 1978DE1K, 1978KR19, 1978PE09, 1978SC19, 1978ZE07, 1979MA27, 1979PE16, 1979SA41, 1979SA43, 1980BR09, 1980FU1G, 1980KU05, 1980MI1G).

*Special states:* (1977MA2P, 1978BR21, 1978MA2H, 1978PE09, 1978PI1E, 1978SC19, 1978TA1A, 1978ZE07, 1979DA15, 1979FO03, 1979LA10, 1979SA41, 1979SA43, 1980BR21, 1980FU1G, 1981ER03, 1982KI02).

*Astrophysical questions:* (1977TR1D, 1978DI1D, 1978ME1D, 1978OR1A, 1978WO1E, 1979CH1T, 1979LA1H, 1979RA1C, 1980CO1R, 1980GO1D).

*Applied topics:* (1979FO1F, 1979GR1E, 1980DE1U, 1980DI03, 1980KR1C, 1980MC1H).

*Complex reactions involving <sup>19</sup>F:* (1978CA1N, 1978GO1N, 1978OB01, 1978VO1D, 1978YO01, 1979BE31, 1979GA04, 1979GO11, 1979MO17, 1979SA26, 1979NA1F, 1980GR10, 1981CI03, 1981GR08, 1981NA1E, 1982SH1E).

*Muon and neutrino capture and reactions:* (1978LE04, 1978SC13, 1981MU1E).

*Pion capture and reactions:* (1977BE2P, 1977MA35, 1977ST27, 1978OL04, 1978SI1D, 1979BE31, 1979BO1N, 1979KI1G, 1979NA1F, 1980LI1J, 1980ST25, 1981BE63, 1981FR17, 1981LI1Z, 1981NI03, 1982BI1H).

*Kaon capture:* (1978AT01).

*Other topics:* (1977GR16, 1978AN15, 1978BR21, 1978DE1K, 1978MA2H, 1979BE1H, 1979BE2L, 1979CO09, 1979DE10, 1979DE18, 1979HE1F, 1979LA10, 1979MA27, 1979PE16, 1979SA41, 1979SA43, 1980BR21, 1980DE1F, 1980KO1U, 1980TA1L, 1981AD1E, 1981AR1D, 1981CA1H, 1981DU1D, 1981ER03, 1982KI02).

*Ground state of <sup>19</sup>F:* (1976MC1G, 1977MA35, 1977MA2P, 1977NO07, 1978AN07, 1978BR21, 1978CH26, 1978HE1D, 1978MA54, 1978ZA1D, 1978ZE07, 1979MA27, 1979SA41, 1979SA43, 1980BR09, 1980BR13, 1980HA41, 1981AR1D).

$$\mu_{\text{g.s.}} = +2.628866 \pm 0.000008 \text{ nm (1978LEZA);}$$

$$\mu_{0.197} = +3.607 \pm 0.008 \text{ nm (1978LEZA)};$$

$$Q_{0.197} = -0.12 \pm 0.02 \text{ b (1978LEZA)}.$$

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^+; \frac{1}{2}$	$\frac{1}{2}^+$	stable		2, 3, 6, 8, 12, 13, 15, 19, 20, 21, 23, 24, 26, 27, 28, 29, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66
$0.109894 \pm 0.0005$	$\frac{1}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 0.853 \pm 0.010 \text{ nsec}$	$\gamma$	13, 20, 21, 26, 29, 35, 36, 41, 43, 44, 48, 58, 62, 64, 65
$0.197143 \pm 0.004$	$\frac{5}{2}^+$	$\frac{1}{2}^+$	$128.8 \pm 1.5 \text{ nsec}$	$\gamma$	13, 15, 19, 20, 21, 26, 27, 29, 35, 36, 42, 43, 44, 45, 48, 50, 62, 64
$1.34567 \pm 0.13$	$\frac{5}{2}^-$	$\frac{1}{2}^-$	$4.4 \pm 0.6 \text{ psec}$	$\gamma$	13, 15, 19, 20, 21, 26, 29, 35, 36, 43, 44, 45, 48
$1.4587 \pm 0.3$	$\frac{3}{2}^-$	$\frac{1}{2}^-$	$90 \pm 20 \text{ fsec}$	$\gamma$	13, 15, 20, 21, 26, 29, 35, 41, 43, 44, 45, 48, 52, 62
$1.554038 \pm 0.009$	$\frac{3}{2}^+$	$\frac{1}{2}^+$	$5 \pm 3 \text{ fsec}$	$\gamma$	13, 19, 20, 21, 26, 27, 29, 34, 35, 36, 42, 43, 44, 45, 48, 50, 58, 62
$2.779849 \pm 0.034$	$\frac{9}{2}^+$	$\frac{1}{2}^+$	$280 \pm 30 \text{ fsec}$	$\gamma$	4, 5, 7, 10, 11, 13, 15, 17, 19, 20, 21, 24, 26, 27, 29, 34, 35, 42, 43, 44, 45, 48, 61, 62

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
$3.90817 \pm 0.20$	$\frac{3}{2}^+$	$\frac{3}{2}^+$	$9 \pm 5$ fsec	$\gamma$	20, 21, 26, 29, 35, 36, 41, 44, 48, 62
$3.9987 \pm 0.7$	$\frac{7}{2}^-$	$\frac{1}{2}^-$	$19 \pm 7$ fsec	$\gamma$	13, 20, 21, 26, 29, 34, 35, 44, 48, 62
$4.0325 \pm 1.2$	$\frac{9}{2}^-$	$\frac{1}{2}^-$	$67 \pm 15$ fsec	$\gamma$	13, 15, 17, 19, 20, 21, 26, 29, 34, 36, 44, 48, 62
$4.377700 \pm 0.042$	$\frac{7}{2}^+$	$\frac{3}{2}^+$	$< 11$ fsec	$\gamma$	13, 19, 20, 21, 26, 27, 29, 34, 35, 36, 42, 44, 48, 62
$4.5499 \pm 0.8$	$\frac{5}{2}^+$	$\frac{3}{2}^+$	$< 50$ fsec	$\gamma$	13, 20, 21, 26, 29, 44, 48, 62
$4.5561 \pm 0.5$	$\frac{3}{2}^-$		$17_{-8}^{+10}$ fsec	$\gamma$	20, 21, 34, 35, 44, 48, 62
$4.648 \pm 1$	$\frac{13}{2}^+$	$\frac{1}{2}^+$	$2.2 \pm 0.3$ psec	$\gamma$	13, 19, 20, 21, 24, 26, 27, 29, 42, 48, 62
$4.6825 \pm 0.7$	$\frac{5}{2}^-$		$15.4 \pm 3.0$ fsec	$\gamma, \alpha$	13, 20, 29, 34, 35, 44, 48, 62
$5.1066 \pm 0.9$	$\frac{5}{2}^+$		$< 30$ fsec	$\gamma, \alpha$	13, 20, 21, 26, 29, 34, 35, 44, 48, 62
$5.337 \pm 2$	$\frac{1}{2}^{(+)}$		$\leq 0.1$ fsec	$\gamma, \alpha$	13, 20, 21, 26, 29, 35, 44, 48, 62
$5.418 \pm 1$	$\frac{7}{2}^-$		$\leq 0.9$ fsec	$\gamma, \alpha$	13, 20, 26, 29, 35, 44, 48
$5.4635 \pm 1.5$	$\frac{7}{2}^+$	$\frac{1}{2}^+$	$\leq 0.26$ fsec	$\gamma, \alpha$	13, 15, 19, 20, 21, 26, 27, 29, 42, 44, 48
$5.5007 \pm 1.7$	$\frac{3}{2}^+$		$\Gamma = 4 \pm 1$ keV	$\gamma, \alpha$	13, 14, 21, 29, 44, 48
$5.535 \pm 2$	$\frac{5}{2}^+$			$\gamma, \alpha$	13, 26, 29, 44, 48, 62
$5.621 \pm 1$	$\frac{5}{2}^-$		$\tau_m < 1.3$ fsec	$\gamma, \alpha$	13, 26, 29, 34, 35, 44, 48, 61, 62

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
$5.938 \pm 1$	$\frac{1}{2}^+$			$\gamma, \alpha$	13, 29, 34, 35, 44, 62
$6.070 \pm 1$	$\frac{7}{2}^+$		$\Gamma = 1.2$	$\gamma, \alpha$	13, 14
$6.088 \pm 1$	$\frac{3}{2}^-$		4	$\gamma, \alpha$	13, 14, 15, 20, 21, 29
$6.100 \pm 2$	$\frac{9}{2}^-$			$\gamma$	29
$6.1606 \pm 0.9$	$\frac{7}{2}^-$			$\gamma, \alpha$	13, 29, 44, 62
$6.255 \pm 1$	$\frac{1}{2}^+$		8	$\gamma, \alpha$	14, 29, 34, 35, 44
$6.282 \pm 2$	$\frac{5}{2}^+$		2.4	$\gamma, \alpha$	13, 14, 19, 29, 34, 44
$6.330 \pm 2$	$\frac{7}{2}^+$		2.4	$\gamma, \alpha$	13, 14, 15, 44
$6.429 \pm 8$	$\frac{1}{2}^-$		280	$\alpha$	14
$6.4967 \pm 1.4$	$\frac{3}{2}^+$			$\gamma, \alpha$	13, 21, 29, 35
$6.5000 \pm 0.9$	$\frac{11}{2}^+$	$\frac{3}{2}^+$		$\gamma, \alpha$	13, 21, 27, 29
$6.5275 \pm 1.4$	$\frac{3}{2}^+$		4	$\gamma, \alpha$	13, 14, 19, 21, 29
$6.554 \pm 2$	$\frac{7}{2}^-$		1.6	$\gamma, \alpha$	13, 14
$6.592 \pm 2$	$\frac{9}{2}^+$	$\frac{3}{2}^+$		$\gamma, \alpha$	13, 19, 29, 35
$6.787 \pm 2$	$\frac{3}{2}^-$		2.4	$\gamma, \alpha$	13, 14, 29, 35, 62
$6.8384 \pm 0.9$	$\frac{5}{2}^+$		1.2	$\gamma, \alpha$	13, 14, 29
$6.891 \pm 4$	$\frac{3}{2}^-$		28	$\gamma, \alpha$	13, 14, 21
$6.9265 \pm 1.7$	$\frac{7}{2}^-$		2.4	$\gamma, \alpha$	13, 14, 15, 19, 20, 29, 35
$6.989 \pm 3$	$\frac{1}{2}^-$		51	$\alpha$	14, 29
$7.114 \pm 6$	$\frac{7}{2}^+ \text{ b}$		32	$\alpha$	14, 35
$7.1662 \pm 0.7$	$\frac{11}{2}^-$			$\gamma, \alpha$	13
$7.262 \pm 2$	$\frac{1}{2}^-, \frac{3}{2}^-$		$\lesssim 6$	$\alpha$	14, 19, 20, 21, 29, 34, 35
$7.364 \pm 4$	$\frac{1}{2}^+$			$\alpha$	14, 21, 34, 35
$7.5396 \pm 0.9$	$\frac{5}{2}^+, \frac{3}{2}^-$			$\gamma, \alpha$	13, 15, 19, 29, 35
$7.56 \pm 10$	$\frac{7}{2}^+$		$\lesssim 90$	$\alpha$	14
$7.6606 \pm 0.9$	$\frac{3}{2}^+, \frac{3}{2}^-$			$\gamma, \alpha$	13, 29, 35, 41, 63
$7.702 \pm 5$	$\frac{1}{2}^-$		$\lesssim 30$	$\alpha$	14, 19, 35

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
7.79			$\lesssim 6$	$\alpha$	14
7.90			$\lesssim 200$	$\alpha$	14
$7.929 \pm 3$	$\frac{7}{2}^+, \frac{9}{2}$			$\gamma, \alpha$	13, 14, 19, 21
$7.937 \pm 3$	$\frac{11}{2}^+$			$\gamma, \alpha$	13, 27
$8.015 \pm 2$	$\frac{5}{2}^+$				29, 35
$8.0838 \pm 2.6$			$\leq 3$	$\text{p}, \alpha$	33, 35
$8.1368 \pm 1.0$	$\frac{1}{2}^+$		$\leq 0.3$	$\gamma, \text{p}, \alpha$	14, 29, 33, 34, 35
(8.16)			$\lesssim 50$	$\alpha$	14
$8.1980 \pm 0.8$	$(\frac{5}{2}^+)$		$\leq 1$	$\gamma, \text{p}, \alpha$	29, 33, 35
$8.2535 \pm 2.6$	$(\frac{5}{2}^+)$		$\leq 1.5$	$\gamma, \text{p}$	29, 35
$8.288 \pm 2$	$\frac{13}{2}^-$	$(\frac{1}{2}^-)$	$< 1$	$\gamma, \alpha$	13, 15, 16, 17, 19, 20, 29
$8.310 \pm 1$	$\frac{5}{2}^+$		$0.047 \pm 0.019$	$\gamma, \text{p}, \alpha$	13, 29, 33, 35
$8.370 \pm 4$	$\frac{7}{2}, \frac{5}{2}^+$		$7.5 \pm 1.5$	$\gamma, \alpha$	13
$8.581 \pm 2$	$\frac{5}{2}^+$		$\leq 0.5$	$\gamma, \text{p}, \alpha$	13, 29
$8.5891 \pm 1.0$	$\frac{3}{2}^-$		$2.0 \pm 0.1$	$\gamma, \text{p}, \alpha$	13, 19, 29, 31, 33, 35
$8.629 \pm 4$	$\frac{7}{2}^-$		$< 1$	$\gamma, \alpha$	13
8.65	$\frac{1}{2}^+$		$\approx 300$	$\gamma, \text{p}, \alpha$	29, 31, 33
$8.793 \pm 2$	$\frac{1}{2}^+, \frac{3}{2}$		$46 \pm 2$	$\gamma, \text{p}$	29, 35
$8.864 \pm 4$	$< \frac{9}{2}$		$\approx 1$	$\gamma, \alpha$	13
$8.919 \pm 2$	$\frac{3}{2}$		$10 \pm 2$	$\gamma, \text{p}$	29
$8.9280 \pm 0.8$	$\frac{3}{2}^-$		$3.6 \pm 0.2$	$\text{p}, \alpha$	31, 33
$8.953 \pm 3$	$\frac{11}{2}^-$		$4.2 \pm 1$	$\gamma, \alpha$	13, 15, 16, 17, 19, 20, 31, 33
$9.030 \pm 5$	$\frac{5}{2}, \frac{7}{2}$			$\gamma, \alpha$	13
$9.0988 \pm 0.6$	$\frac{7}{2}^-$		$0.57 \pm 0.03$	$\gamma, \text{p}, \alpha$	13, 29, 31, 33
$9.101 \pm 4$	$\frac{7}{2}^+, \frac{9}{2}^+$		$\approx 1$	$\gamma, \alpha$	13, 35
$9.167 \pm 1.1$	$\frac{1}{2}^+$		$6.2 \pm 0.5$	$\gamma, \text{p}, \alpha$	13, 31, 33, 35
$9.204 \pm 7$	$\frac{3}{2}$		$10.2 \pm 1.5$	$\gamma, \alpha$	13
$9.267 \pm 4$	$\frac{11}{2}^+, \frac{9}{2}^+$		$2 \pm 1$	$\gamma, \alpha$	13
$9.280 \pm 5$	$\frac{7}{2}, \frac{9}{2}$		$< 1.5$	$\gamma, \alpha$	13



Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
9.318 $\pm$ 2	$\frac{3}{2}^+$		3.4 $\pm$ 0.7	$\gamma, p, \alpha$	13, 19, 29
9.321 $\pm$ 1.1	$\frac{1}{2}^+$		5.0 $\pm$ 0.2	$p, \alpha$	31, 33
9.329 $\pm$ 4	$< \frac{5}{2}$		$\approx 6$	$\gamma, \alpha$	13
9.509 $\pm$ 4	$\frac{5}{2}^+, \frac{7}{2}^+$		$< 1$	$\gamma, \alpha$	13
9.527 $\pm$ 6	$(\frac{5}{2})$		28	$p, \alpha$	31, 33
9.537 $\pm$ 2	$\frac{5}{2}^+$		6.3 $\pm$ 1.5	$\gamma, \alpha$	13, 21, 29, 31, 33
9.565 $\pm$ 3	$\frac{3}{2}^-$		26 $\pm$ 3	$\gamma, p$	29
9.574 $\pm$ 4	$\frac{3}{2}^-$		67 $\pm$ 3	$\gamma, p, \alpha$	29, 31, 33
9.586 $\pm$ 3	$\frac{7}{2}$		8.9 $\pm$ 1.2	$\gamma, p, \alpha$	13, 29, 35
9.642 $\pm$ 6	$\frac{3}{2}, \frac{5}{2}$		$\approx 8$	$\gamma, \alpha$	13
9.654 $\pm$ 6	$\frac{3}{2}, \frac{5}{2}$		$\approx 6$	$\gamma, \alpha$	13
9.6676 $\pm$ 1.3	$\frac{3}{2}^+$		3.6 $\pm$ 0.4	$\gamma, p, \alpha$	13, 29, 31, 32, 35
9.710 $\pm$ 4	$\frac{9}{2}^+, \frac{11}{2}^-$		$< 1$	$\gamma, \alpha$	13, 19
9.819 $\pm$ 0.8	$\frac{5}{2}^-$		0.3 $\pm$ 0.05	$\gamma, p, \alpha$	13, 29, 31, 33
9.834 $\pm$ 3	$\frac{11}{2} \rightarrow \frac{15}{2}$		$< 1$	$\gamma, \alpha$	13
9.8734 $\pm$ 1.7	$\frac{11}{2}^-$		$\lesssim 1.5$	$\gamma, p, \alpha$	13, 19, 20, 29
9.886 $\pm$ 3	$\frac{1}{2}^+$		25 $\pm$ 2	$\gamma, p, \alpha$	29, 31, 33
9.926 $\pm$ 3	$\frac{9}{2}^+$		$\approx 1$	$\gamma, \alpha$	13, 15
10.088 $\pm$ 5	$\frac{5}{2}^-, \frac{7}{2}^-$		$< 1.5$	$\gamma, \alpha$	13
10.136 $\pm$ 0.8	$\frac{3}{2}^-$		4.3 $\pm$ 0.6	$\gamma, p, \alpha$	13, 29, 33
10.161 $\pm$ 3	$\frac{1}{2}^+$		31	$p, \alpha$	31, 33
10.231 $\pm$ 3	$\frac{1}{2}^+$		$< 1$	$p, \alpha$	14, 31, 33
10.253 $\pm$ 3	$\frac{1}{2}^+$		22	$p, \alpha$	31, 33
10.308 $\pm$ 3	$\frac{3}{2}^+$		9.2	$p, \alpha$	14, 21, 31, 33
10.365 $\pm$ 4	$\frac{7}{2} \rightarrow \frac{11}{2}$		3 $\pm$ 1.5	$\gamma, \alpha$	13, 35
10.411 $\pm$ 3	$\frac{13}{2}^+$	$\frac{3}{2}^+$	$< 1.5$	$\gamma, \alpha$	13, 15, 19, 21, 29, 61
10.469 $\pm$ 4			11.0 $\pm$ 1.2	$p, \alpha$	14
10.488 $\pm$ 4			4.8 $\pm$ 0.8	$p, \alpha$	14
10.4964 $\pm$ 1.0	$\frac{3}{2}^+$		5.7 $\pm$ 0.6	$n, p, \alpha$	14, 30, 31, 33
10.521 $\pm$ 4			14 $\pm$ 2	$p, \alpha$	14, 35

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
10.5423 $\pm$ 1.1			2.5 $\pm$ 0.2	n, p, $\alpha$	14, 30
10.555 $\pm$ 3	$\frac{3}{2}^+; (\frac{3}{2})$		4.0 $\pm$ 1.2	p, $\alpha$	14, 31, 33
10.5656 $\pm$ 1.1			4.6 $\pm$ 0.7	n, p, $\alpha$	14, 30
10.580 $\pm$ 4	$(\frac{5}{2}^+)$		22 $\pm$ 3	p, $\alpha$	31, 33
10.613 $\pm$ 1.6	$\frac{5}{2}^+; \frac{3}{2}$		4.7 $\pm$ 0.5	n, p, $\alpha$	30, 31, 33
10.762 $\pm$ 3	$\frac{1}{2}^-$		6 $\pm$ 3	n, p, $\alpha$	19, 30, 31, 33
10.8588 $\pm$ 1.8	$\frac{5}{2}^+$		24.0 $\pm$ 1.5	n, p, $\alpha$	30, 31, 33
10.974 $\pm$ 3	$(\frac{3}{2}, \frac{5}{2})^+$		14 $\pm$ 2	n, p, $\alpha$	30, 31, 33
10.989 $\pm$ 2.5			7 $\pm$ 2	n, p	30
11.071 $\pm$ 2.5	$\frac{1}{2}^+$		35 $\pm$ 4	n, p, $\alpha$	30, 31, 33
11.187 $\pm$ 4	$(\frac{1}{2}^-)$		17 $\pm$ 4	n, p, $\alpha$	30, 31, 33
11.272 $\pm$ 3			7 $\pm$ 2	n, p	30
11.285 $\pm$ 8	$\frac{5}{2}^+$		22 $\pm$ 5	n, p, $\alpha$	30, 31, 33
11.35 $\pm$ 25	$\frac{1}{2}^+$		272 $\pm$ 31	p	31
11.451 $\pm$ 4	$\frac{1}{2}^-$		38 $\pm$ 7	n, p, ( $\alpha$ )	19, 30, 31, 33
11.478 $\pm$ 5			7 $\pm$ 3	n, p	30
11.502 $\pm$ 5	$(\frac{3}{2}^-)$		4 $\pm$ 2	n, p, $\alpha$	30, 31, 33
11.540 $\pm$ 8	$\frac{5}{2}^+$		22 $\pm$ 5	n, p, $\alpha$	30, 31, 33
11.568 $\pm$ 7	$(T = \frac{3}{2})$		15 $\pm$ 10	n, p	30
11.602 $\pm$ 12	$\frac{3}{2}^-$		63 $\pm$ 7	n, p	30, 31
11.652 $\pm$ 4	$\frac{3}{2}^+; (\frac{3}{2})$		33 $\pm$ 6	n, p, ( $\alpha$ )	15, 19, 30, 31, 33
11.84 $\pm$ 10			< 50	n, p	30
11.93 $\pm$ 10			90	n, p	30
12.04 $\pm$ 21	$\frac{1}{2}^-$		71 $\pm$ 24	p, $\alpha$	15, 31, 33
12.14 $\pm$ 10	$\frac{3}{2}^-; \frac{3}{2}$		105 $\pm$ 14	n, p, ( $\alpha$ )	30, 31, 33
12.221 $\pm$ 12	$\frac{3}{2}^+$		74 $\pm$ 1	n, p, $\alpha$	16, 17, 30, 31, 33
12.521 $\pm$ 7	$\frac{1}{2}^-$		15 $\pm$ 4	p	31
12.576 $\pm$ 10	$\frac{5}{2}^+$		48 $\pm$ 10	p, $\alpha$	31, 33
12.58 $\pm$ 25	$\frac{1}{2}^-; \frac{3}{2}$		285 $\pm$ 48	p	31
12.78 $\pm$ 10	$\frac{5}{2}^+; \frac{3}{2}$		95 $\pm$ 38	n, p, ( $\alpha$ )	19, 30, 31, 33
12.86 $\pm$ 30	$\frac{3}{2}^+; \frac{3}{2}$		276 $\pm$ 38	p	31

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
12.94 $\pm$ 25	$\frac{5}{2}^+$		71 $\pm$ 24	p, $\alpha$	31, 33
12.98 $\pm$ 50	$\frac{1}{2}^-$		124 $\pm$ 38	p	31
13.068 $\pm$ 4	$\frac{1}{2}^+$		$\leq 10$	n, p, t	18, 20
13.09 $\pm$ 75	$\frac{3}{2}^-$		285 $\pm$ 71	p	31
13.17 $\pm$ 15			70	n, p	30
13.245 $\pm$ 10	$\frac{1}{2}^-$		7	t	18
13.270 $\pm$ 10	$\frac{1}{2}^+$		4.5	t	18
13.317 $\pm$ 8	$\frac{7}{2}^-; (\frac{3}{2})$		28 $\pm$ 6	n, p, $\alpha$	30, 31, 33
13.36 $\pm$ 25	$\frac{3}{2}^-$		38 $\pm$ 19	p	21, 31
13.532 $\pm$ 10	$\frac{1}{2}^+$		22	t	18
13.731 $\pm$ 11	$\frac{7}{2}^-; \frac{3}{2}$		52 $\pm$ 10	n, p, ( $\alpha$ )	15, 20, 30, 31, 33
13.878 $\pm$ 15	$\frac{1}{2}^+$		101	t	18
14.04 $\pm$ 20	$\frac{5}{2}^+$		141 $\pm$ 28	p	31
14.10 $\pm$ 21	$\frac{3}{2}^-$		84 $\pm$ 28	p	15, 20, 31
14.147 $\pm$ 20	$\frac{1}{2}^+$		21	t	18
14.24 $\pm$ 15			350	n, p	30
14.255 $\pm$ 15	$\frac{3}{2}^+$		51	t	18
14.32 $\pm$ 20	$\frac{3}{2}^-$		76 $\pm$ 28	p	21, 31
14.352 $\pm$ 10	$\frac{1}{2}^+$		154	t	18
14.46 $\pm$ 25	$\frac{3}{2}^+$		179	t	18
14.46 $\pm$ 25	$\frac{5}{2}^+$		46	t	18
14.70 $\pm$ 20	$\frac{3}{2}^-$		124 $\pm$ 38	p	31
14.72 $\pm$ 70	$\frac{1}{2}^-$		257 $\pm$ 67	$\alpha$	33
14.74 $\pm$ 50	$\frac{1}{2}^+$		361 $\pm$ 67	p, $\alpha$	31, 33
14.78 $\pm$ 20	$\frac{5}{2}^+$			n, p	30, 31
14.92 $\pm$ 30	$\frac{7}{2}^-$			p	15, 16, 20, 31
15.00 $\pm$ 20				n, p	30
15.35 $\pm$ 20	$\frac{1}{2}^-$			p	31
15.40 $\pm$ 30	$\frac{5}{2}^+$			p	31
15.56 $\pm$ 30					20
15.77 $\pm$ 21	$\frac{3}{2}^-$		150	n, p	30, 31

Table 19.6: Energy levels of  $^{19}\text{F}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau_m$ or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
16.09 $\pm$ 50					15
16.20 $\pm$ 40	$\frac{3}{2}^+$			p	30
16.23 $\pm$ 30	$\frac{7}{2}^-$			p	31
16.27 $\pm$ 20	$\frac{3}{2}^-$		200	n, p	31
16.45 $\pm$ 50					15
16.80 $\pm$ 30				n, p	30
17.05 $\pm$ 40	$\frac{3}{2}^-$		$331 \pm 67$	p	31
17.16 $\pm$ 40	$\frac{7}{2}^-$		$323 \pm 67$	p	31
17.45 $\pm$ 30	$\frac{3}{2}^-$		$32 \pm 19$	p	15, 16, 31
17.65 $\pm$ 60	$\frac{7}{2}^-$		$95 \pm 57$	p	31
17.93 $\pm$ 40	$\frac{3}{2}^-$		$255 \pm 57$	p	31
18.02 $\pm$ 60	$\frac{7}{2}^-$		$365 \pm 57$	p	31
18.2 $\pm$ 50					15
18.92 $\pm$ 30					15, 20
19.07 $\pm$ 60	$\frac{3}{2}^-$		$555 \pm 143$	p	31
19.83 $\pm$ 150	$\frac{5}{2}^-$		$369 \pm 57$	p	31
19.87 $\pm$ 40	$\frac{3}{2}^-$		$473 \pm 57$	p	31
19.93 $\pm$ 50					15
20.81 $\pm$ 50	$\frac{1}{2}^-$		$412 \pm 57$	p	31
20.93 $\pm$ 50	$\frac{3}{2}^-$		$317 \pm 48$	p	31
21.05 $\pm$ 40	$\frac{7}{2}^-$		$448 \pm 29$	p	31

<sup>a</sup> See also Tables 19.7 and 19.8.

<sup>b</sup> See also (1979FO03).

1. (a)  $^9\text{Be}(^{10}\text{B}, 2\alpha)^{11}\text{B}$   $Q_m = 9.882$
- (b)  $^9\text{Be}(^{10}\text{B}, \alpha n)^{14}\text{N}$   $Q_m = 10.040$

The total reaction cross section has been measured for  $E(^{10}\text{B}) = 2.20$  to 10.43 MeV (1979CH22).

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$ 

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$
0.110	$\frac{1}{2}^-$	0	100 <sup>a</sup>	
0.197	$\frac{5}{2}^+$	0	100 <sup>a</sup>	
		0.110	< 0.06	
1.35	$\frac{5}{2}^-$	0.110	$96.8 \pm 1$	$0.0 \pm 0.7$
		0.197	$3.2 \pm 1$	
1.46 <sup>b</sup>	$\frac{3}{2}^-$	0	$20.5 \pm 0.7$ <sup>c</sup>	$0.01 \pm 0.03$ <sup>d</sup>
		0.110	$68.8 \pm 0.9$ <sup>c</sup>	$0.248 \pm 0.020$ <sup>d</sup>
		0.197	$10.7 \pm 0.5$ <sup>c</sup>	
1.55 <sup>b</sup>	$\frac{3}{2}^+$	0	$2.55 \pm 0.10$ <sup>a</sup>	
		0.110	$4.85 \pm 0.12$ <sup>a</sup>	
		0.197	$92.6 \pm 0.2$ <sup>a</sup>	
2.78 <sup>b</sup>	$\frac{9}{2}^+$	0.197	100 <sup>a</sup>	
3.91 <sup>b,e</sup>	$\frac{3}{2}^+$	0	$48 \pm 2$ <sup>a</sup>	
		0.110	$17 \pm 2$ <sup>a</sup>	
		0.197	$14 \pm 2$ <sup>a</sup>	
		1.55	$21 \pm 3$ <sup>a</sup>	
4.00 <sup>b</sup>	$\frac{7}{2}^-$	0.197	$18 \pm 4$	
		1.35	$70 \pm 4$	
		1.46	$12 \pm 6$	
4.03 <sup>b</sup>	$\frac{9}{2}^-$	1.35	100	
4.38 <sup>f,g</sup>	$\frac{7}{2}^+$	0	< 5	
		0.110	< 2	
		0.197	$80.5 \pm 2.0$ <sup>a</sup>	$0.155 \pm 0.022$
		2.78	$19.5 \pm 1.0$ <sup>a</sup>	$-0.16 \pm 0.07$
4.55 <sup>b,h,i</sup>	$\frac{5}{2}^+$	0.197	$69 \pm 7$	
		1.35	$5 \pm 3$	
		1.46	$8 \pm 3$	
		1.55	$18 \pm 4$	
4.56 <sup>b</sup>	$\frac{3}{2}^-$	0	$36 \pm 4$	
		0.110	$45 \pm 5$	
		0.197	$9 \pm 3$	
		1.35	$4 \pm 3$	
		1.46	< 4	
		1.55	$6 \pm 3$	
4.65	$\frac{13}{2}^+$	2.78	100	$ M ^2 = 5.5 \pm 1.8 \text{ W.u.}$
4.68 <sup>b,c,j</sup>	$\frac{5}{2}^-$	0.197	$5.6 \pm 0.9$	$0 < \delta < 2.0$
		1.35	$63.1 \pm 3.8$	$-0.22^{+0.14}_{-0.24}$
		1.46	$31.3 \pm 2.2$	$0.0 \pm 0.24 \text{ or } 2.0^{+1.5}_{-0.6}$

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$	
5.11	$\frac{5}{2}^+$	0.197	80	$\Gamma_\gamma/\Gamma = 0.83 \pm 0.10$	
		1.46	20		
5.34	$\frac{1}{2}^{(+)}$	0	$37 \pm 4$		
		0.110	$42 \pm 4$		
		1.46	$20 \pm 2$		
5.42	$\frac{7}{2}^-$	1.35	70		
		1.46	13		
		4.00	10		
		4.03	6		
5.46	$\frac{7}{2}^+$	0.197	4		
		1.35	32		
		1.55	5		
		2.78	59		
5.50	$\frac{3}{2}^+$	0.110	25		
		0.197	49		
		1.35	16		
		1.55	11		
5.54	$\frac{5}{2}^+$	0	7		
		0.197	47		
		1.46	45		
5.62 <sup>1</sup>	$\frac{3}{2}^-$	0.197	$39 \pm 4$		
		1.35	$61 \pm 4$		
5.94	$\frac{1}{2}^+$	0	$7 \pm 4$		
		0.110	$20 \pm 6$		
		0.197	$2 \pm 1$		
		1.46	$63 \pm 6$		$0.25 \pm 0.02$
		1.55	$< 2$		
6.07 <sup>m</sup>	$\frac{7}{2}^+$	3.91	$8 \pm 3$	$0.28 \pm 0.09$	
		0.197	$54 \pm 5$	$-0.26 \pm 0.02$	
		1.35	$19 \pm 2$		
		1.55	$1_{-0.5}^{+1}$	$0.035 \pm 0.023$	
		2.78	$23 \pm 3$	$0.06 \pm 0.08$	
		4.38	$4 \pm 1$		
6.09 <sup>n</sup>	$\frac{3}{2}^-$	0	$25 \pm 4$	$-0.021 \pm 0.014$	
		0.110	$61 \pm 5$	$0.045 \pm 0.021$	
		0.197	$14 \pm 3$	$0.014 \pm 0.043$	
6.16 <sup>o</sup>	$\frac{7}{2}^-$	0.197	$31 \pm 3$	$-0.045 \pm 0.025$	
		1.35	$65 \pm 4$	$0.077 \pm 0.007$	

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$	
6.28	$\frac{5}{2}^+$	1.46	$1.3 \pm 0.6$	$-0.05 \pm 0.07$	
		4.00	$1.6 \pm 0.6$		
		4.03	$2.3 \pm 0.3$		
		0	$14 \pm 2$		
		0.197	$4.2 \pm 1.0$		
		1.35	$36 \pm 2$		$-0.01 \pm 0.09$
		1.46	$26 \pm 2$		$-0.02 \pm 0.04$
6.33	$\frac{7}{2}^+$	1.55	$20 \pm 2$	$0.11 \pm 0.06$	
		0.197	$56 \pm 3$	$-0.27 \pm 0.24$	
		1.35	$17 \pm 2$	$-0.02 \pm 0.03$	
		1.55	$8.5 \pm 1.5$	$0.00 \pm 0.14$	
6.497	$\frac{3}{2}^+$	4.38	$18 \pm 2$	$0.04 \pm 0.20$	
		0	$38 \pm 2$	$-0.06 \pm 0.04$ or $2.00 \pm 0.17$	
		0.110	$14 \pm 2$	$0.00 \pm 0.03$	
		0.197	$9 \pm 2$	$0.3 \rightarrow 1.8$	
		1.35	$14 \pm 2$	$-0.11 \pm 0.09$	
		1.46	$25 \pm 2$	$0.00 \pm 0.07$	
6.500 <sup>p</sup>	$\frac{11}{2}^+$	2.78	55		
		4.65	45		
		0	$29 \pm 2$	$0.32 \pm 0.04$ or $0.90 \pm 0.06$	
6.53	$\frac{3}{2}^+$	0.110	$59 \pm 3$	$0.00 \pm 0.02$	
		4.55	$12 \pm 2$	$-0.23 \pm 0.13$	
		0.197	$19 \pm 2$	$0.03 \pm 0.05$	
6.55	$\frac{7}{2}$	1.35	$55 \pm 4$	$0.01 \pm 0.03$	
		2.78	$26 \pm 3$	$0.05 \pm 0.07$	
		0.197	$13 \pm 2$	$-0.13 \pm 0.13$	
6.59 <sup>q</sup>	$\frac{9}{2}^+$	2.78	$63 \pm 3$	$-0.20 \pm 0.20$	
		4.38	$24 \pm 2$	$0.02 \pm 0.07$	
		0	$15 \pm 2$	$-0.08 \pm 0.03$	
6.79	$\frac{3}{2}^-$	0.110	$39 \pm 2$	$0.11 \pm 0.02$	
		0.197	$13 \pm 2$	$0.05 \pm 0.06$	
		1.35	$5.3 \pm 0.8$		
		1.46	$25 \pm 2$	$-0.13 \pm 0.08$	
		3.91	$2.6 \pm 1.0$		
		0	$9 \pm 5$		
6.84	$\frac{5}{2}^+$	0.110	$9 \pm 5$		
		0.197	$27 \pm 6$	$-0.5 \pm 0.5$	
		1.35	$10 \pm 7$		
		0.197	$27 \pm 6$		
		1.35	$10 \pm 7$		

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$
6.89 <sup>r</sup>	$\frac{3}{2}^-$	1.46	$45 \pm 8$	$-0.02 \pm 0.11$
		0	$9 \pm 2$	
		1.35	$61 \pm 5$	$0.22 \rightarrow 2.2$
6.93	$\frac{7}{2}^-$	1.46	$30 \pm 5$	$0.15 \pm 0.12$
		0.197	$73 \pm 3$	$-0.01 \pm 0.03$
		1.35	$22 \pm 2$	$0.01 \pm 0.02$
		2.78	$2.4 \pm 0.5$	$0.00 \pm 0.16$
		4.00	$1.3 \pm 0.5$	
7.17	$\frac{11}{2}^-$	4.03	$1.3 \pm 0.5$	
		4.00	$5 \pm 1$	
		4.03	$91 \pm 1$	
7.54	$\frac{5}{2}^+; T = \frac{3}{2}$	4.65	$4 \pm 1$	
		0.197	$29 \pm 3$	$0.09 \pm 0.04$
		1.35	$1.2 \pm 0.4$	
		1.55	$41 \pm 3$	$0.017 \pm 0.015$
		4.38	$27 \pm 3$	$0.042 \pm 0.030$
7.66 <sup>b,t</sup>	$\frac{3}{2}^+; T = \frac{3}{2}$	5.11	$1.7 \pm 0.4$	
		0	$38 \pm 4$	$0.06 \pm 0.02$
		0.197	$13 \pm 2$	$0.06 \pm 0.07$ or $3.5 \pm 1.1$
		1.55	$36 \pm 2$	$0.06 \pm 0.04$
		3.91	$(3_{-2}^{+3})$	
		4.55	$5.1 \pm 0.3$	$-0.11 \pm 0.13$
		5.11	$5.9 \pm 0.5$	$-0.04 \pm 0.16$
7.93	$\frac{7}{2}^+, \frac{9}{2}$	0.197	4	
		2.78	96	
7.94 <sup>u</sup>	$\frac{11}{2}^+$	2.78	10	
		4.65	90	
8.14 <sup>v</sup>	$\frac{1}{2}^+$	0	$8 \pm 1$	
		0.11	$24 \pm 2$	
		0.197	$8 \pm 1$	
		1.55	$2 \pm 1$	
		3.91	$54 \pm 2$	$\Gamma_\gamma = 1.3 \text{ eV}$
		5.94	$1.0 \pm 0.5$	
		6.26	$3 \pm 1$	
8.25 <sup>v</sup>	$(\frac{5}{2}^+)$	0.197	$18 \pm 7$	
		1.35	$33 \pm 10$	
		1.46	$24 \pm 8$	
		3.91	$25 \pm 8$	



Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$
8.29 <sup>w</sup>	$\frac{13}{2}^-$	4.03	$93 \pm 4$	$\Gamma_\gamma = 72 \pm 8 \text{ meV}$
8.31 <sup>x</sup>	$\frac{5}{2}^+$	4.65	$7 \pm 4$	$\Gamma_\gamma = 0.71 \pm 0.17 \text{ eV}$
		0	$9 \pm 3$	
		1.55	$12 \pm 1^A$	
		4.38	$48 \pm 6$	
		4.38	$48 \pm 2^A$	
8.37 <sup>w</sup>	$\frac{7}{2}, \frac{5}{2}^+$	4.38	$43 \pm 6$	$\delta = 0.02 \pm 0.05 \text{ or } 2.2 \pm 0.6$
		0.197	$40 \pm 2^A$	
		1.35	$13 \pm 2$	
		2.78	$39 \pm 3$	
8.58	$\frac{5}{2}^+$	2.78	$30 \pm 3$	$\delta = -0.14 \pm 0.07$
		4.00	$18 \pm 3$	
		0	$4 \pm 1^A$	
		0.197	$44 \pm 2$	
		0.197	$38 \pm 5^A$	
		1.35	$24 \pm 2$	
		1.35	$23 \pm 3^A$	
		1.55	$20 \pm 2$	
		1.55	$20 \pm 3^A$	
		4.00	$4 \pm 1^w$	
		4.55	$2.0 \pm 0.7^A$	
		5.42	$6 \pm 1$	
		5.42	$4 \pm 1^A$	
		5.46	$2.0 \pm 0.5^A$	
		5.62	$2.2 \pm 0.5^A$	
5.94	$1.8 \pm 0.5^A$			
6.16	$2.5 \pm 0.5^A$			
6.93	$0.5 \pm 0.3^A$			
8.59 <sup>b,x</sup>	$\frac{3}{2}^-$	0	$5 \pm 2^A$	$\Gamma_\gamma = 0.85 \pm 0.17 \text{ eV}$
		0.11	$3 \pm 1^A$	
		0.197	$59 \pm 2$	
		0.197	$42 \pm 2^A$	
		1.35	$7 \pm 1^A$	
		1.55	$34 \pm 2$	
		1.55	$28 \pm 3^A$	
		3.91	$7 \pm 1$	
		3.91	$8 \pm 1^A$	
		4.55	$3.6 \pm 0.6^A$	

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$			
8.63 <sup>w</sup>	$\frac{7}{2}^-$	5.11	$1.0 \pm 0.5^A$				
		5.50	$1.5 \pm 0.5^A$				
		6.28	$0.6 \pm 0.2^A$				
		6.79	$0.3 \pm 0.1^A$				
		0.197	$34 \pm 2$				
		1.35	$6 \pm 1$				
		1.46	$6 \pm 1$				
		2.78	$38 \pm 2$				
		4.00	$13 \pm 1$				
		4.03	$3 \pm 1$				
8.65 <sup>v</sup>	$\frac{1}{2}^+$	0.11	$53 \pm 6$				
		1.46	$23 \pm 6$				
		3.91	$24 \pm 6$				
8.79 <sup>b,v</sup>	$\frac{1}{2}^+; T = \frac{3}{2}$	0	$1.2 \pm 0.4$				
		0.11	$30 \pm 1$				
		0.197	$0.3 \pm 0.2$				
		1.46	$22 \pm 1$				
		1.55	$8 \pm 1$				
		3.91	$22 \pm 1$				
		5.34	$0.5 \pm 0.1$				
		5.94	$1.8 \pm 0.2$				
		6.09	$1.7 \pm 0.2$				
		6.26	$0.2 \pm 0.1$				
		6.49	$6 \pm 1$				
		6.53	$2.1 \pm 0.2$				
		6.79	$1.2 \pm 0.3$				
		6.99	$0.5 \pm 0.1$				
		7.26	$1.7 \pm 0.2$				
		7.36	$0.6 \pm 0.1$				
		7.66	$0.2 \pm 0.1$				
		8.86 <sup>w</sup>	$\frac{3}{2}$		1.35	100	
					8.92 <sup>v</sup>	$\frac{3}{2}$	
		8.92 <sup>v</sup>	$\frac{3}{2}$		0.11	$10 \pm 2$	$0.20 \pm 0.04$ or $2.9 \pm 0.4$
0.197	$24 \pm 7$			$1.0 \pm 0.8$			
1.46	$25 \pm 7$			$3.0 \pm 2.5$			
1.55	$23 \pm 7$			$0.30 \pm 0.06$ or $\infty$			
3.91	$13 \pm 7$						
8.95 <sup>w</sup>	$\frac{11}{2}^-$			2.78	$50 \pm 2$	$\Gamma_\gamma(\text{tot}) = 230 \pm 30$ meV	

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$	
9.03 <sup>w</sup>	$\frac{5}{2}, \frac{7}{2}$	4.00	$26 \pm 2$	$\delta = 0.0 \pm 0.2$ or $2.5 \pm 0.6$ $-0.1 \pm 0.3$ or $\infty$  $-0.09 \pm 0.10$ $0.3 \pm 0.3$ or $-2.2 \pm 0.9$  $-0.08 \pm 0.01$ or $\infty$ $-0.09 \pm 0.34$ or $\infty$ $0.0 \pm 0.2$ or $3.0 \pm 1.6$  $0.25 \pm 0.10$ or $-6.0 \pm 5.5$ $0.1 \pm 0.3$ $0.17 \pm 0.10$ $0.0 \pm 0.3$	
		4.03	$9 \pm 1$		
		4.65	$10 \pm 2$		
		5.42	$5 \pm 1$		
		0.197	$44 \pm 5$		
		4.38	$30 \pm 5$		
9.098 <sup>x</sup>	$\frac{7}{2}^-$	6.07	$26 \pm 4$		
		0.197	$2.0 \pm 0.3^A$		
		1.35	$2.7 \pm 0.3^A$		
		2.78	$71 \pm 2$		
			$47 \pm 2^A$		
		4.00	$2.5 \pm 0.3^A$		
		4.03	$9 \pm 1$		
			$7.0 \pm 0.5^A$		
		4.68	$2.0 \pm 0.3^A$		
		5.11	$1.2 \pm 0.2^A$		
		5.42	$20 \pm 2$		
			$19 \pm 2^A$		
9.101 <sup>w</sup>	$\frac{7}{2}^+, \frac{9}{2}^+$	5.54	$1.3 \pm 0.7^A$		
		5.62	$3.3 \pm 0.3^A$		
		6.10	$12 \pm 1^A$		
		2.78	$11 \pm 2$		
		4.00	$24 \pm 2$		
		4.38	$24 \pm 2$		
		6.07	$15 \pm 2$		
		6.33	$10 \pm 2$		
		9.17 <sup>w</sup>	$\frac{1}{2}^+$	0.197	$51 \pm 2$
				1.55	$30 \pm 2$
				4.56	$19 \pm 2$
		9.20 <sup>w</sup>	$\frac{3}{2}$	0	$18 \pm 2$
0.110	$46 \pm 3$				
0.197	$10 \pm 4$				
1.35	$26 \pm 3$				
9.27 <sup>w</sup>	$\frac{11}{2}^+, \frac{9}{2}^+$	2.78	$27 \pm 2$		
		4.38	$18 \pm 2$		
		4.65	$55 \pm 3$		
9.28 <sup>w</sup>	$\frac{7}{2}, \frac{9}{2}$	4.00	$58 \pm 3$		
		4.03	$42 \pm 3$		

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$		
9.32 <sup>b,x</sup>	$\frac{1}{2}^+$	0	$29 \pm 2$ $30 \pm 1^A$	$0.10 \pm 0.08$ or $1.4 \pm 0.3$		
		0.197	$9 \pm 1$ $12 \pm 1^A$	$0.1 \pm 0.4$ or $\geq 0.6$		
		1.46	$41 \pm 3$ $28 \pm 1^A$	$0.1 \pm 0.2$		
		1.55	$21 \pm 3$ $17 \pm 1^A$	$-0.2 \pm 0.3$ or $\leq 0.9$		
		3.91	$3.0 \pm 0.3^A$	$0.40 \pm 0.05$ or $\geq 2.3$		
		4.56	$3.2 \pm 0.3^A$	$0.2 \pm 0.3$		
		4.68	$6.8 \pm 0.5^A$	$0.1 \pm 0.2$		
		9.33 <sup>w</sup>	$< \frac{5}{2}^+$	1.55	100	
		9.51 <sup>w</sup>		$\frac{5}{2}^+, \frac{7}{2}^+$	1.35	$14 \pm 2$
		9.54 <sup>v</sup>	$\frac{5}{2}^+$	1.55	$14 \pm 2$	
2.78	$72 \pm 3$					
1.35	$100^w$ $26 \pm 2^A$			$0.3 \pm 1.1$		
4.56	$15 \pm 1$			$0.7 \pm 0.4$		
4.68	$12 \pm 1$			$0.3 \pm 0.3$		
5.11	$29 \pm 2$			$0.3 \pm 0.2$		
7.54	$10 \pm 1$			$0.7 \pm 0.3$		
7.66	$6 \pm 1$			$0.4 \pm 0.3$ or $1.0 \pm 0.4$		
9.565 <sup>v</sup>	$\frac{3}{2}^-$	8.02	$2 \pm 1$			
9.574 <sup>v</sup>	$\frac{3}{2}^-$	0.197	$77 \pm 10$			
		6.26	$23 \pm 6$			
9.59 <sup>x</sup>	$\frac{7}{2}$	1.46	$26 \pm 2$	$-0.1 \pm 0.2$		
		3.91	$4 \pm 1$	$-6 \pm 7$		
		4.55	$17 \pm 2$			
		6.09	$38 \pm 2$	$1.8 \pm 1.0$		
		7.54	$11 \pm 2$	$-0.3 \pm 0.8$		
		7.66	$4 \pm 1$	$-0.1 \pm 1.3$		
9.59 <sup>x</sup>	$\frac{7}{2}$	0.197	$24 \pm 2^w$			
		1.35	$17 \pm 2$ $32 \pm 4^A$	$0.0 \pm 0.5$ or $3.7 \pm 2.5$		
		2.78	$33 \pm 3$ $30 \pm 2^A$	$0.1 \pm 0.2$ or $11 \pm 5$		
		4.00	$15 \pm 2$ $17 \pm 2^A$	$-0.7 \pm 1.1$		

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$		
9.64 <sup>w</sup>	$\frac{3}{2}, \frac{5}{2}$	4.03	$11 \pm 1^w$			
		4.55	$21 \pm 2^A$			
		0.197	$13 \pm 3$			
		1.35	$61 \pm 7$			
		4.55	$26 \pm 6$			
9.65 <sup>w</sup>	$\frac{3}{2}, \frac{5}{2}$	1.35	$41 \pm 9$			
		1.55	$59 \pm 9$			
		0	$34 \pm 5$			
9.67 <sup>b,x</sup>	$\frac{3}{2}^+$	0	$22 \pm 2^A$	$-0.72 \pm 0.04$ or $-10 \pm 4$		
		0.11	$36 \pm 5$			
			$20 \pm 2^A$	$0.00 \pm 0.05$		
		0.197	$9 \pm 1^A$	$0.30 \pm 0.03$ or $1.7 \pm 0.3$		
		1.35	$9 \pm 1^A$	$0.00 \pm 0.03$		
		1.46	$5 \pm 1^A$	$0.00 \pm 0.07$		
		1.55	$30 \pm 6$			
			$10 \pm 1^A$	$0.00 \pm 0.06$ or $-4.2 \pm 1.3$		
		3.91	$5.5 \pm 0.5^A$	$0.12 \pm 0.03$ or $-7.5 \pm 2.0$		
		4.38	$0.5 \pm 0.2^A$			
		4.55	$8 \pm 1^A$	$0.00 \pm 0.03$ or $4.7 \pm 0.5$		
		5.11	$1.5 \pm 0.3^A$	$0.00 \pm 0.05$		
		5.34	$1.0 \pm 0.2^A$	$-0.22 \pm 0.03$ or $3.3 \pm 0.2$		
		6.84	$1.0 \pm 0.3^A$	$0.05 \pm 0.02$ or $3.3 \pm 0.2$		
		7.54	$4.0 \pm 0.3^A$	$0.02 \pm 0.03$		
		7.66	$3.5 \pm 0.3^A$	$0.14 \pm 0.04$		
		9.71 <sup>w</sup>	$\frac{9}{2}^+, \frac{11}{2}^-$	2.78	$19 \pm 3$	
				4.03	$80 \pm 4$	
				4.65	$1 \pm 1$	
				0.11	$0.7 \pm 0.2^A$	
0.197	$41 \pm 2$					
9.82 <sup>b,x</sup>	$\frac{5}{2}^-$		$41 \pm 2^A$	$0.00 \pm 0.05$		
		1.35	$2.4 \pm 0.5^A$	$-0.6 \pm 0.2$		
		1.46	$10 \pm 1$			
			$8 \pm 1^A$	$-0.07 \pm 0.05$ or $2.7 \pm 0.7$		
		1.55	$34 \pm 2$			
			$30 \pm 2^A$	$0.01 \pm 0.04$		
		4.00	$1.0 \pm 0.2^A$	$0.0 \pm 0.2$ or $\infty$		
		4.55	$0.5 \pm 0.1^A$	$0.30 \pm 0.15$		
		4.68	$4 \pm 1$			

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$
9.83 <sup>w</sup> 9.87 <sup>x</sup>	$\frac{11}{2} \rightarrow \frac{15}{2}$ $\frac{11}{2}^-$		$4.8 \pm 0.3^A$	$0.0 \pm 0.1$ or $-1.7 \pm 0.4$
		5.11	$0.3 \pm 0.2^A$	$0.4 \pm 0.5$ or $\infty$
		5.42	$11 \pm 1$	
			$10 \pm 1^A$	$-0.04 \pm 0.05$ or $\infty$
		5.54	$0.6 \pm 0.2^A$	$0.0 \pm 0.2$
		5.62	$0.7 \pm 0.2^A$	$0.33 \pm 0.15$ or $-3.4 \pm 1.2$
		4.65	100	
		2.78	$68 \pm 4$	
			$63 \pm 3^A$	$0.0 \pm 0.2$
		4.00	$5 \pm 1$	
			$4.2 \pm 1.0^A$	
		4.03	$24 \pm 3$	
			$24 \pm 2^A$	$-0.43 \pm 0.05$ or $2.2 \pm 0.2$
		4.65	$3 \pm 1$	
			$2.1 \pm 0.8^A$	
9.89 <sup>v</sup>	$\frac{1}{2}^+$	6.10	$3.8 \pm 0.8^A$	$0.2 \pm 0.1$ or $2.7 \pm 1.0$
		6.50	$1.9 \pm 0.7^A$	$-0.4 \pm 0.7$
		8.29	$1.0 \pm 0.3^A$	
		0.197	$15 \pm 8$	
		1.46	$15 \pm 5$	
		3.91	$32 \pm 2$	
		5.94	$4 \pm 1$	
		6.09	$13 \pm 3$	
		6.53	$16 \pm 2$	
		7.66	$5 \pm 1$	
9.93 <sup>w</sup>	$\frac{9}{2}^+$	0.197	$1 \pm 1$	
		2.78	$19 \pm 1$	
		5.46	$10 \pm 1$	
		6.07	$7 \pm 1$	
		6.33	$8 \pm 1$	
		6.50	$54 \pm 2$	
		10.09 <sup>w</sup>	$\frac{5}{2}^-, \frac{7}{2}^-$	0.197
1.35	$35 \pm 2$			
4.00	$19 \pm 2$			
5.42	$26 \pm 2$			
6.07	$10 \pm 1$			
10.14 <sup>w</sup>	$\frac{3}{2}^-$	1.35	$29 \pm 4$	
		1.46	$71 \pm 4$	

Table 19.7: Radiative transitions in  $^{19}\text{F}^\dagger$  (continued)

$E_i$ (MeV)	$J_i^\pi$	$E_f$ (MeV)	Branching ratio (%)	$\delta$
10.37 <sup>w</sup>	$\frac{7}{2} \rightarrow \frac{11}{2}$	4.03	100	
10.41 <sup>w</sup>	$\frac{13}{2}^+$	2.78	$3 \pm 1$	
		4.68	$88 \pm 1$	
		6.50	$9 \pm 1$	

A = adopted.

<sup>†</sup> For references see Table 19.7 in (1978AJ03). See also Tables 19.9, 19.15 and 19.16.

<sup>a</sup> (1981OL1E).

<sup>b</sup> See also (1978DI13).

<sup>c</sup> Revised to sum to 100%: see (1978AJ03).

<sup>d</sup> (1980DI12).

<sup>e</sup> Transitions to  $^{19}\text{F}^*(1.35, 1.46, 2.78)$  are  $< 4$ ,  $< 4$  and  $< 2\%$ .

<sup>f</sup>  $\Gamma_\gamma/\Gamma = 0.91 \pm 0.05$  (1976RO07).

<sup>g</sup> Transitions to  $^{19}\text{F}^*(1.35 + 1.46, 1.55)$  are each  $< 0.8\%$ .

<sup>h</sup>  $\Gamma_\gamma/\Gamma = 0.76 + 0.15$  (1976RO07).

<sup>i</sup> Transitions to  $^{19}\text{F}^*(0, 0.11)$  are each  $< 5\%$ .

<sup>j</sup> Transitions to  $^{19}\text{F}^*(0, 0.11, 1.55, 2.78)$  are  $< 0.5$ ,  $< 1.5$ ,  $< 5$  and  $< 2\%$ .

<sup>k</sup> Transitions to  $^{19}\text{F}^*(0.197, 1.35, 1.55)$  are  $< 1$ ,  $< 1.5$  and  $< 2\%$ .

<sup>l</sup> Transitions to  $^{19}\text{F}^*(0, 0.11, 1.46, 1.55)$  are  $< 5$ ,  $< 2$ ,  $< 25$  and  $< 25\%$ .

<sup>m</sup> Transitions to  $^{19}\text{F}^*(4.00, 4.03)$  are  $< 2$  and  $< 1\%$ .

<sup>n</sup> Transitions to  $^{19}\text{F}^*(1.35, 1.46, 1.55)$  are  $< 0.5$ ,  $< 1.5$  and  $< 1\%$ .

<sup>o</sup> Transitions to  $^{19}\text{F}^*(2.78, 4.38, 4.68)$  are  $< 1$ ,  $< 1$  and  $< 2\%$ .

<sup>p</sup> Transitions to  $^{19}\text{F}^*(4.00, 4.03, 4.38, 5.47)$  are  $< 3$ ,  $< 3$ ,  $< 3$  and  $< 2\%$ .

<sup>q</sup> Transitions to  $^{19}\text{F}^*(4.00, 4.03, 4.55, 4.65, 5.43, 5.47)$  are  $< 2$  to  $< 8\%$ : see (1978AJ03).

<sup>r</sup> Transitions to  $^{19}\text{F}^*(0.11, 0.197)$  are  $< 8$  and  $< 5\%$ .

<sup>s</sup> Transitions to other states are  $< 0.2$  to  $< 2\%$ : see (1978AJ03).

<sup>t</sup>  $\Gamma_\gamma = 4.7$  eV,  $\Gamma_\gamma/\Gamma = 0.65 \pm 0.10$ ; see Table 19.9 in (1972AJ02). Transitions to  $^{19}\text{F}^*(0.11, 1.35, 1.46, 4.38)$  are  $< 0.4$ ,  $< 1.3$ ,  $< 1$  and  $< 1.3\%$ .

<sup>u</sup> Transitions to other states are  $< 7$  to  $< 10\%$ .

<sup>v</sup> (1980WI17).


<sup>w</sup> (1978SY01): branching ratios are relative intensities at  $\theta = 55^\circ$ .

<sup>x</sup> First branching ratio value shown for each transition is from (1978SY01); second is from (1980WI17). Where only one value is shown it is from (1980WI17), except when footnoted.




$$Q_m = 13.2739$$

See (1972AJ02).



$$Q_m = 16.396$$



$$Q_m = 5.964$$

$$E_b = 16.396$$

(c) $^{12}\text{C}(^7\text{Li}, \text{p})^{18}\text{O}$	$Q_{\text{m}} = 8.402$
(d) $^{12}\text{C}(^7\text{Li}, \text{d})^{17}\text{O}$	$Q_{\text{m}} = 2.582$
(e) $^{12}\text{C}(^7\text{Li}, \text{t})^{16}\text{O}$	$Q_{\text{m}} = 4.695$
(f) $^{12}\text{C}(^7\text{Li}, \alpha)^{15}\text{N}$	$Q_{\text{m}} = 12.3817$
(g) $^{12}\text{C}(^7\text{Li}, ^7\text{Li})^{12}\text{C}$	

For reaction (a) see (1978AJ03). The yield of  $^{18}\text{F}$  [reaction (b)] has been determined for  $E(^7\text{Li}) = 2.5$  to  $3.5$  MeV (1961NO05). The cross sections for reaction (c), (d), (e) and (f) vary strongly over the range  $E(^7\text{Li}) = 4$  to  $14$  MeV but with little, if any, cross correlation. Strong fluctuations continue to  $E(^7\text{Li}) = 25$  MeV and there is some evidence of structures at higher energies: for a listing of the earlier references see (1978AJ03). The excitation functions for several groups in  $^{16}\text{O}$  have been studied for  $E(^{12}\text{C}) = 54$  to  $62$  MeV, as has the total cross section (for residues of  $A > 8$ ) for  $E(^7\text{Li}) = 10$  to  $38$  MeV (1981DE15). See (1976PO02, 1978DR07) for reaction (g). See also (1981XE01; theor.),  $^{18}\text{O}$  here,  $^{16}\text{O}$  and  $^{17}\text{O}$  in (1982AJ01),  $^{15}\text{N}$  in (1981AJ01) and  $^{12}\text{C}$  in (1980AJ01).

4.  $^{12}\text{C}(^9\text{Be}, \text{d})^{19}\text{F}$   $Q_{\text{m}} = -0.3005$

At  $E(^9\text{Be}) = 12 - 27$  MeV angular distributions are reported to  $^{19}\text{F}^*(2.78)$  and to several unresolved groups (1979JA22, 1981JA09). The excitation function for  $^{19}\text{F}^*(2.78)$  has been measured for  $E(^9\text{Be}) = 10.5$  to  $25.9$  MeV (1980HU1E).

5.  $^{12}\text{C}(^{11}\text{B}, \alpha)^{19}\text{F}$   $Q_{\text{m}} = 7.7303$

At  $E(^{11}\text{B}) = 18.0$  to  $34.1$  MeV excitation functions are reported for the transitions to  $^{19}\text{F}^*(0 + 0.11 + 0.20, 1.35 + 1.46 + 1.56, 2.78)$  (1979FR05).

6.  $^{12}\text{C}(^{12}\text{C}, \alpha\text{p})^{19}\text{F}$   $Q_{\text{m}} = -8.2266$

For the yield of  $^{19}\text{F}$  for  $E(^{12}\text{C}) = 24$  to  $62$  MeV see (1980KO02).

7.  $^{12}\text{C}(^{14}\text{N}, ^7\text{Be})^{19}\text{F}$   $Q_{\text{m}} = -11.419$

Angular distributions are reported at  $E(^{14}\text{N}) = 78.8$  MeV for the transitions to  $^{19}\text{F}^*(0 + 0.11 + 0.20, 1.35 + 1.46 + 1.55, 2.78)$  (1977MO1A, 1979MO14).



8.  $^{12}\text{C}(^{20}\text{Ne}, ^{13}\text{N})^{19}\text{F}$   $Q_{\text{m}} = -10.901$

See (1979OR01).

9. (a)  $^{13}\text{C}(^6\text{Li}, \text{t})^{16}\text{O}$   $Q_{\text{m}} = 6.9994$   $E_{\text{b}} = 18.6997$   
 (b)  $^{13}\text{C}(^6\text{Li}, \alpha)^{15}\text{N}$   $Q_{\text{m}} = 14.6859$   
 (c)  $^{13}\text{C}(^6\text{Li}, ^6\text{Li})^{13}\text{C}$

Excitation functions for tritons to a number of states of  $^{16}\text{O}$  have been measured for  $E(^6\text{Li}) = 20$  to 32 MeV [correlated structures are not observed (1980CU03)] and 24 to 26 MeV (1982AB02). Excitation functions for  $\alpha_0$  have been measured for  $E(^6\text{Li}) = 7.7$  to 16.8 MeV: structures are observed but they are not correlated (1974CO13). For reaction (c) see (1976PO02, 1978DR07). For fusion cross sections see (1981DE1H, 1981DE2K). See also (1981ME1F; theor.).

10.  $^{13}\text{C}(^9\text{Be}, \text{t})^{19}\text{F}$   $Q_{\text{m}} = 1.0105$

Angular distributions have been studied at  $E(^{13}\text{C}) = 27.9$  MeV to  $^{19}\text{F}^*(0 + 0.11 + 0.20, 2.78)$  (1980BO21).

11. (a)  $^{14}\text{N}(^6\text{Li}, \text{p})^{19}\text{F}$   $Q_{\text{m}} = 11.1491$   
 (b)  $^{14}\text{N}(^7\text{Li}, \text{d})^{19}\text{F}$   $Q_{\text{m}} = 6.123$

See (1978AJ03).

12.  $^{14}\text{N}(^{10}\text{B}, \alpha\text{p})^{19}\text{F}$   $Q_{\text{m}} = 6.6886$

Cross sections have been measured for  $E(^{14}\text{N}) = 7$  to 18 MeV (1978WU1C).

13.  $^{15}\text{N}(\alpha, \gamma)^{19}\text{F}$   $Q_{\text{m}} = 4.0138$

Resonances in the yield of  $\gamma$ -rays are observed below  $E_{\alpha} = 10.4$  MeV: the parameters for these are displayed in Table 19.9. Branching ratios are shown in Table 19.7 and  $\tau_{\text{m}}$  measurements (1977DI18, 1980AN02) in Table 19.8. The  $J^{\pi}$  values are shown in Table 19.9 are based on correlation and angular distribution measurements and on branching ratio determinations.

Table 19.8: Lifetimes of some  $^{19}\text{F}$  states <sup>a</sup>

$^{19}\text{F}^*$ (MeV)	$\tau_m$	Refs.
0.110	$0.853 \pm 0.010$ nsec	mean: see (1972AJ02)
0.197	$128.8 \pm 1.5$ nsec	mean: see (1978AJ03)
1.35	$3.7 \pm 0.7$ psec	(1980AN02)
	$4.4 \pm 0.6$ psec <sup>A</sup>	see (1980AN02)
1.46	$140 \pm 15$ fsec	(1980AN02)
	$90 \pm 20$ fsec <sup>A</sup>	see <sup>e</sup>
1.55	$5 \pm 3$ fsec	(1980AN02)
2.78	$370 \pm 25$ fsec	(1980AN02)
	$280 \pm 30$ fsec <sup>A</sup>	see <sup>e</sup>
3.91 <sup>b</sup>	$9 \pm 5$ fsec	(1977DI18)
4.00 <sup>b</sup>	$19 \pm 7$ fsec	(1980AN02)
4.03	$63 \pm 19$ fsec	(1980AN02)
	$67 \pm 15$ fsec <sup>A</sup>	see (1980AN02)
4.38 <sup>c</sup>	$< 11$ fsec	(1975LE16)
4.55 <sup>d</sup>	$< 50$ fsec	(1976RO07)
4.56	$17_{-8}^{+10}$ fsec <sup>A</sup>	(1975LE16)
	$< 30$ fsec	(1976BH03)
4.65	$2.2 \pm 0.3$ psec	mean: see (1978AJ03)
4.68 <sup>d</sup>	$15.4 \pm 3.0$ fsec	(1972RO01)
5.11 <sup>d</sup>	$< 30$ fsec	(1976RO07)
5.34	$\leq 0.1$ fsec	see <sup>f</sup>
5.42	$\leq 0.9$ fsec	see <sup>g</sup>
5.46	$\leq 0.26$ fsec	see <sup>f</sup>
5.62	$< 1.3$ fsec	see <sup>f</sup>

A = adopted.

<sup>a</sup> See also Tables 19.10 in (1972AJ02) and in (1978AJ03). I am greatly indebted to Dr. D.W.O. Rogers for his comments and criticisms of the 1978 table.

<sup>b</sup> See also (1976BH03).

<sup>c</sup> See also (1976RO07).

<sup>d</sup> See also (1975LE16).

<sup>e</sup> P.M. Endt, private communication; based on reassessment of uncertainties in  $\tau_m$  measurements.

<sup>f</sup> Using the rule  $\Gamma > 4\Gamma_\alpha\Gamma_\gamma/\Gamma$  (P.M. Endt, private communication). See also (1980AN02, 1975LE16).

<sup>g</sup> From  $\omega\gamma$  and  $\Gamma_\gamma/\Gamma$  (P.M. Endt, private communication). See also (1980AN02).

Table 19.9: Resonances in  $^{15}\text{N}(\alpha, \gamma)^{19}\text{F}$  <sup>a</sup>

$E_\alpha$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\omega\gamma$ (eV)	$J^\pi$	$E_x$ (MeV $\pm$ keV)
0.85	$(42.8 \pm 8.5) \times 10^{-6}$ <sup>b</sup>	$(6.0 \pm 1.0) \times 10^{-3}$	$\frac{5}{2}^-$	$4.681 \pm 1$
$1.385 \pm 3$		$(13 \pm 8) \times 10^{-3}$ <sup>c</sup>	$\frac{5}{2}^+$	$5.105 \pm 2$
$1.678 \pm 3$		$1.64 \pm 0.16$	$\frac{1}{2}^{(+)}$	$5.337 \pm 2$
1.790		$0.42 \pm 0.09$ <sup>c</sup>	$\frac{7}{2}^-$	5.427
$1.839 \pm 2$	$< 1$	$2.5 \pm 0.4$ <sup>c</sup>	$\frac{7}{2}^+$	5.465
$1.883 \pm 3$	$4 \pm 1$	$4.2 \pm 1.1$ <sup>c</sup>	$\frac{3}{2}^+$	5.500
1.930		$0.48 \pm 0.11$ <sup>c</sup>	$\frac{5}{2}^+$	5.54
$2.035 \pm 4$		$0.37 \pm 0.09$	$\frac{3}{2}^-$	5.620
$2.441 \pm 4$		$0.53 \pm 0.13$	$\frac{1}{2}^+$	$5.938 \pm 3$
$2.608 \pm 2$		$2.70 \pm 0.54$	$\frac{7}{2}^+$	$6.070 \pm 1$
$2.631 \pm 4$		$4.50 \pm 0.90$	$\frac{3}{2}^-$	$6.088 \pm 3$
$2.722 \pm 2$		$2.40 \pm 0.60$	$\frac{7}{2}^-$	$6.160 \pm 1$
$2.873 \pm 3$		$1.0 \pm 0.2$	$\frac{5}{2}^+$	$6.282 \pm 2$
$2.935 \pm 3$		$0.76 \pm 0.15$	$\frac{7}{2}^+$	$6.330 \pm 2$
$3.1468 \pm 1.5$		$1.7 \pm 0.3$	$\frac{3}{2}^+$	$6.4976 \pm 1.5$
$3.1498 \pm 1.5$		$2.3 \pm 0.4$	$\frac{11}{2}^+$	$6.5000 \pm 1.5$
$3.183 \pm 2$		$2.4 \pm 0.4$	$\frac{3}{2}^+$	$6.526 \pm 2$
$3.218 \pm 2$		$0.63 \pm 0.13$	$\frac{7}{2}^-$	$6.554 \pm 2$
$3.267 \pm 2$		$1.6 \pm 0.3$	$\frac{9}{2}^+$	$6.592 \pm 2$
$3.511 \pm 3$		$10.9 \pm 1.5$	$\frac{3}{2}^-$	$6.785 \pm 2$
$3.576 \pm 3$		$1.0 \pm 0.2$	$\frac{5}{2}^-$	$6.836 \pm 2$
$3.645 \pm 5$		$6.1 \pm 1.3$	$\frac{3}{2}^-$	$6.891 \pm 4$
$3.688 \pm 3$		$9.7 \pm 1.4$	$\frac{7}{2}^-$	$6.925 \pm 2$
$3.993 \pm 2$		$1.00 \pm 0.12$	$\frac{11}{2}^-$	$7.1662 \pm 0.7$
4.465		$17.0 \pm 2.7$	$\frac{5}{2}^+; T = \frac{3}{2}$	$7.538 \pm 2$
4.618		$3.7 \pm 0.9$	$\frac{3}{2}^+; T = \frac{3}{2}$	$7.659 \pm 2$
$4.96 \pm 3$		$2.3 \pm 0.4$	$\frac{7}{2}^+, \frac{9}{2}$	7.929
$4.97 \pm 3$		$3.1 \pm 0.5$	$\frac{11}{2}^+$	7.937
$5.413 \pm 5$	$< 1$	$0.53 \pm 0.08$	$\frac{13}{2}^-$	$8.288 \pm 2$
5.438 <sup>e</sup>	$< 1$	$2.1 \pm 0.5$ <sup>d</sup>	$\frac{5}{2}^+$	$8.306 \pm 4$
5.519 <sup>e</sup>	$7.5 \pm 1.5$	$0.54 \pm 0.2$	$\frac{7}{2}, \frac{5}{2}^+$	$8.370 \pm 4$

Table 19.9: Resonances in  $^{15}\text{N}(\alpha, \gamma)^{19}\text{F}$  <sup>a</sup> (continued)

$E_\alpha$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\omega\gamma$ (eV)	$J^\pi$	$E_x$ (MeV $\pm$ keV)
5.784	$\approx 1$	$5.1 \pm 1.3$	$\frac{5}{2}$	$8.579 \pm 4$
5.794		$1.6 \pm 0.35$ <sup>f</sup>	$\frac{3}{2}$	$8.587 \pm 3$
5.847 <sup>e</sup>	$< 1$	$2.5 \pm 0.4$	$\frac{7}{2}^-$	$8.629 \pm 4$
6.145	$< 1$	$0.2 \pm 0.05$	$< \frac{9}{2}$	$8.864 \pm 4$
6.259 <sup>e</sup>	$\approx 1$	$0.85 \pm 0.2$	$\frac{11}{2}^-, (\frac{9}{2}^+)$	$8.953 \pm 3$
6.356	$4.2 \pm 1$	$0.53 \pm 0.26$	$\frac{5}{2}, \frac{7}{2}$	$9.030 \pm 5$
6.442		$0.48 \pm 0.15$ <sup>g</sup>	$\frac{7}{2}^+$	$9.098 \pm 4$
6.445	$\approx 1$	$0.40 \pm 0.1$	$\frac{7}{2}, \frac{9}{2}$	$9.101 \pm 4$
6.526	$9.9 \pm 1.5$	$1.4 \pm 1$	$\frac{1}{2}, \frac{3}{2}$	$9.165 \pm 5$
6.576	$10.2 \pm 1.5$	1.5	$\frac{3}{2}$	$9.204 \pm 7$
6.656	$2 \pm 1$	$0.15 \pm 0.04$	$\frac{11}{2}^+, \frac{9}{2}^+$	$9.267 \pm 4$
6.672	$< 1.5$	$0.38 \pm 0.09$	$\frac{7}{2}, \frac{9}{2}$	$9.280 \pm 5$
6.723 <sup>e</sup>	$3.4 \pm 1$	$3.4 \pm 1.7$	$\frac{1}{2}^+$	$9.320 \pm 4$
6.735	$\approx 6$		$< \frac{5}{2}$	$9.329 \pm 4$
6.963	$< 1$	$0.7 \pm 0.2$	$\frac{5}{2}^+, \frac{7}{2}^+$	$9.509 \pm 4$
6.993	$6.3 \pm 1.5$	0.5	$\frac{3}{2} \rightarrow \frac{7}{2}$	$9.533 \pm 6$
7.057	$9.6 \pm 1.5$	$5.2 \pm 3$	$\frac{7}{2}$	$9.584 \pm 4$
7.131	$\approx 8$	$\approx 1$	$\frac{3}{2}, \frac{5}{2}$	$9.642 \pm 6$
7.146	$\approx 6$	$\approx 2$	$\frac{3}{2}, \frac{5}{2}$	$9.654 \pm 6$
7.179	$\approx 4$	$\approx 1$	$\frac{1}{2}, \frac{3}{2}$	$9.680 \pm 6$
7.217	$< 1$	$4 \pm 0.7$	$\frac{9}{2}^+, \frac{11}{2}$	$9.710 \pm 4$
7.349	$< 1.5$	$3.5 \pm 0.8$ <sup>h</sup>	$\frac{5}{2}^+$	$9.814 \pm 4$
7.375 <sup>e</sup>	$< 1$	$0.51 \pm 0.1$	$\frac{11}{2} \rightarrow \frac{15}{2}$	$9.834 \pm 3$
7.422	$\approx 1.5$	$3.6 \pm 0.6$	$\frac{9}{2}^+, \frac{11}{2}^-$	$9.872 \pm 3$
7.491	$\approx 1$	$19.3 \pm 3.0$	$\frac{9}{2}^+$	$9.926 \pm 3$
7.696	$< 1.5$	$2.37 \pm 0.5$	$\frac{5}{2}, \frac{7}{2}$	$10.088 \pm 5$
7.749	$3.2 \pm 1$	$1.3 \pm 0.4$	$\frac{3}{2}, \frac{5}{2}$	$10.130 \pm 6$
8.047	$3 \pm 1.5$	$0.9 \pm 0.4$	$\frac{7}{2} \rightarrow \frac{11}{2}$	$10.365 \pm 4$
8.105	$< 1.5$	$15.0 \pm 3.0$	$\frac{11}{2}^+, \frac{13}{2}^+$	$10.411 \pm 3$

<sup>a</sup> For references see Table 19.8 in (1978AJ03). For branching ratios see Table 19.7 here. Resonances with  $E_\alpha > 5.4$  MeV are from (1978SY01).  $\omega\gamma = \omega\Gamma_\gamma\Gamma_\alpha/\Gamma$ .

<sup>b</sup>  $\Gamma_\alpha = 2.1 \pm 0.7$  meV,  $\Gamma_\gamma = 40.7 \pm 8.1$  meV (1972RO01).

<sup>c</sup> See also Table 19.7 in (1972AJ02).

<sup>d</sup>  $\omega\gamma$  ( $55^\circ$ ) for this value and all values below (1978SY01).


<sup>e</sup> Value recalculated by reviewer from  $E_x$ .

<sup>f</sup>  $\Gamma_\alpha/\Gamma_p = 0.026 \pm 0.008$  (1978SY01).

<sup>g</sup>  $\Gamma_\alpha/\Gamma_p = 0.1 \pm 0.04$ . Using  $\Gamma = 0.57 \pm 0.03$  keV (Table 19.18),  $\Gamma_\alpha = 0.052 \pm 0.03$ ,  $\Gamma_p = 0.52 \pm 0.03$  keV (1978SY01).

<sup>h</sup>  $\Gamma_\alpha/\Gamma_p = 0.55 \pm 0.16$  (1978SY01).

The  $E_x$  of states involved in cascade decays are  $4377 \pm 1$  and  $4548 \pm 2$  keV (1976RO07),  $3999.6 \pm 1.2$  and  $4031.9 \pm 0.4$  keV (1973RO09). The  $K^\pi = \frac{1}{2}^+$  band involves  $^{19}\text{F}^*(0.110 [\frac{1}{2}^-], 1.46 [\frac{3}{2}^-], 1.35 [\frac{5}{2}^-], 4.00 [\frac{7}{2}^-], 4.03 [\frac{9}{2}^-], 7.16 [\frac{11}{2}^-])$  (1973RO09) and possibly  $^{19}\text{F}^*(8.29 [\frac{13}{2}^-])$  (1974UN01) [ $J^\pi$  in brackets]. The situation concerning the other bands is not clear: see (1972AJ02) for a discussion of the evidence for other assignments of  $J^\pi$  and  $K^\pi$ .  $^{19}\text{F}^*(10.41)$  is likely to be the second  $\frac{13}{2}^+$  ( $2s, 1d$ )<sup>3</sup> state in  $^{19}\text{F}$  (1976SY01, 1978SY01). See also (1978TA1U; astrophys.).



Resonances observed in the  $(\alpha, \alpha'\gamma)$  and  $(\alpha, p\gamma)$  reactions (1978SY01) and in the elastic scattering (reaction (b)) [see (1978AJ03) for references] are displayed in Table 19.10. The elastic excitation function has been measured for  $E_\alpha = 16$  to  $23.7$  MeV by (1977FE1C). See also (1979LE18; theor.).



This reaction has been studied at  $E(^7\text{Li}) = 40$  MeV: see Table 19.11 (1979MA26). See also (1978AJ03).



At  $E(^{11}\text{B}) = 115$  MeV,  $^{19}\text{F}^*(8.29, 8.96)$  [ $J^\pi = \frac{13}{2}^-, \frac{11}{2}^-$ , respectively] are populated as are states at  $10.4 \pm 0.05$ ,  $12.26 \pm 0.07$ ,  $12.67 \pm 0.08$ ,  $14.9 \pm 0.09$ ,  $15.4 \pm 0.09$ ,  $17.4 \pm 0.08$ ,  $18.6 \pm 0.08$  and  $19.7 \pm 0.08$  MeV (1979BR04).

Table 19.10: Levels of  $^{19}\text{F}$  from  $^{15}\text{N}(\alpha, \text{p})$  and  $^{15}\text{N}(\alpha, \alpha)^{\text{a}}$ 

$E_{\alpha}$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	$J^{\pi}$	$E_{\text{x}}$ (MeV $\pm$ keV)
$1.878 \pm 10$	4	$\frac{3}{2}^{+}$	5.496
$2.614 \pm 10$	1.5	$\frac{5}{2}^{+}$	6.077
$2.635 \pm 10$	5	$\frac{5}{2}^{-}$	6.094
$2.833 \pm 10$	10	$\frac{1}{2}^{+}$	6.250
$2.883 \pm 10$	3	$\frac{5}{2}^{+}$	6.289
$2.944 \pm 10$	3	$\frac{7}{2}^{+}$	6.338
$3.060 \pm 10$	360	$\frac{1}{2}^{-}$	$6.429 \pm 8$
$3.194 \pm 10$	5	$\frac{1}{2}^{+}$	6.535
$3.229 \pm 10$	2	$\frac{5}{2}^{+}$	6.563
$3.525 \pm 10$	3	$\frac{3}{2}^{-}$	6.796
$3.587 \pm 10$	1.5	$(\frac{5}{2}, \frac{3}{2})^{+}$	6.845
$3.648 \pm 10$	35	$\frac{5}{2}^{-}$	6.893
$3.705 \pm 10$	3	$(\frac{9}{2}, \frac{7}{2})^{-}$	6.938
$3.770 \pm 10$	64	$\frac{1}{2}^{-}$	$6.989 \pm 8$
$3.930 \pm 10$	40	$\frac{7}{2}^{+}$	$7.116 \pm 8$
4.127 <sup>c</sup>	$\lesssim 8$		7.271
4.23	$\lesssim 82$	$\frac{7}{2}^{+}$	7.35
4.49	$\lesssim 110$	$\frac{7}{2}^{+}$	7.56
4.53	$\lesssim 50$	$\frac{5}{2}^{+}$	7.59
4.710	$\lesssim 40$	$\frac{1}{2}^{-}$	7.731
4.780	$\lesssim 8$		7.787
4.93	$\lesssim 260$		7.90
(5.005)	( $\lesssim 8$ )		(7.964)
(5.018)	( $\lesssim 5$ )		(7.974)
5.116	$\lesssim 8$		8.052
5.203	$\lesssim 8$		8.120
5.232	$\lesssim 6$		8.143
5.25	$\lesssim 65$		8.16
5.284	$\lesssim 10$		8.184
5.481	$\lesssim 10$		9.340
$7.877^{\text{d}}$	$< 1$	$\frac{1}{2}^{+}$	$10.231 \pm 4$

Table 19.10: Levels of  $^{19}\text{F}$  from  $^{15}\text{N}(\alpha, \text{p})$  and  $^{15}\text{N}(\alpha, \alpha)$  <sup>a</sup> (continued)

$E_\alpha$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	$J^\pi$	$E_x$ (MeV $\pm$ keV)
7.977 <sup>d</sup>		$\frac{3}{2}^+$	10.308 $\pm$ 4
8.179 <sup>d</sup>	13.8 $\pm$ 1.5		10.469 $\pm$ 4
8.205 <sup>d</sup>	6.0 $\pm$ 1.0		10.488 $\pm$ 4
8.220	5.4 $\pm$ 1.0	$\frac{3}{2}^+$	10.501 $\pm$ 4
8.245	18 $\pm$ 2		10.521 $\pm$ 4
8.277	2.5 $\pm$ 1		10.546 $\pm$ 4
8.287 <sup>d</sup>	5.0 $\pm$ 1.5	$\frac{3}{2}^+$	10.554 $\pm$ 4
8.307 <sup>d</sup>	3.7 $\pm$ 1		10.560 $\pm$ 4

<sup>a</sup> For references see Table 19.9 in (1978AJ03).

<sup>b</sup> Resonances below  $E_\alpha = 5.5$  MeV are observed in  $(\alpha, \alpha_0)$ ; resonances above that energy are observed in  $(\alpha, \text{p}\gamma)$  and  $(\alpha, \alpha'\gamma)$  (1978SY01).

<sup>c</sup> I am indebted to Prof. C. Rolfs for his comments on the resonances with  $4 < E_\alpha < 5.5$  MeV.

<sup>d</sup> Value recalculated by reviewer from  $E_x$ .

17.  $^{15}\text{N}(^{13}\text{C}, ^9\text{Be})^{19}\text{F}$   $Q_m = -6.6341$

At  $E(^{13}\text{C}) = 105$  MeV,  $^{19}\text{F}^*(4.02, 8.29, 8.95, 12.26 \pm 0.1, 12.65 \pm 0.1, 14.8 \pm 0.12)$  are strongly populated.  $^{19}\text{F}^*(2.78)$  is also observed. It is suggested that  $^{19}\text{F}^*(8.29, 8.95, 12.26, 12.65)$  have  $J^\pi = \frac{13}{2}^-, \frac{11}{2}^-, (\frac{17}{2}^-), (\frac{15}{2}^-)$ , respectively, and that  $^{19}\text{F}^*(12.26, 12.66)$  are members of the  $2N + L = 8$   $\alpha$ -cluster band in  $^{19}\text{F}$  (1979BR04). See also (1977FO1E).

18. (a)  $^{16}\text{O}(\text{t}, \gamma)^{19}\text{F}$   $Q_m = 11.7003$   
 (b)  $^{16}\text{O}(\text{t}, \text{n})^{18}\text{F}$   $Q_m = 1.2690$   $E_b = 11.7003$   
 (c)  $^{16}\text{O}(\text{t}, \text{p})^{18}\text{O}$   $Q_m = 3.7069$   
 (d)  $^{16}\text{O}(\text{t}, \text{t})^{16}\text{O}$   
 (e)  $^{16}\text{O}(\text{t}, \alpha)^{15}\text{N}$   $Q_m = 7.6865$

For reaction (a) see (1978AJ03). The excitation function for reaction (b) has been measured for  $E_t = 0.3$  to 3.7 MeV: there is evidence for a maximum at  $E_t = 2.5$  MeV. For resonances in the yields of  $\text{p}_0, \text{p}_1, \alpha_0, \alpha_{1+2}$ , see (1978AJ03). The elastic yield [reaction (d)] shows a large number

Table 19.11: States in  $^{19}\text{F}$  from  $^{15}\text{N}(^7\text{Li}, t)$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$E_x$ (MeV $\pm$ keV)	$E_x$ (MeV $\pm$ keV)
$0.19 \pm 15$	$7.54 \pm 15$	$13.78 \pm 30$
$1.34 \pm 15$	$8.29 \pm 15$	$14.12 \pm 30$
$1.46 \pm 15$	$8.95 \pm 15$ <sup>b</sup>	$14.50 \pm 30$ <sup>b</sup>
$2.78 \pm 15$	$9.35 \pm 30$	$14.92 \pm 30$ <sup>b</sup>
$4.02 \pm 15$	$9.92 \pm 30$ <sup>b</sup>	$16.09 \pm 50$
$4.56 \pm 15$	$10.40 \pm 30$	$16.45 \pm 50$
$5.46 \pm 15$	$11.5 \pm 30$	$17.4 \pm 50$
$6.10 \pm 15$	$11.7 \pm 30$	$18.2 \pm 50$
$6.32 \pm 15$	$12.01 \pm 30$	$18.7 \pm 50$
$6.94 \pm 15$	$12.5$ <sup>c</sup>	$19.93 \pm 50$

<sup>a</sup> (1979MA26):  $E(^7\text{Li}) = 40$  MeV.

<sup>b</sup> Relatively strongly populated at  $\theta = 15^\circ$ .

<sup>c</sup> Unresolved.

Table 19.12: Resonances in  $^{16}\text{O}(t, t)$  <sup>a</sup>

$E_{\text{c.m.}}$ (MeV)	$E_x$ (MeV $\pm$ keV)	$J^\pi$	$\Gamma_{\text{c.m.}}$ (keV)
1.368	$13.068 \pm 4$	$\frac{1}{2}^+$	$\leq 10$
1.545	$13.245 \pm 10$	$\frac{1}{2}^-$	7
1.570	$13.270 \pm 10$	$\frac{1}{2}^+$	4.5
1.832	$13.532 \pm 10$	$\frac{1}{2}^+$	22
2.018	$13.718 \pm 20$	$\frac{3}{2}^-$	128
2.178	$13.878 \pm 15$	$\frac{1}{2}^+$	101
2.447	$14.147 \pm 20$	$\frac{1}{2}^+$	21
2.555	$14.255 \pm 15$	$\frac{3}{2}^+$	51
2.652	$14.352 \pm 10$	$\frac{1}{2}^+$	154
2.759	$14.459 \pm 25$	$\frac{3}{2}^+$	179
2.763	$14.463 \pm 25$	$\frac{5}{2}^+$	46

<sup>a</sup> (1973WE11): resonance parameters used to fit elastic scattering data. See also (1978AJ03).



of resonances: their parameters are displayed in Table 19.12. See also (1977CI1D), (1981AO01; theor.) and  $^{18}\text{O}$  here.

$$19. \text{}^{16}\text{O}(\alpha, \text{p})^{19}\text{F} \quad Q_{\text{m}} = -8.1137$$

Angular distributions have been measured at  $E_{\alpha} = 20.1$  to  $40$  MeV [see (1978AJ03)] and at  $E_{\alpha} = 40$  MeV (1979VA08: to  $^{19}\text{F}^*(0 + 0.20, 2.78, 4.65)$ ): the  $3\text{p } \frac{13}{2}^+$  strength is split between  $^{19}\text{F}^*(4.65, 10.42)$ . States observed in this reaction are displayed in Table 19.12 of (1978AJ03).

$$20. \text{}^{16}\text{O}({}^6\text{Li}, {}^3\text{He})^{19}\text{F} \quad Q_{\text{m}} = -4.094$$

This reaction (and its mirror reaction  $^{16}\text{O}({}^6\text{Li}, \text{t})^{19}\text{Ne}$  [see reaction 7 in  $^{19}\text{Ne}$ ]) have been studied at  $E({}^6\text{Li}) = 24$  MeV [see (1978AJ03)] and at  $46$  MeV (1979MA26). Members of the  $K^{\pi} = \frac{1}{2}^+$  and  $\frac{1}{2}^-$  rotational bands have been identified: see Table 19.13. In addition  ${}^3\text{He}$  groups are reported to states with  $E_{\text{x}} = 6.10, 6.92, 7.25, 8.29, 8.96, 9.7, 9.88, 10.4$  [all  $\pm 0.015$ ], to unresolved states near  $11.5, 12.7$  and to states at  $E_{\text{x}} = 13.76, 14.10, 15.00, 15.56$  and  $18.92$  [ $\pm 0.03$ ] MeV (1979MA26). See also (1977FO1E).

$$21. \text{}^{16}\text{O}({}^7\text{Li}, \alpha)^{19}\text{F} \quad Q_{\text{m}} = 9.234$$

Many states have been populated in this reaction: see Table 19.14 in (1978AJ03). See also (1977FO1E).

$$22. \text{}^{16}\text{O}({}^{10}\text{B}, {}^7\text{Be})^{19}\text{F} \quad Q_{\text{m}} = -6.968$$

This reaction, as well as the analog reaction [ $^{16}\text{O}({}^{10}\text{B}, {}^7\text{Li})^{19}\text{Ne}$ ] have been studied at  $E({}^{10}\text{B}) = 100$  MeV: an attempt is made to match analog states [see reaction 8 in  $^{19}\text{Ne}$ ] but problems of energy resolution are evident (1976HA06). See also (1977FO1E).

$$23. \text{}^{16}\text{O}({}^{11}\text{B}, {}^8\text{Be})^{19}\text{F} \quad Q_{\text{m}} = 0.4765$$

The angular distribution for the transition to  ${}^8\text{Be}_{\text{g.s.}} + {}^{19}\text{F}_{\text{g.s.}}$  has been measured at  $E({}^{16}\text{O}) = 60$  MeV (1972SC17).

Table 19.13: Levels of  $^{19}\text{F}$  and  $^{19}\text{Ne}$  from  $^{16}\text{O}(^6\text{Li}, ^3\text{He})$  and  $^{16}\text{O}(^6\text{Li}, t)$  <sup>a</sup>

$J^\pi$ <sup>c</sup>	$E_x$ in $^{19}\text{F}$ (MeV) <sup>b</sup>			$E_x$ in $^{19}\text{Ne}$ (MeV) <sup>b</sup>		
	$K^\pi = \frac{1}{2}^+$	$K^\pi = \frac{1}{2}^-$	other	$K^\pi = \frac{1}{2}^+$	$K^\pi = \frac{1}{2}^-$	other
$\frac{1}{2}^+$	0			0		
$\frac{3}{2}^+$	1.56			1.54 <sup>e</sup>		
$\frac{5}{2}^+$	0.20			0.24		
$\frac{7}{2}^+$	5.47			5.42		
$\frac{9}{2}^+$	2.78			2.79 <sup>e</sup>		
$\frac{11}{2}^+$	(6.50) <sup>d</sup>					
$\frac{13}{2}^+$	4.65			4.64		
$\frac{1}{2}^-$		0.11			0.28	
$\frac{3}{2}^-$		1.46			1.62 <sup>e</sup>	
$\frac{5}{2}^-$		1.35			1.51 <sup>e</sup>	
$\frac{7}{2}^-$		4.00			4.20 <sup>g</sup>	
$\frac{9}{2}^-$		4.03			4.14 <sup>g</sup>	
$\frac{3}{2}^+$			3.91 <sup>e</sup>			4.03 <sup>e</sup>
$\frac{7}{2}^+$			4.38			4.38 <sup>e</sup>
$\frac{5}{2}^+$			4.55			4.55 <sup>e</sup>
$\frac{3}{2}^-, (\frac{1}{2}^-)$			4.56			$4.593 \pm 0.006$
$\frac{5}{2}^-$			4.68			4.71
$\frac{5}{2}^- (-)$			5.11			5.09 <sup>f</sup>
$\frac{5}{2}^+$			5.34			
$\frac{7}{2}^-$			5.43			

<sup>a</sup> (1972BI14, 1972GA08, 1973BI02, 1979MA26).

<sup>b</sup> Energies are nominal.

<sup>c</sup>  $J^\pi$  assignments based on similarities in angular distributions, and on known spin of one of the analog states.

<sup>d</sup> Not strongly populated at  $E(^6\text{Li}) = 24$  MeV.

<sup>e</sup>  $J^\pi$  assignments based on similarities in  $\sigma_{\text{max}}$  in both reactions, and on known spin of analog state.

<sup>f</sup>  $J^\pi = (\frac{5}{2}^-, \frac{7}{2}^-)$  (1973BI02); a state at 4.78 MeV is also reported (1973BI02).

<sup>g</sup> See, however, reaction 10 in  $^{19}\text{Ne}$ .

$$24. \text{ (a) } {}^{16}\text{O}({}^{12}\text{C}, 2\alpha\text{p}){}^{19}\text{F} \quad Q_{\text{m}} = -15.3886$$

$$\text{ (b) } {}^{16}\text{O}({}^{13}\text{C}, {}^{10}\text{B}){}^{19}\text{F} \quad Q_{\text{m}} = -12.1763$$

For reaction (a) see (1978CH15). At  $E({}^{13}\text{C}) = 105$  MeV  ${}^{19}\text{F}^*(2.78, 4.68)$  are preferentially populated (1979GO17, 1979RA10).

$$25. \text{ (a) } {}^{17}\text{O}(\text{d}, \text{n}){}^{18}\text{F} \quad Q_{\text{m}} = 3.382 \quad E_{\text{b}} = 13.813$$

$$\text{ (b) } {}^{17}\text{O}(\text{d}, \text{p}){}^{18}\text{O} \quad Q_{\text{m}} = 5.8199$$

$$\text{ (c) } {}^{17}\text{O}(\text{d}, \text{t}){}^{16}\text{O} \quad Q_{\text{m}} = 2.1130$$

$$\text{ (d) } {}^{17}\text{O}(\text{d}, {}^3\text{He}){}^{16}\text{N} \quad Q_{\text{m}} = -8.287$$

$$\text{ (e) } {}^{17}\text{O}(\text{d}, \alpha){}^{15}\text{N} \quad Q_{\text{m}} = 9.7995$$

For reaction (a) see  ${}^{18}\text{F}$  in (1972AJ02); for reaction (b) see  ${}^{18}\text{O}$ ; for reaction (e) see  ${}^{15}\text{N}$  in (1970AJ04). At  $E_{\text{d}} = 52$  MeV polarization measurements are reported involving the groups to  ${}^{16}\text{O}^*(6.13, 8.87, 19.19)$  and  ${}^{16}\text{N}^*(0, 0.297, 6.17 + 6.36)$  (1981MA1E). See also  ${}^{16}\text{N}$  and  ${}^{16}\text{O}$  in (1982AJ01).

$$26. {}^{17}\text{O}({}^3\text{He}, \text{p}){}^{19}\text{F} \quad Q_{\text{m}} = 8.320$$

States studied in this reaction at  $E({}^3\text{He}) = 18$  MeV are displayed in Table 19.14.

$$27. {}^{17}\text{O}(\alpha, \text{d}){}^{19}\text{F} \quad Q_{\text{m}} = -10.033$$

At  $E_{\alpha} = 47.5$  MeV angular distributions have been studied to the  $\frac{1}{2}^+ \rightarrow \frac{9}{2}^+$  and the  $\frac{13}{2}^+$  members of the  $K = \frac{1}{2}^+$  band [ ${}^{19}\text{F}^*(0, 0.197, 1.55, 2.78, 4.65, 5.47)$ ], to two  $\frac{11}{2}^+$  states  ${}^{19}\text{F}^*(6.49, 7.94)$  [both of which are strongly populated] and to the  $\frac{7}{2}^+$  state at 4.38 MeV. The reaction populates strongly only those positive-parity states that predominantly (sd)<sup>3</sup> (1978FO11, 1975FO07).

$$28. {}^{17}\text{O}({}^{13}\text{C}, {}^{11}\text{B}){}^{19}\text{F} \quad Q_{\text{m}} = -4.865$$

See (1977CH22).

Table 19.14: States of  $^{19}\text{F}$  from  $^{17}\text{O}(^3\text{He}, \text{p})^a$ 

$E_x$ (MeV $\pm$ keV)	$L$	$E_x$ (MeV $\pm$ keV)	$L$
0.0 <sup>b</sup>	2 (+4)	4.373 $\pm$ 4 <sup>b</sup>	0 + 2
0.106 $\pm$ 10	(1 + 3)	4.545 $\pm$ 6 <sup>b</sup>	2
0.188 $\pm$ 5 <sup>b</sup>	0	4.644 $\pm$ 6 <sup>b</sup>	4
1.346 $\pm$ 10	3 (+5)	5.099 $\pm$ 4 <sup>d</sup>	0 + 2
1.460 $\pm$ 8	1 + 3	5.332 $\pm$ 10	
1.556 $\pm$ 3 <sup>b</sup>	0	5.414 $\pm$ 8	3 (+5)
2.783 $\pm$ 3 <sup>b</sup>	2	5.465 $\pm$ 3 <sup>b</sup>	2
3.902 $\pm$ 9 <sup>c</sup>	2 + 0	5.533 $\pm$ 8	2 + 4
3.993 $\pm$ 5	3 (+5)	5.621 $\pm$ 7	1 $\pm$ 3
4.026 $\pm$ 12			

<sup>a</sup> (1978FO22, 1979BI05, 1979FO06):  $E(^3\text{He}) = 18$  MeV.

<sup>b</sup> Dominant (sd)<sup>3</sup> character.

<sup>c</sup>  $\frac{3}{2}^+$ ; predominantly 5p–2h.

<sup>d</sup> Second (sd)<sup>3</sup>  $\frac{5}{2}^+$  level.

29.  $^{18}\text{O}(\text{p}, \gamma)^{19}\text{F}$

$$Q_m = 7.9934$$

This reaction has recently been studied for  $E_p = 80$  to 2200 keV by (1980WI17). A large number of resonances have been investigated and  $E_{\text{res}}$ , total and partial widths, branching and mixing ratios and  $\omega\gamma$  values are reported. Transition strength arguments as well as analyses of  $\gamma$ -ray angular distribution data lead to  $J^\pi$  assignments: see Tables 19.7, 19.15 and 19.16 for a display of the results (1980WI17).

In addition absolute cross sections measured for direct capture leads to  $C^2S$  values for a number of states of  $^{19}\text{F}$ . Reduced widths and  $J^\pi$  determinations lead to (1980WI17) to postulate  $^{19}\text{F}^*(3.91, 4.55, 4.38, 6.59, 6.50, 10.43)$  as the  $J^\pi = \frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+, \frac{9}{2}^+, \frac{11}{2}^+, \frac{13}{2}^+$  of the  $K^\pi = \frac{3}{2}^+$  rotational band;  $^{19}\text{F}^*(7.73 \text{ or } 7.26, 6.09, 9.81, 6.92, 9.87)$  as the  $J^\pi = \frac{1}{2}^-, \frac{3}{2}^-, \frac{5}{2}^-, \frac{7}{2}^-$  and  $\frac{11}{2}^-$  members of the excited  $K^\pi = \frac{1}{2}^-$  rotational band; and  $^{19}\text{F}^*(4.56, 4.68, 5.42, 6.10, 7.17)$  as the  $J^\pi = \frac{3}{2}^-, \frac{5}{2}^-, \frac{7}{2}^-, \frac{9}{2}^-$  and  $\frac{11}{2}^-$  members of the  $K^\pi = \frac{3}{2}^-$  rotational band. The direct capture transition to  $^{19}\text{F}^*(7.54)$  indicates some isospin mixing in this  $\frac{5}{2}^+$ , first  $T = \frac{3}{2}$  state in  $^{19}\text{F}$  (1980WI17).

Stellar reaction rates have also been calculated: the data cover  $T_9 = 0.01 - 5.0$ . The consequences for the final termination of the CNO tri-cycle are discussed (1980WI17). See also (1975ZI1A; astrophys.). For the earlier work see (1978AJ03). See also (1978DI13).

30.  $^{18}\text{O}(\text{p}, \text{n})^{18}\text{F}$ 

$$Q_{\text{m}} = -2.4379$$

$$E_{\text{b}} = 7.9934$$

Table 19.15: Some bound states of  $^{19}\text{F}$  involved in the capture  $\gamma$ -rays from  $^{18}\text{O} + \text{p}$  <sup>a</sup>

$E_{\text{x}}$ (keV)	$E_{\text{x}}$ (keV)	$E_{\text{x}}$ (keV)
$4648 \pm 1$	$6088 \pm 1$	$6839 \pm 1$
$5107 \pm 1$	$6100 \pm 2$ <sup>c</sup>	$6930 \pm 3$
$5338 \pm 4$	$6163 \pm 2$	$6989 \pm 3$
$5418 \pm 1$	$6255 \pm 1$	$7262 \pm 2$ <sup>d</sup>
$5462 \pm 2$	$6283 \pm 3$	$7364 \pm 4$ <sup>e</sup>
$5501 \pm 2$	$6493 \pm 3$	$7540 \pm 1$
$5535 \pm 2$	$6500 \pm 1$	$7661 \pm 1$
$5621 \pm 1$ <sup>b</sup>	$6529 \pm 2$	
$5938 \pm 1$	$6789 \pm 2$	

<sup>a</sup> (1980WI17). See also Tables 19.7 and 19.16.<sup>b</sup>  $J^{\pi} = \frac{5}{2}^{-}$ .<sup>c</sup>  $J^{\pi} = \frac{9}{2}^{-}$ .<sup>d</sup>  $J^{\pi} = \frac{1}{2}^{-}, \frac{3}{2}$ .<sup>e</sup>  $J^{\pi} = \frac{1}{2}^{+}$ .

Yield measurements are reported from  $E_{\text{p}} = 2.5$  to 13.5 MeV [see (1978AJ03) for the references]. Table 19.17 displays the observed resonances. See also  $^{18}\text{F}$ .

31.  $^{18}\text{O}(\text{p}, \text{p})^{18}\text{O}$ 

$$E_{\text{b}} = 7.9934$$

Scattering studies have been carried out for  $E_{\text{p}} = 0.6$  to 16.3 MeV [see (1978AJ03)], and for  $E_{\text{p}} = 3.05$  to 3.5 MeV (1978DI13;  $\text{p}_1\gamma$ ) and  $E_{\text{p}} = 6.1$  to 16.6 MeV (1979MU05;  $\text{p}_0, \text{p}_1$ ). Pronounced resonant structure is evident up to  $E_{\text{p}} = 14$  MeV (1979MU05). Observed resonances are displayed in Table 19.18. For other polarization measurements see (1981GL1B;  $E_{\text{p}} = 800$  MeV) and (1978AJ03). See also (1980FA06) and  $^{18}\text{O}$ .

32. (a)  $^{18}\text{O}(\text{p}, \text{d})^{17}\text{O}$ 

$$Q_{\text{m}} = -5.820$$

$$E_{\text{b}} = 7.9934$$

(b)  $^{18}\text{O}(\text{p}, \text{t})^{16}\text{O}$ 

$$Q_{\text{m}} = -3.7069$$

See (1978AJ03) and  $^{16}\text{O}, ^{17}\text{O}$  in (1982AJ01). See also (1979AM03; theor.).

Table 19.16: Resonances in  $^{18}\text{O}(p, \gamma)^{19}\text{F}$  <sup>a</sup>

$E_p$ (keV)	$\Gamma_{\text{lab}}$ (keV)	$\omega\gamma$ (eV)	$J^\pi$	$E_x$ (MeV)
$151 \pm 2$	$\leq 0.3$	$(1.0 \pm 0.1) \times 10^{-3}$	$\frac{1}{2}^+$	8.136 <sup>e</sup>
$216 \pm 1$	$\leq 1$	$> 0.8 \times 10^{-5}$		8.198
$274 \pm 3$	$\leq 1.5$	$(3.7 \pm 0.5) \times 10^{-5}$	$\leq \frac{7}{2}$	8.253
$334 \pm 2$	$\leq 1$	$(0.95 \pm 0.08) \times 10^{-3}$	$\frac{5}{2}^+$	8.310 <sup>f</sup>
$622 \pm 2$	$\leq 0.5$	$(10 \pm 2) \times 10^{-3}$	$\frac{5}{2}^+$	8.582
$629.6 \pm 0.3$	$2.0 \pm 0.3$	$0.10 \pm 0.02$	$\frac{3}{2}^3-$	8.5896 <sup>g</sup>
$\approx 680$	300		$\frac{3}{2}^3$	8.637
$841 \pm 2$	$48 \pm 2$	$1.4 \pm 0.2$	$\frac{1}{2}^+$ <sup>b</sup> $T = \frac{3}{2}$	8.790 <sup>h</sup>
$977 \pm 2$	$10 \pm 2$	$(1.5 \pm 0.2) \times 10^{-2}$	$\frac{3}{2}^3$	8.919
$1166.5 \pm 0.4$		$0.31 \pm 0.10$	$\frac{7}{2}^-$	9.0980 <sup>i</sup>
$1398 \pm 2$	$3.6 \pm 0.8$	$0.08 \pm 0.01$	$\frac{3}{2}^3+$	9.317
$1630 \pm 2$ <sup>c</sup>	$7 \pm 2$	$0.025 \pm 0.005$	$\frac{5}{2}^+$	9.537
$1660 \pm 3$	$27 \pm 3$	$0.041 \pm 0.010$	$\frac{3}{2}^3-$	9.565
$1670 \pm 4$	$70 \pm 3$	$0.06 \pm 0.01$	$\frac{3}{2}^3-$	9.575
$1684 \pm 4$	$8 \pm 2$	$0.025 \pm 0.004$	$\frac{7}{2}^-$	9.588
$1768 \pm 1.4$	$3.8 \pm 0.4$	$1.2 \pm 0.2$	$\frac{3}{2}^3+$	9.668
$1928.4 \pm 0.6$ <sup>d</sup>	$0.3 \pm 0.05$	$2.8 \pm 0.7$	$\frac{5}{2}^-$	9.819
$1986 \pm 2$	$< 1.5$	$0.13 \pm 0.04$	$\frac{11}{2}^-$	9.874
$1996 \pm 4$	$26 \pm 2$	$0.14 \pm 0.05$	$\frac{1}{2}^+$	9.883
$2263.0 \pm 0.7$	$5.0 \pm 1.0$		$\frac{3}{2}^-$	10.136
$> 2300$ <sup>d</sup>				

<sup>a</sup> Mostly from (1980WI17). For earlier references see Table 19.15 in (1978AJ03). See also Tables 19.7 and 19.15 here.

<sup>b</sup> Supported by direct capture into this state with a  $\sin^2 \theta$  distribution of the d.c.  $\gamma$ -rays and by interference patterns near the resonance (1980WI17).

<sup>c</sup> Decays partly (see Table 19.7) via a state at  $8015 \pm 2$  keV with  $J^\pi = \frac{5}{2}^+$  (1980WI17).

<sup>d</sup> See Table 19.15 in (1978AJ03).

<sup>e</sup>  $\Gamma_p = 0.17$  eV,  $\Gamma_\alpha = 220$  eV,  $\Gamma_\gamma = 1.3$  eV: see (1980WI17) for this footnote and the ones below.

<sup>f</sup>  $\Gamma_\gamma = 0.71 \pm 0.17$  eV,  $\Gamma_p = 0.019 \pm 0.009$  eV,  $\Gamma_\alpha = 46 \pm 19$  eV,  $\Gamma_{\text{total}} = 47 \pm 19$  eV.

<sup>g</sup>  $\Gamma_\gamma = 0.85 \pm 0.17$  eV,  $\Gamma_p = 224 \pm 43$  eV,  $\Gamma_\alpha = 3410 \pm 1220$  eV.

<sup>h</sup> The strength of the transition to  $^{19}\text{F}^*(7.262)$  [see Table 19.7] limits  $J$  to  $\frac{1}{2}$  or  $\frac{3}{2}$  for that state.

<sup>i</sup> The angular distribution of the  $\gamma$ -ray from this state to  $^{19}\text{F}^*(5.62)$  and branching ratio arguments lead to  $J = \frac{5}{2}$  for that state.

Table 19.17: Resonances in  $^{18}\text{O}(p, n)^{18}\text{F}$  <sup>a</sup>

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Res. in yield of <sup>b</sup>	$J\pi$	$E_x$ in $^{19}\text{F}$ (MeV)
$2.643 \pm 1.0$	$6.2 \pm 0.5$	n	$(\frac{3}{2})$	10.496
$2.691 \pm 1.0$	$2.5 \pm 0.2$	n		10.542
$2.717 \pm 1.0$	$5.2 \pm 0.5$	n		10.566
$2.767 \pm 1.5$	$4.7 \pm 0.5$	n	$\frac{5}{2}^{(+)}$	10.613
$2.923 \pm 4$	$6 \pm 3$	n		10.761
$3.025 \pm 2.0$	$24.0 \pm 1.5$	n	$\frac{3}{2}$	10.858
$(3.08 \pm 20)$	$\approx 60$	n		(10.91)
$3.148 \pm 3$	$14 \pm 2$	n		10.974
$3.164 \pm 2.5$	$7 \pm 2$	n		10.989
$3.250 \pm 2.5$	$35 \pm 4$	n	$\frac{3}{2}$	11.071
$3.370 \pm 4$	$17 \pm 4$	n		11.184
$3.463 \pm 3$	$7 \pm 2$	n		11.272
$3.470 \pm 15$	$70 \pm 20$	n		11.279
$3.653 \pm 4$	$40 \pm 10$	n, n <sub>1</sub>		11.452
$3.680 \pm 5$	$7 \pm 3$	n		11.478
$3.705 \pm 5$	$4 \pm 2$	n, n <sub>1</sub>		11.502
$3.748 \pm 15$	$50 \pm 15$	n		11.542
$3.775 \pm 7$	$15 \pm 10$	n, n <sub>2</sub>	$(T = \frac{3}{2})$	11.568
$(3.79 \pm 20)$	$60 \pm 20$	n		(11.58)
$3.863 \pm 4$	$45 \pm 10$	n, n <sub>1</sub>		11.651
4.00		n <sub>1</sub> , n <sub>3</sub>		(11.78)
$4.06 \pm 10$ <sup>c</sup>	$< 50$	n, n <sub>1</sub>		11.84
4.11		n <sub>1</sub>		(11.89)
$4.16 \pm 10$	90	n, n <sub>1</sub>		11.93
4.33		n <sub>1</sub> , n <sub>3</sub>		(12.09)
$4.37 \pm 10$	100	n, n <sub>1</sub> , n <sub>2</sub>		12.13
4.47	50	n, n <sub>1</sub> , n <sub>2</sub> , n <sub>3</sub>		12.23
$4.58 \pm 10$ <sup>d</sup>		n <sub>1</sub>		(12.33)
4.70		n <sub>3</sub>		(12.44)
4.83		n <sub>1</sub> , n <sub>2</sub> , n <sub>3</sub>		(12.57)
4.90		n <sub>2</sub>		(12.63)

Table 19.17: Resonances in  $^{18}\text{O}(\text{p}, \text{n})^{18}\text{F}$  <sup>a</sup> (continued)

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{c.m.}}$ (keV)	Res. in yield of <sup>b</sup>	$J^\pi$	$E_x$ in $^{19}\text{F}$ (MeV)
5.05 $\pm$ 10	200	n, n <sub>1</sub> , n <sub>2</sub>		12.77
5.10		n <sub>1</sub> , n <sub>2</sub>		(12.82)
5.20		n <sub>2</sub> , n <sub>3</sub>		(12.92)
5.35		n, n <sub>1</sub> , n <sub>2</sub> , n <sub>3</sub>		13.06
5.47 $\pm$ 15	70	n, n <sub>1</sub>		13.17
5.622 $\pm$ 15	30	n, n <sub>1</sub> , n <sub>2</sub>	( $T = \frac{3}{2}$ )	13.317
5.76		n <sub>1</sub> , n <sub>3</sub>		(13.45)
6.061 $\pm$ 15	50	n, n <sub>1</sub> , n <sub>2</sub>	( $T = \frac{3}{2}$ )	13.73
6.60 $\pm$ 15	350	n		14.24
(6.70 $\pm$ 15)		n		(14.34)
7.17 $\pm$ 20	300	n		14.78
7.40 $\pm$ 20		n		15.00
(7.8)		n		(15.4)
(7.98)		n		(15.55)
8.19 $\pm$ 25	150	n		15.75
8.74 $\pm$ 25	200	n		16.27
9.30 $\pm$ 30		n		16.80

<sup>a</sup> See Table 19.16 in (1978AJ03) for the references.

<sup>b</sup> n means total yield.

<sup>c</sup> Errors here and below are estimated from published data of (1964BA16) by H.B. Willard, private communication.

$$33. \text{}^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$$

$$Q_m = 3.9796$$

$$E_b = 7.9934$$

Yield measurements have been studied for  $E_p = 0.50$  to 14 MeV [see (1972AJ02)] and at  $E_p = 72$  to 935 keV (1979LO01;  $\alpha_0$ ), 235 to 698 keV (1978MA30;  $\alpha_0$ ) and 6.6 to 10.4 MeV (1979WI09;  $\alpha_0$ ): observed resonances are displayed in Table 19.18. The stellar reaction rates have been determined for  $T_9 = 0.008$  to 2.5 (1978RO1D). See also (1975ZI1A, 1978MA30; astrophys.) and (1978AJ03).



Table 19.18: Energy levels of  $^{19}\text{F}$  from  $^{18}\text{O}(\text{p}, \text{p})^{18}\text{O}$  and  $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$  <sup>a</sup>

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	Particles out	$\Gamma_p^b$ (keV)	$\Gamma_\alpha^b$ (keV)	$J^\pi$	$E_x$ (MeV)
$0.095 \pm 3^c$	$\leq 3$	$\alpha_0$	$\omega\gamma = (1.6 \pm 0.5) \times 10^{-7} \text{ eV}$			8.083
$0.152 \pm 1^c$	$\leq 0.5$	$\alpha_0$	$0.17 \pm 0.02 \text{ eV}$			8.137
$0.216 \pm 1^c$	$\leq 1$	$\alpha_0$	$(2.3 \pm 0.6) \times 10^{-3} \text{ eV}$			8.198
$0.334 \pm 1^c$	$\leq 1$	$\alpha_0$	$0.057 \pm 0.010 \text{ eV}$			8.310
$0.6326 \pm 0.4^d$	$2.1 \pm 0.1$	$p_0, \alpha_0$	$0.065 \pm 0.006$	$2.0 \pm 0.2$	$\frac{3}{2}^-$	8.5925
$\approx 0.695^c$	$\approx 340$	$p_0, \alpha_0$	$5^e$	$95^e$	$\frac{1}{2}^+$	8.65
$0.846 \pm 1.5$	$47 \pm 1$	$p_0, \alpha_0$	$26 \pm 1.5$	$21 \pm 1$	$\frac{1}{2}^+; T = \frac{3}{2}$	8.795
$0.9870 \pm 0.7$	$3.8 \pm 0.2$	$p_0, \alpha_0$	$0.080 \pm 0.007$	$3.7 \pm 0.3$	$\frac{3}{2}^-$	8.928
(1.135)	140					(9.068)
$1.1685 \pm 0.5$	$0.60 \pm 0.03$	$p_0, \alpha_0$	$0.005 \pm 0.0006$	$0.595 \pm 0.08$	$\frac{7}{2}^+$	9.0999
$1.2390 \pm 1$	$6.1 \pm 0.3$	$p_0, (\alpha_0)$	$0.40 \pm 0.03$	$5.7 \pm 0.4$	$\frac{1}{2}^+$	9.167
$1.4025 \pm 1$	$5.2 \pm 0.2$	$p_0, \alpha_0$	$0.23 \pm 0.02$	$5.0 \pm 0.4$	$\frac{1}{2}^+$	9.321
$1.620 \pm 6$	30	$p_0, \alpha_0$			$(\frac{5}{2})$	9.527
$1.668 \pm 6$	27	$p_0, \alpha_0$			$\frac{3}{2}^+$	9.573
$1.766 \pm 3$	3.6	$p_0, \alpha_0$	2.1	1.5	$\frac{3}{2}^+$	9.666
$1.928 \pm 3$	0.16	$p_0, \alpha_0$	0.09	0.07	$(\frac{5}{2}, \frac{7}{2})^-$	9.819
$2.001 \pm 4$	31	$p_0, \alpha_0$	12	19	$\frac{1}{2}^+$	9.888
$2.2630 \pm 0.7$	$5.0 \pm 1.0$	$\alpha_0, \alpha_1, \alpha_2$	$\approx 5$	$0.004^c$	$\frac{3}{2}^-$	10.136
$2.289 \pm 3$	33	$p_0, \alpha_0$	2.3	(1.0)	$\frac{1}{2}^+$	10.161
$2.363 \pm 3$	4.5	$p_0, \alpha_0$	2.8	1.7	$\frac{1}{2}^+$	10.231
$2.387 \pm 3$	24	$p_0, \alpha_0$	11	13	$\frac{3}{2}^+$	10.253
$2.443 \pm 4$	9.7	$p_0, \alpha_0$	5.2	4.5	$\frac{3}{2}^+$	10.307
$2.644 \pm 3$	4.6	$p_0, p_1, \alpha_0, \alpha_{1+2}$	2.4	(1.0)	$\frac{3}{2}^+$	10.497
$2.705 \pm 3$	$8 \pm 2$	$p_1, \alpha_0$			$\frac{3}{2}^{(+)}; (T = \frac{3}{2})$	10.555
$2.732 \pm 4$	$23 \pm 3$	$p_1, \alpha_0$			$(\frac{5}{2}^+)$	10.580
$2.768 \pm 3$	4.0	$p_0, p_1, \alpha_0, \alpha_{1+2}$	0.7	(1.0)	$\frac{5}{2}^+; T = \frac{3}{2}^a$	10.614
$2.925 \pm 3$	5.7	$p_0, p_1, \alpha_0, \alpha_{1+2}$	4.5	1.2	$\frac{1}{2}^-$	10.763
$3.029 \pm 4$	19.5	$p_0, p_1, \alpha_0, \alpha_{1+2}$	13.0		$\frac{5}{2}^+$	10.862
(3.06)		$\alpha_0$				(10.89)
$3.148 \pm 4$	(14)	$p_0, p_1, \alpha_0, \alpha_{1+2}$	(4.5)	(4.5)	$(\frac{3}{2}, \frac{5}{2})^+$	10.974
$3.266 \pm 9$	35	$p_0, p_1, \alpha_0, \alpha_{1+2}$			$\frac{1}{2}^+$	11.086
$3.386 \pm 9$	20	$p_0, p_1, \alpha_0, \alpha_{1+2}$			$(\frac{1}{2}^-)$	11.200
$3.479 \pm 8$	$23 \pm 5$	$p_0, p_1, \alpha_0, \alpha_{1+2}$	$4.3 \pm 1$		$\frac{5}{2}^+$	11.288
$3.547 \pm 25$	$286 \pm 33$	$p_0$	$241 \pm 2$		$\frac{1}{2}^+$	11.35
$3.643 \pm 9$	$40 \pm 7$	$p_0, (\alpha_{1+2})$	$17 \pm 3$		$\frac{1}{2}^-$	11.443
$3.694 \pm 9$	$29 \pm 6$	$p_0, p_1, \alpha_0, (\alpha_{1+2})$	$12 \pm 2$		$\frac{3}{2}^-$	11.491
$3.744 \pm 8$	$23 \pm 5$	$p_0, p_1, \alpha_0$	$3.7 \pm 1$		$\frac{5}{2}^+$	11.539

Table 19.18: Energy levels of  $^{19}\text{F}$  from  $^{18}\text{O}(\text{p}, \text{p})^{18}\text{O}$  and  $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$  <sup>a</sup> (continued)

$E_p$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	Particles out	$\Gamma_p^b$ (keV)	$\Gamma_\alpha^b$ (keV)	$J^\pi$	$E_x$ (MeV)
3.811 $\pm$ 12	66 $\pm$ 7	p <sub>0</sub>	30 $\pm$ 12		$\frac{3}{2}^-$	11.602
3.869 $\pm$ 8	28 $\pm$ 7	p <sub>0</sub> , p <sub>1</sub> , ( $\alpha_{1+2}$ )	12 $\pm$ 2		$\frac{3}{2}^+$ ; ( $T = \frac{3}{2}$ )	11.657
4.290 $\pm$ 30	75 $\pm$ 25	p <sub>0</sub> , $\alpha_0$ , $\alpha_{1+2}$	10 $\pm$ 3		$\frac{1}{2}^-$	12.06
4.390 $\pm$ 15	110 $\pm$ 15	p <sub>0</sub> , p <sub>1</sub> , ( $\alpha_0$ , $\alpha_{1+2}$ )	60 $\pm$ 10		$\frac{3}{2}^-$ ; $T = \frac{3}{2}$	12.150
4.465 $\pm$ 12	78 $\pm$ 1	p <sub>0</sub> , p <sub>1</sub> , $\alpha_0$ , $\alpha_{1+2}$	48 $\pm$ 6		$\frac{3}{2}^+$	12.221
4.782 $\pm$ 7	16 $\pm$ 4	p <sub>0</sub> , p <sub>1</sub>	2.4 $\pm$ 1		$\frac{1}{2}^-$	12.521
4.840 $\pm$ 10	50 $\pm$ 10	p <sub>0</sub> , p <sub>1</sub> , $\alpha_{1+2}$	6.4 $\pm$ 2		$\frac{5}{2}^+$	12.576
4.848 $\pm$ 25	300 $\pm$ 50	p <sub>0</sub>	80 $\pm$ 25		$\frac{1}{2}^-$ ; $T = \frac{3}{2}$	12.58
5.074 $\pm$ 30	100 $\pm$ 40	p <sub>0</sub> , p <sub>1</sub> , ( $\alpha_0$ )	13 $\pm$ 5		$\frac{5}{2}^+$ ; $T = \frac{3}{2}$	12.80
5.135 $\pm$ 30	290 $\pm$ 40	p <sub>0</sub> , p <sub>1</sub>	114 $\pm$ 17		$\frac{3}{2}^+$ ; $T = \frac{3}{2}$	12.86
5.225 $\pm$ 25	75 $\pm$ 25	p <sub>0</sub> , p <sub>1</sub> , $\alpha_{1+2}$	3 $\pm$ 1.5		$\frac{5}{2}^+$	12.94
5.27 $\pm$ 50	130 $\pm$ 40	p <sub>0</sub>	20 $\pm$ 8		$\frac{1}{2}^-$	12.98
5.38 $\pm$ 75	300 $\pm$ 75	p <sub>0</sub>	75 $\pm$ 25		$\frac{3}{2}^-$	13.09
5.622 $\pm$ 8	30 $\pm$ 6	p <sub>0</sub> , p <sub>1</sub> , $\alpha_0$ , $\alpha_{1+2}$	10 $\pm$ 3		$\frac{7}{2}^-$	13.317
5.670 $\pm$ 25	40 $\pm$ 20	p <sub>0</sub>	2 $\pm$ 2		$\frac{3}{2}^-$	13.36
6.060 $\pm$ 11	55 $\pm$ 10	p <sub>0</sub> , p <sub>1</sub> , ( $\alpha_{1+2}$ )	13 $\pm$ 3		$\frac{7}{2}^-$ ; $T = \frac{3}{2}$	13.731
6.390 $\pm$ 20 <sup>f</sup>	148 $\pm$ 30	p <sub>0</sub>	12 $\pm$ 3		$\frac{5}{2}^+$	14.04
6.428 $\pm$ 30	88 $\pm$ 30	p <sub>0</sub>	8 $\pm$ 3		$\frac{3}{2}^-$	14.08
6.687 $\pm$ 20	80 $\pm$ 30	p <sub>0</sub>	9 $\pm$ 3		$\frac{3}{2}^-$	14.32
7.080 $\pm$ 20	130 $\pm$ 40	p <sub>0</sub>	21 $\pm$ 5		$\frac{3}{2}^-$	14.70
7.10 $\pm$ 70 <sup>g</sup>	270 $\pm$ 70	$\alpha_0$			$\frac{1}{2}^-$	14.72
7.125 $\pm$ 50 <sup>f,g</sup>	380 $\pm$ 70	p <sub>0</sub> , $\alpha_0$	100 $\pm$ 25		$\frac{1}{2}^+$	14.74
7.167 $\pm$ 40	210 $\pm$ 50	p <sub>0</sub>	21 $\pm$ 6		$\frac{5}{2}^+$	14.78
7.337 $\pm$ 40	208 $\pm$ 30	p <sub>0</sub>	20 $\pm$ 4		$\frac{7}{2}^-$	14.94
7.775 $\pm$ 20	70 $\pm$ 10	p <sub>0</sub>	6 $\pm$ 2		$\frac{1}{2}^-$	15.35
7.820 $\pm$ 30	84 $\pm$ 25	p <sub>0</sub>	7 $\pm$ 2		$\frac{5}{2}^+$	15.40
8.282 $\pm$ 40	102 $\pm$ 25	p <sub>0</sub>	8 $\pm$ 3		$\frac{3}{2}^-$	15.83
8.670 $\pm$ 40	180 $\pm$ 30	p <sub>0</sub>	16 $\pm$ 4		$\frac{3}{2}^+$	16.20
8.695 $\pm$ 30	234 $\pm$ 40	p <sub>0</sub>	13 $\pm$ 4		$\frac{7}{2}^-$	16.23
8.747 $\pm$ 30	176 $\pm$ 30	p <sub>0</sub>	13 $\pm$ 4		$\frac{3}{2}^-$	16.27
9.563 $\pm$ 40	348 $\pm$ 70	p <sub>0</sub>	39 $\pm$ 8		$\frac{3}{2}^-$	17.05
9.679 $\pm$ 40	340 $\pm$ 70	p <sub>0</sub>	30 $\pm$ 8		$\frac{7}{2}^-$	17.16
9.986 $\pm$ 30	34 $\pm$ 20	p <sub>0</sub>	3 $\pm$ 2		$\frac{3}{2}^-$	17.45
10.200 $\pm$ 60	100 $\pm$ 60	p <sub>0</sub>	5 $\pm$ 3		$\frac{7}{2}^-$	17.65
10.496 $\pm$ 40	268 $\pm$ 60	p <sub>0</sub>	23 $\pm$ 5		$\frac{3}{2}^-$	17.93
10.596 $\pm$ 60	384 $\pm$ 60	p <sub>0</sub>	32 $\pm$ 7		$\frac{7}{2}^-$	18.02
11.698 $\pm$ 60	584 $\pm$ 150	p <sub>0</sub>	22 $\pm$ 7		$\frac{3}{2}^-$	19.07

Table 19.18: Energy levels of  $^{19}\text{F}$  from  $^{18}\text{O}(\text{p}, \text{p})^{18}\text{O}$  and  $^{18}\text{O}(\text{p}, \alpha)^{15}\text{N}$ <sup>a</sup> (continued)

$E_{\text{p}}$ (MeV $\pm$ keV)	$\Gamma_{\text{lab}}$ (keV)	Particles out	$\Gamma_{\text{p}}$ <sup>b</sup> (keV)	$\Gamma_{\alpha}$ <sup>b</sup> (keV)	$J^{\pi}$	$E_{\text{x}}$ (MeV)
12.499 $\pm$ 150	388 $\pm$ 60	p <sub>0</sub>	13 $\pm$ 6		$\frac{5}{2}^{-}$	19.83
12.547 $\pm$ 40	498 $\pm$ 60	p <sub>0</sub>	39 $\pm$ 8		$\frac{3}{2}^{-}$	19.87
13.542 $\pm$ 50	434 $\pm$ 60	p <sub>0</sub>	32 $\pm$ 5		$\frac{1}{2}^{-}$	20.81
13.662 $\pm$ 50	334 $\pm$ 50	p <sub>0</sub>	12 $\pm$ 4		$\frac{3}{2}^{-}$	20.93
13.791 $\pm$ 40	472 $\pm$ 30	p <sub>0</sub>	25 $\pm$ 5		$\frac{7}{2}^{-}$	21.05

<sup>a</sup> See also Tables 19.17 in (1978AJ03) and 19.14 in (1972AJ02) for the earlier work and references.

<sup>b</sup> See also Table 19.16.

<sup>c</sup> (1979LO01).

<sup>d</sup>  $\omega\gamma = 420 \pm 80$  eV (1979LO01).

<sup>e</sup> Widths not in accord with  $\Gamma$  measured by (1979LO01) who calculate also  $\omega\gamma \approx 1.2 \times 10^5$  eV.

<sup>f</sup> The parameters of this resonance and all the ones below [except for the two footnoted <sup>g</sup>] are from a phase-shift analysis by (1979MU05) of the elastic scattering for  $E_{\text{p}} = 6.1$  to 16.6 MeV. Other structures have also been observed but parameters for those have not been obtained.

<sup>g</sup> (1979WI09).

$$34. \text{ (a) } ^{18}\text{O}(\text{d}, \text{n})^{19}\text{F} \quad Q_{\text{m}} = 5.7685$$

$$\text{ (b) } ^{18}\text{O}(\text{d}, \text{n}\alpha)^{15}\text{N} \quad Q_{\text{m}} = 1.7546$$

Angular distributions of neutron groups corresponding to  $^{19}\text{F}$  states with  $E_{\text{x}} < 8.2$  MeV have been measured at  $E_{\text{d}} = 3$  and 4 MeV: see Table 19.18 in (1978AJ03). Gamma-ray measurements are reported here in Table 19.19 and  $\tau_{\text{m}}$  measurements in Table 19.8. For reaction (b) see (1972AJ02).

$$35. ^{18}\text{O}(\text{}^3\text{He}, \text{d})^{19}\text{F} \quad Q_{\text{m}} = 2.4998$$

Angular distributions of the deuterons corresponding to many states of  $^{19}\text{F}$  have been analyzed by DWBA: the results are shown in Table 19.19. The spectroscopic factors obtained for  $^{19}\text{F}^*(7.54, 8.80)$ , the  $T = \frac{3}{2}$ ,  $J^{\pi} = \frac{5}{2}^{+}$  and  $\frac{1}{2}^{+}$  analogs of  $^{19}\text{O}^*(0, 1.47)$  are in good agreement with those obtained for the  $^{19}\text{O}$  states in the  $^{18}\text{O}(\text{d}, \text{p})^{19}\text{O}$  reaction (1971FO18). See also (1979HA1M) and (1978AJ03).

$$36. ^{19}\text{O}(\beta^{-})^{19}\text{F} \quad Q_{\text{m}} = 4.819$$

Table 19.19: Energy levels of  $^{19}\text{F}$  from  $^{18}\text{O}(\text{d}, \text{n})^{19}\text{F}$  and  $^{18}\text{O}(^3\text{He}, \text{d})^{19}\text{F}$  <sup>a</sup>

$E_x$ <sup>b</sup> (MeV $\pm$ keV)	$l$ <sup>b</sup>	$C^2S(2J_f + 1)$ <sup>b</sup>	$J\pi$ <sup>b</sup>
0	0	0.42 <sup>a</sup>	$\frac{1}{2}^+$
$0.112 \pm 3$	1	0.224	$\frac{1}{2}^-$
$0.199 \pm 3$	2	2.45 <sup>a</sup>	$\frac{5}{2}^+$
$1.347 \pm 5$			
$1.460 \pm 5$	1	0.098	$\frac{3}{2}^-$
$1.5544 \pm 0.6$ <sup>c</sup>	2	1.01	$\frac{3}{2}^+$
$2.784 \pm 5$	4	0.027	$\frac{9}{2}^+$
$3.912 \pm 5$			
$3.999 \pm 1$ <sup>c</sup>	(3)	(0.019)	$(\frac{7}{2}^-)$
$4.036 \pm 10$			
$4.3761 \pm 0.8$ <sup>c</sup>	(4)	(0.048)	$(\frac{7}{2}^+)$
$4.5557 \pm 0.5$ <sup>c</sup>	2	0.31	<sup>a</sup>
$4.684 \pm 1$ <sup>c</sup>			
$5.113 \pm 5$	(2, 3)		$\frac{5}{2}^-, \frac{7}{2}^-$ <sup>a</sup>
$5.34 \pm 5$	(2, 3)	0.0065	$\frac{5}{2}^+$
$5.428 \pm 8$	(2, 3)	(0.042)	$(\frac{3}{2}^+)$
$5.492 \pm 5$			
$5.54 \pm 5$	3	0.14	$\frac{7}{2}^-$
$5.625 \pm 4$ <sup>c</sup>			
$5.943 \pm 5$	0	0.014	$\frac{1}{2}^+$
$6.095 \pm 5$	1	0.12	$\frac{1}{2}^-$
$6.167 \pm 5$			
$6.255 \pm 8$	(0)	0.19 <sup>a</sup>	$\frac{1}{2}^+$ <sup>a</sup>
$6.503 \pm 5$ <sup>e</sup>	2	0.133	$\frac{3}{2}^+$
$6.595 \pm 10$			
$6.792 \pm 5$	1	0.29 <sup>a</sup>	$\frac{3}{2}^-$
$6.93 \pm 5$	(2, 3)		$(\frac{5}{2}^+, \frac{7}{2}^-)$
$7.112 \pm 8$ <sup>e</sup>	2	0.087	$\frac{5}{2}^+$
$7.26 \pm 5$			
$7.364 \pm 5$	0	0.091	$\frac{1}{2}^+$
$7.540 \pm 3$	2	0.665	$\frac{5}{2}^+; T = \frac{3}{2}$

Table 19.19: Energy levels of  $^{19}\text{F}$  from  $^{18}\text{O}(\text{d}, \text{n})^{19}\text{F}$  and  $^{18}\text{O}(\text{}^3\text{He}, \text{d})^{19}\text{F}$  <sup>a</sup>  
(continued)

$E_x$ <sup>b</sup> (MeV $\pm$ keV)	$l$ <sup>b</sup>	$C^2S(2J_f + 1)$ <sup>b</sup>	$J\pi$ <sup>b</sup>
$7.665 \pm 5$	(2)	0.035 <sup>a</sup>	$(\frac{3}{2}^+)$
$7.702 \pm 5$	(0, 1)	(0.052)	$(\frac{3}{2}^-)$
$8.015 \pm 5$	2	0.26	$\frac{5}{2}^+$
$8.086 \pm 5$	(2, 3)	0.097	$(\frac{5}{2}^+)$
$8.135 \pm 5$	(0, 1)	0.156	$\frac{1}{2}^+$ <sup>a</sup>
$8.198 \pm 5$	(2, 3)	0.035	$(\frac{5}{2}^+)$
$8.255 \pm 5$	(2)	0.035	$(\frac{5}{2}^+)$
$8.31 \pm 5$			
$8.592 \pm 10$	(2, 3)		
$8.795 \pm 15$	0	(0.13)	$\frac{1}{2}^+; T = \frac{3}{2}$
$9.113 \pm 10$			
$9.18 \pm 15$			
$9.596 \pm 10$			
$9.682 \pm 15$			
$10.275 \pm 15$			
$10.33 \pm 15$			
$10.525 \pm 15$			

<sup>a</sup> See also Table 19.18 in (1978AJ03). Column 3 should refer to footnote <sup>c</sup>.

<sup>b</sup>  $^{18}\text{O}(\text{}^3\text{He}, \text{d})$ :  $E(\text{}^3\text{He}) = 16$  MeV (1970SC25), except where footnote is shown.

<sup>c</sup>  $^{18}\text{O}(\text{d}, \text{n}\gamma)$  (1975LE16).

<sup>d</sup>  $5.106 \pm 3$  (1975LE16).

<sup>e</sup> Unresolved.

The decay is primarily by allowed transitions to  $^{19}\text{F}^*(0.197, 1.55)$ ,  $J^\pi = \frac{5}{2}^+, \frac{3}{2}^+$ . Very weak branches are also observed to  $^{19}\text{F}^*(0.11, 1.35, 3.91, 4.39)$ ,  $J^\pi = \frac{1}{2}^-, \frac{5}{2}^-, \frac{3}{2}^+, \frac{7}{2}^+$ : see Table 19.20. The half-life is  $26.91 \pm 0.08$  sec: see reaction 1 in  $^{19}\text{O}$ . The character of the allowed decay to the  $\frac{5}{2}^+$  and  $\frac{3}{2}^+$  states, and the forbiddenness of the decay to the ground state of  $^{19}\text{F}$  are consistent with  $J^\pi = \frac{5}{2}^+$  for the ground state of  $^{19}\text{O}$ , and then with  $(\frac{7}{2}^+)$  for  $^{19}\text{F}^*(4.39)$ : see (1966OL01). Gamma-ray branching ratios are displayed in Table 19.7.

Time differential  $\beta - \gamma$  angular correlation measurements lead to a ratio  $+0.030 \pm 0.015$  for the Fermi to Gamow-Teller matrix elements ( $C_V M_F / C_A M_{GT}$ ) for the allowed  $\beta^-$  decay ( $\frac{5}{2}^+ \rightarrow \frac{5}{2}^+$ ) from  $^{19}\text{O}_{\text{g.s.}}$  to  $^{19}\text{F}_{0.197}$ . This leads to a value for  $M_F$  (isospin-forbidden fermi matrix element)

Table 19.20: Branching in  $^{19}\text{O}(\beta^-)^{19}\text{F}$  <sup>a</sup>

Decay to $^{19}\text{F}^*$ (keV) <sup>b</sup>	$J^\pi$	Branch (%)	$\log ft$
0	$\frac{1}{2}^+$	$\leq 4$	$\geq 6.5$
110	$\frac{1}{2}^-$	$0.055^{+0.013}_{-0.038}$	$8.34^{+0.30}_{-0.10}$
$197.143 \pm 0.004$	$\frac{5}{2}^+$	$45.4 \pm 1.5$	$5.384 \pm 0.014$
1346	$\frac{5}{2}^-$	$0.017 \pm 0.002$	$8.25 \pm 0.05$
1459	$\frac{3}{2}^-$	$< 0.010$	$> 8.4$
$1554.038 \pm 0.009$	$\frac{3}{2}^+$	$54.4 \pm 1.2$	$4.625 \pm 0.010$
$2779.849 \pm 0.034$	$\frac{9}{2}^+$	$< 0.002$	$> 8.2$
$3908.17 \pm 0.20$	$\frac{3}{2}^+$	$0.0081 \pm 0.0005$	$6.133 \pm 0.027$
3999	$\frac{7}{2}^-$	$< 0.001$	$> 6.9$
4033	$\frac{9}{2}^-$	$< 0.001$	$> 6.8$
$4377.700 \pm 0.042$	$\frac{7}{2}^+$	$0.0984 \pm 0.0030$	$3.859 \pm 0.017$
4550	$\frac{5}{2}^+$	$< 0.001$	$> 5.1$

<sup>a</sup> (1981OL1E). See Table 19.19 in (1978AJ03) for the earlier work.

<sup>b</sup>  $E_x$  shown with uncertainties were determined by (1981OL1E).

of  $(4.7 \pm 2.4) \times 10^{-3}$ . The admixture of the analog of the  $^{19}\text{O}_{\text{g.s.}}$  in the  $^{19}\text{F}$  excited state  $\alpha = (2.7 \pm 1.4) \times 10^{-3}$  and the effective charge-dependent matrix element  $|H_{\text{CD}}| = 20 \pm 10$  keV (1978PE14).

37. (a)  $^{19}\text{F}(\gamma, n)^{18}\text{F}$   $Q_m = -10.4313$   
 (b)  $^{19}\text{F}(\gamma, 2n)^{17}\text{F}$   $Q_m = -19.5819$   
 (c)  $^{19}\text{F}(\gamma, 2np)^{16}\text{O}$   $Q_m = -20.1823$

The cross section for  $(\gamma, Tn)$  has been measured for  $E_\gamma = 10.5$  to 28 MeV: it shows a clear resonance at  $E_\gamma \approx 12$  MeV and unresolved structures at higher energies. The integrated cross section to 29 MeV is  $108 \pm 7$  MeV·mb (1974VE06). For reports of other structures see (1972AJ02, 1978AJ03). See also (1979TH05) and  $^{18}\text{F}$ . For reactions (b) and (c) see (1978AJ03).

38.  $^{19}\text{F}(\gamma, p)^{18}\text{O}$   $Q_m = -7.9934$

The integrated cross sections for the  $p_0$  and  $p_1$  processes at  $90^\circ$  for  $E_\gamma = 13.3 \rightarrow 25.4$  and  $15.2 \rightarrow 26.0$  MeV are, respectively,  $1.80 \pm 0.27$  and  $0.50 \pm 0.45$  MeV · mb/sr. The  $(\gamma, p_0)$  cross section at  $90^\circ$  shows broad structures at  $E_\gamma = 15.0, 17.0$  and  $23$  MeV (1975TS03). For a report of other structures see (1978AJ03). See also (1979KE1B, 1979TH05) and  $^{18}\text{O}$ .

39.  $^{19}\text{F}(\gamma, t)^{16}\text{O}$   $Q_m = -11.7003$

This reaction has been studied for the transition to  $^{16}\text{O}_{\text{g.s.}}$  for  $E_\gamma = 18$  to  $23$  MeV. Two peaks are observed at  $E_\gamma = 18.8$  and  $20.1$  MeV: the angular distribution of  $t_0$  indicates  $J^\pi = \frac{1}{2}^-$  or  $\frac{3}{2}^-$ ,  $T = \frac{1}{2}$ . The triton GDR contributes  $\approx 1\%$  or the total GDR (1974SK04).

40.  $^{19}\text{F}(\gamma, \alpha)^{15}\text{N}$   $Q_m = -4.0138$

See (1979TH05) and  $^{15}\text{N}$  in (1981AJ01).

41.  $^{19}\text{F}(\gamma, \gamma)^{19}\text{F}$

The energy of the first excited state is  $109.894 \pm 0.005$  keV; its width is  $(5.1 \pm 0.7) \times 10^{-7}$  eV.  $^{19}\text{F}^*(1.46, 3.91, 7.66)$  are also excited.  $E_x = 7663 \pm 4$  keV for the latter. The scattering cross section is relatively small and structureless for  $E_\gamma = 14$  to  $30$  MeV. See (1978AJ03) for references.

42. (a)  $^{19}\text{F}(\text{e}, \text{e})^{19}\text{F}$   
 (b)  $^{19}\text{F}(\text{e}, \text{en})^{18}\text{F}$   $Q_m = -10.4313$   
 (c)  $^{19}\text{F}(\text{e}, \text{ep})^{18}\text{O}$   $Q_m = -7.9934$

The rms radius of  $^{19}\text{F} = 2.885 \pm 0.015$  fm (1973HA13). At  $E_e = 400$  MeV,  $0.5 \leq q \leq 2.5$  fm $^{-1}$ , the squared form factors have been determined for  $^{19}\text{F}^*(0, 1.55, 0.197, 5.46, 2.78, 4.65)$  [the  $J^\pi = \frac{1}{2}^+, \frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+, \frac{9}{2}^+, \frac{13}{2}^+$  members of the  $K = \frac{1}{2}^+$  ground-state rotational band] as well as for  $^{19}\text{F}^*(4.38)$  [ $J^\pi = \frac{7}{2}^+$ ]. The data are consistent for  $q < 1.8$  fm $^{-1}$  with transition strengths which were derived from a variation-after-projection, Hartree-Fock approximation (VPHF). However for the transitions to  $^{19}\text{F}^*(1.55, 0.197)$  ( $q > 1.8$  fm $^{-1}$ ) the relative magnitudes of the transitions require a strong M3 component in the  $\frac{5}{2}^+$  state and a weak component in the  $\frac{3}{2}^+$  state, consistent with predominant  $\mathbf{L} \cdot \mathbf{S}$  coupling. Including M5 and M7 strength improves the agreement with the data for  $^{19}\text{F}^*(2.78, 4.65)$  (1978WI01). The deformation parameters for the ground-state band are  $\beta_2 = 0.43 \pm 0.02$ ,  $\beta_4 = 0.12 \pm 0.02$  (1975OY01). Values of  $|M|^2$  adopted

by P.M. Endt (private communication) are  $8.9 \pm 0.5$  (E3),  $6.9 \pm 0.5$  (E2) and  $6.1 \pm 2.4$  W.u. (M5) for the g.s. transitions of  $^{19}\text{F}^*(1.35, 1.55, 2.78)$ : for other values see Table 19.20 in (1978AJ03). For reactions (b) and (c) see (1978AJ03).

43.  $^{19}\text{F}(n, n)^{19}\text{F}$

Angular distributions of neutron groups have been reported at  $E_n = 2.6, 14.1$  and  $14.2$  MeV: see (1972AJ02).  $E_x$  for the first six excited states of  $^{19}\text{F}$ , derived from  $\gamma$ -ray measurements, are  $109.8 \pm 0.2, 197.2 \pm 0.2, 1345.4 \pm 0.3, 1456.9 \pm 1.1, 1554.0 \pm 0.3$  and  $2775.1 \pm 3.5$  keV (1968SP01). See also (1978CO18, 1980CO1U), (1978AJ03), (1977NO07; theor.) and  $^{20}\text{F}$  here.

44. (a)  $^{19}\text{F}(p, p)^{19}\text{F}$

(b)  $^{19}\text{F}(p, 2p)^{18}\text{O} \quad Q_m = -7.9934$

Table 19.21 displays energy levels of  $^{19}\text{F}$  derived from this reaction. Angular distributions of various proton groups have been measured from  $E_p = 4.3$  to  $30$  MeV [see (1978AJ03)] and at  $35.2$  MeV (1980FA07;  $p_0$ ). The ground-state rotational band is characterized by  $\beta_2 = 0.44 \pm 0.04$ ,  $\beta_4 = 0.14 \pm 0.04$  (1973DE06, 1974DE46). The  $g$  of  $^{19}\text{F}^*(0.197) = 1.47 \pm 0.03$  (1978KR05). The mixing ratio for the  $1.46 \rightarrow 0.11$  transition ( $\frac{3}{2}^- \rightarrow \frac{1}{2}^-$ ;  $K = \frac{1}{2}^-$  band),  $\delta(\text{E2/M1})$  has been remeasured by (1980DI12) to be  $0.224 \pm 0.040$ . A re-analysis of earlier experiments leads to a “best” value of  $0.248 \pm 0.020$  for the ratio and to an E2 strength of  $18.7 \pm 1.9$  W.u. The  $1.46 \rightarrow 0$  transition is pure E1 ( $\delta = 0.01 \pm 0.03$ ) (1980DI12). See Table 2 in the Introduction in this publication, and Table 19.7. For reaction (b) see  $^{18}\text{O}$ . See also (1981KE1E, 1982FI1C), (1978SH1U, 1980KR1C; applied), (1977PH02, 1979ZE04, 1980KO1V; theor.) and  $^{20}\text{Ne}$ .

45.  $^{19}\text{F}(d, d)^{19}\text{F}$

Angular distributions of elastically scattered deuterons has been measured for  $E_d = 2.0$  to  $15$  MeV [see (1972AJ02)] for the earlier references] and for  $1.6$  to  $2.4$  MeV (1978AS07). Angular distributions have also been measured for  $d_1 \rightarrow d_5$  and  $B(\text{E}\lambda)$  have been derived (1970DE06;  $E_d = 15$  MeV). For polarization measurements see (1978AJ03). See also (1979AN35).

46.  $^{19}\text{F}(t, t)^{19}\text{F}$

Elastic angular distributions have been measured at  $E_t = 2$  and  $7.2$  MeV: see (1972AJ02).



Table 19.21: States of  $^{19}\text{F}$  from  $^{19}\text{F}(\text{p}, \text{p}')^{19}\text{F}^*$

$E_x$ (keV) <sup>a</sup>	$L$ <sup>b</sup>	$\beta_L$ <sup>b</sup>	$J^\pi$
$197.6 \pm 0.6$	2	0.55	$\frac{5}{2}^+$
$1345.8 \pm 0.2$	3	0.33	$\frac{5}{2}^-$
$1458.8 \pm 0.3$			$\frac{3}{2}^-$
$1554.0 \pm 0.4$	2	0.58	$\frac{3}{2}^+$
$2779.8 \pm 0.6$	4	0.22	$\frac{9}{2}^+$
$3907.1 \pm 1.0$			$\frac{3}{2}^+$
$3998.5 \pm 0.8$			$\frac{7}{2}^-$
$4032.5 \pm 2$			$\frac{9}{2}^-$
$4377.7 \pm 1.0$			$\frac{7}{2}^+$
$4548.8 \pm 1.0$	2	0.20	$\frac{5}{2}^+$
$4557.5 \pm 1.0$			$\frac{3}{2}^-, (\frac{1}{2}^-)$
$4682.5 \pm 1.2$	<sup>c</sup>		
$5110 \pm 10$	2	0.15 <sup>d</sup>	$\frac{5}{2}^{(-)}$
$5340 \pm 10$			
$5420 \pm 10$	3	0.45	$\frac{7}{2}^-$
$5470 \pm 10$			
$5500 \pm 10$			
$5540 \pm 10$			
$5630 \pm 10$	<sup>e</sup>		
$5940 \pm 10$			
(6080)			
$6090 \pm 10$			
$6170 \pm 10$			
$6250 \pm 10$			
$6290 \pm 10$			
$6330 \pm 10$			

<sup>a</sup> For references see Table 19.21 in (1978AJ03).

<sup>b</sup> (1974DE46):  $E_p = 30$  MeV.

<sup>c</sup> (1974DE46) report excitation of a state with  $E_x = 4.69$  MeV,  $J^\pi = \frac{3}{2}^-$ ,  $L = 3$ ,  $\beta_L = 0.17$ .

<sup>d</sup> If  $L = 2$ .

<sup>e</sup> (1974DE46) report excitation of a state with  $E_x = 5.63$  MeV,  $J^\pi = \frac{5}{2}^-$ ,  $L = 3$ ,  $\beta_L = 0.33$ .

47. (a)  $^{19}\text{F}(^3\text{He}, ^3\text{He})^{19}\text{F}$   
 (b)  $^{19}\text{F}(^3\text{He}, ^6\text{He})^{16}\text{F}$   $Q_m = -14.842$

Elastic angular distributions have been measured for  $E(^3\text{He}) = 4.0$  to 29 MeV: see (1972AJ02, 1978AJ03). For reaction (b) see (1978AJ03) and  $^{16}\text{F}$  in (1982AJ01). See also (1979GR11; theor.).

48. (a)  $^{19}\text{F}(\alpha, \alpha)^{19}\text{F}$   
 (b)  $^{19}\text{F}(\alpha, \alpha p)^{18}\text{O}$   $Q_m = -7.9934$   
 (c)  $^{19}\text{F}(\alpha, 2\alpha)^{15}\text{N}$   $Q_m = -4.0138$   
 (d)  $^{19}\text{F}(\alpha, \alpha t)^{16}\text{O}$   $Q_m = -11.7003$

Angular distributions of elastically scattered  $\alpha$ -particles have been measured at  $E_\alpha = 19.9$  to 23.3 MeV and at 38 MeV [see (1972AJ02)]. Many inelastic groups have also been studied: see Table 19.22 in (1978AJ03).

The energy of the  $\gamma$ -ray from  $1.35 \rightarrow 0.11$  transition is  $1235.8 \pm 0.2$  keV. Using  $E_x = 109.894 \pm 0.005$  keV for the energy of the first excited state,  $E_x$  for  $^{19}\text{F}^*$  is then  $1345.7 \pm 0.2$  keV (1967WA13). At  $E_\alpha = 12.7$  MeV, a state at 4.648 MeV is populated which is then observed to  $\gamma$ -decay to the  $\frac{9}{2}^+$  state at 2.78 MeV. The angular distribution of the cascade  $\gamma$ -rays and the lifetime of  $^{19}\text{F}^*(4.65)$ , set  $J^\pi = \frac{13}{2}^+$  for  $^{19}\text{F}^*(4.65)$  (1969JA09). For reaction (b) see  $^{18}\text{O}$  (1979NA06). For reactions (c) and (d) see (1972CH18;  $E_\alpha = 60.2$  MeV).

49. (a)  $^{19}\text{F}(^6\text{Li}, ^6\text{Li})^{19}\text{F}$   
 (b)  $^{19}\text{F}(^7\text{Li}, ^7\text{Li})^{19}\text{F}$

Elastic angular distributions have been reported in both reactions at  $E(\text{Li}) = 20$  and 34 MeV: see (1978AJ03).

50. (a)  $^{19}\text{F}(^{10}\text{B}, ^{10}\text{B})^{19}\text{F}$   
 (b)  $^{19}\text{F}(^{12}\text{C}, ^{12}\text{C})^{19}\text{F}$

For reaction (a) see (1971KN05). Angular distributions [reaction (b)] involving  $^{19}\text{F}^*(0, 0.197, 1.55)$  have been studied at  $E(^{12}\text{C}) = 40.6$  MeV (1980KA1T). See also (1978AJ03). Fusion cross section measurements are reported by (1976SP07, 1977KO38, 1979KO20). See also (1980CO08, 1981MA1Q) and (1978BI1G, 1978HO13, 1978VA1A, 1979NA03, 1980LO02; theor.).

51. (a)  $^{19}\text{F}(^{14}\text{N}, ^{14}\text{N})^{19}\text{F}$   
(b)  $^{19}\text{F}(^{15}\text{N}, ^{15}\text{N})^{19}\text{F}$

Elastic scattering angular distributions have been studied at  $E(^{14}\text{N}) = 19.5$  MeV (1977KU06) and at  $E(^{15}\text{N}) = 23, 26$  and  $29$  MeV (1973GA14).

52. (a)  $^{19}\text{F}(^{16}\text{O}, ^{16}\text{O})^{19}\text{F}$   
(b)  $^{19}\text{F}(^{18}\text{O}, ^{18}\text{O})^{19}\text{F}$

Elastic angular distributions have been studied at  $E(^{16}\text{O}) = 21.4$  and  $25.8$  MeV and at  $E(^{19}\text{F}) = 27, 30, 33$  and  $36$  MeV (reaction (a)) [also to  $^{19}\text{F}^*(1.46)$  at the two higher energies] and at  $27, 30$  and  $33$  MeV [reaction (b)]: see (1978AJ03). See also (1979GA1H) and (1980OH05; theor.).

53.  $^{19}\text{F}(^{20}\text{Ne}, ^{20}\text{Ne})^{19}\text{F}$

See (1972AJ02) and (1979SI1K; theor.).

54.  $^{19}\text{F}(^{23}\text{Na}, ^{23}\text{Na})^{19}\text{F}$

For fusion cross sections see (1978HO20). See also (1978HO13, 1980LE11; theor.).

55.  $^{19}\text{F}(^{24}\text{Mg}, ^{24}\text{Mg})^{19}\text{F}$

For fusion cross sections see (1981LE04).

56.  $^{19}\text{F}(^{27}\text{Al}, ^{27}\text{Al})^{19}\text{F}$

For fusion measurements see (1981CH32). See also (1980SA1L).

57. (a)  $^{19}\text{F}(^{28}\text{Si}, ^{28}\text{Si})^{19}\text{F}$   
(b)  $^{19}\text{F}(^{30}\text{Si}, ^{30}\text{Si})^{19}\text{F}$

See (1980KU04) for the lelastic scattering at  $E(\text{Si}) = 60$  MeV. For fusion measurements see (1981CH32). See also (1979GO13; theor.).

58.  $^{19}\text{Ne}(\beta^+)^{19}\text{F}$   $Q_m = 3.2383$

See  $^{19}\text{Ne}$ .

59.  $^{20}\text{Ne}(\gamma, \text{p})^{19}\text{F}$   $Q_m = -12.8447$

See (1972AJ02).

60.  $^{20}\text{Ne}(\text{n}, \text{d})^{19}\text{F}$   $Q_m = -10.6200$

See (1978AJ03).

61.  $^{20}\text{Ne}(\text{d}, ^3\text{He})^{19}\text{F}$   $Q_m = -7.3515$

At  $E_d = 52$  MeV  $^3\text{He}$  angular distributions are reported corresponding to some unresolved states in  $^{19}\text{F}$  and to  $^{19}\text{F}^*(2.78, 5.62, 10.41)$  [the first two involve  $l = 4$  and  $l = 1$ ]: see (1978AJ03).

62.  $^{20}\text{Ne}(\text{t}, \alpha)^{19}\text{F}$   $Q_m = 6.9694$

See Table 19.22.

63.  $^{21}\text{Ne}(\text{p}, ^3\text{He})^{19}\text{F}$   $Q_m = -11.888$

At  $E_p = 45$  MeV,  $^3\text{He}$  groups are observed to some  $T = \frac{1}{2}$  states in  $^{19}\text{F}$  and to  $\frac{3}{2}^+$ ,  $T = \frac{3}{2}$  analog of  $^{19}\text{O}^*(0.095)$ :  $E_x = 7.660 \pm 0.035$  MeV (1969HA38). At  $E_p = 40$  MeV comparison of the ground-state angular distributions in this reaction and in the mirror (p, t) reaction [see reaction 18 in  $^{19}\text{Ne}$ ] shows a suppression of the  $S = 1, T = 0$  component of the (p,  $^3\text{He}$ ) cross section: this is observed to occur for all  $T_Z = \frac{1}{2}$  nuclei with  $A < 40$  (1976NA18).

Table 19.22: States of  $^{19}\text{F}$  and  $^{19}\text{Ne}$  from  $^{20}\text{Ne}(t, \alpha)$  and  $^{20}\text{Ne}(^3\text{He}, \alpha)$

$E_x$ in $^{19}\text{F}$ (MeV $\pm$ keV) <sup>a</sup>	$l_p$ <sup>b</sup>	$J^\pi$	$C^2S$ <sup>b,c</sup>		$E_x$ in $^{19}\text{Ne}$ (MeV)
			(t, $\alpha$ )	( $^3\text{He}, \alpha$ )	
0	0	$\frac{1}{2}^+$	0.12	0.20	0
0.11	1	$\frac{1}{2}^-$	1.7	1.8	0.28
0.20	2	$\frac{5}{2}^+$	1.6	0.95	0.24
1.46	1	$\frac{3}{2}^-$	0.30	0.21	1.62
1.55	2	$\frac{3}{2}^+$	0.31 <sup>d</sup>	0.70	1.54
$2.794 \pm 15$					
$3.917 \pm 15$		$\frac{3}{2}^+$	$\leq 0.04$	$\leq 0.1$	4.03
4.00					
$4.032 \pm 15$					
$4.385 \pm 15$					
$4.55 + 4.56$	1	$\frac{3}{2}^-$	0.69	0.57	4.55
$4.65 + 4.68$					
$5.102 \pm 15$					
$5.343 \pm 15$					
$5.481 \pm 15$					
$5.539 \pm 15$					
$5.628 \pm 15$					
$5.937 \pm 20$					
$6.092 \pm 15$	1	$\frac{3}{2}^-$	1.0	1.4	6.01
$6.169 \pm 30$					
$6.247 \pm 25$					
$6.501 \pm 25$					
6.79	1	$\frac{3}{2}^-$	0.96	1.5	6.74

<sup>a</sup> For references see Table 19.23 in (1978AJ03).  $E_x$  for which errors are not shown are nominal.

<sup>b</sup> (1974GA28):  $E_t = 20$  MeV.

<sup>c</sup> Calculated using finite range and non-local corrections. The ( $^3\text{He}, \alpha$ ) results are from (1970GA18). The absolute DWBA normalization factors were 4.6 for (t,  $\alpha$ ) and 10.2 for ( $^3\text{He}, \alpha$ ).

<sup>d</sup> Poor DWBA fit.

64.  $^{21}\text{Ne}(\text{d}, \alpha)^{19}\text{F}$   $Q_{\text{m}} = 6.465$

See (1978AJ03).

65.  $^{22}\text{Ne}(\text{p}, \alpha)^{19}\text{F}$   $Q_{\text{m}} = -1.675$

The parity-non-conserving asymmetry of the 110 keV  $\gamma$ -rays emitted by polarized  $^{19}\text{F}^*$  nuclei is  $\delta = -(1.8 \pm 0.9) \times 10^{-4}$  (1975AD01). See also (1978AJ03).

66.  $^{23}\text{Na}(\text{n}, \text{n}'\alpha)^{19}\text{F}$   $Q_{\text{m}} = -10.467$

See (1978AJ03).

<sup>19</sup>Ne  
(Figs. 7 and 8)

GENERAL: (See also (1978AJ03).)

*Nuclear models:* (1978MA2H, 1978PE09, 1978PI06, 1979DA15, 1979MA27, 1979PE16, 1982KI02).

*Electromagnetic transitions:* (1978PE09, 1978SC19, 1979MA27, 1979PE16).

*Special states:* (1978MA2H, 1978PE09, 1978PI06, 1978SC19, 1979DA15, 1980OK01, 1982KI02).

*Astrophysical questions:* (1977SIID, 1978WO1E, 1979RA1C).

*Applied topics:* (1979AL1Q).

*Complex reactions involving <sup>19</sup>Ne:* (1978SH18, 1981GR08).

*Other topics:* (1978MA2H, 1979BE1H, 1979MA27, 1979PE16, 1982KI02).

*Ground state of <sup>19</sup>Ne:* (1978MA54, 1979MA27, 1979SA41, 1979SA43).

$$\mu_{\text{g.s.}} = -1.88542 \pm 0.00006 \text{ nm (1980MA1G, 1982MAZY; prelim. results).}$$

$$\mu_{0.239} = -0.740 \pm 0.008 \text{ nm (1978LEZA).}$$

1. <sup>19</sup>Ne( $\beta^+$ )<sup>19</sup>F  $Q_m = 3.2383$

The half-life of <sup>19</sup>Ne is  $17.22 \pm 0.02$  sec: see (1978AJ03). The decay is principally to <sup>19</sup>F<sub>g.s.</sub> [ $\log ft = 3.237 \pm 0.001$  (1976AL07)]. The allowed nature of the decay to the ground state of <sup>19</sup>F sets  $J^\pi = \frac{1}{2}^+$  for <sup>19</sup>Ne<sub>g.s.</sub>. Very weak branches are also observed to <sup>19</sup>F\*(0.110) [ $J^\pi = \frac{1}{2}^-$ ],  $(1.20 \pm 0.20) \times 10^{-2}\%$ ,  $\log ft = 7.05 \pm 0.08$  (1981AD05) and to <sup>19</sup>F\*(1.55) [ $J^\pi = \frac{3}{2}^+$ ],  $(2.1 \pm 0.3) \times 10^{-3}\%$ ,  $\log ft = 5.72 \pm 0.06$ . The transition <sup>19</sup>F\*(1.55  $\rightarrow$  0.20) involves  $E_\gamma = 1356.92 \pm 0.15$  keV (1976AL07). See also (1982LOZZ) and (1978AJ03).

The <sup>19</sup>Ne decay to <sup>19</sup>F\*(0.11) [ $J^\pi = \frac{1}{2}^+ \rightarrow \frac{1}{2}^-$ ] proceeds by vector and axial vector weak currents, with the former making a negligible contribution. The measured decay rate is roughly an order of magnitude smaller than predicted using standard wave functions (1981AD05). See also (1978KL1D, 1979SC1K, 1982SC1C, 1982SCZZ), (1977GA1E, 1977TE1B, 1978AJ03, 1978CA1H, 1978RA2A, 1978WE1J, 1979CA1K) and (1977KL09, 1980AF1A, 1980OK01, 1981HA1Q; theor.).

2. <sup>12</sup>C(<sup>12</sup>C,  $\alpha$ n)<sup>19</sup>Ne  $Q_m = -12.2473$

See (1980KO02).

Table 19.23: Energy levels of  $^{19}\text{Ne}$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
0	$\frac{1}{2}^+; \frac{1}{2}$	$\frac{1}{2}^+$	$\tau_{1/2} = 17.22 \pm 0.02$ sec	$\beta^+$	1, 2, 3, 5, 6, 7, 11, 12, 13, 14, 15, 16, 17, 18
$0.23827 \pm 0.11$	$\frac{5}{2}^+$	$\frac{1}{2}^+$	$\tau_m = 26.0 \pm 0.8$ nsec $g = -0.296 \pm 0.003$	$\gamma$	6, 7, 9, 11, 12, 13, 17, 18
$0.27509 \pm 0.13$	$\frac{1}{2}^-$	$\frac{1}{2}^-$	$\tau_m = 61.4 \pm 3.0$ psec	$\gamma$	6, 7, 11, 17
$1.50756 \pm 0.3$	$\frac{5}{2}^-$	$\frac{1}{2}^-$	$1.4^{+0.5}_{-0.6}$ psec	$\gamma$	6, 7, 11, 17
$1.5360 \pm 0.4$	$\frac{3}{2}^+$	$\frac{1}{2}^+$	$28 \pm 11$ fsec	$\gamma$	6, 7, 9, 11, 12, 13, 17
$1.6156 \pm 0.5$	$\frac{3}{2}^-$	$\frac{1}{2}^-$	$143 \pm 31$ fsec	$\gamma$	6, 7, 11, 17
$2.7947 \pm 0.6$	$\frac{9}{2}^+$	$\frac{1}{2}^+$	$140 \pm 35$ fsec	$\gamma$	4, 6, 7, 8, 9, 11, 12, 13, 17, 18
$4.0329 \pm 2.4$	$\frac{3}{2}^+$		$< 50$ fsec	$\gamma$	7, 10, 17, 18
$4.140 \pm 4$	$(\frac{9}{2})^-$	$(\frac{1}{2})^-$	$< 0.3$ psec	$\gamma$	7, 10, 17
$4.1971 \pm 2.4$	$(\frac{7}{2})^-$	$(\frac{1}{2})^-$	$< 0.35$ psec	$\gamma$	6, 7, 10, 17
$4.3791 \pm 2.2$	$\frac{7}{2}^+$		$< 0.12$ psec	$\gamma$	7, 10, 13, 17
$4.549 \pm 4$	$(\frac{1}{2}, \frac{3}{2})^-$		$< 80$ fsec	$\gamma$	7, 10, 17
$4.600 \pm 4$	$(\frac{5}{2})^+$		$< 0.16$ psec	$\gamma$	7, 10
$4.635 \pm 4$	$\frac{13}{2}^+$	$\frac{1}{2}^+$	$> 1$ psec	$\gamma$	4, 6, 7, 8, 9, 10, 17
$4.712 \pm 10$	$(\frac{5}{2})^-$				7
$4.783 \pm 20$					17
$5.092 \pm 6$	$\frac{5}{2}^+$			$\gamma$	7, 10, 17, 18
$5.351 \pm 10$	$\frac{1}{2}^+$				17
$5.424 \pm 7$	$(\frac{7}{2})^+$	$(\frac{1}{2})^+$			6, 7, 17
$5.463 \pm 20$					17
$5.539 \pm 9$					17
$5.832 \pm 9$					17
$6.013 \pm 7$	$(\frac{3}{2}, \frac{1}{2})^-$				17
$6.092 \pm 8$					7, 17
$6.149 \pm 20$					17
$6.288 \pm 7$					7, 17
$6.437 \pm 9$					17
$6.742 \pm 7$	$(\frac{3}{2}, \frac{1}{2})^-$				17
$6.861 \pm 7$					7, 17
$7.067 \pm 9$					17
$7.21 \pm 20$					7, 17



Table 19.23: Energy levels of  $^{19}\text{Ne}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
7.253 $\pm$ 10					17
(7.326 $\pm$ 15)					17
(7.531 $\pm$ 15)					17
7.616 $\pm$ 16	$\frac{3}{2}^+; \frac{3}{2}$				6, 17, 18
7.700 $\pm$ 10					17
(7.788 $\pm$ 10)					17
7.994 $\pm$ 15					17
8.069 $\pm$ 12					7, 17
8.236 $\pm$ 10 <sup>b</sup>					17
8.442 $\pm$ 9					6, 7, 17
8.523 $\pm$ 10					17
(8.810 $\pm$ 25)					17
8.920 $\pm$ 9					6, 7, 8, 17
9.013 $\pm$ 10					17
9.100 $\pm$ 20					17
9.240 $\pm$ 20					6, 17
9.489 $\pm$ 25					17
9.81 $\pm$ 20					6, 7, 8, 9, 17
10.01 $\pm$ 20					7
			$\Gamma_{\text{c.m.}} =$ (keV)		
10.407 $\pm$ 30 <sup>b</sup>	$\frac{3}{2}^+$		45	p, $^3\text{He}$ , $\alpha$	5, 6, 17
10.46	$\frac{1}{2}^+$		355	p, $^3\text{He}$ , $\alpha$	5
10.613 $\pm$ 20					17
11.08 $\pm$ 20					6, 7, 8
11.24 $\pm$ 20					7
11.40 $\pm$ 20					7
11.51 $\pm$ 50	$\frac{3}{2}^-, (\frac{1}{2}^-)$		25	$^3\text{He}$ , $\alpha$	5
12.23 $\pm$ 50	$\frac{5}{2}^+$		200 $\pm$ 25	$^3\text{He}$ , $\alpha$	5, 8, 9
12.40 $\pm$ 50	$\frac{7}{2}^+$		180 $\pm$ 25	$^3\text{He}$ , $\alpha$	5
12.56 $\pm$ 20					7
12.69 $\pm$ 50	$\frac{1}{2}^+$		180 $\pm$ 40	p, $^3\text{He}$	5
13.1 $\pm$ 30					7
13.22 $\pm$ 30					7

Table 19.23: Energy levels of  $^{19}\text{Ne}$  <sup>a</sup> (continued)

$E_x$ (MeV $\pm$ keV)	$J^\pi; T$	$K^\pi$	$\tau$ or $\Gamma_{\text{c.m.}}$ (keV)	Decay	Reactions
13.8 $\pm$ 250			670 $\pm$ 250	$\gamma, ^3\text{He}$	5
14.18 $\pm$ 30					7, 8
14.44 $\pm$ 30					7
14.78 $\pm$ 30			620 $\pm$ 130	$\gamma, ^3\text{He}$	5, 7
16.23 $\pm$ 130			400 $\pm$ 130	$\gamma, \text{n}, ^3\text{He}$	5
18.4 $\pm$ 500			4400 $\pm$ 500	$\gamma, ^3\text{He}$	5

<sup>a</sup> See also Table 19.24.


<sup>b</sup> Broad or unresolved states.



See (1978WU1C).



At  $E(^{12}\text{C}) = 115$  MeV  $^{19}\text{Ne}^*(2.79, 4.64)$  are populated (1979PA06).



Excitation functions at  $\theta = 40^\circ$  and  $90^\circ$  have been measured for  $\gamma_{0 \rightarrow 2}$ ,  $\gamma_{3 \rightarrow 5}$  and  $\gamma_6$  [reaction (a)] for  $E(^3\text{He}) = 3$  to 19 MeV: see Table 19.25 for a listing of the observed structures (1981WA1R). A resonance-like structure is reported at  $E(^3\text{He}) \approx 9.5$  MeV in the yield of neutron groups [reaction (b)] (1970AD02).

Table 19.24: Radiative decay of  $^{19}\text{Ne}$  levels <sup>a</sup>

$E_i$ (MeV) <sup>b</sup>	$J_i^\pi$	$E_f$ (MeV)	$J_f^\pi$	Branch (%)	$\tau_m$
0.24	$\frac{5}{2}^+$	0	$\frac{1}{2}^+$	100	$26.0 \pm 0.8$ nsec
0.28	$\frac{1}{2}^-$	0	$\frac{1}{2}^+$	(100) <sup>c</sup>	$61.4 \pm 3.0$ psec
1.51	$\frac{5}{2}^-$	0.24	$\frac{5}{2}^+$	$12 \pm 3$	$1.4^{+0.5}_{-0.6}$ psec
		0.28	$\frac{1}{2}^-$	$88 \pm 3$ <sup>d</sup>	
1.54	$\frac{3}{2}^+$	0.24	$\frac{5}{2}^+$	$95 \pm 3$ <sup>d</sup>	$28 \pm 11$ fsec
		0.28	$\frac{1}{2}^-$	$5 \pm 3$	
1.62	$\frac{3}{2}^-$	0	$\frac{1}{2}^+$	$20 \pm 3$ <sup>d</sup>	$143 \pm 31$ fsec
		0.24	$\frac{5}{2}^+$	$10 \pm 3$	
2.79	$\frac{9}{2}^+$	0.24	$\frac{5}{2}^+$	100 <sup>d</sup>	$140 \pm 35$ fsec
		4.03	$\frac{3}{2}^+$	$80 \pm 15$	
4.03	$\frac{3}{2}^+$	0.28	$\frac{1}{2}^-$	$5 \pm 5$	$< 50$ fsec
		1.54	$\frac{3}{2}^+$	$15 \pm 5$	
4.14	$(\frac{9}{2})^-$	1.51	$\frac{5}{2}^-$	100	$< 0.3$ psec
4.20	$(\frac{7}{2})^-$	0.24	$\frac{5}{2}^+$	$20 \pm 5$	$< 0.35$ psec
		1.51	$\frac{5}{2}^-$	$80 \pm 5$	
4.38	$\frac{7}{2}^+$	0.24	$\frac{5}{2}^+$	$85 \pm 4$	$< 0.12$ psec
		2.79	$\frac{9}{2}^+$	$15 \pm 4$	
4.55	$(\frac{1}{2}, \frac{3}{2})^-$	0	$\frac{1}{2}^+$	$35 \pm 25$	$< 80$ fsec
		0.28	$\frac{1}{2}^-$	$65 \pm 25$	
4.60	$(\frac{5}{2}^+)$	0.24	$\frac{5}{2}^+$	$90 \pm 5$	$< 0.16$ psec
		1.54	$\frac{3}{2}^+$	$10 \pm 5$	
4.64	$\frac{13}{2}^+$	2.79	$\frac{9}{2}^+$	100	$> 1$ psec

<sup>a</sup> See Table 19.26 in (1978AJ03) for additional data and for references.

<sup>b</sup>  $E_x = 238.27 \pm 0.11, 275.09 \pm 0.13, 1507.56 \pm 0.3, 1536.0 \pm 0.4, 1615.6 \pm 0.5$  and  $2794.7 \pm 0.6$  keV from  $E_\gamma$  measurements: see Table 19.25 in (1978AJ03).

<sup>c</sup>  $B(E1) = (1.06 \pm 0.05) \times 10^{-3}$  W.u. (1970BH02)

<sup>d</sup>  $\Gamma_\gamma = 0.17 \pm 0.08, 24^{+27}_{-8}, 3.7^{+1.8}_{-0.9}$  and  $2.0^{+1.3}_{-0.6}$  meV: see Table 19.26 in (1978AJ03).

Table 19.25: Resonances reported in  $^{16}\text{O} + ^3\text{He}$  <sup>a</sup>

$E(^3\text{He})$ (MeV)	Resonance in	$\Gamma_{\text{c.m.}}$ (MeV)	$E_x$ (MeV)	$J^\pi$
2.400	$p_{1 \rightarrow 4}, p_{5,6,7}, \alpha_0$	0.355	10.46	$\frac{1}{2}^+$
2.425	$p_{1 \rightarrow 4}, p_{5,6,7}, \alpha_0$	0.045	10.48	$\frac{3}{2}^+$
3.65	$p\gamma, ^3\text{He}, \alpha_0$	0.025	$11.51 \pm 0.05$	$\frac{3}{2}^-, (\frac{1}{2}^-)$
4.50	$^3\text{He}, \alpha_0$	$0.200 \pm 0.025$	$12.23 \pm 0.05$	$\frac{5}{2}^+$
4.70	$^3\text{He}, \alpha_0$	$0.180 \pm 0.025$	$12.40 \pm 0.05$	$\frac{7}{2}^+$
5.05	$p_0, p_1, p_5, ^3\text{He}$	$0.18 \pm 0.04$	$12.69 \pm 0.05$	$\frac{1}{2}^+$
6.37 <sup>b</sup>	$\gamma_0, \gamma_{1+2}$	$0.67 \pm 0.25$	$13.8 \pm 0.25$	
7.65 <sup>b</sup>	$\gamma_{1+2}$	$0.62 \pm 0.13$	$14.88 \pm 0.13$	
9.26 <sup>b</sup>	$\gamma_{1+2}, \mathbf{n}$	$0.40 \pm 0.13$	$16.23 \pm 0.13$	
11.8 <sup>b</sup>	$\gamma_{0 \rightarrow 2}$	$4.4 \pm 0.5$	$18.4 \pm 0.5$	

<sup>a</sup> See reaction 2,  $^{19}\text{Ne}$ , in (1978AJ03) for references.

<sup>b</sup>  $(2J+1)\Gamma_{^3\text{He}}\Gamma_\gamma = 30 \pm 16, 89 \pm 40, 18 \pm 2$  and  $17000 \pm 4000 \text{ keV}^2$  for  $^{19}\text{Ne}^*(13.8, 14.9, 16.2, 18.4)$  (1981WA1R).

Cross sections for production of 0.94, 1.04 and 1.08 MeV  $\gamma$ -rays [reaction (c)] have been measured for  $E(^3\text{He}) = 2.6$  to 4.0 MeV (1980HE06). The earlier work on reactions (c), (e) and (f) is summarized in (1978AJ03): reported resonances are shown in Table 19.25.

Analyzing powers for reactions (d) [to  $^{17}\text{F}^*(0, 0.5)$ ] and (f) [to  $^{15}\text{O}^*(0, 6.18)$ ] have been measured at  $E(^3\text{He}) = 33$  MeV (1980LU02, 1980LU03, 1981KA1L). Polarization measurements in reaction (g) are reported by (1981LE1F; 41 MeV).

See also (1979KA1G, 1981RO1H) and  $^{18}\text{F}$ ,  $^{18}\text{Ne}$ ,  $^{16}\text{O}$  and  $^{17}\text{F}$  in (1982AJ01),  $^{15}\text{O}$  in (1981AJ01) and  $^{12}\text{C}$  in (1980AJ01).

$$6. \ ^{16}\text{O}(\alpha, \text{n})^{19}\text{Ne} \quad Q_{\text{m}} = -12.1344$$

Gamma transitions have been observed from the first six excited states of  $^{19}\text{Ne}$ : see Table 19.25 in (1978AJ03) and Table 19.24 here. At  $E_\alpha = 41$  MeV angular distributions are reported for the  $n_{0+1}$  group and for neutron groups to  $^{19}\text{Ne}^*(1.55, 2.78, 4.20, 4.63, 5.43, 6.2, 6.80, 7.61, 8.42, 8.95, 9.23, 9.88, 10.40, 11.09, 12.49)$ , many of which correspond to unresolved states. The relative spectroscopic factors, from a DW analysis, are 0.83, 1.62, 1.0, 2.92, 1.63 for  $^{19}\text{Ne}^*(0, 0.24, 1.54, 2.79, 4.62, 5.4)$  (1981OV01).

$$7. \ ^{16}\text{O}(^6\text{Li}, \text{t})^{19}\text{Ne} \quad Q_{\text{m}} = -7.351$$

This reaction (and the mirror reaction  $^{16}\text{O}(^6\text{Li}, ^3\text{He})^{19}\text{F}$ ) have been studied at  $E(^6\text{Li}) = 24, 35$  and  $36$  MeV [see (1978AJ03)] and  $46$  MeV (1979MA26). Table 19.13 displays the analog states observed in the two reactions. In addition triton groups are reported to states with  $E_x = 6.08, 6.28, 6.85, 7.21, 8.08, 8.45, 8.94, 9.81, 10.01, 11.08, 11.24, 11.40, 12.56$  [all  $\pm 0.02$ ],  $13.1, 13.22, 14.18, 14.44, 14.78$  [remaining,  $\pm 0.03$ ] MeV (1979MA26).

$$\begin{aligned} 8. \text{ (a) } & ^{16}\text{O}(^{10}\text{B}, ^7\text{Li})^{19}\text{Ne} & Q_m &= -9.344 \\ & \text{(b) } & ^{16}\text{O}(^{11}\text{B}, ^8\text{Li})^{19}\text{Ne} & Q_m &= -18.767 \end{aligned}$$

Reaction (a) as well as the analog reaction [ $^{16}\text{O}(^{10}\text{B}, ^7\text{Be})^{19}\text{F}$ ] have been studied at  $E(^{10}\text{B}) = 100$  MeV. On the basis of similar yields and  $E_x$ , and in addition to the low-lying analogs, it is suggested that the following pairs of states are analogs in  $^{19}\text{F}$ -( $^{19}\text{Ne}$ ):  $8.98$  ( $8.94$ ),  $11.33$  ( $11.09$ ),  $12.79$  ( $12.48$ ),  $14.15$  ( $14.17$ ),  $14.99$  ( $14.61$ ) and  $15.54$  ( $15.40$ ) [ $\pm 100$  keV]; however, problems of energy resolution are evident (1976HA06).

$^{19}\text{Ne}^*(4.63)$  is relatively strongly populated at  $E(^{11}\text{B}) = 115$  MeV [reaction (b)].  $^{19}\text{Ne}^*(2.79, 5.4, 9.8, 12.27)$  are also excited (1979RA10, 1981GO11) [see latter for  $S_{\text{rel}}$ ].

$$9. \ ^{16}\text{O}(^{12}\text{C}, ^9\text{Be})^{19}\text{Ne} \quad Q = -11.4297$$

At  $E(^{12}\text{C}) = 115$  MeV  $^{19}\text{Ne}^*(4.63)$  is strongly populated.  $^{19}\text{Ne}^*(0.24, 1.54, 2.79, 9.8, 12.27)$  are also observed (1979GO17, 1979RA10, 1981GO11) [see the latter for  $S_{\text{rel}}$ ]. See also (1978CH15).

$$10. \ ^{17}\text{O}(^3\text{He}, n)^{19}\text{Ne} \quad Q_m = 4.299$$

Neutron- $\gamma$  coincidence measurements lead to the determination of excitation energies [ $E_x = 4032.9 \pm 2.4, 4140 \pm 4, 4197.1 \pm 2.4, 4379.1 \pm 2.2, 4549 \pm 4, 4605 \pm 5, 4635 \pm 4$  and  $(5097 \pm 10)$  keV],  $\tau_m$  and branching ratios (see Table 19.24). On the basis of these it is suggested that  $^{19}\text{Ne}^*(4.14, 4.20)$  are the analogs of  $^{19}\text{F}^*(4.03, 4.00)$  [ $J^\pi = \frac{9}{2}^-, \frac{7}{2}^-$ , respectively] and that  $^{19}\text{Ne}^*(4.55, 4.60)$  are the analogs of  $^{19}\text{F}^*(4.558, 4.555)$  [ $J^\pi = \frac{5}{2}^+, \frac{3}{2}^-$ , respectively]. There is no evidence for a reported state at  $E_x = 4.78$  MeV (1973DA31).

$$11. \ ^{19}\text{F}(p, n)^{19}\text{Ne} \quad Q_m = -4.0207$$

Excited states of  $^{19}\text{Ne}$  determined from  $\gamma$ -spectra are displayed in Table 19.25 of (1978AJ03). Branching ratio and  $\tau_m$  measurements are summarized in Table 19.24 here. Neutron measurements are shown in Table 19.24 of (1972AJ02). For the  $g$ -factor of  $^{19}\text{Ne}^*(0.24)$  see Table 19.23. See also (1979WI1N, 1980HU1J) and  $^{20}\text{Ne}$ .

$$12. \ ^{19}\text{F}(^3\text{He}, \text{t})^{19}\text{Ne} \quad Q_m = -3.2570$$

At  $E(^3\text{He}) = 26$  MeV angular distributions of the triton groups to  $^{19}\text{Ne}^*(0.24, 1.54, 2.79)$  have been obtained by (1970SC05); those to  $^{19}\text{Ne}^*(0, 0.24)$  appear to proceed primarily via  $L = 0$  and  $L = 2$ , respectively.

$$13. \ ^{19}\text{F}(^6\text{Li}, ^6\text{He})^{19}\text{Ne} \quad Q_m = -6.745$$

At  $E(^6\text{Li}) = 34$  MeV the transitions to  $^{19}\text{Ne}^*(0, 0.24, 1.54, 2.79, 4.368 \pm 0.010)$  have been studied (1974GA11).

$$14. \ ^{20}\text{Ne}(\gamma, \text{n})^{19}\text{Ne} \quad Q_m = -16.8653$$

See  $^{20}\text{Ne}$ .

$$15. \ ^{20}\text{Ne}(\text{p}, \text{d})^{19}\text{Ne} \quad Q_m = -14.6407$$

See (1972AJ02).

$$16. \ ^{20}\text{Ne}(\text{d}, \text{t})^{19}\text{Ne} \quad Q_m = -10.6080$$

See (1978AJ03).

$$17. \ ^{20}\text{Ne}(^3\text{He}, \alpha)^{19}\text{Ne} \quad Q_m = 3.7125$$

Table 19.26:  $^{19}\text{Ne}$  levels from  $^{20}\text{Ne}(^3\text{He}, \alpha)$  <sup>a</sup>

$E_x$ (MeV $\pm$ keV)	$l_n$	$J^\pi$	$C^2S$	$E_x$ (MeV $\pm$ keV)
0	0	$\frac{1}{2}^+$	0.12	6.862 $\pm$ 7
0.23834 $\pm$ 0.15	2	$\frac{5}{2}^+$	1.04	7.067 $\pm$ 9
0.27530 $\pm$ 0.2	1	$\frac{1}{2}^-$	1.96	(7.178 $\pm$ 15)
1.5040 $\pm$ 3		$(\frac{5}{2}^-)$		7.253 $\pm$ 10
1.5324 $\pm$ 3		$(\frac{3}{2}^+)$	(0.73)	(7.326 $\pm$ 15)
1.6115 $\pm$ 3	1	$(\frac{3}{2})^-$	0.21	(7.531 $\pm$ 15)
2.7917 $\pm$ 3	4, 5	$(\frac{9}{2}^+)$		7.614 $\pm$ 20
4.036 $\pm$ 10	2	$(\frac{3}{2}, \frac{5}{2})^+$		7.700 $\pm$ 10
4.142 $\pm$ 10				(7.788 $\pm$ 10)
4.200 $\pm$ 10				7.994 $\pm$ 15
4.379 $\pm$ 10				8.063 $\pm$ 15
4.551 $\pm$ 10	1	$(\frac{1}{2}, \frac{3}{2})^-$		8.236 $\pm$ 10 <sup>b</sup>
4.625 $\pm$ 10				8.440 $\pm$ 10
4.712 $\pm$ 10				8.523 $\pm$ 10
4.783 $\pm$ 20				(8.810 $\pm$ 25)
5.089 $\pm$ 7				8.915 $\pm$ 10
5.351 $\pm$ 10	0	$\frac{1}{2}^+$	0.01	9.013 $\pm$ 10
5.424 $\pm$ 7				9.100 $\pm$ 20
5.463 $\pm$ 20				9.240 $\pm$ 20
5.539 $\pm$ 9				9.489 $\pm$ 25
5.832 $\pm$ 9				9.886 $\pm$ 50 <sup>b</sup>
6.013 $\pm$ 7	1	$(\frac{3}{2}, \frac{1}{2})^-$	(3.62)	10.407 $\pm$ 30 <sup>b</sup>
6.094 $\pm$ 8				10.613 $\pm$ 20
6.149 $\pm$ 20				
6.289 $\pm$ 7				
6.437 $\pm$ 9				
6.742 $\pm$ 7	1	$(\frac{3}{2}, \frac{1}{2})^-$		

<sup>a</sup> See Table 19.27 of (1978AJ03) for additional results and for a listing of the references.

<sup>b</sup> Unresolved states.

Alpha groups have been observed to  $^{19}\text{Ne}$  states with  $E_x < 10.6$  MeV: see Tables 19.22 and 19.26. Angular distributions have been measured for  $E(^3\text{He}) = 10$  to 35 MeV: see (1972AJ02). DWBA analysis of the strongest transitions leads to the  $l$  and  $J^\pi$  values shown in Table 19.26. Relative spectroscopic factors were also extracted.  $^{19}\text{Ne}^*(0, 0.24, 1.54, 2.79)$  are identified as members of the  $K = \frac{1}{2}^+$  rotational band [with  $^{19}\text{Ne}^*(4.38)$  as the  $\frac{7}{2}^+$  member: see, however, Table 19.23] and  $^{19}\text{Ne}^*(0.28, 1.51, 1.62)$  with the  $K = \frac{1}{2}^-$  band. Candidates for the  $\frac{7}{2}^-$  and  $\frac{9}{2}^-$  members of the  $K = \frac{1}{2}^-$  band are thought to be  $^{19}\text{Ne}^*(4.15, 4.20)$ . Possible matching of other  $^{19}\text{Ne}$  states with those in  $^{19}\text{F}$  is also discussed (1970GA18). For lifetime and radiative measurements see Table 19.24.

18.  $^{21}\text{Ne}(p, t)^{19}\text{Ne}$   $Q_m = -15.145$

At  $E_p = 40$  MeV the angular distributions to  $^{19}\text{Ne}^*(0.24, 4.03, 5.09)$  are well described by  $L = 2, 0$  and  $4$ , respectively.  $^{19}\text{Ne}^*(4.03)$ ,  $J^\pi = \frac{3}{2}^+$ , has dominant 5p-2h configuration.  $^{19}\text{Ne}^*(5.09)$  has  $\pi = +$  and its  $J$  is consistent with the previous value of  $\frac{5}{2}$  (1978FO26, 1979FO06). At  $E_p = 45$  MeV the triton group to a state with  $E_x = 7.620 \pm 0.025$  has an angular distribution [ $L = 0$ ] which is similar to that for  $^{19}\text{F}^*(7.66)$ : both are thought to be analogs of the  $(J^\pi; T) = (\frac{3}{2}^+; \frac{3}{2})$  0.096 MeV first excited state of  $^{19}\text{O}$ . The ground state of  $^{19}\text{O}$  has  $J^\pi = \frac{5}{2}^+$ ;  $L$  for the analog state should be 2 (1969HA38).  $^{19}\text{Ne}^*(0, 2.79)$  are also populated: see (1978AJ03).

**$^{19}\text{Na}$**   
(Fig. 8)

A study of this nucleus via the  $^{24}\text{Mg}(^3\text{He}, ^8\text{Li})^{19}\text{Na}$  reaction at  $E(^3\text{He}) = 76.3$  MeV leads to an atomic mass excess of  $12.928 \pm 0.012$  MeV for  $^{19}\text{Na}$ ; it is then unstable with respect to breakup into  $^{18}\text{Ne} + p$  by  $320 \pm 13$  keV. An excited state at  $E_x = 120 \pm 10$  keV is also reported (1975BE38). See also (1978AJ03, 1978GU10, 1979BE1H).

**$^{19}\text{Mg}$**   
(Not illustrated)

$^{19}\text{Mg}$  has not been observed: see (1977CE05).



## References

(Closed 01 May 1982)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1961NO05 E. Norbeck, Phys. Rev. 121 (1961) 824
- 1964BA16 J.K. Bair, C.M. Jones and H.B. Willard, Nucl. Phys. 53 (1964) 209
- 1964DO08 T.R. Donoghue, A.F. Behof and S.E. Darden, Nucl. Phys. 54 (1964) 33
- 1965AL13 J.P. Allen, A.J. Howard, D.A. Bromley and J.W. Olness, Nucl. Phys. 68 (1965) 426
- 1965VA03 F.J. Vaughn, H.A. Grench, W.L. Imhof, J.H. Rowland and M. Walt, Nucl. Phys. 64 (1965) 336
- 1966OL01 J.W. Olness and D.H. Wilkinson, Phys. Rev. 141 (1966) 966
- 1967ST28 T. Stambach, S.E. Darden, P. Huber and I. Sick, Helv. Phys. Acta 40 (1967) 915
- 1967WA13 E.K. Warburton, J.W. Olness and A.R. Poletti, Phys. Rev. 160 (1967) 938
- 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, Nucl. Phys. A113 (1968) 395
- 1969HA38 J.C. Hardy, H. Brunnader, J. Cerny and J. Janecke, Phys. Rev. 183 (1969) 854
- 1969JA09 K.P. Jackson, K. Bharuth Ram, P.G. Lawson, N.G. Chapman and K.W. Allen, Phys. Lett. B30 (1969) 162
- 1970AD02 E.G. Adelberger and A.B. McDonald, Nucl. Phys. A145 (1970) 497
- 1970AJ04 F. Ajzenberg-Selove, Nucl. Phys. A152 (1970) 1
- 1970BH02 K. Bharuth-Ram, R.D. Gill, K.P. Jackson, B. Povh and E.K. Warburton, Phys. Rev. C2 (1970) 1210
- 1970DE06 D. Dehnhard and N.M. Hintz, Phys. Rev. C1 (1970) 460
- 1970GA18 J.D. Garrett, R. Middleton and H.T. Fortune, Phys. Rev. C2 (1970) 1243
- 1970SC05 J.J. Schwartz and B.A. Watson, Phys. Rev. Lett. 24 (1970) 322
- 1970SC25 C. Schmidt and H.H. Duhm, Nucl. Phys. A155 (1970) 644
- 1971FO18 H.T. Fortune and C.M. Vincent, Phys. Rev. C4 (1971) 1994
- 1971KN05 H. Knoth, P.H. Barker, A. Huber, U. Matter, P.M. Cockburn and P. Marmier, Nucl. Phys. A172 (1971) 25
- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1

- 1972BI14 H.G. Bingham and H.T. Fortune, Phys. Rev. C6 (1972) 1900
- 1972CH18 M. Chevallier, P. Gaillard, J.Y. Grossiord, M. Gusakow, J.R. Pizzi and C. Ruhla, J. Phys. (Paris) 33 (1972) 177
- 1972GA08 J.D. Garrett, H.G. Bingham, H.T. Fortune and R. Middleton, Phys. Rev. C5 (1972) 682
- 1972RO01 D.W.O. Rogers, J.G. Aitken and A.E. Litherland, Can. J. Phys. 50 (1972) 268
- 1972SC17 U.C. Schlotthauer-Voos, R. Bock, H.G. Bohlen, H.H. Gutbrod and W. von Oertzen, Nucl. Phys. A186 (1972) 225
- 1973BI02 H.G. Bingham, H.T. Fortune, J.D. Garrett and R. Middleton, Phys. Rev. C7 (1973) 60
- 1973DA31 J.M. Davidson and M.L. Roush, Nucl. Phys. A213 (1973) 332
- 1973DE06 R. de Swiniarski, A. Genoux-Lubain, G. Bagieu, J.F. Cavaignac, D.H. Worledge and J. Raynal, Phys. Lett. B43 (1973) 27
- 1973DO05 T.R. Donoghue, Phys. Rev. C7 (1973) 1270
- 1973GA14 A. Gamp, W. von Oertzen, H.G. Bohlen, M. Feil, R.L. Walter and N. Marquardt, Z. Phys. 261 (1973) 283
- 1973HA13 P.L. Hallowell, W. Bertozzi, J. Heisenberg, S. Kowalski, X. Maruyama, C.P. Sargent, W. Turchinets, C.F. Williamson, S.P. Fivozinsky, J.W. Lightbody, Jr. et al., Phys. Rev. C7 (1973) 1396
- 1973RO09 D.W.O. Rogers, W.R. Dixon and R.S. Storey, Can. J. Phys. 51 (1973) 1
- 1973WA10 E.K. Warburton, J.W. Olness, G.A.P. Engelbertink and T.K. Alexander, Phys. Rev. C7 (1973) 1120
- 1973WE11 M. Wery, Nucl. Phys. A210 (1973) 329
- 1974BO05 J.D. Bowman, A.M. Poskanzer, R.G. Korteling and G.W. Butler, Phys. Rev. C9 (1974) 836
- 1974CO13 W.J. Courtney, K.W. Kemper, G.E. Moore and R.L. White, Phys. Rev. C9 (1974) 1273
- 1974DE46 R. de Swiniarski, A. Genoux-Lubain, G. Bagieu and J.F. Cavaignac, Can. J. Phys. 52 (1974) 2422
- 1974GA11 C. Gaarde and T. Kammuri, Nucl. Phys. A221 (1974) 238
- 1974GA28 J.D. Garrett and O. Hansen, Nucl. Phys. A229 (1974) 204
- 1974SK04 D.M. Skopik and Y.M. Shin, Nucl. Phys. A223 (1974) 409
- 1974UN01 B.Y. Underwood, M.R. Wormald, N. Anyas-Weiss, N.A. Jelley and K.W. Allen, Nucl. Phys. A225 (1974) 253
- 1974VE06 A. Veyssiere, H. Beil, R. Bergere, P. Carlos, A. Lepretre and A. de Miniac, Nucl. Phys. A227 (1974) 513

- 1975AD01 E.G. Adelberger, H.E. Swanson, M.D. Cooper, J.W. Tape and T.A. Trainor, Phys. Rev. Lett. 34 (1975) 402
- 1975BE38 W. Benenson, A. Guichard, E. Kashy, D. Mueller, H. Nann and L.W. Robinson, Phys. Lett. B58 (1975) 46
- 1975FO07 H.T. Fortune, L.R. Medsker, W.S. Chien, H. Nann and B.H. Wildenthal, Phys. Rev. C12 (1975) 359
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, Phys. Rev. C11 (1975) 2049
- 1975LE16 C. Lebrun, F. Guilbault, P. Avignon and Y. Deschamps, J. Phys. Lett. (Paris) 36 (1975) L201; Erratum J. Phys. Lett. (Paris) 37 (1976) L57
- 1975OY01 M. Oyamada, T. Terasawa, K. Nakahara, Y. Endo, H. Saito and E. Tanaka, Phys. Rev. C11 (1975) 1578
- 1975TS03 H. Tsubota, N. Kawamura, S. Oikawa and J. Uegaki, J. Phys. Soc. Jpn. 38 (1975) 299
- 1975ZI1A Zimmerman, Fowler and Caughlan, OAP-399 (1975)
- 1976AL07 D.E. Alburger, Phys. Rev. C13 (1976) 2593
- 1976BH03 K. Bharuth-Ram, K.P. Jackson, P.G. Lawson, N.A. Jelley and K.W. Allen, Nucl. Phys. A269 (1976) 327
- 1976GO09 G. Goldring, B. Richter, Z. Shkedi and Y. Wolfson, Nucl. Phys. A262 (1976) 214
- 1976HA06 M. Hamm, C.W. Towsley, R. Hanus, K.G. Nair and K. Nagatani, Phys. Rev. Lett. 36 (1976) 846
- 1976MC1G McGrory, in 1974 Gamma Ray Trans. Prob. (U. 3F Delhi Press, 1976) 469
- 1976NA18 H. Nann and B.H. Wildenthal, Phys. Rev. Lett. 37 (1976) 1129
- 1976PO02 J.E. Poling, E. Norbeck and R.R. Carlson, Phys. Rev. C13 (1976) 648
- 1976RO07 D.W.O. Rogers, A.L. Carter, T.J.M. Symons, S.P. Dolan, N. Anyas-Weiss and K.W. Allen, Can. J. Phys. 54 (1976) 938
- 1976SP07 P. Sperr, T.H. Braid, Y. Eisen, D.G. Kovar, F.W. Prosser, Jr., J.P. Schiffer, S.L. Tabor and S. Vigdor, Phys. Rev. Lett. 37 (1976) 321
- 1976SY01 T.J.M. Symons, L.K. Fifield, M.J. Hurst, A. Pakkanen, F. Watt, C.H. Zimmerman and K.W. Allen, Phys. Lett. B63 (1976) 409
- 1977BE2P Berr et al., in Proc., Zurich-Sin (1977) 284
- 1977BI1D Bizzeti, Nucl. Instrum. Meth. Phys. Res. 146 (1977) 285
- 1977BU22 B. Buck, H. Friedrich and A.A. Pilt, Nucl. Phys. A290 (1977) 205
- 1977CE05 J. Cerny and J.C. Hardy, Ann. Rev. Nucl. Sci. 27 (1977) 333
- 1977CH22 R. Chechik, Z. Fraenkel, H. Stocker and Y. Eyal, Nucl. Phys. A287 (1977) 353
- 1977CI1D Ciric et al., Sb. Rad. Prir.-Mat. Fak. Ser. Fiz. Univ. 7 (1977) 61; Phys. Abs. 37359 (1978)

- 1977CL03 C.F. Clement, A.M. Lane and J. Kopecky, Phys. Lett. B71 (1977) 10
- 1977DI18 W.R. Dixon, T.J.M. Symons, A.A. Pilt, K.W. Allen, C.H. Zimmerman, F. Watt and S.P. Dolan, Phys. Lett. A62 (1977) 479
- 1977FE1C Fecfilov et al., Bull. Acad. Sci. USSR, Phys. Ser. 41 (1977) 153
- 1977FO10 H.T. Fortune and H.G. Bingham, Nucl. Phys. A293 (1977) 197
- 1977FO1E Fortune, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 99
- 1977GA1E Garvey, AIP Conf. Proc. No. 37 (Indiana Univ.), Weak Int. Phys., 1977 (Ed. Lichtenberg; AIP, New York 1977) 104
- 1977GR16 R. Gross and I. Talmi, Nucl. Phys. A286 (1977) 211
- 1977HE1L Henley, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 812
- 1977KA1Y Kalinsky, Melnik, Smilansky and Trautner, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 263
- 1977KA21 D. Kalinsky, D. Melnik, U. Smilansky, N. Trautner, Y. Horowitz and S. Mordechai, Nucl. Phys. A289 (1977) 205
- 1977KL09 W.E. Kleppinger, F.P. Calaprice and B.R. Holstein, Nucl. Phys. A293 (1977) 46
- 1977KO38 B. Kohlmeyer, W. Pfeffer and F. Puhlhofer, Nucl. Phys. A292 (1977) 288
- 1977KU06 H. Kumpf, J. Mosner, W. Neubert and G. Schmidt, Yad. Fiz. 25 (1977) 481; Sov. J. Nucl. Phys. 25 (1977) 259
- 1977MA2P Mankoc-Borstnik, Jang and Brut, Fiz. (Yugoslavia) 9, Suppl. 1 (1977) 11, 13; Phys. Abs. 12974, 13022 (1978)
- 1977MA35 V.B. Mandelzweig, Nucl. Phys. A292 (1977) 333
- 1977MO1A T. Motobayashi, Rept. Inst. Phys. Chem. Res. 53 (1977) 81.
- 1977NO07 J.M. Normand, Nucl. Phys. A291 (1977) 126
- 1977OK1A Oka and Kubodera, in Proc., Zurich-Sin. (1977) 329
- 1977PH02 D.-L. Pham and R. de Swiniarski, Nuovo Cim. A41 (1977) 543
- 1977SH18 J. Shurpin, H. Muther, T.T.S. Kuo and A. Faessler, Nucl. Phys. A293 (1977) 61
- 1977SI1D Silberberg and Tsao, Astrophys. J. Suppl. 35 (1977) 129
- 1977ST27 M.M. Sternheim and R.R. Silbar, Phys. Rev. C16 (1977) 2059
- 1977TE1B Telegdi, in High Energy Phys. Nucl. Struct., Zurich, 1977 (Birkhauser Verlag, 1977) 367
- 1977TR1D Truran, in CNO Isotopes in Astrophys., Ed. J. Audouze (D. Reidel, Holland, 1977) 145

- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1
- 1978AN07 I. Angeli and M. Csatlos, Atomki Kozlem. 20 (1978) 1
- 1978AN15 N. Anyas-Weiss and D. Strottman, Nucl. Phys. A306 (1978) 201
- 1978AS07 F. Asfour, I. Bondouk, M.H.S. Bakr, M.S. El-Tahawy and K. Omar, Rev. Roum. Phys. 23 (1978) 535
- 1978AT01 M. Atarashi, K. Hira and H. Narumi, Prog. Theor. Phys. 60 (1978) 209
- 1978BI1G J.R. Birkelund and J.R. Huizenga, Phys. Rev. C17 (1978) 126
- 1978BR21 F. Brut, N. Mankoc-Borstnik and S. Jang, Nucl. Phys. A304 (1978) 429
- 1978CA1H Calaprice, Hyperfine Int. 4 (1978) 25
- 1978CA1N Cassagnou, in Clustering, Winnipeg 1978, AIP Conf. Proc. 47 (1978) 434
- 1978CH15 Y.-D. Chan, H. Bohn, R. Vandenbosch, K.G. Bernhardt, J.G. Cramer, R. Sielemann and L. Green, Nucl. Phys. A303 (1978) 500
- 1978CH16 R. Chechik, Z. Fraenkel, Y. Eyal and H. Stocker, Nucl. Phys. A304 (1978) 243
- 1978CH26 W. Chung, J. van Hienen, B.H. Wildenthal and C.L. Bennett, Phys. Lett. B79 (1978) 381
- 1978CO18 V. Corcalciuc, B. Holmqvist, A. Marcinkowski and G.A. Prokopets, Nucl. Phys. A307 (1978) 445
- 1978DA1N Darema-Rogers and Millener, Bull. Amer. Phys. Soc. 23 (1978) 938
- 1978DE1K B. Desplanques and J. Missimer, Nucl. Phys. A300 (1978) 286
- 1978DI13 G.U. Din, Aust. J. Phys. 31 (1978) 267
- 1978DI1D Dietrich and Simpson, Astrophys. J. 225 (1978) L41
- 1978DR07 W. Dreves, P. Zupranski, P. Egelhof, D. Kassen, E. Steffens, W. Weiss and D. Fick, Phys. Lett. B78 (1978) 36
- 1978FO11 H.T. Fortune, L.R. Medsker, H. Nann and B.H. Wildenthal, Nucl. Phys. A301 (1978) 441
- 1978FO22 H.T. Fortune, J.N. Bishop, L.R. Medsker and B.H. Wildenthal, Phys. Rev. Lett. 41 (1978) 527
- 1978FO26 H.T. Fortune, H. Nann and B.H. Wildenthal, Phys. Rev. C18 (1978) 1563
- 1978GO1N Goldberg et al., Hyperfine Interactions 4 (1978) 262
- 1978GU10 K. Gul, J. Phys. Soc. Jpn. 44 (1978) 353
- 1978HE1D Heitzmann, Atomkernenergie 31 (1978) 262
- 1978HO13 D. Horn and A.J. Ferguson, Phys. Rev. Lett. 41 (1978) 1529
- 1978HO20 D. Horn, A.J. Ferguson and O. Hausser, Nucl. Phys. A311 (1978) 238
- 1978KL1D Kleppinger, Calaprice and Mueller, Bull. Amer. Phys. Soc. 23 (1978) 603

- 1978KO01 D.G. Kovar, W. Henning, B. Zeidman, Y. Eisen, J.R. Erskine, H.T. Fortune, T.R. Ophel, P. Sperr and S.E. Vigdor, *Phys. Rev. C* 17 (1978) 83
- 1978KR05 W. Kreishe, H. Niedrig, K. Reuter, K. Roth and K. Thomas, *Phys. Rev. C* 17 (1978) 2006
- 1978KR19 K.S. Krane, *At. Data Nucl. Data Tables* 22 (1978) 269
- 1978LE04 H.C. Lee, *Nucl. Phys.* A294 (1978) 473
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, *Table of Isotopes 7th Ed.* (John Wiley and Sons, Inc., New York, 1978)
- 1978MA2H Mavromatis, Muther and Faessler, *Z. Phys.* A284 (1978) 195
- 1978MA30 H.-B. Mak, H.C. Evans, G.T. Ewan and J.D. Macarthur, *Nucl. Phys.* A304 (1978) 210
- 1978MA54 N. Mankoc-Borstnik, *Fiz. (Yugoslavia) Suppl.* 10 (1978) 55
- 1978ME1D Meyer, *Nature* 272 (1978) 675
- 1978OB01 F.E. Obenshain, R.L. Ferguson, M.L. Halbert, D.C. Hensley, H. Nakahara, F. Plasil, F. Pleasonton, A.H. Snell and R.G. Stokstad, *Phys. Rev. C* 18 (1978) 764
- 1978OL04 A. Olin, G.A. Beer, D.A. Bryman, M.S. Dixit, J.A. Macdonald, G.R. Mason, R.M. Pearce and P.R. Poffenberger, *Nucl. Phys.* A312 (1978) 361
- 1978OR1A Orth, Buffington, Smoot and Mast, *Astrophys. J.* 226 (1978) 1147
- 1978PE09 T. Pedersen and E. Osnes, *Nucl. Phys.* A303 (1978) 345
- 1978PE14 D.M. Perlman, L. Grodzins and C.E. Thorn, *Phys. Rev. C* 18 (1978) 2333
- 1978PI06 A.A. Pilt and H.T. Fortune, *J. Phys.* G4 (1978) L77
- 1978PI1E Pilt, in *Clustering, Winnipeg 1978*, AIP Conf. Proc. 47 (1978) 173
- 1978RA2A S. Raman, C.A. Houser, T.A. Walkiewicz and I.S. Towner, *At. Data Nucl. Data Tables* 21 (1978) 567; Erratum *At. Data Nucl. Data Tables* 22 (1978) 369
- 1978RO1D Rolfs and Trautvetter, *Ann. Rev. Nucl. & Part. Sci.* 28 (1978) 115
- 1978SC13 L.A. Schaller, T. Dubler, K. Kaeser, G.A. Rinker, Jr., B. Robert-Tissot, L. Schellenberg and H. Schneuwly, *Nucl. Phys.* A300 (1978) 225; Erratum *Nucl. Phys.* A312 (1978) 524
- 1978SC19 K.W. Schmid and G. Do Dang, *Phys. Rev. C* 18 (1978) 1003
- 1978SH18 T. Shibata, H. Ejiri, J. Chiba, S. Nagamiya and K. Nakai, *Nucl. Phys.* A308 (1978) 513
- 1978SH1U Shroy et al., *Nucl. Instrum. Meth. Phys. Res.* 149 (1978) 313
- 1978SI1D R.R. Silbar, J.N. Ginocchio and M.M. Sternheim, *Phys. Rev. C* 18 (1978) 2785
- 1978SY01 T.J.M. Symons, L.K. Fifield, M.J. Hurst, F. Watt, C.H. Zimmerman and K.W. Allen, *J. Phys.* G4 (1978) 411

- 1978TA1A Tang, Lemere and Thompson, Phys. Rept. 47 (1978) 167
- 1978TA1U R.E. Taam and R.E. Picklum, Astrophys. J. 224 (1978) 210
- 1978TH1A Thompson, in Clustering, Winnipeg 1978, AIP Conf. Proc. 47 (1978) 69
- 1978VA1A L.C. Vaz and J.M. Alexander, Phys. Rev. C18 (1978) 2152
- 1978VO1D Volkov, Phys. Rept. 44 (1978) 93
- 1978WE1J Weinberg, Proc. 19th Int. Conf. High Energy Phys., Tokyo (1978) 907
- 1978WI01 C.F. Williamson, F.N. Rad, S. Kowalski, J. Heisenberg, H. Crannell, J.T. O'Brien and H.C. Lee, Phys. Rev. Lett. 40 (1978) 1702
- 1978WO1E Woosley, Fowler, Holmes and Zimmerman, At. Data Nucl. Data Tables 22 (1978) 371
- 1978WU1C Wu, Overley, Barnes and Switkowski, LAP-164 (1978)
- 1978YO01 J. Yonnet, M. Louvel, C. Lebrun, G. Roche, G. Landaud, A. Devaux, J. Castor, A. Baldit and J.P. Alard, Phys. Rev. Lett. 40 (1978) 164
- 1978ZA1D Zalm, Van Hienen and Glaudemans, Z. Phys. A287 (1978) 255
- 1978ZE07 L. Zetta and L. Zuffi, Nuovo Cim. A47 (1978) 141
- 1979AL1Q Alonso, Chatterjee and Tobias, Bull. Amer. Phys. Soc. 24 (1979) 142
- 1979AL22 Y. Alhassid, R.D. Levine, J.S. Karp and S.G. Steadman, Phys. Rev. C20 (1979) 1789
- 1979AM03 H. Amakawa and K.-I. Kubo, Phys. Rev. C19 (1979) 2403
- 1979AN35 G.B. Andreev and A.P. Tomozov, Ukr. Fiz. Zh. (USSR) 24 (1979) 1888; Phys. Abs. 32288 (1980)
- 1979BA31 G.C. Ball, W.G. Davies, J.S. Forster, H.R. Andrews, D. Horn and W. McLatchie, Nucl. Phys. A325 (1979) 305
- 1979BE1H Benenson and Kashy, Rev. Mod. Phys. 51 (1979) 527
- 1979BE2L Benkoula et al., Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York, 1979) 371
- 1979BE31 W. Benenson, G. Bertsch, G.M. Crawley, E. Kashy, J.A. Nolen, Jr., H. Bowman, J.G. Ingersoll, J.O. Rasmussen, J. Sullivan, M. Koike et al., Phys. Rev. Lett. 43 (1979) 683; Erratum Phys. Rev. Lett. 44 (1980) 56
- 1979BI05 J.N. Bishop, L.R. Medsker, H.T. Fortune and B.H. Wildenthal, Phys. Rev. C20 (1979) 1221
- 1979BO1N Bobchenko et al., Yad. Fiz. 30 (1979) 1553
- 1979BR04 H.S. Bradlow, W.D.M. Rae, P.S. Fisher, N.S. Godwin, G. Proudfoot and D. Sinclair, Nucl. Phys. A314 (1979) 207
- 1979CA1K Calaprice, Bull. Amer. Phys. Soc. 24 (1979) 39
- 1979CH1T Chance and Harris, Astron. Astrophys. 74 (1979) 247

- 1979CH22 M.L. Chatterjee, H.C. Cheung and B. Cujec, Nucl. Phys. A323 (1979) 461
- 1979CO09 A. Cortes and A.P. Zuker, Phys. Lett. B84 (1979) 25
- 1979DA15 B.J. Dalton, W.J. Baldrige and J.P. Vary, Phys. Rev. C20 (1979) 1908
- 1979DE10 B. Desplanques, Nucl. Phys. A316 (1979) 244
- 1979DE18 B. Desplanques and J. Missimer, Phys. Lett. B84 (1979) 363
- 1979FO03 H.T. Fortune, J. Phys. G5 (1979) 381
- 1979FO06 H.T. Fortune, J.N. Bishop, H. Nann and B.H. Wildenthal, Phys. Rev. C19 (1979) 1147
- 1979FO1F Fou, Rasmussen, Swann and Van Patter, IEEE Trans. on Nucl. Sci. 26 (1979) 1378
- 1979FR05 A.D. Frawley, J.F. Mateja, A. Roy and N.R. Fletcher, Phys. Rev. C19 (1979) 2215
- 1979GA04 J.A. Gaidos, L.J. Gutay, A.S. Hirsch, R. Mitchell, T.V. Ragland, R.P. Scharenberg, F. Turkot, R.B. Willmann and C.L. Wilson, Phys. Rev. Lett. 42 (1979) 82
- 1979GA1H Gazis et al., Bull. Amer. Phys. Soc. 24 (1979) 695
- 1979GO11 J. Gomez del Campo, R.G. Stokstad, J.A. Biggerstaff, R.A. Dayras, A.H. Snell and P.H. Stelson, Phys. Rev. C19 (1979) 2170
- 1979GO13 M.B. Golin and S. Kubono, Phys. Rev. C20 (1979) 1347
- 1979GO17 N.S. Godwin, W.D. Rae, B. Cooke, N.J. Eyre, P.S. Fisher, G. Proudfoot and D. Sinclair, Nucl. Phys. A331 (1979) 237
- 1979GR11 J.M. Greben and F.S. Levin, Nucl. Phys. A325 (1979) 145
- 1979GR1E Greenwood, in Proc. 3rd Symp. on Neutron Capture Gamma Rays, 1978 (Plenum Press, New York 1979) 441
- 1979HA1M Harwood and Crawley, Bull. Amer. Phys. Soc. 24 (1979) 817
- 1979HE1F K.T. Hecht and W. Zahn, Nucl. Phys. A313 (1979) 77
- 1979JA22 L. Jarczyk, B. Kamys, J. Okolowicz, J. Sromicki, A. Strzalkowski, H. Witala, Z. Wrobel, M. Hugi, J. Lang, R. Muller et al., Nucl. Phys. A325 (1979) 510
- 1979KA1G Karban, Microscopic Optical Potentials. Hamburg, Germany (Springer-Verlag 1979) p. 264; Phys. Abs. 97712 (1979)
- 1979KE1B Kerkhove et al., Int. Conf. Nucl. Phys. with Electromagnetic Interactions, Mainz (1979) 4.16
- 1979KI1G Kim and Primakoff, in "Mesons in Nucl.", Eds. Rho and Wilkenson (North-Holland, Amsterdam, 1979) 69
- 1979KO20 D.G. Kovar, D.F. Geesaman, T.H. Braid, Y. Eisen, W. Henning, T.R. Ophel, M. Paul, K.E. Rehm, S.J. Sanders, P. Sperr et al., Phys. Rev. C20 (1979) 1305
- 1979KO26 L. Koester, K. Knopf and W. Waschowski, Z. Phys. A292 (1979) 95
- 1979LA10 R.D. Lawson, Phys. Rev. C20 (1979) 842



- 1979LA1H Lazareff, Audouze, Starrfield and Truran, *Astrophys. J.* 228 (1979) 875
- 1979LE18 M. LeMere and Y.C. Tang, *Phys. Rev. C*20 (1979) 2003
- 1979LO01 H. Lorenz-Wirzba, P. Schmalbrock, H.P. Trautvetter, M. Wiescher, C. Rolfs and W.S. Rodney, *Nucl. Phys.* A313 (1979) 346
- 1979MA26 Martz, Sanders, Parker and Dover, *Phys. Rev. C*20 (1979) 1340
- 1979MA27 N. Mankoc-Borstnik, F. Brut and S. Jang, *Nucl. Phys.* A325 (1979) 100
- 1979MO14 T. Motobayashi, I. Kohno, T. Ooi and S. Nakajima, *Nucl. Phys.* A331 (1979) 193
- 1979MO17 D.J. Morrissey, L.F. Oliveira, J.O. Rasmussen, G.T. Seaborg, Y. Yariv and Z. Fraenkel, *Phys. Rev. Lett.* 43 (1979) 1139
- 1979MU05 G. Murillo, M. Fernandez, P. Perez, J. Ramirez, S.E. Darden, M.C. Cobian-Rozak and L. Montestruque, *Nucl. Phys.* A318 (1979) 352
- 1979NA03 K. Nagatani and J.C. Peng, *Phys. Rev. C*19 (1979) 747
- 1979NA06 A. Nadasen, T.A. Carey, P.G. Roos, N.S. Chant, C.W. Wang and H.L. Chen, *Phys. Rev. C*19 (1979) 2099
- 1979NA1F Nagamiya, in BNL-51115 (1979) p. 131
- 1979OR01 M.E. Ortiz, A. Dacal, A. Menchaca-Rocha, M. Buenerd, D.L. Hendrie, J. Mahoney, C. Olmer and D.K. Scott, *Phys. Lett.* B84 (1979) 63
- 1979PA06 N. Paras, A.M. Bernstein, K.I. Blomqvist, G. Franklin, M. Pauli, B. Schoch, J. LeRose, K. Min, D. Rowley, P. Stoler et al., *Phys. Rev. Lett.* 42 (1979) 1455
- 1979PE16 T. Pedersen, E. Osnes and M. Guttormsen, *Nucl. Phys.* A332 (1979) 1
- 1979RA10 Rae et al., *Nucl. Phys.* A319 (1979) 239
- 1979RA1C Ramaty, Kozlovsky and Lingenfelter, *Astrophys. J. Suppl. Series* 40 (1979) 487
- 1979SA26 F. Saint-Laurent, M. Conjeaud, S. Harar, J.M. Loiseaux, J. Menet and J.B. Viano, *Nucl. Phys.* A327 (1979) 517
- 1979SA41 T. Sakuda and F. Nemoto, *Prog. Theor. Phys.* 62 (1979) 1274
- 1979SA43 T. Sakuda and F. Nemoto, *Prog. Theor. Phys.* 62 (1979) 1606
- 1979SC1K Schreiber et al., *Bull. Amer. Phys. Soc.* 24 (1979) 51
- 1979SI1K P.J. Siemens and J.I. Kapusta, *Phys. Rev. Lett.* 43 (1979) 1486
- 1979ST21 E.J. Stephenson, B.P. Hichwa and J.D. Hutton, *Nucl. Phys.* A331 (1979) 269
- 1979TH05 J.E.M. Thomson and M.N. Thompson, *Nucl. Phys.* A330 (1979) 66
- 1979VA08 K. van der Borg, R.J. de Meijer, A. van der Woude and H.T. Fortune, *Phys. Lett.* B84 (1979) 51
- 1979WI09 L.W.J. Wild and B.M. Spicer, *Aust. J. Phys.* 32 (1979) 187
- 1979WI1N Wilkerson, Clegg and Ludwig, *Bull. Amer. Phys. Soc.* 24 (1979) 107

- 1979ZE04 L. Zetta, L. Zuffi and P.L. Ottaviani, *Nuovo Cim.* A54 (1979) 421
- 1980AF1A Afzal, Ansari and Sergal, *Lett. Nuovo Cim.* 29 (1980) 350
- 1980AJ01 F. Ajzenberg-Selove and C.L. Busch, *Nucl. Phys.* A336 (1980) 1
- 1980AL1F Alburger, *Transactions of the N.Y. Acad. of Sci.* 40 (1980) 1
- 1980AN02 A. Anttila, S. Brandenburg, J. Keinonen and M. Bister, *Nucl. Phys.* A334 (1980) 205
- 1980BO21 K. Bodek, M. Hugi, L. Jarczyk, B. Kamys, J. Lang, R. Muller, M. Porebska, J. Sromicki, A. Strzalkowski, E. Ungricht et al., *J. Phys.* G6 (1980) 1017
- 1980BR09 B.A. Brown, W. Chung and B.H. Wildenthal, *Phys. Rev.* C21 (1980) 2600
- 1980BR13 B.A. Brown, W. Chung and B.H. Wildenthal, *Phys. Rev.* C22 (1980) 774
- 1980BR21 B.A. Brown, W.A. Richter and N.S. Godwin, *Phys. Rev. Lett.* 45 (1980) 1681
- 1980CO08 A.J. Cole, N. Longequeue, J. Menet, J.J. Lucas, R. Ost and J.B. Viano, *Nucl. Phys.* A341 (1980) 284
- 1980CO1R Cook, Stone and Vogl, *Astrophys. J.* 238 (1980) L97
- 1980CO1U Corcalciuc, *Stud. Cercet. Fiz.* 32 (1980) 1135, 1166
- 1980CU03 A. Cunsolo, A. Foti, G. Imme, G. Pappalardo, G. Raciti, F. Rizzo and N. Saunier, *Phys. Rev.* C21 (1980) 952
- 1980DE1F B. Desplanques, *Nucl. Phys.* A335 (1980) 147
- 1980DE1U DeConninck, *Bull. Amer. Phys. Soc.* 25 (1980) 757
- 1980DI03 D. Dieumegard, B. Maurel and G. Amsel, *Nucl. Instrum. Meth. Phys. Res.* 168 (1980) 93
- 1980DI12 W.R. Dixon, R.S. Storey and B.A. Logan, *Phys. Rev.* C22 (1980) 1279
- 1980FA06 E. Fabrici, S. Micheletti, M. Pignanelli, F.G. Resmini, R. De Leo, G. D'Erasmus, A. Pantaleo, J.L. Escudie and A. Tarrats, *Phys. Rev.* C21 (1980) 830
- 1980FA07 E. Fabrici, S. Micheletti, M. Pignanelli, F.G. Resmini, R. De Leo, G. D'Erasmus and A. Pantaleo, *Phys. Rev.* C21 (1980) 844
- 1980FU1G Furutani et al., *Suppl. Prog. Theor. Phys.* 68 (1980) 193
- 1980GO1D Goret, *Bull. Amer. Phys. Soc.* 25 (1980) 563
- 1980GR10 R.E.L. Green and R.G. Korteling, *Phys. Rev.* C22 (1980) 1594
- 1980HA41 W.C. Haxton, B.F. Gibson and E.M. Henley, *Phys. Rev. Lett.* 45 (1980) 1677
- 1980HE06 J.C.P. Heggie, Z.E. Switkowski and G.J. Clark, *Nucl. Instrum. Meth. Phys. Res.* 168 (1980) 125
- 1980HU1E Hugi et al., *Proc. Int. Conf. on Nucl. Phys. (Berkeley)* (1980) 508
- 1980HU1J Hugg and Hanna, *Polariz., Santa Fe* (1980) 294
- 1980KA1T Kato et al., *Proc. Int. Conf. Nucl. Phys., Berkeley* (1980) 462

- 1980KO02 J.J. Kolata, R.M. Freeman, F. Hass, B. Heusch and A. Gallmann, Phys. Rev. C21 (1980) 579
- 1980KO1U Kolb and Brenn, Hyperfine Inter. 7 (1980) 471, 480
- 1980KO1V Koptev, Maev, Makarov and Khanzadeev, Yad. Fiz. 31 (1980) 1501
- 1980KR1C Kraner, Bull. Amer. Phys. Soc. 25 (1980) 758
- 1980KU04 S. Kubono, D.A. Lewis and D. Dehnhard, Nucl. Phys. A334 (1980) 336
- 1980KU05 T.T.S. Kuo and E. Osnes, J. Phys. G6 (1980) 335
- 1980LE11 S.M. Lee, T. Matsuse and A. Arima, Phys. Rev. Lett. 45 (1980) 165
- 1980LI1J Lieb et al., Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 715
- 1980LO02 M. Lozano and G. Madurga, Phys. Lett. B90 (1980) 50
- 1980LU02 Y.-W. Lui, O. Karban, S. Roman, R.K. Bhowmik, J.M. Nelson and E. C. Pollacco, Nucl. Phys. A333 (1980) 205
- 1980LU03 Y.-W. Lui, O. Karban, S. Roman, R.K. Bhowmik, J.M. Nelson and E.C. Pollacco, Nucl. Phys. A333 (1980) 221
- 1980MA1G MacArthur et al., Bull. Amer. Phys. Soc. 25 (1980) 485
- 1980MC1H McKlveen, Bull. Amer. Phys. Soc. 25 (1980) 771
- 1980MC1L McGrory and Wildenthal, Ann. Rev. Nucl. Part. Sci. 30 (1980) 383
- 1980MI1G D.J. Millener, Phys. Rev. C22 (1980) 1355
- 1980NA12 H. Nann, K.K. Seth, S.G. Iversen, M.O. Kaletka, D.B. Barlow and D. Smith, Phys. Lett. B96 (1980) 261
- 1980OH05 M. Ohta and S. Okai, Prog. Theor. Phys. 63 (1980) 881
- 1980OK01 M. Oka and K. Kubodera, Phys. Lett. B90 (1980) 45
- 1980RO13 H.A. Roth, J.E. Christiansson and J. Dubois, Nucl. Phys. A343 (1980) 148
- 1980SA1L Sato et al., Proc. Int. Conf. on Nucl. Phys. (Berkeley) (1980) 536
- 1980SH1H Shen and Zhang, Phys. Energ. Fortis Phys. Nucl. 4 (1980) 43; Phys. Abs. 47420 (1980)
- 1980ST25 K. Stricker, J.A. Carr and H. McManus, Phys. Rev. C22 (1980) 2043
- 1980TA1L Tadic, Rept. Prog. Phys. 43 (1980) 67
- 1980WI17 M. Wiescher, H.W. Becker, J. Gorres, K.-U. Kettner, H.P. Trautvetter, W.E. Kieser, C. Rolfs, R.E. Azuma, K.P. Jackson and J.W. Hammer, Nucl. Phys. A349 (1980) 165
- 1981AD05 E.G. Adelberger, M.M. Hindi, C.D. Hoyle, H.E. Swanson and R.D. von Lintig, Phys. Rev. C24 (1981) 313
- 1981AD1E Adelberger, Santa Fe 1980, AIP Conf. Proc. 69 (1981) 1367
- 1981AJ01 F. Ajzenberg-Selove, Nucl. Phys. A360 (1981) 1.

1981AO01 K. Aoki and H. Horiuchi, *Prog. Theor. Phys.* 66 (1981) 1508  
 1981AR1D A. Arima, *Nucl. Phys.* A354 (1981) 19  
 1981BE63 V.B. Belyaev and O.P. Solovtsova, *Yad. Fiz.* 33 (1981) 699; *Sov. J. Nucl. Phys.* 33 (1981) 363  
 1981CA1H Cahn and Glashow, *Science* 213 (1981) 607  
 1981CH32 M.S. Chiou, M.W. Wu, N. Easwar and J.V. Maher, *Phys. Rev.* C24 (1981) 2507  
 1981CI03 M.A. Cirit and F. Yazici, *Phys. Rev.* C23 (1981) 2627  
 1981DE15 L.C. Dennis, A. Roy, A.D. Frawley and K.W. Kemper, *Nucl. Phys.* A359 (1981) 455  
 1981DE1H Dennis, Abdo, Frawley and Kemper, *Bull. Amer. Phys. Soc.* 26 (1981) 554  
 1981DE2K Dennis, Abdo, Frawley and Kemper, *Bull. Amer. Phys. Soc.* 26 (1981) 1236  
 1981DU1D Dubovik and Tosunyan, *Proc. Versailles Conf.* (1981) p. 471  
 1981ER03 M. Erculisse, *J. Phys.* G7 (1981) 951  
 1981FR17 E. Friedman, *Phys. Lett.* B104 (1981) 357  
 1981GL1B Glashausser et al., *Proc. Versailles Conf.* (1981) p. 156  
 1981GO11 N.S. Godwin, W.D.M. Rae, B. Cooke, A. Etchegoyen, N.J. Eyre, P.S. Fisher, G. Proudfoot and D. Sinclair, *Nucl. Phys.* A363 (1981) 493  
 1981GR06 S.M. Grimes and S.D. Bloom, *Phys. Rev.* C23 (1981) 1259  
 1981GR08 K. Grotowski, P. Belery, Th. Delbar, Y. El Masri, Gh. Gregoire, R. Janssens, J. Vervier, G. Paic, M. Albinska, J. Albinski et al., *Phys. Rev.* C23 (1981) 2513  
 1981HA1Q Haxton, *Proc. Versailles Conf.* (1981) p. 481  
 1981JA09 L. Jarczyk, B. Kamys, A. Magiera, J. Sromicki, A. Strzalkowski, G. Willim, Z. Wrobel, D. Balzer, K. Bodek, M. Hugi et al., *Nucl. Phys.* A369 (1981) 191  
 1981KA1L Karban, Basak, Morrison, Nelson and Roman, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 703  
 1981KE1E Kenny, *Australian J. Phys.* 34 (1981) 35  
 1981LE04 S.M. Lee, Y. Higashi, Y. Nagashima, S. Hanashima, M. Sato, H. Yamaguchi, M. Yamanouchi and T. Matsuse, *Phys. Lett.* B98 (1981) 418  
 1981LE1F Lezoch, Trost, Rahman and Strohmusch, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 745  
 1981LI1Z Lieb et al., *Bull. Amer. Phys. Soc.* 26 (1981) 1125  
 1981MA1E Mairle et al., Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 685  
 1981MA1Q Maguire et al., *Bull. Amer. Phys. Soc.* 26 (1981) 1221  
 1981ME1F Meyer and Elbaz, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 1083  
 1981MU1E Mukhopadhyay and Hintermann, Santa Fe 1980, *AIP Conf. Proc.* 69 (1981) 1068

- 1981MUZQ S.F. Mughabghab, M. Divadeenam and N.E. Holden, Neutron Cross Sections Part A, Z=1-60 (Academic Press, New York, 1981)
- 1981NA1E S. Nagamiya, M.-C. Lemaire, E. Moeller, S. Schnetzer, G. Shapiro, H. Steiner and I. Tanihata, Phys. Rev. C24 (1981) 971
- 1981NI03 T. Nishi, I. Fujiwara, N. Imanishi, H. Moriyama, K. Otozai, R. Arakawa, S. Saito, T. Tsuneyoshi, N. Takahashi, S. Iwata et al., Nucl. Phys. A352 (1981) 461
- 1981OL1E Olness, Warburton and Alburger, Unknown Source (1981)
- 1981OV01 D.J. Overway and W.C. Parkinson, Nucl. Phys. A363 (1981) 93
- 1981RO1H Roman, Santa Fe 1980, AIP Conf. Proc.69 (1981) 282
- 1981WA1R Waltham et al., Private Communication
- 1981XE01 A.C. Xenoulis, A.E. Aravantinos, C.J. Lister, J.W. Olness and R.L. Kozub, Phys. Lett. B106 (1981) 461
- 1982AB02 K.M. Abdo, L.C. Dennis, A.D. Frawley and K.W. Kemper, Nucl. Phys. A377 (1982) 281
- 1982AJ01 F. Ajzenberg-Selove, Nucl. Phys. A375 (1982) 1
- 1982BI1H Birbrair and Kalachnikov, in Kiev (1982) 172, 173
- 1982FI1C E. Fiorini, Nucl. Phys. A374 (1982) 577
- 1982KI02 M.M. King Yen, S.T. Hsieh, H.C. Chiang and D.S. Chuu, J. Phys. G8 (1982) 245
- 1982LOZZ M.M. Lowry, R.T. Kouzes, F.P. Calaprice and C.L. Bennett, Bull. Amer. Phys. Soc. 27 (1982) 492, DXa4
- 1982MAZY D.W. Macarthur, F.P. Calaprice, A.L. Hallin, M.B. Schneider and D.F. Schreiber, Bull. Amer. Phys. Soc. 27 (1982) 493, DXa11
- 1982NA08 F. Naulin, C. Detraz, M. Roy-Stephan, M. Bernas, J. de Boer, D. Guillemaud, M. Langevin, F. Pougheon and P. Roussel, J. Phys. Lett. 43 (1982) L29
- 1982SC1C Schneider et al., Bull. Amer. Phys. Soc. 27 (1982) 492
- 1982SCZZ D. Schreiber, F.P. Calaprice, A. Hallin, D. MaCarthur and M. Schneider, Bull. Amer. Phys. Soc. 27 (1982) 492, DXa6
- 1982SH1E Sharaf, Lett. Nuovo Cim. 33 (1982) 299