

Energy Levels of Light Nuclei $A = 20$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 18-20$ was published in *Nuclear Physics A300* (1978), p. 1. This version of $A = 20$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed November 1, 1977)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-AC02-76-ER02785]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 20$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below.

A. Nuclides: [\$^{20}\text{B}\$](#) , [\$^{20}\text{C}\$](#) , [\$^{20}\text{N}\$](#) , [\$^{20}\text{O}\$](#) , [\$^{20}\text{F}\$](#) , [\$^{20}\text{Ne}\$](#) , [\$^{20}\text{Na}\$](#) , [\$^{20}\text{Mg}\$](#) , [\$^{20}\text{Al}\$](#)

B. Tables of Recommended Level Energies:

[Table 20.1](#): Energy levels of ^{20}O

[Table 20.4](#): Energy levels of ^{20}F

[Table 20.18](#): Energy levels of ^{20}Ne

[Table 20.39](#): Energy levels of ^{20}Na

C. [References](#)

D. Figures: [\$^{20}\text{O}\$](#) , [\$^{20}\text{F}\$](#) , [\$^{20}\text{Ne}\$](#) , [\$^{20}\text{Na}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

²⁰B
(Not illustrated)

²⁰B has not been observed. The mass excess is predicted to be 69.08 MeV (1974TH01). ²⁰B is then unstable with respect to breakup into ¹⁹B + n by 0.9 MeV [see ¹⁹B]. See also (1976JA23, 1976WA18) and (1975BE31; theor.).

²⁰C
(Not illustrated)

²⁰C has not been observed in the 4.8 GeV proton irradiation of a uranium target (1974BO05). The mass excess is predicted to be 37.17 MeV (1974TH01), 37.41 MeV (1976JA23, 1976WA18). Assuming the mass excess of ²⁰C to be 37.3 MeV, ²⁰C is then stable with respect to ¹⁹C + n and ¹⁸C + 2n by 3.2 and 4.2 MeV, respectively [see ¹⁸C and ¹⁹C]. See also (1972TH13) and (1972ST1C, 1975BE31, 1976BE1G; theor.).

²⁰N
(Not illustrated)

²⁰N has been observed. It is particle stable: see (1972AJ02). Recent calculations of the atomic mass excess of ²⁰N are 21.67 MeV (1974TH01), 21.60 (1975JE02; transverse form of IMME), 21.88 (1976JA23) and 22.2 MeV (1977WA08). Assuming that the atomic mass excess is 22.0 MeV, ²⁰N is then stable with respect to ¹⁹N + n by 1.9 MeV (see ¹⁹N). See also (1972TH13, 1973TO16, 1975VO09, 1976WA18, 1977AR06, 1977BH1B) and (1973WI15, 1975BE31; theor.).

²⁰O
(Figs. 9 and 13)

GENERAL: (See also (1972AJ02).)

Shell model: (1972LE13, 1973JU1A, 1973LA1D, 1973MA1K, 1973MC06, 1974CO40, 1975BA81).

Cluster, collective and deformed models: (1973AB01).

Electromagnetic transitions: (1976VO1C).

Special states: (1972LE13, 1972SA04, 1973JU1A, 1973MC06, 1975BA81).

Complex reactions involving ²⁰O: (1973BA81, 1973VO1G, 1973WI15, 1975BA1Q, 1975VO09, 1976VA29, 1977AR06).

Other topics: (1972CA37, 1972SA04, 1972ST1C, 1973GR11, 1973SA24, 1973SP1A, 1974CO40, 1974HA17, 1974SE1B, 1975BA81, 1977DA10, 1977SH13).

Table 20.1: Energy levels of ^{20}O

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
0	$0^+; 2$	$\tau_{1/2} = 13.57 \pm 0.1$ sec	β^-	1, 2, 4, 5
1.67368 ± 0.15	2^+	$\tau_m = 9.8 \pm 0.7$ psec	γ	2
		$g = -0.39 \pm 0.04$		
3.568 ± 5	4^+		(γ)	2
4.065 ± 5	2^+		γ	2
4.446 ± 7	0^+		γ	2
4.838 ± 7			(γ)	2
4.997 ± 7			γ	2
5.220 ± 7			γ	2
5.298 ± 7			(γ)	2
5.382 ± 7	(0^+)		γ	2
5.603 ± 7			(γ)	2
(5.83 ± 20)			(γ)	2

Table 20.2: Branching in $^{20}\text{O}(\beta^-)^{20}\text{F}$ (1970MA42)

Decay to $^{20}\text{F}^*$ (MeV)	J^π	Branch (%)	$\log ft$
0.98	1^-	< 0.6	> 6.0
1.06	1^+	100.0	3.73
1.31	2^-	< 0.8	> 5.7
1.84	2^-	< 1.9	> 4.8
1.97	(3^-)	< 1.4	> 4.8
2.04	2^+	< 2.0	> 4.6
2.19	(3^+)	< 1.0	> 4.7

Table 20.3: Energy levels of ^{20}O from $^{18}\text{O}(t, p)^{20}\text{O}$

E_x (MeV \pm keV) (1962HI06)	L	J^π
0	0 ^{a,b}	0 ⁺
1.672 \pm 5 ^c	2 ^{a,b}	2 ⁺
3.568 \pm 5	4 ^a	4 ⁺
4.065 \pm 5 ^d	2 ^{a,b}	2 ⁺
4.446 \pm 7 ^d	0 ^a	0 ⁺
4.838 \pm 7		
4.997 \pm 7		
5.220 \pm 7 ^d		
5.298 \pm 7 ^d		
5.382 \pm 7 ^d		
5.603 \pm 7		
(5.83 \pm 20)		

^a (1964MI05): $E_t = 10.0$ MeV.

^b (1965MO19): $E_t = 5.55$ MeV.

^c E_γ measurements lead to $E_x = 1.67368 \pm 0.15$ (1973WA19),
1.6750 \pm 1.0 (1975BE15).

^d Preliminary results by K. Young (private communication) show that $^{20}\text{O}^*(4.07)$ decays to $^{20}\text{O}^*(0, 1.67)$ with branching ratios of 26 ± 4 and $74 \pm 4\%$: $\delta(E2/M1) = -0.18 \pm 0.10$; the work also favors 0^+ for $^{20}\text{O}^*(4.45, 5.38)$ (decay is to $^{20}\text{O}^*(1.67)$). $^{20}\text{O}^*(5.22, 5.30)$ appear to decay predominantly through $^{20}\text{O}^*(1.67)$ also.

Ground state of ^{20}O : (1973MC06, 1973SP1A, 1974CO40, 1974MC1F, 1974SHYR, 1975BE31, 1976BE1G).

1. $^{20}\text{O}(\beta^-)^{20}\text{F}$ $Q_m = 3.816$

^{20}O decays to $^{20}\text{F}^*(1.06)$ [$J^\pi = 1^+$] with a half-life of 13.57 ± 0.1 sec (1970MA42: $\log ft = 3.73$). See also (1972EY01). Upper limits for the branching to other states of ^{20}F are shown in Table 20.2. See also (1972WI1C, 1973LA03, 1973WI11, 1974WI1L, 1975NA20, 1975NA21, 1977AZ02; theor.).

2. $^{18}\text{O}(t, p)^{20}\text{O}$ $Q_m = 3.079$

Observed proton groups are displayed in Table 20.3 (1962HI06, 1964MI05, 1965MO19). The excitation energy of $^{20}\text{O}^*(1.67)$ is 1673.68 ± 0.15 keV (1973WA19), 1675.0 ± 1.0 keV (1975BE15); $J = 2$ (1970NI03); $\tau_m = 14.2 \pm 0.8$ psec (1975BE15), 9.8 ± 0.7 psec (1977HE12, 1977HE1D); $g = -0.39 \pm 0.04$ (1975BE15, 1976GE01). See also (1972AJ02).

3. $^{18}\text{O}(^{14}\text{C}, ^{12}\text{C})^{20}\text{O}$ $Q_m = -1.562$

See (1972EY01).

4. $^{18}\text{O}(^{16}\text{O}, ^{14}\text{O})^{20}\text{O}$ $Q_m = -17.327$

See (1976HI10).

5. $^{18}\text{O}(^{18}\text{O}, ^{16}\text{O})^{20}\text{O}$ $Q_m = -0.628$

At $E(^{18}\text{O}) = 52$ MeV angular distributions have been measured to four states of ^{20}O (1976KU1F; abstract). See also (1972EY01).

²⁰F
(Figs. 10 and 13)

GENERAL: (See also (1972AJ02).)

Shell model: (1972LE13, 1972WI13, 1973LA1D, 1973MA1K, 1973MC06, 1974CO39, 1975BA81).

Electromagnetic transitions: (1970HE1B, 1974MC1F).

Special states: (1972LE13, 1973MC06, 1975BA81, 1975MI03).

Complex reactions involving ²⁰F: (1972MI11, 1973BA81, 1973WI15, 1974HA61, 1975BA1Q, 1976HI05, 1977AR06).

Muon and pion capture and reactions: (1974LI1D).

Other topics: (1972CA37, 1972NA11, 1972WI13, 1973GR11, 1974CO39, 1974HA17, 1974MA1E, 1975BA81, 1977DA10, 1977SH13).

Ground state of ²⁰F: (1971SH26, 1972AC03, 1973AC1A, 1973MC06, 1973SU1B, 1974CO39, 1974MC1F, 1974MI21, 1974SHYR, 1976CH1T).

$$\mu = 2.094 (2) \text{ nm ((1976FU06) and V. Shirley, private communication);}$$

$$Q = 0.064 \pm 0.012 \text{ b (1974ST10).}$$

1. $^{20}\text{F}(\beta^-)^{20}\text{Ne}$ $Q_m = 7.0259$

Recent values of the half-life of ²⁰F are 11.03 ± 0.06 (1970WI05), 10.996 ± 0.020 (1975AL27) and 11.18 ± 0.01 sec (1976GE08). We adopt $\tau_{1/2} = 11.00 \pm 0.02$ sec. See also (1970AS1C, 1974AL11, 1975SA1D) and Table 20.5 in (1972AJ02) for the earlier values. ²⁰F decays principally to ²⁰Ne*(1.63): see ²⁰Ne and Table 20.36. See also (1976CA1E) and (1971WI1C, 1972BE1E, 1972EM02, 1972WI1C, 1973BE35, 1973LA03, 1973WI11, 1974WI1L, 1975NA20, 1975NA21, 1975WI1E, 1977AZ02, 1977CA11; theor.).

2. $^{12}\text{C}(^9\text{Be}, p)^{20}\text{F}$ $Q_m = 4.0761$

See (1974HA25).

3. $^{14}\text{N}(^7\text{Li}, p)^{20}\text{F}$ $Q_m = 12.054$

Table 20.7 shows the ^{20}F states observed in this reaction at $E(^7\text{Li}) = 16$ MeV. The cross sections for forming states of known J^π are linear with $2J_f + 1$ with slopes which are different for the even- and the odd-parity states. Extrapolation of these relationships to states of unknown J^π leads to the assignments shown in Table 20.7: $^{20}\text{F}^*(2.86, 2.97, 3.59, 3.68, 4.52, 4.76)$ and one or both of the unresolved states at 4.2, 4.6 and 4.9 MeV, must, on the basis of the σ_t for their population, have large values of J (1975BI04). See also (1973FO1A, 1975HO1L).

Table 20.4: Energy levels of ^{20}F ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
0	$2^+; 1$	$\tau_{1/2} = 11.0 \pm 0.02$ sec	β^-	1, 2, 3, 13, 16, 17, 25, 28, 31, 34, 37, 38, 39
0.65594 ± 0.15	3^+	$\tau_m = 0.39 \pm 0.03$ psec	γ	3, 13, 17, 25, 28, 34, 38
0.82288 ± 0.20	4^+	79 ± 6 psec	γ	3, 12, 13, 17, 25, 34, 38
0.98371 ± 0.20	1^-	2.03 ± 0.20 psec	γ	3, 12, 13, 17, 25, 34, 38
1.05693 ± 0.20	1^+	45 ± 13 fsec	γ	3, 13, 17, 25, 30, 34, 38
1.30923 ± 0.20	2^-	1.16 ± 0.20 psec	γ	3, 12, 13, 17, 25, 34, 38
1.8244 ± 1.2	5^+	≤ 65 fsec	γ	3, 12, 13, 25, 38
1.84337 ± 0.30	2^-	30 ± 20 fsec	γ	3, 13, 17, 25, 34, 38
1.9707 ± 0.4	(3^-)		γ	3, 12, 13, 17, 38
2.04400 ± 0.30	2^+	37 ± 16 fsec	γ	3, 13, 17, 25, 28, 34, 38
2.1948 ± 0.4	(3^+)	< 12 fsec	γ	3, 13, 17, 25, 34, 38
2.8649 ± 1.5	$(2, 3, 4)$		γ	3, 13, 25, 38
2.9661 ± 0.4	3^+	60 ± 40 fsec	γ	3, 13, 17, 25, 38
3.1740 ± 1.5	(1^+)		γ	13, 25, 38
3.48843 ± 0.25	1^+	44 ± 11 fsec	γ	13, 17, 25, 38
3.5260 ± 0.4	0^+	30 ± 15 fsec	γ	13, 17, 25
3.5871 ± 0.3	$(1, 2, 3)^+$	≤ 60 fsec	γ	3, 13, 17, 25, 38
3.6810 ± 0.4	$(1, 2, 3)^+$		γ	3, 13, 17, 25, 38
3.7611 ± 1.9			γ	13, 25, 38
3.9660 ± 1.5	1^+		γ	13, 17, 25, 38
4.0823 ± 0.4	$(1)^+$		γ	13, 17, 25, 38
4.1989 ± 2.7			(γ)	3, 25, 38
4.2077 ± 2.6			(γ)	3, 25, 38
4.2766 ± 0.5	$(1, 2, 3)^+$		γ	17, 25, 38
4.313 ± 3	$(0, 1)^+$		(γ)	25, 38
4.372 ± 4			(γ)	38
4.518 ± 4			(γ)	3, 38

Table 20.4: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
4.5838 \pm 3.0			(γ)	3, 25, 38
4.5922 \pm 2.9			(γ)	3, 25, 38
4.7302 \pm 2.9			(γ)	25, 38
4.7638 \pm 2.7			(γ)	3, 25, 38
4.8916 \pm 2.8			(γ)	3, 25, 38
4.8982 \pm 2.8			(γ)	3, 25, 38
5.0402 \pm 3.1	(0, 1, 2) ⁻		(γ)	25, 38
5.0655 \pm 3.1			(γ)	25, 38
5.131 \pm 5			(γ)	38
5.2240 \pm 2.8	(0, 1, 2) ⁻		(γ)	25, 38
5.279 \pm 3	(0, 1, 2) ⁻		(γ)	25, 38
5.3171 \pm 2.7			(γ)	25, 38
5.3445 \pm 3.3			(γ)	25, 38
5.4131 \pm 0.6			γ	17, 38
5.4503 \pm 3.8			(γ)	25, 38
5.4554 \pm 3.2			(γ)	25, 38
5.4634 \pm 3.3			(γ)	25
5.556 \pm 4			γ	17, 38
5.574 \pm 6			(γ)	38
5.621 \pm 3			(γ)	25, 38
5.713 \pm 2			γ	17, 38
5.7640 \pm 2.5			(γ)	25, 38
5.8104 \pm 2.5			(γ)	25, 38
5.9361 \pm 0.3	(0, 1, 2) ⁻		γ	17, 25, 38
6.0174 \pm 0.3	(0, 1, 2) ⁻		γ	17, 25, 38
6.0446 \pm 0.4			γ	17, 25, 38
6.163 \pm 6			(γ)	38
6.205 \pm 6			(γ)	38
6.240 \pm 7			(γ)	38
6.300 \pm 5			(γ)	38
6.337 \pm 5			(γ)	38
6.370 \pm 6			(γ)	38
6.407 \pm 12			(γ)	38

Table 20.4: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
6.441 \pm 9			(γ)	38
6.480 \pm 5			(γ)	38
6.519 \pm 3	0 ⁺ ; 2		γ	13, 37
6.588 \pm 5			(γ)	38
6.6013 \pm 0.3	0 ⁺ , 1 ⁺		γ	17
6.6269 \pm 0.6	2 ⁻	0.310 \pm 0.020	γ , n	17, 18
6.6425 \pm 0.6	(3, 4)	< 0.08	γ , n	17, 38
6.6474 \pm 0.7	1 ⁻	1.59 \pm 0.10	γ , n	17, 18, 38
6.6933 \pm 0.8	1 ⁻	13.8 \pm 0.8	γ , n	17, 18, 38
6.7660 \pm 1.1		\leq 0.6	γ , n	17, 38
6.829			n	19
6.8566 \pm 1.2	2	10 \pm 2	γ , n	17, 19, 38
(6.858 \pm 8)	1		γ , n	17
6.905 \pm 8				38
6.9677 \pm 1.2	1 ⁻	5 \pm 1	γ , n	17, 19
(7.0670 \pm 1.2)	0 ⁻	(2.4 \pm 0.6)	γ , n	17, 19
7.076	(1 ⁺)	24	n	18
7.166 \pm 2	2 ⁽⁺⁾	8 \pm 1	γ , n	17, 18, 19
7.311	(1)	33	γ , n	17, 18
7.361	(1)	19	n	18, 19
7.410	(2 ⁺)	10	γ , n	17, 18, 19
7.50	(2)	80	γ , n	17, 18
7.67	(2 ⁺)	65	γ , n	17, 18, 19
7.79		140	n	18, 19
(7.831 \pm 12)	1 ⁻	(50 \pm 10)	γ , n	17
7.988 \pm 3	1	14 \pm 2	γ , n	17
8.05 \pm 100	2 ⁺ ; 2			37
8.13		195	γ , n	17, 18, 19
8.163		15	n	19
8.421		27	n	19
8.50		140	n	18
8.728		\leq 30	n	18, 19
8.77		76	n	18

Table 20.4: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
8.942		73	n	18, 19
9.165			n	19
9.521		110	n	19
9.654		100	n	18, 19
9.830		33	n	19
9.85		120	n	18
(9.886 \pm 10)			n	18
9.901		≤ 30	n	19
(9.929 \pm 10)			n	18
(9.981 \pm 10)			n	18
10.024 \pm 10		150	n, α	18, 19, 24
10.10 \pm 50			n, α	24
10.228 \pm 10	$0^-, 1$	≈ 200	n, α	18, 24
10.480 \pm 10		≈ 10	n, α	18, 24
10.641 \pm 10	1, 2	70	n	18, 19
10.807 \pm 10	$0^-, 1$	≈ 310	n, α	18, 24
10.988		190	n	19
(11.045 \pm 10)		≈ 30	n	18
(11.130 \pm 10)		< 25	n	18
(11.244 \pm 10)		< 25	n	18, 19
(11.287 \pm 10)			n	18
11.49 \pm 50			n, α	24
12.0			n, α	24
12.2 \pm 100			n, α	24
12.39			n, α	24
12.82			n, α	24
13.2			n, α	24
13.66			n, α	19, 24
14.0			n, α	24

^a See also Tables 20.5 and 20.6.

Table 20.5: Radiative transitions in ^{20}F

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	δ	Refs.
0.66	3^+	0	2^+	100	0.10 ± 0.05	^a , (1973HA14)
0.82	4^+	0	2^+	41 ± 4		(1969HE20, 1970QU04)
		0.66	3^+	60 ± 5		(1969HE20, 1970QU04)
0.98	1^-	0	2^+	≥ 96	^b	(1973PR01)
		0.66	3^+	≤ 2		(1973PR01)
				< 1		(1968SP01)
		0.82	4^+	≤ 2		(1973PR01)
				< 1		(1968SP01)
1.06 ^c	1^+	0	2^+	≥ 96		(1973PR01)
				100		(1968SP01, 1970QU04)
		0.66	3^+	≤ 2		(1973PR01)
				< 1		(1968SP01)
		0.82	4^+	≤ 2		(1973PR01)
				< 1		(1968SP01)
1.31	2^-	0	2^+	100	^b	^a
		0.66	3^+	≤ 14		(1969HE20)
		0.82	4^+	< 2		(1968SP01)
		0.98	1^-	< 1		(1968SP01)
		1.06	1^+	< 1		(1968SP01)
1.82	5^+	0	2^+	< 3		(1972AL26, 1973PR01)
		0.66	3^+	< 3		(1972AL26)
		0.82	4^+	≥ 95	-0.03 ± 0.07	(1973PR01)
1.84 ^d	2^-	0	2^+	≥ 94		(1973PR01)
		0.66	3^+	≤ 6		(1973PR01)
		0.82	4^+	< 4		(1973PR01)
				< 3		(1972AL26)
		0.98	1^-	< 4		(1973PR01)
		1.31	2^-	< 4		(1973PR01)
1.97 ^d	(3^-)	0	2^+	16 ± 4	-0.06 ± 0.14	(1973PR01)
		0.66	3^+	< 3		(1973PR01)

Table 20.5: Radiative transitions in ^{20}F (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	δ	Refs.
2.04 ^e	2 ⁺	0.82	4 ⁺	55 ± 3	+0.27 ± 0.30	(1973PR01)
		0.98	1 ⁻	< 3		(1973PR01)
		1.31	2 ⁻	29 ± 3		(1973PR01)
		0	2 ⁺	8 ± 4		(1973PR01)
		0.66	3 ⁺	92 ± 4		0.08 ^{+0.06} _{-0.1}
2.19 ^e	(3 ⁺)	0.82	4 ⁺	< 5	0 ± 0.09	(1973PR01)
		0	2 ⁺	58 ± 4		(1973PR01)
		0.66	3 ⁺	< 5		(1973PR01)
2.86	3 ⁺	0.82	4 ⁺	42 ± 4	+0.07 ± 0.10	(1973PR01)
		0	2 ⁺	(100)		(1970QU04)
2.97 ^e	3 ⁺	0	2 ⁺	24 ± 3		(1969HE20)
				19 ± 5		(1973PR01)
		0.66	3 ⁺	14 ± 2		(1969HE20)
				17 ± 5		(1973PR01)
		0.82	4 ⁺	62 ± 6		(1969HE20)
				35 ± 4		(1973PR01)
		1.97	(3 ⁻)	29 ± 4		(1973PR01)
3.17 ^f	(1 ⁺)	0	2 ⁺	< 5		(1970QU04)
		0.98	1 ⁻	> 95		(1970QU04)
3.49	1 ⁺	0	2 ⁺	68 ± 4		(1969HO20)
		0.98	1 ⁻	7 ± 1		(1969HO20)
		1.06	1 ⁺	7 ± 1		(1969HO20)
		1.31	2 ⁻	10 ± 2		(1969HO20)
		1.84	2 ⁻	8 ± 2		(1969HO20)
		3.53	0 ⁺	1.06		1 ⁺
3.59	(1, 2, 3) ⁺	0	2 ⁺	66		(1968SP01)
				60		(1969HO20)
		2.04	2 ⁺	34		(1968SP01)
3.68	(1, 2, 3) ⁺			40		(1969HO20)
		0	2 ⁺	33		(1968SP01)

Table 20.5: Radiative transitions in ^{20}F (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	δ	Refs.
3.76	1^+	0.66	3^+	67		(1968SP01)
		0.66	3^+	observed		(1973PR01)
3.97 ^g	1^+	0.98	1^-	22 ± 7		(1973PR01)
4.08 ^h	$(1)^+$	1.31	2^-	78 ± 7		(1973PR01)
		0	2^+	35 ± 5		(1969HO20)
				45 ± 7		(1973PR01)
		1.06	1^+	65 ± 5		(1969HO20)
4.28	$(1, 2, 3)^+$			55 ± 7		(1973PR01)
		1.06	1^+	100		(1968SP01)
5.41		2.04	2^+	100		(1968SP01)
5.56		0	2^+	24		(1968SP01)
		1.31	2^-	76		(1968SP01)
5.71		1.06	1^+	60		(1969HA04)
		4.28	$(1, 2)^+$	40		(1969HA04)
5.94 ^g	$(0, 1, 2)^-$	0	2^+	7		(1968SP01)
		0.66	3^+	35		(1968SP01)
		0.98	1^-	4		(1968SP01)
		1.31	2^-	1		(1968SP01)
		1.84	2^-	7		(1968SP01)
		1.97	(3^-)	31		(1968SP01)
		2.19	(3^+)	5		(1968SP01)
		3.49	1^+	10		(1968SP01)
		0	2^+	29		(1968SP01)
		40				(1968BL1C)
6.02 ^g	$(0, 1, 2)^-$	0.66	3^+	3		(1968SP01)
		0.98	1^-	19		(1968SP01)
				23		(1968BL1C)
		1.06	1^+	1		(1968BL1C)
		1.31	2^-	2		(1968BL1C)
		1.84	2^-	6		(1968SP01)

Table 20.5: Radiative transitions in ^{20}F (continued)

E_i (MeV)	J_i^π	E_f (MeV)	J_f^π	Branching (%)	δ	Refs.
6.04				6		(1968BL1C)
		2.19	(3 ⁺)	3		(1968SP01)
		2.97	3 ⁺	8		(1968SP01)
		3.49	1 ⁺	19		(1968SP01)
				18		(1968BL1C)
		3.59	(1, 2, 3) ⁺	10		(1968SP01, 1968BL1C)
		4.08		3		(1968SP01)
		1.31	2 ⁻	46		(1968SP01)
1.84	2 ⁻	54		(1968SP01)		
6.60	0 ⁺ , 1 ⁺	see Table 20.10				
6.63	2 ⁻	see Tables 20.9 and 20.10				
6.65	1 ⁻	see Tables 20.9 and 20.10				

^a See Table 20.6 in (1972AJ02).

^b Pure E1 (1973HA14).

^c See also (1973HA14).

^d See also (1970QU04, 1974KE18).

^e See also (1970QU04).

^f See also (1972AL26).

^g See also (1969HA04).

^h See also (1974KE18).

4. $^{16}\text{O}(^7\text{Li}, ^3\text{He})^{20}\text{F}$ $Q_m = -4.743$

See (1974BI1D).

5. $^{17}\text{O}(t, p)^{19}\text{O}$ $Q_m = 3.520$ $E_b = 14.157$

See ^{19}O .

Table 20.6: Lifetime measurements of some ^{20}F states ^a

$^{20}\text{F}^*$ (MeV)	τ_m	Refs.
0.66	0.36 ± 0.07 psec	(1969HE20)
	0.37 ± 0.06 psec	(1970HO05)
	0.42 ± 0.05 psec	(1973WA10)
	0.39 ± 0.04 psec	(1975SE04)
	0.39 ± 0.03 psec	mean
0.82	79 ± 6 psec ^A	(1971PR10)
0.98	$1.3^{+0.6}_{-0.4}$ psec	(1969HE20)
	1.8 ± 0.4 psec	(1970HO05)
	2.30 ± 0.25 psec	(1973WA19)
	2.03 ± 0.20 psec	mean
1.06	45 ± 13 fsec ^A	(1970HO05)
1.31	$1.1^{+0.4}_{-0.3}$ psec	(1969HE20)
	0.8 ± 0.3 psec	(1970HO05)
	1.45 ± 0.25 psec	(1973WA19)
	1.16 ± 0.20 psec	mean
1.82	≤ 65 fsec ^A	(1973PR01)
1.84	30 ± 20 fsec ^A	(1970HO05)
1.97 ^b	1.4 ± 0.4 psec ^A	(1973WA19)
2.04	37 ± 16 fsec ^A	(1970HO05)
2.19	< 12 fsec ^A	(1970HO05)
2.97	60 ± 40 fsec ^A	(1970HO05)
3.49	44 ± 11 fsec ^A	(1970HO05)
3.53	30 ± 15 fsec ^A	(1970HO05)
3.59	30 ± 30 fsec ^A	(1970HO05)

A = Adopted.

^a See also Table 20.14 in (1972AJ02).

^b See also (1973PR01).

Table 20.7: States of ^{20}F from $^{14}\text{N}(^7\text{Li}, \text{p})^{20}\text{F}$ ^a

E_x ^b (MeV)	J^π	σ_t (μb)	E_x (MeV)	J^π	σ_t (μb)
0	2^+	23.8	2.86	$2^-, 3, 4^+$ ^d	40.7
0.66	3^+	37.4	2.97	$4^-, 5, 6^+, 7^+$ ^d	68.2
0.82	4^+	48.2	3.50 ^c	$1^+ + 0^+$	19.6
0.98	1^-	20.2	3.59	$2^-, 3, 4, 5^+$ ^d	47.6
1.06	1^+	15.0	3.68	$3^-, 4, 5, 6^+$ ^d	58.6
1.31	2^-	33.8	4.20 + 4.21	^e	130
1.83 ^c	$5^+ + 2^-$	85.7	4.52	$3^-, 4^-, 5, 6^+, 7^+$ ^d	63.8
1.97	(3^-)	48.9	4.58 + 4.59		80
2.04	2^+	24.5	4.76	$3^-, 4^-, 5, 6^+$ ^d	61.4
2.19	3^+	35.4	4.89 + 4.90		54

^a (1975BI04): $E(^7\text{Li}) = 16$ MeV.

^b Nominal energy.

^c Unresolved.

^d Based on σ_t measurements: see text.

^e One of these two states has $J^\pi \geq 4^-$ or 5^+ .

6. $^{17}\text{O}(\alpha, \text{p})^{20}\text{F}$ $Q_m = -5.657$

Not reported.

7. $^{17}\text{O}(^6\text{Li}, ^3\text{He})^{20}\text{F}$ $Q_m = -1.637$

See (1977MA2G).

8. $^{18}\text{O}(\text{d}, \text{n})^{19}\text{F}$ $Q_m = 5.7685$ $E_b = 12.3699$

See ^{19}F .

9. $^{18}\text{O}(\text{d}, \text{p})^{19}\text{O}$ $Q_m = 1.732$ $E_b = 12.3699$

Vector analyzing power measurements have been carried out at $E_{\bar{d}} = 14.8$ MeV for the proton groups to $^{19}\text{O}^*(0, 1.47, 3.15, 3.95, 4.12, 4.58, 4.71, 5.00 + 5.09, 5.15, 5.46, 5.71, 6.28)$ (1974SE01): see ^{19}O . Measurements at $E_{\bar{d}} = 10$ MeV are reported by (1973ST1B; abstract). See also (1974FO1J, 1976DA1K).

10. $^{18}\text{O}(\text{d}, \text{d})^{18}\text{O}$ $E_{\text{b}} = 12.3699$

Vector analyzing power measurements involving the elastic group have been carried out at $E_{\bar{d}} = 14.8$ MeV (1974SE01).

11. $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$ $Q_{\text{m}} = 4.246$ $E_{\text{b}} = 12.3699$

Excitation functions have been measured for $E_{\text{d}} = 0.9$ to 3.0 MeV and 9.6 to 11.5 MeV: see (1972AJ02). At the lower energies a number of sharp structures are reported.

12. $^{18}\text{O}(\text{t}, \text{n})^{20}\text{F}$ $Q_{\text{m}} = 6.1126$

Measurements of τ_{m} of several ^{20}F states are reported by (1971PR10, 1973PR01, 1973WA19): see Table 20.6.

13. $^{18}\text{O}(^3\text{He}, \text{p})^{20}\text{F}$ $Q_{\text{m}} = 6.8764$

Proton groups have been observed to states of ^{20}F with $E_{\text{x}} < 4.1$ MeV (1970RO06, 1974CR04): see Table 20.8. Angular distribution measurements at $E(^3\text{He}) = 18$ (1974CR04) and 19 MeV (1976ME14), γ -ray polarization data (1973LO13) and branching ratio and life-time measurements (1972AL26, 1973PR01) lead to the J^{π} values shown in Table 20.8 [see also Tables 20.5 and 20.6]. At $E(^3\text{He}) = 5$ MeV, a state is populated at $E_{\text{x}} = 6519 \pm 3$ keV (1976MI01), which decays principally ($> 90\%$) to $^{20}\text{F}^*(1.06)$: the γ -rays are isotropic [$\Gamma_{\gamma} = 3.6 \pm 0.6$ eV, based on the analog decay in ^{20}Ne]. $^{20}\text{F}^*(6.52)$ is the 0^+ ; $T = 2$ analog of the ground state of ^{20}O (1976MI01, 1977BA50). See also (1971NE1E).

14. $^{18}\text{O}(\alpha, \text{d})^{20}\text{F}$ $Q_{\text{m}} = -11.4769$

Not reported.

Table 20.8: States in ^{20}F from $^{18}\text{O}(^3\text{He}, \text{p})^{20}\text{F}$

E_x (keV)		L^b	J^π^b
(1970RO06)	(1974CR04)		
0	0	2	2_1^+
657.2 ± 1.3	656	$2 + 4$	3_1^+ ^d
823.5 ± 1.5	822.6 ± 1.9	4	4_1^+ ^d
982.9 ± 1.3	983.3 ± 5.3	c	1^- ^e
1058.1 ± 1.4	1057.5 ± 2.4	$0 + 2$	1_1^+
1309.1 ± 1.4	1310.2 ± 3.1	c	2^- ^e
1824.4 ± 1.6^a	1824.1 ± 3.6	4	5_1^+ ^d
1843.0 ± 1.7^a			(2^-) ^e
1971.9 ± 1.6	1978.0 ± 2.8	c	(3^-) ^g
2044.0 ± 1.6	2044.9 ± 2.2	2	2_2^+
2195.5 ± 2.0	2194.7 ± 2.8		(3^+) ^g
2868.2 ± 2.3	2863.6 ± 3.9	c	
2967.1 ± 2.0	2961.4 ± 3.5		see ^b
	3167.2 ± 3.8^i	$(0 + 2)^h$	$(1^+)^h$
3487.8 ± 2.2	3485.9 ± 2.3	$0 + 2$	1_2^+
	3.53 ^j	c	$(0^+)^j$
3586.3 ± 2.2	3583.1 ± 2.7		see ^b
3681.0 ± 2.5	3669.4 ± 4.9		see ^b
3761.0 ± 3.1^f	3760 ± 10	c	
3966.9 ± 2.8		$0 + 2^h$	1^+^h
4083.7 ± 2.9			
6519 ± 3^k			$0^+; T = 2^k$

^a (1967QU01) find $E_x = 1824.4 \pm 2.1$ and 1843.0 ± 2.2 keV.

^b $E(^3\text{He}) = 18$ MeV (1974CR04): predominant L -values.

^c Weakly populated.

^d Also (1973LO13, 1973PR01).

^e (1973LO13).

^f $E_x = 3765 \pm 6$ keV, based on $E_x = 657 \pm 1$ keV (1973PR01).

^g Suggested by (1973PR01).

^h (1976ME14): $E(^3\text{He}) = 19$ MeV.

ⁱ $E_x = 3175 \pm 6$ keV (1972AL26), based on $E_x = 983 \pm 1$ keV.

^j See (1971FO14).

^k (1976MI01).

15. (a) $^{18}\text{O}(^6\text{Li}, \alpha)^{20}\text{F}$ $Q_m = 10.896$
 (b) $^{18}\text{O}(^7\text{Li}, \alpha n)^{20}\text{F}$ $Q_m = 3.646$

See (1972AJ02).

16. $^{18}\text{O}(^{18}\text{O}, ^{15}\text{N})^{20}\text{F}$ $Q_m = -1.6505$

See (1972EY01).

17. $^{19}\text{F}(n, \gamma)^{20}\text{F}$ $Q_m = 6.6012$
 $Q_0 = 6600.4 \pm 0.5 \text{ keV}$ (1974SP04).

The thermal capture cross section is $9.8 \pm 0.7 \text{ mb}$ (1974SH1E). See also (1973MU14). A number of resonances have been observed for $E_n \leq 1.65 \text{ MeV}$: see Table 20.9 (1959GA08, 1971NY02, 1973MA14). The primary γ -rays resulting from capture at thermal energies ($^{20}\text{F}^*(6.60)$; $J^\pi = 0^+$, 1^+) and at $E_n = 27, 44$ and 49 keV ($^{20}\text{F}^*(6.63, 6.643, 6.647)$; $J^\pi = 2^-, (3, 4)$ and 1^- , respectively) have been studied by several groups: see (1972AJ02) and Table 20.10 here (1967BE36, 1968SP01, 1969HA04, 1974KE18). It appears that the decay of $^{20}\text{F}^*(6.60)$ is dominated by two intense transitions (probably E1) to $^{20}\text{F}^*(5.94, 6.02)$ [thus $J^\pi = 1^-, 2^-$]. If the ground-state transition is mainly M1, these two E1 transitions are (in terms of W.u.) about 150 times stronger than the M1 transition (1968SP01). It appears also that at $^{20}\text{F}^*(6.63, 6.64, 6.65)$ [$J^\pi = 2^-, (3, 4)$ and 1^- , respectively] the E1 transitions to the ground state are very weak, even though other E1 transitions in the decay of these two states have approximately normal strengths (1967BE36, 1974KE18). The strongest transitions from the 27 keV resonance appear to be M1. On the basis of the J^π of the final states involved in the decay of the 44 keV resonance, $J = 3$ or 4 , assuming dipole transitions (1974KE18). Branching ratios for other ^{20}F states involved in this reaction are shown in Table 20.5.

Table 20.11 displays excitation energies for ^{20}F states involved in cascade and in primary γ transitions (1968SP01, 1969HA04, 1972OP01).

18. $^{19}\text{F}(n, n)^{19}\text{F}$ $E_b = 6.6012$

The scattering amplitude (bound) is $a = 5.66 \pm 0.02 \text{ fm}$ (1975KO1K). The value of the coherent scattering cross section recommended by (1973MU14) is $4.0 \pm 0.1 \text{ b}$. (1974DI1D) find $3.58 \pm 0.02 \text{ b}$ and $a = 5.60 \pm 0.01 \text{ fm}$. The spin-dependent part of the scattering amplitude $\beta \equiv a_+ - a_- = -0.135 \pm 0.002 \text{ fm}$ (1972AB1A).

Table 20.9: Resonances in $^{19}\text{F}(n, \gamma)^{20}\text{F}$

E_n (keV)	J^π ^a	Γ_γ (eV)	$\Gamma_{\text{c.m.}}$ (keV)	E_x in ^{20}F (MeV)	Refs.
27.07 ± 0.05	2^-	1.4 ± 0.3	0.355 ± 0.03	6.6269	(1973MA14, 1971NY02)
43.5 ± 0.1	$(3, 4)$ ^f	^b	< 0.08	6.6425	(1973MA14)
48.7 ± 0.3	1^-	1.6 ± 0.3	1.96 ± 0.3	6.6474	(1973MA14, 1971NY02)
97.0 ± 0.5	1^-	6.0 ± 1.8 ^c	13.5 ± 1.5	6.6933	(1973MA14)
173.5 ± 0.9		^d	≤ 0.6	6.7660	(1973MA14)
269 ± 1	2	3.5 ± 0.8	10 ± 2	6.8566	(1973MA14)
(270 ± 8)	1	≤ 4.4		(6.858)	(1973MA14)
386 ± 1	1^-	2.4 ± 0.8 ^g	5 ± 1	6.9677	(1973MA14)
(490.5 ± 1)	0^-	$(\geq 10 \pm 3)$	(2.4 ± 0.6)	(7.0670)	(1973MA14)
595 ± 2	2	6.3 ± 1.2 ^g	8 ± 1	7.166	(1973MA14)
760		2.9	60	7.32	(1959GA08)
865			60	7.42	(1959GA08)
950		2.8	95	7.50	(1959GA08)
1125		3.9	80	7.67	(1959GA08)
(1295 ± 12)	1^-	8.6 ^g	(50 ± 10)	(7.831)	(1973MA14, 1959GA08)
1460 ± 3	1	$\geq 11 \pm 3$	14 ± 2	7.988	(1973MA14)
1635		11 ± 3 ^g	180	8.15	(1959GA08)

^a Assumed: (1973MA14).

^b $g\Gamma_n = 0.086 \pm 0.02$ eV (1973MA14).

^c May be two resonances.

^d $g\Gamma_n = 0.35 \pm 0.1$ eV (1973MA14).

^e See also Table 20.8 in (1972AJ02) and (1973MU14).

^f (1974KE18).

^g (1973MU14).

Table 20.10: Primary capture transitions in $^{19}\text{F}(n, \gamma)^{20}\text{F}^{\text{a}}$

Final state $^{20}\text{F}^*$ (MeV)	I_γ^{b} from $^{20}\text{F}^*(6.60)$		I_γ^{b} from $^{20}\text{F}^*(6.63)$		I_γ^{b} from $^{20}\text{F}^*(6.64)$	I_γ^{b} from $^{20}\text{F}^*(6.65)$
	(1968SP01)	(1969HA04)	(1974KE18)	(1967BE36)	(1974KE18)	(1974KE18)
0	11	10	2.0 ± 0.5			
0.66	< 0.1		6 ± 1	6 ± 2	42 ± 7	
0.82	< 0.1				23 ± 7	
0.98	1.5	2				18 ± 4
1.06 ^c	5	6				9 ± 4
1.31	3	3	31 ± 2	32 ± 2		
1.84	2	2	8 ± 2			
1.97	0.1		46 ± 4	50 ± 2		
2.04	6	6	1.5 ± 1			59 ± 6
2.19						
2.87						
2.97	< 0.2				35 ± 9	
3.49	2.5	2	3 ± 1			14 ± 5
3.53	2			8 ± 1		
3.59	4.4	4				
3.68	1	1				
3.97		1				
4.08		1	2.5 ± 1	5 ± 2		
4.28	1	1				
5.04						
5.41	0.5					
5.55	3					
5.94	17	13				
6.02 ^c	38	48				
6.04	2					

^a See also Tables 20.5 and 20.9.

^b In units of photons/100 captures.

^c E_γ for the transitions (6.60 \rightarrow 0), (6.60 \rightarrow 1.06) and (6.60 \rightarrow 6.02) are, respectively, 6599.8 ± 3.0 , 5534.9 ± 2.0 and 583.6 ± 0.5 keV (1967VA08).

Table 20.11: States of ^{20}F involved in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

E_x (keV)	
(1968SP01)	(1969HA04)
0	0
656.3 ± 0.3 ^b	656 ± 1
822.9 ± 0.3	
983.8 ± 0.3 ^b	984 ± 1
1057.2 ± 0.3	1057 ± 1
1309.1 ± 0.3	1309 ± 1
1843.4 ± 0.3	1843 ± 1
1970.6 ± 0.3	
2044.2 ± 0.4	2044 ± 1
2194.5 ± 0.6	2194 ± 2
2965.8 ± 0.5	2966 ± 2
3488.3 ± 0.3	3488 ± 2
3526.0 ± 0.5	
3587.3 ± 0.3	3588 ± 2
3681.0 ± 0.4	3681 ± 2
	3967 ± 2
4082.2 ± 0.5	4085 ± 2
4276.7 ± 0.5	4275 ± 2
5413.1 ± 0.6	
5554.7 ± 6	
	5713 ± 2
5936.0 ± 0.3	5937 ± 1
6017.3 ± 0.3	6018 ± 1
6044.6 ± 0.4	
6601.1 ± 0.3	6602.0 ± 0.6

^a For measurements of I_γ , see (1968SP01, 1969HA04) and Table 20.10.

^b 656.1 ± 0.3 and 983.4 ± 0.4 keV (1972OP01).

The total cross section has been measured for $E_n = 0.5$ to 29.1 MeV: see the listings in (1972AJ02, 1973MU14), the display in (1976GAYV) and (1974SI27: $E_n = 0.5$ to 200 keV), (1974JO1H: $E_n = 20$ keV to 10 MeV; abstract). Observed resonances are displayed in Table 20.12. For angular distribution measurements see ¹⁹F. For polarization measurements see (1972AJ02). See also (1971KO1W) and (1973BA1U, 1974HU07; theor.).

$$19. \text{}^{19}\text{F}(n, n')\text{}^{19}\text{F}^* \qquad E_b = 6.6012$$

Recent excitation functions have been measured for the 0.11 and 0.20 MeV γ -rays for E_n from threshold to 19.4 and 18.6 MeV, respectively: the differential cross section at 92° for the 0.11 MeV γ -ray shows a number of resonances the most pronounced of which are displayed in Table 20.13. The 0.20 MeV γ -ray yield does not show clear resonance structure (1976MO13). See also (1974RO03: $E_n = 0.2$ to 1.7 MeV). (1973MA14) has studied the inelastic yield involving the states of ¹⁹F near 1.5 MeV: observed resonances are also displayed in Table 20.13. See also (1969RO1F, 1977DI1A), (1975FU1D) and (1972AJ02) for the earlier work.

$$20. \text{}^{19}\text{F}(n, 2n)\text{}^{18}\text{F} \qquad Q_m = -10.4313 \qquad E_b = 6.6012$$

Cross sections have been measured for $E_n = 10$ to 37 MeV: see (1972AJ02) for the earlier work, (1972MO15: $E_n = 14.1$ MeV) and (1973RO29: $E_n = 14.78$ MeV) and the summary in (1976GAYV). See also (1973BO1K, 1974BO1E, 1974CA1J, 1974KO35).

$$21. \text{}^{19}\text{F}(n, p)\text{}^{19}\text{O} \qquad Q_m = -4.036 \qquad E_b = 6.6012$$

The differential cross section at 92° for production of the 96 keV γ -ray has been studied by (1976MO13: $E_n = 4.0$ to 18.6 MeV): the cross section increases sharply at $E_n = 6$ MeV and then gradually decreases beyond $E_n = 12$ MeV. Cross sections have also been measured for $E_n = 12.6$ to 21 MeV: see (1972AJ02) and the summary in (1976GAYV). See also (1972BA1T, 1973BA1T; abstracts) and (1972ED01, 1974BO1E, 1974CA1J, 1976SL2A, 1977DI1A).

$$22. \text{}^{19}\text{F}(n, d)\text{}^{18}\text{O} \qquad Q_m = -5.7688 \qquad E_b = 6.6012$$

See (1972AJ02) and (1976PR08, 1976SL2A).

$$23. \text{}^{19}\text{F}(n, t)\text{}^{17}\text{O} \qquad Q_m = -7.556 \qquad E_b = 6.6012$$

Table 20.12: Resonances in $^{19}\text{F}(n, n)^{19}\text{F}$

E_n (keV)	Γ_{lab} (keV)	J^π	$^{20}\text{F}^*$ (MeV)	Refs.
26.99	0.325 ± 0.020	2^-	6.6268	(1974JO1H, 1974SI27)
48.78	1.67 ± 0.10	1^-	6.6475	(1974JO1H, 1974SI27)
97.50	14.5 ± 0.8	1^-	6.6938	(1974JO1H, 1974SI27)
a				
500	25^b	(1^+)	7.076	(1966CA14)
600	15^b	(2^+)	7.171	(1966CA14)
747	35^b	(1)	7.311	(1966CA14)
794	20	(1)	(7.355)	(1966CA14)
852	11^b	(2^+)	7.410	(1966CA14)
935	60	(2)	7.489	(1958WI36, 1966CA14)
1100	50	(2^+)	7.65	(1958WI36, 1966CA14)
1250	150		7.79	(1958WI36)
1620	220		8.14	(1958WI36)
2000	150		8.50	(1958WI36)
2250	≤ 30		8.74	(1958WI36)
2280	80		8.77	(1958WI36)
2520	150		8.99	(1958WI36)
3250	150		9.69	(1958WI36)
3420	130		9.85	(1958WI36, 1960TS02)
3460 ± 10			(9.886)	(1960TS02)
3505 ± 10			(9.929)	(1960TS02)
3560 ± 10			(9.981)	(1960TS02)
3605 ± 10	200		10.024	(1958WI36, 1960TS02)
3820 ± 10	≈ 200	$0^-, 1$	10.228	(1960TS02)
4085 ± 10	≈ 10		10.480	(1960TS02)
4255 ± 10	≈ 60	1, 2	10.641	(1960TS02)
4430 ± 10	≈ 330	$0^-, 1$	10.807	(1960TS02)
4680 ± 10	≈ 30		11.045	(1960TS02)
4770 ± 10	< 25		11.130	(1960TS02)
4890 ± 10	< 25		11.244	(1960TS02)
(4935)			(11.287)	(1960TS02)

^a See (1976GAYV) for resonances in σ_t derived unpublished work.

^b $\Gamma_\gamma = 3.3 \pm 1.0, 6.3 \pm 1.2, 2.4 \pm 0.8$ and 1.5 ± 0.5 eV for $^{20}\text{F}^*(7.08, 7.17, 7.31, 7.41)$ (1973MU14).

Table 20.13: States of ^{20}F from resonances in $^{19}\text{F}(n, n'\gamma)^{19}\text{F}$

E_n (keV)	Γ_{lab} (keV)	Resonance in		E_x in ^{20}F (MeV)
		$\gamma_{0.11}$ ^a	$\gamma_{1.5}$ ^b	
240		r		6.829
270		r		6.858
386		r		6.968
420		r		7.000
490		r		7.066
620		r		7.190
800		r		7.361
860		r		7.418
1150 ^c		r		7.693
1250		r		7.788
1580		r		8.101
1645	15	r	r	8.163
1916	28		r	8.421
2240	45		r	8.728
2465	75	r	r	8.942
2700		r		9.165
3075	120		r	9.521
3215	80		r	9.654
3400	35		r	9.830
3475	≤ 30		r	9.901
3620	120	r	r	10.038
4240	90	r	r	10.627
4620	200		r	10.988
4900	≤ 50		r	11.254
7300		r		13.532

r = resonant.

^a Resonances in yield of 0.11 MeV γ -rays at $\theta = 92^\circ$: values for E_n read by reviewer from differential cross section tables (1976MO13).

^b Resonances in yields to states of ^{19}F with $E_x \approx 1.5$ MeV: see (1973MA14).

^c Appears to be unresolved.

Table 20.14: Resonances in $^{19}\text{F}(n, \alpha)^{16}\text{N}$

E_n (MeV \pm keV)			E_x (MeV)
(1955MA1D)	(1960SM03)	(1961DA16)	
3.4			9.8
3.61 ± 50			10.03
3.69 ± 50			10.10
3.77 ± 50	3.75 ± 50	3.85	10.17
4.11 ± 50	4.08 ± 50	4.1	10.49
4.42 ± 50	4.36 ± 50	4.35	10.77
	4.52 ^a		10.89
4.86 ± 50	4.79 ± 50	4.80	11.18
	5.15 ± 50		11.49
	5.40 ^a		11.73
		5.7	12.0
5.9 ± 100	5.9 ^a		12.2
		6.10	12.39
		6.55	12.82
		6.9	13.2
		7.44	13.66
		7.8	14.0

^a Not resolved.

See ^{17}O in (1977AJ02) and (1976SL2A).

24. $^{19}\text{F}(n, \alpha)^{16}\text{N}$

$$Q_m = -1.522$$

$$E_b = 6.6012$$

Observed resonances are shown in Table 20.14: see graph in (1976GAYV). See also (1972FO21: $E_n = 5.85$ MeV, $\sigma_{n,\alpha} = 135 \pm 27$ mb). See also (1972BA1T) and (1974BO1E, 1974CA1J, 1976SL2A).

25. $^{19}\text{F}(d, p)^{20}\text{F}$

$$Q_m = 4.3765$$

Measurements of proton groups and of γ -rays have led to the identification of a number of ^{20}F states: see Table 20.15. Angular distributions have been measured for $E_d = 0.6$ to 13 MeV (see (1972AJ02)) and at $E_d = 1.00 - 1.50$ MeV (1976BE51), 1.20 to 2.20 MeV (1975CO18: p₁), 3 MeV (1972LA18; l_n and $(2J + 1)S$ values for a number of states) and at 12 (1974FO21) and 16 MeV (1972FO11). The J^π and $(2J + 1)S$ values derived from the experiments at 12 and 16 MeV are displayed in Table 20.15. (1975MC1J) have studied the $l_n = 2$ transitions to $^{20}\text{F}^*(0.66, 2.04, 2.19, 2.97)$ with $E_d = 11$ MeV while (1973JO10) report a study at $E_d = 12.3$ MeV involving $^{20}\text{F}_{\text{g.s.}}$. (1970QU1A), also using polarized deuterons, have looked at the protons corresponding to $^{20}\text{F}^*(0.66, 2.04)$. These two states are found to be populated by a $j = \frac{5}{2}$ neutron transfer. This result, together with (p- γ) correlation data, provides a unique $J^\pi = 3^+$ assignment for $^{20}\text{F}^*(0.66)$ and is in agreement with 2^+ for $^{20}\text{F}^*(2.04)$. Transitions to $^{20}\text{F}^*(2.19, 2.97)$ appear to involve admixed $j = \frac{3}{2}, \frac{5}{2}$ transfer implying $J = 2$ for both these states (1970QU1A).

Table 20.15: States in ^{20}F from $^{19}\text{F}(d, p)^{20}\text{F}$

E_x (keV)			l_n ^{d,e}	J^π ^{d,e}	$(2J + 1)S$		
(1970RO06) ^a	(1969HO20) ^{b,c}	(1972FO11) ^a			$E_d = 12$ MeV ^d	16 MeV ^e	n, l, j ^h
0			2	2^+	0.054	≤ 0.06	$1d_{5/2}$
654.9 ± 1.0	655.9 ± 0.2	655 ± 1	2	3^+	2.32	2.59	$1d_{5/2}$
821.6 ± 1.0	823.0 ± 0.3	823 ± 5	i	4^+	0.32	0.36	$1g_{9/2}$
983.3 ± 1.0	983.9 ± 0.3	983 ± 5	i	1^-	0.014	0.016	$1p_{1/2}$
1056.3 ± 1.0	1057.0 ± 0.2	1056 ± 3	$0 + 2$	1^+	0.013	0.019	$2s_{1/2}$
1310.8 ± 1.1	1309.3 ± 0.2	1309 ± 5	i	2^-	0.017	0.013	$1p_{3/2}$
		1820 ± 10	i	(5^+)	0.35	0.32	$1g_{9/2}$
1843.4 ± 1.2	1843.5 ± 0.7	1845 ± 4	i	2^-	0.007	0.03	$2p_{3/2}$
		1970 ± 10	i	(3^-)	0.038	0.042	$1f_{7/2}$
	2043.7 ± 0.5	2044 ± 1	2	2^+	2.32	2.32	$1d_{5/2}$
2195.1 ± 1.5	2194.5 ± 0.6	2196 ± 1	2	3^+	0.55	0.50	$1d_{5/2}$
2863.7 ± 1.6		2871 ± 5	i		0.044		$1f_{7/2}$
2966.6 ± 1.7	2966.8 ± 0.6	2966 ± 1	2	3^+	0.38	0.36	$1d_{3/2}$
3171.8 ± 2.2	3175.6 ± 1.3	3176 ± 5	i		0.019	0.014	$1d_{5/2}$
	3488.5 ± 0.3	3489 ± 1	0	1^+		1.20	$2s_{1/2}$
3525.5 ± 2.6	3525.9 ± 0.5	3531 ± 3	0	0^+		0.28	$2s_{1/2}$
3586.4 ± 2.7	3586.5 ± 0.6	3590 ± 1	2	$\pi = +$	0.038	0.42	$1d_{3/2}$
3681.0 ± 2.5		3686 ± 4	2	$\pi = +$	0.031	≤ 0.04	$1d_{5/2}$
3760.8 ± 2.7			i		see ^d		
3964.5 ± 2.5		3977 ± 5	2	$\pi = +$	0.036	0.043	$1d_{5/2}$
4080.9 ± 2.5	4082.5 ± 0.8	4089 ± 3	$0 + 2$	$\pi = +$	0.13	0.18	$1s_{1/2}$
4198.9 ± 2.7							
4207.7 ± 2.6			i		0.083	0.10	$1d_{3/2}$
4276.3 ± 2.8		4282 ± 5	2	$\pi = +$			
4311.5 ± 2.6		4318 ± 5	0	$(0, 1)^+$			

Table 20.15: States in ^{20}F from $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$ (continued)

E_x (keV)			l_n ^{d,e}	J^π ^{d,e}	$(2J + 1)S$		
(1970RO06) ^a	(1969HO20) ^{b,c}	(1972FO11) ^a			$E_d = 12$ MeV ^d	16 MeV ^e	n, l, j ^h
4583.8 ± 3.0							
4592.2 ± 2.9							
4730.2 ± 2.9							
4763.8 ± 2.7							
4891.6 ± 2.8							
4898.2 ± 2.8							
5040.2 ± 3.1			1 ^j	(0, 1, 2) ⁻			
5065.5 ± 3.1							
5224.0 ± 3.1			1 ^j	(0, 1, 2) ⁻			
5281.0 ± 3.3			1 ^j	(0, 1, 2) ⁻			
5317.1 ± 2.7							
5344.5 ± 3.3							
5450.3 ± 3.8							
5455.4 ± 3.2							
5463.4 ± 3.3							
5620.3 ± 3.3							
5762.8 ± 3.4							
5809.1 ± 2.9							
5933.9 ± 3.3 ^f			1 ^j	(0, 1, 2) ⁻			
6015.0 ± 3.8 ^f			1 ^f	(0, 1, 2) ⁻			
6043.3 ± 3.7		g					

^a From measurements of proton groups.

^b From measurements of γ -rays.

^c (1969HE20) find $E_x = 655.4 \pm 0.5, 822.6 \pm 0.7, 983.4 \pm 0.7, 1055.2 \pm 0.6, 1308.0 \pm 0.9$ and 2964.5 ± 2.0 keV.

^d (1974FO21): $E_d = 12$ MeV.

^e (1972FO11): $E_d = 16$ MeV.

^f (1971RO06) find $E_x = 5932 \pm 5$ and 6013 ± 5 keV, respectively.

^g (1956EL1A) find additional transitions to states with $E_x = 6250, 6520, 6630, 6810, 6980$ and $7200 [\pm 20]$ keV: the last four involve $l_n = 1$.

^h For $(2J + 1)S$ values for other n, l, j , see (1974FO21).

ⁱ Weak states: see (1972FO11, 1974FO21).

^j (1956EL1A).

Branching ratio, angular correlation and γ -ray polarization measurements, as well as τ_m determinations, permit a unique choice of J^π in many cases from among the J^π values stemming from direct interaction analyses of angular distributions: see Table 20.5 and (1973HA14, 1973WA10). See also (1971BE2F, 1972BA1V) and (1973DO02; theor.).

26. $^{19}\text{F}(t, d)^{20}\text{F}$ $Q_m = 0.3438$

See (1972AJ02).

27. $^{19}\text{F}(^9\text{Be}, ^8\text{Be})^{20}\text{F}$ $Q_m = 4.9360$

See (1977SW05).

28. $^{19}\text{F}(^{13}\text{C}, ^{12}\text{C})^{20}\text{F}$ $Q_m = 1.6548$

Total cross section measurements for formation of $^{20}\text{F}^*(0, 0.66)$ (1974LA27) and of $^{20}\text{F}^*(0.66)$ (1975SE03) have been measured for $E(^{13}\text{C}) = 10$ to 25 MeV and $E(^{19}\text{F}) = 13$ to 24 MeV, respectively. In the latter experiment σ_t is compared with DWBA assuming $S_1 S_2 = 0.54$; the population of $^{20}\text{F}^*(2.04)$ is also reported (1975SE03). τ_m for $^{20}\text{F}^*(0.66) = 390 \pm 40$ fsec (1975SE04): see Table 20.6.

29. $^{19}\text{F}(^{19}\text{F}, ^{18}\text{F})^{20}\text{F}$ $Q_m = -3.830$

See (1972AJ02).

30. $^{20}\text{O}(\beta^-)^{20}\text{F}$ $Q_m = 3.816$

See ^{20}O .

31. $^{20}\text{Ne}(n, p)^{20}\text{F}$ $Q_m = -6.2435$

See (1971KA18, 1976KI1D).

32. $^{20}\text{Ne}(t, ^3\text{He})^{20}\text{F}$ $Q_m = -7.0073$

Not reported.

Table 20.16: Analog states of $A = 20$ observed in $^{21}\text{Ne}(d, ^3\text{He})^{20}\text{F}$ and $^{21}\text{Ne}(d, t)^{20}\text{Ne}$ ^a

$^{20}\text{F}^*$ (MeV) ^b	$J\pi$	$^{20}\text{Ne}^*$ (MeV \pm keV)	l	C^2S			
				^{20}F		^{20}Ne	
0	2^+	10.27 ^b	$0 + 2$	0.24 + 0.58		0.08 + 0.25	
0.66	3^+	10.880 ± 10	2	0.66		0.42	
0.82	4^+	11.086 ± 10	2	0.26		0.18	
0.98	1^-		1		0.84		0.52
		11.27					
1.06	1^+		$0 + 2$	0.08 + 0.25		0.03 + 0.18	
1.31	2^-	11.601 ± 10	1		0.86		0.50
1.84	2^-	12.100 ± 10	1		0.69		0.43
2.04	2^+		2	0.15			
2.19	(3^+)		2	0.16			
			sums:	$l = 0 + 2$ 2.38	$l = 1$ 2.39	$l = 0 + 2$ 1.14	$l = 1$ 1.45

^a (1974MI13); $E_d = 26$ MeV; DWBA analysis of angular distributions. See Table 20.38 for $T = 0$ states in ^{20}Ne observed in the (d, t) reaction.

^b E_x are nominal.

33. $^{21}\text{Ne}(n, d)^{20}\text{F}$ $Q_m = -10.780$

Not reported.

34. $^{21}\text{Ne}(d, ^3\text{He})^{20}\text{F}$ $Q_m = -7.511$

The ^{20}F states observed, at $E_d = 26$ MeV, in this reaction and analog [$T = 1$] states observed in ^{20}Ne in the (d, t) reaction are displayed in Table 20.16. The spectroscopic factors of analog states are consistent to within 20% for states excited by a single l -transfer (1974MI13).

35. $^{21}\text{Ne}(t, \alpha)^{20}\text{F}$ $Q_m = 6.809$

Not reported.

Table 20.17: States of ^{20}F from $^{22}\text{Ne}(\text{d}, \alpha)^{20}\text{F}$ ^a

E_x (keV)	E_x (keV)	E_x (keV)	E_x (keV)
655 ± 2	3680 ± 3	5224 ± 6	6205 ± 6
824 ± 3	3762 ± 4	5276 ± 5	6240 ± 7
983 ± 3	3964 ± 4	5321 ± 4 ^b	6300 ± 5
1056 ± 4	4083 ± 4	5405 ± 5	6337 ± 5
1307 ± 3	4206 ± 4 ^b	5451 ± 5 ^b	6370 ± 6
1830 ± 7 ^b	4279 ± 4 ^b	5557 ± 5	6407 ± 12
1970 ± 4	4372 ± 4	5574 ± 6	6441 ± 9
2042 ± 3	4518 ± 4	5623 ± 4	6480 ± 5
2192 ± 3	4597 ± 4 ^b	5710 ± 11	6588 ± 5
2864 ± 3	4728 ± 8	5765 ± 3	6645 ± 5 ^b
2967 ± 3	4764 ± 7	5813 ± 4	6711 ± 5 ^b
3174 ± 3	4888 ± 4 ^b	5940 ± 5	6772 ± 6
3491 ± 4 ^{b,c}	5048 ± 4 ^b	6040 ± 4 ^b	6860 ± 13
3589 ± 3	5131 ± 5	6163 ± 6	6905 ± 8

^a (1976FO16): $E_d = 10$ MeV.

^b Unresolved.

^c Earlier results showed that $^{20}\text{F}^*(3.49)$ is strongly populated and that $^{20}\text{F}^*(3.53)$ is weakly populated, and has an angular distribution which is roughly symmetric about 90° . These results are consistent with $J^\pi = 1^+$ and 0^+ , respectively (1971FO14).

36. $^{21}\text{Ne}(\text{n}, \text{t})^{20}\text{F}$

$$Q_m = -14.8875$$

Not reported.

37. $^{22}\text{Ne}(\text{p}, ^3\text{He})^{20}\text{F}$

$$Q_m = -15.6513$$

At $E_p = 43.7$ to 45.0 MeV analog states have been studied in ^{20}F and ^{20}Ne [the latter via $^{22}\text{Ne}(\text{p}, \text{t})^{20}\text{Ne}$]. The experimental cross-section ratio, R , for the transitions to $^{20}\text{Ne}^*(10.28)$ and $^{20}\text{F}(0)$ [2^+ ; $T = 1$] is 2.00 ± 0.20 ; R for the transitions to $^{20}\text{Ne}^*(12.25 \pm 0.03)$ and $^{20}\text{F}^*(1.82)$ [5^+ ; $T = 1$] is 1.40 ± 0.15 (1969HA19). Angular distributions for the ^3He ions and the tritons corresponding to the first $T = 2$ states ($J^\pi = 0^+$) [$^{20}\text{Ne}^*(16.722 \pm 0.025)$ and $^{20}\text{F}^*(6.513 \pm 0.033)$]

have also been compared. There is indication also for the excitation of the 2^+ ; $T = 2$ states [at $E_x = 8.05$ MeV in ^{20}F and at 18.5 MeV in ^{20}Ne (estimated errors ± 0.1 MeV)] (1964CE05, 1969HA38).

$$38. \text{}^{22}\text{Ne}(d, \alpha)^{20}\text{F} \quad Q_m = 2.7019$$

Angular distributions have been obtained at $E_d = 10$ MeV to all ^{20}F states with $E_x < 4.4$ MeV: they are generally featureless. The cross sections for $^{20}\text{F}^*(3.68, 3.97)$ are consistent with their being the second 4^+ and the third 1^+ states in ^{20}F . A listing of all the states observed in this experiment is displayed in Table 20.17 (1976FO16). See also (1971MO40: $E_d = 2$ and 3 MeV; $\alpha_0 \rightarrow \alpha_7$).

$$39. \text{}^{23}\text{Na}(n, \alpha)^{20}\text{F} \quad Q_m = -3.866$$

See (1975SA1D) and (1972AJ02).

^{20}Ne
(Figs. 11 and 13)

GENERAL: (See also (1972AJ02).)

Shell model: (1970CR1A, 1971DE56, 1971RA1B, 1971ZO1A, 1972AB12, 1972AR1F, 1972AS13, 1972BO38, 1972BR1G, 1972JA24, 1972KA39, 1972KA67, 1972KH08, 1972KR1D, 1972KU1F, 1972LE13, 1972LE38, 1972MA07, 1972NI14, 1972RE03, 1972SA1B, 1972VO09, 1972WH04, 1973CO03, 1973DH1A, 1973EL04, 1973EN1C, 1973GI09, 1973HA05, 1973HE1F, 1973IC01, 1973IR01, 1973MA1K, 1973MC06, 1973MC1E, 1973ME1D, 1973PA05, 1973PU1A, 1973SM1C, 1973SV1A, 1973YA1A, 1974AY02, 1974KH03, 1974KR17, 1975BA81, 1975DR01, 1975GI03, 1975GR1M, 1975HA1U, 1975HA1V, 1975MP01, 1975MU13, 1975TA1H, 1976NG04, 1976PO01, 1976SC08, 1976SH07, 1977CA08, 1977FE07, 1977HA33, 1977MA09, 1977NE1D, 1977RO1V, 1977SH11).

Collective and deformed models: (1970CR1A, 1970FL1A, 1971RA1B, 1972BO38, 1972DE1F, 1972HA1Q, 1972HI17, 1972HO1D, 1972HO56, 1972JA24, 1972KU1F, 1972TE02, 1972VO09, 1973CA16, 1973CU03, 1973DE06, 1973EL04, 1973GO26, 1973HO1H, 1973HO40, 1973HO41, 1973IC01, 1973IR01, 1973KO1F, 1973KR1E, 1973LA35, 1973MC1E, 1973MI1F, 1973NG1B, 1973NO1B, 1973RE11, 1973SE15, 1973ST25, 1973SV1A, 1973VA1H, 1973YA1A, 1974AR04, 1974KR17, 1974MA11, 1974MP01, 1974NO14, 1974RI1B, 1974SC34, 1974TA19, 1975AB04, 1975BU1L, 1975DR01, 1975GR1M, 1975HA1U, 1975HE1L, 1975HE10, 1975HO1H, 1975MA26, 1975NE04, 1975TA1H, 1976BA11, 1976CH23, 1976JA1F, 1976KE03, 1976MA36, 1976NO10, 1976PA02, 1976SH07, 1976ST09, 1976UL1D, 1977CA08, 1977FE07, 1977HO1F, 1977KA2F, 1977MA09, 1977PA07, 1977RO13, 1977ST21, 1977TH03).

Cluster and α -particle models: (1971AB1B, 1971AR1V, 1971RI1D, 1971SA1A, 1971WO1C, 1972AR12, 1972AR1F, 1972BA59, 1972FR1B, 1972GO16, 1972GR42, 1972HI17, 1972HO56, 1972IK1A, 1972KA39, 1972KA67, 1972LA08, 1972LI1E, 1972NE1B, 1972VO09, 1973AR1C, 1973CH1E, 1973CU03, 1973HO1H, 1973HO40, 1973IC01, 1973KR1E, 1973LA35, 1973MA50, 1973SU1D, 1973VA21, 1973YA1A, 1974AB05, 1974GO12, 1974MO01, 1974YA11, 1975AR1J, 1975BU03, 1975BU1L, 1975HA1V, 1975HE1L, 1975HE10, 1975IK1A, 1975MA26, 1975NE04, 1976BA11, 1976BA1N, 1976FU1K, 1976KI18, 1976SA1F, 1977FL1E, 1977FU1P, 1977HE10, 1977HO1E, 1977HO1F, 1977KA2F, 1977RO13, 1977ST21, 1977TO1L).

Astrophysical questions: (1972CL1A, 1972KO1A, 1973AR1E, 1973AR1H, 1973AU1D, 1973CA1B, 1973CO1J, 1973HO1G, 1973RA1D, 1974AR1G, 1974AR1H, 1974PA1E, 1975BA2D, 1975EN1A, 1975IB1A, 1975LA1E, 1975MA1R, 1975PE1E, 1975RA1M, 1975RO08, 1975TA1G, 1975TR1A, 1975WO1D, 1976GI1C, 1976NO1C, 1976RO1J, 1976RU1C, 1976SI1D, 1977AR1H, 1977BU1Q, 1977CA1N, 1977DW1B, 1977KI1M, 1977PA1J, 1977VO1A).

Giant resonance (See also reaction 46.): (1973HA1Q, 1974HA1C, 1975AB04, 1975GR1M, 1976BE1P, 1976SC08, 1977AB08, 1977KN02, 1977SC08).

Electromagnetic transitions: (1971DE56, 1972AB12, 1972AS13, 1972BE1E, 1972BO38, 1972DE1F,

Table 20.18: Energy levels of ^{20}Ne ^a

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$0^+; 1$	0_1^+		stable	2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 14, 19, 20, 22, 23, 25, 26, 27, 29, 30, 31, 32, 33, 34, 40, 41, 42, 43, 44, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 59, 62, 64, 65, 66, 67, 68
1.6337 ± 0.3	$2^+; 0$	0_1^+	$\tau_m = 1.05 \pm 0.06$ psec $ g = 0.54 \pm 0.04$	γ	2, 3, 7, 8, 9, 10, 12, 14, 19, 20, 22, 23, 25, 27, 29, 31, 32, 33, 34, 40, 41, 42, 43, 45, 47, 49, 50, 52, 53, 56, 57, 58, 59, 62, 64, 65, 66, 67, 68
4.2477 ± 1.1	$4^+; 0$	0_1^+	$\tau_m = 93 \pm 9$ fsec	γ	2, 3, 7, 8, 9, 14, 19, 20, 22, 23, 24, 25, 27, 29, 31, 32, 33, 40, 41, 42, 43, 49, 50, 52, 53, 56, 59, 62, 64, 65, 66, 67, 68
4.9679 ± 0.7	$2^-; 0$	2^-	4.8 ± 0.5 psec	γ	2, 3, 7, 8, 9, 14, 19, 22, 25, 31, 34, 40, 41, 42, 43, 45, 49, 59, 62, 64
5.6214 ± 1.7	$3^-; 0$	2^-	200 ± 50 fsec	γ, α	2, 3, 7, 8, 14, 19, 29, 40, 41, 42, 43, 49, 60, 62, 64, 66, 67
5.784 ± 2	$1^-; 0$	0^-		γ, α	2, 3, 7, 8, 14, 19, 20, 22, 29, 31, 32, 41, 42, 43, 49, 60, 62, 66, 67
6.724 ± 5	$0^+; 0$	0_2^+	$\Gamma = 15 \pm 7$ keV	γ, α	3, 7, 8, 14, 15, 19, 31, 40, 41, 43, 47, 49, 63
7.004 ± 3.6	$4^-; 0$	2^-	$\tau_m = 440 \pm 90$ fsec	γ	2, 3, 7, 8, 14, 19, 25, 41, 42, 49, 63
7.168 ± 5	$3^-; 0$	0^-	$\Gamma = 8$ keV	α	2, 3, 7, 8, 15, 19, 20, 25, 31, 32, 40, 41, 42, 43, 49
7.191 ± 3	$0^+; 0$	0_3^+	4	γ, α	3, 7, 14, 15, 22, 40, 42, 47, 49
7.4214 ± 1.0	$2^+; 0$	0_2^+	8	γ, α	2, 3, 5, 7, 14, 15, 19, 40, 41, 43, 47, 49, 58, 60
7.8290 ± 2.0	$2^+; 0$	0_3^+	2.4	γ, α	2, 3, 7, 14, 15, 31, 40, 41, 47, 49, 58, 60
≈ 8.3	$0^+; 0$	0_4^+	> 800	α	3, 15, 41
8.4486 ± 2.3	$5^-; 0$	2^-	0.013 ± 0.004	γ, α	2, 3, 6, 7, 14, 15, 19, 25, 41, 49
8.694 ± 6	$1^-; 0$	(1_1^-)	2.5		3, 7, 15, 40, 41, 49

Table 20.18: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
8.7767 ± 2.3	$6^+; 0$	0_1^+	0.11 ± 0.02	γ, α	2, 3, 6, 7, 9, 14, 15, 19, 20, 22, 23, 24, 25, 31, 41, 49
≈ 8.8	$2^+; 0$	0_4^+	> 800	α	15, 41, 58
8.848 ± 5	$1^-; 0$	(1_1^-)	19	α	7, 15, 32, 41, 60
9.030 ± 5	$4^+; 0$	0_3^+	3.2	γ, α	2, 3, 6, 7, 14, 15, 31, 41, 49, 60
9.115 ± 4	$3^-; 0$		3.2	α	2, 7, 15, 40, 41, 49
9.318 ± 6					5, 7, 41, 49, 60
9.508 ± 12	$2^+; 0$		29 ± 15	γ, α	3, 7, 14, 15, 40, 41, 49, 58
9.873 ± 4	$3^+; 0$			γ	7, 41, 58
9.92 ± 20	$(1^+); 0$		$\tau_m < 35$ fsec	γ	3, 31, 41, 60
9.99 ± 20	$4^+; 0$	0_2^+	$\Gamma = 155 \pm 30$	γ, α	2, 7, 14, 15, 41
10.261 ± 4	$5^-; 0$	0^-	145 ± 40	α	2, 3, 7, 15, 19, 20, 22, 23, 25, 32
10.2724 ± 2.0	$2^+; 1$		0.4 ± 0.2	γ, α	2, 3, 14, 15, 31, 41, 58, 60, 62
10.403 ± 5	$3^-; 0$		80	α	7, 15, 32, 41, 60
10.548 ± 5	$4^+; 0$		16	α	3, 7, 15, 19, 41
10.583 ± 6	$2^+; 0$		24	α	15, 41, 58
10.609 ± 6	$6^-; 0$	2^-	$\tau_m = 23 \pm 7$ fsec	γ	2, 3, 6, 7
10.694 ± 6	$4^-, 3^+; 0$			γ	6, 7
10.79 ± 100	$4^+; 0$	0_4^+	$\Gamma = 350$	α	15, 24
10.838	3^-		45	α	7, 15, 60
10.840 ± 5	$2^+; 0$		13	α	7, 15, 41, 58
10.89 ± 10	$3^+; 1$		$\tau_m < 30$ fsec	γ	6, 7, 41, 58, 60
10.97	$0^+; 0$		$\Gamma = 580$	α	15
11.015 ± 6	$4^+; 0$		24	α	6, 7, 10
11.073 ± 8	$4^+; 1$		≤ 0.5	γ, α	14, 15, 41, 60
11.23 ± 30	$1^-; 0$		170	α	15
11.23 ± 10	$1^+; 1$			γ	34, 47, 58, 60
11.256 ± 8	$1^-; 1$		≤ 0.3	γ, α	14, 15
11.322 ± 7	$2^+; 0$		40 ± 10	α	15, 41, 58
11.528 ± 6	$3^+, 4^-; 0$		$\tau_m \leq 30$ fsec	γ	7, 31
11.552 ± 8	$(2^+, 0^+)$		$\Gamma = 1.0 \pm 0.5$ keV	γ, α	14, 15, 31, 40, 41
11.555 ± 6	$1^+, 2^-, 3^+$			γ	7, 31
11.601 ± 10	$2^-; 1$				31, 60
11.656	$(3^+); 0$			γ	6, 7
11.866 ± 9	$2^+; 0$		60	α	7, 15, 41, 58
11.926 ± 5	$4^+; 0$		0.44 ± 0.15	$(\gamma), \alpha$	14, 15, 40

Table 20.18: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
11.949 \pm 5	8 ⁺ ; 0	0 ₁ ⁺	0.035 \pm 0.010	γ, α	6, 7, 8, 9, 14, 15, 19, 20, 22, 23, 41
11.962 \pm 8	1 ⁻ ; 0		30 \pm 5	α	15
12.100 \pm 10	2 ⁻ ; 1				60
12.134 \pm 10	6 ⁺ ; 0	0 ₃ ⁺	0.13 \pm 0.07	(γ), α	6, 7, 8, 14, 19
12.215 \pm 5	2 ⁺ ; 1		< 0.1	γ, α	14, 34, 41, 62
12.24 \pm 30	4 ⁺ ; 0		148 \pm 20	α	15
12.254 \pm 6	3 ⁻ , 2 ⁺		\approx 5	γ, α	14, 40
(12.35 \pm 100)	(2 ⁺)		\approx 500	α	15
12.39 \pm 10	3 ⁻ ; (1)		33 \pm 4	γ, α	6, 7, 14, 15, 40, 41
12.412 \pm 5	(0 ⁺); 0		\leq 8	α	7, 15, 31, 41
12.49 \pm 30				γ, α	14, 41
12.591 \pm 10	6 ⁺ ; 0	0 ₄ ⁺	88 \pm 10	α	6, 7, 15, 19, 20, 22
12.683 \pm 15	5 ⁻ ; 0		97	α	15
12.730 \pm 10	4 ⁺ ; 0		100	α	6, 7, 15
12.83 \pm 30			55	α	15, 31, 41
12.919 \pm 10				γ	7
13.010 \pm 10	(4 ⁺ ; 0)		60	α	7, 15
13.049 \pm 10	(4 ⁺ ; 0)		70	α	6, 7, 15
13.060 \pm 3.5	2 ⁻		1.0	p, α	37, 40, 41
13.1680 \pm 0.6	1 ⁺ ; (1)		2.3 \pm 0.2	γ, p, α	34, 35, 37, 40, 41
13.190 \pm 10	(4 ⁺ ; 0)		60	α	6, 7, 15
13.225	1 ⁻		95	p, α	37
13.225	0 ⁺		95	p, α	37
13.3038 \pm 0.7	1 ⁺		0.9 \pm 0.1	γ, p, α	7, 34, 35, 37, 41
13.334 \pm 6	7 ⁻ ; 0	2 ⁻	(8 \pm 3) \times 10 ⁻⁴	α	6, 7, 8, 15, 31
13.343 \pm 6	4 ⁺ ; 0		20 \pm 5	α	15, 31
13.412 \pm 1	2 ⁻		29 \pm 3	γ, p, α	15, 34, 35
(13.42 \pm 140)	(4 ⁺ ; 0)		110	α	15
13.462 \pm 20	1 ⁻		190	p, α	37
13.482 \pm 1	1 ⁺ ; 1		5.7 \pm 0.7	γ, p, α	31, 34, 35, 40
13.519	(1 ⁻)		33	p, α	37
13.569 \pm 15	2 ⁺		63	p, α	7, 31, 35, 37
13.583	2 ⁺		\approx 10	p, α	37
13.64 \pm 15	0 ⁺ ; 1		22	p, α	7, 31, 35, 37, 40
(13.66)	(1 ⁻)		110	p, α	37
13.6729 \pm 0.7	2 ⁻		4.5 \pm 0.2	γ, p, α	6, 7, 34, 35, 37
13.7 \pm 400	(3, 7) ⁻		320	α	15

Table 20.18: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
(13.73)	(0 ⁺)		≈ 170	p, α	37
13.733 \pm 1.4	1 ⁺		7.7 \pm 0.5	γ , p, α	34, 35, 37
13.845 \pm 15	(1 ⁻)		≈ 190	p, α	7, 37
13.886 \pm 15	2 ⁻		0.8	γ , p	6, 7, 8, 31, 34, 35, 40
13.904	2 ⁺		47	p, α	37
13.904 \pm 20	6 ⁺		≈ 100	α	15, 19, 20
13.946	0 ⁺		≈ 70	p, α	37
14.017	1 ⁻		≈ 70	p, α	37
14.03	2 ⁺		≈ 140	p, α	37
14.124 \pm 1.2	2 ⁻		4.7 \pm 0.7	γ , p, α	34, 35, 37
14.144 \pm 15	2 ⁺		50	p, α	7, 15, 37
14.148 \pm 1.2	2 ⁻		11.8 \pm 1.0	γ , p, α	34, 35, 37
14.195	1 ⁺		14 \pm 1	γ , p	34, 35
14.307 \pm 10	6 ⁺		< 100		6, 7, 19, 20, 24
14.44 \pm 20	0 ⁺		110	p, α	6, 7, 35, 37
14.453 \pm 1.8			33 \pm 3	p, α	37
14.6 \pm 300	(4 ⁺)		240	α	15
14.60 \pm 20	1 ⁻		140	p, α	7, 35, 37
14.695 \pm 2.6			36 \pm 10	p, α	37
14.772 \pm 3.0			110 \pm 20	p, α	37
14.812 \pm 15	(2 ⁺ , 4 ⁺)		≈ 100	p, α	6, 7, 15, 37
15.034 \pm 15	(2 ⁺)		≈ 100	p, α	7, 15, 37
15.16 \pm 15	6 ⁺			α	6, 7
15.23			28	p, α	37
15.27	(1 ⁻)		285	p, α	37
15.30	(0 ⁺)		285	p, α	37
15.336 \pm 15	7 ⁻	0 ⁻	380 \pm 60	α	6, 7, 15, 19, 20, 22, 32
15.39			85	p, α	37
15.47			55	p, α	37
b					
15.50 \pm 20	(8 ⁻)	(2 ⁻)			7, 32
15.70 \pm 15	(6 ⁺)			α	6, 7, 15
15.879 \pm 15	(8 ⁺)		< 250	α	6, 7, 19
(15.97)	(6 ⁺)			α	15
16.01 \pm 25	(2 ⁺ ; 1)		100	p, α	31, 37
16.139 \pm 15			38	p, α	6, 7, 15, 37
16.25				α	6, 15
16.326 \pm 15	4 ⁺		43	p, α	15, 37

Table 20.18: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
16.434 \pm 15	(0, 2, 4) ⁺		34	p, α	15, 37
16.510 \pm 10	(2, 4, 6) ⁺		23	α	6, 7, 15
16.52	7 ⁻			α	7
16.63 \pm 20	7 ⁻		190 \pm 40	α	7, 15, 19, 20, 22
16.719 \pm 15	(1, 3, 7) ⁻		10	α	15
16.730 \pm 3	0 ⁺ ; 2		2.0 \pm 0.5	γ , p, α	6, 31, 34, 35, 37, 62
16.8	7 ⁻			α	7
16.854 \pm 15	5 ⁻		10	α	15
16.98			100	p, α	37
17.162 \pm 15	5 ⁻ , (7 ⁻)		37	α	15
17.30 \pm 20	8 ⁺		220 \pm 40	α	15, 19, 20, 22
17.38 \pm 15	9 ⁻	2 ⁻	< 10	α	6, 7, 8, 15, 23
17.542 \pm 15	6 ⁺		136	α	15
17.55 \pm 10	(2 ⁺ ; 1)		19	n, p, α	31, 36, 37
17.752 \pm 15	4 ⁺ , (0 ⁺)		50	p, α	15, 37
17.91 \pm 20	(0 ⁺)			n, p	31, 36
18.002 \pm 15	7 ⁻		< 10	α	15
18.022 \pm 15	(2 ⁺ , 5 ⁻ , 6 ⁺)		45	α	15
18.113 \pm 15	7 ⁻		33	α	6, 7, 8, 15
18.32 \pm 20	(6 ⁺)		240	α	6, 15
18.427 \pm 7	2 ⁺ ; 2		9.5 \pm 3	γ , n, p, α	34, 35, 36, 37, 47
18.7 \pm 100	(6 ⁺ , 7 ⁻)		600	α	15, 19
19.16 \pm 250	(6 ⁺)		200	α	8, 15
19.40 \pm 350	6 ⁺		280	α	15
19.4 \pm 100	7 ⁻		400		19
19.84 \pm 350	6 ⁺		280	α	15
20.0 \pm 100	7 ⁻		300	α	15, 19
20.4 \pm 180	6 ⁺		360	α	7, 15
20.4 \pm 100	7 ⁻		200	α	7, 15
20.67 \pm 40	9 ⁻		120	α	7, 15, 20
20.8 \pm 100	7 ⁻ , (6 ⁺)			α	19
21.0 \pm 100	7 ⁻		200	α	15
21.08 \pm 30	9 ⁻		100 \pm 50	α	15, 20, 22, 23
21.3 \pm 100	7 ⁻ , 8 ⁺		300	α	15, 19
21.8 \pm 100	7 ⁻ , 8 ⁺		300	α	15, 19
22.3 \pm 100	7 ⁻ , 8 ⁺		500	α	15, 19
22.7 \pm 250	9 ⁻		500	α	15
22.87 \pm 40	9 ⁻		225 \pm 40	α	15, 20, 22

Table 20.18: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
23.70 ± 30	$9^-, (8^+)$		≤ 200	α	15, 19, 20, 24
24.21 ± 25	8^+		≈ 500	α	15, 20
25.10 ± 50	8^+		≤ 200	α	15, 20
25.67 ± 50			≈ 500	α	15, 20
27.1 ± 100	(9^-)		700	α	15, 19
28	8^+		1600	α	15, 28
28.1 ± 100	(10^+)		700	α	15, 19

^a See also Tables 20.19 and 20.20.

^b For other states with $E_x > 15.5$ MeV see Tables 20.30, 20.31, 20.32, 20.33 and reactions 46 and 47. It is clear that there are many states with low angular momentum and with unnatural parity which have not been located at high E_x .

1972HA1Q, 1972HI17, 1972LE38, 1972LO1D, 1972MA20, 1972NA05, 1973CA16, 1973CU03, 1973EL04, 1973GO26, 1973HA1N, 1973HA1Q, 1973HA1V, 1973HO41, 1973LE1E, 1973PU1A, 1973ST25, 1973UN01, 1974AB05, 1974HA1G, 1974HA1C, 1974KH03, 1974MA11, 1974MC1F, 1974NO14, 1974SC34, 1975BU03, 1975BU1L, 1975MA26, 1975NA21, 1975NA20, 1975TA1H, 1976BA11, 1976FU1K, 1976SC08, 1976VO1C, 1977HO1F, 1977KN02, 1977MA09, 1977NE1D, 1977RO1V, 1977SC08).

Special levels: (1972AR12, 1972BE1E, 1972FO29, 1972HA1R, 1972HI17, 1972JA24, 1972KH08, 1972LE13, 1972MA07, 1972MA20, 1972NI14, 1972SA04, 1972TE02, 1973CH1E, 1973CU03, 1973DH1A, 1973EN1C, 1973HO40, 1973IR01, 1973JU1A, 1973KO42, 1973MA50, 1973MC06, 1973MI1F, 1973NO1B, 1973PU1A, 1973SU1D, 1973UN01, 1973YA1A, 1974AR04, 1974GO12, 1974GO37, 1974HA1G, 1974IT1A, 1974KH03, 1974KR17, 1974MA1K, 1974MO01, 1975AR1J, 1975BA81, 1975BU03, 1975BU1L, 1975HA1V, 1975MA10, 1975MA1W, 1975MI03, 1975NA21, 1975NE04, 1976BA11, 1976SA21, 1976SC08, 1977CA08, 1977FU1P, 1977HA33, 1977NE1D, 1977SC08, 1977SH11, 1977ST21).

Applied work: (1976CH1P).

Special reactions: (1972BO1H, 1973WI15, 1974HA61, 1975AR17, 1975AR14, 1975DR01, 1975HO1K, 1975KU1K, 1975RE08, 1975VO09, 1976CO1W, 1976EG02, 1976HI05, 1977CO14, 1977LE1T, 1977NA03, 1977RA1D, 1977TA07).

Muon and neutrino capture and reactions: (1972VA1F, 1973BE53, 1974EN10, 1974GR1M, 1974KA1Q, 1975DO10, 1975PF01).

Pion and kaon capture and reactions: (1971JH1A, 1971MO1P, 1972BU1G, 1972EC1A, 1974HA61, 1974HU14, 1974LI1D, 1974TA18, 1974UL02, 1975BU1K, 1975EL1A, 1975HU1D, 1975SC1N, 1975TA1C, 1976KA1M, 1976RO14, 1977BA2H, 1977MC1K, 1977WA1W).

Other topics: (1969FU1A, 1971RA1B, 1971ZO1A, 1972AB12, 1972BR1G, 1972EL1C, 1972FR1B, 1972GR1G, 1972GR1H, 1972JA24, 1972KA67, 1972KH08, 1972KR11, 1972KR1D, 1972LA08,

1972MU08, 1972NI14, 1972PA36, 1972PL1C, 1972RE03, 1972SA04, 1972SA1B, 1972SH24, 1972TE02, 1973BE35, 1973CA32, 1973CO03, 1973CU03, 1973EL04, 1973ER10, 1973FO1F, 1973GI09, 1973GO26, 1973GR11, 1973GR36, 1973HA05, 1973HA1W, 1973HE1F, 1973HO1H, 1973KE1C, 1973KO26, 1973KO42, 1973KR1B, 1973LA35, 1973MA1K, 1973MA48, 1973ME1G, 1973MI1F, 1973NG1A, 1973NG1B, 1973PA05, 1973RA1E, 1973SE17, 1973SE15, 1973SU1D, 1973SV1A, 1973UL01, 1973UN01, 1973VA01, 1973VA21, 1974AB05, 1974AR04, 1974AY02, 1974CO39, 1974GO12, 1974HA17, 1974MA1K, 1974MP01, 1974NG01, 1974RE03, 1974SC10, 1974SC11, 1974TI03, 1974TI04, 1974WO02, 1974ZU1A, 1975BA81, 1975GI03, 1975HE10, 1975MA10, 1975MA1W, 1975MI02, 1975MU13, 1975SO04, 1976VOID, 1976HA1V, 1976MA04, 1976SA21, 1976SM1D, 1976ST09, 1976UL1D, 1977FL1E, 1977HE10, 1977PA07, 1977SH11, 1977SH13).

Ground-state properties: (1971RU14, 1971SH26, 1972AB12, 1972AS13, 1972BR1G, 1972CA37, 1972GR1H, 1972GR42, 1972KR11, 1972KR1D, 1972LE38, 1972NI14, 1972OL02, 1972SC27, 1972YO1B, 1973AR1C, 1973CA16, 1973CU03, 1973ER10, 1973FO1F, 1973GO26, 1973GR36, 1973HO41, 1973KO26, 1973LE1E, 1973MA1K, 1973MC06, 1973ME1G, 1973NG1A, 1973PA05, 1973PU1A, 1973RE11, 1973SC1H, 1973SV1A, 1973UN01, 1973VA01, 1974AB05, 1974AR04, 1974CO39, 1974DE1E, 1974EN10, 1974KH03, 1974KR17, 1974MA11, 1974MC1F, 1974MP01, 1974RE03, 1974RI1B, 1974SC34, 1974SHYR, 1974TI03, 1975AL19, 1975BU03, 1975HO1H, 1975LE05, 1975MA26, 1975MI02, 1975MP01, 1976CH1T, 1976FU06, 1976KE03, 1976PO01, 1976ST09, 1977AN12, 1977AN21, 1977KN02, 1977MP01, 1977PA07, 1977SU04, 1977TH03).

$$Q_{1.63} = -0.20 \pm 0.05 \text{ b (1974OL01);}$$

$$B(E2) \uparrow [0 \rightarrow 1.63] = 0.032 \pm 0.003 e^2 \cdot \text{b}^2 \text{ (1977SC36);}$$

$$= 0.037 \pm 0.003 e^2 \cdot \text{b}^2 \text{ (1974OL01);}$$

$$= 0.028 \pm 0.004 e^2 \cdot \text{b}^2 \text{ (1973SI31);}$$

$$|M|^2 = 23.0 \pm 1.9 \text{ W.u. (1974OL01).}$$

1. (a) $^{10}\text{B}(^{10}\text{B}, ^{10}\text{B})^{10}\text{B}$

$$E_b = 31.1464$$

(b) $^{10}\text{B}(^{10}\text{B}, \alpha)^{16}\text{O}$

$$Q_m = 26.4155$$

The total reaction cross section has been measured for $E(^{10}\text{B}) = 3.7$ to 7.3 MeV (1976HI05), and the elastic excitation function has been studied for 8 to 30 MeV (1975DI08). Excitation functions for reactions (a) and (b) have been measured for $E_{c.m.} = 3$ to 10 MeV: large resonant structures are observed in most channels of reaction (b) (1978MA07): see (1978EN06). See also ^{10}B in (1979AJ01).

2. $^{10}\text{B}(^{14}\text{N}, \alpha)^{20}\text{Ne}$

$$Q_m = 19.5332$$

At $E(^{14}\text{N}) = 25$ and 35 MeV angular distributions of the α -particles to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62, 5.78, 7.00, 7.17, 7.42, 7.83, 8.45, 8.78, 9.03 + 9.12, 9.99, 10.26, 10.61)$ have been measured by (1974MA38, 1975LE23). The average behavior of the cross section is generally well described by a statistical mechanism but the reaction mechanism is not purely statistical (1975LE23). For excitation functions see (1972VO02, 1975LE23, 1977MA2V, 1977MA2W): see also (1978EN06). See also (1973ST1A) and (1975KL05; theor.).

3. $^{10}\text{B}(^{16}\text{O}, ^6\text{Li})^{20}\text{Ne} \quad Q_m = 0.270$

At $E(^{16}\text{O}) = 19.5, 24.3, 31.5, 35.9$ and 42 MeV angular distributions have been measured for the ^6Li ions corresponding to transitions to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62 + 5.78, 6.7 - 7.2)$. Hauser-Feshbach calculations are generally in good agreement with the data (1976LO03). Differential cross sections at 7° have been measured at $E(^{16}\text{O}) = 45.6$ MeV for many states of ^{20}Ne with $E_x < 10.62$ MeV (1974FO15). For fusion cross sections see (1977MO1X). See also (1974ST1L, 1978EN06).

4. $^{11}\text{B}(^{16}\text{O}, ^7\text{Li})^{20}\text{Ne} \quad Q_m = -3.934$

This reaction has been studied at $E(^{16}\text{O}) = 60$ MeV (1972SC17) and at $E(^{11}\text{B}) = 114$ MeV (1973ST24): a number of states of ^{20}Ne are populated but they are not excited strongly at either energy. See also (1973SC1J).

5. $^{12}\text{C}(^9\text{Be}, n)^{20}\text{Ne} \quad Q_m = 10.3196$

At $E(^9\text{Be}) = 16, 20$ and 24 MeV, angular distributions have been measured for $^{20}\text{Ne}^*(7.4, 9.3, 11.0, 12.2)$ (1977EL1E).

6. (a) $^{12}\text{C}(^{10}\text{B}, d)^{20}\text{Ne} \quad Q_m = 5.9589$
 (b) $^{12}\text{C}(^{11}\text{B}, t)^{20}\text{Ne} \quad Q_m = 0.7610$

At $E(^{12}\text{C}) = 45$ MeV the population of states of ^{20}Ne with $E_x = 8.46, 8.78, 9.03, 10.61, 10.67, 10.99, 11.01, 11.66, 11.94, 12.14, 12.39, 12.58, 12.73, 13.05, 13.17, 13.34 [7^-], 13.69, 13.91, 14.29, 14.36, 14.81, 15.17 [6^+], 15.38 [7^-], 15.71 [(7, 8)], 15.89 [(7)], 16.16, 16.22, 16.51 [(8)], 16.73, 17.39 [9^-], 18.18$ and 18.32 MeV is reported. [Values in brackets are J^π suggested on basis of Hauser-Feshbach calculations. The states in italics are well resolved: the authors indicate ± 20 keV for such states.] The relative intensities of the groups to $^{20}\text{Ne}^*(17.39, 15.38) [J^\pi = 9^-, 7^-]$ argue against the existence of a superband (1974KL05, 1976KL03). See also (1973MI1E, 1974ME1C, 1977HI01), (1973BR1C, 1973FO1A, 1973FO1E, 1974FO1J, 1977KL1G), (1975KL05, 1975KL08; theor.) and ^{22}Na in (1978EN06). For reaction (b) see (1977HI01).

Table 20.19: Radiative decays in ^{20}Ne ^a

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)	$ M ^2$ (W.u.)	Refs.
1.63	$2^+; 0$	0	100		17.8 ± 2.5 (E2)	(1971HA26)
4.25	$4^+; 0$	1.63	≈ 100		21.9 ± 2.1 (E2)	(1971HA26)
4.97	$2^-; 0$	0	0.6 ± 0.2		$(2.5 \pm 0.9) \times 10^{-3}$ (M2)	(1967BR22, 1971HA26)
		1.63	99		$(7.0 \pm 1.0) \times 10^{-6}$ (E1)	(1971HA26)
5.62	$3^-; 0$				$(1.7 \pm 0.7) \times 10^{-2}$ (M2) ^b	(1971HA26)
					6 ± 5 (E3) ^c	(1971HA26)
		0	7.6 ± 1.0		10.6 ± 2.7 (E3)	(1964BR18, 1971HA26)
		1.63	87.6 ± 1.0		$(6.6 \pm 1.5) \times 10^{-6}$ (E1)	(1964BR18, 1971HA26)
5.78	$1^-; 0$	4.97	4.8 ± 1.6		35 ± 9 (E2)	(1971HA26)
		0	18 ± 5		7.2×10^{-6} (E1)	(1965VA14)
6.72	$0^+; 0$	1.63	82 ± 5	4.0	9.2×10^{-5} (E1)	(1965VA14)
		0			7.4 ± 2.0 fm ² ^o	(1972MI06)
7.00	$4^-; 0$	1.63	100	33	3.8 (E2)	(1965VA14)
		1.63	0.5 ± 0.2			(1967BR22)
7.19	$0^+; 0$	4.25	63.5		$(1.7 \pm 0.4) \times 10^{-5}$ (E1)	(1967BR22, 1971HA26)
		4.97	11		10.4 ± 1.8 (E2)	(1967BR22, 1971HA26)
		5.62	25		24 ± 5 (E2)	(1967BR22, 1971HA26)
		0		$\Gamma_\pi = 3.9 \times 10^{-2}$	6.9 ± 1.4 fm ² ^o	(1972MI06, 1973SI31)
7.42	$2^+; 0$	1.63	100	4.35 ± 0.75	0.37 ± 0.07 (E2)	(1972AL32)
		0	$\leq 9.4 \pm 1.4$	$\leq 3.0 \pm 0.6$	$\leq 0.05 \pm 0.01$ (E2) ^p	(1972AL32)
7.83	$2^+; 0$	1.63	$\geq 90.6 \pm 1.4$ ^d	29 ± 4	1.65 ± 0.27 (E2)	(1972AL32)
					$(1.0 \pm 0.3) \times 10^{-4}$ (M1)	(1972AL32)
		4.25	≤ 7.6		< 2.6 (E2)	(1972AL32)
		0	83 ± 1	57 ± 7	0.73 ± 0.09 (E2) ^p	(1972AL32)

Table 20.19: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)	$ M ^2$ (W.u.)	Refs.
		1.63	17 ± 1	11.7 ± 1.6	0.48 ± 0.07 (E2)	(1972AL32)
		4.25	< 3	< 2	< 1.3 (E2)	(1972AL32)
8.45	$5^-; 0$	5.62	100	13 ± 3	26 ± 6 (E2)	(1971RO33)
8.78	$6^+; 0$	4.25	100	100 ± 15	20.4 ± 2.4 (E2)	(1971DI08, 1971RO13)
9.03	$4^+; 0$	1.63	100	340 ± 42	5.8 ± 0.7 (E2)	(1972AL32)
		4.25	< 2	< 6.8	< 1 (E2)	(1972AL32)
9.51	$2^+; 0$	0		$\lesssim 60$	$\lesssim 0.3$ (E2)	(1964PE05)
		1.63	(100)	260 ± 100	3.2 (E2)	(1964PE05)
					0.03 (M1)	(1964PE05)
9.87	$3^+; 0$	0	< 0.5			(1977MA07)
		1.63	78	e		(1977MA07)
		4.25	12 ± 3			(1977MA07)
		4.97	≤ 5			(1977MA07)
		5.63	≈ 7			(1977MA07)
		7.42	≈ 3			(1977MA07)
9.92	$(1^+); 0$	1.63	78 ± 5			(1976FI10)
		4.97	22 ± 5			(1976FI10)
9.99	$4^+; 0$	0		$\lesssim 70$		(1964PE05)
		1.63	(100)	900 ± 400	6.9 (E2)	(1964PE05)
10.27 ^f	$2^+; 1$	0	0.65 ± 0.14	29 ± 8	0.10 ± 0.03 (E2)	(1977FI08)
		1.63	88.9 ± 0.5	4080 ± 440	0.31 ± 0.03 (M1)	(1977FI08)
		4.97	1.3 ± 0.1	60 ± 8	$(8.3 \pm 1.0) \times 10^{-4}$ (E1)	(1977FI08)
		5.62	2.1 ± 0.2	97 ± 14	$(2.0 \pm 0.3) \times 10^{-3}$ (E1)	(1977FI08)
		7.42	6.9 ± 0.4	310 ± 40	0.65 ± 0.008 (M1)	(1977FI08)

Table 20.19: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)	$ M ^2$ (W.u.)	Refs.
10.61 ^g	$6^-; 0$	7.83	0.22 ± 0.06	8 ± 2	0.027 ± 0.006 (M1)	(1977FI08)
		7.00	95.5 ± 1.2		17_{-4}^{+7} (E2)	(1971HA26)
		8.45	4.5 ± 1.2		10 ± 4 (E2)	(1971HA26)
10.69	$4^-, 3^+; 0$	4.25	25 ± 4			(1976FI10)
		4.97	75 ± 4			(1976FI10)
10.89	$3^+; 1$	1.63	26 ± 3			(1971HA26, 1976FI10)
		4.25	74 ± 3	h		(1971HA26, 1976FI10)
11.07	$4^+; 1$	1.63	≤ 2	$\lesssim 80$	$\lesssim 0.4$ (E2)	(1964PE05)
		4.25	(100)	4800 ± 500	0.72 (M1), 7.3 (E2)	(1964PE05)
				3400 ± 300	0.50 (M1)	(1978ST08)
11.26 ^f	$1^-; 1$	7.00		$\lesssim 1400$	$\lesssim 0.04$	(1964PE05)
		0	≈ 70	190 ± 20	2.8×10^{-4} (E1)	(1978ST08)
		1.63	≈ 30			(1978ST08)
11.53 ⁱ	$3^+, 4^-; 0$	4.25	30 ± 3 ⁿ			(1971HA26, 1976FI10)
		4.97	70 ± 3 ⁿ			(1971HA26, 1976FI10)
		7.00	j			
11.55	$(2^+, 0^+)$	1.63	≈ 70	80 ± 10 ⁿ	3.9×10^{-3} (M1)	(1978ST08)
		4.25	≈ 30			(1978ST08)
11.56 ⁱ	$1^+, 2^-, 3^+; 0$	1.63	j,n			(1976FI10)
		7.00	j,n			(1976FI10)
11.66	(3^+)	1.63	14 ± 3			(1976FI10)
		4.25	86 ± 3			(1976FI10)
11.95	$8^+; 0$	8.78	100	12 ± 3	7.5 ± 2.5 (E2)	(1972AL05)
12.22	$2^+; 1$	1.63	100	k		(1964PE05, 1977MA07)

Table 20.19: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)	$ M ^2$ (W.u.)	Refs.	
12.39	$3^-; (1)$	0	≈ 1	280 ± 20	1.1×10^{-3} (E1)	(1978ST08)	
		1.63	≈ 29			(1978ST08)	
		4.25	≈ 70			(1978ST08)	
12.49	$1^+; 1$	1.63	100			(1978ST08)	
13.48		1.63	95			(1961GO21)	
		4.97	5			(1961GO21)	
13.89		1.63	20			(1961GO21)	
		4.97	80			(1961GO21)	
16.73	$0^+; 2$	1.63		≈ 5000 ¹	< 1.3 (E2) ¹	(1976MA01)	
		5.78			$< 4.2 \times 10^{-3}$ (E1) ¹	(1976MA01)	
		11.23	(100)			(1967KU06)	
18.43	$2^+; 2$	0		m	$< 1.4 \times 10^{-3}$ (E2)	(1976MA01)	
		1.63		m	$< 5.4 \times 10^{-4}$ (M1)	(1976MA01)	
		4.25			$< 2.9 \times 10^{-2}$ (E2)	(1976MA01)	
		4.97			$< 5.1 \times 10^{-5}$ (E1)	(1976MA01)	
		5.62			$< 6.6 \times 10^{-5}$ (E1)	(1976MA01)	
		5.78			$< 5.8 \times 10^{-5}$ (E1)	(1976MA01)	
		7.19			$< 9.9 \times 10^{-5}$ (E1)	(1976MA01)	
		7.42			$< 3.7 \times 10^{-3}$ (M1)	(1976MA01)	
		12.22	(100)		≈ 300		(1972KU24)

- ^a See also Table 20.17 in (1972AJ02) and Tables 20.22 and 20.26 here.
- ^b From $\delta(M2/E1) = 0.076 \pm 0.011$ (1967BR22).
- ^c From $\delta(E3/E1) = 0.043 \pm 0.016$ (1967BR22).
- ^d $\delta(E2/M1) = -8.36^{+1.0}_{-1.5}$.
- ^e $\Gamma_\gamma(\text{total})/\Gamma = 0.82 \pm 0.27$. See also (1976FI10).
- ^f See also (1964PE05, 1976IN05).
- ^g See also (1976FI10).
- ^h $\Gamma_\gamma(\text{total})/\Gamma < 0.3$ (1977MA07).
- ⁱ See also (1977MA07).
- ^j See discussion in (1976FI10).
- ^k $\Gamma_\gamma(\text{total})/\Gamma \geq 0.25$; upper limits for transitions to $^{20}\text{Ne}^*(0, 4.25, 4.97)$ are 1.5, 3 and 4% (1977MA07). See also (1978ST08).
- ^l See, however, footnote ^a in Table 2 of (1976MA01).
- ^m See also (1972KU24).
- ⁿ If $J = 2$.
- ^o Monopole matrix element.
- ^p See also (1972MI06).

7. $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$

$$Q_m = 4.6181$$

Particle group and γ -ray energy measurements have been made for many states of ^{20}Ne : see Table 20.21. Angular correlation and γ -ray branching measurements [see Table 20.19] lead to the J^π assignments shown in Table 20.21, which also show level assignments to rotational bands. Angular distributions have been reported at $E(^{12}\text{C}) = 5.6$ to 28.3 MeV [see (1972AJ02)] and at $E(^{12}\text{C}) = 4.9$ to 9.8 MeV (1973MA09: $\alpha_1 \rightarrow \alpha_3, \alpha_{4+5}, \alpha_6, \alpha_7$), 8.5 to 12.5 MeV (1977GA04, 1977GA05: α_0, α_3), 13 to 22 MeV (1976ER03, 1977ER1F: $\alpha_0 \rightarrow \alpha_3, \alpha_6$), 13 to 24 MeV (1977VO05: $\alpha_0 \rightarrow \alpha_{11}$), 15 MeV (1976BA53: α to ^{20}Ne states with $E_x < 8$ MeV), 36.6, 36.8 and 37.4 MeV (1976FO15: α_0), 37 MeV (1976EB01: α_0), 37 MeV (1975ME04: see Table 20.21), 37, 41, 45, 48 and 51 MeV (1975GR21: α to $^{20}\text{Ne}^*(7.42, 7.83, 9.04)$), 38 MeV (1976FO09: α_0, α_6), 45 MeV (1975GR21: α to many states of ^{20}Ne with $E_x < 13.4$ MeV) and 50 MeV (1977PA19: α -particles to $^{20}\text{Ne}^*(0, 1.63, 4.25, 5.62, 5.78, 7.42, 7.83, 8.45)$). Both compound nucleus and direct reaction mechanisms appear to contribute to this reaction in the range $E(^{12}\text{C}) = 36$ to 51 MeV. Strong fluctuations in the yield require the presence of narrow compound nuclear levels while the selective population of particular states indicates the presence of a strong direct component. Hauser-Feshbach theory predicts lower cross sections for low- J states than are experimentally observed (1975GR21).

For lifetime measurements see Table 20.20. (1975HO15) have determined $|g| = 0.54 \pm 0.04$ for $^{20}\text{Ne}^*(1.63)$.

Studies of yields of particular α -groups are also reported for $E(^{12}\text{C}) = 8$ to 16 MeV (1977GA04, 1977GA05: α_0, α_3), 11 to 13 MeV (1973VO01, 1974VO09: α to $^{20}\text{Ne}^*(6.72, 7.2)$), 11.6 to 16 MeV (1977ER1F: α_1, α_6), 13 to 24 MeV (1977VO05: $\alpha_0 \rightarrow \alpha_6, \alpha_{11}$) and 25 to 27.1 MeV (1972CO30: $\alpha_0 \rightarrow \alpha_3, \alpha_{11}$); only the yields of $\alpha_1, \alpha_2, \alpha_6$ and α_{11} show strong fluctuations above $E(^{12}\text{C}) = 20$ MeV.

The yields of α -particle groups corresponding to ^{20}Ne states with $E_x < 17.5$ MeV have been studied in the range $E(^{12}\text{C}) = 32$ to 51 MeV by (1974FO20, 1974GR27, 1974SH16, 1975GR21, 1977FO01, 1977PA19). Many strong structures are observed which are interpreted in terms of states in the compound nucleus, ^{24}Mg : see (1978EN06). See also (1976BA22, 1976BA53, 1976EB01, 1976ER03, 1976FO15, 1976FO09, 1977ER1E, 1977HI04, 1977KE02, 1977MA1M, 1977VA1G).

See also (1972FI1B, 1973CO1K, 1973FE03, 1974KL1B, 1976NA11, 1977ER1E), (1973BR1C, 1973FO1E, 1973FO1A, 1973GO1M, 1973SC1B, 1973ST1A, 1974FO25, 1974FO1J), (1973AR1F, 1973MA09, 1973TR1B, 1975FO19, 1977DA07; astrophys. considerations) and (1973SC1K, 1976DA1P, 1977DE2A, 1977MA2X; theor.).

8. $^{12}\text{C}(^{14}\text{N}, ^6\text{Li})^{20}\text{Ne}$

$$Q_m = -4.181$$

Angular distributions of the ^6Li ions to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62 + 5.78, 6.7 - 7.2)$ have been measured at $E(^{14}\text{N}) = 30$ and 36 MeV (1976LO03), 52 and 60 MeV (1971MA23), 52 and 78 MeV (1974HA11: additionally $^{20}\text{Ne}^*(8.45 - 9.03)$), and at $E(^{14}\text{N}) = 76.1$ MeV and

Table 20.20: Lifetime measurements of some ^{20}Ne states ^a

$^{20}\text{Ne}^*$ (MeV)	τ_m	Refs.
1.63	1.04 ± 0.07 psec	(1974OL01)
	1.15 ± 0.20 psec	(1971HA26)
	0.8 ± 0.2 psec	(1975HO15)
4.25	1.05 ± 0.06 psec	“best” value ^b
	134 ± 12 fsec	(1965EV03)
	150 ± 25 fsec	(1969AN08)
4.97	93 ± 9 fsec	(1971HA26)
	93 ± 9 fsec	“best” value
	$4.8^{+1.7}_{-1.1}$ psec	(1965EV03)
5.62	4.8 ± 0.5 psec	(1971HA26)
	4.8 ± 0.5 psec	“best” value
	350 ± 75 fsec	(1965EV03)
7.00	200 ± 50 fsec	(1971HA26)
	200 ± 50 fsec	“best” value
	405^{+110}_{-90} fsec	(1962BR35)
9.92	470 ± 90 fsec	(1971HA26)
	440 ± 90 fsec	mean value
10.61	< 35 fsec	(1971HA26)
10.89	23 ± 7 fsec	(1971HA26)
11.53	< 30 fsec	(1971HA26)
	< 30 fsec	(1971HA26)

^a See also Table 20.18 in (1972AJ02).

^b P.M. Endt, private communication.

$E(^{12}\text{C}) = 67.2$ MeV (1973BE11: not $^{20}\text{Ne}^*(4.97)$ but additionally $^{20}\text{Ne}^*(8.45 - 9.03, 10.26 - 11.02, 11.95 + 12.13, 13.33, 13.92, 16.67, 17.38)$). The angular distributions are symmetric about 90° and compound nucleus formation appears to be dominant (1973BE11, 1974HA11). In the work of (1973BE11) states which are particularly strongly populated are $^{20}\text{Ne}^*(16.67, 17.38, 18.11, 19.16, 19.6)$. The excitation functions for many states of ^{20}Ne with $E_x < 10$ MeV have been measured by (1975VO02) for $E(^{14}\text{N}) = 20$ to 60 MeV and by (1977MO1X) for $E(^{14}\text{N}) = 30$ to 90 MeV. In addition, see (1976ST07) and ^{26}Al in (1978EN06). See also (1975ZE1C), (1971BA2V, 1973BR1C, 1973FO1E, 1973SC1J, 1973SC1B, 1974ST1L) and (1974KL13, 1975KL05, 1975KL08, 1976KL03; theor.).

9. $^{12}\text{C}(^{18}\text{O}, ^{10}\text{Be})^{20}\text{Ne}$ $Q_m = -6.3476$

At $E(^{18}\text{O}) = 80$ MeV, the population of $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 8.78, 11.95)$ is observed (1974BA15).

10. $^{12}\text{C}(^{19}\text{F}, ^{11}\text{B})^{20}\text{Ne}$ $Q_m = -3.1122$

At $E(^{19}\text{F}) = 40, 68$ and 68.8 MeV angular distributions involving $^{11}\text{B}_{\text{g.s.}}$ and $^{20}\text{Ne}^*(0, 1.63)$ are reported (1972SC03). See also (1975PU02) and ^{31}P in (1978EN06) and (1972BO21; theor.).

11. $^{13}\text{C}(^{12}\text{C}, n\alpha)^{20}\text{Ne}$ $Q_m = -0.3283$

See (1973SA1G, 1976DA13) and ^{25}Mg in (1978EN06).

12. $^{14}\text{C}(^{18}\text{O}, ^{12}\text{Be})^{20}\text{Ne}$ $Q_m = -15.75$

See (1974BA15, 1974CE1A).

13. (a) $^{14}\text{N}(^6\text{Li}, \text{p})^{19}\text{F}$ $Q_m = 11.1491$ $E_b = 23.994$
 (b) $^{14}\text{N}(^6\text{Li}, \text{d})^{18}\text{F}$ $Q_m = 2.942$
 (c) $^{14}\text{N}(^6\text{Li}, \alpha)^{16}\text{O}$ $Q_m = 19.2628$
 (d) $^{14}\text{N}(^6\text{Li}, \alpha n)^{15}\text{O}$ $Q_m = 3.5990$

Yield curves have been measured for $E(^6\text{Li}) = 4.1$ to 6.3 MeV for the protons to $^{19}\text{F}^*(2.78)$ and for the d_0 and α_0 groups (1968RI13) and at 5 to 9.2 MeV for the $p_0, p_1, d_0, d_{1+2}, \alpha_0, \alpha_{1+2}$ and α_{3+4} groups (1977LE1N): no structure is apparent.

Table 20.21: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ ^a

E_x (MeV \pm keV)		J^π ^d	K^π ^d
(1971HA26) ^b	(1975ME04) ^c		
1.6329 ± 1.0		2^+	0_1^+
4.2456 ± 2.5		4^+	0_1^+
4.9663 ± 2.5	4.968	2^-	2^-
5.618 ± 4	5.618	3^-	2^-
	5.774	1^-	0^-
	6.725 ^e	0^+	0_2^+
7.004 ± 4	7.004 ^e	4^-	2^-
	7.169 ^e	3^-	0^-
	7.196 ^e	0^+	0_3^+
	7.435 ^e	2^+	0_2^+
	7.835 ^e	2^+	0_3^+
8.446 ± 9	8.451	5^-	2^-
	8.694	1^-	(1_1^-)
	8.779	6^+	0_1^+
	8.85	1^-	(1_1^-)
	9.033	4^+	0_3^+ ^f
	9.110	d	
	9.318	d	
	9.533		
	9.872	$1^+, 2^-, 3^+$ ^{d,g}	
9.950 ± 6		$1^+, 2^-, 3^+$ ^g	
	10.024		
	10.264	5^-	0^-
	10.407	d	
	10.545		
10.609 ± 7	10.609	6^-	2^-
	10.694	$4^-, 3^+$ ^g	
	10.840	d	
10.920 ± 7	10.915	3^+ ^g	
	11.013		

Table 20.21: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ ^a (continued)

E_x (MeV \pm keV)		J^π ^d	K^π ^d	
(1971HA26) ^b	(1975ME04) ^c			
11.528 ± 6 ^g	11.555	$1^+, 2^-, 3^+$ ^{d,g}		
	11.656	(3^+) ^{d,g}		
	11.871	d		
	11.949	8^+	0_1^+	
	12.134 ^h	6^+	0_3^+ ^f	
	12.381			
	12.436 ± 5 ⁱ	12.594		
		12.730		
		12.919		
		13.010		
13.049				
13.190				
13.277				
13.335		7^-	2^-	
13.441		e		
13.569				
13.631				
13.679				
13.845				
13.886				
14.144				
14.305				
14.44				
14.60				
14.812				
15.034	d			
15.155 ^j	6^+ ^j			
15.359	d			

Table 20.21: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ ^a (continued)

E_x (MeV \pm keV)		J^π ^d	K^π ^d
(1971HA26) ^b	(1975ME04) ^c		
	15.50	(8 ⁻)	(2 ⁻)
	15.691		
	15.879	(8 ⁺) ^{d,k,l}	(2 ⁻)
	16.139		
	(16.50)		
	16.52 ^m	7 ⁻ ^m	
	16.68 ^m	7 ⁻ ^m	
	17.372 ⁿ	9 ⁻ ⁿ	2 ⁻ ⁿ
	18.11	(7 ⁻) ^{l,m}	
	20.5 ^o		
	20.67 ^m	9 ⁻ ^m	

^a See Table 20.16 in (1972AJ02) for the earlier work.

^b From measurements of γ -rays.

^c From measurements of α -groups: approximately ± 6 keV for $E_x < 13.5$ MeV and ± 15 keV for higher E_x . Angular distributions have also been obtained at $E(^{12}\text{C}) = 37$ MeV.

^d See discussion in (1975ME04).

^e Previously reported as 6722, 7007, 7159, 7195, 7421 and 7833 (± 4) keV (1971MI09).

^f (1974FO20, 1974FO25).

^g (1976FI10): see Table 20.19.

^h Alpha decay is by α_2 to $^{16}\text{O}^*(6.13)$, $\Gamma_{\alpha'}/\Gamma = (6.0 \pm 0.15)\%$: assuming $\Gamma_\alpha \Gamma_{\alpha'}/\Gamma = 7.7 \pm 3.8$ eV this leads to $\Gamma_\alpha = 0.128 \pm 0.072$ keV for this 6⁺ state (1972BA97).

ⁱ (1977BA3W, 1977BA3X): $\Gamma_{\text{lab}} = 29 \pm 1$ keV.

^j 15.18 ± 0.02 MeV. Alpha decay is $3 \pm 3\%$ by α_0 , $49 \pm 3\%$ by $\alpha_1 + \alpha_2$ (mainly α_2 , to $^{16}\text{O}^*(6.13)$) and $48 \pm 3\%$ by $\alpha_3 + \alpha_4$ (mainly α_3 to $^{16}\text{O}^*(6.92)$) (1973FI12, 1973ZU1A).

^k Angular correlations suggest 8⁺ for this state which decays $\approx 10\%$ by α_0 and $\approx 90\%$ by $\alpha_1 + \alpha_2$ (mainly the latter) (K. Young, private communication).

^l See also (1972PA16).

^m Alpha- α and $\alpha - \gamma$ correlations (1977YO1H) and (K. Young, private communication).

ⁿ Decays $> 99\%$ to $^{16}\text{O}^*(6.13)$ by a pure $L = 6$ transition (1973FI12, 1973ZU1A).

^o Reported by (1977CO2C: preliminary results).

Table 20.22: Resonances in $^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}$

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\omega\gamma^{\text{a}}$ (eV)	E_x (MeV \pm keV)	$J^\pi; T$	Refs.
1.116 \pm 4	2.6×10^{-6} ^a	$(1.7 \pm 0.3) \times 10^{-3}$	5.624	$3^-; 0^{\text{d}}$	(1965VA14, 1971TO06, 1971TO1C)
1.319 \pm 3	$> 1.3 \times 10^{-2}$ ^a	$(14 \pm 3) \times 10^{-3}$	5.786	$1^-; 0^{\text{e}}$	(1965VA14, 1971TO06, 1971TO1C)
2.490 \pm 8	15 ± 7 ^a	$(38 \pm 10) \times 10^{-3}$	6.722	$0^+; 0$	(1965VA14, 1971TO06, 1971TO1C)
3.074	4	$(4.35 \pm 0.75) \times 10^{-3}$	7.189 ± 3	$0^+; 0$	(1972AL32)
3.363	8	0.146 ± 0.019	7.421 ± 1	$2^+; 0$	(1972AL32)
3.872	2.4	0.343 ± 0.035	7.828 ± 3	$2^+; 0$	(1972AL32)
(4.647 \pm 3)			(8.447)	$(5^-; 0)$	(1971RO33)
5.06	< 3	1.35 ± 0.15	8.776 ± 3.2	$6^+; 0$	(1967LI07, 1971DI08, 1971RO13)
5.368	3.2	3.05 ± 0.38	9.024 ± 3	$4^+; 0$	(1964PE05, 1972AL32)
5.94 ± 30	29 ± 15	1.3 ± 0.5	9.48	$2^+; 0$	(1964PE05)
6.61 ± 30	155 ± 30	8 ± 3	10.02	$(4^+); 0$	(1964PE05)
6.924 ± 7	≤ 1	$19.5 \pm 1.5^{\text{g}}$	$10.2712 \pm 7^{\text{f}}$	$2^+; 1$	(1964PE05, 1976IN05, 1977FI08, 1977MA07, 1978ST08)
7.932 ± 10	≤ 3	30.4 ± 3	11.074	$(4^+; 1)$	(1964PE05, 1978ST08)
8.161 ± 10	≤ 3	0.58 ± 0.05	11.257	$1^-; 1$	(1964PE05, 1978ST08)
8.53 ± 10	≤ 5	0.41 ± 0.05	11.55	$(2^+, 0^+)$	(1964PE05, 1978ST08)
i		0.104 ± 0.035	11.948	$8^+; 0^{\text{h}}$	(1972AL05)
(9.05 \pm 50)	< 40		11.97		(1978ST08)
(9.15 \pm 50)	< 40		12.05		(1978ST08)
$9.359 \pm 5^{\text{b}}$	< 2		12.215	$2^+; 1$	(1977MA07)
$9.407 \pm 6^{\text{c}}$	≈ 5		12.254	$3^-, 2^+$	(1977MA07)
9.57 ± 10	33 ± 4	1.94 ± 0.15	12.38	$3^-; (1)$	(1978ST08)
9.70 ± 30	≤ 10	0.17 ± 0.05	12.49		(1978ST08)

^a This is also Γ_α .

^b $\Gamma_{\alpha_0}\Gamma_{\gamma_1}/\Gamma = 0.292 \pm 0.044$ eV; $0.25 \leq \Gamma_{\alpha_0} \leq 1.34$ eV, strong support for the $T = 1$ assignment (1977MA07). See also (1978ST08).

^c $(2J + 1)\Gamma_{\alpha_0}\Gamma_{\gamma_1}/\Gamma = 3.9 \pm 0.8$ eV (1977MA07). See also (1978ST08).

^d $K^\pi = 2^-$ (1973HA63).

^e $K^\pi = 0^-$ (1973HA63).

^f From E_γ measurements (1977FI08). The measurements of the decay of this state lead to $E_x = 4247.9 \pm 1.3, 4966.0 \pm 1.9, 5621.0 \pm 3.5, 7423.1 \pm 3.0, 7828.1 \pm 3.8$ and 8776.7 ± 2.3 keV (1977FI08).

^g $\omega\gamma = 19.2 \pm 1.9$ eV (1978ST08); $\Gamma_\alpha = 116 \pm 20$ eV (1976IN05); $\Gamma_\gamma = 4.26 \pm 0.23$ eV (1977FI08) [summary of several measurements].

^h $K^\pi = 0_1^+$ (1972AL05).

ⁱ $E(^{16}\text{O}) \approx 36.05$ MeV.

14. $^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}$

$$Q_m = 4.7309$$

Observed resonances in the yield of capture γ -rays over the range $E_\alpha = 0.8$ to 10 MeV are displayed in Table 20.22. Information on the character of the radiative decay is shown in Table 20.19. For τ_m , see Table 20.20.

No resonances have been observed below $E_\alpha = 1$ MeV: see (1971TO06, 1971TO1C) for the data and for the astrophysical implications of this reaction. See also (1971BA1A, 1973CL1E).

The $J^\pi = 5^-$ state at $E_x = 8.45$ MeV [$E_\alpha = 4.65$ MeV] decays by an E2 transition [$|M|^2 = 26 \pm 6$ W.u.] to the 3^- state at $E_x = 5.62$ MeV (1971RO33). The $J^\pi = 6^+$ state at $E_x = 8.78$ MeV [$E_\alpha = 5.06$ MeV] decays by an E2 transition [$|M|^2 = 20.4 \pm 2.4$ W.u.] to the 4^+ state at $E_x = 4.25$ MeV (1971DI08, 1971RO13). See also (1971RO1C). The $J^\pi = 8^+$ state at $E_x = 11.95$ MeV decays by an E2 transition to the $E_x = 8.78$ MeV [$J^\pi = 6^+$] state which then decays via the 4^+ and 2^+ members of the ground-state rotational band. The transition probability of the $8^+ \rightarrow 6^+$ transition is 7.5 ± 2.5 W.u. which establishes $^{20}\text{Ne}^*(11.95)$ as the 8^+ member of the ground-state band. The experimental E2 transition probabilities in the ground-state band deviate strongly from those predicted by the pure rotational model but agree reasonably well with simple shell-model predictions (1972AL05). The $T = 0$ isospin impurity of $^{20}\text{Ne}^*(10.27)$ [$2^+; 1$] is quite large: see (1977FI08) for a discussion of this point and for CVC predictions of the weak magnetism form factors from the radiative widths for the transitions to $^{20}\text{Ne}^*(1.63, 7.42)$ [see Table 20.19]. See also (1977WA1V), (1973AL1C, 1974HA1C) and (1973GA23; theor.).

15. (a) $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$

$$E_b = 4.7309$$

(b) $^{16}\text{O}(\alpha, 2\alpha)^{12}\text{C}$

$$Q_m = -7.1620$$

Excitation functions have been measured over a wide energy range for elastically and inelastically scattered α -particles, and γ -rays from the decay of $^{16}\text{O}^*(6.13, 6.92, 7.12)$: see Table 20.20 in (1972AJ02) for the earlier work and (1973HA63: 4.6 MeV and 14.1 to 17.1 MeV; α_0, α_{1+2} ; the latter in the higher energy range), (1978ST08: 6.9 to 10.2 MeV; $\alpha_0, \alpha_2\gamma$), (1977MA07: 9.2 to 9.7 MeV; $\gamma_{6.13}$), (1975CE01: 12.8 to 14.8 MeV; α_0), (1975BI1H: 14.0 to 18.1 MeV; $\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$) and (1972BU1H: 22.75 to 28.40 MeV). See also (1974BE41) and ^{16}O in (1977AJ02).

A number of anomalies are observed: they are displayed in Table 20.23. Identifying various of the observed states as members of different rotational bands is discussed by (1972HA07, 1973HA63). The reduced α -widths in the first $K^\pi = 0^+, 2^-$ and 0^- bands are small, indicating that the bands could be described in terms of the spherical shell model. The reduced widths within bands are found to decrease sharply with increasing spin (1972HA07, 1973HA63).

For reaction (b) see (1972AJ02). For spallation reactions involving the emission of ^6Li , ^7Li , ^7Be , ^8Be and ^9Be ions see (1972RU03, 1974JE1A, 1974RA11). See also (1973AL1C) and (1972HO1E, 1972HO56, 1972SU10, 1973AR1C, 1973EB1A, 1973HO1L, 1973IC01, 1974CA31, 1974FR06, 1974KA1P, 1974TA17, 1974WA1D, 1975BA05, 1975CU1B, 1975KO28, 1975KU1K, 1975MA26, 1975NE04, 1975TA1A, 1976CU07, 1976HE19, 1976JA1F, 1976LE20, 1976PA1J, 1977AB03, 1977AN1B, 1977BA12, 1977FL13, 1977IK1C, 1977OH01; theor.).

Table 20.23: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%) ^a	E_x (MeV)	J^π	Refs.
2.490 \pm 10	19	α_0	22	6.722	0 ⁺ i	(1953CA44)
3.045 \pm 10	8	α_0	36	7.166	3 ⁻ g	(1953CA44)
3.090 \pm 10	4	α_0	1.1	7.202	0 ⁺ j	(1953CA44)
3.380 \pm 10	8	α_0	4.7	7.434	2 ⁺ i	(1953CA44)
3.885 \pm 10	2	α_0	0.6	7.838	2 ⁺ j	(1953CA44)
4.653 \pm 5	0.013 \pm 0.004	α_0	0.07	8.452	5 ⁻ h	(1973HA63)
\approx 4.9	> 800	α_0	\approx 70	\approx 8.6	0 ⁺ k	(1960MC09)
5.002	2.5	α_0	0.23	8.731	1 ⁻	(1960MC09)
5.058 \pm 3	0.11 \pm 0.02	α_0	8.5 \pm 1.5	8.776	6 ⁺ f	(1972HA07)
\approx 5.1	> 800	α_0	\approx 95	\approx 8.8	2 ⁺ k	(1960MC09)
5.11	< 1	α_0		8.82	(5 ⁻)	(1960MC09)
5.152 \pm 5	19	α_0	1.1	8.851	1 ⁻	(1960MC09, 1969JO18)
5.395 \pm 5	3	α_0	3.9	9.046	4 ⁺ j	(1960MC09, 1969JO18)
5.486 \pm 5	3.2	α_0	0.49	9.118	3 ⁻	(1960MC09, 1969JO18)
5.955 \pm 10	24	α_0	1.4	9.493	2 ⁺	(1960MC09, 1967HU06, 1969JO18)
6.569 \pm 10	97	α_0	17	9.984	4 ⁺ i	(1967HU06, 1969JO18)
6.912 \pm 5	141	α_0	66	10.259	5 ⁻ g	(1967HU06, 1969JO18)
6.92 \pm 10	\leq 0.3	α_0	$\leq 1.3 \times 10^{-3}$	10.27	(2 ⁺)	(1978ST08)
7.092 \pm 5	81	α_0	4.8	10.403	3 ⁻	(1967HU06, 1969JO18)
7.276 \pm 5	16	α_0	1.8	10.550	4 ⁺	(1969JO18)
7.314 \pm 10	24	α_0	0.85	10.580	2 ⁺	(1965MC02, 1967HU06, 1969JO18)
7.580 \pm 100	349	α_0	33	10.79	4 ⁺ k	(1967HU06, 1969JO18)
7.635 \pm 5	13	α_0	0.42	10.837	2 ⁺	(1965MC02, 1967HU06, 1969JO18)
7.636	45	α_0	2.1	10.838	3 ⁻	(1969JO18)

Table 20.23: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%) ^a	E_x (MeV)	J^π	Refs.
(7.75)	80	α_0		(10.93)		(1967HU06)
7.80 ± 150	576	α_0	14	10.97	0^+	(1969JO18)
7.860 ± 10	24	α_0	2.0	11.017	4^+	(1965MC02, 1967HU06, 1969JO18)
7.93 ± 10	≤ 0.5	α_0	≤ 0.05	11.07	(4^+)	(1978ST08)
8.132 ± 30	172	α_0	4.2	11.234	1^-	(1969JO18)
8.16 ± 10	≤ 0.3	α_0	≤ 0.009	11.26	(1^-)	(1978ST08)
8.24 ± 10	40 ± 10	α_0	1.4	11.32	2^+	(1965MC02, 1967HU06, 1969JO18, 1978ST08)
8.528 ± 10	1.0 ± 0.5	α_0	0.03, 0.02	11.551	($2^+, 0^+$)	(1978ST08)
(≈ 8.6)	≈ 500	α_0		(≈ 11.6)	(2^+)	(1967HU06)
8.930 ± 20	46	α_0	1.1	11.872	2^+	(1969JO18)
8.997 ± 5	0.44 ± 0.15	$\alpha_0, \gamma_{6.13}$	0.04 ± 0.01	11.926	4^+	(1972HA07)
9.026 ± 5	$(35 \pm 10) \times 10^{-3}$	α_0	1.0 ± 0.3	11.949	8^+ f	(1972HA07)
9.043 ± 10	30 ± 5	α_0	0.72	11.962	1^-	(1967HU06, 1969JO18, 1972HA07, 1978ST08)
(9.37 ± 20)	≤ 20	α_0	≤ 0.5	(12.22)	(2^+)	(1978ST08)
9.39 ± 30	148 ± 20	α_0	7.7	12.24	4^+	(1964PE05, 1967HU06, 1969JO18, 1978ST08)
9.530 ± 100	≈ 500	α_0	≈ 13	12.35	2^+	(1969JO18, 1978ST08)
$9.58 \pm 10 \text{ b,c}$	37 ± 5	$\alpha_0, \gamma_{6.13}$	1.2	12.39	3^-	(1964PE05, 1969JO18, 1977MA07, 1978ST08)
9.605 ± 5	≤ 8	α_0	< 0.15	12.412	0^+ e	(1969JO18)
9.790 ± 10	88 ± 10	α_0	28	12.560	6^+ k	(1967HU06, 1969JO18, 1978ST08)
(9.860 ± 100)		α_0		(12.62)		(1969JO18)
9.944 ± 15	97	α_0	7.3	12.683	5^-	(1969JO18)

Table 20.23: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%) ^a	E_x (MeV)	J^π	Refs.
10.050 \pm 100 ^d	100	α_0		12.77	4 ⁺	(1967ME10, 1969JO18)
10.14 \pm 70	55	$\alpha_0, \gamma_{6.13}$		12.84		(1967ME10)
10.32 \pm 75	60	$\alpha_0, \gamma_{6.13}$		12.98	(4 ⁺)	(1967ME10)
10.43 \pm 90	70	$\alpha_0, \gamma_{6.13}$		13.07	(4 ⁺)	(1967ME10)
10.57 \pm 75	60	$\alpha_0, \gamma_{6.13}$		13.18	(4 ⁺)	(1967ME10)
10.759 \pm 6	$(80 \pm 30) \times 10^{-5}$	α_0	0.08 \pm 0.03	13.334	7 ⁻ ^h	(1972HA07)
10.770 \pm 6	20 \pm 5	$\alpha_0, \gamma_{6.13}$	0.07 \pm 0.03	13.343	4 ⁺	(1967ME10, 1972HA07)
10.83 \pm 50	40	$\gamma_{6.13}$		13.39		(1967ME10)
10.87 \pm 140	110	$\alpha_0, \gamma_{6.13}$		13.42	(4 ⁺)	(1967ME10)
11.20 \pm 400	320	$\alpha_0, \gamma_{6.13}$		13.7	(3, 7) ⁻	(1967ME10)
11.51 \pm 125	400	$\alpha_0, \gamma_{6.13}$		13.93	(6 ⁺)	(1967ME10)
11.77		$\alpha_0, \gamma_{6.9+7.1}$		14.14		(1967ME10)
11.97 \pm 300	240	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		14.3	6 ⁺	(1967ME10)
(12.06)		$\alpha_0, \gamma_{6.9+7.1}$		(14.37)		(1967ME10)
12.31 \pm 300	240	$\alpha_0, \gamma_{6.9+7.1}$		14.6	(4 ⁺)	(1967ME10)
12.66 \pm 150	120	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		14.85		(1967ME10)
12.86 \pm 150	120	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.01		(1967ME10)
13.165 \pm 150	120	$\alpha_0, \gamma_{6.13}$		15.26		(1967ME10)
13.22		α_0		15.30		(1967ME10)
13.37 \pm 470	380	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.4	7 ⁻ ^g	(1967ME10, 1975CE01)
13.58		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.59		(1967ME10)
13.73		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.71	(6 ⁺)	(1967ME10, 1975CE01)
14.05		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		15.97	(6 ⁺)	(1967ME10, 1975CE01)
14.26		$\gamma_{6.13}, \gamma_{6.9+7.1}$		16.13		(1967ME10)

Table 20.23: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%) ^a	E_x (MeV)	J^π	Refs.
14.40		$\gamma_{6.13}$		16.25		(1967ME10)
14.501 \pm 15	43	α_0, α_{1+2}		16.326	4 ⁺	(1973HA63)
14.636 \pm 15	34	α_0, α_{1+2}		16.434	(0, 2, 4) ⁺	(1973HA63)
14.732 \pm 15	23	α_0, α_{1+2}		16.510	(2, 4, 6) ⁺	(1973HA63)
14.935 \pm 15	110	α_0		16.673	(0, 2) ⁺	(1973HA63)
14.993 \pm 15	10	α_0, α_{1+2}		16.719	(1, 3, 7) ⁻	(1973HA63)
15.162 \pm 15	10	α_0, α_{1+2}		16.854	5 ⁻	(1973HA63)
15.547 \pm 15	37	α_0, α_{1+2}		17.162	5 ⁻ , (7 ⁻)	(1973HA63)
15.695 \pm 15	32	α_0, α_{1+2}		17.280	1 ⁻ , 3 ⁻ , 4 ⁺	(1973HA63)
15.828 \pm 15	< 10	α_{1+2}		17.387		(1973HA63)
16.023 \pm 15	136	α_0, α_{1+2}		17.542	6 ⁺	(1973HA63)
16.285 \pm 15	36	α_0, α_{1+2}		17.752	4 ⁺ , (0 ⁺)	(1973HA63)
16.598 \pm 15	< 10	α_0, α_{1+2}		18.002	7 ⁻	(1973HA63)
16.623 \pm 15	45	α_0, α_{1+2}		18.022	(2 ⁺ , 5 ⁻ , 6 ⁺)	(1973HA63)
16.737 \pm 15	33	α_0, α_{1+2}		18.113	7 ⁻	(1973HA63)
16.98 \pm 300	240	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$		18.31	(6 ⁺)	(1967ME10)
17.45	600	$\alpha_0, \gamma_{6.13}$		18.7	(6 ⁺)	(1967ME10)
18.05 \pm 250	200	$\alpha_0, \gamma_{6.9+7.1}$		19.16	(6 ⁺)	(1967ME10)
18.35 \pm 350	280	α_0		19.40	6 ⁺	(1967ME10)
18.90 \pm 350	280	α_0		19.84	6 ⁺	(1967ME10)
19.30 \pm 120	250	α_0		20.16	7 ⁻	(1971BE17)
19.6 \pm 180	360	α_0		20.4	6 ⁺	(1971BE17)
19.6 \pm 100	200	α_0		20.4	7 ⁻	(1971BE17)
19.95 \pm 60	120	α_0		20.68	9 ⁻	(1971BE17)

Table 20.23: Resonances in $^{16}\text{O}(\alpha, \alpha)^{16}\text{O}$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	θ^2 (%) ^a	E_x (MeV)	J^π	Refs.
20.18		α_0		20.9		(1971BE17)
20.4 \pm 100	200	α_0		21.0	7 ⁻	(1971BE17)
20.45 \pm 40	80	α_0		21.08	9 ⁻	(1971BE17)
20.70	300	α_0		21.3	7 ⁻	(1962JO14, 1971BE17, 1971TA05)
21.3 \pm 200	300	α_0		21.8	7 ⁻	(1971BE17, 1971TA05)
22.0 \pm 200	500	α_0		22.3	7 ⁻	(1971BE17, 1971TA05)
22.5 \pm 250	500	α_0		22.7	9 ⁻	(1971BE17)
22.65 \pm 125	250	α_0		22.84	9 ⁻	(1971BE17)
23.3 \pm 250	500	α_0		23.4	8 ⁺	(1971BE17, 1971TA05)
24.24 \pm 150	350	α_0		24.11	8 ⁺	(1971BE17, 1971TA05)
25.4 \pm 300	600	α_0		25.0	8 ⁺	(1971BE17)
26.2 \pm 200	400	α_0		25.7		(1971BE17)
28.1 \pm 350	700	α_0		27.2		(1971BE17)
29	1600	α_0		28	8 ⁺	(1969CO19, 1970CO13)
29.4 \pm 350	700	α_0		28.2		(1971BE17)

^a See also discussion and Table 2 in (1973HA63).

^b $^{20}\text{Ne}^*(12.39)$ decays by α_2 to $^{16}\text{O}^*(6.13)$ with $\omega\Gamma_{\alpha}\Gamma'_{\alpha}/\Gamma = 3 \pm 1$ keV (1964PE05), 2.0 ± 0.1 keV (1978ST08).

^c $\omega\gamma = 0.8 \pm 0.2$ and 1.94 ± 0.15 eV to $^{20}\text{Ne}^*(1.63, 4.25)$, respectively; $\omega\gamma_0 < 0.02$ eV (1978ST08).

^d Values quoted are taken preferentially from the elastic scattering results (1967ME10).

^e (1977BA3W, 1977BA3X) report a probable 0^+ state at $E_x = 12.436 \pm 0.005$ MeV, $\Gamma_{\text{lab}} = 29 \pm 1$ keV, $\Gamma_{\alpha_0} = 22.7 \pm 2.0$ keV, $\Gamma_{\alpha_1} = 6.3 \pm 2.0$ keV. These data are preliminary and are not used in Table 20.18.

^f $K^\pi = 0_1^+$ (1972HA07, 1973HA63).

^g $K^\pi = 0^-$ (1973HA63).

^h $K^\pi = 2^-$ (1972HA07, 1973HA63).

ⁱ $K^\pi = 0_2^+$ (1973HA63).

^j $K^\pi = 0_3^+$ (1973HA63).

^k $K^\pi = 0_4^+$ (1973HA63).

$$16. \text{}^{16}\text{O}(\alpha, n)^{19}\text{Ne} \quad Q_m = -12.1344 \quad E_b = 4.7309$$

The excitation function (activation measurements) has been measured from threshold to $E_\alpha = 26.8$ MeV (1973GR29). See also (1974LO1B, 1977LI19) and ^{19}Ne .

$$17. \text{}^{16}\text{O}(\alpha, p)^{19}\text{F} \quad Q_m = -8.1137 \quad E_b = 4.7309$$

See ^{19}F .

$$18. \text{}^{16}\text{O}(\alpha, \text{}^3\text{He})^{17}\text{O} \quad Q_m = -16.434 \quad E_b = 4.7309$$

See ^{17}O in (1977AJ02) and (1972RI03; theor.).

$$19. \text{}^{16}\text{O}(\text{}^6\text{Li}, d)^{20}\text{Ne} \quad Q_m = 3.257$$

Deuteron groups have been observed to many states of ^{20}Ne : see Table 20.24. Angular distributions have been measured at a number of energies for $E(^6\text{Li}) = 5.5$ to 25.8 MeV [see (1972AJ02)] and at 32 MeV (1973GO1K, 1973GO1L; abstracts; to most states shown in Table 20.24 with $E_x < 12.2$ MeV), (1977GU01: d to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62, 7.00)$) and at 35.3 to 45 MeV (1975AR20, 1975AR25, 1976AR04: see Table 20.24). (1977GU01) report that the forward peaks of the deuterons to $^{20}\text{Ne}^*(4.97, 5.62)$ [$J^\pi = 2^-$ and 4^-] cannot be explained solely in terms of zero-range coupled-channel Born approximation α -transfer. The α -particle spectroscopic strengths for the ground state $K^\pi = 0^+$ states have been studied by (1973FO06, 1975AN13). (1973FO06) find 0.22 and 0.15 for the α reduced widths for $^{20}\text{Ne}^*(8.78, 11.95)$ in agreement with their assignments as the 6^+ and 8^+ members of the $K^\pi = 0_1^+$ band. See also (1972PA16, 1977BE2R). ($d\gamma$) coincidences are being studied by (1977SU1M; abstract).

For excitation functions see (1976HO1K) and ^{22}Na in (1978EN06). See also (1972BA1P, 1973ST1E), (1971BA2V, 1972GA1E, 1973FO1E, 1973FO1A, 1973OG1A, 1975GO15, 1976OG1A) and (1973YA1A, 1974DO03, 1974NE18, 1976KO1L; theor.).

$$20. \text{}^{16}\text{O}(\text{}^7\text{Li}, t)^{20}\text{Ne} \quad Q_m = 2.264$$

A number of ^{20}Ne states have been studied in this reaction: see Table 20.24 (1976CO23) and P.D. Parker, private communication. Angular distributions have been reported at $E(^7\text{Li}) = 15$ to 30.3 MeV [see (1972AJ02)] and at 38 MeV (1976CO23). Correlations between the tritons and the α -particles to ^{16}O have been studied for a number of states of ^{20}Ne at $E(^7\text{Li}) = 20$ and 24 MeV: it is suggested that the reaction mechanism is mainly direct (1974PA16). See also (1977DA1J). See also (1972BA1P, 1972CA1D, 1972PA16, 1973ST1E, 1977SA1X), (1971BA2V, 1972GA1E, 1973BR1C, 1973FO1A, 1973OG1A, 1975GO15) and (1972KU12, 1972KU13, 1973AR1C, 1973IC01, 1973MC17, 1974DO03, 1974NE18, 1975BIIJ; theor.).

Table 20.24: States of ^{20}Ne from $^{16}\text{O}(^6\text{Li}, d)$, $^{16}\text{O}(^7\text{Li}, t)$ and $^{16}\text{O}(^{12}\text{C}, ^8\text{Be})$

E_x (MeV \pm keV)			$\Gamma_{\text{c.m.}}$ (keV)	$\Gamma_{\alpha_0}/\Gamma^g$	J^π	K^π
$(^6\text{Li}, d)^a$	$(^7\text{Li}, t)^b$	$(^{12}\text{C}, ^8\text{Be})^g$				
0	0				$0^+{}^c$	0_1^+
1.63	1.63				$2^+{}^c$	0_1^+
4.25	4.25				$4^+{}^c$	0_1^+
4.97					2^-	2^-
5.62					3^-	2^-
5.78	5.78				$1^-{}^d$	0^-
6.72						
7.00					4^-	2^-
7.17	7.17	7.17			$3^-{}^d$	0^-
7.42						
8.45						
8.78	8.78	8.78			6^+	0_1^+
10.3 ± 100	10.26	10.26	145 ± 40	1	5^-	0^-
10.7 ± 100					4^+	
11.95	11.95	11.95		0.85 ± 0.15	8^+	0_1^+
12.15						
12.6 ± 100	12.591 ± 10	12.59	110 ± 40	0.80 ± 0.10	6^+	$0^+{}^e$
13.9	13.904 ± 20		≈ 100		6^+	
14.3	14.310 ± 20		< 100		6^+	
15.35 ± 100	15.336 ± 15	15.34	380 ± 60	0.90 ± 0.10	7^-	$0^-{}^e$
15.9 ± 100			< 250		7^-	
16.7 ± 100	16.63 ± 20	16.63	190 ± 40	0.90 ± 0.10	7^-	
17.35 ± 100	17.30 ± 20	17.30	220 ± 40	0.40 ± 0.10	8^+	
18.7 ± 100					7^-	
19.4 ± 100			400		7^-	
19.9 ± 100			400		7^-	
	20.67 ± 40					
20.8 ± 100					$7^- (6^+)$	
	21.08 ± 30	21.08	100 ± 50	0.65 ± 0.15	9^-	
21.3 ± 100			300		8^+	

Table 20.24: States of ^{20}Ne from $^{16}\text{O}(^6\text{Li}, \text{d})$, $^{16}\text{O}(^7\text{Li}, \text{t})$ and $^{16}\text{O}(^{12}\text{C}, ^8\text{Be})$ (continued)

E_x (MeV \pm keV)			$\Gamma_{\text{c.m.}}$ (keV)	$\Gamma_{\alpha_0}/\Gamma^g$	J^π	K^π
$(^6\text{Li}, \text{d})^a$	$(^7\text{Li}, \text{t})^b$	$(^{12}\text{C}, ^8\text{Be})^g$				
21.8 \pm 100			300		8 ⁺	
22.3 \pm 100			300		8 ⁺	
	22.87 \pm 40	22.87	225 \pm 40	0.90 \pm 0.10	9 ⁻	
23.5 \pm 100	23.70 \pm 30 ^f		\leq 200		9 ⁻ (8 ⁺)	
	24.21 \pm 25 ^f		\approx 500			
	25.10 \pm 50 ^f		\lesssim 200			
	25.67 \pm 50 ^f		\approx 500			
27.1 \pm 100					(9 ⁻)	
28.1 \pm 100					(10 ⁺)	

^a Levels with energy uncertainties shown are from (1975AR20, 1975AR25, 1976AR04: $E(^6\text{Li}) = 35.3$ to 45 MeV). The other states have been reported by other groups: see text.

^b (1976CO23). $E(^7\text{Li}) = 38$ MeV and P.D. Parker (private communication).

^c Relative α -cluster spectroscopic factors for $^{20}\text{Ne}^*(0, 1.63, 4.25)$ are 1.00, 0.81, 0.36 (FRDWBA), 1.00, 1.00, 0.75 (FRCCBA) (1976CO23).

^d Spectroscopic factors are 0.30 and 0.15 for $^{20}\text{Ne}^*(5.78, 7.17)$ (FRDWBA) (1976CO23).

^e (1974PA16).

^f $E(^7\text{Li}) = 60$ MeV (P.D. Parker, private communication).

^g (1977SA1X, 1977SA2A, 1977SA2F, 1977SA2G) and P.D. Parker (private communication): $E(^{12}\text{C}) = 78$ MeV.

$$21. \ ^{16}\text{O}(^9\text{Be}, \alpha\text{n})^{20}\text{Ne} \quad Q_m = 3.1576$$

See (1974VA31).

$$22. \text{ (a) } ^{16}\text{O}(^{12}\text{C}, 2\alpha)^{20}\text{Ne} \quad Q_m = -2.5439$$

$$\text{ (b) } ^{16}\text{O}(^{12}\text{C}, ^8\text{Be})^{20}\text{Ne} \quad Q_m = -2.6358$$

In reaction (a) at $E(^{16}\text{O}) = 58.3$ MeV, $^{20}\text{Ne}^*(0, 1.63, 4.25 + 4.97)$ are strongly populated (1974WI05). See also ^{24}Mg in (1978EN06). For fusion studies see (1976KO24, 1976SW02, 1976WE15, 1977CH10) and for other yield measurements see (1976EY01, 1977SW01) and ^{28}Si in (1978EN06). See also (1975SC2A).

Angular distributions in reaction (b) have been studied at $E(^{16}\text{O}) = 27.1$ to 37.1 MeV (1977HU1C; g.s.), 30.2 , 30.7 , 31.2 and 32.1 MeV (1976VI05, 1977BR24; g.s.) and 46.4 MeV (1976JA19; g.s.), at $E(^{12}\text{C}) = 56$ MeV (1976MA12: to $^{20}\text{Ne}^*(0, 1.63, 4.25, 5.8, 7.2, 8.8, 10.34, \approx 12)$) and at 78 MeV (1977SA1X, 1977SA2A, 1977SA2F, 1977SA2G). At $E(^{12}\text{C}) = 56$ MeV the strongest groups are those to the 6^+ and 5^- states $^{20}\text{Ne}^*(8.78, 10.26)$ (1976MA12). See also (1972WO10). At $E(^{12}\text{C}) = 78$ MeV, $^8\text{Be}-\alpha$ correlations have led to a number of J^π determinations and to measurements of Γ_{α_0}/Γ : see Table 20.24 (P.D. Parker, private communication). For yield measurements see (1976JA19, 1976VI05, 1977BR24, 1977FL1F, 1977HU1C) and ^{28}Si in (1978EN06). See also (1972PA1G, 1974ME09) and (1973YO03, 1974KR1E, 1975BA2J, 1976BU22; theor.).

$$23. \ ^{16}\text{O}(^{13}\text{C}, ^9\text{Be})^{20}\text{Ne} \quad Q_m = -5.9170$$

At $E(^{13}\text{C}) = 105$ MeV the cross sections for formation of the $K^\pi = 0^+$ states $^{20}\text{Ne}^*(0, 1.63, 4.25, 8.78, 11.95)$ are in excellent agreement with shell-model predictions. $^{20}\text{Ne}^*(10.26, 17.38, 21.1)$, as well as the 4^+ , 6^+ and 8^+ states above, are strongly populated (1976PI16).

$$24. \ ^{16}\text{O}(^{14}\text{N}, ^{10}\text{B})^{20}\text{Ne} \quad Q_m = -6.8823$$

At $E(^{14}\text{N}) = 155$ MeV states with high angular momentum appear to be preferentially excited (1976NA22).

$$25. \ ^{16}\text{O}(^{16}\text{O}, ^{12}\text{C})^{20}\text{Ne} \quad Q_m = -2.4310$$

Angular distributions have been reported at $E(^{16}\text{O}) = 23.9$ MeV (1974SP06; g.s.), $44.3 - 54.5$ MeV (1972SI17, 1973SI1D; to $^{20}\text{Ne}^*(0, 1.63, 4.25)$), 51.5 MeV (1974RO04; to $^{20}\text{Ne}^*(0, 1.63, 4.3, 5.6 - 6.1, 8.5 - 8.9)$) and 95.2 MeV (1977MO1H; to $^{20}\text{Ne}^*(0, 1.63)$). At 55.5 MeV (1973ER1B, 1974ER1A; prelim.), $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 7.00)$ are populated. For yield measurements see (1972SI17, 1973PE1D, 1974ER1A, 1974RO04, 1974SP06, 1977KO16) and ^{32}S in (1978EN06). In particular the fusion cross section shows evidence of periodic structure between $E_{\text{c.m.}} = 14$ and 28 MeV, as predicted by the optical model (1977KO16): a resonance ($\Gamma \leq 1$ MeV) at $E_{\text{c.m.}} = 26.5$ MeV is suggested to be an α -exchange resonance. Polarization measurements are reported by (1977PO1E). See also (1973ST1A, 1973ST1F) and (1972AR11, 1973AR18; theor.).

$$26. \ ^{16}\text{O}(^{18}\text{O}, ^{14}\text{C})^{20}\text{Ne} \quad Q_m = -1.4970$$

Angular distributions are reported at $E(^{18}\text{O}) = 28$ and 32 MeV (1971BA68).

27. $^{16}\text{O}(^{19}\text{F}, ^{15}\text{N})^{20}\text{Ne}$ $Q_m = 0.7171$

At $E(^{19}\text{F}) = 36$ MeV (1973GA14) and $E(^{16}\text{O}) = 46, 58$ and 68 MeV (1975PO01) angular distributions involving $^{20}\text{Ne}^*(0, 1.63)$ are reported. (1975PO01) have also studied $^{20}\text{Ne}^*(4.25)$: the strong population of the 4^+ state may indicate evidence for a two-step process. See also (1976LO06; theor.).

28. (a) $^{17}\text{O}(^3\text{He}, n)^{19}\text{Ne}$ $Q_m = 4.299$ $E_b = 21.164$
 (b) $^{17}\text{O}(^3\text{He}, \alpha)^{16}\text{O}$ $Q_m = 16.4335$

For reaction (a) see ^{19}Ne . The ground state excitation energy for reaction (b) has been measured for $E(^3\text{He}) = 7.0$ to 10.0 MeV: it shows a resonance corresponding to $^{20}\text{Ne}^*(28.)$. This resonance is also observed in the $^{16}\text{O}(\alpha, \alpha)$ elastic scattering. It is interpreted in terms of a quasi-molecular α -particle cluster model (1969CO19).

29. $^{17}\text{O}(\alpha, n)^{20}\text{Ne}$ $Q_m = 0.587$

Angular distributions have been measured at $E_\alpha = 9.8$ to 12.3 MeV by (1967HA14: n_1, n_2, n_{4+5}). The neutron decay of the lowest $T = \frac{3}{2}$ state in ^{21}Ne to $^{20}\text{Ne}^*(1.63)$ has been studied by (1976MC12). The total neutron yield has been measured for $E_\alpha = 1.0$ to 5.3 MeV by (1973BA10). The astrophysical import of this reaction is discussed by (1973CL1E, 1975FO19). See also ^{21}Ne in (1978EN06).

30. $^{17}\text{O}(^6\text{Li}, t)^{20}\text{Ne}$ $Q_m = 5.370$

See (1973ST1E, 1977MA2G). See also (1974KU07; theor.).

31. $^{18}\text{O}(^3\text{He}, n)^{20}\text{Ne}$ $Q_m = 13.1199$

Neutron groups have been observed to a number of ^{20}Ne states: see Table 20.25 (1969AD02, 1970GU08, 1977EV01). Angular distributions have been measured at $E(^3\text{He}) = 2.8$ to 7.3 MeV (see (1972AJ02)) and at 18 (1977EV01) and 18.3 MeV (1975PE11).

32. $^{18}\text{O}(^{11}\text{B}, ^9\text{Li})^{20}\text{Ne}$ $Q_m = -10.027$

At $E(^{11}\text{B}) = 114$ MeV, $^{20}\text{Ne}^*(4.25, 8.9, 10.39, 15.43)$ [± 60 keV] are relatively strongly populated. $^{20}\text{Ne}^*(0, 1.63, 5.78, 7.17)$ are also excited (1973ST24). See also (1973SC1J).

Table 20.25: States of ^{20}Ne from $^{18}\text{O}(^3\text{He}, n)^{20}\text{Ne}$

E_x (MeV \pm keV)		L^a	$J^\pi; T$
(1977EV01)	(1970GU08)		
0	0	0	0^+
1.65 ± 15	1.63 ± 160	2	2^+
4.21 ± 30	4.22 ± 150	4	4^+
	4.96 ± 150		
5.71 ± 30	5.73 ± 120		
6.72 ± 90	6.72 ± 100		
7.15 ± 20			
	7.86 ± 100		
(8.74 ± 150)	8.79 ± 60		
	9.05 ± 60		
	9.98 ± 50		
10.24 ± 30	10.25 ± 50	2	$2^+; (1)$
10.83 ± 90	10.88 ± 50		
	11.27 ± 50		
11.48 ± 60		(0)	(0^+)
	11.59 ± 40		
12.21 ± 15	12.20 ± 30	2	2^+
	12.41 ± 30	0	0^+
	12.83 ± 30		
	13.10 ± 30	0	0^+
	13.34 ± 30		
	13.48 ± 30		
13.57 ± 20	13.63 ± 30	(2)	(2^+)
13.93 ± 30	13.88 ± 30	(2)	(2^+)
	14.22 ± 30		
(15.1)			
15.52 ± 15		(2)	($2^+; 1$)
16.01 ± 25		(2)	($2^+; 1$)
16.73 ± 10	16.730 ± 6^b	0	$0^+; T = 2$
(17.03)			

Table 20.25: States of ^{20}Ne from $^{18}\text{O}(^3\text{He}, n)^{20}\text{Ne}$ (continued)

E_x (MeV \pm keV)		L^a	$J^\pi; T$
(1977EV01)	(1970GU08)		
17.55 ± 10		(2)	(2 ⁺ ; 1)
17.91 ± 20		(0)	(0 ⁺)
19.33 ± 15			

^a From analysis of angular distributions at $E(^3\text{He}) = 3.1$ MeV (1970GU08), 18 MeV (1977EV01) and 18.3 MeV (1975PE11).

^b $\Gamma < 20$ keV. This state is reported by (1969AD02).

$$33. \ ^{18}\text{O}(^{12}\text{C}, ^{10}\text{Be})^{20}\text{Ne} \quad Q_m = -4.7816$$

At $E(^{12}\text{C}) = 46$ MeV angular distributions to $^{20}\text{Ne}^*(0, 1.63, 4.25)$ have been studied: the 2p spectroscopic factors are 0.58, 0.24 and 0.20, respectively (1975CO15).

$$34. \ ^{19}\text{F}(p, \gamma)^{20}\text{Ne} \quad Q_m = 12.8447$$

Over the range $E_p = 2.9$ to 12.8 MeV, the γ_0 and γ_1 yields are dominated by the E1 giant resonance ($\Gamma \approx 6$ MeV) with the γ_1 giant resonance displayed upward in energy. Strong, well correlated structure is observed with a characteristic $\Gamma \approx 175$ keV. Angular distributions taken over the energy range do not vary greatly with energy. They are incompatible with γ_0 and γ_1 coming from the same levels in ^{20}Ne (1967SE02). See also (1973AV1B).

The yield curve for 11.2 MeV γ -rays [from the decay of $^{20}\text{Ne}^*(11.23)$, $J^\pi = 1^+$; $T = 1$, to the ground state] displays a resonance at $E_p = 4.090 \pm 0.005$ MeV [$^{20}\text{Ne}^*(16.73)$]. The 11.2 MeV γ -rays are isotropic which is consistent with the presumed 0^+ character of this lowest $T = 2$ state in ^{20}Ne : $\Gamma_p\Gamma_\gamma/\Gamma \approx 0.5$ eV. Since Γ_p/Γ (from the elastic scattering) is ≈ 0.1 , $\Gamma_\gamma \approx 5$ eV (1967KU06). For $E_p = 5.65$ to 6.21 MeV, the γ_0 and γ_1 yields are not resonant but the yield of 10.6 MeV γ -rays is resonant at 5.879 ± 0.007 MeV [$\Gamma_{c.m.} = 9.5 \pm 3$ keV, $\Gamma_{p0}\Gamma_\gamma/\Gamma \approx 0.05$ eV; $\Gamma_\gamma \approx 0.3$ eV]. The 10.6 MeV γ -ray is due to the cascade decay of $^{20}\text{Ne}^*(18.43)$, $J^\pi = 2^+$; $T = 2$ via $^{20}\text{Ne}^*(12.22)$ to the 2^+ state at 1.63 MeV (1972KU24). (1976MA01) have determined the upper limits to the strengths of the transitions to various states of ^{20}Ne from the 0^+ and 2^+ $T = 2$ states: these are displayed in Table 20.19. No evidence is found for an isotensor transition amplitude (1976MA01).

Resonances observed in this capture reaction are displayed in Table 20.26. See also (1973GL1B, 1973GL1C, 1973HA1X, 1974AD1B, 1974HA1G, 1974HA1N) and (1977SC02, 1977SC08; theor.).

Table 20.26: Resonances in $^{19}\text{F}(p, \gamma)^{20}\text{Ne}$ ^a

E_p (keV)	Γ_{lab} (keV)	Γ_{γ_0} (eV)	Γ_{γ_1} (eV)	$^{20}\text{Ne}^*$ (MeV)	$J^\pi; T$	Refs.
340		< 0.07	0.28 ± 0.06	13.168		(1962KE03)
484		≈ 0.05	0.42	13.304		(1963BE19)
597 ± 1	30 ± 3	< 0.6	12	13.412		(1963BE19)
671 ± 1	6.0 ± 0.7	1.0×10^{-2}	2.2^{b}	13.482	1^+^{d}	A
874				13.675		(1955FA1A)
935				13.733		(1955FA1A)
980				13.775		(1955FA1A)
1091	0.8			13.881	2^-^{e}	A, (1975SU1E)
1280				14.060		(1955FA1A)
1320^{b}	4.0			14.098		A
1350				14.127		(1955FA1A)
1370				14.146		(1955FA1A)
1420	15.7			14.193		A
$4090 \pm 5^{\text{e}}$		$\Gamma_\gamma \approx 5 \text{ eV}^{\text{c}}$		16.728	$0^+; 2$	(1967KU06, 1976MA01)
$5879 \pm 7^{\text{d}}$	10 ± 3	$\Gamma_\gamma \approx 0.3 \text{ eV}^{\text{c}}$		18.427	$2^+; 2$	(1972KU24, 1976MA01)

A: See references listed for this state in (1972AJ02) and in Table 20.13 of (1959AJ76) [see (1955HU1A)].

^a See also Table 20.19.

^b Γ_γ to $^{20}\text{Ne}^*(4.97) = 0.12 \text{ eV}$ (1961GO21), 0.24 eV (1960KA18).

^c See text of reaction 34.

^d Decays predominantly to $^{20}\text{Ne}^*(1.63)$ via an M1 transition: see Table 20.19.

^e Based on non-isotropic distribution of γ -rays from $13.88 \rightarrow 4.97$ transition, which leads to odd parity for $^{20}\text{Ne}^*(13.88)$ assuming p-wave protons (1975SU1E; preliminary).

Table 20.27: Levels of ^{20}Ne from $^{19}\text{F}(p, p_0)^{19}\text{F}$ (1955BA1C)

E_p (keV)	Γ_{lab} (keV)	l	$J^\pi; T$	Γ_p/Γ	θ_p^2 (%)	$^{20}\text{Ne}^*$ (MeV)
340	2.9	0	1^+	0.016	3.8	13.168
483 ^b			1^+			13.303
598	37	1	2^-	0.0012	0.38	13.413
669	7.5	0	1^+	0.98	9.6	13.480
843	23	0	0^+	0.996	10.8	13.645
873	5.2	1	2^- ^a	0.21	1.5	13.674
935	8.0	0	1^+	0.17	0.44	13.733
1346	4.5	1	2^- ^a	0.067	0.07	14.123
1372	15	1	2^- ^a	0.17	0.52	14.148
1422	14.6	0	1^+	0.85	0.92	14.195
1694 ^c						14.453
1940 ^c		(0)	($0^+, 1^+$)			14.687
2030 ^c						14.772
4094 ± 3 ^d	2.1 ± 0.5	0	$0^+; 2$	0.062 ± 0.004		16.732
5879 ± 7 ^e	10 ± 3	2	$2^+; 2$	≈ 0.2		18.427

^a 1^- not excluded by elastic scattering alone.

^b (1963BE19).

^c (1956DE33).

^d (1967BL19, 1967KU06).

^e (1972KU24). Resonance also observed in p_1, p_3, p_4 and p_5 yields.

Table 20.28: Resonance parameters in $^{19}\text{F} + \text{p}$ (1955BA1C)

E_p (keV)	J^π	θ^2 ^a					
		p ₀	p ₁	p ₂	α_1	α_2	α_3
340	1 ⁺	3.8	< 15		18.8	1.0	7.2
598	2 ⁻	0.38	< 28	< 145	31	< 0.5	< 5.1
669	1 ⁺	9.6	0.6	< 0.4	0.26	0.005	0.27
843	0 ⁺	10.8	≈ 0.14	< 0.92			
873	2 ⁻	1.5	< 0.07	2.7	1.05 ^b	1.45 ^b	3.4
935	1 ⁺	0.44	5.0	< 0.8	3.3	0.34	2.3
1346	2 ⁻	0.07 ^b	0.92	0.24	0.36	0.21 ^b	2.1
1372	2 ⁻	0.52 ^b	1.93	0.56	1.7 ^b	0.34 ^b	0.86
1422	1 ⁺	0.92	0.56	< 0.11	total < 0.034		

^a p₀, p₁, p₂ represent transitions to $^{19}\text{F}(0)$, (0.1), (0.2). α_1 , α_2 , α_3 represent transitions to $^{16}\text{O}(6.1)$, (6.9), (7.1).

^b Assuming lowest possible values of l ; see (1957MA1A).

35. (a) $^{19}\text{F}(\text{p}, \text{p})^{19}\text{F}$

$$E_b = 12.8447$$

(b) $^{19}\text{F}(\text{p}, \text{p}')^{19}\text{F}^*$

The elastic scattering has been studied in the range $E_p = 500$ to 2000 keV by (1954DE1A, 1954PE1A, 1955BA1C, 1955WE1A, 1956DE33, 1963BE19). Parameters for the observed resonances are exhibited in Tables 20.27 and 20.28 taken mainly from (1955BA1C). Some unresolved structure is observed at $E_p = 900$, 1092 and 1137 keV, in addition to a broad structure near $E_p = 1700$ keV (1955WE1A). A sharp anomaly is observed in the elastic scattering at $E_p = 4.096 \pm 0.003$ MeV (1967BL19), 4.090 ± 0.005 MeV (1967KU06). It is an s-wave resonance corresponding to the 0⁺; $T = 2$ state at $E_x = 16.73$ MeV (1967BL19, 1967KU06). There is no indication of this resonance in the p₃, p₄ or p₅ yields (1967KU06). The amplitude of the $T = 1$ or $T = 0$ impurity in this state is $\approx 1.5\%$ (1967BL19). In the range $E_p = 5.65$ to 6.21 MeV, a single anomaly is seen in the elastic scattering at $E_p = 5.879 \pm 0.007$ MeV. The interference patterns show that the scattering is d-wave, corresponding to the excitation of the 2⁺; $T = 2$ state at $E_x = 18.43$ MeV (1972KU24). The 5.88 MeV anomaly also appears in the p₁, p₃, p₄ and p₅ yields (1972KU24). The parameters of these two $T = 2$ states are shown in Table 20.27. The elastic scattering has also been studied for $E_p = 4.2$ to 7.5 MeV by (1967TH06). The total reaction cross section is reported for $E_p = 24.9$ to 46.3 MeV by (1974MC19: 8 energies in that range).

Resonances for inelastic scattering involving $^{19}\text{F}^*(0.11)$ ($J^\pi = \frac{1}{2}^-$) and $^{19}\text{F}^*(0.197)$ ($J^\pi = \frac{5}{2}^+$) [p₁ and p₂] are listed in Table 20.29 (1955BA94, 1963BE19). In general the resonances observed

are identical with those reported from other $^{19}\text{F} + \text{p}$ reactions, although the relative intensities differ greatly. The p_2 scattering has been measured at $E_p = 5.6$ to 6.3 MeV (1967TH06), and at $E_p = 3.5$ to 7.0 MeV the yields of p_2, p_3, p_4 and p_5 show structures (1973BE1H; prelim.). The analyzing power has been measured for the p_0, p_2 and p_5 groups with polarized protons in the range 4 to 10 MeV: strong structures are observed for p_0 (1975EN1C, 1976EN1C; prelim. work; and R. Kaita, private communication). See also (1972AJ02, 1974HA1G) and ^{19}F .

$$36. \ ^{19}\text{F}(\text{p}, \text{n})^{19}\text{Ne} \qquad Q_m = -4.0207 \qquad E_b = 12.8447$$

Yield measurements have been reported for $E_p = 4.23$ to 11 MeV: see (1972AJ02) and (1972KU24: 5.78 to 5.96 MeV; σ_t, n_0, n_{1+2}). Observed resonances are displayed in Table 20.30. A narrow anomaly is reported in the n_0 and n_{1+2} yields at $E_p = 5.879 \pm 0.007$ MeV, corresponding to the 2^+ ; $T = 2$ state of ^{20}Ne at 18.43 MeV (1972KU24). See also ^{19}Ne .

$$37. \ ^{19}\text{F}(\text{p}, \alpha)^{16}\text{O} \qquad Q_m = 8.1137 \qquad E_b = 12.8447$$

Many resonances occur in this reaction. They are displayed in Tables 20.31, 20.32 and 20.33 depending on whether they are observed in the α_0 yield [Table 20.31], in the α_1 [or α_π] yield to $^{16}\text{O}^*(6.05)$ [Table 20.32] or in the α_2, α_3 and α_4 yields [or in the yield of the γ -rays from $^{16}\text{O}^*(6.13, 6.92, 7.12)$] [Table 20.33]. Resonances for α_0 and α_1 are required to have even J , even π or odd J , odd π , while the α_2, α_3 and α_4 resonances are all odd-even or even-odd, with the exception of the $T = 2$ resonance discussed below.

A listing of the earlier yield studies is given in (1972AJ02). A detailed discussion of the evidence leading to many of the J^π assignments shown in Table 20.33 is given in (1959AJ76). Recent measurements are reported by (1974CA22: $E_p = 0.6$ to 1.8 MeV; $\alpha_0 \rightarrow \alpha_4$), (1972HE39: 1.3 to 2.2 MeV; $\gamma_{6.13}, \gamma_{6.92}, \gamma_{7.12}$) and (1972KU24: 5.65 to 6.20 MeV; $\alpha_0 \rightarrow \alpha_5, \alpha_7 \rightarrow \alpha_9$). The width of the forbidden decay of the 1^+ state at 13.17 MeV [see Table 20.33] by $\alpha_0, \Gamma_{\gamma_0} \lesssim 7 \times 10^{-6}$ eV (1974KR1D; prelim. results). Anomalies are observed in $\alpha_0, \alpha_1, \alpha_2, \alpha_4$ and α_8 but not in $\alpha_3, \alpha_5, \alpha_7$ and α_9 corresponding to the formation of the 2^+ ; $T = 2$ state at 18.43 MeV [$E_p = 5.88$ MeV] (1972KU24). The analyzing power for $\alpha_0, \alpha_{1+2}, \alpha_3$ and α_4 have been measured for $E_p = 4$ to 10 MeV: strong structures are observed (1975EN1C, 1976EN1C; prelim. results; and R. Kaita, private communication). See also (1974HA1G, 1974LO1B, 1976CA1P), (1976GA1K; theor.), (1977CL1F; astrophys.) and ^{16}O in (1977AJ02).

$$38. \ ^{19}\text{F}(\text{p}, \ ^8\text{Be})^{12}\text{C} \qquad Q_m = 0.8599 \qquad E_b = 12.8447$$

The excitation curves show strong resonant behavior (cross sections up to 1.5 mb/sr) for $E_x = 15.3$ to 18.7 MeV, over which region 28 angular distributions have been measured. Twelve states with $J^\pi \leq 4^+$ have been observed. The strongly populated states are in better agreement with those reported in the (p, α_1) yield to $^{16}\text{O}^*(6.05)$ [$J^\pi = 0^+$] than those reported in the (p, α_0) yield

Table 20.29: Resonances in $^{19}\text{F}(p, p')^{19}\text{F}^*$ (1955BA94)

E_p (keV)	J^π	Γ_{lab} (keV)	Γ_{p_1} (eV)	Γ_{p_2} (eV)	$\theta_{p_1}^2$ ^a (%)	$\theta_{p_2}^2$ ^a (%)	E_x in ^{20}Ne (MeV)
340	1^+	2.9	< 0.5	< 0.1	< 15		13.168
483 ^b	1^+	2.2	< 1.3	< 1.2			13.303
598 ^b	2^-	37	< 100	< 60	< 28	< 145	13.413
669	1^+	7.5	46	< 0.5	0.6	< 0.4	13.480
720		≈ 30	< 10000	< 10000			13.528
780		≈ 10	< 400	≈ 9000			13.585
831		8.3	< 6	≈ 2300			13.634
845	0^+	23	≈ 50	< 10	≈ 0.14	< 0.92	13.647
873	2^-	5.2	< 2	570	< 0.07	2.7	13.674
900		4.8	< 30	≈ 2200			13.699
935	1^+	8.0	3000	< 20	5.0	< 0.8	13.733
1092		< 1.2					13.882
1137		3.7	< 40	≈ 2100			13.924
≈ 1250		≈ 80	≈ 70000	< 4000			14.03
1290		19	< 600	≈ 900			14.070
1346	2^-	4.5	300	600	0.92	0.24	14.123
1372	2^-	15	700	1400	1.93	0.56	14.148
1422	1^+	14.6 ± 1	2200	≤ 35	0.56	≤ 0.11	14.195
1610		≈ 5					14.374
1660							14.421
1700							14.459
5879 ^c	$2^+; 2$		r				18.427

r = resonant.

^a (1955BA1C).

^b (1963BE19).

^c (1972KU24). Resonance also observed in p_3 , p_4 and p_5 yields.

Table 20.30: Resonances in $^{19}\text{F}(p, n)^{19}\text{Ne}$

E_p (MeV)		Γ_{lab} (keV)	$^{20}\text{Ne}^*$ (MeV)
(1968RI08) ^a	(1963JE04) ^b	(1952WI1A)	
4.30		45	16.93
4.46		80	17.08
4.52		20	17.14
4.61		60	17.22
4.72		25	17.33
4.75		45	17.35
4.87			17.47
4.95	4.96	20	17.55
5.03	5.03		17.62
5.11	5.11		17.70
5.26			17.81
5.25	5.26		17.84
5.37	5.37		17.94
(5.44)			(18.01)
5.50			18.07
5.57			18.13
(5.62)			(18.18)
(5.69)			(18.25)
5.72	5.73		18.28
5.77			18.32
5.84			18.39
	5.879 ± 0.007^c	10 ± 3^c	18.427
5.90			18.45
6.00	6.03		18.54
	6.15		18.68
	6.35		18.87
	6.53		19.04
	6.81		19.31
	7.14		19.62
	7.27		19.75

Table 20.30: Resonances in $^{19}\text{F}(p, n)^{19}\text{Ne}$ (continued)

E_p (MeV)		Γ_{lab} (keV) (1952W11A)	$^{20}\text{Ne}^*$ (MeV)
(1968RI08) ^a	(1963JE04) ^b		
	7.41		19.88
	7.52		19.98
	7.74		20.19
	8.02		20.46
	8.15		20.58
	8.28		20.71
	8.37		20.79
	8.70		21.10
	8.82		21.22
	9.08		21.47
	9.2		21.6
	9.5		21.9
	9.8		22.1
	10.2		22.5

^a ± 5 keV.

^b ± 20 keV, except for the last four values.

^c (1972KU24): 2^+ ; $T = 2$.

Table 20.31: Resonances for ground-state α -particles (α_0) in $^{19}\text{F}(p, \alpha_0)^{16}\text{O}$


E_p (keV)	Γ_{lab} (keV)	θ_{α}^2 ^a (%)	J^{π}	$^{20}\text{Ne}^*$ (MeV)
400 ^b	100		1^-	13.225
400 ^b	100		0^+	13.225
650 ± 20 ^b	200		1^-	13.462
710 ^{a,b}	35	0.6	(1^-)	13.519
733	66	1.0	2^+	13.541
778	≈ 10	0.02	2^+	13.583
843	23	0.16 ^j	(2^+) ^g	13.645
≈ 860	120	2.1	1^-	13.66
≈ 930	≈ 180	2.9	0^+	13.73

Table 20.31: Resonances for ground-state α -particles (α_0) in $^{19}\text{F}(p, \alpha_0)^{16}\text{O}$ (continued)

E_p (keV)	Γ_{lab} (keV)	θ_{α}^2 ^a (%)	J^{π}	$^{20}\text{Ne}^*$ (MeV)
≈ 1080	≈ 200	3.4	1^-	13.87
1115	50	0.55	2^+	13.904
1160	≈ 70	1.1	0^+	13.946
1235 ^{a,c}	≈ 70	1.2	1^-	14.017
≈ 1250 ^a	≈ 150	2.7	2^+	14.03
1358 ^{a,c,d}	54	0.49	2^+	14.134
1640 ^c	< 115			14.402
1709 ^{c,d}	140		0^+	14.468
1853 ^{c,d}	132		1^-	14.604
2110 ^{c,d,e}	75		$(2^+, 4^+)$	14.85
2310 ^{c,d,e}	90		(2^+)	15.04
2550 ^e	300		(1^-)	15.27
2390 ^{c,d,h}	300		(0^+)	15.30
2680 ^{c,h}	80			15.39
2730 ^e	60			15.44
2820 ^e	160			15.52
2940 ^h				(15.64)
3120 ^h	170			(15.81)
3340	105			16.02
3680	(100)			16.34
3860				16.51
3980	135			16.62
(4090) ^k			$0^+; T = 2$	16.73
4130	100			16.77
4360	100			16.98
4460	95			17.08
4690	65			17.30
4900	90			17.50
4990	40			17.58
5879 ± 7 ^f	10 ± 3		$2^+; T = 2$	18.427

- ^a (1958IS10, 1958IS11).
^b (1959BR67).
^c (1958RA15).
^d (1957CL42).
^e (1964BR12).
^f (1972KU24): $\Gamma_{\alpha_0} \approx 0.3$ keV.
^g $J = 0$ from $^{19}\text{F}(p, p)^{19}\text{F}$; possibly $T = 1$ (1955BA94, 1955BA1C).
^h See, however, (1964BR12).
ⁱ (1967KU06).
^j $\Gamma_{\alpha_0} \approx 0.06$ keV (1974CA22).
^k See (1972AJ02): not published.

(1969GO1B, 1971GO1R). It is suggested that most of the observed states are of 8p-4h and 12p-8h configurations (1971GO1R). [(1969GO1B, 1971GO1R) are preliminary reports.]



For reaction (a) see ^{17}F in (1977AJ02) and (1974NE03). The polarization analyzing powers for the reaction to $^{17}\text{O}^*(0, 0.87)$ have been measured at $E_p = 49.5$ MeV (1974NE03).



Levels of ^{20}Ne derived from reported neutron groups are displayed in Table 20.34. Angular distributions have been measured at $E_d = 0.5$ to 6.1 MeV: see (1972AJ02) and at $E_d = 8$ MeV (1976BE2E: n_1, n_6). The branching ratio for the γ -decay of the lowest 1^+ ; $T = 1$ state [$^{20}\text{Ne}^*(11.23)$] to the ground state to $^{20}\text{Ne}^*(1.63)$ is 0.53 ± 0.07 (1973KU1E: $E_d = 5.5$ MeV). See also (1975LE1K; applied) and (1972AJ02).



Levels of ^{20}Ne observed in this reaction are displayed in Table 20.35. Deuteron angular distributions have been measured at $E(^3\text{He}) = 9.5$ and 10.0 MeV (1965SI18), 13.0 MeV (1963JA01), 16 MeV (1974VE03, 1976SE10), 18 MeV (1975BE02, 1976FO05) and 21 MeV (1973OB04). Spectroscopic factors obtained by (1973OB04, 1975BE02, 1976FO05, 1976SE10) are shown in Table 20.35. The angular distribution of the deuterons to $^{20}\text{Ne}^*(4.25)$ is explained in terms of inelastic effects (1974VE03). See also (1965SI18) and (1977SU04; theor.).

Gamma-ray measurements are summarized in Tables 20.19 (1977MA07), 20.20 (1969AN08) and 20.35 (1969AN08, 1977MA07). See also (1972FO16, 1973FO1A, 1973FO1E, 1974FO1J).

Table 20.32: Nuclear pair resonances (α_π) in $^{19}\text{F}(\text{p}, \alpha_\pi)^{16}\text{O}$

E_p (keV)	Γ_{lab} (keV)	σ^c (mb)	θ_α^2 (%)	J^π	$^{20}\text{Ne}^*$ (MeV)
710 ^a	35	≈ 0.2	2	1^-	13.519
780	≈ 10	≈ 0.2	0.15	2^+	13.585
842	23	3.4	0.27	2^+ ^d	13.644
1115	50	1.5	3.6	2^+	13.904
1236 ^{a,b}	≈ 70	3	1.0	1^-	14.018
1367 ^{a,b}	30	6.0	0.29	2^+	14.143
1630 ^b	60				14.39
1720	95	≈ 18			14.48
1880	170				14.63
2170	95				14.91
2330	≈ 100				15.06
2600	100				15.31
2680	100				15.39
2820	125				15.52
3120	145				15.81
3340	100				16.02
(3500)	(80)				(16.17)
(3590)	(115)				(16.25)
3960	200				16.60
4360	95				16.98
4690	< 150				17.30
4900	115				17.50
4990	40				17.58
5170	220				17.75

^a See (1972AJ02).

^b (1958RA15).

^c (1958IS11).

^d See footnote ^g in Table 20.31.

Table 20.33: Resonances for 6 – 7 MeV γ -rays ($\alpha_2, \alpha_3, \alpha_4$) in $^{19}\text{F}(p, \alpha)^{16}\text{O}$

E_p^\dagger (keV)	Γ_{lab} (keV)	Γ_{α_2} (eV)	Γ_{α_3} (eV)	Γ_{α_4} (eV)	J^π	$^{20}\text{Ne}^*$ (MeV)
$226.9 \pm 3.4^{a,b}$	1.0	1000	< 2.5	< 2.5	2^-	13.060
$340.46 \pm 0.04^{c,d}$	2.4 ± 0.2	2800	16	75	1^+	13.1680
483.8 ± 0.3^e	0.9 ± 0.1	700	19	190	1^+	13.3038
$872.11 \pm 0.20^{f,g}$	4.7 ± 0.2	2200	620	180	2^-	13.6729
935.4 ± 1.3^g	8.1 ± 0.5	2900	110	720	1^+	13.733
$1347.7 \pm 1.0^{h,m}$	4.9 ± 0.7	2250	650	1200	2^-	14.124
$1373.0 \pm 1.0^{i,m}$	12.4 ± 1.0	6650	700	300	2^-	14.148
$1694 \pm 1.7^{j,m}$	35 ± 3					14.453
$1949 \pm 2.5^{j,m}$	40 ± 10					14.695
$2030 \pm 3.0^{j,m}$	120 ± 20					14.722
2320 ^k	85					15.05
2510	30					15.23
2630	90					15.34
2800	60					15.50
3020	30					15.71
3190	80					15.87
3490	40					16.16
3920	30					16.57
4000	110					16.64
4090 ^l					$0^+; T = 2$	16.73
4290	50					16.92
4490	30					17.11
4570	30					17.18
4710	30					17.32
4780	35					17.38
4990	20					17.58
5070	35					17.66
5200	70					17.78

- ^a (1959KU79; assignment to this reaction probable but not certain). See also (1959AJ76).
- ^b (1962KE03). $\Gamma_{\alpha_0} < 100$ eV.
- ^c (1955BA94, 1959BO14, 1964SE1A, 1974KR1D). $\Gamma_{\alpha_0} \lesssim 7 \times 10^{-6}$ eV (1974KR1D).
- ^d (1950AR1A, 1950BA1A, 1950CH1A). Values listed for E_p and Γ are those recommended by (1966MA60).
- ^e (1959BO14, 1959KU79, 1960HU11, 1963BE19). $\Gamma_{\alpha_0} < 25$ eV (1963BE19).
- ^f Values listed for E_p and Γ are those recommended by (1966MA60). Other values are $E_{\text{res}} = 872.4 \pm 0.4$ keV (1959BO14), 873.5 ± 0.7 keV (1960HU11), 872.3 ± 0.5 keV (1961BE13), 871.80 ± 0.25 keV (1962RY01), 872.5 ± 1.1 keV (1965AS07). See also (1974CA22).
- ^g (1965AS07).
- ^h Other reported resonance energies are $E_p = 1344.5 \pm 1.0$ keV (1959LI51), 1347.7 ± 1.0 keV (1960HU11), 1347.7 ± 1.8 keV (1965AS07).
- ⁱ Other reported resonance energies are $E_p = 1373.0 \pm 1.0$ keV (1959LI51), 1373.7 ± 1.2 keV (1960HU11), 1374.5 ± 1.8 keV (1965AS07).
- ^j (1952WI1A, 1955HU1A).
- ^k (1952WI1A); these values should be reduced by about 0.2%: see (1955KI28).
- ^l Resonance in α_2 yield: see text (1967KU06).
- ^m See also (1972HE39). The 1.69 MeV resonance may correspond to more than one state.

[†] *Footnote added in proof.* A number of other resonances shown in Table 20.13 of (1959AJ76) are not listed here because questions arose about their identification. However, (1978GO1F) report the observation of resonances at $E_p = 597, 672, 835, 902, 1090, 1140, 1189$ and 1283 keV: see (1959AJ76). I am indebted to F.C. Young for his comments.

$$42. \ ^{19}\text{F}(\alpha, t)^{20}\text{Ne} \quad Q_m = -6.9694$$

At $E_\alpha = 28.5$ MeV angular distributions are reported for the tritons to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62 + 5.78, 7.00 + 7.17 + 7.19)$ (1967HA23; DWBA). The distributions of the tritons to $^{20}\text{Ne}^*(0, 1.63, 4.25)$ [$J^\pi = 0^+, 2^+$ and 4^+ , respectively] have been reanalyzed in terms of the collective-model coupled-channels Born-approximation theory: C^2S for these three states are 0.08, 0.16 and 0.0 (CCBA) [the DWBA results are nearly the same]. Agreement with the values obtained in the (d, n) and (^3He , d) reactions is poor (1974OB02). Angular distributions have also been reported at $E_\alpha = 18.5$ and 28.4 MeV: see (1972AJ02).

$$43. \ ^{19}\text{F}(^7\text{Li}, ^6\text{He})^{20}\text{Ne} \quad Q_m = 2.867$$

Angular distributions have been studied at $E(^7\text{Li}) = 34$ MeV for the transitions to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97$ (partial), $5.62, 5.78, 6.72, 7.1, 7.42)$. The spectroscopic factors, C^2S , for $^{20}\text{Ne}^*(0, 1.63, 4.25, 5.78, 6.72, 7.42)$ are 0.36, 0.54, 0.06, 0.20 and 0.22, respectively, in good agreement with those reported in the (d, n) and (^3He , d) reactions (1975WI30).

Table 20.34: Neutron groups from $^{19}\text{F}(\text{d}, \text{n})^{20}\text{Ne}$ ^a

E_x (MeV \pm keV)		l_p ^a	$J^\pi; T$
(1958MO02, 1968LA03) ^b	(1969RI01)		
0		0	0^+
1.74 \pm 30		2	2^+
4.20 \pm 40			
4.96 \pm 50			
5.62 \pm 40			
6.80 \pm 10		0	0^+
7.16 \pm 90			
7.41 \pm 50			
7.90 \pm 40			
(8.71 \pm 10)			
9.15 \pm 40			
(9.50 \pm 40)			
10.01 \pm 30			
10.30 ^c			
	^d		
	10.59		
10.853 \pm 10	10.879 \pm 40	2	$T = 1$ ^f
	11.03 \pm 80 ^{d,h}		
11.233 \pm 10	11.26 \pm 40	0	$1^+; (1)$ ^{f,g}
11.549 \pm 10	11.568 \pm 35	2	$(T = 1)$ ^f
	11.915 \pm 30		
12.086 \pm 10		e	$(T = 1)$ ^f
12.150 \pm 10		e	$(T = 0)$ ^f
	12.179 \pm 25		
12.200 \pm 10		e	$(T = 1)$ ^f
12.245 \pm 10		2	$T = 1$ ^f
12.379 \pm 10	12.397 \pm 20	0	$T = 0$ ^f
	13.086 \pm 15		
	13.170 \pm 15	0	$1^+; (1)$ ^g
	13.481 \pm 15	0	$1^+; 1$ ^g
	13.650 \pm 15	0	$(0^+); 1$ ^g

Table 20.34: Neutron groups from $^{19}\text{F}(d, n)^{20}\text{Ne}$ ^a (continued)

E_x (MeV \pm keV)		l_p ^a	$J^\pi; T$
(1958MO02, 1968LA03) ^b	(1969RI01)		
	13.882 ± 15		

^a See also Table 20.31 in (1972AJ02).

^b States below 10.1 MeV are from (1958MO02) who also report evidence for some other states; higher states are from (1968LA03). Data of (1968LA03) are adjusted downward by 26 keV: see (1969RI01).

^c $E_x = 10.31 \pm 0.07$ (1964SA09), 10.33 ± 0.05 MeV (1966RI05).

^d This state decays to $^{20}\text{Ne}^*(1.63)$ (1966RI05).

^e Weak group.

^f (1968LA03).

^g (1969RI01).

^h A study of $^{19}\text{F}(d, n)$ thresholds by (1960BU07) suggested states at $E_x = 11.08, 11.31, 11.66, 11.84, 12.16, 12.24$ and 12.48 MeV [± 20 keV]: see Table 20.32 in (1972AJ02).

 Table 20.35: States of ^{20}Ne from $^{19}\text{F}(^3\text{He}, d)^{20}\text{Ne}$

E_x ^a (MeV \pm keV)	Γ ^a (keV)	nlj ^{a,b,c}	$J^\pi; T$ ^{a,b}	K^π ^d	$(2J + 1)C^2S$		
					(1975BE02) ^e	(1976SE10) ^f	(1973OB04) ^f
0 ^b		$2s_{1/2}$	0^+	0_1^+		0.30	0.43
1.6353 ± 1.8 ^h		$1d_{5/2}$	2^+	0_1^+		1.43	1.90
4.249 ± 2.5 ^h		n.s.	4^+	0_1^+		0.08	0.0
4.968 ± 3 ^h		$1p_{3/2}$	2^-	2^-	(0.03)	0.021	0.040
5.623 ± 3 ^h		$1f_{7/2}$	3^-	2^-	(0.09)	0.10	0.028
5.785 ± 3		$2p_{3/2}$	1^-	0^-	0.16	0.10	0.12
6.722 ± 3		$2s_{1/2}$	0^+	0_2^+	0.52	0.39	0.22
7.00 ^b		$1f_{7/2}$	4^-	2^-		0.12	0.11
7.156 ± 8		$1f_{7/2}$	3^-	0^-	0.42	0.12	
7.422 ± 3		$1d_{5/2}$	2^+	0_2^+	0.79	0.39	0.60
7.829 ± 10		$1d_{5/2}$	2^+	0_3^+	0.06	0.046	0.045
≈ 8.3	≈ 800	$2s_{1/2}$	0^+	0_4^+	0.13		
8.45 ^b		n.s.	5^-	2^-			
8.70 ^b		n.s.	1^-				
8.769 ± 10		n.s.	6^+	0_1^+			

Table 20.35: States of ^{20}Ne from $^{19}\text{F}(^3\text{He}, d)^{20}\text{Ne}$ (continued)

E_x^a (MeV \pm keV)	Γ^a (keV)	$nlj^{a,b,c}$	$J^\pi; T^{a,b}$	K^π^d	$(2J+1)C^2S$			
					(1975BE02) ^e	(1976SE10) ^f	(1973OB04) ^f	
8.8 ^g	broad	$1d_{5/2}$	2^+	0_3^+	0.21 ^g			
8.841 ± 10		$2p_{3/2}$	1^-		(0.01)			
9.03 ^b		n.s.	4^+					
9.12 ^b		n.s.	3^-					
9.305 ± 10		$1d_{5/2}$	$(1, 2, 3)^+$		0.04			
9.469 ± 10		$1d_{5/2}$	2^+		0.03			
9.859 ± 3		$1d_{5/2}$	$3^+ i$		2.37			
9.92 ^b		n.s.	(1^+)					
9.99 ^b		n.s.	4^+			0_2^+		
10.257 ± 15		$1d_{5/2}$	$2^+; 1$		0.07			
10.40 ^b	27							
10.55 ^b								
10.568 ± 15		$1d_{5/2}$	2^+	0.05				
10.815 ± 15		12	$1d_{5/2}$	2^+	0.05			
10.860 ± 15		$1d_{5/2}$	$3^+; 1^i$	2.82				
10.951 ± 15								
11.067 ± 15		n.s.	$(4^+; 1)$					
11.239 ± 15					see ^a			
11.27 ± 15		73	n.s.					
11.549 ± 15		$1d_{5/2}$	$3^+ i$	1.00				
11.83 ± 15	81	$1d_{5/2}$		0.10				
11.992 ± 15	n.s.	(8^+)		0_1^+				
12.082 ± 15	$1d_{5/2}$			0.35				
12.190 ± 15	$< 0.1^i$	$1d_{5/2}$	$(1, 2, 3)^i$	2.10				
12.367 ± 15^j	$< 200^i$		$3^- i$	see ^{a,i}				
12.423 ± 15	160	$1d_{5/2}$	(2^+)	0.19				
12.503 ± 15	$1d_{5/2}$			0.02				
12.823 ± 15	$2s_{1/2}$			0.15				
13.037 ± 15	$1d_{5/2}$							
13.135 ± 15								
13.270 ± 15								

n.s. = not stripping.

^a (1975BE02): $E(^3\text{He}) = 18$ MeV. The E_x measured by (1975BE02) appear to be systematically low by 14 – 30 keV: see (1977MA07).

^b (1976SE10): $E(^3\text{He}) = 16$ MeV.

^c Orbital for direct transfer.

^d (1973OB04): $E(^3\text{He}) = 21$ MeV.

^e DWBA.

^f CCBA.

^g (1976FO05): $E(^3\text{He}) = 18$ MeV.

^h Gamma-ray measurements (1969AN08).

ⁱ Gamma-ray measurements (1977MA07): $E_x = 9.88 \pm 0.03, 10.89 \pm 0.03, 11.59 \pm 0.03, 12.22, 12.40 \pm 0.04$ MeV.

^j α -decays to $^{16}\text{O}^*(6.13)$ (1977MA07).

$$44. \text{}^{19}\text{F}(^{10}\text{B}, \text{}^9\text{Be})^{20}\text{Ne} \quad Q_m = 6.259$$

See (1973ME1F).

$$45. \text{}^{20}\text{F}(\beta^-)^{20}\text{Ne} \quad Q_m = 7.0259$$

The decay is principally to $^{20}\text{Ne}^*(1.63)$ with a half-life of 11.00 ± 0.02 sec: see reaction 1 in ^{20}F and Table 20.36 for the branching to various ^{20}Ne states. The 0.02% branching to $^{20}\text{Ne}^*(4.97)$ [$J^\pi = 2^-$] is consistent with the assignment $J^\pi = 2^+$ to the ground state of ^{20}F (1969GA05), as are measurements of the β - γ circularly polarized correlation (1961FR02, 1965MA28). When the β - γ correlation is described by $W(\theta) = 1 + p \cos^2 \theta$, p has a value slightly different from zero and it varies linearly with E : $dp/dE = +0.09 \pm 0.05\%$ MeV^{-1} (1977MC1J). (1977TR1H) report $dp/dE = +0.05 \pm 0.07\%$ MeV^{-1} and (1977RO1Y) find $+0.02 \pm 0.02$ MeV^{-1} . The β -decay asymmetry, $(ft)^+/(ft)^- = 1.044 \pm 0.010$ (1975AL27), 1.025 ± 0.013 (1976GE08): see also reaction 58 and (1974ST10). (1977RO1Y) find, by comparison with ^{20}Na , a vanishing second class current: the ratio of the second-class axial matrix element to the main first-class one is 1.1 ± 1.5 and a first-class second-forbidden axial contribution is 40 ± 8 .

The energy of the γ -ray from $^{20}\text{Ne}^*(1.63)$ is 1634.8 ± 0.6 keV (1967VA08), 1632.6 ± 0.8 keV (1966AL12), 1633.7 ± 0.3 keV (1968SP01), 1633.6 ± 0.3 keV (1972OP01). The γ -ray from the $(4.97 \rightarrow 1.63)$ transition has $E_\gamma = 3334.3 \pm 0.7$ keV, $\Delta E_x = 3334.6 \pm 0.7$ keV, and using the E_x for $^{20}\text{Ne}^*(1.63)$ shown in Table 20.18, E_x for $^{20}\text{Ne}^*(4.97) = 4968.4 \pm 0.8$ keV (1969GA05).

$$46. (a) \text{}^{20}\text{Ne}(\gamma, n)^{19}\text{Ne} \quad Q_m = -16.8653$$

- (b) $^{20}\text{Ne}(\gamma, 2n)^{18}\text{Ne}$ $Q_m = -14.419$
(c) $^{20}\text{Ne}(\gamma, p)^{19}\text{F}$ $Q_m = -12.8447$
(d) $^{20}\text{Ne}(\gamma, \alpha)^{16}\text{O}$ $Q_m = -4.7309$

Table 20.36: Branching in $^{20}\text{F}(\beta^-)^{20}\text{Ne}$

Decay to $^{20}\text{Ne}^*$ (MeV)	J^π	Branch (%)	$\log ft^a$	Refs.
0	0^+	< 0.032		(1954WO23)
1.63	2^+	99.98	4.97	(1975AL27)
4.25	4^+	< 0.015		(1969GA05)
4.97	2^-	0.017 ± 0.003	6.88	(1969GA05)
5.62	3^-	< 0.048		(1963GL01)
5.78	1^-	< 0.1		(1963GL01)
6.72	0^+	< 0.67		(1963GL01)
7.00	4^-	< 0.2		(1963GL01)

^a Based on $\tau_{1/2} = 11.00 \pm 0.02$ sec (1975AL27).

Bremsstrahlung measurements show peaks in the neutron yield (reaction (a)) at 17.8, 18.8, 19.1, 20.1, (22.0), (23.0) and (24.8) MeV [numbers in parentheses refer to relatively weak and broad structures]. The giant dipole resonance is centered at ≈ 20 MeV and the integral cross section to 28 MeV exhausts half of the dipole sum rule (1975WO06). The cross section for (γ, Tn) using monoenergetic photons shows a structure at 18 MeV and some fluctuations atop the broad giant dipole resonance, $\sigma_{\text{max}} \approx 7$ mb. The double photoneutron cross section, $\sigma(\gamma, 2n)$, is dominated by a single peak at ≈ 20.5 MeV, $\sigma_{\text{max}} \approx 1.1$ mb (1974VE06). See also (1972AJ02, 1975BR1F, 1975WO06, 1977DA1B) and (1973ROYN, 1975SC05; theor.). For reaction (c) see (1962DO1A, 1963FI1B, 1969HO16) and (1975SC2B; theor.). For reaction (d) see (1959HA1C) and reaction 47.

47. (a) $^{20}\text{Ne}(e, e)^{20}\text{Ne}$
(b) $^{20}\text{Ne}(e, \text{ep})^{19}\text{F}$ $Q_m = -12.8447$
(c) $^{20}\text{Ne}(e, e\alpha)^{16}\text{O}$ $Q_m = -4.7309$

The ^{20}Ne charge radius, $r_{\text{rms}} = 3.116 \pm 0.025$ fm (using a Born approx.) (1971MO15). See also the discussion in (1973SI31).

Form factors for many of the excited states of ^{20}Ne with $E_x < 8$ MeV have been reported: see (1973SI31; $E_e = 77.0, 81.3, 102.4, 110.5, 114.4$ MeV) and (1972MI06; 200 and 250 MeV). The monopole matrix elements for the $0^+ \rightarrow 0^+$ transitions are 7.4 ± 2.0 and 6.9 ± 1.4 fm² for $^{20}\text{Ne}^*(6.72, 7.19)$ (1972MI06) while (1973SI31) calculate $\Gamma_\pi = 3.9 \times 10^{-5}$ eV for $^{20}\text{Ne}^*(6.72)$. $B(E2)\uparrow$ is 0.028 ± 0.004 e² · b² for the transition to $^{20}\text{Ne}^*(1.63)$. The transitions to $^{20}\text{Ne}^*(4.25, 7.83)$ correspond to 23 ± 5 W.u. and 0.77 ± 0.25 W.u. (1973SI31). (1972MI06) report $B(E2)\uparrow = 0.13 \pm 0.03$ and 0.83 ± 0.13 W.u. for the transitions to $^{20}\text{Ne}^*(7.42, 7.83)$.

At $E_e = 39$ and 56 MeV, the 180° inelastic scattering is dominated by the transition to a $J^\pi = 1^+$; $T = 1$ state at $E_x = 11.22 \pm 0.05$ MeV with $\Gamma_{\gamma_0} = 11.2_{-1.8}^{+2.1}$ eV. A subsidiary peak is observed corresponding to a state at an $E_x = 0.35 \pm 0.03$ MeV higher [if $J^\pi = 1^+$ or 2^+ , $\Gamma_{\gamma_0} = 0.65 \pm 0.18$ or 0.40 ± 0.13 eV]. A number of small peaks are also reported corresponding to $E_x \approx 12.0, 12.9, 13.9, 15.8, 16.9, 18.0$ and 19.0 MeV ((1971BE18), and W.L. Bendel, private communication). See also (1975CH41, 1975SZ1B).

A study of reaction (b) at $E_e = 30$ MeV shows strong resonances (assuming ground-state transitions) at $E_x = 17.70, 18.87, 19.87$ and 21.02 MeV, respectively, as well as some weaker structures (1962DO1A).

Reaction (c) has been studied in order to obtain the (γ, α_0) cross section in the giant resonance region: the cross section at 90° for $E_x = 15$ to 24 MeV is dominated by an E1 resonance [1^- ; $T = 1$, with an admixture of $T = 0$ which permits the α_0 decay] at $E_x = 20$ MeV; lesser E1 structures are reported at $E_x = 16.7, 17.1, 21$ and 22 MeV. A relatively strong 2^+ ; $T = 0$ resonance appears at $E_x = 18.5$ MeV, and evidence is reported for increasing E2 strength below 16 MeV (1975SK06).

See also (1972RI1B, 1972THZF, 1973TH1B, 1974DE1E, 1974HA1G, 1975BE1G, 1975FA1A, 1976HA1Q) and (1972HA25, 1972HI17, 1972HO1D, 1972SA1C, 1973AN30, 1973CH1F, 1973CU03, 1973HO41, 1973LE1D, 1974AB05, 1974DZ05, 1974HA1C, 1975AB04, 1975DO10, 1976FU1K, 1976NO10, 1977SU04; theor.).

48. $^{20}\text{Ne}(n, n)^{20}\text{Ne}$

See ^{21}Ne in (1978EN06) and (1976ZA1E; theor.).

49. (a) $^{20}\text{Ne}(p, p)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(p, p\alpha)^{16}\text{O}$ $Q_m = -4.7309$

(c) $^{20}\text{Ne}(p, 2p)^{19}\text{F}$ $Q_m = -12.8447$

Angular distributions of elastically scattered protons and of a number of inelastic groups have been measured for $E_p = 2.15$ to 41.8 MeV: see (1972AJ02) for the earlier work and (1969MA48: 3.66 to 5.90 MeV; p_0, p_1), (1976DE12, 1976SW1C: 24.5 MeV; p to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62, 5.78, 6.72, 7.00, 7.17 + 7.19, 7.42, 7.83, 8.45, 8.69, 8.78, 9.03, 9.12, 9.31 + 0.02, 9.51)$),

(1973DE06, 1974DE46: 30 MeV; p_0, p_1, p_2) and (1977CO1G: 35.2 MeV; p_0). The $E_p = 24.5$ and 30 MeV angular distributions for the $0^+, 2^+$ and 4^+ members of the ground-state $K^\pi = 0^+$ band are well fitted using coupled-channels calculations and deformation parameters of $\beta_2 = +0.47 \pm 0.04$ and $\beta_4 = +0.28 \pm 0.05$. When the 6^+ state is included [$^{20}\text{Ne}^*(8.78)$], the fit is improved if $\beta_6 = -0.10$ is included (1973DE06, 1974DE46, 1976DE12, 1976SW1C). The state at $E_x = 9.31 \pm 0.02$ MeV is suggested to have $J^\pi = (4^+)$ and to be the 4^+ member of the $K^\pi = 0^+$ band based on $^{20}\text{Ne}^*(6.72)$: $\beta_2 = 0.15$ (1976SW1C): see, however, Table 20.35.

For $p\text{-}\gamma$ angular correlations see (1972AJ02). See also (1969MA48). For polarization measurements see (1972DE18, 1976DE12, 1977PL1C) and ^{21}Na in (1978EN06). For yield measurements see (1972HI1C, 1976SW1C) and ^{21}Na in (1978EN06). For reaction (b) see (1971EP03). See also (1970QU1C). For a study of spallation see (1974PA10). See also (1972AS13, 1976AM04, 1976ES1B; theor.). For reaction (c) see (1964KA1A; theor.).

50. $^{20}\text{Ne}(d, d)^{20}\text{Ne}$

Angular distributions have been reported at $E_d = 10.95$ to 11.8 MeV and at 52 MeV [see (1972AJ02)], at $E_d = 10.0$ to 12.0 MeV (1975DA1F; d_1), 11.6 MeV (1973BR15; d_0) and 11.66 MeV (1975DA1F; d_2) and at $E_d = 40$ MeV (1974EP1B; d_1 and d_2). See ^{22}Na in (1978EN06) for polarization and yield measurements. See also (1971SI24; theor.).

51. $^{20}\text{Ne}(t, t)^{20}\text{Ne}$

The elastic scattering has been studied at $E_t = 1.80$ and 2.00 MeV (1969HE08). See also (1972SC10; theor.).

52. $^{20}\text{Ne}(^3\text{He}, ^3\text{He})^{20}\text{Ne}$

Angular distributions have been measured at $E(^3\text{He}) = 10$ to 35 MeV [see (1972AJ02)] and at 68 MeV (1974DE05: ^3He to $^{20}\text{Ne}^*(0, 1.63, 4.25)$). The angular distributions at $E(^3\text{He}) = 68$ MeV are well fitted by a coupled channels calculation with $\beta_2 = 0.47$ and $\beta_4 = 0.17$ (1974DE05). See also (1975MA25, 1976MA36; theor.).

53. (a) $^{20}\text{Ne}(\alpha, \alpha)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(\alpha, 2\alpha)^{16}\text{O}$ $Q_m = -4.7309$

(c) $^{20}\text{Ne}(\alpha, ^{12}\text{C})^{12}\text{C}$ $Q_m = -4.6181$

Angular distributions have been measured to low-lying states of ^{20}Ne at $E_\alpha = 3.8$ to 155 MeV: see (1972AJ02) and (1975AB1H, 1975DA1E: α_0 ; 3.8 to 8 MeV), (1977EN01: α_0 ; 20.2 – 23.0 MeV), (1975CO1K: α_0 ; 26.6, 27.2 and 27.8 MeV), (1972RE05: $\alpha_0, \alpha_1, \alpha_2$; 104 MeV) and (1976KN05: α_1 and α to $^{20}\text{Ne}^*(18.1 - 28.2)$; 155 MeV). A coupled-channel analysis of the work at $E_\alpha = 104$ MeV leads to $\beta_2 = +0.35 \pm 0.01$, $\beta_4 = +0.11 \pm 0.01$, $Q_{20} = +0.46 \pm 0.02$ b and $Q_{40} = +0.026 \pm 0.002$ b² (1972RE05). At $E_\alpha = 155$ MeV (1976KN05) find that the strength concentrated in the giant quadrupole resonance exhausts more than 30% of the isoscalar energy weighted sum rule. See also (1975MO04, 1976YO02). Yield measurements are reported by (1974BA76: $E_\alpha = 10.25$ to 14.82 MeV; most α -groups to $^{20}\text{Ne}^*$ states with $E_x < 8$ MeV) and by (1975AB1H, 1975CO1K, 1975CO1M, 1975DA1E). See also (1974HA1C, 1976HA1Q) and (1971FE1F, 1972MA65, 1972RE01, 1972SH09, 1973RE11, 1974PI11, 1974MA11, 1975GI10, 1976BE1P, 1976MA19, 1976MA36, 1977KN02; theor.).

For reaction (b) see ^{16}O in (1977AJ02), (1974EP01: $E_\alpha = 78.6$ MeV) and (1977WA1M: $E_\alpha = 140$ MeV). See also (1974MU1D), (1975RO1B) and (1975MI11; theor.). For reaction (c) see (1963LA08). See also ^{24}Mg in (1978EN06).

54. $^{20}\text{Ne}(^7\text{Li}, ^7\text{Li})^{20}\text{Ne}$

The elastic angular distribution has been measured at $E(^7\text{Li}) = 36$ MeV (1976CO23).

55. $^{20}\text{Ne}(^{12}\text{C}, ^{12}\text{C})^{20}\text{Ne}$

Elastic angular distributions have been obtained at $E(^{12}\text{C}) = 22.2, 27.2, 32.3, 37.3$ and 42.7 MeV [see (1974VA18)] and at $E(^{20}\text{Ne}) = 65.7$ MeV (1975DO06). For a fusion study, see (1977CO1Q). See also (1977PR1F) and (1976VA12, 1977OS02; theor.).

56. $^{20}\text{Ne}(^{16}\text{O}, ^{16}\text{O})^{20}\text{Ne}$

Angular distributions have been studied at $E(^{20}\text{Ne}) = 50$ MeV (1976ST18) and 94.8 MeV (1977MO1H) involving $^{16}\text{O}_{\text{g.s.}}$ and $^{20}\text{Ne}^*(0, 1.63, 4.25)$: qualitative agreement is found with the assumption of an α -cluster exchange process dominating at backward angles (1976ST18). See also (1975ZI1C).

57. $^{20}\text{Ne}(^{32}\text{S}, ^{32}\text{S})^{20}\text{Ne}$

The static quadrupole moment of $^{20}\text{Ne}^*(1.63)$, $Q_{20} = +0.94 \pm 0.38$ b (1969SC08).

Table 20.37: Decay of ^{20}Na

Decay to $^{20}\text{Ne}^*$ (MeV \pm keV)	$J^\pi; T$	Branching ratio (%) ^a		log ft
		(1973TO08)	(1976IN06) ^d	
1.633 ± 2 ^b	$2^+; 0$	79.47 ± 1.57	79.18 ± 1.58	4.988 ± 0.009 ^f
7.415 ± 5 ^e	$2^+; 0$	16.37 ± 1.28		4.19 ± 0.05
7.826 ± 7 ^e	$2^+; 0$	0.674 ± 0.055		5.417 ± 0.033
8.82 ± 10 ^e		0.034 ± 0.007		6.27 ± 0.08
9.481 ± 7 ^e	$2^+; 0$	0.247 ± 0.020		5.064 ± 0.034
9.873 ± 5 ^b	$3^+; 0$		0.0272 ± 0.0138	5.78 ± 0.18 ^f
10.274 ± 3 ^{b,c}	$2^+; 1$ ^g	2.89 ± 0.23	2.944 ± 0.224	3.471 ± 0.033 ^f
10.584 ± 7 ^e	$2^+; 0$	0.087 ± 0.009		4.76 ± 0.05
10.848 ± 7 ^e	$2^+; 0$	0.193 ± 0.016		4.179 ± 0.035
10.884 ± 3 ^b	$3^+; 1$		0.0392 ± 0.0139	4.84 ± 0.13 ^f
11.261 ± 5 ^b	$1^+; 1$		0.203 ± 0.026	3.73 ± 0.05
11.320 ± 15 ^e	$2^+; 0$	0.036 ± 0.004		4.41 ± 0.05
11.856 ± 20 ^e	$2^+; 0$	0.0016 ± 0.0004		4.98 ± 0.10

^a For upper limits to other ^{20}Ne states see Table 20.34 in (1972AJ02) and (1973TO08, 1976IN06). For earlier values see (1972AJ02).

^b (1976IN06).

^c 10.278 ± 5 (1973TO08).

^d Electron capture + β^+ .

^e (1973TO08).

^f Includes radiative, nuclear size, lepton wavelength, electron screening and electron capture corrections (1976IN06).

^g Assuming $\Gamma_\gamma = 5.6 \pm 0.6$ eV, $\Gamma_{\text{total}} = 356 \pm 230$ eV (1973TO08).

58. $^{20}\text{Na}(\beta^+)^{20}\text{Ne}$

$$Q_m = 13.887$$

^{20}Na has a half-life of 446 ± 3 msec: see reaction 1 in ^{20}Na . It decays to a number of states of ^{20}Ne , principally $^{20}\text{Ne}^*(1.63)$: see Table 20.37. The ratio of the mirror decays $^{20}\text{Na} \xrightarrow{\beta^+} ^{20}\text{Ne}^*(1.63)$ and $^{20}\text{F} \xrightarrow{\beta^-} ^{20}\text{Ne}^*(1.63)$, $(ft)^+/(ft)^- = 1.026 \pm 0.024$ (1973TO08), 1.033 ± 0.022 (1976IN06). (1977RO20) obtain a correlation $W_+(\theta) = 1 + \cos^2 \theta [(-4.0 \pm 0.7) \times 10^{-3} E + (1.3 \pm 0.9) \times 10^{-4} E^2](p/E)^2$: assuming the validity of CVC this leads to a vanishing second-class current. See also reaction 45. The β - α angular correlation involving the allowed transition to $^{20}\text{Ne}^*(7.42)$ is consistent with the predictions of the CVC theory (1977FR08). See also (1972TO08, 1973IN01, 1974AL11, 1975RO1R, 1977RO1W) and (1971WI1C, 1972EM02, 1972WI1C, 1973LA03, 1974WI1L, 1975WI1E, 1977CA11, 1977OK1E, 1977WI02; theor.).

59. $^{21}\text{Ne}(p, d)^{20}\text{Ne}$

$$Q_m = -4.537$$

Angular distributions have been measured for the first four deuteron groups at $E_p = 14.1$ MeV (1972HE24) and 20 MeV (1970HO19). (1972HE24) report spectroscopic factors of 0.50 and 0.14 for $^{20}\text{Ne}^*(1.63, 4.97)$: the other two distributions do not show direct reaction features.

60. $^{21}\text{Ne}(d, t)^{20}\text{Ne}$

$$Q_m = -0.504$$

The $T = 1$ states observed in this reaction, and the analog states observed in ^{20}F in the $(d, ^3\text{He})$ reaction, are displayed in Table 20.16. The spectroscopic factors of analog states are consistent to within 20% for states excited by a single l -transfer. $T = 0$ states are displayed in Table 20.38 (1974MI13).

61. $^{21}\text{Ne}(^3\text{He}, \alpha)^{20}\text{Ne}$

$$Q_m = 13.816$$

See (1973FO1E).

62. $^{22}\text{Ne}(p, t)^{20}\text{Ne}$

$$Q_m = -8.644$$

Angular distributions have been reported at $E_p = 26.9, 35.1$ and 42.4 MeV (1971FA07: $t_0, t_1, t_2, t_3, t_{4+5}, t_6$) and at 43.7 MeV (1964CE05). At the higher energy the distributions of the tritons to the ground state of ^{20}Ne and to the first 0^+ ; $T = 2$ state [$E_x = 16.722 \pm 0.025$ MeV (1969HA38)] have been fitted by $L = 0$ and the tritons to $^{20}\text{Ne}^*(18.5)$ by $L = 2$. The latter is the first 2^+ ;

Table 20.38: $T = 0$ states of ^{20}Ne from $^{21}\text{Ne}(d, t)^{20}\text{Ne}$ (1974MI13) ^a

E_x (MeV \pm keV)	l	nlj ^b	C^2S	J^π ^c	K^π
$\equiv 5.622$	1	$1p_{3/2}$	0.02	3^-	
5.785 ± 4	1	$1p_{1/2}$	0.03	1^-	
$\equiv 7.424$	$0 + 2$	$2s_{1/2}$	0.05		
		$1d_{5/2}$	0.07	2^+	
7.827 ± 9	$0 + 2$	$2s_{1/2}$	0.005		
		$1d_{5/2}$	0.023	2^+	
8.839 ± 8	1	$1p_{1/2}$	0.33	1^-	(1^-)
9.084 ± 21 ^d	2	$1d_{5/2}$	≤ 0.12		
9.357 ± 17 ^d	1	$1p_{1/2}$	≤ 0.1	^e	(1^-)
9.913 ± 19 ^d	2	$1d_{5/2}$	< 0.16		
10.385 ± 12	1	$1p_{3/2}$	0.08	3^-	
10.880 ± 10 ^d	1	$1p_{3/2}$	0.13		

^a For $T = 1$ states see Table 20.16.

^b Values used in DWBA calculations.

^c From Table 20.18.

^d Unresolved.

^e See, however, discussion in (1974MI13).

$T = 2$ state (1974CE1A). The 0^+ ; $T = 2$ state [$^{20}\text{Ne}^*(16.73)$] decays by α_0 [$-6 \pm 5\%$], $\alpha_1 + \alpha_2$ [$35 \pm 12\%$], $\alpha_3 + \alpha_4$ [$29 \pm 12\%$], $p_0 + p_1 + p_2$ [$14 \pm 9\%$] and $p_3 + p_4 + p_5$ [$13 \pm 8\%$] [measured branching ratios in percent are given in the brackets] to final states in ^{16}O and ^{19}F (1970MC04). The ratios of the cross section for formation of the analog states $^{20}\text{Ne}^*(10.27)/^{20}\text{F}^*(0)$ and $^{20}\text{Ne}^*(12.25 \pm 0.03)/^{20}\text{F}^*(1.85)$ are 2.00 ± 0.20 and 1.40 ± 0.15 , respectively, at $E_p = 45$ MeV (1969HA19). See also (1971GO1Q).

At $E_p = 40$ MeV angular distributions of the tritons to $^{20}\text{Ne}^*(4.97, 5.62, 7.00)$ [$J^\pi = 2^-, 3^-$ and 4^- , respectively] have been measured. Coupled-channels calculations reproduce the distributions to the 2^- and 3^- states, but the distribution to the 4^- states cannot be explained entirely in terms of multistep inelastic processes (1975CH17). See also (1974VO11).

See also (1973FO1E) and (1972FE1A, 1972OL04, 1972SC10, 1973BA18, 1973OL03, 1973UD01, 1976KI09; theor.).

63. $^{22}\text{Ne}(\alpha, ^6\text{He})^{20}\text{Ne}$

$Q_m = -16.155$

See (1972AJ02).

$$64. \text{}^{23}\text{Na}(\text{p}, \alpha)\text{}^{20}\text{Ne} \quad Q_{\text{m}} = 2.377$$

The first five states of ^{20}Ne have been observed in this reaction, and angular distributions have been measured at $E_{\text{p}} = 10.0$ and 45.5 MeV: see (1972AJ02).

For work dealing with resonances in the compound nucleus see (1972AJ02), (1974VO09) and ^{24}Mg in (1978EN06). See also (1973AR1F, 1973CL1E; astrophys. considerations).

$$65. \text{}^{23}\text{Na}(\text{}^3\text{He}, \text{}^6\text{Li})\text{}^{20}\text{Ne} \quad Q_{\text{m}} = -1.643$$

At $E(^3\text{He}) = 40.7$ MeV, angular distributions have been measured to $^{20}\text{Ne}^*(0, 1.63, 4.25)$ and analyzed using zero-range DWBA (1972OH01).

$$\begin{aligned} 66. \text{(a) } &^{24}\text{Mg}(\gamma, \alpha)\text{}^{20}\text{Ne} & Q_{\text{m}} &= -9.3125 \\ \text{(b) } &^{24}\text{Mg}(\text{p}, \text{p}\alpha)\text{}^{20}\text{Ne} & Q_{\text{m}} &= -9.3125 \\ \text{(c) } &^{24}\text{Mg}(\alpha, 2\alpha)\text{}^{20}\text{Ne} & Q_{\text{m}} &= -9.3125 \end{aligned}$$

For reaction (a) see (1973CL1E; astrophys.). For reaction (b) see (1970LI08, 1975ST1Q). See also (1975BA2H), (1975RO1B) and (1975PA1K, 1977CH02; theor.). At $E_{\alpha} = 90$ MeV ^{20}Ne states at $E_{\text{x}} = 0, 1.63 \pm 0.035, 4.25 \pm 0.06, 5.70 \pm 0.06, 6.72 \pm 0.05$ and 11.41 ± 0.05 MeV are populated: the ground states S_{α} (DWIA) = 1.3 ± 0.2 (1976SH02).

$$67. \text{}^{24}\text{Mg}(\text{d}, \text{}^6\text{Li})\text{}^{20}\text{Ne} \quad Q_{\text{m}} = -7.839$$

Angular distributions have been studied involving $^{20}\text{Ne}^*(0, 1.63, 4.25)$ at $E_{\text{d}} = 28$ MeV (1972CO23), 35 MeV (1976CO15; also $^{20}\text{Ne}^*(5.62, 5.78)$), 54.3 MeV (1977TA1D; also $^{20}\text{Ne}^*(5.62+5.78)$) and 80 MeV (1974DJ01; also $^{20}\text{Ne}^*(5.62)$). Using zero-range DWBA, (1976CO15) report $S_{\alpha} = 0.21, 0.19, 0.87, 0.26, 0.04$ for $^{20}\text{Ne}^*(0, 1.63, 4.25, 5.62, 5.78)$. See also (1973FO1A, 1973FO1E, 1974FO1J).

$$68. \text{}^{24}\text{Mg}(\text{}^3\text{He}, \text{}^7\text{Be})\text{}^{20}\text{Ne} \quad Q_{\text{m}} = -7.726$$

Angular distributions involving ${}^7\text{Be}_{\text{g.s.}}$ and ${}^7\text{Be}_{0.43}^*$ and ${}^{20}\text{Ne}^*(0, 1.63, 4.25)$ have been measured at $E({}^3\text{He}) = 25.5$ MeV (1976PI10) and 70 MeV (1976ST11; also ${}^{20}\text{Ne}^*(4.97, 5.62)$). Spectroscopic factors from a FRDWBA analysis of the 25.5 MeV data give 1.0, 15.4 and 3.5 [from ${}^7\text{Be}_{\text{g.s.}}$ data] and 1.0, 23.4 and 4.3 [from ${}^7\text{Be}_{0.43}^*$ data] for ${}^{20}\text{Ne}^*(0, 1.63, 4.25)$. The absolute spectroscopic factors for ${}^{20}\text{Ne}_{\text{g.s.}}$ are 0.67 [${}^7\text{Be}_{\text{g.s.}}$] and 0.63 [${}^7\text{Be}_{0.43}^*$] (1976PI09, 1976PI10). See also (1973BR1C).

$$69. {}^{25}\text{Mg}(\text{d}, {}^7\text{Li}){}^{20}\text{Ne} \quad Q_{\text{m}} = -7.920$$

See (1971MC04).

$$70. {}^{26}\text{Mg}(\text{p}, \text{t}_\alpha){}^{20}\text{Ne} \quad Q_{\text{m}} = -19.255$$

See (1967MC06).

^{20}Na
(Figs. 11, 12 and 13)

GENERAL: (See also (1972AJ02).)

(1973HA77, 1973SU1B, 1974HA17, 1976CH1T, 1977SH13).

$$J = 2 \text{ (1975SC20);}$$

$$\mu = 0.3694 \pm 0.0002 \text{ nm (1975SC20).}$$

1. $^{20}\text{Na}(\beta^+)^{20}\text{Ne}$ $Q_m = 13.887$

^{20}Na decays by positron emission to $^{20}\text{Ne}^*(1.63)$ and to a number of other excited states of ^{20}Ne : see Table 20.37. The half-life of ^{20}Na is 442 ± 5 msec (1971GO18, 1971WI07), 446 ± 8 msec (1972MO08), 448 ± 4 msec (1973TO08): the weighted mean of these measurements is 446 ± 3 msec. The character of the β^+ decay sets $J^\pi = 2^+$ for the ground state of ^{20}Na . $J = 2$ is confirmed also by the work of (1975SC20) who studied the asymmetry in the β^+ decay of polarized ^{20}Na nuclei to detect r.f. transitions between h.f.s. Zeeman levels of the atomic ground state [see the “GENERAL” section here]. See also reaction 58 in ^{20}Ne .

2. $^{20}\text{Ne}(p, n)^{20}\text{Na}$ $Q_m = -14.669$

Observed neutron groups at $E_p = 22.9$ MeV are displayed in Table 20.40 (1971MO34). See also (1972BA1U) and (1972AJ02).

3. $^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}$ $Q_m = -13.906$

At $E(^3\text{He}) = 32$ MeV, triton groups are observed to nine states of ^{20}Na : see Table 20.40 (1965DO04, 1965PE04).

4. $^{24}\text{Mg}(^3\text{He}, ^7\text{Li})^{20}\text{Na}$ $Q_m = -20.751$

See (1976BE1L).

Table 20.39: Energy levels of ^{20}Na

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
0	$2^+; 1$	$\tau_{1/2} = 446 \pm 3$ msec	β^-	1, 2, 3, 4
0.591 ± 12				2, 3
0.768 ± 8				2, 3
(0.85 ± 50)				3
0.958 ± 8				2, 3
(1.010 ± 14)				2
1.310 ± 10				2, 3
1.92 ± 40				3
2.89 ± 50		a		3
4.33 ± 100		a		3

^a Broad or unresolved.

Table 20.40: States of ^{20}Na from $^{20}\text{Ne}(p, n)^{20}\text{Na}$ and $^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}$

E_x (MeV \pm keV)	
$^{20}\text{Ne}(p, n)^{20}\text{Na}$ (1971MO34)	$^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}$ (1965DO04, 1965PE04)
0	0
0.591 ± 12	0.65 ± 50 ^a
0.768 ± 8	0.75 ± 50 ^a
	0.85 ± 50 ^a
0.958 ± 8	0.95 ± 50 ^a
(1.010 ± 14)	
1.310 ± 10	1.27 ± 50
	1.92 ± 40
	2.89 ± 50
	4.33 ± 100

^a These states are not fully resolved.

²⁰Mg
(Fig. 13)

²⁰Mg has been populated in the ²⁴Mg(α , ⁸He) reaction at $E_\alpha = 126.9$ MeV (1976TR03) and 156 MeV (1974RO17) with differential cross sections (lab) of 3 ± 1 nb/sr ($\theta = 5^\circ$, lab) and ≈ 7 nb/sr (2°), respectively. Assuming the mass of ⁸He to be 31.601 ± 0.013 MeV, the mass excess of ²⁰Mg is 17.57 ± 0.03 MeV. (1977WA08) adopt a mass excess of 17.568 ± 0.027 MeV, and so do we. ²⁰Mg is then stable with respect to breakup into ¹⁹Na + p [see ¹⁹Na] and ¹⁸Ne + 2p by 2.65 ± 0.03 and 2.33 ± 0.03 MeV, respectively. See (1976WA18) for a display of calculations of the mass of ²⁰Mg and (1972AJ02) for the earlier work. The mass excess of ²⁰Mg appears to involve a deviation from the quadratic IMME prediction (1976TR03). Using the wave function predictions of (1973LA03) and the mass excess of ²⁰Mg (1976TR03) calculate the partial $\tau_{1/2}$ for the allowed Fermi transition from the 0^+ ; $T = 2$ ground state of ²⁰Mg to the 0^+ ; $T = 2$ state in ²⁰Na [not observed] to be ≈ 3.4 sec, corresponding to a β^+ branching of $\approx 4\%$. See also (1976BE1L, 1977SH13).

²⁰Al
(Not illustrated)

²⁰Al has not been observed: see (1966KE16).

References

(Closed 01 November 1977)

References are arranged and designated by the year of publication followed by the first two letters of the first-mentioned author's name and then by two additional characters. Most of the references appear in the National Nuclear Data Center files (Nuclear Science References Database) and have NNDC key numbers. Otherwise, TUNL key numbers were assigned with the last two characters of the form 1A, 1B, etc. In response to many requests for more informative citations, we have, when possible, included up to ten authors per paper and added the authors' initials.

- 1950AR1A Arnold, Phys. Rev. 79 (1950) 170; Erratum Phys. Rev. 80 (1950) 34
1950BA1A Barnes, French and Devons, Nature 166 (1950) 145
1950CH1A Chao, Phys. Rev. 80 (1950) 1035
1952WI1A Willard et al., Phys. Rev. 85 (1952) 849
1953CA44 J.R. Cameron, Phys. Rev. 90 (1953) 839
1954DE1A Dearnaley, Phil. Mag. 45 (1954) 1213
1954PE1A Peterson, Barnes, Fowler and Lauritsen, Phys. Rev. 94 (1954) 1075
1954WO23 C. Wong, Phys. Rev. 95 (1954) 761
1955BA94 C.A. Barnes, Phys. Rev. 97 (1955) 1226
1955BA1C Baranger, Phys. Rev. 99 (1955) 145
1955FA1A Farney, Givin, Kern and Hahn, Phys. Rev. 97 (1955) 720
1955HU1A Hunt and Firth, Phys. Rev. 99 (1955) 786
1955KI28 J.D. Kington, J.K. Bair, H.O. Cohn and H.B. Willard, Phys. Rev. 99 (1955) 1393
1955MA1D Marion and Brugger, Phys. Rev. 100 (1955) 69
1955WE1A Webb, Hagecorn, Fowler and Lauritsen, Phys. Rev. 99 (1955) 138
1956DE33 G. Dearnaley, Phil. Mag. 1 (1956) 821
1956EL1A El-Bedewi, Proc. Phys. Soc. (London) A69 (1956) 221
1957CL42 R.L. Clarke and E.B. Paul, Can. J. Phys. 35 (1957) 155
1957MA1A Martin, Fowler, Lauritsen and Lauritsen, Phys. Rev. 106 (1957) 1260
1958IS10 A. Isoya, H. Ohmura and T. Momota, Nucl. Phys. 7 (1958) 116
1958IS11 A. Isoya, Nucl. Phys. 7 (1958) 126
1958MO02 S. Morita and K. Takeshita, J. Phys. Soc. Jpn. 13 (1958) 1241
1958RA15 W.A. Ranken, T.W. Bonner and J.H. McCrary, Phys. Rev. 109 (1958) 1646
1958WI36 J.E. Wills, Jr., J.K. Bair, H.O. Cohn and H.B. Willard, Phys. Rev. 109 (1958) 891
1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1

1959BO14 R.O. Bondelid and C.A. Kennedy, Phys. Rev. 115 (1959) 1601
 1959BR67 G. Breuer, Z. Phys. 154 (1959) 339
 1959GA08 F. Gabbard, R.H. Davis and T.W. Bonner, Phys. Rev. 114 (1959) 201
 1959HA1C Hay and Warren, Can. J. Phys. 37 (1959) 1153
 1959KU79 J. Kuperus, P.J.M. Smulders and P.M. Endt, Physica 25 (1959) 600
 1959LI51 A. Lippold, J.W. Muller, H.H. Staub and H. Winkler, Helv. Phys. Acta 32 (1959) 268
 1960BU07 J.W. Butler, Phys. Rev. 118 (1960) 222
 1960HU11 S.E. Hunt, R.A. Pope, D.V. Freck and W.W. Evans, Phys. Rev. C12 (1960) 1740
 1960KA18 J.V. Kane, R.E. Pixley and D.H. Wilkinson, Phys. Rev. 120 (1960) 952
 1960MC09 L.C. McDermott, K.W. Jones, H. Smotrlich and R.E. Benenson, Phys. Rev. 118 (1960) 175
 1960SM03 D.M. Smith, N.A. Bostrom and E.L. Hudspeth, Phys. Rev. 117 (1960) 514
 1960TS02 K. Tsukada and T. Fuse, J. Phys. Soc. Jpn. 15 (1960) 1994
 1961BE13 E.H. Beckner, R.L. Bramblett, G.C. Phillips and T.A. Eastwood, Phys. Rev. 123 (1961) 2100
 1961DA16 E.A. Davis, F. Gabbard, T.W. Bonner and R. Bass, Nucl. Phys. 27 (1961) 448
 1961FR02 E. Freiberg and V. Soergel, Z. Phys. 162 (1961) 114
 1961GO21 H.E. Gove, A.E. Litherland and A.J. Ferguson, Phys. Rev. 124 (1961) 1944
 1962BR35 C. Broude, M.A. Clark and A.E. Litherland, Phys. Lett. 3 (1962) 118
 1962DO1A Dodge and Barber, Phys. Rev. 127 (1962) 1746
 1962HI06 S. Hinds, H. Marchant and R. Middleton, Nucl. Phys. 38 (1962) 81
 1962JO14 J.C. Jodogne, P.C. Macq and J. Steyaert, Phys. Lett. 2 (1962) 325
 1962KE03 L. Keszthelyi, I. Berkes, I. Demeter and I. Fodor, Nucl. Phys. 29 (1962) 241
 1962RY01 A. Rytz, H.H. Staub, H. Winkler and W. Zych, Helv. Phys. Acta 35 (1962) 341
 1963BE19 I. Berkes, I. Dezs, I. Fodor and L. Keszthelyi, Nucl. Phys. 43 (1963) 103
 1963FI1B Finck et al., Z. Phys. 174 (1963) 337
 1963GL01 S.S. Glickstein and R.G. Winter, Phys. Rev. 129 (1963) 1281
 1963JA01 R. Jahr, Phys. Rev. 129 (1963) 320
 1963JE04 J.G. Jenkin, L.G. Earwaker and E.W. Titterton, Nucl. Phys. 44 (1963) 453
 1963LA08 N.O. Lassen and J.S. Olsen, Kgl. Danske Videnskab.Selskab., Mat.-Fys.Medd. 33, No.13 (1963)
 1964BR12 G. Breuer and L. Jahnke, Z. Naturforsch. A19 (1964) 471
 1964BR18 C. Broude, A.E. Litherland and J.D. Pearson, Phys. Lett. 11 (1964) 321

1964CE05 J. Cerny, R.H. Pehl and G.T. Garvey, Phys. Lett. 12 (1964) 234
 1964KA1A Kawai, Terasawa and Izumo, Nucl. Phys. 59 (1964) 289
 1964MI05 R. Middleton and D.J. Pullen, Nucl. Phys. 51 (1964) 63
 1964PE05 J.D. Pearson and R.H. Spear, Nucl. Phys. 54 (1964) 434
 1964SA09 D.G. Sargood and G.D. Putt, Aust. J. Phys. 17 (1964) 56; Phys. Abstr. 67, 1526, Abstr. 17040 (1964)
 1964SE1A Seagrave, Brolley and Beery, Rev. Sci. Instrum. 35 (1964) 1290
 1965AS07 L. Ask, Ark. Fys. 29 (1965) 195
 1965DO04 P.F. Donovan and P.D. Parker, Phys. Rev. Lett. 14 (1965) 147
 1965EV03 H.C. Evans, M.A. Eswaran, H.E. Gove, A.E. Litherland and C. Broude, Can. J. Phys. 43 (1965) 82
 1965MA28 L.G. Mann and S.D. Bloom, Phys. Rev. 139 (1965) B540
 1965MC02 G.J. McCallum and A.J. Ferguson, Can. J. Phys. 43 (1965) 111
 1965MO19 R. Moreh, Nucl. Phys. 70 (1965) 293
 1965PE04 R.H. Pehl and J. Cerny, Phys. Lett. 14 (1965) 137
 1965SI18 R.H. Siemssen, L.L. Lee, Jr. and D. Cline, Phys. Rev. 140 (1965) B1258
 1965VA14 C. van der Leun, D.M. Sheppard and P.J.M. Smulders, Phys. Lett. 18 (1965) 1344
 1966AL12 D.E. Alburger and K.W. Jones, Phys. Rev. 149 (1966) 743
 1966CA14 J. Cabe, M. Laurat and P. Yvon, J. Phys. (Paris) 27 (1966) 755
 1966KE16 I. Kelson and G.T. Garvey, Phys. Lett. 23 (1966) 689
 1966MA60 J.B. Marion, Rev. Mod. Phys. 38 (1966) 660
 1966RI05 R.C. Ritter, Nucl. Phys. 80 (1966) 377
 1967BE36 I. Bergqvist, J.A. Biggerstaff, J.H. Gibbons and W.M. Good, Phys. Rev. 158 (1967) 1049
 1967BL19 R. Bloch, R.E. Pixley and P. Truol, Phys. Lett. B25 (1967) 215
 1967BR22 C. Broude, A.E. Litherland, R.W. Ollerhead and T.K. Alexander, Can. J. Phys. 45 (1967) 3837
 1967HA14 L.F. Hansen, J.D. Anderson, J.W. McClure, B.A. Pohl, M.L. Stelts, J.J. Wesolowski and C. Wong, Nucl. Phys. A98 (1967) 25
 1967HA23 L.F. Hansen, H.F. Lutz, M.L. Stelts, J.G. Vidal and J.J. Wesolowski, Phys. Rev. 158 (1967) 917
 1967HU06 W.E. Hunt, M.K. Mehta and R.H. Davis, Phys. Rev. 160 (1967) 782
 1967KU06 H.M. Kuan, D.W. Heikkinen, K.A. Snover, F. Riess and S.S. Hanna, Phys. Lett. B25 (1967) 217

- 1967LI07 A.E. Litherland, R.W. Ollerhead, P.J.M. Smulders, T.K. Alexander, C. Broude, A.J. Ferguson and J.A. Kuehner, *Can. J. Phys.* 45 (1967) 1901
- 1967MC06 R.L. McGrath, S.W. Cospser and J. Cerny, *Phys. Rev. Lett.* 18 (1967) 243
- 1967ME10 M.K. Mehta, W.E. Hunt and R.H. Davis, *Phys. Rev.* 160 (1967) 791
- 1967QU01 P.A. Quin, A.A. Rollefson, G.A. Bissinger, C.P. Browne and P.R. Chagnon, *Phys. Rev.* 157 (1967) 991
- 1967SE02 R.E. Segel, Z. Vager, L. Meyer-Schutzmeister, P.P. Singh and R.G. Allas, *Nucl. Phys.* A93 (1967) 31
- 1967TH06 W.J. Thompson, S. Edwards and T. Tamura, *Nucl. Phys.* A105 (1967) 678
- 1967VA08 G. van Middelkoop, *Nucl. Phys.* A97 (1967) 209
- 1968BL1C Blatchley and Thomas, ANL 7282 (1968) 71
- 1968LA03 B.T. Lawergren, A.T.G. Ferguson and G.C. Morrison, *Nucl. Phys.* A108 (1968) 325
- 1968RI08 F.L. Riffle, J.D. Goss, D.R. Parsignault and J.C. Harris, *Nucl. Phys.* A115 (1968) 120
- 1968RI13 L.R. Rice and R.T. Carpenter, *Nucl. Phys.* A120 (1968) 220
- 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, *Nucl. Phys.* A113 (1968) 395
- 1969AD02 E.G. Adelberger, A.B. McDonald and C.A. Barnes, *Nucl. Phys.* A124 (1969) 49
- 1969AN08 J.H. Anderson and R.C. Ritter, *Nucl. Phys.* A128 (1969) 305
- 1969CO19 A.A. Cowley and G. Heymann, *Phys. Lett.* B30 (1969) 618
- 1969FU1A Fukushima, Kamimura and Takaca, *Mem. Fac. Sci. Kyushu U. (Japan)* B4 (1969)39
- 1969GA05 A. Gallmann, F. Jundt, E. Aslanides and D.E. Alburger, *Phys. Rev.* 179 (1969) 921
- 1969GO1B Gorodetzky, Rudolf, Scheibling and Chevallier, *Contrib., Montreal* (1969) 302
- 1969HA04 R. Hardell and A. Hasselgren, *Nucl. Phys.* A123 (1969) 215
- 1969HA19 J.C. Hardy, H. Brunnader and J. Cerny, *Phys. Rev. Lett.* 22 (1969) 1439
- 1969HA38 J.C. Hardy, H. Brunnader, J. Cerny and J. Janecke, *Phys. Rev.* 183 (1969) 854
- 1969HE08 G.H. Herling, L. Cohen and J.D. Silverstein, *Phys. Rev.* 178 (1969) 1551
- 1969HE20 R.L. Hershberger, M.J. Wozniak, Jr. and D.J. Donahue, *Phys. Rev.* 186 (1969) 1167
- 1969HO16 W. Hofmann, R. Kosiek, G. Kraft and R. Mundhenke, *Z. Phys.* 225 (1969) 303
- 1969HO20 T. Holtebekk, S. Tryti and G. Vamraak, *Nucl. Phys.* A134 (1969) 353
- 1969JO18 J. John, J.P. Aldridge and R.H. Davis, *Phys. Rev.* 181 (1969) 1455
- 1969MA48 N. Martalogu, *Stud. Cercet. Fiz. (Romania)* 21 (1969) 1115
- 1969RI01 R.C. Ritter, J.T. Parson and D.L. Bernard, *Phys. Lett.* B28 (1969) 588
- 1969RO1F Roturier, *Ann. Phys.* 4 (1969) 289

1969SC08 D. Schwalm and B. Povh, Phys. Lett. B29 (1969) 103

1970AS1C Aslanides, Frnc Th 116 (1970)

1970CO13 A.A. Cowley and G. Heymann, Nucl. Phys. A146 (1970) 465

1970CR1A Craig, Thesis, Yale Univ. (1970); Phys. Abs. 75453 (1971)

1970FL1A Flowers, Meth. and Problems of Theor. Phys. (North-Holland, 1970) 289

1970GU08 K. Gul, B.H. Armitage and B.W. Hooton, Nucl. Phys. A153 (1970) 390

1970HE1B Hershberger, Thesis, Univ. Arizona (1970); Phys. Abs. 63693 (1971)

1970HO05 T. Holtebekk, R. Stromme and S. Tryti, Nucl. Phys. A142 (1970) 251

1970HO19 A.J. Howard, J.G. Pronko and R.G. Hirko, Nucl. Phys. A150 (1970) 609

1970LI08 R.A. Lindgren, J.G. Pronko, A.J. Howard and D.A. Bromley, Phys. Lett. B32 (1970) 103

1970MA42 H.B. Mak, H. Spinka and H. Winkler, Phys. Rev. C2 (1970) 1729

1970MC04 R.L. McGrath, J. Cerny, J.C. Hardy, G. Goth and A. Arima, Phys. Rev. C1 (1970) 184

1970NI03 R.W. Nightingale, J.A. Becker, R.E. McDonald and D. Kohler, Phys. Rev. C1 (1970) 893

1970QU04 P.A. Quin, G.A. Bissinger and P.R. Chagnon, Nucl. Phys. A155 (1970) 495

1970QU1A Quin and Vigdor, Bull. Amer. Phys. Soc. 15 (1970) 1686

1970QU1C Quinn, Thesis, Univ. California, Los Angeles (1970); Phys. Abs. 60773 (1971)

1970RO06 A.A. Rollefson, P.F. Jones and R.J. Shea, Phys. Rev. C1 (1970) 1761

1970WI05 D.H. Wilkinson and D.E. Alburger, Phys. Rev. Lett. 24 (1970) 1134

1971AB1B Abgrall and Caurier, Suppl. J. Phys. 32 (1971) C6-63

1971AR1V Arima, Suppl. J. Phys. 32 (1971) C6-33

1971BA1A Barnes, Advances in Nucl. Phys., Eds. Baranger and Vogt, Vol. 4 (Plenum Press, 1971) p. 133

1971BA2V Bassani, CEA N 1474 (1971)

1971BA68 P.H. Barker, P.M. Cockburn, A. Huber, H. Knoth, U. Matter, H.-P. Seiler and P. Marmier, Ann. Phys. (New York) 66 (1971) 705

1971BE17 C. Bergman and R.K. Hobbie, Phys. Rev. C3 (1971) 1729

1971BE18 W.L. Bendel, L.W. Fagg, S.K. Numrich, E.C. Jones, Jr. and H.F. Kaiser, Phys. Rev. C3 (1971) 1821

1971BE2F Beckert, Zentral. Kernf. Rossendorf Bei Dresden, Rept. No. Zfk 222 (1971)

1971DE56 D.R. de Oliveira, Rev. Bras. Fis. (Brasil) 1 (1971) 403; Phys. Abs. 38483 (1972)

1971DI08 W.T. Diamond, T.K. Alexander and O. Hausser, Can. J. Phys. 49 (1971) 1589

- 1971EP03 M.B. Epstein, J.R. Quinn, S.N. Bunker, J.W. Verba and J.R. Richardson, Nucl. Phys. A169 (1971) 337
- 1971FA07 W.R. Falk, P. Kulisic and A. McDonald, Nucl. Phys. A167 (1971) 157
- 1971FE1F Fernandez, Ruiz and Bauza, An. Fisica 67 (1971) 387
- 1971FO14 H.T. Fortune, J.D. Garrett, J.R. Powers and R. Middleton, Phys. Rev. C4 (1971) 850; Erratum Phys. Rev. C5 (1972) 1438
- 1971GO18 D.R. Goosman, K.W. Jones, E.K. Warburton and D.E. Alburger, Phys. Rev. C4 (1971) 1800
- 1971GO1Q Goth, LBL224 (1971)
- 1971GO1R Gorodetzky, Rudolf, Scheibling and Chevallier, Suppl. J. Phys. 32 (1971) C6-197
- 1971HA26 O. Hausser, T.K. Alexander, A.B. McDonald, G.T. Ewan and A.E. Litherland, Nucl. Phys. A168 (1971) 17
- 1971JH1A Jhaveri, Thesis, Univ. Tennessee (1971); Phys. Abs. 70460 (1972)
- 1971KA18 S. Kardonsky, H.L. Finston and E.T. Williams, Phys. Rev. C4 (1971) 846
- 1971KO1W Kopsch and Cierjacks, Proc. Int. Conf. Statistical Properties of Nuclei, Albany, New York, 1971 (New York: Plenum Press 1972) p. 455; Phys. Abs. 46270 (1972)
- 1971MA23 N. Marquardt, W. Von Oertzen and R.L. Walter, Phys. Lett. B35 (1971) 37
- 1971MC04 R.L. McGrath, D.L. Hendrie, E.A. McClatchie, B.G. Harvey and J. Cerny, Phys. Lett. B34 (1971) 289
- 1971MI09 R. Middleton, J.D. Garrett and H.T. Fortune, Phys. Rev. Lett. 27 (1971) 950
- 1971MO15 J.R. Moreira, R.P. Singhal and H.S. Caplan, Can. J. Phys. 49 (1971) 1434
- 1971MO1P Moulder et al., Nucl. Phys. B35 (1971) 332
- 1971MO34 C.E. Moss and A.B. Comiter, Nucl. Phys. A178 (1971) 241
- 1971MO40 C. Morand, H. Beaumevieille, A. Dauchy, G. Dumazet, M. Lambert and C. Meynadier, Nuovo Cim. A6 (1971) 380
- 1971NE1E Nettles, Thesis, C.I.T. (1971); Phys. Abs. 78443 (1971)
- 1971NY02 G. Nystrom, B. Lundberg and I. Bergqvist, Phys. Scr. 4 (1971) 95
- 1971PR10 J.G. Pronko and R.W. Nightingale, Phys. Rev. C4 (1971) 1023
- 1971RA1B Rabbat and Do Dang, Rept. Lpthe 71/18, Lab. Phys. Theor. Hautes Ener., Orsay (1971); Phys. Abs. 42493 (1972)
- 1971RI1D Ripka, Suppl. J. Phys. 32 (1971) C6-261
- 1971RO06 A.A. Rollefson and J.A. Aymar, Phys. Rev. C3 (1971) 1704
- 1971RO13 D.W.O. Rogers, J.H. Aitken, A.E. Litherland, W.R. Dixon and R.S. Storey, Can. J. Phys. 49 (1971) 1397

- 1971RO1C Rogers, Thesis, Univ. Toronto (1971)
- 1971RO33 D.W.O. Rogers, K.W. Allen, H.C. Evans, N.A. Jelley, A.E. Litherland and B.Y. Underwood, Phys. Lett. B37 (1971) 65
- 1971RU14 F.R. Ruehl, Jr., Indian J. Phys. 45 (1971) 149
- 1971SA1A Satpathy, Friedrich and Weiguny, Suppl. J. Phys. 32 (1971) C6-269
- 1971SH26 V.S. Shirley, Proc. Int. Conf. Hyperfine Interactions Detected by Nucl. Radiation, Israel, 1970 (London, Gordon & Breach, 1971) 1255
- 1971SI24 M.H. Simbel and A.Y. Abul-Magd, Nucl. Phys. A177 (1971) 322
- 1971TA05 M. Takeda, S. Kato and T. Yamazaki, J. Phys. Soc. Jpn. 30 (1971) 56
- 1971TO06 J.W. Toevs, Nucl. Phys. A172 (1971) 589
- 1971TO1C Toevs, Fowler, Barnes and Lyons, Astrophys. J. 169 (1971) 421
- 1971WI07 D.H. Wilkinson, D.E. Alburger, D.R. Goosman, K.W. Jones, E.K. Warburton, G.T. Garvey and R.L. Williams, Nucl. Phys. A166 (1971) 661.
- 1971WI1C Wilkinson, Nucl. Phys. A178 (1971) 65
- 1971WO1C Wong and Zuker, Suppl. J. Phys. 32 (1971) C6-51
- 1971ZO1A Zofka, Czech. J. Phys. B21 (1971) 1051
- 1972AB12 Y. Abgrall, B. Morand and E. Caurier, Nucl. Phys. A192 (1972) 372
- 1972AB1A A. Abragam, G.L. Bacchella, C. Long, P. Meriel, J. Peisvaux and M. Pinot, Phys. Rev. Lett. 28 (1972) 805, 1225; Erratum Phys. Rev. Lett. 29 (1972) 894
- 1972AC03 H. Ackermann, D. Dubbers, M. Grupp, P. Heitjans, G.Z. Putlitz and H.-J. Stockmann, Phys. Lett. B41 (1972) 143; Erratum Phys. Lett. B44 (1973) 257
- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1
- 1972AL05 T.K. Alexander, O. Hausser, A.B. McDonald, A.J. Ferguson, W.T. Diamond and A.E. Litherland, Nucl. Phys. A179 (1972) 477
- 1972AL26 L.A. Alexander, W.K. Collins, B.P. Hichwa, J.C. Lawson, D.S. Longo, E.D. Berners and P.R. Chagnon, Phys. Rev. C6 (1972) 817
- 1972AL32 T.K. Alexander, B.Y. Underwood, N. Anyas-Weiss, N.A. Jelley, J. Szucs, S.P. Dolan, M.R. Wormald and K.W. Allen, Nucl. Phys. A197 (1972) 1
- 1972AR11 A. Arima, G. Scharff-Goldhaber and K.W. McVoy, Phys. Lett. B40 (1972) 7
- 1972AR12 A. Arima and S. Yoshida, Phys. Lett. B40 (1972) 15
- 1972AR1F Arima, Proc. 1971 Mont Tremblant Summer School Meeting on Dynamic Struct. of Nucl. States, Canada (Toronto, Canada, Univ. Toronto Press, 1972) 292; Phys. Abs. 82868 (1972)
- 1972AS13 R.J. Ascuitto, R.C. Braley and W.F. Ford, Nucl. Phys. A192 (1972) 97

- 1972BA1P Bassani et al., Communications, Proc. Aix-En-Provence Conf. 2 (1972) 68
- 1972BA1T Bartle and Guin, Bull. Amer. Phys. Soc. 17 (1972) 900
- 1972BA1U Baghuis, Ned. Tijdschr. Natuurk. (Netherlands) 38 (1972) 171; Phys. Abs. 49796 (1972)
- 1972BA1V Backe, Holtebekk and Tryti, Phys. Norv. (Norway) 6 (1972) 61
- 1972BA59 M.K. Basu, Phys. Rev. C6 (1972) 476
- 1972BA97 D.P. Balamuth, J.W. Noe, H.T. Fortune and R.W. Zurmuhle, Phys. Rev. C6 (1972) 1694
- 1972BE1E Bertsch and Mekjian, Ann. Rev. Nucl. Sci., 1972 (Ann. Rev. Inc., Palo Alto, California) 25
- 1972BO1H Bond et al., Bull. Amer. Phys. Soc. 17 (1972) 76
- 1972BO21 R. Bock and H. Yoshida, Nucl. Phys. A189 (1972) 177
- 1972BO38 M. Bouten and M.C. Bouten, Nucl. Phys. A193 (1972) 49
- 1972BR1G Braley, Ford, Becker and Patterson, NASA TN D 6834 (1972)
- 1972BU1G W.M. Bugg, G.T. Condo, E.L. Hart, M.R. Jhaveri, H.O. Cohn and R.D. McCulloch, Phys. Rev. C5 (1972) 2142
- 1972BU1H Budzanowski et al., INP 824 (1972)
- 1972CA1D Carpenter and van Staden, Bull. Amer. Phys. Soc. 17 (1972) 532
- 1972CA37 P. Camiz, E. Olivieri, M. Scalia and A. D'Andrea, Nuovo Cim. A12 (1972) 71
- 1972CL1A Clayton, Encyclopedia of the Twentieth Century (1972)
- 1972CO23 J.R. Comfort, W.J. Braithwaite, J.R. Duray, H.T. Fortune, W.J. Courtney and H.G. Bingham, Phys. Lett. B40 (1972) 456
- 1972CO30 W.J. Courtney and H.T. Fortune, Phys. Rev. C6 (1972) 1453
- 1972DE18 R. de Swinarski, A.D. Bacher, F.G. Resmini, G.R. Plattner and D.L. Hendrie, Phys. Rev. Lett. 28 (1972) 1139
- 1972DE1F de Oliveira and Mizrahi, Rev. Bras. Fis. (Brazil) 2 (1972) 311; Phys. Abs. 49156 (1973)
- 1972EC1A Eckhause et al., Nucl. Phys. B44 (1972) 83
- 1972ED01 G. Eder, G. Winkler and P. Hille, Z. Phys. 253 (1972) 335
- 1972EL1C Elliott, Proc. Roy. Soc. 326 (1972) 199
- 1972EM02 B. Eman, D. Tadic, F. Krmpotic and L. Szybisz, Phys. Rev. C6 (1972) 1
- 1972EY01 Y. Eyal, I. Dostrovsky and Z. Fraenkel, Nucl. Phys. A180 (1972) 545
- 1972FE1A Feng, Thesis, Univ. Minnesota (1972); Phys. Abs. 19373 (1974)
- 1972FI1B Fischer, Nucl. Instrum. Meth. 105 (1972) 413

- 1972FO11 H.T.Fortune, G.C.Morrison, R.C.Bearse, J.L.Yntema and B.H.Wildenthal, Phys. Rev. C6 (1972) 21
- 1972FO16 H.T. Fortune, R. Middleton and R.R. Betts, Phys. Rev. Lett. 29 (1972) 738
- 1972FO21 F. Foroughi and J. Rossel, Helv. Phys. Acta 45 (1972) 439
- 1972FO29 H.T. Fortune, Part. Nucl. 4 (1972) 245
- 1972FR1B Friedrich, Husken and Weiguny, Communications, Proc. Aix-En-Provence Conf. 2 (1972) 85
- 1972GA1E Garrett, Symp. on Two-Nucleon Transfer and Pairing Excitations, Argonne (1972) 232; (CONF-720309)
- 1972GO16 A.L. Goodman, Nucl. Phys. A186 (1972) 475
- 1972GR1G Grin and Leinson, Sov. J. Nucl. Phys. 14 (1972) 55
- 1972GR1H Grin, Kochetev and Anankin, Sov. J. Nucl. Phys. 14 (1972) 534
- 1972GR42 F. Grummer and A. Faessler, Z. Phys. 255 (1972) 112
- 1972HA07 O. Hausser, A.J. Ferguson, A.B. McDonald, I.M. Szoghy, T.K. Alexander and D.L. Disdier, Nucl. Phys. A179 (1972) 465
- 1972HA1Q Harvey, 1970 4th Symp. Struct. of Low-Medium Mass Nuclei (Univ. Press of Kansas, 1972) 1
- 1972HA1R Hanna, 1970 4th Symp. Struct. of Low-Medium Mass Nuclei (Univ. Press of Kansas, 1972) 30
- 1972HA25 G.R. Hammerstein, D. Larson and B.H. Wildenthal, Phys. Lett. B39 (1972) 176
- 1972HE24 D.W. Heikkinen, H.F. Lutz and W. Bartolini, Nucl. Phys. A193 (1972) 372
- 1972HE39 R. Hellborg and L. Ask, Phys. Scr. 6 (1972) 47
- 1972HI17 J. Hiura, F. Nemoto and H. Bando, Suppl. Prog. Theor. Phys. 52 (1972) 173
- 1972HI1C Hiddleston, Aymar and Darden, Bull. Amer. Phys. Soc. 17 (1972) 563
- 1972HO1D Horikawa, Prog. Theor. Phys. 47 (1972) 867
- 1972HO1E Horiuchi, Prog. Theor. Phys. 47 (1972) 1058
- 1972HO56 H. Horiuchi, K. Ikeda and Y. Suzuki, Suppl. Prog. Theor. Phys. 52 (1972) 89
- 1972IK1A Ikeda, Marumori, Tamagaki and Tanaka, Suppl. Prog. Theor. Phys. 52 (1972) 1
- 1972JA24 A. Jaffrin, Nucl. Phys. A196 (1972) 577
- 1972KA39 M. Kamimura, T. Matsuse and K. Takada, Prog. Theor. Phys. 47 (1972) 1537
- 1972KA67 M. Kamimura, T. Marumori and K. Takada, Suppl. Prog. Theor. Phys. 52 (1972) 282
- 1972KH08 S.B. Khadkikar and D.R. Kulkarni, Phys. Rev. C6 (1972) 866
- 1972KO1A Kocharov, Izv. Akad. Nauk SSSR Ser. Fiz. 36 (1972) 2052

- 1972KR11 S.J. Krieger and S.A. Moszkowski, Phys. Rev. C5 (1972) 1990
- 1972KR1D Krieger, 1970 4th Symp. on the Struct. of Low-Medium Mass Nuclei (Univ. Press of Kansas, 1972) 146
- 1972KU12 K.-I. Kubo and M. Hirata, Nucl. Phys. A187 (1972) 186
- 1972KU13 K.-I. Kubo, Nucl. Phys. A187 (1972) 205
- 1972KU1F Kulkarni and Pandya, Nucl. Phys. Solid State Phys. Symp. Ab., Bombay, 1972 (Bombay, India, Bhabha Atomic Res. Centre 1972); Phys. Abs. 38484 (1972)
- 1972KU24 H.M. Kuan, G.L. Latshaw, W.J. O'Connell, D.W. Heikkinen, E.G. Adelberger, A.V. Nero and S.S. Hanna, Nucl. Phys. A193 (1972) 497
- 1972LA08 K.R. Lassey and A.B. Volkov, Phys. Lett. B39 (1972) 169
- 1972LA18 S. Laribi, H. Beaumevieille, N. Bendjaballah, D. Lalanne, J.F. Allard and B. Faid, Nucl. Phys. A191 (1972) 368
- 1972LE13 T.Y. Lee, S.T. Hsieh and C.M. Yang, Phys. Rev. C5 (1972) 2013
- 1972LE38 H.C. Lee and R.Y. Cusson, Phys. Rev. Lett. 29 (1972) 1525; Erratum Phys. Rev. Lett. 30 (1973) 153
- 1972LI1E Litherland, Proc. 1971 Mont Tremblant Summer School Meeting on Dynamic Struct. of Nucl. States, Mont Tremblant, Canada, 1972 (Toronto, Canada, Univ. Toronto Press, 1972) 310; Phys. Abs. 82884 (1973)
- 1972LO1D Lobov, Izv. Akad. Nauk SSSR Ser. Fiz. 36 (1972) 881
- 1972MA07 N. Macdonald, I. Morrison and A. Watt, Nucl. Phys. A182 (1972) 183
- 1972MA20 S. Maripuu and B.H. Wildenthal, Phys. Lett. B38 (1972) 464
- 1972MA65 R.S. Mackintosh, Nucl. Phys. A198 (1972) 343
- 1972MI06 S. Mitsunobu and Y. Torizuka, Phys. Rev. Lett. 28 (1972) 920
- 1972MI11 G.H. Miller, M. Eckhause, P. Martin and R.E. Welsh, Phys. Rev. C6 (1972) 487
- 1972MO08 C.E. Moss, C. Detraz, C.S. Zaidins and D.J. Frantsvog, Phys. Rev. C5 (1972) 1122
- 1972MO15 R. Mogharrab and H. Neuert, Atomkernenergie 19 (1972) 107
- 1972MU08 H. Muther, K. Goeke and A. Faessler, Z. Phys. 253 (1972) 61
- 1972NA05 P.T. Nang, Nucl. Phys. A185 (1972) 413
- 1972NA11 K. Nakagawa, T. Obinata and K. Sasaki, Nucl. Phys. A191 (1972) 535
- 1972NE1B Nemoto and Bando, Prog. Theor. Phys. 47 (1972) 1210
- 1972NI14 R.A. Nisley and M.H. Hull, Jr., Nucl. Phys. A198 (1972) 561
- 1972OH01 I.K. Oh, C.S. Zaidins, C.D. Zafiratos and S.I. Hayakawa, Nucl. Phys. A178 (1972) 497
- 1972OL02 D.K. Olsen, W.R. Phillips and A.R. Barnett, Phys. Lett. B39 (1972) 201

- 1972OL04 D.K. Olsen, T. Udagawa, T. Tamura and R.E. Brown, Phys. Rev. Lett. 29 (1972) 1178; Erratum Phys. Rev. Lett. 30 (1973) 940
- 1972OP01 A.M.F. Op den Kamp and A.M.J. Spits, Nucl. Phys. A180 (1972) 569
- 1972PA16 A.D. Panagiotou, H.E. Gove and S. Harar, Phys. Rev. C5 (1972) 1995
- 1972PA1G Panagiotou et al., Communications, Proc. of Aix-En-Provence Conf. 2 (1972) 77
- 1972PA36 J.C. Parikh, Phys. Lett. B41 (1972) 468
- 1972PL1C Plasil, Communications, Proc. of Aix-En-Provence Conf. 2 (1972) 51
- 1972RE01 H. Rebel, Nucl. Phys. A180 (1972) 332
- 1972RE03 N.E. Reid, G.J. Stephenson and Jr., M.K. Banerjee, Phys. Rev. C5 (1972) 287
- 1972RE05 H. Rebel, G.W. Schweimer, G. Schatz, J. Specht, R. Lohken, G. Hauser, D. Habs and H. Klewe-Nebenius, Nucl. Phys. A182 (1972) 145
- 1972RI03 A.S. Rinat, Phys. Lett. B38 (1972) 281
- 1972RI1B Ripka, Sendai (1972) 7
- 1972RU03 C. Rudy, R. Vandenbosch, P. Russo and W.J. Braithwaite, Nucl. Phys. A188 (1972) 430.
- 1972SA04 L. Satpathy, K. Goeke and A. Faessler, Nucl. Phys. A179 (1972) 177
- 1972SA1B Satpathy, Schmid and Faessler, Communications, Proc. of Aix-Ev-Provence Conf. 2 (1972) 87
- 1972SA1C Sakakura, Sebe and Arima, Proc. Int. Conf. on Nucl. Struct. Studies Using Electron Scattering & Photoreaction, Sendai, Japan (1972)
- 1972SC03 U.C. Schlotthauer-Voos, H.G. Bohlen, W. von Oertzen and R. Bock, Nucl. Phys. A180 (1972) 385
- 1972SC10 L.R. Scherk and W.R. Falk, Nucl. Phys. A183 (1972) 240
- 1972SC17 U.C. Schlotthauer-Voos, R. Bock, H.G. Bohlen, H.H. Gutbrod and W. von Oertzen, Nucl. Phys. A186 (1972) 225
- 1972SC27 D. Schwalm, A. Bamberger, P.G. Bizzeti, B. Povh, G.A.P. Engelbertink, J.W. Olness and E.K. Warburton, Nucl. Phys. A192 (1972) 449
- 1972SH09 M.A. Sharaf, Phys. Lett. B40 (1972) 5
- 1972SH24 R.K. Sheline, Nucl. Phys. A195 (1972) 321
- 1972SI17 P.P. Singh, D.A. Sink, P. Schwandt, R.E. Malmin and R.H. Siemssen, Phys. Rev. Lett. 28 (1972) 1714
- 1972ST1C Steshenko and Filippov, Sov. J. Nucl. Phys. 14 (1972) 403
- 1972SU10 W. Sunkel and K. Wildermuth, Phys. Lett. B41 (1972) 439
- 1972TE02 S.N. Tewari, Phys. Rev. C6 (1972) 179

- 1972TH13 C. Thibault and R. Klapisch, Phys. Rev. C6 (1972) 1509
- 1972THZF H. Theissen, Springer Tracts in Modern Phys., Ed. Hohler, Vol. 65 (Berlin, Germany-Springer Verlag, 1972) 1
- 1972TO08 D.F. Torgerson, K. Wien and R.D. Macfarlane, Phys. Lett. B40 (1972) 203
- 1972VA1F Valamov et al., JETP Lett. 16 (1972) 224
- 1972VO02 H. Voit, G. Ischenko, F. Siller and H.-D. Helb, Nucl. Phys. A179 (1972) 23
- 1972VO09 E. Vogt, Phys. Lett. B40 (1972) 345
- 1972WH04 R.E. Whitehead, Nucl. Phys. A182 (1972) 290
- 1972WI13 F.C. Williams, Jr., G. Chan and J.R. Huizenga, Nucl. Phys. A187 (1972) 225
- 1972WI1C Wilkinson, Few Particle Problems, UCLA, 1972 (North-Holland, 1972) 191
- 1972WO10 G.J. Wozniak, H.L. Harney, K.H. Wilcox and J. Cerny, Phys. Rev. Lett. 28 (1972) 1278; Erratum Phys. Rev. Lett. 29 (1972) 760
- 1972YO1B Yoshida and Zamick, Ann. Rev. Nucl. Sci. (Ann. Rev. Inc., Palo Alto, Calif., 1972) 121
- 1973AB01 S.M. Abecasis, Nucl. Phys. A205 (1973) 475
- 1973AC1A Ackermann et al., J. Phys. Soc. Jpn. Suppl. 34 (1973) 215
- 1973AL1C Alexander, Fifth Symp. Struct. Low-Medium Mass Nuclei (Univ. Press of Kentucky, 1973) 272
- 1973AN30 A.N. Antonov and E.V. Inopin, Ukr. Fiz. Zh. 18 (1973) 1585; Phys. Abs. 78313 (1973)
- 1973AR18 A. Arima, R.A. Broglia, M. Ichimura and K. Schafer, Nucl. Phys. A215 (1973) 109
- 1973AR1C Arima, in Munich (1973) 184
- 1973AR1E Arnett, Ann. Rev. Astron. Astrophys. 11 (1973) 73
- 1973AR1F Arnett, in Asilomar (1973) 1083
- 1973AR1H Arnett, in Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 236
- 1973AU1D Audouze and Fricke, Astrophys. J. 186 (1973) 239
- 1973AV1B Avida, Calarco, Glavish and Hanna, Bull. Amer. Phys. Soc. 18 (1973) 677
- 1973BA10 J.K. Bair and F.X. Haas, Phys. Rev. C7 (1973) 1356
- 1973BA18 B.F. Bayman and D.H. Feng, Nucl. Phys. A205 (1973) 513
- 1973BA1T Bartle, Bull. Amer. Phys. Soc. 18 (1973) 1412
- 1973BA1U Baudinet-Robinet and Mahaux, Asilomar (1973) Paper 2A12
- 1973BA81 L.K. Batist, E.E. Berlovich, Y.S. Blinnikov, Y.V. Elkin, Y.N. Novikov, B.M. Ovchinnikov and V.K. Tarasov, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1944; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 124

- 1973BE11 T.A. Belote, N. Anyas-Weiss, J.A. Becker, J.C. Cornell, P.S. Fisher, P.N. Hudson, A. Menchaca-Rocha, A.D. Panagiotou and D.K. Scott, Phys. Rev. Lett. 30 (1973) 450
- 1973BE1H Berinde et al., in Munich 1 (1973) 555
- 1973BE35 G.F. Bertsch and B.H. Wildenthal, Phys. Rev. C8 (1973) 1023
- 1973BE53 A. Bertin, A. Vitale and A. Placci, Phys. Rev. A7 (1973) 2214
- 1973BO1K Body and Csikai, At. Energy Rev. 11 (1973) 153
- 1973BR15 R.C. Brown, J.A.R. Griffith, O. Karban, L. Mesko, J.M. Nelson and S. Roman, Nucl. Phys. A207 (1973) 456
- 1973BR1C Bromley, in Munich 2 (1973) 22
- 1973CA16 E. Caurier, B. Bourotte-Bilwes and Y. Abgrall, Phys. Lett. B44 (1973) 411
- 1973CA1B Cameron, Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 3
- 1973CA32 F. Catara and J.M.G. Gomez, Nucl. Phys. A215 (1973) 85
- 1973CH1E Chong and Tomusiak, Asilomar (1973) Paper 5E8
- 1973CH1F Chong and Tomusiak, Phys. in Canada 29 (1973) 5
- 1973CL1E Clayton and Woosley, in Munich 2 (1973) 718
- 1973CO03 M. Conze, H. Feldmeier and P. Manakos, Phys. Lett. B43 (1973) 101
- 1973CO1J Couch and Arnett, in Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 179
- 1973CO1K Conjeud, Harar, Ca Silveira and Volant, in Munich (1973) 474
- 1973CU03 R.Y. Cusson and H.C. Lee, Nucl. Phys. A211 (1973) 429
- 1973DE06 R. de Swiniarski, A. Genoux-Lubain, G. Bagieu, J.F. Cavaignac, D.H. Worledge and J. Raynal, Phys. Lett. B43 (1973) 27
- 1973DH1A Dhar, Kulkarni and Bhatt, in Munich 1 (1973) 75
- 1973DO02 E.I. Dolinsky, P.O. Dzhamalov and A.M. Mukhamedzhanov, Nucl. Phys. A202 (1973) 97
- 1973EB1A Eberhard, Eck, Schiele and Trombik, in Munich 1 (1973) 507
- 1973EL04 E. Elbaz, C. Fayard, G-H. Lamot, J. Meyer, R.S. Nahabetian, J. Pigeon and P. Boschán, Phys. Rev. C7 (1973) 1445
- 1973EN1C Engeland and Strottman, in Munich 1 (1973) 69
- 1973ER10 D.J. Ernst, C.M. Shakin and R.M. Thaler, Phys. Rev. C8 (1973) 440
- 1973ER1B Erb et al., Bull. Amer. Phys. Soc. 18 (1973) 1415
- 1973FE03 M. Feil, W. von Oertzen, H.G. Bohlen, A. Gamp, R.L. Walter and B. Kohlmeyer, Z. Phys. 260 (1973) 271
- 1973FI12 L.K. Fifield, R.W. Zurmuhle, D.P. Balamuth and J.W. Noe, Phys. Rev. C8 (1973) 2203

- 1973FO06 H.T. Fortune, R.R. Betts, J.D. Garrett and R. Middleton, Phys. Lett. B44 (1973) 65
- 1973FO1A Fortune, in Heavy Ion Lecture Series, Kansas State Univ. (1973) 1
- 1973FO1E Fortune, in Symp. on Heavy Ion Transfer Reactions, ANL Phys. B1 (1973) 287
- 1973FO1F Ford, Braley, Becker and Patterson, in Nucl. Many-Body Problem, Vol. 2, 1972 (Ed. Compositori, Bologna, 1973) 241
- 1973GA14 A. Gamp, W. von Oertzen, H.G. Bohlen, M. Feil, R.L. Walter and N. Marquardt, Z. Phys. 261 (1973) 283
- 1973GA23 M. Gari and A.H. Huffman, Nucl. Phys. A215 (1973) 532
- 1973GI09 J.N. Ginocchio, Phys. Rev. Lett. 31 (1973) 1260
- 1973GL1B Glavish, in Fifth Symp. Struct. Low-Medium Mass Nuclei, Univ. Press of Kentucky (1973) 233
- 1973GL1C Glavish, in Asilomar (1973) 755
- 1973GO1K Gove, Jundt and Lindgren, in Munich (1973) 166
- 1973GO1L Gove, Jundt, Lindgren and Trentelman, Bull. Amer. Phys. Soc. 18 (1973) 1391
- 1973GO1M Goldring, in Lect. Notes in Phys., 23 (Springer-Verlag, 1973) 178
- 1973GO26 K. Goeke, J. Garcia and A. Faessler, Nucl. Phys. A208 (1973) 477
- 1973GR11 R. Gross and I. Talmi, Phys. Lett. B44 (1973) 147
- 1973GR29 W. Gruhle, T. Bauer, T.H. Seligman and H.H. Hackenbroich, Z. Phys. 262 (1973) 271
- 1973GR36 Y.T. Grin and A.B. Kochetov, Yad. Fiz. 18 (1973) 283; Sov. J. Nucl. Phys. 18 (1974) 145
- 1973HA05 M. Harvey, Nucl. Phys. A202 (1973) 191
- 1973HA14 K.A. Hardy and Y.K. Lee, Phys. Rev. C7 (1973) 1441
- 1973HA1N Harvey and Khanna, in Munich, 1 (1973) 277
- 1973HA1Q Hanna, in Asilomar (1973) 417
- 1973HA1V Hausser, J. Phys. Soc. Jpn. Suppl. 34 (1973) 135
- 1973HA1W Harvey, in Munich 1 (1973) 44
- 1973HA1X Halpern, in Asilomar (1973) 909
- 1973HA63 O. Hausser, T.K. Alexander, D.L. Disdier, A.J. Ferguson, A.B. McDonald and I.S. Towner, Nucl. Phys. A216 (1973) 617
- 1973HA77 J.C. Hardy, Nucl. Data Tables A11 (1973) 327
- 1973HE1F Hecht, in Ann. Rev. Nucl. Sci., Vol. 23 (1973); Ann. Rev. Inc., Palo Alto (1973) 123
- 1973HO1G Howard, in Explosive Nucleosynthesis (Univ. Texas Press, 1973) p. 60
- 1973HO1H Horiuchi et al., in Munich 1 (1973) 121

- 1973HO1L Horiuchi, Prog. Theor. Phys. 50 (1973) 529
- 1973HO40 H. Horiuchi and Y. Suzuki, Prog. Theor. Phys. 49 (1973) 1974
- 1973HO41 Y. Horikawa, A. Nakada and Y. Torizuka, Prog. Theor. Phys. 49 (1973) 2005
- 1973IC01 M. Ichimura, A. Arima, E.C. Halbert and T. Terasawa, Nucl. Phys. A204 (1973) 225
- 1973IN01 P.D. Ingalls, Phys. Rev. C7 (1973) 464
- 1973IR01 J.M. Irvine, G.S. Mani, V. Pucknell, A. Watt and R.R. Whitehead, Phys. Lett. B44 (1973) 16
- 1973JO10 R.C. Johnson, F.D. Santos, R.C. Brown, A.A. Debenham, G.W. Greenlees, J.A.R. Griffith, O. Karban, D.C. Kocher and S. Roman, Nucl. Phys. A208 (1973) 221
- 1973JU1A Junkin and Suen, in Munich 1 (1973) 77
- 1973KE1C A.K. Kerman and M.S. Weiss, Phys. Rev. C8 (1973) 408
- 1973KO1F Kohler and Bethgz, in Munich 1 (1973) 134
- 1973KO26 D. Kolb, R.Y. Cusson and M. Harvey, Nucl. Phys. A215 (1973) 1
- 1973KO42 V.Y. Kostin, E.G. Kopanets, A.A. Koval, A.N. Lvov, V.Y. Migaleny and S.P. Tsytko, Izv. Akad. Nauk SSSR Ser. Fiz. 37 (1973) 1846; Bull. Acad. Sci. USSR Phys. Ser. 37 (1974) 48
- 1973KR1B Kravtsov and Skachkov, At. Data Nucl. Data Tables 12 (1973) 143
- 1973KR1E Kramer, Wildermuth and Beam, Part. Nucl. 5 (1973) 145
- 1973KU1E Kuhlmann, Mamis and Reiss, in Asilomar (1973) Paper 5E4
- 1973LA03 W.A. Lanford and B.H. Wildenthal, Phys. Rev. C7 (1973) 668
- 1973LA1D Lanford and Wildenthal, Bull. Amer. Phys. Soc. 18 (1973) 578
- 1973LA35 K.R. Lassey, M.R.P. Manning and A.B. Volkov, Can. J. Phys. 51 (1973) 2522
- 1973LE1D Lee, in Fifth Symp. Struct. Low-Medium Mass Nuclei, (Univ. Press of Kentucky, 1973) 20
- 1973LE1E Lee, J. Phys. Soc. Jpn. Suppl. 34 (1973) 445
- 1973LO13 D.S. Longo, J.C. Lawson, L.A. Alexander, B.P. Hichwa and P.R. Chagnon, Phys. Rev. C8 (1973) 1347
- 1973MA09 M.G. Mazarakis and W.E. Stephens, Phys. Rev. C7 (1973) 1280
- 1973MA14 R.L. Macklin and R.R. Winters, Phys. Rev. C7 (1973) 1766
- 1973MA1K Maripuu, in 5th Symp. Struct. Low-Medium Mass Nuclei, Univ. Press of Kentucky (1973) 63
- 1973MA48 F. Malaguti and P.E. Hodgson, Nucl. Phys. A215 (1973) 243
- 1973MA50 T. Matsuse and M. Kamimura, Prog. Theor. Phys. 49 (1973) 1765
- 1973MC06 J.B. McGrory and B.H. Wildenthal, Phys. Rev. C7 (1973) 974

1973MC17 C.A. McMahan and W. Tobocman, Nucl. Phys. A212 (1973) 465
 1973MC1E McGrory, in Munich 2 (1973) 146
 1973ME1D Meijer, Bull. Amer. Phys. Soc. 18 (1973) 578
 1973ME1F Menchaca-Rocha et al., in Munich 1 (1973) 477
 1973ME1G Meyer et al., in Nucl. Many-Body Problem, Vol. 1, 1972 (Ed. Compositori, Bologna, 1973) 176
 1973MI1E Middleton, Betts and Fortune, Bull. Amer. Phys. Soc. 18 (1973) 79
 1973MI1F Mihailovic, in Nucl. Many-Body Problem, Vol. 1, 1972 (Ed. Compositori, Bologna, 1973) 605
 1973MU14 S.F. Mughabghab and D.I. Garber, BNL 325, 3rd Edition, Vol. 1 (1973)
 1973NG1A Ng and Trainor, in Munich 1 (1973) 160
 1973NG1B Ng and Trainor, in Munich 1 (1973) 147
 1973NO1B Noto, Abe, Hiura and Tanaka, in Munich 1 (1973) 136
 1973OB04 A.W. Obst and K.W. Kemper, Phys. Rev. C8 (1973) 1682
 1973OG1A Ogloblin, Sov. J. Part. Nucl. 3 (1973) 467
 1973OL03 D.K. Olsen, T. Udagawa, T. Tamura and R.E. Brown, Phys. Rev. C8 (1973) 609
 1973PA05 J.C. Parikh, Ann. Phys. 76 (1973) 202
 1973PE1D Pelte, in Munich 2 (1973) 256
 1973PR01 J.G. Pronko, Phys. Rev. C7 (1973) 127
 1973PU1A Pucknell, Mani and Irvine, in Munich 1 (1973) 65
 1973RA1D Ramaty, Borner and Cohen, Astrophys. J. 181 (1973) 891
 1973RA1E Rayet, Nucl. Phys. B57 (1973) 269
 1973RE11 H. Rebel and G.W. Schweimer, Z. Phys. 262 (1973) 59
 1973RO29 J.C. Robertson, B. Audric and P. Kolkowski, J. Nucl. Energy 27 (1973) 531
 1973ROYN D.J. Rowe, in Asilomar (1973) 717
 1973SA1G Sandoval, Homyer and Duhn, in Munich 1 (1973) 553
 1973SA24 T.S. Sandhu and M.L. Rustgi, Phys. Rev. C8 (1973) 1152
 1973SC1B Scott, in Munich 2 (1973) 216
 1973SC1H Schwalm, Olness, Warburton and Engelbertink, in Munich 1 (1973) 282
 1973SC1J Scott, Symp. on Heavy Ion transfer Reactions, ANL Phys. B1 (1973) 97
 1973SC1K Scheid, Fink and Muller, in Proc. Europhys. 1972 Plitvice Lakes Conf. (Springer-Verlag, 1973) 144
 1973SE15 A. Sevgen, Z. Phys. 262 (1973) 221

- 1973SE17 A. Sevgen, Nucl. Phys. A216 (1973) 429
- 1973SI1D Sink, Schwandt and Singh, Bull. Amer. Phys. Soc. 18 (1973) 601
- 1973SI31 R.P. Singhal, H.S. Caplan, J.R. Moreira and T.E. Drake, Can. J. Phys. 51 (1973) 2125
- 1973SM1C Smirnov, Shitikova and Orlova, Moscow Univ. Phys. Bull. 28:5 (1973) 32
- 1973SP1A Sprung, in Nucl. Many-Body Problem, Vol. 2, 1972 (Ed. Compositori, Bologna, 1973) 123
- 1973ST1A Stokstad, in Proc. Europhys. 1972 Lakes Conf. (Springer-Verlag, 1973) 179
- 1973ST1B Stephenson, Hichwa and Meyer, Bull. Amer. Phys. Soc. 18 (1973) 1390
- 1973ST1E Staden and Bethge, in Munich 1 (1973) 473
- 1973ST1F Stokstad, in Symp. on Heavy Ion Transfer Reactions, ANL Phys. B1 (1973) 325
- 1973ST24 D. Strottman, N. Anyas-Weiss, J.C. Cornell, P.S. Fisher, P.N. Hudson, A. Menchaca-Rocha, A.D. Panagiotou and D.K. Scott, Phys. Lett. B47 (1973) 16
- 1973ST25 D. Strottman, Phys. Lett. B47 (1973) 215
- 1973SU1B Sugimoto, J. Phys. Soc. Jpn. Suppl. 34 (1973) 197
- 1973SU1D Suzuki, Ikeda and Horiuchi, in Munich 1 (1973) 119
- 1973SV1A Svenne, Reid and Davison, in Munich 1 (1973) 45
- 1973TH1B Theissen, in Asilomar (1973) 871
- 1973TO08 D.F. Torgerson, K. Wien, Y. Fares, N.S. Oakey, R.D. Macfarlane and W.A. Lanford, Phys. Rev. C8 (1973) 161
- 1973TO16 L. Tomlinson, At. Data Nucl. Data Tables 12 (1973) 179
- 1973TR1B Truran, in Cosmochem., Ed. Cameron (Reidel Pub. Co., 1973) 23
- 1973UD01 T. Udagawa and D.K. Olsen, Phys. Lett. B46 (1973) 285
- 1973UL01 N. Ullah and K.R. Sandhya Devi, Phys. Rev. C8 (1973) 1167
- 1973UN01 T. Une, Prog. Theor. Phys. 49 (1973) 1587
- 1973VA01 D. Vautherin, Phys. Rev. C7 (1973) 296
- 1973VA1H Vautherin, in Munich 2 (1973) 108
- 1973VA21 J.P. Vary and C.B. Dover, Phys. Rev. Lett. 31 (1973) 1510
- 1973VO01 H. Voit, G. Ischenko and F. Siller, Phys. Rev. Lett. 30 (1973) 564
- 1973VO1G Vorobyov et al., in Munich 1 (1973) 716
- 1973WA10 E.K. Warburton, J.W. Olness, G.A.P. Engelbertink and T.K. Alexander, Phys. Rev. C7 (1973) 1120
- 1973WA19 E.K. Warburton, P. Gorodetzky and J.A. Becker, Phys. Rev. C8 (1973) 418
- 1973WI11 D.H. Wilkinson, Nucl. Phys. A209 (1973) 470

- 1973WI15 J. Wilczynski, Phys. Lett. B47 (1973) 124
- 1973YA1A Yazaki, Prog. Theor. Phys. 49 (1973) 1205
- 1973YO03 H. Yoshida, Phys. Lett. B47 (1973) 411
- 1973ZU1A R.W. Zurmuhle, D.P. Balamuth, L.K. Fifield and J.W. Noe, Phys. Lett. B44 (1973) 453
- 1974AB05 Y. Abgrall, P. Gabinski and J. Labarsouque, Nucl. Phys. A232 (1974) 235
- 1974AD1B Adelberger, Int. Conf. Nucl. Struct. Spectrosc., Pt. II, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 641
- 1974AL11 D.E. Alburger and D.R. Goosman, Phys. Rev. C10 (1974) 912
- 1974AR04 A. Arima and S. Yoshida, Nucl. Phys. A219 (1974) 475
- 1974AR1G Arnett, Astrophys. J. 194 (1974) 373
- 1974AR1H Arnett, Astrophys. J. 193 (1974) 169
- 1974AY02 S. Ayik and J.N. Ginocchio, Nucl. Phys. A234 (1974) 13
- 1974BA15 G.C. Ball, J.G. Costa, W.G. Davies, J.S. Forster, J.C. Hardy and A.B. McDonald, Phys. Lett. B49 (1974) 33
- 1974BA76 R. Ballini, N. Cindro, J.P. Fouan, C. Kalbach, M. Lepareux and N. Saunier, Nucl. Phys. A234 (1974) 33
- 1974BE41 F.E. Bertrand, R.W. Peelle and C. Kalbach-Cline, Phys. Rev. C10 (1974) 1028
- 1974BI1D Bishop and Fortune, Bull. Amer. Phys. Soc. 19 (1974) 16
- 1974BO05 J.D. Bowman, A.M. Poskanzer, R.G. Korteling and G.W. Butler, Phys. Rev. C9 (1974) 836
- 1974BO1E Bormann, Neuert and Scobel, IAEA, STI/DOC/10/156 (1974) 87
- 1974CA1J Calamand, IAEA, STI/DOC/10/156 (1974) 273
- 1974CA22 R. Caracciolo, P. Cuzzocrea, A. De Rosa, G. Inghima, E. Perillo, M. Sandoli and G. Spadaccini, Lett. Nuovo Cim. 11 (1974) 33
- 1974CA31 E. Casal, A. Garcia and F. Senent, An. Fis. 70 (1974) 18
- 1974CE1A Cerny, in Nashville Vol. 2 (1974) 483
- 1974CO39 B.J. Cole, A. Watt and R.R. Whitehead, J. Phys. A7 (1974) 1374
- 1974CO40 B.J. Cole, A. Watt and R.R. Whitehead, J. Phys. A7 (1974) 1399
- 1974CR04 D.J. Crozier and H.T. Fortune, Phys. Rev. C10 (1974) 1697
- 1974DE05 R. de Swiniarski, G. Bagieu, A.J. Cole, P. Gaillard, A. Guichard, J.Y. Grossiord, M. Gusakow and J.R. Pizzi, J. Phys. (Paris), Suppl. Lett. 35 (1974) L25
- 1974DE1E De Jager, De Vries and De Vries, At. Data Nucl. Data Tables 14 (1974) 479

- 1974DE46 R. de Swinarski, A. Genoux-Lubain, G. Bagieu and J.F. Cavaignac, *Can. J. Phys.* 52 (1974) 2422
- 1974DI1D Dilg, *Z. Naturf. A29* (1974) 1750
- 1974DJ01 A. Djaloeis, D. Ingham, H. Kelleter, O. Aspelund and C. Mayer-Boricke, *Z. Phys.* 269 (1974) 133
- 1974DO03 E.I. Dolinskii, V.V. Turovtsev and R. Yarmukhamedov, *Yad. Fiz.* 19 (1974) 1003; *Sov. J. Nucl. Phys.* 19 (1975) 514
- 1974DZ05 R.I. Dzhibuti and R.Y. Kezerashvili, *Yad. Fiz.* 20 (1974) 340; *Sov. J. Nucl. Phys.* 20 (1975) 181
- 1974EN10 R. Engfer, H. Schneuwly, J.L. Vuilleumier, H.K. Walter and A. Zehnder, *At. Data Nucl. Data Tables* 14 (1974) 509
- 1974EP01 M.B. Epstein, D.J. Margaziotis, N.S.P. King and T.A. Cahill, *Phys. Rev. C9* (1974) 581
- 1974EP1B Epstein et al., *Bull. Amer. Phys. Soc.* 19 (1974) 453
- 1974ER1A Erb et al., in *Proc. Conf. Reactions Between Complex Nuclei*, 1 (North-Holland, 1974) 103
- 1974FO15 J.L.C. Ford, Jr., J. Gomez del Campo, R.L. Robinson, P.H. Stelson and S.T. Thornton, *Z. Phys.* 269 (1974) 147
- 1974FO1J Fortune, *Proc. Int. Conf. Nucl. Struct. Spectroscopy*, Amsterdam, 1974 (Scholar's Press, Amsterdam, 1974) 341
- 1974FO20 H.T. Fortune, L.R. Greenwood, R.E. Segel and J.R. Erskine, *Phys. Lett.* B52 (1974) 51
- 1974FO21 H.T. Fortune and R.R. Betts, *Phys. Rev. C10* (1974) 1292
- 1974FO25 H.T. Fortune, R.R. Betts and R. Middleton, *Phys. Rev. C10* (1974) 2135
- 1974FR06 H. Friedrich, *Nucl. Phys. A224* (1974) 537; *Erratum Nucl. Phys. A239* (1975) 531
- 1974GO12 V.Z. Goldberg, V.P. Rudakov and V.A. Timofeev, *Yad. Fiz.* 19 (1974) 503; *Sov. J. Nucl. Phys.* 19 (1974) 253
- 1974GO37 V.Z. Goldberg, K.A. Gridnev and V.M. Semenov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2524; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 52
- 1974GR1M Griffith, Haff and Tombrello, *Ann. Phys.* 87 (1974) 1
- 1974GR27 L.R. Greenwood, H.T. Fortune, R.E. Segel and J.R. Erskine, *Phys. Rev. C10* (1974) 1211
- 1974HA11 D.L. Hanson, R.G. Stokstad, K.A. Erb, C. Olmer and D.A. Bromley, *Phys. Rev. C9* (1974) 929
- 1974HA17 R.U. Haq and J.C. Parikh, *Nucl. Phys. A220* (1974) 349

- 1974HA1C Hanna, Int. Conf. Nucl. Struct. Spectrosc., Pt. II, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 249
- 1974HA1G Hanna, Nukelonika (Poland) 19 (1974) 655
- 1974HA1N Hanna, Private Communication (1974)
- 1974HA25 D.L. Hanson, R.G. Stokstad, K.A. Erb, C. Olmer, M.W. Sachs and D.A. Bromley, Phys. Rev. C9 (1974) 1760
- 1974HA61 G.D. Harp, Phys. Rev. C10 (1974) 2387
- 1974HU07 J.R. Huizenga, A.N. Behkami, J.S. Sventek and R.W. Atcher, Nucl. Phys. A223 (1974) 577
- 1974HU14 J. Hufner, L. Tauscher and C. Wilkin, Nucl. Phys. A231 (1974) 455
- 1974IT1A Itoh and Anderson, Private Communication (1974)
- 1974JE1A Jelley, Wozniak, Nagarajan and Cerny, Bull. Amer. Phys. Soc. 19 (1974) 431
- 1974JO1H Johnson, Larson and Harvey, Bull. Amer. Phys. Soc. 19 (1974) 1030
- 1974KA1P Kamimura and Matsuse, Prog. Theor. Phys. 51 (1974) 438
- 1974KA1Q B. Kayser, G.T. Garvey, E. Fischbach and S.P. Rosen, Phys. Lett. B52 (1974) 385
- 1974KE18 M.J. Kenny, P.W. Martin, L.E. Carlson and J.A. Biggerstaff, Aust. J. Phys. 27 (1974) 759
- 1974KH03 S.B. Khadkikar and D.R. Kulkarni, Phys. Rev. C10 (1974) 1189
- 1974KL05 H.V. Klapdor, H. Reiss, G. Rosner and M. Schrader, Phys. Lett. B49 (1974) 431
- 1974KL13 H.V. Klapdor, H. Reiss and G. Rosner, Phys. Lett. B53 (1974) 147
- 1974KL1B Klapdor, Reiss and Rosner, Proc. Conf. Reactions between Complex Nuclei, Vol. I (North-Holland, 1974) 92
- 1974KO35 E. Kondaiah, J. Phys. 7A (1974) 1457
- 1974KR17 S. Krewald, K.W. Schmid, A. Faessler and J.B. McGrory, Nucl. Phys. A228 (1974) 524
- 1974KR1D Krimmelbein, Schober and Waffler, Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974) 434
- 1974KR1E Krause, Scheid and Greiner, Nukleonika (Poland) 19 (1974) 267; Phys. Abs. 36360 (1975)
- 1974KU07 K.I. Kubo, F. Nemoto and H. Bando, Nucl. Phys. A224 (1974) 573
- 1974LA27 H. Laumer and G.G. Seaman, Phys. Rev. C10 (1974) 2159
- 1974LI1D Lieb et al., Bull. Amer. Phys. Soc. 19 (1974) 594
- 1974LO1B Lorenzen and Brune, IAEA, STI/DOC/10/156 (1974) 325
- 1974MA11 R.S. Mackintosh and L.J. Tassie, Nucl. Phys. A222 (1974) 187

- 1974MA1E Maguin, *Nuovo Cim.* A19 (1974) 638
- 1974MA1K Mankoc-Borstnik and Mihailovic, *Fizika Suppl. (Yugoslavia)* 6 (1974) 3
- 1974MA38 N. Marquardt, R. Volders, C. Cardinal and J. L'Ecuyer, *Phys. Rev. Lett.* 33 (1974) 1389
- 1974MC19 W.F. McGill, R.F. Carlson, T.H. Short, J.M. Cameron, J.R. Richardson, I. Slaus, W.T.H. van Oers, J.W. Verba, D.J. Margaziotis and P. Doherty, *Phys. Rev.* C10 (1974) 2237
- 1974MC1F McGrory, *Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam, 1974 (Amsterdam, Scholars Press, 1974)* 73
- 1974ME09 A. Menchaca-Rocha, *Nucl. Instrum. Meth.* 114 (1974) 425
- 1974ME1C Medsker et al., *Bull. Amer. Phys. Soc.* 19 (1974) 471
- 1974MI13 G.F. Millington, J.R. Leslie, W. McLatchie, G.C. Ball, W.G. Davies and J.S. Forster, *Nucl. Phys.* A228 (1974) 382
- 1974MI21 T. Minamisono, Y. Nojiri, A. Mizobuchi and K. Sugimoto, *Nucl. Phys.* A236 (1974) 416
- 1974MO01 C.A. Mosley and Jr., H.T. Fortune, *Phys. Rev.* C9 (1974) 775
- 1974MP01 P. Mpanias and M.L. Rustgi, *Phys. Rev.* C9 (1974) 2261
- 1974MU1D Murphy and Benson, *Bull. Amer. Phys. Soc.* 19 (1974) 1013
- 1974NE03 J.M. Nelson and W.R. Falk, *Nucl. Phys.* A218 (1974) 441
- 1974NE18 V.G. Neudachin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 38 (1974) 2010; *Bull. Acad. Sci. USSR Phys. Ser.* 38 (1974) 1
- 1974NG01 W.Y. Ng and L.E.H. Trainor, *Can. J. Phys.* 52 (1974) 541
- 1974NO14 H. Noto, Y. Abe, J. Hiura and H. Tanaka, *Prog. Theor. Phys.* 52 (1974) 339
- 1974OB02 A.W. Obst and K.W. Kemper, *Phys. Rev.* C9 (1974) 1643
- 1974OL01 D.K. Olsen, A.R. Barnett, S.F. Biagi, N.H. Merrill and W.R. Phillips, *Nucl. Phys.* A220 (1974) 541
- 1974PA10 L.M. Panggabean, S.M. Austin and H. Laumer, *Phys. Rev.* C10 (1974) 1605
- 1974PA16 A.D. Panagiotou, J.C. Cornell, N. Anyas-Weiss, P.N. Hudson, A. Menchaca Rocha, D.K. Scott and B.E.F. Macefield, *J. Phys. (London)* A7 (1974) 1748
- 1974PA1E Pardo, Couch and Arnett, *Astrophys. J.* 191 (1974) 711
- 1974PI11 C.F. Pineda, M.L. de Toledo and C. Ruiz Bauza, *An. Fis.* 70 (1974) 30
- 1974RA11 J.R. Radin, A.R. Smith and N. Little, *Phys. Rev.* C9 (1974) 1718
- 1974RE03 N.E. Reid, N.E. Davison and J.P. Svenne, *Phys. Rev.* C9 (1974) 1882

- 1974RI1B Ripka and Kassis, in Proc. Conf. Reactions between Complex Nuclei, Vol. 1 (North-Holland, 1974) 182
- 1974RO03 V.C. Rogers, Phys. Rev. C9 (1974) 527
- 1974RO04 H.H. Rossner, G. Hinderer, A. Weidinger and K.A. Eberhard, Nucl. Phys. A218 (1974) 606
- 1974RO17 R.G.H. Robertson, S. Martin, W.R. Falk, D. Ingham and A. Djalois, Phys. Rev. Lett. 32 (1974) 1207
- 1974SC10 K.W. Schmid, L. Satpathy and A. Faessler, Z. Phys. 267 (1974) 337
- 1974SC11 K.W. Schmid, L. Satpathy and A. Faessler, Z. Phys. 267 (1974) 345
- 1974SC34 K.W. Schmid, S. Krewald, A. Faessler and L. Satpathy, Z. Phys. 271 (1974) 149
- 1974SE01 S. Sen, S.E. Darden, H.R. Hiddleston and W.A. Yoh, Nucl. Phys. A219 (1974) 429
- 1974SE1B Serov, Izv. Akad. Nauk SSSR Ser. Fiz. 38 (1974) 84
- 1974SH16 D. Shapira, R.G. Stokstad and D.A. Bromley, Phys. Rev. C10 (1974) 1063
- 1974SH1E Sher, IAEA, STI/DOC/10/156 (1974) 1
- 1974SHYR V.S. Shirley and C.M. Lederer, LBL-3450 (1974)
- 1974SI27 U.N. Singh, H.I. Liou, J. Rainwater, G. Hacken and J.B. Garg, Phys. Rev. C10 (1974) 2147
- 1974SP04 A.M.J. Spits and J. de Boer, Nucl. Phys. A224 (1974) 517
- 1974SP06 H. Spinka and H. Winkler, Nucl. Phys. A233 (1974) 456
- 1974ST10 H.-J. Stockmann, H. Ackermann, D. Dubbers, M. Grupp and P. Heitjans, Z. Phys. 269 (1974) 47
- 1974ST1L Stokstad, in Nashville, 2 (1974) 327
- 1974TA17 F. Tanabe and F. Nemoto, Prog. Theor. Phys. 51 (1974) 2009
- 1974TA18 L. Tauscher and W. Schneider, Z. Phys. 271 (1974) 409
- 1974TA19 K. Takada and S. Tazaki, Prog. Theor. Phys. 52 (1974) 1205
- 1974TH01 C. Thibault and R. Klapisch, Phys. Rev. C9 (1974) 793
- 1974TI03 C. Titin-Schnaider and P. Quentin, Phys. Lett. B49 (1974) 213
- 1974TI04 C. Titin-Schnaider and P. Quentin, Phys. Lett. B49 (1974) 397
- 1974UL02 H. Ullrich, E.T. Boschitz, H.D. Engelhardt and C.W. Lewis, Phys. Rev. Lett. 33 (1974) 433
- 1974VA18 R. Vandenbosch, M.P. Webb and M.S. Zisman, Phys. Rev. Lett. 33 (1974) 842
- 1974VA31 V. Valkovic, R.B. Liebert and G.C. Phillips, Nucl. Instrum. Meth. 122 (1974) 533
- 1974VE03 T. Vertse, A. Dudek-Ellis, P.J. Ellis, T.A. Belote and D. Roaf, Nucl. Phys. A223 (1974) 207

- 1974VE06 A. Veyssiere, H. Beil, R. Bergere, P. Carlos, A. Lepretre and A. de Miniac, Nucl. Phys. A227 (1974) 513
- 1974VO09 H. Voit, P. Duck, W. Galster, E. Haindl, G. Hartmann, H.-D. Helb, F. Siller and G. Ischenko, Phys. Rev. C10 (1974) 1331
- 1974VO11 G. Vourvopoulos, M.B. Greenfield, J.B. Ball, S. Raman and W.K. Dagenhart, Phys. Lett. B52 (1974) 187
- 1974WA1D W.L. Wang, Phys. Lett. B52 (1974) 143
- 1974WI05 R. Wieland, R. Stokstad, A. Gobbi, D. Shapira, L. Chua, M.W. Sachs and D.A. Bromley, Phys. Rev. C9 (1974) 1474
- 1974WI1L Wilkinson, Private Communication (1974)
- 1974WO02 S.S.M. Wong, D.J. Rowe and J.C. Parikh, Phys. Lett. B48 (1974) 403
- 1974YA11 Y. Yamamoto, Prog. Theor. Phys. 52 (1974) 471
- 1974ZU1A Zuker, Proc. Int. Conf. Nucl. Struct. Spectroscopy, Amsterdam. 1974 (Amsterdam, Scholars Press, 1974) 115
- 1975AB04 Y. Abgrall and E. Caurier, Phys. Lett. B56 (1975) 229
- 1975AB1H Abegg and Cavis, Bull. Amer. Phys. Soc. 20 (1975) 1181
- 1975AL19 J.M. Alexander, H. Delagrange and A. Fleury, Phys. Rev. C12 (1975) 149
- 1975AL27 D.E. Alburger and F.P. Calaprice, Phys. Rev. C12 (1975) 1690
- 1975AN13 N. Anantaraman, C.L. Bennett, J.P. Draayer, H.W. Fulbright, H.E. Gove and J. Toke, Phys. Rev. Lett. 35 (1975) 1131
- 1975AR14 A.G. Artyukh, V.V. Volkov, G.F. Gridnev and V.L. Mikheev, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 2; Bull. Acad. Sci. USSR Phys. Ser. 39 (1975) 1
- 1975AR17 O. Artun, Y. Cassagnou, R. Legrain, N. Lisbona, L. Roussel, J.P. Alard, A. Baldit, J.P. Costilhes, J. Fargeix, G. Roche et al., Phys. Rev. Lett. 35 (1975) 773
- 1975AR1J Arima, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 38
- 1975AR20 K.P. Artemov, V.Z. Goldberg, I.P. Petrov, V.P. Rudakov, I.N. Serikov and V.A. Timofeev, Yad. Fiz. 21 (1975) 1157; Sov. J. Nucl. Phys. 21 (1976) 596
- 1975AR25 K.P. Artemov, V.Z. Goldberg, I.P. Petrov, I.N. Serikov, V.P. Rudakov, V.A. Timofeev, R. Wolski and J. Szmider, Yad. Fiz. 22 (1975) 242; Sov. J. Nucl. Phys. 22 (1976) 125
- 1975BA05 S. Barshay, C.B. Dover and J.P. Vary, Phys. Rev. C11 (1975) 360
- 1975BA1Q Batist et al., Sov. J. Nucl. Phys. 20 (1975) 453
- 1975BA2D Barket, Ann. Rev. Astron. Astrophys. 13 (1975) 45
- 1975BA2H Bachelier et al., in Clustering Phenomena in Nuclei, II, ORO-4856-26 (1975) 313
- 1975BA2J Baltz, Dover and Vary, Bull. Amer. Phys. Soc. 20 (1975) 1168

- 1975BA81 B.R. Barrett, E.C. Halbert and J.B. McGrory, *Ann. Phys.* 90 (1975) 321
- 1975BE02 R.R. Betts, H.T. Fortune and R. Middleton, *Phys. Rev. C*11 (1975) 19
- 1975BE15 Z. Berant, C. Broude, G. Engler, M. Hass, R. Levy and B. Richter, *Nucl. Phys.* A243 (1975) 519
- 1975BE1G Bertozzi, in *LASL, AIP Conf. Proc.* 26 (1975) 409
- 1975BE31 M. Beiner, R.J. Lombard and D. Mas, *Nucl. Phys.* A249 (1975) 1
- 1975BI04 J.N. Bishop and H.T. Fortune, *Phys. Rev. Lett.* 34 (1975) 1350
- 1975BI1H Billen, *Bull. Amer. Phys. Soc.* 20 (1975) 1181
- 1975BI1J Bishop, *Bull. Amer. Phys. Soc.* 20 (1975) 1174
- 1975BR1F Brown, *At. Data Nucl. Data Tables* 15 (1975) 111
- 1975BU03 B. Buck, C.B. Dover and J.P. Vary, *Phys. Rev. C*11 (1975) 1803
- 1975BU1K Busza, in *Lasl, AIP Conf. Proc.* 26 (1975) 211
- 1975BU1L Buck, in *Lect. Notes in Phys.* 33 (Springer-Verlag, 1975) 152
- 1975CE01 R. Ceuleneer, F. Michel, M. Bosman, J. Lega, P. Leleux, P.C. Macq, J.P. Meulders and C. Pirart, *Phys. Rev. C*11 (1975) 631
- 1975CH17 W.S. Chien, C.H. King, J.A. Nolen, Jr. and M.A.M. Shahabuddin, *Phys. Rev. C*12 (1965) 332
- 1975CH41 I.I. Chkalov, N.G. Shevchenko, A.Y. Buki, A.A. Khomich, A.S. Litvinenko and V.N. Polishchuk, *Yad. Fiz.* 22 (1975) 893; *Sov. J. Nucl. Phys.* 22 (1976) 464
- 1975CO15 M. Conjeaud, S. Harar, E.F. Da Silveira and C. Volant, *Nucl. Phys.* A250 (1975) 182
- 1975CO18 M.N.H. Comsan, A.A. El-Kamhawy, F. Asfour and M. Abd-El-Fattah, *Atomkernen.* 26 (1975) 37
- 1975CO1K Cowley et al., in *Clustering Phenomena in Nuclei, II, ORO-4856-26* (1975) 238
- 1975CO1M Cowley et al., in *JUL-CONF-16* (1975) 102
- 1975CU1B Cusson and Hilko, *Bull. Amer. Phys. Soc.* 20 (1975) 67
- 1975DA1E Davis and Abegg, *Bull. Amer. Phys. Soc.* 20 (1975) 1181
- 1975DA1F Datta, berg and Quin, *Bull. Amer. Phys. Soc.* 20 (1975) 1162
- 1975DI08 G. Dietl, G. Gruber, H. Schmidt-Bocking and K. Bethge, *Nucl. Phys.* A250 (1975) 322
- 1975DO06 H. Doubre, E. Plagnol, J.C. Roynette, J.M. Loiseaux, P. Martin and P. de Saintignon, *J. Phys. (Paris), Suppl. Lett.* 36 (1975) L-113
- 1975DO10 T.W. Donnelly, J. Dubach and W.C. Haxton, *Nucl. Phys.* A251 (1975) 353
- 1975DR01 J.P. Draayer, *Nucl. Phys.* A237 (1975) 157

- 1975EL1A J.R. Elliott, L.R. Fortney, A.T. Goshaw, J.W. Lamsa, J.S. Loos, W.J. Robertson, W.D. Walker, W.M. Yeager and M. E. Binkley, Phys. Rev. Lett. 34 (1975) 607; Erratum Phys. Rev. Lett. 34 (1975) 920
- 1975EN1A Endal, Astrophys. J. 195 (1975) 187
- 1975EN1C Eng et al., Bull. Amer. Phys. Soc. 20 (1975) 694
- 1975FA1A Fagg, Rev. Mod. Phys. 47 (1975) 683
- 1975FO19 W.A. Fowler, G.R. Caughlan and B.A. Zimmerman, Ann. Rev. Astron. Astrophys. 13 (1975) 69
- 1975FU1D Fu, Nucl. Cross Sections and Tech. Pt. I, Washington, D.C. 1975 (Washington D.C., Nat. Bur. Standards 1975) 328; Phys. Abs. 23045 (1977)
- 1975GI03 J.N. Ginocchio and M.M. Yen, Nucl. Phys. A239 (1975) 365
- 1975GI10 H.J. Gils and H. Rebel, Z. Phys. A274 (1975) 259
- 1975GO15 V.Z. Goldberg, V.P. Rudakov and V.A. Timofeev, Yad. Fiz. 21 (1975) 1001; Sov. J. Nucl. Phys. 21 (1975) 513
- 1975GR1M B. Grammaticos, Phys. Lett. B57 (1975) 306
- 1975GR21 L.R. Greenwood, R.E. Segel, K. Raghunathan, M.A. Lee, H.T. Fortune and J.R. Erskine, Phys. Rev. C12 (1975) 156
- 1975HA1U Harvey, Ann. Phys. 94 (1975) 47
- 1975HA1V Harvey, in Clustering Phenomena in Nuclei, II, ORO-4856-26 (1975) p. 549
- 1975HE10 K.T. Hecht and D. Braunschweig, Nucl. Phys. A244 (1975) 365
- 1975HE1L Hecht and Sato, in Clustering Phenomena in Nuclii, II, ORO-4856-26 (1975) 454
- 1975HO15 R.E. Horstman, J.L. Eberhardt, H.A. Doubt, C.M.E. Otten and G. Van Middelkoop, Nucl. Phys. A248 (1975) 291
- 1975HO1H Hodgson, Nature 257 (1975) 446
- 1975HO1K Hornyak, Chang and Walters, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 354
- 1975HO1L Hodgson, Nature 257 (1975) 181
- 1975HU1D Hufner, Phys. Rept. C21 (1975) 1
- 1975IB1A Ibragimov and Kocharov, Izv. Akad. Nauk SSSR Ser. Fiz. 39 (1975) 287
- 1975IK1A Ikeda and Suzuki, in Clustering Phenomena in Nuclii, II, ORO-4856-26 (1975) 543
- 1975JE02 N.A. Jelley, J. Cerny, D.P. Stahel and K.H. Wilcox, Phys. Rev. C11 (1975) 2049
- 1975KL05 H.V. Klapdor, G. Rosner, H. Reiss and M. Schrader, Nucl. Phys. A244 (1975) 157
- 1975KL08 H.V. Klapdor, H. Reiss and G. Rosner, Phys. Lett. B58 (1975) 279
- 1975KO1K Koester and Nistler, Z. Phys. A272 (1975) 189

- 1975KO28 Y. Kondo, S. Nagata, S. Ohkubo and O. Tanimura, *Prog. Theor. Phys.* 53 (1975) 1006
- 1975KU1K Kuznetsov and Chernov, *Sov. J. Nucl. Phys.* 20 (1975) 340
- 1975LA1E Lavrukhina, *Izv. Akad. Nauk SSSR Ser. Fiz.* 39 (1975) 395
- 1975LE05 G. Leander and S.E. Larsson, *Nucl. Phys.* A239 (1975) 93
- 1975LE1K Lefevre, Davis and Anderson, *Bull. Amer. Phys. Soc.* 20 (1975) 584
- 1975LE23 J. L'Ecuyer, R. Volders, C. Cardinal, L. Deschenes and N. Marquardt, *Phys. Rev.* C12 (1975) 1878
- 1975MA10 N. Mankoc-Borstnik, M.V. Mihailovic and M. Rosina, *Nucl. Phys.* A239 (1975) 321
- 1975MA1R Maehl, Fisher, Hagan and Ormes, *Astrophys. J.* 202 (1975) L119
- 1975MA1W Mankoc-Borstnik, Mihailovic and Rosina, *Fizika Suppl. (Yugoslavia)* 7 (1975) 12
- 1975MA25 R.S. Mackintosh and R. De Swinarski, *Phys. Lett.* B57 (1975) 139
- 1975MA26 T. Matsuse, M. Kamimura and Y. Fukushima, *Prog. Theor. Phys.* 53 (1975) 706
- 1975MC1J McDaniel and Quin, *Bull. Amer. Phys. Soc.* 20 (1975) 1162
- 1975ME04 L.R. Medsker, H.T. Fortune, R.R. Betts and R. Middleton, *Phys. Rev.* C11 (1975) 1880
- 1975MI02 M.V. Mihailovic and M. Rosina, *Nucl. Phys.* A237 (1975) 221
- 1975MI03 M.V. Mihailovic and M. Rosina, *Nucl. Phys.* A237 (1975) 229
- 1975MI11 B. Mithra and R. Laverriere, *Lett. Nuovo Cim.* 14 (1975) 13
- 1975MO04 J.M. Moss, C.M. Rozsa, D.H. Youngblood, J.D. Bronson and A.D. Bacher, *Phys. Rev. Lett.* 34 (1975) 748
- 1975MP01 P. Mpanias and M.L. Rustgi, *Phys. Rev.* C12 (1975) 2072
- 1975MU13 H. Muther, A. Faessler and R.K. Tripathi, *Nucl. Phys.* A255 (1975) 132
- 1975NA20 Y.V. Naumov and O.E. Kraft, *Izv. Akad. Nauk SSSR Ser. Fiz.* 39 (1975) 1656; *Bull. Acad. Sci. USSR Phys. Ser.* 39 (1975) 76
- 1975NA21 Y.V. Naumov and O.E. Kraft, *Fiz. Elem. Chastits At. Yad. (USSR)* 6 (1975) 892; *Sov. J. Part. Nucl.* 6 (1976) 361
- 1975NE04 F. Nemoto, Y. Yamamoto, H. Horiuchi, Y. Suzuki and K. Ikeda, *Prog. Theor. Phys.* 54 (1975) 104
- 1975PA1K Park, in *Clustering Phenomena in Nuclei, II*, ORO-4856-26 (1975) 293
- 1975PE11 J.F. Petersen, *Nucl. Phys.* A255 (1975) 307
- 1975PE1E Peimbert, *Ann. Rev. Astron. Astrophys.* 13 (1975) 113
- 1975PF01 H.-J. Pfeiffer, K. Springer and H. Daniel, *Nucl. Phys.* A254 (1975) 433
- 1975PO01 F. Pougheon, G. Rotbard, P. Roussel and J. Vernotte, *Phys. Rev. Lett.* 34 (1975) 158

- 1975PU02 F. Puhlhofer, W. Pfeffer, B. Kohlmeyer and W.F.W. Schneider, Nucl. Phys. A244 (1975) 329
- 1975RA1M Ramaty, Bull. Amer. Phys. Soc. 20 (1975) 680
- 1975RE08 L.P. Remsberg and D.G. Perry, Phys. Rev. Lett. 35 (1975) 361
- 1975RO08 C. Rolfs, W.S. Rodney, M.H. Shapiro and H. Winkler, Nucl. Phys. A241 (1975) 460
- 1975RO1B Roos and Chant, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 242, 265
- 1975RO1R Rolin et al., Bull. Amer. Phys. Soc. 20 (1975) 714
- 1975SA1D Sanchez, Casanova and Casanova, An. Fis. (Spain) 71 (1975) 119
- 1975SC05 H. Schier and B. Schoch, Lett. Nuovo Cim. 12 (1975) 334
- 1975SC1N Schiffer, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 329
- 1975SC20 H. Schweickert, J. Dietrich, R. Neugart and E.W. Otten, Nucl. Phys. A246 (1975) 187
- 1975SC2A Scheurer et al., in Clustering Phenomena in Nucleii, II, ORO-4856-26 (1975) 545
- 1975SC2B Schmid and Do Dang, in JUL-CONF-16 (1975) 12
- 1975SE03 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 151
- 1975SE04 H.P. Seiler, R. Kulesa, P.M. Cockburn, P. Marmier and P.H. Barker, Nucl. Phys. A241 (1975) 159
- 1975SK06 D.M. Skopik, Y.M. Shin and J.J. Murphy, II, Can. J. Phys. 53 (1975) 1398
- 1975SO04 B. Sommer, J.E. Galonska and A. Faessler, Nucl. Phys. A241 (1975) 1
- 1975ST1Q Steinberg et al., in Clustering Phenomena in Nuclei, II, ORO-4856-26 (1975) 315
- 1975SU1E Subotic, Stepanic and Stanojevic, Fizika Suppl. (Yugoslavia) 7 (1975) 60
- 1975SZ1B Szalata et al., Bull. Amer. Phys. Soc. 20 (1975) 567
- 1975TA1A Tang and Thompson, in Clustering Phenom. in Nucl., II, ORO-4856-26 (1975) 119
- 1975TA1C Tauscher, in LASL, AIP Conf. Proc. 26 (1975) 541
- 1975TA1G Tassoul, Astrophys. J. 202 (1975) 755
- 1975TA1H Takahashi, Prog. Theor. Phys. 53 (1975) 461
- 1975TR1A Trimble, Rev. Mod. Phys. 47 (1975) 877
- 1975VO02 C. Volant, M. Conjeaud, S. Harar, S.M. Lee, A. Lepine and E.F. Da Silveira, Nucl. Phys. A238 (1975) 120
- 1975VO09 V.V. Volkov, Fiz. Elem. Chastits At. Yad. (USSR) 6 (1975) 1040; Sov. J. Part. Nucl. 6 (1976) 420
- 1975WI1E Wilkinson, Proc. Int. Symp. on Inter. Studies in Nucl., Germany, 1975 (Amsterdam, North-Holland, 1975) 147

- 1975WI30 M.E. Williams-Norton, G.M. Hudson, K.W. Kemper, G.E. Moore, G.A. Norton, R.J. Puigh and A.F. Zeller, Phys. Rev. C12 (1975) 1899
- 1975WO06 J.G. Woodworth, J.W. Jury, K.H. Lokan and N.K. Sherman, Can. J. Phys. 53 (1975) 795
- 1975WO1D Woosley, Fowler, Holmes and Zimmermna, OAP - 422 (1975)
- 1975ZE1C Zeller et al., Bull. Amer. Phys. Soc. 20 (1975) 718
- 1975ZI1C Zisman, Vandenbosch and Webb, Bull. Amer. Phys. Soc. 20 (1975) 718
- 1976AM04 K. Amos, P. Nesci and R. Smith, Aust. J. Phys. 29 (1976) 233
- 1976AR04 K.P. Artemov, V.Z. Goldberg, I.P. Petrov, V.P. Rudakov, I.N. Serikov and V.A. Timofeev, Yad. Fiz. 23 (1976) 489; Sov. J. Nucl. Phys. 23 (1976) 257
- 1976BA11 H. Bando, Nucl. Phys. A261 (1976) 269
- 1976BA1N Baz, Goldberg, Gridnev and Semenov, in Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) 96
- 1976BA22 Z. Basrak, F. Auger, B. Fernandez, J. Gastebois and N. Cindro, J. Phys. Lett. (Paris) 37 (1976) L131
- 1976BA53 Z. Basrak, F. Auger, B. Fernandez, J. Gastebois and N. Cindro, Phys. Lett. B65 (1976) 119
- 1976BE1G Berlovich, Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 756
- 1976BE1L Benenson, Kashy, Mueller and Nann, in Cargese Conf., CERN 76-13 (1976) 235
- 1976BE1P Bertrand, Ann. Rev. Nucl. Sci. 26 (1976) 457
- 1976BE2E Berg, Datta, Tenhaken and Quin, Bull. Amer. Phys. Soc. 21 (1976) 991
- 1976BE51 H. Beaumevielle, N. Bendjaballah, A. Dauchy, S. Laribi and C. Morand, Nuovo Cim. A33 (1976) 535
- 1976BU22 A. Buta, Rev. Roum. Phys. (Romania) 21 (1976) 869
- 1976CA1E Calprice, Bull. Amer. Phys. Soc. 21 (1976) 507
- 1976CA1P Casanova and Casanova, An. Fis. (Spain) 72 (1976) 174
- 1976CH1P Chatterjee, Tobias and Lyman, in Spallation Nucl. Reactions and Their Applications, Eds., Shen and Merker (D. Reidel Pub. Co., The Netherlands, 1976) 169
- 1976CH1T Chung and Wildenthal, MSUCL-214 (1976)
- 1976CH23 H. Chandra and U. Mosel, Phys. Lett. B64 (1976) 373
- 1976CO15 J.D. Cossairt, R.D. Bent, A.S. Broad, F.D. Becchetti and J. Janecke, Nucl. Phys. A261 (1976) 373
- 1976CO1W Compani-Tabrizi and Malik, Bull. Amer. Phys. Soc. 21 (1976) 682
- 1976CO23 M.E. Cobern, D.J. Pisano and P.D. Parker, Phys. Rev. C14 (1976) 491

- 1976CU07 R.Y. Cusson, R. Hilko and D. Kolb, Nucl. Phys. A270 (1976) 437
- 1976DA13 R.A. Dayras, R.G. Stokstad, Z.E. Switkowski and R.M. Wieland, Nucl. Phys. A263 (1976) 153
- 1976DA1K Darden and Haerberli, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 229
- 1976DA1P Da Silveira, Proc. Int. School of Phys. "Enrico Fermi", Course LXII. Nucl. Spectroscopy and Nucl. Reactions with Heavy Ions, Varenna, 1974 (Amsterdam, North-Holland, 1976) 588; Phys. Abs. 18607 (1977)
- 1976DE12 R. de Swinarski, F.G. Resmini, D.L. Hendrie and A.D. Bacher, Nucl. Phys. A261 (1976) 111
- 1976EB01 K.A. Eberhard and K.G. Bernhardt, Phys. Rev. C13 (1976) 440
- 1976EG02 R. Eggers, M.N. Namboodiri, P. Gonthier, K. Geoffroy and J.B. Natowitz, Phys. Rev. Lett. 37 (1976) 324
- 1976EN1C Eng et al., in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 783
- 1976ER03 K.A. Erb, R.R. Betts, D.L. Hanson, M.W. Sachs, R.L. White, P.P. Tung and D.A. Bromley, Phys. Rev. Lett. 37 (1976) 670
- 1976ES1B Escusie, Fabroco, Pignanelli and Resmini, in Polariz., Zurich, 1975 (Birkhauser Verlag, 1976) 721
- 1976EY01 Y. Eyal, M. Beckerman, R. Chechik, Z. Fraenkel and H. Stocker, Phys. Rev. C13 (1976) 1527
- 1976FI10 L.K. Fifield, R.W. Zurmuhle and D.P. Balamuth, Phys. Rev. C14 (1976) 1010
- 1976FO05 H.T. Fortune, R.R. Betts and R. Middleton, Phys. Lett. B62 (1976) 287
- 1976FO09 H.T. Fortune, T.H. Braid, R.E. Segel and K. Raghunathan, Phys. Lett. B63 (1976) 403
- 1976FO15 H.T. Fortune, S.C. Headley, L.R. Medsker, T.H. Braid, R.E. Segel and K. Raghunathan, Phys. Rev. C14 (1976) 1271
- 1976FO16 H.T. Fortune and J.D. Garrett, Phys. Rev. C14 (1976) 1695
- 1976FU06 G.H. Fuller, J. Phys. Chem. Ref. Data 5 (1976) 835
- 1976FU1K Fukushima, Kamimura and Matsuse, Prog. Theor. Phys. 55 (1976) 1310
- 1976GA1K Gadioli, Gadioli Erba and Iori, Lett. Nuovo Cim. 15 (1976) 83
- 1976GAYV D.I. Garber and R.R. Kinsey, BNL 325, Vol. 2 (1976)
- 1976GE01 J. Gerber, M.B. Goldberg and K.-H. Speidel, Phys. Lett. B60 (1976) 338
- 1976GE08 H. Genz, A. Richter, B.M. Schmitz and H. Behrens, Nucl. Phys. A267 (1976) 13
- 1976GI1C Ginzburg and Ptjskin, Rev. Mod. Phys. 48 (1976) 161
- 1976HA1Q Hanna, in Proc. Dubna Conf., JINR D-9920 Vol. II (1976) 195

- 1976HA1V Harvey and Towner, in Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct. Vol. 1, JINR D-9682 (1976) 17
- 1976HE19 P.-H. Heenen, Nucl. Phys. A272 (1976) 399
- 1976HI05 M.D. High and B. Cujec, Nucl. Phys. A259 (1976) 513
- 1976HI10 G.T. Hickey, D.C. Weisser, J. Cerny, G.M. Crawley, A.F. Zeller, T.R. Ophel and D.F. Hebbard, Phys. Rev. Lett. 37 (1976) 130
- 1976HO1K House and Kemper, Bull. Amer. Phys. Soc. 21 (1976) 1005
- 1976IN05 P.D. Ingalls, Nucl. Phys. A265 (1976) 93
- 1976IN06 P.D. Ingalls, Phys. Rev. C14 (1976) 254
- 1976JA19 D.R. James, G.R. Morgan, N.R. Fletcher and M.B. Greenfield, Nucl. Phys. A274 (1976) 177
- 1976JA1F H.R. Jaqaman and A.Z. Mekjian, Nucl. Phys. A259 (1976) 157
- 1976JA23 J. Janecke, At. Data Nucl. Data Tables 17 (1976) 455
- 1976KA1M Kanofsky, Allen and Lazo, Meson-Nucl. Phys., 1976 (AIP, 1976) 346
- 1976KE03 I. Kelson and Y. Shoshani, Nucl. Phys. A268 (1976) 332
- 1976KI09 C.H. King, M.A.M. Shahabuddin and B.H. Wildenthal, Nucl. Phys. A270 (1976) 399
- 1976KI18 V.S. Kinchakov, Izv. Akad. Nauk SSSR Ser. Fiz. 40 (1976) 150; Bull. Acad. Sci. USSR Phys. Ser. 40 (1976) 126
- 1976KI1D King et al., in Lowell Conf., CONF-760715-P2 (1976) 1344
- 1976KL03 H.V.Klapdor, H.Reiss and G.Rosner, Nucl. Phys. A262 (1976) 157
- 1976KN05 K.T. Knopfle, G.J. Wagner, A. Kiss, M. Rogge, C. Mayer-Boricke and T. Bauer, Phys. Lett. B64 (1976) 263
- 1976KO1L Koshel and Nagel, Bull. Amer. Phys. Soc. 21 (1976) 512
- 1976KO24 J.J. Kolata, R.M. Freeman, F. Haas, B. Heusch and A. Gallmann, Phys. Lett. B65 (1976) 333
- 1976KU1F Kubono et al., Bull. Amer. Phys. Soc. 21 (1976) 578
- 1976LE20 M. LeMere, Y.C. Tang and D.R. Thompson, Phys. Rev. C14 (1976) 23
- 1976LO03 M. Lowry, J.S. Schweitzer, R. Dayras and R.G. Stokstad, Nucl. Phys. A259 (1976) 122
- 1976LO06 K.S. Low, T. Tamura and T. Udagawa, Phys. Rev. C13 (1976) 2579
- 1976MA01 R.E.Marrs, E.G.Adelberger, K.A.Snover and M.D.Cooper, Nucl. Phys. A256 (1976) 1
- 1976MA04 F. Malaguti and P.E. Hodgson, Nucl. Phys. A257 (1976) 37

- 1976MA12 E. Mathiak, K.A. Eberhard, J.G. Cramer, H.H. Rossner, J. Stettmeier and A. Weidinger, Nucl. Phys. A259 (1976) 129
- 1976MA19 R.S. Mackintosh, Z. Phys. A276 (1976) 25
- 1976MA36 R.S. Mackintosh, Nucl. Phys. A266 (1976) 379
- 1976MC12 A.B. McDonald, H.B. Mak, H.C. Evans, G.T. Ewan and H.B. Trautvetter, Nucl. Phys. A273 (1976) 477
- 1976ME14 R. Medoff, L.R. Medsker, S.C. Headley and H.T. Fortune, Phys. Rev. C14 (1976) 1
- 1976MI01 G.F. Millington, R.M. Hutcheon, J.R. Leslie and W. McLatchie, Phys. Rev. C13 (1976) 879
- 1976MO13 G.L. Morgan and J.K. Dickens, Nucl. Sci. Eng. 60 (1976) 36
- 1976NA11 M.N. Namboodiri, E.T. Chulick and J.B. Natowitz, Nucl. Phys. A263 (1976) 491
- 1976NA22 K. Nagatani, C.W. Towsley, K.G. Nair, R. Hanus, M. Hamm and D. Strottman, Phys. Rev. C14 (1976) 2133
- 1976NG04 W.Y. Ng and S.S.M. Wong, Can. J. Phys. 54 (1976) 2367
- 1976NO10 H. Noto, Y. Abe, J. Hiura and H. Tanaka, Prog. Theor. Phys. 55 (1976) 1432
- 1976NO1C Norgaard and Fricke, Astron. Astrophys. 49 (1976) 337
- 1976OG1A Ogloblin, in Proc. Dubna Conf. JINR D-9920, Vol. II (1976) 31
- 1976PA02 K.H. Passler and U. Mosel, Nucl. Phys. A257 (1976) 242
- 1976PA1J Partridge, Tang, Thompson and Brown, Private Communication (1976)
- 1976PI09 D.J. Pisano, Phys. Rev. C14 (1976) 468
- 1976PI10 D.J. Pisano and P.D. Parker, Phys. Rev. C14 (1976) 475
- 1976PI16 A.A. Pilt, D.J. Millener, H. Bradlow, O. Dietzsch, P.S. Fisher, W.J. Naude, W.D.M. Rae and D. Sinclair, Nucl. Phys. A273 (1976) 189
- 1976PO01 V. Potbhare and S.P. Pandya, Nucl. Phys. A256 (1976) 253
- 1976PR08 G.A. Prokopets, B. Holmqvist and A.V. Murzin, Yad. Fiz. 23 (1976) 935; Sov. J. Nucl. Phys. 23 (1976) 492
- 1976RO14 C. Rogers, Nucl. Phys. A271 (1976) 447
- 1976RO1J Ross and Aller, Science 191 (1976) 1223
- 1976RU1C Ruge and Walker, Astrophys. J. 203 (1976) L139
- 1976SA1F Sasakawa, Proc. 7th Int. Conf. on Few Body Problems in Nucl. Particle Phys., Delhi, 1975 (Amsterdam, North-Holland, 1976) 577
- 1976SA21 L. Satpathy, K.W. Schmid, S. Krewald and A. Faessler, Phys. Rev. C14 (1976) 1995
- 1976SC08 K.W. Schmid and G. Do Dang, Z. Phys. A276 (1976) 233
- 1976SE10 H.M. Sen Gupta, T.A. Belote and D. Roaf, Nucl. Phys. A267 (1976) 205

- 1976SH02 J.D. Sherman, D.L. Hendrie and M.S. Zisman, Phys. Rev. C13 (1976) 20
- 1976SH07 S.K. Sharma, L. Satpathy, S.B. Khadkikar and S.C.K. Nair, Phys. Lett. B61 (1976) 122
- 1976SI1D Silberberg, Tsao and Shapiro, in Spallation Nucl. Reactions & Their Appl., Eds., Shen and Mercer (D. Reidel, Netherlands, 1976) p. 49
- 1976SL2A I. Slaus, in Lowell Conf., CONF-760715-P1 (1976) 272
- 1976SM1D Smirnov and Filippov, in Contrib. to Dubna Conf. on Selected Topics in Nucl. Struct., Vol. 1, JINR D-9682 (1976) 8
- 1976ST07 R.G. Stokstad, J. Gomez del Campo, J.A. Biggerstaff, A.H. Snell and P.H. Stelson, Phys. Rev. Lett. 36 (1976) 1529
- 1976ST09 A.P. Stamp, Nucl. Phys. A266 (1976) 119
- 1976ST11 W.F. Steele, P.A. Smith, J.E. Finck and G.M. Crawley, Nucl. Phys. A266 (1976) 424
- 1976ST18 R. Stock, U. Jahnke, D.L. Hendrie, J. Mahoney, C.F. Maguire, W.F.W. Schneider, D.K. Scott and G. Wolschin, Phys. Rev. C14 (1976) 1824
- 1976SW02 Z.E. Switkowski, R.G. Stokstad and R.M. Wieland, Nucl. Phys. A274 (1976) 202
- 1976SW1C Swiniarski et al., Helv. Phys. Acta 49 (1976) 241
- 1976TR03 R.E. Tribble, J.D. Cossairt and R.A. Kenefick, Phys. Lett. B61 (1976) 353
- 1976UL1D Ullah, J. Phys. G2 (1976) 33
- 1976VA12 R. Vandenbosch and K.G. Bernhardt, J. Phys. Lett. (Paris) 37 (1976) L-161
- 1976VA29 G.V. Valskii, Yad. Fiz. 24 (1976) 270; Sov. J. Nucl. Phys. 24 (1976) 140
- 1976VI05 D.A. Viggars, T.W. Conlon, I. Naqib and A.T. McIntyre, J. Phys. G2 (1976) L55
- 1976VO1C Voinova, INDC(CCP)-93/N (1976)
- 1976VO1D von Charzewski and Dreizler, Z. Phys. A278 (1976) 35
- 1976WA18 A.H. Wapstra and K. Bos, At. Data Nucl. Data Tables 17 (1976) 474
- 1976WE15 A. Weidinger, F. Busch, G. Gaul, W. Trautmann and W. Zipper, Nucl. Phys. A263 (1976) 511
- 1976YO02 D.H. Youngblood, J.M. Moss, C.M. Rozsa, J.D. Bronson, A.D. Bacher and D.R. Brown, Phys. Rev. C13 (1976) 994
- 1976ZA1E Zaichenko and Ol'khovzkii, Ukr. Fiz. Zh. 21 (1976) 1107; Phys. Abs. 75110 (1976)
- 1977AB03 A.Y. Abul-Magd, H.M. Khalil and M.M. Shalaby, J. Phys. G3 (1977) 381
- 1977AB08 Y. Abgrall, B. Morand, E. Caurier and B. Grammaticos, Phys. Rev. Lett. 39 (1977) 922
- 1977AJ02 F. Ajzenberg-Selove, Nucl. Phys. A281 (1977) 1
- 1977AN12 G.S. Anagnostatos, Atomkernenergie 29 (1977) 207

1977AN1B Ando, Ikeda and Tohsaki, in Tokyo (1977), Contrib. Papers p. 587

1977AN21 I. Angeli and M. Csatos, Nucl. Phys. A288 (1977) 480

1977AR06 A.G. Artukh, G.F. Gridnev, V.L. Mikheev and V.V. Volkov, Nucl. Phys. A283 (1977) 350

1977AR1H Arnett, Astrophys. J. (Suppl. Series) 35 (1977) 145

1977AZ02 G. Azuelos and J.E. Kitching, Nucl. Phys. A285 (1977) 19

1977BA12 A.I. Baz, V.Z. Goldberg, N.Z. Darwisch, K.A. Gridnev, V.M. Semjonov and E.F. Hefter, Lett. Nuovo Cim. 18 (1977) 227

1977BA2H Bacher et al., in Tokyo (1977), Contrib. Papers p. 812

1977BA3W Balamuth et al., Bull. Amer. Phys. Soc. 22 (1977) 527

1977BA3X Balamuth et al., in Tokyo (1977), Contrib. Papers p. 194

1977BA50 D.P. Balamuth and E.G. Adelberger, Phys. Rev. C16 (1977) 928

1977BE2R Bennett, Fulbright, Gove and Draayer, Bull. Amer. Phys. Soc. 22 (1977) 995

1977BH1B Bhatia et al., in Tokyo (1977), Contrib. Papers p. 181

1977BR24 F.P. Brady, D.A. Viggars, T.W. Conlon and D.J. Parker, Phys. Rev. Lett. 39 (1977) 870

1977BU1Q Butler, Newman and Talbot, OAP-492 (1977)

1977CA08 E. Caurier and B. Grammaticos, Nucl. Phys. A279 (1977) 333

1977CA11 F.P. Calaprice, W. Chung and B.H. Wildenthal, Phys. Rev. C15 (1977) 2178

1977CA1N Casse and Goret, OAP-499 (1977)

1977CH02 N.S. Chant and P.G. Roos, Phys. Rev. C15 (1977) 57

1977CH10 P.R. Christensen, Z.E. Switkowski and R.A. Dayras, Nucl. Phys. A280 (1977) 189

1977CL1F Clayton, Dwek and Woosley, Astrophys. J. 214 (1977) 300

1977CO14 T.M. Cormier, P. Braun-Munzinger, P.M. Cormier, J.W. Harris and L.L. Lee, Jr., Phys. Rev. C16 (1977) 215

1977CO1G Colombo et al., in Tokyo (1977), Contrib. Papers p. 490

1977CO1Q Conjeaud et al., in Tokyo (1977), Contrib. Papers p. 663

1977CO2C Cobern et al., Bull. Amer. Phys. Soc. 22 (1977) 40

1977DA07 R. Dayras, Z.E. Switkowski and S.E. Woosley, Nucl. Phys. A279 (1977) 70

1977DA10 B.J. Dalton, J.P. Vary and W.J. Baldrige, Phys. Rev. Lett. 38 (1977) 1348

1977DA1B Davydov, Naumov and Shcherbachenko, INDC(CCP)-104/LN (1977)

1977DA1J Da Silveira et al., in Tokyo (1977), Contrib. Papers p. 624

1977DE2A Dennis, Thcrnton and Cordell, in Tokyo (1977), Contrib. Papers p. 644

- 1977DI1A Dickens et al., Nucl. Instrum. Meth. Phys. Res. 62 (1977) 515
- 1977DW1B Dwyer and Meyer, Astrophys. J. 216 (1977) 646
- 1977EL1E Elmore, Sugarbaker, Boyd and Gove, Bull. Amer. Phys. Soc. 22 (1977) 1010
- 1977EN01 J.B.A. England, E. Casal, A. Garcia, T. Picazo, J. Aguilar and H.M. Sen Gupta, Nucl. Phys. A284 (1977) 29
- 1977ER1E Erb et al., Bull. Amer. Phys. Soc. 22 (1977) 38
- 1977ER1F Erb et al., in Tokyo (1977), Contrib. Papers p. 638, 640
- 1977EV01 D. Evers, C. Ley, E. Spindler, W. Assmann, K. Rudolph, P. Konrad and P. Sperr, Nucl. Phys. A275 (1977) 363
- 1977FE07 H. Feldmeier and P. Manakos, Z. Phys. A281 (1977) 379
- 1977FI08 L.K. Fifield, F.P. Calaprice, C.H. Zimmerman, M.J. Hurst, A. Pakkanen, T.J.M. Symons, F. Watt and K.W. Allen, Nucl. Phys. A288 (1977) 57
- 1977FL13 T. Fliessbach and P. Manakos, J. Phys. G3 (1977) 643
- 1977FL1E T. Fliessbach, Nucl. Phys. A285 (1977) 262
- 1977FL1F Fletcher and James, Bull. Amer. Phys. Soc. 22 (1977) 1010
- 1977FO01 H.T. Fortune, L.R. Greenwood, R.E. Segel and J.R. Erskine, Phys. Rev. C15 (1977) 439
- 1977FR08 S.J. Freedman, R.D. Cousins, Jr., C.A. Gagliardi, G.T. Garvey and J.F. Greenhalgh, Phys. Lett. B67 (1977) 165
- 1977FU1P Fujiwara, Horiuchi and Tamagaki, in Tokyo (1977), Contrib. Papers p. 196
- 1977GA04 W. Galster, W. Treu, P. Duck, H. Frohlich and H. Voit, Phys. Rev. C15 (1977) 950
- 1977GA05 W. Galster, W. Treu, P. Duck, H. Frohlich and H. Voit, Phys. Lett. B67 (1977) 262
- 1977GU01 G.D. Gunn, R.N. Boyd, N. Anantaraman, D. Shapira, J. Toke and H.E. Gove, Nucl. Phys. A275 (1977) 524
- 1977HA33 C.S. Han, D.S. Chuu, M.C. Wang and S.T. Hsieh, Phys. Rev. C16 (1977) 1645
- 1977HE10 K.T. Hecht, Nucl. Phys. A283 (1977) 223
- 1977HE12 J.A.J. Hermans, G.A.P. Engelbertink, L.P. Ekstrom, H.H. Eggenhuisen and M.A. Van Driel, Nucl. Phys. A284 (1977) 307
- 1977HE1D Hermans, Thesis Univ. Utrecht (1977)
- 1977HI01 M.D. High and B. Cujec, Nucl. Phys. A278 (1977) 149
- 1977HI04 M.D. High and B. Cujec, Nucl. Phys. A282 (1977) 181
- 1977HO1E Horiuchi, Invited Talk at Int. Symp. Nucl. Collisions & Their Microsc. Descr., Bled, Yugoslavia (1977)

- 1977HO1F Horiuchi, Proc. Int. Conf. Nucl. Struct., Tokyo, (1977); J. Phys. Soc. Jpn. Suppl. 44 (1978) 85
- 1977HU1C Hurd et al., Bull. Amer. Phys. Soc. 22 (1977) 1010
- 1977IK1C Ikeda, in Tokyo (1977), Contrib. Papers p. 218
- 1977KA2F KAto and Banco, in Tokyo (1977), Contrib. Papers p. 195
- 1977KE02 K.-U. Kettner, H. Lorenz-Wirzba, C. Rolfs and H. Winkler, Phys. Rev. Lett. 38 (1977) 337
- 1977KI1M Kido et al., Bull. Amer. Phys. Soc. 22 (1977) 567
- 1977KL1G Klapdor, Rosner and Willmes, Proc. Int. Workshop V on Gross Properties of Nuclei and Nucl. Excitations, Hirschegg (1977)
- 1977KN02 W. Knupfer, K. Knauss and M.G. Huber, Phys. Lett. B66 (1977) 305
- 1977KO16 J.J. Kolata, R.C. Fuller, R.M. Freeman, F. Haas, B. Heusch and A. Gallmann, Phys. Rev. C16 (1977) 891
- 1977LE1N Lecuyer et al., Nucl. Instrum. Meth. 140 (1977) 305
- 1977LE1T Lebrun et al., in Tokyo (1977), Contrib. Papers p. 521
- 1977LI19 H. Liskien and A. Paulsen, Atomkernenergie 30 (1977) 59
- 1977MA07 R.E.Marrs, E.G.Adelberger and K.A.Snover, Nucl. Phys. A277 (1977) 429
- 1977MA09 R.S. Mackintosh, Nucl. Phys. A280 (1977) 86
- 1977MA1M Mausner et al., Bull. Amer. Phys. Soc. 22 (1977) 547
- 1977MA2G Martz et al., Bull. Amer. Phys. Soc. 22 (1977) 634
- 1977MA2V Marquardt, Hoppe, Buttlar and Sprengel, Contrib. Int. Conf. on Resonances in Heavy Ion Reactions, Hvar, Yugoslavia (1977)
- 1977MA2W Marquardt, Hoppe, Buttlar and Sprengel, in Tokyo (1977), Contrib. Papers p. 632
- 1977MA2X Matsuse, in Tokyo (1977), Contrib. Papers p. 637
- 1977MC1J McKeown, Calaprice and Alburger, Bull. Amer. Phys. Soc. 22 (1977) 28
- 1977MC1K P.J. McNulty, G.E. Farrell, R.C. Filz, W. Schimmerling and K.G. Vosburgh, Phys. Rev. Lett. 38 (1977) 1519
- 1977MO1H Motobayashi, Kohno, Odi and Nakajima, in Tokyo (1977), Contrib. Papers p. 628
- 1977MO1X Motobayashi et al., in Tokyo (1977), Contrib. Papers p. 662
- 1977MP01 P. Mpanias and M.L. Rustgi, Phys. Rev. C16 (1977) 816
- 1977NA03 J.B. Natowitz, M.N. Namboodiri, R. Eggers, P. Gonthier, K. Geoffroy, R. Hanus, C. Towsley and K. Das, Nucl. Phys. A277 (1977) 477; Erratum Nucl. Phys. A285 (1977) 532
- 1977NE1D Nesci and Amos, in Tokyo (1977), Contrib. Papers p. 491

- 1977OH01 S. Ohkubo, Y. Kondo and S. Nagata, *Prog. Theor. Phys.* 57 (1977) 82
- 1977OK1E Oka and Kubodera, in *Tokyo (1977)*, *Contrib. Papers* p. 792
- 1977OS02 F. Osterfeld, V. Hnizdo and C. Toepffer, *Phys. Lett.* B68 (1977) 319
- 1977PA07 K.H. Passler and U. Mosel, *Phys. Lett.* B66 (1977) 323
- 1977PA19 A.D. Panagiotou, *Nuovo Cim.* A37 (1977) 370
- 1977PA1J Parkinson, *Astron. Astrophys.* 57 (1977) 185
- 1977PL1C Plavko, *Izv. Akad. Nauk SSSR Ser. Fiz.* 41 (1977) 1300
- 1977PO1E Pougheon et al., in *Tokyo (1977)*, *Contrib. Papers* p. 627
- 1977PR1F Proudfoot et al., *Int. Conf. on Resonances in Heavy Ion Reactions*, Hvar, Yugoslavia (1977)
- 1977RA1D Rasmussen, Oliveira, Donangelo and Idannou, in *Tokyo (1977)*, *Contrib. Papers* p. 771
- 1977RO13 N. Rowley, *Phys. Lett.* B69 (1977) 25
- 1977RO1V G. Rosensteel and D.J. Rowe, *Phys. Rev. Lett.* 38 (1977) 10
- 1977RO1W Robertson, Bowles and Freedman, MSUCL-241 (1977)
- 1977RO1Y Rolin et al., in *Tokyo (1977)*, *Contrib. Papers* p. 781
- 1977RO20 N. Rolin, J.P. Deutsch, D. Favart, M. Lebrun and R. Prieels, *Phys. Lett.* B70 (1977) 23
- 1977SA1X Sanders, Martz and Parker, in *Tokyo (1977)*, *Contrib. Papers* p. 625
- 1977SA2A Sanders, Ph. D. Dissertation, Yale Univ. (1977)
- 1977SA2F Sanders, Martz and Parker, *Bull. Amer. Phys. Soc.* 22 (1977) 552
- 1977SA2G Sanders, Martz and Parker, in *Tokyo (1977)*, *Contrib. Papers* p. 626
- 1977SC02 K.W. Schmid and G. Do Dang, *Phys. Lett.* B66 (1977) 5
- 1977SC08 K.W. Schmid and G. Do Dang, *Phys. Rev.* C15 (1977) 1515
- 1977SC36 D. Schwalm, E.K. Warburton and J.W. Olness, *Nucl. Phys.* A293 (1977) 425
- 1977SH11 J. Shurpin, D. Strottman, T.T.S. Kuo, M. Conze and P. Manakos, *Phys. Lett.* B69 (1977) 395
- 1977SH13 R. Sherr, *Phys. Rev.* C16 (1977) 1159
- 1977ST21 A. Strazzeri, *Lett. Nuovo Cim.* 19 (1977) 597
- 1977SU04 T. Suzuki, A. Arima and K.-I. Kubo, *Nucl. Phys.* A288 (1977) 493
- 1977SU1M Sugarbaker et al., *Bull. Amer. Phys. Soc.* 22 (1977) 1003
- 1977SW01 Z.E. Switkowski, H. Winkler and P.R. Christensen, *Phys. Rev.* C15 (1977) 449

- 1977SW05 Z.E. Switkowski, Shiu-Chin Wu, J.C. Overley and C.A. Barnes, Nucl. Phys. A289 (1977) 236
- 1977TA07 S.L. Tabor, Y. Eisen, D.G. Kovar and Z. Vager, Phys. Rev. C16 (1977) 673
- 1977TA1D Takimoto et al., in Tokyo (1977), Contrib. Papers p. 567
- 1977TH03 W.J. Thompson and J.S. Eck, Phys. Lett. B67 (1977) 151
- 1977TO1L Tonozuka, in Tokyo (1977), Contrib. Papers p. 197
- 1977TR1H Tribble and May, in Tokyo (1977), Contrib. Papers p. 780
- 1977VA1G van Bibber et al., Bull. Amer. Phys. Soc. 22 (1977) 39
- 1977VO05 H. Voit, W. Galster, W. Treu, H. Frohlich and P. Duck, Phys. Lett. B67 (1977) 399
- 1977VO1A von Rosenvinge, McDonald, Trainor and Webber, Bull. Amer. Phys. Soc. 22 (1977) 567
- 1977WA08 A.H. Wapstra and K. Bos, At. Data Nucl. Data Tables 19 (1977) 175; Erratum At. Data Nucl. Data Tables 20 (1977) 1
- 1977WA1M Wang, Chant, Cowley and Roos, in Tokyo (1977), Contrib. Papers p. 568
- 1977WA1V Watt et al., in Tokyo (1977), Contrib. Papers p. 525
- 1977WA1W Walker, Bull. Amer. Phys. Soc. 22 (1977) 1253
- 1977WI02 D.H. Wilkinson, Phys. Lett. B66 (1977) 105
- 1977YO1H Young, Lind, Balamuth and Zurmuhle, Bull. Amer. Phys. Soc. 22 (1977) 527
- 1978EN06 P.M. Endt and C. van der Leun, Nucl. Phys. A310 (1978) 96
- 1978GO1F J. Golden, R.A. Mahaffey, J.A. Pasour, F.C. Young and C.A. Kapetanacos, Rev. Sci. Instrum. 49 (1978) 1384
- 1978MA07 N. Marquardt, D. Sprengel, H. Buttler and W. Hoppe, Z. Phys. A285 (1978) 89
- 1978ST08 D.J. Steck, Phys. Rev. C17 (1978) 1034
- 1979AJ01 F. Ajzenberg-Selove, Nucl. Phys. A320 (1979) 1