

Energy Levels of Light Nuclei $A = 20$

F. Ajzenberg-Selove

University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396

Abstract: An evaluation of $A = 18-20$ was published in *Nuclear Physics A475* (1987), p. 1. This version of $A = 20$ differs from the published version in that we have corrected some errors discovered after the article went to press. Figures and introductory tables have been omitted from this manuscript. [Reference](#) key numbers have been changed to the NNDC/TUNL format.

(References closed June 1, 1987)

The original work of Fay Ajzenberg-Selove was supported by the US Department of Energy [DE-FG02-86ER40279]. Later modification by the TUNL Data Evaluation group was supported by the US Department of Energy, Office of High Energy and Nuclear Physics, under: Contract No. DEFG05-88-ER40441 (North Carolina State University); Contract No. DEFG05-91-ER40619 (Duke University).

Table of Contents for $A = 20$

Below is a list of links for items found within the PDF document. Figures from this evaluation have been scanned in and are available on this website or via the link below. The introductory [Table 2](#) is also available via the link.

A. Nuclides: [\$^{20}\text{n}\$](#) , [\$^{20}\text{He}\$](#) , [\$^{20}\text{Li}\$](#) , [\$^{20}\text{Be}\$](#) , [\$^{20}\text{B}\$](#) , [\$^{20}\text{C}\$](#) , [\$^{20}\text{N}\$](#) , [\$^{20}\text{O}\$](#) , [\$^{20}\text{F}\$](#) , [\$^{20}\text{Ne}\$](#) , [\$^{20}\text{Na}\$](#) , [\$^{20}\text{Mg}\$](#) , [\$^{20}\text{Al}\$](#) , etc.

B. Tables of Recommended Level Energies:

[Table 20.1](#): Energy levels of ^{20}O

[Table 20.2](#): Energy levels of ^{20}F

[Table 20.13](#): Energy levels of ^{20}Ne

[Table 20.27](#): Energy levels of ^{20}Na

C. [References](#)

D. Figures: [\$^{20}\text{O}\$](#) , [\$^{20}\text{F}\$](#) , [\$^{20}\text{Ne}\$](#) , [\$^{20}\text{Na}\$](#) , [Isobar diagram](#)

E. Erratum to the Publication: [PS](#) or [PDF](#)

^{20}n , ^{20}He , ^{20}Li , ^{20}Be
(Not observed)

See (1983ANZQ, 1983BE55; theor.).

^{20}B
(Not observed)

The mass excess of ^{20}B is predicted to be 69.08 MeV. ^{20}B is then unstable with respect to breakup into $^{19}\text{B} + \text{n}$ by 0.9 MeV: see ^{19}B and (1978AJ03). See also (1983ANZQ; theor.).

^{20}C
(Not illustrated)

^{20}C has been observed in the fragmentation of 60 MeV/A argon ions: its mass excess is 37.20 ± 1.13 MeV (1987GI1E). It is then stable with respect to $^{19}\text{C} + \text{n}$ and $^{18}\text{C} + 2\text{n}$ by 3.3 and 3.9 MeV, respectively. See also (1978AJ03, 1983AJ01). The half-life of ^{20}C is calculated to be 9.3×10^{-3} sec (1984KL06). See also (1985AN1B, 1985LA03, 1986AN07, 1986GU1D) and (1982AV1A, 1983ANZQ, 1987SA15; theor.).

^{20}N
(Not illustrated)

^{20}N is particle stable. Its atomic mass excess is 21.64 ± 0.26 MeV (1986VI09), 22.20 ± 0.36 MeV (1986GI10), 21.62 ± 0.14 MeV (1987GI1E). We adopt 21.62 ± 0.14 MeV. ^{20}N is then stable with respect to $^{19}\text{N} + \text{n}$ by 2.32 MeV (see ^{19}N). The half-life of ^{20}N is 100_{-20}^{+30} msec, $P_n \approx 61\%$ (1987MU1J; prelim.). See also (1984KL06; theor.). See also (1985PIZZ, 1986PI09), (1983WI1A, 1984HI1A, 1986AN07, 1986GU1D) and (1983ANZQ; theor.).

^{20}O
(Figs. 10 and 13)

GENERAL: (See also (1983AJ01).)

Model calculations: (1978WI1B, 1982SH30, 1984CH1V, 1984HA14, 1984RA13, 1984SA37, 1985HA15, 1985HU08, 1985LE1L, 1986COZZ, 1986HE13, 1986HU1G, 1986VO07, 1986WA1R, 1987IA1B).

Complex reactions involving ^{20}O : (1983FR1A, 1983WI1A, 1984HI1A, 1985HA1N, 1985PO11, 1986HA1B, 1986IR01, 1986PO06, 1986PO15, 1987RI03).

Other topics: (1978WI1B, 1983SH32, 1984PO11, 1984SA37, 1985AN28, 1985MU10, 1986AN07).

Ground state of ^{20}O : (1978WI1B, 1983ANZQ, 1984FR13, 1987SA15).

Mass of ^{20}O : From the Q -value of the $^{18}\text{O}(t, p)^{20}\text{O}$ reaction the atomic mass excess of ^{20}O is stated to be 3796.3 ± 1 keV (1985AN17). See also (1982AN12, 1985WA02, 1986GU1D).

For $B(E2)$ of $^{20}\text{O}^*(1.67)$ and other parameters see (1987RA01) and Table 2 of the Introduction in this publication.

1. $^{20}\text{O}(\beta^-)^{20}\text{F}$ $Q_m = 3.814$

^{20}O decays with a half-life of 13.51 ± 0.05 sec to the 1^+ states $^{20}\text{F}^*(1.06, 3.49)$ with branching ratios (99.973 ± 0.003) and $(0.027 \pm 0.003)\%$, $\log f_0 t = 3.740 \pm 0.006$ and 3.65 ± 0.06 , respectively (1987AL06). Upper limits for branching to other states of ^{20}F are shown in Table II of (1987AL06). See also (1985BR29).

2. $^{18}\text{O}(t, p)^{20}\text{O}$ $Q_m = 3.082$

$Q_0 = 3082.4 \pm 1.3$ keV (1985AN17). See also (1982AN12).

Observed proton groups are displayed in Tables 20.2 of (1983AJ01) and 20.1 here. $^{20}\text{O}^*(4.07)$ decays to $^{20}\text{O}^*(0, 1.67)$ with branchings of (26 ± 4) and $(74 \pm 4)\%$. The p - γ angular correlations lead to $J = 2$; the strength of the transition favors $\pi = +[\delta(E2/M1) = -0.18 \pm 0.08$ for the $2^+ \rightarrow 2^+$ transition]. $^{20}\text{O}^*(4.46)$ and $^{20}\text{O}^*(5.39)$ decay primarily via $^{20}\text{O}^*(1.67)$; the direct ground-state decay is $< 4\%$ for the first and $< 7\%$ for the second of these states. The angular correlations are essentially isotropic, favoring $J^\pi = 0^+$. The transition $^{20}\text{O}^*(5.39 \rightarrow 4.07)$ is not observed: the upper limit is 8% . See also (1978AJ03, 1983AJ01). For a discussion of $A = 20$ isobaric states see (1982AN12, 1985AN17).

3. $^{18}\text{O}(\alpha, 2p)^{20}\text{O}$ $Q_m = -16.732$

See (1983AJ01).

4. $^{18}\text{O}(^{18}\text{O}, ^{16}\text{O})^{20}\text{O}$ $Q_m = -0.624$

See (1983AJ01).

Table 20.1: Energy levels of ^{20}O

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
0	$0^+; 2$	$\tau_{1/2} = 13.57 \pm 0.1$ sec	β^-	1, 2, 3, 4
1.67368 ± 0.15	2^+	$\tau_m = 10.5 \pm 0.4$ psec $g = -0.352 \pm 0.015$	γ	2, 3, 4
3.570 ± 7	4^+		(γ)	2, 3, 4
4.072 ± 4	2^+		γ	2, 4
4.456 ± 5	0^+		γ	2, 4
4.850 ± 15	4^+		(γ)	2
5.002 ± 6			(γ)	2
5.234 ± 5	2^+		(γ)	2
5.304 ± 6	2^+		(γ)	2
5.387 ± 6	0^+		γ	2
5.614 ± 3	(3^-)		(γ)	2
6.555 ± 8	(2)		(γ)	2
7.252 ± 8	5^-		(γ)	2
7.622 ± 7	$3^- + 4^+$			2
7.754 ± 5	4^+			2, 3
7.855 ± 6	(5^-)			2, 3
8.554 ± 8	4^+			2
8.804 ± 9	3^-			2, 3
8.962 ± 21	(0^+)			2
9.770 ± 8	0^+			2
10.125 ± 11	2^+			2, 3

²⁰F
(Figs. 11 and 13)

GENERAL (See also (1983AJ01).)

Model calculations: (1978WI1B, 1982HA43, 1983BR29, 1984FO16, 1984RA13, 1986CA27, 1986COZZ, 1986VO05, 1986WA1R, 1987HA08, 1987IA1B).

Complex reactions involving ²⁰F: (1983BE02, 1983DE26, 1983WI1A, 1984GR08, 1984HO23, 1984KO25, 1985BE40, 1985HA1N, 1985PO11, 1986GA1P, 1986HA1B, 1986ME06, 1986PO06, 1987RI03, 1987RO10).

Hypernuclei: (1984AS1D).

Other topics: (1978WI1B, 1983AR1J, 1983BR29, 1985AN28, 1986AN07, 1986VO05, 1987HA08).

Ground state of ²⁰F: (1978WI1B, 1983ANZQ, 1983AR1J, 1984KO25, 1986CA27).

$$\mu = +2.094 (2) \text{ nm (1978LEZA)}$$

$$Q = 0.070 (13) \text{ b (1978LEZA)}$$

1. $^{20}\text{F}(\beta^-)^{20}\text{Ne}$ $Q_m = 7.0289$

The half-life of ²⁰F is 11.00 ± 0.02 sec: see (1978AJ03). ²⁰F decays principally to ²⁰Ne*(1.63): see ²⁰Ne, reaction 33. See also (1984KO25, 1985HE08).

2. $^{12}\text{C}(^9\text{Be}, \text{p})^{20}\text{F}$ $Q_m = 4.0760$

For excitation curves involving ²⁰F*(0, 1.82 + 1.84 + 1.97 + 2.04 + 2.19) see (1982HU06, 1983JA09). At $E(^9\text{Be}) = 12$ to 27 MeV angular distributions are reported for p_0 and $p_{1+2+3+4}$: see (1983AJ01).

3. $^{13}\text{C}(^7\text{Li}, ^7\text{Li})^{13}\text{C}$ $E_b = 18.0492$

For fusion cross sections see (1982DE30). See also ¹³C in (1986AJ01) and (1983AJ01).

4. $^{13}\text{C}(^9\text{Be}, \text{d})^{20}\text{F}$ $Q_m = 1.3542$

See (1983AJ01).

Table 20.2: Energy levels of ^{20}F ^a

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
0	$2^+; 1$	$\tau_{1/2} = 11.00 \pm 0.02$ sec	β^-	1, 2, 4, 6, 7, 9, 10, 11, 14, 16, 17, 19, 20
0.65600 ± 0.04	3^+	$\tau_m = 0.39 \pm 0.03$ psec	γ	6, 7, 9, 10, 11, 14, 19
0.82268 ± 0.08	4^+	79 ± 6 psec	γ	5, 6, 7, 9, 10, 11, 14, 17, 19
0.98371 ± 0.05	1^-	2.0 ± 0.2 psec	γ	6, 7, 9, 11, 14, 17, 19
1.056818 ± 0.004	1^+	45 ± 13 fsec	γ	6, 7, 9, 10, 11, 14, 15, 16, 17, 19
1.30934 ± 0.05	2^-	1.6 ± 0.3 psec	γ	6, 7, 9, 11, 14, 16, 17, 19
1.8244 ± 1.2	5^+	≤ 65 fsec	(γ)	2, 6, 9, 10, 14, 19
1.84397 ± 0.08	2^-	30 ± 20 fsec	γ	2, 7, 9, 11, 14, 17
1.97080 ± 0.07	(3^-)		γ	2, 5, 6, 7, 9, 11, 14, 19
2.04405 ± 0.06	2^+	37 ± 16 fsec	γ	2, 6, 7, 9, 11, 14, 17, 19
2.19436 ± 0.08	(3^+)	< 12 fsec	γ	2, 6, 7, 9, 10, 11, 14, 17, 19
2.8649 ± 1.5	(3^-)		(γ)	6, 7, 9, 14, 19
2.96616 ± 0.08	3^+	60 ± 40 fsec	γ	6, 7, 9, 11, 14, 19
2.968 ± 1.5	(4^-)		(γ)	5, 6, 7, 19
3.17258 ± 0.42	(1^+)		γ	6, 7, 9, 11, 14, 19
3.48849 ± 0.06	1^+	44 ± 11 fsec	γ	6, 7, 9, 11, 14, 15, 19
3.52628 ± 0.07	0^+	30 ± 15 fsec	γ	9, 11, 14
3.58656 ± 0.09	($1, 2$) ⁺	≤ 60 fsec	γ	6, 7, 9, 11, 14, 19
3.68013 ± 0.06	1, 2		γ	6, 7, 9, 11, 14, 19
3.7611 ± 1.9	($2^-, 3^+$)		(γ)	6, 7, 9, 14, 19
3.96519 ± 0.16	1^+		γ	6, 7, 9, 11, 14, 19
4.08208 ± 0.11	(1) ⁺		γ	6, 7, 9, 11, 14, 19

Table 20.2: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
4.1989 \pm 2.7			(γ)	6, 14
4.2077 \pm 2.6			(γ)	7, 14, 19
4.27722 \pm 0.14	(1, 2) ⁺		γ	6, 7, 11, 14, 19
4.3154 \pm 2.0	(0, 1) ⁺		(γ)	14
4.37138 \pm 0.12	(2 ⁺)		γ	6, 7, 11, 14, 19
4.5087 \pm 0.4	1 ⁺ (2)		γ	6, 7, 11, 14, 19
4.5808 \pm 1.8			(γ)	6, 7, 14
4.5922 \pm 2.9			(γ)	14, 19
4.7310 \pm 2.0	(3 ⁻ , 4 ⁻ , 4 ⁺ , 5 ⁺)		(γ)	6, 7, 14, 19
4.7656 \pm 2.0			(γ)	6, 7, 14, 19
4.8916 \pm 2.8			(γ)	6, 14, 19
4.8982 \pm 2.8			(γ)	7, 14
5.047 \pm 4	(2) ⁻		(γ)	6, 14, 19
5.068 \pm 3	(1 ⁻ , 2, 3 ⁺)		(γ)	6, 14
5.1310 \pm 2.5	(2 ⁻ , 3, 4 ⁺)		(γ)	6, 14, 19
5.2239 \pm 2.3	(1, 2) ⁻		(γ)	6, 7, 14, 19
5.2819 \pm 2.5			(γ)	6, 14, 19
5.31887 \pm 0.17	0, 1, 2		γ	6, 11, 14, 19
5.349 \pm 0.4	(3) ⁺		(γ)	6, 14
5.4131 \pm 0.6			γ	6, 7, 14, 19
5.4503 \pm 3.8			(γ)	14, 19
5.4554 \pm 3.2			(γ)	14
5.463 \pm 3	(1, 2, 3) ⁺		(γ)	14
5.55534 \pm 0.13	1, 2 ⁺		γ	7, 11, 14, 19
5.5881 \pm 1.5			(γ)	14
5.620 \pm 3			(γ)	7, 14, 19
5.713 \pm 2			γ	6, 14, 19
5.7640 \pm 2.5	(3) ⁺		(γ)	6, 14, 19
5.8104 \pm 2.5	(1 ⁺)		(γ)	6, 14, 19
5.93609 \pm 0.05	2 ⁻		γ	11, 14, 19
6.01777 \pm 0.03	2 ⁻		γ	11, 14

Table 20.2: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
6.04498 \pm 0.08	0, 1, 2		γ	11, 14, 19
6.090 \pm 7	(0 ⁻)		(γ)	6
6.161 \pm 4	(2, 3 ⁺)		(γ)	6, 19
6.200 \pm 4	(2 ⁻ , 3, 4 ⁺)		(γ)	6, 19
6.240 \pm 7			(γ)	19
6.299 \pm 4			(γ)	6, 19
6.339 \pm 4			(γ)	6, 19
6.375 \pm 4			(γ)	6, 19
6.416 \pm 4			(γ)	6, 19
6.441 \pm 9			(γ)	19
6.474 \pm 3			(γ)	6, 19
6.519 \pm 3	0 ⁺ ; $T = 2$		γ	9, 18
6.588 \pm 5			(γ)	19
6.6270 \pm 0.3	2 ⁻	0.31 \pm 0.02	γ, n	11, 12
6.6426 \pm 0.3	(3, 4)	< 0.08	γ, n	11
6.6475 \pm 0.4	1 ⁻	1.59 \pm 0.10	γ, n	11, 12
6.6934 \pm 0.6	1 ⁻	13.8 \pm 0.8	γ, n	6, 11, 12
6.7661 \pm 0.9	(2 ⁻ , 3, 4 ⁺)	\leq 0.6	γ, n	6, 11, 19
6.825 \pm 5			n	6, 12, 19
6.8567 \pm 1.0	2	10 \pm 2	γ, n	11
6.905 \pm 8				19
6.936 \pm 4				6
6.9678 \pm 1.0	1 ⁻	5 \pm 1	γ, n	6, 11, 12
(7.0670 \pm 1.2)	0 ⁻	(2.4 \pm 0.6)	γ, n	11, 12
7.08	(1 ⁺)	24	n	6, 12
7.166 \pm 2	2 ⁽⁺⁾	8 \pm 1	γ, n	6, 11, 12, 13
7.232 \pm 7				6
7.283 \pm 4				6
7.319 \pm 8	(1)	33	γ, n	6, 11, 12
7.37 \pm 20	(1)	19	n	6, 12
7.42 \pm 20	(2 ⁺)	10	γ, n	6, 11, 12

Table 20.2: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
7.495 \pm 5	(2)	80	γ, n	6, 11, 12
7.655 \pm 5	(2 ⁺)	65	γ, n	6, 11, 12
7.734 \pm 6		140	n	6, 12
7.843 \pm 11	1 ⁻	(50 \pm 10)	γ, n	6, 11
7.985 \pm 4	1	14 \pm 2	γ, n	6, 11
8.05 \pm 100	2 ⁺ ; $T = 2$			18
8.062 \pm 8				6
8.113 \pm 4		195	γ, n	6, 11, 12
8.147 \pm 6		15	n	6, 12
8.268 \pm 12				6
8.349 \pm 4				6
8.421		27	n	12
8.50		140	n	12
8.72		≤ 30	n	6, 12
8.77		76	n	6, 12
8.94		73	n	6, 12
9.01				6
9.2			n	10, 12
9.52		110	n	12
9.65		100	n	12
9.83		33	n	12
9.85		120	n	12
(9.886 \pm 10)			n	12
9.90		≤ 30	n	12
(9.929 \pm 10)			n	12
(9.981 \pm 10)			n	12
10.024 \pm 10		150	n, α	12, 13
10.10 \pm 50			n, α	13
10.228 \pm 10	0 ⁻ , 1	≈ 200	n, α	12, 13
10.480 \pm 10		≈ 10	n, α	12, 13
10.641 \pm 10	1, 2	70	n	12

Table 20.2: Energy levels of ^{20}F ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	τ	Decay	Reactions
10.807 \pm 10	$0^-, 1$	≈ 310	n, α	12, 13
10.99		190	n	12
(11.045 \pm 10)		≈ 30	n	12
(11.130 \pm 10)		< 25	n	12
(11.244 \pm 10)		< 25	n	12
(11.287 \pm 10)			n	12
11.49 \pm 50			n, α	13
12.0			n, α	13
12.2 \pm 100			n, α	13
12.4			n, α	13
12.7			n, α	10, 13
13.2			n, α	13
13.7			n, α	12, 13
14.0			n, α	13

^a See also Tables 20.3, 20.4 and 20.5.

5. $^{13}\text{C}(^{11}\text{B}, \alpha)^{20}\text{F}$ $Q_m = 9.3854$

The upper of the two states at 2.97 MeV has an excitation energy of 2968 ± 1.5 keV and γ branching ratios of (61 ± 4) and $(39 \pm 4)\%$, respectively, to $^{20}\text{F}^*(1.97, 0.82)$ [$J^\pi = (3^-, 4^+)$]: this is consistent with $J^\pi = (4^-)$ for $^{20}\text{F}^*(2.968)$.

6. $^{14}\text{N}(^7\text{Li}, \text{p})^{20}\text{F}$ $Q_m = 10.4985$

Tables 20.5 here and in (1983AJ01) display ^{20}F states reported in this reaction.

7. $^{16}\text{O}(^7\text{Li}, ^3\text{He})^{20}\text{F}$ $Q_m = -4.7442$

For reported states see Table 20.6 in (1983AJ01).

Table 20.3: Radiative transitions in ^{20}F ^a

E_i (MeV)	J_i^π	E_f (MeV)	Branching (%)	δ
0.66	3^+	0	100	0.10 ± 0.05
0.82	4^+	0	36 ± 3	
		0.66	64 ± 3 ^d	
0.98	1^-	0	100	f
1.06	1^+	0	100	
1.31	2^-	0	100	f
1.82	5^+	0.82	≥ 95	-0.03 ± 0.07
1.84	2^-	0	100	
1.97	(3^-)	0	16 ± 4	-0.06 ± 0.14
		0.82	55 ± 3	$+0.27 \pm 0.30$
		1.31	29 ± 3	
2.04	2^+	0	8.1 ± 1.9	
		0.66	91.9 ± 1.9	$0.08^{+0.06}_{-0.1}$
2.19	3^+	0	53.7 ± 2.1	0 ± 0.09
		0.82 ^d	46.3 ± 2.1	$+0.07 \pm 0.10$
2.86 ^b	(3^-)	0	(100)	
2.966 ^c	3^+	0	23 ± 3	
		0.66	21 ± 3	
		0.82	56 ± 3	
2.968	(4^-)	0.82	39 ± 4	
		1.97	61 ± 4	
3.17 ^b	(1^+)	0.98	> 95	
3.49 ^c	1^+	0	63 ± 4	
		1.06	22 ± 4	
		1.31	8 ± 2	
		1.84	7 ± 2	
3.53	0^+	1.06	100	
3.59	$(1, 2)^+$	0	30.5 ± 2.4	
		0.66	9.8 ± 1.2	
		0.98	3.5 ± 1.0	
		2.04	50 ± 3	

Table 20.3: Radiative transitions in ^{20}F ^a (continued)

E_i (MeV)	J_i^π	E_f (MeV)	Branching (%)	δ
		2.19	7 ± 3	
3.68	1, 2	0 ^d	66 ± 4	
		1.06	34 ± 4	
3.97	1 ⁺	0.98	24 ± 6	
		1.31	76 ± 6	
4.08	(1) ⁺	0	34 ± 3	
		1.06	66 ± 3	
4.28	(1, 2) ⁺	1.06	100	
4.37	(2 ⁺)	0.82	7 ± 3	
		0.98	34 ± 5	
		2.97	59 ± 5	
4.51	1 ⁺ , 2	0.66	100	
5.32	0 – 2	0	34 ± 7	
		0.98	66 ± 7	
5.56	1 ⁻	0	29 ± 4	
		1.31	37 ± 5	
		3.53	34 ± 7	
5.94	2 ⁻	0	6.4 ± 0.4	
		0.66	22.7 ± 1.4	
		0.98	11 ± 3	
		1.97	24.2 ± 1.4	
		2.04	2.5 ± 0.5	
		2.19	4.1 ± 0.6	
		2.97	2.5 ± 0.7	
		3.49	9.1 ± 1.2	
		3.59	7.8 ± 1.6	
		3.68	7.2 ± 0.8	
		3.97	(2.7 ± 0.9)	
6.02	2 ⁻	0	24.3 ± 1.2	
		0.66	3.05 ± 0.17	
		0.98	15.1 ± 0.7	

Table 20.3: Radiative transitions in ^{20}F ^a (continued)

E_i (MeV)	J_i^π	E_f (MeV)	Branching (%)	δ
6.05	0 - 2	1.31	0.83 ± 0.21	
		1.84	5.3 ± 0.4	
		1.97	1.04 ± 0.23	
		2.04	0.68 ± 0.15	
		2.19	3.8 ± 0.4	
		2.97	8.9 ± 0.5	
		3.49	20.8 ± 1.1	
		3.59	13.6 ± 2.4	
		4.08	2.6 ± 0.4	
		6.60 ^g	0 ⁺ , 1 ⁺	1.31
1.84	36 ± 3			
3.49	15 ± 5			
3.53	(28 ± 3)			
0	8.5 ± 0.6			
0.98	1.29 ± 0.10			
1.06	3.4 ± 0.3			
1.31	2.21 ± 0.13			
1.84	1.62 ± 0.11			
2.04	4.9 ± 0.3			
3.49	2.31 ± 0.18			
3.53	2.05 ± 0.21			
3.59	4.7 ± 0.3			
3.68	0.99 ± 0.18			
3.97	0.82 ± 0.12			
4.08	(0.90 ± 0.13)			
4.28	1.42 ± 0.20			
4.37	0.83 ± 0.14			
4.51	0.53 ± 0.17			
5.32	1.5 ± 0.5			
5.56	2.4 ± 0.4			
5.94	15.0 ± 1.5			

Table 20.3: Radiative transitions in ^{20}F ^a (continued)

E_i (MeV)	J_i^π	E_f (MeV)	Branching (%)	δ
e		6.02	40 ± 3	
		6.05	4.8 ± 0.5	

^a Branching ratios from (1983HU12) and from the earlier work displayed in Tables 20.5 in (1978AJ03) and 20.4 in (1983AJ01). Branching ratios renormalized to add to 100%, except for $^{20}\text{F}^*(6.60)$.

^b The population of $^{20}\text{F}^*(2.86)$ and the γ -decay of $^{20}\text{F}^*(3.17)$ are not observed by (1983HU12) [(n, γ)]. See also (1987AL06).

^c See, however, Table 20.5 in (1978AJ03).

^d Transition not observed by (1983HU12) because of a background problem.

^e For higher states see Tables 20.6 and 20.7.

^f Pure E1.

^g See Table 20.8.

8. (a) $^{18}\text{O}(\text{d}, \text{n})^{19}\text{F}$	$Q_m = 5.7697$	$E_b = 12.3710$
(b) $^{18}\text{O}(\text{d}, \text{p})^{19}\text{O}$	$Q_m = 1.732$	
(c) $^{18}\text{O}(\text{d}, \text{d})^{18}\text{O}$		
(d) $^{18}\text{O}(\text{d}, ^3\text{He})^{17}\text{N}$	$Q_m = -10.449$	
(e) $^{18}\text{O}(\text{d}, \alpha)^{16}\text{N}$	$Q_m = 4.247$	

See (1983AJ01) for a listing of the polarization measurements. For VAP measurements at $E_d^- = 52$ MeV (reaction (e)), see (1982MA25). See also ^{19}O and ^{19}F here, and ^{16}N and ^{17}N in (1986AJ04). See also (1986SE1B).

9. $^{18}\text{O}(^3\text{He}, \text{p})^{20}\text{F}$	$Q_m = 6.8774$
--	----------------

Proton groups have been observed to states of ^{20}F with $E_x < 4.1$ MeV: see Table 20.8 in (1978AJ03). Angular distributions, γ -ray polarization data and branching ratios lead to the J^π values shown in that table. A state at $E_x = 6519 \pm 3$ keV is also populated. It decays principally (> 90%) to $^{20}\text{F}^*(1.06)$ [$J^\pi = 1^+$]: the γ -rays are isotropic. $^{20}\text{F}^*(6.52)$ is the 0^+ ; $T = 2$ analog of the ground state of ^{20}O : see (1978AJ03).

Table 20.4: Lifetime measurements of some ^{20}F states ^a

$^{20}\text{F}^*$ (MeV)	τ_m
0.66	0.39 ± 0.03 psec
0.82	79 ± 6 psec
0.98	2.03 ± 0.20 psec ^a
	1.8 ± 0.3 psec ^b
	2.0 ± 0.2 psec ^A
1.06	45 ± 13 fsec
1.31	1.16 ± 0.20 psec ^a
	1.9 ± 0.3 psec ^b
	1.6 ± 0.3 psec ^A
1.82	≤ 65 fsec
1.84	30 ± 20 fsec
1.97	1.4 ± 0.4 psec
2.04	37 ± 16 fsec
2.19	< 12 fsec
2.97	60 ± 40 fsec
3.49	44 ± 11 fsec
3.53	30 ± 15 fsec
3.59	30 ± 30 fsec

A = adopted.

^a For references see Table 20.6 in (1978AJ03).

^b (1983K001): $^7\text{Li}(^{18}\text{O}, \alpha n)^{20}\text{F}$; $E_x = 983.58 \pm 0.16$ and 1309.19 ± 0.20 keV.

Table 20.5: Some states of ^{20}F reported in $^{14}\text{N}(^7\text{Li}, \text{p})^a$

E_x (keV)	J^π	E_x (keV)	J^π
4512 ± 4	$(3^-, 4^-, 5^+, 6^+)$	6695 ± 3	^c
4579 ± 4^b		6756 ± 3	$(2^-, 3, 4^+)$
4728 ± 5	$(3^-, 4^-, 4^+, 5^+)$	6823 ± 3	
4760 ± 5	$(4 \rightarrow 6^-, 6 \rightarrow 8^+)$	6936 ± 4	
4889 ± 4^b	^c	6968 ± 4	
5032 ± 4	2^-	6991 ± 7	
5064 ± 5	$(1^-, 2, 3^+)$	7034 ± 9^d	
5128 ± 5	$(2^-, 3, 4^+)$	7080 ± 7	
5222 ± 4	$(1, 2)^-$	7154 ± 5	
5282 ± 11	^c	7232 ± 7	
5316 ± 7	^c	7283 ± 4	
5350 ± 5	3^+	7319 ± 8	
5405 ± 4	^c	7370 ± 20	
5448 ± 6^b		7419 ± 20	
5560 ± 6^b		7495 ± 5	
5612 ± 5^b	^c	7655 ± 5	
5725 ± 10	$(2 \rightarrow 5)$	7734 ± 6	
5765 ± 8	3^+	7865 ± 16	
5803 ± 7	1^+	7975 ± 5	
5940 ± 5	^c	8062 ± 8	
6021 ± 4^b		8113 ± 4	
6090 ± 7	(0^-)	8147 ± 6	
6160 ± 5	$((1^-), 2, 3^+)$	8268 ± 12	
6193 ± 6	$(2^-, 3, 4^+)$	8349 ± 4	
6297 ± 5^b	^c	8573	
6344 ± 9^b	^c	8697	
6379 ± 5^b	^c	8754	
6417 ± 4	$(3^-, 4, 5, (6^+))$	8792	
6470 ± 4	^c	8907	
6565 ± 6^b	^c	8946	
6600 ± 8^b	^c	9022	
6633 ± 3^b			

^a (1985FO07); $E(^7\text{Li}) = 16$ MeV. For the low-lying states reported in this reaction see Table 20.6 in (1983AJ01). Please note that the density of states is very high and that when J^π assignments are made [based on cross sections and the $2J_f + 1$ relationship, with slopes which are different for even- and odd-parity states], these depend on the states having been resolved.

^b Unresolved [based on data from other reactions and on spectrum shown].

^c See (1985FO07).

^d All the observed groups for $E_x \gtrsim 7.0$ MeV appear to be due to unresolved states. See (1985FO07) for $\sigma_{\text{tot}}(0^\circ - 90^\circ)$ and J^π .

$$10. \text{}^{18}\text{O}(\alpha, \text{d})^{20}\text{F} \quad Q_m = -11.4758$$

At $E_\alpha = 64.4$ MeV angular distributions have been reported to $^{20}\text{F}^*(0, 0.66, 0.82, 1.06, 1.82, 2.20, 2.97, 4.24, 4.54, 5.07, 5.44, 5.80, 6.67, 7.29, 7.75, 8.34, 8.75, 9.00, 9.24, 9.78, 10.01, 10.51, 10.85, 11.56, 12.32, 12.72)$: L assignments are made [the groups above $E_x \approx 2.9$ MeV are probably unresolved] (1986KA36).

$$11. \text{}^{19}\text{F}(\text{n}, \gamma)^{20}\text{F} \quad Q_m = 6.6013$$

$$Q_0 = 6601.33 \pm 0.14 \text{ keV (1983HU12)}$$

$$Q_0 = 6601.344 \pm 0.055 \text{ keV (1986KE15)}$$

The thermal capture cross section is 9.8 ± 0.7 mb. A number of resonances have been observed: see Table 20.6. The primary γ -rays resulting from capture at thermal energies ($^{20}\text{F}^*(6.60)$; $J^\pi = 1^+$) and at $E_n = 27, 44$ and 49 keV ($^{20}\text{F}^*(6.63, 6.643, 6.647)$; $J^\pi = 2^-, (3, 4)$ and 1^-) have been studied by several groups: see (1972AJ02) and Table 20.7 here. It appears that the thermal capture [$^{20}\text{F}^*(6.60)$] is dominated by two intense transitions (E1) to $^{20}\text{F}^*(5.94, 6.02)$ [thus $J^\pi = 1^-, 2^-$]. If the ground-state transition is mainly M1, these two E1 transitions are (in terms of W.u.) about 150 times stronger than the M1 transition (1968SP01). See also (1983HU12). It appears also that at $^{20}\text{F}^*(6.63, 6.64, 6.65)$ [$J^\pi = 2^-, (3, 4)$ and 1^-] the E1 transitions to the ground state are very weak, even though other E1 transitions in the decay of these three states have approximately normal strengths. The strongest transitions from the 27 keV resonance appear to be M1. On the basis of J^π of the final states involved in the decay of the 44 keV resonance $J = 3$ or 4 , assuming dipole transitions. Branching ratios for other ^{20}F states involved in this reaction are shown in Table 20.3.

Table 20.6: Resonances in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

E_n (keV)	J^π ^b	Γ_γ (eV)	$\Gamma_{\text{c.m.}}$ (keV)	E_x in ^{20}F (MeV)
27.07 ± 0.05	2^-	1.4 ± 0.3	0.355 ± 0.03	6.6270
43.5 ± 0.1	(3, 4)	^c	< 0.08	6.6426
48.7 ± 0.3	1^-	1.6 ± 0.3	1.96 ± 0.3	6.6475
97.0 ± 0.5	1^-	6.0 ± 1.8 ^d	13.5 ± 1.5	6.6934
173.5 ± 0.9		^e	≤ 0.6	6.7661
269 ± 1	2	3.5 ± 0.8	10 ± 2	6.8567
(270 ± 8)	1	≤ 4.4		(6.859)
386 ± 1	1^-	2.4 ± 0.8	5 ± 1	6.9678
(490.5 ± 1)	0^-	($\geq 10 \pm 3$)	(2.4 ± 0.6)	(7.0671)
595 ± 2	2	6.3 ± 1.2	8 ± 1	7.166
760		2.9	60	7.32
865			60	7.42
950		2.8	95	7.50
1125		3.9	80	7.67
(1295 ± 12)	1^-	8.6	(50 ± 10)	(7.831)
1460 ± 3	1	$\geq 11 \pm 3$	14 ± 2	7.988
1635		11 ± 3	180	8.15

^a For complete references see Table 20.9 in (1978AJ03).

^b Assumed.

^c $g\Gamma_n = 0.086 \pm 0.02$ eV.

^d May be two resonances.

^e $g\Gamma_n = 0.35 \pm 0.1$ eV.

Table 20.7: Primary capture transitions in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

Final state $^{20}\text{F}^*$ (MeV)	I_γ ^b from		
	$^{20}\text{F}^*(6.63)$	$^{20}\text{F}^*(6.64)$	$^{20}\text{F}^*(6.65)$
0	2.0 ± 0.5		
0.66	6 ± 1	42 ± 7	
0.82		23 ± 7	
0.98			18 ± 4
1.06			9 ± 4
1.31	31 ± 2		
1.84	8 ± 2		
1.97	46 ± 4		
2.04	1.5 ± 1		59 ± 6
2.97		35 ± 9	
3.49	3 ± 1		14 ± 5
3.53	8 ± 1		
4.08	2.5 ± 1		

^a For complete references see Table 20.10 in (1978AJ03). See also Tables 20.3 and 20.6 here.

^b In units of photons/100 captures.

Table 20.8: States of ^{20}F involved in $^{19}\text{F}(n, \gamma)^{20}\text{F}$ ^a

E_x (keV)	J^π	E_x (keV)	J^π
0	2^+	3586.56 ± 0.09	$(1, 2)^+$
656.00 ± 0.04	3^+	3680.13 ± 0.06	1, 2
822.68 ± 0.08	4^+	3965.19 ± 0.16	1^+
983.71 ± 0.05	1^-	4082.08 ± 0.11	$(1)^+$
1057.02 ± 0.04	1^+	4277.22 ± 0.14	$(1, 2)^+$
1309.34 ± 0.05	2^-	4371.38 ± 0.12	(2^+)
1843.97 ± 0.08	2^-	4508.7 ± 0.4	$1^+, 2$
1970.80 ± 0.07	(3^-)	5318.87 ± 0.17	0 – 2
2044.05 ± 0.06	2^+	5555.34 ± 0.13	1, 2^+
2194.36 ± 0.08	(3^+)	5936.09 ± 0.05	2^-
2966.16 ± 0.08	3^+	6017.77 ± 0.03	2^-
3488.49 ± 0.06	1^+	6044.98 ± 0.08	0 – 2
3526.28 ± 0.07	0^+	6601.33 ± 0.04	$0^+, 1^+$ ^b

^a (1983HU12). For the earlier work see Table 20.11 in (1978AJ03). A state at 5713 ± 2 keV reported earlier is not seen here. (1987AL06) suggest that a 3428.4 ± 0.4 keV γ -ray reported by (1983HU12) feeds $^{20}\text{F}^*(3.17)$ and that its excitation energy is then 3172.58 ± 0.42 keV [for the decay of this state see Table 20.3].

^b The transition $C \rightarrow 3.53$ [$J^\pi = 0^+$] is observed.

Table 20.8 displays excitation energies for ^{20}F states involved in cascade and in primary γ -transitions (1983HU12). For the earlier references see (1978AJ03). See also (1984VO1H) and (1986KR16; theor.).

12. (a) $^{19}\text{F}(\text{n}, \text{n})^{19}\text{F}$ $E_{\text{b}} = 6.6013$
 (b) $^{19}\text{F}(\text{n}, \text{n}')^{19}\text{F}^*$
 (c) $^{19}\text{F}(\text{n}, 2\text{n})^{18}\text{F}$ $Q_{\text{m}} = -10.4320$

The scattering amplitude (bound) $a = 5.654 \pm 0.010$ fm, $\sigma_{\text{free}} = 3.641 \pm 0.010$ b (1979KO26). The difference in the bound-state scattering lengths, $b^+ - b^- = -0.019 \pm 0.002$ (1979GL12). The total cross section has been measured for $E_{\text{n}} = 0.5$ to 29.1 MeV: see (1978AJ03). Observed resonances are displayed in Table 20.9.

Observed resonances in the excitation functions involving $^{19}\text{F}^*(0.11, 1.5(\text{u}))$ are displayed in Table 20.10. For reaction (c) see (1983CSZX). See also (1986BAYL) and (1986SA40; theor.).

13. $^{19}\text{F}(\text{n}, \alpha)^{16}\text{N}$ $Q_{\text{m}} = -1.523$ $E_{\text{b}} = 6.6013$

Reported resonances are shown in Table 20.11.

14. $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$ $Q_{\text{m}} = 4.3767$

States of ^{20}F observed in this reaction are displayed in Table 20.12. See (1978AJ03) for a discussion of the earlier work. See also (1983JI04).

15. $^{20}\text{O}(\beta^-)^{20}\text{F}$ $Q_{\text{m}} = 3.8136$

The decay is to $^{20}\text{F}^*(1.06, 3.49)$, $J^\pi = 1^+$: see ^{20}O . The E_{x} of $^{20}\text{F}^*(1.06)$ is 1056.848 ± 0.004 keV. The β branch to $^{20}\text{F}^*(3.17)$ (1^+) is $< 0.012\%$, $\log f_0 t > 5.1$ (1987AL06). [See also for general discussion of the low-lying states of ^{20}F].

16. $^{20}\text{Ne}(\pi^-, \gamma)^{20}\text{F}$ $Q_{\text{m}} = 132.505$

The branching ratio to $^{20}\text{F}^*(1.06)$ [$J^\pi = 1^+$] is compared to the analogous M1 decay width $^{20}\text{Ne}^*(11.24)$ [$J^\pi = 1^+$] \rightarrow $^{20}\text{Ne}_{\text{g.s.}}$. The M1 amplitude contains $(47 \pm 16)\%$ spin-flip, in agreement with shell-model calculations. The population of $^{20}\text{F}^*(0, 1.31, 1.84)$ [$J^\pi = 2^+, 2^-, 2^-$] is also reported (1981MA04). See also (1986BA16) and (1983KN05; theor.).

Table 20.9: Resonances in $^{19}\text{F}(n, n)^{19}\text{F}$ ^a

E_n (keV)	Γ_{lab} (keV)	J^π	$^{20}\text{F}^*$ (MeV)
26.99	0.325 ± 0.020	2^-	6.6269
48.78	1.67 ± 0.10	1^-	6.6476
97.50	14.5 ± 0.8	1^-	6.6939
500	25^{b}	(1^+)	7.076
600	15^{b}	(2^+)	7.171
747	35^{b}	(1)	7.311
794	20	(1)	(7.355)
852	11^{b}	(2^+)	7.410
935	60	(2)	7.489
1100	50	(2^+)	7.65
1250	150		7.79
1620	220		8.14
2000	150		8.50
2250	≤ 30		8.74
2280	80		8.77
2520	150		8.99
3250	150		9.69
3420	130		9.85
3460 ± 10			(9.886)
3505 ± 10			(9.929)
3560 ± 10			(9.981)
3605 ± 10	200		10.024
3820 ± 10	≈ 200	$0^-, 1$	10.228
4085 ± 10	≈ 10		10.480
4255 ± 10	≈ 60	1, 2	10.641
4430 ± 10	≈ 330	$0^-, 1$	10.807
4680 ± 10	≈ 30		11.045
4770 ± 10	< 25		11.130
4890 ± 10	< 25		11.244
(4935)			(11.287)

^a For references see Table 20.12 in (1978AJ03).

^b $\Gamma_\gamma = 3.3 \pm 1.0, 6.3 \pm 1.2, 2.4 \pm 0.8$ and 1.5 ± 0.5 eV for $^{20}\text{F}^*(7.08, 7.17, 7.31, 7.41)$.

Table 20.10: States of ^{20}F from resonances in $^{19}\text{F}(n, n'\gamma)^{19}\text{F}$

E_n (keV)	Γ_{lab} (keV)	Resonance in		E_x in ^{20}F (MeV)
		$\gamma_{0.11}$ ^a	$\gamma_{1.5}$ ^b	
240		r		6.829
270		r		6.858
386		r		6.968
420		r		7.000
490		r		7.066
620		r		7.190
800		r		7.361
860		r		7.418
1150 ^c		r		7.693
1250		r		7.788
1580		r		8.101
1645	15	r	r	8.163
1916	28		r	8.421
2240	45		r	8.728
2465	75	r	r	8.942
2700		r		9.165
3075	120		r	9.521
3215	80		r	9.654
3400	35		r	9.830
3475	≤ 30		r	9.901
3620	120	r	r	10.038
4240	90	r	r	10.627
4620	200		r	10.988
4900	≤ 50		r	11.254
7300		r		13.532

r = resonant.

^a Resonances in yield of 0.11 MeV γ -rays at $\theta = 92^\circ$: values for E_n read by reviewer from differential cross section tables. See Table 20.13 in (1978AJ03) for references.

^b Resonances in yields of ^{19}F with $E_x \approx 1.5$ MeV: see (1973MA14).

^c Appears to be unresolved.

Table 20.11: Resonances in $^{19}\text{F}(n, \alpha)^{16}\text{N}$ ^a

E_n (MeV \pm keV)	E_x (MeV)
3.4	9.8
3.61 ± 50	10.03
3.69 ± 50	10.10
3.76 ± 40	10.17
4.09 ± 40	10.48
4.39 ± 40	10.77
4.52^b	10.89
4.82 ± 40	11.18
5.15 ± 50	11.49
5.40^b	11.73
5.7	12.0
5.9 ± 100^b	12.2
6.10	12.39
6.55	12.82
6.9	13.2
7.44	13.66
7.8	14.0

^a For references see Table 20.14 in (1978AJ03). See also graph in (1976GAYV).

^b Not resolved.

17. $^{21}\text{Ne}(\text{d}, ^3\text{He})^{20}\text{F}$

$$Q_m = -7.514$$

The ^{20}F states observed at $E_d = 26$ MeV in this reaction and analog [$T = 1$] states observed in ^{20}Ne in the (d, t) reaction are displayed in Table 20.16 of (1978AJ03). The spectroscopic factors of analog states are consistent to within 20% for states excited by a single l -transfer.

Table 20.12: States in ^{20}F from $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$ ^a

E_x (keV) ^b	l_n ^c	J^π	$(2J + 1)S$	n, l, j ^c
0	2	2^+	0.054	$1d_{5/2}$
655.9 ± 0.2	2	3^+	2.32	$1d_{5/2}$
823.0 ± 0.3	d	4^+	0.32	$1g_{9/2}$
983.9 ± 0.3	d	1^-	0.014	$1p_{1/2}$
1057.0 ± 0.2	$0 + 2$	1^+	0.013	$2s_{1/2}$
1309.3 ± 0.2	d	2^-	0.017	$1p_{3/2}$
1820 ± 10	d	(5^+)	0.35	$1g_{9/2}$
1843.5 ± 0.7	d	2^-	0.007	$2p_{3/2}$
1970 ± 10	d	(3^-)	0.038	$1f_{7/2}$
2043.7 ± 0.5	2	2^+	2.32	$1d_{5/2}$
2194.5 ± 0.6	2	3^+	0.55	$1d_{5/2}$
2863.7 ± 1.6	d		0.044	$1f_{7/2}$
2966.8 ± 0.6	2	3^+	0.38	$1d_{3/2}$
3175.6 ± 1.3	d		0.019	$1d_{5/2}$
3488.5 ± 0.3	0	1^+	1.20 ^e	$2s_{1/2}$
3525.9 ± 0.5	0	0^+	0.28 ^e	$2s_{1/2}$
3586.5 ± 0.6	2	$\pi = +$	0.038	$1d_{3/2}$
3681.0 ± 2.5	2	$\pi = +$	0.031	$1d_{5/2}$
3760.8 ± 2.7	d		^c	
3964.5 ± 2.5	2	$\pi = +$	0.036	$1d_{5/2}$
4082.5 ± 0.8	$0 + 2$	$\pi = +$	0.13	$1s_{1/2}$
4198.9 ± 2.7	d		0.083	$1d_{3/2}$
4207.7 ± 2.6				
4279.0 ± 2.0	2	$\pi = +$	0.087	$1d_{5/2}$
4315.4 ± 2.0	0	$(0, 1)^+$	0.20	$2s_{1/2}$

Table 20.12: States in ^{20}F from $^{19}\text{F}(\text{d}, \text{p})^{20}\text{F}$ ^a (continued)

E_x (keV) ^b	l_n ^c	J^π	$(2J + 1)S$	n, l, j ^c
4374.5 \pm 2.0				
4509.5 \pm 3.0				
4583.8 \pm 3.0			0.02	2p _{3/2}
	1	(0 - 2) ⁻		
4592.2 \pm 2.9			(< 0.05)	(1f _{7/2})
4730.2 \pm 2.9	2, 3			
4763.8 \pm 2.7	2, 3			
4891.6 \pm 2.8				
4898.2 \pm 2.8				
5048.7 \pm 1.5				
5069.0 \pm 3	2	(1, 2, 3) ⁺	0.09	1d _{5/2}
5132.4 \pm 3.5				
5225.0 \pm 3	1, 3		0.09	2p _{3/2}
5284.0 \pm 3	0	(1, 0) ⁺	0.34	2s _{1/2}
5318.0 \pm 3	2 or 1 + 3	(1, 2, 3) ⁺ or 2 ⁻	0.10	1d _{5/2}
5349.0 \pm 4	2	(1, 2, 3) ⁺	0.06	1d _{5/2}
5408.2 \pm 2.5				
5450.3 \pm 3.8				
5455.4 \pm 3.2				
5463.0 \pm 3	2	(1, 2, 3) ⁺	0.27	1d _{5/2}
5562.9 \pm 2.0	1	(0, 1, 2) ⁻	0.03	2p _{3/2}
5588.1 \pm 1.5				
5620.0 \pm 3	d			
5710.8 \pm 6.0	d			
5764.0 \pm 3	2	(1, 2, 3) ⁺	0.15	1d _{5/2}
5810.0 \pm 3	0 + 2, or 1 + 3	(2 ⁻ , 1 ⁺)		
5935.0 \pm 3	1(+3)	(1 ⁻ , 2 ⁻)	0.43	2p _{3/2}
6015.0 \pm 3.8	1 + 3	(2 ⁻)	0.68	2p _{3/2}
			1.40	1f _{7/2}
6043.3 \pm 3.7				

^a For complete references see Table 20.15 in (1978AJ03) and see also Table 20.14 in (1983AJ01).

^b Best values.

^c Assumed in analysis; $E_d = 12$ MeV.

^d Weak groups.

^e At $E_d = 16$ MeV.

18. $^{22}\text{Ne}(\text{p}, \text{}^3\text{He})^{20}\text{F}$ $Q_m = -15.6516$

At $E_p = 43.7$ to 45.0 MeV analog states have been studied in ^{20}F and ^{20}Ne [the latter via $^{22}\text{Ne}(\text{p}, \text{t})^{20}\text{Ne}$]. Angular distributions for the ^3He ions and the tritons corresponding to the first $T = 2$ states ($J^\pi = 0^+$) [$^{20}\text{Ne}^*(16.722 \pm 0.025)$ and $^{20}\text{F}^*(6.513 \pm 0.033)$] have been compared. There is indication also for the excitation of the 2^+ ; $T = 2$ states [at $E_x = 8.05$ MeV in ^{20}F and at 18.5 MeV in ^{20}Ne (estimated ± 0.1 MeV)]: see (1978AJ03).

19. $^{22}\text{Ne}(\text{d}, \alpha)^{20}\text{F}$ $Q_m = 2.702$

Angular distributions have been obtained at $E_d = 10$ MeV to ^{20}F states with $E_x < 4.4$ MeV: they are generally featureless. Observed states are displayed in Table 20.17 of (1978AJ03).

20. $^{27}\text{Al}(^{20}\text{Ne}, ^{27}\text{Si})^{29}\text{F}$ $Q_m = -11.840$

The Δ resonance is very strongly excited in the reaction at $E(^{20}\text{Ne}) = 950$ MeV/A (1986BA16).

^{20}Ne
(Figs. 11 and 13)

GENERAL: (See also (1983AJ01).)

Shell model: (1978WI1B, 1982BR08, 1982FL04, 1982RA1N, 1982SH30, 1983BR29, 1983DR04, 1983DR03, 1984JA15, 1984PA04, 1984RA13, 1985AN16, 1985HA15, 1985HU08, 1985MI23, 1985MU10, 1986CH28, 1986COZZ, 1986HU1G, 1986WA1R, 1987PR01).

Collective, deformed and rotational models: (1981OK02, 1982BR08, 1982RA1N, 1982RO06, 1982SC20, 1983DR04, 1983DR03, 1983LO05, 1983MA29, 1983MA68, 1983SC08, 1983VA16, 1983WA04, 1984DR01, 1984GR29, 1984KA11, 1984LO05, 1984PA04, 1984PR09, 1984RI01, 1984SA37, 1984SC41, 1985HA15, 1985LE1H, 1985OM01, 1985RO1G, 1985ST23, 1985WA08, 1986BA3P, 1986BE1D, 1986CEZW, 1986LE16, 1986SN1B, 1986STZT, 1986SU01, 1987MI07, 1987ZH1B).

Cluster and α -particle models: (1982AO06, 1982LE1P, 1982SU1B, 1983BI1F, 1983FU1D, 1983PI03, 1983MA68, 1984KA11, 1984SE20, 1984ZH07, 1985VO1E, 1986AB1H, 1986CO15, 1986GR1Q, 1986KA17, 1986KAZP, 1986LE06, 1986MAZC, 1986SU01).

Special states: (1978WI1B, 1981OK02, 1982AO06, 1982BR08, 1982FL04, 1982RA1N, 1982SC20, 1982SH30, 1982VA1E, 1983AD1D, 1983AR07, 1983BA50, 1983BI1C, 1983BR29, 1983LO05, 1983MA23, 1983VA21, 1983WA04, 1984AD1E, 1984AM03, 1984BY02, 1984DR01, 1984GR29, 1984HA14, 1984IA1A, 1984KA11, 1984RA13, 1984SA37, 1984SC01, 1984SC41, 1984SE20, 1984ZH07, 1985AD1A, 1985HA18, 1985HA1K, 1985HA15, 1985HU08, 1985LI19, 1985MI10, 1985MU10, 1985OM01, 1985RO1G, 1985VO1E, 1985WA08, 1986AN10, 1986AN07, 1986BA3P, 1986CA27, 1986CEZW, 1986CH28, 1986GR1Q, 1986HU1G, 1986KA17, 1986MAZC, 1986STZT, 1986SU01, 1986WI1P, 1987HA08, 1987IA1B, 1987KA18, 1987MI07, 1987PR01, 1987YA03).

Electromagnetic transitions and giant resonances: (1978WI1B, 1981OK02, 1982BR24, 1982RO06, 1983BA50, 1983BR29, 1983BR1P, 1983LO05, 1983VA16, 1984DR01, 1984HA14, 1984KA11, 1984LO05, 1984NA21, 1984SC01, 1984WE13, 1985AL21, 1985VE04, 1985WI17, 1986AN10, 1986CA15, 1986CA27, 1986CH28, 1986KA17, 1986KA2G, 1986KAZP, 1986SU01).

Astrophysical questions: (1981WA1Q, 1982CA1A, 1982HI1E, 1982NO1D, 1982WO1A, 1983AL23, 1983HA1P, 1983SI1B, 1983WE1A, 1984BA34, 1984CO1H, 1984TR1C, 1985AR1A, 1985DW1A, 1985NO1H, 1986MA1E, 1986TH1E, 1986TR1C, 1987BO1B, 1987CU1A, 1987MA1R, 1987ME1B, 1987MU1B, 1987PR1A).

Applications: (1984WI1E, 1986BO1L, 1986FO1D, 1986TR1B, 1987NA1D).

Complex reactions involving ^{20}Ne : (1982FA1D, 1982HO10, 1982KA33, 1982VI01, 1983BA2J, 1983DE26, 1983FR1G, 1983GA01, 1983HU1E, 1983IS1E, 1983JA05, 1983LE1F, 1983ME1R, 1983RA1E, 1983RO1H, 1983SC1M, 1983TO1H, 1983WE1C, 1983WI1A, 1984BE22, 1984DE1Q, 1984FI17, 1984GR08, 1984HO23, 1984KA1J, 1984NA12, 1984TS03, 1985AG1A, 1985BE40, 1985GAZT, 1985HA1N, 1985HO05, 1985KA1E, 1985KA1G, 1985KAZQ, 1985MC03, 1985MO08,

1985OS05, 1985SA40, 1985ST20, 1986AI1A, 1986BA1E, 1986BE2M, 1986BL06, 1986CH2G, 1986GR1A, 1986GR1B, 1986HA1B, 1986HE1A, 1986IK03, 1986MA19, 1986ME06, 1986PL02, 1986PO06, 1986SA2N, 1986SA30, 1986SC28, 1986SC29, 1986SH07, 1986SH1F, 1986SO10, 1986TR1B, 1986VA10, 1986VA18, 1986VA23, 1986WE1C, 1987FA09, 1987KO15, 1987NI04, 1987RI03, 1987RO10).

Table 20.13: Energy levels of ^{20}Ne ^a

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m ^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
0	$0^+; 0$	0_1^+		stable	2, 3, 7, 8, 12, 15, 16, 18, 20, 24, 25, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 56, 57, 58, 59, 60, 61, 62, 63, 64
1.633674 ± 0.015	$2^+; 0$	0_1^+	$\tau_m = 1.05 \pm 0.06$ psec $g = +0.54 \pm 0.04$	γ	2, 3, 7, 8, 9, 11, 12, 15, 16, 18, 19, 20, 23, 24, 25, 29, 30, 31, 32, 33, 35, 36, 38, 40, 41, 42, 43, 46, 47, 49, 52, 53, 54, 55, 56, 57, 60, 61
4.2477 ± 1.1	$4^+; 0$	0_1^+	$\tau_m = 93 \pm 9$ fsec $g = +0.13 \pm 0.15$	γ	2, 3, 7, 8, 9, 12, 15, 16, 18, 19, 20, 23, 24, 29, 30, 31, 32, 33, 36, 38, 41, 42, 47, 54, 57, 60, 61
4.96651 ± 0.20	$2^-; 0$	2^-	$\tau_m = 4.8 \pm 0.5$ psec	γ	2, 3, 7, 8, 9, 12, 15, 24, 25, 29, 30, 31, 32, 33, 54, 56, 57, 60, 61
5.6214 ± 1.7	$3^-; 0$	2^-	200 ± 50 fsec	γ, α	2, 3, 7, 8, 12, 15, 29, 30, 32, 33, 55, 56, 57, 60, 61
5.7877 ± 2.6	$1^-; 0$	0^-	$\Gamma = (2.8 \pm 0.3) \times 10^{-2}$	γ, α	2, 3, 7, 8, 12, 14, 15, 16, 18, 30, 32, 33, 52, 55, 60
6.725 ± 5	$0^+; 0$	0_2^+	19.0 ± 0.9	γ, α	8, 12, 14, 15, 24, 29, 30, 32, 33, 36, 52, 60
7.004 ± 3.6	$4^-; 0$	2^-	$\tau_m = 440 \pm 90$ fsec	γ	2, 7, 8, 15, 30, 33, 56, 60
7.1563 ± 0.5	$3^-; 0$	0^-	$\Gamma = 8.2 \pm 0.3$	γ, α	2, 4, 7, 8, 12, 14, 15, 16, 18, 20, 23, 24, 29, 30, 52
7.191 ± 3	$0^+; 0$	0_3^+	3.4 ± 0.2	γ, α	5, 6, 7, 12, 14, 36, 60
7.4219 ± 1.2	$2^+; 0$	0_2^+	15.1 ± 0.7	γ, α	2, 5, 6, 7, 12, 14, 15, 29, 30, 32, 36, 53, 55, 60
7.829 ± 2.4	$2^+; 0$	0_3^+	2	γ, α	2, 6, 7, 12, 14, 24, 30, 36, 53, 55, 60
8.453 ± 4	$5^-; 0$	2^-	0.013 ± 0.004	γ, α	2, 6, 7, 12, 14, 15, 30, 60
≈ 8.7	$0^+; 0$	0_4^+	> 800	α	14

Table 20.13: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
8.708 ± 7	$1^-; 0$		2.1 ± 0.8	γ, α	7, 12, 14, 30, 60
8.7776 ± 2.2	$6^+; 0$	0_1^+	0.11 ± 0.02	γ, α	2, 4, 6, 7, 12, 14, 15, 16, 18, 19, 20, 23, 24, 30, 52, 60
≈ 8.8	$2^+; 0$	0_4^+	> 800	α	14, 30
8.82	$(5^-); 0$		< 1	α	14
8.854 ± 5	$1^-; 0$	1^-	19	α	7, 14, 55
9.031 ± 7	$4^+; 0$	0_3^+	3	γ, α	2, 6, 7, 12, 14, 24, 30, 60
9.116 ± 3	$3^-; 0$		3.2	γ, α	2, 7, 12, 14, 29, 30, 60
9.318 ± 2	$(2^-); 0$			γ	7, 12, 30, 60
9.487 ± 5	$2^+; 0$		29 ± 15	γ, α	12, 14, 53, 60
9.873 ± 4	$3^+; 0$			γ	7, 30, 53
9.935 ± 12	$(1^+); 0$		$\tau_m < 35$ fsec	γ	7, 30, 60
9.990 ± 8	$4^+; 0$	0_2^+	$\Gamma = 155 \pm 30$	γ, α	2, 7, 12, 14, 29, 30, 60
10.262 ± 5	$5^-; 0$	0^-	145 ± 40	α	2, 4, 7, 14, 15, 16, 18, 20, 30, 52
10.274 ± 3	$2^+; 1$		≤ 0.3	γ, α	12, 14, 53, 55
10.406 ± 5	$3^-; 0$	1^-	80	α	7, 14, 30, 60
10.553 ± 5	$4^+; 0$		16	α	7, 14, 30
10.584 ± 5	$2^+; 0$		24	α	14, 30, 53, 60
10.609 ± 6	$6^-; 0$	2^-	$\tau_m = 23 \pm 7$ fsec	γ	2, 6, 7
10.694 ± 6	$4^-, 3^+; 0$			γ	6, 7
10.80 ± 75	$4^+; 0$	0_4^+	$\Gamma = 350$	α	14, 15, 30
10.843 ± 4	$2^+; 0$		13	α	14, 53, 60
10.840 ± 6	$3^-; 0$		45	γ, α	7, 14
10.884 ± 3	$3^+; 1$		$\tau_m < 30$ fsec	γ	53, 55
10.917 ± 6	$3^+; 0$			γ	7
10.97 ± 120	$0^+; 0$	0_5^+	$\Gamma = 580$	α	14
11.020 ± 8	$4^+; 0$		24	α	6, 7, 14, 60
11.090 ± 3	$4^+; 1$		≤ 0.5	γ, α	12, 14, 30, 55
11.24 ± 23	$1^-; 0$		175	α	14, 30
11.2623 ± 1.9	$1^+; 1$			γ	12, 35, 36, 38, 53
11.270 ± 5	$1^-; 1$		≤ 0.3	γ, α	12, 14
11.320 ± 9	$2^+; 0$		40 ± 10	α	14, 53
11.528 ± 6	$3^+, 4^-; 0$		$\tau_m \leq 30$ fsec	γ	7, 30
11.555 ± 6	$(3^+); 0$			γ	7, 30
11.558 ± 4	$0^+; 0$	0_6^+	$\Gamma = 1.1 \pm 0.4$	γ, α	12, 14
11.601 ± 10	$2^-; 1$				55
11.653 ± 5	$(3^+); 0$			γ	6, 7, 36

Table 20.13: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
11.885 \pm 7	2 ⁺ ; 0		46	γ, α	7, 14, 30, 53, 60
11.928 \pm 4	4 ⁺ ; 0		0.44 \pm 0.15	γ, α	12, 14, 60
11.951 \pm 4	8 ⁺ ; 0	0 ₁ ⁺	(3.5 \pm 1.0) $\times 10^{-2}$	γ, α	4, 6, 7, 8, 12, 14, 15, 16, 18, 19, 23, 30, 52
11.985 \pm 16	1 ⁻ ; 0		30 \pm 5	γ, α	7, 12, 14
12.098 \pm 6	2 ⁻ ; 1			γ	7, 30, 38, 55
12.137 \pm 5	6 ⁺ ; 0	0 ₃ ⁺		α	5, 6, 7, 8, 14, 15
12.221 \pm 4	2 ⁺ ; 1		< 1	γ, α	7, 12
12.253 \pm 10	4 ⁺ ; 0		155 \pm 15	α	14
12.256 \pm 3	3 ⁻ ; 1		< 1	γ, α	12, 14
12.327 \pm 10	2 ⁺ ; 0	0 ₅ ⁺	390 \pm 50	α	14
12.401 \pm 5	3 ⁻ ; (1)	0 ₇ ⁺	37.3 \pm 0.9	γ, α	6, 7, 12, 14, 29, 60
12.433 \pm 5	0 ⁺ ; 0		24.4 \pm 0.5	γ, α	7, 12, 14
12.472 \pm 10	(2 ⁺); 0		124 \pm 6	α	14
12.585 \pm 5	6 ⁺ ; 0	(0 ₂ ⁺)	72 \pm 9	α	6, 7, 14, 15, 16, 18, 19
12.592 \pm 15	(2 ⁺); 0		145 \pm 25	α	14
12.713 \pm 5	5 ⁻ ; 0	1 ⁻	84 \pm 8	α	6, 7, 14
12.743 \pm 10	(2 ⁺); 0		61 \pm 12	α	6, 7, 14
12.836 \pm 5	1 ⁻ ; 0		30 \pm 5	α	7, 14
12.957 \pm 5	2 ⁺ ; 0	(0 ₇ ⁺)	38 \pm 4	α	7, 14, 60
13.048 \pm 5	4 ⁺ ; 0		18 \pm 3	α	6, 7, 14
13.0607 \pm 2.1	2 ⁻		1.0	p, α	28
13.099 \pm 10	(0 ⁺); 0		53 \pm 24	α	14
13.105 \pm 5	6 ⁺ ; 0	(0 ₂ ⁺)	102 \pm 5	α	14
13.137 \pm 5	3 ⁻ ; 0		48 \pm 4	α	14
13.1713 \pm 2.1	1 ⁺ ; (1)		2.3 \pm 0.2	γ, p, α	25, 26, 28, 29
13.222 \pm 10	0 ⁺ ; 0		40 \pm 13	α	7, 14, 28
13.224 \pm 15	1 ⁻ ; 0		80	p, α	14, 28
13.226 \pm 5	3 ⁻ ; 0		53 \pm 4	α	14
13.3075 \pm 2.1	1 ⁺		0.9 \pm 0.1	γ, p, α	25, 26, 28
13.338 \pm 5	7 ⁻ ; 0	2 ⁻	(8 \pm 3) $\times 10^{-2}$	α	6, 7, 8, 14
13.341 \pm 5	4 ⁺ ; 0		26 \pm 3	α	14
13.414 \pm 2	3 ⁻ ; 0		24 \pm 3	α	14, 25, 26, 28
13.426 \pm 5	(5 ⁻); 0		49 \pm 7	α	14
13.461 \pm 10	1 ⁻		195 \pm 25	p, α	14, 28
13.484 \pm 2	1 ⁺ ; 1		6.4 \pm 0.3	γ, p, α	25, 26, 28, 38
13.507 \pm 5	1 ⁻ ; 0		24 \pm 8	p, α	14, 26, 28
13.529 \pm 5	2 ⁺ ; 0		61 \pm 8	α	14

Table 20.13: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
13.530 \pm 15	(0 ⁺); 0		76 \pm 32	α	14
13.573 \pm 5	2 ⁺ ; 0		12 \pm 5	α	7, 14, 28
13.586 \pm 3	2 ⁺		9 \pm 1	p, α	26, 28
13.642 \pm 3	0 ⁺ ; 1		17 \pm 1	p, α	7, 26, 28, 29
13.676 \pm 2.3	(2 ⁻)		4.5 \pm 0.2	γ , p, α	25, 26, 28
13.677 \pm 5	5 ⁻ ; 0		11 \pm 2	α	6, 14
13.692 \pm 10	7 ⁻ ; 0	0 ⁻	310 \pm 30	α	14
13.736 \pm 2.5	1 ⁺		7.7 \pm 0.5	γ , p, α	25, 26, 28
13.744 \pm 20	0 ⁺ ; 0		\approx 80	α	14
13.827 \pm 10	3 ⁻ ; 0		136 \pm 15	α	7, 14
13.866 \pm 30	1 ⁻ ; 0		\approx 175	p, α	7, 14, 28
13.881 \pm 2.3	2 ⁺ ; 1		0.14 \pm 0.05	γ , p, α	7, 8, 25, 26, 28, 29
13.908 \pm 5	2 ⁺ ; 0		74 \pm 10	α	14, 28
13.926 \pm 2.3	(0 ⁺)		3.5 \pm 0.4	p, α	28
13.928 \pm 5	6 ⁺ ; 0		65 \pm 3	α	14, 15, 16
13.948 \pm 10	0 ⁺ ; 0		79 \pm 15	α	14
13.965 \pm 5	4 ⁺ ; 0	(0 ₆ ⁺)	8.1 \pm 1	α	14
14.02	1 ⁻		\approx 70	p, α	28
14.063 \pm 2.3	2 ⁺		\approx 140	p, α	26, 28
14.115 \pm 5	2 ⁺ ; 0		42 \pm 6	α	14
14.128 \pm 2	2 ⁻		4.7 \pm 0.7	γ , p, α	25, 26, 28
14.150 \pm 2.3	2 ⁻		11.8 \pm 1.0	γ , p, α	25, 26, 28
14.20	1 ⁺		14 \pm 1	γ , p	25, 26
14.270 \pm 10	4 ⁺ ; 0		92 \pm 9	α	14
14.304 \pm 10	(6 ⁺); 0		60 \pm 13	α	6, 7, 14
14.311 \pm 5	6 ⁺ ; 0		117 \pm 8	α	6, 7, 14, 15, 16, 18
14.313 \pm 15	(3 ⁻); 0		\approx 45	α	14
14.370 \pm 3			\approx 5	p, α	26, 28
14.454 \pm 5	5 ⁻ ; 0		\approx 15	α	14
14.455 \pm 3	(0 ⁺ , 2 ⁺); 0		33 \pm 3	p, α	14, 26, 28
14.475 \pm 6	0 ⁺		68 \pm 2	p, α	26, 28
14.597 \pm 7	1 ⁻ ; 0		116 \pm 5	p, α	14, 28
14.593 \pm 10	4 ⁺ ; 0		260 \pm 25	α	14
14.653 \pm 10	(0 ⁺)		25	p, α	26, 28
14.699 \pm 3.3	(1 ⁺)		36 \pm 10	p, α	14, 26, 28
14.731 \pm 10	(4 ⁺); 0		60 \pm 25	α	14
14.761 \pm 5	6 ⁺ ; 0		7.3 \pm 4.8	α	14
14.776 \pm 4	(1 ⁻)		110 \pm 20	p, α	26, 28

Table 20.13: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
14.807 \pm 5	6 ⁺ ; 0		86 \pm 7	α	6, 14, 28
14.816 \pm 5	5 ⁻ ; 0		117 \pm 13	α	6, 14
14.839 \pm 10	(4 ⁺); 0		79 \pm 15	α	14
14.888 \pm 10	2 ⁺ ; 0		100 \pm 30	p, α	14, 28
15.047 \pm 10	2 ⁺ ; 0		66 \pm 20	p, α	7, 14, 28
15.073 \pm 10	5 ⁻ ; 0		160 \pm 25	α	14
15.142 \pm 15	(2 ⁺); 0		\approx 60	α	14
15.174 \pm 10	5 ⁻ ; 0		230 \pm 25	α	6, 14
15.23			28	p, α	28
15.27	(1 ⁻)		285	p, α	28
15.319 \pm 25	7 ⁻ ; 0		280 \pm 40	α	4, 6, 7, 14, 15, 16, 18
15.330 \pm 5	4 ⁺ ; 0		34 \pm 10	α	4, 6, 7, 14
15.366 \pm 5	7 ⁻ ; 0		110 \pm 10	α	14, 15, 16, 18, 19
15.436 \pm 15	(3 ⁻); 0		90 \pm 20	p, α	7, 14, 28
15.5			55	p, α	14, 28
15.70 \pm 15	(8 ⁻); 0	(2 ⁻)		α	6, 7, 14
15.874 \pm 9	8 ⁺		100 \pm 15	α	5, 6, 7, 15, 18, 19
15.97	(6 ⁺); 0			α	14
16.01 \pm 25	(2 ⁺ ; 1)		100	p, α	28
16.139 \pm 15			38	α	6, 7, 14, 28
16.25				α	6, 14
16.329 \pm 11	4 ⁺ ; 0		45	p, α	14, 28
16.437 \pm 11	(0, 2, 4) ⁺ ; 0		35	α	14
16.505 \pm 15	6 ⁺ ; 0	(0 ₆ ⁺)	24 \pm 4	α	6, 14
16.559 \pm 15	5 ⁻ ; 0		90 \pm 30	α	14
16.581 \pm 15	7 ⁻ ; 0	1 ⁻	92 \pm 8	α	7, 14
16.628 \pm 20	3 ⁻ ; 0		80 \pm 25	α	14
16.63 \pm 20	(7 ⁻)			α	15, 16, 18
16.667 \pm 15	4 ⁺ ; 0		100 \pm 25	α	14
16.717 \pm 15	5 ⁻ ; 0		\approx 25	α	6, 7, 14
16.732 \pm 5	0 ⁺ ; 2		2.0 \pm 0.5	γ , p, α	24, 25, 26, 28, 56
16.746 \pm 25	8 ⁺ ; 0		160 \pm 50	α	14
16.847 \pm 15	5 ⁻ ; 0		16 \pm 8	α	14
16.871 \pm 20	6 ⁺ ; 0		350 \pm 50	α	14
17.072 \pm 20	4 ⁺ ; 0		180 \pm 30	α	14
17.155 \pm 15	5 ⁻ ; 0		26 \pm 5	α	14
17.213 \pm 15	4 ⁺ ; 0		225 \pm 30	α	14
17.284 \pm 15	3 ⁻ ; 0		86 \pm 25	α	14

Table 20.13: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
17.295 \pm 15	8 ⁺ ; 0		200 \pm 25	α	4, 14, 15, 16, 18, 19
17.390 \pm 15			< 10	α	14
17.430 \pm 15	9 ⁻ ; 0	(0 ⁻)	220 \pm 25	α	6, 7, 8, 14
17.541 \pm 15	6 ⁺ ; 0		86 \pm 9	α	14
17.55 \pm 10	(2 ⁺ ; 1)		19	n, p, α	27, 28
17.606 \pm 15	5 ⁻ ; 0		140 \pm 20	α	14
17.769 \pm 20	4 ⁺ ; 0		\approx 125	p, α	14, 28
17.851 \pm 15	5 ⁻ ; 0		200 \pm 30	α	14
17.91 \pm 20	(0 ⁺)			n, p	27
18.005 \pm 15	7 ⁻ ; 0		< 10	α	14
18.024 \pm 5	5 ⁻ ; 0		34 \pm 7	α	14
18.083 \pm 25	4 ⁺ ; 0		140 \pm 60	α	14
18.125 \pm 5	7 ⁻ ; 0		29 \pm 6	α	6, 7, 8, 14
18.286 \pm 10	6 ⁺ ; 0		190 \pm 30	α	6, 14
18.430 \pm 7	2 ⁺ ; 2		9.5 \pm 3	γ , n, p, α	25, 26, 27, 28, 56
18.430 \pm 20	7 ⁻ ; 0		185 \pm 40	α	14
18.494 \pm 20	5 ⁻ ; 0		130 \pm 30	α	14
18.621 \pm 20	8 ⁺ ; 0	(0 ₆ ⁺)	185 \pm 30	α	14
18.745 \pm 25	6 ⁺ ; 0		140 \pm 50	α	14
18.768 \pm 20	7 ⁻ ; 0		140 \pm 35	α	14, 15
18.960 \pm 25	8 ⁺ ; 0		200 \pm 60	α	14
19.051 \pm 15	5 ⁻ ; 0		\approx 90	α	14
19.15 \pm 20	6 ⁺ ; 0		200 \pm 50	α	8, 14
19.284 \pm 15	6 ⁺ ; 0		140 \pm 25	α	14
19.298 \pm 25	7 ⁻ ; 0		430 \pm 60	α	14, 15
19.443 \pm 10	6 ⁺ ; 0	(0 ₇ ⁺)	130 \pm 15	α	14
19.536 \pm 25	6 ⁺ ; 0		250 \pm 60	α	14
19.655 \pm 20	6 ⁺ ; 0		140 \pm 35	α	14
19.731 \pm 20	8 ⁺ ; 0		330 \pm 60	α	14
19.845 \pm 40	6 ⁺ ; 0		360 \pm 120	α	14
19.859 \pm 10	5 ⁻ ; 0		170 \pm 25	α	14
19.884 \pm 40	7 ⁻ ; 0		\approx 120	α	14, 15
19.991 \pm 30	4 ⁺ ; 0		130 \pm 100	α	14
20.027 \pm 15	6 ⁺ ; 0		80 \pm 35	α	14
20.106 \pm 25	7 ⁻ ; 0		190 \pm 35	α	14
20.15 \pm 150			broad	γ , n	34
20.168 \pm 35	6 ⁺ ; 0		285 \pm 100	α	14
20.296 \pm 15	7 ⁻ ; 0		255 \pm 40	α	14

Table 20.13: Energy levels of ^{20}Ne ^a (continued)

E_x (MeV \pm keV)	$J^\pi; T$	K^π	τ_m ^b or $\Gamma_{c.m.}$ (keV)	Decay	Reactions
20.341 \pm 20	5 ⁻ ; 0		190 \pm 40	α	14
20.344 \pm 15	7 ⁻ ; 0		135 \pm 35	α	14
20.419 \pm 30	6 ⁺ ; 0		215 \pm 90	α	14
20.445 \pm 25	6 ⁺ ; 0		370 \pm 55	α	14
20.468 \pm 30	5 ⁻ ; 0		280 \pm 70	α	14
20.686 \pm 6	9 ⁻ ; 0	(1 ⁻)	78 \pm 11	α	7, 14, 16
20.76 \pm 30	7 ⁻ ; 0		240 \pm 50	α	14, 15
20.800 \pm 25	5 ⁻ ; 0		170 \pm 60	α	14
20.95 \pm 40	7 ⁻ ; 0		300 \pm 50	α	7, 14
21.062 \pm 6	9 ⁻ ; 0	(1 ⁻)	60 \pm 6	α	4, 7, 14, 16, 18, 19
21.3 \pm 100	7 ⁻ ; 0		300	α	14, 15
21.8 \pm 100	7 ⁻ ; 0		300	α	7, 14, 15
22.3 \pm 100	7 ⁻ ; 0		500	α	7, 14, 15
22.6 \pm 300			broad	γ, n	34
22.8 \pm 60	9 ⁻ ; 0		500	α	7, 14
22.87 \pm 40	9 ⁻ ; 0		225 \pm 40	α	4, 7, 14, 16, 18
23.4 \pm 200	8 ⁺ ; 0		500	α	14
23.70 \pm 30	(9 ⁻)		\leq 200	α	15, 16
24.21 \pm 25	8 ⁺ ; 0		350	α	14, 16
24.9 \pm 500			broad	γ, n	34
25.10 \pm 50	8 ⁺ ; 0		\approx 200	α	14, 16
25.67 \pm 50			\approx 400	α	14, 16
27.1 \pm 100	(9 ⁻)		700	α	14, 15, 18
27.5			broad	γ, n	34
28	8 ⁺ ; 0		1600	α	14
28.2 \pm 300			700	α	14

^a See also Tables 20.14 and 20.15. For other states with $E_x > 15.5$ MeV see Tables 20.30 in (1978AJ03) and 20.23, 20.24 and 20.25 here and reactions 1, 34 and 36. It is clear that there are many states with low angular momentum and with unnatural parity which have not been located at high E_x

^b See Table 20.20 in (1978AJ03).

Muon and neutrino capture and reactions: (1983RO1E, 1984EL1D, 1984GR03, 1985MI1D, 1987SU06).

Pion and kaon capture and reactions: (1982BE1M, 1982BI08, 1982LE1P, 1982NA10, 1982OL03, 1982WO1E, 1983BA22, 1983CO08, 1983GE12, 1983GM1A, 1983MA16, 1983RA1J, 1984EF03, 1984EL1D, 1985AI1C, 1985AZ1A, 1985BI01, 1985CU1F, 1985SE21, 1986AN40, 1986BA3E, 1986LE06, 1986LE22, 1986RO03, 1986TK1C, 1987GI01, 1987SC03, 1987TE01).

Antiproton interactions: (1985BA2T, 1985CU1F, 1986BA22, 1986BA3Q, 1986BA2W, 1986VA04).

Hypernuclei: (1984AS1D, 1985OS1C).

Other topics: (1978WI1B, 1980KR24, 1981BL1K, 1981CL05, 1982BR08, 1982DR1E, 1982FL04, 1982GO1U, 1982VA1E, 1982VE02, 1983AR07, 1983AR1J, 1983BI1C, 1983BR29, 1983BR1P, 1983FU1D, 1983GR26, 1983IS03, 1983LO05, 1983MA23, 1983SH32, 1983VA21, 1984AM03, 1984JA15, 1984JE02, 1984KA11, 1984LO05, 1984SC41, 1985AD1A, 1985AL21, 1985AN28, 1985IV1B, 1985KA01, 1985LI19, 1985MI10, 1985MU10, 1985ST19, 1985VE04, 1986SA02, 1986TA1A, 1987HA08, 1987PR01).

Ground state of ^{20}Ne : (1978WI1B, 1980KR24, 1982BR24, 1982FL04, 1982LO13, 1982OL03, 1982SC20, 1982ZE1A, 1983ANZQ, 1983AR07, 1983AR1J, 1983BA50, 1983BR1P, 1983DR03, 1983MA68, 1983SP04, 1983TO1L, 1983VA16, 1984AN1B, 1984BR25, 1984DR01, 1984HA14, 1984JA15, 1984KA11, 1984LO05, 1984OK04, 1984SP06, 1984WE04, 1985AN16, 1985AN28, 1985CL1A, 1985HA18, 1985HU09, 1985KR1G, 1985MI23, 1985SP06, 1986CA27, 1986LE16, 1986RO03, 1986SA02, 1986TA1X, 1987AB03).

$$Q_{1.63} = -0.27 \pm 0.03 \text{ e} \cdot \text{b} \text{ (1978GR06)}$$

$$g_{1.63} = +0.54 \pm 0.04 \text{ (1975HO15)}$$

$$B(E2) \uparrow [0 \rightarrow 1.63] = 0.0330 \pm 0.0015 \text{ e}^2 \cdot \text{b}^2 \text{ (1978GR06)}. \text{ See also (1987RA01).}$$

$$Q_{4.25} = 0.022 \pm 0.003 \text{ e}^2 \cdot \text{b}^2 \text{ (1978GR06)}$$

$$g_{4.25} = +0.13 \pm 0.15 \text{ (1986TR08)}. \text{ See also (1984BR15).}$$

Isotopic abundance: $(90.51 \pm 0.09)\%$ (1984DE53).

1. (a) $^{10}\text{B}(^{10}\text{B}, ^{10}\text{B})^{10}\text{B}$

$$E_b = 31.148$$

(b) $^{10}\text{B}(^{10}\text{B}, \alpha)^{16}\text{O}$

$$Q_m = 26.4137$$

Excitation functions have been measured for $E(^{10}\text{B}) = 6$ to 30 MeV (reaction (a)) and 6 to 20 MeV (reaction (b)). Large resonant structures are observed in reaction (b), particularly at $E_x \approx 38$ MeV (α_0) and 38.6 MeV (α to $^{16}\text{O}^*(7.0, 10.3, 16.2(\text{u}))$), $\Gamma \approx 0.6$ MeV. See also (1983KAZF) and (1978AJ03).

2. $^{10}\text{B}(^{14}\text{N}, \alpha)^{20}\text{Ne}$

$$Q_m = 19.535$$

Angular distributions of α -particles to many states of ^{20}Ne below $E_x = 10.7$ MeV have been measured at $E(^{14}\text{N}) = 23.5$ to 35 MeV. See also (1978AJ03, 1983AJ01).

3. $^{10}\text{B}(^{16}\text{O}, ^6\text{Li})^{20}\text{Ne}$ $Q_m = 0.274$

At $E(^{16}\text{O}) = 19.5$ to 42 MeV angular distributions for the ^6Li ions corresponding to transitions to $^{20}\text{Ne}^*(0, 1.63, 4.25, 4.97, 5.62 + 5.79, 6.7-7.2)$ are in good agreement with Hauser-Feshbach calculations. See also ([1978AJ03](#), [1985ST1B](#)).

4. $^{11}\text{B}(^{16}\text{O}, ^7\text{Li})^{20}\text{Ne}$ $Q_m = -3.930$

At $E(^{11}\text{B}) = 115$ MeV, angular distributions are reported to $^{20}\text{Ne}^*(7.16, 8.78, 10.26, 11.95, 15.4)$. $^{20}\text{Ne}^*(8.78, 15.4, 17.3, 21.0 \pm 0.07, 22.78 \pm 0.06)$ are particularly strongly populated. It is suggested that these five states have $J^\pi = 6^+, 7^-, (8^+), 9^-$ and 9^- : see ([1983AJ01](#)).

5. $^{12}\text{C}(^9\text{Be}, n)^{20}\text{Ne}$ $Q_m = 10.323$

At $E(^9\text{Be}) = 16$ and 24 MeV, angular distributions have been measured for $^{20}\text{Ne}^*(7.3 \pm 0.4, 9.2 \pm 0.4, 10.9 \pm 0.3, 12.2 \pm 0.3, 15.7 \pm 0.3)$: see ([1983AJ01](#)).

6. (a) $^{12}\text{C}(^{10}\text{B}, d)^{20}\text{Ne}$ $Q_m = 5.961$

(b) $^{12}\text{C}(^{11}\text{B}, t)^{20}\text{Ne}$ $Q_m = 0.764$

At $E(^{12}\text{C}) = 45$ MeV the population of states of ^{20}Ne with $E_x = 8.45, \underline{8.78}, 9.03, 10.61, 10.67, 10.99, 11.01, 11.66, \underline{11.94}, \underline{12.14}, \underline{12.39}, \underline{12.58}, 12.73, 13.05, 13.17, \underline{13.34} [7^-], \underline{13.69}, 13.91, 14.29, 14.36, 14.81, \underline{15.17} [6^+], \underline{15.38} [7^-], 15.71 [(7, 8)], \underline{15.89} [(7)], 16.16, 16.22, \underline{16.51} [(8)], 16.73, \underline{17.39} [9^-], 18.18$ and $\underline{18.32}$ MeV is reported. [Values in brackets are J^π suggested on basis of Hauser-Feshbach calculations. The underlined states are well resolved: the authors indicate ± 20 keV for such states.] The relative intensities of the groups to $^{20}\text{Ne}^*(17.39, 15.38)$ [$J^\pi = 9^-, 7^-$] argue against the existence of a superband: see ([1978AJ03](#)). See also ([1983AJ01](#)).

7. $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ $Q_m = 4.621$

Double and triple (α, α, γ) correlations and γ -ray branching measurements [see Table [20.14](#)] lead to the J^π assignments shown in Table [20.16](#). See Table [20.15](#) for assignments to rotational bands. Angular distributions have been reported at $E(^{12}\text{C}) = 4.9$ to 51 MeV [see ([1978AJ03](#), [1983AJ01](#))] and at 10.0 and 10.5 MeV ([1982DA28](#); α_0), 11.38 to 13.23 MeV ([1985BA43](#); α_0), 21 to 30 MeV ([1984DH02](#), [1983FOZV](#); α_0), 29.5 to 30.5 MeV ([1982KA17](#); many states) and 35.6 to 41.2 MeV ([1984LE19](#); many states). At $E(^{12}\text{C}) = 38$ to 64 MeV, $^{20}\text{Ne}^*(7.17, 7.83, 8.54, 8.78,$

9.03, 11.95, 12.13, 12.59, 13.90) are strongly populated and decay to $^{16}\text{O}_{\text{g.s.}}$ (1987RA02). For γ -decay measurements see (1987FI01), Table 20.16 and (1978AJ03).

The yields of various groups of α -particles and their relevance to states of ^{24}Mg , and fusion cross sections, have been studied by many groups: see (1978AJ03, 1983AJ01) for the earlier work and (1982DA28, 1982KA17, 1982SA27, 1983FOZV, 1984AR20, 1984DH02, 1984LE19, 1985BA43, 1986WU01).

Table 20.14: Radiative decays in ^{20}Ne ^a

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)
1.63	$2^+; 0$	0	100	$0.63 \pm 0.04^{\text{b}}$
4.25	$4^+; 0$	1.63	≈ 100	$7.1 \pm 0.7^{\text{b}}$
4.97	$2^-; 0$	0	0.6 ± 0.2	$(8 \pm 3) \times 10^{-4}^{\text{b}}$
		1.63	99	$0.14 \pm 0.02^{\text{b}}$
				$\delta(\text{M2/E1}) = 0.076 \pm 0.011$
				$\delta(\text{E3/E1}) = 0.043 \pm 0.016$
5.62	$3^-; 0$	0	7.6 ± 1.0	0.018 ± 0.006
		1.63	87.6 ± 1.0	0.21 ± 0.06
		4.97	4.8 ± 1.6	0.012 ± 0.005
5.79	$1^-; 0$	0	18 ± 5	0.8 ± 0.3
		1.63	82 ± 5	3.8 ± 0.8
6.73	$0^+; 0$	0		$ M ^2 = 7.4 \pm 2.0 \text{ fm}^2^{\text{d}}$
		1.63	100	33
7.00	$4^-; 0$	1.63	0.5 ± 0.2	$(7 \pm 3) \times 10^{-3}^{\text{b}}$
		4.25	63.5	0.95^{b}
		4.97	11	0.16^{b}
		5.62	25	0.37^{b}
7.16	$3^-; 0$	4.25	60 ± 5	0.97 ± 0.11
		5.79	40 ± 5	0.64 ± 0.10
7.20	$0^+; 0$	0		$\Gamma_\pi = 3.9 \times 10^{-2}$
				$6.9 \pm 1.4 \text{ fm}^2^{\text{d}}$
		1.63	100	4.35 ± 0.75
7.42	$2^+; 0$	0	$\leq 9.4 \pm 1.4$	$\leq 3.0 \pm 0.6$
		1.63	$\geq 90.6 \pm 1.4^{\text{f}}$	29 ± 4
		4.25	≤ 7.6	

Table 20.14: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)
7.83	$2^+; 0$	0	83 ± 1	57 ± 7
		1.63	17 ± 1	11.7 ± 1.6
		4.25	< 3	< 2
8.46	$5^-; 0$	5.62	100	13 ± 3
8.71	$1^-; 0$	0	87 ± 8	61 ± 16
		1.63	13 ± 8	9 ± 6
8.78	$6^+; 0$	4.25	100	100 ± 15
9.03	$4^+; 0$	1.63	100	340 ± 42
		4.25	< 2	< 6.8
9.12	$3^-; 0$	1.63	50 ± 5	13 ± 2
		4.97	33 ± 5	8.6 ± 1.7
		5.62	17 ± 4	4.4 ± 1.1
9.32 ¹	$(2^-; 0)$	1.63		
9.49	$2^+; 0$	0		$\lesssim 60$
		1.63	(100)	260 ± 100
9.87	$3^+; 0$	0	< 0.5	
		1.63	78	^g
		4.25	12 ± 3	
		4.97	≤ 5	
		5.62	≈ 7	
		7.43	≈ 3	
9.94	$(1^+); 0$	1.63	78 ± 5	
		4.97	22 ± 5	
9.99	$4^+; 0$	0		$\lesssim 70$
		1.63	(100)	900 ± 400
10.27	$2^+; 1$	0	0.65 ± 0.14	29 ± 8
		1.63	88.9 ± 0.5	4080 ± 440
		4.97	1.3 ± 0.1	60 ± 8
		5.62	2.1 ± 0.2	97 ± 14
		7.43	6.9 ± 0.4	310 ± 40

Table 20.14: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)
10.61	$6^-; 0$	7.83	0.22 ± 0.06	8 ± 2
		7.00	95.5 ± 1.2	29 ± 9^b
		8.46	4.5 ± 1.2	1.3 ± 0.4
10.69	$4^-, 3^+; 0$	4.25	25 ± 4	
		4.97	75 ± 4	
10.88	$3^+; 1$	1.63	77 ± 5	^h
		4.25	23 ± 5	
11.09 ^c	$4^+; 1$	1.63	0.5 ± 0.25	2 ± 1
		4.25 ⁱ	99.5 ± 0.25	338 ± 40
11.26 ^j	$1^+; 1$	0	84 ± 5	$(11.2 \pm 2.0) \times 10^3$
		1.63	16 ± 5	$(2.1 \pm 0.7) \times 10^3$
11.27 ^c	$1^-; 1$	0	55 ± 2	390 ± 47
		1.63	2.5 ± 1	18 ± 7
		4.97	6.5 ± 1	46 ± 9
		8.85	27 ± 1.5	189 ± 24
		9.32	9 ± 1	63 ± 10
		4.25	30 ± 3	
		4.97	70 ± 3	
11.53	$3^+, 4^-; 0$	7.00	^f	
		1.63		
		7.00		
11.555	$(3^+; 0)$	1.63		
11.558	$0^+; 0$	7.00		
		1.63	100	
11.65	$(3^+); 0$	4.25	< 8	
		1.63	14 ± 3	
11.93	$4^+; 0$	4.25	86 ± 3	
		1.63	21 ± 11	5.5 ± 3.0
11.95	$8^+; 0$	4.25	79 ± 11	20.5 ± 5.5
		8.78	100	7.7 ± 1.1
12.22 ^k	$2^+; 1$	1.63	(100)	
12.26	$3^-; 1$	1.63	63 ± 1.5	

Table 20.14: Radiative decays in ^{20}Ne ^a (continued)

E_i (MeV)	$J_i^\pi; T$	E_f (MeV)	Branch (%)	Γ_γ (meV)
12.40	$3^-; (1)$	5.62	37 ± 1.5	80
		0	≈ 1	
		1.63	≈ 29	
12.43	$0^+; 0$	4.25	≈ 70	200
		1.63	100	170 ± 50
13.48	$1^+; 1$	1.63	95	
		4.97	5	
13.88	$2^+; 1$	1.63	20	
		4.97	80	
16.73	$0^+; 2$	1.63	^e	
		5.79	^e	
		11.23	(100)	
18.43	$2^+; 2$	12.22	(100)	≈ 300

^a For earlier references see Tables 20.19 in (1978AJ03) and 20.18 in (1983AJ01). See also Tables 20.17 and 20.20 here.

^b From τ_m : see Table 20.20 in (1978AJ03) and branching ratios.

^c See also Table 20.19 in (1978AJ03).

^d Monopole matrix element.

^e See footnote (a) in Table 2 of (1976MA01).

^f $\delta(E2/M1) = -8.36_{-1.5}^{+1.0}$.

^g $\Gamma_\gamma(\text{total})/\Gamma = 0.82 \pm 0.27$.

^h $\Gamma_\gamma(\text{total})/\Gamma < 0.3$ (1977MA07). See also (1987FI01).

ⁱ $\delta = +0.01 \pm 0.06$.

^j (1983BE19): see reaction 35.

^k (1984CA08).

^l (1987FI01).

See also (1982BLZT, 1983SH1Z, 1984HU1E, 1985XI1E, 1986TR08), (1981DA13, 1982CO1X, 1983SI1L, 1984CU1B), (1986SZ02; applied), (1982BA1D, 1982SA1A, 1984FO1A, 1985AR1A, 1986TH1E; astrophysics) and (1982HO05, 1982MO1V, 1982SU06, 1983AH1A, 1984DA1B, 1986FA12; theor.).

8. (a) $^{12}\text{C}(^{14}\text{N}, ^6\text{Li})^{20}\text{Ne}$

$$Q_m = -4.176$$

Table 20.15: K^π assignments to states of ^{20}Ne ^a

K^π	J^π	E_x (MeV)	K^π	J^π	E_x (MeV)
0_1^+	0^+	0	0_7^+ ^b	6^+	(16.51)
	2^+	1.63		8^+	(18.62)
	4^+	4.25		0^+	12.43
	6^+	8.78		2^+	(12.96)
	8^+	11.95		6^+	(19.44)
0_2^+	0^+	6.73	0^-	1^-	5.79
	2^+	7.42		3^-	7.16
	4^+	9.99		5^-	10.26
	6^+	(12.59, 13.11)		7^-	13.69
0_3^+	0^+	7.20	1^-	9^-	(17.43)
	2^+	7.83		1^-	8.85
	4^+	9.03		3^-	10.41
	6^+	12.14		5^-	12.71
0_4^+	0^+	8.7	2^-	7^-	16.58
	2^+	8.8		9^-	(20.69, 21.06)
	4^+	10.80		2^-	4.97
	6^+ ^c	(12.59)		3^-	5.62
	8^+ ^c	(17.30)		4^-	7.00
0_5^+	0^+	10.97	5^-	5^-	8.46
	2^+ ^d	12.33		6^-	10.61
0_6^+ ^b	0^+	11.55	7^-	7^-	13.34
	4^+	(13.97)		8^-	(15.70) ^e
				9^-	17.43


^a See Tables 20.19, 20.20, 20.21, 20.22 and 20.23 in (1983AJ01) and (1984RI01, 1984RI07, 1985MU14, 1986MA48). I am greatly indebted to Prof. H.T. Richards for his comments on this table.

^b See also (1985LAZZ; prelim.).

^c However (1987MI07) predict the $J^\pi = 6^+, 8^+$ and 10^+ members of the 0_4^+ band to be at $E_x \approx 14 - 15$ MeV [$\Gamma \approx 1 - 2$ MeV], ≈ 21 MeV [$\Gamma \approx 2$ MeV], and ≈ 29 MeV [$\Gamma \approx 29$ MeV], suggesting that the 0_4^+ band has a moment of inertia which is very similar to that of the 0^- band.

^d For the location of higher J^π members of this band see (1984RI01).

^e See (1970PA08) and (1984RI01).



Angular distributions of the ^6Li ions to many states of ^{20}Ne below 17.5 MeV have been reported for $E(^{14}\text{N}) = 30$ to 78 MeV and $E(^{12}\text{C}) = 67.2$ MeV. At the latter energy $^{20}\text{Ne}^*(16.67, 17.38, 18.11, 19.16, 19.6)$ are particularly strongly populated: see (1978AJ03). For reaction (b) to $^{20}\text{Ne}_{\text{g.s.}}$, see (1984AR20, 1986WU01). See also (1982HO1E, 1985ST1B, 1986AR04) and ^6Li in (1988AJ01).

Table 20.16: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ ^a

E_x (MeV \pm keV) ^b	J^π ^c	Γ_γ/Γ ^d	$\Gamma_{\text{c.m.}}$ (keV)	θ_α^2 ^e
1.6329 \pm 1.0	2 ⁺			
4.2456 \pm 2.5	4 ⁺			
4.9663 \pm 2.5	2 ⁻			
5.618 \pm 4	3 ⁻			
5.774 \pm 6	1 ⁻			
6.725 \pm 6	0 ⁺			
7.004 \pm 4	4 ⁻			
7.169 \pm 6	3 ⁻			
7.196 \pm 6	0 ⁺			0.026 ^q
7.435 \pm 6	2 ⁺			
7.835 \pm 6	2 ⁺			0.015 ^q
8.449 \pm 6	5 ⁻			$(1.6 \pm 0.5) \times 10^{-3}$ ^r
8.694 \pm 6	1 ⁻			0.0027 ^q
8.779 \pm 6	6 ⁺			
8.85	1 ⁻			0.0179 ^q
9.033 \pm 6	4 ⁺			0.033 ^q , 0.022 ^r
9.110 \pm 6				
9.318 \pm 6	2 ⁻	> 0.90		
9.533 \pm 6				
9.872 \pm 6	1 ⁺ , 2 ⁻ , 3 ⁺	> 0.8		
9.948 \pm 5 ^d	1 ⁺ , 2 ⁻ , 3 ⁺	> 0.7		
10.024 \pm 6				
10.264 \pm 6	5 ⁻			
10.407 \pm 6	(3 ⁻)			0.078 ^q
10.545 \pm 6				

Table 20.16: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ ^a (continued)

E_x (MeV \pm keV) ^b	J^π ^c	Γ_γ/Γ ^d	$\Gamma_{\text{c.m.}}$ (keV)	θ_α^2 ^e
10.609 \pm 5	6 ⁻	$\equiv 1$		
10.693 \pm 5	4 ⁻ , 3 ⁺	> 0.95		
10.840 \pm 6	(3 ⁻)			0.0099 ^q
10.917 \pm 6	3 ⁺ ; $T = 0$	> 0.7		
11.013 \pm 6				
11.528 \pm 5 ^d	(3 ⁺ , 4 ⁻)	> 0.90		
11.568 \pm 10 ^d	(3 ⁺ ; $T = 0$)	0.75 \pm 0.10		
11.653 \pm 5 ^d	(3 ⁺)	> 0.90		
11.892 \pm 8 ^d		0.16 \pm 0.02		
11.949 \pm 6	8 ⁺			(7.6 \pm 2.2) $\times 10^{-3}$ ^r
12.014 \pm 10 ^d		> 0.10		
12.097 \pm 8 ^d		> 0.20		
12.135 \pm 5 ^f	6 ⁺			(4.9 \pm 2.6) $\times 10^{-4}$ ^{r,t}
12.172 \pm 8 ^d		> 0.45		
12.219 \pm 10 ^d	2 ⁺ ; $T = 1$	> 0.45		
12.379 \pm 8 ^d		0.005 \pm 0.001		
12.436 \pm 5	0 ⁺ ^s		24 \pm 1	^{r,s}
12.596 \pm 5	6 ⁺		50 \pm 10	0.09 \pm 0.02 ^r
12.730 \pm 6	(5 ⁻)			0.129 ^q
12.919 \pm 6				
13.010 \pm 6				
13.049 \pm 6				
13.190 \pm 6				
13.277 \pm 6				
13.335 \pm 6	7 ⁻			(2.4 \pm 1.0) $\times 10^{-4}$ ^{r,u}
13.441 \pm 6	(5 ⁻)			≤ 0.023 ^q
13.569 \pm 15				
13.631 \pm 15				
13.679 \pm 15				
13.845 \pm 15				
13.886 \pm 15				

Table 20.16: Excited states of ^{20}Ne from $^{12}\text{C}(^{12}\text{C}, \alpha)^{20}\text{Ne}$ ^a (continued)

E_x (MeV \pm keV) ^b	J^π ^c	Γ_γ/Γ ^d	$\Gamma_{\text{c.m.}}$ (keV)	θ_α^2 ^e
13.927 \pm 5	6 ⁺		113 \pm 7	0.10 \pm 0.01 ^r
14.144 \pm 15				
14.308 \pm 10	6 ⁺		< 50 ^r	< 0.45 ^r
14.60				
14.812 \pm 15				
15.034 \pm 15	^a			
15.159 \pm 5 ^g	6 ⁺		60 \pm 15	< 8 \times 10 ⁻⁴ ^{r,v}
15.364 \pm 14 ^h	7 ⁻		410 \pm 130	
15.438 \pm 10 ⁱ			100 \pm 20	
15.691 \pm 15				
15.874 \pm 8 ^j	8 ⁺		100 \pm 15	0.047 \pm 0.013 ^{r,w}
16.139 \pm 15				
16.600 \pm 15 ^k	7 ⁻		160 \pm 30	0.10 \pm 0.02 ^{r,x}
16.717 \pm 10			37 \pm 10	
17.259 \pm 11 ^l	7 ⁻ (9 ⁻)		162 \pm 20	0.019 \pm 0.004 ^{r,y}
18.153 \pm 10 ^m	7 ⁻			
18.538 \pm 7 ⁿ	8 ⁺		138 \pm 33	(3.2 \pm 1.5) \times 10 ⁻³ ^{r,z}
20.478 \pm 11 ^o	(8 ⁺)		250 \pm 30	0.11 \pm 0.04 ^{r,aa}
20.704 \pm 11 ^p	(9 ⁻)		\approx 120	^r
20.89 \pm 30				
21.05 \pm 20			140 \pm 50	
21.65 \pm 100	(7 ⁻ , 9 ⁻)		240 \pm 50	
22.03 \pm 70	(8 ⁺)		630 \pm 80	
22.7 \pm 70			490 \pm 110	
23.2 \pm 100			300 \pm 100	
23.74 \pm 100			230 \pm 100	
24.374 \pm 30	7 ⁻ (5 ⁻)		210 \pm 50	

- ^a For complete references see Table 20.21 in (1978AJ03). Table 20.19 in (1983AJ01) has a number of errors.
- ^b Uncertainties shown for $E_x > 5.7$ MeV are approximate, except for states flagged (d): see footnote ^c in Table 20.21 in (1978AJ03).
- ^c See discussions in (1975ME04), (1983HI06), (1984LE19) and (1987FI01). See also Table 20.14 here.
- ^d (1987FI01). $^{20}\text{Ne}^*(11.89, 12.38)$ also decay via α_2 .
- ^e See also (1984LE19).
- ^f Alpha decay is by α_2 to $^{16}\text{O}^*(6.13)$: $\Gamma'_\alpha/\Gamma = (6.0 \pm 0.15)\%$: assuming $\Gamma_\alpha \Gamma'_\alpha/\Gamma = 7.7 \pm 3.8$ eV this leads to $\Gamma_\alpha = 0.128 \pm 0.072$ keV for this 6^+ state: see (1978AJ03). (1983HI06) report an α_0 branching ratio of $(90 \pm 6)\%$.
- ^g Alpha decay is $(2 \pm 2)\%$ by α_0 , $(46 \pm 2)\%$ via α_{1+2} (mainly α_2) and $(52 \pm 2)\%$ via α_{3+4} (mainly α_3) (1979YO04).
- ^h Alpha decay is $(32 \pm 2)\%$ by α_0 , $(58 \pm 2)\%$ via α_{1+2} (mainly α_2) and $(10 \pm 2)\%$ via α_{3+4} (mainly α_3); $\Gamma_{\alpha_0}/\Gamma = 0.3 \pm 0.02$, assuming a single state. The state may correspond to a doublet (1979YO04). See also (1983HI06).
- ⁱ Alpha decay is $(20 \pm 5)\%$ by α_0 , $(57 \pm 7)\%$ by α_{1+2} and $(23 \pm 4)\%$ by α_{3+4} (1983HI06).
- ^j Alpha decay is $(9 \pm 2)\%$ by α_0 , $(79 \pm 2)\%$ via α_{1+2} (mainly α_2) and $(12 \pm 4)\%$ via α_{3+4} (mainly α_3) (1979YO04); $(24 \pm 5)\%$ via α_0 , $(51 \pm 7)\%$ via α_{1+2} , $(25 \pm 5)\%$ via α_{3+4} (1983HI06).
- ^k Alpha decay is $(72 \pm 3)\%$ via α_0 , $(20 \pm 3)\%$ via α_{1+2} (mainly α_2) and $(8 \pm 3)\%$ via α_{3+4} (mainly α_3) (1979YO04); $(60 \pm 5)\%$ via α_0 , $(20 \pm 5)\%$ via α_{1+2} and $(20 \pm 5)\%$ via α_{3+4} (1983HI06).
- ^l Alpha decay is $(15 \pm 2)\%$ via α_0 , $(50 \pm 6)\%$ via α_{1+2} and $(35 \pm 7)\%$ via α_{3+4} (1983HI06). See also (1979YO04).
- ^m Alpha decay is $(71 \pm 6)\%$ via α_0 and $(29 \pm 6)\%$ via α_{1+2} (mainly α_2) (1979YO04).
- ⁿ Alpha decay is $(1.8 \pm 0.9)\%$ via α_0 , $(60 \pm 8)\%$ via α_{1+2} and $(26 \pm 4)\%$ via α_{3+4} . Decay to $^{12}\text{C}_{g.s.} + ^8\text{Be}_{g.s.}$ is also observed: the branching ratio is $(12 \pm 1.2)\%$. This state may be a member of an excited $8p$ - $4h$ ($K^\pi = 0_6^+$) band of which $^{20}\text{Ne}^*(12.44)$ is the 0^+ band head (1983HI06).
- ^o Decay is $(66 \pm 26)\%$ via α_0 , $(14 \pm 7)\%$ via α_{1+2} , $(13.2 \pm 2.5)\%$ via $^{12}\text{C} + ^8\text{Be}$ (1983HI06).
- ^p Decay is $\lesssim 14\%$ via α_0 , $(25 \pm 15)\%$ via α_{1+2} , $(46 \pm 22)\%$ via α_{3+4} and $(4.5 \pm 0.9)\%$ via $^{12}\text{C} + ^8\text{Be}$ (1983HI06). See also (1979YO04).
- ^q (1979YO04).
- ^r θ_α^2 shown are $\theta_{\alpha_0}^2$ (1983HI06). See also (1987FI01).
- ^s See footnote ^f in Table 20.21 in (1978AJ03).
- ^t $\theta_{\alpha_2}^2 = 0.66 \pm 0.36$ (1983HI06).
- ^u $\theta_{\alpha_2}^2 = 0.025 \pm 0.010$ (1983HI06).
- ^v $\theta_{\alpha_2}^2 = 0.05 \pm 0.013$, $\theta_{\alpha_3}^2 = 0.91 \pm 0.23$ (1983HI06).
- ^w $\theta_{\alpha_2}^2 = 0.94 \pm 0.14$, $\theta_{\alpha_3}^2 = 4.2 \pm 0.9$ (1983HI06).
- ^x $\theta_{\alpha_2}^2 = 0.048 \pm 0.013$, $\theta_{\alpha_3}^2 = 0.44 \pm 0.12$ (1983HI06).
- ^y $\theta_{\alpha_2}^2 = 0.071 \pm 0.013$, $\theta_{\alpha_3}^2 = 0.32 \pm 0.08$ [all θ_α^2 assume $J^\pi = 7^-$] (1983HI06).
- ^z $\theta_{\alpha_2}^2 = 0.085 \pm 0.014$, $\theta_{\alpha_3}^2 = 0.24 \pm 0.04$, $\theta^2(^{12}\text{C}) = 1.50 \pm 0.21$ (1983HI06).
- ^{aa} $\theta_{\alpha_2}^2 = 0.016 \pm 0.008$, $\theta^2(^{12}\text{C}) = 0.24 \pm 0.05$ (1983HI06).

9. $^{13}\text{C}(^9\text{Be}, 2n)^{20}\text{Ne}$

$Q_m = 5.376$

For cross sections see (1986CU02).

$$10. \text{}^{14}\text{N}(\text{}^{12}\text{C}, \text{}^6\text{Li})\text{}^{20}\text{Ne} \quad Q_{\text{m}} = -4.176$$

See reaction 8.

$$11. \text{}^{14}\text{N}(\text{}^{14}\text{N}, 2\alpha)\text{}^{20}\text{Ne} \quad Q_{\text{m}} = 7.923$$

For yields of 1.63 MeV γ -rays see (1982DE39).

$$12. \text{}^{16}\text{O}(\alpha, \gamma)\text{}^{20}\text{Ne} \quad Q_{\text{m}} = 4.734$$

Observed resonances in the yield of capture γ -rays over the range $E_{\alpha} = 0.8$ to 10 MeV are displayed in Table 20.17. For a discussion of $^{20}\text{Ne}^*(11.28)$ [$J^{\pi} = 1^+$; $T = 1$] whose excitation is parity forbidden see (1983FI02). See also (1984BU01). Total cross sections have been measured in the range $E_{\text{c.m.}} = 1.7$ to 2.35 MeV. Assuming that S does not vary with energy over that interval, the astrophysical factor for non-resonant capture to $^{20}\text{Ne}_{\text{g.s.}}$ is 0.26 ± 0.07 MeV \cdot b. An estimate of 0.7 ± 0.3 MeV \cdot b for S at 300 keV is deduced (1987HA24). See also Table 20.17. For other papers on astrophysical considerations see (1982BA1D, 1982SA1A, 1984LA18, 1984NO1B, 1984TR1C, 1985AR1A, 1985TA1A, 1986DE27, 1986FI15, 1986KH1J). See also (1981KH1H, 1983BA50, 1983DE32, 1983LA25, 1984LA18, 1985BA2W, 1985BA1Q, 1986DE27; theor.).

$$13. \text{}^{16}\text{O}(\alpha, \text{d})\text{}^{18}\text{F} \quad Q_{\text{m}} = -16.3211 \quad E_{\text{b}} = 4.734$$

See (1986KA36).

$$14. \text{(a) } \text{}^{16}\text{O}(\alpha, \alpha)\text{}^{16}\text{O} \quad E_{\text{b}} = 4.734$$

$$\text{(b) } \text{}^{16}\text{O}(\alpha, 2\alpha)\text{}^{12}\text{C} \quad Q_{\text{m}} = -7.16195$$

Excitation functions have been measured over a wide range of energies for elastically and inelastically scattered α -particles and γ -rays from the decay of $^{16}\text{O}^*(6.13, 6.92, 7.13)$ [see (1978AJ03, 1983AJ01)] and (1986LE23; 1.8 to 4.8 MeV; α_0), (1985JA17; 2.0 to 3.6 MeV; α_0), (1983CA1F, 1985CA09; 9.2 to 13.5 MeV; α_0), (1984LA1N, 1986LAZZ; 10.8 to 18 MeV; α_1 ; prelim.) and (1979BI10, 1984RI06; 14.6 to 20.4 MeV; $\alpha_0 \rightarrow \alpha_5$). See also (1983FR14, 1985ISZU) and ^{16}O in (1986AJ04).

Table 20.17: Resonances in $^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}$ ^a

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	$\omega\gamma$ ^b (eV)	E_x (MeV \pm keV)	$J^\pi; T$
1.116 \pm 4	2.6×10^{-6} ^d	$(1.7 \pm 0.3) \times 10^{-3}$	5.627 \pm 4	3 ⁻ ; 0
1.3174 \pm 2.2 ^c	$(2.8 \pm 0.3) \times 10^{-2}$ ^d	$(1.7 \pm 0.3) \times 10^{-2}$ ^l	5.7877 \pm 3.0	1 ⁻ ; 0
2.490 \pm 8	20 ± 3 ^{d,m}	$(7.1 \pm 1.2) \times 10^{-2}$ ^m	6.726 \pm 6	0 ⁺ ; 0
3.0359 \pm 2.3 ^c	8.2 ± 0.3 ^l		7.1563 \pm 0.5	3 ⁻ ; 0
3.069	4	$(4.4 \pm 0.8) \times 10^{-3}$	7.189 \pm 3	0 ⁺ ; 0
3.359	8	0.146 \pm 0.019	7.421 \pm 1	2 ⁺ ; 0
3.868	2.4	0.343 \pm 0.035	7.828 \pm 3	2 ⁺ ; 0
(4.647 \pm 3)			(8.451 \pm 3)	(5 ⁻ ; 0)
4.969 \pm 9	2.1 ± 0.8	0.21 \pm 0.05	8.708 \pm 7	1 ⁻ ; 0
5.05	< 3	1.35 \pm 0.15	8.776 \pm 3.2	6 ⁺ ; 0
5.364	3.2	3.05 \pm 0.38	9.024 \pm 3	4 ⁺ ; 0
5.477 \pm 4	< 4	0.18 \pm 0.02	9.114 \pm 3	3 ⁻ ; 0
5.94 \pm 30	29 ± 15	1.3 \pm 0.5	9.48 \pm 24	2 ⁺ ; 0
6.61 \pm 30	155 ± 30	8 \pm 3	10.02 \pm 24	(4 ⁺); 0
6.924 \pm 7 ^k	≤ 1	19.5 \pm 1.5 ^e	10.271 \pm 7 ^f	2 ⁺ ; 1
7.948 \pm 4	< 1	30.2 \pm 3.5	11.090 \pm 3	4 ⁺ ; 1
8.180 \pm 5 ^g	< 1	2.06 \pm 0.25 ^h	11.276 \pm 4	1 ⁻ ; 1
8.535 \pm 6	1.3 ± 0.8	0.41 \pm 0.05	11.559 \pm 6	0 ⁺ ; 0 ^j
8.994 \pm 8	< 1	0.23 \pm 0.05 ⁱ	11.926 \pm 6	4 ⁺ ; 0
9.02		0.131 \pm 0.018	11.950 \pm 4	8 ⁺ ; 0
(9.05 \pm 50)	< 40		(11.97)	
(9.15 \pm 50)	< 40		(12.05)	
9.362 \pm 5	< 1	1.41 \pm 0.23	12.221 \pm 4	2 ⁺ ; 1
9.406 \pm 4	< 1	6.6 \pm 0.8 ^g	12.256 \pm 3	3 ⁻ ; 1
9.57 \pm 10	33 ± 4	1.94 \pm 0.15	12.39	3 ⁻ ; (1)
9.70 \pm 30	≤ 10	0.17 \pm 0.05	12.49	

- ^a For complete references see Tables 20.22 in (1978AJ03) and 20.20 in (1983AJ01). See also Table 20.18 here.
- ^b $\omega\gamma = (2J + 1)\Gamma_\alpha\Gamma_\gamma/\Gamma$.
- ^c The strength of the γ -decay of $^{20}\text{Ne}^*(7.16)$ to $^{20}\text{Ne}^*(5.79)$ (see Table 20.14) is strong evidence that these two states are members of the $K^\pi = 0^-$ band.
- ^d This is also Γ_α .
- ^e Other values are $\omega\gamma = 19.2 \pm 1.9$ eV; $\Gamma_\alpha = 116 \pm 20$ eV; $\Gamma_\gamma = 4.26 \pm 0.23$ eV: see (1983AJ01).
- ^f The measurements of the decay of this state lead to $E_x = 4247.9 \pm 1.3, 4966.0 \pm 1.9, 5621.0 \pm 3.5, 7423.1 \pm 3.0, 7828.1 \pm 3.8$ and 8776.7 ± 2.3 keV.
- ^g See also Table 20.20 in (1983AJ01).
- ^h The γ -decay is partly (see Table 20.14) to a state at $E_x = 9318 \pm 2$ keV. The strength of this transition and the subsequent decay to $^{20}\text{Ne}^*(1.63)$ (and not to the ground state) favor 2^- for $^{20}\text{Ne}^*(9.32)$. The other M1 transition [1.27 \rightarrow 8.85] is also strong suggesting similar structures for $^{20}\text{Ne}^*(8.85, 9.32)$ (1980FI01).
- ⁱ Also observed as a resonance in the yield of 6.13 MeV γ -rays with $(2J + 1)\Gamma_{\alpha_0}\Gamma_{\alpha_2}/\Gamma = 5.2 \pm 0.9$ eV (1980FI01).
- ^j From (α, α_0) : see (1984RI07).
- ^k See also (1984RO04).
- ^l Best value including the recent work by (1987HA24).
- ^m (1987HA24).

A number of anomalies are observed: see Table 20.18. K^π parameter assignments derived from this and other work are displayed in Table 20.15 (1984RI07). See also (1983MI22). For reaction (b) see ^{12}C in (1985AJ01). For cross sections of relevance to astrophysics for both reactions see (1985DY05; $E_\alpha = 10$ to 26 MeV). For other channels see (1983AJ01). See also (1981BU27, 1983SA07, 1984BUZO, 1984SA28), (1984RE14), (1982WA1K, 1985ME1N; applications) and (1981SH1A, 1982AO06, 1982LE23, 1982SC16, 1983AO03, 1983BI1F, 1983BR1V, 1983CS02, 1983HO1D, 1983HO1F, 1983MA29, 1983MA68, 1983SM1B, 1983WA04, 1984BY02, 1984GO04, 1984OK04, 1984PR09, 1984SE20, 1984SU02, 1985MI11, 1985WA08, 1986ALZZ, 1986CEZW, 1986DE31, 1986HO33, 1986MA35, 1986SH07, 1986SU06, 1987HO1B, 1987PR01, 1987WA1B; theor.).

$$15. \ ^{16}\text{O}(^6\text{Li}, d)^{20}\text{Ne} \quad Q_m = 3.259$$

Deuteron groups have been observed to many states of ^{20}Ne : see Table 20.19. Angular distributions have been measured for $E(^6\text{Li}) = 5.5$ to 75.4 MeV: see (1978AJ03, 1983AJ01). See also (1984MO08). Angular correlations [(d, α_0) to $^{16}\text{O}_{\text{g.s.}}$] have been measured at $E(^6\text{Li}) = 60, 75$ and 95 MeV (1982AR20). See also (1986BE1D) and (1980GR1L, 1982ES1A, 1982GY02, 1983COZV, 1983OS03, 1983SU11, 1984GR29, 1986GR1R; theor.).

$$16. \ ^{16}\text{O}(^7\text{Li}, t)^{20}\text{Ne} \quad Q_m = 2.266$$

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
1.3174 ± 2.2	$(2.8 \pm 0.3) \times 10^{-2}$ ^b	α_0			5.7877 ± 2.6	1^-
2.522 ± 2.5 ^c	19.0 ± 0.9	α_0		22	6.751 ± 3	0^+
3.0382 ± 2.0 ^{a,c}	8.1 ± 0.3 ^b	α_0		36	7.164	3^-
3.082 ± 3.1 ^c	3.4 ± 0.2 ^c	α_0		1.1	7.199 ± 3	0^+
3.372 ± 3.4 ^c	15.1 ± 0.7 ^c	α_0		4.7	7.431 ± 3	2^+
3.885 ± 10	2	α_0		0.6	7.841 ± 8	2^+
4.653 ± 5	0.013 ± 0.004	α_0		0.07	8.455 ± 5	5^-
≈ 4.9	> 800	α_0		≈ 70	≈ 8.7	0^+
5.002	2.5	α_0		0.23	8.734	1^-
5.058 ± 3	0.11 ± 0.02	α_0		8.5 ± 1.5	8.779 ± 3	6^+
≈ 5.1	> 800	α_0		≈ 95	≈ 8.8	2^+
5.11	< 1	α_0			8.82	(5^-)
5.152 ± 5	19	α_0		1.1	8.854 ± 5	1^-
5.395 ± 5	3	α_0		3.9	9.049 ± 5	4^+
5.486 ± 5	3.2	α_0		0.49	9.121 ± 5	3^-
5.955 ± 10	24	α_0		1.4	9.496 ± 8	2^+
6.569 ± 10	97	α_0		17	9.987 ± 8	4^+
6.912 ± 5	141	α_0		66	10.262 ± 5	5^-
6.92 ± 10	≤ 0.3	α_0		$\leq 1.3 \times 10^{-3}$	10.27 ± 10	(2^+)
7.092 ± 5	81	α_0		4.8	10.406 ± 5	3^-
7.276 ± 5	16	α_0		1.8	10.553 ± 5	4^+
7.314 ± 10	24	α_0		0.85	10.583 ± 8	2^+

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
7.580 \pm 100	349	α_0		33	10.80 \pm 75	4 ⁺
7.635 \pm 5	13	α_0		0.42	10.840 \pm 5	2 ⁺
7.636	45	α_0		2.1	10.841	3 ⁻
(7.75)	80	α_0			(10.93)	
7.80 \pm 150	576	α_0		14	10.97 \pm 113	0 ⁺
7.860 \pm 10	24	α_0		2.0	11.020 \pm 8	4 ⁺
7.93 \pm 10	≤ 0.5	α_0		≤ 0.05	11.08 \pm 10	(4 ⁺)
8.132 \pm 30	172	α_0		4.2	11.24 \pm 23	1 ⁻
8.16 \pm 10	≤ 0.3	α_0		≤ 0.009	11.26 \pm 10	(1 ⁻)
8.24 \pm 10	40 \pm 10	α_0		1.4	11.32 \pm 10	2 ⁺
8.528 \pm 10	1.0 \pm 0.5	α_0		0.03	11.551 \pm 8	0 ⁺ ⁱ
(\approx 8.6)	\approx 500	α_0			(\approx 11.6)	(2 ⁺)
8.930 \pm 20	46	α_0		1.1	11.875 \pm 15	2 ⁺
8.997 \pm 5	0.44 \pm 0.15	$\alpha_0, \gamma_{6.13}$		0.04 \pm 0.01	11.929 \pm 5	4 ⁺
9.026 \pm 5	(35 \pm 10) $\times 10^{-3}$	α_0		1.0 \pm 0.3	11.952 \pm 5	8 ⁺
9.043 \pm 10	30 \pm 5	α_0		0.72	11.966 \pm 8	1 ⁻
9.25 ^d		$\alpha_0, \gamma_{6.13}$		^e	12.137 \pm 5	6 ⁺
9.403 \pm 9	155 \pm 15	α_0	0.89 \pm 0.05	6.8	12.253 \pm 10	4 ⁺
9.406 \pm 4 ^f	< 1	$\gamma_{6.13}$		^e	12.256 \pm 4	3 ⁻ ; 1
9.495 \pm 13	390 \pm 50	α_0	0.92 \pm 0.04	8	12.327 \pm 10	2 ⁺
9.587 \pm 2	37.3 \pm 0.9	$\alpha_0, \gamma_{6.13}$	1.00 \pm 0.04	1.2	12.401 \pm 5	3 ⁻
9.628 \pm 5	24.4 \pm 0.5	α_0, α_1	0.62 \pm 0.15	0.3	12.433 \pm 5	0 ⁺

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
9.677 \pm 8	124 \pm 6	α_0	0.88 \pm 0.05	2.4	12.472 \pm 10	(2 ⁺)
9.818 \pm 6	72 \pm 9	α_0	0.68 \pm 0.05	14	12.585 \pm 5	6 ⁺
9.827 \pm 14	145 \pm 25	α_0	0.78 \pm 0.09	2.5	12.592 \pm 15	(2 ⁺)
9.978 \pm 6	84 \pm 8	α_0	1.00 \pm 0.05	7.3	12.713 \pm 5	5 ⁻
10.015 \pm 7	61 \pm 12	α_0	0.72 \pm 0.09	0.9	12.743 \pm 10	(2 ⁺)
10.132 \pm 2	30 \pm 5	$\alpha_0, \gamma_{6.13}$	0.83 \pm 0.09	0.45	12.836 \pm 5	1 ⁻
(10.27)	(580)	(α_0)	(0.92)	(21)	(12.95)	(4 ⁺)
10.283 \pm 2	38 \pm 4	$\alpha_0, \gamma_{6.13}$	1.00 \pm 0.08	0.8	12.957 \pm 5	2 ⁺
10.397 \pm 1	18 \pm 3	$\alpha_0, \gamma_{6.13}$	0.55 \pm 0.05	0.4	13.048 \pm 5	4 ⁺
(10.419 \pm 15)	(305 \pm 55)	(α_0)	(0.42 \pm 0.03)	(3.2)	(13.066 \pm 15)	(3 ⁻ , 5 ⁻)
10.461 \pm 12	53 \pm 24	α_0	0.22 \pm 0.07	0.5	13.099 \pm 10	(0 ⁺)
10.468 \pm 5	102 \pm 5	α_0	0.52 \pm 0.04	11	13.105 \pm 5	6 ⁺
10.508 \pm 2	48 \pm 4	α_0	1.00 \pm 0.05	1.2	13.137 \pm 5	3 ⁻
10.614 \pm 7	40 \pm 13	α_0	0.55 \pm 0.13	0.4	13.222 \pm 10	0 ⁺
10.617 \pm 19	\approx 80	α_0	0.22 \pm 0.07	0.3	13.224 \pm 15	1 ⁻
10.620 \pm 2	53 \pm 4	α_0	1.00 \pm 0.04	1.3	13.226 \pm 5	3 ⁻
10.759 \pm 6 ^f	(8 \pm 3) \times 10 ⁻²	α_0		0.08 \pm 0.03	13.338 \pm 5	7 ⁻
10.763 \pm 1	26 \pm 3	$\alpha_0, \gamma_{6.13}$	0.70 \pm 0.05	0.6	13.341 \pm 5	4 ⁺
10.854 \pm 3	34 \pm 5	$\alpha_0, \gamma_{6.13}$	0.46 \pm 0.05	0.4	13.414 \pm 5	3 ⁻
10.857 \pm 4	\approx 16	α_0	0.16 \pm 0.06	0.06	13.416 \pm 5	(3 ⁻)
10.870 \pm 4	49 \pm 7	α_0	0.38 \pm 0.04		13.426 \pm 5	(5 ⁻)
10.913 \pm 8	195 \pm 25	α_0	0.99 \pm 0.05	3.2	13.461 \pm 10	1 ⁻

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
10.971 \pm 4	24 \pm 8	α_0	0.36 \pm 0.07	0.15	13.507 \pm 5	1 ⁻
10.999 \pm 4	61 \pm 8	α_0	0.72 \pm 0.05	0.8	13.529 \pm 5	2 ⁺
11.000 \pm 15	76 \pm 32	α_0	0.52 \pm 0.13	0.6	13.530 \pm 15	(0 ⁺)
11.054 \pm 3	12 \pm 5	α_0	0.19 \pm 0.06	0.04	13.573 \pm 5	2 ⁺
11.183 \pm 1	11 \pm 2	α_0	0.33 \pm 0.05	0.2	13.677 \pm 5	5 ⁻
11.202 \pm 12	310 \pm 30	$\alpha_0, \gamma_{6.13}$	0.51 \pm 0.03	84	13.692 \pm 10	7 ⁻
11.267 \pm 26	\approx 80	α_0	0.33 \pm 0.12	0.4	13.744 \pm 20	0 ⁺
11.371 \pm 9	136 \pm 15	α_0	0.73 \pm 0.04	2.1	13.827 \pm 10	3 ⁻
11.420 \pm 34	\approx 175	α_0	0.21 \pm 0.06	0.6	13.866 \pm 30	1 ⁻
11.473 \pm 5	74 \pm 10	α_0	0.75 \pm 0.06	1.0	13.908 \pm 5	2 ⁺
11.498 \pm 5	65 \pm 3	α_0	0.86 \pm 0.04	6.9	13.928 \pm 5	6 ⁺
11.522 \pm 7	79 \pm 15	α_0	1.0 \pm 0.1	1.3	13.948 \pm 10	0 ⁺
11.544 \pm 2	8.1 \pm 1	α_0	0.46 \pm 0.05	0.11	13.965 \pm 5	4 ⁺
(11.607 \pm 19)	(\approx 80)	(α_0)	(0.19 \pm 0.05)	(0.25)	(14.015 \pm 15)	(1 ⁻)
(11.663 \pm 19)	(150 \pm 50)	(α_0)	(0.24 \pm 0.05)	(0.6)	(14.060 \pm 15)	(2 ⁺)
11.732 \pm 4	42 \pm 6	$\alpha_0, \gamma_{6.9+7.1}$	0.71 \pm 0.06	0.5	14.115 \pm 5	2 ⁺
11.925 \pm 7	92 \pm 9	α_0	0.64 \pm 0.04	1.6	14.270 \pm 10	4 ⁺
11.968 \pm 8	60 \pm 13	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$	0.31 \pm 0.05	1.9	14.304 \pm 10	(6 ⁺)
11.977 \pm 6	117 \pm 8	α_0	0.82 \pm 0.04	9.6	14.311 \pm 5	6 ⁺
11.979 \pm 15	\approx 45	α_0	0.13 \pm 0.06	0.1	14.313 \pm 15	(3 ⁻)
12.148 \pm 28	\approx 95	α_0	0.18 \pm 0.06 ^e	0.3	14.448 \pm 25	(0 ⁺ , 2 ⁺)
12.156 \pm 4	\approx 15	α_0	0.09 \pm 0.04	0.05	14.454 \pm 5	5 ⁻

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
12.322 \pm 25	140 \pm 50	α_0	0.45 \pm 0.08	0.9	14.587 \pm 20	1 ⁻
12.329 \pm 13	260 \pm 25	$\alpha_0, \gamma_{6.9+7.1}$	0.79 \pm 0.04	5.3	14.593 \pm 10	4 ⁺
12.447 \pm 11	90 \pm 30	α_0	0.35 \pm 0.06	0.6	14.687 \pm 10	(3 ⁻)
12.502 \pm 10	60 \pm 25	α_0	0.25 \pm 0.06	0.4	14.731 \pm 10	(4 ⁺)
12.539 \pm 2	7.3 \pm 4.8	α_0	0.18 \pm 0.05	0.1	14.761 \pm 5	6 ⁺
12.597 \pm 4	86 \pm 7	α_0	0.95 \pm 0.04	6.5	14.807 \pm 5	6 ⁺
12.608 \pm 5	117 \pm 13	α_0	0.69 \pm 0.04	3.1	14.816 \pm 5	5 ⁻
12.637 \pm 8	79 \pm 15	α_0	0.45 \pm 0.05	0.9	14.839 \pm 10	(4 ⁻)
12.699 \pm 12	100 \pm 30	α_0	0.44 \pm 0.06	0.7	14.888 \pm 10	2 ⁺
12.897 \pm 10	66 \pm 20	α_0	0.31 \pm 0.06	0.3	15.047 \pm 10	2 ⁺
12.930 \pm 12	160 \pm 25	α_0	0.40 \pm 0.04	2.3	15.073 \pm 10	5 ⁻
13.016 \pm 20	\approx 60	α_0	\approx 0.12	0.11	15.142 \pm 15	(2 ⁺)
13.056 \pm 10	230 \pm 25	α_0	0.70 \pm 0.04	5.5	15.174 \pm 10	5 ⁻
(13.238 \pm 10)	(130 \pm 20)	(α_0)	(0.99 \pm 0.08)		(15.319 \pm 10)	(1 ⁻)
(13.266 \pm 12)	(50 \pm 25)	(α_0)	(0.69 \pm 0.17)		(15.342 \pm 10)	(0 ⁺)
13.237 \pm 29	280 \pm 40	α_0	0.39 \pm 0.04	20	15.319 \pm 25	7 ⁻
13.251 \pm 6	34 \pm 10	α_0	0.29 \pm 0.05	0.2	15.330 \pm 5	4 ⁺
13.296 \pm 5	110 \pm 10	$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$	0.71 \pm 0.04	14	15.366 \pm 5	7 ⁻
13.384 \pm 15 ^d	85 \pm 35	α_0	0.26 \pm 0.05	0.4	15.436 \pm 15	(3 ⁻)
13.58		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$			15.59	
13.73		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$			15.71	(6 ⁺)
14.05		$\alpha_0, \gamma_{6.13}, \gamma_{6.9+7.1}$			15.97	(6 ⁺)

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
14.26		$\gamma_{6.13}, \gamma_{6.9+7.1}$			16.14	
14.40		$\gamma_{6.13}$			16.25	
14.501 \pm 15	45	α_0, α_{1+2}			16.329 \pm 11	4 ⁺
14.636 \pm 15 ^g	35	$\alpha_0, \alpha_{1+2}, \alpha_3$			16.437 \pm 11	(0, 2, 4) ⁺
14.721 \pm 15	24 \pm 4	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.36 \pm 0.03	0.38 \pm 0.07	16.505 \pm 15	6 ⁺
14.789 \pm 18	90 \pm 30	α_0	0.16 \pm 0.03	0.37 \pm 0.13	16.559 \pm 15	5 ⁻
14.816 \pm 15	92 \pm 8	α_0, α_3	0.45 \pm 0.03	4.1 \pm 0.5	16.581 \pm 15	7 ⁻
14.875 \pm 22	80 \pm 25	α_0	0.18 \pm 0.04	0.22 \pm 0.08	16.628 \pm 20	3 ⁻
14.924 \pm 20	100 \pm 25	$\alpha_0, (\alpha_3)$	0.23 \pm 0.03	0.42 \pm 0.11	16.667 \pm 15	4 ⁺
14.987 \pm 18	\approx 25	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.08 \pm 0.03	\approx 0.05	16.717 \pm 15	5 ⁻
15.023 \pm 33	160 \pm 50	α_0	0.10 \pm 0.02	4.8 \pm 1.9	16.746 \pm 25	8 ⁺
15.149 \pm 16	16 \pm 8	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.11 \pm 0.02	0.04 \pm 0.02	16.847 \pm 15	5 ⁻
15.179 \pm 25	350 \pm 50	α_0	0.28 \pm 0.03	3.9 \pm 0.7	16.871 \pm 20	6 ⁺
15.430 \pm 21	180 \pm 30	α_0	0.32 \pm 0.03	1.0 \pm 0.2	17.072 \pm 20	4 ⁺
15.535 \pm 15	26 \pm 5	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.22 \pm 0.02	0.13 \pm 0.03	17.155 \pm 15	5 ⁻
15.607 \pm 19	225 \pm 30	α_0	0.32 \pm 0.02	1.2 \pm 0.2	17.213 \pm 15	4 ⁺
15.696 \pm 20	86 \pm 25	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.16 \pm 0.03	0.20 \pm 0.07	17.284 \pm 15	3 ⁻
15.710 \pm 17	200 \pm 25	α_0	0.26 \pm 0.02	11.6 \pm 1.4	17.295 \pm 15	8 ⁺
15.828 \pm 15 ^f	< 10	α_{1+2}			17.390 \pm 15	
15.878 \pm 18	220 \pm 25	α_0	0.24 \pm 0.01	48 \pm 6	17.430 \pm 15	9 ⁻
16.017 \pm 16	86 \pm 9	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.45 \pm 0.03	1.3 \pm 0.2	17.541 \pm 15	6 ⁺
16.099 \pm 17	140 \pm 20	α_0, α_4	0.36 \pm 0.03	1.05 \pm 0.15	17.606 \pm 15	5 ⁻

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
16.302 \pm 23	\approx 125	α_0	0.13 \pm 0.03	\approx 0.3	17.769 \pm 20	4 ⁺
16.405 \pm 17	200 \pm 30	α_0	0.38 \pm 0.03	1.6 \pm 0.3	17.851 \pm 15	5 ⁻
16.598 \pm 15 ^f	< 10	α_0, α_{1+2}			18.005 \pm 15	7 ⁻
16.622 \pm 6	34 \pm 7	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.34 \pm 0.04	0.23 \pm 0.06	18.024 \pm 5	5 ⁻
16.695 \pm 30	140 \pm 60	α_0	0.20 \pm 0.05	0.4 \pm 0.2	18.083 \pm 25	4 ⁺
16.748 \pm 6	29 \pm 6	$\alpha_0, \alpha_{1+2}, \alpha_3, \alpha_4$	0.46 \pm 0.06	0.8 \pm 0.2	18.125 \pm 5	7 ⁻
16.949 \pm 13	190 \pm 30	α_0, α_4	0.32 \pm 0.02	1.7 \pm 0.3	18.286 \pm 10	6 ⁺
17.129 \pm 24	185 \pm 40	$\alpha_0, (\alpha_{1+2}), \alpha_3, \alpha_4$	0.19 \pm 0.02	1.8 \pm 0.4	18.430 \pm 20	7 ⁻
17.210 \pm 21	130 \pm 30	α_0, α_3	0.21 \pm 0.03	0.5	18.494 \pm 20	5 ⁻
17.368 \pm 23	185 \pm 30	α_0, α_4	0.24 \pm 0.03	5.5 \pm 1.1	18.621 \pm 20	8 ⁺
17.524 \pm 29	140 \pm 50	α_0, α_{1+2}	0.17 \pm 0.04	0.6 \pm 0.3	18.745 \pm 25	6 ⁺
17.552 \pm 24	140 \pm 35	α_0	0.22 \pm 0.03	1.5 \pm 0.4	18.768 \pm 20	7 ⁻
17.793 \pm 29	200 \pm 60	α_0	0.15 \pm 0.02	3.2 \pm 1.1	18.960 \pm 25	8 ⁺
17.906 \pm 18	\approx 90	α_0, α_{1+2}	0.18 \pm 0.03	\approx 0.3	19.051 \pm 15	5 ⁻
18.03 \pm 20	200 \pm 50	$\alpha_0, \alpha_1, (\alpha_2), \alpha_4, \alpha_5$	0.38 \pm 0.04 ^d	\approx 2	19.15 \pm 20	6 ⁺
18.198 \pm 17	140 \pm 25	$\alpha_1, (\alpha_5)$	0.12 \pm 0.02 ^h		19.284 \pm 15	6 ⁺
18.216 \pm 30	430 \pm 60	α_0	0.36 \pm 0.03	6.4 \pm 1.1	19.298 \pm 25	7 ⁻
18.397 \pm 11	130 \pm 15	$\alpha_0, \alpha_3, \alpha_4$	0.38 \pm 0.01 ^h		19.443 \pm 10	6 ⁺
18.514 \pm 29	250 \pm 60	$\alpha_0, \alpha_2, \alpha_3$	0.27 \pm 0.04	1.6 \pm 0.4	19.536 \pm 25	6 ⁺
(18.563 \pm 25)	(140 \pm 50)	(α_1)	(0.09 \pm 0.02) ^h		(19.576 \pm 20)	(7 ⁻)
18.662 \pm 23	140 \pm 35	α_1	0.14 \pm 0.02 ^h		19.655 \pm 20	6 ⁺
18.757 \pm 28	330 \pm 60	$\alpha_0, (\alpha_2), \alpha_3$	0.23 \pm 0.02	6.3 \pm 1.2	19.731 \pm 20	8 ⁺

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
18.900 \pm 48	360 \pm 120	α_0	0.18 \pm 0.03	1.4 \pm 0.5	19.845 \pm 40	6 ⁺
18.918 \pm 11	170 \pm 25	α_1	0.26 \pm 0.02 ^h		19.859 \pm 10	5 ⁻
18.949 \pm 52	\approx 120	α_0	0.08 \pm 0.03	\approx 0.35	19.884 \pm 40	7 ⁻
19.083 \pm 39	130 \pm 100	$\alpha_0, \alpha_2, (\alpha_5)$	0.11 \pm 0.04	0.19 \pm 0.04	19.991 \pm 30	4 ⁺
19.128 \pm 16	80 \pm 35	α_1, α_4	0.10 \pm 0.04 ^h		20.027 \pm 15	6 ⁺
19.227 \pm 28	190 \pm 35	α_1	0.29 \pm 0.03 ^h		20.106 \pm 25	7 ⁻
19.304 \pm 47	285 \pm 100	α_0, α_3	0.18 \pm 0.04	1.1 \pm 0.4	20.168 \pm 35	6 ⁺
19.464 \pm 19	255 \pm 40	α_1, α_5	0.28 \pm 0.03 ^h		20.296 \pm 15	7 ⁻
19.521 \pm 22	190 \pm 40	α_1	0.26 \pm 0.03 ^h		20.341 \pm 20	5 ⁻
19.524 \pm 16	135 \pm 35	α_0, α_3	0.25 \pm 0.04	1.1 \pm 0.3	20.344 \pm 15	7 ⁻
19.618 \pm 39	215 \pm 90	α_0	0.14 \pm 0.03	0.6 \pm 0.3	20.419 \pm 30	6 ⁺
19.651 \pm 32	370 \pm 55	α_1	0.32 \pm 0.03 ^h		20.445 \pm 25	6 ⁺
19.679 \pm 35	280 \pm 70	α_0, α_2	0.20 \pm 0.03	0.86 \pm 0.25	20.468 \pm 30	5 ⁻
19.952 \pm 8	78 \pm 11	$\alpha_0, \alpha_1, \alpha_2, \alpha_3$	0.33 \pm 0.03 ^j	4.5 \pm 0.8	20.686 \pm 6	9 ⁻
20.04	240 \pm 50	$\alpha_0, \alpha_1, \alpha_4$	0.2 ^j	1.8 \pm 0.5	20.76 \pm 30	7 ⁻
20.095 \pm 32	170 \pm 60	α_1	0.11 \pm 0.02 ^h		20.800 \pm 25	5 ⁻
20.28	300 \pm 50	α_0, α_1	0.23 \pm 0.03 ^j	2.1 \pm 0.6	20.95 \pm 40	7 ⁻
20.423 \pm 8 ^g	60 \pm 6	α_0, α_3	0.46 \pm 0.03	4.1 \pm 0.5	21.062 \pm 6	9 ⁻
20.7	300	α_0			21.3	7 ⁻
21.3 \pm 200	300	α_0			21.8 \pm 150	7 ⁻
22.0 \pm 200	500	α_0			22.3 \pm 150	7 ⁻
22.5 \pm 250	500	α_0			22.7 \pm 200	9 ⁻

Table 20.18: Resonances in $^{16}\text{O}(\alpha, \alpha)$ ^a (continued)

E_α (MeV \pm keV)	$\Gamma_{\text{c.m.}}$ (keV)	Outgoing particles	Γ_{α_0}/Γ	θ^2 (%)	E_x (MeV \pm keV)	J^π
22.65 \pm 125	250	α_0			22.84 \pm 100	9 ⁻
23.3 \pm 250	500	α_0			23.4 \pm 200	8 ⁺
24.24 \pm 150	350	α_0			24.11 \pm 100	8 ⁺
25.4 \pm 300	600	α_0			25.0 \pm 250	8 ⁺
26.2 \pm 200	400	α_0			25.7 \pm 150	
28.1 \pm 350	700	α_0			27.2 \pm 300	
29	1600	α_0			28	8 ⁺
29.4 \pm 350	700	α_0			28.2 \pm 300	

59

^a For earlier references see Tables 20.23 in (1978AJ03) and 20.21 in (1983AJ01). For K^π assignments see Table 20.15 here. The uncertainties in the excitation energies are calculated by taking the uncertainty in the E_α in the c.m. [$\frac{3}{4} \times$ uncertainty in the lab] and adding the uncertainty in E_b [2 keV], in quadrature, rounding upwards. I am indebted to Prof. H.T. Richards for many very useful comments.

^b $\Gamma_{\text{c.m.}} = \Gamma_\alpha$.

^c (1985JA17).

^d Resonances with $9.25 \leq E_\alpha \leq 13.39$ MeV are from (1985CA09), except for the states labelled ^f. Certain values are rounded upwards. See also (1983CA1F) and Table 20.21 in (1983AJ01).

^e $(2J+1)\Gamma_{\alpha_0}\Gamma_{\alpha_2}/\Gamma_g = 81 \pm 12$ eV and 14 ± 2 eV, respectively, for $^{20}\text{N}^*(12.14, 12.25)$ [for the latter see Table 20.17] (1980FI01).

^f See Table 20.21 in (1983AJ01).

^g Resonances with $14.6 < E_\alpha < 20.4$ MeV are from the re-analysis of the data of (1979BI10) by (1984RI06). Certain values are rounded upwards.

^h $(\Gamma_{\alpha_0}\Gamma_{\alpha_1})^{1/2}/\Gamma$.

ⁱ (1984RI07).

^j For information on the α_1 strength see (1984RI06).

Table 20.19: States of ^{20}Ne from $^{16}\text{O}(^6\text{Li}, \text{d})$, $^{16}\text{O}(^7\text{Li}, \text{t})$ and $^{16}\text{O}(^{12}\text{C}, ^8\text{Be})$
^a

E_x (MeV \pm keV)			$\Gamma_{\text{c.m.}}$ (keV)	Γ_{α_0}/Γ	S^b	J^π
($^6\text{Li}, \text{d}$)	($^7\text{Li}, \text{t}$)	($^{12}\text{C}, ^8\text{Be}$)				
0	0	0			1.00	0 ⁺
1.63	1.63	1.63			0.41	2 ⁺
4.25	4.25	4.25			0.22	4 ⁺
4.97						2 ⁻
5.62					0.06	3 ⁻
5.79	5.79	5.79			0.54	1 ⁻
6.73					0.56	0 ⁺
7.00						4 ⁻
7.16	7.16	7.16			0.26	3 ⁻
7.43					0.13	2 ⁺
8.46					0.04	5 ⁻
8.78	8.78	8.78			0.20	6 ⁺
10.3 \pm 100	10.26	10.26	145 \pm 40	1	0.15	5 ⁻
10.7 \pm 100						4 ⁺
11.95	11.95	11.95		0.85 \pm 0.15	0.51	8 ⁺
12.14					0.05	6 ⁺
12.6 \pm 100	12.591 \pm 10	12.59	110 \pm 40	0.80 \pm 0.10		6 ⁺
13.9	13.904 \pm 20		\approx 100			6 ⁺
14.3	14.310 \pm 20	14.3 ^d	< 100			6 ⁺
15.35 \pm 100	15.336 \pm 15	15.34	380 \pm 60	0.90 \pm 0.10		7 ⁻
15.9 \pm 100		15.87	< 250			7 ⁻
16.7 \pm 100	16.63 \pm 20	16.63	190 \pm 40	0.90 \pm 0.10		7 ^{-e}
17.35 \pm 100	17.30 \pm 20	17.30	220 \pm 40	\geq 0.40 \pm 0.10		8 ^{+e}
18.7 \pm 100						7 ⁻
19.4 \pm 100			400			7 ⁻
19.9 \pm 100			400			7 ⁻
	20.67 \pm 40	20.5 ^d				
20.8 \pm 100						7 ^{-(6⁺)}
	21.08 \pm 30	21.08	100 \pm 50	0.65 \pm 0.15		9 ⁻

Table 20.19: States of ^{20}Ne from $^{16}\text{O}(^6\text{Li}, \text{d})$, $^{16}\text{O}(^7\text{Li}, \text{t})$ and $^{16}\text{O}(^{12}\text{C}, ^8\text{Be})$
^a (continued)

E_x (MeV \pm keV)			$\Gamma_{\text{c.m.}}$ (keV)	Γ_{α_0}/Γ	S^{b}	J^π
$(^6\text{Li}, \text{d})$	$(^7\text{Li}, \text{t})$	$(^{12}\text{C}, ^8\text{Be})$				
21.3 \pm 100			300			8 ⁺
21.8 \pm 100			300			8 ⁺
22.3 \pm 100			300			8 ⁺
	22.87 \pm 40	22.87	225 \pm 40	0.90 \pm 0.10		9 ⁻
23.5 \pm 100	23.70 \pm 30		\leq 200			9 ⁻ (8 ⁺)
	24.21 \pm 25		\approx 500			
	25.10 \pm 50		\leq 200			
	25.67 \pm 50		\approx 500			
27.1 \pm 100 ^c		27.0 ^d				9 ⁻
28.1 \pm 100 ^c						10 ⁺
(29.4) ^c						(10 ⁺)
((33.4))						((10 ⁺))

^a For complete references see Tables 20.24 in (1978AJ03) and 20.22 in (1983AJ01).

^b Relative α -particle spectroscopic factors (DWBA). Other S_α values have also been reported.

^c (1982AR20).

^d (1983SH26).

^e An admixture of 6⁺ or 8⁺ in the d- α angular correlation involving $^{20}\text{Ne}^*(16.6)$ and a doublet (8⁺ + 7⁻) at $E_x = 17.4$ MeV have been suggested. See also Table 20.18.

States observed in this reaction are displayed in Table 20.19. Angular distributions have been measured at $E(^7\text{Li}) = 15$ to 68 MeV: see (1978AJ03, 1983AJ01). See also (1986CO15; theor.).

$$17. \ ^{16}\text{O}(^9\text{Be}, ^5\text{He})^{20}\text{Ne} \quad Q_{\text{m}} = 2.27$$

See (1985CU1A).

$$18. \ (a) \ ^{16}\text{O}(^{12}\text{C}, ^8\text{Be})^{20}\text{Ne} \quad Q_{\text{m}} = -2.633$$

$$(b) \ ^{16}\text{O}(^{12}\text{C}, 2\alpha)^{20}\text{Ne} \quad Q_{\text{m}} = -2.541$$

$$(c) \ ^{16}\text{O}(^{12}\text{C}, \alpha^{12}\text{C})^{12}\text{C} \quad Q_{\text{m}} = -7.16195$$

Angular distributions in reaction (a) have been measured for $E(^{16}\text{O}) = 27.1$ to 46.4 MeV and for $E(^{12}\text{C}) = 22.7$ to 78 MeV [see (1978AJ03, 1985AJ01)] as well as at $E(^{12}\text{C}) = 109$ MeV (1984MU04, 1985MU14; $^{20}\text{Ne}^*(1.63, 4.25, 5.79, 7.16, 8.78, 10.26, 11.95, 12.59, 15.34, 15.87, 17.30, 21.08, 22.87)$; $\sigma(\theta)$ at several angles; EFR-DWBA analysis). Γ_{α_0}/Γ are displayed in Table 20.19: see (1983AJ01) and (1983SH26). Preliminary evidence for 10^+ strength at $E_x \approx 28$ MeV is reported by (1986ALZN). See also (1983DEZW). For reaction (b) see (1978AJ03) and (1986CA19). For reaction (c) and for a discussion of ^{24}Mg states reached in this reaction see (1983SH26, 1984MU04). See also (1985BE37, 1987SU03) and (1986BE19; theor.).

$$19. \text{}^{16}\text{O}(^{13}\text{C}, \text{}^9\text{Be})^{20}\text{Ne} \quad Q_m = -5.914$$

At $E(^{13}\text{C}) = 105$ MeV angular distributions to $^{20}\text{Ne}^*(1.63, 4.25, 8.78, 11.95, 15.34, 21.0)$ have been studied by (1979BR03): the first four states are the 2^+ , 4^+ , 6^+ and 8^+ members of the 0_1^+ band; the two higher states [$J^\pi = 7^-, 9^-$] belong to the 0^- band whose band head is $^{20}\text{Ne}^*(5.79)$. In addition distributions are reported to $^{20}\text{Ne}^*(12.59, 15.9, 17.3)$ [$J^\pi = 6^+, 8^+, 8^+$] (1979BR03). See also (1985MU14). For fusion cross sections see (1986PA10).

$$20. \text{}^{16}\text{O}(^{16}\text{O}, \text{}^{12}\text{C})^{20}\text{Ne} \quad Q_m = -2.428$$

Angular distributions have been reported to a number of states of ^{20}Ne at $E(^{16}\text{O}) = 23.9$ to 95.2 MeV [see (1978AJ03, 1983AJ01)] and recently at $E(^{16}\text{O}) = 26, 28$ and 30 MeV (1986CA24). (1983ME13) have studied the quasi-elastic spectrum at $E(^{16}\text{O}) = 50, 60, 68$ and 72 MeV. For excitation functions see (1986CA24; $^{20}\text{Ne}^*(0, 1.63)$). See also (1982KO1C, 1984ME10, 1985ST1B) and (1982KO1D, 1984AP03, 1984KO13; theor.).

$$21. \text{(a) } ^{17}\text{O}(^3\text{He}, ^3\text{He})^{17}\text{O} \quad E_b = 21.168$$

$$\text{(b) } ^{17}\text{O}(^3\text{He}, \alpha)^{16}\text{O} \quad Q_m = 16.4341$$

The excitation function for α_0 shows a resonance corresponding to $^{20}\text{Ne}^*(28.)$: see (1978AJ03). At $E(^3\vec{\text{He}}) = 33$ MeV A_y measurements have been reported for the elastic scattering [reaction (a)] (1983LE03) and for many α -groups [see ^{16}O in (1986AJ04)] (1982KA12). For the earlier work and for other channels see (1983AJ01). See also (1985HA11, 1987CO07; theor.).

$$22. \text{}^{17}\text{O}(\alpha, \text{n})^{20}\text{Ne} \quad Q_m = 0.590$$

See (1978AJ03).

23. (a) $^{17}\text{O}(^{11}\text{B}, ^8\text{Li})^{20}\text{Ne}$ $Q_m = -6.041$
 (b) $^{17}\text{O}(^{12}\text{C}, ^9\text{Be})^{20}\text{Ne}$ $Q_m = -5.111$

At $E = 115$ MeV the 8^+ state at $E_x = 11.95$ MeV is particularly strongly populated in both reactions: see (1983AJ01).

24. $^{18}\text{O}(^3\text{He}, n)^{20}\text{Ne}$ $Q_m = 13.124$

Angular distributions have been measured for $E(^3\text{He}) = 2.8$ to 18.3 MeV. States of ^{20}Ne observed in this reaction are displayed in Table 20.23 of (1983AJ01). These include a state at $E_x = 16.730 \pm 0.006$ MeV, $\Gamma < 20$ keV: $J^\pi = 0^+$, $T = 2$.

25. $^{19}\text{F}(p, \gamma)^{20}\text{Ne}$ $Q_m = 12.848$

Over the range $E_p = 2.9$ to 12.8 MeV, the γ_0 and γ_1 yields are dominated by the E1 giant resonance ($\Gamma \approx 6$ MeV) with the γ_1 giant resonance displaced upward in energy. Strong well-correlated structure is observed with a characteristic $\Gamma \approx 175$ keV. Angular distributions taken over the energy range do not vary greatly with energy. They are incompatible with γ_0 and γ_1 coming from the same levels in ^{20}Ne . The 90° γ_0 yield for $E_{\bar{p}}$ and $E_p = 3.5$ to 10 MeV has been measured: the results are interpreted in terms of four primary doorway states at $E_x = 16.7, 17.8, 19.1$ and 20.2 MeV. See also (1985WAZV; $E_{\bar{p}} = 5.9$ to 10.3 MeV; E2 strength; prelim.). See also (1986OUZZ).

The yield curve for 11.2 MeV γ -rays [from the decay of $^{20}\text{Ne}^*(11.23)$, $J^\pi = 1^+$, $T = 1$, to the ground state] displays a resonance at $E_p = 4.090 \pm 0.005$ MeV [$^{20}\text{Ne}^*(16.73)$]. The 11.2 MeV γ -rays are isotropic which is consistent with the presumed 0^+ character of this lowest $T = 2$ state in ^{20}Ne : $\Gamma_p \Gamma_\gamma / \Gamma \approx 0.5$ eV. Since Γ_p / Γ (from the elastic scattering) is ≈ 0.1 , $\Gamma_\gamma \approx 5$ eV. For $E_p = 5.65$ to 6.21 MeV, the γ_0 and γ_1 yields are not resonant but the yield of 10.6 MeV γ -rays is resonant at 5.879 ± 0.007 MeV [$\Gamma_{\text{c.m.}} = 9.5 \pm 3$ keV, $\Gamma_{p_0} \Gamma_\gamma / \Gamma \approx 0.05$ eV; $\Gamma_\gamma \approx 0.3$ eV]. The 10.6 MeV γ -ray is due to the cascade decay of $^{20}\text{Ne}^*(18.43)$, $J^\pi = 2^+$; $T = 2$ via $^{20}\text{Ne}^*(12.22)$ to the 2^+ state at 1.63 MeV. For the upper limits to the strengths of the transitions to various states of ^{20}Ne from the 0^+ and 2^+ $T = 2$ states, see (1983AJ01). Resonances observed in this capture reaction are displayed in Table 20.20. For references see (1978AJ03, 1983AJ01). See also (1987RO25; astrophysics).

26. (a) $^{19}\text{F}(p, p)^{19}\text{F}$ $E_b = 12.848$
 (b) $^{19}\text{F}(p, p')^{19}\text{F}^*$
 (c) $^{19}\text{F}(p, d)^{18}\text{F}$ $Q_m = -8.2074$

Table 20.20: Resonances in $^{19}\text{F}(p, \gamma)^{20}\text{Ne}$ ^a

E_p (keV)	Γ_{lab} (keV)	Γ_{γ_0} (eV)	Γ_{γ_1} (eV)	$^{20}\text{Ne}^*$ (MeV)	$J^\pi; T$
340		< 0.07	0.28 ± 0.06	13.171	
484		≈ 0.05	0.42	13.308	
597 ± 1	30 ± 3	< 0.6	12	13.415	
671 ± 1	6.0 ± 0.7	1.0×10^{-2}	2.2	13.485	1^+
874				13.678	
935				13.736	
980				13.779	
1091	0.8		1.1	13.884	$2^+; 1$
1280				14.063	
1320	4.0			14.101	
1350				14.130	
1370				14.149	
1420	15.7			14.196	
4090 ± 5		$\Gamma_\gamma \approx 5 \text{ eV}$		16.732	$0^+; 2$
5879 ± 7	10 ± 3	$\Gamma_\gamma \approx 0.3 \text{ eV}$		18.430	$2^+; 2$

^a For earlier references see Table 20.26 in (1978AJ03) and 20.24 in (1983AJ01). See also Table 20.14 here.

The elastic scattering has been studied in the range $E_p = 0.5$ to 7.5 MeV and 24.9 to 46.3 MeV [see (1978AJ03)] and at $E_p = 1.5$ to 3.5 MeV (1985OU01, 1986OUZZ, 1986OU01). The observed anomalies are displayed in Table 20.21.

Resonances for inelastic scattering [p_1 and p_2] are listed in Table 20.22. In general the resonances observed are identical with those reported from other $^{19}\text{F}+p$ reactions, although the relative intensities differ greatly. Cross sections for production of 110 and 197 keV γ -rays are reported for $E_p = 0.5$ to 4.3 MeV by (1986CHYY). See also (1983LE28; astrophysics) and (1986BA88). For reaction (c) see (1986KA1U; applied) and ^{18}F .

$$27. \ ^{19}\text{F}(p, n)^{19}\text{Ne} \qquad Q_m = -4.0207 \qquad E_b = 12.848$$

Observed resonances are displayed in Table 20.30 of (1978AJ03). See also (1984BA1R).

$$28. \ ^{19}\text{F}(p, \alpha)^{16}\text{O} \qquad Q_m = 8.1137 \qquad E_b = 12.848$$

Many resonances occur in this reaction. They are displayed in Tables 20.23, 20.24 and 20.25 depending on whether they are observed in the α_0 yield [20.23], in the α_1 [or α_π] yield to $^{16}\text{O}^*(6.05)$ [20.24] or in the α_2 , α_3 and α_4 yields [or in the yield of the γ -rays from $^{16}\text{O}^*(6.13, 6.92, 7.12)$ [20.25]]. Resonances for α_0 and α_1 are required to have even J , even π or odd J , odd π , while the α_2 , α_3 and α_4 resonances are all odd-even or even-odd, with the exception of the $T = 2$ resonance.

Listings of the earlier yield studies are given in (1972AJ02, 1978AJ03, 1983AJ01). A detailed discussion of the evidence leading to many of the J^π assignments is given in (1959AJ76). For values of θ^2 see Table 20.28 in (1978AJ03). Recent measurements are reported by (1985OU01; 1.15 to 2.1 MeV; $\alpha_0 \rightarrow \alpha_3$) and (1984IN04; 4.15 to 13 MeV; $\alpha_0 \rightarrow \alpha_5$). In the latter work there are no marked correlations between the different channels.

Longitudinally and transversely polarized protons with $E_p \approx 0.67$ MeV have been used to study $^{20}\text{Ne}^*(13.48)$ [$J^\pi = 1^+$; $T = 1$] via a parity (and isospin) forbidden α -transition. The state is not excited. The upper limits for the process, and their significance in the determination of f_π , the weak pion-nucleon coupling constant, are discussed by (1983KN01, 1986KN1C). See also (1983AJ01, 1984KN1A).

See also (1985ISZU), (1982MA1Q, 1982MA1V, 1983DE2J, 1986SI1L; applications), (1987RO25; astrophysics), (1986SI1D; computing) and (1983IN1B, 1985LO1C, 1986LO1E, 1986SI1K; theor.).

$$29. \ ^{19}\text{F}(d, n)^{20}\text{Ne} \qquad Q_m = 10.623$$

Levels of ^{20}Ne derived from this reaction are displayed in Tables 20.31 in (1972AJ02) and 20.34 in (1978AJ03). See also (1983LIZW).

Table 20.21: Levels of ^{20}Ne from $^{19}\text{F}(p, p_0)$ ^a

E_p (keV)	Γ_{lab} (keV)	l	$J^\pi; T$	Γ_p/Γ	θ_p^2 (%)	$^{20}\text{Ne}^*$ (MeV)
340	2.9	0	1^+	0.016	3.8	13.171
483			1^+			13.307
598	37	1	2^-	0.0012	0.38	13.416
669	7.5	0	1^+	0.98	9.6	13.483
843	23	0	0^+	0.996	10.8	13.649
873	5.2	1	2^- ^b	0.21	1.5	13.677
935	8.0	0	1^+	0.17	0.44	13.736
1346	4.5	1	2^- ^b	0.067	0.07	14.126
1372	15	1	2^- ^b	0.17	0.52	14.151
1422	14.6	0	1^+	0.85	0.92	14.198
1710 ^c	90	0	0^+	0.8		14.472
1896 ^c	25	0	0^+	0.3		14.648
1943 ^c	40	0	(1^+)	0.5		14.693
2030 ^c	70	1	(1^-)	0.75		14.776
2.763 ^c		2				15.472
2.970 ^c		2				15.668
4094 ± 3	2.1 ± 0.5	0	$0^+; 2$	0.062 ± 0.004		16.735
5879 ± 7 ^d	10 ± 3	2	$2^+; 2$	≈ 0.2		18.430

^a For references see Table 20.27 in (1978AJ03). For θ^2 see Table 20.28 in (1978AJ03).

^b 1^- not excluded by elastic scattering alone.

^c (1985OU01, 1986OU01; *R*-matrix analysis). Weak resonances at $E_p = 1.75$ and 1.78 MeV are also suggested.

^d Resonance also observed in p_1, p_3, p_4 and p_5 yields.

Table 20.22: Resonances in $^{19}\text{F}(\text{p}, \text{p}')^{19}\text{F}^*$ ^a

E_p (keV)	J^π	Γ_{lab} (keV)	Γ_{p_1} (eV)	Γ_{p_2} (eV)	$\theta_{p_1}^2$ (%)	$\theta_{p_2}^2$ (%)	E_x in ^{20}Ne (MeV)
340	1 ⁺	2.9	< 0.5	< 0.1	< 15		13.171
483	1 ⁺	2.2	< 1.3	< 1.2			13.307
598	2 ⁻	37	< 100	< 60	< 28	< 145	13.416
669	1 ⁺	7.5	46	< 0.5	0.6	< 0.4	13.483
720		≈ 30	< 10000	< 10000			13.532
780		≈ 10	< 400	≈ 9000			13.589
831		8.3	< 6	≈ 2300			13.637
845	0 ⁺	23	≈ 50	< 10	≈ 0.14	< 0.92	13.650
873	2 ⁻	5.2	< 2	570	< 0.07	2.7	13.677
900		4.8	< 30	≈ 2200			13.703
935	1 ⁺	8.0	3000	< 20	5.0	< 0.8	13.736
1092 ^b	2 ⁺	0.8	173	592			13.885
1137		3.7	< 40	≈ 2100			13.928
≈ 1250		≈ 80	≈ 70000	< 4000			14.03
1290		19	< 600	≈ 900			14.073
1346	2 ⁻	4.5	300	600	0.92	0.24	14.126
1372	2 ⁻	15	700	1400	1.93	0.56	14.151
1422	1 ⁺	14.6 ± 1	2200	≤ 35	0.56	≤ 0.11	14.198
1610		≈ 5					14.377
1660							14.424
1700							14.462
2763 ^c							15.472
2970 ^c							15.668
5879 ^d	2 ⁺ ; 2		r				18.430

r = resonant.

^a For references see Tables 20.29 in (1978AJ03) and 20.26 in (1983AJ01).

^b $\Gamma_{p_0} = 29$ eV.

^c Reported in $p_{1 \rightarrow 4}$ yield (1986OU01).

^d Resonance also observed in p_3 , p_4 and p_5 yields.

Table 20.23: Resonances for ground-state α -particles (α_0) in $^{19}\text{F}(p, \alpha_0)$ ^a

E_p (keV)	Γ_{lab} (keV)	θ_α^2 (%) ^a	$J\pi$	$^{20}\text{Ne}^*$ (MeV)
400	100		1^-	13.228
400	100		0^+	13.228
650 ± 20	200		1^-	13.465
710	35	0.6	(1^-)	13.522
733	66	1.0	2^+	13.544
777 ± 2	9 ± 1	0.02	2^+	13.586
842 ± 2	18 ± 1	0.16^b	$(2^+)^c$	13.648
≈ 860	120	2.1	1^-	13.66
≈ 930	≈ 180	2.9	0^+	13.73
≈ 1080	≈ 200	3.4	1^-	13.87
1115	50	0.55	2^+	13.907
1160	≈ 70	1.1	0^+	13.950
1235	≈ 70	1.2	1^-	14.021
≈ 1250	≈ 150	2.7	2^+	14.03
1350 ± 3	36 ± 1		2^+	14.130
1652 ± 5	90 ± 5		1^-	14.417
1713 ± 6	72 ± 2		0^+	14.475
1842 ± 7	122 ± 5		1^-	14.597
1901 ± 10	25^d		0^+	14.653
2110	75		$(2^+, 4^+)$	14.85
2310	90		(2^+)	15.04
2550	300		(1^-)	15.27
2590	300		(0^+)	15.31
2680	80			15.39
2730	60			15.44
2820	160			15.53
2940				(15.64)
3120	170			(15.81)
3340	105			16.02
3680	(100)			16.34
3860				16.51

Table 20.23: Resonances for ground-state α -particles (α_0) in $^{19}\text{F}(\text{p}, \alpha_0)$ ^a
(continued)

E_p (keV)	Γ_{lab} (keV)	θ_{α}^2 (%) ^a	J^{π}	$^{20}\text{Ne}^*$ (MeV)
3980	135			16.63
4130	100			16.77
4360	100			16.99
4460	95			17.08
4690	65			17.30
4900	90			17.50
4990	40			17.59
5879 ± 7	10 ± 3	^d	$2^+; T = 2$	18.430

^a For earlier references and additional comments see Tables 20.31 in (1978AJ03) and 20.28 in (1983AJ01). See also (1985OU01, 1986OU01).

^b $\Gamma_{\alpha_0} \approx 0.06$ keV.

^c $J = 0$ from $^{19}\text{F}(\text{p}, \text{p})$; possibly $T = 0$.

^d $\Gamma_{\alpha_0} \approx 0.3$ keV.

$$30. \ ^{19}\text{F}(^3\text{He}, \text{d})^{20}\text{Ne} \quad Q_m = 7.354$$

Levels of ^{20}Ne observed in this reaction are displayed in Table 20.35 of (1978AJ03). Deuteron angular distributions have been studied at $E(^3\text{He}) = 9.5$ to 21 MeV: see (1978AJ03).

The ΔE_x between the 1^+ and 1^- , $T = 1$ states $^{20}\text{Ne}^*(11.26, 11.27)$ is 11.1 ± 0.7 keV (1983FI02). $\Gamma_{\gamma}/\Gamma_{\alpha} = 0.88 \pm 0.05$ for $^{20}\text{Ne}^*(12.22)$ [$2^+; T = 1$] (1984CA08). Using $(2J + 1)\Gamma_{\alpha}\Gamma_{\gamma}/\Gamma = 1.41 \pm 0.23$ eV (1980FI01), $\Gamma_{\alpha} = 0.32 \pm 0.06$ eV for $^{20}\text{Ne}^*(12.22)$ (1984CA08). The Γ_{γ}/Γ of $^{20}\text{Ne}^*(12.22)$ implies $B(\text{M1}) = 0.07$ W.u. for the transition from $^{20}\text{Ne}^*(18.43)$ [$2^+; T = 2$]. This is much weaker than other isovector M1 transitions in ^{20}Ne and a factor of five lower than predicted by shell model calculations: see (1984CA08).

$$31. \ ^{19}\text{F}(\alpha, \text{t})^{20}\text{Ne} \quad Q_m = -6.966$$

Angular distributions have been measured at $E_{\alpha} = 18.5$ and 28.5 MeV: see (1978AJ03, 1983AJ01).

Table 20.24: Nuclear pair resonances (α_π) in $^{19}\text{F}(\text{p}, \alpha_\pi)$ ^a

E_p (keV)	Γ_{lab} (keV)	σ (mb)	θ_α^2 (%)	J^π	$^{20}\text{Ne}^*$ (MeV)
710	35	≈ 0.2	2	1^-	13.522
780	≈ 10	≈ 0.2	0.15	2^+	13.589
842	23	3.4	0.27	2^+ ^c	13.648
1115	50	1.5	3.6	2^+	13.907
1236	≈ 70	3	1.0	1^-	14.022
1367	30	6.0	0.29	2^+	14.146
1640	60			1^-	14.41
1720	95	≈ 18		0^+	14.48
1850	170			1^-	14.60
1896	25			0^+	14.65
2080 ^b	60	12.1		(2^+)	14.82
2170 ^b	70	12.2		(0^+)	14.91
2330 ^b	70	17.0		(2^+)	15.06
2600	100				15.32
2680	100				15.39
2820	125				15.53
3120	145				15.81
3340	100				16.02
(3500)	(80)				(16.17)
(3590)	(115)				(16.26)
3960	200				16.61
4360	95				16.99
4690	< 150				17.30
4900	115				17.50
4990	40				17.59
5170	220				17.76

^a For references see Tables 20.32 in (1978AJ03) and 20.29 in (1983AJ01). See also (1985OU01, 1986OU01).

^b (1980CU09): see also for partial widths.

^c See footnote ^c in Table 20.23.

Table 20.25: Resonances for 6 – 7 MeV γ -rays ($\alpha_2, \alpha_3, \alpha_4$) in $^{19}\text{F}(\text{p}, \alpha)$ ^a

E_p (keV)	Γ_{lab} (keV)	Γ_{α_2} (eV)	Γ_{α_3} (eV)	Γ_{α_4} (eV)	$J\pi$	$^{20}\text{Ne}^*$ (MeV)
223.99 ± 0.07 ^b	0.99 ± 0.02	1000	< 2.5	< 2.5	2^-	13.0607
340.46 ± 0.04 ^{b,c}	2.34 ± 0.04	2800	16	75	1^+	13.1713
483.91 ± 0.10 ^b	0.90 ± 0.03	700	19	190	1^+	13.3075
594 ± 3	25 ± 3					13.412
667.5 ± 2	6.7 ± 0.3					13.482
832.1 ± 1						13.638
872.11 ± 0.20 ^d	4.53 ± 0.16	2200	620	180	2^-	13.6762
935.4 ± 1.3	8.1 ± 0.5	2900	110	720	1^+	13.736
1087.7 ± 1	0.15 ± 0.05					13.881
1135.6 ± 1						13.926
1280 ± 1						14.063
1347.7 ± 1	4.9 ± 0.7	2250	650	1200	2^-	14.128
1371.0 ± 1	12.4 ± 1.0	6650	700	300	2^-	14.150
1603 ± 2						14.370
1692 ± 2	35 ± 3				$(1, 2)^-$	14.455
1949 ± 2.5	40 ± 10				$(0, 1)^+$	14.699
2030 ± 3.0	120 ± 20					14.776
2320	85					15.05
2510	30					15.23
2630	90					15.35
2800	60					15.51
3020	30					15.72
3190	80					15.88
3490	40					16.16
3920	30					16.57
4000	110					16.65
4090					$0^+; T = 2$	16.73
4290	50					16.92
4490	30					17.11
4570	30					17.19
4710	30					17.32

Table 20.25: Resonances for 6 – 7 MeV γ -rays ($\alpha_2, \alpha_3, \alpha_4$) in $^{19}\text{F}(\text{p}, \alpha)$ ^a
(continued)

E_p (keV)	Γ_{lab} (keV)	Γ_{α_2} (eV)	Γ_{α_3} (eV)	Γ_{α_4} (eV)	J^π	$^{20}\text{Ne}^*$ (MeV)
4780	35					17.39
4990	20					17.59
5070	35					17.66
5200	70					17.79

^a See Tables 20.33 in (1978AJ03) and 20.30 in (1983AJ01) for earlier references and for additional comments. See also (1985OU01, 1986OU01).

^b (1985UH01). See also (1977FR20).

^c (1982BE29): $\sigma = 88 \pm 3$ mb, $\omega\gamma = 22.3 \pm 0.8$ eV.

^d (1982BE29): $\sigma = 440 \pm 13$ mb, $\omega\gamma = 570 \pm 30$ eV.

$$32. \ ^{19}\text{F}(^7\text{Li}, ^6\text{He})^{20}\text{Ne} \quad Q_m = 2.873$$

Angular distributions have been studied at $E(^7\text{Li}) = 34$ MeV to a number of states of ^{20}Ne . C^2S values are consistent with those reported in the (d, n) and (^3He , d) reactions: see (1978AJ03).

$$33. \ ^{20}\text{F}(\beta^-)^{20}\text{Ne} \quad Q_m = 7.029$$

The decay is primarily to $^{20}\text{Ne}^*(1.63)$ with a half-life of 11.00 ± 0.02 sec: see reaction 1 in ^{20}F . Besides the principal decay to $^{20}\text{Ne}^*(1.63)$ [$\log f_0t = 4.97$], ^{20}F also decays to $^{20}\text{Ne}^*(4.97)$ [$J^\pi = 2^-$] with a branching ratio of $(0.0082 \pm 0.0006)\%$ (1987AL06) [$\log f_0t = 7.20 \pm 0.03$; D.E. Alburger and E.K. Warburton, private communication]. The upper limit for the ground-state decay is 0.001% [$\log f_0t > 10.5$]. For other values and earlier references see Table 20.36 in (1978AJ03). The energy of the γ -ray from $^{20}\text{Ne}^*(1.63)$ is 1633.602 ± 0.015 keV. E_γ for the $4.97 \rightarrow 1.63$ transition is 3332.54 (19) keV which gives $E_x = 4966.51$ (20) keV based on $E_x = 1633.674$ (15) keV for the first excited state. The shape of the β -spectrum is not inconsistent with the predictions of CVC. $\beta - \gamma$ correlation measurements lead to an upper limit for the second-class current contribution to the correlation which is consistent with zero. For references see (1983AJ01); for the earlier work see (1978AJ03). See also (1985BR29, 1985GR1A) and (1983KA32, 1985YA1J; theor.).

$$34. \text{ (a) } ^{20}\text{Ne}(\gamma, \text{n})^{19}\text{Ne} \quad Q_m = -16.869$$

$$\text{ (b) } ^{20}\text{Ne}(\gamma, 2\text{n})^{18}\text{Ne} \quad Q_m = -28.508$$

$$\text{ (c) } ^{20}\text{Ne}(\gamma, \alpha)^{16}\text{O} \quad Q_m = -4.734$$

The photoneutron cross section (bremsstrahlung photons) shows peaks at $E_x = 17.78 \pm 0.05$, 19.00 ± 0.05 , 20.15 ± 0.15 [main peak of the GDR], 22.6 ± 0.3 , 24.9 ± 0.5 and 27.5 MeV [the latter three states are broad]: the integrated cross section to 28.5 MeV is 58 ± 6 MeV · mb [exhausting $\approx 20\%$ of the dipole sum]. The cross section for (γ, Tn) using monoenergetic photons shows a structure at 18 MeV and some fluctuations atop the broad giant resonance, $\sigma_{\text{max}} \approx 7$ mb. The double photo-neutron cross section, $\sigma(\gamma, 2\text{n})$, is dominated by a single peak at ≈ 20.5 MeV, $\sigma_{\text{max}} \approx 1.1$ mb. For references see (1978AJ03, 1983AJ01). The significance of reaction (c) to astrophysics is discussed by (1982SA1A, 1984FO1A).

35. $^{20}\text{Ne}(\gamma, \gamma)^{20}\text{Ne}$

$E_x = 11262.3 \pm 1.9$ keV for $^{20}\text{Ne}^*(11.26)$, the first 1^+ ; $T = 1$ state. The branchings to $^{20}\text{Ne}^*(0, 1.63)$ are (84 ± 5) and $(16 \pm 5)\%$, respectively (1983BE19). See also (1984BE26).

36. (a) $^{20}\text{Ne}(e, e)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(e, \text{ep})^{19}\text{F}$ $Q_m = -12.848$

(c) $^{20}\text{Ne}(e, e\alpha)^{16}\text{O}$ $Q_m = -4.734$

The ^{20}Ne charge radius, $\langle r^2 \rangle^{1/2} = 3.004 \pm 0.025$ fm. Form factors for many excited states of ^{20}Ne with $E_x < 8$ MeV have been reported: see (1978AJ03).

At $E_e = 39$ and 56 MeV, the 180° inelastic scattering is dominated by the transition to a $J^\pi = 1^+$, $T = 1$ state at $E_x = 11.22 \pm 0.05$ MeV with $\Gamma_{\gamma_0} = 11.2^{+2.1}_{-1.8}$ eV. A subsidiary peak is observed corresponding to a state 0.35 ± 0.03 MeV higher [if $J^\pi = 1^+$ or 2^+ , $\Gamma_{\gamma_0} = 0.65 \pm 0.18$ or 0.40 ± 0.13 eV]. A number of small peaks are also reported corresponding to $E_x \approx 12.0, 12.9, 13.9, 15.8, 16.9, 18.0$ and 19.0 MeV. Prominent electric dipole peaks are reported at $E_x = 17.7, 19.1, 20.2$ and 23 MeV, in addition to weaker structures between 12.5 and 15 MeV; and prominent electric-quadrupole peaks are observed at $E_x = 13.0, 13.7, 14.5, 15.0, 15.4$ and 16.2 MeV and there is a broad quadrupole excitation between 16 and 25 MeV. The GDR cross section integrated from 11 to 25 MeV contains about 65% of the dipole EWSR while over 90% of the isoscalar quadrupole EWSR is exhausted by the strength in the region $10 - 25$ MeV.

For $11 < E_x < 24$ MeV only two isovector M2 transitions appear: These are to $^{20}\text{Ne}^*(11.62, 12.10)$ with $B(\text{M2}, k) \uparrow = 64 \pm 13$ and $56 \pm 13 \mu_{\text{N}}^2 \text{fm}^2$ [orbital contributions are non-negligible]. The M1 transition to $^{20}\text{Ne}^*(11.26)$ is also observed but that to $^{20}\text{Ne}^*(13.48)$ is not: it is $< 0.2 \mu_{\text{N}}^2$ (1985RA08). For reaction (b) see (1978AJ03).

Reaction (c) has been studied in order to obtain the (γ, α_0) cross section in the giant resonance region: the cross section at 90° for $E_x = 15$ to 24 MeV is dominated by an E1 resonance [1^- ; $T = 1$, with an admixture of $T = 0$ which permits the α_0 decay] at $E_x = 20$ MeV; lesser E1 structures are reported at $E_x = 16.7, 17.1, 21$ and 22 MeV. A relatively strong 2^+ ; $T = 0$ resonance appears at $E_x = 18.5$ MeV, and evidence is reported for increasing E2 strength below 16 MeV.

For references see (1978AJ03). See also (1987DE43) and (1978WI1B, 1982RU05, 1983BR1P, 1983KN05, 1984AM03, 1984AM07, 1985LE1H, 1985LE14, 1985ST23, 1986BR1X, 1986KA17, 1986SH16, 1986TA1W, 1986VA12, 1987VA12; theor.).

37. $^{20}\text{Ne}(n, n)^{20}\text{Ne}$

See (1978AJ03).

38. (a) $^{20}\text{Ne}(p, p)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(p, p\alpha)^{16}\text{O}$ $Q_m = -4.734$

Angular distributions of elastically scattered protons and of a number of inelastic groups have been measured for $E_p = 2.15$ to 65 MeV [see (1978AJ03, 1983AJ01)] and at $E_{\bar{p}} = 0.8$ GeV (1984BL14; to $^{20}\text{Ne}^*(0, 1.63, 4.25, 8.7(u); \text{also } A_y)$). The latter work confirms the large hexadecapole deformation of ^{20}Ne . At $E_p = 201$ MeV, probable 1^+ states at $E_x = 11.25 \pm 0.01$, 13.51 ± 0.03 and 15.72 ± 0.05 MeV are reported by (1987WI03): There does not appear to be any quenching of the M1 strength. In addition 2^- states are observed at 11.58 and 12.08 MeV with $B(M2) = 64 \pm 13$ and $56 \pm 13 \mu_N^2$ as is a state of unknown J^π at $E_x \approx 17$ MeV (1987WI03). See also the earlier work in (1978AJ03). For reaction (b) see (1984CA09; $E_p = 101.5$ MeV). See also (1983CE1A, 1986BA2U), (1986BA88) and (1982SC20, 1983SM04, 1984AM03, 1984AM07, 1985KN04; theor.).

39. $^{20}\text{Ne}(\bar{p}, \bar{p})^{20}\text{Ne}$

For a study of antiproton interactions see (1986BA22).

40. (a) $^{20}\text{Ne}(d, d)^{20}\text{Ne}$

(b) $^{20}\text{Ne}(t, t)^{20}\text{Ne}$

Angular distributions of deuterons have been reported at $E_d = 10.0$ to 52 MeV [see (1978AJ03, 1983AJ01)] and at $E_{\bar{d}} = 52$ MeV (1987NU01). For reaction (b) see (1978AJ03).

41. $^{20}\text{Ne}(^3\text{He}, ^3\text{He})^{20}\text{Ne}$

Angular distributions have been measured at $E(^3\text{He}) = 10$ to 35 MeV and at 68 MeV: see (1978AJ03). See also (1987TR01; theor.).

42. (a) $^{20}\text{Ne}(\alpha, \alpha)^{20}\text{Ne}$
 (b) $^{20}\text{Ne}(\alpha, 2\alpha)^{16}\text{O}$ $Q_m = -4.734$

Angular distributions have been measured at $E_\alpha = 3.8$ to 155 MeV [see (1978AJ03, 1983AJ01)] and at 54.1 MeV (1987AB03; g.s.). See also (1986BU1L; $E_\alpha = 50.5$ MeV). For a spallation study see (1984AN1H); for cross sections of astrophysical interest see (1984SE02). For reaction (b) see (1983AJ01). See also (1984SA28, 1986TR08), (1982VO04, 1983CH1B) and (1982JA07, 1982WA13, 1984BU1R, 1984SR01, 1985UM01, 1986GA1N, 1986KA17; theor.).

43. $^{20}\text{Ne}(^7\text{Li}, ^7\text{Li})^{20}\text{Ne}$

Angular distributions have been studied at $E(^7\text{Li}) = 36, 68$ and 89 MeV: see (1983AJ01).

44. $^{20}\text{Ne}(^9\text{Be}, ^9\text{Be})^{20}\text{Ne}$

For pion production see (1985FR13).

45. (a) $^{20}\text{Ne}(^{10}\text{B}, ^{10}\text{B})^{20}\text{Ne}$
 (b) $^{20}\text{Ne}(^{11}\text{B}, ^{11}\text{B})^{20}\text{Ne}$

Elastic angular distributions have been measured at $E(^{10}\text{B}) = 65.9$ and $E(^{11}\text{B}) = 115$ MeV: see (1983AJ01).

46. (a) $^{20}\text{Ne}(^{12}\text{C}, ^{12}\text{C})^{20}\text{Ne}$
 (b) $^{20}\text{Ne}(^{12}\text{C}, \alpha^8\text{Be})^{20}\text{Ne}$ $Q_m = -7.3665$

Elastic angular distributions have been obtained at $E(^{12}\text{C}) = 22.2$ to 42.7 MeV and 77.4 MeV and at $E(^{20}\text{Ne}) = 65.9, 74$ and 75.2 MeV [see (1978AJ03, 1983AJ01)] as well as at $E(^{20}\text{Ne}) = 72.6, 74.0$ and 75.2 MeV (1982SH29). For yield, fusion, total reaction cross section and fragmentation studies see (1982KO29, 1982MO15, 1982SH29, 1984AN1H, 1984KO12, 1985FL1B, 1985OS05, 1986BL1K, 1987KO12) and (1983AJ01). For pion production see (1982AN1H, 1983AN1L).

For reaction (b) see (1983AJ01), (1984RA10, 1987SI06) and ^{12}C in (1985AJ01). See also (1985MU18), (1983HE1B, 1984FR1A, 1984HA53, 1984NA1D, 1984ST1B, 1985BE1A, 1985CU1A, 1985ST1B, 1986IK03, 1986ST1J, 1987LA05, 1987SC1D) and (1982LO13, 1983CI08, 1983TO1L, 1984HA43, 1984IN03, 1984MAZT, 1984SH1T, 1985CH11, 1985GA1G, 1985GU1J, 1985HU04, 1985KO1J, 1986GA1F, 1986GI03, 1986HA13, 1986HE1A; theor.).

47. $^{20}\text{Ne}(^{16}\text{O}, ^{16}\text{O})^{20}\text{Ne}$

Angular distributions have been studied at $E(^{20}\text{Ne}) = 50$ and 94.8 MeV involving $^{16}\text{O}_{\text{g.s.}}$ and $^{20}\text{Ne}^*(0, 1.63, 4.25)$ [see (1983AJ01)], at $E(^{16}\text{O}) = 25.6$ to 44.5 MeV (1984GA22; elastic; also to $^{20}\text{Ne}^*(1.63)$ at $31.3, 33.3$ and 44.5 MeV) and at $E(^{20}\text{Ne}) = 66.8, 115, 137$ and 156 MeV (1983SH25; elastic). Yield and fusion cross section measurements are reported by (1982SC13, 1982SH1N, 1983BR1R, 1983SH25, 1984GA22) and in (1983AJ01). See also (1983DU13, 1986BA69, 1986ST1J, 1987SC1D) and (1983KO31, 1983MA29, 1984NI1D, 1985CH11, 1985GU1J, 1985IC01, 1985KO43, 1985KO38, 1986GA1F, 1986HE1A, 1986KO1C, 1986TA1A; theor.).

48. $^{20}\text{Ne}(^{20}\text{Ne}, ^{20}\text{Ne})^{20}\text{Ne}$

Elastic angular distributions are reported at $E(^{20}\text{Ne}) = 68, 117, 140$ and 156 MeV (1983SH25). For yield and fusion measurements see (1983PO09, 1983SH25) and (1983AJ01). See also (1983RA1J, 1987SC1D) and (1983AB1G, 1983GO13, 1983SI01, 1984TO02, 1985GA1T, 1985IV1C, 1985IV1B, 1987BA1V; theor.).

49. (a) $^{20}\text{Ne}(^{24}\text{Mg}, ^{24}\text{Mg})^{20}\text{Ne}$

(b) $^{20}\text{Ne}(^{26}\text{Mg}, ^{26}\text{Mg})^{20}\text{Ne}$

Elastic angular distributions for reaction (a) have been measured at $E(^{20}\text{Ne}) = 50, 60, 80, 90$ and 100 MeV [see (1983AJ01)] and at 40 MeV (1983NA04; S_α for the system $^{20}\text{Ne} + ^{24}\text{Mg} = 0.08 \pm 0.02$). For yield and fusion cross sections see (1982GR1T, 1983AL01, 1984CHZU [also reaction (b)], 1984LE06) and (1983AJ01). See also (1983BI13, 1983DU13, 1984FR1A, 1986BL08) and (1982LO13, 1983OH1C, 1985NI1C, 1987BA01; theor.).

50. $^{20}\text{Ne}(^{27}\text{Al}, ^{27}\text{Al})^{20}\text{Ne}$

Elastic angular distributions are reported at $E(^{20}\text{Ne}) = 55.7, 63, 125$ and 151 MeV (1983NG01). For yield, fusion and evaporation residue studies see (1982MO15, 1983MO13, 1983NG01, 1985MO08,

1986GRZK, 1987JA06) and (1983AJ01). See also (1983BI13, 1983HE1B, 1984FR1A, 1984HA53, 1985ST1B, 1985TU01) and (1982LO13, 1983CI08, 1983GO13, 1983OH1B, 1984BI06, 1985HU04; theor.).

51. (a) $^{20}\text{Ne}(^{28}\text{Si}, ^{28}\text{Si})^{20}\text{Ne}$
 (b) $^{20}\text{Ne}(^{29}\text{Si}, ^{29}\text{Si})^{20}\text{Ne}$

See (1983DU13).

52. $^{20}\text{Ne}(^{40}\text{Ca}, ^{40}\text{Ca})^{20}\text{Ne}$

Angular distributions have been studied at $E(^{20}\text{Ne}) = 44.1$ to 70.4 MeV and at 151 MeV: see (1983AJ01). For an evaporation residue study see (1982MO15). For yield and fusion measurements see (1983AJ01). The breakup of ^{20}Ne at $E(^{20}\text{Ne}) = 92, 149$ and 213 MeV involves $^{20}\text{Ne}^*(5.79, 6.73, 7.16, 8.78, 10.26, 11.95)$ (1986SH30). See also (1983BI13, 1983HE1B, 1984FR1A) and (1982ST08, 1983SI01, 1984GU09, 1984SH2D, 1985AN16, 1985GU08; theor.).

53. $^{20}\text{Na}(\beta^+)^{20}\text{Ne} \quad Q_m = 13.887$

^{20}Na has a half-life of 447.9 ± 2.3 msec: see reaction 1 in ^{20}Na . It decays to a number of states of ^{20}Ne , principally $^{20}\text{Ne}^*(1.63)$: see Table 20.26. The ratio of the mirror decays $^{20}\text{Na} \xrightarrow{\beta^+} ^{20}\text{Ne}^*(1.63)$ and $^{20}\text{F} \xrightarrow{\beta^-} ^{20}\text{Ne}^*(1.63)$, $(ft)^+/(ft)^- = 1.03 \pm 0.02$. $\beta - \gamma$ correlation measurements, as in the decay of ^{20}F , lead to an upper limit for the second-class contribution to the correlation which is consistent with zero: see (1983AJ01). $\beta - \nu - \alpha$ triple correlation coefficient measurements for the transitions via the α -unstable 2^+ states shown in Table 20.26 lead to values of the isospin mixing amplitudes [and to a determination of the vector weak coupling constant] (1983CL01). See also (1987ROZZ) and (1983HA1V, 1985BR29, 1985GR1A, 1985HA1T).

54. $^{21}\text{Ne}(p, d)^{20}\text{Ne} \quad Q_m = -4.536$

See (1978AJ03).

55. $^{21}\text{Ne}(d, t)^{20}\text{Ne} \quad Q_m = -0.503$

Table 20.26: Decay of ^{20}Na ^a

Decay to $^{20}\text{Ne}^*$ (MeV \pm keV)	$J^\pi; T$	Branching ratio (%)	$\log ft$
1.633 ± 2	$2^+; 0$	79.33 ± 1.11	4.988 ± 0.009
7.415 ± 5	$2^+; 0$	16.37 ± 1.28	4.19 ± 0.05
7.826 ± 7	$2^+; 0$	0.674 ± 0.055	5.417 ± 0.033
8.82 ± 10		0.034 ± 0.007	6.27 ± 0.08
9.481 ± 7	$2^+; 0$	0.247 ± 0.020	5.064 ± 0.034
9.873 ± 5	$3^+; 0$	0.0272 ± 0.0138	5.78 ± 0.18
10.275 ± 3	$2^+; 1$	2.868 ± 0.039 ^b	3.476 ± 0.009 ^b
10.584 ± 7	$2^+; 0$	0.087 ± 0.009	4.76 ± 0.05
10.848 ± 7	$2^+; 0$	0.193 ± 0.016	4.179 ± 0.035
10.884 ± 3	$3^+; 1$	0.0392 ± 0.0139	4.84 ± 0.13
11.261 ± 5	$1^+; 1$	0.203 ± 0.026	3.73 ± 0.05
11.320 ± 15	$2^+; 0$	0.036 ± 0.004	4.41 ± 0.05
11.856 ± 20	$2^+; 0$	0.0016 ± 0.0004	4.98 ± 0.10

^a For additional comments and references see Table 20.37 in (1978AJ03).

^b (1983CL01).

The $T = 1$ states observed in this reaction, and their analog states observed in ^{20}F in the (d, ^3He) reaction, are displayed in Table 20.16 of (1978AJ03). $T = 0$ states are presented in Table 20.38 of (1978AJ03).

$$56. \ ^{22}\text{Ne}(\text{p}, \text{t})^{20}\text{Ne} \quad Q_{\text{m}} = -8.641$$

Angular distributions have been reported at $E_{\text{p}} = 26.9$ to 43.7 MeV: see (1978AJ03, 1983AJ01). The angular distributions of the tritons to the ground state of ^{20}Ne and to the first 0^+ , $T = 2$ state [$E_{\text{x}} = 16.722 \pm 0.025$ MeV] have been fitted by $L = 0$ and the tritons to $^{20}\text{Ne}^*(18.4)$ by $L = 2$. The latter is the first 2^+ , $T = 2$ state. The 0^+ , $T = 2$ state [$^{20}\text{Ne}^*(16.73)$] decays by $\alpha_0[(-6 \pm 5)\%]$, $\alpha_1 + \alpha_2[(35 \pm 12)\%]$, $\alpha_3 + \alpha_4[(29 \pm 12)\%]$, $\text{p}_0 + \text{p}_1 + \text{p}_2[(14 \pm 9)\%]$ and $\text{p}_3 + \text{p}_4 + \text{p}_5[(13 \pm 8)\%]$ [measured branching ratios in percent are given in the brackets] to final states in ^{16}O and ^{19}F . See (1978AJ03) for references and additional information.

$$57. \ ^{23}\text{Na}(\text{p}, \alpha)^{20}\text{Ne} \quad Q_{\text{m}} = 2.379$$

Angular distributions have been measured at $E_{\text{p}} = 10.0$ and 45.5 MeV: see (1972AJ02). For astrophysical considerations see (1982SA1A, 1984FO1A, 1985AR1A) and (1983AJ01). See also (1985RH1A) and (1986TU1B; applied).

$$58. \ ^{23}\text{Na}(^3\text{He}, ^6\text{Li})^{20}\text{Ne} \quad Q_{\text{m}} = -1.639$$

See (1978AJ03).

$$59. \ ^{24}\text{Mg}(\text{p}, \text{p}\alpha)^{20}\text{Ne} \quad Q_{\text{m}} = -9.312$$

See (1984CA09). See also (1978AJ03).

$$60. \ ^{24}\text{Mg}(\text{d}, ^6\text{Li})^{20}\text{Ne} \quad Q_{\text{m}} = -7.837$$

Angular distributions have been studied to many states of ^{20}Ne at $E_{\text{d}} = 28$ to 80 MeV [see (1978AJ03, 1983AJ01)] and at $E_{\text{d}} = 54.2$ MeV (1984UM04; to $^{20}\text{Ne}^*(0, 1.63, 4.25, 5.62)$). Table 20.35 in (1983AJ01) displays the observed states and S_{α} obtained from several analyses. For newer values of S_{α} see (1984UM04, 1986OE01). See also (1984PA18, 1986PAZJ; theor.).

$$61. {}^{24}\text{Mg}({}^3\text{He}, {}^7\text{Be}){}^{20}\text{Ne} \quad Q_m = -7.724$$

Angular distributions have been studied at $E({}^3\text{He}) = 25.5$ and 70 MeV: see (1978AJ03). See also (1983AJ01) and (1986RA15; theor.).

$$62. {}^{24}\text{Mg}(\alpha, {}^8\text{Be}){}^{20}\text{Ne} \quad Q_m = -9.404$$

See (1983AJ01).

$$63. {}^{24}\text{Mg}({}^{12}\text{C}, {}^{16}\text{O}){}^{20}\text{Ne} \quad Q_m = -2.150$$

The angular distribution for the ground state transition has been measured at $E({}^{12}\text{C}) = 40$ MeV (1982LI16). See also (1983AJ01).

$$64. {}^{28}\text{Si}(\alpha, {}^{12}\text{C}){}^{20}\text{Ne} \quad Q_m = -12.021$$

See (1983AJ01).

²⁰Na
(Figs. 11, 12 and 13)

GENERAL: (See also (1983AJ01).)

(1981WA1Q, 1983ANZQ, 1983BR29, 1985AN28, 1985HA1N, 1985RO1N, 1986AN07, 1986GA1P).

1. $^{20}\text{Na}(\beta^+)^{20}\text{Ne}$ $Q_m = 13.887$

²⁰Na decays by positron emission to ²⁰Ne*(1.63) and to a number of other excited states of ²⁰Ne: see Table 20.26 and reaction 53 in ²⁰Ne. The half-life of ²⁰Na is 447.9 ± 2.3 msec [weighted mean of values quoted in (1978AJ03) and in (1983CL01)]; $J^\pi = 2^+$: see (1978AJ03). See also (1986HO35).

2. $^{19}\text{F}(p, \pi^-)^{20}\text{Na}$ $Q_m = -140.608$

Angular distributions and A_y have been studied at $E_p = 199.6$ MeV to ²⁰Na*(0.74, 1.85, 3.01, 4.11) [probably unresolved]: it is suggested that the latter two have $J = 6$ or 7 (1987CA05).

3. $^{19}\text{Ne}(p, \gamma)^{20}\text{Na}$ $Q_m = 2.199$

(1986LA07) have calculated the reaction rates at stellar energies. They are higher than those previously estimated. The higher rate implies a greater production of intermediate A elements in the r-p process in stellar evolution. See also (1987LA14).

4. $^{20}\text{Ne}(p, n)^{20}\text{Na}$ $Q_m = -14.670$

For observed neutron groups see Tables 20.40 in (1978AJ03) and 20.27 here. For preliminary work at $E_p = 120$ MeV see (1983DEZT). See also (1986BA16) and (1983KN05; theor.).

5. $^{20}\text{Ne}(^3\text{He}, t)^{20}\text{Na}$ $Q_m = -13.906$

States derived from triton groups listed in Tables 20.40 in (1978AJ03) and 20.27 here: see, in particular, (1987LA14). [I am grateful to Professor M. Wiescher for communicating these results to me].

Table 20.27: Energy levels of ^{20}Na

E_x (MeV \pm keV)	$J^\pi; T$	$\tau_{1/2}$ or $\Gamma_{\text{c.m.}}$	Decay	Reactions
0	$2^+; 1$	$\tau_{1/2} = 447.9 \pm 2.3$ msec	β^-	1, 4, 5
0.591 ± 12			(γ)	4, 5
0.768 ± 8			(γ)	2, 4, 5
(0.85 ± 50)			(γ)	5
0.958 ± 8			(γ)	4, 5
(1.010 ± 14)			(γ)	4
1.310 ± 10			(γ)	4, 5
1.82 ± 20				5
1.91 ± 20				2, 5
1.98 ± 20				5
2.57 ± 20				5
2.66 ± 20				5
2.88 ± 40				2, 5
2.96 ± 40				5
3.06 ± 40				5
3.16 ± 40				5
4.33 ± 100		^a		2, 5
6.57 ± 50	$0^+; 2$		p	6

^a Broad or unresolved. See also reaction 2.

6. $^{20}\text{Mg}(\beta^+)^{20}\text{Na}$ $Q_m = 10.731$

^{20}Mg decays to $^{20}\text{Na}^*(6.57 \pm 0.05)$ [$J^\pi = 0^+$; $T = 2$]. That state decays by proton emission: see ^{20}Mg .

7. $^{27}\text{Al}(^{20}\text{Ne}, ^{27}\text{Mg})^{20}\text{Na}$ $Q_m = -16.498$

The Δ resonance is very strongly excited in this reaction at $E(^{20}\text{Ne}) = 950 \text{ MeV}/A$ (1986BA16).

^{20}Mg
(Figs. 12 and 13)

^{20}Mg has been populated in the $^{24}\text{Mg}(\alpha, ^8\text{He})$ reaction at $E_\alpha = 127$ and 156 MeV and in the $^{20}\text{Ne}(^3\text{He}, 3n)$ reaction at $E(^3\text{He}) = 70 \text{ MeV}$. The super-allowed decay of ^{20}Mg to the first $T = 2$ ($J^\pi = 0^+$) state of ^{20}Na [$E_x = 6.57 \pm 0.05 \text{ MeV}$] has been reported from observations of the subsequent decay of that state by proton emission [see Fig. 12]. The partial half-life is 95_{-50}^{+80} msec leading to a branching ratio of $(3 \pm 2)\%$ for the super-allowed decay; $\log ft = 3.18$. The results for $A = 20$ are in agreement with the quadratic form of the IMME: see (1978AJ03, 1983AJ01). See also (1985AN28, 1986AN07) and (1983ANZQ; theor.).

^{20}Al , etc.
(Not observed)

See (1972AJ02, 1986AN07) and (1983ANZQ; theor.).

References

(Closed 01 June 1987)

- 1959AJ76 F. Ajzenberg and T. Lauritsen, Nucl. Phys. 11 (1959) 1
- 1968SP01 P. Spilling, H. Gruppelaar, H.F. De vries and A.M.J. Spits, Nucl. Phys. A113 (1968) 395
- 1970PA08 A.D. Panagiotou, Phys. Lett. B31 (1970) 361
- 1972AJ02 F. Ajzenberg-Selove, Nucl. Phys. A190 (1972) 1
- 1973MA14 R.L.Macklin and R.R.Winters, Phys. Rev. C7 (1973) 1766
- 1975HO15 R.E. Horstman, J.L. Eberhardt, H.A. Doubt, C.M.E. Otten and G. Van Middelkoop, Nucl. Phys. A248 (1975) 291
- 1975ME04 L.R. Medsker, H.T. Fortune, R.R. Betts and R. Middleton, Phys. Rev. C11 (1975) 1880
- 1976GAYV D.I. Garber and R.R. Kinsey, BNL 325 (1976)
- 1976MA01 R.E.Marrs, E.G.Adelberger, K.A.Snover and M.D.Cooper, Nucl. Phys. A256 (1976) 1
- 1977FR20 T. Freye, H. Lorenz-Wirzba, B. Cleff, H.P. Trautvetter, C. Rolfs, Z. Phys. A281 (1977) 211
- 1977MA07 R.E.Marrs, E.G.Adelberger and K.A.Snover, Nucl. Phys. A277 (1977) 429
- 1978AJ03 F. Ajzenberg-Selove, Nucl. Phys. A300 (1978) 1
- 1978GR06 E.E. Gross, T.P. Cleary, J.L.C. Ford, D.C. Hensley and K.S. Toth, Phys. Rev. C17 (1978) 1665
- 1978LEZA C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki, R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine, J.K. Tuli and A.B. Buyrn, Table of Isotopes 7th Ed. (1978)
- 1978WI1B Wildenthal, Nukelonika 23 (1978) 459
- 1979BI10 J.H. Billen, Phys. Rev. C20 (1979) 1648
- 1979BR03 H.S. Bradlow, W.D.M. Rae, P.S. Fisher, N.S. Godwin, G. Proudfoot and D. Sinclair, Nucl. Phys. A314 (1979) 171
- 1979GL12 H. Glattli, G.L. Bacchella, M. Fourmond, A. Malinovski, P. Meriel, M. Pinot, P. Roubeau and A. Abragam, J. Phys. (Paris) 40 (1979) 629
- 1979KO26 L. Koester, K. Knopf and W. Waschkowski, Z. Phys. A292 (1979) 95
- 1979YO04 K.C. Young Jr., R.W. Zurmuhle, J.M. Lind, D.P. Balamuth, Nucl. Phys. A330 (1979) 452

- 1980CU09 P. Cuzzocrea, A. De Rosa, G. Inghima, E. Perillo, E. Rosato, M. Sandoli and G. Spadaccini, *Lett. Nuovo Cim.* 28 (1980) 515
- 1980FI01 L.K. Fifield, M.J. Hurst, E.F. Garman, T.J.M. Symons, F. Watt and K.W. Allen, *Nucl. Phys.* A334 (1980) 109
- 1980GR1L Gridnev et al., *Izv. Akad. Nauk SSSR Ser. Fiz.* 44 (1980) 2320
- 1980KR24 V.A. Krutov, *Izv. Akad. Nauk SSSR Ser. Fiz.* 44 (1980) 2239; *Bull. Acad. Sci. USSR Phys. Ser.* 44 (1980) 12
- 1981BL1K Blokhintsev, *Sov. J. Nucl. Phys.* 34 (1981) 525
- 1981BU27 M.W. Buser, *Helv. Phys. Acta* 54 (1981) 439
- 1981CL05 C.F. Clement and S.M. Perez, *J. Phys. (London)* G7 (1981) L293
- 1981DA13 C.A. Davis, *Phys. Rev.* C24 (1981) 1891
- 1981KH1H Khairozzaman, *Proc. Indian Natl. Sci. Acad.* A47 (1981) 264
- 1981MA04 Martoff et al., *Phys. Rev. Lett.* 46 (1981) 891
- 1981OK02 I.P. Okhrimenko and A.I. Steshenko, *Yad. Fiz.* 34 (1981) 873
- 1981SH1A C.S. Shastry and R.K. Satpathy, *Proc. Indian Natl. Sci. Acad.* A47 (1981) 373
- 1981WA1Q Wallace and Woosley, *Astrophys. J. Suppl.* 45 (1981) 389
- 1982AN12 M.S. Antony, *J. Phys. (London)* G8 (1982) 1659
- 1982AN1H Anikina et al., *JETP Lett.* 36 (1982) 331
- 1982AO06 K. Aoki and H. Horiuchi, *Prog. Theor. Phys.* 68 (1982) 2028
- 1982AR20 K.P. Artemov, V.Z. Goldberg, I.P. Petrov, V.P. Rudakov, I.N. Serikov and V.A. Timofeev, *Yad. Fiz.* 36 (1982) 1345; *Sov. J. Nucl. Phys.* 36 (1982) 779
- 1982AV1A I.K. Averyanov, A.I. Golubev, A.A. Sadovoi, *Yad. Fiz.* 35 (1982) 833; *Sov. J. Nucl. Phys.* 35 (1982) 484
- 1982BA1D Barnes, *Essays in Nucl. Astrophys.* (1982) 193
- 1982BE1M Benenson, *Pion Production and Absorption in Nucl.*, Indiana, 1981; *AIP Conf. Proc.* 79 (1982) 381
- 1982BE29 H.W. Becker, W.E. Kieser, C. Rolfs, H.P. Trautvetter and M. Wiescher, *Z. Phys.* A305 (1982) 319
- 1982BI08 B.L. Birbrair and Yu.A. Kalashnikov, *J. Phys. (London)* G8 (1982) 1531
- 1982BLZT L.C. Bland, M. Carchidi, K.S. Dhuga, H.T. Fortune, J. Gilfoyle, R. Gilman, G.S. Stephans, J. Sweet and R. Zurmuhle, *Bull. Amer. Phys. Soc.* 27 (1982) 698, AD8
- 1982BR08 B.A. Brown, *J. Phys. (London)* G8 (1982) 679
- 1982BR24 B.A. Brown, B.H. Wildenthal, W. Chung, S.E. Massen, M. Bernas, A.M. Bernstein, R. Miskimen, V.R. Brown and V. A.Madsen, *Phys. Rev.* C26 (1982) 2247

- 1982CA1A Cameron, *Essays in Nucl. Astrophys.* (1982) 23
- 1982CO1X Cormier, *Ann. Rev. Nucl. Part. Sci.* 32 (1982) 271
- 1982DA28 B. Dasmahapatra, B. Cujec and F. Lahlou, *Lett. Nuovo Cim.* 35 (1982) 345
- 1982DE30 L.C. Dennis, K.M. Abdo, A.D. Frawley and K.W. Kemper, *Phys. Rev. C* 26 (1982) 981
- 1982DE39 P.A. DeYoung, J.J. Kolata, L.J. Satkowiak and M.A. Xapsos, *Phys. Rev. C* 26 (1982) 1482
- 1982DR1E J.P. Draayer and G. Rosensteel, *Nucl. Phys.* A386 (1982) 189
- 1982ES1A Eswaran and Chakrabarty, in *MSU* (1982) 65
- 1982FA1D G. Fai and J. Randrup, *Nucl. Phys.* A381 (1982) 557
- 1982FL04 H. Flocard, P.H. Heenen, S.J. Krieger and M.S. Weiss, *Nucl. Phys.* A391 (1982) 285
- 1982GO1U Gomez, *Proc. 4th Topical School, Granada, 1981* (1982) 163
- 1982GR1T Grotowski et al., *Nukleonika* 27 (1982) 265
- 1982GY02 B. Gyarmati, K.F. Pal and T. Vertse, *Phys. Rev. C* 26 (1982) 2674
- 1982HA43 J.A. Hall and I.P. Johnstone, *Phys. Rev. C* 26 (1982) 1248
- 1982HI1E Hillebrandt and Ober, *Naturwiss.* 69 (1982) 205
- 1982HO05 H.M. Hofmann, W. Zahn, *Czech. J. Phys.* B32 (1982) 288
- 1982HO10 H. Homeyer, M. Burgel, M. Clover, Ch. Egelhaaf, H. Fuchs, A. Gamp, D. Kovar and W. Rauch, *Phys. Rev. C* 26 (1982) 1335
- 1982HO1E Hodgson, *Contemp. Phys.* 23 (1982) 495
- 1982HU06 M. Hugi, J. Lang, R. Muller, J. Sromicki, E. Ungricht, K. Bodek, L. Jarczyk, B. Kamys, A. Strzalkowski and H. Witala, *Phys. Rev. C* 25 (1982) 2403
- 1982JA07 B.K. Jain and N.R. Sharma, *Nucl. Phys.* A388 (1982) 243
- 1982KA12 O. Karban, A.K. Basak, P.M. Lewis and S. Roman, *Phys. Lett.* B112 (1982) 433
- 1982KA17 A. Kapuscik, R. Zybert, M. Pajek, J. Ploskonka and L. Zybert, *Nucl. Phys.* A383 (1982) 530
- 1982KA33 S.B. Kaufman, M.S. Freedman, D.J. Henderson, E.P. Steinberg, B.D. Wilkins, A. Baden, H.H. Gutbrod, M.R. Maier, J. Peter, H.G. Ritter et al., *Phys. Rev. C* 26 (1982) 2694
- 1982KO1C J.J. Kolata, *Proc. Fifth Oaxtepec Symp. on Nucl. Phys., Oaxtepec, Mexico, Jan. 6-8, 1982, UNAM* 5 (1982) 99
- 1982KO1D Y. Kondo and T. Tamura, *Proc. Symp. on Resonances in Heavy Ion Reactions, Bad Honnef, West Germany; 12-15 Oct. 1981* (1982) 314
- 1982KO29 S. Kox, R. Ost and A.J. Cole, *Phys. Lett.* B119 (1982) 303

- 1982LE1P Leeb and Schmid, Czech. J. Phys. B32 (1982) 213
- 1982LE23 M. LeMere, Y. Fujiwara, Y.C. Tang and Q.K.K. Liu, Phys. Rev. C26 (1982) 1847
- 1982LI16 R. Lichtenthaler Jr., A. Lepine-Szily, A.C.C. Villari, W. Mittig, V.J.G. Porto, C.V. Acquadro, Phys. Rev. C26 (1982) 2487
- 1982LO13 M. Lozano, J.I. Escudero and G. Madurga, J. Phys. (London) G8 (1982) 1259
- 1982MA1Q Maurel, Amsel and Nadai, Nucl. Instrum. Meth. Phys. Res. 197 (1982) 1
- 1982MA1V Mach and Rogers, Bull. Amer. Phys. Soc. 27 (1982) 770
- 1982MA25 G. Mairle, G.J. Wagner, P. Grabmayr, K.T. Knopfle, Liu Ken Pao, H. Riedesel, K. Schindler, V. Bechtold, L. Friedrich and P. Ziegler, Nucl. Phys. A382 (1982) 173
- 1982MO15 H. Morgenstern, W. Bohne, K. Grabisch, D.G. Kovar and H. Lehr, Phys. Lett. B113 (1982) 463
- 1982MO1V Mosel, Tanimura and Wolf, Lecture Notes in Phys. 168 (1982) 328
- 1982NA10 S. Nagamiya, H. Hamagaki, P. Hecking, S. Kadota, R. Lombard, Y. Miake, E. Moeller, S. Schnetzer, H. Steiner, I. Tanihata et al., Phys. Rev. Lett. 48 (1982) 1780
- 1982NO1D Norman, Astrophys. J. 258 (1982) 41
- 1982OL03 B.H. Olaniyi, G.A. Beer, A. Fry, J.A. MacDonald, G.R. Mason, A. Olin, R.M. Pearce and P.R. Poffenberger, Nucl. Phys. A384 (1982) 345
- 1982RA1N Ragnarsson and Aberg, Proc. Nucl. Phys. Workshop I.C.T.P. Trieste, Italy, 1981 (1982) 175
- 1982RO06 D.J. Rowe and G. Rosensteel, Phys. Rev. C25 (1982) 3236
- 1982RU05 G. Rufa, Nucl. Phys. A384 (1982) 273
- 1982SA1A D.G. Sargood, Phys. Rept. 93 (1982) 61
- 1982SA27 L.J. Satkowiak, P.A. DeYoung, J.J. Kolata and M.A. Xapsos, Phys. Rev. C26 (1982) 2027
- 1982SC13 J. Schimizu, W. Yokota, T. Nakagawa, Y. Fukuchi, H. Yamaguchi, M. Sato, S. Hanashima, Y. Nagashima, K. Furuno, K. Katori et al., Phys. Lett. B112 (1982) 323
- 1982SC16 E.W. Schmid, S. Saito and H. Fiedeldey, Z. Phys. A306 (1982) 37
- 1982SC20 K.W. Schmid and R. Smith, Nucl. Phys. A381 (1982) 195
- 1982SH1N Shimizu, J. Phys. Soc. Jpn. 51 (1982) 2068
- 1982SH29 D. Shapira, J.L.C. Ford, Jr. and J. Gomez del Campo, Phys. Rev. C26 (1982) 2470
- 1982SH30 S.K. Sharma, V. Potbhare and V.K.B. Kota, Phys. Rev. C26 (1982) 2706
- 1982ST08 Fl. Stancu and D.M. Brink, Phys. Rev. C25 (1982) 2450
- 1982SU06 Y. Suzuki and K.T. Hecht, Nucl. Phys. A388 (1982) 102
- 1982SU1B Y. Suzuki, E.J. Reske and K.T. Hecht, Nucl. Phys. A381 (1982) 77

- 1982VA1E Vanagas, Sabaliauskas and Eriksonas, *Sov. Phys. -Collect.* 22 (1982) 8
- 1982VE02 J.J.M. Verbaarschot, G.A. Timmer and P.J. Brussaard, *Nucl. Phys.* A378 (1982) 280
- 1982VI01 V.E. Viola, Jr., B.B. Back, K.L. Wolf, T.C. Awes, C.K. Gelbke and H. Breuer, *Phys. Rev. C* 26 (1982) 178
- 1982VO04 W. von Oertzen, *Nucl. Phys.* A387 (1982) 93c
- 1982WA13 G.J. Wagner, P. Grabmayr and H.R. Schmidt, *Phys. Lett.* B113 (1982) 447
- 1982WA1K Wang, Westendorp and Saris, *FOM-Inst. for Atomic and Molecular Phys.* (1982) 37
- 1982WO1A Woosley and Weaver, *Essays in Nucl. Astrophys.* (1982) 377
- 1982WO1E K.L. Wolf, H.H. Gutbrod, W.G. Meyer, A.M. Poskanzer, A. Sandoval, R. Stock, J. Gosset, J.-C. Jourdain, C.H. King, G. King et al., *Phys. Rev. C* 26 (1982) 2572
- 1982ZE1A Zeng, Lin and Yang, *Chin. Phys.* 2 (1982) 707
- 1983AB1G Abutalybova, Kiselev and Pokrovskii, *Sov. J. Nucl. Phys.* 38 (1983) 865
- 1983AD1D Adelberger, *Proc. Int. Conf. on Nucl. Phys., Florence, Aug.-Spet. 1983, Vol. 2, Eds., P. Blasi and R.A. Ricci* (1983) 499
- 1983AH1A Ahmed and Beres, *Phys. Rev. C* 28 (1983) 660
- 1983AJ01 F. Ajzenberg-Selove, *Nucl. Phys.* A392 (1983) 1; *Erratum Nucl. Phys.* A413 (1984) 168
- 1983AL01 M. Albinska, P. Belery, Th. Delbar, Y. El Masri, Gh. Gregoire, C. Michel, J. Vervier, J. Albinski, K. Grotowski, S. Kopta et al., *Phys. Rev. C* 27 (1983) 207
- 1983AL23 J. Almeida and F. Kappeler, *Astrophys. J.* 265 (1983) 417
- 1983AN1L Anikina et al., *Sov. J. Nucl. Phys.* 38 (1983) 901; *Erratum Sov. J. Nucl. Phys.* 39 (1984) 1024
- 1983ANZQ Y. Ando, M. Uno and M. Yamada, *JAERI-M-83-025* (1983)
- 1983AO03 K. Aoki, H. Horiuchi, *Prog. Theor. Phys. (Kyoto)* 69 (1983) 857
- 1983AR07 F. Arickx, J. Broeckhove, E. Caurier, E. Deumens and P. Van Leuven, *Nucl. Phys.* A398 (1983) 467
- 1983AR1J Arima et al., in *Florence* (1983) 207
- 1983BA22 H.W. Barz, T.S. Biro, B. Lukacs and J. Zimanyi, *Z. Phys.* A311 (1983) 311
- 1983BA2J Baldo et al., *Proc. Int. Conf. on Heavy Ion Phys. and Nucl. Phys., Eds., Catania, A. Agodi and C. Villi* (1983) 77
- 1983BA50 D. Baye and P. Descouvemont, *Nucl. Phys.* A407 (1983) 77
- 1983BE02 E. Betak and V.D. Toneev, *J. Phys. (London)* G9 (1983) L47
- 1983BE19 U.E.P. Berg, K. Ackermann, K. Bangert, R. Stock and K. Wienhard, *Phys. Rev. C* 27 (1983) 2981

- 1983BE55 J.J. Bevelacqua, *Indian J. Phys.* A57 (1983) 26
- 1983BI13 J.R. Birkelund and J.R. Juizenga, *Ann. Rev. Nucl. Part. Sci.* 33 (1983) 265
- 1983BI1C Bizzeti, *Riv. Nuovo Cim.* 6 (1983) 1
- 1983BI1F Bistrenko, in *Moscow* (1983) 186
- 1983BR1P Brown, Radhi and Wildenthal, *Phys. Rept.* 101 (1983) 313
- 1983BR1R P. Braun-Munzinger, *Nucl. Phys.* A409 (1983) 31
- 1983BR1V Breitschaft, Canto and Schechter, *Rev. Bras. Fis.* 13 (1983) 328
- 1983BR29 B.A. Brown and B.H. Wildenthal, *Phys. Rev.* C28 (1983) 2397
- 1983CA1F G.T. Caskey, PhD. Thesis, Univ. Wisconsin (1983)
- 1983CE1A Cereda, de Leo, Micheletti and Pignanelli, in *Florence* (1983) 234
- 1983CH1B Chant, *AIP Conf. Proc.* 97 (1983) 205
- 1983CI08 O. Civitarese, B.V. Carlson, M.S. Hussein and A. Szanto de Toledo, *Phys. Lett.* B125 (1983) 22
- 1983CL01 E.T.H. Clifford, J.C. Hardy, H. Schmeing, R.E. Azuma, H.C. Evans, T. Faestermann, E. Hagberg, K.P. Jackson, V.T. Koslowsky, U.J. Schrewe et al., *Phys. Rev. Lett.* 50 (1983) 23
- 1983CO08 J. Cohen, *J. Phys.* G9 (1983) 621
- 1983COZV J. Cook, *Bull. Amer. Phys. Soc.* 28 (1983) 662, AY7b
- 1983CS02 J. Cseh, *Phys. Rev.* C27 (1983) 2991
- 1983CSZX J. Csikai, in *Florence* (1983) 451
- 1983DE26 P.A. De Young, J.J. Kolata, L.J. Satkowiak and M.A. Xapsos, *Phys. Rev.* C28 (1983) 692
- 1983DE2J Deconnick and Vanoystaeyen, *Nucl. Instrum. Meth. Phys. Res.* 218 (1983) 165
- 1983DE32 P. Descouvemont and D. Baye, *Phys. Lett.* B127 (1983) 286
- 1983DEZT R.P. DeVito, C.C. Foster, C.D. Goodman, T.N. Taddeucci, S.M. Austin, A. Galonsky and U.E.P. Berg, *Bull. Amer. Phys. Soc.* 28 (1983) 733, HG1
- 1983DEZW L.C. Dennis, A.D. Frawley and J.F. Mateja, *Bull. Am. Phys. Soc.* 28 (1983) 669, BF1
- 1983DR03 J.P. Draayer and G. Rosensteel, *Phys. Lett.* B124 (1983) 281
- 1983DR04 J.P. Draayer and G. Rosensteel, *Phys. Lett.* B125 (1983) 237
- 1983DU13 G.G. Dussel, A.O. Gattone and E.E. Maqueda, *Phys. Rev. Lett.* 51 (1983) 2366
- 1983FI02 L.K. Fifield, W.N. Catford, S.H. Chew, E.F. Garman, D.M. Pringle, K.W. Allen and J. Lowe, *Nucl. Phys.* A394 (1983) 1
- 1983FOZV H.T. Fortune, K.S. Dhuga, M. Carchidi, G.P. Gilfoyle, R. Gilman, J. Sweet and A.H. Wuosmaa, *Bull. Amer. Phys. Soc.* 28 (1983) 995, EB7

- 1983FR14 A.D. Frawley, J.D. Fox, L.C. Dennis, K.W. Kemper and N.R. Fletcher, Phys. Rev. C27 (1983) 2482
- 1983FR1A Friedman and Lynch, Phys. Rev. C28 (1983) 950
- 1983FR1G Friedman, Phys. Rev. C27 (1983) 569
- 1983FU1D Fujiwara, Tang and Horiuchi, Prog. Theor. Phys. 70 (1983) 809
- 1983GA01 A. Gavron, J.R. Beene, B. Cheynis, R.L. Ferguson, F.E. Obenshain, F. Plasil, G.R. Young, G.A. Petitt, C.F. Maguire, D.G. Sarantites et al., Phys. Rev. C27 (1983) 450
- 1983GE12 P.M. Gensini, Lett. Nuovo Cim. 38 (1983) 469
- 1983GM1A Gmitro, Kissener, Truol and Eramzhyan, Sov. J. Part. Nucl. 14 (1983) 323
- 1983GO13 J. Gomez del Campo and G.R. Satchler, Phys. Rev. C28 (1983) 952
- 1983GR26 D.H.E. Gross and M.C. Nemes, Phys. Lett. B130 (1983) 131
- 1983HA1P Hashimoto, Hanawa and Sugimoto, Publ. Astron. Soc. Jpn. 35 (1983) 1
- 1983HA1V Hardy, Report AECL-8258 (1983)
- 1983HE1B W. Henning, Nucl. Phys. A400 (1983) 295
- 1983HI06 M.M. Hindi, J.H. Thomas, D.C. Radford and P.D. Parker, Phys. Rev. C27 (1983) 2902
- 1983HO1D Horiuchi, KUNS 656 (1983)
- 1983HO1F Horiuchi, Prog. Theor. Phys. 69 (1983) 886
- 1983HU12 Hungerford et al., Z. Phys. A313 (1983) 339
- 1983HU1E Huie et al., Phys. Rev. C27 (1983) 439
- 1983IN1B Inoue and Isoya, Proc. RCNP Int. Symp. on Light Ion Reaction Mechanism, Osaka (1983) 482
- 1983IS03 B.S. Ishkhanov, I.M. Kapitonov and A.V. Shumakov, Nucl. Phys. A394 (1983) 131
- 1983IS1E M. Ishihara, Nucl. Phys. A400 (1983) 153
- 1983JA05 U. Jahnke, G. Ingold, H. Homeyer, M. Burgel, Ch. Egelhaaf, H. Fuchs and D. Hilscher, Phys. Rev. Lett. 50 (1983) 1246
- 1983JA09 L. Jarczyk, B. Kamys, Z. Rudy, A. Strzalkowski, H. Witala, M. Hugi, J. Lang, R. Muller, J. Sromicki and H.H. Wolter. Phys. Rev. C28 (1983) 700
- 1983JI04 Jiang Chenglie, Han Shukui, Guo Qingjiang and Li Qingli, Chin. Phys. 3 (1983) 675; Chin. J. Nucl. Phys. 5 (1983) 8
- 1983KA32 K. Kar, Phys. Rev. C28 (1983) 2446
- 1983KAZF J.V. Kane, Bull. Am. Phys. Soc. 28 (1983) 661, AY5a
- 1983KN01 N. Kniest, E. Huttel, J. Gunzl, G. Clausnitzer, P.G. Bizzeti, P.R. Maurenzig and N. Taccetti, Phys. Rev. C27 (1983) 906
- 1983KN05 Knupfer and Metsch, Phys. Rev. C27 (1983) 2487

- 1983KO01 R.L. Kozub, J. Lin, J.F. Mateja, C.J. Lister, D.J. Millener, J.W. Olness and E.K. Warburton, Phys. Rev. C27 (1983) 158
- 1983KO31 Y. Kondo, B.A. Robson and R. Smith, Nucl. Phys. A410 (1983) 289
- 1983LA25 K. Langanke, Phys. Lett. B131 (1983) 21
- 1983LE03 P.M. Lewis, A.K. Basak, J.D. Brown, P.V. Drumm, O. Karban, E.C. Pollacco and S. Roman, Nucl. Phys. A395 (1983) 204
- 1983LE1F Letaw, Phys. Rev. C28 (1983) 2178
- 1983LE28 Letaw, Silberberg and Tsao, Astrophys. J. 51 (1983) 271
- 1983LIZW P.M. Lister, F.D. Brooks and J.M. Nelson, Bull. Am. Phys. Soc. 28 (1983) 658, AX5
- 1983LO05 J. Lomnitz-Adler and P. Van Isacker, Phys. Rev. C27 (1983) 1356
- 1983MA16 C.J. Martoff, J.A. Bistirlich, C.W. Clawson, K.M. Crowe, M. Koike, J.P. Miller, S.S. Rosenblum, W.A. Zajc, H.W. Baer, A.H. Wapstra et al., Phys. Rev. C27 (1983) 1621
- 1983MA23 P. Manakos, T. Fliessbach and H. Walliser, Phys. Rev. C27 (1983) 2930
- 1983MA29 N. Marquardt, Phys. Rev. C28 (1983) 202
- 1983MA68 S. Marcos, H. Flocard and P.H. Heenen, Nucl. Phys. A410 (1983) 125
- 1983ME13 M.C. Mermaz, F. Auger and B. Fernandez, Phys. Rev. C28 (1983) 1587
- 1983ME1R Mewalst and Wiedenbeck, 18th Int. Cosmic Ray Conf., India (1983) 206
- 1983MI22 F. Michel, J. Albinski, P. Belery, Th. Delbar, Gh. Gregoire, B. Tasiaux and G. Reide-meister, Phys. Rev. C28 (1983) 1904
- 1983MO13 H. Morgenstern, W. Bohne, K. Grabisch, H. Lehr and W. Stoffler, Z. Phys. A313 (1983) 39
- 1983NA04 W.J. Naude, H.S. Bradlow, O. Dietzsch, A.A. Pilt, W.D.M. Rae and D. Sinclair, Z. Phys. A311 (1983) 297
- 1983NG01 van Sen et al., Phys. Rev. C27 (1983) 194
- 1983OH1B Ohta, Hatogai and Okai, in Florence (1983) 558
- 1983OH1C Ohta, Hatogai, Okai and Abe, RIFP-524 (1983)
- 1983OS03 A. Osman, Int. J. Theor. Phys. 22 (1983) 341
- 1983PI03 A.A. Pilt, Nuovo Cim. A74 (1983) 185
- 1983PO09 N. Poffe, N. Rowley and R. Lindsay, Nucl. Phys. A410 (1983) 498
- 1983RA1E Radi et al., Phys. Rev. C27 (1983) 606
- 1983RA1J J.O. Rasmussen, Nucl. Phys. A400 (1983) 383
- 1983RO1E Romanov and Grechukhin, in Moscow (1983) 249
- 1983RO1H E. Roeckl, Nucl. Phys. A400 (1983) 131

- 1983SA07 M. Sasagase, M. Sato, S. Hanashima, K. Furuno, Y. Nagashima, Y. Tagishi, S.M. Lee and T. Mikumo, Phys. Rev. C27 (1983) 2630
- 1983SC08 H. Schultheis and R. Schultheis, Phys. Rev. C27 (1983) 1367
- 1983SC1M Scott, Int. Conf. on Heavy Ion Phys. and Nucl. Phys., Catania, Italy (1983); MSUCL-422
- 1983SH1Z Shimoura et al., in Florence (1983) 661
- 1983SH25 D. Shapira, D. DiGregorio, J. Gomez del Campo, R.A. Dayras, J.L.C. Ford, Jr., A.H. Snell, P.H. Stelson, R.G. Stokstad and F. Pougheon, Phys. Rev. C28 (1983) 1148
- 1983SH26 T. Shimoda, S. Shimoura, T. Fukuda, M. Tanaka, H. Ogata, I. Miura, E. Takada, M.-K. Tanaka, K. Takimoto and K. Katori, J. Phys. (London) G9 (1983) L199
- 1983SH32 J. Shurpin, T.T.S. Kuo and D. Strottman, Nucl. Phys. A408 (1983) 310
- 1983SI01 B. Sinha, Phys. Rev. Lett. 50 (1983) 91
- 1983SI1B Simpson, Ann. Rev. Nucl. Part. Sci. 33 (1983) 323
- 1983SI1L R.H. Siemssen, Nucl. Phys. A409 (1983) 73
- 1983SM04 Yu.F. Smirnov and Yu.M. Tchuvilsky, Czech. J. Phys. B33 (1983) 1215
- 1983SM1B Smida and Fellah, in Florence (1983) 542
- 1983SP04 K.-H. Speidel, V. Mertens, M. Knopp, W. Trolenberg, H. Neuburger, J. Gerber and K. Bharuth-Ram, Hyperfine Interactions 15/16 (1983) 55
- 1983SU11 V.B. Subbotin, V.M. Semjonov, K.A. Gridnev and E.F. Hefter, Phys. Rev. C28 (1983) 1618; Erratum Phys. Rev. C30 (1984) 738
- 1983TO1H V.D. Toneev and K.K. Gudima, Nucl. Phys. A400 (1983) 173
- 1983TO1L Townsend, Can. J. Phys. 61 (1983) 93
- 1983VA16 M.G. Vassanji and D.J. Rowe, Phys. Lett. B127 (1983) 1
- 1983VA21 V.V. Vanagas, L.Yu. Sabalyauskas and K.M. Eriksonas, Izv. Akad. Nauk SSSR Ser. Fiz. 47 (1983) 963
- 1983WA04 H. Walliser and T. Fliessbach, Nucl. Phys. A394 (1983) 387
- 1983WE1A Webber, Brautigam, Kish and Schrier, Bull. Amer. Phys. Soc. 28 (1983) 754
- 1983WE1C Webber, Brautigam, Kish and Schrier, 18th Int. Cosmic Ray Conf., India (1983) 202
- 1983WI1A Wilczynski, Proc. Int. Conf. on Nucl. Phys., Aug.-Sept. 1983, Florence; Eds., P. Blasi and R.A. Ricci; Vol. 2 (1983) 305
- 1984AD1E Adelberger, AIP Conf. Proc. 123 (1984) 300
- 1984AM03 K. Amos, I. Morrison and R. Smith, J. Phys. (London) G10 (1984) 331
- 1984AM07 K. Amos and W. Bauhoff, Nucl. Phys. A424 (1984) 60
- 1984AN1B Anagnostatos, in Panic (1984) I56

- 1984AN1H Anikina et al., Sov. J. Nucl. Phys. 40 (1984) 311
- 1984AP03 B. Apagyi and W. Scheid, J. Phys. (London) G10 (1984) 791
- 1984AR20 Artemov K.P. Artemov, V.Z. Goldberg, M.S. Golovkov, I.P. Petrov, V.P. Rudakov, I.N. Serikov, V.A. Timofeev, H.U. Gersch and E. Hentschel, Phys. Lett. B149 (1984) 325
- 1984AS1D Asai, Bando and Sano, Phys. Lett. B145 (1984) 19
- 1984BA1R V.N. Baturin, A.V. Khanzadeev, V.P. Koptev, E.M. Maev, M.M. Makarov, V.V. Nelyubin, G.V. Shcherbakov and V.V. Sulimov, in Panic (1984) I11
- 1984BA34 F. Baros and S. Regnier, J. Phys. (Paris) 45 (1984) 855
- 1984BE22 C. Beck, F. Haas, R.M. Freeman, B. Heusch, J.P. Coffin, G. Guillaume, F. Rami and P. Wagner, Phys. Rev. C29 (1984) 1942
- 1984BE26 U.E.P. Berg, K. Ackermann, K. Bangert, C. Blasing, W. Naatz, R. Stock, K. Wienhard, M.K. Brussel, T.E. Chapuran and B.H. Wildenthal, Phys. Lett. B140 (1984) 191
- 1984BI06 J.R. Birkelund and J.R. Huizenga, Phys. Rev. C30 (1984) 401
- 1984BL14 G.S. Blanpied, G.A. Balchin, G.E. Langston, B.G. Ritchie, M.L. Barlett, G.W. Hoffmann, J.A. McGill, M.A. Franey, M. Gazzaly and B.H. Wildenthal, Phys. Rev. C30 (1984) 1233
- 1984BR15 T. Bright, D. Ballon, R.J. Saxena, Y. Niv and N. Benczer-Koller, Phys. Rev. C30 (1984) 696
- 1984BR25 B.A. Brown, C.R. Bronk and P.E. Hodgson, J. Phys. (London) G10 (1984) 1683
- 1984BU01 V.V. Burov, V.M. Dubovik, S.G. Kadmsky, Yu.M. Tchuvilsky and L.A. Tosunyan, J. Phys. (London) G10 (1984) L21
- 1984BU1R Burtebaev, Duisebaev and Kniazkov, in Alma Ata (1984) 349
- 1984BUZO I.E. Burkova and A.A. Yatis, in Alma Ata (1984) 564
- 1984BY02 A.V. Bystrenko, Yad. Fiz. 40 (1984) 389; Sov. J. Nucl. Phys. 40 (1984) 247
- 1984CA08 W.N. Catford, E.F. Garman and L.K. Fifield, Nucl. Phys. A417 (1984) 77
- 1984CA09 T.A. Carey, P.G. Roos, N.S. Chant, A. Nadasen and H.L. Chen, Phys. Rev. C29 (1984) 1273
- 1984CH1V W.H. Chung and J.H. Lee, New Phys. (Korea) 24 (1984) 267
- 1984CHZU Y. Chan, C. Albiston, M. Bantel, A. Budzanowski, P. Countryman, D. Digregorio, K. Grotowski, R. Planeta, R. Stokstad, S. Wald et al., Bull. Amer. Phys. Soc. 29 (1984) 1048, DB7
- 1984CO1H Cook, Stone and Vogt, Astrophys. J. 279 (1984) 827
- 1984CU1B Cujec, Fizika 16 (1984) 3
- 1984DA1B Danilin and Zhukov, in Alma Ata (1984) 460
- 1984DE1Q Denisov, Sov. J. Nucl. Phys. 39 (1984) 522

- 1984DE53 P. De Bievre, M. Gallet, N.E. Holden and I.L. Barnes, *J. Phys. Chem. Ref. Data* 13 (1984) 809
- 1984DH02 K.S. Dhuga, H.T. Fortune, M. Carchidi, G.P. Gilfoyle, R. Gilman, J.W. Sweet and A. Wuosmaa, *Phys. Rev. C* 29 (1984) 1566
- 1984DR01 J.P. Draayer, K.J. Weeks and G. Rosensteel, *Nucl. Phys. A* 413 (1984) 215
- 1984EF03 V.P. Efrosinin and D.A. Zaikin, *Yad. Fiz.* 39 (1984) 1135; *Sov. J. Nucl. Phys.* 39 (1984) 717
- 1984EL1D Eliseev and Yuldashev, *Sov. J. Nucl. Phys.* 40 (1984) 601
- 1984FI17 D.J. Fields, W.G. Lynch, C.B. Chitwood, C.K. Gelbke, M.B. Tsang, H. Utsunomiya and J. Aichelin, *Phys. Rev. C* 30 (1984) 1912
- 1984FO16 H.T. Fortune and B.H. Wildenthal, *Phys. Rev. C* 30 (1984) 1063
- 1984FO1A W.A. Fowler, *Rev. Mod. Phys.* 56 (1984) 149
- 1984FR13 H. Friedrich, *Phys. Lett. B* 146 (1984) 135
- 1984FR1A P. Frobrich, *Phys. Rept.* 116 (1984) 337
- 1984GA22 M. Gai, G.M. Berkowitz, P. Braun-Munzinger, C.M. Jachcinski, C.E. Ordonez, T.R. Renner and C.D. Uhlhorn, *Phys. Rev. C* 30 (1984) 925
- 1984GO04 A. Gokmen, G.J. Mathews and V.E. Viola, Jr., *Phys. Rev. C* 29 (1984) 1606
- 1984GR03 D.P. Grechukhin and S.V. Romanov, *Yad. Fiz.* 39 (1984) 48; *Sov. J. Nucl. Phys.* 39 (1984) 28
- 1984GR08 R.E.L. Green, R.G. Korteling and K.P. Jackson, *Phys. Rev. C* 29 (1984) 1806
- 1984GR29 K.A. Gridnev, Kh.M. Omer, V.M. Semenov and V.B. Subbotin, *Izv. Akad. Nauk SSSR Ser. Fiz.* 48 (1984) 963
- 1984GU09 S.K. Gupta and S. Kailas, *Z. Phys. A* 317 (1984) 75
- 1984HA14 P. Halse, J.P. Elliott and J.A. Evans, *Nucl. Phys. A* 417 (1984) 301
- 1984HA43 Q. Haider and B. Cujec, *Nucl. Phys. A* 429 (1984) 116
- 1984HA53 Q. Haider and F.B. Malik, *At. Data Nucl. Data Tables* 31 (1984) 185
- 1984HI1A A.S. Hirsch, A. Bujak, J.E. Finn, L.J. Gutay, R.W. Minich, N.T. Porile, R.P. Scharenberg and B.C. Stringfellow, *Phys. Rev. C* 29 (1984) 508
- 1984HO23 H. Homeyer, M. Burgel, Ch. Egelhaaf, H. Fuchs and G. Thoma, *Z. Phys. A* 319 (1984) 143
- 1984HU1E Hunyadi, Szoghy and Cujec, *Nucl. Tracks Radiat. Meas.* 8 (1984) 525
- 1984IA1A Iachello, *Nuovo Cim.* A81 (1984) 130
- 1984IN03 M. Inoue and S.E. Koonin, *Phys. Rev. C* 30 (1984) 175
- 1984IN04 T. Inoue, *J. Phys. Soc. Jpn.* 53 (1984) 4158

- 1984JA15 H.R. Jaqaman and L. Zamick, Phys. Rev. C30 (1984) 1719
- 1984JE02 A.S. Jensen, P.G. Hansen and B. Jonson, Nucl. Phys. A431 (1984) 393
- 1984KA11 H. Kazama, K. Kato and H. Tanaka, Prog. Theor. Phys. 71 (1984) 215
- 1984KA1J Karol, Science 226 (1984) 1425
- 1984KL06 H.V. Klapdor, J. Metzinger and T. Oda, At. Data Nucl. Data Tables 31 (1984) 81
- 1984KN1A N. Kniest, E. Huttel, E. Pfaff, G. Reiter and G. Clausnitzer, in Panic (1984) H9
- 1984KO12 S. Kox, A. Gamp, R. Cherkaoui, A.J. Cole, N. Longequeue, J. Menet, C. Perrin and J.B. Viano, Nucl. Phys. A420 (1984) 162
- 1984KO13 Y. Kondo and T. Tamura, Phys. Rev. C30 (1984) 97
- 1984KO25 I. Koenig, D. Fick, S. Kossionides, P. Egelhof, K.-H. Mobius and E. Steffens, Z. Phys. A318 (1984) 135
- 1984LA18 K. Langanke, Z. Phys. A317 (1984) 325
- 1984LA1N Laymon, Balamuth and Garman, Bull. Amer. Phys. Soc. 29 (1984) 630
- 1984LE06 H. Lehr, W. Bohne, K. Grabisch, H. Morgenstern and W. Von Oertzen, Nucl. Phys. A415 (1984) 149
- 1984LE19 R.J. Ledoux, C.E. Ordonez, M.J. Bechara, H.A. Al-Juwair, G. Lavelle and E.R. Cosman, Phys. Rev. C30 (1984) 866
- 1984LO05 N. Lo Iudice and F. Palumbo, Phys. Rev. C30 (1984) 360
- 1984MAZT F.B. Malik and Q. Haider, Bull. Am. Phys. Soc. 29 (1984) 1047
- 1984ME10 M.C. Mermaz, Nuovo Cim. A81 (1984) 291
- 1984MO08 S. Mordechai and H.T. Fortune, Phys. Rev. C29 (1984) 1765
- 1984MU04 T. Murakami, E. Ungricht, N. Takahashi, Y.-W. Lui, Y. Mihara, R.E. Neese, E. Takada, D.M. Tanner, R.E. Tribble and K. Nagatani, Phys. Rev. C29 (1984) 847
- 1984NA12 Y. Nagame, H. Nakahara, K. Sueki, H. Kudo, I. Kohno and M. Yanokura, Z. Phys. A317 (1984) 31
- 1984NA1D Nagamiya, Randrup and Symons, Ann. Rev. Nucl. Part. Sci. 34 (1984) 155
- 1984NA21 R. Nayak, A. Faessler, H. Muther and A. Watt, Nucl. Phys. A427 (1984) 61
- 1984NI1D Ning, Gao, He and Chen, Chin. J. Nucl. Phys. 6 (1984) 105
- 1984NO1B Nomoto, Thielemann and Wheeler, Astrophys. J. 279 (1984) L23
- 1984OK04 S. Okabe, Nucl. Phys. A427 (1984) 87
- 1984PA04 P. Park, J. Carvalho, M. Vassanji, D.J. Rowe and G. Rosensteel, Nucl. Phys. A414 (1984) 93
- 1984PA18 G. Palla and W. Oelert, Phys. Rev. C30 (1984) 1331
- 1984PO11 D.N. Poenaru and M. Ivascu, J. Phys. (Paris) 45 (1984) 1099

- 1984PR09 D. Provoost, F. Grummer, K. Goeke, P.-G. Reinhard, Nucl. Phys. A431 (1984) 139
- 1984RA10 W.D.M. Rae, A.J. Cole, B.G. Harvey and R.G. Stokstad, Phys. Rev. C30 (1984) 158
- 1984RA13 P.K. Rath, H. Muther, A. Polls, A. Faessler, B. Metsch and D. Zwarts, Nucl. Phys. A427 (1984) 511
- 1984RE14 S.M. Read and V.E. Viola, Jr., At. Data Nucl. Data Tables 31 (1984) 359
- 1984RI01 H.T. Richards, Phys. Rev. C29 (1984) 276
- 1984RI06 S.R. Riedhauser, Phys. Rev. C29 (1984) 1961
- 1984RI07 H.T. Richards, G. Caskey, J.H. Billen, S.R. Riedhauser and D.J. Steck, Phys. Rev. C29 (1984) 2332
- 1984RO04 R.G.H. Robertson, P. Dyer, R.C. Melin, T.J. Bowles, A.B. McDonald, G.C. Ball, W.G. Davies and E.D. Earle, Phys. Rev. C29 (1984) 755
- 1984SA28 V.S. Sadkovsky, G.A. Feofilov, A.E. Denisov, R.P. Kolalis and L. Peres Tamaio, Izv. Akad. Nauk SSSR Ser. Fiz. 48 (1984) 995
- 1984SA37 M. Sakai, At. Data Nucl. Data Tables 31 (1984) 399
- 1984SC01 K.W. Schmid, F. Grummer and A. Faessler, Phys. Rev. C29 (1984) 308
- 1984SC41 K.W. Schmid, F. Grummer and A. Faessler, Nucl. Phys. A431 (1984) 205
- 1984SE02 A.G. Seamster, E.B. Norman, D.D. Leach, P. Dyer and D. Bodansky, Phys. Rev. C29 (1984) 394
- 1984SE20 V.M. Semjonov, K.A. Gridnev, E.F. Hefter, H.M. Omar, S. Saad and V.B. Subbotin, Nuovo Cim. A84 (1984) 89
- 1984SH1T Shen, Qiao, Zhu and Zhan, Phys. Energ. Fortis Phys. Nucl. 8 (1984) 732
- 1984SH2D Shalaby, El-Degheidy, Khalil and Morgan, Acta Phys. Slovaca 34 (1984) 3
- 1984SP06 K.-H. Speidel, Nuovo Cim. A81 (1984) 231
- 1984SR01 D.K. Srivastava and H. Rebel, Z. Phys. A316 (1984) 225
- 1984ST1B Stokstad, Comments Nucl. Part. Phys. 13 (1984) 231
- 1984SU02 P. Suebka, C.K. Chan, Z.C. Kang and P. Lu, Phys. Rev. C29 (1984) 1088
- 1984TO02 M. Tohyama, R. Kaps, D. Masak and U. Mosel, Phys. Lett. B136 (1984) 226
- 1984TR1C Truran, Ann. Rev. Nucl. Part. Sci. 34 (1984) 53
- 1984TS03 M.B. Tsang, D.R. Klesch, C.B. Chitwood, D.J. Fields, C.K. Gelbke, W.G. Lynch, H. Utsunomiya, K. Kwiatkowski, V.E. Viola, Jr. and M. Fatyga, Phys. Lett. B134 (1984) 169
- 1984UM04 K. Umeda, T. Yamaya, T. Suehiro, K. Takimoto, R. Wada, E. Takada, S. Shimoura, A. Sakaguchi, S. Murakami, M. Fukada et al., Nucl. Phys. A429 (1984) 88
- 1984VO1H von Egidy et al., in Konxville (1984) 37

- 1984WE04 E. Wesolowski, *J. Phys. (London)* G10 (1984) 321
- 1984WE13 E. Wesolowski, *Acta Phys. Pol.* B15 (1984) 559
- 1984WI1E Wilson et al., *Health Phys.* 46 (1984) 1101
- 1984ZH07 M.V. Zhukov, *Yad. Fiz.* 40 (1984) 892; *Sov. J. Nucl. Phys.* 40 (1984) 568
- 1985AD1A Adelberger and Haxton, *Ann. Rev. Nucl. Part. Sci.* 35 (1985) 501
- 1985AG1A M.M. Aggarwal and P.L. Jain, *Phys. Rev.* C31 (1985) 1233
- 1985AI1C Aichelin and Ko, *Phys. Rev. Lett.* 55 (1985) 2661
- 1985AJ01 F. Ajzenberg-Selove, *Nucl. Phys.* A433 (1985) 1; Erratum *Nucl. Phys.* A449 (1986) 155
- 1985AL21 T.K. Alexander, B. Castel and I.S. Towner, *Nucl. Phys.* A445 (1985) 189
- 1985AN16 M.V. Andres, J.M. Quesada, M. Lozano and G. Madurga, *Nucl. Phys.* A443 (1985) 380
- 1985AN17 M.S. Antony, J. Britz, J.B. Bueb and V.B. Ndocko-Ndongue, *Nuovo Cim.* A88 (1985) 265
- 1985AN1B Anne et al., in *Visby* (1985) 31
- 1985AN28 M.S. Antony, J. Britz, J.B. Bueb and A. Pape, *At. Data Nucl. Data Tables* 33 (1985) 447
- 1985AR1A Arnett and Thielemann, *Astrophys. J.* 295 (1985) 589
- 1985AZ1A Azimov et al., *Czech. J. Phys.* 35 (1985) 832
- 1985BA1Q Baye and Descouvemont, *Ann. Phys.* 165 (1985) 115
- 1985BA2T Balestra et al., *Nucl. Instrum. Meth. Phys. Res.* A234 (1985) 30
- 1985BA2W Baye and Descouvemont, *Lecture Notes in Phys.* 219 (1985) 138
- 1985BA43 Z. Basrak, W. Tiereth, N. Bischof, H. Frohlich, B. Nees, E. Nieschler and H. Voit, *Phys. Rev.* C32 (1985) 910
- 1985BE1A M. Beckerman, *Phys. Rept.* 129 (1985) 145
- 1985BE37 C. Beck, R.M. Freeman, F. Haas, B. Heusch and J.J. Kolata, *Nucl. Phys.* A443 (1985) 157
- 1985BE40 C. Beck, F. Haas, R.M. Freeman, B. Heusch, J.P. Coffin, G. Guillaume, F. Rami and P. Wagner, *Nucl. Phys.* A442 (1985) 320
- 1985BI01 B.L. Birbrair, V.N. Fomenko, A.B. Gridnev and Yu.A. Kalashnikov, *J. Phys. (London)* G11 (1985) 471
- 1985BR29 B.A. Brown and B.H. Wildenthal, *At. Data Nucl. Data Tables* 33 (1985) 347
- 1985CA09 G. Caskey, *Phys. Rev.* C31 (1985) 717
- 1985CH11 Chengqun Gao, Pingzhi Ning and Guozhu He, *Nucl. Phys.* A438 (1985) 281

1985CL1A F.E. Close, Nucl. Phys. A446 (1985) 273
 1985CU1A B. Cujec, Lecture Notes in Phys. 219 (1985)108
 1985CU1F J. Cugnon and J. Vandermeulen, Nucl. Phys. A445 (1985) 717
 1985DW1A Dwyer and Meyer, Astrophys. J. 294 (1985) 441
 1985DY05 P. Dyer, D. Bodansky, D.D. Leach, E.B. Norman and A.G. Seamster, Phys. Rev. C32 (1985) 1873
 1985FL1B Fletcher et al., Bull. Amer. Phys. Soc. 30 (1985) 1768
 1985FO07 H.T.Fortune and R.Eckman, Phys. Rev. C31 (1985) 2076
 1985FR13 K.A. Frankel, J.A. Bistirlich, R. Bossingham, H.R. Bowman, K.M. Crowe, C.J. Martoff, D.L. Murphy, J.O. Rasmussen, J.P. Sullivan, E. Yoo et al., Phys. Rev. C32 (1985) 975
 1985GA1G Gao, Ning and He, Chin. Phys. Lett. 2 (1985) 125
 1985GA1T Gale and Das Gupta, Phys. Lett. B162 (1985) 35
 1985GAZT S.B. Gazes, C.R. Albiston, Y. Chan, H.R. Schmidt, R.G. Stokstad and R. Kamermans, Bull. Amer. Phys. Soc. 30 (1985) 1280, EB8
 1985GR1A Grenacs, Ann. Rev. Nucl. Part. Sci. 35 (1985) 455
 1985GU08 F. Guzman Martinez and R. Reif, Nucl. Phys. A436 (1985) 294
 1985GU1J K.K. Gudima and V.D. Toneev, Yad. Fiz. 42 (1985) 645; Sov. J. Nucl. Phys. 42 (1985) 409
 1985HA11 J.S. Hanspal, R.J. Griffiths and N.M. Clarke, Phys. Rev. C31 (1985) 1138
 1985HA15 P. Halse, Phys. Lett. B156 (1985) 1
 1985HA18 S.S. Hanna and J.W. Hugg, Hyperfine Interactions 21 (1985) 59
 1985HA1K J.H. Hamilton, P.G. Hansen and E.F. Zganjar, Rept. Prog. Phys. 48 (1985) 631
 1985HA1N B.G. Harvey, Nucl. Phys. A444 (1985) 498
 1985HA1T Hardy, Science 227 (1985) 993
 1985HE08 A.R. Heath and G.T. Garvey, Phys. Rev. C31 (1985) 2190
 1985HO05 J.J. Hogan, J. Asher and D.J. Parker, Phys. Rev. C31 (1985) 477
 1985HU04 M.S. Hussein, B.V. Carlson, O. Civitarese and A. Szanto De Toledo, Phys. Rev. Lett. 54 (1985) 2659
 1985HU08 Weizhi Huang, Zixing Wang, Hongqiu Song and Yanhuang Cai, Chin. J. Nucl. Phys. 7 (1985) 140
 1985HU09 Jimin Hu, Chunkai Zheng, Chin. J. Nucl. Phys. 7 (1985) 1
 1985IC01 A. Ichimura and M. Ichimura, Nucl. Phys. A432 (1985) 475

- 1985ISZU B.I. Islamov, V.N. Kadushkin, N.A. Kakurina, G.A. Radyuk, A.N. Sukmanov and I.I. Trinkin, in Leningrad, (1985) 323
- 1985IV1B Yu.B. Ivanov and L.M. Satarov, Nucl. Phys. A446 (1985) 727
- 1985IV1C Yu.B. Ivanov, I.N. Mishustin and L.M. Satarov, Nucl. Phys. A433 (1985) 713
- 1985JA17 R.A. Jarjis. Nucl. Instrum. Meth. Phys. Res. B12 (1985) 331
- 1985KA01 M. Kawai and Y. Iseri, Phys. Rev. C31 (1985) 400
- 1985KA1E Kadmenskii and Chuvilskii, in Leningrad (1985) 437, 440
- 1985KA1G Kadmenskii, Furman and Chuvilskii, in Leningrad (1985) 439
- 1985KAZQ S.G. Kadmenskii, S.D. Kurgalin and Yu.M. Chuvilskii, in Leningrad (1985) 438
- 1985KN04 O.M. Knyazkov, Yu.V. Maksimachev and Yu.V. Khrisanfov, Izv. Akad. Nauk SSSR Ser. Fiz. 49 (1985) 1020; Bull. Acad. Sci. USSR Phys. Ser. 49 (1985) 173
- 1985KO1J Koonin, Lecture Notes in Phys. 219 (1985) 129
- 1985KO38 Y. Kondo, B.A. Robson, R. Smith and H.H. Wolter, Phys. Lett. B162 (1985) 39
- 1985KO43 Y. Kondo, B.A. Robson and R. Smith, Nucl. Phys. A437 (1985) 117
- 1985KR1G Kroger and Perne, Can. J. Phys. 63 (1985) 366
- 1985LA03 M. Langevin, E. Quiniou, M. Bernas, J. Galin, J.C. Jacmart, F. Naulin, F. Pougheon, R. Anne, C. Detraz, D. Guerreau, D. Guillemaud-Mueller and A.C. Mueller, Phys. Lett. B150 (1985) 71
- 1985LAZZ C.M. Laymon and D.P. Balamuth, Bull. Amer. Phys. Soc. 30 (1985) 724, CE1
- 1985LE14 E.B. Levshin, P.V. Skorobogatov and V.K. Tartakovskiy, Ukr. Fiz. SSSR 30 (1985) 828
- 1985LE1H Levshin, Skorobogatov and Tartakovskii, in Leningrad (1985) 219
- 1985LE1L J.H. Lee and W.-H. Chang, New Phys. (Korea) 25 (1985) 381
- 1985LI19 H. Liu and L. Zamick, Phys. Rev. C32 (1985) 1754
- 1985LO1C Lobov, AIP Conf. Proc. 125 (1985) 900
- 1985MC03 M.A. McMahan, L.G. Moretto, M.L. Padgett, G.J. Wozniak, L.G. Sobotka and M.G. Mustafa, Phys. Rev. Lett. 54 (1985) 1995
- 1985ME1N Mezey et al., Acta Phys. Hung. 58 (1985) 39
- 1985MI10 R.E. Mischke, Nucl. Phys. A434 (1985) 505c
- 1985MI11 F. Michel and G. Reidemeister, J. Phys. (London) G11 (1985) 835
- 1985MI1D J. Missimer and L.M. Simons, Phys. Rept. 118 (1985) 179
- 1985MI23 T.A. Minelli, A. Pascolini and C. Villi, Nuovo Cim. A90 (1985) 185
- 1985MO08 M. Morjean, J.L. Charvet, J.L. Uzureau, Y. Patin, A. Peghaire, Y. Pranal, L. Sinopoli, A. Billerey, A. Chevarier, N. Chevarier et al., Nucl. Phys. A438 (1985) 547

- 1985MU10 H. Muther, A. Polls, P.K. Rath and A. Faessler, Nucl. Phys. A442 (1985) 68
- 1985MU14 T. Murakami, E. Ungricht, Y.-W. Lui, Y.Mihara, E. Takada and R.E. Tribble, Phys. Rev. C32 (1985) 1558
- 1985MU18 T. Murakami, N. Takahashi, Y.-W. Lui, E. Takada, D.M. Tanner, R.E. Tribble, E. Ungricht and K. Nagatani, Phys. Rev. C32 (1985) 2161
- 1985NI1C Ning, Gao and He, Phys. Energ. Fortis Phys. Nucl. 9 (1985) 101; Chin. Phys. 5 (1985) 899
- 1985NO1H Nomoto et al., Astrophys. J. 297 (1985) 531
- 1985OM01 K.M. Omar, V.M. Semenov, V.B. Subbotin and K.A. Gridnev, Izv. Akad. Nauk SSSR Ser. Fiz. 49 (1985) 170; Bull. Acad. Sci. USSR Phys. Ser. 49 (1985) 179
- 1985OS05 R. Ost, A. Gamp, S. Kox, C. Perrin, N. Longequeue and F. Merchez, Phys. Rev. C32 (1985) 1927
- 1985OS1C E. Oset and L.L. Salcedo, Nucl. Phys. A443 (1985) 704
- 1985OU01 S. Ouichaoui, H. Beaumevieuille, N. Bendjaballah, A.C. Chami, A. Dauchy, B. Chambon, D. Drain and C. Pastor, Nuovo Cim. A86 (1985) 170
- 1985PIZZ C. Pillai, L.W. Swenson, D.J. Vieira, G.W. Butler, J.M. Wouters, K. Vaziri, S.H. Rokni and L.P. Remsberg, Bull. Amer. Phys. Soc. 30 (1985) 1161, GX8
- 1985PO11 D.N. Poenaru, M. Ivascu, A. Sandulescu and W. Greiner, Phys. Rev. C32 (1985) 572
- 1985RA08 C. Rangacharyulu, E.J. Ansaldo, D. Stockhausen, D. Bender, S. Muller, A. Richter, N. Lo Iudice and F. Palumbo, Phys. Rev. C31 (1985) 1656
- 1985RH1A Rha et al., New Phys. (Korea) 25 (1985) 394
- 1985RO1G D.J. Rowe, Rept. Prog. Phys. 48 (1985) 1419
- 1985RO1N M. Roy-Stephan, Nucl. Phys. A446 (1985) 499
- 1985SA40 A. Sandulescu, Yu.S. Zamyatnin, I.A. Lebedev, B.F. Myasoedov, S.P. Tretyakova and D. Hasegan, Izv. Akad. Nauk. SSSR Ser. Fiz. 49 (1985) 2104; Bull. Acad. Sci. USSR Phys. Ser. 49 (1985) 20
- 1985SE21 R. Seki, Prog. Theor. Phys. 74 (1985) 511
- 1985SP06 K.-H. Speidel, Hyperfine Interactions 25 (1985) 477
- 1985ST19 B. Strohmaier, S.M. Grimes and S.D. Bloom, Phys. Rev. C32 (1985) 1397
- 1985ST1B R.G.Stokstad, Treatise on Heavy-Ion Sci. 3 (1985) 83
- 1985ST20 G.S.F. Stephans, D.G. Kovar, R.V.F. Janssens, G. Rosner, H. Ikezoe, B. Wilkins, D. Henderson, K.T. Lesko, J.J. Kolata, C.K. Gelbke et al., Phys. Lett. B161 (1985) 60
- 1985ST23 A.I. Steshenko, Nucl. Phys. A445 (1985) 462
- 1985TA1A Taam, Ann. Rev. Nucl. Part. Sci. 35 (1985) 1

- 1985TU01 L.E. Tubbs, J.R. Birkelund, J.R. Huizenga, D. Hilscher, U. Jahnke, H. Rossner and B. Gebauer, Phys. Rev. C32 (1985) 214
- 1985UH01 M. Uhrmacher, K. Pampus, F.J. Bergmeister, D. Purschke and K.P. Lieb, Nucl. Instrum. Meth. Phys. Res. B9 (1985) 234
- 1985UM01 A.S. Umar, M.R. Strayer, R.Y. Cusson, P.-G. Reinhard and D.A. Bromley, Phys. Rev. C32 (1985) 172
- 1985VE04 J.J.M. Verbaarschot and P.J. Brussaard, Z. Phys. A321 (1985) 125
- 1985VO1E Voinova-Eliseeva and Mitropolskii, in Leningrad (1985) 251
- 1985WA02 A.H. Wapstra and G. Audi, Nucl. Phys. A432 (1985) 1
- 1985WA08 H. Walliser, T. Fliessbach and Y.C. Tang, Nucl. Phys. A437 (1985) 367
- 1985WAZV T.R. Wang, W. Haeberli, S.W. Wissink, S.S. Hanna, Bull. Am. Phys. Soc. 30 (1985) 1248, AC8
- 1985WI17 B.H. Wildenthal, B.A. Brown and I. Sick, Phys. Rev. C32 (1985) 2185
- 1985XI1E Xie et al., Chin. Phys. 5 (1985) 893
- 1985YA1J Yalcin, Aksahin and Koru, Doga Bilim Dergisi Ser. A1 (Turkey) 9 (1985) 86
- 1986AB1H Abe, Kato and Uegaki, RIFP-683 (1986)
- 1986AI1A Aichelin and Stocker, Phys. Lett. B176 (1986) 14
- 1986AJ01 F. Ajzenberg-Selove, Nucl. Phys. A449 (1986) 1
- 1986AJ04 F. Ajzenberg-Selove, Nucl. Phys. A460 (1986) 1
- 1986ALZN S.C. Allcock, W.D.M. Rae, P.R. Keeling, S. Marsh, A.E. Smith, B.R. Fulton and D.W. Banes, in Harrogate (1986) 46; B20
- 1986ALZZ M.M. Alam and F.B. Malik, Bull. Amer. Phys. Soc. 31 (1986) 770, AG11
- 1986AN07 M.S. Antony, J. Britz and A. Pape, AT. Data Nucl. Data Tables 34 (1986) 279
- 1986AN10 M.S. Antony, J. Britz and J.B. Bueb, Nuovo Cim. A91 (1986) 283
- 1986AN40 M. Anikina, A. Golokhvastov, K. Iovchev, S. Khorozov, E. Kuznetsova, J. Lukstins, E. Okonov, T. Ostanovich, V. Toneev, G. Vardenga et al., Phys. Rev. C33 (1986) 895
- 1986AR04 K.P. Artemov, M.S. Golovkov, V.Z. Goldberg, I.P. Petrov, V.P. Rudakov, I.N. Serikov and V.A. Timofeev, Yad. Fiz. 43 (1986) 529; Sov. J. Nucl. Phys. 43 (1986) 335
- 1986BA16 D. Bachelier, J.L. Boyard, T. Hennino, J.C. Jourdain, M. Roy-Stephan, D. Contardo, J.Y. Grossiord, A. Guichard, J.R. Pizzi, P. Radvanyi et al., Phys. Lett. B172 (1986) 23
- 1986BA1E Baur and Bertulani, Phys. Rev. C34 (1986) 1654
- 1986BA22 F. Balestra, S. Bossolasco, M.P. Bussa, L. Busso, L. Ferrero, A. Grasso, D. Panzieri, G. Piragino, T. Tosello, G. Bendiscioli et al., Nucl. Phys. A452 (1986) 573
- 1986BA2U Baker, Bull. Am. Phys. Soc. 31 (1986) 1282

- 1986BA2W Balestra et al., AIP Conf. Proc. 150 (1986) 526
- 1986BA3E Bartke, Phys. Lett. B174 (1986) 32
- 1986BA3P Baktash, Bull. Amer. Phys. Soc. 31 (1986) 1224
- 1986BA3Q Balestra et al., Czech. J. Phys. 36 (1986) 340
- 1986BA69 D. Baye, Nucl. Phys. A460 (1986) 581
- 1986BA88 W. Bauhoff, At. Data Nucl. Data Tables 35 (1986) 429
- 1986BAYL M. Baba, M. Ono, N. Yabuta, T. Kikuti and N. Hirakawa, in Santa Fe 1985 (1986) 223
- 1986BE19 B. Bendyk, L. Jarczyk, B. Kamys, A. Strzalkowski and H. Witala, Phys. Rev. C34 (1986) 753
- 1986BE1D Betts and Rae, Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., No. 68, Vol. 2 (1986) 189
- 1986BE2M Bertulani and Bauer, Nucl. Phys. A458 (1986) 725
- 1986BL06 Y. Blumenfeld, Ph. Chomaz, N. Frascaria, J.P. Garron, J.C. Jacmart, J.C. Roynette, D. Ardouin and W. Mittig, Nucl. Phys. A455 (1986) 357
- 1986BL08 K. Blatt, K. Becker, B. Heck, H. Jansch, H. Leucker, D. Fick, R. Caplar, R. Butsch, D. Kramer, K.-H. Mobius et al., Phys. Rev. Lett. 57 (1986) 819
- 1986BL1K Blinov et al., P7-86-322, Dubna (1986) 40
- 1986BO1L Borzonov et al., P7-86-322, Dubna (1986) 66
- 1986BR1X Brown, Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., No. 68, Vol. 2 (1986) 119
- 1986BU1L Burtebaev, Duisebaev, Sadvoskii and Feofilov, in Kharkov (1986) 334
- 1986CA15 B. Castel and Y.K. Ho, Phys. Rev. C34 (1986) 408
- 1986CA19 W.N. Catford, D.M. Pringle, D.G. Lewis, A.E. Smith, E.F. Garman, I.F. Wright and J. Lukasiak, Nucl. Instrum. Meth. Phys. Res. A247 (1986) 367
- 1986CA24 B.O. Carragher, J. Carter, R.G. Clarkson, V. Hnizdo and J.P.F. Sellschop, Nucl. Phys. A460 (1986) 341
- 1986CA27 M. Carchidi, B.H. Wildenthal and B.A. Brown, Phys. Rev. C34 (1986) 2280
- 1986CEZW R. Ceuleneer, F. Michel, G. Reidemeister and S. Ohkubo, in Harrogate (1986) 239
- 1986CH28 L. Chaves and A. Poves, Phys. Rev. C34 (1986) 1137
- 1986CH2G Chbihi et al., J. Phys. (Paris) 47 (1986) C4-87
- 1986CHYY M.A. Chaudhri, in Santa Fe 1985 (1986) 819
- 1986CO15 S.G. Cooper, J. Phys. (London) G12 (1986) 371
- 1986COZZ D.F. Collinson, K.-X. Wang, D.H. Feng and C.-L. Wu, Bull. Amer. Phys. Soc. 31 (1986) 816

- 1986CU02 B. Cujec, B. Dasmahapatra, Q. Haider, F. Lahlou and R.A. Dayras, Nucl. Phys. A453 (1986) 505
- 1986DE27 P. Descouvemont and D. Baye, Nucl. Phys. A459 (1986) 374
- 1986DE31 P. Descouvemont, Phys. Lett. B181 (1986) 199
- 1986FA12 A.R. Farhan, B.J. Stoyanov, A. Nagl, H. Uberall and M. de Llano, Phys. Rev. C34 (1986) 2134
- 1986FI15 Filippone, Ann. Rev. Nucl. Part. Sci. 36 (1986) 717
- 1986FO1D Fowler, Bull. Amer. Phys. Soc. 31 (1986) 1267
- 1986GA1F M. Gazdzicki, K. Iovchev, E. Kladnitskaya, E. Okonov and E.Z. Skrzypczak, Z. Phys. C31 (1986) 549
- 1986GA1N Gadioli and Hodgson, Rept. Prof. Phys. 49 (1986) 951
- 1986GA1P Gaarde, Proc. Int. Nucl. Phys. Conf., Harrogate, U.K., No. 68, Vol. 2 (1986) 173
- 1986GI03 R. Gissler, D. Provoost, F. Grummer and K. Goeke, Phys. Lett. B166 (1986) 385
- 1986GI10 A. Gillibert, L. Bianchi, A. Cunsolo, B. Fernandez, A. Foti, J. Gastebois, Ch. Gregoire, W. Mittig, A. Peghaire, Y. Schutz et al., Phys. Lett. B176 (1986) 317
- 1986GR1A C. Gregoire and B. Tamain, Ann. Phys. (France) 11 (1986) 323
- 1986GR1B Grashin et al., Izv. Akad. Nauk SSSR Ser. Fiz. 50 (1986) 944
- 1986GR1Q Gridnev et al., in Kharkov (1986) 456
- 1986GR1R Gridnev, Darvich, Subbotin and Fadeev, in Kharkov (1986) 486
- 1986GRZK K.A. Griffioen, E.A. Bakkum, C.P.M. van Engelen, R.J. Meijer and R. Kamermans, in Harrogate (1986) 219; C32
- 1986GU1D D. Guerreau, J. Phys. (Paris) 47 (1986) C4-207
- 1986HA13 Q. Haider and F.B. Malik, J. Phys. (London) G12 (1986) 537
- 1986HA1B B.G. Harvey, J. Phys. (Paris) 47 (1986) C4-29
- 1986HE13 K. Heyde, J. Jolie, J. Moreau, J. Ryckebusch, M. Waroquier and J.L. Wood, Phys. Lett. B176 (1986) 255
- 1986HE1A He et al., in Harrogate (1986) C51
- 1986HO33 H. Horiuchi, T. Wada, K. Yabana, Prog. Theor. Phys. (Kyoto) 76 (1986) 837
- 1986HO35 J. Honkanen, J. Aysto and K. Eskola, Phys. Scr. 34 (1986) 608
- 1986HU1G Huang, Wang, Song and Cai, Chin. Phys. 6 (1986) 132
- 1986IK03 H. Ikezoe, N. Shikazono, Y. Tomita, K. Ideno, Y. Sugiyama, E. Takekoshi, T. Tachikawa and T. Nomura, Nucl. Phys. A456 (1986) 298
- 1986IR01 M. Iriondo, D. Jerrestam and R.J. Liotta, Nucl. Phys. A454 (1986) 252
- 1986KA17 H. Kazama, K. Kato and H. Tanaka, Prog. Theor. Phys. 75 (1986) 571

- 1986KA1U Kamanin et al., P7-86-322, Dubna (1986) 55
- 1986KA2G Kachan, in Kharkov (1986) 171
- 1986KA36 Y. Kadota, K. Ogino, K. Obori, Y. Taniguchi, T. Tanabe, M. Yasue and J. Schimizu, Nucl. Phys. A458 (1986) 523
- 1986KAZP H. Kazama, K. Kato and H. Tanaka, in Harrogate (1986) 86
- 1986KE15 T.J. Kennett, W.V. Prestwich and J.S. Tsai, Nucl. Instrum. Meth. Phys. Res. A247 (1986) 420
- 1986KH1J Khairozzaman, Astron. Nachr. 307 (1986) 235
- 1986KN1C N. Kniest, E. Huttel, E. Pfaff, G. Reiter, S. Tharraketta, A. Hofmann, G. Clausnitzer, J. Phys. Soc. Jpn. Suppl. 55 (1986) 1034
- 1986KO1C Kondo et al., 11th Ains Nucl. Phys. Conf., Melbourne (1986) 93
- 1986KR16 B. Krusche and K.P. Lieb, Phys. Rev. C34 (1986) 2103
- 1986LA07 K. Langanke, M. Wiescher, W.A. Fowler and J. Gorres, Astrophys. J. 301 (1986) 629
- 1986LAZZ C.M. Laymon and D.P. Balamuth, Bull. Amer. Phys. Soc. 31 (1986) 771, AH5
- 1986LE06 H. Leeb, Czech. J. Phys. 36 (1986) 253
- 1986LE16 S.-J. Lee, J. Fink, A.B. Balantekin, M.R. Strayer, A.S. Umar, P.-G. Reinhard, J.A. Maruhn and W. Greiner, Phys. Rev. Lett. 57 (1986) 2916; Erratum Phys. Rev. Lett. 59 (1987) 1171
- 1986LE22 H. Leeb and E.W. Schmid, Few-Body Systems 1 (1986) 203
- 1986LE23 J.A. Leavitt, P. Stoss, D.B. Cooper J., L. Seerveld, L.C. McIntyre J., R.E. Davis, S. Gutierrez and T.M. Reith, Nucl. Instrum. Meth. Phys. Res. B15 (1986) 296
- 1986LO1E Lobov, J. Phys. Soc. Jpn. Suppl. 55 (1986) 1002
- 1986MA19 J.F. Mateja, A.D. Frawley, L.C. Dennis and K. Sartor, Phys. Rev. C33 (1986) 1649
- 1986MA1E Matteucci, Astrophys. J. 305 (1986) L81
- 1986MA35 C. Mahaux, H. Ngo and G.R. Satchler, Nucl. Phys. A456 (1986) 134
- 1986MA48 D.M. Manley, B.L. Berman, W. Bertozzi, J.M. Finn, F.W. Hersman, C.E. Hyde-Wright, M.V. Hynes, J.J. Kelly, M.A. Kovash, S. Kowalski et al., Phys. Rev. C34 (1986) 1214
- 1986MAZC S. Marsh and W.D.M. Rae, in Harrogate (1986) 285; C98
- 1986ME06 M.C. Mermaz, T. Suomijarvi, R. Lucas, B. Berthier, J. Matuszek, J.P. Coffin, G. Guillaume, B. Heusch, F. Jundt and F. Rami, Nucl. Phys. A456 (1986) 186
- 1986OE01 W. Oelert, Nucl. Phys. A449 (1986) 395
- 1986OU01 S. Ouichaoui, H. Beaumevielle, N. Bendjaballah and A. Genoux-Lubain, Nuovo Cim. A94 (1986) 133

- 1986OUZZ S. Ouichaoui, H. Beaumevieille, N. Bendjaballah and G.J. Costa, in Harrogate (1986) 105
- 1986PA10 C.T. Papadopoulos, R. Vlastou, E.N. Gazis, P.A. Assimakopoulos, C.A. Kalfas, S. Kossionides and A.C. Xenoulis, Phys. Rev. C34 (1986) 196
- 1986PAZJ G. Palla, in Harrogate (1986) 407
- 1986PI09 Pillai et al., in Santa Fe (1985) 1383
- 1986PL02 R. Planeta, P. Belery, J. Brzychczyk, P. Cohilis, Y. El Masri, Gh. Gregoire, K. Gro-towski, Z. Majka, S. Micek, M. Szczodrak et al., Phys. Rev. C34 (1986) 512
- 1986PO06 D.N. Poenaru, W. Greiner, K. Depta, M. Ivascu, D. Mazilu and A. Sandulescu, At. Data Nucl. Data Tables 34 (1986) 423
- 1986PO15 D.N. Poenaru, W. Greiner, M. Ivascu, D. Mazilu and I.H. Plonski, Z. Phys. A325 (1986) 435
- 1986RA15 Md.A. Rahman and H.M. Sen Gupta, Nuovo Cim. A93 (1986) 236
- 1986RO03 R. Rockmore and B. Saghai, Phys. Rev. C33 (1986) 576
- 1986SA02 M. Sato, S.A. Coon, H.J. Pirner, J.P. Vary, Phys. Rev. C33 (1986) 1062
- 1986SA2N Saini et al., in Harrogate (1986) C35
- 1986SA30 H. Sato and Y. Okuhara, Phys. Rev. C34 (1986) 2171
- 1986SA40 G.S. Samosvat, Fiz. Elem. Chastits At.Yadra 17 (1986) 713; Sov. J. Part. Nucl. 17 (1986) 313
- 1986SC28 C.J.S. Scholz, L. Ricken and E. Kuhlmann, Z. Phys. A325 (1986) 203
- 1986SC29 H.R. Schmidt, S.B. Gazes, Y. Chan, R. Kamermans and R.G. Stokstad, Phys. Lett. B180 (1986) 9
- 1986SE1B N. Seichert, W. Assmann, H. Clement, G. Graw, C. Hategan, H. Kader, F. Merz and P. Schiemenz, J. Phys. Soc. Jpn. Suppl. 55 (1986) 646
- 1986SH07 A.C. Shotter, J. Phys. (London) G12 (1986) L93
- 1986SH16 N.G. Shevchenko, Izv. Akad. Nauk SSSR Ser. Fiz. 50 (1986) 121; Bull. Acad. Sci. USSR Phys. Ser. 50 (1986) 117
- 1986SH1F W.Q. Shen, S.Z. Yin, Z.Y. Guo, W.L. Zhan, Y.T. Zhu, G.M. Jin, W.M. Qiao, E.C. Wu and C.L. Jiang, Chin. Phys. 6 (1986) 80
- 1986SH30 T. Shimoda, K. Katori, T. Fukuda, H. Ogata, S. Shimoura, M. Tanaka and E. Takada, J. Phys. Soc. Jpn. 55 (1986) 3021
- 1986SI1D Simpson and Earwaker, Nucl. Instrum. Meth. Phys. Res. B15 (1986) 502
- 1986SI1K Simonius, J. Phys. Soc. Jpn. Suppl. 55 (1986) 523
- 1986SI1L Sie, McKenzie, Smith and Ryan, Nucl. Instrum. Meth. Phys. Res. B15 (1986) 525
- 1986SN1B Snover, Ann. Rev. Nucl. Part. Sci. 36 (1986) 545

- 1986SO10 L.G. Sobotka, D.G. Sarantites, H. Puchta, F.A. Dilmanian, M. Jaaskelainen, M.L. Halbert, J.H. Barker, J.R. Beene, R.L. Ferguson, D.C. Hensley et al., Phys. Rev. C34 (1986) 917
- 1986ST1J Stock, Phy. Rept. 135 (1986) 259
- 1986STZT A.P. Stamp, in Harrogate (1986) 128; B102
- 1986SU01 Y.Suzuki, Nucl. Phys. A448 (1986) 395
- 1986SU06 Y. Suzuki and K.T. Hecht, Nucl. Phys. A455 (1986) 315
- 1986SZ02 I.M. Szoghy, Q. Haider and R. Ouellet, Nucl. Instrum. Meth. Phys. Res. A242 (1986) 277
- 1986TA1A Tanimura, Phys. Lett. B181 (1986) 13
- 1986TA1W Tartakovskii and Isupov, in Kharkov (1986) 168
- 1986TA1X Tarasov et al., in Kharkov (1986) 207
- 1986TH1E Thielemann, Nomoto and Yokoi, Astron. Astrophys. 158 (1986) 17
- 1986TK1C Tkaczyk et al., Z. Phys. C33 (1986) 33
- 1986TR08 W. Trolenberg, F. Hagelberg, H.J. Simonis, P.N. Tandon, K.-H. Speidel, M. Knopp and J. Gerber, Nucl. Phys. A458 (1986) 95
- 1986TR1B Tretiakova et al., P7-86-322, Dubna (1986) 19
- 1986TR1C Truran and Livio, Astrophys. J. 308 (1986) 721
- 1986TU1B Tungate et al., J. Phys. Soc. Jpn. Suppl. 55 (1986) 1116
- 1986VA04 J. Vandermeulen, Phys. Rev. C33 (1986) 1101
- 1986VA10 C.P.M. van Engelen, E.A. Bakkum, R.J. Meijer and R. Kamermans, Phys. Rev. C33 (1986) 1851
- 1986VA12 M.G. Vassanji and D.J. Rowe, Nucl. Phys. A454 (1986) 288
- 1986VA18 C.P.M. van Engelen, E.A. Bakkum, R.J. Meijer and R. Kamermans, Z. Phys. A324 (1986) 121
- 1986VA23 C.P.M. van Engelen, E.A. Bakkum, R.J. Meijer and R. Kamermans, Nucl. Phys. A457 (1986) 375
- 1986VI09 D.J. Vieira, J.M. Wouters, K. Vaziri, R.H. Kraus, Jr., H. Wollnik, G.W. Butler, F.K. Wohn and A.H. Wapstra, Phys. Rev. Lett. 57 (1986) 3253
- 1986VO05 T. von Egidy, A.N. Behkami and H.H. Schmidt, Nucl. Phys. A454 (1986) 109
- 1986VO07 N.A. Voinova-Elseeva and I.A. Mitropolsky, Izv. Akad. Nauk SSSR, Ser. Fiz. 50 (1986) 14; Bull. Acad. Sci. USSR, Phys. Ser. 50 (1986) 12
- 1986WA1R Wang et al., Bull. Amer. Phys. Soc. 31 (1986) 816
- 1986WE1C Westfall, Nucl. Phys. A447 (1986) 591c

- 1986WI1P D. Wilkinson, J. Phys. Soc. Jpn. Suppl. 55 (1986) 347
- 1986WU01 A.H. Wuosmaa, S. Saini, P.H. Kutt, S.F. Pate and R.W. Zurmuhle, Phys. Lett. B172 (1986) 297
- 1987AB03 H. Abele, H.J. Hauser, A. Korber, W. Leitner, R. Neu, H. Plappert, T. Rohwer, G. Staudt, M. Strasser, S. Welte et al., Z. Phys. A326 (1987) 373
- 1987AL06 D.E. Alburger, G. Wang and E.K. Warburton, Phys. Rev. C35 (1987) 1479
- 1987BA01 D. Bandyopadhyay, S.R. Samaddar, K. Krishan and J.N. De, Nucl. Phys. A462 (1987) 587
- 1987BA1V Bauer, Bertsch and Das Gupta, Phys. Rev. Lett. 58 (1987) 863
- 1987BO1B Bond and Luck, Astrophys. J. 312 (1987) 203
- 1987CA05 Z.-J. Cao, R.D. Bent, H. Nann, T.E. Ward, Phys. Rev. C35 (1987) 625
- 1987CO07 J. Cook, Nucl. Phys. A465 (1987) 207
- 1987CU1A Cummings and Stone, Bull. Amer. Phys. Soc. 32 (1987) 1066
- 1987DE43 H. De Vries, C.W. De Jager and C. De Vries, At. Data Nucl. Data Tables 36 (1987) 495
- 1987FA09 M. Fatyga, K. Kwiatkowski, V.E. Viola, W.G. Wilson, M.B. Tsang, J. Pochodzalla, W.G. Lynch, C.K. Gelbke, D.J. Fields, C.B. Chitwood et al., Phys. Rev. Lett. 58 (1987) 2527
- 1987FI01 L.K. Fifield, D.M. Pringle and W.J. Vermeer, Nucl. Phys. A463 (1987) 644
- 1987GI01 K.L. Giovanetti, G. de Chambrier, P.F.A. Goudsmit, H.J. Leisi, B. Jeckelmann, Th. Karapiperis, Phys. Lett. B186 (1987) 9
- 1987GI1E Gillibert et al., Ganil P.87-05 (1987)
- 1987HA08 P. Halse and Z.Y. Pan, Phys. Rev. C35 (1987) 774; Erratum Phys. Rev. C35 (1987) 1964
- 1987HA24 K.H. Hahn, K.H. Chang, T.R. Donoghue and B.W. Filippone, Phys. Rev. C36 (1987) 892
- 1987HO1B Horiuchi, Wada and Yabana, Kuns 849; Workshop on Relations between Struct. and Reactions in Nucl. Phys., Philadelphia (1986)
- 1987IA1B Iachello and Talmi, Rev. Mod. Phys. 59 (1987) 339
- 1987JA06 B.V. Jacak, G.D. Westfall, G.M. Crawley, D. Fox, C.K. Gelbke, L.H. Harwood, B.E. Hasselquist, W.G. Lynch, D.K. Scott, H. Stocker et al., Phys. Rev. C35 (1987) 1751
- 1987KA18 K. Kato, H. Kazama and H. Tanaka, Prog. Theor. Phys. 77 (1987) 185
- 1987KO12 S. Kox, A. Gamp, C. Perrin, J. Arvieux, R. Bertholet, J.F. Bruandet, M. Buenerd, R. Cherkaoui, A.J. Cole, Y. El-Masri et al., Phys. Rev. C35 (1987) 1678

- 1987KO15 T. Kozik, J. Buschmann, K. Grotowski, H.J. Gils, N. Heide, J. Kiener, H. Klewe-Nebenius, H. Rebel, S. Zagromski, A.J. Cole et al., *Z. Phys.* A326 (1987) 421
- 1987LA05 M. Lattuada, D. Vinciguerra, C.M. Sutura, G. Inghima and M. Sandoli, *Phys. Rev.* C35 (1987) 818
- 1987LA14 L.O. Lamm, C.P. Browne, J. Gorres, M. Wiescher and A.A. Rollefson, *Z. Phys.* A327 (1987) 239
- 1987MA1R Marti and Craig, *Nature* 325 (1987) 335
- 1987ME1B Mewaldt and Stone, *Bull. Amer. Phys. Soc.* 32 (1987) 1037
- 1987MI07 F. Michel, G. Reidemeister and S. Ohkubo, *Phys. Rev.* C35 (1987) 1961
- 1987MU1B Murphy, Dermer and Ramaty, *Astrophys. J. Suppl.* 63 (1987) 721
- 1987MU1J Mueller, GANIL P.87-06; XXV Int. Winter Meeting on Nucl. Phys., Bormio, Italy (1987)
- 1987NA1D Nagata et al., *Nucl. Instrum. Meth. Phys. Res.* B18 (1987) 515
- 1987NI04 W. Nitsche, G.J. Wagner, K.T. Knopfle, P. Grabmayr and Y. Kawazoe, *Z. Phys.* A326 (1987) 435
- 1987NU01 J. Nurzynski, T. Kihm, K.T. Knopfle, G. Mairle and H. Clement, *Nucl. Phys.* A465 (1987) 365
- 1987PR01 D. Provoost, F. Grummer and K. Goeke, *Ann. Phys.* 174 (1987) 202
- 1987PR1A Prapkos, Arnould and Arcoragi, *Astrophys. J.* 315 (1987) 209
- 1987RA01 S. Raman, C.H. Malarkey, W.T. Milner, C.W. Nestor, Jr. and P.H. Stelson, *At. Data Nucl. Data Tables* 36 (1987) 1
- 1987RA02 W.D.M. Rae, P.R. Keeling and S.C. Allcock, *Phys. Lett.* B184 (1987) 133
- 1987RI03 J. Richert and P. Wagner, *Nucl. Phys.* A466 (1987) 132
- 1987RO10 G. Royer, Y. Raffray, A. Oubahadou and B. Remaud, *Nucl. Phys.* A466 (1987) 139
- 1987RO25 C. Rolfs, H.P. Trautvetter and W.S. Rodney, *Rept. Prog. Phys.* 50 (1987) 233
- 1987ROZZ R.D. Rosa, W.W. Daehnick, S.K. Saha and P.C. Li, *Bull. Amer. Phys. Soc.* 32 (1987) 1062, EG10
- 1987SA15 H. Sagawa and H. Toki, *J. Phys. (London)* G13 (1987) 453
- 1987SC03 B. Schurmann and W. Zwermann, *Phys. Lett.* B183 (1987) 31
- 1987SC1D Schurmann, Zwermann and Malfliet, *Phys. Rept.* 147 (1987) 1
- 1987SI06 K. Siwek-Wilczynska, J. Wilczynski, C.R. Albiston, Y. Chan, E. Chavez, S.B. Gazes, H.R. Schmidt and R.G. Stokstad, *Phys. Rev.* C35 (1987) 1316
- 1987SU03 T. Sugimitsu, H. Inoue, H. Fujita, N. Kato, K. Kimura, T. Tachikawa, K. Anai, Y. Ikeda and Y. Nakajima, *Nucl. Phys.* A464 (1987) 415

- 1987SU06 T. Suzuki, D.F. Measday and J.P. Roalsvig, Phys. Rev. C35 (1987) 2212
- 1987TE01 A. Tellez-Arenas, R.J. Lombard and J.P. Maillet, J. Phys. (London) G13 (1987) 311
- 1987TR01 H.-J. Trost, P. Lezoch and U. Strohmusch, Nucl. Phys. A462 (1987) 333
- 1987VA12 M.G. Vassanji and D.J. Rowe, Nucl. Phys. A466 (1987) 227
- 1987WA1B Wada and Horiuchi, Phys. Rev. Lett. 58 (1987) 2190
- 1987WI03 A. Willis, M. Morlet, N. Marty, C. Djalali, G.M. Crawley, A. Galonsky, V. Rotberg and B.A. Brown, Nucl. Phys. A464 (1987) 315
- 1987YA03 B.-J. Yang and Z. Hwang, Phys. Rev. C35 (1987) 851
- 1987ZH1B Zheng, Ragnarsson and Bengtsson, Proc. Beijing Int. Symp. on Phys. at Tandem, 1986 (1987) 538
- 1988AJ01 F. Ajzenberg-Selove, Nucl. Phys. A490 (1988) 1

